

Hyperbolic contractivity and the Hilbert metric on probability measures

Samuel N. Cohen* Eliana Fausti†

September 3, 2023

Abstract

This paper gives a self-contained introduction to the Hilbert projective metric \mathcal{H} and its fundamental properties, with a particular focus on the space of probability measures. We start by defining the Hilbert pseudo-metric on convex cones, focusing mainly on dual formulations of \mathcal{H} . We show that linear operators on convex cones contract in the distance given by the hyperbolic tangent of \mathcal{H} , which in particular implies Birkhoff’s classical contraction result for \mathcal{H} . Turning to spaces of probability measures, where \mathcal{H} is a metric, we analyse the dual formulation of \mathcal{H} in the general setting, and explore the geometry of the probability simplex under \mathcal{H} in the special case of discrete probability measures. Throughout, we compare \mathcal{H} with other distances between probability measures. In particular, we show how convergence in \mathcal{H} implies convergence in total variation, p -Wasserstein distance, and any f -divergence. Finally, we derive a novel sharp bound for the total variation between two probability measures in terms of their Hilbert distance.

MSC: 47A35, 15B51, 60B10, 52B12, 62B11, 94A15

Keywords: Hilbert projective metric, probability simplex, contraction of positive linear operators.

1 Introduction

The main goal of this paper is to provide a clear, self-contained guide to the Hilbert projective (pseudo-)metric and its merits in applied probability theory. In particular, as we discuss in more detail below, this metric may serve as a powerful tool for understanding whether linear operators on probability measures contract, in what sense, and at what rate. This, of course, is part of a much bigger story concerning the contraction of linear operators on positive cones—of which the space of (non-negative) measures is a prominent example.

In [7], Garrett Birkhoff¹ proved that positive linear operators on a positive convex cone contract in the Hilbert projective (pseudo-)metric (introduced by Hilbert in [20]). Starting from this result, he easily derives a proof of the famous Perron theorem, by essentially

*Mathematical Institute, University of Oxford and Alan Turing Institute, cohens@maths.ox.ac.uk.

†Mathematical Institute, University of Oxford, fausti@maths.ox.ac.uk.

¹The son of George D. Birkhoff, who is better known in probabilistic circles for his proof of the ergodic theorem.

reducing it to a special case of the Banach fixed point theorem. Similarly, he also immediately proves several of its generalizations: first, the extension of the Perron theorem to non-negative matrices due to Frobenius, then the extensions to infinite dimensional function spaces and positive integral and compact operators, originally due to Jentzsch [23] and Krein and Rutman [26]. In the same year as Birkhoff, Samelson [35] also published a proof of the Perron–Frobenius theorem using projective geometry, and similarly a few years later Hopf [21] presented a contraction result for positive linear integral operators and an alternative proof of Jentzsch’s theorem (apparently without being aware of Birkhoff’s previous results). The combination of the Hilbert projective metric with the contraction mapping theorem has inspired a vast amount of further work on the Perron–Frobenius theorem and its various extensions, see Lemmens and Nussbaum [30] and references therein for a detailed overview of the topic. We highlight the early works of Thompson [38], Bushell [11], and Kohlberg and Pratt [25], as well as the recent works by Rugh [34] and Dubois [17] on operators on complex cones which have strongly influenced the presentation in this paper.

While the impact on Perron–Frobenius theory is certainly the most significant consequence of Birkhoff’s work, we became interested in the Hilbert metric and Birkhoff’s contraction result due to a different (although related) application: the study of the ergodicity of non-negative linear operators. Work in this direction was presented by Birkhoff himself in [8, Chapter XVI, Sec.7-8], Seneta and Sheridan [36, 37] and Le Gland and Mevel [27]. Clearly, the ergodic theory of Markov processes with linear transition kernels can be reduced to a particular case of this broader topic. Nevertheless, the Hilbert metric is not strictly necessary for developing the standard ergodic theory of Markov processes. For example, in [36, Chapter 3] Seneta uses the Hilbert metric to study the ergodicity of inhomogenous products of non-negative matrices; on the other hand, in the following chapter [36, Chapter 4], which treats specifically the ergodicity of discrete Markov chains, the Hilbert metric is not used, since the analysis simplifies by virtue of the generators being stochastic matrices.

Thus, one of the main avenues of application for the Hilbert projective metric in probability theory is, in fact, relatively unknown, which might in part explain why, despite being quite powerful tools, neither the Hilbert metric nor Birkhoff’s contraction results seem to have found widespread use in the probability community². There are of course a few notable exceptions: Birkhoff’s contraction theorem was employed, for example, in an elegant proof of the linear convergence of the Sinkhorn algorithm by Franklin and Lorenz [19]. It also proved fundamental in the qualitative and quantitative analysis of the stability of hidden Markov processes, where the use of the Hilbert metric was introduced by Atar and Zeitouni [3, 4], and, following their work, became a well-established approach to the problem, see e.g. [5, 14, 28, 29]. More recently, the Hilbert metric has found computational applications in entropic interpolation [12] and nonlinear embeddings [32].

In writing this paper, we had in mind applied probabilists and statisticians who find themselves curious, as we were, about the merits and limitations of working with the Hilbert projective metric in a probabilistic context. Consequently, our main focus is the study of the space of probability measures when equipped with the Hilbert projective metric.

Nevertheless, we shall start in greater generality, by defining the Hilbert projective (pseudo-)metric on a proper convex cone in a locally convex topological vector space. Our main contribution in Section 2 is the introduction of a new (pseudo-)metric, the hyperbolic tangent

²To illustrate this point, no mention is made of this approach in any of the books by Ethier and Kurtz [18], Meyn and Tweedie [31], Brémaud [10], Kallenberg [24], or Jacod and Shiryaev [22].

of the Hilbert metric (which we call the \mathcal{T} -distance), under which we prove linear operators also contract (see Definition 2.5 and Theorem 2.6). The advantage of using the \mathcal{T} -distance compared to the Hilbert projective metric is that \mathcal{T} stays bounded, while the Hilbert metric (easily) diverges to infinity. As far as we are aware, the formulation of this contraction result in Theorem 2.6 has not been given before (but we note that our proof is inspired by Dubois' proof of [17, Thm. 2.3], so we do not wish to claim full credit for the result).

When introducing the Hilbert projective metric, we especially insist on its definition through duality, which we first came across in [34]. This 'dual' definition, in particular through a predual space, is natural in the context of probability measures, where distances are often defined by considering measures as integrators dual to particular classes of functions. In Section 3 we provide a careful analysis of the Hilbert metric and the topology it induces on the space of probability measures using duality, and we show that convergence of measures in the Hilbert metric (or in the \mathcal{T} -distance) is stronger than convergence in total variation (as was already shown in [3, Lemma 1]), convergence in p -Wasserstein distance and convergence in any f -divergence.

In Section 4 we study the geometry of the probability simplex under the Hilbert metric. Using our dual approach, we give an easy derivation of the explicit formula for the contraction rate of a linear operator (Proposition 4.1), which can also be extended to infinite settings (Proposition 4.2). As the probability simplex is finite dimensional, it has a natural manifold structure, however the Hilbert metric does not induce a hyperbolic (in the sense of Gromov) geometry on the simplex (since its boundary is not differentiable, see [6]), nor a Riemannian structure. The Hilbert geometry is far more curious: we find an explicit characterizations of Hilbert balls as 'hexagonal' convex polytopes, extending to the n -dimensional case work by Phadke [33] and de la Harpe [16]. Finally, in the last section of the paper we use these geometric observations to prove a sharp bound for the total variation distance between two probability measures in terms of their \mathcal{T} -distance (see Theorem 5.1 and Cor. 5.2.1).

2 The Hilbert projective metric: definitions and contractivity

Let us start with the definition of the Hilbert projective distance, in the sense of Birkhoff [7,8], on a cone in a (real) locally convex topological vector space (LCS). Note that Birkhoff works more specifically with cones in real Banach spaces (lattices). However, as we will see shortly, the definition of the metric does not require the space to be Banach, so for the sake of generality we work with an LCS.

Let X be an LCS. Let $C \subset X$ be a proper closed convex cone, meaning that C is closed and satisfies

$$C + C \subseteq C, \quad \mathbb{R}_+ C = C, \quad C \cap -C = \{0\}.$$

Following [34, Sec. 4] (or extrapolating directly from [7] or [8, Chapter XVI]), we give Birkhoff's definition of the Hilbert projective distance.

Definition 2.1 (Hilbert projective pseudo-metric). For $x, y \in C \setminus \{0\}$, where C is a proper closed convex cone in a real LCS, let $\beta(x, y) \in (0, \infty]$ be given by

$$\beta(x, y) = \inf\{r > 0 : rx - y \in C\} = \sup\{r > 0 : rx - y \notin C\}.$$

Then the Hilbert projective distance is defined by

$$\mathcal{H}(x, y) = \log(\beta(x, y)\beta(y, x)) \in [0, \infty], \quad \forall x, y \in C \setminus \{0\}. \quad (1)$$

It is worth spending a few moments to properly understand these definitions. Since C is closed, and $-y \notin C$ for all $y \in C \setminus \{0\}$, we have $\beta(x, y) > 0$. It is then straightforward to see that $\beta(x, y)\beta(y, x) \geq 1$, by noting that

$$\begin{aligned} \frac{1}{\beta(x, y)} &= \inf\{1/r > 0 : rx - y \notin C\} = \inf\{r > 0 : x - ry \notin C\} = \inf\{r > 0 : ry - x \notin -C\} \\ &\leq \inf\{r > 0 : ry - x \in C\} = \beta(y, x), \end{aligned} \quad (2)$$

and $\beta(x, y) = \frac{1}{\beta(y, x)}$ if and only if $y = cx$ for some $c \in \mathbb{R}_+$ (in this case, x and y are said to be *collinear*). Hence (1) is well-defined with $\mathcal{H}(x, y) = 0$ if and only if x and y are collinear. Symmetry of \mathcal{H} is clear from the definition, and one can verify that the triangle inequality is also satisfied (see Remark 2): then, \mathcal{H} is a pseudo-metric for C (see also [8, Chapter XVI]).

Example 2.2. The finite non-negative measures on $[0, 1]$ form a proper convex (non-negative) cone in the space of the signed measures on $[0, 1]$. Since the signed measures on $[0, 1]$ equipped with the total variation distance form a Banach space (and therefore an LCS), the definition above applies. The Hilbert pseudo-metric can then be restricted to the probability measures on $[0, 1]$, to give a true metric. We will explore in detail the Hilbert distance on the space of measures in Section 3.

Remark 1. The only role of the specific choice of LCS topology on X , in defining the \mathcal{H} -metric, is to have the notion of C being closed in X . Two different topologies on X for which C is closed in X will both give rise to the same Hilbert projective metric. In this sense, the Hilbert metric on C is independent of the topology on the ambient space X .

By closure of C in X , if $\beta(x, y) < \infty$, we must have that $\beta(x, y)x - y \in \partial C$, where ∂C is the boundary of C . If $x \in \partial C$ and $y \in \overset{\circ}{C}$, where $\overset{\circ}{C}$ denotes the interior of C , then $\beta(x, y) = \infty$. However, if both $x, y \in \partial C$, then $\beta(x, y)$ might be finite. Using Birkhoff's choice of terminology [8, Chapter XVI], $x, y \in C$ are *comparable* if $\beta(x, y)$ and $\beta(y, x)$ are both finite. We observe that two boundary elements $x, y \in \partial C$ might still be comparable.

Especially when X is infinite dimensional, it can be useful to understand the Hilbert projective distance through duality (see [17, 34], where this is exploited in the analysis of complex cones). Let X^* be the (topological) dual of X , and let $\langle \cdot, \cdot \rangle$ be the natural bilinear form $X^* \times X \rightarrow \mathbb{R}$. One can define the dual cone C^* as

$$C^* = \{f \in X^* : f|_C \geq 0\}. \quad (3)$$

Proposition 2.3. *An equivalent definition of the Hilbert pseudo-metric is given by*

$$\mathcal{H}(x, y) = \sup_{\substack{f, g \in C^* \\ \langle f, x \rangle, \langle g, y \rangle \neq 0}} \left\{ \log \frac{\langle f, y \rangle \langle g, x \rangle}{\langle f, x \rangle \langle g, y \rangle} \right\}. \quad (4)$$

Proof. Let $\tilde{C} := \{x \in X : \langle f, x \rangle \geq 0, \forall f \in C^*\}$. One can confirm that \tilde{C} is a proper closed convex cone. We clearly have that $C \subseteq \tilde{C}$. Now consider $x \notin C$. Since C is convex and closed, by the Geometric Hahn–Banach theorem (see e.g. [15, Thm. IV.3.9 & Cor. IV.3.10]) there exists a continuous linear functional $g \in X^*$ and an $\alpha \in \mathbb{R}$ such that $\langle g, x \rangle < \alpha$ and $\langle g, y \rangle \geq \alpha$ for all $y \in C$. Since the image of the cone C under the functional g can only be one of \mathbb{R}_+ , \mathbb{R}_- , \mathbb{R} or $\{0\}$, we must have $\alpha = 0$. Hence $g \in C^*$ but $\langle g, x \rangle < 0$, which implies that $x \notin \tilde{C}$. Therefore $\tilde{C} \subseteq C$ also, and so $C = \tilde{C}$.

Take $x, y \in C$ and let $r \in \mathbb{R}_+$ with $rx - y \notin C$. Since $C = \tilde{C}$, there exists some $f \in C^*$ such that $\langle f, rx - y \rangle < 0$. Consequently,

$$\begin{aligned} \beta(x, y) &= \sup\{r > 0 : rx - y \notin C\} = \sup\{r > 0 : \langle f, rx - y \rangle < 0 \text{ for some } f \in C^*\} \\ &= \sup\{r > 0 : r\langle f, x \rangle < \langle f, y \rangle \text{ for some } f \in C^*\} = \sup\left\{\frac{\langle f, y \rangle}{\langle f, x \rangle} : f \in C^*, \langle f, x \rangle \neq 0\right\}, \end{aligned} \quad (5)$$

and similarly for $\beta(y, x)$. Then (4) is indeed equivalent to (1). \square

Remark 2. Given $\log \frac{\langle f, y \rangle \langle g, x \rangle}{\langle f, x \rangle \langle g, y \rangle} = \log \frac{\langle f, z \rangle \langle h, x \rangle}{\langle f, x \rangle \langle h, z \rangle} + \log \frac{\langle h, z \rangle \langle g, x \rangle}{\langle h, x \rangle \langle g, z \rangle}$, for any $f, g, h \in C^*$, it is easy to verify the triangle inequality for \mathcal{H} using the representation (4).

Remark 3. Once more, since the topology on X does not change the Hilbert metric on C , when working with (4) one can choose the topology, and therefore the dual space, cleverly: a coarser topology, with a smaller corresponding dual space, will almost always be preferable.

2.1 Contraction properties

Positive linear operators on a positive closed convex cone contract in the Hilbert projective distance: this result is again due to Birkhoff [7]. More generally, for proper closed convex cones $C \subset X$, Birkhoff's contraction theorem can be stated as follows:

Theorem 2.4 (Birkhoff). *Let X be a LCS, take $L : X \rightarrow X$ to be a linear transformation, and suppose that $L(C \setminus \{0\}) \subseteq C \setminus \{0\}$. If the diameter $\Delta(L) = \sup_{x, y \in C \setminus \{0\}} \mathcal{H}(Lx, Ly)$ is finite, then we have*

$$\mathcal{H}(Lx, Ly) \leq \tau(L)\mathcal{H}(x, y), \quad \forall x, y \in C \setminus \{0\}, \quad (6)$$

and $\tau(L) = \tanh\left(\frac{\Delta(L)}{4}\right)$ is called the *Birkhoff contraction coefficient*.

The theorem holds equivalently if one drops finiteness of $\Delta(L)$ as a condition and extends the definition of the contraction coefficient to $\tau(L) = 1$ when $\Delta(L) = \infty$. In other words, any bounded linear operator L is non-expansive in C under the Hilbert distance, but it is *strictly contracting* if and only if the diameter $\Delta(L)$ of C under L in the Hilbert metric is finite, i.e. $\tau(L) < 1$.

We will now show that a stronger result than Theorem 2.4 is possible.

Definition 2.5 (\mathcal{T} -distance). For $x, y \in C \setminus \{0\}$, where C is a proper closed convex cone in a real LCS, we define the *hyperbolic tangent of the Hilbert pseudo-metric* as

$$\mathcal{T}(x, y) := \tanh\left(\frac{\mathcal{H}(x, y)}{4}\right), \quad (7)$$

where $\tanh(\infty) := 1$. For simplicity we refer to (7) as the \mathcal{T} -distance.

Note that the \mathcal{T} -distance is a pseudo-metric for C : one can easily check that the triangle inequality and symmetry properties are inherited from the Hilbert distance. However, the metric \mathcal{T} makes the cone C into a *bounded* space, while \mathcal{H} gives an infinite distance between any points in $\overset{\circ}{C}$ and ∂C . Borrowing ideas from the proof of [17, Thm. 2.3], we obtain the following theorem.

Theorem 2.6. *Let X be a LCS, and $L : X \rightarrow X$ a linear transformation. Suppose that $L(C \setminus \{0\}) \subseteq C \setminus \{0\}$. Then we have*

$$\mathcal{T}(Lx, Ly) \leq \tau(L)\mathcal{T}(x, y), \quad \forall x, y \in C \setminus \{0\}, \quad (8)$$

where $\tau(L) = \sup_{x, y \in C \setminus \{0\}} \mathcal{T}(Lx, Ly)$ is the diameter of C under L in the \mathcal{T} -distance, and is equal to the Birkhoff contraction coefficient.

Proof. For all $x, y \in C \setminus \{0\}$, define the set

$$E_C(x, y) := \{r > 0 : rx - y \notin C\}.$$

Using the same notation as in (1), we have $\mathcal{H}(x, y) = \log(\beta(x, y)\beta(y, x))$, where

$$\beta(x, y) := \sup E_C(x, y) \in (0, \infty], \quad \beta(y, x) := \sup E_C(y, x) \in (0, \infty].$$

Fix $x, y \in C \setminus \{0\}$. If Lx and Ly are collinear, then $\mathcal{T}(Lx, Ly) = 0$ and the claim holds trivially. Similarly, if $\mathcal{H}(x, y) = \infty$, then $\mathcal{T}(x, y) = 1$; as $\mathcal{T}(Lx, Ly)$ is certainly less than its supremum over x and y , the claim holds. It remains to consider the case $\mathcal{H}(x, y) < \infty$ and Lx and Ly not collinear (so $\mathcal{H}(Lx, Ly) \neq 0$).

For $r > 0$, consider $rx - y \in C$. By linearity of L , we also have $rLx - Ly \in C$, and in particular $E_C(Lx, Ly) \subset E_C(x, y)$. This implies that

$$\beta(x, y) \geq \beta(Lx, Ly) > \frac{1}{\beta(Ly, Lx)} \geq \frac{1}{\beta(y, x)},$$

where the strict inequality in the middle is due to (2) and the assumption that $\mathcal{H}(Lx, Ly) \neq 0$. We now approximate $\beta(x, y)$ and $\beta(y, x)$ from above (since $\mathcal{H}(x, y) < \infty$, also $\beta(x, y), \beta(y, x) < \infty$), and $\beta(Lx, Ly)$ and $\beta(Ly, Lx)$ from below, i.e. take $M, m > 0$ and $M', m' > 0$ such that

$$M > \beta(x, y), \quad m > \beta(y, x), \quad \frac{1}{\beta(Ly, Lx)} < \left\{ \frac{1}{m'}, M' \right\} < \beta(Lx, Ly).$$

By definition of $E_C(Lx, Ly)$ and $E_C(Ly, Lx)$, we have $M'Lx - Ly \notin C$, and $m'Ly - Lx \notin C$. Similarly, $Mx - y \in C$ and $my - x \in C$. For $r > 0$, note that

$$rL(Mx - y) - L(my - x) = (rM + 1)Lx - (r + m)Ly \in C \iff \frac{rM + 1}{r + m}Lx - Ly \in C.$$

Letting $h_1(r) = (rM + 1)/(r + m)$, this implies in particular that

$$\begin{aligned} E_C(L(Mx - y), L(my - x)) &= \left\{ r > 0 : h_1(r)Lx - Ly \notin C \right\} \\ &= \left\{ h_1^{-1}(w) > 0 : wLx - Ly \notin C \right\} \\ &= h_1^{-1} \left(\left\{ \frac{1}{m} < w < M : wLx - Ly \notin C \right\} \right) \subset h_1^{-1} \left(E_C(Lx, Ly) \right), \end{aligned}$$

where $h_1^{-1}(r) = (rm - 1)/(M - r)$. Since $M' \in \left(\frac{1}{\beta(Ly, Lx)}, \beta(Lx, Ly) \right) \subset \left(\frac{1}{m}, M \right)$ by assumption, and $M'Lx - Ly \notin C$, we have that $h_1^{-1}(M') \in E_C(L(Mx - y), L(my - x))$.

Analogously, for $r > 0$,

$$rL(my - x) - L(Mx - y) \in C \iff \frac{rm + 1}{r + M}Ly - Lx \in C.$$

Letting $h_2(r) = (rm + 1)/(r + M)$ and $h_2^{-1}(r) = (rM - 1)/(m - r)$, this implies that

$$E_C(L(my - x), L(Mx - y)) = h_2^{-1} \left(\left\{ \frac{1}{M} < w < m : wLy - Lx \notin C \right\} \right) \subset h_2^{-1} \left(E_C(Ly, Lx) \right).$$

Since $\frac{1}{m'} \in (\frac{1}{m}, M)$ by assumption, which implies that $m' \in (\frac{1}{M}, m)$, and $m'Ly - Lx \notin C$, we have that $h_2^{-1}(m') \in E_C(L(my - x), L(Mx - y))$.

We know $Mx - y, my - x \in C$ by assumption, and $\Delta(L) = \sup_{x,y \in C \setminus \{0\}} \mathcal{H}(Lx, Ly)$. Therefore,

$$\begin{aligned} h_1^{-1}(M')h_2^{-1}(m') &\leq \beta(L(Mx - y), L(my - x))\beta(L(my - x), L(Mx - y)) \\ &\leq \sup_{\tilde{x}, \tilde{y} \in C \setminus \{0\}} \beta(L\tilde{x}, L\tilde{y})\beta(L\tilde{y}, L\tilde{x}) = e^{\Delta(L)}, \end{aligned}$$

which yields the inequality

$$\frac{(M'm - 1)(m'M - 1)}{(M - M')(m - m')} \leq e^{\Delta(L)}.$$

Now let $\log(Mm) = D$ and $\log(M'm') = d$. Substituting $m' = \frac{e^d}{M'}$ in the above yields

$$f(M') := \frac{(M'm - 1)(e^d M - M')}{(M - M')(mM' - e^d)} \leq e^{\Delta(L)}.$$

Differentiating the left-hand side, we find that the minimum of $f(M')$ is attained when $M' = e^{d/2} \sqrt{\frac{M}{m}}$. Substituting into the expression above, we get

$$f(M'^*) = \frac{(e^{d/2} \sqrt{Mm} - 1) \left(e^d - e^{d/2} \frac{1}{\sqrt{Mm}} \right)}{\left(1 - e^{d/2} \frac{1}{\sqrt{Mm}} \right) (e^{d/2} \sqrt{Mm} - e^d)} = \frac{\sinh^2 \left(\frac{D+d}{4} \right)}{\sinh^2 \left(\frac{D-d}{4} \right)}.$$

Taking square-roots yields

$$\frac{\sinh \left(\frac{D+d}{4} \right)}{\sinh \left(\frac{D-d}{4} \right)} \leq \sqrt{f(M')} \leq e^{\Delta(L)/2}.$$

Using the identity $\sinh(a \pm b) = \sinh(a) \cosh(b) \pm \sinh(b) \cosh(a)$ and the fact that $\frac{x-1}{x+1}$ is increasing for $x > 0$, we obtain the final expression

$$\tanh \left(\frac{d}{4} \right) \leq \tanh \left(\frac{\Delta(L)}{4} \right) \tanh \left(\frac{D}{4} \right).$$

Taking limits as $M, m \rightarrow \beta(x, y), \beta(y, x)$ and $M', m' \rightarrow \beta(Lx, Ly), \beta(Ly, Lx)$, we are done. \square

Remark 4. Concavity and monotonicity of $\tanh(x)$ for $x \geq 0$ immediately yield Birkhoff's Theorem 2.4. The advantage of using \mathcal{T} instead of \mathcal{H} is negligible when the distances are small, since $\mathcal{T}(x, y)$ is equivalent to $\mathcal{H}(x, y)$ asymptotically as $\mathcal{H}(x, y)$ approaches 0. However, we can find points $x, y \in C$ such that $\mathcal{H}(x, y) = \infty$, such as when comparing an element $x \in \overset{\circ}{C}$ with an element $y \in \partial C$. In these cases, the \mathcal{T} -distance is preferable, since $\mathcal{T}(x, y)$ stays finite and the inequality (8) remains meaningful.

Example 2.7. As we mentioned in the introduction, an immediate application of Birkhoff's theorem is in the ergodic theory of Markov processes, since transition operators are positive linear operators that map probability distributions to probability distributions, and so the assumptions of Theorem 2.4 are satisfied. We discuss explicit forms of Birkhoff's contraction coefficient for stochastic matrices and a class of transition kernels in Section 4.

3 \mathcal{H} -metric on the space of probability measures

From general LCS we now move to the space of probability measures, and consider the Hilbert projective distance in this context specifically. In [8, Chapter XVI] Birkhoff works with positive cones in a Banach lattice. Since the probability measures are a subset of the positive measures, which form the positive cone in the space of signed measures, the definition of the Hilbert distance on probability measures can be easily deduced from Birkhoff's work (see e.g. [3, Eq. 9] and [29, Def. 3.3]). In this section we choose to derive the Hilbert distance in the framework of duality instead, drawing a parallel with the works on convex cones [17, 34]. The purpose of this exercise is to gain an understanding of the Hilbert metric in terms of functions acting on probability measures, and to investigate how a change in the test-functions affects the distance itself.

Notation. For any σ -algebra \mathcal{F} and space F , let $L^0(\mathcal{F}, F)$ denote the space of \mathcal{F} -measurable functions, valued in F , and let $B(\mathcal{F}, F)$ be the subspace of bounded \mathcal{F} -measurable functions. For any two spaces E, F , let $C_b(E, F)$ denote the space of bounded continuous functions $E \rightarrow F$. If E and F are metric spaces, let $C_{bL}(E, F)$ be the space of bounded F -valued Lipschitz functions. By $\|f\|_\infty$ we denote the L^∞ -norm of f and by $\|f\|_{Lip}$ its Lipschitz coefficient.

Let (E, \mathcal{F}) be a measurable space, and consider the space $\mathcal{M}(E)$ of finite signed measures on (E, \mathcal{F}) . A natural approach is to make $\mathcal{M}(E)$ into a Banach space by equipping it with the total variation norm $\|\cdot\|_{TV}$. The total variation norm is defined, as usual, by

$$\|\mu\|_{TV} := |\mu|(E) = \mu^+(E) + \mu^-(E), \quad \text{for } \mu \in \mathcal{M}(E), \quad (9)$$

where $\mu = \mu^+ - \mu^-$ is the Hahn–Jordan decomposition of μ . It can be expressed equivalently in terms of μ acting on elements of $B(\mathcal{F}, \mathbb{R})$ as

$$\|\mu\|_{TV} := \sup \left\{ \int_E f \, d\mu : f \in B(\mathcal{F}, \mathbb{R}), \|f\|_\infty \leq 1 \right\}. \quad (10)$$

Now, the subset of positive measures $\mathcal{M}_+(E)$ is a proper closed convex cone in $\mathcal{M}(E)$. The probability measures on (E, \mathcal{F}) , denoted by $\mathcal{P}(E)$, are a subset of $\mathcal{M}_+(E)$. In (4), following ideas from [17, 34], we provided an equivalent definition of the Hilbert metric using duality. This is not a convenient approach when viewing $\mathcal{M}(E)$ as the Banach space $(\mathcal{M}(E), \|\cdot\|_{TV})$: for one thing, when dealing with signed measures, one usually prefers to work with a *predual* space instead of the dual.

Taking the predual point of view, we could consider $\mathcal{M}(E)$ as a subset of $X = C_b(E, \mathbb{R})^*$, which is a real Banach space under the operator norm. When E is a Polish space with the Borel σ -algebra $\mathcal{B}(E)$, this amounts to a linear isometric embedding that is weak*-dense. Equipping X with the weak*-topology, rather than the operator norm, we get that X is a LCS and $X^* = C_b(E, \mathbb{R})$, so one expects a predual characterisation of the cone of positive measures

in terms of $C_b(E, \mathbb{R})$. This, however, does not immediately follow from the procedure that led to (4). Instead, we give here a direct argument for the desired characterization (11), where one can think of $C_b(E, \mathbb{R}_+)$ as the ‘predual cone’, in analogy with (3). In fact, we can further restrict the space to bounded Lipschitz functions.

Proposition 3.1. *Let E be a Polish space with Borel σ -algebra $\mathcal{B}(E)$ and let $C_{bL}(E, \mathbb{R}_+)$ be the space of bounded Lipschitz functions $E \rightarrow \mathbb{R}_+$. Then we can characterize the space of positive measures $\mathcal{M}_+(E)$ as*

$$\mathcal{M}_+(E) = \{\mu \in \mathcal{M}(E) : \langle \mu, f \rangle \geq 0, \forall f \in C_{bL}(E, \mathbb{R}_+)\}. \quad (11)$$

If E is only metrizable, then replace $C_{bL}(E, \mathbb{R}_+)$ with $C_b(E, \mathbb{R}_+)$.

Proof. Let M denote the right-hand side of (11). We want to show that $\mathcal{M}_+(E) = M$. By non-negativity of the elements of $\mathcal{M}_+(E)$ and $C_{bL}(E, \mathbb{R}_+)$ (resp. $C_b(E, \mathbb{R}_+)$), it is obvious that $\mathcal{M}_+(E) \subseteq M$. For the opposite inclusion, suppose $\mu \notin \mathcal{M}_+(E)$. By the Hahn–Jordan decomposition, there exist disjoint sets $P, N \subset E$ such that $P \cup N = E$, and (Borel) measures μ^+ and μ^- such that μ^+ is supported on P and μ^- is supported on N . Since $\mu \notin \mathcal{M}_+$, we have $\mu^-(N) > 0$. Recall that μ, μ^+ and μ^- are regular, as they are Borel measures on a metric space E (see e.g [9, Thm. 7.1.7]). Take $0 < \varepsilon < \mu^-(N)/4$. By regularity of μ^- , we can find a closed set $A_\varepsilon^- \subset N$ such that $\mu^-(N \setminus A_\varepsilon^-) < \varepsilon$. Likewise, there exists a closed set $A_\varepsilon^+ \subset P$ such that $\mu^+(P \setminus A_\varepsilon^+) < \varepsilon$. Note that $A_\varepsilon^+ \cap A_\varepsilon^- = \emptyset$, since they are respectively the subsets of disjoint sets P and N . For E Polish, we can take the sets $A_\varepsilon^-, A_\varepsilon^+$ to be compact (again, [9, Thm. 7.1.7]), so the Lipschitz version of Urysohn’s Lemma [13, Prop. 2.1.1] (resp. Urysohn’s Lemma [13, Thm. 1.2.10]) guarantees that there exists $f \in C_{bL}(E, \mathbb{R}_+)$ (resp. $C_b(E, \mathbb{R}_+)$) taking values in $[0, 1]$ with

$$f(x) = \begin{cases} 0 & \text{for } x \in A_\varepsilon^+, \\ 1 & \text{for } x \in A_\varepsilon^-. \end{cases}$$

Integrating f against μ we have

$$\int_E f d\mu = \int_E f d\mu^+ - \int_E f d\mu^- \leq \mu^+(P \setminus A_\varepsilon^+) - \mu^-(A_\varepsilon^-) \leq 2\varepsilon - \mu^-(N) \leq -\frac{\mu^-(N)}{2} < 0.$$

Consequently, we have $f \in C_{bL}(E, \mathbb{R}_+)$ (resp. $C_b(E, \mathbb{R}_+)$), but $\langle \mu, f \rangle < 0$, so $\mu \notin M$. Therefore $M \subseteq \mathcal{M}_+(E)$, and the two sets are equal. \square

When E is a Polish space, the above proposition is all we need to define the Hilbert projective (pseudo-)metric on $\mathcal{M}_+(E)$ in terms of bounded positive Lipschitz functions in $C_{bL}(E, \mathbb{R}_+)$, analogously to (4). On the other hand, if we do not want to assume E to be Polish, or even metric, we need to enlarge the set of test-functions for the construction of the Hilbert metric to still make sense. Similar to Proposition 3.1, we find the following (trivial) characterization of $\mathcal{M}_+(E)$ in terms of bounded positive measurable functions.

Proposition 3.2. *Let (E, \mathcal{F}) be a measurable space and let $B(\mathcal{F}, \mathbb{R}_+)$ be the space of \mathcal{F} -measurable bounded functions taking values in \mathbb{R}_+ . Then we have*

$$\mathcal{M}_+(E) = \{\mu \in \mathcal{M}(E) : \langle f, \mu \rangle \geq 0, \forall f \in B(\mathcal{F}, \mathbb{R}_+)\}. \quad (12)$$

Proof. Let M' be the right-hand side of (12). By non-negativity of the functions in $B(\mathcal{F}, \mathbb{R}_+)$, $\mu \in \mathcal{M}_+(E)$ implies $\mu \in M'$. Conversely, assume $\mu \notin \mathcal{M}_+(E)$. Using the Hahn–Jordan decomposition, take $N \in \mathcal{F}$ such that $\mu(N) = -\mu^-(N) < 0$. Let $f := \mathbf{1}_N \in B(\mathcal{F}, \mathbb{R}_+)$. Then $\langle f, \mu \rangle < 0$, but $\langle f, \nu \rangle \geq 0$ for all $\nu \in \mathcal{M}_+(E)$. Hence $\mu \notin M'$, and we are done. \square

Proposition 3.3. *Let (E, \mathcal{F}) be a measurable space. Write $\mathcal{S} = C_{bL}(E, \mathbb{R}_+)$ if E is Polish (with \mathcal{F} the corresponding Borel σ -algebra), $\mathcal{S} = C_b(E, \mathbb{R}_+)$ if E is metrizable, or $\mathcal{S} = B(\mathcal{F}, \mathbb{R}_+)$ otherwise. Then the Hilbert projective pseudo-metric can be written as follows: for $\mu, \nu \in \mathcal{M}_+(E) \setminus \{0\}$,*

$$\mathcal{H}(\mu, \nu) = \sup_{\substack{f, g \in \mathcal{S} \\ \langle f, \mu \rangle, \langle g, \nu \rangle \neq 0}} \left\{ \log \frac{\langle f, \nu \rangle \langle g, \mu \rangle}{\langle f, \mu \rangle \langle g, \nu \rangle} \right\}. \quad (13)$$

Proof. Consider any $\mu, \nu \in \mathcal{M}_+(E)$, and take $r \in \mathbb{R}_+$ such that $r\mu - \nu \notin \mathcal{M}_+(E)$. Using Proposition 3.1 and Proposition 3.2, there is an $f \in \mathcal{S}$ such that $\langle f, r\mu - \nu \rangle < 0$. Then an argument analogous to (5) gives equivalence between (13) and the original definition of the Hilbert metric (1). \square

Now, a natural question to ask is under which conditions $\mathcal{H}(\mu, \nu)$ is finite. For $\mu, \nu \in \mathcal{M}_+(E)$, let

$$\beta(\mu, \nu) = \sup_{\substack{f \in B(\mathcal{F}, \mathbb{R}_+) \\ \langle f, \mu \rangle \neq 0}} \left\{ \frac{\langle f, \nu \rangle}{\langle f, \mu \rangle} \right\}. \quad (14)$$

Clearly, $\mathcal{H}(\mu, \nu) < \infty$ if and only if $\beta(\mu, \nu), \beta(\nu, \mu) < \infty$. We see immediately that if there exists an unbounded measurable function $h \in L^0(\mathcal{F}, \mathbb{R}_+)$ such that $\langle h, \mu \rangle < \infty$ but $\langle h, \nu \rangle = \infty$, then we can take a sequence of bounded functions $h_n \in B(\mathcal{F}, \mathbb{R}_+)$ such that $h_n \rightarrow h$, and the right-hand side of (14) is infinite. Consequently, if, for example, ν has a strictly higher number of finite moments than μ , then $\beta(\mu, \nu) = \infty$, and conversely if ν has (strictly) fewer finite moments, then $\beta(\nu, \mu) = \infty$. Thus, to have $\mathcal{H}(\mu, \nu) < \infty$, we need a condition on μ, ν that is quite a lot stronger than simple equivalence of measures (which we denote as usual by \sim). This condition, which we call *comparability* again, in accordance with Birkhoff, and denote by $\overset{\text{cmp}}{\sim}$, has already been stated in [29, Def. 3.1] and [3]. Here we derive it directly from the ‘predual’ formulation (13).

Let $\mu \sim \nu$, with Radon-Nikodym derivatives $\frac{d\mu}{d\nu}$ and $\frac{d\nu}{d\mu}$. For all $\varphi \in B(\mathcal{F}, \mathbb{R}_+)$ we have $\langle \varphi, \nu \rangle = \langle \varphi \frac{d\nu}{d\mu}, \mu \rangle \leq \left\| \frac{d\nu}{d\mu} \right\|_{L^\infty(\mu)} \langle \varphi, \mu \rangle$, so

$$M_\varphi := \sup_{\substack{\varphi \in B(\mathcal{F}, \mathbb{R}_+) \\ \langle \varphi, \mu \rangle \neq 0}} \left\{ \frac{\langle \varphi, \nu \rangle}{\langle \varphi, \mu \rangle} \right\} \leq \left\| \frac{d\nu}{d\mu} \right\|_{L^\infty(\mu)}.$$

On the other hand,

$$\left\langle \varphi \frac{d\nu}{d\mu}, \mu \right\rangle = \langle \varphi, \nu \rangle \leq \langle \varphi, \mu \rangle M_\varphi,$$

so $\frac{d\nu}{d\mu} \leq M_\varphi$ μ -a.e., and in particular $\left\| \frac{d\nu}{d\mu} \right\|_{L^\infty(\mu)} \leq M_\varphi$. Therefore, $\beta(\mu, \nu) = \left\| \frac{d\nu}{d\mu} \right\|_{L^\infty(\mu)}$, and hence we can state our comparability condition as follows:

Definition 3.4. Let (E, \mathcal{F}) be a measurable space. Two positive measures $\mu, \nu \in \mathcal{M}_+(E)$ are *comparable* if $\mu \sim \nu$ and their Radon-Nikodym derivatives $\frac{d\mu}{d\nu}$ and $\frac{d\nu}{d\mu}$ are essentially bounded, i.e. $\frac{d\mu}{d\nu} \in L^\infty(\nu)$ and $\frac{d\nu}{d\mu} \in L^\infty(\mu)$. Equivalently, μ and ν are comparable if there exists scalars $q, r > 0$ such that

$$q\mu(A) \leq \nu(A) \leq r\mu(A), \quad \forall A \in \mathcal{F}. \quad (15)$$

Then the Hilbert projective pseudo-metric for $\mu, \nu \in \mathcal{M}_+(E)$ can be defined as

$$\mathcal{H}(\mu, \nu) = \log \left(\left\| \frac{d\mu}{d\nu} \right\|_\infty \left\| \frac{d\nu}{d\mu} \right\|_\infty \right) = \sup_{\substack{A, B \in \mathcal{F} \\ \nu(A) > 0, \mu(B) > 0}} \left\{ \log \frac{\nu(B)\mu(A)}{\mu(B)\nu(A)} \right\}, \quad \text{if } \mu \stackrel{\text{cmp}}{\sim} \nu, \quad (16)$$

and $\mathcal{H}(\mu, \nu) = \infty$ otherwise, and these definitions are equivalent to (13). The right-most formulation of (16) is the definition chosen, for example, by Le Gland and Oudjane in [29, Def. 3.3] and Atar and Zeitouni in [3]. Note that if $\mu, \nu \in \mathcal{P}(E)$, then $\left\| \frac{d\mu}{d\nu} \right\|_\infty, \left\| \frac{d\nu}{d\mu} \right\|_\infty \geq 1$ since μ and ν must integrate to 1, and hence also $\left\| \frac{d\mu}{d\nu} \right\|_\infty, \left\| \frac{d\nu}{d\mu} \right\|_\infty \leq e^{\mathcal{H}(\mu, \nu)}$ by (16). Then, for $\mu, \nu \in \mathcal{P}(E)$, $\mu \sim \nu$, and an arbitrary $f \in B(\mathcal{F}, \mathbb{R})$ such that $\|f\|_\infty \leq 1$, we have

$$\begin{aligned} |\langle f, \mu - \nu \rangle| &\leq \int_{\left\{ \frac{d\mu}{d\nu} \geq 1 \right\}} |f| \left(\frac{d\mu}{d\nu} - 1 \right) d\nu + \int_{\left\{ \frac{d\mu}{d\nu} < 1 \right\}} |f| \left(1 - \frac{d\mu}{d\nu} \right) d\nu \\ &\leq \left(\left\| \frac{d\mu}{d\nu} \right\|_\infty - 1 \right) \nu(\left\{ \frac{d\mu}{d\nu} \geq 1 \right\}) + \left(1 - \left\| \frac{d\nu}{d\mu} \right\|_\infty^{-1} \right) \nu(\left\{ \frac{d\mu}{d\nu} < 1 \right\}) \leq e^{\mathcal{H}(\mu, \nu)} - 1. \end{aligned}$$

Together with the fact that the total variation distance between two probability measures is at most 2, this yields the following bound, first shown by Atar and Zeitouni [3, Lemma 1]:

$$\|\mu - \nu\|_{\text{TV}} \leq \frac{2}{\log 3} \mathcal{H}(\mu, \nu). \quad (17)$$

This bound is clearly not sharp, since the right-hand side can easily be much larger than 2. We will improve it in Corollary 5.2.1. For now, we use Atar and Zeitouni's result to prove the following Lemma.

Lemma 3.5. *Let (E, \mathcal{F}) be a measurable space. Then $(\mathcal{P}(E), \mathcal{H})$ is a complete metric space.*

Proof. Note that two probability measures $\mu, \nu \in \mathcal{P}(E)$ which are collinear must be necessarily equal, so \mathcal{H} is a metric on $\mathcal{P}(E)$. Let $(\mu_n) \in \mathcal{P}(E)$ be a Cauchy sequence for the Hilbert metric \mathcal{H} . Then (μ_n) is also Cauchy in total variation norm by (17), so $\mu_n \rightarrow \mu \in \mathcal{P}(E)$ in $\|\cdot\|_{\text{TV}}$ since $\mathcal{P}(E)$ is complete as it is a closed subset of $\mathcal{M}(E)$. Since $\langle f, \mu_n \rangle \rightarrow \langle f, \mu \rangle$ for $f \in B(E, \mathbb{R})$ if $\mu_n \rightarrow \mu$ in total variation norm, (13) gives that \mathcal{H} is lower semi-continuous with respect to $\|\cdot\|_{\text{TV}}$, as a supremum over continuous functions. Hence $\mathcal{H}(\mu_n, \mu) \leq \liminf_{k \rightarrow \infty} \mathcal{H}(\mu_n, \mu_k)$, where the right-hand side goes to 0 as $n \rightarrow \infty$ by the Cauchy assumption. \square

Corollary 3.5.1. *Let (E, \mathcal{F}) be a measurable space. Then $(\mathcal{P}(E), \mathcal{T})$ is a complete metric space.*

We have seen in Section 2 that many of the interesting properties of the Hilbert metric also hold for its transformation \mathcal{T} . The following gives a key reason why the classic Hilbert pseudo-metric is also of interest: \mathcal{H} turns the space of probability measures comparable to a reference measure ρ into a normed vector space (with a modified algebra).

Proposition 3.6. *Let (E, \mathcal{F}) be a measurable space and fix a reference measure $\rho \in \mathcal{M}_+(E)$. We let essential infima and suprema be defined with respect to the nullsets of ρ , and consider*

- (i) *the space of measures comparable to ρ , namely $\mathcal{M}_\rho := \{\mu \in \mathcal{M}_+(E) : \frac{d\mu}{d\rho}, \frac{d\rho}{d\mu} \in L^\infty(\rho)\}$, and $\mathcal{P}_\rho := \mathcal{M}_\rho \cap \mathcal{P}(E)$;*

- (ii) the equivalence relation \sim_{coll} on \mathcal{M}_ρ given by collinearity, that is $\mu \sim_{\text{coll}} \nu \Leftrightarrow \mu = c\nu$ for some $c > 0$; note that \mathcal{P}_ρ is isomorphic to $\mathcal{M}_\rho / \sim_{\text{coll}}$ (as it is a selection of a unique element from each equivalence class);
- (iii) the equivalence relation \sim_{const} on $L^\infty(\rho)$ given by $f \sim_{\text{const}} g \Leftrightarrow f = g + c$ ρ -a.e. for some $c \in \mathbb{R}$; and the associated quotient space $\Theta_\rho := L^\infty(\rho) / \sim_{\text{const}}$.

Then the map

$$\|\cdot\|_\Theta : L^\infty(\rho) \rightarrow \mathbb{R}; \quad f \mapsto \operatorname{ess\,sup}_{x \in E} f(x) - \operatorname{ess\,inf}_{x \in E} f(x),$$

defines a seminorm on $L^\infty(\rho)$, and a norm on Θ_ρ . Moreover, the map

$$\theta : \mathcal{M}_\rho \rightarrow L^\infty(\rho); \quad \mu \mapsto \log(d\mu/d\rho)$$

is an isomorphism of the pseudo-metric spaces $(\mathcal{M}_\rho, \mathcal{H})$ and $(L^\infty(\rho), \|\cdot\|_\Theta)$, satisfying

$$\mathcal{H}(\mu, \nu) = \|\theta(\mu) - \theta(\nu)\|_\Theta, \quad \text{for all } \mu, \nu \in \mathcal{M}_\rho,$$

and it is an isomorphism of the metric spaces $(\mathcal{P}_\rho, \mathcal{H})$ and $(\Theta_\rho, \|\cdot\|_\Theta)$. In particular, $(\mathcal{P}_\rho, \mathcal{H})$ is a normed vector space, when endowed with the algebra of (renormalized) addition and scalar multiplication of log-densities.

Proof. It is easy to see that $\|\cdot\|_\Theta$ is absolutely homogeneous. From (16), we know that

$$\begin{aligned} \mathcal{H}(\mu, \nu) &= \log \left(\left\| \frac{d\mu}{d\nu} \right\|_\infty \right) + \log \left(\left\| \frac{d\nu}{d\mu} \right\|_\infty \right) \\ &= \operatorname{ess\,sup} \left\{ \log \left(\frac{d\mu}{d\nu} \right) \right\} - \operatorname{ess\,inf} \left\{ \log \left(\frac{d\mu}{d\nu} \right) \right\} \\ &= \operatorname{ess\,sup} \left\{ \theta(\mu) - \theta(\nu) \right\} - \operatorname{ess\,inf} \left\{ \theta(\mu) - \theta(\nu) \right\} \\ &= \|\theta(\mu) - \theta(\nu)\|_\Theta. \end{aligned}$$

From this it follows that $\|\cdot\|_\Theta$ is sublinear (as \mathcal{H} satisfies the triangle inequality), and is therefore a seminorm. It is easy to check that $\|\theta\|_\Theta = 0$ iff $\theta \sim_{\text{const}} 0$, so $\|\cdot\|_\Theta$ is a norm on the vector space $\Theta_\rho = L^\infty(\rho) / \sim_{\text{const}}$.

For $f \in \Theta_\rho$, the inverse of $\theta : \mathcal{M}_\rho \rightarrow L^\infty(\rho)$ is given by

$$\theta^{-1}(f)(A) = \int_A \exp(f(x)) d\rho, \quad \forall A \in \mathcal{F},$$

so θ is clearly a bijection, and hence an isomorphism of $(\mathcal{M}_\rho, \mathcal{H})$ and $(L^\infty(\rho), \|\cdot\|_\Theta)$. Similarly, taking account of the equivalence relation, the inverse of $\theta : \mathcal{P}_\rho \rightarrow \Theta_\rho$ is given, for $f \in \Theta_\rho$, by

$$\theta^{-1}(f + c)(A) = \frac{\int_A \exp(f(x)) d\rho}{\int_E \exp(f(x)) d\rho}, \quad \forall A \in \mathcal{F},$$

which clearly does not depend on the choice of $c \in \mathbb{R}$ (and hence is well defined on $\Theta_\rho = L^\infty(\rho) / \sim_{\text{const}}$). It follows that θ is an isomorphism of $(\mathcal{P}_\rho, \mathcal{H})$ and $(\Theta_\rho, \|\cdot\|_\Theta)$.

Finally, as $(\Theta_\rho, \|\cdot\|_\Theta)$ is a normed vector space, we simply observe that addition and scalar multiplication in Θ_ρ correspond to (renormalized) addition and scalar multiplication of log-densities. \square

Remark 5. We will see in Section 4 that, when E is finite, we can avoid the equivalence relation above by selecting the unique representative $\theta_0(\mu)$ which satisfies $\theta_0(\mu)(x_0) = 0$ for a fixed $x_0 \in E$ (see Remark 10). This does not work as cleanly in infinite state spaces, as the value at a single point is typically not well defined when functions are only specified ρ -a.e.

Remark 6. Proposition 3.6 also helps us to understand the topology of $(\mathcal{P}(E), \mathcal{H})$. For every $\rho \in \mathcal{P}(E)$, we have the corresponding vector space \mathcal{P}_ρ (and any $\rho' \in \mathcal{P}_\rho$ will yield $\mathcal{P}_{\rho'} = \mathcal{P}_\rho$). As they are normed vector spaces (with an appropriate algebra), these sets are both closed and open, and give a disconnected partition of $\mathcal{P}(E)$. In other words, $(\mathcal{P}(E), \mathcal{H})$ has the topology of a disjoint union of normed vector spaces (which may have different dimensions).

We conclude this (rather lengthy) section about the Hilbert metric on probability measures with a few important observations, which motivate why we started looking carefully at the ‘predual’ formulation of the Hilbert metric in the first place.

Remark 7. Take $\mu, \nu \in \mathcal{M}_+(E)$, with E Polish. Consider distances of the form

$$D(\mu, \nu) = \sup \left\{ \left| \int_E f \, d(\mu - \nu) \right| : f \in C_{bL}(E, \mathbb{R}_+) \right\}.$$

Different conditions on $\|f\|_\infty$ and $\|f\|_{Lip}$ yield different metrics: the total variation norm (10) if one imposes $\|f\|_\infty \leq 1$, the bounded-Lipschitz distance by taking $\|f\|_\infty + \|f\|_{Lip} \leq 1$, or the 1-Wasserstein distance \mathcal{W}_1 (when μ and ν are additionally taken to have finite first moment) by imposing $\|f\|_{Lip} \leq 1$. This differentiation based on the choice of test-functions is *completely lost* when we work with the Hilbert metric. If we restricted our space to $\mathcal{P}_1(E)$ (the probability measures on E with finite first moment), for example, we could of course characterize our ‘predual’ cone of test-functions using (unbounded) positive Lipschitz functions $Lip(E, \mathbb{R}_+)$, analogously to the Kantorovich–Rubinstein dual formulation of \mathcal{W}_1 . However, this would not yield a different metric from (13). Since any Lipschitz function can be approximated from below by bounded Lipschitz functions, if $\mu \overset{\text{cmp}}{\sim} \nu$ and $\mu, \nu \in \mathcal{P}_1(E)$, taking the supremum over $Lip(E, \mathbb{R}_+)$ or $C_{bL}(E, \mathbb{R}_+)$ does not change the Hilbert distance.

In the wake of the above remark, we deduce that the Hilbert metric is stronger than the p -Wasserstein distance \mathcal{W}_p . Let (E, d) be a metric space and $\{\mu_n\} \in \mathcal{P}_p(E)$ a sequence of probability measures with finite p^{th} -moment such that $\mu_n \rightarrow \mu$ in Hilbert metric. By (17), convergence in Hilbert metric implies convergence in total variation norm, which in turn implies $\mu_n \rightarrow \mu$ weakly. Moreover, $\mu \in \mathcal{P}_p(E)$, by definition of the Hilbert metric and comparability of measures (15). Fix an arbitrary $x_0 \in E$. Then an argument similar to the one that lead to (17) yields that, for all $q \leq p$,

$$\left| \int_E d(x_0, x)^q \, d(\mu_n - \mu) \right| \leq K_q (e^{\mathcal{H}(\mu_n, \mu)} - 1), \quad (18)$$

where $K_q < \infty$ is the q^{th} -moment of $\mu \in \mathcal{P}_p(E)$. So convergence of moments is preserved under convergence in the Hilbert metric, and thus convergence in the Hilbert metric implies convergence in \mathcal{W}_p .

The Kantorovich–Rubinstein dual formulation in particular yields the following bound for the \mathcal{W}_1 distance with respect to \mathcal{H} . For $\mu, \nu \in \mathcal{P}_1(E)$, and an arbitrary $x_0 \in E$, we have

$$\mathcal{W}_1(\mu, \nu) \leq (e^{\mathcal{H}(\mu, \nu)} - 1) \int_E d(x_0, x) \, d\mu, \quad (19)$$

or analogously in terms of the first moment of ν .

Finally, our work so far, the definition of the Hilbert metric and of comparability of measures (15), all clearly point to emphasise that convergence in the Hilbert metric is a very strong form of convergence. The Hilbert projective metric not only dominates TV and \mathcal{W}_p , but also the Kullback–Leibler divergence (or relative entropy):

$$D_{KL}(\mu\|\nu) := \int_E \log \frac{d\mu}{d\nu} d\mu \leq \log \left\| \frac{d\mu}{d\nu} \right\|_\infty \leq \mathcal{H}(\mu, \nu). \quad (20)$$

In fact, one can show that:

Proposition 3.7. *Let (E, \mathcal{F}) be a measurable space, and let $\{\mu_n\} \in \mathcal{P}(E)$ be a sequence of probability measures converging to $\mu \in \mathcal{P}(E)$ in \mathcal{H} . Then, for any f -divergence D_f , $D_f(\mu_n\|\mu) \rightarrow 0$ and $D_f(\mu\|\mu_n) \rightarrow 0$ as $n \rightarrow \infty$ also.*

Proof. Let $f : \mathbb{R}_+ \rightarrow (-\infty, \infty]$ be a convex function with $f(1) = 0$, and $f(u) < \infty$ for all $u > 0$. Let $\mu, \nu \in \mathcal{P}(E)$, with $\mu \ll \nu$. Then the f -divergence of μ from ν is given by

$$D_f(\mu\|\nu) = \int_E f\left(\frac{d\mu}{d\nu}\right) d\nu.$$

Note that the f -divergence is unchanged if we add a linear term to f , i.e. let $\bar{f}(u) = f(u) + c(u - 1)$, then $D_{\bar{f}}(\mu\|\nu) = D_f(\mu\|\nu)$. Moreover, by taking $c \in -\partial f(1)$ (where by ∂ we denote the subgradient of f), we have $0 \in \partial \bar{f}(1)$, so without loss of generality we can restrict our attention to convex functions f such that $f(1) = 0 \in \partial f(1)$. Consider a sequence $\{\mu_n\} \in \mathcal{P}(E)$ such that $\lim_{n \rightarrow \infty} \mathcal{H}(\mu_n, \mu) = 0$. Then there exists $N > 0$ such that for all $n \geq N$, $\mu_n \sim \nu$. Since f must be decreasing for $x < 1$ and increasing for $x > 1$ by virtue of being convex, we have, for all $n \geq N$,

$$\begin{aligned} D_f(\mu_n\|\mu) &= \int_{\left\{\frac{d\mu_n}{d\mu} \leq 1\right\}} f\left(\frac{d\mu_n}{d\mu}\right) d\mu + \int_{\left\{\frac{d\mu_n}{d\mu} > 1\right\}} f\left(\frac{d\mu_n}{d\mu}\right) d\mu \\ &\leq f\left(\operatorname{ess\,inf}_{x \in E} \frac{d\mu_n}{d\mu}\right) \mu(\left\{\frac{d\mu_n}{d\mu} \leq 1\right\}) + f\left(\operatorname{ess\,sup}_{x \in E} \frac{d\mu_n}{d\mu}\right) \mu(\left\{\frac{d\mu_n}{d\mu} > 1\right\}) \\ &\leq \max \left\{ f\left(\left\| \frac{d\mu}{d\mu_n} \right\|_\infty^{-1}\right), f\left(\left\| \frac{d\mu_n}{d\mu} \right\|_\infty\right) \right\} \leq \max \left\{ f(e^{-\mathcal{H}(\mu_n, \mu)}), f(e^{\mathcal{H}(\mu_n, \mu)}) \right\}, \end{aligned}$$

where we have used that $\left\| \frac{d\mu}{d\mu_n} \right\|_\infty, \left\| \frac{d\mu_n}{d\mu} \right\|_\infty \geq 1$, and $\left\| \frac{d\mu}{d\mu_n} \right\|_\infty^{-1} \geq e^{-\mathcal{H}(\mu_n, \mu)}$ and $\left\| \frac{d\mu_n}{d\mu} \right\|_\infty \leq e^{\mathcal{H}(\mu_n, \mu)}$. Since $f(1) = 0$ by assumption, the right-hand side of the above goes to 0 as $\mathcal{H}(\mu_n, \mu) \rightarrow 0$, so $D_f(\mu_n\|\mu)$ converges. The argument for $D_f(\mu\|\mu_n)$ is analogous by symmetry, and we are done. \square

Remark 8. Total variation distance, Kullback–Leibler divergence, Jensen–Shannon divergence, squared Hellinger distance and χ^2 -divergence are all examples of f -divergences.

4 Hilbert projective geometry on the probability simplex

In this section we consider the Hilbert metric on the probability measures with finite state-space $E \cong \{0, \dots, n\}$, which form the *probability simplex*. In this case, the form of the Hilbert

metric simplifies, and there exists an explicit expression for Birkhoff's contraction coefficient (see Section 4 of [36, Chapter 3]). We briefly remark on this, and present a short derivation of Birkhoff's coefficient using duality. Then we move on to studying the geometry of the probability simplex under the Hilbert projective metric: using a coordinate transformation inspired by information geometry [1, 2], we describe the Hilbert balls as convex polytopes in the probability simplex, in an extension of the work in [33] to the n -dimensional case.

Let $E \cong \{0, \dots, n\}$. The probability measures $\mathcal{P}(E)$ are represented by the set

$$\mathcal{P}(E) \cong \mathcal{S}^n = \left\{ x \in \mathbb{R}^{n+1} : \sum_{i=0}^n x_i = 1 \right\} \subset \mathbb{R}^{n+1},$$

and \mathcal{S}^n is called the n -dimensional probability simplex. It is given by the intersection of the convex cone of non-negative vectors \mathbb{R}_+^{n+1} with the plane $\sum_i x_i = 1$.

Consider the Hilbert distance on the non-negative orthant $C = \mathbb{R}_+^n$. Recall the duality expression for \mathcal{H} given in (4) and the equality (5). The dual cone C^* is again \mathbb{R}_+^n . Take $x, y \in \mathbb{R}_+^n \setminus \{0\}$ such that $\beta(x, y) < \infty$. Note that $\beta(x, y) < \infty$ if and only if there exists a scalar $b > 0$ such that $y^i \leq bx^i$ for all $i = 1, \dots, n$, which in particular implies that if $y^i = 0$ for some i , then $x^i = 0$. We have

$$\beta(x, y) = \sup_{\substack{w \in \mathbb{R}_+^n \setminus \{0\} \\ w^\top x > 0}} \frac{w^\top y}{w^\top x} = \sup_{r \in [0, 1]^{n+1} \setminus \{0\}} \sum_{j: x^j > 0} r^j \frac{y^j}{x^j} = \max_{j: x^j > 0} \frac{y^j}{x^j} = \max_{j: e_j^\top x > 0} \frac{e_j^\top y}{e_j^\top x}, \quad (21)$$

where the second equality follows by setting $r^j = w^j x^j / w^\top x$, $0 \leq r^j \leq 1$ for all $j = 1, \dots, n$ and at least one $r^j > 0$, and $\{e_j\}_{j=1}^n$ are the basis vectors of \mathbb{R}_+^n . So $\sup_w w^\top y / w^\top x$ is attained when w is a basis vector. By symmetry, we have that $\beta(y, x) < \infty$ if there exists $b' > 0$ such that $x^i \leq b' y^i$ for all $i = 1, \dots, n$. Then two elements $x, y \in \mathbb{R}_+^n$ are *comparable* (denoted again by $\overset{\text{cmp}}{\sim}$) if there exist constants $a, b > 0$ such that $ax \leq y \leq bx$, where the inequalities hold component-wise (this is the definition of comparability given in [8, Chapter XVI]). Then the definition (1) of the Hilbert projective distance for $x, y \in \mathbb{R}_+^n$ simplifies to

$$\mathcal{H}(x, y) = \begin{cases} \log \left(\frac{\max_{i: y^i > 0} \frac{x^i}{y^i}}{\min_{j: y^j > 0} \frac{x^j}{y^j}} \right), & x \overset{\text{cmp}}{\sim} y, \\ \infty, & x \not\overset{\text{cmp}}{\sim} y. \end{cases} \quad (22)$$

Remark 9. In this finite-state context, the comparability condition $\overset{\text{cmp}}{\sim}$ reduces to *equivalence of measures* \sim on \mathcal{S}^n . Recall that $(\mathcal{S}^n, \mathcal{H})$ is a complete metric space by Lemma 3.5.

Using (21) we can now easily derive an explicit expression for Birkhoff's contraction coefficient of a linear operator $\mathbb{R}_+^n \setminus \{0\} \rightarrow \mathbb{R}_+^n \setminus \{0\}$. Define a matrix $A = (A_{ij})$ to be *allowable* if A is non-negative (i.e. $A_{ij} \geq 0$ for all i, j) and if every row and every column of A has at least one strictly positive element (this definition is given by Seneta in [36, Def. 3.1]). Clearly any linear operator $\mathbb{R}_+^n \setminus \{0\} \rightarrow \mathbb{R}_+^n \setminus \{0\}$ can be represented as an allowable $n \times n$ matrix. We prove the following result, which was already stated by Birkhoff without proof in Corollary 2 of [8, Chapter XVI, Section 3] and obtained by Seneta in Section 4 of [36, Chapter 3], although the derivation there is significantly more involved.

Proposition 4.1. *Let $A = (A_{ij})$ be an allowable $n \times n$ matrix. Birkhoff's contraction coefficient $\tau(A)$ can be written as*

$$\tau(A) = \frac{1 - \sqrt{\phi(A)}}{1 + \sqrt{\phi(A)}}, \quad \text{with } \phi(A) = \min_{i,j,k,l} \frac{A_{ik}A_{jl}}{A_{jk}A_{il}}, \quad (23)$$

(with the convention that $0/0 = 1$).

Proof. Consider the diameter of \mathbb{R}_+^n under the matrix A , i.e.

$$\Delta(A) = \sup_{x,y \in \mathbb{R}_+^n \setminus \{0\}} \mathcal{H}(Ax, Ay).$$

Assume that $\Delta(A) < \infty$, so $\beta(Ax, Ay), \beta(Ay, Ax) < \infty$ for all $x, y \in \mathbb{R}_+^n \setminus \{0\}$. Note that $\Delta(A) < \infty$ if and only if A is strictly positive (i.e. $A_{ij} > 0$ for all i, j), so in particular Ax has strictly positive entries for all $x \in \mathbb{R}_+^n$, which implies that $w^\top Ax > 0$ for all $x, w \in \mathbb{R}_+^n$. Using (21) in the second and fourth equalities below, we get

$$\begin{aligned} e^{\Delta(A)} &= \sup_{x,y \in \mathbb{R}_+^n \setminus \{0\}} \sup_{w,z \in \mathbb{R}_+^n \setminus \{0\}} \left\{ \frac{w^\top Ay \ z^\top Ax}{w^\top Ax \ z^\top Ay} \right\} = \sup_{x,y \in \mathbb{R}_+^n \setminus \{0\}} \max_{i,j} \left\{ \frac{e_i^\top Ay \ e_j^\top Ax}{e_i^\top Ax \ e_j^\top Ay} \right\} \\ &= \max_{i,j} \sup_{x,y \in \mathbb{R}_+^n \setminus \{0\}} \left\{ \frac{y^\top A^\top e_i \ x^\top A^\top e_j}{y^\top A^\top e_j \ x^\top A^\top e_i} \right\} = \max_{i,j} \max_{k,l} \left\{ \frac{e_k^\top A^\top e_i \ e_l^\top A^\top e_j}{e_k^\top A^\top e_j \ e_l^\top A^\top e_i} \right\} = \max_{i,j,k,l} \frac{A_{ik}A_{jl}}{A_{jk}A_{il}}, \end{aligned}$$

so finally Birkhoff's contraction coefficient is given by

$$\tau(A) = \tanh \left(\frac{\Delta(A)}{4} \right) = \frac{e^{\frac{\Delta(A)}{2}} - 1}{e^{\frac{\Delta(A)}{2}} + 1} = \frac{1 - \sqrt{\phi(A)}}{1 + \sqrt{\phi(A)}}, \quad \text{with } \phi(A) = \min_{i,j,k,l} \frac{A_{ik}A_{jl}}{A_{jk}A_{il}}. \quad (24)$$

We can check that if A is not strictly positive, so $\Delta(A) = \infty$, the formula above still holds with the convention that $\tanh(\infty) := 1$. \square

In fact, we obtain a similar representation of Birkhoff's contraction coefficient for a class of transition kernels on more general spaces. Let E be a Polish space with Borel σ -algebra $\mathcal{B}(E)$, and consider a positive kernel K on $\mathcal{B}(E) \times E$. Then there exists an associated positive linear operator (again denoted by K) such that $K : \mathcal{M}_+(E) \rightarrow \mathcal{M}_+(E)$ and

$$K\mu(da) = \int_E K(da, x) d\mu(x). \quad (25)$$

In the statement below, we restrict our attention to kernels that have a density with respect to a reference measure $\rho \in \mathcal{M}_+(E)$. In other words, let $K(da, x) = \kappa(a, x) d\rho(a)$ for some positive function $\kappa : \text{Supp}(\rho) \times E \rightarrow \mathbb{R}_+$. Then (25) reduces to

$$K\mu(A) = \int_{a \in A} \int_{x \in E} \kappa(a, x) d\mu(x) d\rho(a), \quad \forall A \in \mathcal{B}(E), \quad (26)$$

(where we have swapped the order of integration using Tonelli's theorem). We then have the following infinite dimensional counterpart to Proposition 4.1.

Proposition 4.2. *Let E be a Polish space with Borel σ -algebra $\mathcal{B}(E)$ and reference measure $\rho \in \mathcal{M}_+(E)$ with support $\text{Supp}(\rho) \subset E$. Consider a kernel operator $K : \mathcal{M}_+(E) \rightarrow \mathcal{M}_+(E)$ of the form (26) defined by a density*

$$\frac{d(K\mu)}{d\rho}(a) := \int_E \kappa(a, x) d\mu(x),$$

for $\kappa : \text{Supp}(\rho) \times E \rightarrow (0, \infty)$. Assume κ is a bounded continuous function. Then the Birkhoff coefficient of K is given by

$$\tau(K) = \frac{1 - \sqrt{\phi(K)}}{1 + \sqrt{\phi(K)}}, \quad \text{with} \quad \phi(K) = \inf_{\substack{a, b \in \text{Supp}(\rho) \\ x, y \in E}} \left\{ \frac{\kappa(a, x)\kappa(b, y)}{\kappa(a, y)\kappa(b, x)} \right\}. \quad (27)$$

Proof. From the structure of the operator, we know that the Radon–Nikodym derivative of $K\mu$ and $K\nu$ (for $\nu \neq 0$) is given by

$$\frac{d(K\mu)}{d(K\nu)}(a) = \frac{\int_E \kappa(a, x) d\mu(x)}{\int_E \kappa(a, x) d\nu(x)}, \quad \forall a \in \text{Supp}(\rho).$$

As κ is continuous and bounded, by dominated convergence we have that $a \mapsto \frac{d(K\mu)}{d(K\nu)}(a)$ is continuous, and so the ρ -essential supremum of $\frac{d(K\mu)}{d(K\nu)}(a)$ is equal to its (pointwise) supremum on $\text{Supp}(\rho)$. Using the definition of \mathcal{T} and (16), we know that

$$\begin{aligned} \mathcal{T}(K\mu, K\nu) &= \tanh \left(\frac{1}{4} \log \left(\sup_a \left\{ \frac{\int_E \kappa(a, x) d\mu(x)}{\int_E \kappa(a, x) d\nu(x)} \right\} \sup_b \left\{ \frac{\int_E \kappa(b, y) d\nu(y)}{\int_E \kappa(b, y) d\mu(y)} \right\} \right) \right) \\ &= \sup_{a, b} \left\{ \tanh \left(\frac{1}{4} \log \left(\frac{\int_E \kappa(a, x) d\mu(x)}{\int_E \kappa(b, y) d\mu(y)} \frac{\int_E \kappa(b, y) d\nu(y)}{\int_E \kappa(a, x) d\nu(x)} \right) \right) \right\}. \end{aligned}$$

From the definition of τ in Theorem 2.6 and monotonicity of \tanh , we know that

$$\begin{aligned} \tau(K) &= \sup_{\mu, \nu \in \mathcal{M}_+(E)} \mathcal{T}(K\mu, K\nu) \\ &= \sup_{a, b} \left\{ \tanh \left(\frac{1}{4} \log \left(\sup_{\mu \in \mathcal{M}_+(E)} \left\{ \frac{\int_E \kappa(a, x) d\mu(x)}{\int_E \kappa(b, y) d\mu(y)} \right\} \sup_{\nu \in \mathcal{M}_+(E)} \left\{ \frac{\int_E \kappa(b, y) d\nu(y)}{\int_E \kappa(a, x) d\nu(x)} \right\} \right) \right) \right\}. \end{aligned} \quad (28)$$

In order to compute the inner suprema, we observe that for all $\mu \in \mathcal{M}_+(E)$ and $a, b \in E$,

$$\frac{\int_E \kappa(a, x) d\mu(x)}{\int_E \kappa(b, y) d\mu(y)} = \int_E \frac{\kappa(b, x)}{\int_E \kappa(b, y) d\mu(y)} \frac{\kappa(a, x)}{\kappa(b, x)} d\mu(x) = \int_E \frac{\kappa(a, x)}{\kappa(b, x)} d\tilde{\mu}_b(x)$$

where $\tilde{\mu}_b \in \mathcal{P}(E)$ is defined by $\frac{d\tilde{\mu}_b}{d\mu}(x) = \frac{\kappa(b, x)}{\int_E \kappa(b, y) d\mu(y)}$. Therefore, as κ is continuous,

$$\sup_{\mu \in \mathcal{M}_+(E)} \frac{\int_E \kappa(a, x) d\mu(x)}{\int_E \kappa(b, y) d\mu(y)} = \sup_{\tilde{\mu} \in \mathcal{P}(E)} \int_E \frac{\kappa(a, x)}{\kappa(b, x)} d\tilde{\mu}(x) = \sup_{x \in E} \frac{\kappa(a, x)}{\kappa(b, x)}.$$

Substituting in (28) yields

$$\tau(K) = \tanh \left(\frac{\Delta(K)}{4} \right), \quad \text{with} \quad \Delta(K) = \sup_{\substack{a, b \in \text{Supp}(\rho) \\ x, y \in E}} \left\{ \log \left(\frac{\kappa(a, x)\kappa(b, y)}{\kappa(a, y)\kappa(b, x)} \right) \right\},$$

and essentially the same calculation as (24) gives the form (27). \square

4.1 Hexagonal polytopes in Hilbert projective geometry

We now introduce a change of coordinates from the interior of \mathcal{S}^n to \mathbb{R}^n which allows us to build an understanding of the geometry of the probability simplex as a metric space equipped with the Hilbert distance \mathcal{H} . We have already employed this coordinate transformation in [14] in the setting of nonlinear filtering, and it proved crucial for the analysis of the Hilbert distance between two probability measures $\mu, \nu \in \mathcal{S}^n$.

Notation. For simplicity, let $N := \{0, 1, \dots, n\}$ throughout this section.

Let $\mathring{\mathcal{S}}^n$ be the interior of the n -dimensional probability simplex $\mathcal{S}^n \subset \mathbb{R}^{n+1}$. Following Amari [1, 2], we map any discrete distribution $\mu \in \mathring{\mathcal{S}}^n$ to its natural parameters $\theta \in \mathbb{R}^n$. In other words, for all $k = 0, \dots, n$, let $\theta_k : \mathring{\mathcal{S}}^n \rightarrow \mathbb{R}^n$ such that

$$\theta_k^i(\mu) = \log \frac{\mu^i}{\mu^k}, \quad \forall i \in N \setminus \{k\}. \quad (29)$$

Since we exclude the k^{th} component, $\theta_k(\mu) = \{\theta_k^i(\mu)\}_{i \neq k}$ is an n -dimensional vector. The inverse mapping $\theta_k^{-1} : \mathbb{R}^n \rightarrow \mathring{\mathcal{S}}^n$ is given by

$$\mu^i(\theta_k) = \frac{e^{\theta_k^i}}{\sum_{i=0}^n e^{\theta_k^i}}, \quad \forall i \in N \quad (\text{where for notational simplicity } \theta_k^k \equiv 0). \quad (30)$$

Then θ_k is a diffeomorphism $\mathring{\mathcal{S}}^n \rightarrow \mathbb{R}^n$, and a global chart for $\mathring{\mathcal{S}}^n$. Note that we have $n + 1$ choices for k , so in fact we have a family of $n + 1$ coordinate transformations.

As we already noted in [14, Eq. 19], we have the convenient equivalence

$$\mathcal{H}(\mu, \nu) = \max_k \|\theta_k(\mu) - \theta_k(\nu)\|_{\ell^\infty} = \max_{i,k} (\theta_k^i(\mu) - \theta_k^i(\nu)), \quad (31)$$

where by ℓ^∞ we denote the standard supremum norm between vectors.

Remark 10. It is informative to compare this with Proposition 3.6. We fix $k = 0$ for simplicity, and ρ as the counting measure on E . Writing $\mathbf{1} \in \mathbb{R}^n$ for the vector of ones, we then have

$$\begin{aligned} \theta_0(\mu) &= (\log(\mu^1/\mu^0), \log(\mu^2/\mu^0), \dots, \log(\mu^n/\mu^0)) \\ &= (\log(\mu^1), \log(\mu^2), \dots, \log(\mu^n)) - \log(\mu^0)\mathbf{1}. \end{aligned}$$

On the other hand, in the notation of Proposition 3.6, we have the equivalence class

$$\theta(\mu) = \left\{ (\log(\mu^0), \log(\mu^1), \dots, \log(\mu^n)) + c\mathbf{1}; c \in \mathbb{R} \right\} \in \Theta_\rho \cong \mathbb{R}^{n+1} / \sim_{\text{const}}.$$

Of course, we can identify $\theta_0(\mu)$ with $(0, \theta_0(\mu)) \in \theta(\mu)$. Therefore, we see that our θ_0 -coordinates (29) simply choose the representative element in $\theta(\mu)$ with $c = -\log(\mu^0)$, or equivalently, the (unique) element with 0 in the first entry. Consequently, Proposition 3.6 gives the representation of the metric (in θ_0 -coordinates)

$$\mathcal{H}(\mu, \nu) = \left(\max_{i \neq 0} \{\theta_0^i(\mu) - \theta_0^i(\nu)\} \right)^+ + \left(\min_{i \neq 0} \{\theta_0^i(\mu) - \theta_0^i(\nu)\} \right)^-,$$

with $x^+ = \max\{0, x\}$ and $x^- = \max\{0, -x\}$. As this representation shows the Hilbert metric is given by a norm in θ_k -coordinates (for any k), we know that translation in θ_k -coordinates will not change the size or shape of a ball.

The θ_k -coordinates allow us to investigate in detail the shape of Hilbert balls in \mathcal{S}^n . The main idea is as follows: let $\mu, \nu \in \mathring{\mathcal{S}}^n$, so μ and ν are equivalent, and for all $k \in N$ consider the transformations $\mu \mapsto \theta_k(\mu)$ and $\nu \mapsto \theta_k(\nu)$. Let $\mathcal{H}(\mu, \nu) = R < \infty$. Fixing ν and $k = 0$, we prove that the \mathcal{H} -ball of radius R around $\theta_0(\nu)$ is a convex polytope $\mathcal{C} \subset \mathbb{R}^n$. Mapping \mathcal{C} to the simplex \mathcal{S}^n , we find that the image of \mathcal{C} through the inverse transformation θ_0^{-1} is also a convex polytope.

We start with the following lemma.

Lemma 4.3. *Consider two probability measures $\mu, \nu \in \mathring{\mathcal{S}}^n$ such that $\mathcal{H}(\mu, \nu) = R > 0$. Let $\theta_0(\mu), \theta_0(\nu) \in \mathbb{R}^n$ be their natural parameters under the mapping θ_0 given by (29). Then $\theta_0(\mu)$ belongs to the boundary $\partial\mathcal{C}$ of an n -dimensional convex polytope $\mathcal{C} \subset \mathbb{R}^n$ centred at $\theta_0(\nu)$, with $2(2^n - 1)$ vertices at the points*

$$v_{\mathcal{I}}^+ := \theta_0(\nu) + R \sum_{i \in \mathcal{I}} e_i, \quad v_{\mathcal{I}}^- := \theta_0(\nu) - R \sum_{i \in \mathcal{I}} e_i, \quad (32)$$

where $\{e_i\}_{i=1}^n$ denote the basis vectors of \mathbb{R}^n and $\mathcal{I} \subseteq \{1, 2, \dots, n\}$, $\mathcal{I} \neq \emptyset$.

Proof. The first thing we do, to simplify our calculations, is to translate $\theta_0(\nu) \in \mathbb{R}^n$ to the origin $\mathbf{0}$. Now let $\theta_k^i := \log(\mu^i/\mu^k) \in \mathbb{R}$ for all $i, k = 0, \dots, n$. We fix the coordinate system in \mathbb{R}^n to be given by $(x^1, \dots, x^n) \equiv (\theta_0^1, \dots, \theta_0^n)$, so that the basis vectors e_i are the unit vectors in the θ_0^i -direction. We look for a representation of the \mathcal{H} -ball of radius R around the origin in this coordinate system.

By (31), clearly $|\theta_k^i| \leq R$ for all pairs $(i, k) \in N \times N$. We consider all these inequalities, noting that, since $|\theta_k^i| = |\theta_i^k|$ by properties of log, we can avoid needless repetitions by restricting our consideration to all pairs of indices $(i, k) \in I^n := \{(i, k) \in N \times N : i > k\}$. Then we have in total $n(n+1)/2$ unique inequalities. Recalling that $\theta_k^i = \theta_0^i - \theta_0^k$, for $(i, k) \in I^n$ we define the $(n-1)$ -dimensional hyperplanes

$$h_{i,k}^\pm := \begin{cases} \{(x^1, \dots, x^n) \in \mathbb{R}^n : x^i - x^k = \pm R\} & k \neq 0, \\ \{(x^1, \dots, x^n) \in \mathbb{R}^n : x^i = \pm R\} & k = 0, \end{cases} \quad (33)$$

and denote by $p_{i,k}^+ := \{x \in \mathbb{R}^n : x^i - x^k \leq R\}$ the half-spaces bounded by $h_{i,k}^+$, and similarly by $p_{i,k}^- := \{x \in \mathbb{R}^n : x^i - x^k \geq -R\}$ those bounded by $h_{i,k}^-$ (and equivalently when $k = 0$).

We let $\mathcal{C}^n = \bigcap_{(i,k) \in I^n} p_{i,k}^+ \cap p_{i,k}^-$. By standard results in n -dimensional geometry, $\mathcal{C}^n \subset \mathbb{R}^n$ is a polyhedron, since it is the intersection of a finite number of closed half-spaces. We claim \mathcal{C}^n is bounded.

Note that the intersection $\mathcal{C}_0^n = \bigcap_{i=1}^n p_{i,0}^+ \cap p_{i,0}^-$ is the n -cube with side-length $2R$ centred at $\mathbf{0}$ with 2^n vertices at all possible positive/negative combinations of the coordinates $(\pm R, \dots, \pm R)$. Then

$$\mathcal{C}^n = \bigcap_{\substack{(i,k) \in I^n \\ k \neq 0}} p_{i,k}^+ \cap p_{i,k}^- \cap \mathcal{C}_0^n, \quad (34)$$

and the intersection of a hypercube with closed half-spaces is bounded, so \mathcal{C}^n is a bounded polyhedron, and therefore a convex polytope. (Note that \mathcal{C}^n is non-empty, since one can easily check that $\mathbf{0} \in \mathcal{C}^n$.)

We now would like to find the vertices of \mathcal{C}^n . We look for all the points in \mathbb{R}^n where exactly n of the hyperplanes (33) intersect uniquely.

Consider a linear system S given by n equations from (33). We say indices i, j are *linked* if an equation of the form $x^i - x^j = \pm R$ appears in the system S , and extend this definition by transitivity to partition the indices appearing in S into *linked classes*. We say an index i is a *base case* if $x_i = \pm R$ appears in S .

1. Consider a class not containing a base case. Then we can add a constant $r \in \mathbb{R}$ to each component x^i in the class without altering the equations of the form $x^i - x^j = \pm R$. Therefore the subsystem of S containing all the equations for this class cannot have a unique solution, so S cannot give a vertex.
2. For any class containing a base case, let's say x^i , note that $(x^i - x^j)/R \in \mathbb{Z}$ for any x^j linked to x^i . By transitivity and additive closure of \mathbb{Z} , we observe that all indices in a class containing a base case must have $x^j/R \in \mathbb{Z}$.

By combining the above observations, all vertices of \mathcal{C}^n must have coordinates that are integer multiples of R , with at least one coordinate given by a base-case, i.e. any point $v \in \mathbb{R}^n$ that solves uniquely S must be of the form (m_1R, \dots, m_nR) for $m_i \in \mathbb{Z}$ with $-n \leq m_i \leq n$ for all $i = 1, \dots, n$, and at least one $m_i \in \{\pm 1\}$.

Now, by (34) we must have that all the vertices of \mathcal{C}^n belong to \mathcal{C}_0^n . Thus, any point $v \in \mathbb{R}^n$ that uniquely solves S and is potentially a vertex of \mathcal{C}^n must be of the form (m_1R, \dots, m_nR) with $m_i \in \{-1, 0, 1\}$ for all $i = 1, \dots, n$. Moreover, assume that $m_i = 1$ and $m_k = -1$ for $i > k$. Then $m_iR - m_kR = 2R \geq R$, so $v \notin p_{i,k}^+$ and $v \notin \mathcal{C}^n$. Similarly, if $m_i = -1$ and $m_k = 1$ for $i > k$, then $m_iR - m_kR \leq -2R$ so $v \notin p_{i,k}^-$ and $v \notin \mathcal{C}^n$.

Thus we must have that any vertex of \mathcal{C}^n is of the form (m_1R, \dots, m_nR) with either $m_i \in \{-1, 0\}$ for all $i = 1, \dots, n$ or $m_i \in \{0, 1\}$ for all $i = 1, \dots, n$. Conversely, consider any point $v \in \mathbb{R}^n$ of this form. It is easy to construct a system S for a choice of n equations in $\{x^i - x^k = \pm R\} \cup \{x^j = \pm R\}$ such that v solves S . Then the points $(m_1R, \dots, m_nR) \in \mathbb{R}^n$ with either $m_i \in \{-1, 0\}$ for all $i = 1, \dots, n$ or $m_i \in \{0, 1\}$ for all $i = 1, \dots, n$ (and not all m_i identically 0) are in fact all the vertices of \mathcal{C}^n . Letting $\mathcal{C} = \mathcal{C}^n + \theta_0(\nu)$, we are done. \square

Remark 11. As can be deduced by (31), the \mathcal{H} -ball is, in a way, nothing but the intersection of $n+1$ skewed ℓ_∞ -balls, or, geometrically speaking, the intersection of $n+1$ skewed hypercubes. Recall the notation from our proof above. Define the following intersections

$$\mathcal{C}_k^n := \left[\bigcap_{j=k+1}^n p_{j,k}^+ \cap p_{j,k}^- \right] \cap \left[\bigcap_{j=0}^{k-1} p_{k,j}^+ \cap p_{k,j}^- \right], \quad \forall k = 1, \dots, n.$$

Then for each $k = 1, \dots, n$, \mathcal{C}_k^n is the image of the hypercube centred at $\theta_k(\nu)$ with side-length $2R$ under the linear transformation $\theta_k \mapsto \theta_0$.

We now map our convex polytope from \mathbb{R}^n to S^n through the inverse transformation θ_0^{-1} given by (30). Note that if θ_0^{-1} were an affine transformation, then Lemma 4.4 stated below would be trivially true. However, as θ_0^{-1} is not affine, a bit more work is required to prove that convexity, linearity of the boundary, and intersections are preserved. We give a depiction of this result in Figure 1.

Lemma 4.4. *The \mathcal{H} -ball of radius R around $\theta_0(\nu)$, given by the convex polytope \mathcal{C} of Lemma 4.2, maps to a convex polytope \mathcal{D} centred at ν in \hat{S}^n under the inverse transformation $\theta_0^{-1} : \mathbb{R}^n \rightarrow \hat{S}^n$. The vertices of \mathcal{D} are given by the images of the vertices of \mathcal{C} under the same transformation.*

Proof. We use the same notation as in Lemma 4.3. Recall that we centred our \mathcal{H} -ball at $\hat{\theta}_0 = \theta_0(\nu)$, the image of ν under the mapping $\theta_0 : \mathcal{S}^n \rightarrow \mathbb{R}^n$. Similarly, we will fix the centre of \mathcal{D} , the representation of the \mathcal{H} -ball in \mathcal{S}^n , at ν , and proceed as if ν were known.

We start by noting that the bounds $|\theta_k^i - \hat{\theta}_k^i| \leq R$, which correspond to the linear constraints (33), are equivalent to linear constraints in \mathcal{S}^n . Recalling the notation of the proof of Lemma 4.3, consider the half-spaces $p_{i,k}^+$ under the transformation $\theta_0^{-1} : \mathbb{R}^n \rightarrow \mathcal{S}^n$. We compute

$$\begin{aligned} \theta_0^{-1}(p_{i,k}^+) &= \{\theta_0^{-1}(\theta_0) \in \mathcal{S}^n : \theta_0^i - \theta_0^k \leq \hat{\theta}_0^i - \hat{\theta}_0^k + R\} \\ &= \left\{ \mu \in \mathcal{S}^n : \log \frac{\mu^i}{\mu^k} \leq \log \frac{\nu^i}{\nu^k} + R \right\} = \left\{ \mu \in \mathcal{S}^n : \mu^i - \mu^k \frac{\nu^i}{\nu^k} e^R \leq 0 \right\}, \end{aligned}$$

where we have used bijectivity of θ_0^{-1} and the fact that \exp is increasing. Similarly,

$$\theta_0^{-1}(p_{i,k}^-) = \left\{ \mu \in \mathcal{S}^n : \mu^i - \mu^k \frac{\nu^i}{\nu^k} e^{-R} \geq 0 \right\}.$$

Note that $\theta_0^{-1}(p_{i,k}^+)$ and $\theta_0^{-1}(p_{i,k}^-)$ are $(n-1)$ -dimensional flat subspaces of \mathcal{S}^n . In particular, since \mathcal{S}^n is a subset of an n -dimensional affine space $A \cong \mathbb{R}^n$, we see that $\theta_0^{-1}(p_{i,k}^+)$ and $\theta_0^{-1}(p_{i,k}^-)$ are closed half-spaces of A , bounded by the hyperplanes $\ell_{i,k}^+ := \{\mu \in A : \mu^i = \mu^k \frac{\nu^i}{\nu^k} e^R\}$ and $\ell_{i,k}^- := \{\mu \in A : \mu^i = \mu^k \frac{\nu^i}{\nu^k} e^{-R}\}$, which are the images (extended to A) of respectively $h_{i,k}^+$ and $h_{i,k}^-$ under θ_0^{-1} . Then, recalling that $I^n := \{(i, k) \in N \times N : i > k\}$, the intersection

$$\mathcal{D} := \bigcap_{(i,k) \in I^n} \theta_0^{-1}(p_{i,k}^+) \cap \theta_0^{-1}(p_{i,k}^-)$$

is a convex polyhedron in A , and in particular, since θ_0^{-1} is a bijection from \mathbb{R}^n into $\hat{\mathcal{S}}^n$,

$$\theta_0^{-1}(\mathcal{C}) = \theta_0^{-1} \left(\bigcap_{(i,k) \in I^n} p_{i,k}^+ \cap p_{i,k}^- \right) = \mathcal{D} \subset \mathcal{S}^n.$$

Finally, both boundedness of \mathcal{D} in \mathcal{S}^n (in the sense that \mathcal{D} is bounded away from $\partial\mathcal{S}^n$), so that \mathcal{D} is a convex polytope in \mathcal{S}^n , and the fact that vertices are preserved under the mapping follow easily from θ_0 being an homeomorphism between $\hat{\mathcal{S}}^n$ and \mathbb{R}^n . \square

Remark 12. Using Figure 1, we can build an intuition of how the transformation θ_0^{-1} deforms \mathcal{C} by considering what happens to the parallel pairs of hyperplanes $h_{i,k}^+$ and $h_{i,k}^-$ when mapped into \mathcal{S}^n . For each pair $(i, k) \in N \times N$, $\ell_{i,k}^+ = \theta_0^{-1}(h_{i,k}^+)$ and $\ell_{i,k}^- = \theta_0^{-1}(h_{i,k}^-)$ are not parallel in \mathcal{S}^n , but meet at the $(n-2)$ -face of the simplex given by $f_{i,k} = \{\mu \in \mathcal{S}^n : \mu^i = 0, \mu^k = 0\}$. In other words, the ‘point at infinity’ at which $h_{i,k}^+$ and $h_{i,k}^-$ meet in \mathbb{R}^n is mapped to the boundary of the simplex, and in particular to $f_{i,k}$, under θ_0^{-1} .

For example, in dimension 2, $f_{i,k}$ are vertices of \mathcal{S}^2 : when mapping $\mathcal{C} \in \mathbb{R}^2$ to \mathcal{S}^2 , we can think of squeezing together the ∞ -extremities of each pair of parallel lines $h_{i,k}^\pm$ (for $(i, k) \in \{(1, 0), (2, 0), (2, 1)\}$) so that they meet at an angle of $\alpha = a(e^R - e^{-R})/(1 + a^2)$, where $a = \nu^i/\nu^k$. Then we place the intersection point at the vertex $f_{i,k}$ of \mathcal{S}^2 , so that, intuitively, the strip of plane between $h_{i,k}^+$ and $h_{i,k}^-$ is mapped to a slice of \mathcal{S}^2 of width α bounded by $l_{i,k}^+$




Figure 1: On the left: representation in θ_0 -coordinates of a 2-dimensional \mathcal{H} -ball \mathcal{C} of radius R around $\hat{\theta}_0 = \theta_0(\nu) = (0, 0)$. On the right, the image of \mathcal{C} under θ_0^{-1} , which gives the \mathcal{H} -ball \mathcal{D} around $\nu = (\frac{1}{3}, \frac{1}{3}, \frac{1}{3})$ as a hexagonal polygon in the simplex \mathcal{S}^2 .

and $l_{i,k}^-$. Then it is easy to visualize how the straight lines that compose the boundary of \mathcal{C} are mapped to straight lines, and how intersections are preserved, making $\theta_0^{-1}(\mathcal{C})$ into a polytope as well. However, these lines (and those parallel to them) are in fact the only straight lines in θ_0 -coordinates that map to straight lines in \mathcal{S}^2 (as expected, since θ_0 and its inverse are nonlinear). We illustrate this in Figure 2 below.




Figure 2: Straight lines through the origin in θ_0 -coordinates on the left, and their images in \mathcal{S}^2 under the inverse mapping θ_0^{-1} on the right. In red the lines parallel to the axes and the diagonal in \mathbb{R}^2 , which remain straight in \mathcal{S}^2 .

Remark 13. The regularity of the \mathcal{H} -balls, when represented in θ_0 -coordinates, has other

surprising consequences³—for example, Hilbert balls of constant radius naturally tile the space, as illustrated in Figure 3.




Figure 3: Tiling of the 2-dimensional probability simplex \mathcal{S}^2 with Hilbert balls of radius 0.5, starting from the ball around the center $(\frac{1}{3}, \frac{1}{3}, \frac{1}{3})$ (corresponding to the origin in the θ_0 -coordinates on the left).

5 Metric-comparisons for probability measures: TV and \mathcal{H}

We now exploit the geometric intuition we gathered in the previous section to derive a bound, sharper than (17), for the total variation norm with respect to the Hilbert projective metric. We start by working with discrete probabilities in the simplex \mathcal{S}^n and then extend our result to probability measures on a general measurable space. Perhaps unsurprisingly, the distance $\mathcal{T}(\mu, \nu) = \tanh(\mathcal{H}(\mu, \nu)/4)$, which we defined in Section 2.1, plays a role once more in the computations below.

5.1 Probabilities on finite state-space

First of all, recall that for a measurable space (E, \mathcal{F}) the total variation distance (10) between two probability measures $\mu, \nu \in \mathcal{P}(E)$ is equivalent to

$$\|\mu - \nu\|_{\text{TV}} = 2 \sup_{A \in \mathcal{F}} |\mu(A) - \nu(A)|, \quad (35)$$

which, in the case of $E \cong \{0, \dots, n\}$ and $\mu, \nu \in \mathcal{S}^n$, reduces to

$$\|\mu - \nu\|_{\text{TV}} = \sum_{i=0}^n |\mu^i - \nu^i| = \|\mu - \nu\|_{\ell^1}. \quad (36)$$

³In a more artistic vein, Figure 3 also illustrates that the map $\theta_0 : \mathcal{S}^n \rightarrow \mathbb{R}^n$, for $n = 2$, corresponds to the classical transformation between parallel oblique perspective (θ_0 -coordinates) and three-point perspective (by viewing \mathcal{S}^n with its vertices at the three vanishing points), linking back well beyond Birkhoff (1957) and Hilbert (1895), at least as far as the work of Jean Pelegrin (Viator) in *De Artificiali Perspectiva* (1505).

Remark 14. Note that the factor of 2 in (35) is sometimes dropped, in which case (36) would be stated as $\|\mu - \nu\|_{\text{TV}} = \frac{1}{2}\|\mu - \nu\|_{\ell^1}$. We keep the factor of 2 in analogy with Atar and Zeitouni [3].

Theorem 5.1. *Given two probability measures $\mu, \nu \in \mathcal{S}^n$, we have that*

$$\|\mu - \nu\|_{\text{TV}} = \|\mu - \nu\|_{\ell^1} \leq 2 \tanh \frac{\mathcal{H}(\mu, \nu)}{4}. \quad (37)$$

Equivalently, we have

$$\sup_{A \subseteq \{0, \dots, n\}} \sum_{i \in A} |\mu^i - \nu^i| \leq \mathcal{T}(\mu, \nu). \quad (38)$$

From Lemma 4.4 we know that if $\mu, \nu \in \hat{\mathcal{S}}^n$ with $\mathcal{H}(\mu, \nu) = R < \infty$, then μ belongs to the boundary of a convex polytope $\mathcal{D} \in \mathcal{S}^n$, centred at ν and with vertices $\{\theta_0^{-1}(v_{\mathcal{I}}^+), \theta_0^{-1}(v_{\mathcal{I}}^-) : \mathcal{I} \subseteq \{1, \dots, n\}, \mathcal{I} \neq \emptyset\}$. Finding an upper bound for $\|\mu - \nu\|_{\ell^1}$ is now a simple convex optimization problem: we know the ℓ^1 -distance between ν and μ is maximized when μ is at one of the vertices of \mathcal{D} , so we compute the ℓ^1 -distance between ν and each of these vertices, and then maximize over the choice of vertex.

Lemma 5.2. *Assume $\nu \in \hat{\mathcal{S}}^n$ is known. If $\mathcal{H}(\mu, \nu) = R$, the ℓ^1 -distance between μ and ν is bounded by*

$$\|\mu - \nu\|_{\ell^1} \leq 2 \max_{\substack{\mathcal{I} \subseteq \{1, \dots, n\} \\ \mathcal{I} \neq \emptyset}} \left\{ \frac{\mathbf{S}_{\mathcal{I}}(1 - \mathbf{S}_{\mathcal{I}})(e^R - 1)}{1 + \mathbf{S}_{\mathcal{I}}(e^R - 1)} \vee \frac{\mathbf{S}_{\mathcal{I}}(1 - \mathbf{S}_{\mathcal{I}})(1 - e^{-R})}{1 + \mathbf{S}_{\mathcal{I}}(e^{-R} - 1)} \right\}, \quad (39)$$

where $\mathbf{S}_{\mathcal{I}} := \sum_{i \in \mathcal{I}} \nu^i$ for any subset \mathcal{I} (not \emptyset) of the indices.

Proof. Recall (32). Consider a vertex $\mathbf{v}_{\mathcal{I}^+} = \theta_0^{-1}(v_{\mathcal{I}}^+)$ of $\mathcal{D} \in \mathcal{S}^n$. Compute

$$\begin{aligned} \|\nu - \mathbf{v}_{\mathcal{I}^+}\|_{\ell^1} &= \sum_{i=0}^n |\nu^i - \mathbf{v}_{\mathcal{I}^+}^i| \\ &= \left| \nu^0 - \frac{1}{1 + \sum_{i \in \mathcal{I}} \exp\{\theta_0^i(\nu) + R\} + \sum_{i \in N_0 \setminus \mathcal{I}} \exp\{\theta_0^i(\nu)\}} \right| \\ &\quad + \sum_{k \in \mathcal{I}} \left| \nu^k - \frac{\exp\{\theta_0^k(\nu) + R\}}{1 + \sum_{i \in \mathcal{I}} \exp\{\theta_0^i(\nu) + R\} + \sum_{i \in N_0 \setminus \mathcal{I}} \exp\{\theta_0^i(\nu)\}} \right| \\ &\quad + \sum_{k \in N_0 \setminus \mathcal{I}} \left| \nu^k - \frac{\exp\{\theta_0^k(\nu)\}}{1 + \sum_{i \in \mathcal{I}} \exp\{\theta_0^i(\nu) + R\} + \sum_{i \in N_0 \setminus \mathcal{I}} \exp\{\theta_0^i(\nu)\}} \right| \\ &= \left| \nu^0 - \frac{1}{1 + e^R \sum_{i \in \mathcal{I}} \frac{\nu^i}{\nu^0} + \sum_{i \in N_0 \setminus \mathcal{I}} \frac{\nu^i}{\nu^0}} \right| + \sum_{k \in \mathcal{I}} \left| \nu^k - \frac{e^R \frac{\nu^k}{\nu^0}}{1 + e^R \sum_{i \in \mathcal{I}} \frac{\nu^i}{\nu^0} + \sum_{i \in N_0 \setminus \mathcal{I}} \frac{\nu^i}{\nu^0}} \right| \\ &\quad + \sum_{k \in N_0 \setminus \mathcal{I}} \left| \nu^k - \frac{\frac{\nu^k}{\nu^0}}{1 + e^R \sum_{i \in \mathcal{I}} \frac{\nu^i}{\nu^0} + \sum_{i \in N_0 \setminus \mathcal{I}} \frac{\nu^i}{\nu^0}} \right|. \end{aligned}$$

Letting $\mathbf{S}_{\mathcal{I}} := \sum_{i \in \mathcal{I}} \nu^i$ and $\check{\mathbf{S}}_{\mathcal{I}} := \sum_{i \in N_0 \setminus \mathcal{I}} \nu^i$, so that $\mathbf{S}_{\mathcal{I}} + \check{\mathbf{S}}_{\mathcal{I}} + \nu^0 = 1$, some algebra yields

$$\|\nu - \mathbf{v}_{\mathcal{I}^+}\|_{\ell^1} = 2(e^R - 1) \frac{\mathbf{S}_{\mathcal{I}}(1 - \mathbf{S}_{\mathcal{I}})}{1 + \mathbf{S}_{\mathcal{I}} e^R - \mathbf{S}_{\mathcal{I}}} =: g_R^+(\mathbf{S}_{\mathcal{I}})$$

Equivalently, if we let $\mathbf{v}_{\mathcal{I}^-} = \theta_0^{-1}(v_{\mathcal{I}^-})$, we obtain

$$\|\nu - \mathbf{v}_{\mathcal{I}^-}\|_{\ell^1} = 2(1 - e^{-R}) \frac{\mathbf{S}_{\mathcal{I}}(1 - \mathbf{S}_{\mathcal{I}})}{1 + \mathbf{S}_{\mathcal{I}} e^{-R} - \mathbf{S}_{\mathcal{I}}} =: g_R^-(\mathbf{S}_{\mathcal{I}})$$

Then the ℓ^1 -distance between μ and ν is bounded by the maximum between $\|\nu - \mathbf{v}_{\mathcal{I}^+}\|_{\ell^1}$ and $\|\nu - \mathbf{v}_{\mathcal{I}^-}\|_{\ell^1}$ over all choices of vertices, which yields the lemma. \square

Proof of Theorem 5.1. Let $\mu, \nu \in \mathring{\mathcal{S}}^n$ be such that $\mathcal{H}(\mu, \nu) = R$. Recall the notation from the proof of Lemma 5.2. Note that $g_R^+(x)$ and $g_R^-(x)$ for $x \in [0, 1]$ are symmetric around $x = \frac{1}{2}$. By standard calculus, we find that the maximum of g_R^+ is attained at $x_+^* = \frac{1}{1+e^{R/2}}$; while g_R^- is maximized at $x_-^* = 1 - x_+^* = \frac{e^{R/2}}{1+e^{R/2}}$. Evaluating g_R^+ and g_R^- at their respective maximizers gives the upper bound

$$\begin{aligned} \|\nu - \mathbf{v}\|_{\ell^1} &\leq \max_{\substack{\mathcal{I} \subseteq \{1, \dots, n\} \\ \mathcal{I} \neq \emptyset}} \{ \|\nu - \mathbf{v}_{\mathcal{I}^+}\|_{L^1} \vee \|\nu - \mathbf{v}_{\mathcal{I}^-}\|_{L^1} \} \\ &\leq \max_{\nu \in \mathring{\mathcal{S}}^n} \max_{\substack{\mathcal{I} \subseteq \{1, \dots, n\} \\ \mathcal{I} \neq \emptyset}} \{ \|\nu - \mathbf{v}_{\mathcal{I}^+}\|_{L^1} \vee \|\nu - \mathbf{v}_{\mathcal{I}^-}\|_{L^1} \} \\ &\leq \max_{\mathbf{S}_{\mathcal{I}} \in [0, 1]} \{ g_R^+(\mathbf{S}_{\mathcal{I}}) \vee g_R^-(\mathbf{S}_{\mathcal{I}}) \} \\ &\leq 2 \tanh \frac{R}{4}. \end{aligned}$$

Finally, note that the statement is trivial if $\mu \approx \nu$. Moreover, if $\mu \sim \nu$ and $\mu, \nu \in \partial \mathcal{S}^n$, then they must belong to the same $(n-d)$ -face of \mathcal{S}^n , which is also a probability simplex \mathcal{S}^d with $1 \leq d < n$. In particular, $\mu, \nu \in \mathring{\mathcal{S}}^d$, which is the same as the case we considered originally, so we are done. \square

Remark 15. For low dimensions, Lemma 5.2 provides a way to compute a bound for $\|\mu - \nu\|_{\ell^1}$ which is tighter than Theorem 5.1. While this still holds true in higher dimensions, the improvement gained by computing explicitly the bound (39) instead of using (37) can be negligible, since it is likely that at least one of the possible combinations for $\mathbf{S}_{\mathcal{I}}$ comes very close to the maximizer.

5.2 Probabilities on a general measurable space

We now use the discrete result of Theorem 5.1 to give bounds for general probability measures.

Corollary 5.2.1. *Let (E, \mathcal{F}) be a measurable space. Consider $\mu, \nu \in \mathcal{P}(E)$. We have*

$$\frac{1}{2} \|\mu - \nu\|_{\text{TV}} = \sup_{A \in \mathcal{F}} |\mu(A) - \nu(A)| \leq \mathcal{T}(\mu, \nu).$$

Proof. Note that if $\mu \approx \nu$, then the statement follows trivially, so assume that $\mu \sim \nu$. By definition of the total variation distance (35), for all $n \in \mathbb{N}$ there exists a set $A_n \in \mathcal{F}$ such that

$$\frac{1}{2} \|\mu - \nu\|_{\text{TV}} - \frac{1}{n} \leq |\mu(A_n) - \nu(A_n)| \leq \frac{1}{2} \|\mu - \nu\|_{\text{TV}}. \quad (40)$$

Let \mathcal{F}_n be the σ -algebra generated by A_n , i.e. $\mathcal{F}_n = \{A_n, A_n^c, E, \emptyset\}$, and let π_n be the partition of E given by $\pi_n = \{A_n, A_n^c\}$. Consider the probability measures μ_n, ν_n on the space (E, \mathcal{F}_n) given by

$$\mu_n = \mu|_{\pi_n}, \quad \nu_n = \nu|_{\pi_n}.$$

Then μ_n and ν_n are probabilities on the finite state space $\{A_n, A_n^c\}$, and in particular $\mu_n, \nu_n \in \mathcal{S}^1$. Therefore it holds that

$$\begin{aligned} \sup_{A \in \mathcal{F}_n} |\mu_n(A) - \nu_n(A)| &= |\mu_n(A_n) - \nu_n(A_n)| \\ &\leq \tanh \frac{\mathcal{H}(\mu_n, \nu_n)}{4} = \tanh \frac{\left| \log \frac{\mu_n(A_n)}{\nu_n(A_n)} - \log \frac{\mu_n(A_n^c)}{\nu_n(A_n^c)} \right|}{4} \\ &\leq \tanh \frac{\sup_{A, B \in \mathcal{F}} \left(\log \frac{\mu(A)}{\nu(A)} - \log \frac{\mu(B)}{\nu(B)} \right)}{4} = \tanh \frac{\mathcal{H}(\mu, \nu)}{4}. \end{aligned}$$

As for the left-hand side, we have that

$$\sup_{A \in \mathcal{F}_n} |\mu_n(A) - \nu_n(A)| = |\mu_n(A_n) - \nu_n(A_n)| = |\mu(A_n) - \nu(A_n)| \leq \sup_{A \in \mathcal{F}} |\mu(A) - \nu(A)|.$$

But by (40) we also have that

$$\sup_{A \in \mathcal{F}_n} |\mu_n(A) - \nu_n(A)| = |\mu_n(A_n) - \nu_n(A_n)| = |\mu(A_n) - \nu(A_n)| \geq \sup_{A \in \mathcal{F}} |\mu(A) - \nu(A)| - \frac{1}{n}.$$

Therefore, putting together the two above inequalities, we arrive at the final expression

$$\lim_{n \rightarrow \infty} \sup_{A \in \mathcal{F}_n} |\mu_n(A) - \nu_n(A)| = \sup_{A \in \mathcal{F}} |\mu(A) - \nu(A)|,$$

and the result follows. \square

Remark 16. Note that the bounds in Theorem 5.1 and Corollary 5.2.1 can be attained, and therefore are sharp. In particular, $2\mathcal{T}$ is nothing but the maximum TV (or ℓ^1) norm between two probability measures which are a fixed \mathcal{H} -distance apart. Theorem 2.6 then tells us that this quantity contracts under (positive) linear transformations.

Remark 17. For $\mu, \nu \in \mathcal{P}(E)$, we can also find an upper bound for $\mathcal{T}(\mu, \nu)$ in terms of TV. For all $A \in \mathcal{F}$, we have

$$\tanh \left(\frac{1}{2} \log \frac{\mu(A)}{\nu(A)} \right) = \frac{\mu(A) - \nu(A)}{\mu(A) + \nu(A)},$$

therefore

$$\mathcal{T}(\mu, \nu) \leq \tanh \left(\frac{1}{2} \log \left(\sup_{A \in \mathcal{F}} \frac{\mu(A)}{\nu(A)} \vee \sup_{B \in \mathcal{F}} \frac{\mu(B)}{\nu(B)} \right) \right) \leq \frac{\sup_{A \in \mathcal{F}} |\mu(A) - \nu(A)|}{2(\inf_{B \in \mathcal{F}} \mu(B) \wedge \inf_{B \in \mathcal{F}} \nu(B))}. \quad (41)$$

However, we note that the denominator of the right hand side may become arbitrarily small, which is unsurprising given \mathcal{T} is a stronger metric than TV on S^n .

Acknowledgements The research of EF was supported by the EPSRC under the award EP/L015811/1. SC acknowledges the support of the UKRI Prosperity Partnership Scheme (FAIR) under EPSRC Grant EP/V056883/1, the Alan Turing Institute and the Office for National Statistics (ONS), and the Oxford–Man Institute for Quantitative Finance.

References

- [1] S. Amari. *Differential-geometrical methods in statistics*, volume 28 of *Lecture Notes in Statistics*. Springer-Verlag, New York, 1985.
- [2] S. Amari. *Information geometry and its applications*, volume 194 of *Applied Mathematical Sciences*. Springer, Tokyo, 2016.
- [3] R. Atar and O. Zeitouni. Exponential stability for nonlinear filtering. *Ann. Inst. H. Poincaré Probab. Statist.*, 33(6):697–725, 1997.
- [4] R. Atar and O. Zeitouni. Lyapunov exponents for finite state nonlinear filtering. *SIAM J. Control Optim.*, 35(1):36–55, 1997.
- [5] P. Baxendale, P. Chigansky, and R. Liptser. Asymptotic stability of the Wonham filter: ergodic and nonergodic signals. *SIAM J. Control Optim.*, 43(2):643–669, 2004.
- [6] Y. Benoist. Convexes hyperboliques et fonctions quasisymétriques. *Publ. Math. Inst. Hautes Études Sci.*, (97):181–237, 2003.
- [7] G. Birkhoff. Extensions of Jentzsch’s theorem. *Trans. Amer. Math. Soc.*, 85:219–227, 1957.
- [8] G. Birkhoff. *Lattice theory*. American Mathematical Society Colloquium Publications, Vol. XXV. American Mathematical Society, Providence, R.I., third edition, 1967.
- [9] V. I. Bogachev. *Measure theory. Vol. I, II*. Springer-Verlag, Berlin, 2007.
- [10] P. Brémaud. *Discrete probability models and methods*. Springer, 2017.
- [11] P. J. Bushell. Hilbert’s metric and positive contraction mappings in a Banach space. *Arch. Rational Mech. Anal.*, 52:330–338, 1973.
- [12] Y. Chen, T. Georgiou, and M. Pavon. Entropic and displacement interpolation: a computational approach using the Hilbert metric. *SIAM J. Appl. Math.*, 76(6):2375–2396, 2016.
- [13] Ş. Cobzaş, R. Miculescu, and A. Nicolae. *Lipschitz functions*, volume 2241 of *Lecture Notes in Mathematics*. Springer, Cham, 2019.
- [14] S. N. Cohen and E. Fausti. Exponential contractions and robustness for approximate Wonham filters. *arXiv preprint arXiv:2305.02256*, 2023.
- [15] J. B. Conway. *A course in functional analysis*, volume 96 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1990.
- [16] P. de la Harpe. On Hilbert’s metric for simplices. In *Geometric group theory, Vol. 1 (Sussex, 1991)*, volume 181 of *London Math. Soc. Lecture Note Ser.*, pages 97–119. Cambridge Univ. Press, Cambridge, 1993.
- [17] L. Dubois. Projective metrics and contraction principles for complex cones. *J. Lond. Math. Soc. (2)*, 79(3):719–737, 2009.

- [18] S.N. Ethier and T.G. Kurtz. *Markov processes: characterization and convergence*. John Wiley & Sons, 2009.
- [19] Joel Franklin and Jens Lorenz. On the scaling of multidimensional matrices. *Linear Algebra Appl.*, 114/115:717–735, 1989.
- [20] D. Hilbert. Ueber die gerade linie als kürzeste verbindung zweier punkte. *Mathematische Annalen*, (46):91–96, 1895.
- [21] E. Hopf. An inequality for positive linear integral operators. *J. Math. Mech.*, 12:683–692, 1963.
- [22] J. Jacod and A.N. Shiryaev. *Limit theorems for stochastic processes*. Springer, 2nd edition, 2013.
- [23] R. Jentzsch. Über Integralgleichungen mit positivem Kern. *J. Reine Angew. Math.*, 141:235–244, 1912.
- [24] O. Kallenberg. *Foundations of modern probability*. Springer, 1997.
- [25] E. Kohlberg and J. W. Pratt. The contraction mapping approach to the Perron-Frobenius theory: why Hilbert’s metric? *Math. Oper. Res.*, 7(2):198–210, 1982.
- [26] M. G. Kreĭn and M. A. Rutman. Linear operators leaving invariant a cone in a Banach space. *Uspehi Matem. Nauk (N.S.)*, 3(1(23)):3–95, 1948.
- [27] F. Le Gland and L. Mevel. Basic properties of the projective product with application to products of column-allowable nonnegative matrices. *Math. Control Signals Systems*, 13(1):41–62, 2000.
- [28] F. Le Gland and L. Mevel. Exponential forgetting and geometric ergodicity in hidden Markov models. *Math. Control Signals Systems*, 13(1):63–93, 2000.
- [29] F. Le Gland and N. Oudjane. Stability and uniform approximation of nonlinear filters using the Hilbert metric and application to particle filters. *Ann. Appl. Probab.*, 14(1):144–187, 2004.
- [30] B. Lemmens and R. Nussbaum. *Nonlinear Perron-Frobenius theory*, volume 189 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2012.
- [31] S.P. Meyn and R.L. Tweedie. *Markov chains and stochastic stability*. Cambridge, 2nd edition, 2009.
- [32] F. Nielsen and K. Sun. Non-linear embeddings in hilbert simplex geometry. *arXiv preprint arXiv:2203.11434*, 2022.
- [33] B. B. Phadke. A triangular world with hexagonal circles. *Geometriae Dedicata*, 3:511–520, 1974/75.
- [34] H. H. Rugh. Cones and gauges in complex spaces: spectral gaps and complex Perron-Frobenius theory. *Ann. of Math. (2)*, 171(3):1707–1752, 2010.
- [35] H. Samelson. On the Perron-Frobenius theorem. *Michigan Math. J.*, 4:57–59, 1957.

- [36] E. Seneta. *Non-negative matrices and Markov chains*. Springer Series in Statistics. Springer, New York, 2006. Revised reprint of the second (1981) edition [Springer-Verlag, New York; MR0719544].
- [37] E. Seneta and S. Sheridan. Strong ergodicity of nonnegative matrix products. *Linear Algebra Appl.*, 37:277–292, 1981.
- [38] A. C. Thompson. On certain contraction mappings in a partially ordered vector space. *Proc. Amer. Math. Soc.*, 14:438–443, 1963.