Binomial expansion for a linearized map with memory. A final comment on: "Criticality and the fractal structure of -5/3 turbulent cascades"

Juan S. Medina-Alvarez^{a,b,*}

September 7, 2023

^aDeeptikus Ltd., 10A Tasman Ave., Mount Albert, Auckland 1025, New Zealand

^bAvda. Rafael Cabrera 10, 2ºM, Las Palmas de Gran Canaria 35002, Spain
* *E-mail address*: tlazcala@yahoo.es

Keywords

fractals, nonlinear, stochastic, maps, complex, Fibonacci sequence

Abstract

In a recent paper by Cabrera et al. [*Chaos, Solitons and Fractals* 2021;146:110876], a linearization of DRM differences equation, (Delayed Regulation Model), has been proposed as a scheme to explain transfer of energy through different scales in turbulence. They claim that this apparently simple model, by replication of Kolmogorov power law of $k^{-5/3}$ scaling, remarks a key mechanism of behaviour for more complex systems. Their proposal requires computation of several products of random

matrices, nevertheless they only offer an onset of time evolution or an asymptotic approximation to all them. Also it is suggested a fractal nature in the process of calculating the successive characteristic polynomials of these products or the eigenvalues of their associated self-adjoint matrices. Both questions are addressed in this comment and are answered positively. A general formula for the evolution in every step of mentioned stochastic linear approximation to DRM is found as well a map from a binomial expansion of these matrices key products to a well described fractal object.

1 Introduction

The Delayed Regulation Model (DRM) [2], a well known workbench of population dynamics with delay, have been proposed to replicate the transfer of energy in multi-scale cascades as described in the Kolmogorov's -5/3 power spectrum model of turbulence [1]. Some of the exceptional characteristics of this equation in differences, as the occurrence of the limit cycles and the inner chaotic dynamics the populations near to such geometric locus suffer [3], are attributable to the quadratic and discrete nature of the equation. However as emergence of those behaviours is controlled with parameter r in equation

$$x_{g+1} = rx_g(1 - x_{g-1}) \tag{1}$$

 $g=0,1,2,\ldots,+\infty, x\in[0,1]$, it is required to reach certain value of it to qualitatively change such conducts which range from quenching around a stable point to circulate around a cycle [3]. And once the threshold for the existence of a limit set has been trespassed the role of control parameter becomes the habitual of a logistic equation [2]. It will determine duplications of period and transition to chaos on variable x_g . This could be looked as the linear part of equation, -describing the whole time-evolution near an unstable fixed point-, losing dynamic significance with respect to the quadratic part. It would simply attest how and at what strength population points in configuration space are injected towards the limit set. All that because period bifurcation subtleties are much more associated with the non-linear part.

Precisely Cabrera et al. [1] seem to set out from the opposite as they do not attribute a major role in the transfer of energy through different scales in the turbulence to the nonlinear part of map (1). The statistics of variable x_g in the vicinity of a fixed point is equated in their study to the recounted energy transport. (See reference [1]). They assign great significance to the linear part of a modified version of DRM in which parameter r is subjected to a random variation. The differences equation now it is read

$$x_{g+1} = r_g x_g (1 - x_{g-1}), (2)$$

where $r_g = b + av_g$, with b > 1, $a \ge 0$ real numbers, and $v_g \in [0, 1]$ is a random variable distributed uniformly. After linearization of eq. (2) around fixed point¹ of eq. (1), -this is with $r = \langle r_g \rangle \equiv 1/(1 - \alpha)$, the average value of stochastic control parameter-, a linear affine equation in differences for a two dimensional vectorial space $E \simeq \mathbb{R}^2$ is obtained:

$$\vec{X}_{g+1} = \boldsymbol{A}_{\boldsymbol{g}} \vec{X}_g + \vec{B}_g, \tag{3}$$

¹Indeed Cabrera *et al.* [1] have linearized around the other fixed point of equation (1) in plane \mathbb{R}^2 , this is the origin (0,0). Nevertheless this choice is irrelevant for our description as the equations have the same form and described dynamics is the same. Our option makes the affine part of equation (3) less significative as its mean value is zero remarking so the homothetic part of it.

where

$$\boldsymbol{A}_{\boldsymbol{g}} = \left(\begin{array}{cc} r_g(1-\alpha) & -r_g\alpha \\ 1 & 0 \end{array} \right), \ \vec{B}_g = \left(\begin{array}{cc} r_g(1-\alpha)\alpha - \alpha \\ 0 \end{array} \right)$$

which are called the homothetic matrix and the affine vector respectively. Besides at the beginning of the series some initial conditions vector \vec{X}_0 should be picked up preferably in a neighborhood of $(\alpha, \alpha)^T$, the unstable after Hopf's bifurcation fixed point of eq.(1). In this way its components, and equation's evolution will start isotropically in a disc centered in (0,0), now the new coordinates of mentioned reference point after proper shift. Finally initial conditions, near or not to zero, will take form

$$\vec{X}_0 = \begin{pmatrix} x_0 - \alpha \\ y_0 - \alpha \end{pmatrix}, \quad x_0, y_0 \in [0, 1].$$

Iterating formula (3) for $g \ge 0$, it is easy to see evolution of \vec{X}_{g+1} via expression

$$\vec{X}_{g+1} = P_{g,-1}\vec{X}_0 + \sum_{j=0}^{g} P_{g,j}\vec{B}_j$$
 (4)

being operators $P_{g,j}$, j = -1, 0, ..., g the following descending in index ordered products of matrices $\{A_i\}_{i=0,1,...,g}$:

$$P_{g,-1} = \prod_{i=0}^{g} A_{g-i}$$

$$P_{g,j} = \prod_{i=0}^{g-j-1} A_{g-i}, \ j = 0, 1, \dots, g-1$$

 $P_{g,g} = \prod_{i=0}^{-1} A_{g-i} \equiv 1.$

In this case eq. (4) will give rise to a very complicate formula with products that are not reducible to a simple one operation of diagonalized matrices as we don't posses a common base for all of them. That is the result of introducing a random value in every matrix A_i . Everything we can aspire to is obtaining a significantly close bound to vector \vec{X}_{g+1} or simulate several times its evolution through multiple and different realizations $\{r_0, \ldots, r_g\}$. Both cases, though, will require an accurate knowledge of self-adjoint matrices $M_{g,j} = P_{g,j}^{\dagger}P_{g,j}$ [1] to make a correct exposition of the evolution of \vec{X}_0 with the linear equation (3) since they will lavishly appear in evaluation of $||\vec{X}_{g+1}||_2$.

Upon reaching this state of affairs, Cabrera et al. [1] made a couple of insightful observations. First of them that traces of matrices $M_{g,j}$ are dominant as to determine their greatest eigenvalues, being the latters part and parcel of the bound we are looking for. Second that traces on generation g, depending on coefficients of A_i 's as well as on stochastic parameters r_i 's, are deductible through a recurrence relation of previous findings of the parameters over generations $i = 0, 1, \ldots, g - 1$. (See equations 49-50 in reference [1]). Furthermore they put forward a step down tree of relations among coefficients of polynomials in $\{r_i\}_{i=0,\ldots,g-1}$ constituting such traces, and that gives account of deleting and making of coefficients from each level to the next of series given by eq. (4). Also they hinted for a fractal structure subjacent to this recursion or tree but no clue about dimension was expressed. Interesting and significative these questions as they are, were answered with a combination of partial simulations and approximations that rely too heavily on intuition as formal calculations were stopped in a not so far step, g = 5, and sets of stochastic realizations were not fully developed [1].

We will address in following section the issue of writing mathematically as far as possible matrices $M_{g,j}$, in hope that such formulation will give an efficient tool to answer more precisely the former results and descriptive intuitions.

2 Product of matrices A_i

The evaluation of how fast \vec{X}_0 is sinking or sourcing, amidst noise jolts, from the fixed point $(\alpha, \alpha)^T$ along generations $g = 0, 1, \ldots$ would be quite directly estimated by mean of euclidean distance $||\vec{X}_{g+1}||_2$. Unfortunately due to the random nature of equation (3) a simple formula yielding a precise number it is not possible, we must be settled with a statistical distribution of points or an upper bound to the temporal series of vectors. To achieve this last option we'll recur to some properties of norms as triangular inequality as well as definition of an operator's norm, -in this case the supreme of values attained over unit ball in E_{-} , and following expression shall be gotten

$$\|\vec{X}_{g+1}\|_{2} \leq \|\boldsymbol{P}_{g,-1}\|_{2} \|\vec{X}_{0}\|_{2} + \sum_{i=0}^{g} (\|\boldsymbol{P}_{g,i}\|_{2} \|\vec{B}_{i}\|_{2}).$$
(5)

As operator's P supreme norm definition is $||P||_2 \equiv \sup \frac{||P\vec{v}||_2}{||\vec{v}||_2}$, $\vec{0} \neq \vec{v} \in E$ [4] all our efforts will focus on establishing the eigenvalues of matrices $M_{g,j} \equiv P_{g,j}^{\dagger}P_{g,j}$. To this end a general formula for $P_{g,j}$, $j = -1, 0, \ldots, g$ should be deduced as a first step and that endeavour will start considering the definition of every A_i , -components of products $P_{g,j}$ -, split in two parts one fixed and the other associated to noise and its realizations. Every matrix A_i is

$$\boldsymbol{A_i} = \begin{pmatrix} r_i(1-\alpha) & -r_i\alpha \\ 1 & 0 \end{pmatrix} = s_i \begin{pmatrix} 1 & -\beta \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$
$$\equiv s_i \boldsymbol{M} + \boldsymbol{N}$$

with $s_i = r_i(1 - \alpha)$ and $\beta = \alpha/(1 - \alpha) \in (0, \infty)$. Consequently a general ordered in descending indexes product of matrices can be marked as

$$\begin{aligned} \boldsymbol{J_m} &\equiv \boldsymbol{J_m}(s_1, \dots, s_m) = \prod_{i=1}^m (s_i \boldsymbol{M} + \boldsymbol{N}) \\ &= (s_m \boldsymbol{M} + \boldsymbol{N}) \cdot (s_{m-1} \boldsymbol{M} + \boldsymbol{N}) \cdot \dots \cdot (s_1 \boldsymbol{M} + \boldsymbol{N}) \end{aligned}$$

staying any of previous operators as $P_{g,j} = J_{g-j}(r_{j+1}(1-\alpha), \dots, r_g(1-\alpha)),$ $j = -1, 0, \dots, g-1$ and $P_{g,g} \equiv \mathbf{1} \equiv J_0$ the identity matrix.

It is possible to write products J_m in a polynomial form by means of a binomial-like expansion and prove that this form contains any possible variation of two elements, M and N, taken mtimes. It is a simple proof, left to the reader as induction exercise, made easier when symbols 1 and 0 are arbitrary and respectively assigned to matrices M, N. In this manner such notation allows to index products $Y_m \cdot Y_{m-1} \cdot \ldots \cdot Y_1$, -where $Y_i = \{M, N\}$ -, by binary numbers of m digits. Namely

$$\boldsymbol{J_m} = \sum_{i=0}^{2^m-1} c_i(s_1,\ldots,s_m) \{\boldsymbol{Y_m} \cdot \boldsymbol{Y_{m-1}} \cdot \ldots \cdot \boldsymbol{Y_1}\}_i,$$

where $c_i(s_1,\ldots,s_m) = \prod_{k=1}^m t_{k,i}$, with

$$t_{k,i} = \left\{ \begin{array}{cc} s_k & & 1 \\ & , \text{ if k-th bit of i is} & \\ 1 & & 0 \end{array} \right\},$$

and

$$oldsymbol{Y_k} = \left\{ egin{array}{cc} oldsymbol{M} &, ext{ if k-th bit of i is } & 1 \ oldsymbol{N} & 0 \end{array}
ight\}$$

The summation J_m can be grouped in m + 1 sets of indexes *i* according to the number of zeroes the binary representation each one has. A particular set will

have then $\binom{m}{j}$ elements where now $j = 0, 1, \ldots, m$ denotes the number of zeroes of any index i which belongs to it. An additional partition of every one of these sets in four parts, for $m \ge 2$, can be done having in mind that both ends of any m-tuple which is a binary representation of some i has values 0 or 1. Writing f11, f01, f10 and f00 for the usually non void m-tuples outfits of respective type $(1, \ldots, 1), (0, \ldots, 1), (1, \ldots, 0)$ and $(0, \ldots, 0)$, it is obtained the following formula

$$J_{m} = \sum_{j=0}^{m} \{ \sum_{**\in\{11,01,10,00\}} \sum_{i\in f^{**}} c_{i} \cdot \{Y_{m} \cdot Y_{m-1} \cdot \ldots \cdot Y_{1}\}_{i} \}_{\#0's \text{ in } i=j}.$$
 (6)

3 Sorting the products $Y_m \cdot Y_{m-1} \cdot \ldots \cdot Y_1$

It may seem that the previous formula (6) is just one of many multiple possible outcomes after shuffling summand in J_m , but really is the proper grouping of products to reduce them to a minimum of calculations. From there on the complexity of determining and gathering explicitly summands of J_m will rely on how hard is to write down coefficients c_i 's. Yet to see that this classification of products $Y_m \cdot Y_{m-1} \cdot \ldots \cdot Y_1$ is really optimal, we must first delve into the behaviour of products by pairs of matrices M and N.

When naming $\mathbf{Q} \equiv \mathbf{M} \cdot \mathbf{N} = \begin{pmatrix} -\beta & 0 \\ 0 & 0 \end{pmatrix}$ and $\mathbf{R} \equiv \mathbf{N} \cdot \mathbf{M} = \begin{pmatrix} 0 & 0 \\ 1 & -\beta \end{pmatrix}$, we get a quartet of matrices, $\{\mathbf{M}, \mathbf{R}, \mathbf{Q}, \mathbf{N}\}$, that forms a linearly independent set in the four dimensional vectorial space of 2x2 matrices over reals, and in consequence any product $\mathbf{Y}_m \cdot \mathbf{Y}_{m-1} \cdot \ldots \cdot \mathbf{Y}_1$ will be a linear combination of them with real coefficients. Besides these four have a very nice property as they are a set closed under matrix product which will allow to set up simple recursion formulas. To this end we write the following table (1) for the matrix products

r*c	Q	M	N	R	
Q	$-eta oldsymbol{Q}$	$-\beta M$	0	0	
M	Q	M	Q	$-\beta M$	
N	$-\beta N$	R	0	0	
R	$-\beta N$	R	$-\beta N$	-eta R	

Table 1: Matrix products of the basis $\{M, R, Q, N\}$; rows multiplies columns by the left.

of the selected basis, it represents the multiplications in a row times column convention.

From table (1) are easily deduced by induction the following matrix equations

$$oldsymbol{N}^2 = oldsymbol{0} \implies oldsymbol{N}^n = oldsymbol{0}, \ n \ge 2,$$

 $oldsymbol{M}^2 = oldsymbol{M} \implies oldsymbol{M}^n = oldsymbol{M}, \ n \ge 2,$
 $oldsymbol{Q}^2 = -eta oldsymbol{Q} \implies oldsymbol{Q}^j = (-eta)^{j-1} oldsymbol{Q}, \ j \ge 1,$
 $oldsymbol{R}^2 = -eta oldsymbol{R} \implies oldsymbol{R}^j = (-eta)^{j-1} oldsymbol{R}, \ j \ge 1.$

This result will serve to a further reduction of every product $Y_m \cdot Y_{m-1} \cdot \ldots \cdot Y_1$. Each one is no other thing than an arbitrary succession of M's or N's that can be rephrased as an alternating product of powers of N's and M's, according to how many neighbours of same nature remain in a row separated by others of different nature. Being the total of multiplicands m the sum of all exponents for these clusters will amount naturally to this number. At this point a fork to classify all the products in types is obvious due to reductions implied in the previous table and derived subsequent equations. First alternative gives a class of products whose result is a Zero matrix, **0**. A particular $\prod_{k=0}^{m} Y_k$ will be in this one if any of the exponents of all its N^k powers is greater than one. The second option will comprise non null results and can be subdivided in four other classes. As each selection of m matrices now is an alternating array of M's and N's due to coalescence of every power of M to M itself, effectively four subtypes of arrays will be found depending on possibilities the ends of the product allow. These are a) $M \cdot N \cdot \ldots \cdot M$, b) $N \cdot M \cdot \ldots \cdot M$, c) $M \cdot N \cdot \ldots \cdot N$ and d) $N \cdot M \cdot \ldots \cdot N$.

Precisely the sets of binary indexes f11, f01, f10 and f00, at every stage of $j \leq m$ zeroes, represent products $Y_m \cdot Y_{m-1} \cdot \ldots \cdot Y_1$ that will respectively give arrays of type a), b), c) and d) after symbol redundancy is resolved. As long as, of course, no two consecutive zeroes can be found in the inspected index belonging to f * *. In what follows we will understand that teams of indices f11, f01, f10 and f00, at every level j, are already purged of those i whose binary representation have two or more adjacent 0's.

We are now in a position to calculate the product of matrices inside J_m classified into classes f11, f01, f10 or f00 and levels $0 \le j \le m$.

In the *j*-th stage of equation (6), where *j* denotes the number of zeroes of binary representation of indices *i* tagging products $\{Y_m \cdot Y_{m-1} \cdot \ldots \cdot Y_1\}_i$, we will have the following results for

a) $\underbrace{M \cdot N \cdot \ldots \cdot M \cdot N}_{j \text{ pairs}, j \ge 1} \cdot M = Q^j \cdot M = (-\beta)^{j-1} Q \cdot M = (-\beta)^{j-1} (-\beta) M =$

 $(-\beta)^{j}M$, if j = 0 no pair $M \cdot N$ would be present although the array being just M also fulfill $M = (-\beta)^{0}M$; result is valid then for $j \ge 0$,

- b) $\underbrace{N \cdot M \cdot \ldots \cdot N \cdot M}_{j \text{pairs}, j \ge 1} = \mathbf{R}^{j} = (-\beta)^{j-1} \mathbf{R}$, this time no j = 0 case is possible since array begins with N,
 - c) $\underbrace{\mathbf{M} \cdot \mathbf{N} \cdot \ldots \cdot \mathbf{M} \cdot \mathbf{N}}_{j \text{pairs}, j \ge 1} = \mathbf{Q}^{j} = (-\beta)^{j-1} \mathbf{Q}$, also case j = 0 is forbidden as \mathbf{N}

ends the sequence,

d)
$$\underbrace{N \cdot M \cdot \ldots \cdot N \cdot M}_{j-1 \text{ pairs}, j \ge 2} \cdot N = R^{j-1} \cdot N = (-\beta)^{j-2} R \cdot N = (-\beta)^{j-2} (-\beta) N = (-\beta)^{j-1} N, \quad j \ge 2 \text{ is necessary as array starts and ends by } N.^2$$

 $(-\beta)^{j-1}N, j \ge 2$ is necessary as array starts and ends by N.² ²Formula is also valid for j = 1 as $R^0 \cdot N = (-\beta)^0 N = N$, and is required to formally

3.1 Void f11, f01, f10, f00 populations

Except for those few cases in which the use of binomial coefficients implies a negative factorial, –a zero result then–, the original populations of sets f11, f01, f10, f00, at level j, contain respectively $\binom{m-2}{j}$, $\binom{m-2}{j-1}$, $\binom{m-2}{j-1}$, and $\binom{m-2}{j-2}$ elements, $(j \ge 2, m-j \ge 2)$. Nevertheless this quantities will be deploted by for large j.

and $\binom{m-2}{j-2}$ elements, $(j \ge 2, m-j \ge 2)$. Nevertheless this quantities will be depleted in a fractal look as greater values of m and j are considered due to the ruling out of binary sequences with adjacent zeros. Not being interested by now on the geometry of such decimation, which is a problem to pose in next sections, we just will count in next paragraphs those stages with too many zeros as to have no population at all in sets f * * since their wiping out makes the formula (6) clearer to write.

For example, we observe that if 2j > m the outfits f01 and f10 would never give rise to any non null product of type b) or c) since the number of matrices N is greater than M's and j pairs of $M \cdot N$ products, or $N \cdot M$, cannot be formed. This will lighten up the second summation in equation (6) from a particular index of all possible $j \in \{0, 1, ..., m\}$. With the intent of figuring it out we must discriminate two possible situations: m is even, (m = 2[m/2]), or m is odd, (m = 2[m/2] + 1), so elimination condition it will be read now

$$j > \left[\frac{m}{2}\right] + \left\{ \begin{array}{ll} 0 & \text{if } m \text{ is even} \\ \frac{1}{2} & \text{if } m \text{ is odd} \end{array} \right\}$$

though as j must be an entire number the latter simplifies to j > [m/2].

Also populations f_{11} and f_{00} , -which after simplification drive to corre-

extent this treatment to m = 1, the trivial case for J_m . In this situation only a) y d) sets, and j = 0, 1 levels, are present in eq. (6). In the nontrivial cases $m \ge 2$, when j = 0, 1, d) set is empty.

sponding product types a) and d)–, will be restricted based on the number of zeroes, j, in their binary representations.

In case $f00 \rightarrow d$) the number of zeroes minus one, j-1, which represents the number of matrices N paired to the right with at least one matrix M, must be less or equal to the number of ones, m-j, to exist. This is, case d) is obliterated from our accountancy if j-1 > m-j, and again a condition with multiples of j it is not suitable for use in equation (6), so 2j > m+1 will be expressed as

$$j > \left[\frac{m}{2}\right] + \left\{ \begin{array}{ll} 0 & \text{if } m \text{ is even} \\ 1 & \text{if } m \text{ is odd} \end{array} \right\}.$$

In case $f11 \rightarrow a$) the number of M's minus one must be at least equal to the number of N's to exist, although can be greater of course. This is $m - j - 1 \ge j$ and consequently case a) cannot be possible if m - 1 < 2j, or what is the same

$$j > \left[\frac{m}{2}\right] + \left\{ \begin{array}{cc} -1 & \text{if } m \text{ is even} \\ 0 & \text{if } m \text{ is odd} \end{array} \right\}.$$

The first summation in eq. (6) will be reduced from $\sum_{j=0}^{m}$ to $\sum_{j=0}^{\left[\frac{m}{2}\right]+1}$ for being null summands $j \ge [m/2] + 2$, (cases a), b), c) or d) are not possible from elements in any f * *). From the remaining summands we have already discussed that indices $2 \le j \le [m/2] - 1$ will raise to cases a), b), c) and d), yet j = 0only to a) and j = 1 to a), b), c) cases but not to d) one.³ And just as at the beginning of the count all cases are not present and it is required for summands fulfill $j \ge 2$ in order to contain the four cases, the end of summation is not abrupt at j = [m/2] + 1 either. Terms with j = [m/2] or [m/2] + 1 not always will have all four, yet depends on parity of m to know which ones survive.

If m is even and:

³It is an exception m = 2 with j = 1 as a) case is (1,1) and contains no zeroes.

• $j = \left[\frac{m}{2}\right]$, a) it does not ride out, but b), c) and d) do,

•
$$j = \left[\frac{m}{2}\right] + 1$$
, neither of a), b), c) or d) rides out.

If m is odd and:

- $j = \left\lceil \frac{m}{2} \right\rceil$, all cases, (a, b, c, d), survive,
- $j = \left[\frac{m}{2}\right] + 1$, a), b), and c) they don't ride out the cut but d) does.

This is all there is to consider in relationship to which product sequences $Y_m \cdot Y_{m-1} \cdot \ldots \cdot Y_1$ disappear due to nilpotency of N and the canonical types the survivors fall into. A closure in the characterization of J_m summands requires to write properly remaining coefficients $c_i(s_1, \ldots, s_m)$ in function of indices i and j and families f00, f01, f10, and f11 they belong to.

4 Coefficients $c_i(s_1, \ldots, s_m)$

We saw that c_i 's are productories with m terms $t_{k,i}$, (k = 1, ..., m), the latter being one two choices: the random variable s_k or 1 depending upon the binary representation of $i=0, ..., 2^m - 1$. At the moment the best description for them since all numbers i of m binary digits where included, nevertheless the suitable c_i 's for the equation (6) are obtained after two selective processes, a sort and a purging one. The first is a grouping of summands according to the number of zeroes, j, in the binary representation of i; the second is a overriding of those same sequences when two or more adjacent zeroes exist in them. As such it looks as convenient a change of notation in c_i 's to reflect all this and make an indexed use of them easier.

To this end let us define as auxiliary functions the products of consecutive variables x_s , $s \in \{1, \ldots, m\}$, specifically they shall be written in this guise $\pi_l^n(x_{l+1}, \ldots, x_{l+n}) = \prod_{k=1}^n x_{l+k}$, for $n \ge 1$, or $\pi_l^0 \equiv 1$, for n = 0. Also the binary representation of i will be additionally tagged again. Originally to a unique index i for each product of matrices was added a second index, j, due to operative reasons as shown in formula (6), however outfits f * * need also a more descriptive and possibly more efficient third system, since it will facilitate a systematic writing of those valid c_i coefficients.

Two additional marked positions, m+1 and 0, will be added to left and right of a binary m-tuple, $i \in \{0, \ldots, 2^m - 1\}$, with the purpose of accounting for loci of zeros and clustering of ones. As it is highlighted the number of zeros of each sequence, \neg or binary representation \neg , with number j, they will numbered with indices $k = 1, \ldots, j$ counting them from right to left, and their locations logged with indices l_k . In this way $1 \leq l_1 < \cdots < l_j \leq m$, and $l_0 \equiv 0$, $l_{j+1} \equiv m + 1$ always, what allows to write the number of ones between consecutive, and not necessarily adjacent, zeros as $n_k = l_{k+1} - l_k - 1$, with $0 \leq k \leq j$. Obviously the total number of 1's in the representation of i is obtained without ambiguity, *i.e.* $m - j = \sum_{k=0}^{j} n_k$, and with the zeroes as milestones, neither their positions are subjected to confusion. Now, \neg having in mind it has been abridged every $l_k(i)$ to l_k , we can rewrite all coefficients as

$$c_i(s_1,\ldots,s_m) = \prod_{k=0}^{j} \pi_{l_k}^{n_k}(s_{l_k+1},\ldots,s_{l_k+n_k}).$$

This wouldn't be a great change of notation but for we have previously rejected use of all c_i 's associated with null products of matrices, (this is *i*'s with adjacent 0's). Since from now on $n_k \ge 1$, for k = 1, 2, ..., j-1, the latter will be the most compact form of writing such coefficients for families f11, f01, f10 and f00, since it will involve a minimal use of functions $\pi_{l_k}^{n_k}$ containing random variables $\{s_1, ..., s_m\}$. Besides any padding 1's is excluded of $\prod_{k=1}^{j-1}$ unlike when use of symbols $t_{k,i}$ in c_i was habitual.

The described set of new indexes based on numbers m, j, and grouping of

ones $\{l_k\}$, as well as a forementioned conditions on n_{l_k} will be denoted with symbol

$$\mathcal{P}_{m,j} = \left\{ (l_j, \dots, l_1) | l_k \in \{1, \dots, m\}; \{n_k \ge 1\}_{k=1,\dots,j-1}; \sum_{k=0}^j n_k = m - j \right\}.$$

5 The binomial expansion of J_m

We rephrase eq. (6) for each level $j \in \{0, 1, \dots, [m/2]+1\}$ as follows, each element i from any of four outfits f * *, j will be substituted by the corresponding j-tuple of $\mathcal{P}_{m,j}$ as counting index in the inner summations. In this way

$$a, b, c, d) \sum_{\substack{i \in f^{**} \\ \# 0's \text{ in } i=j}} c_i \to \left\{ \sum_{\substack{l \in \mathcal{P}_{m,j} \\ \mathcal{C}^{**}(n_0, n_j)}} \prod_{k=0}^j \pi_{l_k}^{n_k}(s_{l_k+1}, \dots, s_{l_k+n_k}) \right\} \equiv p_{**,j},$$

where if a) **=11 then $C11(n_0, n_j) = \{n_0, n_j \ge 1\}$, if b) **=01 then $C01(n_0, n_j) = \{n_0 \ge 1, n_j = 0\}$, if c) ** = 10 then $C10(n_0, n_j) = \{n_0 = 0, n_j \ge 1\}$ and if d) **=00 then $C00(n_0, n_j) = \{n_0 = n_j = 0\}$.

Finally we obtain the coveted expression for equation

$$\boldsymbol{J_m} = \sum_{j=0}^{\left[\frac{m}{2}\right]+1} \left\{ p_{11,j}(-\beta)^j \boldsymbol{M} + p_{01,j}(-\beta)^{j-1} \boldsymbol{R} + p_{10,j}(-\beta)^{j-1} \boldsymbol{Q} + p_{00,j}(-\beta)^{j-1} \boldsymbol{N} \right\},$$

with $p_{01,j} = p_{10,j} \equiv 0$, for j = 0, and $p_{00,j} \equiv 0$ for j = 0 or j = 1. Also is possible to write again this formula by extracting common matricial factors out of summations, we will obtain then

$$J_{m}(\vec{s}_{m}) = g_{12}(\vec{s}_{m})M + g_{22}(\vec{s}_{m})R + g_{11}(\vec{s}_{m})Q + g_{21}(\vec{s}_{m})N,$$
(7)

r*c	Q	M	N	R	
Q^{\dagger}	$-eta oldsymbol{Q}$	-eta M	0	0	
M^\dagger	$-eta M^\dagger$	$M - \beta R$	0	0	
N^{\dagger}	0	0	$-rac{1}{eta} Q$	M	
R^{\dagger}	0	0	M^{\dagger}	$oldsymbol{M}-etaoldsymbol{R}$	

Table 2: Products of transposes of $\{M, R, Q, N\}$ by original ones; rows multiplies columns by the left.

where \vec{s}_m stands for (s_1, \ldots, s_m) . Obviously functions g_{xy} will be

$$g_{12}(\vec{s}_m) = \sum_{j=0}^{\left[\frac{m}{2}\right]+1} p_{11,j}(-\beta)^j,$$

$$g_{22}(\vec{s}_m) = \sum_{j=1}^{\left[\frac{m}{2}\right]+1} p_{01,j}(-\beta)^{j-1},$$
$$g_{11}(\vec{s}_m) = \sum_{j=1}^{\left[\frac{m}{2}\right]+1} p_{10,j}(-\beta)^{j-1},$$
$$g_{21}(\vec{s}_m) = \sum_{j=2}^{\left[\frac{m}{2}\right]+1} p_{00,j}(-\beta)^{j-1}.$$

6 Eigenvalues of $M_{g,j}$ and norm of $P_{g,j}$

Once the structure of J_m have been clarified it is straightforward to get $J_m^{\dagger} J_m$. We skip details here as the result pops up after easy and laborious calculations facilitated by inspection of table (2). The outcome it is read as

$$J_{m}^{\dagger}J_{m} = (-\beta g_{11}^{2} - \frac{1}{\beta}g_{21}^{2})Q + (-\beta g_{11}g_{12} + g_{22}g_{21})(M^{\dagger} + M) + (g_{12}^{2} + g_{22}^{2})(M - \beta R).$$
(8)

However as matrices and coefficients in previous formula both show a dependence in parameter β , it is tidier and more efficient operationally to split this formula in a scalar part depending on β and a vectorial one not doing so. We will resort to use three additional matrices to make this possible. They are $K = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, $L = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, $S = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$, and with them the matrices involved in $J_m^{\dagger} J_m$ are written as $Q = -\beta K$, $M^{\dagger} + M = 2K - \beta L$ and $M - \beta R = K - \beta L + \beta^2 S$. Introducing these identities in the equation, expanding parenthesis and rearranging all terms in function of matrices K, L and S a new version of eq. (8) is achieved

$$J_{m}^{\dagger}J_{m} = \left\{ (\beta g_{11} - g_{12})^{2} + (g_{21} + g_{22})^{2} \right\} K$$
$$+ \beta \left\{ g_{12}(\beta g_{11} - g_{12}) - g_{22}(g_{21} + g_{22}) \right\} L$$
$$+ \beta^{2} \left\{ g_{12}^{2} + g_{22}^{2} \right\} S. \quad (9)$$

That symmetric matrix is shorten to $J_m^{\dagger} J_m = \begin{pmatrix} h_{11} & h_{12} \\ h_{21} & h_{22} \end{pmatrix}_m$ to point out the functional look of its two real and positive eigenvalues

$$\lambda_{m,\pm} = \left(\frac{h_{11} + h_{22} \pm \sqrt{(h_{11} - h_{22})^2 + 4h_{12}h_{21}}}{2}\right)_m,\tag{10}$$

of which the greater is, by the definition given in section 2, the square of operator's, J_m , norm, (*i.e.* $||J_m||_2^2 = \lambda_{m,+}(\vec{s}_m)$ [4]).

Finally all requirements to describe dynamics of equation (4) end here, as formula (7) provides the elements needed to follow its evolution in time. However we have continued a little further in the search of an mathematical expression for $M_{g,j}$ as it allows for an unidimensional picture of the linearized DRM difference system. These matrices, we saw, are built in a natural way as self-adjoint operators derived from corresponding $P_{g,j}$'s, though. And in section 2 was shown that $P_{g,j} = J_{g-j}(\vec{s}_{g-j})$, with $s_k = r_{j+k}(1 - \alpha)$, $j = -1, 0, \ldots, g - 1$, and $k = 1, \ldots, g - j$, so accordingly to equation (5) the former eigenvalues in eq. (10), considered as functions of random variables, furnish all is needed for making up every one of the norms required to bound closely $||\vec{X}_{g+1}||_2$, once is given a realization $\{r_0, \ldots, r_g\}$.

7 Subjacent fractal distribution of matrices' products

The question now is to ascertain how many matrices' products $\prod_{k=1}^{m} \mathbf{Y}_{k}$ among those initial 2^m have survived after being purged by nilpotence of N, inasmuch as its answer will make easier and efficient the writing of partition sets $\mathcal{P}_{m,j}$ and consequently that of functions $g_{\{1,2\}\{1,2\}}(\vec{s}_m)$ in equation (7). To take one step back it is needed then, and all products null or ending in classes a) to d) must be again considered. Explicitly the index i attributed to each one will be written down $i = \sum_{k=0}^{m-1} i_k 2^k$, with $i_k \in \{0,1\}$, or in *m*-tuple form (i_{m-1},\ldots,i_1,i_0) , this latter is an ever growing set of indices, –increasing with the number of generations m tried, that is hard to apprehend in a geometric picture. Nevertheless it is always possible mapping all indexes to the finite interval $[0,1] \subset \mathbb{R}$ by mean of inversion and study the distribution of survivors. Henceforth at every generation the interval [0,1] will be divided in 2^m equal subintervals and they will be numbered following binary notation from $(0, \ldots, 0)$ to $(1, \ldots, 1)$, these will be associated to indices *i*'s as it was done before but this time reading the m-tuples in reverse bit order. Every index with this idea in mind will be paired with a subinterval of extent $1/2^m$ whose left end in a fractional binary representation is the new m-tuple. This is, if a matrix product was indexed with an $i \in \{0, \ldots, 2^m - 1\}$, as just described above now it will be assigned to subinterval starting at boundary post $I_i = \sum_{k=1}^m i_{k-1} 2^{-k}$. Such indexation has a tremendous advantage, since as generations run they are drawn in a stack and on it each subinterval at layer m-1 is split in two which will be put just beneath their parent in the next layer, also a consecutive numbering is hold among all members of the new generation m. Besides this method, or image, reflects exactly how products constituting J_{m-1} will sire those new elements of J_m , and how to neglect those they are null and record the survivors with a huge economy of means that avoids effectively count all the 2^m products.

The procedure as told contains all elements necessary for registering in a descending tree of decisions all cases. A simple two steps system is all what is needed to depict correctly the geometry of every layer as well as the limit set. We observe when m = 1 a simple division in two of segment [0, 1], subsegments are numbered 0 and 1. The next layer, m = 2, split each one of the previous in two and results are tagged as 00, 01, 10 and 11. This pattern is crucial since every two layers, from m even to m + 2, every subsegment will be split in four and these same mentioned tags will be added to the sequence of ones and zeroes each interval already has assigned. And this is the first hint of a fractal structure in the limit set $m \rightarrow \infty$.

The Cantor set is the result of deleting indefinitely the middle third at every turn the remains of interval [0, 1] which is that was started with. It has a fractal dimension of 0.631 and it is the classical example of self-similarity when these objects are introduced. We face here something lookalike though different. As we only allow chains of symbols with no consecutive zeroes one quarter is wiped out of present segments every two steps of duplicating segments and the process is iterated also indefinitely. Anyone can argue this is a bad and nonsymmetric copy of Cantor's set, but no argument against its fractality can be issued. Nevertheless there is still more, the three segments 01, 10 and 11 alive at step m once duplicated they become in 010, 011,100, 101, 110 and 111 at step m + 1 and five of them survive for an already explained further pruning

$\ln \mathcal{M}_{\delta}$	1	3	5	8	13	21	34	55	89	144
m	1	2	3	4	5	6	7	8	9	10

Table 3: Number of boxes \mathcal{M}_{δ} at division scale $\delta = 2^{-m} vs$. number of generations m in the splitting process of interval [0, 1].

at m + 2. As deduced from inspection after adding 0's or 1's to the binary fractional numbers which divide interval [0, 1] at each step, no other patterns of elimination of subintervals are visible. Always at each layer there are groupings of two or three neighboring intervals of scale 2^{-m} and after a splitting the first type of groupings sizes three subintervals of scale 2^{-m-1} and the second class five. These are then all the rules for characterizing the fractal we observe.

While these guidelines allow to describe a fractal, \mathcal{F} , in an iterative mode we simply can in a first instance to use the easy original rule of no "adjacent zeros" to examine and count filled boxes, –or not neglected intervals–, along a few steps m with intend of esteeming a fractal dimension which tells how many surviving intervals are found at each scale of division δ . The implicit model to account for content of a fractal would be $\mathcal{M}_{\delta}(\mathcal{F}) \sim \mathcal{C}\delta^{-s}$, where \mathcal{M}_{δ} is a function which answer how much *matter* of the object \mathcal{F} is found at scale δ and s is the fractal dimension of it. In our description of \mathcal{F} we have counted remaining intervals of length $\delta = 1/2^m$ after m layers of pruning, that is our \mathcal{M}_{δ} then. In this way of things a formula to work out dimension s is [5]

$$s = \lim_{\delta \to 0} \frac{\ln \mathcal{M}_{\delta}(\mathcal{F})}{-\ln \delta}.$$

In table (3) are data needed to justify the following regression results of formula $\ln \mathcal{M}_{\delta} \sim s(m \ln 2) + \ln \mathcal{C}$: correlation coefficient, $\rho = 0.9999229$, constant term, $\ln \mathcal{C} = 0.173 \pm 0.013$, slope $s = 0.691 \pm 0.003$. This ends our problem of counting matrices in J_m .

7.1 The Fibonacci Sequence

However a final remark must be added on the light of explained "2-3 grouping to 3-5 survivors" rule used for reproducing non void subintervals and a remarkable fact, shown in table (3), which is that the number of boxes recorded follows a Fibonacci sequence. With the final purpose of giving an homogeneous treatment to both kinds of groupings we start a description of their spawning at the level m = 3 where only the intervals 010, 011, -both contiguous-, and 101, 110, 111, -also neighbors-, have survived to the purge. This level is chosen as is the first time the mentioned groups of two or three subintervals are present together and henceforth they will be appearing persistently in the ongoing levels. In contrast when m = 0 is the whole interval [0, 1] which stands, level m = 1 have a unique grouping of two subintervals, and m = 2 consists of an array of three adjacent surviving intervals of length $\delta = 0.25$ each.

It is easy to see from our rule of non adjacent zeros that the set of two splits in three elements with endings already known and the set of three produces five intervals. A surprise comes now as those five are not contiguous ones but they form a group of two and a group of three whose endings, –last three ciphers–, are equal to those recently written so that the process of division repeats indefinitely and only groupings of two or three elements contribute to the total of surviving intervals. Hence a formula can be written, be $m \geq 3$, D_m the number of groups of two subintervals and T_m the number of groups of three elements so grand total will be

$$\#Boxes_m = 2 \times D_m + 3 \times T_m$$

and as a consequence of explained reproduction scheme on next level, m + 1, they will result in

$$\#Boxes_{m+1} = 2 \times T_m + 3 \times (D_m + T_m)$$

In this way coefficients fulfill a recurrence ratio

$$D_{3} = 0, T_{3} = 1,$$

$$\begin{cases} D_{m+1} = T_{m} \\ T_{m+1} = T_{m} + D_{m} \end{cases}, m \ge 3$$

which is no other than the recurrence which defines Fibonacci sequence though with index shifted by two. On the other hand we have just shown by a hand calculation that the number of non discarded subintervals at level m follows also a Fibonacci sequence shifted by minus two for $0 \le m \le 10$. If this pattern continued indefinitely the formula would be

$$F_{m+2} = 2 \times F_{m-2} + 3 \times F_{m-1}, \ m \ge 3, \tag{11}$$

being F_n , $n \ge 0$ the *n*-th term of Fibonacci sequence.

In conclusion if we can prove such formula the problem of counting filled boxes which defines the content at each level of fractal $\mathcal{F}_m \to \mathcal{F}$ and of working out its dimension would be solved.

7.1.1 A recurrence quadratic formula

Starting with the solutions to quadratic equation $x^2 - x - 1 = 0$ two independent sequences of powers of them can be built which satisfy the Fibonacci's recurrence $x_{\pm}^{n+1} = x_{\pm}^n + x_{\pm}^{n-1}$, $n \ge 0$. Neither of both will be sequences of entire numbers, nevertheless. Though as they are $x_{\pm} = (1 + \sqrt{5})/2 \equiv \phi$ and $x_{\pm} = (1 - \sqrt{5})/2 \equiv \psi$ some linear combination of powers is expected to yield entire numbers at every step, and indeed so it is. The well known class of linear combinations, (Binet's formula) [6],

$$F_n = \frac{\phi^n - \psi^n}{\phi - \psi}, \ n \ge 0,$$

fulfills the recurrence

$$F_{n+1} = F_n + F_{n-1}, \ n \ge 1,$$

and as $F_0=0$ and $F_1=1$ results in a sequence of natural numbers.

Once the general term is presented in this form it is quite easy to prove [6, 7]

$$F_l F_n + F_{l+1} F_{n+1} = F_{l+n+1}, \, l, n \ge 0.$$
(12)

Let's see it. On one side

$$F_{l}F_{n} + F_{l+1}F_{n+1} = \frac{(\phi^{l+n} - \psi^{l}\phi^{n} - \phi^{l}\psi^{n} + \psi^{l+n}) + (\phi^{l+n+2} - \psi^{l+1}\phi^{n+1} - \phi^{l+1}\psi^{n+1} + \psi^{l+n+2})}{(\phi - \psi)^{2}}$$
$$= \frac{\phi^{l+n}(1 + \phi^{2}) - (\psi^{l}\phi^{n} + \phi^{l}\psi^{n})(1 + \psi\phi) + \psi^{l+n}(1 + \psi^{2})}{(\phi - \psi)^{2}}$$
$$= \frac{\phi^{l+n}(1 + \phi^{2}) + \psi^{l+n}(1 + \psi^{2})}{(\phi - \psi)^{2}},$$

since $1 + \psi \phi = 0$, and we hold apart this in mind for a moment. On the other hand if we take into account that $\phi(1 + \psi^2) = (\phi - \psi) = -\psi(1 + \phi^2)$, we will get

$$F_{l+n+1} = \frac{(\phi - \psi)(\phi^{l+n+1} - \psi^{l+n+1})}{(\phi - \psi)^2}$$
$$= \frac{-\psi \phi^{l+n+1}(1 + \phi^2) - \phi \psi^{l+n+1}(1 + \psi^2)}{(\phi - \psi)^2}$$
$$= \frac{\phi^{l+n}(1 + \phi^2) + \psi^{l+n}(1 + \psi^2)}{(\phi - \psi)^2}.$$

As both expressions are the same one, identity (12) is proven.⁴ Recalling that $F_3 = 2$ and $F_4 = 3$ we can substitute l = 3 and n = m - 2 in formula (12) to

 $^{^{4}}$ Though not so well known, formula (12) has also been baptized by practitioners in the field, this time as Honsberger identity [6, 7]. Nevertheless this last one can be also be derived straightforwardly from another more familiar one, d'Ocagne identity. Although negative indices, not only positives, must be taken into consideration.

obtain formula (11).

7.2 Fractal dimension

Therefore it has been proved that the number of non-empty boxes of length δ = $1/2^m$ in \mathcal{F}_m , $m \geq 3$ is F_{m+2} according to equation (11) secured by the spawn of boxes present in step m - 1. (In fact, as a digression, we are compelled to acknowledge validity of formula (11) also for $0 \leq m < 3$ as a direct calculation shows.⁵) So we are in a position to work out precisely the already mentioned fractal dimension. Now taking into account that $\ln \mathcal{M}_{\delta}(\mathcal{F}) = \ln F_{m+2}$ a seamless recast of such quantity as $s(m \ln 2) + \ln \mathcal{C}$, embodies like

$$\ln F_{m+2} = \ln \phi^{m+2} + \ln(1 - \left(\frac{\psi}{\phi}\right)^{m+2}) - \ln(\phi - \psi)$$
$$= (m \ln 2) \left(\frac{\ln \phi + \frac{1}{m} \ln(1 - (\psi/\phi)^{m+2})}{\ln 2}\right) + \ln\left(\frac{1 + \phi}{\phi - \psi}\right).$$

And as it also happens that $|\psi/\phi| = |\frac{1-\sqrt{5}}{1+\sqrt{5}}| < 1$ results in $\lim_{m\to\infty} (1-(\psi/\phi)^{m+2})$ = 1, hence as expected a simple asymptotic formula is obtained

$$\mathcal{M}_{\delta}(\mathcal{F}) = F_{m+2} \sim \mathcal{C}\delta^{-s} \tag{13}$$

for $\delta \to 0$, (*i.e.* $m \to \infty$), which yield results $s = \frac{\ln \phi}{\ln 2} = 0.694241...$ and $C = \frac{1+\phi}{\phi-\psi} = 1.17082...$ as fractal dimension and content respectively.

We would like to finish our exposition of current problem with a remark highlighting how much information formula (13) conveys. Values of m not near of being suspicious for justifying formula as m = 0, 1, 2, 3 gives respective countings of 1.171, 1.894, 3.065 and 4.960 instead of correct answers 1, 2, 3 and 5. Really not a very bad closeness, although responses improve to give at

 $^{{}^5\}mathrm{It}$ is needed then to extent the use of Fibonacci sequence to negative subindices. In such cases $F_{-n}=(-1)^{n+1}F_n.$

m = 20 a number of boxes of 17711.00001 being $F_{22} = 17711$.

8 Conclusions

In this work has been shown how to proceed from the linearization of a canonical class of maps with memory describing population dynamics, -DRM and its manipulations in eqs. (1), (2) and (3)-, to an expression bounding its evolution in eq. (5) in which the key point to focus attention on was the product of a random matrices set. These are A_i 's in eq. (3) and nearby expressions, and while watching at their structure a strategy springs for calculation of those products in a binomial way, one who breaks up them in a stochastic part and another that accounts for memory. Stripping in a first step the deterministic matricial skeleton out of scalar random values it is possible to classify and calculate the many products that have been put forward. It is also possible without lose of information formulate the products of random variables and sum up them as well as to assign such probabilistic functions to corresponding matrices' products, or more properly speaking to classes of products. These algorithm-like procedure is condensed and used in sections 5 and 6 to write needed norms of eq. (5) in an operative way, eqs. (9) and (10) in the light of the general formula for above mentioned products, summarized in eq. (7).

Once is finished this quasi-computational narrative about products of matrices through a binomial like expansion of them a discussion in section 7 is carried out to evaluate the number of useful products which contributes to the result. The representation chosen suggests a fractal distribution of those surviving contributors over the whole set of all possible products in the expansion. A criterion, among many available in the field, is given to characterize the fractality, (*i.e.* dimension), of this *binomial set* and two equivalent ways to count elements are implemented. First an empirical one which possesses the purpose of methodology goodness illustration, (in the very optimistic aim of not being this problem an isolated one but the tip of iceberg for similar random maps or dynamical models), and second a extension of the former originated in number theory which gives an exact answer to the question of fractal dimension and content. Effectivity of a few steps approach is granted when compared both results for dimension quest, since $s_e = 0.691 \pm 0.003$ and $s_t = 0.694$.

The original motivation of this study was a careful reading of a paper from Cabrera *et al.* [1] in which they employ linearization of DRM to explain the origin of power spectrum of some theories of turbulence. At some point of their research they face how $P_{g,-1}$ behaves and $||P_{g,-1}||_2$ grows when g increases, as well in what way eigenvalues of $M_{g,j} \equiv P_{g,-1}^{\dagger}P_{g,-1}$ reflect a subjacent fractal structure generated by all coefficients of the former product $P_{g-1,-1}$. We here have just suggested a procedure which should allow them to go ahead with a full stochastic simulation and a complete description of equation (5), for arbitrary g. That is, one formulation and run involving also $P_{g,i}$, $i = 0, \ldots, g$ and not only the operator $P_{g,-1}$ attached to initial conditions \vec{X}_0 .

Acknowledgement

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

 Cabrera JL, Gutiérrez ED, Rodríguez Márquez M. Criticality and the fractal structure of -5/3 turbulent cascades. Chaos, Solitons and Fractals 2021;146:110876. https://doi.org/10.1016/j.chaos.2021.110876.

- [2] Maynard Smith J. Mathematical Ideas in Biology. Cambridge University Press, 1968.
- [3] Pounder JR, Roger TD. The geometry of chaos: Dynamics of a nonlinear second-order difference equation. Bulletin of Mathematical Biology 1980;42(4):551-597.
- [4] Kolmogorov AN, S. V. Fomin SV. Elements of the Theory of Functions and Functional Analysis. Mir, 1978.
- [5] Falconer K. Fractal Geometry: Mathematical Foundations and Applications.
 Wiley & Sons, 1990. ISBN 0-471-92287-0
- [6] Honsberger R. A second look at the Fibonacci and Lucas numbers, in Mathematical Gems III. Mathematical Association of America, Washington, 1985. ISBN 0-88385-313-2
- [7] Basin SL, Hoggatt VE Jr. A Primer on The Fibonacci Sequence Part II. The Fibonacci Quarterly 1963;1(2):61-68.