

Technical University of Munich

Department of Mathematics

Master's Thesis in Mathematics

Variations on Reinforced Random Walks

Fabian Michel

Supervisor: Prof. Dr. rer. nat. habil. Nina Gantert

Submission Date: 16/09/2022

Technical University of Munich

Department of Mathematics

Master's Thesis in Mathematics

Variations on Reinforced Random Walks

Abwandlungen selbstverstärkender Irrfahrten

Fabian Michel

Supervisor: Prof. Dr. rer. nat. habil. Nina Gantert

Submission Date: 16/09/2022

final digital version

I confirm that this master's thesis is my own work and I have documented all sources and material used.

Ich erkläre hiermit, dass ich diese Arbeit selbständig und nur mit den angegebenen Hilfsmitteln angefertigt habe.

Fabian Michel, München, 16/09/2022

Abstract

This thesis examines edge-reinforced random walks with some modifications to the standard definition. An overview of known results relating to the standard model is given and the proof of recurrence for the standard linearly edge-reinforced random walk on bounded degree graphs with small initial edge weights is repeated. Then, the edge-reinforced random walk with multiple walkers influencing each other is considered. The following new results are shown: on a segment of three nodes, the edge weights resemble a Pólya urn and the fraction of the edge weights divided by the total weight forms a converging martingale. On **Z**, the behavior is the same as for a single walker – either all walkers have finite range or all walkers are recurrent. Finally, edge-reinforced random walks with a bias in a certain direction are analysed, in particular on **Z**. It is shown that the bias can introduce a phase transition between recurrence and transience, depending on the strength of the bias, thus fundamentally altering the behavior in comparison to the standard linearly reinforced random walk.

Zusammenfassung

Diese Masterarbeit betrachtet (kanten-)selbstverstärkende Irrfahrten mit einigen Veränderungen im Vergleich zur gängigen Definition. Es wird eine Übersicht über bekannte Ergebnisse zum gängigen Modell gegeben und der Beweis, dass die linear selbstverstärkende Irrfahrt auf Graphen mit beschränktem Grad und hinreichend kleinen anfänglichen Kantengewichten rekurrent ist, wird wiederholt. Danach werden selbstverstärkende Irrfahrten mit mehreren Walkern, die sich gegenseitig beeinflussen, untersucht. Die folgenden neuen Resultate werden bewiesen: Auf einer Strecke mit drei Knoten ähneln die Kantengewichte einer Pólya-Urne und der Anteil der Kantengewichte am Gesamtgewicht bildet ein konvergierendes Martingal. Auf **Z** ist das Verhalten das gleiche wie für einen einzelnen Walker – entweder besuchen alle Walker nur einen endlichen Teil des Graphen oder alle sind rekurrent. Zum Schluss wird die selbstverstärkende Irrfahrt mit Bias in eine bestimmte Richtung betrachtet, vor allem auf **Z**. Es wird gezeigt, dass der Bias einen Phasenübergang zwischen Rekurrenz und Transienz verursachen kann, der von der Stärke des Bias abhängt, und damit das Verhalten im Vergleich zur normalen linear selbstverstärkenden Irrfahrt grundlegend ändert.

Contents

List of Figures

List of Tables

[1 Overview of results and conjectures on modified reinforced walks](#page-61-1) 62

A note on cross-references in this thesis

Cross-references and citations are marked in red, and they are clickable and directly link to the referenced object in the digital version. Titles, definitions, theorems, etc. are colored in blue.

1 Introduction

The central topic of this thesis is the edge-reinforced random walk (ERRW), a special type of random walk on a graph. The edges in the graph are weighted, and the probability to leave a node via one of the incident edges is proportional to the respective edge weight compared to the weights of the other incident edges. Each time an edge is crossed, its weight is increased according to some reinforcement scheme. Thus, it becomes more likely to visit parts of the graph which have already been visited before again. Next to reviewing results on this model, two modifications are considered: introducing multiple random walkers which influence each other and introducing a bias in a certain direction.

The ERRW is harder to analyze compared to random processes which possess the Markov property. In contrast to Markov chains (MCs), where the transition probabilities from one state to another only depend on the current state, but not on the past, the transition probabilities of the ERRW change over time as the edge weights are reinforced. The Markov property is lost. However, for a certain class of ERRWs, namely the linearly edge-reinforced random walk (LERRW), where the edge weights are increased by a constant increment upon every traversal, the reinforced random walk is equal in law to a mixture of MCs, i.e. a MC with random transition probabilities. This equivalence was the basis for many of the known results for ERRWs, since it allows us to use tools developed for MCs on the ERRW as well.

The new variations introduced in this thesis, multiple walkers and a bias, break this connection to MCs as well (at least in most of the considered cases, if not in all of them). The walk can no longer be represented as a mixture of MCs, and has to be analyzed using different techniques. The results presented here indicate that multiple walkers do not fundamentally change the behavior of the LERRW, while a bias is strong enough to do so.

1.1 Literature

The ERRW, as well as its counterpart, the vertex-reinforced random walk (VRRW), have been studied extensively, the first papers dating back to 1987, when the model was introduced by Coppersmith and Diaconis. Even before, [\[9\]](#page-63-1) showed that the LERRW (if certain assumption are satisfied) has a representation as a mixture of MCs. Much later, [\[20\]](#page-64-0) showed that this representation can be used for the LERRW on any graph, and [\[19\]](#page-64-1) even gave a formula for the so-called mixing measure on finite graphs. The mixing measure simply is the distribution of the random transition probabilities in the mixture of MCs.

Relatively early, results for the LERRW on trees were obtained. [\[14,](#page-64-2) [15,](#page-64-3) [29\]](#page-65-0) showed that there is a phase transition between recurrence and transience in the initial edge weights. The interesting case of \mathbb{Z}^d remained open much longer until [\[4,](#page-63-2) [5,](#page-63-3) [6\]](#page-63-4) showed that for $d \geq 3$, there is again a transition from recurrence to transience in the initial weights. [\[7\]](#page-63-5) proved that this transition is sharp, i.e. there is a certain critical initial edge weight such that for smaller initial weights, the random walk is recurrent, and transient for larger weights.

In parallel, the VRRW was analyzed, but with completely different tools (there is no representation as a mixture of MCs). [\[8\]](#page-63-6) analyzed the behavior of the VRRW on finite graphs with the help of a so-called stochastic approximation: the evolution of the vertex weights is approximated by a differential equation. [\[12,](#page-63-7) [13\]](#page-63-8) proved that the VRRW (with linear reinforcement) gets stuck on 5 nodes on **Z**, a result which could in part be generalized to arbitrary graphs: [\[11\]](#page-63-9) showed that the VRRW gets stuck with positive probability on certain finite subgraphs for almost any graph. The behavior of the VRRW is thus largely different from the behavior of the ERRW, and it was therefore unclear how the variations of the ERRW considered here would

affect the random walk.

An overview of results on reinforced processes in general can be found in [\[1,](#page-63-10) [21\]](#page-64-4). These surveys also show that reinforced random walks are closely related to urn processes, which have a very similar reinforcement component to the linearly reinforced walks: in most urn models, when a ball of a certain color is drawn, a fixed number of balls of the same color is added to the urn. This is also a type of linear reinforcement, and urns have been used on many occasions to analyze reinforced walks (see, for example, [\[14,](#page-64-2) [16\]](#page-64-5)).

Reinforced walks with multiple walkers or with a bias have (to the best of our knowledge) not been studied yet. There is however, another relevant part of the literature which deals with Markovian random walks. Markov chains are still a vital tool to understand the more complicated reinforced walks. [\[3,](#page-63-11) [10\]](#page-63-12) are two very good resources for results and techniques which can be applied to MCs. Finally, the references list additional sources which look at various aspects of random walks, including more and different models for reinforced walks.

1.2 Main Results

The following results are presented in this thesis. After covering preliminary definitions in [Section 2,](#page-9-0) some of the most important results on reinforced random walks, in particular in relation to this thesis, are presented in [Section 3,](#page-12-0) without proof. [Section 4](#page-17-0) repeats the proof of one such result: the recurrence of the LERRW on bounded-degree graphs for small initial weights (see [Theorem 4.4\)](#page-26-0).

[Section 5](#page-29-0) covers the first new variation to the LERRW: multiple walkers influencing each other. [Corollary 5.9](#page-36-0) shows that edge weights of the LERRW for 2 walkers on a segment of 3 nodes behave similarly to the proportion of balls in a Pólya urn, and it is conjectured in [Conjecture 5.10](#page-37-2) that the limit of the edge weight proportions will have similar properties to the limit of the proportion of balls in the urn. [Theorem 5.17](#page-44-0) considers the ERRW with more general reinforcement and multiple walkers on **Z** and shows that its behavior is similar to the case with a single walker: either all walkers are recurrent, or all walkers have finite range.

In [Section 6,](#page-49-0) the LERRW on **Z**, but with an additional bias is considered. [Lemma 6.2](#page-50-0) proves that, for a multiplicative bias, the probability to move in the direction of bias converges to 1 for nodes which are visited infinitely often. This might hint at transience, as noted in [Conjec](#page-51-1)[ture 6.5.](#page-51-1) For an additive bias, the representation as a mixture of MCs is recovered, allowing us to show a phase transition between recurrence and transience in the bias in [Theorem 6.7,](#page-58-1) and it is even possible to show positive speed if the bias is strong enough in [Theorem 6.8.](#page-58-2) Finally, [Section 6.3](#page-58-0) considers the LERRW with a strong bias in the initial edge weights: the initial edge weight to the right of a node $z \in \mathbb{Z}$ is set to λ^z for some parameter λ . The unfinished calculation in [Section 6.3](#page-58-0) seems to indicate that the reinforced walk will be transient, i.e. that the initially biased environment dominates the reinforcement. This is similar in spirit to [\[7,](#page-63-5) Theorem 4], where it was shown that if the initial weights are set to the conductances of a recurrent MC, then the LERRW is also recurrent. In [Section 6.3,](#page-58-0) we have an initially transient environment. However, there cannot be a theorem like [\[7,](#page-63-5) Theorem 4] for transience, as can already be seen by looking at [Theorem 4.4.](#page-26-0)

Finally, [Section 7](#page-61-0) reviews the results on the new variations of the ERRW obtained in this thesis, and looks at the many open questions which still remain.

2 Preliminaries

Throughout this thesis, random walks on finite and infinite connected graphs will be considered. All considered graphs will be locally finite, that is, the number of edges incident to a node is always finite. The following notation is used in connection with graphs:

Definition 2.1 Graph

A (possibly infinite) graph *G* is a tuple (V, E) of vertices (or nodes) and edges, with $E \subseteq$ $\{\{u, v\} \mid u, v \in V \land u \neq v\}$. Sometimes, it is useful to consider directed edges, hence we also define the set $\vec{E} := \{(u, v) \mid u, v \in V \land \{u, v\} \in E\}$ of directed edges. For $e = (u, v) \in$ \vec{E} , we call \vec{e} := (v, u) the corresponding reversed edge, and we write $\vec{e} = u, \hat{e} = v$, so $e = (\tilde{e}, \tilde{e})$. We write $u \sim v$ if $\{u, v\} \in E$. dist (u, v) is the edge length of the shortest path between the two nodes and dist(v, e) is the minimum of the distances of v to the two endpoints of *e*.

2.1 Random Walks

One type of random walk on a graph is the RWRE, which is used as a tool to analyze reinforced random walks. The RWRE is basically a Markov chain, but with random transition probabilities, which are also referred to as the environment. The formal definition is as follows:

Definition 2.2 Random Walk in Random Environment

Consider a graph $G = (V, E)$. Let $\mathbf{c} := (c_e)_{e \in E}$ be a collection of random variables over some probability space with $c_e > 0$. For $v \in V$, we further denote by $c_v := \sum_{u \in V: u \sim v} c_{\{u,v\}}$ the sum of the conductances of all adjacent edges, and assume $c_v > 0$. The **environment** P_c : $V \times V \rightarrow [0, 1]$ on the graph *G* is defined by

$$
P_{\mathbf{c}}\left(u,v\right):=\frac{c_{\{u,v\}}}{c_u}
$$

The Markov chain (MC) $(X_n)_{n\geq 0}$, conditioned on the knowledge of **c**, and starting at $v \in V$, is now defined by

$$
\mathbb{P}_{v}^{\mathbf{c}}\left[X_{0}=v\right]=1 \qquad \mathbb{P}_{v}^{\mathbf{c}}\left[X_{n+1}=t \mid X_{n}=u\right]=P_{\mathbf{c}}\left(u,t\right)
$$

Xⁿ is a random walk taking values in *V*.

The distribution of **c** is called **P**, and is called the mixing measure, and the probability measure \mathbb{P}_v is defined by setting

$$
\mathbb{P}_{v}\left[\cdot\right]:=\int\mathbb{P}_{v}^{\mathbf{c}}\left[\cdot\right]\ \mathbf{P}\left(\mathrm{d}\mathbf{c}\right)
$$

 $(X_n)_{n\geq 0}$, distributed according to \mathbb{P}_v , is called **random walk in random environment** (RWRE), or mixture of MCs.

For more background on conductances as well as electrical networks and their relation to random walks, [\[10\]](#page-63-12) gives an excellent introduction (in Chapters 1 and 2).

We shall also use the following terminology:

- (1) $\pi_c : V \to [0, \infty)$ is called a **stationary measure** for the MC given by the conductances **c** if it satisfies $\pi_c(v) = \sum_{u \in V: u \sim v} \pi_c(u) P_c(u, v)$ for all $v \in V$. π_c is further called a stationary distribution if it is a probability measure. Setting $\pi_c(v) := c_v$ defines a stationary (and reversible) measure.
- (2) The MC given by **c** is called **recurrent** if $\mathbb{P}^{\mathbf{c}}_v$ [$X_n = v$ for infinitely many n] = 1, and **transient** otherwise. In the transient case, it holds that $\mathbb{P}_{v}^{\mathbf{c}}[X_{n} = v$ for infinitely many $n] = 0$, so the starting node *v* is only visited finitely often a.s., and $\mathbb{P}_{v}^{\mathbf{c}}\left[\exists n>0: X_{n}=v\right] < 1$. In the recurrent case, it holds that $\mathbb{P}^{\mathbf{c}}_v$ $[\exists n > 0 : X_n = v] = 1$ (obviously). Since the considered MCs are a.s. irreducible (every node can be reached from every other node with positive probability), this corresponds to the standard notion of recurrence and transience.

The MC is further called **positive recurrent** if the expected return time to the root is finite, i.e. $\mathbb{E}^{\mathbf{c}}_v$ [min $\{n > 0 : X_n = v\}$] $< \infty$ (this implies recurrence). Positive recurrence is equivalent to the existence of a (unique) stationary distribution. Note that the existence of a stationary measure does not imply the existence of a stationary distribution since the stationary measure may be infinite (if it is finite, it can always be normalized). Indeed, in the null recurrent case (recurrent, but not positive recurrent), there exists a unique stationary measure which is infinite.

We next define the ERRW, which is one of the main topics of this thesis. The ERRW is not a MC since the transition probabilities change over time.

Edge-Reinforced Random Walk Definition 2.3

Consider a graph $G = (V, E)$ and a counting function $z : \mathbb{N}_{\geq 0} \times E \to \mathbb{N}_{\geq 0}$ where we set *z* (0,*e*) = 0 for all *e* ∈ *E*. Let W_e : $\mathbb{N}_{\geq 0}$ → (0, ∞) be weight functions for *e* ∈ *E*. We now define the evolution of *z* as well as the **edge-reinforced random walk** (ERRW) $(X_n)_{n\geq 0}$ on *G* as follows:

for
$$
n \ge 1
$$
: $z(n, \{u, v\}) = \sum_{i=1}^{n} 1\!\!1_{(X_{i-1} = u \wedge X_i = v) \vee (X_{i-1} = v \wedge X_i = u)}$

i.e. *z* counts the number of edge traversals, and

$$
\mathbb{P}_{v} [X_{0} = v] = 1
$$

$$
\mathbb{P}_{v} [X_{n+1} = t | X_{n} = u, ..., X_{0} = x_{0}] = \frac{W_{\{u,t\}} (z (n, \{u, t\}))}{\sum_{s \in V : s \sim u} W_{\{u,s\}} (z (n, \{u, s\}))}
$$

i.e. the probability to make a transition from u to t at time n is proportional to the weight $W_{\{u,t\}}(z(n,\{u,t\}))$ associated to the edge $\{u,t\}$, which depends on the number of traversals of the edge. We set

$$
w(n, \{u, t\}) := W_{\{u, t\}}(z(n, \{u, t\}))
$$

The random variables *w* thus describe the evolution of the edge weights to which the transition probabilities are proportional.

It is also possible to make the edge weight depend on more than just the number of edge traversals and the edge, but we restrict ourselves to this case in this thesis which is already quite general. If W_e is the same for every $e \in E$, then we will write *W* instead of W_e . One important choice of *W* is *W* (n) = 1 + n , which corresponds to starting with initial edge weights 1 everywhere and incrementing the weight of the traversed edge by 1 at every step. We define the **linearly edge-reinforced random walk** (LERRW) in a way which is a little more general: the LERRW is the reinforced random walk where $W_e(n) = a_e + n$ for some choice of initial weights $(a_e)_{e \in E} \in (0, \infty)^E$.

Later, it will be useful to count the number of directed edge traversals. We therefore also define the following counting function:

$$
\overrightarrow{z} (n,(u,v)) = \sum_{i=1}^{n} \mathbb{1}_{X_{i-1} = u \wedge X_i = v}
$$

Another type of random walk is obtained by using vertex weights instead of edge weights for reinforcing.

Definition 2.4 Vertex-Reinforced Random Walk

Consider a graph $G = (V, E)$ and a counting function $z : \mathbb{N}_{\geq 0} \times V \to \mathbb{N}_{\geq 0}$ where we set *z* (0, *v*) = 0 for all *v* ∈ *V*. Let W_v : $\mathbb{N}_{\geq 0}$ → (0, ∞) be weight functions for *v* ∈ *V*. We now define the evolution of *z* as well as the **vertex-reinforced random walk** (VRRW) $(X_n)_{n>0}$ on *G* as follows:

for
$$
n \ge 1
$$
: $z(n, v) = \sum_{i=1}^{n} 1 \mathbb{1}_{X_i = v}$

i.e. *z* counts the number of vertex visits, and

$$
\mathbb{P}_{v}[X_{0}=v]=1 \qquad \mathbb{P}_{v}[X_{n+1}=t | X_{n}=u,\ldots,X_{0}=x_{0}]=\frac{W_{t}(z(n,t))}{\sum_{s\in V:s\sim u}W_{s}(z(n,s))}
$$

i.e. the probability to go from *u* to *t* at time *n* is proportional to the weight $W_t(z(n, t))$ associated to the node *t*, which depends on the number of visits to the node. We set

$$
w(n,t) := W_t(z(n,t))
$$

3 Selected Results

Here, we present known results on reinforced random walks to put the following sections into context.

3.1 Reinforced Random Walks

3.1.1 Linearly Edge-Reinforced Random Walk

The following relation between LERRW and RWRE is a very important tool to analyze the LERRW. Many previous results on the LERRW have been proved using the fact that the LERRW is equal in distribution to a mixture of Markov chains, i.e. a MC with random transition probabilities. MCs are well understood, thus this theorem allows us to use the tools we have for MCs, and apply them to the non-Markovian LERRW.

Reinforced Walk as Mixture of Markov Chains ([\[4,](#page-63-2) Theorem 4], [\[20,](#page-64-0) Theorem 2.2], [\[9\]](#page-63-1)) **Theorem 3.1**

For a finite graph G with initial weights $(a_e)_{e \in E}$, the LERRW is equal in law to a RWRE with a uniquely determined mixing measure **P**. For any $e \in E$, **P** $[c_e > 0] = 1$. In particular, if the probability measure corresponding to the LERRW is denoted by **P***v*, then

$$
\mathbb{P}_{v}\left[\cdot\right]=\int\mathbb{P}_{v}^{\mathbf{c}}\left[\cdot\right]\ \mathbf{P}\left(\mathrm{d}\mathbf{c}\right)
$$

with the notation from [Definition 2.2.](#page-9-2)

We next turn to the LERRW on trees. Using [Theorem 3.1,](#page-12-3) one can show that

Phase Transition of Reinforced Walk on Trees ([\[14,](#page-64-2) [16\]](#page-64-5)) **Theorem 3.2**

Let *T* be a an infinite tree. Consider the LERRW on *T* with initial weights $a > 0$. Let Δ_0 be the solution to the equation

$$
\frac{\Gamma\left(\frac{2+\Delta}{4\Delta}\right)^2}{\Gamma\left(\frac{1}{2\Delta}\right)\Gamma\left(\frac{1+\Delta}{2\Delta}\right)}=\frac{1}{\text{br}\left(T\right)}
$$

where br(T) is the branching number of the tree T (a measure of the number of children per node, the branching number is equal to the number of children for regular trees; see [\[16,](#page-64-5) Definition 2.6] for the exact definition). Then the LERRW on *T* is

- (i) a.s. transient if *a* > $\frac{1}{\Delta_0}$
- (ii) a.s. recurrent if $a < \frac{1}{\Delta_0}$.

The expected return time to the root is always infinite.

The same holds for Galton-Watson trees with br(*T*) replaced by the mean number of children, if the mean is > 1 and conditioned upon non-extinction.

On **Z**, the LERRW is recurrent for any choice of initial weight *a*.

In this thesis, a recurrence result for the LERRW on general graphs will be shown. It is a part of the following theorem, for which we also need to define the Cheeger constant. A graph is called nonamenable if there is a constant $h > 0$ such that for any finite set of nodes $A \subseteq V$, $|\partial A|$ ≥ *h* |*A*| where $\partial A = \{v \in V \setminus A : \text{dist}(v, A) = 1\}$, i.e. ∂A is the set of nodes which are outside of *A*, but direct neighbors to a node within *A*. The Cheeger constant is the largest such constant *h*, and can be defined as follows:

$$
h(G) := \inf \left\{ \frac{|\partial A|}{|A|} : A \subseteq V \text{ finite} \right\}
$$

Theorem 3.3 LERRW on General Graphs ([\[4,](#page-63-2) [5,](#page-63-3) [6\]](#page-63-4))

Let *G* be any graph. Consider the LERRW with initial edge weight *a* for every edge. Then:

- (i) for every constant *K*, there exists some $a_0 > 0$ such that if all degrees of *G* are bounded by *K*, the LERRW on *G* is recurrent if $a \in (0, a_0)$.
- (ii) for every constant *K*, and any $c_0 > 0$, there exists some $a_0 > 0$ such that if all degrees of *G* are bounded by *K*, and if $h(G) \ge c_0$, then the LERRW on *G* is transient if $a > a_0$.
- (iii) for $G = \mathbb{Z}^3$ and $G = \mathbb{Z}^d$ with any $d \geq 3$, there exists some $a_0 > 0$ such that the LERRW on *G* is transient if $a > a_0$.

Part [\(i\)](#page-13-0) of the theorem above is shown in this thesis in [Theorem 4.4.](#page-26-0) Next to the above, and next to other results which are not listed here, the following is known about the LERRW:

- On \mathbb{Z}^3 and \mathbb{Z}^d with any $d \geq 3$, there is a sharp phase transition between recurrence and transience of the LERRW in the initial edge weights. If the initial edge weights are all set to a constant $a > 0$, then there is a critical threshold a_0 such that the LERRW on \mathbb{Z}^d is recurrent if $a < a_0$ and transient if $a > a_0$. See [\[7,](#page-63-5) Theorem 3].
- On any locally finite graph, if a random walk with given conductances on the edges is recurrent, then so is the LERRW if these conductances are used as initial edge weights. See [\[7,](#page-63-5) Theorem 4].
- The VRJP (vertex-reinforced jump process) is a continuous-time version of the LERRW. The process $(Y_t)_{t\geq0}$ takes values in the vertices of a graph with conductances W_e on the edges. If we are at vertex v at time t , then, conditional on $(Y_s, s \leq t)$, the process jumps to a neighbor *u* of *v* at rate $W_{\{v,u\}}\cdot L_u\left(t\right)$ with

$$
L_u(t) = 1 + \int_0^t \mathbb{1}_{Y_s = u} ds
$$

Up to an exponential time change, and using independent Gamma-distributed conductances, this has the same distribution as the following model of a continuous-time LERRW when looked at at jump times. In order to make the LERRW continuous, we can add clocks on every edge. The time of an edge runs only when the process is in a node adjacent to it, and the alarm of an edge rings at exponential times with the rate of the edge weight (i.e. the rates are increasing). When an alarm rings, then the process immediately traverses the corresponding edge.

The VRJP is distributed as a mixture of time-changed Markov jump processes, just as the LERRW is distributed as a mixture of Markov chains.

The results on the LERRW basically also apply to the VRJP (or the other way round). See

[\[5\]](#page-63-3) for more details.

• For the LERRW on finite graphs (with arbitrary initial weights), there is a "magic formula": The LERRW is a mixture of Markov chains where the mixing measure can be given explicitly: if the edge weights are normalized to sum to one, they converge to a random limit on the unit simplex. The law can be given as a density w.r.t. the Lebesgue measure on the unit simplex, and the LERRW is distributed as a mixture of (reversible) MCs with the weights chosen according to this same measure (the weights are not independent across edges). See [\[19\]](#page-64-1).

The main reason why this is true and also the basis of the proof is partial exchangeability, i.e. two finite paths have the same probability if the edges are traversed the same number of times (this is easy to see: the probability of a path is a product of fractions; for nominators, consider the edges, for denominators, the vertices, to see that the order of traversals is irrelevant).

For recurrent LERRWs in general, we can use a de Finetti theorem on MCs, by considering blocks of excursions from the starting vertex (these are iid conditioned on some parameter) instead of balls drawn as for the most basic de Finetti theorem. See [\[9\]](#page-63-1).

• The RWRE representation for the LERRW also exists for infinite graphs and in particular transient LERRW (no explicit formula for the mixing measure is known). See [\[20\]](#page-64-0).

Just in order to mention that there is more than just linear reinforcement: there is stronger than linear reinforcement which can lead to being stuck on a single edge, and there is the oncereinforced random walk (where the edge weight is only increased on the first visit and then stays constant) which is even more difficult to analyze, little is known about it.

This thesis adapts the LERRW and adds multiple walkers or a bias to the model. [Section 5](#page-29-0) and [Section 6](#page-49-0) add some results to the list above.

3.1.2 Vertex-Reinforced Random Walk

There are some changes in behavior if vertex weights instead of edge weights are reinforced. A short overview of known results:

- On any complete graph, the linearly vertex-reinforced random walk converges to uniform vertex occupation (meaning that the time spent at each vertex, divided by the total time, will converge to a uniform distribution, so the vertex weights will all be of the same order). On any finite graph fulfilling a non-degeneracy condition, vertex occupation converges to a rest point of a differential equation approximating the evolution of the vertex weights. See [\[8\]](#page-63-6).
- On **Z**, the VRRW (linear reinforcement) is eventually trapped on an interval of 5 vertices, the middle vertex is visited with frequency $\frac{1}{2}$, its neighbors with positive frequency, and the two outer vertices are visited infinitely often but at frequency 0. See [\[13\]](#page-63-8).
- On \mathbb{Z}^2 , the VRRW (linear reinforcement) is trapped with positive probability on 13 vertices (a diamond with $\frac{1}{2}$ -frequency core and with 0-frequency boundary) and with positive probability on 12 vertices (square with 0-frequency boundary). See [\[11\]](#page-63-9). It still remains an open question whether the VRRW is trapped with probability 1.
- On any graph of bounded degree, the VRRW (linear reinforcement) is trapped with positive probability on finitely many vertices which form a complete *n*-partite graph with outer boundary such that every vertex in the outer boundary is not connected to one of the partitions plus one extra node. See [\[11\]](#page-63-9).
- On any tree of bounded degree, the VRRW (linear reinforcement) is trapped with proba-bility 1 on a finite subgraph. See [\[11\]](#page-63-9).
- On a tree with K_n children for every node on level *n*, where $\sum K_n^{-1} < \infty$, there is a positive probability for the VRRW (linear reinforcement) to move away from the root on *every single* step (transience), but also a positive probability to get trapped (finite range), so there is no 0-1-law. See [\[11\]](#page-63-9).

These results show that an apparently small change in the model can dramatically affect the behavior of the random walk. Instead of being recurrent or transient, as the LERRW, the VRRW with linear reinforcement has finite range (gets stuck on a finite subgraph) with positive probability. This is one of the reasons why it was interesting to study variations to the model of the ERRW, in order to better understand which changes in the model cause which effects.

3.2 Urn Models

Urn models are closely related to reinforced random walks since they are also an example of a reinforced random process: usually, balls are drawn uniformly at random, and more balls of the drawn color are added. Hence, the fraction of balls of a certain color is reinforced upon drawing a ball of the same color.

3.2.1 Pólya Urn

Consider a Pólya urn, starting with *w* white and *b* black balls. At every time step, draw a ball from the urn, and put it back along with Δ more balls of the same color. Call w_n and b_n the number of white and black balls after the *n*-th draw, so $w_0 = w$ and $b_0 = b$. Then:

Theorem 3.4 Pólya Urn ([\[14,](#page-64-2) Lemma 1])

The fraction of white balls $\frac{w_n}{w_n+b_n}$ is a martingale, and it thus converges to a limit random variable *L* which is distributed according to a Beta distribution:

$$
L \sim B\left(\frac{w}{\Delta}, \frac{b}{\Delta}\right)
$$

The sequence of balls drawn is equal in law to a mixture of iid sequences. Indeed, conditioned on *L*, the sequence of balls drawn is distributed as an iid sequence where a white ball is drawn with probability *L*.

The fact that the sequence of draws is distributed as a mixture of iid sequences follows from de Finetti's theorem, which is applicable since the draws are exchangeable: the probability of a certain sequence of draws occurring depends only on the number of white and black balls in the sequence, but not on their order. This is very similar to what we already saw in [Theorem 3.1,](#page-12-3) and indeed, the same arguments are used in the proof. Another thing to note is that [Theorem 3.4](#page-15-3) does not only hold when *w*, *b* and ∆ are integers, but also when they are replaced by any positive real numbers. In other words, it is also possible to have only a fraction of a ball in the urn.

In [Section 5,](#page-29-0) [Corollary 5.9](#page-36-0) and [Conjecture 5.10](#page-37-2) show that something similar to a two-player Pólya urn (where two players draw balls and do not put them back immediately) exhibits very similar properties to that of the standard one-player Pólya urn.

3.2.2 Randomly Reinforced Urn

Consider the following urn model: initially, the urn contains b_0 black and w_0 white balls. The numbers b_n and w_n of balls contained in the urn in the *n*-th step is determined as follows. Draw a ball uniformly at random from the urn with *bn*−¹ black balls and *wn*−¹ white balls. If the ball is black, replace it with *Mⁿ* black balls. If the ball is white, replace it with *Nⁿ* white balls. *Mⁿ* and N_n are series of independent random variables on the nonnegative natural numbers and all *Mⁿ* have the common distribution *µ*, all *Nⁿ* have the common distribution *ν*. It is assumed that *µ* and *v* both have finite support. Call $z_n = \frac{b_n}{b_n + w_n}$ the fraction of black balls in the urn. Denote by B_n , W_n , Z_n the corresponding random variables.

Randomly Reinforced Urn ([\[18,](#page-64-6) Theorem 4.1]) **Theorem 3.5**

If the following assumptions hold (and μ and ν both have finite support, as stated above)

(i) $b_0 > 0$ (at least one black ball in the urn at the beginning)

- (ii) $\mu({0}) = 0$ (if a black ball is drawn, it is replaced by at least one black ball)
- (iii) $\mu \geq_{st} \nu$ (*µ* stochastically dominates *ν*)
- (iv) $\mathbb{E}[M_n] = \int_{\mathbb{N}} x \mu(dx) > \int_{\mathbb{N}} x \nu(dx) = \mathbb{E}[N_n]$ (the expected number of black balls added is bigger than the expected number of white balls added)

Then $\lim_{n\to\infty} Z_n = \lim_{n\to\infty} \frac{B_n}{B_n + W_n}$ (the fraction of black balls converges almost surely to 1).

[Theorem 3.5](#page-16-0) covers a large class of urns, including urns with deterministic reinforcement where a different number of balls is added depending on the color.

4 Recurrence of Edge-Reinforced Random Walk

In this section, we repeat the proof of the known result that the LERRW is recurrent on any bounded-degree graph for small initial weights. We already know that the LERRW has an equivalent representation as a RWRE with random conductances *c^e* on the edges. If we further normalize these conductances by setting $c_{v_0} = 1$ where v_0 is the starting vertex of the LERRW (we will assume this normalization in the following sections), then we can give a more precise statement of what we want to show. We consider the weight function $W(n) = a + n$ and are interested in the following result, given in [\[4,](#page-63-2) Theorem 2]: if *G* is a graph with degree at most *K*, if the LERRW is recurrent on *G* for any choice of initial weights, and if $s \in \left(0, \frac{1}{4}\right)$, then there is some $a_0 > 0$, depending on *s* and *K*, such that for initial weights $a \in (0, a_0)$ we have $\mathbb{E}_{v_0}[c_e^s] \leq 2K$ (*C* (*s*, *K*) $\sqrt{ }$ \overline{a})^{dist(v_0 , e) where *C* (*s*, *K*) solely depends on *K* and *s*. In particular, for} *a* small enough, we have exponential decay in the expected conductances *c^e* of the random environment corresponding to the LERRW (exponential decay with increasing distance, and with the conductances raised to the *s*-th power). As a consequence, we can prove under almost the same assumptions, but without requiring a priori that the random walk is recurrent, that the LERRW is recurrent for *a* small enough.

The proof given here will follow the proof given in [\[4,](#page-63-2) Section 2] with only minor modifications, which hopefully make a proof which was already really well presented in [\[4\]](#page-63-2) even easier to understand. So the mathematics presented here was really done by [\[4\]](#page-63-2). The main idea is to estimate the ratios $\frac{c_e}{c_f}$ for edges *e* and *f* which are incident on the same vertex by comparing with a corresponding ratio of numbers of edge traversals of the LERRW, under the assumption that the random walk is recurrent. We then bound the estimated ratio as well as the error caused by the estimation to conclude.

4.1 Estimating Conductance Ratios

The first step is to construct an estimate of conductance ratios. Our final goal will be to bound the expectation of c_e^s for any edge *e*. We therefore choose an arbitrary directed edge $e \in \overrightarrow{E}$ to begin with. Next, we will define a path from the starting vertex v_0 to e and estimate the conductance ratios along this path, which will result in an estimate for *c^e* . The construction of the path is as follows: for the edge $e = (e, e)$, we can find the directed edge e' through which \check{e} ^{*} was first reached by the LERRW (or the RWRE, depending on your point of view). If e is traversed before its corresponding inverse \overleftarrow{e} , then *e'* \neq *e* as undirected edges. Since *e'* must have been traversed before its inverse, by its definition, it is possible to iterate this construction until the starting vertex v_0 is reached. This results in a (random) path $\gamma_e = (\ldots, e'', e', e)$ starting from v_0 and ending with *e* (the path can be a loop if $\hat{e} = v_0$). For a deterministic path γ , we call *D*_{*γ*} the event that for the last edge *e* of *γ*, we have $\gamma_e = \gamma$, and that *e* is traversed before ϵ .

For *e* such that $e' \neq \overleftarrow{e}$, we now estimate the ratio $\frac{c_e}{c_{e'}}$ (i.e. the ratio of the conductances of two edges along one of the paths constructed above) by

$$
Q(e) := \frac{M_e}{M_f} \quad \text{where } f := \overleftarrow{e'}, \quad \text{an estimate for } R(e) := \frac{c_e}{c_{e'}} \tag{4.1}
$$

where *M^e* counts the number of times the directed edge *e* was crossed in the right direction before the edge *f* was crossed in the right direction by the random walk, if *e* is crossed before *f* . In this case, M_f is set to 1. Otherwise, M_f counts the number of times f was crossed before *e*, and *M^e* is 1. In other words, *M^e* and *M^f* count the numbers of directed edge crossings of *e*

and *f* until both edges have been traversed by the random walk. Since both edges lead out of the same vertex, this can also be seen as counting the number of departures from \check{e} until both edges were used.

Figure 1: Estimating conductance ratios

We next define the deterministic set of simple paths (or loops, if one of the endpoints of *e* is *v*¹ *v*_{*e*} as the set of paths which end with *e* or [←]*e*. So, Γ_{*e*} is the set of possible values which the random path γ_e (or $\gamma_{\overline{e}}^+$, if \overline{e}^+ is traversed before *e*) can take. We now have

$$
\mathbb{E}_{v_0}\left[c_e^s\right] = \sum_{\gamma \in \Gamma_e} \mathbb{E}_{v_0}\left[c_e^s \cdot \mathbb{1}_{D_\gamma}\right] \le \sum_{\gamma \in \Gamma_e} \mathbb{E}_{v_0}\left[\left(\frac{c_e}{c_{\gamma_1}}\right)^s \cdot \mathbb{1}_{D_\gamma}\right]
$$

where γ_1 is the first edge of γ . Recall that we normalized the environment such that $c_{v_0} = 1$, so in particular, $c_{\gamma_1} \leq 1$ since γ_1 leads out of v_0 .

One of the assumptions we started with was that we already know that the LERRW is recurrent. This implies that *e* will be visited a.s. and therefore the partition into possible paths *γ* ∈ Γ*^e* does indeed cover the whole probability space. Furthermore, by [Theorem 3.1](#page-12-3) and [\[9\]](#page-63-1), the conductances in the random environment corresponding to the LERRW are uniquely defined and we can therefore look at $\frac{c_e}{c_{\gamma_1}}$.

We now want to compare the estimated conductance ratios along the path *γe* with the real conductance ratios. If we fix a path γ ending in *e* and D_{γ} occurs, then

$$
\frac{c_e}{c_{\gamma_1}} = \prod_{f \in \gamma, f \neq \gamma_1} R(f) = \underbrace{\prod_{f \in \gamma, f \neq \gamma_1} \frac{R(f)}{Q(f)}}_{\text{error}} \underbrace{\prod_{f \in \gamma, f \neq \gamma_1} Q(f)}_{\text{estimate}}
$$

By the Cauchy-Schwarz inequality, we have

$$
\mathbb{E}_{v_0}\left[\left(\frac{c_e}{c_{\gamma_1}}\right)^s \cdot \mathbb{1}_{D_{\gamma}}\right] \leq \mathbb{E}_{v_0}\left[\prod_{f \in \gamma, f \neq \gamma_1} \left(\frac{R\left(f\right)}{Q\left(f\right)}\right)^{2s} \cdot \mathbb{1}_{D_{\gamma}}\right]^{\frac{1}{2}} \mathbb{E}_{v_0}\left[\prod_{f \in \gamma, f \neq \gamma_1} Q\left(f\right)^{2s} \cdot \mathbb{1}_{D_{\gamma}}\right]^{\frac{1}{2}}
$$

The above equation allows us to bound the expectation of the error introduced by the estimate separately from the expectation of the estimate. The following two sections will bound the two terms on the right hand side. Then, we will be able to show exponential decay of the edge conductances in the distance to the starting vertex for graphs on which the LERRW is recurrent, a result which can subsequently actually be used to show recurrence of the LERRW

itself.

4.2 Bounding the Error

Lemma 4.1 Error Bound ([\[4,](#page-63-2) Lemma 7])

Let *G* be a graph, $v_0 \in V$ the starting vertex, and $a \in (0, \infty)$ such that the LERRW on *G* with initial weights equal to *a* is recurrent. Then, for any $e \in E$, any $\gamma \in \Gamma_e$, and any $s \in (0,1)$:

$$
\mathbb{E}_{v_0} \left[\prod_{f \in \gamma, f \neq \gamma_1} \left(\frac{R(f)}{Q(f)} \right)^s \cdot \mathbb{1}_{D_{\gamma}} \right] \leq C(s)^{|\gamma|-1}
$$

where *C* (*s*) is a constant depending solely on *s* and $|\gamma|$ is the length of the path γ .

Note that the bound of the error given above shows that we have at most an exponential increase of our error in the length of the path *γ*. The corresponding bound of the estimated ratios will balance this potentially exponentially increasing error.

Proof R and *Q* depend in general on the random path *γ^e* (which was defined recursively as the path of first entrance edges in the section above). For a deterministic path *γ*, we can define *R^γ* and Q_γ for edges $f \in \gamma$, $f \neq \gamma_1$ as we defined *R* and *Q* for the random path: if $f \in \gamma$, and if *f*^{$′$} ∈ $γ$ is the preceding edge in the path, then $R_γ$ (*f*) = $c_f c_{f'}^{-1}$, and $Q_γ$ (*f*) is the ratio of exits from \check{f} along f and ←− *f* ′ until both edges have been visited (see [Equation 4.1\)](#page-17-2). We have, for fixed *γ*

$$
\mathbb{E}_{v_0} \left[\prod_{f \in \gamma, f \neq \gamma_1} \left(\frac{R(f)}{Q(f)} \right)^s \cdot \mathbb{1}_{D_\gamma} \right] \leq \mathbb{E}_{v_0} \left[\prod_{f \in \gamma, f \neq \gamma_1} \left(\frac{R_\gamma(f)}{Q_\gamma(f)} \right)^s \right]
$$

This inequality holds since R_γ (*f*) = $R(f)$ and Q_γ (*f*) = $Q(f)$ if D_γ occurs. The inequality is a crude bound but sufficient for our purposes.

We want to show a bound on the error of estimating the conductance ratios. It is therefore necessary to look at the RWRE description of the random walk, and we will now condition on a particular vector **c** of conductances being chosen. We then show that for every such vector, the bound above holds, and this will give the desired result. So from now on, assume we have conductances **c**. $R_\gamma(f)$ is then also a fixed number, since it only depends on the conductances. The $Q_{\gamma}(f)$, on the other hand, are independent for any two different edges if we condition on the conductances c . This is because we just count directed exits from \check{f} , and we can couple these with iid sequences of random variables $(Z_n^v)_{n\geq 1}$ for every node v where Z_n^v takes values in the edges incident to node *v*, and is distributed such that an edge *e* appears with probability $c_e \left(\sum_{f \ni v} c_f \right)^{-1}$. These Z_n^v are iid for every node, and independent for different nodes.

By the independence of the $Q_\gamma(f)$ (conditioned on **c**), we have

$$
\mathbb{E}_{v_0} \left[\prod_{f \in \gamma, f \neq \gamma_1} \left(\frac{R_{\gamma}(f)}{Q_{\gamma}(f)} \right)^s \middle| \mathbf{c} \right] = \prod_{f \in \gamma, f \neq \gamma_1} \mathbb{E}_{v_0} \left[\left(\frac{R_{\gamma}(f)}{Q_{\gamma}(f)} \right)^s \middle| \mathbf{c} \right]
$$
\n
$$
= \prod_{f \in \gamma, f \neq \gamma_1} \mathbb{E}_{v_0} \left[\left(\frac{c_f}{c_{f'}} \cdot \frac{M_{\overline{f'} , \gamma}}{M_{f, \gamma}} \right)^s \middle| \mathbf{c} \right]
$$

where f' is the edge preceding f in γ and $M_{f,\gamma}$ is defined as M_f in [Equation 4.1](#page-17-2) but with the

deterministic path *γ* instead of the random path *γe* . To finish the proof, it is therefore sufficient to show that, for any edge $f \in \gamma$ preceded by f' ,

$$
\mathbb{E}_{v_0}\left[\left(\frac{c_f}{c_{f'}}\cdot\frac{M_{\overline{f'},\gamma}}{M_{f,\gamma}}\right)^s \middle| \mathbf{c}\right] \leq C\left(s\right)
$$

where *C* (*s*) does not depend on the conductance vector **c** we condition on.

Figure 2: Estimating conductance ratios along a deterministic path *γ*

Write $v = \check{f}$, $e =$ ←− *f* ′ (see [Figure 2](#page-20-0) for reference). The law of *Me*,*^γ* and *M^f* ,*^γ* is determined by the iid sequence $(Z_n^v)_{n\geq 1}$ (since we condition on **c**), and does not depend on other edges incident to *v*: it is only relevant with which probability the two edges appear relative to each other in the sequence. We can therefore assume w.l.o.g. that only *e* and *f* appear in the sequence Z_n^v with probabilities *p* and *q* = 1 − *p*, respectively. Then

$$
\mathbb{E}_{v_0} \left[\left(\frac{c_f}{c_e} \cdot \frac{M_{e,\gamma}}{M_{f,\gamma}} \right)^s \middle| \mathbf{c} \right] = \left(\frac{c_f}{c_e} \right)^s \left(\sum_{k \ge 1} k^s \cdot \mathbb{P}_{v_0} \left[k \text{ exits from } v \text{ along } e, \text{ then exit } v \text{ along } f \right] \right)
$$

+
$$
\sum_{k \ge 1} k^{-s} \cdot \mathbb{P}_{v_0} \left[k \text{ exits from } v \text{ along } f, \text{ then exit } v \text{ along } e \right] \right)
$$

=
$$
\left(\frac{q}{p} \right)^s \left(\sum_{k \ge 1} k^s p^k q + \sum_{k \ge 1} k^{-s} q^k p \right)
$$

$$
\stackrel{\circledast}{\le} \left(\frac{q}{p} \right)^s \left(q^{-s} p + C \left(s \right) q p^s \right) = p^{1-s} + C \left(s \right) q^{1+s} \le 2C \left(s \right)
$$

Since the bound is independent of *p* and *q* and thereby independent of the choice of the conductances **c**, this completes the proof, up to showing ⊛. Note that the bound can be improved, dependent on the value of *s*. We show ⊛ using the following observations. If $X \sim \text{Geo}_{\geq 0}(q)$ is a random variable with geometric distribution of parameter q , i.e. $\mathbb{P}\left[X=k\right]=p^k q$, then we can write

$$
\mathbb{E}\left[X^{s}\right] = \sum_{k\geq 1} k^{s} p^{k} q \leq pq^{-s} \iff \mathbb{E}\left[\left(qX\right)^{s}\right] \leq p \tag{4.2}
$$

Now, proving that the inequality in [Equation 4.2](#page-20-1) holds will show the bound on the first sum in ⊛. Note that the function $f(s) = x^s$ is convex for arbitrary choice of $x > 0$. Therefore, $g(s) = \mathbb{E}[(qX)^s]$ is convex in *s*. As *X* takes values in the integers, we have that $\mathbb{E}[(qX)^s] =$ $q^s \mathbb{E}[X^s] \leq \mathbb{E}[X] < \infty$ for $s \in (0,1)$. By dominated convergence, $g(s)$ is therefore continuous

in *s*, and can be continued continuously to the interval $s \in [0,1]$. For $s = 1$, we have $g(1) =$ $qE[X] = q \cdot \frac{p}{q} = p$, and for $s = 0$, we have

$$
\lim_{s \to 0} g(s) = \lim_{s \to 0} \mathbb{E} \left[(qX)^s \right] = \mathbb{E} \left[\lim_{s \to 0} (qX)^s \right] = \mathbb{E} \left[\mathbb{1}_{\{X > 0\}} \right] = p
$$

By convexity of *g*, we conclude that $g(s) \leq p$ for all $s \in (0,1)$, which proves [Equation 4.2.](#page-20-1)

For bounding the second sum, let *X* \sim Geo_{>1} (*p*) be a random variable with geometric distribution of parameter *p*, but which takes values in $\mathbb{N} \setminus \{0\}$, so $\mathbb{P}[X = k] = q^{k-1}p$ where $k \geq 1$. Then we can write

$$
q \cdot \mathbb{E}\left[X^{-s}\right] = q \cdot \sum_{k \ge 1} k^{-s} q^{k-1} p = \sum_{k \ge 1} k^{-s} q^k p \le C\left(s\right) q p^s \iff \mathbb{E}\left[\left(pX\right)^{-s}\right] \le C\left(s\right) \tag{4.3}
$$

Now, $g(p) = \mathbb{E}\left[\left(pX\right)^{-s}\right]$ is decreasing in p (to see this, note that $\mathbb{E}\left[\left(pX\right)^{-s}\right] = \frac{p^{1-s}}{1-n}$ 1−*p* ·Li*^s* (1 − *p*) where Li denotes the polylogarithm, derive: [https://bit.ly/3BOmLjP,](https://www.wolframalpha.com/input?i=d%2Fdx+of+x%5E%281-s%29+%2F+%281-x%29+*+PolyLog%5Bs%2C+1-x%5D) and analyze the derivative: [https://bit.ly/3RVn5CA\)](https://www.wolframalpha.com/input?i=plot+%28s+*+%28x+-+1%29+%2B+1%29+*+PolyLog%5Bs%2C+1-x%5D+-+x+*+PolyLog%5Bs+-+1%2C+1-x%5D+for+x+from+0+to+1+and+s+from+0+to+1), so it is sufficient to show that $\lim_{p\to 0} g(p) =: C(s)$ exists and is finite. Note that for $p \to 0$, pX converges in distribution to a random variable *Y* which is Exp (1)-distributed (we have that $\mathbb{P}[pX > t] \to e^{-t}$ for $p \to 0$). Therefore, $\lim_{p\to 0} g(p) =$ $\mathbb{E}[Y^{-s}] = \Gamma(1-s) < \infty$ for $s \in (0,1)$. Thus, we can choose $C(s) = \Gamma(1-s)$ in [Equa](#page-21-1)[tion 4.3.](#page-21-1) \Box

4.3 Bounding the Estimate

Lemma 4.2 Estimate Bound ([\[4,](#page-63-2) Lemma 8])

Let *G* be a graph with degree at most *K*, $v_0 \in V$ the starting vertex, and $a \in (0, \infty)$ such that the LERRW on *G* with initial weights equal to *a* is recurrent. Then, for any $e \in E$, any $\gamma\in\Gamma$ e, and any $s\in\left(0,\frac{1}{2}\right)$:

$$
\mathbb{E}_{v_0}\left[\prod_{f\in\gamma,f\neq\gamma_1}Q\left(f\right)^s\cdot\mathbb{1}_{D_{\gamma}}\right]\leq \left(C\left(s,K\right)a\right)^{|\gamma|-1}
$$

where *C* (*s*) is a constant depending solely on *s* and *K*.

Choosing *a* small enough, the bound above shows that the estimated ratios decrease exponentially and fast enough to balance the potential error caused by the estimation given in [Lemma 4.1.](#page-19-1)

Proof The idea of the proof is to find iid random variables $\overline{Q}(f)$ which stochastically dominate the variables $Q(f)$ on all edges of the deterministic path γ , provided that D_{γ} occurs. It is then much easier to bound the expectation of the random variables $\overline{Q}(f)$. We start by fixing γ and now want to define $\overline{Q}(f)$ such that $Q(f) \le \overline{Q}(f)$ for all $f \in \gamma$, $f \ne \gamma_1$ on the event D_γ .

For every such edge *f*, we define two independent series of Bernoulli random variables (also independent of the corresponding series for different edges) $(Y_j)_{j\geq 0}$ and (Y'_j) *j*≥0 :

$$
\mathbb{P}\left[Y_j=1\right] = \frac{a}{j+1+2a} = 1 - \mathbb{P}\left[Y_j=0\right] \qquad \mathbb{P}\left[Y'_j=1\right] = \frac{1+a}{2j+1+Ka} = 1 - \mathbb{P}\left[Y'_j=0\right]
$$

Intuitively, and up to some technical details, these variables are used as follows. Recall that

f' was the edge through which $\check{f} = v$ was first reached (which is the edge preceding f in γ on the event D_{γ}) and we set again $e =$ ←− \overleftarrow{f} ^{*f*}. We want to bound $Q(f) = \frac{M_f}{M_f}$ $\frac{M_{\text{e}}}{M_{\text{e}}}$, hence we want to upper bound *M^f* , the number of exits along *f* until both edges were used, and we want to lower bound M_e . First, $Y'_0 = 1$ represents the event that on the first visit to *v*, we depart along *e*, and sometimes also if $\hat{Y}'_0 = 0$. $\hat{Y}'_0 = 1$ is thus a lower bound for the departure along *e*. Now,

- if $Y'_0 = 0$, i.e. if the first departure was not necessarily along *e*, and *e* was not used yet: then, for $n \geq 1$, on the *n*-th visit to *v*, we will couple the random walk with the variables *Y*^{\prime} such that we depart along *e* if $Y'_{n-1} = 1$, and sometimes also if $Y'_{n-1} = 0$. $Y'_{n-1} = 1$ lower bounds the event that *e* is crossed.
- if $Y'_0 = 1$, i.e. if the first departure was along *e*, and if *f* was not used yet: then, for $n \ge 1$, on the *n*-th visit to v where v is left along \hat{f} or e , we will couple the random walk with the variables *Y* such that we depart along *e* if $Y_{n-1} = 0$, and sometimes also if $Y_{n-1} = 1$. *Y*_{*n*−1} = 1 upper bounds the event that *f* is crossed.

We now set

$$
\overline{Q}(f) = \frac{M_f}{\overline{M}_e} \text{ where}
$$
\n
$$
\overline{M}_f = \min \{ j \ge 1 : Y'_j = 1 \} \text{ and } \overline{M}_e = 1 \text{ if } Y'_0 = 0
$$
\n
$$
\overline{M}_e = \min \{ j \ge 1 : Y_j = 1 \} \text{ and } \overline{M}_f = 1 \text{ if } Y'_0 = 1
$$

Figure 3: Bounding *Q* with the help of Bernoulli random variables

The figure above demonstrates why the indicated coupling is possible. We are in the situation where *v* has not yet been exited along both *e* and *f* . On the left, the first exit from *v* was not necessarily along *e*, and we want to lower bound the probability to exit along *e* on the *n*-th visit to *v*. *v* was first reached by f' , so the weight of the edge *e* must already be at least $1 + a$, which yields the desired lower bound of $\mathbb{P}\left[Y_{n-1}'=1\right]$. On the right, the first exit from *v* was along *e*, and we want to upper bound the probability to exit along *f* , if we already know that we exit

along one of the two edges *e* and *f* . Since we look at the situation that not both edges have been used yet, *f* cannot have been used yet (also not in reverse direction, at least not on the event D_{γ}), so its weight is still *a*. The weight of *e*, on the other hand must be at least *n* + *a* if this is the *n*-th time at which we exit along one of these two edges (it could be higher, since *e* might also have been used in the reverse direction). This yields the upper bound of $\mathbb{P}[Y_{n-1} = 1]$.

The preceding paragraphs hopefully gave enough details to intuitively understand how the random walk and the variables Y and Y' should be coupled, and why \overline{Q} is an upper bound for *Q*. If not, the technical details will be given at the end of the proof. Assuming that we know that $Q(f) \le \overline{Q}(f)$ for all $f \in \gamma$, $f \ne \gamma_1$ on the event D_γ , we can now finish the prove as follows. Let $s \in \left(0, \frac{1}{2}\right)$. We have $\mathbb{E}\left[\overline{\mathcal{Q}}\left(f\right)^s\right] = \mathbb{E}\left[\overline{\mathcal{Q}}\left(f\right)^s 1\!\!1_{Y_0'=0}\right]$ $\Big] + \mathbb{E}\left[\overline{\mathcal{Q}}\left(f\right) ^{s} 1\!\!1_{Y_{0}^{\prime}=1}\right]$ i .

If $Y'_0 = 0$, then $\overline{M}_e = 1$. In addition,

$$
\mathbb{P}\left[Y'_0 = 0, \overline{M}_f = n\right] = \mathbb{P}\left[Y'_0 = 0\right] \cdot \mathbb{P}\left[Y'_n = 1\right] \cdot \prod_{j=1}^{n-1} \mathbb{P}\left[Y'_j = 0\right]
$$
\n
$$
= \underbrace{\frac{(K-1)a}{1+Ka}}_{\leq \frac{Ka}{1+Ka}} \cdot \underbrace{\frac{1+a}{2n+1+Ka}}_{\leq \frac{1+Ka}{2n}} \cdot \prod_{j=1}^{n-1} \left(1 - \frac{1+a}{2j+1+Ka}\right)
$$
\n
$$
\leq \frac{Ka}{2n} \cdot \prod_{j=1}^{n-1} \left(1 - \frac{1+a}{2j+1+Ka}\right)
$$

For the terms in the product, we have the following estimate, since $a > 0$:

$$
\frac{1+a}{2j+1+Ka} \ge \min\left\{\frac{1}{2j+1}, \frac{a}{Ka}\right\} = \min\left\{\frac{1}{2j+1}, \frac{1}{K}\right\}
$$

$$
\implies 1 - \frac{1+a}{2j+1+Ka} \le \exp\left(-\frac{1+a}{2j+1+Ka}\right) \le \exp\left(-\min\left\{\frac{1}{2j+1}, \frac{1}{K}\right\}\right)
$$

Note that $-\frac{1}{2j+1} = -\frac{1}{2j} + \frac{1}{j(4j+2)} \leq -\frac{1}{2j} + \frac{1}{4j^2}$. Therefore,

$$
\prod_{j=1}^{n-1} \left(1 - \frac{1+a}{2j+1+Ka} \right) \le \exp\left(-\sum_{j=1}^{n-1} \frac{1}{2j} + \sum_{j=1}^{n-1} \frac{1}{4j^2} + C_1(K) \right)
$$

where $0 < C_1(K) < \infty$ accounts for the fact that for only finitely many $j \in \mathbb{N}$, we have that $\min\left\{\frac{1}{2j+1},\frac{1}{K}\right\}=\frac{1}{K}$. We conclude, since $\sum_{j=1}^{\infty}\frac{1}{4j}$ $\frac{1}{4j^2}<\infty$,

$$
\prod_{j=1}^{n-1} \left(1 - \frac{1+a}{2j+1+Ka} \right) \le \exp\left(-\sum_{j=1}^{n} \frac{1}{2j} + C_2(K) \right) \le e^{C_2(K)} \exp\left(-\frac{1}{2} \ln(n) \right)
$$

We can therefore conclude

$$
\mathbb{P}\left[Y'_0=0,\overline{M}_f=n\right]\leq \frac{Ka}{2n}\cdot e^{C_2(K)}\cdot n^{-\frac{1}{2}}=C(K) \,an^{-\frac{3}{2}}
$$

This yields, noting that $s\in\left(0,\frac{1}{2}\right)$ and therefore $s-\frac{3}{2}<-1$:

$$
\mathbb{E}\left[\overline{Q}\left(f\right)^s \mathbb{1}_{Y'_0=0}\right] = \sum_{n\geq 1} n^s \mathbb{P}\left[Y'_0=0, \overline{M}_f=n\right]
$$

\$\leq \sum_{n\geq 1} C(K) an^{s-\frac{3}{2}} = C(K) a \sum_{n\geq 1} n^{s-\frac{3}{2}} \leq C(s, K) a\$

If $Y'_0 = 1$, then $\overline{M}_f = 1$. In addition,

$$
\mathbb{P}\left[Y'_0 = 1, \overline{M}_e = n\right] \le \mathbb{P}\left[Y_n = 1\right] = \frac{a}{n+1+2a} \le \frac{a}{n}
$$
\n
$$
\mathbb{E}\left[\overline{Q}\left(f\right)^s \mathbb{1}_{Y'_0 = 1}\right] = \sum_{n \ge 1} n^{-s} \mathbb{P}\left[Y'_0 = 1, \overline{M}_e = n\right] \le \sum_{n \ge 1} a n^{-1-s} < C\left(s\right) a
$$

Thus (recall that the random variables \overline{Q} were constructed to be iid),

$$
\mathbb{E}_{v_0} \left[\prod_{f \in \gamma, f \neq \gamma_1} Q(f)^s \cdot \mathbb{1}_{D_{\gamma}} \right] \leq \mathbb{E}_{v_0} \left[\prod_{f \in \gamma, f \neq \gamma_1} \overline{Q}(f)^s \cdot \mathbb{1}_{D_{\gamma}} \right] \leq \mathbb{E}_{v_0} \left[\prod_{f \in \gamma, f \neq \gamma_1} \overline{Q}(f)^s \right]
$$

\n
$$
= \prod_{f \in \gamma, f \neq \gamma_1} \mathbb{E}_{v_0} \left[\overline{Q}(f)^s \right]
$$

\n
$$
= \prod_{f \in \gamma, f \neq \gamma_1} \left(\mathbb{E} \left[\overline{Q}(f)^s \mathbb{1}_{Y'_0 = 0} \right] + \mathbb{E} \left[\overline{Q}(f)^s \mathbb{1}_{Y'_0 = 1} \right] \right)
$$

\n
$$
\leq (C(s, K) a)^{|\gamma| - 1}
$$

This finishes the proof up to the technical details of the coupling of the random variables *Y*,*Y* ′ and the domination of *Q* by *Q*.

The intuition for how the random variables *Y*,*Y* ′ should be coupled with the LERRW was already given in [Figure 3:](#page-22-0) the events $Y = 1$ and $Y' = 1$ upper and lower bound the events that a certain edge in the deterministic path γ (which was fixed at the beginning of the proof) is crossed. We now give the details of the coupling by constructing the LERRW depending on the values taken by *Y*,*Y* ′ , and some additional randomness. If the random walker already did *t* steps and is at vertex *v*, then the LERRW proceeds as follows ($v \in \gamma$ shall denote that *v* is in the interior of the path γ , i.e. not one of its endpoints):

- If $v \notin \gamma$, then the next edge is chosen as always in the LERRW: according to the current reinforced edge weights after the first *t* steps. The choice of edge is independent of the variables *Y*, *Y*^{*T*}.
- If $v \in \gamma$ and the first *t* steps of the LERRW show that D_{γ} does not occur, then, as in the previous case, the next edge is chosen according to the edge weights independently of *Y*,*Y* ′ . We can ignore the values of the variables *Y*,*Y* ′ here because we only claimed $Q \leq \overline{Q}$ if *D*_{*γ*} occurs. The first *t* steps of the LERRW are inconsistent with *D*_{*γ*} if an edge in *γ* is traversed only after its inverse or if the first arrival to some node in *γ* was not through the preceding edge in *γ*.
- If $v \in \gamma$ and the value of *Q* is already determined by the first *t* steps of the LERRW, then we again ignore *Y*, *Y'*. By *Q* being determined we mean that $v = \check{f}$ with $f \in \gamma$ and both f as well as *e* = ←− *f* ′ have already been traversed.

In all remaining cases, we thus have that the first *t* steps of the LERRW are are consistent with

*D*_{*γ*} occurring, and that $v \in \gamma$, i.e. $v = \check{f}$ with $\gamma_1 \neq f \in \gamma$. Again, call $e = \overleftarrow{f'}$ *f* ′ . As we are not in the last case, we have that *f* and *e* have not both been traversed yet. The following cases remain:

- The walker is at *v* for the first time. All incident edges thus have weight *a*, except for *e*, which has weight $1 + a$. The probability to take *e* is $\frac{1+a}{1+\deg(v)a} \ge \frac{1+a}{1+Ka} = \mathbb{P}[Y_0^0 = 1]$. The LERRW will be coupled as follows: whenever $Y'_0 = 1$, the walk exits along *e*, and sometimes also if $Y'_0 = 0$.
- Later visits to *v*, $Y'_0 = 0$ (the walk did not necessarily exit along *e* after the first visit). If *v* is visited for the *n*-th time, then the weight of *e* is still at least $1 + a$, and the total weight of all incident edges is 2*n* − 1 + deg (*v*) *a*. The probability to take *e* is therefore

$$
\frac{z(t, e) + a}{2n - 1 + \deg(v) a} \ge \frac{1 + a}{2n - 1 + Ka} = \mathbb{P}\left[Y'_{n-1} = 1\right]
$$

where $z(t, e)$ was the number of undirected traversals of e up to time t . The LERRW will now be coupled such that it takes *e* if $Y'_{n-1} = 1$, and sometimes also if $Y'_{n-1} = 0$.

• Later visits to *v*, $Y'_0 = 1$ (the walk exited along *e* after the first visit). *f* has not been traversed yet since not both *f* and *e* have been traversed already. As the first *t* steps are consistent with D_{γ} , \overline{f} has also not been traversed yet, and therefore, the weight of f is *a*. We now decide, independently of *Y*,*Y* ′ and with probabilities corresponding to the current edge weights, whether one of the edges *f* ,*e* will be used or not. In the latter case, the variables *Y*,*Y* ′ are ignored again. If *f* or *e* is used, and this happens for the *n*-th time, then $z(t, e) \ge n$ since f was not traversed yet, and since *e* was also traversed in the other direction at least once. The probability to choose *f* from *f* ,*e* is

$$
\frac{a}{z(t,e)+2a} \le \frac{a}{n+2a} = \mathbb{P}\left[Y_{n-1} = 1\right]
$$

We couple the LERRW such that if $Y_{n-1} = 0$, then *e* is chosen, and sometimes also if $Y_{n-1} = 1$.

To finish the proof, we give some details on why $Q(f) \le \overline{Q}(f)$ on D_{γ} (fix some edge $\gamma_1 \ne$ *f* $\in \gamma$ with the usual notation). Hence we assume that D_{γ} occurs.

• If $Y'_0 = 0$ (the walk did not necessarily exit along *e* after the first visit), then only the variables *Y'* are relevant. At every visit to *v* (until *e* is used), one variable *Y'* is considered. By the coupling given above, *e* is used at the latest when $Y'_{n-1} = 1$ for the first time, during the *n*-th visit to *v*. At this point, *f* can have been taken at most *n* − 1 times. We have

$$
Q(f) \le M_f \le n - 1 = \overline{M}_f = \overline{Q}(f)
$$

• If $Y'_0 = 1$ (the walk exited along *e* after the first visit), then only the variables *Y* are relevant. At every visit to *v* where *f* or *e* is used, one variable *Y* is considered. *f* is used at the earliest when $Y_{n-1} = 1$ for the first time, during the *n*-th visit where one of *f*,*e* is used. At this point, *e* will have been taken at least *n* − 1 times. We have

$$
Q(f) = \frac{1}{M_e} \le \frac{1}{n-1} = \frac{1}{\overline{M}_e} = \overline{Q}(f)
$$

4.4 Proof of Recurrence

Before we arrive at the final result, we are now able to finish the last intermediate step:

Exponential Conductance Decay ([\[4,](#page-63-2) Theorem 2]) **Theorem 4.3**

Let *G* be a graph with degree at most *K* such that the LERRW on *G* is recurrent for any choice of initial weights, let $v_0\in V$ be the starting vertex, and let $s\in \left(0,\frac14\right).$ Then, there is some $a_0 > 0$, depending on *s* and *K*, such that for initial weights $a \in (0, a_0)$ we have

$$
\mathbb{E}_{v_0}\left[c_e^s\right] \leq 2K\left(C\left(s,K\right)\sqrt{a}\right)^{\text{dist}(v_0,e)}
$$

where *C* (*s*, *K*) solely depends on *K* and *s*.

We have *Proof*

$$
\mathbb{E}_{v_0} [c_e^s] = \sum_{\gamma \in \Gamma_e} \mathbb{E}_{v_0} \left[c_e^s \cdot \mathbb{1}_{D_\gamma} \right] \leq \sum_{\gamma \in \Gamma_e} \mathbb{E}_{v_0} \left[\left(\frac{c_e}{c_{\gamma_1}} \right)^s \cdot \mathbb{1}_{D_\gamma} \right]
$$
\n
$$
\leq \sum_{\gamma \in \Gamma_e} \mathbb{E}_{v_0} \left[\prod_{f \in \gamma, f \neq \gamma_1} \left(\frac{R(f)}{Q(f)} \right)^{2s} \cdot \mathbb{1}_{D_\gamma} \right]^{\frac{1}{2}} \mathbb{E}_{v_0} \left[\prod_{f \in \gamma, f \neq \gamma_1} Q(f)^{2s} \cdot \mathbb{1}_{D_\gamma} \right]^{\frac{1}{2}}
$$
\n
$$
\leq \sum_{\gamma \in \Gamma_e} \left(C(2s)^{|\gamma|-1} \right)^{\frac{1}{2}} \left((C(2s,K)a)^{|\gamma|-1} \right)^{\frac{1}{2}}
$$
\n
$$
= \sum_{\gamma \in \Gamma_e} \left(\frac{C(2s)^{\frac{1}{2}} C(2s,K)^{\frac{1}{2}}}{\pi^{s}} \sqrt{a} \right)^{|\gamma|-1}
$$

where \circledast holds by [Lemma 4.1](#page-19-1) and [Lemma 4.2.](#page-21-2) Now, choose a_0 such that $KC_0\sqrt{a_0} = \frac{1}{2}$ (1). Then, for $a < a_0$ (2), we have, since any $\gamma \in \Gamma_e$ has length at least $|\gamma| = dist(v_0, e) + 1$ (3), and since there are at most K^l paths of length l $(\mathcal{4})$:

$$
\mathbb{E}_{v_0}[c_e^s] \leq \sum_{\gamma \in \Gamma_e} (C_0 \sqrt{a})^{|\gamma|-1} \stackrel{\text{(3)}}{=} \sum_{l \geq \text{dist}(v_0, e) + 1} \sum_{\gamma \in \Gamma_e, |\gamma| = l} (C_0 \sqrt{a})^{l-1}
$$
\n
$$
\stackrel{\text{(4)}}{\leq} \sum_{l \geq \text{dist}(v_0, e) + 1} K \cdot (KC_0 \sqrt{a})^{l-1} = K \cdot (KC_0 \sqrt{a})^{\text{dist}(v_0, e)} \cdot \sum_{l \geq 0} (KC_0 \sqrt{a})^l
$$
\n
$$
\stackrel{\text{(1)}\text{(2)}}{<} K \cdot (KC_0 \sqrt{a})^{\text{dist}(v_0, e)} \cdot \sum_{l \geq 0} \left(\frac{1}{2}\right)^l = 2K (KC_0 \sqrt{a})^{\text{dist}(v_0, e)}
$$

We can now finally prove the result we are actually interested in:

Recurrence on Bounded Degree Graphs ([\[4,](#page-63-2) Theorem 1]) **Theorem 4.4**

Let $K \in \mathbb{N}_{\geq 1}$. Then, there exists $a_0 > 0$ such that if the graph *G* has degree at most *K*, then the LERRW on *G* with initial weights set to $a \in (0, a_0)$ is recurrent. The corresponding RWRE is a.s. positive recurrent, but this does not necessarily imply that the expected return time to the starting vertex is finite.

27

Proof The idea is to apply [Theorem 4.3](#page-26-1) to the LERRW on finite balls of the infinite graph *G*. Call the starting vertex v_0 and let $B_R(v_0) := \{v \in V : dist(v_0, v) \le R\}$ be the set of vertices of distance at most *R* to the starting vertex. Consider the LERRW $X^{(R)}$ on the finite ball $B_R(v_0)$. Since $B_R(v_0)$ is finite, $X^{(R)}$ is recurrent for any choice of initial weights, and [Theorem 4.3](#page-26-1) is applicable. By [Theorem 3.1,](#page-12-3) *X*(*R*) is equal in law to a RWRE with random conductances $\mathbf{c}^{(R)} = \left(c_e^{(R)} \right)$ *e*∈*B_{<i>R*}(*v*₀)</sub>. Denote the mixing measure (giving the distribution of $\mathbf{c}^{(R)}$) by $\mu^{(R)}$. The measures $\mu^{(R)}$ are a sequence of measures on the set of possible conductance vectors.

The following calculation will, next to other results, show that for fixed initial weights *a*, the measures $\mu^{(R)}$ are tight (see end of the proof). So, by [Prokhorov's theorem,](https://en.wikipedia.org/wiki/Prokhorov%27s_theorem) there is a subsequence converging to some measure *µ*. The first *R* steps of the LERRW on the whole graph *G* are equal in law to the first *R* steps of the LERRW on $B_R(v_0)$. So the LERRW on *G* has the same distribution as a RWRE with mixing measure *µ*.

Fix $s \in \left(0, \frac{1}{4}\right)$ and let $e \in B_R(v_0)$ be an edge in the finite ball of radius R. By Markov's inequality, and by [Theorem 4.3,](#page-26-1) for initial weights *a* small enough, we have

$$
\mu^{(R)}\left(c_{e}^{(R)} > Q\right) = \mu^{(R)}\left(\left(c_{e}^{(R)}\right)^{s} > Q^{s}\right) \leq \frac{\mathbb{E}\left[\left(c_{e}^{(R)}\right)^{s}\right]}{Q^{s}} \leq \frac{2K\left(C\left(s, K\right) \sqrt{a}\right)^{\text{dist}(v_{0}, e)}}{Q^{s}}
$$

Choose $Q = (2K)^{-\text{dist}(v_0, e)}$ and note that there are at most K^{l+1} edges at distance *l* from v_0 . Then

$$
\mu^{(R)} \left(\exists e \in B_R \left(v_0 \right) : \text{dist} \left(v_0, e \right) = l \text{ and } c_e^{(R)} > (2K)^{-l} \right)
$$
\n
$$
\leq \sum_{e \in B_R \left(v_0 \right) : \text{dist} \left(v_0, e \right) = l} \mu^{(R)} \left(c_e^{(R)} > (2K)^{-l} \right)
$$
\n
$$
\leq K^{l+1} 2K \left(C \left(s, K \right) \sqrt{a} \right)^l (2K)^{sl} = 2K^2 \left(2^s K^{1+s} C \left(s, K \right) \sqrt{a} \right)^l
$$
\n(4.4)

If we now choose a_0 such that $2^{s}K^{1+s}C(s,K)\sqrt{a_0} \leq \frac{1}{2}$ and such that a_0 is smaller than the bound given by [Theorem 4.3,](#page-26-1) then, for intital weights $a \in (0, a_0)$,

$$
\mu^{(R)}\left(\exists e \in B_R(v_0) : \text{dist}(v_0, e) = l \text{ and } c_e^{(R)} > (2K)^{-l}\right) < 2K^2\left(\frac{1}{2}\right)^l = K^2 2^{1-l}
$$

This bound is uniform in *R*, so it holds for all measures $\mu^{(R)}$, and in consequence also for the limit *µ* (by the [Portmanteau theorem\)](https://en.wikipedia.org/wiki/Convergence_of_measures#Weak_convergence_of_measures). Since *K* 22 1−*l* is summable in *l*, Borel-Cantelli implies that the probability that infinitely many of the events $\left\{c_e > (2K)^{-\text{dist}(v_0,e)}\right\}$ occur is 0. In other words, for all but a finite number of edges, $c_e \leq \left(2K\right)^{-\operatorname{dist}(v_0,e)}$ a.s. But then,

$$
\sum_{e \in E} c_e = \sum_{l \ge 0} \sum_{e \in E: \text{dist}(v_0, e) = l} c_e \le A + \sum_{l \ge 0} K^{l+1} (2K)^{-l} = A + K \cdot \sum_{l \ge 0} 2^{-l} < \infty
$$

where *A* accounts for the finitely many exceptions to the *c^e* bound. So the RWRE is a.s. positive recurrent.

To finish the proof, we will show how tightness follows from the above calculation for fixed initial weights *a*. Let $\varepsilon > 0$. We have to show there is a compact set $K_{\varepsilon} \subseteq \mathbb{R}_{\geq 0}^E$ such that for all *R*, the measure of the complement $\mu^{(R)}(K_{\varepsilon}^C)$ is smaller than *ε*. If we choose $Q = (\lambda K)^{-\text{dist}(v_0, e)}$

in [Equation 4.4,](#page-27-0) we get

$$
\mu^{(R)}\left(\exists e \in B_R\left(v_0\right) : \text{dist}\left(v_0, e\right) = l \text{ and } c_e^{(R)} > (\lambda K)^{-l}\right)
$$

$$
\leq 2K^2 \left(\lambda^s K^{1+s} C\left(s, K\right) \sqrt{a}\right)^l
$$

Thus,

$$
\mu^{(R)}\left(\exists e \in B_R\left(v_0\right):c_e^{(R)} > (\lambda K)^{-\operatorname{dist}(v_0,e)}\right)
$$
\n
$$
\leq \sum_{l\geq 0} \mu^{(R)}\left(\exists e \in B_R\left(v_0\right): \operatorname{dist}\left(v_0, e\right) = l \text{ and } c_e^{(R)} > (\lambda K)^{-l}\right)
$$
\n
$$
\stackrel{\circledast}{=} \sum_{l\geq 1} \mu^{(R)}\left(\exists e \in B_R\left(v_0\right): \operatorname{dist}\left(v_0, e\right) = l \text{ and } c_e^{(R)} > (\lambda K)^{-l}\right)
$$
\n
$$
\leq 2K^2 \sum_{l\geq 1} \left(\lambda^s K^{1+s} C\left(s, K\right) \sqrt{a}\right)^l = \frac{2K^2 \lambda^s K^{1+s} C\left(s, K\right) \sqrt{a}}{1 - \lambda^s K^{1+s} C\left(s, K\right) \sqrt{a}}
$$

For \circledast , recall that we normalized the conductances to $c_{v_0} = 1$, so for any edge *e* with distance 0 to v_0 , it holds that $c_e \leq 1$. We can now choose λ such that

$$
\lambda < \left(\frac{\varepsilon}{K^{1+s}C\left(s,K\right)\sqrt{a}\left(2K^2+\varepsilon\right)}\right)^{\frac{1}{s}}
$$

Then

$$
\mu^{(R)}\left(\exists e \in B_R(v_0):c_e^{(R)} > (\lambda K)^{-\operatorname{dist}(v_0,e)}\right) < \varepsilon
$$

which holds independently of *R*. So we can choose

$$
K_{\varepsilon} = \prod_{e \in E} \left[0, (\lambda K)^{-\operatorname{dist}(v_0, e)} \right]
$$

which is compact. Note that *K*^{*ε*} grows when *ε* gets smaller, because $λ$ must also be chosen smaller in this case. smaller in this case.

5 Reinforced Random Walk with Multiple Walkers

Here, we consider multiple walkers on a single environment influencing each other.

5.1 A Two-Player Urn

We start with a very simple model, the linearly edge-reinforced random walk with two walkers on a segment of **Z** with three nodes.

Figure 4: Edge weights of the line segment at time *n*

5.1.1 Alternating Players

We first define the following dynamics:

- There are two walkers $X^{(1)}$ and $X^{(2)}$, which both start at the node in the center, i.e. $X_0^{(1)}$ = $0, X_0^{(2)} = 0.$
- Initially, both edge weights are 1. We denote the edge weight of the left edge at time *n* by $w(n,0)$, the weight of the right edge by $w(n,1)$.
- Whenever an edge is crossed by either of the walkers, its weight is increased by 1, so we set $W(n) = 1 + n$ in terms of [Definition 2.3.](#page-10-0)
- The walkers move alternately, i.e. at odd time steps, walker 1 moves (in particular, walker 1 moves first at step 1) and at even time steps, walker 2 moves. This implies that the walkers will meet at the node in the center every four steps.
- • When a walker at the node in the center is about to move, he chooses the edge to traverse with probability proportional to the respective edge weight.

Proof We can calculate:

$$
\mathbb{E}\left[\frac{w(4n+4,0)}{w(4n+4,0)+w(4n+4,1)}\middle| w(4n,0) = a, w(4n,1) = b\right]
$$
\n
$$
=\frac{a}{a+b}\frac{a+1}{a+b+1}\frac{a+4}{a+b+4} + \frac{a}{a+b}\frac{b}{a+b+1}\frac{a+2}{a+b+4} + \frac{b}{a+b}\frac{a+2}{a+b+1}\frac{a+b+4}{a+b+4} + \frac{b}{a+b}\frac{b+1}{a+b+1}\frac{a}{a+b+4}
$$
\n
$$
=\frac{a}{a+b}\cdot\frac{(a+1)(a+4)+2b(a+2)+b(b+1)}{(a+b+1)(a+b+4)} = \frac{a}{a+b}
$$

 \Box

 $\frac{w(n,0)}{0\right)+w(n,1)}$ converges almost surely for *n* → ∞ **Corollary 5.2** $\frac{w(n,0)+w(n,1)}{w(n,0)}$ We immediately get from [Lemma 5.1](#page-29-4) that $\frac{w(4n,0)}{w(4n,0)+w(4n,1)}$ converges a.s. and then, it is easy to Proof see that also $\frac{w(n,0)}{w(n,0)+w(n,1)}$ converges. Define the random variable $Y := \lim_{n \to \infty} \frac{w(n,0)}{w(n,0) + w(n,1)}$. Then $Y \in [0,1]$ has a density Conjecture 5.3 w.r.t. the Lebesgue measure on [0, 1]. *Y* is not Beta-distributed.

5.1.2 Random Player Selection

Consider next the case where at every step, we choose uniformly at random (independent of all other steps) which of the two walkers moves. [Lemma 5.4](#page-30-1) shows that the expected time to meet again in the middle, if both walkers start in the center, is 4, just as in the previous case. Of course, the difference now is that the next meeting time is random. We call $\tau_0 = 0$ and $\tau_n = \inf \Big\{ k > \tau_{n-1} : X_k^{(1)} = X_k^{(2)} = 0 \Big\}.$

For any $n \ge 0$ and any $l \ge 1$, it holds that $\mathbb{P}[\tau_{n+1} - \tau_n = 2l] = 2^{-l}$ and $\mathbb{E}[\tau_{n+1} - \tau_n] = 4$. Lemma 5.4

We consider a MC consisting of three states and coupled with the edge-reinforced random *Proof* walk. The MC is in state *s*center if both walkers are in the center, in state *s*_{mixed} if one walker is in the center and the other in either of the two outer nodes, and in state *s*_{none} if none of the walkers is in the center. It is easy to verify that this is indeed a MC with the following transition probabilities:

Let $n \geq 0$. At time τ_n , both walkers are in the center and the MC is therefore in state *s*_{center}. The time $\tau_{n+1} - \tau_n$ corresponds to the time needed to return again to the state s_{center} . It is now a standard calculation to show $\mathbb{P}[\tau_{n+1} - \tau_n = 2l] = 2^{-l}$. Consequently,

$$
\mathbb{E}\left[\tau_{n+1}-\tau_n\right] = \sum_{l\geq 1} 2^{-l} \cdot 2l = \sum_{l\geq 1} l \cdot \left(\frac{1}{2}\right)^{l-1} = \frac{1}{\left(1-\frac{1}{2}\right)^2} = 4
$$

We next want to prove, in [Lemma 5.7,](#page-35-0) that the fraction of the left edge weight, $\frac{w(\tau_n,0)}{w(\tau_n,0)+w(\tau_n,1)}$, is a martingale, as it was in the previous case and as in the one-player urn with the difference that we look at the fraction not at every time step, but at certain stopping times. We use two lemmas for the proof and introduce the following notation:

(1) We look at the expectation of the proportion of the left edge weight multiplied by the

indicator of the event that the walkers need 2*l* steps to meet again:

$$
\mathbb{E}_{a,b,l} := \mathbb{E}\left[\frac{w\left(\tau_{n+1},0\right)}{w\left(\tau_{n+1},0\right)+w\left(\tau_{n+1},1\right)}\cdot \mathbb{1}_{\{\tau_{n+1}-\tau_n=2l\}}\;\middle|\; w\left(\tau_n,0\right)=a,w\left(\tau_n,1\right)=b\right]
$$

(2) We also consider the probability that the last walker which returns to the center comes from the left node, again intersected with the event that the walkers need 2*l* steps to meet again:

*L*_{*n*} := {the walker returning to the center at time τ_{n+1} comes from the left node} $q_{a,b,l} := \mathbb{P} [L_n \text{ occurs and } \tau_{n+1} - \tau_n = 2l \mid w(\tau_n, 0) = a, w(\tau_n, 1) = b]$

Lemma 5.5 We have the following recursive equations:

$$
\mathbb{E}_{a,b,l+1} = \frac{1}{2} \mathbb{E}_{a,b,l} + \frac{1}{(a+b+2l-1)(a+b+2l+2)} (\mathbb{E}_{a,b,l} - q_{a,b,l})
$$

$$
q_{a,b,l+1} = \frac{1}{2} q_{a,b,l} + \frac{1}{4} \cdot \frac{a+b+2l}{a+b+2l-1} (\mathbb{E}_{a,b,l} - q_{a,b,l})
$$

Proof The following notation will be useful. We define a path of the two walkers as a sequence of the symbols 1_l , 1_r , 2_l , 2_r which correspond to the first (respectively second) walker moving left and right, where we assume that both walkers start in the center. The set Path_{2l} contains all the paths of length 2*l* (a sequence of 2*l* symbols) such that the first time at which both walkers are in the center at the same time again is at the end of the path. Note that any such path must be of even length since each walker can only be in the center after having made an even number of movements.

For, $\rho \in \text{Path}_{2l}$, we set $x(\rho)$ to be the number of traversals of the left edge when the path ρ is taken and we can now write

$$
\mathbb{E}_{a,b,l} = \sum_{\rho \in \text{Path}_{2l}} \underbrace{\mathbb{P}\left[\text{the walksers move according to } \rho \mid w\left(\tau_n, 0\right) = a, w\left(\tau_n, 1\right) = b\right]}_{=: \mathbb{P}_{\rho,a,b}} \cdot \frac{a + x\left(\rho\right)}{a + b + 2l}
$$

new step: center walker goes left new step: center walker goes right

To prove the recurrence relation, we build paths of length $2(l + 1)$ out of paths of length 2*l*.

 -1 0 1 *a* + *x* (*ρ*) − 1 $\left(\begin{array}{cc} b + 2l - x \\ 0 \end{array}\right)$ *a* + *x* (*ρ*) − 1 $\left(\begin{array}{cc} b + 2l - x \\ 0 \end{array}\right)$ -1 0 1 or or -1 0 1 $a + x(\rho)$ $\lambda b + 2l - x(\rho) - 1$ -1 0 1 $a + x(\rho) \longrightarrow b + 2l - x(\rho)$

Figure 5: Possible walker locations after 2*l* − 1 steps of a path *ρ* of length 2*l*

For any possible path of the walkers, at any uneven time step, there will be one walker

which is in the center and one which is in one of the outer nodes. The paths in Path_{2*l*} have the additional condition that at any even time step (except for the beginning and end), both walkers have to be in the outer nodes, not necessarily the same one. We can therefore construct all paths in Path_{2(*l*+1)} as follows: for every $\rho \in$ Path_{2*l*}, we take the path ρ up to time 2*l* − 1. The possible walker locations and edge weights after 2*l* − 1 steps are depicted in [Figure 5.](#page-31-0) The path ρ now continues with the walker in the outer node moving back to the center. To get a path of length $2(l+1)$, the walker in the center node has to move instead in the next step, and then both walkers will return to the center in either order. Out of ρ , we can thus construct 4 new paths of length $2(l + 1)$: we have two choices for which way the center walker moves in step 2*l*, and then two choices for the order in which the walkers return. Note that we do not construct any path twice, since any two different paths in Path_{2*l*} must already differ somewhere in the first 2*l* − 1 steps.

Now, we look at how this changes the outcome (the final edge weights) and the probability of the path. Let $\rho \in \text{Path}_{2l}$.

- If the outer walker is on the left after $2l 1$ steps (this is a condition on ρ):
	- **–** If the center walker should move left in step 2*l* (this is one choice for creating the new path): the probability $\mathbb{P}_{\rho,a,b}$ of ρ is a product over the probability that the walker indicated in ρ is chosen at the respective step (which is always $\frac{1}{2}$) and the probability that the walker moves in the direction indicated by ρ (this can either be 1, if the walker moves back to the center, or a fraction depending on the edge weights). In our new modified path of length $2(l+1)$, the first difference is that we choose a different walker to move in step 2*l*. This event has probability $\frac{1}{2}$, but this is the same as chosing the original walker, hence this part is already included in the product giving the probability $\mathbb{P}_{\rho,a,b}$ of ρ .

Next, the probability for the center walker to go left, if he is chosen to move at step 2*l*, is given by $\frac{a+x(\rho)-1}{a+b+2l-1}$ (compare with [Figure 5\)](#page-31-0). This factor is new and has to be added to the product. In the next step, one of the two walkers is chosen and will move back to the center. It is irrelevant which one moves, so we get only a new factor of 1. Finally, in step $2(l+1)$, the walker which is still in an outer node has to be chosen to move back into the center. This happens with probability $\frac{1}{2}$, and this is the final factor to be added to the product. Our new path therefore has probability

$$
\mathbb{P}_{\rho,a,b} \cdot \frac{1}{2} \cdot \frac{a+x(\rho)-1}{a+b+2l-1}
$$

The new path will end with the following ratio of the left edge weight divided by the total edge weights:

$$
\frac{a+x(\rho)+2}{a+b+2l+2}
$$

since the left edge will be traversed twice more by the center walker.

– If the center walker should move right in step 2*l*:

new path probability:
$$
\mathbb{P}_{\rho,a,b} \cdot \frac{1}{2} \cdot \frac{b+2l-x(\rho)}{a+b+2l-1}
$$
, new outcome: $\frac{a+x(\rho)}{a+b+2l+2}$

• If the outer walker is on the right after 2*l* − 1 steps:

– If the center walker should move left in step 2*l*:

new path probability: $\mathbb{P}_{\rho,a,b} \cdot \frac{1}{2}$ $\frac{1}{2} \cdot \frac{a+x(\rho)}{a+b+2l}$ $\frac{a + x(\rho)}{a + b + 2l - 1}$, new outcome: $\frac{a + x(\rho) + 2}{a + b + 2l + 2}$ $a + b + 2l + 2$

– If the center walker should move right in step 2*l*:

new path probability: $\mathbb{P}_{\rho,a,b} \cdot \frac{1}{2}$ $\frac{1}{2} \cdot \frac{b+2l-x(\rho)-1}{a+b+2l-1}$ $\frac{a+2l-x(\rho)-1}{a+b+2l-1}$, new outcome: $\frac{a+x(\rho)}{a+b+2l-1}$ $a + b + 2l + 2$

For $\rho \in \text{Path}_{2l}$, we write left $(\rho) = 1$ if the outer walker after step 2*l* − 1 is on the left, and left (ρ) = 0 otherwise. Using the recursive path construction above, we get that

$$
\mathbb{E}_{a,b,l+1} = \sum_{\rho \in \text{Path}_{2(l+1)}} \mathbb{P}_{\rho,a,b} \cdot \frac{a + x(\rho)}{a + b + 2l + 2} =
$$
\n
$$
\sum_{\rho \in \text{Path}_{2l}} \mathbb{P}_{\rho,a,b} \cdot \frac{1}{2} \cdot \left(\text{left}(\rho) \cdot \left(\frac{a + x(\rho) - 1}{a + b + 2l - 1} \cdot \frac{a + x(\rho) + 2}{a + b + 2l + 2} + \frac{b + 2l - x(\rho)}{a + b + 2l - 1} \cdot \frac{a + x(\rho)}{a + b + 2l + 2} \right) + (1 - \text{left}(\rho)) \cdot \left(\frac{a + x(\rho)}{a + b + 2l - 1} \cdot \frac{a + x(\rho) + 2}{a + b + 2l + 2} + \frac{b + 2l - x(\rho) - 1}{a + b + 2l - 1} \cdot \frac{a + x(\rho)}{a + b + 2l + 2} \right) \right)
$$

Using this expression for $\mathbb{E}_{a,b,l+1}$, we can calculate (a simple, but longer calculation which we skip here, see [https://bit.ly/3trqm2r](https://www.wolframalpha.com/input?i=simplify+%28a%2Bx-1%29%28a%2Bx%2B2%29%28a%2Bb%2B2l%29+%2B+%28b%2B2l-x%29%28a%2Bx%29%28a%2Bb%2B2l%29+-+%28a%2Bx%29%28a%2Bb%2B2l-1%29%28a%2Bb%2B2l%2B2%29) and [https://bit.ly/391ma2o\)](https://www.wolframalpha.com/input?i=simplify+%28a%2Bx%29%28a%2Bx%2B2%29%28a%2Bb%2B2l%29+%2B+%28b%2B2l-x-1%29%28a%2Bx%29%28a%2Bb%2B2l%29+-+%28a%2Bx%29%28a%2Bb%2B2l-1%29%28a%2Bb%2B2l%2B2%29):

$$
\mathbb{E}_{a,b,l+1} - \frac{1}{2} \mathbb{E}_{a,b,l} = \sum_{\rho \in \text{Path}_{2l}} \mathbb{P}_{\rho,a,b} \cdot \frac{1}{(a+b+2l-1) (a+b+2l+2)} \cdot \left(\text{left}(\rho) \cdot \frac{x(\rho) - b - 2l}{a+b+2l} + (1 - \text{left}(\rho)) \cdot \frac{a+x(\rho)}{a+b+2l} \right) \right)
$$
\n
$$
= \frac{1}{(a+b+2l-1) (a+b+2l+2)} \cdot \sum_{\rho \in \text{Path}_{2l}} \mathbb{P}_{\rho,a,b} \cdot \left(\text{left}(\rho) \cdot \frac{-a-b-2l}{a+b+2l} + \frac{a+x(\rho)}{a+b+2l} \right) \right)
$$
\n
$$
= \frac{1}{(a+b+2l-1) (a+b+2l+2)} \cdot \left(\mathbb{E}_{a,b,l} - \sum_{\rho \in \text{Path}_{2l}} \mathbb{P}_{\rho,a,b} \cdot \text{left}(\rho) \right) \right)
$$
\n
$$
= \frac{1}{(a+b+2l-1) (a+b+2l+2)} \cdot (\mathbb{E}_{a,b,l} - q_{a,b,l})
$$

This proves the first equation. For the second equation, we use the same strategy. For our newly constructed paths, we have to calculate the probability that the last walker to move to the center comes from the left node. Let $\rho \in \text{Path}_{2l}$.

- If the outer walker is on the left after $2l 1$ steps (this is a condition on ρ):
	- **–** If the center walker should move left in step 2*l* (this is one choice for creating the new path):

new path probability: $\mathbb{P}_{\rho,a,b} \cdot \frac{1}{2}$ $\frac{1}{2} \cdot \frac{a+x(\rho)-1}{a+b+2l-1}$ $\frac{a+b+2l-1}{a+b+2l-1}$, probability of *L_n*: 1

– If the center walker should move right in step 2*l*:

new path probability:
$$
\mathbb{P}_{\rho,a,b} \cdot \frac{1}{2} \cdot \frac{b+2l-x(\rho)}{a+b+2l-1}
$$
, probability of L_n : $\frac{1}{2}$

 \Box

- If the outer walker is on the right after 2*l* − 1 steps:
	- **–** If the center walker should move left in step 2*l*:

new path probability:
$$
\mathbb{P}_{\rho,a,b} \cdot \frac{1}{2} \cdot \frac{a+x(\rho)}{a+b+2l-1}
$$
, probability of L_n : $\frac{1}{2}$

– If the center walker should move right in step 2*l*:

new path probability:
$$
\mathbb{P}_{\rho,a,b} \cdot \frac{1}{2} \cdot \frac{b+2l-x(\rho)-1}{a+b+2l-1}
$$
, probability of L_n : 0

Therefore

$$
q_{a,b,l+1} - \frac{1}{2}q_{a,b,l} = \sum_{\rho \in \text{Path}_{2l}} \mathbb{P}_{\rho,a,b} \cdot \frac{1}{2} \cdot \frac{1}{a+b+2l-x} \cdot \text{left}(\rho) + \frac{1}{2} \cdot \frac{b+2l-x(\rho)}{a+b+2l-1} \cdot \text{left}(\rho) + \frac{1}{2} \cdot \frac{a+x(\rho)}{a+b+2l-1} \cdot (1 - \text{left}(\rho)) - \text{left}(\rho) \right) \right) \newline + \frac{1}{2} \cdot \frac{a+x(\rho)}{a+b+2l-1} \cdot (1 - \text{left}(\rho)) - \text{left}(\rho) \right) \newline = \frac{1}{2} \cdot \sum_{\rho \in \text{Path}_{2l}} \mathbb{P}_{\rho,a,b} \cdot \left(-\frac{1}{2} \cdot \frac{a+b+2l}{a+b+2l-1} \cdot \text{left}(\rho) + \frac{1}{2} \cdot \frac{a+x(\rho)}{a+b+2l-1} \right) \newline = \frac{1}{4} \cdot \frac{a+b+2l}{a+b+2l-1} \cdot \sum_{\rho \in \text{Path}_{2l}} \mathbb{P}_{\rho,a,b} \cdot \left(\frac{a+x(\rho)}{a+b+2l} - \text{left}(\rho) \right) \newline = \frac{1}{4} \cdot \frac{a+b+2l}{a+b+2l-1} \cdot (\mathbb{E}_{a,b,l} - q_{a,b,l})
$$

The expectation of the proportion of the left edge weight and the probability that the last **Corollary 5.6** walker returning to the center comes from the left coincide, and:

$$
\mathbb{E}_{a,b,l}=q_{a,b,l}=\frac{1}{2^l}\cdot\frac{a}{a+b}
$$

We use [Lemma 5.5.](#page-31-1) Let us first calculate $\mathbb{E}_{a,b,1}$ and $q_{a,b,1}$. There are four possible paths of *Proof* length 2 which end again with both walkers in the center: first, we choose which of the two walker moves, and this walker can than either move left or right and then back to the center. Since it is irrelevant which walker we choose in the beginning, we can disregard which walker moves. We thus get:

- With probability $\frac{a}{a+b} \cdot \frac{1}{2}$ the walker which moves in the first step moves left and is then chosen again to move back to the center in the next step. In this case, *Lⁿ* occurs and the resulting edge weight ratio is $\frac{a+2}{a+b+2}$.
- With probability $\frac{b}{a+b} \cdot \frac{1}{2}$ the walker which moves in the first step moves right and is then chosen again to move back to the center in the next step. In this case, *Lⁿ* does not occur and the resulting edge weight ratio is $\frac{a}{a+b+2}$.

We see directly that $q_{a,b,1} = \frac{1}{2} \cdot \frac{a}{a+b}$ and that

$$
\mathbb{E}_{a,b,1} = \frac{1}{2} \cdot \left(\frac{a}{a+b} \cdot \frac{a+2}{a+b+2} + \frac{b}{a+b} \cdot \frac{a}{a+b+2} \right) = \frac{1}{2} \cdot \frac{a}{a+b}
$$

The remaining proof is now a simple induction using [Lemma 5.5,](#page-31-1) where it should be noted that $\mathbb{E}_{a,b,l} - q_{a,b,l} = 0$ under the induction assumption. \Box

Lemma 5.7 The random variables
$$
\frac{w(\tau_n, 0)}{w(\tau_n, 0) + w(\tau_n, 1)}
$$
 for $n \ge 0$ form a martingale.

Proof We have by [Corollary 5.6:](#page-34-0)

$$
\mathbb{E}\left[\frac{w\left(\tau_{n+1},0\right)}{w\left(\tau_{n+1},0\right)+w\left(\tau_{n+1},1\right)}\cdot\mathbb{1}_{\{\tau_{n+1}-\tau_n=2l\}}\middle|\, w\left(\tau_n,0\right)=a,w\left(\tau_n,1\right)=b\right]=2^{-l}\cdot\frac{a}{a+b} \implies \mathbb{E}\left[\frac{w\left(\tau_{n+1},0\right)}{w\left(\tau_{n+1},0\right)+w\left(\tau_{n+1},1\right)}\middle|\, w\left(\tau_n,0\right)=a,w\left(\tau_n,1\right)=b\right]=\frac{a}{a+b}\cdot\sum_{l\geq 1}2^{-l}=\frac{a}{a+b} \quad \square
$$

Lemma 5.8 Assume that $w(0,0) = a > 0, w(0,1) = b > 0$. Then, the random variables $M_n :=$ *w*(*τn*,0) $\frac{w(\tau_n,\sigma)}{w(\tau_n,0)+w(\tau_n,1)}$ take values in the set

$$
R := \left\{ \frac{a+2x}{a+b+2l} \middle| x \in \mathbb{N}_{\geq 0}, l \in \mathbb{N}_{\geq 1}, x \leq l \right\}
$$

and, for every $r \in R$ and every $n \in \mathbb{N}_{\geq 1}$, we have that $\mathbb{P}[M_n = r] > 0$. The set R is dense in $[0, 1]$.

Proof It is immediately clear that M_n can take only values in the set *R*: $w(\tau_n, 0)$ will be equal to *a* plus the number of crossings of the left edge until time *τn*. Since both walkers are in the center at time τ_n , this number of edge crossings has to be even, hence we can write $w(\tau_n, 0) = a + 2x$ for some $x \in \mathbb{N}_{\geq 0}$. On the other hand, the total edge weight also must have increased by an even number since both walkers have to do an even number of steps before meeting again in the center.

We now want to construct a sequence of walker steps leading to any possible outcome $\frac{a+2x}{a+b+2l}$ with positive probability. First, note that $w(\tau_n, 0) + w(\tau_n, 1) = a + b + \tau_n$. Since $\tau_n \geq 2n$, we cannot have a sequence of length 2*l* leading to our desired outcome when *l* < *n*. In this case, we have to reach $\frac{a+2x}{a+b+2l}$ by a longer sequence. We have, for any *k*,

$$
\frac{a+2x}{a+b+2l} = \frac{k(a+2x)}{k(a+b+2l)} = \frac{a+2(kx+\frac{1}{2}(k-1)a)}{a+b+2(kx+\frac{1}{2}(k-1)(a+b))}
$$

To reach our desired fraction, we thus want a sequence of length 2 $\left(kl+\frac{1}{2}\left(k-1\right)\left(a+b\right)\right)\geq$ 2*n*. In addition, $(k-1)(a+b)$ and $(k-1)a$ should both be even. It thus suffices to choose an uneven *k* which is large enough such that $kl + \frac{1}{2}(k-1)(a+b) \geq n$. This allows us to assume w.l.o.g. that $l \geq n$.

What remains to show is that we can construct a sequence of length 2*l* such that after all 2*l* steps, the walkers meet again in the center for the *n*-the time and such that the left edge is traversed 2*x* times. Since any finite sequence of possible walker movements has positive probability, this concludes the proof. Finding such a sequence is easy: in the first 2 (*n* − 1) steps, one walker moves min $\{x, n-1\}$ times to the left node and back, and in the remaining steps (if there are any), the same walker moves to the right node and back. Then, in the $(2n - 1)$ -th step, the same walker moves to the left node, if the left edge was not yet crossed 2*x* times, and otherwise to the right node. The walker then remains there until returning to the center in the 2*l*-th step. In the remaining steps of the complete sequence of length 2*l*, the other walker moves in such a way as to reach a total number of 2*x* left edge crossings. This leads to the desired fraction at time *τn*.

It is easy to see that *R* is dense in [0, 1]. Indeed, choosing $x = k \cdot p$ and $l = k \cdot q$ for increasing *k* shows that we can find sequences in *R* converging to any rational number, and these are already dense in [0, 1]. П

 $\frac{w(n,0)}{w(n,0)+w(n,1)}$ converges almost surely for *n* → ∞, and the limit is identical to the limit of **Corollary 5.9** *w*(*τn*,0) $\overline{w(\tau_n,0)+w(\tau_n,1)}$.

We use the following notation: *Proof*

$$
F_n := \frac{w(n,0)}{w(n,0) + w(n,1)} \qquad M_n := F_{\tau_n} = \frac{w(\tau_n,0)}{w(\tau_n,0) + w(\tau_n,1)} \qquad M_\infty := \lim_{n \to \infty} M_n
$$

Let $\varepsilon > 0$ and consider the events

$$
A_n := \{ |F_k - M_\infty| > \varepsilon \text{ for some } k \in [\tau_n, \tau_{n+1}]\}
$$

It is sufficient to show that only finitely many of the events A_n can occur a.s. Further set

$$
B_n := \left\{ |F_k - M_n| > \frac{\varepsilon}{2} \text{ for some } k \in [\tau_n, \tau_{n+1}] \right\}
$$

Since M_n converges a.s. by [Lemma 5.7,](#page-35-0) there is some (random) $N \in \mathbb{N}$ such that for all $n \geq N$, it holds that $|M_n - M_\infty| < \frac{\varepsilon}{2}$. For $n \geq N$, the occurrence of A_n implies that B_n occurs as well, so it is sufficient to show that only finitely many of the events B_n can occur.

Now, at time τ_n , the random walkers must have moved at least 2*n* times, so $w(\tau_n, 0)$ + $w(\tau_n,1) \ge 2n$. If $|F_k - M_n| = |F_k - F_{\tau_n}| > \frac{\varepsilon}{2}$ for some $k \in [\tau_n, \tau_{n+1}]$, it is therefore necessary that at least εn steps were made by the walkers between time τ_n and time k , since every step changes the value of F_k by at most $\frac{1}{2n}$. Thus

$$
B_n\subseteq \{\tau_{n+1}-\tau_n\geq \varepsilon n\}
$$

We have

$$
\mathbb{P}\left[\tau_{n+1} - \tau_n \ge \varepsilon n\right] = \sum_{l \ge \lceil \frac{\varepsilon n}{2} \rceil} \mathbb{P}\left[\tau_{n+1} - \tau_n = 2l\right] \xrightarrow{\text{Lemma 5.4}} \sum_{l \ge \lceil \frac{\varepsilon n}{2} \rceil} 2^{-l}
$$
\n
$$
\le 2^{-\frac{\varepsilon n}{2}} \sum_{l \ge 0} 2^{-l} = 2^{1 - \frac{\varepsilon n}{2}}
$$
\n
$$
\implies \sum_{n \ge 1} \mathbb{P}\left[\tau_{n+1} - \tau_n \ge \varepsilon n\right] \le \sum_{n \ge 1} 2^{1 - \frac{\varepsilon n}{2}} = 2 \cdot \sum_{n \ge 1} \left(2^{-\frac{\varepsilon}{2}}\right)^n < \infty
$$

By Borel-Cantelli, it follows that only finitely many of the events $\{\tau_{n+1} - \tau_n \geq \varepsilon n\}$ can occur, and therefore also only finitely many of the events B_n . This concludes the proof. \Box

The same model with only one random walker results in a standard Pólya urn where two balls of the drawn color are added after every draw. For the Pólya urn, the fraction of balls of one of the colors is a martingale and converges to a limit which is Beta-distributed. We have shown now for the two-player urn that the fraction of the left edge weight also has a limit, even if it is a martingale only if looked at at certain stopping times. Simulations and intuition suggest that the limit should also have a density. However, we couldn't prove this so far, so we only give the following conjecture. Simulations also suggest that the limiting distribution is not a Beta distribution.

Conjecture 5.10 Define the random variable $Y := \lim_{n \to \infty} \frac{w(n,0)}{w(n,0) + w(n)}$ $\frac{w(n,0)}{w(n,0)+w(n,1)}$. Then $Y \in [0,1]$ has a density w.r.t. the Lebesgue measure on [0, 1]. *Y* is not Beta-distributed, and its distribution is different from the distribution of the limit in [Conjecture 5.3.](#page-30-2)

5.2 Model on Z

So far, we have looked at the LERRW with multiple walkers only on a very simple graph and only with 2 walkers. We now look at the edge-reinforced random walk on **Z** with an arbitrary finite number of random walkers. Formally, we have *k* sequences (for *k* walkers) $\left(X_n^{(m)}\right)$ $n \geq 0$, $1 \leq m \leq k$ of random variables with the following dynamics. The transition probabilities depend on the edge weights $w(n, j) > 0$ for $n \ge 0, j \in \mathbb{Z}$ where *j* corresponds to the edge from j to $j + 1$.

Figure 6: Edge weights on **Z** at time *n*

If \mathcal{G}_n denotes $\sigma\left(\left\{X^{(m_2)}_{m_1}: 0 \leq m_1 \leq n, 1 \leq m_2 \leq k\right\} \cup \left\{w\left(m, j\right): 0 \leq m \leq n, j \in \mathbb{Z}\right\}\right)$ i.e. the history of the random walkers and edge weights up to and including time *n*, then we define, conditioned on \mathcal{G}_n , the following transition probabilities:

- A random walker $m (1 \le m \le k)$ which is going to jump is chosen uniformly at random (independent of \mathcal{G}_n) amongst the *k* walkers.
- If the chosen random walker is at position *j* (i.e. $X_n^{(m)} = j$), he then jumps
	- **−** to the right (i.e. $X_{n+1}^{(m)} = j + 1$) with probability $\frac{w(n,j)}{w(n,j-1)+w(n,j)}$ **−** to the left (i.e. $X_{n+1}^{(m)} = j - 1$) with probability $\frac{w(n,j-1)}{w(n,j-1)+w(n,j)}$

i.e. the jump probabilities are proportional to the corresponding edge weights.

• If *j*^{*} is the traversed edge (*j*^{*} = *j* if the walker jumps to the right, *j*^{*} = *j* − 1 if he jumps to the left), then for $i \neq j^*$, $w(n,i) = w(n+1,i)$ and $w(n,j^*) \leq w(n+1,j^*)$, i.e. the weight of the traversed edge may be increased according to some reinforcement scheme.

We consider schemes where the increment $w(n+1,j^*) - w(n,j^*)$ may solely depend on *j* ∗ , and the number of times the edge was crossed up to time *n*. In other words, *w* (*n*, *j*) can still be written in terms of the weight function *W^e* (*k*) as defined in [Definition 2.3.](#page-10-0) Some generalizations are possible, such as making the weight also depend on *n* (i.e. at which times the edge was crossed), but will not be considered here.

- The initial edge weights can be chosen arbitrarily, but all of them must be positive.
- • The initial positions of the *k* walkers can be chosen arbitrarily.

5.3 Recurrence or Finite Range on Z

We say that one of the walkers m ($1 \le m \le k$) **Definition 5.11** • is **transient**, if he visits every integer only finitely often, that is, every integer appears only finitely often in the sequence $\left(X_n^{(m)}\right)$ *n*≥1 • is **recurrent**, if he visits every integer infinitely often, that is, every integer appears infinitely often in the sequence $\left(X_n^{(m)}\right)$ *n*≥1 • has **finite range**, if he only visits finitely many integers, that is, the number of distinct integers appearing in the sequence $\left(X_n^{(m)}\right)$ $\sum_{n\geq 1}$ is finite Assume the edge-reinforced random walk with *k* walkers starts with an initial configura- **Lemma 5.12** tion of the weights $w(0, j)$ such that all but finitely many of them are 1. Then, for every random walker m ($1 \leq m \leq k$) with $X_0^{(m)} \geq 0$, we have the following: $\mathbb{P}\left[X_n^{(m)}=0 \text{ for some } n\geq 0\right]$ $A + \mathbb{P}\left[X_n^{(m)} \neq 0 \text{ for all } n \geq 0 \text{ and } X^{(m)} \text{ only visits finitely many nodes}\right]$ which have not been visited before by any other walker $\big] = 1$

We follow the proof of [\[2,](#page-63-13) Lemma 3.0]. Consider a fixed random walker *m* with $X_0^{(m)} \geq 0$. *Proof* We now define:

$$
F(n, j) := \begin{cases} \sum_{i=0}^{j-1} \frac{1}{w(n,i)} & \text{if } j > 0 \\ 0 & \text{if } j \le 0 \end{cases} \qquad \tau^{(m)} := \inf \left\{ n \ge 0 : X_n^{(m)} \le 0 \right\}
$$

$$
M_n^{(m)} := H\left(n \wedge \tau^{(m)}, X_{n \wedge \tau^{(m)}}^{(m)}\right)
$$

$$
H_n^{(m)} := M_n^{(m)} + \sum_{i=1}^n \left(\frac{1}{w\left(i-1, X_{i-1}^{(m)}\right)} - \frac{1}{w\left(i, X_{i-1}^{(m)}\right)}\right) \cdot 1_{X_i^{(m)} > X_{i-1}^{(m)}, i \le \tau^{(m)}}
$$

$$
+ \sum_{i=1}^n \sum_{l=1, l \ne m}^k \sum_{j=0}^\infty \left(\frac{1}{w\left(i-1, j\right)} - \frac{1}{w\left(i, j\right)}\right) \cdot 1_{\left\{X_{i-1}^{(l)}, X_i^{(l)}\right\} = \left\{j, j+1\right\}, i \le \tau^{(m)}, j < X_i^{(m)}}
$$

= 1 for at most one pair of l,j

 $M_n^{(m)}$ is nonnegative by definition of *F*, and $H_n^{(m)}\,\geq\,M_n^{(m)}\,\geq\,0$ since edge weights can only increase and therefore, all terms in the sums in the definition of $H_n^{(m)}$ are nonnegative. $H_n^{(m)}$ is a martingale: let

$$
d_n^{(m)} :=
$$
\n
$$
H_n^{(m)} - H_{n-1}^{(m)} = \underbrace{M_n^{(m)} - M_{n-1}^{(m)}}_{:=e_n^{(m)}} + \underbrace{\left(\frac{1}{w(n-1, X_{n-1}^{(m)})} - \frac{1}{w(n, X_{n-1}^{(m)})}\right) \cdot \mathbb{1}_{X_n^{(m)} > X_{n-1}^{(m)}, n \leq \tau^{(m)}}}_{:=f_n^{(m)}} + \underbrace{\sum_{l=1, l \neq m}^k \sum_{j=0}^\infty \left(\frac{1}{w(n-1,j)} - \frac{1}{w(n,j)}\right) \cdot \mathbb{1}_{\left\{X_{n-1}^{(l)}, X_n^{(l)}\right\} = \{j, j+1\}, n \leq \tau^{(m)}, j < X_n^{(m)} \text{ is } e_n^{(m)}}}_{:=g_n^{(m)}}.
$$

then we have to show that $\mathbb{E}\left[d_n^{(m)}\ \middle|\ \mathcal{G}_{n-1}\right]=0$. We have:

- if $n-1 \ge \tau^{(m)}$, then $d_n^{(m)} = 0$. Hence, it suffices to consider the case $X_{n-1}^{(m)} = j > 0$ and $\tau^{(m)} \geq n$.
- with probability $\frac{1}{k}$, the walker *m* jumps at time *n* − 1. In this case, $g_n^{(m)} = 0$ since no *k* other walker can jump and the indicator variable in $g_n^{(m)}$ is therefore 0. If he jumps to the right (with probability $\frac{1}{k} \cdot \frac{w(n-1,j)}{w(n-1,j-1)+w(n)}$ $\frac{w(n-1,j)}{w(n-1,j-1)+w(n-1,j)}$), then $e_n^{(m)} = w(n,j)^{-1}$ and $f_n^{(m)} =$ $w(n-1,j)^{-1} - w(n,j)^{-1}$, hence $d_n^{(m)} = w(n-1,j)^{-1}$. If he jumps left (with probability $\frac{1}{k} \cdot \frac{w(n-1,j-1)}{w(n-1,j-1)+w(n)}$ $\frac{w(n-1,j-1)}{w(n-1,j-1)+w(n-1,j)}$, then $e_n^{(m)} = -w(n-1,j-1)^{-1}$ and $f_n^{(m)} = 0$, hence $d_n^{(m)} = 0$ $-w(n-1,j-1)^{-1}$.
- with probability $\frac{k-1}{k}$, the walker *m* does not jump. In this case, $f_n^{(m)} = 0$ since the indicator variable in $f_n^{(m)}$ is therefore 0. The value of $M_{n-1}^{(m)}$ *^{(<i>m*}</sup>) now changes (that is, $e_n^{(m)} \neq 0$) if one of the other *k* − 1 walkers crosses one of the edges between the nodes 0 and *j*. At the same time, $g_n^{(m)} \neq 0$ only in this exact case. Now assume the walker $l \neq m$ crosses the edge *i* with $0\leq i < j$. Then $e_n^{(m)}=\frac{1}{w(n,i)}-\frac{1}{w(n-1,i)}$ and $g_n^{(m)}=\frac{1}{w(n-1,i)}-\frac{1}{w(n,i)}$, hence $d_n^{(m)}=0$.
- conditioned on $X_{n-1}^{(m)} = j > 0$ and $\tau^{(m)} \ge n$ (both events measurable w.r.t. \mathcal{G}_{n-1}), we can therefore conclude

$$
\mathbb{E}\left[d_n^{(m)}\middle|\mathcal{G}_{n-1}\right] = \frac{1}{k} \cdot \frac{1}{w(n-1,j-1)+w(n-1,j)} \cdot \left(\frac{w(n-1,j)}{w(n-1,j)} - \frac{w(n-1,j-1)}{w(n-1,j-1)}\right)
$$

$$
= \frac{1}{k} \cdot \frac{1}{w(n-1,j-1)+w(n-1,j)} \cdot (1-1) = 0
$$

(By the same arguments, but only considering $e_n^{(m)}$, we can show that $M_n^{(m)}$ is a supermartingale.)

As a nonnegative martingale*,* $H_n^{(m)}$ converges almost surely.

Figure 7: Evolution of the supermartingale during 4 steps of the ERRW with multiple walkers In the figure above, we consider $k = 4$ walkers whose positions are indicated by the circled nodes. The value of $M_n^{(3)}$ is shown for the first 4 steps, and the corresponding walker $X^{(3)}$ is highlighted in blue.

We just showed this for all walkers m $(1 \le m \le k)$ with $X_0^{(m)} \ge 0$. Further observe that for such a walker *m*, we have, on the event $B_n^{(m)} = \left\{X_n^{(m)} > X_{n-1}^{(m)}\right\}$ *n*−1 , *n* ≤ *τ* (*m*) , *w n* − 1, *X* (*m*) $\binom{m}{n-1} = 1$, that $e_n^{(m)} = w\left(n, X_{n-1}^{(m)}\right)$ $\binom{m}{n-1}$ ⁻¹, $f_n^{(m)} = w \left(n-1, X_{n-1}^{(m)} \right)$ $\binom{m}{n-1}^{-1} - w(n, X_{n-1}^{(m)})$ $\binom{m}{n-1}^{-1}$, $g_n^{(m)} = 0$ and hence $d_n^{(m)} = 1$. Thus, by convergence, only a finite number of the events $B_n^{(m)}$ can occur for every such walker *m*.

Now define Γ to be the set of edges between two nonnegative integers to the right of the integer max $\left\{X_0^{(m)}\right\}$ $\binom{m}{0}: 1 \leq m \leq k \Big\}$ for which the initial weight was 1 (all but finitely many edges meet the latter criterion), and further define the event

 $D_n = \left\{\exists l: X_0^{(l)} \geq 0 \text{, an edge in } \Gamma \text{ is crossed between time } n-1 \text{ and } n \text{ for the first time by }\right.$ any walker, the crossing walker is *l* and $n \leq \tau^{(l)}$

Clearly, $D_n \subseteq B_n^{(m)}$ for some random walker *m* with $X_0^{(m)} \geq 0$, hence only a finite number of the events D_n can occur.

Now the proof cannot be continued along [\[2,](#page-63-13) Lemma 3.0] since the walkers starting to the left of 0 and the walkers which reach 0 can later cross edges to the right of 0 without triggering D_n and the other walkers can then follow them without triggering D_n . So, we only proved that walkers which never go to 0 and start to the right of 0 cannot visit infinitely many edges which have not been visited before by any other walker. \Box

Corollary 5.13 Assume the edge-reinforced random walk with *k* walkers starts with an initial configuration of the weights $w(0, i)$ such that all but finitely many of them are 1. Then, we have the following:

$$
\mathbb{P}\left[\exists m : X_n^{(m)} = 0 \text{ for some } n \ge 0\right]
$$

+
$$
\mathbb{P}\left[\forall m : X_n^{(m)} \ne 0 \text{ for all } n \ge 0 \text{ and } X^{(m)} \text{ has finite range}\right] = 1
$$

Proof It is sufficient to show that, conditional on $A^C = \{ \nexists m : X_n^{(m)} = 0 \text{ for some } n \geq 0 \}$, we have that the second event $B = \{ \forall m : X_n^{(m)} \neq 0 \text{ for all } n \geq 0 \text{ and } X^{(m)} \text{ has finite range} \}$ occurs a.s. Now, by [Lemma 5.12,](#page-38-1) every random walker *m* separately either reaches 0 or some other event *E* (*m*) occurs (by symmetry, [Lemma 5.12](#page-38-1) can also be applied to random walkers which start to the left of 0). But now, if A^C occurs, then no random walker reaches 0, hence the event $E^{(m)}$ occurs for every random walker *m* where

 $E^{(m)} = \left\{ X_n^{(m)} \neq 0 \text{ for all } n \geq 0 \text{ and } X^{(m)} \text{ only visits finitely many nodes} \right\}$

which have not been visited before by any other walker

So every walker visits only finitely many nodes not visited before by any of the other walkers. But this implies that all walkers together can only visit finitely many nodes. \Box

Corollary 5.14 Assume the edge-reinforced random walk with *k* walkers starts with an initial configuration of the weights $w(0, j)$ such that all but finitely many of them are 1. Then, we have the following:

> **P** $\forall j$ ∈ **Z** : *j* is visited ∞ often by at least one of the walkers $+$ **P** [all walkers have finite range] $= 1$

Proof It suffices to show that for every *j* $\in \mathbb{Z}$, we have

P $\vert j \rangle$ is visited ∞ often by at least one of the random walkers $+$ **P** $\left| i \right|$ is visited only finitely often and all random walkers have finite range = 1 (5.1)

To see this, assume that not all random walkers have finite range. Then, if we have proved [Equation 5.1,](#page-41-0) we know that for every $j \in \mathbb{Z}$, *j* is visited infinitely often by one of the random walkers. Thus, conditional on not all random walkers having finite range, all nodes are visited infinitely often, which is equivalent to [Corollary 5.14.](#page-41-1)

To show [Equation 5.1,](#page-41-0) it suffices in turn to show, for every *j* $\in \mathbb{Z}$ and $\forall n \geq 0$:

P [*j* is visited by at least one walker at a time $t \geq n$] $+$ **P** [no walker visits *j* at a time $t > n$ and all walkers have finite range] = 1 (5.2)

since the given events are decreasing and increasing respectively, and their limits correspond to the events in [Equation 5.1.](#page-41-0)

But now, consider the random walkers $\left(X_{n+i}^{(m)}\right)$ $\binom{m}{n+i}$. These form again an edge-reinforced $$ random walk with *k* walkers, and since until time *n*, only a finite number of edge weights can have changed, all but finitely many edges will still have weight 1 at time *n*. Thus, we can apply

[Corollary 5.13](#page-41-2) to the walkers $(X_{n+i}^{(m)})$ $\binom{m}{n+i}$ *i*≥0' which directly proves [Equation 5.2](#page-41-3) [\(Corollary 5.13](#page-41-2) only proves [Equation 5.2](#page-41-3) for $j = 0$ but of course we can relabel the nodes such that any other node gets the label 0, hence [Corollary 5.13](#page-41-2) is valid for any choice of node). \Box

Assume the edge-reinforced random walk with $k \ge 2$ walkers starts with an arbitrary ini-
Lemma 5.15 tial configuration of the weights $w(0, i)$. Assume further that $X^{(1)}$ and $X^{(2)}$ meet infinitely often. Then (almost surely):

- (i) if at least one of the walkers $X^{(1)}$ and $X^{(2)}$ does not have finite range, then both $X^{(1)}$ and $X^{(2)}$ do not have finite range.
- (ii) if some integer *z* is visited infinitely often by at least one of the walkers $X^{(1)}$ and $X^{(2)}$, then both $X^{(1)}$ and $X^{(2)}$ visit *z* infinitely often.
- (iii) if every integer is visited infinitely often by at least one of the walkers $X^{(1)}$ and $X^{(2)}$, then both $X^{(1)}$ and $X^{(2)}$ are recurrent.

The proof idea is the following: whenever $X^{(1)}$ and $X^{(2)}$ meet, we can randomly exchange *Proof* their labels, i.e. we can randomly decide whether we want to rename *X*(1) to *X*(2) and vice versa, and the law of the edge-reinforced random walk with the two walkers is invariant under such relabelings because the only distinguishing feature of a random walker is his position. But now, to construct counterexamples to the two statements in [Lemma 5.15,](#page-42-0) we would have to choose a fixed labeling for infinitely many times at which the walkers meet. But if we randomize the labeling with a sequence of independent Bernoulli random variables, then the probability of choosing a certain fixed labeling at infinitely many points in the sequence is 0, and since the law was invariant under random relabeling, it follows that the probability of any such counterexample is 0. We continue with the formal proof.

Set $\tau_1 := \inf \left\{ n \geq 0 : X_n^{(1)} = X_n^{(2)} \right\}$ and $\tau_{i+1} := \inf \left\{ n > \tau_i : X_n^{(1)} = X_n^{(2)} \right\}$. If $X^{(1)}, X^{(2)}$ meet infinitely often, then $\forall n : \tau_n < \infty$, but the construction also works if this is not the case. Let $(\omega_i)_{i\geq 1}$ be a sequence of iid random variables with $\mathbb{P}[\omega_i = 1] = \frac{1}{2} = \mathbb{P}[\omega_i = 0]$ (the ω_i are also independent of \mathcal{G}_n for all *n*, i.e. independent of the edge-reinforced random walk). Define $\widetilde{X}_n^{(1)}$ and $\widetilde{X}_n^{(2)}$ as follows (with $\omega_0 = 0$ and $\tau_0 = -1$):

$$
\widetilde{X}_n^{(1)} := X_n^{(1,\omega)} = \sum_{i \ge 0} \left((1 - \omega_i) X_n^{(1)} + \omega_i X_n^{(2)} \right) \cdot 1_{\tau_i < n \le \tau_{i+1}}
$$
\n
$$
\widetilde{X}_n^{(2)} := X_n^{(2,\omega)} = \sum_{i \ge 0} \left((1 - \omega_i) X_n^{(2)} + \omega_i X_n^{(1)} \right) \cdot 1_{\tau_i < n \le \tau_{i+1}}
$$

Note that the sums collapse to a single term. $\omega_i = 1$ means that we switch the labels of $X^{(1)}$ and $X^{(2)}$ during the time interval $(\tau_i, \tau_{i+1}]$. If we consider $\left(X^{(1)}, X^{(2)}\right)$ and $\left(\widetilde{X}^{(1)}, \widetilde{X}^{(2)}\right)$ as sequences of pairs of integers, then we have

$$
\left(\widetilde{X}_n^{(i)}\right)_{1\leq i\leq 2,n\geq 0} \stackrel{\text{d}}{=} \left(X_n^{(i)}\right)_{1\leq i\leq 2,n\geq 0} \tag{5.3}
$$

The equality in distribution follows from the mentioned invariance of the law of the random walk under relabelings at meeting times which is quite intuitive, and could be proved formally by looking at cylinder events, for example.

Figure 8: Label exchange lemma

The figure above illustrates how the "label exchange" of the two walkers works. Two sample paths for the two walkers are drawn, together with meeting points and the result of the label exchange.

We now show that any counterexamples to statements [\(i\)](#page-42-1) or [\(iii\)](#page-42-2) have probability 0:

(i) let *A* be the event that one of the walkers $X^{(1)}$, $X^{(2)}$ has finite range while the other one has infinite range, and that they meet infinitely often. It suffices to show that $\mathbb{P}[A] =$ 0. Denote by **P** the probability measure induced by the edge reinforced random walk alone and by $\bf Q$ the probability measure induced by the sequence $(\omega_i)_{i\geq 1}$ alone. Then, by [Equation 5.3,](#page-42-3) we have

> $\mathbb{P}\left[A\right] = \int \int \mathbb{1}_B \, \mathrm{d} \mathbf{Q} \, \mathrm{d} \mathbf{P}$ where $B := \left\{ \text{one of } \widetilde{X}^{(1)}, \widetilde{X}^{(2)} \text{ has finite range while} \right.$

> > the other has infinite range, they meet infinitely often $\{$

We have to show that the inner integral is 0 almost surely with respect to **P**. Consider fixed walker sequences $X^{(1)}$ and $X^{(2)}$. If one of $\widetilde{X}^{(1)}$, $\widetilde{X}^{(2)}$ should have finite range while the other has infinite range, then, by definition of $\widetilde{X}^{(1)}$ and $\widetilde{X}^{(2)}$, at least one of $X^{(1)}$, $X^{(2)}$ must have infinite range. Of course, by definition, we also have that $\widetilde{X}^{(1)}$, $\widetilde{X}^{(2)}$ meet infinitely often if, and only if, $X^{(1)}$, $X^{(2)}$ meet infinitely often. Hence, the indicator variable in the integral above can only be 1 in the case where one of the walkers $X^{(1)}$, $X^{(2)}$ has infinite range and the two walkers meet infinitely often, so we only need to show that in this particular case, the inner integral is still 0 almost surely.

Assume *X*(1) does not have finite range (w.l.o.g.). Then, for every *n*, one can find *i* such that between times τ_i and τ_{i+1} (all τ_i are finite if the two walkers meet infinitely often), *X*⁽¹⁾) visits a node at distance at least *n* from the integer 0. Call these times τ_{i_n} with i_n strictly increasing in *n* (w.l.o.g.).

Now consider the walkers $\widetilde{X}^{(1)}$, $\widetilde{X}^{(2)}$. One of them can have finite range only if the following holds. The same argument works for both walkers, we do it here for $\tilde{X}^{(1)}$ w.l.o.g. $\widetilde{X}^{(1)}$ can only have finite range if there exists *N* such that for all $n \ge N$ we have $\omega_{i_n} = 1$. Assume to the contrary that no such *N* exists. Then we can find arbitrarily large *n* such that $\omega_{i_n} = 0$ which means that the labels of $X^{(1)}$ and $X^{(2)}$ are *not* exchanged in the interval $(\tau_{i_n}, \tau_{i_n+1}]$. Since $X^{(1)}$ visits a node at distance at least *n* from 0 in this time interval, the same holds then for $\widetilde{X}^{(1)}$, so $\widetilde{X}^{(1)}$ would not have finite range.

But the probability that the sequence ω_{i_n} is 1 for all $n \geq N$ is 0 for any N (since the choice of i_n only depends on the edge-reinforced random walk, i.e. is independent of the ω_i , and since the probability of *ω* being constantly 1 on any fixed infinite subset of the integers is 0 by the choice of ω). Hence, the probability that such *N* exists is 0, and therefore the probability that $\widetilde{X}^{(1)}$ has finite range is 0 as well, and the same arguments give that the probability for $\widetilde{X}^{(2)}$ having finite range is 0 as well (both with respect to the measure **Q**).

So the indicator variable in the integral above is 0 almost surely w.r.t. **Q**, and hence the inner integral is always 0, which implies that the outer integral is also 0 and hence $\mathbb{P}[A] =$ Ω

(ii) similar. Let *A* now be the event that the integer *z* is visited infinitely often by at least one of the walkers $X^{(1)}$, $X^{(2)}$, that they meet infinitely often, and that one of them does *not* visit *z* infinitely often. Then, we have again:

$$
\mathbb{P}[A] = \int \int \mathbb{1}_B \, d\mathbf{Q} \, d\mathbf{P}
$$

where $B := \{z \text{ visited } \infty \text{ often by at least one of } \widetilde{X}^{(1)}, \widetilde{X}^{(2)},$
they meet ∞ often, one of them visits z only finitely often

We see that the indicator variable can be 1 only if at least one of $X^{(1)}$, $X^{(2)}$ visits z infinitely often, and w.l.o.g. assume that this holds for $X^{(1)}$. As before, we can construct a stricly increasing sequence i_n such that in the time interval $(\tau_{i_n}, \tau_{i_n+1}]$, $X^{(1)}$ visits *z*. Again as before, one of $\widetilde{X}^{(1)}$, $\widetilde{X}^{(2)}$, take $\widetilde{X}^{(1)}$ w.l.o.g., can visit *z* only finitely often only if $\omega_{i_n} = 1$ for all $n > N$ for some *N*, an event which has again probability 0 w.r.t. **Q**.

(iii) apply [\(ii\)](#page-42-4) to every integer.

 \Box

Consider the edge-reinforced random walk with *k* walkers. Define, for $1 \le m \le k$: **Definition 5.16** • $\overline{X}^{(m)} := \limsup_{n \to \infty} X_n^{(m)}$ • $\underline{X}^{(m)} := \liminf_{n \to \infty} X_n^{(m)}$ Assume the edge-reinforced random walk with *k* walkers starts with an initial configura- **Theorem 5.17** tion of the weights $w(0, j)$ such that all but finitely many of them are 1. Then, we have the following: $\mathbb{P}\left[\forall m : X^{(m)} \text{ is recurrent} \right] + \mathbb{P}\left[\forall m : X^{(m)} \text{ has finite range} \right] = 1$

We present two variants of the proof. The first is a little less formal than the second, but *Proof* hopefully easier to understand. The second variant is as formal as possible without becoming totally incomprehensible.

Proof Variant 1 (less formal):

We have to show the following: if at least one of the walkers does not have finite range, then, almost surely, all of them are recurrent. Showing recurrence of all walkers is equivalent to showing that $\overline{X}^{(m)}=\infty$, $\underline{X}^{(m)}=-\infty$ for all walkers $m.$ Now, if at least one walker does not have finite range, then we know by [Corollary 5.14](#page-41-1) that every integer is visited infinitely often by at least one of the walkers. Hence, there must be walkers m_1 and m_2 with $\overline{X}^{(m_1)}=\infty$, $\underline{X}^{(m_2)}=-\infty$. For a contradiction, assume there is some walker which is not recurrent. w.l.o.g. we assume that there is some walker m_3 with $\underline{X}^{(m_3)} > -\infty$ (the proof is the same if the other condition on the lim sup is not met). Since we have $\underline{X}^{(m_2)} = -\infty$, we can partition the set of walkers into two non-empty sets:

$$
P_1 := \left\{ m : 1 \le m \le k \text{ and } \underline{X}^{(m)} = -\infty \right\}
$$

$$
P_2 := \left\{ m : 1 \le m \le k \text{ and } \underline{X}^{(m)} > -\infty \right\}
$$

 $\left\{\text{Choose } m_4 \in \argmax_{m \in P_1} \overline{X}^{(m)} \text{ and } m_5 \in \argmin_{m \in P_2} \underline{X}^{(m)}. \text{ Let } y := \min\left\{1, \underline{X}^{(m_5)}\right\} - 1 \in \mathbb{Z}.$ By choice of *m*5, all walkers in *P*² visit *y* only finitely often. Therefore, by [Lemma 5.12,](#page-38-1) each of them can only visit finitely many nodes not visited before by any other walker. We know that all nodes are visited infinitely often by at least one of the walkers, so we must have $\overline{X}^{(m_4)}=\infty$. If this was not the case, the walkers in P_1 would only visit nodes to the left of some fixed integer, and since the walkers in *P*² together only visit finitely many new nodes, this would imply that there is some largest visited integer, a contradiction to the fact that every integer is visited infinitely often (since at least one walker does not have finite range).

Now consider $P_3 := \left\{m \in P_1 : \overline{X}^{(m)} = \infty \right\} \, \neq \, \varnothing$ (since $m_4 \, \in \, P_3$) and the walker m_5 . If m_5 has finite range, then it is clear that m_5 will meet any walker in P_3 infinitely often, but this is a contradiction to [Lemma 5.15](#page-42-0) [\(i\).](#page-42-1) So $m₅$ must have infinite range. This is only possible if $\overline{X}^{(m_{5})} = \infty$. But since the walkers in P_2 , including m_5 , only visit finitely many nodes not visited before by any other walker, at least one walker m_6 in P_3 must be to the right of m_5 infinitely often in order to "free the path" for m_5 . As the walkers in P_3 are all recurrent, and as m_5 only visits nodes to the right of *y*, this implies that m_5 meets this walker m_6 infinitely often, which is a contradiction to [Lemma 5.15](#page-42-0) [\(iii\).](#page-42-2) Hence, we almost surely arrive at a contradiction.

To summarize: assuming that at least one walker does not have finite range and that least one walker is not recurrent at the same time leads (almost surely) to a contradiction. Therefore, the assumption that this can happen must be wrong (almost surely). This implies that if at least one walker does not have finite range, then (almost surely) all walkers must be recurrent.

Figure 9: Illustration: proof that all walkers recurrent or all have finite range

Above, the two possible behaviors of the walker $m₅$ (which are both, in fact, almost surely impossible) are shown: either m_5 has finite range (indicated by the light green background), but then it would meet m_4 infinitely often, or m_5 does not have finite range, but then m_6 would have to "free the path" for m_5 (indicated by the light orange background) and m_5 would meet $m₆$ infinitely often. The relevant meeting points are circled.

Proof Variant 2 (more formal):

Our goal is to show the following:

$$
\mathbb{P}\left[\underbrace{\exists m_0, m_3 : X^{(m_0)} \text{ does not have finite range and } X^{(m_3)} \text{ is not recurrent}}_{=:A}\right] = 0
$$

The indices of *m* are chosen in such a way that they agree with proof variant 1, and they therefore do not appear in any logical order in this proof variant. The proof proceeds by showing that the above event *A* is subset of a null set. We first apply [Corollary 5.14.](#page-41-1) Since the intersection of *A* and the event {all walkers have finite range} is empty, we can conclude that

$$
A \subseteq (A \cap \{ \text{every integer is visited } \infty \text{ often by at least one of the walksers} \}) \cup N
$$

where *N* is a null set

Hence, it suffices to show that $P[B] = 0$. *B* can only occur if there are walkers m_1 and m_2 with $\overline{X}^{(m_1)} = \infty$, $\underline{X}^{(m_2)} = -\infty$ since every integer is visited infinitely often (when *B* occurs). Furthermore, for the non-recurrent walker m_3 , we necessarily have $\overline{X}^{(m_3)}\neq\infty$ or $\underline{X}^{(m_3)}\neq-\infty$ (when *B* occurs). Hence,

$$
B \subseteq \underbrace{\{ \exists m_1, m_2, m_3 : \overline{X}^{(m_3)} < \infty, \overline{X}^{(m_1)} = \infty, \underline{X}^{(m_2)} = -\infty \}}_{=:C}
$$
\n
$$
\cup \underbrace{\{ \exists m_1, m_2, m_3 : \underline{X}^{(m_3)} > -\infty, \overline{X}^{(m_1)} = \infty, \underline{X}^{(m_2)} = -\infty \}}_{=:D}
$$

We show w.l.o.g. that $P[D] = 0$ since the proof for $P[C] = 0$ is the same. When *D* occurs, the \mathcal{L} two sets $P_1 := \left\{m: 1 \leq m \leq k \text{ and } \underline{X}^{(m)}=-\infty\right\}$ and $P_2 := \left\{m: 1 \leq m \leq k \text{ and } \underline{X}^{(m)} > -\infty\right\}$ are both non-empty (these are both random set). Choosing $m_5 \in \arg \min_{m \in P_2} \underline{X}^{(m)}$ as well as $y := \min\left\{1, \underline{X}^{(m_5)}\right\} - 1 \in \mathbb{Z}$, we see that there is a random integer y which is only visited finitely often by the walkers in the set P_2 . Hence,

$$
D \subseteq \bigcup_{y \in \mathbb{Z}} \bigcup_{n \in \mathbb{N}} \overline{\left\{P_1 \neq \emptyset, P_2 \neq \emptyset, y \text{ not visited by walkers in } P_2 \text{ after time } n, \right\}}
$$

$$
\exists m_1, m_2 : \overline{X}^{(m_1)} = \infty, \underline{X}^{(m_2)} = -\infty \left\}
$$

It thus suffices to show $\mathbb{P}\left[E_{y,n}\right] = 0$. If we set $[k] := \{1,\ldots,k\}$ and $\mathfrak{P} := \mathcal{P}\left([k]\right) \setminus \{\varnothing,[k]\}$ where P denotes the power set, then we can further write

=:*Fy*,*n*,*^P*

$$
E_{y,n} \subseteq \bigcup_{P \in \mathfrak{P}} \overbrace{\left\{P_1 = P = [k] \setminus P_2, y \text{ not visited by walksers in } P_2 \text{ after time } n, \right\}}^{\text{max}} \exists m_1, m_2 : \overline{X}^{(m_1)} = \infty, \underline{X}^{(m_2)} = -\infty \right\}
$$

and it suffices to show $\mathbb{P}\left[F_{y,n,P}\right] = 0$ for all $y \in \mathbb{Z}$, $n \in \mathbb{N}$, $P \in \mathfrak{P}$. We are now ready to apply [Lemma 5.12](#page-38-1) for every walker in the fixed set $[k] \setminus P = P_2$ (equality if $F_{\nu,n,P}$ occurs). Since the intersection of $F_{v,n,P}$ and the event that any walker in $[k] \setminus P = P_2$ returns to *y* at a time larger than *n* is empty, we can conclude by [Lemma 5.12](#page-38-1) (as in the proof of [Corollary 5.13,](#page-41-2) only with the random walk started at time $n + 1$, and with a subset of the walkers) that

 $F_{y,n,P} \subseteq$ =:*Gy*,*n*,*^P* ${P_1 = P, P_2 = [k] \setminus P, y \text{ not visited by walksers in } P_2 \text{ after time } n \text{ and the walksers}}$

in *P*₂ only visit finitely many nodes not visited before by any other walker,

$$
\exists m_1, m_2 : \overline{X}^{(m_1)} = \infty, \underline{X}^{(m_2)} = -\infty
$$

∪ \widehat{N} where \widehat{N} is a null set

and it suffices to show $\mathbb{P}\left[G_{y,n,P}\right]=0.$ Choose $m_4\in\argmax_{m\in P_1}\overline{X}^{(m)}.$ If $G_{y,n,P}$ occurs, then we necessarily have $\overline{X}^{(m_4)}=\infty.$ If this was not the case, the walkers in P_1 would only visit nodes to the left of some fixed integer, and since the walkers in P_2 together only visit finitely many new nodes (when *Gy*,*n*,*^P* occurs), this would imply that there is some largest visited integer, a contradiction to the fact that $\overline{X}^{(m_1)} = \infty$ when $G_{y,n,P}$ occurs.

Now consider the random set of recurrent walkers $P_3:=\left\{m\in P_1:\overline{X}^{(m)}=\infty\right\}\neq\varnothing$ (since *m*₄ ∈ *P*₃ when *G*_{*y*,*n*,*P* occurs) and the walker m_5 ∈ arg min_{*m*∈*P*₂} $\frac{X^{(m)}}{2}$. If m_5 has finite range,} then it is clear that m_5 will meet any walker in P_3 infinitely often. On the other hand, if m_5 has infinite range, then $\overline{X}^{(m_5)} = \infty$ since $m_5 \in P_2$. But since the walkers in P_2 , including m_5 , only visit finitely many nodes not visited before by any other walker when *Gy*,*n*,*^P* occurs, at least one walker m_6 in P_3 must be to the right of m_5 infinitely often in order to "free the path" for m_5 . As the walkers in P_3 are all recurrent, and as m_5 only visits nodes to the right of γ , this implies that m_5 meets this walker m_6 infinitely often (when $G_{\nu,n,P}$ occurs). Therefore, regardless of whether m_5 has finite range or not, there is always a recurrent walker $m_6 \in P_3$ which meets m_5 infinitely often when *Gy*,*n*,*^P* occurs.

We now apply [Lemma 5.15](#page-42-0) [\(iii\)](#page-42-2) to see that $G_{V,n,P}$ is a null set (which concludes the proof): as done repeatedly in this proof, we find that $G_{V,n,P}$ is a subset of a countable union of events of the form "some fixed walker is not recurrent (corresponding to *m*5) and meets some other fixed walker which is recurrent (corresponding to m_6) infinitely often". However, any event of this form is a null set by [Lemma 5.15](#page-42-0) [\(iii\).](#page-42-2) \Box

6 Biased Reinforced Random Walk

Here, we consider biased random walks with reinforcement. We could prove some results, but there are also still open questions. This section therefore presents proven results next to open questions and conjectures, and sometimes also just possible ways to tackle a question, which would have to be pursued further. In order to limit the length of this thesis, this section will often only provide proof sketches or skip some proof altogether.

6.1 *λ* [∗]**-Biased Edge-Reinforced Random Walk**

The *λ* ∗ -biased edge-reinforced random walk uses the same linearly reinforced edge weights as the LERRW, but introduces an additional bias in a certain direction. Here, we consider a biased reinforced random walk on **Z**, so the bias can be either to the left or to the right. Formally, define the *λ* ∗ -biased edge-reinforced random walk on **Z** as the sequence *Xⁿ* of random variables with

$$
X_0 = 0
$$

\n
$$
\mathbb{P}[X_{n+1} = x_n + 1 | X_n = x_n, ..., X_0 = x_0] = \frac{\lambda \cdot w(n, x_n)}{w(n, x_n - 1) + \lambda \cdot w(n, x_n)}
$$

\n
$$
w(n, z) = \underbrace{w(0, z)}_{=1 \text{ here}} + \sum_{i=0}^{n-1} 1 \mathbb{1}_{X_i = z, X_{i+1} = z+1} + 1 \mathbb{1}_{X_i = z+1, X_{i+1} = z}
$$

Figure 10: Edge weights and transition probabilities for the biased walk on **Z** at time *n*

In terms of [Definition 2.3,](#page-10-0) we take $W(n) = 1 + n$, so the edge weights are being linearly reinforced, but then we introduce an additional bias by multiplying the right edge weight with the parameter *λ*.

Now, consider $\lambda \in \mathbb{Q}_{>0}$ (the result can later be extended to all $\lambda \in \mathbb{R}_{>0}$) and a single node, say 0, the root. Initially, the weights on both adjacent edges are 1. Whenever the random walk leaves 0 along one edge, it can only return to 0 by the same edge. The weight of this edge will then have increased by 2. If $\lambda = 1$, then this process is equivalent to drawing balls from an urn which initially contains 1 black and 1 white ball and where a ball is replaced with 3 balls of the same color as the ball which was drawn, i.e. the urn will contain two more balls of the respective color than before (white balls corresponding to the left edge, black balls to the right edge). In general, the process is biased, and the number of balls which are put into the urn has to be adapted:

Let $\lambda = \frac{p}{q}$. Consider a node $z \in \mathbb{Z}$ with initial edge weights $w(0, z - 1) = l_0$ and $w(0, z) = \mathbb{Z}$ Lemma 6.1 r_0 . If the λ^* -biased edge-reinforced random walk is started in *z*, then the sequence of left turns and right turns of the random walk at *z* has the same distribution as the sequence of white and black balls drawn from the following urn:

- Initially, the urn contains $a_0 = q \cdot l_0$ white and $b_0 = p \cdot r_0$ black balls
- If a white ball is drawn, it is replaced with $2q + 1$ white balls (i.e. 2q more white balls than before)
- If a black ball is drawn, it is replaced with $2p + 1$ black balls (i.e. $2p$ more black balls than before)

The urn and the edge weights can be coupled such that, writing *τⁿ* for the time at which *z* is visited for the *n*-th time, $w(\tau_n, z - 1) = \frac{a_n}{q}$ and $w(\tau_n, z) = \frac{b_n}{p}$.

Note that, after the *n*-th draw and the *n*-th visit to the node *z*, we have with the coupling *Proof Sketch* given in [Lemma 6.1:](#page-50-1)

$$
\mathbb{P}\left[X_{\tau_n+1} = z+1 \mid X_{\tau_n} = z, \dots, X_0 = x_0\right] = \frac{\lambda \cdot w(\tau_n, z)}{w(\tau_n, z-1) + \lambda \cdot w(\tau_n, z)} = \frac{\frac{p}{q} \cdot \frac{b_n}{p}}{\frac{a_n}{q} + \frac{p}{q} \cdot \frac{b_n}{p}}
$$

$$
= \frac{b_n}{q} \cdot \frac{q}{a_n + b_n} = \frac{b_n}{a_n + b_n}
$$

Consider the λ^* -biased edge-reinforced random walk at node $z \in \mathbb{Z}$ with $\lambda \in \mathbb{Q}_{>0}$. If *z* is Lemma 6.2 visited infinitely often, then

- (i) If $\lambda > 1$, then $\frac{\lambda w(n,z)}{w(n,z-1)+\lambda w(n,z)}$ converges almost surely to 1
- (ii) If $\lambda < 1$, then $\frac{w(n,z-1)}{w(n,z-1)+\lambda w(n,z)}$ converges almost surely to 1

Use [Lemma 6.1](#page-50-1) and [Theorem 3.5](#page-16-0) with $\mu({p}) = 1$ and $\nu({q}) = 1$ (or vice versa if $p < q$). \Box Proof Sketch

The *λ*^{*}-biased edge-reinforced random walk is said to be **γ 12 • 12 • 12 • 12 • 13 • 13 • 13 • 13 • 13 • 13 • 13 • 13 • 13 • 13 • 13 • 13 • 13 •** 13

- **transient** if it a.s. visits every node only finitely often
- **recurrent** otherwise, i.e. if there is at least one node which is visited infinitely often

If the λ^* -biased edge-reinforced random walk is recurrent, then all nodes are visited in-
 finitely often almost surely.

If the random walk is recurrent, then there is at least one node which is visited infinitely *Proof* often, say *z*. To show that all nodes are visited infinitely often, it suffices to show that *z* being visited infinitely often implies that both neighbors of *z* are visited infinitely often (one can then continue by induction). Assume for a contradiction that one of the neighbors, say *y*, is visited only finitely often, and let *t* be the time of the last visit to γ (at time $t + 1$, the random walk then

necessarily is at *z*). We can assume w.l.o.g. $y = z + 1$ (in the other case, reflect \mathbb{Z} at *z*, set a new $\lambda' = \frac{1}{\lambda}$ and continue as below).

Let $\tau_1 = t + 1$, τ_2 , τ_3 , ... be the times at which *z* is visited after the last visit to *y*. Then, at each time *τn*, we have

$$
\mathbb{P}\left[X_{\tau_n+1} = z+1 = y \mid X_{\tau_n} = z, \dots, X_0 = x_0\right] = \frac{\lambda \cdot w\left(\tau_n, z\right)}{w\left(\tau_n, z-1\right) + \lambda \cdot w\left(\tau_n, z\right)}
$$
\n
$$
= \frac{\lambda \cdot w\left(t+1, z\right)}{\lambda \cdot w\left(t+1, z\right) + \underbrace{w\left(t+1, z-1\right)}_{w_l} + n - 1}
$$
\n
$$
\implies \mathbb{P}\left[\forall n : X_{\tau_n+1} \neq y\right] = \prod_{n \geq 1} \frac{w_l + n - 1}{\lambda w_r + w_l + n - 1} \stackrel{\textcircled{\tiny{\#}}}{\leq} \exp\left(\sum_{n \geq 1} \left(\frac{w_l + n - 1}{\lambda w_r + w_l + n - 1} - 1\right)\right)
$$
\n
$$
= \exp\left(-\sum_{n \geq 1} \frac{\lambda w_r}{\lambda w_r + w_l + n - 1}\right) = 0
$$
\nwhere

\n
$$
\text{b holds since } \forall x \in \mathbb{R} : x \leq \exp\left(x - 1\right)
$$

Hence, the event *y* being visited only finitely often has probability 0.

 \Box

Based on [Lemma 6.2,](#page-50-0) it seems reasonable to conjecture that the λ^* -biased edge-reinforced random walk is transient whenever $\lambda \neq 1$, since at every node, the probability to go in the direction of bias would converge to 1 if the node was visited infinitely often. This intuitively seems to be a contradiction: if the probability to go in one direction goes to 1, then every node should be visited only finitely often. However, we did not manage to prove the following conjecture.

Conjecture 6.5 ^{*}-biased edge-reinforced random walk is transient for $\lambda \neq 1$.

Note that for $\lambda = 1$, the walk is recurrent since this case corresponds to the standard LERRW on **Z**.

6.1.1 Some Simulations

Below, the result of some simulations is presented. The λ^* -biased edge-reinforced random walk was simulated in 100 simulations for 1 000 000 steps each, in 1 000 simulations for 100 000 steps each and in 10 000 simulations for 10 000 steps each.

As can be seen in [Figure 11,](#page-52-0) the simulated speed strongly depends on the number of steps for which the random walk was simulated. In this case, the speed decreased with an increasing number of simulated steps. This could either be an indication that the speed has not yet converged and longer simulations would be necessary or that the random walk is actually always recurrent and hence the speed is actually zero (however, it seems intuitively very likely that it will be transient for large values of λ). For $\lambda = 1.1$, the speed was almost indistinguishable from 0. To see if the walk is really recurrent, it is however also of interest when the root was visited for the last time, since a speed of 0 does not necessarily imply recurrence.

Figure 11: Average speed of the random walk

Figure 12: Average last visit to the root by the random walk

Longer or more simulations are necessary to get reliable results (note for example that in [Figure 12,](#page-52-1) the green curve suddenly goes up for $\lambda = 1.6$ which is likely not a property of the model, but a case of variation in the simulation). However, it is already visible that for $\lambda = 1.1$, the last visit to the root occurred on average relatively late, i.e. at a noticeable fraction of the total number of simulated steps. Indeed, the last visit to the root seemed to occur later (compared to the total number of steps) when simulating the walk for longer, which we would not have expected. Of course, this could also just be a case of variation in the simulation.

The simulations do unfortunately not present any evidence which would support [Conjec](#page-51-1)[ture 6.5,](#page-51-1) but to argue about properties like recurrence and transience, which only make sense on an infinite time horizon, is very difficult with the help of only finite simulations.

6.1.2 Stochastic Approximation

To better understand the *λ* ∗ -biased edge-reinforced random walk on **Z**, it might be helpful to approximate it by reinforced walks on finite circular graphs. We superficially present a basic idea how this could be done. However, it is completely unclear if the analysis of the walk on a finite circular graph will provide new insights for the walk on **Z**. Since the easiest circular graph is the triangle, we consider the *λ* ∗ -biased edge-reinforced random walk on the triangle, defined as follows:

$$
r(x) = (x \mod 3) + 1 \text{ the right neighbor of } x \in \{1, 2, 3\}
$$

\n
$$
l(x) = ((x + 1) \mod 3) + 1 \text{ the left neighbor of } x
$$

\n
$$
X_0 = 1
$$

\n
$$
\mathbb{P}[X_{n+1} = r(x_n) | X_n = x_n, ..., X_0 = x_0] = \frac{\lambda \cdot w(n, x_n)}{\lambda \cdot w(n, x_n) + w(n, l(x_n))}
$$

\n
$$
w(n, i) = \underbrace{w(0, i)}_{=1 \text{ here}} + \sum_{k=0}^{n-1} 1_{X_k = i, X_{k+1} = r(i)} + 1_{X_k = r(i), X_{k+1} = i}
$$

\n
$$
c_{n,i} = \frac{w(n, i)}{\sum_{j=1}^{3} w(n, j)}
$$

For simplicity, we will assume $\lambda > 1$.

We call c_n the vector of normalized edge weights at time n , which is located on the unit simplex and has entries $c_{n,i}$ where $i = 1, 2, 3$. If we assume that the normalized edge weights were fixed to some vector *c* (on the unit simplex), then we get a Markov chain with the following stationary distribution π_c (only the value at the node 1 is shown, the others follow from the symmetry of the model):

$$
\pi_c(1) = \left(\lambda^3 c_1 c_2 c_3 + \lambda^2 \left(c_1^2 c_3 + c_2 c_3^2\right) + \lambda \left(c_1^2 c_2 + c_1 c_3^2\right) + c_1 c_2 c_3\right) Z^{-1}
$$

where *Z* is the appropriate normalizing constant.

Figure 13: Transition probabilities of the *λ* ∗ -biased ERRW on the triangle at time *n* (two arrows are missing for clarity)

We further define the stationary distribution on the edges π_c^{edge} simply by looking at the time spent on each edge when the chain is run starting from the stationary distribution *π*. For example,

$$
\pi_c^{\text{edge}}(1) = \frac{\lambda c_1}{\lambda c_1 + c_3} \cdot \pi_c(1) + \frac{c_1}{\lambda c_2 + c_1} \cdot \pi_c(2)
$$

We now approximate the evolution of the time-dependent vector \mathbf{c}_n of the edge weights:

$$
1 \ll k \ll n \implies (n+k) \mathbf{c}_{n+k} \approx n\mathbf{c}_n + k\pi_{\mathbf{c}_n}^{\text{edge}}
$$

\n
$$
\iff \mathbf{c}_{n+k} - \mathbf{c}_n \approx \frac{k}{n+k} \left(\pi_{\mathbf{c}_n}^{\text{edge}} - \mathbf{c}_n \right)
$$

\ntherefore approximate with
$$
\frac{d}{dt} \tilde{\mathbf{c}}(t) = \frac{1}{t} \left(\pi_{\tilde{\mathbf{c}}(t)}^{\text{edge}} - \tilde{\mathbf{c}}(t) \right)
$$

\nexponential time change to get
$$
\frac{d\mathbf{c}}{dt} = \pi_{\mathbf{c}}^{\text{edge}} - \mathbf{c} \quad \text{with } \tilde{\mathbf{c}}(t) = \mathbf{c} \left(\ln(t) \right)
$$

This is called a stochastic approximation: we approximate the random evolution of the vector of normalized edge weights by a differential equation. This is possible because the changes in the vector of normalized edge weights get ever smaller as time increases, and the randomness gets less noticeable by virtue of the law of large numbers. To argue formally, many additional steps would be necessary, but this approximation can already give a good intuition on what is happening.

Without any sort of formal proof by simply looking at [Figure 14,](#page-55-1) we conclude that the normalized edge weights will eventually converge to the uniform distribution. This probably also holds for circles with more than three nodes, showing that the long-run behavior of the biased reinforced random walk on circles is fundamentally different from the long-run behavior of the walk on the integers (where at every vertex the quotient of the right edge weight divided by the left edge weight converges to 1).

Figure 14: The vector field of the differential equation on the unit simplex

In blue, three segments of solutions to the differential equation are shown. In red, orange and green, three actual trajectories obtained by simulating the *λ* ∗ -biased edge-reinforced random walk are shown. The red trajectory results from setting the initial weights to 1 and running the simulation for 10 000 steps. For the orange trajectory, the initial weights were 100, 450, 450 and the simulation was run for 100 000 steps; for the green one, they were 500, 2250, 2250 and the simulation ran for 500 000 steps.

This already gives an indication that approximating the wlak on $\mathbb Z$ with that on a finite circular graph could be very hard. Indeed, [Lemma 6.2](#page-50-0) shows that in the long run, the ratio of the adjacent edge weights at every node behaves fundamentally different for the two models. It could however be possible to analyze the reinforced walk on finite circular graphs after a finite amount of time, increasing with the size of the graph, to make the transition from the circular graphs to **Z**.

6.2 *λ* ⁺**-Biased Edge-Reinforced Random Walk**

Similarly to the multiplicative bias studied above, we can also consider an additive bias. Define the *λ* ⁺-biased edge-reinforced random walk on **Z** as the sequence *Xⁿ* of random variables with

$$
X_0 = 0
$$

\n
$$
\mathbb{P}[X_{n+1} = x_n + 1 | X_n = x_n, ..., X_0 = x_0] = \frac{\lambda + w(n, x_n)}{\lambda + w(n, x_n) + w(n, x_n - 1)}
$$

\n
$$
w(n, z) = \underbrace{w(0, z)}_{=1 \text{ here}} + \sum_{i=0}^{n-1} \mathbb{1}_{X_i = z, X_{i+1} = z+1} + \mathbb{1}_{X_i = z+1, X_{i+1} = z}
$$

(where we require $\lambda \geq 0$). The problem with the definition of the λ^* -biased edge-reinforced random walk was that it could no longer be represented as a mixture of Markov chains since the probability of the occurrence of a certain edge sequence depended on the order in which the edges appeared in the sequence. This is not the case with the new definition given above. In other words, when the edge weights are represented by urns at every node, then the sequence of draws is now exchangeable, while this was not the case before.

Following [\[14,](#page-64-2) Lemma 1 and Lemma 2], the λ^+ -biased edge-reinforced random walk is equivalent to a mixture of MCs (or a RWRE). For shorter notation, we will adapt the notation of [Definition 2.2](#page-9-2) as follows: we call $\mathbb{P}[X_{n+1} = x + 1 | X_n = x] = P_c(x, x + 1) =: \omega_x$. Instead of looking at the distribution of the random conductances, we look directly at the distribution of the ω_x .

Figure 15: A random walk in a random environment on **Z**

The mixture of MCs equivalent to the λ^+ -biased walk is constructed by placing independent urns at every node which are coupled with which edges the random walk takes. Initially, we start with 1 black and $1 + \lambda$ (we can also have half a ball or any positive real number of balls) white balls, the black balls representing the edge to the left of the node, the white balls representing the edge to the right. Now, if a ball is drawn, we take the edge corresponding to the color and in addition to the drawn ball add two more balls of the same color (when the random walk returns to the node, the edge weight will have increased by two). Now, for any nodes to the right of 0, the first time the random walk gets there, there will already be 2 black balls in the urn, since the edge to the left was already traversed once. For nodes to the left of 0, there will already be $2 + \lambda$ white balls. So we have the following urns:

- at nodes > 0 , the urn initially contains 2 black and $1 + \lambda$ white balls
- at 0, the urn initially contains 1 black and $1 + \lambda$ white balls
- at nodes < 0 , the urn initially contains 1 black and $2 + \lambda$ white balls
- whenever a ball of one color is drawn, it is put back together with two more balls of the same color

With [\[14,](#page-64-2) Lemma 1 and Lemma 2], we get that the reinforced random walk with initial bias is equivalent to the mixture of Markov Chains where the ω_x are independent with the following distributions (also compare with [\[16,](#page-64-5) Section 5]):

for
$$
x > 0
$$
: $\omega_x \sim B\left(\frac{1+\lambda}{2}, \frac{2}{2}\right)$
\nfor $x = 0$: $\omega_x \sim B\left(\frac{1+\lambda}{2}, \frac{1}{2}\right)$
\nfor $x < 0$: $\omega_x \sim B\left(\frac{2+\lambda}{2}, \frac{1}{2}\right)$

where B is the beta distribution. We use the following:

Lemma 6.6	\n $\text{Let } A \sim B(\alpha, \beta) \text{ with } \alpha, \beta > 0 \text{ and } t \in \mathbb{R}. \text{ Then:}$ \n
\n $\mathbb{E}\left[\left(\frac{1-A}{A}\right)^t\right] = \begin{cases} \frac{\Gamma(\alpha-t)\Gamma(\beta+t)}{\Gamma(\alpha)\Gamma(\beta)} & \text{if } -\beta < t < \alpha \\ \infty & \text{otherwise} \end{cases}$ \n	\n $\mathbb{E}\left[\frac{1-A}{A}\right] = \begin{cases} \frac{\beta}{\alpha-1} & \text{if } \alpha > 1 \\ \infty & \text{otherwise} \end{cases}$ \n
\n $\mathbb{E}\left[\left(\frac{A}{1-A}\right)^t\right] = \begin{cases} \frac{\Gamma(\alpha+t)\Gamma(\beta-t)}{\Gamma(\alpha)\Gamma(\beta)} & \text{if } -\alpha < t < \beta \\ \infty & \text{otherwise} \end{cases}$ \n	\n $\mathbb{E}\left[\frac{A}{1-A}\right] = \begin{cases} \frac{\alpha}{\beta-1} & \text{if } \beta > 1 \\ \infty & \text{otherwise} \end{cases}$ \n

Proof We have:

$$
\mathbb{E}\left[\left(\frac{1-A}{A}\right)^t\right] = \frac{\Gamma\left(\alpha+\beta\right)}{\Gamma\left(\alpha\right)\Gamma\left(\beta\right)} \int_0^1 \frac{\left(1-x\right)^t}{x^t} \cdot x^{\alpha-1} \left(1-x\right)^{\beta-1} dx
$$
\n
$$
= \frac{\Gamma\left(\alpha+\beta\right)}{\Gamma\left(\alpha\right)\Gamma\left(\beta\right)} \int_0^1 x^{\alpha-t-1} \left(1-x\right)^{\beta+t-1} dx
$$
\n
$$
= \begin{cases} \frac{\Gamma\left(\alpha+\beta\right)}{\Gamma\left(\alpha\right)\Gamma\left(\beta\right)} \cdot \frac{\Gamma\left(\alpha-t\right)\Gamma\left(\beta+t\right)}{\Gamma\left(\alpha+t\right)} = \frac{\Gamma\left(\alpha-t\right)\Gamma\left(\beta+t\right)}{\Gamma\left(\alpha\right)\Gamma\left(\beta\right)} & \text{if } -\beta < t < \alpha \\ \infty & \text{otherwise} \end{cases}
$$

Hence,

$$
\mathbb{E}\left[\frac{1-A}{A}\right] = \begin{cases} \frac{\Gamma(\alpha-1)\Gamma(\beta+1)}{\Gamma(\alpha)\Gamma(\beta)} = \frac{\beta}{\alpha-1} & \text{if } -\beta < 1 < \alpha \\ \infty & \text{otherwise} \end{cases}
$$

The other statements follow from the fact that $1 - A \sim B(\beta, \alpha)$.

We can use [\[17,](#page-64-7) Theorem 1.7] to analyze transience and recurrence. It only applies to the situation where the ω_x are iid. We have only iid variables on the positive half-line as well as iid variables on the negative half-line, but for a moment we just assume that all *ωx* are distributed as the variables on the positive half-line. The σ from [\[17,](#page-64-7) Theorem 1.7] is $\frac{1-\omega_x}{\omega_x}$ in our case. We have, with $C_{\lambda} > 0$ a constant depending on λ :

$$
\mathbb{E}\left[\ln\left(\frac{1-\omega_x}{\omega_x}\right)\right] = C_{\lambda} \int_0^1 \ln\left(\frac{1-x}{x}\right) x^{\frac{\lambda-1}{2}} dx \begin{cases} > 0 & \text{if } 0 \le \lambda < 1 \\ = 0 & \text{if } \lambda = 1 \\ < 0 & \text{if } \lambda > 1 \end{cases}
$$

Unfortunately, this integral cannot be evaluated easily and in a nice way (in the general case; for *λ* = 1 there is a nice representation and probably also for *λ* ∈ **Q**). However, WolframAlpha shows how the integral behaves. To formally conclude here, a bit more work in analyzing the

 \Box

integral would be needed.

 $\lambda < 1$ $\lambda = 1$ $\lambda > 1$ [https://bit.ly/3yrjGCG](https://www.wolframalpha.com/input/?i=integral+from+0+to+1+of+ln%28%281-x%29%2Fx%29+*+x%5E%28%280.9-1%29%2F2%29+dx) [https://bit.ly/3F00yy3](https://www.wolframalpha.com/input/?i=integral+from+0+to+1+of+ln%28%281-x%29%2Fx%29+*+x%5E%28%281-1%29%2F2%29+dx) [https://bit.ly/3IK0jJQ](https://www.wolframalpha.com/input/?i=integral+from+0+to+1+of+ln%28%281-x%29%2Fx%29+*+x%5E%28%281.1-1%29%2F2%29+dx)

Hence, [\[17,](#page-64-7) Theorem 1.7] implies that the random walk in the environment where all ω_x are iid is transient to the right when $\lambda > 1$, recurrent when $\lambda = 1$, and transient to the left when λ < 1. But now, we are actually considering the random environment for the λ^+ -biased edgereinforced random walk. The ω_x with $x \leq 0$ all stochastically dominate the ω_x with $x > 0$ (the probability to go to the right is higher). Hence, for the case $\lambda > 1$, it is immediately (intuitively) clear that the random walk will still be transient to the right for this random environment, and still recurrent for $\lambda = 1$. For the case $\lambda < 1$ we see that the λ^+ -biased edge-reinforced random walk always returns to the root from the positive half-line. Now, we already know (e.g. by [\[16,](#page-64-5) Section 5]) that the edge-reinforced walk without initial bias is recurrent. The initial bias makes it only more likely to go to the right, hence the walk with initial bias will also be recurrent on the negative half-line for $\lambda < 1$ and therefore recurrent as a whole. We therefore get, even though a formal proof would still be needed:

The edge-reinforced random walk on **Z** with initial bias λ is recurrent for $0 \leq \lambda \leq 1$ and transient for $\lambda > 1$.

Next, we can look at the speed in the transient regime. Since the walk is only transient to the right if it is transient at all, it suffices to look at the distributions of ω_x for $x > 0$ (the finite number of steps spent to the left of the root 0 is irrelevant for the speed). We have by [Lemma 6.6:](#page-57-0)

$$
\mathbb{E}\left[\frac{1-\omega_x}{\omega_x}\right] = \begin{cases} \infty & \text{if } \lambda \le 1\\ \frac{2}{\lambda-1} & \text{if } \lambda > 1 \end{cases}
$$

Clearly, $\frac{2}{\lambda - 1}$ is (strictly) decreasing in λ for $\lambda > 1$, and hence, we have $\mathbb{E}\left[\frac{1 - \omega_x}{\omega_x}\right] < 1 \iff$ $\lambda > 3$. We now want to use [\[17,](#page-64-7) Theorem 1.16]. To get a statement of the type "positive speed \iff some condition", we therefore also need to look at the following, where we use again [Lemma 6.6:](#page-57-0)

$$
\mathbb{E}\left[\frac{\omega_x}{1-\omega_x}\right] = \infty \text{ for all choices of } \lambda
$$

Hence, by [\[17,](#page-64-7) Theorem 1.16]:

The edge-reinforced random walk on **Z** with initial bias *λ* has positive speed, if, and only **Theorem 6.8** if, $\lambda > 3$. If this is the case, then $\frac{X_n}{n} \to \frac{\lambda - 3}{\lambda + 1}$ almost surely. Otherwise, $\frac{X_n}{n} \to 0$ almost surely.

6.3 Reinforced Random Walk on Transient Environment

We now look at a special case of the LERRW where the initial edge weights are not all 1. Define the edge-reinforced random walk on **Z** with initially λ -biased environment as the sequence X_n

of random variables with

$$
X_0 = 0
$$

\n
$$
\mathbb{P}\left[X_{n+1} = x_n + 1 \mid X_n = x_n, \dots, X_0 = x_0\right] = \frac{w(n, x_n)}{w(n, x_n) + w(n, x_n - 1)}
$$

\n
$$
w(n, z) = \underbrace{w(0, z)}_{:=\lambda^z} + \sum_{i=0}^{n-1} \mathbb{1}_{X_i = z, X_{i+1} = z+1} + \mathbb{1}_{X_i = z+1, X_{i+1} = z}
$$

where $\lambda > 0$. This random walk can again be represented by a mixture of MCs, but the ω_x are no longer identically distributed (they are still independent, though).

Figure 16: Edge weights of the walk on transient environment at time of first visit to *z* > 0

By [Figure 16,](#page-59-0) we get the following distributions for ω_x :

for
$$
x > 0
$$
: $\omega_x \sim B\left(\frac{\lambda^x}{2}, \frac{1 + \lambda^{x-1}}{2}\right)$
\nfor $x = 0$: $\omega_x \sim B\left(\frac{1}{2}, \frac{1}{2\lambda}\right)$
\nfor $x < 0$: $\omega_x \sim B\left(\frac{1 + \lambda^x}{2}, \frac{\lambda^{x-1}}{2}\right)$

The electrical network corresponding to the random environment puts the following conductances **c** and resistances **r** on the edges:

$$
\begin{aligned}\n\text{for } z > 0: \qquad c_{\{z,z+1\}} = \prod_{x=1}^{z} \frac{\omega_x}{1 - \omega_x} & r_{\{z,z+1\}} = \prod_{x=1}^{z} \frac{1 - \omega_x}{\omega_x} \\
\text{for } z = 0: \qquad c_{\{0,1\}} = 1 & r_{\{0,1\}} = 1 \\
\text{for } z - 1 < 0: \qquad c_{\{z-1,z\}} = \prod_{x=z}^{0} \frac{1 - \omega_x}{\omega_x} & r_{\{z-1,z\}} = \prod_{x=z}^{0} \frac{\omega_x}{1 - \omega_x}\n\end{aligned}
$$

The effective resistance between 0 and ∞ is therefore (using the series law for the negative and positive half-lines and the parallel law to merge these):

$$
\mathbf{R}_{\text{eff}} = \left(\left(1 + \sum_{z \ge 1} \prod_{x=1}^{z} \frac{1 - \omega_x}{\omega_x} \right)^{-1} + \left(\sum_{z \le 0} \prod_{x=z}^{0} \frac{\omega_x}{1 - \omega_x} \right)^{-1} \right)^{-1}
$$

Hence, the random walk is recurrent if, and only if:

$$
\sum_{z\geq 1}\prod_{x=1}^{z}\frac{1-\omega_x}{\omega_x}=\infty=\sum_{z\leq 0}\prod_{x=z}^{0}\frac{\omega_x}{1-\omega_x}
$$

Otherwise, the random walk is transient. To get a better understanding, we calculate expecta-

tions. First, for $x > 0$ using [Lemma 6.6:](#page-57-0)

$$
\mathbb{E}\left[\frac{1-\omega_x}{\omega_x}\right] = \begin{cases} \frac{1+\lambda^{x-1}}{2} \cdot \left(\frac{\lambda^x}{2}-1\right)^{-1} = \frac{\lambda^{x-1}+1}{\lambda^x-2} & \text{if } \frac{\lambda^x}{2} > 1 \iff \lambda^x > 2\\ \infty & \text{otherwise} \end{cases}
$$

And for *x* < 0:

$$
\mathbb{E}\left[\frac{\omega_x}{1-\omega_x}\right] = \begin{cases} \frac{\lambda^x+1}{\lambda^{x-1}-2} & \text{if } \lambda^{x-1} > 2\\ \infty & \text{otherwise} \end{cases}
$$

Hence, if $\lambda > 2$:

$$
\mathbb{E}\left[\prod_{x=1}^{z} \frac{1-\omega_x}{\omega_x}\right] = \prod_{x=1}^{z} \frac{\lambda^{x-1}}{\lambda^{x-1}} \cdot \frac{1+\frac{1}{\lambda^{x-1}}}{\lambda - \frac{2}{\lambda^{x-1}}} = \prod_{x=1}^{z} \frac{1+\frac{1}{\lambda^{x-1}}}{\lambda - \frac{2}{\lambda^{x-1}}}
$$

Therefore, the expectation decreases exponentially for $\lambda > 2$. This seems to indicate transience for $\lambda > 2$ but a few calculations still remain to formally conclude. Indeed, for any $\lambda > 1$, the expectation $\mathbb{E}\left[\frac{1-\omega_x}{\omega_x}\right]$ will eventually be finite and of order $\frac{1}{\lambda}$ for *x* large enough, which can probably be used to show that the walk is transient for all $\lambda > 1$. The final proof is still open, but will hopefully follow in the near future.

7 Conclusion

We quickly look at how the considered modifications changed the behavior of the ERRW.

Table 1: Overview of results and conjectures on modified reinforced walks

There are two main conclusions which can be drawn from this overview. First, for reinforced random walks, having multiple random walkers moving in the same environment which influence each other doesn't seem to fundamentally change the behavior in comparison to a single walker. Indeed, after working on this topic for such a long period of time, the following (vague) conjecture seems reasonable: if, for a given graph and given reinforcement scheme, the reinforced random walk is recurrent for a single walker, then it is also recurrent for any finite number of walkers influencing each other, and vice versa. There is still a lot of work to do to get in any way closer to prove some form of this conjecture, but analyzing various toy models seems to indicate that this could be true. Of course, one would also have to formally define a reinforcement scheme to make a true mathematical claim, but the intuitive meaning should be clear.

The second conclusion is that introducing a bias can change the behavior. The bias, which makes the walk more transient, is sometimes competing with the reinforcement, which makes the walk more recurrent. This result is insofar expected as on regular trees, the reinforcement is also competing with the transient nature of a random walk on a tree. Indeed, depending on the strength of the reinforcement, there is a phase transition between recurrence and transience on trees, a result which we find again for the additive bias on **Z**.

All in all, the methods to prove the results given here are not new. Adding multiple walkers and a bias complicates an already complicated model even more. Therefore, the random walks were only analyzed on very simple graphs, because the methods used here do not work anymore for more general classes of graphs. Adding a bias or multiple walkers often destroys the property of exchangeability which the basic linearly reinforced random walk with a single walker possesses (the probability of taking a certain path only depends on how often each edge in the path is traversed, but not on the order of traversals). This makes the analysis harder and it is no longer possible to represent the reinforced walk as a RWRE, which was often the tool of choice to prove previous results. One goal of this thesis was to better understand how reinforced walks react to variations in the model. Some steps in the right direction were made by analyzing the behavior on simple graphs. At the same time, better tools to analzye these complicated models are still missing. The mathematics here was often adapted from the basic linear reinforcement case and is thus limited to a very restricted set of graphs.

7.1 Outlook

Even though some new results were obtained, many open questions remain. On the one hand, some proofs and calculations in this thesis still lack some details or formal precision. A natural next step would be to fill these gaps to be sure that the unfinished calculations and proof sketches actually show what they are supposed to show. However, the answers to more interesting questions are often only conjectured, and no proof idea has been found yet. The following four main points would be very interesting for future research:

- For the 2 walkers on a 3-node segment, which can also be seen as a kind of modified, 2-player Pólya urn, does the limit of the proportion of the left edge weight really have a density w.r.t. the Lebesgue measure? What is the distribution of the limit?
- For *k* walkers on **Z**, is the behavior for *k* walkers and for a single walker identical if the same reinforcement scheme is used?
- For linear reinforcement and additional multiplicative bias on **Z**, is a single walker transient for any bias $\lambda \neq 1$?
- For general graphs, does the behavior of the reinforced random walk only depend on the graph and the reinforcement scheme, but not on the (finite) number of walkers?

In answering these question, another goal would be to find more general techniques to analyze these types of random walks.

This thesis was really only a starting point in better understanding how changes to the model of the reinforced random walk will affect its behavior. While a definitive answer to the last question listed above still seems a long way off, the other three questions seem easier to deal with and can hopefully be answered in the near future.

8 References

- [1] Robin Pemantle. *A Survey of Random Processes with Reinforcement*. Probability Surveys, Vol. 4, pp. 1-79. Institute of Mathematical Statistics and Bernoulli Society, 2007. Accessible at [https://arxiv.org/abs/math/0610076.](https://arxiv.org/abs/math/0610076)
- [2] Burgess Davis. *Reinforced Random Walk*. Probability Theory and Related Fields, Vol. 84, pp. 203-229. Springer-Verlag, 1990. Accessible at [https://link.springer.com/article/10.](https://link.springer.com/article/10.1007/BF01197845) [1007/BF01197845.](https://link.springer.com/article/10.1007/BF01197845)
- [3] Tom Hutchcroft. *Random Walks and Uniform Spanning Trees*. Lecture notes. University of Cambridge, 2020. Accessible at <http://www.its.caltech.edu/~thutch/> under [https://](https://www.dropbox.com/s/9cuw6ay03vld3nj/RW_and_USTs_Public.pdf?dl=0) [www.dropbox.com/s/9cuw6ay03vld3nj/RW_and_USTs_Public.pdf?dl=0.](https://www.dropbox.com/s/9cuw6ay03vld3nj/RW_and_USTs_Public.pdf?dl=0)
- [4] Omer Angel, Nicholas Crawford and Gady Kozma. *Localization for Linearly Edge Reinforced Random Walks*. Duke Mathematical Journal, Vol. 163, No. 5, pp. 889-921. Duke University Press, 2014. Accessible at [https://arxiv.org/abs/1203.4010.](https://arxiv.org/abs/1203.4010)
- [5] Christophe Sabot and Pierre Tarrès. *Edge-Reinforced Random Walk, Vertex-Reinforced Jump Process and the Supersymmetric Hyperbolic Sigma Model*. Journal of the European Mathematical Society, Vol. 17, pp. 2353-2378. European Mathematical Society, 2015. Accessible at [https://arxiv.org/abs/1111.3991.](https://arxiv.org/abs/1111.3991)
- [6] Margherita Disertori, Christophe Sabot and Pierre Tarrès. *Transience of Edge-Reinforced Random Walk*. Communications in Mathematical Physics, Vol. 339, pp. 121-148. Springer-Verlag, 2015. Accessible at [https://arxiv.org/abs/1403.6079.](https://arxiv.org/abs/1403.6079)
- [7] Rémy Poudevigne-Auboiron. *Monotonicity and phase transition for the VRJP and the ERRW*. Published on arXiv, 2019. Accessible at [https://arxiv.org/abs/1911.02181.](https://arxiv.org/abs/1911.02181)
- [8] Robin Pemantle. *Vertex-Reinforced Random Walk*. Probability Theory and Related Fields, Vol. 92, pp. 117-136. Springer-Verlag, 1992. Accessible at [https://arxiv.org/abs/math/](https://arxiv.org/abs/math/0404041) [0404041.](https://arxiv.org/abs/math/0404041)
- [9] Persi Diaconis and David Freedman. *de Finetti's Theorem for Markov Chains*. The Annals of Probability, Vol. 8, No. 1 (Feb., 1980), pp. 115-130. Institute of Mathematical Statistics, 1980. Accessible at [https://projecteuclid.org/euclid.aop/1176994828.](https://projecteuclid.org/euclid.aop/1176994828)
- [10] Russell Lyons and Yuval Peres. *Probability on Trees and Networks*. Version of 6 August 2019. Cambridge University Press, 2017. Accessible at <https://rdlyons.pages.iu.edu/prbtree/> (online-only corrected edition at [https://rdlyons.pages.iu.edu/prbtree/book_corr.pdf\)](https://rdlyons.pages.iu.edu/prbtree/book_corr.pdf).
- [11] Stanislav Volkov. *Vertex-Reinforced Random Walk on Arbitrary Graphs*. The Annals of Probability, Vol. 29, No. 1 (Feb., 2001), pp. 66-91. Institute of Mathematical Statistics, 2001. Accessible at [https://projecteuclid.org/euclid.aop/1008956322.](https://projecteuclid.org/euclid.aop/1008956322)
- [12] Robin Pemantle and Stanislav Volkov. *Vertex-Reinforced Random Walk on Z has Finite Range*. The Annals of Probability, Vol. 27, No. 3 (Jul., 1999), pp. 1368-1388. Institute of Mathematical Statistics, 1999. Accessible at [https://projecteuclid.org/euclid.aop/1022677452.](https://projecteuclid.org/euclid.aop/1022677452)
- [13] Pierre Tarrès. *Vertex-Reinforced Random Walk on Z Eventually Gets Stuck on Five Points*. The Annals of Probability, Vol. 32, No. 3B (Jul., 2004), pp. 2650-2701. Institute of Mathematical Statistics, 2004. Accessible at [https://arxiv.org/abs/math/0410171.](https://arxiv.org/abs/math/0410171)
- [14] Robin Pemantle. *Phase Transition in Reinforced Random Walk and RWRE on Trees*. The Annals of Probability, Vol. 16, No. 3 (Jul., 1988), pp. 1229-1241. Institute of Mathematical Statistics, 1988. Accessible at [https://projecteuclid.org/euclid.aop/1176991687.](https://projecteuclid.org/euclid.aop/1176991687)
- [15] Russell Lyons and Robin Pemantle. *Random Walk in a Random Environment and First-Passage Percolation on Trees*. The Annals of Probability, Vol. 20, No. 1 (Jan., 1992), pp. 125-136. Institute of Mathematical Statistics, 1992. Accessible at [https://projecteuclid.org/euclid.aop/](https://projecteuclid.org/euclid.aop/1176989920) [1176989920.](https://projecteuclid.org/euclid.aop/1176989920)
- [16] Fabian Michel. *Linearly Edge-Reinforced Random Walks*. Bachelor's Thesis. Technical University of Munich, 2020.
- [17] Fred Solomon. *Random Walks in a Random Environment*. The Annals of Probability, Vol. 3, No. 1 (Feb., 1975), pp. 1-31. Institute of Mathematical Statistics, 1975. Accessible at [https:](https://projecteuclid.org/euclid.aop/1176996444) [//projecteuclid.org/euclid.aop/1176996444.](https://projecteuclid.org/euclid.aop/1176996444)
- [18] Pietro Muliere, Anna Maria Paganoni and Piercesare Secchi. *A two-color, randomly reinforced urn*. Journal of Statistical Planning and Inference, No. 136, pp. 1853-1874. 2006. Accessible at [https://www.mate.polimi.it/biblioteca/add/qmox/mox41.pdf.](https://www.mate.polimi.it/biblioteca/add/qmox/mox41.pdf)
- [19] Franz Merkl, Aniko Öry and Silke Rolles. *The "Magic Formula" for Linearly Edge-Reinforced Random Walks*. Statistica Neerlandica, Vol. 62, No. 3, pp. 345-363. Netherlands Society for Statistics and Operations Research, 2008. Accessible at [https://www-m5.ma.tum.de/](https://www-m5.ma.tum.de/foswiki/pub/M5/Allgemeines/SilkeRollesPublications/magic-snversion.pdf) [foswiki/pub/M5/Allgemeines/SilkeRollesPublications/magic-snversion.pdf.](https://www-m5.ma.tum.de/foswiki/pub/M5/Allgemeines/SilkeRollesPublications/magic-snversion.pdf)
- [20] Franz Merkl and Silke Rolles. *A Random Environment for Linearly Edge-Reinforced Random Walks on Infinite Graphs*. Probability Theory and Related Fields, Vol. 138, pp. 157- 176. Springer-Verlag, 2007. Accessible at [https://link.springer.com/article/10.1007/](https://link.springer.com/article/10.1007/s00440-006-0016-3) [s00440-006-0016-3.](https://link.springer.com/article/10.1007/s00440-006-0016-3)
- [21] Gady Kozma. *Reinforced Random Walk*. Published online. 6th European Congress of Mathematics, Kraków. 2012. Accessible at [https://arxiv.org/abs/1208.0364.](https://arxiv.org/abs/1208.0364)
- [22] Ewa Damek, Nina Gantert and Konrad Kolesko. *Absolute Continuity of the Martingale Limit in Branching Processes in Random Environment*. Electronic Communications in Probability, Vol. 24, pp. 1-13. Institute of Mathematical Statistics, 2019. Accessible at [https://arxiv.org/](https://arxiv.org/abs/1210.7664) [abs/1210.7664.](https://arxiv.org/abs/1210.7664)
- [23] Noam Berger and Eviatar Procaccia. *Mutually excited random walks*. Published on arXiv, 2012. Accessible at [https://arxiv.org/abs/1210.7664.](https://arxiv.org/abs/1210.7664)
- [24] Steven Lalley. *Random Walks on Infinite Discrete Groups*. Lecture notes. Northwestern University Summer School in Probability, 2018. Accessible at [https://sites.math.northwestern.](https://sites.math.northwestern.edu/~auffing/SNAP/rw-northwestern.pdf) [edu/~auffing/SNAP/rw-northwestern.pdf.](https://sites.math.northwestern.edu/~auffing/SNAP/rw-northwestern.pdf)
- [25] Steven Lalley. *One-Dimensional Random Walks*. Lecture notes. University of Chicago, 2016. Accessible at [http://galton.uchicago.edu/~lalley/Courses/312/RW.pdf.](http://galton.uchicago.edu/~lalley/Courses/312/RW.pdf)
- [26] Itai Benjamini and Gady Kozma. *Nonamenable Liouville Graphs*. Published online. 2010. Accessible at [https://arxiv.org/abs/1010.3365.](https://arxiv.org/abs/1010.3365)
- [27] Bruce Hill, David Lane and William Sudderth. *A Strong Law for Some Generalized Urn Processes*. The Annals of Probability, Vol. 8, No. 2 (Apr., 1980), pp. 214-226. Institute of Mathematical Statistics, 1980. Accessible at [https://projecteuclid.org/euclid.aop/1176994772.](https://projecteuclid.org/euclid.aop/1176994772)
- [28] Henrik Renlund. *Reinforced Random Walk*. Department of Mathematics, Uppsala University. 2005. Accessible at [http://www.diva-portal.org/smash/record.jsf?pid=diva2%](http://www.diva-portal.org/smash/record.jsf?pid=diva2%3A305167&dswid=6914) [3A305167&dswid=6914](http://www.diva-portal.org/smash/record.jsf?pid=diva2%3A305167&dswid=6914)
- [29] Russell Lyons and Robin Pemantle. *Correction: Random walk in a random environment and first-passage percolation on trees*. The Annals of Probability, Vol. 31, No. 1 (Jan., 2003), pp. 528- 529. Institute of Mathematical Statistics, 2003. Accessible at [https://projecteuclid.org/](https://projecteuclid.org/euclid.aop/1046294319) [euclid.aop/1046294319.](https://projecteuclid.org/euclid.aop/1046294319)
- [30] Elie Aidékon. *Transient random walks in random environment on a Galton-Watson tree*. Probability Theory and Related Fields, Vol. 142, No. 3-4 (Nov., 2008), pp. 525-559. Springer, 2008. Accessible at [https://arxiv.org/abs/0710.3377.](https://arxiv.org/abs/0710.3377)
- [31] Pablo Lessa. *Recurrence vs Transience: An introduction to random walks*. 2015. Accessible at <http://www.cmat.edu.uy/~lessa/otherwork.html> under [http://www.cmat.edu.uy/](http://www.cmat.edu.uy/~lessa/resource/randomwalknotes.pdf) [~lessa/resource/randomwalknotes.pdf.](http://www.cmat.edu.uy/~lessa/resource/randomwalknotes.pdf)
- [32] Russell Lyons. *The Ising Model and Percolation on Trees and Tree-Like Graphs*. Communications in Mathematical Physics, Vol. 125, No. 2 (Jun., 1989), pp. 337-353. Springer, 1989. Accessible at [https://projecteuclid.org/euclid.cmp/1104179469.](https://projecteuclid.org/euclid.cmp/1104179469)
- [33] Russell Lyons. *Random Walks and Percolation on Trees*. The Annals of Probability, Vol. 18, No. 3 (Jul., 1990), pp. 931-958. Institute of Mathematical Statistics, 1990. Accessible at [https:](https://projecteuclid.org/euclid.aop/1176990730) [//projecteuclid.org/euclid.aop/1176990730.](https://projecteuclid.org/euclid.aop/1176990730)

