

Entropy solutions to macroscopic IPM

Ángel Castro Daniel Faraco Björn Gebhard

Abstract

We investigate maximal potential energy dissipation as a selection criterion for subsolutions (coarse grained solutions) in the setting of the unstable Muskat problem. We show that both, imposing this criterion on the level of convex integration subsolutions, and the strategy of Otto based on a relaxation via minimizing movements, lead to the same nonlocal conservation law. Our main result shows that this equation admits an entropy solution for unstable initial data with an analytic interface.

1 Introduction

An outstanding open problem in hydrodynamics is the description of unstable interface configurations quickly leading to turbulent regimes. Examples are the thoroughly studied Saffman-Taylor, Rayleigh-Taylor and Kelvin-Helmholtz instabilities. In these unstable regimes Eulerian quantities, such as the velocity field, are very irregular and the Lagrangian trajectories typically fail to be uniquely defined. Hence uniqueness is not to be expected at the microscopical level, a phenomenon that in the physics literature is known as spontaneous stochasticity [94], and instead it will be desirable to have a well-defined deterministic evolution at the macroscopical level. The current paper provides such a macroscopical evolution in the context of the incompressible porous medium equation derived from maximal potential energy dissipation.

1.1 IPM and interfaces

Throughout the article, we will consider the incompressible porous media equation (IPM), given by

$$(1.1) \quad \begin{aligned} \partial_t \rho + \operatorname{div}(\rho v) &= 0, \\ \operatorname{div} v &= 0, \\ v &= -\nabla p - \rho e_2, \end{aligned}$$

on the two-dimensional periodic strip $\mathbb{T} \times \mathbb{R}$, \mathbb{T} denotes the flat one-torus of length 2π , and over a time interval $[0, T)$, $T > 0$. Here the fluid density $\rho : [0, T) \times \mathbb{T} \times \mathbb{R} \rightarrow$

\mathbb{R} , the velocity $v : [0, T) \times \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{R}^2$ and the pressure $p : [0, T) \times \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{R}$ are the unknowns, and $e_2 := (0, 1)^T \in \mathbb{R}^2$.

The model describes the evolution of a two-dimensional density dependent incompressible fluid in an overdamped scenario (the porous medium) and under the influence of gravity. It consists of the law for mass conservation, the incompressibility condition for the velocity field and Darcy's law (see [46, 76, 86, 8, 87, 93] for more physical background). Constants such as mobilities (viscosities), permeability of the medium, and gravity have been set to 1. System (1.1) also models the motion of an incompressible and viscous fluid in a Hele-Shaw cell [86], a different physical scenario with the same mathematical formulation.

Concerning initial conditions we are interested in the unstable interface case, i.e.

$$(1.2) \quad \rho_0(x) = \begin{cases} +1, & x_2 > \gamma_0(x_1), \\ -1, & x_2 < \gamma_0(x_1) \end{cases}$$

for a graph $\gamma_0 : \mathbb{T} \rightarrow \mathbb{R}$.

Generally speaking, if the initial data ρ_0 is sufficiently regular it is well-known that the IPM equation has a unique regular local in time solution, see [39, 24]. However, the problem of formation of singularities versus global existence is still open and only partial results are known. For example, the existence of solutions with Sobolev norms unbounded in time has recently been proven in [69].

In the case of discontinuous initial data of the type (1.2) the situation is even more subtle as the following dichotomy shows: If the denser fluid is below the lighter one, then the problem is stable and the existence of solutions is well-known (see Section 2.1). However, if the lighter fluid is below the heavier one, the problem is ill-posed (at least in the Muskat sense, and in the sense of bounded weak solutions).

1.2 Macroscopic IPM

In spite of this difficulty, there have been several attempts to understand the evolution of such an initial configuration at least in the coarse grained picture. Namely, on one hand Felix Otto discovered that in Lagrangian formulation IPM is a gradient flow and he suggested in the unstable situation a relaxation based on the corresponding minimizing movements scheme in the Wasserstein setting [83] (JKO scheme). On the other hand in [37] it was shown that IPM can be recast as a differential inclusion in the Tartar framework and therefore fits the De Lellis-Székelyhidi adaptation of convex integration in hydrodynamics [47, 48]. Subsequently, the full relaxation of the differential inclusion has been computed in [91] leading to a concept of coarse-grained solutions (subsolutions in the convex integration jargon). In Section 2 we present precise definitions and review the historical landmarks of the theory.

Let us remark that it also has been proven in [91] that in the case of a flat interface Otto's relaxation selects a convex integration subsolution, which turns out

to be the global in time entropy solution to a one-dimensional Burgers equation, reconciling both relaxation theories. In the case of a non-flat interface, the theory of convex integration starting from [20] has provided a number of subsolutions [12, 55, 81, 26]. In all these situations the starting point is an ansatz for the coarse grained density $\bar{\rho}$ and no selection criterion is available.

The aim of this paper is to use maximal potential energy dissipation as a selection criterion. Since as discovered by Otto in Lagrangian coordinates IPM is a gradient flow with respect to potential energy, this seems a natural approach. In any case, we first retake the strategy proposed by Otto in [83] in the case of non flat interfaces (the scheme is explained in Section 2.5.1 and Appendix B). We then reconcile it by selecting the subsolution in the convex integration terminology which at each time instant dissipates the most potential energy.

It can be shown (at least formally), that both, the relaxed minimizing movements scheme provided by [83] and imposing maximal potential energy dissipation among convex integration subsolutions, lead to the following equation

$$(1.3) \quad \begin{aligned} \partial_t \rho + \operatorname{div}(\rho v) + \partial_{x_2}(\rho^2) &= 0, \\ \operatorname{div} v &= 0, \\ v &= -\nabla p - \rho e_2, \end{aligned}$$

which will be referred to as *macroscopic IPM*. In Section 2 we explain in detail how the Muskat problem, the theory of subsolutions, convex integration for IPM and Otto's relaxation are connected. In particular, it will be explained the way in which (1.3) can offer a selection criterion for IPM subsolutions. For example, as opposed to other available subsolution, the solutions to macroscopic IPM are subsolutions as long as they exist.

1.3 Existence result and idea of proof

The bulk of the paper is devoted to proving the existence of an entropy solution for (1.3) with (1.2) as initial data. System (1.3) can be written as a single scalar nonlocal hyperbolic conservation law,

$$(1.4) \quad \partial_t \rho + \operatorname{div}(\rho T[\rho]) + \partial_{x_2}(\rho^2) = 0,$$

where $v = T[\rho]$ is a 0-order singular integral operator. Contrary to other nonlocal conservation laws with a more regular nonlocal feedback, see [10, 11, 13, 14, 31] for examples and [68] for a recent overview, a general existence and uniqueness theory for nonlocal terms as in (1.4) is not available. We bypass this by using the structure of the two-phase initial data (1.2). This approach, born out of necessity, not only provides us with the existence of a solution, but in addition allows us to learn about certain properties of it. More precisely, by showing that the Burgers' term $\partial_{x_2}(\rho^2)$ is able to tear up the initial discontinuity of the density even in the presence of the incompressible velocity v , we will prove the existence of a local in time solution which is Lipschitz for $t > 0$. This fact, although expected, is highly

nontrivial and presents many technical difficulties that will be tackled in Sections 4 to 7. Our main theorem itself, containing further properties of the solution, can be found in Section 3.

One main ingredient of our proof is to look at the evolution of level sets of the density ρ in suitably scaled coordinates and to adjust properly to leading order terms of this evolution. These steps, motivated in Section 4 and carried out in Section 5, reduce the initial value problem (1.2), (1.3) to a fixed point problem of the type

$$(1.5) \quad \eta(t, y) = \frac{1}{t^{1+\alpha}} \int_0^t \int_{-2}^2 \int_{\mathbb{T}} (K_s[\eta(s, \cdot)])(y, z) (h_s[\partial_{y_1} \eta(s, \cdot)])(y, z) dz_1 dz_2 ds - \frac{1}{t^\alpha} h_0(y),$$

for functions $\eta : [0, T] \times \mathbb{T} \times (-2, 2) \rightarrow \mathbb{R}$ describing the evolution of the level sets in superlinear order with respect to $t > 0$ small. Here $h_0(y)$ is one of the mentioned leading order terms, in fact the first order term, depending on the initial graph γ_0 , and $\alpha \in (0, 1)$. Moreover, for each $s > 0$, $y \in \mathbb{T} \times (-2, 2)$ and $\xi : \mathbb{T} \times (-2, 2) \rightarrow \mathbb{R}$ fixed, the function $z \mapsto (K_s[\xi])(y, z)$ is an order -1 convolution kernel induced by the Biot-Savart law. The dependence on ξ involves both $\xi(z)$ and $\xi(y)$ in the form of the difference $\xi(y) - \xi(z)$. Similar the function $(y, z) \mapsto h_s[\partial_{y_1} \xi](y, z)$, again considered for a fixed s and ξ , depends on the difference $\partial_{y_1} \xi(y) - \partial_{y_1} \xi(z)$. Thus after integration in z the regularity of the right-hand side of (1.5) with respect to y is the regularity of $\partial_{y_1} \eta$, i.e. the right-hand side when seen as an operator loses one derivative in y_1 .

In addition, as one of the main difficulties, also for potential equivalent reformulations of (1.5) where the above loss of a derivative might be avoided, we like to point out that the kernels $K_s[\xi]$ degenerate as $s \rightarrow 0$ to a one-dimensional kernel with singularity $\sim 1/(y_1 - z_1)$, i.e. to a order 0 integral kernel. Thus estimates for $K_s[\xi](y, \cdot)$ as an order -1 kernel, and thus after integration compensating the loss $\partial_{y_1} \xi(z)$ (but not $\partial_{y_1} \xi(y)$ though), can not be obtained uniformly in s .

Anyhow, in order to keep the paper enjoyable, we deal with (1.5) and its loss of derivative by considering real analytic initial interfaces. This allows us to use an adaptation of the Nirenberg-Nishida abstract Cauchy-Kovalevskaya Theorem. Still, the application of it, even when we continue to ignore the so far not mentioned factor $t^{-(1+\alpha)}$ on the right-hand side, takes quite a lot of effort. It is the second main part of our proof and can be found in Section 6.

Finally, Section 7 puts everything together to give a solution to the macroscopic IPM equation. In the appendices A,B,C we give a proof of a version of the abstract Cauchy-Kovalevskaya Theorem needed for our situation and we give some more details regarding the derivation of the macroscopic IPM equation.

1.4 On the entropy condition

We emphasize that the solution we find is an entropy solution of (1.3), or rather (1.4). The notion of entropy solution is stated in Definition 3.4. This is consistent

with the flat case $\gamma_0 = 0$ where, as said earlier, the relaxed minimizing movements scheme of Otto [83] converges to the entropy solution of (1.4) which in that case reduces to Burgers' equation. For an extended discussion concerning the selection of the entropy solution by the minimizing movements scheme (including other gradient flows as counterexamples where a corresponding selection fails) we refer to [63] where Gigli and Otto have revisited the IPM relaxation in the flat setting of Otto's original work [83]. Concerning the general, non-flat case it was also conjectured by Otto [82] that the convergence of the minimizing movements scheme to an entropy solution remains true.

Moreover, we point out that some sort of choice among solutions of (1.3) is critical in order to have a selection criterion. Indeed, already in the flat case solutions are clearly not unique, and also in the general case nonentropic solutions for (1.3) can be obtained in an easier way, for instance via (2.1) below, cf. Remark 3.3. We believe that the requirement of being an entropy solution leads to uniqueness for the initial value problem (1.2), (1.3), but since the velocity v depends on ρ in a comparably singular nonlocal way, standard methods do not seem to work and uniqueness of entropy solutions to macroscopic IPM stands as an interesting open question. In any case, we emphasize that for the scheme we present there is a unique solution, and therefore our maximal dissipating subsolution is amenable to numerical calculations.

1.5 Further questions

Besides the question of uniqueness of the found entropy solution, our work opens the door to many other questions with various levels of difficulty, such as improving the regularity of the solutions, considering initial interfaces (non analytical or non graphical, as for example in [26]) or other densities as initial data as well. It would be interesting to see whether the JKO scheme does converge rigorously or what happens in the case of different mobilities [73, 83]. On a more general level, there might be other selection criteria, for example based on surface tension [67] or on vanishing diffusion [75]. In any case, we hope our work encourages the research on finding a deterministic coarse grained evolution in presence of instabilities.

2 Ill-posedness and relaxation

The here considered unstable interface initial value problem is highly ill-posed. In this section we explain in which sense this ill-posedness holds, as well as a strategy based on convex integration and the relaxation of Otto [83] to overcome it. This will fully motivate system (1.3). A reader only interested in solving that system can go directly to Section 3.

2.1 The Muskat problem

If one assumes that

$$\rho(x, t) = \begin{cases} \rho_{\text{up}} & x_2 > f(x_1, t) \\ \rho_{\text{down}} & x_2 < f(x_1, t) \end{cases},$$

a closed equation from (1.1) can be obtained for the interface $(x_1, f(x_1, t))$. Indeed,

$$(2.1) \quad \partial_t f(x, t) = \frac{\rho_{\text{down}} - \rho_{\text{up}}}{4\pi} \int_{\mathbb{T}} \frac{\sin(y)(f_x(x, t) - f_x(x - y, t))}{\cosh(f(x, t) - f(x - y, t)) - \cos(y)} dy.$$

This equation is usually known in the literature as the Muskat equation honoring M. Muskat [76].

In the case $\rho_{\text{down}} > \rho_{\text{up}}$ the problem is stable and local existence and regularity of solutions can be proven in different functional settings and situations [38, 35, 36, 28, 43, 51, 3, 78, 5, 4, 7, 6, 18, 53, 72, 1, 27, 88, 57, 58, 30, 2], as well as global for small and medium size initial data [32, 33, 34, 56, 42, 64, 9, 52]. The existence of singularities for large initial data is shown in [22, 21] and also in [40, 41].

However, if $\rho_{\text{down}} < \rho_{\text{up}}$ the Muskat equation is ill-posedness [89, 38] and we can not use it to find solutions of IPM.

Surprisingly, convex integration has allowed us to construct solutions to IPM starting in this kind of unstable situations. They have been called mixing solutions and, in them, the initial interface between the two different densities disappears and a strip arises in which the two densities mix. We elaborate on these mixing solutions in the next sections. For a general picture on convex integration in the context of fluid dynamics we refer to the surveys [17, 49, 50].

2.2 IPM as differential inclusion

The first examples of nonuniqueness of weak solutions for (1.1) using convex integration were given in [37] by Córdoba, Gancedo and the second author for the initial value $\rho_0 = 0$. Their method bypasses the computation of the relaxation by means of so-called $T4$ configurations. After this Székelyhidi established in [91] the explicit relaxation of (1.1) for initial data of two-phase type, enabling a systematic investigation of interface problems in IPM. While the results in [37, 91] established ill-posedness of IPM in the class of essentially bounded solutions, Isett and Vicol could also show the existence of compactly supported $\mathcal{C}_{t,x}^\alpha$ -solutions for $\alpha < 1/9$, [66]. The starting point of our investigation is the relaxation of [91], which we will describe in this subsection.

We fix the notion of weak solutions to initial value problems with $|\rho_0| = 1$ almost everywhere. Note that for such initial data the last condition in Definition 2.1 below is an additional consistency requirement coming from the continuity, or rather transport, equation in (1.1).

Definition 2.1. A pair $\rho \in L^\infty((0, T) \times \mathbb{T} \times \mathbb{R})$, $v \in L^\infty(0, T; L^2(\mathbb{T} \times \mathbb{R}; \mathbb{R}^2))$ is a solution of (1.1), (1.2) provided for any $\varphi \in C_c^\infty([0, T) \times \mathbb{T} \times \mathbb{R})$ there holds

$$\begin{aligned} \int_0^T \int_{\mathbb{T} \times \mathbb{R}} \rho \partial_t \varphi + \rho v \cdot \nabla \varphi \, dx \, dt + \int_{\mathbb{T} \times \mathbb{R}} \rho_0 \varphi(0, \cdot) \, dx &= 0, \\ \int_0^T \int_{\mathbb{T} \times \mathbb{R}} v \cdot \nabla \varphi \, dx \, dt &= 0, \\ \int_0^T \int_{\mathbb{T} \times \mathbb{R}} (v + \rho e_2) \cdot \nabla^\perp \varphi \, dx \, dt &= 0, \end{aligned}$$

and $|\rho(t, x)| = 1$ for almost every $(t, x) \in (0, T) \times \mathbb{T} \times \mathbb{R}$.

A key step in the works [37, 91] is to recast weak solutions as defined above as solutions to a differential inclusion, to be able to use the Murat-Tartar compensated compactness formalism [92].

A pair (ρ, v) is a weak solution if and only if the triple $(\rho, v, m) \in L^\infty((0, T) \times \mathbb{T} \times \mathbb{R}) \times (L^\infty(0, T; L^2(\mathbb{T} \times \mathbb{R})))^2$ satisfies the linear system

$$(2.2) \quad \begin{aligned} \partial_t \rho + \operatorname{div} m &= 0, \\ \operatorname{div} v &= 0, \\ v &= -\nabla p - \rho e_2, \\ \rho(0, \cdot) &= \rho_0 \end{aligned}$$

distributionally, i.e. in analogy to Definition 2.1, together with

$$(2.3) \quad (\rho(t, x), v(t, x), m(t, x)) \in K := \{ (\rho, v, m) \in \mathbb{R}^5 : |\rho| = 1, m = \rho v \}$$

for almost every $(t, x) \in (0, T) \times \mathbb{T} \times \mathbb{R}$.

Then the relaxation of the incompressible porous media equation is understood as the relaxation of the corresponding differential inclusion, i.e. in the pointwise nonlinear constraint (2.3) the set K is replaced by its convex (or more generally Λ -convex) hull. Up to technicalities one can recover highly oscillatory solutions from this set, as the main theorem of Székelyhidi in [91] shows.

Theorem 2.2 ([91]). *Let $\bar{\rho} \in L^\infty((0, T) \times \mathbb{T} \times \mathbb{R})$ and $\bar{v}, \bar{m} \in L^\infty(0, T; L^2(\mathbb{T} \times \mathbb{R}))$ satisfy (2.2) in the sense of distributions. Suppose that there exists a bounded and open set $\mathcal{U} \subset (0, T) \times \mathbb{T} \times \mathbb{R}$ such that there holds (2.3) for almost every $(t, x) \notin \mathcal{U}$, while $(\bar{\rho}, \bar{v}, \bar{m})$ are continuous on \mathcal{U} with*

$$(2.4) \quad \begin{aligned} &(\bar{\rho}(t, x), \bar{v}(t, x), \bar{m}(t, x)) \\ &\in \{ (\rho, v, m) \in \mathbb{R}^5 : |\rho| < 1, |2(m - \rho v) + (1 - \rho^2)e_2| < (1 - \rho^2) \} \end{aligned}$$

for every $(t, x) \in \mathcal{U}$. Then there exist infinitely many weak solutions (ρ, v) of (1.1), (1.2) that coincide with $(\bar{\rho}, \bar{v})$ outside of \mathcal{U} and are arbitrarily close to $(\bar{\rho}, \bar{v})$ in the weak $L^2(\mathcal{U})$ -topology.

In the case of the IPM system the set on the right-hand side of (2.4) is indeed only the interior of the Λ -convex hull of K , which does not coincide with the full convex hull as opposed to the Euler equation, see [91]. Still, (2.4) describes all possible weak limits of solutions to the IPM system, cf. [91]. In view of that one can therefore truly speak about the full relaxation of IPM in the context of two-phase mixtures.

This fact has been quantified in [25], where the relation between solutions and subsolutions has been made precise through an adapted h -principle. In particular, this leads to additional properties of the solutions like a degraded macroscopic behaviour or the turbulent mixing at every time-slice property. The latter means that the solutions (ρ, v) induced by $(\bar{\rho}, \bar{v}, \bar{m})$ satisfy $\rho \in \mathcal{C}^0([0, T]; L^2_{weak}(\mathbb{T} \times (-R, R)))$, where R is some positive number with $\mathcal{U} \subset (0, T) \times \mathbb{T} \times (-R, R)$, and

$$(2.5) \quad \int_B 1 - \rho(t, x) dx \int_B 1 + \rho(t, x) dx > 0$$

for any $t \in (0, T)$ and any ball B fully contained in $\mathcal{U}_t := \{x \in \mathbb{T} \times \mathbb{R} : (t, x) \in \mathcal{U}\}$.

For later purpose we also point out the following possible upgrade of Theorem 2.2 which is obtained by using convex integration as in [25, 48].

Lemma 2.3. *Let $(\bar{\rho}, \bar{v}, \bar{m})$ be as in Theorem 2.2 and $\delta : [0, T] \rightarrow \mathbb{R}$ continuous with $\delta(0) = 0$, $\delta(t) > 0$, $t > 0$. Then there exist infinitely many solutions (ρ, v) as in Theorem 2.2 with the additional property that*

$$\left| \int_{\mathbb{T} \times \mathbb{R}} (\bar{\rho}(t, x) - \rho(t, x)) x_2 dx \right| \leq \delta(t)$$

for almost every $t \in [0, T]$.

Definition 2.4. Any triple $(\bar{\rho}, \bar{v}, \bar{m})$ satisfying the conditions of Theorem 2.2 is called a subsolution of (1.1), (1.2). The set \mathcal{U} , in other papers frequently also denoted by Ω_{mix} , is called the mixing zone of the subsolution.

Theorem 2.2 shifts the focus from a single solution to the investigation of subsolutions which are understood as possible coarse grained or averaged solutions. As subsolutions play the central role also in the present investigation we will frequently omit the bars in notation and instead mark solutions by (ρ_{sol}, v_{sol}) in case there is a chance of confusion.

2.3 Examples of subsolutions

The first examples of non constant subsolutions have been given in the same paper [91] of Székelyhidi for the perfectly flat initial interface, $\rho_0(x) = \text{sign}(x_2)$. Keeping the one-dimensional structure of the initial data one sees that $v = 0$, $m = -\alpha(1 - \rho^2)e_2$, $\alpha \in (0, 1)$ reduces (2.2), (2.3) to the one-dimensional conservation law

$$\partial_t \rho + \alpha \partial_{x_2}(\rho^2) = 0,$$

which has a unique entropy solution given by

$$\rho(t, x) = \begin{cases} 1, & x_2 > 2\alpha t, \\ \frac{x_2}{2\alpha t}, & -2\alpha t < x_2 < 2\alpha t, \\ -1, & x_2 < -2\alpha t. \end{cases}$$

It also has been mentioned in [91] that the limiting case $\alpha = 1$ is in agreement with the relaxation of Otto [83]. It coincides with (1.3) in the flat situation, cf. Section 4.1. In addition this case gives an upper bound for the the mixing zone. More precisely, it has been shown in [91] that the mixing zone at time $t > 0$, \mathcal{U}_t of any one-dimensional subsolution emanating from $\rho_0(x) = \text{sign}(x_2)$ is contained in the strip $[-1, 1] \times (-2t, 2t)$. A similar subsolution in the harder case of different viscosities was studied in [73]. Actually in [73] the Λ -hull of IPM with different viscosities and densities is computed.

In the context of IPM-hulls we also like to mention the article [65] which addresses the stationary, i.e. time-independent, IPM system. In [65] the lamination convex hull of that system is computed, as well as a rigidity result for its subsolutions and an application for long-term limits of (1.1) given.

The first examples of subsolutions giving rise to mixing solutions for IPM starting in a non flat interface $(x_1, f_0(x_1))$ have been provided in [20]. In this paper the density ρ of the subsolution is Lipschitz and the speed of opening of the mixing zone $c(x_1)$ ($= 2\alpha$ in the flat case above), satisfies $1 \leq c < 2$ and, as indicated, might depend on x_1 . The result of [20] holds for initial data $f_0 \in H^5(\mathbb{R})$, i.e. in a regime where the Muskat problem can not be solved. A numerical analysis of these subsolutions can be found in [19] where the formation of fingers can be observed. In [12], the semiclassical viewpoint developed in [20] is taken one step further (using semiclassical Sobolev spaces for example), providing an alternative proof to the main result of [20]. Indeed this later approach improves the subsolutions with respect to their regularity, as the boundary of the mixing zone is in $H^{5-\frac{1}{c(x_1)}}$, where $c(x_1)$ is the local speed of opening of the mixing zone, instead of merely in H^4 .

In [55] the authors constructed mixing solutions with an initial interface $f_0 \in \mathcal{C}^{3+\alpha}$ relaxing the initial regularity needed in [20] but relying on subsolutions with piecewise-constant density instead of Lipschitz. In this case the speed of opening of the mixing zone is $0 < c < 2$ with c uniform in x_1 . Thereafter the same kind of subsolutions have been constructed in [81] with variable speed of opening.

As mentioned before, mixing solutions obtained via convex integration are not unique. There are two reasons for this fact: a) different subsolutions can be found, b) infinitely many solutions, corresponding to different distributions of the density, emanate from every fixed subsolution. In order to deal with the point b), in [25] it has been shown that all the solutions obtained from a fixed subsolution can be chosen in such a way that they share averages over large sets, i.e., they are the same as the subsolution at a macroscopic level. One of the main points of the

present paper is to deal with point a). A particular instance of this multiplicity will be illustrated in Subsection 2.4 below.

The constructions of the subsolutions above seem to rely on the Saffman-Taylor instability (heavy fluid on top of a lighter fluid). In [20] it was observed that there also exist mixing solutions in the fully stable regime (see also [55]) which build on Kelvin-Helmholtz type instabilities (discontinuity of the velocity field, see [73] for a thorough discussion of this phenomena at the level of the hulls). Actually, the analysis in [20] indicates that the mixing can be created around any point of the interface which is not both flat (with zero slope) and stable. We call these kind of points fully stable points. It happens that in an initially overhanging interface there must be always a fully stable point. Partially unstable situations therefore require to find compatibility between the Muskat solution and mixing solutions, see [26]. Remarkably, the construction in [26] allows to answer the question on how to prolongate in time the singular solutions to the Muskat problem found in [22, 21], namely as mixing solutions.

As a last remark we like to point out that the subsolutions constructed in [20, 55, 81, 26] are local in time in the sense that although the involved functions exist over a potentially larger time interval, a small time interval has to be chosen in order to guarantee that they take values inside the convex hull, i.e. that (2.4) holds true. This is in contrast to the flat cases [91, 73] and to the subsolution constructed in the present paper. Although also here we will only prove a local in time existence result, the involved functions take values in (the closure) of the convex hull as long as they exist.

2.4 The subsolution selection problem

As described, the constructions from the previous subsection contain ansatzes for certain properties of the subsolution and hence for the induced mixing solutions of (1.1). To illustrate this freedom in the simplest case let us discuss the flat interface with $\gamma_0(x_1) = 0$ in slightly more detail. As in [91], setting $v \equiv 0$, $m = m_2(t, x_2)e_2$, $\rho = \rho(t, x_2)$ one sees that $(\rho, 0, m)$ is a subsolution if and only if

$$\begin{aligned} \partial_t \rho + \partial_{x_2} m_2 &= 0, & \rho(0, x) &= \text{sign}(x_2), \\ |\rho| &\leq 1, \\ |2m_2 + 1 - \rho^2| &< 1 - \rho^2, & \text{when } |\rho| < 1, \\ m_2 &= 0, & \text{when } |\rho| = 1 \end{aligned}$$

and the required continuity conditions hold true. Thus one could make the ansatz

$$(2.6) \quad m_2 = -\frac{1 - \rho^2}{2} + \frac{1 - \rho^2}{2} \xi_2$$

with $\xi_2 : [0, T) \times \mathbb{R} \rightarrow \mathbb{R}$ satisfying $|\xi_2| < 1$ and for any such ξ_2 solve the conservation law

$$\partial_t \rho + \partial_{x_2} \left((\xi_2(t, x_2) - 1) \frac{1 - \rho^2}{2} \right) = 0$$

with initial data $\rho_0(x_2) = \text{sign}(x_2)$ to get plenty of subsolutions with different mixing zones and density profiles. Note that in that sense ξ_2 , or rather the whole relation (2.6), plays the role of a constitutive law.

Summarizing once more, these examples show that not only each subsolution induces infinitely many solutions of the incompressible porous media equation sharing a common coarse grained, or averaged, behaviour, but that there are also infinitely many possibilities for this averaged evolution via the vast amount of possible subsolutions. This is a common problem in the construction of turbulent solutions emanating from unstable interface initial data, as for instance also for the Kelvin-Helmholtz instability [60, 74, 90] and the Rayleigh-Taylor instability in the context of the Euler equations [61, 59, 62].

2.5 A selection criterion

We now focus in the general, not necessarily flat, case on the selection of subsolutions in terms of choosing an appropriate relation between m , ρ and v such that (2.4) holds true provided $|\rho| \leq 1$.

First we will review the strategy proposed by F. Otto in [83] to relax system (1.1) based on its gradient flow structure in Lagrangian coordinates and we will formally obtain (1.3) from this relaxation. The strategy of Otto does not rely on the notion of a subsolution in the context of differential inclusions as in Section 2. However, the solution of (1.3) will be a (non-strict) subsolution with $m = \rho v - (1 - \rho^2)e_2$.

Thereafter, we will also give an argument to derive (1.3) in Eulerian coordinates directly based on subsolutions. Also here the starting point will be the gradient flow structure of (1.1). This second argument shows that the relaxation of Otto selects among all subsolutions precisely those that maximize the dissipation of potential energy at every time instant.

The relations are summarized in Figure 1.

2.5.1 Otto's relaxation

In this section we give a very brief summary of Otto's 5 step strategy leading to the macroscopic IPM equation (1.3). The discussion is not rigorous and even then we have put most of the explicit calculations to Appendix B. We adapt our notation to the one of [83], which, due to a different normalization, studies the evolution of

$$s(x, t) = \frac{1 - \rho(x, t/2)}{2}$$

instead of $\rho(x, t)$. In these coordinates the IPM system (1.1) reads

$$(2.7) \quad \begin{aligned} \partial_t s + u \cdot \nabla s &= 0, \\ \operatorname{div} u &= 0, \\ u &= -\nabla \Pi + s e_2, \end{aligned}$$

see Appendix B.

The starting point (Step 1) of Otto's relaxation is the vital fact that, when formulated in Lagrangian coordinates, IPM can be seen as a gradient flow with respect to the potential energy

$$E[\Phi] = - \int s(x, 0) \Phi(x) \cdot e_2$$

on the manifold

$$M_0 = \{\Phi \text{ one-to-one and onto, smooth, volume preserving maps}\}.$$

More precisely, if (s, u, Π) is a solution of (2.7) then the flow $\Phi(x, t)$ induced by u satisfies

$$(2.8) \quad \int \partial_t \Phi(\cdot, t) \cdot w = -dE[\Phi(\cdot, t)]w, \quad \forall w \in T_{\Phi(\cdot, t)}M_0,$$

where $dE[\Phi]w$ is the Fréchet derivative of the functional E at the point $\Phi \in M_0$ in the direction

$$w \in T_{\Phi}M_0 = \{w \text{ smooth and such that } \nabla \cdot (w \circ \Phi^{-1}) = 0\}.$$

Fast-forwarding a bit, the next steps of Otto consist of the introduction of a time discretization with stepsize $h > 0$ in form of a minimizing movements scheme (Step 2), the extension of the underlying manifold M_0 to its L^2 -closure in order to turn the potentially ill-posed discrete variational problems emanating from Step 2 to well-posed ones (Step 3), and a translation of the now existing sequence of minimizers back to Eulerian coordinates (Step 4). At this point there exists a sequence of functions $\theta^{(k)}$ corresponding to $s(\cdot, t)$ at time $t = kh$, but of course potentially on a coarse grained or "locally averaged" level, which is characterized by the following JKO scheme: $\theta^{(0)} = s(\cdot, 0)$, and given $\theta^{(k)}$, $\theta^{(k+1)}$ is the minimizer in K of

$$(2.9) \quad \frac{1}{2} \operatorname{dist}^2(\theta^{(k)}, \theta) + \frac{1}{2} \operatorname{dist}^2(1 - \theta^{(k)}, 1 - \theta) - h \int \theta(x) x_2$$

where the set K consists of measurable θ taking values in $[0, 1]$ and such that $\int \theta = \int s(x, 0)$, and $\operatorname{dist}^2(\theta_0, \theta_1)$ is the L^2 -Wasserstein distance

$$\operatorname{dist}^2(\theta_0, \theta_1) = \inf_{\Phi \in I(\theta_0, \theta_1)} \int \theta_0(x) |\Phi(x) - x|^2 dx$$

with

$$I(\theta_0, \theta_1) = \left\{ \Phi : \int \theta_1(y) \zeta(y) dy = \int \theta_0(x) \zeta(\Phi(x)) dx \quad \forall \zeta \in \mathcal{C}_0^0 \right\}.$$

Notice that this indeed is a relaxation of the original problem since the densities are no longer taking values in $\{0, 1\}$ and the transport maps are not necessarily injective.

The fifth and last step consists of passing to the limit $h \rightarrow 0$ whenever this is possible. In [83], Otto proved that this is the case for the unstable flat situation

$$s(x, 0) = \begin{cases} 0 & x_2 > 0 \\ 1 & x_2 < 0 \end{cases},$$

and that the limit of θ_h defined by

$$\theta_h(x, t) := \theta^{(k)}(x), \quad t \in [kh, (k+1)h)$$

is the unique entropy solution of the conservation law

$$\partial_t \theta + \partial_y (\theta(1 - \theta)) = 0.$$

For a different proof of this statement we refer to the work of Gigli and Otto [63] which in particular also contains a further examination of the relation between the minimizing movements scheme and the entropy condition.

In fact, it was conjectured by Otto [82] that the described scheme, if it converges, should also lead to an entropy solution of the macroscopic IPM equation in the general, non-flat case. We refer to Section 3 for the definition of entropy solutions.

In the rest of this section we sketch how at least formally system (1.3), or rather its equivalent reformulation in terms of $s(x, t)$, arises from the JKO-characterization (2.9) of the discrete functions $\theta^{(k)}$ when *assuming* suitable convergence. Our presentation here, as well as in Appendix B which contains some more details, is devoted to convey that the scheme indeed leads to the macroscopic IPM equation, rather than in providing a rigorous proof which we defer to future work. A similar computation was derived by Otto [82].

Fix t and denote for simplicity $\theta^0 := \theta_h(t)$, $\theta^1 := \theta_h(t + h)$. Furthermore, let Φ^h denote the transport map corresponding to $\text{dist}^2(\theta^0, \theta^1)$ and $\bar{\Phi}^h$ the transport map corresponding to $\text{dist}^2(1 - \theta^0, 1 - \theta^1)$. Then it can be shown that there are functions a^h, \bar{a}^h such that

$$\begin{aligned} \Phi^h(x) &= x + (\nabla a^h \circ \Phi^h)(x), \\ \bar{\Phi}^h(x) &= x + (\nabla \bar{a}^h \circ \bar{\Phi}^h)(x). \end{aligned}$$

This in fact is a consequence of Brenier's Theorem [16], still an argument is also provided in Appendix B.

Moreover, it can be deduced from first variations of the functional (2.9) that

$$(2.10) \quad a^h - \bar{a}^h = hx_2.$$

Now, we write $a^h = hp^h$, $\bar{a}^h = h\bar{p}^h$ and make the strong assumption that the introduced functions p^h, \bar{p}^h have a well defined \mathcal{C}^2 limit denoted by p, \bar{p} .

If this is the case we can pass to the limit $h \rightarrow 0$ and obtain

$$(2.11) \quad \partial_t \theta = -\operatorname{div}(\theta \nabla p),$$

$$(2.12) \quad \partial_t \theta = \Delta \bar{p} - \operatorname{div}(\theta \nabla \bar{p}).$$

Now (2.10) yields $p = \bar{p} + x_2$. Thus (2.11), (2.12) imply that

$$(2.13) \quad \Delta \bar{p} = \operatorname{div}((\nabla \bar{p} - \nabla p)\theta) = -\partial_{x_2} \theta.$$

Therefore, from (2.12) and (2.13), we deduce

$$\begin{aligned} \partial_t \theta &= -\partial_{x_2} \theta - \operatorname{div}(\nabla \bar{p} \theta) \\ &= -\partial_{x_2} \theta - \operatorname{div}((\nabla \bar{p} + \theta e_2) \theta) + \operatorname{div}(\theta^2 e_2) \end{aligned}$$

To finish we define $u = \nabla \bar{p} + \theta e_2$, which clearly satisfies $\operatorname{div} u = 0$, to get

$$\begin{aligned} \partial_t \theta + u \cdot \nabla \theta + \partial_{x_2} \theta - 2\theta \partial_{x_2} \theta &= 0, \\ u &= \nabla \bar{p} + \theta e_2, \\ \operatorname{div} u &= 0. \end{aligned}$$

Undoing the change of coordinates from the beginning, i.e., considering

$$\rho(t, x) = 1 - 2s(x, 2t),$$

one obtains (1.3). As said some more details can be found in Appendix B.

2.5.2 Transfer to subsolutions

Now we give an alternative derivation of the macroscopic system (1.3), taking a different route after step 1 of Otto's relaxation. i.e., the starting point is again the gradient flow structure of IPM saying that solutions of (1.1) seek to maximize the dissipation of potential energy at every time instance. However, at this point we do not care in which precise sense the dissipation is maximized (in Lagrangian coordinates with respect to the L^2 -metric on the manifold of area preserving diffeomorphisms). We instead simply extend the principle of maximal energy dissipation for solutions of (1.1) to its relaxation given in Theorem 2.2, i.e. we seek to investigate also subsolutions that decrease the potential energy at every time instant as much as possible.

Suppose that (ρ, v, m) is a subsolution in the sense of Definition 2.4. We define its associated relative potential energy

$$(2.14) \quad E_{rel}(t) := \int_{\mathbb{T} \times \mathbb{R}} (\rho(t, x) - \rho_0(x)) x_2 \, dx$$

and, for now formally, compute

$$(2.15) \quad \partial_t E_{rel}(t) = - \int_{\mathbb{T} \times \mathbb{R}} x_2 \operatorname{div} m(t, x) \, dx = \int_{\mathbb{T} \times \mathbb{R}} m_2(t, x) \, dx.$$

Moreover, similar to (2.6), condition (2.4) implies

$$m = \rho v - \frac{1 - \rho^2}{2} e_2 + \frac{1 - \rho^2}{2} \xi$$

almost everywhere for some $\xi : [0, T) \times \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{R}^2$ satisfying $|\xi| < 1$. Plugging this into (2.15) one deduces

$$\partial_t E_{rel}(t) = \int_{\mathbb{T} \times \mathbb{R}} \rho v_2 - (1 - \rho^2) \frac{1 - \xi_2}{2} \, dx.$$

Hence considering $\rho(t, \cdot)$, and therefore also $v(t, \cdot)$, cf. Section 4.2 below, to be given, one easily sees that the energy dissipation at time t is maximized in the closure of all admissible ξ with the choice $\xi(t, x) = -e_2$.

Hence choosing constantly $\xi = -e_2$, and therefore

$$(2.16) \quad m = \rho v - (1 - \rho^2) e_2$$

we deduce that (non-strict) subsolutions that maximize at each time instant the dissipation of potential energy are characterized as solutions of

$$(2.17) \quad \begin{aligned} \partial_t \rho + \operatorname{div} (\rho v - (1 - \rho^2) e_2) &= 0, \\ \operatorname{div} v &= 0, \\ v &= -\nabla p - \rho e_2. \end{aligned}$$

The above formal computation in (2.15) can be made rigorous under mild decay assumptions, as for instance shown in Appendix C. Here however, we like to state some further remarks.

First of all we emphasize that by choosing m as in (2.16) we do not obtain a subsolution in the sense of Definition 2.4, since (2.4) holds only in a non-strict sense, thus we speak about a non-strict subsolution. By considering instead

$$(2.18) \quad m = \rho v - \mu(1 - \rho^2) e_2,$$

i.e. $\xi = (1 - 2\mu) e_2$ with μ arbitrarily close to 1, but $\mu < 1$, one obtains strict subsolutions, and hence actual mixing solutions via Theorem 2.2, arbitrarily close

to the non-strict ones with maximal energy dissipation. However, in the remaining part of the paper we will solve (2.17) as the most standing out case and remark that a similar analysis leads to a subsolution corresponding to the system with m given by (2.18), cf. also Remark 3.3.

Moreover, we like to point out that in the flat case, where $v = 0$, system (2.17) is exactly the hyperbolic conservation law found in [91], whose entropy solution corresponds to the maximum speed of expansion of the mixing zone, cf. Section 2.3.

Furthermore, we remark that given a strict subsolution (ρ, v, m) with relative potential energy $E_{rel}(t)$ defined in (2.14) one obtains infinitely many mixing solutions (ρ_{sol}, v_{sol}) as in Theorem 2.2 with the additional property that their relative potential energy at almost every time t is arbitrarily close to $E_{rel}(t)$, see Lemma 2.3. In that sense there also exist actual mixing solutions with potential energy decay arbitrarily close to the maximal decay for subsolutions characterized by (2.17).

2.5.3 Comparison to selection criteria in related problems

As mentioned in Section 2.4 the selection of a meaningful subsolution is a general problem when studying hydrodynamic instabilities via differential inclusions. We briefly give an overview of previously applied selection criteria.

In the case of a perfectly flat interface the selection typically is done by reducing the subsolution system to a one-dimensional hyperbolic conservation law and picking the unique entropy solution as a natural candidate. This has been done in the context of the Kelvin-Helmholtz instability for the Euler equations [90], the Rayleigh-Taylor instability for the inhomogeneous Euler equations [62], and as discussed in all detail above for the flat unstable Muskat problem in IPM [91].

Another approach, selecting the subsolution that at initial time maximizes the total energy dissipation, has been applied in the context of the non-flat Kelvin-Helmholtz instability [74] within the class of all subsolutions with vorticity concentrated on a finite number of sheets, and thereafter in the class of one-dimensional self-similar subsolutions emanating from the flat Rayleigh-Taylor instability modelled by the Euler equations in Boussinesq approximation [61]. This strategy has been motivated by Dafermos' entropy rate admissibility criterion [44], which has also been investigated in [29, 54] for convex integration solutions of the compressible Euler equations. In view of Section 2.5.2 also the selection criterion considered in the present paper falls into that category. However, in contrast to [74, 61] the selection applies among all possible subsolutions (with certain natural decay at infinity) and not only within a special subclass, and it applies at all times instead of only the initial time.

Another way to select subsolutions globally in time has been studied in [59] in the context of the flat Rayleigh-Taylor instability for the Euler equations in Boussinesq approximation. Similar as in Section 2.5.2 above, the underlying geo-




Figure 1: Relaxation of IPM in Eulerian coordinates via subsolutions on the left and in Lagrangian coordinates via minimizing movements on the right.

metric principle of the equation, in that case the least action principle, has been imposed on the level of subsolutions leading to a degenerate elliptic variational problem that turns out to be formally equivalent to the direct relaxation of the least action principle by Brenier [15]. However, solutions obtained from this relaxation conserve the total energy, which is inconsistent with anomalous energy dissipation present in turbulent regimes. In view of that, in [59] an additional term, responsible for energy dissipation, but subject to certain choices, has been added in the variational problem. In contrast the here considered relaxation of IPM is not relying on any comparable choices.

3 The main result

According to the previous section we consider on $\mathbb{T} \times \mathbb{R}$ the system

$$(3.1) \quad \begin{aligned} \partial_t \rho + \operatorname{div}(\rho v + \rho^2 e_2) &= 0, \\ \operatorname{div} v &= 0, \\ v &= -\nabla p - \rho e_2 \end{aligned}$$

with initial data (1.2), i.e.

$$\rho_0(x) = \begin{cases} +1, & x_2 > \gamma_0(x_1), \\ -1, & x_2 < \gamma_0(x_1) \end{cases}$$

for a sufficiently regular function $\gamma_0 : \mathbb{T} \rightarrow \mathbb{R}$. In fact we here consider the case of a real analytic initial interface. For completeness we also state the notion of a general weak solution to system (3.1).

Definition 3.1. A pair $\rho \in L^\infty((0, T) \times \mathbb{T} \times \mathbb{R})$, $v \in L^\infty(0, T; L^2(\mathbb{T} \times \mathbb{R}; \mathbb{R}^2))$ is a solution of (3.1), (1.2) provided for any $\varphi \in C_c^\infty([0, T) \times \mathbb{T} \times \mathbb{R})$ there holds

$$\begin{aligned} \int_0^T \int_{\mathbb{T} \times \mathbb{R}} \rho \partial_t \varphi + (\rho v + \rho^2 e_2) \cdot \nabla \varphi \, dx \, dt + \int_{\mathbb{T} \times \mathbb{R}} \rho_0 \varphi(0, \cdot) \, dx &= 0, \\ \int_0^T \int_{\mathbb{T} \times \mathbb{R}} v \cdot \nabla \varphi \, dx \, dt &= 0, \\ \int_0^T \int_{\mathbb{T} \times \mathbb{R}} (v + \rho e_2) \cdot \nabla^\perp \varphi \, dx \, dt &= 0. \end{aligned}$$

Theorem 3.2. *Let $\gamma_0 : \mathbb{T} \rightarrow \mathbb{R}$ be real analytic. Then the initial value problem (3.1), (1.2) has a local in time solution with the following properties*

- (i) ρ and v are continuous on $[0, T) \times \mathbb{T} \times \mathbb{R} \setminus \{(0, x_1, \gamma_0(x_1)) : x_1 \in \mathbb{T}\}$,
- (ii) at positive times $\rho(t, \cdot)$ is Lipschitz continuous, $v(t, \cdot)$ is log-Lipschitz continuous with

$$(3.2) \quad \|\nabla \rho(t, \cdot)\|_{L^\infty(\mathbb{T} \times \mathbb{R})} \leq C_0 t^{-1},$$

$$(3.3) \quad |v(t, x) - v(t, x')| \leq C_0 t^{-1} |(x - x') \log |x - x'||$$

for $t \in (0, T)$, $x, x' \in \mathbb{T} \times \mathbb{R}$, $|x - x'| \leq 1$ and a constant $C_0 > 0$ depending on γ_0 ,

- (iii) for $t \in (0, T)$ there exist two real analytic curves $\gamma_t(\cdot, \pm 1) : \mathbb{T} \rightarrow \mathbb{R}$ such that $\rho(t, x) = 1$ whenever $x_2 \geq \gamma_t(x_1, 1)$ and $\rho(t, x) = -1$ whenever $x_2 \leq \gamma_t(x_1, -1)$. Moreover, $\rho(t, \cdot)$ maps the remaining set into $(-1, 1)$. Also there the level sets $\Gamma_t(h) := \{x \in \mathbb{T} \times \mathbb{R} : \rho(t, x) = h\}$, $h \in (-1, 1)$ are given by

graphs of real analytic functions $\gamma_t(\cdot, h) : \mathbb{T} \rightarrow \mathbb{R}$. Furthermore, the joint map $[0, T) \times \mathbb{T} \times [-1, 1] \rightarrow \mathbb{R}$, $(t, x_1, h) \mapsto \gamma_t(x_1, h)$ belongs to the space $\mathcal{C}^1([0, T); \mathcal{C}^1(\mathbb{T} \times [-1, 1]))$ and there exists a real analytic function $s_0 : \mathbb{T} \rightarrow \mathbb{R}$ such that

$$(3.4) \quad \gamma_t(x_1, h) = \gamma_0(x_1) + t(2h + s_0(x_1)) + o(t)$$

with respect to $\|\cdot\|_{\mathcal{C}^1(\mathbb{T} \times [-1, 1])}$ as $t \rightarrow 0$,

(iv) for any Lipschitz continuous $\eta : \mathbb{R} \rightarrow \mathbb{R}$ there holds the balance

$$(3.5) \quad \partial_t(\eta(\rho)) + \operatorname{div}(\eta(\rho)v + Q(\rho)e_2) = 0,$$

with initial data $\eta(\rho)(0, \cdot) = \eta(\rho_0)$ and flux $Q(\rho) := \int_0^\rho 2\eta'(s)s \, ds$.

Remark 3.3. a) In fact the function $s_0 : \mathbb{T} \rightarrow \mathbb{R}$ appearing in (3.4) is precisely the normal part of the initial velocity when evaluated in $(x_1, \gamma_0(x_1))$. See Section 4.2, in particular equation (4.9), for the definition and further discussion.

b) Note that (iii) implies that ρ is piecewise \mathcal{C}^1 with the exceptional set given by $\{(t, x_1, \gamma_t(x_1, \pm 1)) : t \in [0, T), x_1 \in \mathbb{T}\}$.

c) Equation (3.5) is a priori understood in analogy to Definition 3.1, i.e. in a distributional sense. However, given the regularity of ρ and v it in fact holds pointwise almost everywhere on $(0, T) \times \mathbb{T} \times \mathbb{R}$, cf. Section 7.

d) The balance (3.5) in particular states that ρ is an entropy solution for the conservation law $\partial_t \rho + \operatorname{div}(\rho v + \rho^2 e_2) = 0$, cf. Definition 3.4 below.

e) We notice that, for an analytic initial interface, the Muskat equation (2.1) can be solved for short time in order to find a solution to the macroscopic IPM system (3.1), which at the same time is also a solution for IPM (see [22] and in the case of the vortex-sheet problem [23]). However, this solution is not an entropy solution. Moreover, piece-wise constant solutions of (3.1) also could be constructed but again they would not be entropy solutions.

f) As discussed earlier in Section 2.5.2 the solution (ρ, v) given by Theorem 3.2 induces only a non-strict subsolution by setting $m := \rho v - (1 - \rho^2)e_2$. However, an analogous existence statement remains true when replacing the first equation of (3.1) by

$$\partial_t \rho + \operatorname{div}(\rho v + \mu \rho^2 e_2) = 0$$

corresponding to a choice of m as in (2.18) and thus to strict subsolutions when $\mu < 1$. This can be seen for instance by rescaling time and considering the nonlocal velocity field $\mu^{-1}v$ in Sections 5, 6.

g) Notice that (iii) describes precisely the mixing zone \mathcal{U} of the subsolution, cf. Definition 2.4, where the corresponding solutions develop a mixing behaviour. In particular, from (3.4) one can deduce the initial growth of the mixing zone, which is linear in time. When combined with [25], it also implies the observed degraded mixing property of solutions (the closer to the upper boundary, the bigger the volume fraction of the heavier fluid).

For completeness we add in the following the notion of an entropy solution for equation (3.1). Note that (3.1) is a nonlocal hyperbolic conservation law. As is common for such equations, cf. e.g. [10, 11, 13, 14, 31], the notion of an entropy solution is the one for the corresponding local conservation law where the otherwise nonlocal velocity field is considered as a fixed local one:

Definition 3.4 (Entropy solution). A solution (ρ, v) in the sense of Definition 3.1 is called an entropy solution provided for any $\varphi \in \mathcal{C}_c^\infty([0, T] \times \mathbb{T} \times \mathbb{R})$, $\varphi \geq 0$ and any convex $\eta : \mathbb{R} \rightarrow \mathbb{R}$ with induced flux $Q(\rho) := \int_0^\rho 2\eta'(s)s ds$ there holds

$$\int_0^T \int_{\mathbb{T} \times \mathbb{R}} \eta(\rho) \partial_t \varphi + (\eta(\rho)v + Q(\rho)e_2) \cdot \nabla \varphi dx dt + \int_{\mathbb{T} \times \mathbb{R}} \eta(\rho_0) \varphi(0, \cdot) dx \geq 0.$$

We remark that typically the set of η for which the stated imbalance is required to hold is taken to be a strict subset of all convex functions, such as for instance the Kruřkov family $\{|\cdot - c| : c \in \mathbb{R}\}$, [70], see also [45]. Since our solution anyhow satisfies the stronger condition (iv), we refrain at this point from restricting the set of entropies.

In any case, due to the nature of the nonlocality of our velocity field, which is a zero order singular integral operator with respect to the density ρ (see Section 4.2) the uniqueness of the found entropy solution remains open.

4 Preliminary observations

We begin our investigation with a look at the illustrative example of a perfectly flat initial interface $\gamma_0(x_1) = 0$, and some known facts concerning the nonlocal velocity field v , in particular at initial time, in the non-flat case.

4.1 The flat interface

In the perfectly flat case, $\gamma_0 = 0$, a x_1 -independent solution of equation (3.1) is obtained by observing that $v = 0$ and solving the Riemann problem for Burgers' equation

$$\partial_t \rho + \partial_{x_2}(\rho^2) = 0, \quad \rho(0, x_2) = \text{sign}(x_2).$$

The unique entropy solution is Lipschitz continuous at positive times and explicitly given by

$$\rho(t, x) = \begin{cases} 1, & x_2 > 2t, \\ \frac{x_2}{2t}, & |x_2| \leq 2t, \\ -1, & x_2 < -2t. \end{cases}$$

As discussed earlier, cf. Section 2.3, this solution bounds the mixing zone in the class of all one-dimensional IPM-subolutions.

However, in rescaled coordinates $y \mapsto x$, $x = (y_1, ty_2)$ the solution is given by the stationary profile

$$(4.1) \quad \rho(t, y_1, ty_2) = \phi_0(y) := \begin{cases} 1, & x_2 > 2, \\ \frac{1}{2}y_2, & |y_2| \leq 2, \\ -1, & y_2 < -2, \end{cases}$$

or in other words the level sets $\rho(t, \cdot)^{-1}(\{h\})$, $h \in (-1, 1)$ are given by flat lines $\{x : x_2 = 2ht\}$ that as time evolves are pulled apart with speed $2h$.

Of course these are simple reformulations, but a key point in our analysis is an appropriate extension of this principle to the general, non-flat case where the velocity field does not vanish. This will be done by keeping the profile $\phi_0(y)$ on the right-hand side of (4.1) and allowing the transformation $y \mapsto x$ to be of the type $x = (y_1, ty_2 + f(t, y))$, i.e. we keep the “pulling”-term ty_2 dealing with the Burgers’ term $\partial_{x_2}(\rho^2)$ in the equation and allow the level sets to have a general form reacting to the nonlocal velocity field. The details in terms of induced equations for f are in Section 5.

4.2 Biot-Savart and the initial velocity field

The flat case discussed in the previous subsection is a very special case in the sense that $v = 0$ and the resulting equation is local. In the general case a key feature of both systems, IPM and the relaxation, is the nonlocal relation between the density ρ and the velocity field v . More precisely, the last two equations in (1.1), (3.1) resp., i.e. the incompressibility condition and Darcy’s law, can be understood by means of a 0-order convolution operator. Indeed, taking the curl of Darcy’s law one sees that at each time $v(t, \cdot)$ is an incompressible vectorfield with vorticity given by

$$(4.2) \quad \partial_{x_1} v_2(t, x) - \partial_{x_2} v_1(t, x) = -\partial_{x_1} \rho(t, x).$$

Thus when requiring decay as $|x_2| \rightarrow \infty$ the velocity field v is, at least in the case of our interest, uniquely determined in terms of the Biot-Savart operator

$$(4.3) \quad v(t, x) = (K * (-\partial_{x_1} \rho(t, \cdot)))(x) = \int_{\mathbb{T} \times \mathbb{R}} K(x - z) (-\partial_{x_1} \rho(t, z)) dz.$$

On $\mathbb{T} \times \mathbb{R}$ the kernel K is given by

$$(4.4) \quad K(z) := \frac{1}{4\pi} \frac{(-\sinh(z_2), \sin(z_1))^T}{\cosh(z_2) - \cos(z_1)},$$

and as usual, K is the orthogonal gradient of the corresponding Green’s function

$$(4.5) \quad G(z) := \frac{1}{4\pi} \log(\cosh(z_2) - \cos(z_1)).$$

Relation (4.3) has to be interpreted accordingly at initial time $t = 0$ due to the fact that $-\partial_{x_1}\rho_0$ is only a measure supported on the interface

$$\Gamma_0 := \{ (x_1, \gamma_0(x_1)) : x_1 \in \mathbb{T} \}.$$

Thus the initial velocity field $v_0(x)$ is the one of a vortex-sheet and therefore discontinuous across the interface.

Lemma 4.1. *The unique square integrable solution of*

$$(4.6) \quad v = -\nabla p - \rho_0 e_2, \quad \operatorname{div} v = 0 \quad \text{on } \mathbb{T} \times \mathbb{R}$$

is given by

$$(4.7) \quad v_0(x) = \int_{\mathbb{T}} K \begin{pmatrix} x_1 - z_1 \\ x_2 - \gamma_0(z_1) \end{pmatrix} 2\gamma'_0(z_1) dz_1$$

for $x \notin \Gamma_0$, while the one-sided limits at Γ_0 are given by

$$(4.8) \quad \lim_{\substack{y \rightarrow (x_1, \gamma_0(x_1)) \\ \pm(y_2 - \gamma_0(y_1)) > 0}} v_0(y) = p.v. \int_{\mathbb{T}} K \begin{pmatrix} x_1 - z_1 \\ \gamma_0(x_1) - \gamma_0(z_1) \end{pmatrix} 2\gamma'_0(z_1) dz_1 \\ \mp \frac{\gamma'_0(x_1)}{1 + \gamma'_0(x_1)^2} \begin{pmatrix} 1 \\ \gamma'_0(x_1) \end{pmatrix}.$$

Proof. First of all one can check that the right-hand side of (4.7) defines a locally integrable solution of (4.6) with exponential decay as $|x_2| \rightarrow \infty$. Thus standard elliptic estimates imply that this is the only solution with these properties.

In order to compute the one-sided limits we write

$$K(z) = \frac{1}{2\pi} \frac{z^\perp}{|z|^2} \eta(z_1) + K_{reg}(z),$$

where $\eta : \mathbb{T} \times \mathbb{R}$ is a smooth periodic cutoff function with $\eta(z_1) = 1$ for $|z_1| \leq 1$ and $\eta(z_1) = 0$ for $|z_1| \geq 2$, and the regular part $K_{reg} : \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{R}^2$,

$$K_{reg}(z) := K(z) - \frac{1}{2\pi} \frac{z^\perp}{|z|^2} \eta(z_1)$$

is smooth. In fact K_{reg} is harmonic where $\eta(z_1) = 1$. Furthermore, using complex notation we write $z^\perp/|z|^2 = (1/(iz))^*$ where z^* denotes complex conjugation.

Then denoting by $v_{0,reg}$ the contribution from the regular part K_{reg} we have

$$v_0(y) - v_{0,reg}(y) = \left(\frac{1}{2\pi i} \int_{\mathbb{T}} \frac{2\gamma'_0(z_1)}{y - (z_1 + i\gamma_0(z_1))} dz_1 \right)^* \\ = - \left(\frac{1}{2\pi i} \int_{\Gamma_0} \frac{1}{\xi - y} \frac{2\gamma'_0(\xi_1)}{1 + i\gamma'_0(\xi_1)} d\xi \right)^*$$

for $y \notin \Gamma_0$. Now taking one sided limits $y \rightarrow x \in \Gamma_0$ expression (4.8) follows from the Sokhotski-Plemelj formula, see [77]. \square

Formulas (4.8) show that the initial velocity field is still continuous across the interface in the normal direction. Therefore the (not normalized) normal velocity at the interface $s_0 : \mathbb{T} \rightarrow \mathbb{R}$,

$$(4.9) \quad \begin{aligned} s_0(x_1) &:= v_0(x_1, \gamma_0(x_1)) \cdot \begin{pmatrix} -\gamma_0'(x_1) \\ 1 \end{pmatrix} \\ &= p.v. \int_{\mathbb{T}} K \begin{pmatrix} x_1 - z_1 \\ \gamma_0(x_1) - \gamma_0(z_1) \end{pmatrix} 2\gamma_0'(z_1) dz_1 \cdot \begin{pmatrix} -\gamma_0'(x_1) \\ 1 \end{pmatrix} \end{aligned}$$

is well-defined. It will play an important role in our further analysis as it dictates the motion of Lagrangian particles at the interface to first order when ignoring the Burgers' term $\partial_{x_2}(\rho^2)$.

5 Formal ansatz for general graphs

In this section we will reformulate problem (3.1), (1.2) as a suitable fixed point problem. The reformulation here is understood on a formal level. We will solve the fixed point problem in Section 6 and a posteriori justify the transformations in Section 7.

5.1 Rescaling and level set function

In order to capture the effect of the Burger's part, cf. Section 4.1, we introduce the following ansatz for ρ . Assume that there exists $f : [0, T) \times \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{R}$ sufficiently regular with

$$(5.1) \quad f(0, y) = \gamma_0(y_1)$$

and such that for every $t \in (0, T)$, $y_1 \in \mathbb{T}$ the map $\mathbb{R} \rightarrow \mathbb{R}$, $y_2 \mapsto ty_2 + f(t, y_1, y_2)$ is a monotone diffeomorphism.

Then each of the transformations $X_t : \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{T} \times \mathbb{R}$, $t \in (0, T)$,

$$X_t(y) = \begin{pmatrix} y_1 \\ ty_2 + f(t, y) \end{pmatrix}$$

is a diffeomorphism as well.

We now seek to find a solution of (3.1), (1.2) on $[0, T)$ having the property that

$$(5.2) \quad \rho(t, X_t(y)) = \phi_0(y_2) = \begin{cases} +1, & y_2 \geq 2, \\ \frac{1}{2}y_2, & y_2 \in (-2, +2), \\ -1, & y_2 \leq -2. \end{cases}$$

For $t > 0$ we compute

$$(5.3) \quad DX_t(y) = \begin{pmatrix} 1 & 0 \\ \partial_{y_1} f(t, y) & t + \partial_{y_2} f(t, y) \end{pmatrix},$$

$$(5.4) \quad DX_t(y)^{-1} = \begin{pmatrix} 1 & 0 \\ \frac{-\partial_{y_1} f(t, y)}{t + \partial_{y_2} f(t, y)} & \frac{1}{t + \partial_{y_2} f(t, y)} \end{pmatrix},$$

$$(5.5) \quad \nabla \rho(t, X_t(y)) = \frac{1}{2(t + \partial_{y_2} f(t, y))} \begin{pmatrix} -\partial_{y_1} f(t, y) \\ 1 \end{pmatrix} \mathbb{1}_{(-2, 2)}(y_2),$$

such that the first equation of (3.1), when written in non divergence form, under the ansatz (5.2) is equivalent to

$$0 = \mathbb{1}_{(-2, 2)}(y_2) \begin{pmatrix} -\partial_{y_1} f(t, y) \\ 1 \end{pmatrix} \cdot \left(\begin{pmatrix} 0 \\ y_2 + \partial_t f(t, y) - 2\phi_0(y_2) \end{pmatrix} - v(t, X_t(y)) \right).$$

Since $2\phi_0(y_2) = \mathbb{1}_{(-2, 2)}(y_2) = y_2$, expanding the above equation leads to

$$(5.6) \quad \partial_t f(t, y) = v(t, y_1, ty_2 + f(t, y)) \cdot \begin{pmatrix} -\partial_{y_1} f(t, y) \\ 1 \end{pmatrix}$$

for $(t, y_1, y_2) \in (0, T) \times \mathbb{T} \times (-2, 2)$.

Note that in view of (5.5) the velocity field in (5.6) is always considered in directions normal to the level sets of ρ .

5.2 Transformation of the velocity field

For $t > 0$ we have that $v(t, \cdot)$ (in all reasonable scenarios) is given by the Biot-Savart law (4.3), cf. Section 4.2.

Applying the transformation $X_t(y)$ we compute the velocity field $v(t, y_1, ty_2 + f(t, y)) = v(t, X_t(y))$ occurring in (5.6). First of all formulas (5.3) and (5.5) imply

$$\begin{aligned} v(t, X_t(y)) &= - \int_{\mathbb{T} \times \mathbb{R}} K(X_t(y) - z) \partial_{x_1} \rho(t, z) dz \\ &= - \int_{\mathbb{T} \times \mathbb{R}} K(X_t(y) - X_t(z)) \partial_{x_1} \rho(t, X_t(z)) \det DX_t(z) dz \\ &= \frac{1}{2} \int_{-2}^2 \int_{\mathbb{T}} K(X_t(y) - X_t(z)) \partial_{y_1} f(t, z) dz_1 dz_2. \end{aligned}$$

Next we compute the full right-hand side of (5.6) and exploit the fact that the velocity field $v(t, X_t(y))$ is only needed in normal directions. More precisely, for

$z \neq y$ we have

$$\begin{aligned}
& \partial_{y_1} f(t, z) K(X_t(y) - X_t(z)) \cdot \begin{pmatrix} -\partial_{y_1} f(t, y) \\ 1 \end{pmatrix} \\
&= \partial_{y_1} f(t, z) \nabla G(X_t(y) - X_t(z)) \cdot \begin{pmatrix} 1 \\ \partial_{y_1} f(t, y) \end{pmatrix} \\
&= \partial_{y_1} f(t, y) \nabla G(X_t(y) - X_t(z)) \cdot \begin{pmatrix} 1 \\ \partial_{y_1} f(t, z) \end{pmatrix} \\
&\quad - \partial_1 G(X_t(y) - X_t(z)) (\partial_{y_1} f(t, y) - \partial_{y_1} f(t, z)) \\
&= -\partial_{y_1} f(t, y) \frac{d}{dz_1} (G(X_t(y) - X_t(z))) \\
&\quad - K_2(X_t(y) - X_t(z)) (\partial_{y_1} f(t, y) - \partial_{y_1} f(t, z)).
\end{aligned}$$

Thus after integration we obtain an additional cancelation in the convolution, i.e. there holds

$$\begin{aligned}
(5.7) \quad & v(t, X_t(y)) \cdot \begin{pmatrix} -\partial_{y_1} f(t, y) \\ 1 \end{pmatrix} \\
&= -\frac{1}{2} \int_{-2}^2 \int_{\mathbb{T}} K_2(X_t(y) - X_t(z)) (\partial_{y_1} f(t, y) - \partial_{y_1} f(t, z)) dz_1 dz_2.
\end{aligned}$$

5.3 Equation for f

Combining equation (5.7) with (5.6) we see that (3.1) can after our ansatz be written in the closed form

$$(5.8) \quad \partial_t f(t, y) = -\frac{1}{2} \int_{-2}^2 \int_{\mathbb{T}} K_2(\tilde{\Delta} X_t(y, z)) (\partial_{y_1} f(t, y) - \partial_{y_1} f(t, z)) dz_1 dz_2$$

where

$$\tilde{\Delta} X_t(y, z) := X_t(y) - X_t(z) = \begin{pmatrix} y_1 - z_1 \\ t(y_2 - z_2) + f(t, y) - f(t, z) \end{pmatrix}$$

also depends on f . Via translation in z_1 equation (5.8) can also be written as

$$(5.9) \quad \partial_t f(t, y) = -\frac{1}{2} \int_{-2}^2 \int_{\mathbb{T}} K_2(\Delta X_t(y, z)) \Delta \partial_{y_1} f_t(y, z) dz_1 dz_2,$$

where we have abbreviated

$$\begin{aligned}
\Delta X_t(y, z) &:= \begin{pmatrix} z_1 \\ t(y_2 - z_2) + \Delta f_t(y, z) \end{pmatrix}, \\
\Delta f_t(y, z) &:= \gamma_0(y_1) - \gamma_0(y_1 - z_1) + t(s_0(y_1) - s_0(y_1 - z_1)) \\
&\quad + \frac{t^{1+\alpha}}{2} (\eta(t, y) - \eta(t, y_1 - z_1, z_2)), \\
\Delta \partial_{y_1} f_t(y, z) &:= \gamma'_0(y_1) - \gamma'_0(y_1 - z_1) + t(s'_0(y_1) - s'_0(y_1 - z_1)) \\
&\quad + \frac{t^{1+\alpha}}{2} (\partial_{y_1} \eta(t, y) - \partial_{y_1} \eta(t, y_1 - z_1, z_2)).
\end{aligned}$$

The latter form turns out to be more convenient to work with.

5.4 One more ansatz

One important assumption in the above derivation is the invertibility of the maps $(X_t)_{t>0}$. In order to guarantee that we further make the ansatz

$$(5.10) \quad f(t, y) = \gamma_0(y_1) + ts_0(y_1) + \frac{1}{2}t^{1+\alpha}\eta(t, y),$$

where $\alpha \in (0, 1)$ and the functions $s_0 : \mathbb{T} \rightarrow \mathbb{R}$, $\eta : (0, T) \times \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{R}$ are sufficiently regular. In order to avoid potential confusion we emphasize that the function η has nothing to do with an entropy as for example appearing in Definition 3.4. Furthermore, we remark that the particular choice of $\alpha \in (0, 1)$ is not important, see Section 6.7 for further discussion.

By this ansatz f satisfies (5.1), and the desired invertibility can be assumed to be true for a small time interval (depending on $\|\partial_{y_2}\eta\|_{L^\infty}$). Moreover, since at $t = 0$ there holds $\partial_t f = s_0$, $\partial_{y_1} f = \gamma'_0$, passing formally to the limit on the right-hand side of (5.9) one sees that s_0 necessarily is given by

$$(5.11) \quad -\frac{1}{2} \int_{-2}^2 \int_{\mathbb{T}} K_2(\Delta X_0(y_1, z_1)) \Delta \gamma'_0(y_1, z_1) dz_1 dz_2,$$

where

$$\begin{aligned} \Delta X_0(y_1, z_1) &:= \begin{pmatrix} z_1 \\ \gamma_0(y_1) - \gamma_0(y_1 - z_1) \end{pmatrix}, \\ \Delta \gamma'_0(y_1, z_1) &:= \gamma'_0(y_1) - \gamma'_0(y_1 - z_1). \end{aligned}$$

A quick computation similar to the one in Section 5.2 and comparison with (4.9) shows that the above expression is precisely the normal component of the initial velocity evaluated at $(y_1, \gamma_0(y_1))$, i.e. (5.11). This shows that the function $s_0(y_1)$ is indeed forced to be the normal component of $v_0(y_1, \gamma_0(y_1))$.

Finally we integrate (5.9) in time and use (5.10), (5.11) in order to deduce that for f to be a solution to (5.9) η must be a solution of the following fixed point problem:

$$(5.12) \quad \begin{aligned} \eta(t, y) = & -\frac{1}{t^{1+\alpha}} \int_0^t \int_{-2}^2 \int_{\mathbb{T}} K_2(\Delta X_s(y, z)) \Delta \partial_{y_1} f_s(y, z) \\ & - K_2(\Delta X_0(y_1, z_1)) \Delta \gamma'_0(y_1, z_1) dz_1 dz_2 ds. \end{aligned}$$

6 Existence of a solution for analytic graphs

Our goal is to show that for a real analytic $\gamma_0 : \mathbb{T} \rightarrow \mathbb{R}$ there exists a unique local in time solution η of problem (5.12). The proof relies on the following version of

the abstract Cauchy-Kovalevskaya theorem based on the formulation of Nishida [80], see also [79].

In order to avoid confusion we emphasize that throughout Section 6 every symbol $\rho, \rho', \bar{\rho}, \rho_0$ denotes a positive constant referring to the size of the domain of analyticity. This is done in analogy to [80]. At no time in Section 6 the density function $\rho(t, x)$, which we seek to construct, or the initial density $\rho_0(x)$ are mentioned.

Theorem 6.1. *Let $(B_\rho)_{\rho \in (0, \rho_0)}$, $\rho_0 > 0$ be a scale of Banach spaces with $\|\cdot\|_{\rho'} \leq \|\cdot\|_\rho$ for $0 < \rho' < \rho < \rho_0$ and consider the integral equation*

$$(6.1) \quad u(t) = \frac{1}{a(t)} \int_0^t F(u(s), s) ds$$

for a given continuous function $a : [0, \infty) \rightarrow \mathbb{R}$ with $a(t) > 0$ for $t > 0$. If F is such that

(i) *there exists $R > 0, T > 0$ such that for every $0 < \rho' < \rho < \rho_0$ the map*

$$\left\{ u \in B_\rho : \|u\|_\rho < R \right\} \times [0, T) \rightarrow B_{\rho'}, \quad (u, t) \mapsto F(u, t)$$

is well-defined and continuous,

(ii) *there exists $b : [0, T) \rightarrow [0, \infty)$ continuous such that for any $0 < \rho' < \rho < \rho_0$ and all $u, v \in B_\rho$, $\|u\|_\rho < R$, $\|v\|_\rho < R$, $t \in [0, T)$ there holds*

$$\|F(u, t) - F(v, t)\|_{\rho'} \leq \frac{b(t)}{\rho - \rho'} \|u - v\|_\rho,$$

(iii) *$F(0, \cdot) \in L^1(0, T; B_\rho)$ for any $\rho \in (0, \rho_0)$ and there exists $c : [0, T) \rightarrow [0, \infty)$ continuously differentiable on $(0, T)$, continuous on $[0, T)$ with $c(0) = 0$, as well as $c'(t) > 0$ for $t > 0$, such that for all $\rho \in (0, \rho_0)$, $t \in (0, T)$ there holds*

$$\frac{1}{a(t)} \int_0^t \|F(0, s)\|_\rho ds \leq \frac{c(t)}{\rho_0 - \rho},$$

(iv) *for a constant $K > 0$ the functions $a(t)$, $b(t)$, $c(t)$ appearing in (6.1), (ii), (iii) satisfy the relation*

$$(6.2) \quad \sup_{s \in (0, t)} \left| \frac{b(s)c(s)}{c'(s)} \right| \leq Ka(t)c(t), \quad t \in (0, T),$$

then there exists a constant $\bar{a} = \bar{a}(K, R) > 0$ and a unique $u(t)$ which for any $\rho \in (0, \rho_0)$ maps the interval $\{t \in [0, T) : c(t) < \bar{a}(\rho_0 - \rho)\}$ continuously into the R -ball of B_ρ . Moreover, u satisfies (6.1) and $\|u(t)\|_\rho = O\left(\frac{c(t)}{\rho_0 - \rho}\right)$ as $t \rightarrow 0$. In particular $u(0) = 0$.

For the choices $a(t) = 1$, $b(t) = c_1$, $c(t) = c_2 t$ with some constants $c_1, c_2 > 0$ the above theorem is the abstract Cauchy-Kovalevskaya theorem in the formulation of Nishida [80]. The proof of Theorem 6.1 requires indeed just some minor modifications which are presented in Appendix A. For a related generalization of the abstract Cauchy-Kovalevskaya theorem see also [84, 85].

We will apply Theorem 6.1 in the following situation.

Lemma 6.2. *Let $c_1, c_2 > 0$ and $\alpha \in (0, 1)$. There exists $T = T(\alpha)$, $K = K(c_1, c_2) > 0$ such that $a(t) := t^{1+\alpha}$, $b(t) := c_1 t^{1+\alpha} |\log t|$, $c(t) := c_2 t^{1-\alpha} |\log t|$ satisfy (6.2).*

Proof. Consider $T \in (0, 1)$ such that

$$(1 - \alpha) |\log t| \geq 2, \quad |\log t| t^\alpha \leq 1,$$

for all $t \in (0, T)$. Then for $0 < s < t < T$ there holds

$$\frac{b(s)c(s)}{c'(s)} = c_1 \frac{s^{2+\alpha} |\log s|^2}{(1 - \alpha) |\log s| - 1} \leq c_1 s^2 |\log s| \leq c_1 t^2 |\log t| = \frac{c_1}{c_2} a(t)c(t).$$

Thus (6.2) holds true with $K := c_1 c_2^{-1}$. □

6.1 Banach spaces

Set

$$\Omega_0 := \mathbb{T} \times (-2, 2),$$

as well as

$$U_\rho := \{z \in \mathbb{C} : |\operatorname{Im}(z)| < \rho\}, \quad \Omega_\rho := U_\rho \times (-2, 2)$$

for $\rho > 0$.

We define the space B_ρ to consist of all continuous functions $\eta : \Omega_0 \rightarrow \mathbb{R}$, $y \mapsto \eta(y)$ which satisfy

- (i) for every $y_2 \in (-2, 2)$ the function $\eta(\cdot, y_2)$ extends to a holomorphic function $U_\rho \rightarrow \mathbb{C}$ which is again denoted by $\eta(\cdot, y_2)$,
- (ii) the derivative $\partial_{y_2} \eta : \Omega_\rho \rightarrow \mathbb{C}$ exists, is uniformly continuous and $\partial_{y_2} \eta(\cdot, y_2)$ is holomorphic on U_ρ for every $y_2 \in (-2, 2)$,
- (iii) the norm

$$\|\eta\|_\rho := \|\eta\|_{L^\infty(\Omega_\rho)} + \|\partial_{y_1} \eta\|_{L^\infty(\Omega_\rho)} + \|\partial_{y_2} \eta\|_{L^\infty(\Omega_\rho)}$$

is finite.

For clarification, here $\partial_{y_1}\eta$ denotes the complex derivative in the first component, while $\partial_{y_2}\eta$ is the real partial derivative with respect to the second component. Although the two derivatives are of slightly different nature, we still use a gradient notation $\nabla_y\eta := (\partial_{y_1}\eta, \partial_{y_2}\eta)^T$.

The extension in (i) strictly speaking is the extension of the 2π -periodic function $\eta(\cdot, y_2) : \mathbb{R} \rightarrow \mathbb{R}$.

Clearly each B_ρ is a Banach space and $B_\rho \subset B_{\rho'}$, $\|\cdot\|_{\rho'} \leq \|\cdot\|_\rho$ whenever $\rho' < \rho$. Moreover, for the introduced scale of spaces we have the following lemma, which is a direct consequence of Cauchy integral formula for analytic functions.

Lemma 6.3 (Cauchy). *Let $0 < \rho' < \rho$ and $\eta \in B_\rho$. Then for $j = 1, 2$ there holds*

$$\|\partial_{y_1}\partial_{y_j}\eta\|_{L^\infty(\Omega_{\rho'})} \leq \frac{C}{\rho - \rho'} \|\eta\|_\rho$$

for $C = (2\pi)^{-1}$.

In particular $\partial_{y_1}\eta$ is, as η itself, Lipschitz continuous on Ω_0 . This together with the assumed uniform continuity of $\partial_{y_2}\eta$ implies

Lemma 6.4. *Let $\rho > 0$ and $\eta \in B_\rho$. Then $\eta : \Omega_0 \rightarrow \mathbb{R}$ extends to $\mathcal{C}^1(\overline{\Omega}_0)$ and $\eta(\cdot, y_2), \partial_{y_2}\eta(\cdot, y_2)$ are real analytic for each $y_2 \in [-2, 2]$.*

Also note that $\partial_{y_2}\partial_{y_1}\eta(y) = \partial_{y_1}\partial_{y_2}\eta(y)$ for $\eta \in B_\rho, y \in \Omega_\rho$ for instance by means of Cauchy's integral formula.

6.2 Notation

From now on we fix $\alpha \in (0, 1)$ and a real analytic initial datum $\gamma_0 : \mathbb{T} \rightarrow \mathbb{R}$. Clearly γ_0 can be extended to a holomorphic function defined on $U_{2\rho_0}$ for some $\rho_0 > 0$ small.

Hence all (complex) derivatives are uniformly bounded on U_{ρ_0} , i.e. there exist a constant $C_0 > 0$ such that for instance

$$(6.3) \quad \|\gamma_0'\|_{L^\infty(U_{\rho_0})} \leq C_0.$$

More generally, in the following $C_0 > 0$ always denotes a constant depending solely on the $L^\infty(U_{\rho_0})$ -norm of a finite amount of derivatives of γ_0 . In contrast $C > 0$ usually denotes a constant not depending on γ_0 . Both constants typically change from line to line. Also we point out that distinguishing C_0 from C is not essential for the proof of Theorem 3.2.

For a pair $a = (a_1, a_2) \in \mathbb{R} \times \mathbb{C}$ we denote

$$(6.4) \quad |a|_* := (|a_1|^2 + |a_2|^2)^{\frac{1}{2}} = (a_1^2 + a_2\bar{a}_2)^{\frac{1}{2}}.$$

Moreover, whenever we write $|z_1|$ for $z_1 \in \mathbb{T}$ we mean the absolute value of the unique representant of z_1 in $[-\pi, \pi)$. In particular we will also use $|a|_*$ for pairs $a \in \mathbb{T} \times \mathbb{C}$.

For any function $g : \Omega_{\rho_0} \rightarrow \mathbb{C}^n$ or $h : U_{\rho_0} \rightarrow \mathbb{C}^n$ we abbreviate

$$(6.5) \quad \begin{aligned} \Delta g(y, z) &:= g(y) - g(y_1 - z_1, z_2), \\ \Delta h(y_1, z_1) &:= h(y_1) - h(y_1 - z_1) \end{aligned}$$

for $y = (y_1, y_2) \in \Omega_{\rho_0}$, $z = (z_1, z_2) \in \Omega_0$ and $y_1 \in U_{\rho_0}$, $z_1 \in \mathbb{T}$ resp.. In proofs we will most of the time omit the points (y, z) and simply write Δg , Δh .

Furthermore, for $t \geq 0$ and $\eta \in B_{\rho_0}$ we define $f_t^\eta : \Omega_{\rho_0} \rightarrow \mathbb{C}$, $X_t^\eta : \Omega_{\rho_0} \rightarrow \mathbb{C}^2$,

$$(6.6) \quad f_t^\eta(y) := \gamma_0(y_1) + t s_0(y_1) + \frac{1}{2} t^{1+\alpha} \eta(y), \quad X_t^\eta(y) := \begin{pmatrix} y_1 \\ t y_2 + f_t^\eta(y) \end{pmatrix}.$$

The function $s_0 : U_{\rho_0} \rightarrow \mathbb{C}$ will be introduced in Lemma 6.6 below. At time $t = 0$ we simply write $X_0(y_1)$ instead of $X_0^\eta(y)$. The second component of $X_t^\eta(y)$ is denoted by $X_{t,2}^\eta(y)$. There is no need to distinguish the first component, since it is just given by y_1 .

6.3 Preliminary lemmas

In order to define the function F as a complex extension of the functional appearing in (5.12) we need some preparation.

Recall that the second component K_2 of the Biot-Savart kernel on $\mathbb{T} \times \mathbb{R}$ is given by

$$K_2(a) = K_2(a_1, a_2) = \frac{1}{4\pi} \frac{\sin(a_1)}{\cosh(a_2) - \cos(a_1)}.$$

Thus for fixed $a_1 \in \mathbb{T}$ the canonical extension of $K_2(a_1, \cdot)$ to $a_2 \in \mathbb{C}$ is holomorphic on the open set $\{a_2 \in \mathbb{C} : \cosh(a_2) - \cos(a_1) \neq 0\}$. We define

$$\mathcal{U} := \{a \in \mathbb{T} \times \mathbb{C} : \cosh(a_2) - \cos(a_1) \neq 0\}.$$

Lemma 6.5. *Let $\kappa \in (0, 1/2)$. The sets*

$$\mathcal{U}^\kappa := \{(a_1, a_2) \in \mathbb{T} \times \mathbb{C} : |\operatorname{Im}(a_2)| < \kappa(|a_1| + |\operatorname{Re}(a_2)|), \quad |\operatorname{Im}(a_2)| < \pi/2\}$$

are subsets of \mathcal{U} with $\partial \mathcal{U}^\kappa \cap \partial \mathcal{U} = \{0\}$. Moreover, there exists a constant $C > 0$ depending on κ such that for all $a \in \mathcal{U}^\kappa$, $j = 0, 1, 2$ there holds

$$(6.7) \quad |\partial_{a_2}^j K_2(a)| \leq C |a|_*^{-(1+j)}.$$

Proof. Let $a \in \overline{\mathcal{U}^\kappa} \cap \partial \mathcal{U}$, $a_2 = u + iv$. Then

$$0 = \cosh(a_2) - \cos(a_1) = \cosh(u) \cos(v) - \cos(a_1) + i \sinh(u) \sin(v)$$

implies $v = 0$ and thus $\cosh(u) = \cos(a_1)$, which is only possible for $a_1 = u = 0$; or $u = 0$ and $|v| = |a_1|$, which in the closure of \mathcal{U}^κ is again only possible for $a_1 = u = 0$. Thus $\mathcal{U}^\kappa \subset \mathcal{U}$ and $\partial \mathcal{U}^\kappa \cap \partial \mathcal{U} = \{0\}$.

For the second part it thus remains to look at $a \in \mathcal{U}^\kappa$, a close to 0. Writing again $a = (a_1, u + iv)$ and using that $v^2 \leq 2\kappa^2(a_1^2 + u^2)$ there holds

$$\begin{aligned} |a_1^2 + a_2^2| &= (a_1^4 + 2a_1^2(u^2 - v^2) + (u^2 - v^2)^2 + 4u^2v^2)^{\frac{1}{2}} \\ &\geq ((1 - 4\kappa^2)a_1^4 + 2a_1^2(1 - 2\kappa^2)u^2 + (u^2 + v^2)^2)^{\frac{1}{2}} \\ &\geq (1 - 4\kappa^2)(a_1^4 + |a_2|^4)^{\frac{1}{2}} \geq \frac{1 - 4\kappa^2}{2} |a|_*^2. \end{aligned}$$

Then for $a \in \mathcal{U}^\kappa$, a small it follows that

$$|K_2(a)| \leq \frac{1}{2\pi} \frac{|a_1| + O(|a_1|^3)}{|a_1^2 + a_2^2| - O(|a_1|^4) - O(|a_2|^4)} \leq \frac{1}{\pi} \frac{|a|_* + O(|a|_*^3)}{(1 - 4\kappa^2)|a|_*^2 - O(|a|_*^4)} \leq \frac{C}{|a|_*}.$$

Higher order derivatives follow in the same way. \square

Lemma 6.6. *Let $\rho_0 > 0$ be chosen such that γ_0 extends holomorphically to $U_{2\rho_0}$ with*

$$(6.8) \quad 4 \|\operatorname{Im}(\gamma'_0)\|_{L^\infty(U_{\rho_0})} < 1.$$

Then the complex extension of the initial normal velocity $s_0 : \Omega_0 \rightarrow \mathbb{R}$,

$$s_0(y_1) := -2 \int_{\mathbb{T}} K_2(\Delta X_0(y_1, z_1)) \Delta \gamma'_0(y_1, z_1) dz_1$$

is holomorphic on U_{ρ_0} . Moreover, the $L^\infty(U_{\rho_0})$ -norm of any finite amount of derivatives of s_0 can be bounded by C_0 . In particular, all derivatives of s_0 are given by differentiation under the integral.

Proof. By (6.8) one estimates

$$|\operatorname{Im}(\gamma_0(y_1) - \gamma_0(y_1 - z_1))| < \frac{1}{4} |z_1|$$

for $z_1 \in [-\pi, \pi)$, $z_1 \neq 0$, $y_1 \in U_{\rho_0}$. Thus by Lemma 6.5 the composition of K_2 with $y_1 \mapsto \Delta X_0(y_1, z_1)$ is holomorphic for every $z_1 \neq 0$. Moreover, again by Lemma 6.5 for such z_1 there holds

$$|\partial_{y_1}(K_2(\Delta X_0(y_1, z_1)))| \leq C \left(\frac{|\Delta \gamma'_0(y_1, z_1)|^2}{|\Delta X_0(y_1, z_1)|_*^2} + \frac{|\Delta \gamma''_0(y_1, z_1)|}{|\Delta X_0(y_1, z_1)|_*} \right) \leq C_0.$$

It follows that s_0 is holomorphic and that $\|s'_0\|_{L^\infty(U_{\rho_0})} \leq C_0$. The same can be shown for higher order derivatives. \square

The following two lemmas provide careful estimates needed for the compensation of various terms appearing in the definition of our nonlinear map F below. We are also careful with the uniform integrability as we need to be able to neglect what happens in some small sets.

Lemma 6.7. *Let $\rho_0 > 0$ be as in Lemma 6.6 and let $R > 0$. There exists $T = T(R, C_0, \alpha) \in (0, 1)$ such that for all $\eta \in B_\rho$, $\|\eta\|_\rho < R$, $\rho \in (0, \rho_0)$ and $t \in [0, T)$, $y \in \Omega_\rho$, $z \in \Omega_0$ there holds $\Delta X_t^\eta(y, z) \in \overline{\mathcal{U}^{3/8}}$ and*

$$(6.9) \quad t|y_2 - z_2| \leq C_0 |\Delta X_t^\eta(y, z)|_*.$$

Proof. First of all chose $T \in (0, 1)$ with $T^\alpha R \leq 1$. Then, omiting the (y, z) dependence in notation, cf. Sec. 6.2, there holds

$$\begin{aligned} \frac{t}{2}|y_2 - z_2| &\leq t|y_2 - z_2| - \frac{1}{2}t^{1+\alpha}R|y_2 - z_2| \\ &\leq \left| t(y_2 - z_2) + \frac{1}{2}t^{1+\alpha} \operatorname{Re}(\eta(y) - \eta(y_1, z_2)) \right| \\ &= \left| \operatorname{Re} \left(\Delta X_{t,2}^\eta - \Delta \gamma_0 - t\Delta s_0 - \frac{1}{2}t^{1+\alpha}(\eta(y_1, z_2) - \eta(y_1 - z_1, z_2)) \right) \right| \\ &\leq |\operatorname{Re}(\Delta X_{t,2}^\eta)| + (C_0(1+T) + T^{1+\alpha}R)|z_1| \leq |\operatorname{Re}(\Delta X_{t,2}^\eta)| + C_0|z_1| \end{aligned}$$

This implies (6.9).

In order to see that $\Delta X_t^\eta \in \overline{\mathcal{U}^{3/8}}$ we use (6.8), as well as the just shown inequality to deduce

$$\begin{aligned} |\operatorname{Im}(\Delta X_{t,2}^\eta)| &= \left| \operatorname{Im} \left(\Delta \gamma_0 + t\Delta s_0 + \frac{1}{2}t^{1+\alpha} \Delta \eta \right) \right| \\ &\leq \left(\frac{1}{4} + TC_0 + T^{1+\alpha}R \right) |z_1| + T^\alpha R t |y_2 - z_2| \\ &\leq \left(\frac{1}{4} + T(C_0 + 1) \right) |z_1| + T^\alpha R |\operatorname{Re}(\Delta X_{t,2}^\eta)| + T^\alpha R C_0 |z_1|. \end{aligned}$$

Thus by chosing $T > 0$ even smaller the desired inequality

$$|\operatorname{Im}(\Delta X_{t,2}^\eta)| \leq \frac{3}{8} (|z_1| + |\Delta \operatorname{Re}(X_{t,2}^\eta)|)$$

holds true. □

Lemma 6.8. *Let $\rho_0, R, T > 0$ be as in Lemma 6.7. For $\eta \in B_\rho$, $\|\eta\|_\rho < R$, $\rho \in (0, \rho_0)$ and $y \in \Omega_\rho$, $t \in (0, T)$ there holds*

$$(6.10) \quad \int_{\Omega_0} \frac{1}{|\Delta X_t^\eta(y, z)|_*} dz \leq C_0 |\log t|.$$

The integrability of $|\Delta X_t^\eta(y, \cdot)|_^{-1}$ is uniform with respect to $y \in \Omega_0$ and with respect to t considered on any interval of the form $[t_0, T)$ with $t_0 > 0$.*

Proof. In view of (6.9) we have

$$\begin{aligned} \int_{\Omega_0} \frac{1}{|\Delta X_t^\eta|_*} dz &\leq C_0 \int_{\Omega_0} \frac{1}{|z_1| + t|y_2 - z_2|} dz = C_0 \int_{\mathbb{T}} \int_{y_2-2}^{y_2+2} \frac{1}{|z_1| + t|z_2|} dz_2 dz_1 \\ &\leq C_0 \int_0^\pi \int_0^4 \frac{1}{z_1 + tz_2} dz_2 dz_1 = C_0 \left(\frac{\pi}{t} \log \left(1 + \frac{4t}{\pi} \right) + 4 \log \left(\frac{\pi}{4t} + 1 \right) \right), \end{aligned}$$

which is of order $|\log t|$.

The uniform integrability follows from $\frac{1}{|z_1+t_0z_2|} \in L^1(\mathbb{T} \times (-4, 4))$. \square

6.4 Definition of F

Let us fix $\rho_0 > 0$ as in Lemma 6.6. Take $R = 1$ and a corresponding $T \in (0, 1)$ from Lemma 6.7.

We define the application $(\eta, t) \mapsto F(\eta, t) = F_t(\eta)$ by setting

$$F_t(\eta)(y) := - \int_{\Omega_0} K_2(\Delta X_t^\eta(y, z)) \Delta \partial_{y_1} f_t^\eta(y, z) - K_2(\Delta X_0(y_1, z_1)) \Delta \gamma'_0(y_1, z_1) dz$$

for $t > 0$ and $F_0(\eta)(y) = 0$.

Lemma 6.9. *F when seen as a map*

$$\left\{ \eta \in B_\rho : \|\eta\|_\rho < 1 \right\} \times [0, T) \rightarrow B_{\rho'}$$

is well-defined for all $0 < \rho' < \rho < \rho_0$. Moreover, for $\eta \in B_\rho$, $\|\eta\|_\rho < 1$ the map $[0, T) \rightarrow B_{\rho'}$, $t \mapsto F_t(\eta)$ is continuous.

Proof. Let $\eta \in B_\rho$, $\|\eta\|_\rho < 1$ and $t \in (0, T)$. In view of Lemma 6.6 it remains to look at

$$\tilde{F}_t(\eta)(y) := F_t(\eta)(y) + 2s_0(y_1) = - \int_{\Omega_0} K_2(\Delta X_t^\eta(y, z)) \Delta \partial_{y_1} f_t^\eta(y, z) dz.$$

For $y \in \Omega_\rho$ and $z \in \Omega_0$ with $z_1 \neq 0$ one computes

$$(6.11) \quad \partial_{y_1} (K_2(\Delta X_t^\eta) \Delta \partial_{y_1} f_t^\eta) = \partial_{a_2} K_2(\Delta X_t^\eta) (\Delta \partial_{y_1} f_t^\eta)^2 + K_2(\Delta X_t^\eta) \Delta \partial_{y_1}^2 f_t^\eta,$$

$$(6.12) \quad \begin{aligned} \partial_{y_2} (K_2(\Delta X_t^\eta) \Delta \partial_{y_1} f_t^\eta) &= \partial_{a_2} K_2(\Delta X_t^\eta) \left(t + \frac{1}{2} t^{1+\alpha} \partial_{y_2} \eta(y) \right) \Delta \partial_{y_1} f_t^\eta \\ &\quad + K_2(\Delta X_t^\eta) \frac{1}{2} t^{1+\alpha} \partial_{y_2} \partial_{y_1} \eta(y), \end{aligned}$$

where, as usual, we have omitted the (y, z) dependence in the Δ -notation.

In order to get uniform integrability we use (6.9) to estimate

$$(6.13) \quad \begin{aligned} |\Delta \partial_{y_1}^j f_t^\eta| &\leq C_0 |z_1| + \|\partial_{y_1}^{j+1} \eta\|_{L^\infty(\Omega_{\rho'})} |z_1| + \|\partial_{y_2} \partial_{y_1}^j \eta\|_{L^\infty(\Omega_{\rho'})} t |y_2 - z_2| \\ &\leq C_0 \left(1 + \|\partial_{y_1}^j \nabla_y \eta\|_{L^\infty(\Omega_{\rho'})} \right) |\Delta X_t^\eta|_* \end{aligned}$$

for $y \in \Omega_{\rho'}$, $j = 0, 1, 2$. Now (6.13) and Lemmas 6.5, 6.7 imply

$$(6.14) \quad |K_2(\Delta X_t^\eta) \Delta \partial_{y_1} f_t^\eta| \leq C_0 \left(1 + \|\partial_{y_1} \nabla_y \eta\|_{L^\infty(\Omega_{\rho'})} \right).$$

As a consequence \tilde{F}_t , and thus F_t , maps at least into $L^\infty(\Omega_{\rho'})$. Moreover, combining similarly (6.7), (6.13) for $j = 1, 2$, and (6.9) to estimate (6.11), (6.12) one sees that

$$(6.15) \quad |\partial_{y_1} (K_2(\Delta X_t^\eta) \Delta \partial_{y_1} f_t^\eta)| \leq C_0 \left(1 + \|\partial_{y_1} \nabla_y \eta\|_{L^\infty(\Omega_{\rho'})}^2 + \|\partial_{y_1}^2 \nabla_y \eta\|_{L^\infty(\Omega_{\rho'})} \right),$$

and, recall $t < 1$, $|\partial_{y_2} \eta(y)| < 1$, that

$$(6.16) \quad |\partial_{y_2} (K_2(\Delta X_t^\eta) \Delta \partial_{y_1} f_t^\eta)| \leq C_0 \frac{t}{|\Delta X_t^\eta|_*} \left(1 + \|\partial_{y_1} \nabla_y \eta\|_{L^\infty(\Omega_{\rho'})} \right).$$

It follows that the complex derivative $\partial_{y_1} F_t(\eta)$ exists and is bounded on $\Omega_{\rho'}$. Moreover, in view of Lemma 6.8 the same is true for the (real) derivative $\partial_{y_2} F_t(\eta)$.

Next we turn to the required uniform continuity of $\partial_{y_2} F_t(\eta)$ on $\Omega_{\rho'}$. First of all observe that the corresponding integrant (6.12) as a function of $(z, y) \in \Omega_0 \times \Omega_{\rho'}$ is uniformly continuous on subsets which have their z_1 -component bounded away from 0. Here one uses the Cauchy integral formula and the assumed uniform continuity of $\partial_{y_2} \eta$ on the larger set Ω_ρ in order to conclude the uniform continuity of $\partial_{y_2} \partial_{y_1} \eta(y) = \partial_{y_1} \partial_{y_2} \eta(y)$. This together with the uniform integrability of the majorant given in (6.16) via Lemma 6.8 implies that $\partial_{y_2} F_t(\eta)$ is uniformly continuous on $\Omega_{\rho'}$, cf. also the argument below for continuity in time.

Moreover, in a similar way as above for (6.14), (6.15), (6.16) one can check that for any $y \in \Omega_{\rho'}$, $z_1 \neq 0$ there holds

$$|\partial_{y_1} \partial_{y_2} (K_2(\Delta X_t^\eta) \Delta \partial_{y_1} f_t^\eta)| \leq \frac{C_0 t}{|\Delta X_t^\eta|_*} \left(1 + \|\partial_{y_1} \nabla_y \eta\|_{L^\infty(\Omega_{\rho'})}^2 + \|\partial_{y_1}^2 \nabla_y \eta\|_{L^\infty(\Omega_{\rho'})} \right),$$

which implies that also $\partial_{y_2} F_t(\eta)$ is complex differentiable in y_1 .

In order to conclude $F_t(\eta) \in B_{\rho'}$ it therefore only remains to observe that $F_t(y) \in \mathbb{R}$ for $y \in \Omega_0$.

It remains to prove the continuity of $[0, T) \ni t \mapsto F_t(\eta) \in B_{\rho'}$. Let $t, t_0 \in (0, T)$ and take $\delta > 0$ sufficiently small. For $z \in \Omega_0$ with $|z_1| > \delta$, as well as $y \in \Omega_{\rho'}$ there holds

$$|K_2(\Delta X_t^\eta) \Delta \partial_{y_1} f_t^\eta - K_2(\Delta X_{t_0}^\eta) \Delta \partial_{y_1} f_{t_0}^\eta| \leq \frac{C_0}{\delta^2} |t - t_0|$$

due to Lemmas 6.5, 6.7. On the set $\{z \in \Omega_0 : |z_1| < \delta\}$ one uses the uniform majorant given in (6.14) to conclude the continuity of $(0, T) \ni t \mapsto F_t(\eta)$ with respect to $\|\cdot\|_{L^\infty(\Omega_{\rho'})}$.

For the corresponding continuity of $\partial_{y_1} F_t(\eta)$, $\partial_{y_2} F_t(\eta)$ with respect to $\|\cdot\|_{L^\infty(\Omega_{\rho'})}$ one uses a similar combination of Lipschitz continuity on $|z_1| > \delta$ and uniform integrability on the strip $|z_1| < \delta$ induced by (6.15), (6.16) and Lemma 6.8.

Finally, continuity at $t_0 = 0$ can be shown in the exact same way by noting that compared to Lemma 6.8 the additional factor t in (6.16) for $\partial_{y_2} F_t(\eta)$ renders $t |\Delta X_t^\eta|_*^{-1}$ to be uniformly integrable with respect to t taken from the open interval $t \in (0, T)$. \square

Remark 6.10. The continuity of F as stated in assumption (i) of Theorem 6.1 will follow from Lemma 6.9 when combined with the Lipschitz property of Lemma 6.11 below.

6.5 Contraction property

Next we will verify assumption (ii) of Theorem 6.1 with $b(t) = C_0 t^{1+\alpha} |\log t|$. Let $\rho_0, R, T > 0$ be as in Section 6.4. Recall that $R = 1$ and $T = T(R, C_0, \alpha) < 1$. Without loss of generality we also assume $\rho_0 < 1$.

Lemma 6.11. *For all $0 < \rho' < \rho < \rho_0$, $\eta, \zeta \in B_\rho$, $\|\eta\|_\rho < 1$, $\|\zeta\|_\rho < 1$ and $t \in [0, T)$ there holds*

$$\|F_t(\eta) - F_t(\zeta)\|_{\rho'} \leq \frac{C_0 t^{1+\alpha} |\log t|}{\rho - \rho'} \|\eta - \zeta\|_\rho.$$

For the proof of Lemma 6.11 we first of all state some estimates implied by the Lemmas in Section 6.3.

Lemma 6.12. *Let $0 < \rho' < \rho < \rho_0$ and $\eta, \xi, \zeta \in B_\rho$ with $\|\eta\|_\rho, \|\xi\|_\rho, \|\zeta\|_\rho < 1$. For $y \in \Omega_{\rho'}$, $z \in \Omega_0$, $t \in [0, T)$ there holds*

$$(6.17) \quad t |\Delta \zeta(y, z)| \leq C_0 \|\zeta\|_{\rho'} |\Delta X_t^\xi(y, z)|_*,$$

$$(6.18) \quad t |\Delta \partial_{y_1} \zeta(y, z)| \leq \frac{C_0}{\rho - \rho'} \|\zeta\|_\rho |\Delta X_t^\xi(y, z)|_*,$$

$$(6.19) \quad |\Delta \partial_{y_1} f_t^\eta(y, z)| \leq \frac{C_0}{\rho - \rho'} |\Delta X_t^\xi(y, z)|_*,$$

$$(6.20) \quad |\Delta \zeta(y, z) \Delta \partial_{y_1} f_t^\eta(y, z)| \leq C_0 \|\zeta\|_{\rho'} |\Delta X_t^\xi(y, z)|_*,$$

$$(6.21) \quad |\Delta \zeta(y, z) \Delta \partial_{y_1}^2 f_t^\eta(y, z)| \leq \frac{C_0}{\rho - \rho'} \|\zeta\|_{\rho'} |\Delta X_t^\xi(y, z)|_*.$$

Proof. By (6.9) in Lemma 6.7 one deduces

$$t |\Delta \zeta| \leq \|\nabla_y \zeta\|_{L^\infty(\Omega_{\rho'})} (|z_1| + t |y_2 - z_2|) \leq C_0 \|\zeta\|_{\rho'} |\Delta X_t^\xi(y, z)|_*.$$

This shows (6.17). Inequality (6.18) is obtained in the same way by additionally applying Cauchy's Lemma 6.3.

Next, (6.19) follows from

$$|\Delta \partial_{y_1} f_t^\eta| \leq C_0 |z_1| + t^{1+\alpha} |\Delta \partial_{y_1} \eta|$$

and (6.18), while (6.20) is a consequence of

$$|\Delta\zeta\Delta\partial_{y_1}f_t^\eta| \leq C_0 \|\zeta\|_{L^\infty(\Omega_{\rho'})} |z_1| + t^{1+\alpha} |\Delta\zeta| \|\partial_{y_1}\eta\|_{L^\infty(\Omega_{\rho'})}$$

and (6.17).

Finally, (6.21) is achieved in the same way as (6.20), but with an additional use of Lemma 6.3. \square

Proof of Lemma 6.11. Let $0 < \rho' < \rho < \rho_0$ and $\eta, \zeta \in B_\rho$ be as stated. For $\lambda \in [0, 1]$ define

$$\xi_\lambda := \lambda\eta + (1 - \lambda)\zeta.$$

Then $\xi_\lambda \in B_\rho$ and $\|\xi_\lambda\|_\rho < 1$.

Now for $y \in \Omega_{\rho'}$ we split

$$(6.22) \quad \begin{aligned} |F_t(\eta)(y) - F_t(\zeta)(y)| &\leq \int_{\Omega_0} \left| \left(K_2(\Delta X_t^\eta) - K_2(\Delta X_t^\zeta) \right) \Delta\partial_{y_1}f_t^\eta \right| dz \\ &\quad + \int_{\Omega_0} \left| K_2(\Delta X_t^\zeta) \left(\Delta\partial_{y_1}f_t^\eta - \Delta\partial_{y_1}f_t^\zeta \right) \right| dz. \end{aligned}$$

In order to estimate the first term we first use the fundamental theorem of Calculus to write

$$\begin{aligned} &\int_{\Omega_0} |K_2(\Delta X_t^\eta) - K_2(\Delta X_t^\zeta)| |\Delta\partial_{y_1}f_t^\eta| dz \\ &= \int_{\Omega_0} \left| \int_0^1 \partial_{a_2} K_2(\Delta X_t^{\xi_\lambda}) \frac{1}{2} t^{1+\alpha} (\Delta\eta - \Delta\zeta) d\lambda \right| |\Delta\partial_{y_1}f_t^\eta| dz. \end{aligned}$$

Now, $|\Delta\partial_{y_1}f_t^\eta|$ is dealt with by (6.19) with $\xi = \xi_\lambda$, Lemma 6.5 and its equation (6.7) deal with $\partial_{a_2} K_2(\Delta X_t^{\xi_\lambda})$ and by definition of $\|\cdot\|_{\rho'}$ we arrive to the estimate

$$\begin{aligned} &\int_{\Omega_0} |K_2(\Delta X_t^\eta) - K_2(\Delta X_t^\zeta)| |\Delta\partial_{y_1}f_t^\eta| dz \\ &= \int_{\Omega_0} \left| \int_0^1 \partial_{a_2} K_2(\Delta X_t^{\xi_\lambda}) \frac{1}{2} t^{1+\alpha} (\Delta\eta - \Delta\zeta) d\lambda \right| |\Delta\partial_{y_1}f_t^\eta| dz \\ &\leq C_0 t^{1+\alpha} \|\eta - \zeta\|_{\rho'} \int_{\Omega_0} \int_0^1 \frac{1}{|\Delta X_t^{\xi_\lambda}|_*} \frac{|\Delta X_t^{\xi_\lambda}|_*}{\rho - \rho'} d\lambda dz \leq \frac{C_0 t^{1+\alpha} |\log t|}{\rho - \rho'} \|\eta - \zeta\|_\rho, \end{aligned}$$

where the last inequality is a direct application of Lemma 6.8. Again by Lemmas 6.5, 6.8 the second term in (6.22) is bounded by

$$\begin{aligned} \int_{\Omega_0} \left| K_2(\Delta X_t^\zeta) \left(\Delta\partial_{y_1}f_t^\eta - \Delta\partial_{y_1}f_t^\zeta \right) \right| dz &\leq C \int_{\Omega_0} \frac{1}{|\Delta X_t^\zeta|_*} t^{1+\alpha} |\Delta\partial_{y_1}\eta - \Delta\partial_{y_1}\zeta| dz \\ &\leq C_0 t^{1+\alpha} |\log t| \|\eta - \zeta\|_\rho. \end{aligned}$$

Thus

$$\|F_t(\eta) - F_t(\zeta)\|_{L^\infty(\Omega_{\rho'})} \leq \frac{C_0 t^{1+\alpha} |\log t|}{\rho - \rho'} \|\eta - \zeta\|_\rho.$$

Let us now turn to the corresponding inequality with ∂_{y_1} . In a similar way as before we split

$$\partial_{y_1} F_t(\eta)(y) - \partial_{y_1} F_t(\zeta)(y) = - \int_{\Omega_0} A_1 + A_2 + A_3 + A_4 dz,$$

where

$$\begin{aligned} A_1 &:= \left(\partial_{a_2} K_2(\Delta X_t^\eta) - \partial_{a_2} K_2(\Delta X_t^\zeta) \right) (\Delta \partial_{y_1} f_t^\eta)^2, \\ A_2 &:= \partial_{a_2} K_2(\Delta X_t^\zeta) \left((\Delta \partial_{y_1} f_t^\eta)^2 - (\Delta \partial_{y_1} f_t^\zeta)^2 \right), \\ A_3 &:= \left(K_2(\Delta X_t^\eta) - K_2(\Delta X_t^\zeta) \right) \Delta \partial_{y_1}^2 f_t^\eta, \\ A_4 &:= K_2(\Delta X_t^\zeta) \left(\Delta \partial_{y_1}^2 f_t^\eta - \Delta \partial_{y_1}^2 f_t^\zeta \right), \end{aligned}$$

cf. (6.11). Regarding A_1 we use (6.19), (6.20) to deduce

$$\begin{aligned} \int_{\Omega_0} |A_1| dz &\leq C \int_{\Omega_0} \int_0^1 \frac{1}{|\Delta X_t^{\xi_\lambda}|_*^3} t^{1+\alpha} |\Delta(\eta - \zeta) \Delta \partial_{y_1} f_t^\eta| |\Delta \partial_{y_1} f_t^\eta| dz d\lambda \\ &\leq \frac{C_0 t^{1+\alpha}}{\rho - \rho'} \|\eta - \zeta\|_{\rho'} \int_{\Omega_0} \int_0^1 \frac{1}{|\Delta X_t^{\xi_\lambda}|_*} dz d\lambda \leq \frac{C_0 t^{1+\alpha} |\log t|}{\rho - \rho'} \|\eta - \zeta\|_\rho. \end{aligned}$$

By making use of (6.21) instead of (6.20) one can bound $\int_{\Omega_0} |A_3| dz$ in a similar way. We omit the details.

Next for A_2 , inequality (6.19) implies

$$\begin{aligned} \int_{\Omega_0} |A_2| dz &\leq C \int_{\Omega_0} \frac{1}{|\Delta X_t^\zeta|_*^2} \left| \Delta \partial_{y_1} f_t^\eta + \Delta \partial_{y_1} f_t^\zeta \right| t^{1+\alpha} |\Delta \partial_{y_1} \eta - \Delta \partial_{y_1} \zeta| dz \\ &\leq \frac{C_0 t^{1+\alpha} |\log t|}{\rho - \rho'} \|\eta - \zeta\|_\rho. \end{aligned}$$

Finally, the estimate for $\int_{\Omega_0} |A_4| dz$ is a straightforward consequence of Lemmas 6.5, 6.7, 6.8 and the Cauchy Lemma 6.3.

Summarizing, we have shown

$$\|\partial_{y_1} F_t(\eta) - \partial_{y_1} F_t(\zeta)\|_{L^\infty(\Omega_{\rho'})} \leq \frac{C_0 t^{1+\alpha} |\log t|}{\rho - \rho'} \|\eta - \zeta\|_\rho.$$

It therefore remains to check ∂_{y_2} . Again we write

$$\partial_{y_2} F_t(\eta)(y) - \partial_{y_2} F_t(\zeta)(y) = - \int_{\Omega_0} B_1 + B_2 + B_3 + B_4 dz,$$

where, cf. (6.12),

$$\begin{aligned}
B_1 &:= \left(\partial_{a_2} K_2(\Delta X_t^\eta) - \partial_{a_2} K_2(\Delta X_t^\zeta) \right) \left(t + \frac{1}{2} t^{1+\alpha} \partial_{y_2} \eta(y) \right) \Delta \partial_{y_1} f_t^\eta, \\
B_2 &:= \partial_{a_2} K_2(\Delta X_t^\zeta) \left[\left(t + \frac{1}{2} t^{1+\alpha} \partial_{y_2} \eta(y) \right) \Delta \partial_{y_1} f_t^\eta - \left(t + \frac{1}{2} t^{1+\alpha} \partial_{y_2} \zeta(y) \right) \Delta \partial_{y_1} f_t^\zeta \right], \\
B_3 &:= \left(K_2(\Delta X_t^\eta) - K_2(\Delta X_t^\zeta) \right) \frac{1}{2} t^{1+\alpha} \partial_{y_2} \partial_{y_1} \eta(y), \\
B_4 &:= K_2(\Delta X_t^\zeta) \frac{1}{2} t^{1+\alpha} (\partial_{y_2} \partial_{y_1} \eta(y) - \partial_{y_2} \partial_{y_1} \zeta(y)).
\end{aligned}$$

Since $t < 1$ and $|\partial_{y_2} \eta(y)| < 1$ we get

$$\begin{aligned}
\int_{\Omega_0} |B_1| dz &\leq C t^{1+\alpha} \int_0^1 \int_{\Omega_0} \frac{1}{|\Delta X_t^{\xi_\lambda}|_*^3} t |\Delta \eta - \Delta \zeta| |\Delta \partial_{y_1} f_t^\eta| dz d\lambda \\
&\leq \frac{C_0 t^{1+\alpha} |\log t|}{\rho - \rho'} \|\eta - \zeta\|_\rho
\end{aligned}$$

by (6.17), (6.19).

Moreover,

$$\begin{aligned}
\int_{\Omega_0} |B_2| dz &\leq C \int_{\Omega_0} \frac{1}{|\Delta X_t^\zeta|_*^2} \left[t^{1+\alpha} |\partial_{y_2} \eta(y) - \partial_{y_2} \zeta(y)| |\Delta \partial_{y_1} f_t^\eta| \right. \\
&\quad \left. + (t + t^{1+\alpha} |\partial_{y_2} \zeta(y)|) t^{1+\alpha} |\Delta \partial_{y_1} \eta - \Delta \partial_{y_1} \zeta| \right] dz \\
&\leq \frac{C_0 t^{1+\alpha} |\log t|}{\rho - \rho'} \|\eta - \zeta\|_\rho
\end{aligned}$$

by use of (6.19) in the first term, as well as (6.18) in the second.

The estimate for $\int_{\Omega_0} |B_3| dz$ follows in analogy to $\int_{\Omega_0} |B_1| dz$ by utilizing (6.17) and Cauchy's Lemma 6.3, whereas the estimate for $\int_{\Omega_0} |B_4| dz$ relies solely on Lemma 6.3.

This finishes the proof of Lemma 6.11. \square

6.6 The affine term

In order to complete the list of ingredients of Theorem 6.1, we investigate $F_t(0)$. As usual we consider $\rho_0 \in (0, 1)$ to be fixed according to Lemma 6.6, as well as $R = 1$ and $T = T(R, C_0, \alpha) \in (0, 1)$ given by Lemma 6.7.

Lemma 6.13. *For any $\rho \in (0, \rho_0)$, $t \in (0, T)$ there holds*

$$\|F_t(0)\|_\rho \leq C_0 t |\log t|.$$

Proof. Let $y \in \Omega_\rho$. Recall that

$$\Delta X_t^0 = \begin{pmatrix} z_1 \\ \Delta\gamma_0 + t\Delta s_0 + t(y_2 - z_2) \end{pmatrix} = \Delta X_0 + \begin{pmatrix} 0 \\ t\Delta s_0 + t(y_2 - z_2) \end{pmatrix},$$

$z \in \Omega_0$. In view of Lemmas 6.5, 6.7, 6.8 and the boundedness of Ω_0 there holds

$$\begin{aligned} |F_t(0)(y)| &\leq \int_{\Omega_0} |K_2(\Delta X_t^0) - K_2(\Delta X_0)| |\Delta\gamma'_0| + t |K_2(\Delta X_t^0)| |\Delta s'_0| dz \\ &\leq \int_0^1 \int_{\Omega_0} |\partial_{a_2} K_2(\Delta X_{\lambda t}^0) t(y_2 - z_2 + \Delta s_0)| |\Delta\gamma'_0| dz d\lambda + C_0 t \\ &\leq C_0 t \left(1 + \int_0^1 \int_{\Omega_0} \frac{1}{|\Delta X_{\lambda t}^0|_*} dz d\lambda \right) \leq C_0 t \left(1 + \int_0^1 |\log(\lambda t)| d\lambda \right) \\ &\leq C_0 t |\log t|. \end{aligned}$$

Next we estimate $\partial_{y_1} F_t(0)$. One has

$$\begin{aligned} |\partial_{y_1} F_t(0)(y)| &\leq \int_{\Omega_0} |\partial_{a_2} K_2(\Delta X_t^0) - \partial_{a_2} K_2(\Delta X_0)| |\Delta\gamma'_0|^2 + |K_2(\Delta X_t^0)| t |\Delta s''_0| \\ &\quad + |K_2(\Delta X_t^0) - K_2(\Delta X_0)| |\Delta\gamma''_0| + |\partial_{a_2} K_2(\Delta X_t^0)| (2t\Delta\gamma'_0\Delta s'_0 + t^2 |\Delta s'_0|^2) dz. \end{aligned}$$

The terms appearing on the right-hand side can be dealt with in a similar way as above.

Finally, we also state

$$|\partial_{y_2} F_t(0)(y)| \leq \int_{\Omega_0} |\partial_{a_2} K_2(\Delta X_t^0)| t |\Delta\gamma'_0 + t\Delta s'_0| dz \leq C_0 t |\log t|.$$

This finishes the proof of Lemma 6.13. \square

Remark 6.14. Note that Lemma 6.13 implies

$$\frac{1}{t^{1+\alpha}} \int_0^t \|F_s(0)\|_\rho ds \leq C_0 t^{1-\alpha} |\log t| \leq \frac{C_0}{\rho_0 - \rho} t^{1-\alpha} |\log t|,$$

i.e. assumption (iii) of Theorem 6.1 holds with $a(t) = t^{1+\alpha}$, $c(t) = C_0 t^{1-\alpha} |\log t|$.

6.7 Conclusion and additional remarks

In Sections 6.1-6.6 we have verified all conditions of Theorem 6.1. As a consequence we deduce the following statement.

Proposition 6.15. *Let $\rho_0 > 0$ be as in Lemma 6.6. There exists $\bar{a} = \bar{a}(C_0) > 0$, $T = T(C_0, \alpha) > 0$ and a unique function $t \mapsto \eta_t$ with the properties that for every $\rho \in (0, \rho_0)$ the map*

$$I_\rho := \{ t \in [0, T) : C_0 t^{1-\alpha} |\log t| < \bar{a}(\rho_0 - \rho) \} \ni t \mapsto \eta_t \in B_\rho$$

is continuous with $\|\eta_t\|_\rho < 1$, $t \in I_\rho$, and such that for all $y \in \Omega_\rho$, $t \in I_\rho$ there holds

$$(6.23) \quad \eta_0(y) = 0, \quad \eta_t(y) = \frac{1}{t^{1+\alpha}} \int_0^t F_s(\eta_s)(y) ds.$$

We finish the investigation of the fixed-point problem (5.12) with some accompanying remarks concerning properties of the solution η_t given by Proposition 6.15.

The first addresses regularity. In contrast to the analyticity of η_t in y_1 , we only know that η_t is continuously differentiable in y_2 . Using (6.23) it seems possible to upgrade the regularity with respect to y_2 . However, since $F_s(\eta_s)(y)$ involves the integration over the finite interval $(-2, 2)$ with respect to z_2 , in contrast to \mathbb{T} for the integration in z_1 , the maximal regularity for $\eta_t(y_1, \cdot) : [-2, 2] \rightarrow \mathbb{C}$ is expected to be finite. In any case, since a higher regularity of η_t with respect to y_2 would only improve the regularity of our subsolution inside the mixing zone and not across its boundary, we have not pursued this topic any further.

Next we turn to the role of the parameter α . Suppose that we set up problem (5.12) with respect to two different choices $0 < \alpha < \beta < 1$ leading to two different right-handsides involving $F_t^\alpha(\eta)$, $F_t^\beta(\eta)$. Our previous analysis gives two solutions η_t^α , η_t^β with corresponding intervals $I_\rho^\alpha \subset [0, T^\alpha)$, $I_\rho^\beta \subset [0, T^\beta)$, $\rho \in (0, \rho_0)$. Note that the intervals I_ρ^α , I_ρ^β are defined with the same \bar{a} and recall that $T^\alpha, T^\beta \in (0, 1)$.

Lemma 6.16. *There holds $t^{\beta-\alpha}\eta_t^\beta = \eta_t^\alpha$ on $[0, \min\{T^\alpha, T^\beta\})$.*

Proof. Define $J_\rho^\alpha := I_\rho^\alpha \cap [0, T^\beta)$, $J_\rho^\beta := I_\rho^\beta \cap [0, T^\alpha)$. Then $J_\rho^\beta \subset J_\rho^\alpha$ due to the fact that $\beta > \alpha$ and $t < 1$. Both functions $t^{\beta-\alpha}\eta_t^\beta$, η_t^α are continuous maps from J_ρ^β into the unit ball of B_ρ , $\rho \in (0, \rho_0)$, and they both vanish at $t = 0$. Moreover, it is easy to check that

$$t^{\beta-\alpha}\eta_t^\beta(y) = t^{\beta-\alpha} \frac{1}{t^{1+\beta}} \int_0^t F_s^\beta(\eta_s^\beta)(y) ds = \frac{1}{t^{1+\alpha}} \int_0^t F_s^\alpha(s^{\beta-\alpha}\eta_s^\beta)(y) ds.$$

Thus Proposition 6.15 implies $t^{\beta-\alpha}\eta_t^\beta = \eta_t^\alpha$ for as long as both are defined. \square

Both solutions $t^{\beta-\alpha}\eta_t^\beta$, η_t^α of (6.23) then extend uniquely to a common maximal solution of (6.23) enjoying the properties of Proposition 6.15. Moreover, Lemma 6.16 shows that $t^{1+\alpha}\eta_t^\alpha$ is independent of the considered $\alpha \in (0, 1)$, such that the induced function $f(t, y)$ defined in Section 7 below, is independent of $\alpha \in (0, 1)$.

Finally we remark that for the choice $\alpha = 1$ in ansatz (5.10) a more careful analysis would have been required. In that case the initial value $\eta_0(y)$ is not expected to be given by 0 and the estimate given in Lemma 6.13 does not even lead to boundedness of $t^{-2} \int_0^t \|F_s(0)\|_\rho ds$. However, since this analysis has not been needed in order to prove existence of a Lipschitz solution of (3.1), we leave the case $\alpha = 1$ as a possible future improvement.

7 Justification of ansatzes

We will now verify that η provided by Proposition 6.15 induces when undoing the transformations stated in Section 5 indeed a solution of the macroscopic IPM system (3.1).

Given η from Proposition 6.15 we first of all define $f : [0, T) \times \overline{\Omega}_0 \rightarrow \mathbb{R}$,

$$f(t, y) := f_t^{\eta_t}(y) = \gamma_0(y_1) + ts_0(y_1) + \frac{1}{2}t^{1+\alpha}\eta_t(y),$$

where $T = T(C_0, \alpha) > 0$ can be taken as the endpoint of the interval $I_{\rho_0/2}$ for instance. Also recall Lemma 6.4 if needed for the extension to the closure of Ω_0 .

Lemma 7.1. *There holds $f \in \mathcal{C}^1([0, T); \mathcal{C}^1(\overline{\Omega}_0))$ with*

$$(7.1) \quad \|\partial_{y_2} f(t, \cdot)\|_{L^\infty(\Omega_0)} \leq \frac{1}{2}t^{1+\alpha}, \quad t \in (0, T).$$

Moreover, the functions $f(t, \cdot, y_2)$, $\partial_t f(t, \cdot, y_2)$, $\partial_{y_2} f(t, \cdot, y_2)$, $t \in [0, T)$, $y_2 \in [-2, 2]$ are real analytic and f satisfies the initial value problem $f(0, y) = \gamma_0(y_1)$,

$$(7.2) \quad \partial_t f(t, y) = -\frac{1}{2} \int_{\Omega_0} K_2(\Delta X_t^{\eta_t}(y, z)) (\partial_{y_1} f(t, y) - \partial_{y_1} f(t, y_1 - z_1, z_2)) dz$$

for $t \in [0, T)$, $y \in \overline{\Omega}_0$.

Proof. As a direct consequence of Lemma 6.9 and Proposition 6.15 one sees that f belongs to $\mathcal{C}^1([0, T); B_{\rho_0/2})$ and satisfies (7.1), (7.2) for $t \in [0, T)$, $y \in \Omega_{\rho_0/2}$. The statement follows from the definition of the spaces B_ρ and Lemma 6.4. \square

We are now able to prove our main result.

Proof of Theorem 3.2. Let f be as in Lemma 7.1. Define the open space-time set

$$\mathcal{U} := \{ (t, x) \in (0, T) \times \mathbb{T} \times \mathbb{R} : -2t + f(t, x_1, -2) < x_2 < 2t + f(t, x_1, 2) \},$$

as well as the slices

$$\mathcal{U}_t := \{ x \in \mathbb{T} \times \mathbb{R} : (t, x) \in \mathcal{U} \}, \quad t \in (0, T).$$

As a consequence of (7.1) the maps $X_t : \Omega_0 \rightarrow \mathcal{U}_t$,

$$X_t(y) := \begin{pmatrix} y_1 \\ ty_2 + f(t, y) \end{pmatrix}, \quad t \in (0, T)$$

are \mathcal{C}^1 diffeomorphisms with the property that also the joint maps $(0, T) \times \Omega_0 \rightarrow \mathbb{T} \times \mathbb{R}$, $(t, y) \mapsto X_t(y)$ and $\mathcal{U} \rightarrow \mathbb{T} \times \mathbb{R}$, $(t, x) \mapsto X_t^{-1}(x)$ are of class \mathcal{C}^1 .

In view of (5.2) we thus can indeed define the density $\rho : [0, T) \times \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{R}$ by setting

$$\rho(t, x) := \begin{cases} 1, & x_2 \geq 2t + f(t, x_1, 2), \\ \frac{1}{2}(X_t^{-1}(x))_2, & x \in \mathcal{U}_t, \\ -1, & x_2 \leq -2t + f(t, x_1, -2) \end{cases}$$

for $t \in (0, T)$ and $\rho(0, x) := \rho_0(x)$. Here $(X_t^{-1}(x))_2$ denotes the second component of $X_t^{-1}(x)$. Observe that ρ is continuous except at points $(0, x_1, \gamma_0(x_1))$, $x_1 \in \mathbb{T}$, and piecewise \mathcal{C}^1 with the exceptional set being $\partial\mathcal{U} \subset [0, T) \times \mathbb{T} \times \mathbb{R}$. Moreover, as long as t is positive $\rho(t, \cdot)$ is Lipschitz continuous and there exists a constant $C_0 > 0$ depending on the initial data such that

$$(7.3) \quad |\nabla \rho(t, x)| \leq \frac{C_0}{t} \mathbb{1}_{\mathcal{U}_t}(x)$$

for all $(t, x) \notin \partial\mathcal{U}$.

Moreover, standard elliptic estimates show that v defined through (4.3), (4.7), (4.8) is the unique L^2 solution of the last two equations of (3.1).

The stated log-Lipschitz continuity of $v(t, \cdot)$, $t > 0$ is a consequence of the Biot-Savart operator acting on a compactly supported L^∞ -vorticity, cf. [71]. In addition it is easy to see that also $v : [0, T) \times \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{R}^2$ is continuous except at the one-dimensional set $\{(0, x_1, \gamma_0(x_1)) : x_1 \in \mathbb{T}\}$.

Hence we have shown properties (i), (ii) of Theorem 3.2. Moreover, observe that property (iii) holds by construction with γ_t given by

$$\gamma_t(x_1, h) := t2h + f(t, x_1, 2h), \quad x_1 \in \mathbb{T}, \quad h \in [-1, 1].$$

It thus remains to show that the first equation of (3.1) and the entropy balances (3.5) are satisfied.

The regularity of ρ implies

$$\int_{\mathbb{T} \times \mathbb{R}} \rho(t, \cdot) v(t, \cdot) \cdot \nabla \varphi \, dx = - \int_{\mathbb{T} \times \mathbb{R}} v(t, \cdot) \cdot \nabla \rho(t, \cdot) \varphi \, dx$$

for all $t \in (0, T)$, $\varphi \in \mathcal{C}^\infty(\mathbb{T} \times \mathbb{R})$. It follows that (ρ, v) is a solution in the sense of Definition 3.1 if and only if the transport form

$$(7.4) \quad \partial_t \rho + v \cdot \nabla \rho + 2\rho \partial_{x_2} \rho = 0$$

of the equation is satisfied pointwise in $(0, T) \times \mathbb{T} \times \mathbb{R} \setminus \partial\mathcal{U}$.

At points $(t, x) \notin \overline{\mathcal{U}}$ equation (7.4) trivially holds true. Inside \mathcal{U} one can check that the computations in Section 5.1 are possible showing that (7.4) is equivalent to (5.6). Note that in Section 5 we have formally assumed that X_t are global diffeomorphisms $\mathbb{T} \times \mathbb{R} \rightarrow \mathbb{T} \times \mathbb{R}$, but as the reader can easily see, it is enough to have transformations from Ω_0 to the corresponding \mathcal{U}_t .

Observing that also the computations in Section 5.2 are legal in our scenario one sees that (7.4) on \mathcal{U} is indeed equivalent to (7.2).

Finally, let $\eta : \mathbb{R} \rightarrow \mathbb{R}$ be an arbitrary Lipschitz continuous function and define $Q : \mathbb{R} \rightarrow \mathbb{R}$,

$$Q(u) := \int_0^u 2\eta'(s)s \, ds,$$

which is also Lipschitz continuous when restricted to any compact interval of \mathbb{R} . Consequently we have enough regularity to deduce (3.5) by multiplying (7.4) with $\eta'(\rho(t, x))$ and applying chain rule. This finishes the proof of Theorem 3.2. \square

A The abstract Cauchy-Kovalevskaya Theorem

Proof of Theorem 6.1. As indicated in Section 6 the proof of Theorem 6.1 is a slight modification of the original proof in [80].

Let $a_0 > 0$ and set $a_{k+1} := a_k(1 - (k+2)^{-2})$, $k = 0, 1, \dots$. Then

$$a := \lim_{k \rightarrow \infty} a_k > 0.$$

For $\rho \in (0, \rho_0)$ and $k = 0, 1, \dots$ we define the intervals

$$I_{k,\rho} := \{ t \in [0, T) : c(t) < a_k(\rho_0 - \rho) \}.$$

We also define for a function u with $u : I_{k,\rho} \rightarrow B_\rho$ continuous for any $\rho \in (0, \rho_0)$ the norm

$$M_k[u] := \sup \left\{ \|u(t)\|_\rho \left(\frac{a_k(\rho_0 - \rho)}{c(t)} - 1 \right) : \rho \in (0, \rho_0), t \in I_{k,\rho} \right\}.$$

Note that for $c(t) = t$ one recovers Nishida's set up. Now one recursively constructs the sequence

$$u_0(t) := 0, \quad u_{k+1}(t) := \begin{cases} \frac{1}{a(t)} \int_0^t F(u_k(s), s) \, ds, & t \in (0, T), \\ 0, & t = 0. \end{cases}$$

We claim that for a_0 chosen sufficiently small the recursion is well-defined, that each $u_k : I_{k,\rho} \rightarrow B_\rho$ is continuous with $\|u_k(t)\|_\rho < R/2$ for $t \in I_{k,\rho}$, $\rho \in (0, \rho_0)$, and that

$$(A.1) \quad \lambda_{k-1} := M_k[u_k - u_{k-1}] \leq (4Ka_0)^{k-1} a_0,$$

where $K > 0$ is the constant appearing in (6.2).

We first of all note that $u_1(t)$ exists and satisfies the stated continuity condition due to assumptions (i), (iii). Moreover, for $t \in I_{0,\rho}$ there holds $\|u_1(t)\|_\rho < a_0$. Thus we pick a_0 at least smaller than $R/2$. One also easily checks that $\lambda_0 \leq a_0$.

From now on we proceed by induction. Assume that the recursion with the above stated properties is possible up to some $k \geq 1$. Then it is clear that also $u_{k+1} : I_{k+1,\rho} \rightarrow B_\rho$ is well-defined, as well as continuous on the open interval $I_{k+1,\rho} \setminus \{0\}$ for any $\rho \in (0, \rho_0)$.

If we assume for now that (A.1) also holds true for λ_k , then for $t \in I_{k+1,\rho}$ we obtain in analogy to [80] the estimate

$$\begin{aligned} \|u_{k+1}(t)\|_\rho &\leq \sum_{j=0}^k \lambda_j \left(\frac{a_j(\rho_0 - \rho)}{c(t)} - 1 \right)^{-1} \leq \sum_{j=0}^k \lambda_j \left(\frac{a_j}{a_{j+1}} - 1 \right)^{-1} \\ &\leq a_0 \sum_{j=0}^k (4K a_0)^j (j+2)^2 < R/2 \end{aligned}$$

by choice of a_0 independent of k . Moreover, the first inequality in the above line of estimates applied at times $t > 0$ with $c(t) < \frac{a}{2}(\rho_0 - \rho)$ also gives

$$\|u_{k+1}(t)\|_\rho \leq c(t) \sum_{j=0}^k \frac{\lambda_j}{a_j(\rho_0 - \rho) - c(t)} \leq \frac{2a_0 c(t)}{a(\rho_0 - \rho)} \sum_{j=0}^k (4K a_0)^j,$$

which shows that u_{k+1} is also continuous with respect to $\|\cdot\|_\rho$ at $t = 0$.

To finish the induction it thus remains to show (A.1) for λ_k . The clever move is to use the contraction property with a different Banach space at each time τ inside the integral. Namely exactly as in [80, p. 631], the contraction property of F (Theorem 6.1 part (ii)) with $\rho(\tau) := \frac{1}{2} \left(\rho_0 - \frac{c(\tau)}{a_k} + \rho \right)$ and the definition of λ_{k-1} lead to

$$\|u_{k+1}(t) - u_k(t)\|_\rho \leq \frac{4\lambda_{k-1}a_k}{a(t)} \int_0^t \frac{b(\tau)c(\tau)}{(a_k(\rho_0 - \rho) - c(\tau))^2} d\tau$$

for $t \in I_{k,\rho}$. At this point we use (6.2) and change of variables to obtain

$$\|u_{k+1}(t) - u_k(t)\|_\rho \leq 4\lambda_{k-1}a_k K c(t) \int_0^{c(t)} \frac{1}{(a_k(\rho_0 - \rho) - \xi)^2} d\xi,$$

from where one can conclude $\lambda_k \leq 4K\lambda_{k-1}a_0$ by following [80] again.

Now Theorem 6.1 follows as in [79, 80]. \square

B More on Otto's relaxation

We here add some more details regarding the fifth step of Otto's relaxation [83] in the general non-flat case, which has only been sketched in Section 2.5.1.

Before doing that we will quickly convince ourselves that the setting in [83] is indeed equivalent to the formulation of IPM considered in our paper. Otto

considers the equations

$$(B.1) \quad \begin{aligned} \partial_t s + u \cdot \nabla s &= 0, \\ \nabla \cdot u &= 0, \\ u &= -\nabla p + s e_2, \end{aligned}$$

which correspond with [83]-(1.1)-(1.2) and the first equation on page 875 of [83] with $\lambda = 1$. The parameter λ in [83] is the quotient of the mobilities. In our case, we have taken both mobilities equal to one and then $\lambda = 1$. More importantly, in [83],

$$(B.2) \quad s = \{0, 1\},$$

however

$$(B.3) \quad \rho = \{-1, 1\}$$

in our case.

Let us see how we can go from (1.1),(B.3) to (B.1)-(B.2). Firstly we define

$$\bar{s} = \frac{1 - \rho}{2}, \quad \rho = 1 - 2\bar{s}$$

thus

$$\begin{aligned} \partial_t \bar{s} + v \cdot \nabla \bar{s} &= 0 \\ \nabla \cdot v &= 0 \\ v &= -\nabla(p + x_2) + 2\bar{s}e_2, \end{aligned}$$

with

$$\bar{s} = \{0, 1\}.$$

We define $\bar{u} = v/2$ and $\bar{\Pi} = (p + x_2)/2$ which yields

$$\begin{aligned} \partial_t \bar{s} + 2\bar{u} \cdot \nabla \bar{s} &= 0 \\ \nabla \cdot \bar{u} &= 0 \\ \bar{u} &= -\nabla \bar{\Pi} + \bar{s}e_2, \end{aligned}$$

Finally we take $s(x, t) = \bar{s}(x, t/2)$, $u(x, t/2) = \bar{u}(x, t/2)$ and $\Pi(x, t) = \bar{\Pi}(x, t/2)$ thus

$$\begin{aligned} \partial_t s + u \cdot \nabla s &= 0 \\ \nabla \cdot u &= 0 \\ u &= -\nabla \Pi + s e_2, \end{aligned}$$

with $s = \{0, 1\}$, which agrees with (B.1)-(B.2) (up to a relabeling of the pressure). Therefore, if we show that (B.1)-(B.2) relaxes to

$$(B.4) \quad \begin{aligned} \partial_t s + u \cdot \nabla s + \partial_{x_2} s - 2s \partial_{x_2} s &= 0, \\ \nabla \cdot u &= 0, \\ u &= -\nabla \Pi + s e_2, \end{aligned}$$

with $s \in [0, 1]$, by undoing the previous transformations, we see that (1.1),(B.3) relaxes to

$$(B.5) \quad \begin{aligned} \partial_t \rho + v \cdot \nabla \rho + 2\rho \partial_{x_2} \rho &= 0, \\ \nabla \cdot v &= 0, \\ v &= -\nabla p - \rho e_2, \end{aligned}$$

with $\rho \in [-1, 1]$.

Next, we begin our formal discussion with the outcome of the fourth step of Otto, after which there exists for each $h > 0$ a sequence of ‘‘coarse grained’’ functions $\{\theta^k\}_{k=0}^{N(h)}$ that are characterized by the following JKO scheme (which we understand as a minimizing movements scheme with respect to the Wasserstein distance):

$\theta^{(k+1)}$ is the minimizer in K of

$$(B.6) \quad \frac{1}{2} \text{dist}^2(\theta^{(k)}, \theta) + \frac{1}{2} \text{dist}^2(1 - \theta^{(k)}, 1 - \theta) - h \int \theta(x) x_2$$

where the set K consists of measurable θ taking values in $[0, 1]$ and such that $\int \theta = \int s(x, 0)$, and $\text{dist}^2(\theta_0, \theta_1)$ is the L^2 -Wasserstein distance

$$\text{dist}^2(\theta_0, \theta_1) = \inf_{\Phi \in I(\theta_0, \theta_1)} \int \theta_0(x) |\Phi(x) - x|^2 dx$$

with

$$I(\theta_0, \theta_1) = \left\{ \Phi : \int \theta_1(y) \zeta(y) dy = \int \theta_0(x) \zeta(\Phi(x)) dx \quad \forall \zeta \in \mathcal{C}_0^0 \right\}.$$

In the definition of $I(\theta_0, \theta_1)$, we have been deliberately imprecise and defer the reader to [83] for the proper definition. Even more, in order to make the exposition more clear in the following we will assume that the minimizer exists, that it is smooth and that it satisfies pointwise the corresponding Monge-Ampere equation. I.e.,

$$I(\theta_0, \theta_1) = \{ \Phi \text{ diffeomorphism} : (\theta_1 \circ \Phi)(x) J_\Phi(x) = \theta_0(x) \}.$$

Here J_Φ denotes the Jacobian determinant $\det D\Phi$.

As explained in Section 2.5.1 our goal is to show, on a formal level, that the limit as $h \rightarrow 0$, we will assume that it exists in the first place, of the functions

$$\theta_h(x, t) := \theta^{(k)}(x), \quad t \in [kh, (k+1)h)$$

is characterized by system (B.4).

We begin with the Euler-Lagrange equation of (B.6). For a given $\theta_0 \in K$ let θ_1 be the minimizer in K of

$$F[\theta] \equiv \frac{1}{2} \text{dist}^2(\theta_0, \theta) + \frac{1}{2} \text{dist}^2(1 - \theta_0, 1 - \theta) - h \int \theta(x) x_2.$$

Then we have that

$$D_\theta F[\theta_1] \psi = \left. \frac{d}{d\tau} F[\theta_1 + \tau \psi] \right|_{\tau=0} = 0,$$

where we simply assume that $\theta_1 + \tau \psi \in K$, i.e. we in particular consider ψ with $\int \psi = 0$. In order to compute $D_\theta F[\theta_1] \psi$ we first look at $D_\theta \text{dist}^2(\theta_0, \theta_1) \psi$. Let $\Phi_0^\tau \in I(\theta_0, \theta_1 + \tau \psi)$ be such that

$$\text{dist}^2(\theta_0, \theta_1 + \tau \psi) = \inf_{\Phi \in I(\theta_0, \theta_1 + \tau \psi)} \int \theta_0(x) |\Phi(x) - x|^2 dx = \int \theta_0(x) |\Phi_0^\tau(x) - x|^2 dx.$$

We define

$$w \circ \Phi_0^0 = \left. \frac{d\Phi_0^\tau}{d\tau} \right|_{\tau=0},$$

thus

$$(B.7) \quad \frac{1}{2} D_\theta \text{dist}^2(\theta_0, \theta_1) \psi = \int \theta_0(x) (\Phi_0^0(x) - x) \cdot (w \circ \Phi_0^0)(x) dx.$$

We next compute for which w we have $\Phi_0^\tau \in I(\theta_0, \theta_1 + \tau \psi)$. There holds

$$(B.8) \quad J_{\Phi_0^\tau}(x) ((\theta_1 + \tau \psi) \circ \Phi_0^\tau)(x) = \theta_0(x).$$

Taking a τ -derivative in (B.8) and evaluating at $\tau = 0$ yields,

$$J_{\Phi_0^0} \text{div } w \circ \Phi_0^0 \theta_1 \circ \Phi_0^0 + J_{\Phi_0^0} w \circ \Phi_0^0 \cdot \nabla \theta_1 \circ \Phi_0^0 + J_{\Phi_0^0} \psi \circ \Phi_0^0 = 0$$

which reduces to

$$(B.9) \quad \text{div } (w \theta_1) + \psi = 0.$$

In addition, Φ_0^0 minimizes

$$\int \theta_0(x) |\Phi(x) - x|^2 dx$$

in $I(\theta_0, \theta_1)$. So for every family of flows $(\Phi_\delta^0) \in I(\theta_0, \theta_1)$ we have that

$$\left. \frac{d}{d\delta} \int \theta_0(x) |\Phi_\delta^0(x) - x|^2 dx \right|_{\delta=0} = 0.$$

That is

$$\int \theta_0(x)(\Phi_0^0(x) - x) \cdot (\bar{w} \circ \Phi_0^0)(x) dx = 0,$$

where if Φ_δ is the flow of a vector field \bar{w} ,

$$(B.10) \quad \bar{w} \circ \Phi_0^0 = \left. \frac{d\Phi_\delta^0}{d\delta} \right|_{\delta=0},$$

$$(B.11) \quad \operatorname{div}(\theta_1 \bar{w}) = 0.$$

The condition (B.11), equivalent to $\Phi_\delta \in I(\theta_0, \theta_1)$, is deduced by differentiating

$$(B.12) \quad J_{\Phi_\delta^0}(x)(\theta_1 \circ \Phi_\delta^0)(x) = \theta_0(x)$$

with respect to δ .

Therefore, it holds that

$$0 = \int \theta_0(x)(\Phi_0^0(x) - x) \cdot (\bar{w} \circ \Phi_0^0)(x) dx = \int \theta_1(x) \bar{w}(x) \cdot (x - (\Phi_0^0)^{-1}(x)) dx,$$

where in the last equality we have used the definition of $I(\theta_0, \theta_1)$. Since \bar{w} is an arbitrary vector field, Hodge decomposition implies that

$$(B.13) \quad x - (\Phi_0^0)^{-1}(x) = \nabla a(x)$$

for some function a . In order to avoid technicalities we here have implicitly assumed that θ_1 does not vanish.

From (B.7), (B.9) and (B.13) we see that

$$\begin{aligned} \frac{1}{2} D_\theta \operatorname{dist}^2(\theta_0, \theta_1) \psi &= \int \theta_1(x) w(x) \cdot \nabla a(x) dx = - \int \nabla \cdot (\theta_1 w)(x) a(x) dx \\ &= \int \psi(x) a(x). \end{aligned}$$

We have obtained that

$$\begin{aligned} \frac{1}{2} D_\theta \operatorname{dist}^2(\theta_0, \theta_1) \psi &= \int \psi(x) a(x) \\ \Phi_0^0(x) &= x + (\nabla a \circ \Phi_0^0)(x). \end{aligned}$$

Similar computations yield

$$\begin{aligned} \frac{1}{2} D_\theta \operatorname{dist}^2(1 - \theta_0, 1 - \theta_1) \psi &= - \int \psi(x) \bar{a}(x) \\ \bar{\Phi}_0^0(x) &= x + (\nabla \bar{a} \circ \bar{\Phi}_0^0)(x), \end{aligned}$$

and putting everything together we arrive at

$$(B.14) \quad D_\theta F[\theta_1]\psi = \int (a(x) - \bar{a}(x) - hx_2)\psi(x)dx = 0$$

for all ψ with $\int \psi = 0$. Moreover, since $\Phi_0^0 \in I(\theta_0, \theta_1)$ and $\bar{\Phi}_0^0 \in I(1 - \theta_0, 1 - \theta_1)$,

$$(B.15) \quad \theta_1(x) = J_{(\Phi_0^0)^{-1}}(x)\theta_0(x - \nabla a(x)),$$

$$(B.16) \quad (1 - \theta_1)(x) = J_{(\bar{\Phi}_0^0)^{-1}}(x)(1 - \theta_0)(x - \nabla \bar{a}(x)).$$

Note that so far we have omitted the h -dependence of the functions $a, \bar{a}, \Phi_0^0, \bar{\Phi}_0^0$ in our notation. We continue doing so when introducing $p = \frac{a}{h}, \bar{p} = \frac{\bar{a}}{h}$ which, up to a constant satisfy

$$p - \bar{p} = x_2$$

by (B.14). Note that the constant is irrelevant since only derivatives of p, \bar{p} will play a role. To obtain a formal limit as $h \rightarrow 0$, we will assume in the following that the h -dependent functions p, \bar{p} have a well-defined \mathcal{C}^2 limit, which will again be denoted by p, \bar{p} .

Now we take said limit. On one hand we have from (B.15) that

$$\begin{aligned} \frac{\theta_1(x) - \theta_0(x)}{h} &= \frac{J_{(\Phi_0^0)^{-1}}(x)\theta_0(x - h\nabla p(x)) - \theta_0(x)}{h} \\ &= \frac{(J_{(\Phi_0^0)^{-1}}(x) - 1)\theta_0(x - h\nabla p(x)) + \theta_0(x - h\nabla p(x)) - \theta_0(x)}{h}. \end{aligned}$$

Recall that Φ_0^0 is linked to p via (B.13) and thus since $\Phi_0^0(x) \rightarrow x$, it holds that

$$J_{(\Phi_0^0)^{-1}}(x) - 1 = -h\Delta p(x),$$

at first order in h . Thus when letting $h \rightarrow 0$ in the difference quotient we arrive at

$$(B.17) \quad \partial_t \theta = -\Delta p \theta - \nabla p \cdot \nabla \theta.$$

On the other hand we have from (B.16) that

$$\frac{\theta_1(x) - \theta_0(x)}{h} = \frac{1 - J_{(\bar{\Phi}_0^0)^{-1}}(x) + J_{(\bar{\Phi}_0^0)^{-1}}(x)\theta_0(x - h\nabla \bar{p}(x)) - \theta_0(x)}{h}.$$

Passing to the limit yields

$$(B.18) \quad \partial_t \theta = \Delta \bar{p} - \Delta \bar{p} \theta - \nabla \theta \cdot \nabla \bar{p}.$$

In order for (B.17) and (B.18) to agree, there holds

$$\Delta \bar{p} - \nabla \cdot (\nabla \bar{p} \theta) = -\nabla \cdot (\nabla p \theta)$$

and since $p = \bar{p} + x_2$ we have that

$$(B.19) \quad \Delta \bar{p} = -\nabla \cdot (\nabla x_2 \theta) = -\partial_{x_2} \theta.$$

Therefore, from (B.18) and (B.19),

$$\begin{aligned} \partial_t \theta &= -\partial_{x_2} \theta - \nabla \cdot (\nabla \bar{p} \theta) \\ &= -\partial_{x_2} \theta - \nabla \cdot ((\nabla \bar{p} + \theta e_2) \theta) + \nabla \cdot (\theta^2 e_2) \end{aligned}$$

To finish we define $u = \nabla \bar{p} + \theta e_2$, which clearly satisfies $\nabla \cdot u = 0$, to get

$$\begin{aligned} \partial_t \theta + u \cdot \nabla \theta + \partial_{x_2} \theta - 2\theta \partial_{x_2} \theta &= 0, \\ u &= \nabla \bar{p} + \theta e_2, \\ \nabla \cdot u &= 0, \end{aligned}$$

which agrees with (B.4).

C Rigorous energy dissipation

In Section 2.5.2 equation (2.15) we have formally computed the decay rate of the total potential energy. For completeness we give sufficient conditions when this computation is justified. Also for completeness we show that the subsolution given by Theorem 3.2 indeed satisfies the sufficient conditions.

Lemma C.1. *Let $\rho_0 \in L^1_{loc}(\mathbb{T} \times \mathbb{R})$ be some initial data and suppose that the pair of functions $(\rho, m) \in L^1_{loc}((0, T) \times \mathbb{T} \times \mathbb{R}; \mathbb{R} \times \mathbb{R}^2)$ satisfies*

$$\partial_t \rho + \operatorname{div} m = 0, \quad \rho(0, \cdot) = \rho_0$$

on $(0, T) \times \mathbb{T} \times \mathbb{R}$ in the sense of distributions. If there exists $\alpha > 0$ such that

$$\begin{aligned} m_2, (\rho - \rho_0)x_2, (\rho - \rho_0)|x_2|^{1+\alpha} &\in \mathcal{C}^0([0, T]; L^1(\mathbb{T} \times \mathbb{R})), \\ m_2 |x_2|^\alpha &\in L^1((0, T) \times \mathbb{T} \times \mathbb{R}), \end{aligned}$$

then the relative potential energy defined in (2.14) belongs to $\mathcal{C}^1([0, T])$ and there holds

$$\frac{d}{dt} E_{rel}(t) = \int_{\mathbb{T} \times \mathbb{R}} m_2(t, x) dx.$$

Proof. Let $R > 0$ and $\varphi_R : \mathbb{R} \rightarrow [0, 1]$ be a cutoff function with $\varphi_R(x_2) = 1$ for $|x_2| \leq R$, $\varphi_R(x_2) = 0$ for $|x_2| \geq 2R$ and $|\varphi'_R(x_2)| \leq 2R^{-1}$, $x_2 \in \mathbb{R}$.

We abbreviate $E(t) = E_{rel}(t)$ and define

$$E_R(t) := \int_{\mathbb{T} \times \mathbb{R}} (\rho(t, x) - \rho_0(x)) x_2 \varphi_R(x_2) dx.$$

Note that $E(t)$, $E_R(t)$ are well-defined at every time $t \in [0, T)$ and there holds

$$|E(t) - E_R(t)| \leq \|(\rho(t, \cdot) - \rho_0) |x_2|^{1+\alpha}\|_{L^1(\mathbb{T} \times \mathbb{R})} \frac{1}{R^\alpha} = O(R^{-\alpha})$$

uniformly in time as $R \rightarrow \infty$. Thus

$$h^{-1}(E(t+h) - E(t)) = h^{-1}(E_R(t+h) - E_R(t)) + h^{-1}O(R^{-\alpha}).$$

Moreover, the assumed continuity conditions and approximation of the indicator function of $[t, t+h]$ imply

$$\begin{aligned} E_R(t+h) - E_R(t) &= \int_t^{t+h} \int_{\mathbb{T} \times \mathbb{R}} m \cdot \nabla(x_2 \varphi_R(x_2)) dx ds \\ &= \int_t^{t+h} \int_{\mathbb{T} \times \mathbb{R}} m_2 dx ds + \int_t^{t+h} \int_{\mathbb{T} \times \mathbb{R}} m_2(\varphi_R(x_2) - 1 + x_2 \varphi'_R(x_2)) dx ds. \end{aligned}$$

Now the latter term can be bounded by $5R^{-\alpha} \|m_2 |x_2|^\alpha\|_{L^1((0,T) \times \mathbb{T} \times \mathbb{R})}$ implying that

$$h^{-1}(E(t+h) - E(t)) = h^{-1} \int_t^{t+h} \int_{\mathbb{T} \times \mathbb{R}} m_2 dx ds + h^{-1}O(R^{-\alpha}).$$

The statement follows. \square

Let (ρ, v) be the solution constructed in Theorem 3.2 and set

$$m = \rho v - (1 - \rho^2)e_2.$$

Lemma C.2. *In addition to the properties stated in Theorem 3.2 the velocity field v satisfies*

$$|v(t, x)| \leq C e^{-|x_2|}$$

whenever $|x_2| \geq R$ for constants $C, R > 0$ independent of t . The pair (ρ, m) in particular satisfies the conditions of Lemma C.1.

Proof. Regarding the second component one easily sees that

$$|v_2(t, x)| \leq |\mathcal{U}_t| \|\partial_{x_1} \rho(t, \cdot)\|_{L^\infty(\mathbb{T} \times \mathbb{R})} \|K_2(x - \cdot)\|_{L^\infty(\mathcal{U}_t)}$$

which can be bounded by $C e^{-|x_2|}$ for $|x_2| \geq R$ with constants $C, R > 0$ independent of time.

Regarding the first component we can not exploit the decay of the kernel, since $K_1(z) \rightarrow \mp 1$ as $z_2 \rightarrow \pm\infty$. Still by subtracting vanishing horizontal averages we deduce

$$\begin{aligned} |v_1(t, x)| &= \left| \int_{\mathbb{T} \times \mathbb{R}} \partial_{x_1} \rho(t, y) (K_1(x - y) - K_1(x - (0, y_2))) dy \right| \\ &\leq |\mathcal{U}_t| \|\partial_{x_1} \rho(t, \cdot)\|_{L^\infty(\mathbb{T} \times \mathbb{R})} \|\partial_{z_1} K_1(x - \cdot)\|_{L^\infty(\mathcal{U}_t^*)} \pi, \end{aligned}$$

where \mathcal{U}_t^* is the set of points obtained by taking all segments between $y \in \mathcal{U}_t$ and $(0, y_2)$. It is only important that those sets are bounded uniformly in time, which allows to argue as above for v_2 , since $\partial_{z_1} K_1$ now has the required decay. \square

References

- [1] H. Abels and B.-V. Matioc. Well-posedness of the Muskat problem in sub-critical L_p -Sobolev spaces. *European J. Appl. Math.*, 33(2):224–266, 2022.
- [2] S. Agrawal, N. Patel, and S. Wu. Rigidity of acute angled corners for one phase Muskat interfaces. *Adv. Math.*, 412:Paper No. 108801, 71, 2023.
- [3] T. Alazard and O. Lazar. Paralinearization of the Muskat equation and application to the Cauchy problem. *Arch. Ration. Mech. Anal.*, 237(2):545–583, 2020.
- [4] T. Alazard and Q.-H. Nguyen. On the Cauchy problem for the Muskat equation. II: Critical initial data. *Ann. PDE*, 7(1):Paper No. 7, 25, 2021.
- [5] T. Alazard and Q.-H. Nguyen. On the Cauchy problem for the Muskat equation with non-Lipschitz initial data. *Comm. Partial Differential Equations*, 46(11):2171–2212, 2021.
- [6] T. Alazard and Q.-H. Nguyen. Quasilinearization of the 3D Muskat equation, and applications to the critical Cauchy problem. *Adv. Math.*, 399:Paper No. 108278, 52, 2022.
- [7] T. Alazard and Q.-H. Nguyen. Endpoint Sobolev theory for the Muskat equation. *Comm. Math. Phys.*, 397(3):1043–1102, 2023.
- [8] G. Allaire. Homogenization of the Stokes flow in a connected porous medium. *Asymptotic Anal.*, 2(3):203–222, 1989.
- [9] D. Alonso-Orán and R. Granero-Belinchón. Global existence and decay of the inhomogeneous Muskat problem with Lipschitz initial data. *Nonlinearity*, 35(9):4749–4778, 2022.
- [10] D. Amadori and W. Shen. An integro-differential conservation law arising in a model of granular flow. *J. Hyperbolic Differ. Equ.*, 9(1):105–131, 2012.
- [11] P. Amorim. On a nonlocal hyperbolic conservation law arising from a gradient constraint problem. *Bull. Braz. Math. Soc. (N.S.)*, 43(4):599–614, 2012.
- [12] V. Arnaiz, A. Castro, and D. Faraco. Semiclassical estimates for pseudodifferential operators and the Muskat problem in the unstable regime. *Comm. Partial Differential Equations*, 46(1):135–164, 2021.
- [13] F. Betancourt, R. Bürger, K. H. Karlsen, and E. M. Tory. On nonlocal conservation laws modelling sedimentation. *Nonlinearity*, 24(3):855–885, 2011.
- [14] S. Blandin and P. Goatin. Well-posedness of a conservation law with non-local flux arising in traffic flow modeling. *Numer. Math.*, 132(2):217–241, 2016.

- [15] Y. Brenier. The least action principle and the related concept of generalized flows for incompressible perfect fluids. *J. Amer. Math. Soc.*, 2(2):225–255, 1989.
- [16] Y. Brenier. Polar factorization and monotone rearrangement of vector-valued functions. *Comm. Pure Appl. Math.*, 44(4):375–417, 1991.
- [17] T. Buckmaster and V. Vicol. Convex integration constructions in hydrodynamics. *Bull. Amer. Math. Soc. (N.S.)*, 58(1):1–44, 2021.
- [18] S. Cameron. Global well-posedness for the two-dimensional Muskat problem with slope less than 1. *Anal. PDE*, 12(4):997–1022, 2019.
- [19] A. Castro. Mixing solutions for ipm. *Journées équations aux dérivées partielles*, 2017. talk:3.
- [20] A. Castro, D. Córdoba, and D. Faraco. Mixing solutions for the Muskat problem. *Invent. Math.*, 226(1):251–348, 2021.
- [21] A. Castro, D. Córdoba, C. Fefferman, and F. Gancedo. Breakdown of smoothness for the Muskat problem. *Arch. Ration. Mech. Anal.*, 208(3):805–909, 2013.
- [22] A. Castro, D. Córdoba, C. Fefferman, F. Gancedo, and M. López-Fernández. Rayleigh-Taylor breakdown for the Muskat problem with applications to water waves. *Ann. of Math. (2)*, 175(2):909–948, 2012.
- [23] A. Castro, D. Córdoba, and F. Gancedo. A naive parametrization for the vortex-sheet problem. In *Mathematical aspects of fluid mechanics*, volume 402 of *London Math. Soc. Lecture Note Ser.*, pages 88–115. Cambridge Univ. Press, Cambridge, 2012.
- [24] A. Castro, D. Córdoba, F. Gancedo, and R. Orive. Incompressible flow in porous media with fractional diffusion. *Nonlinearity*, 22(8):1791–1815, 2009.
- [25] A. Castro, D. Faraco, and F. Mengual. Degraded mixing solutions for the Muskat problem. *Calc. Var. Partial Differential Equations*, 58(2):Paper No. 58, 29, 2019.
- [26] A. Castro, D. Faraco, and F. Mengual. Localized mixing zone for Muskat bubbles and turned interfaces. *Ann. PDE*, 8(1):Paper No. 7, 50, 2022.
- [27] K. Chen, Q.-H. Nguyen, and Y. Xu. The Muskat problem with C^1 data. *Trans. Amer. Math. Soc.*, 375(5):3039–3060, 2022.
- [28] C. H. A. Cheng, R. Granero-Belinchón, and S. Shkoller. Well-posedness of the Muskat problem with H^2 initial data. *Adv. Math.*, 286:32–104, 2016.

- [29] E. Chiodaroli and O. Kreml. On the energy dissipation rate of solutions to the compressible isentropic Euler system. *Arch. Ration. Mech. Anal.*, 214(3):1019–1049, 2014.
- [30] S. Choi, D. Jerison, and I. Kim. Regularity for the one-phase Hele-Shaw problem from a Lipschitz initial surface. *Amer. J. Math.*, 129(2):527–582, 2007.
- [31] R. M. Colombo, M. Garavello, and M. Lécureux-Mercier. A class of nonlocal models for pedestrian traffic. *Math. Models Methods Appl. Sci.*, 22(4):1150023, 34, 2012.
- [32] P. Constantin, D. Córdoba, F. Gancedo, L. Rodríguez-Piazza, and R. M. Strain. On the Muskat problem: global in time results in 2D and 3D. *Amer. J. Math.*, 138(6):1455–1494, 2016.
- [33] P. Constantin, D. Córdoba, F. Gancedo, and R. M. Strain. On the global existence for the Muskat problem. *J. Eur. Math. Soc. (JEMS)*, 15(1):201–227, 2013.
- [34] P. Constantin, F. Gancedo, R. Shvydkoy, and V. Vicol. Global regularity for 2D Muskat equations with finite slope. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 34(4):1041–1074, 2017.
- [35] A. Córdoba, D. Córdoba, and F. Gancedo. Interface evolution: the Hele-Shaw and Muskat problems. *Ann. of Math. (2)*, 173(1):477–542, 2011.
- [36] A. Córdoba, D. Córdoba, and F. Gancedo. Porous media: the Muskat problem in three dimensions. *Anal. PDE*, 6(2):447–497, 2013.
- [37] D. Córdoba, D. Faraco, and F. Gancedo. Lack of uniqueness for weak solutions of the incompressible porous media equation. *Arch. Ration. Mech. Anal.*, 200(3):725–746, 2011.
- [38] D. Córdoba and F. Gancedo. Contour dynamics of incompressible 3-D fluids in a porous medium with different densities. *Comm. Math. Phys.*, 273(2):445–471, 2007.
- [39] D. Córdoba, F. Gancedo, and R. Orive. Analytical behavior of two-dimensional incompressible flow in porous media. *J. Math. Phys.*, 48(6):065206, 19, 2007.
- [40] D. Córdoba, J. Gómez-Serrano, and A. Zlatoš. A note on stability shifting for the Muskat problem. *Philos. Trans. Roy. Soc. A*, 373(2050):20140278, 10, 2015.

- [41] D. Córdoba, J. Gómez-Serrano, and A. Zlatoš. A note on stability shifting for the Muskat problem, II: From stable to unstable and back to stable. *Anal. PDE*, 10(2):367–378, 2017.
- [42] D. Córdoba and O. Lazar. Global well-posedness for the 2D stable Muskat problem in $H^{3/2}$. *Ann. Sci. Éc. Norm. Supér. (4)*, 54(5):1315–1351, 2021.
- [43] D. Córdoba Gazolaz, R. Granero-Belinchón, and R. Orive-Illera. The confined Muskat problem: differences with the deep water regime. *Commun. Math. Sci.*, 12(3):423–455, 2014.
- [44] C. M. Dafermos. The entropy rate admissibility criterion for solutions of hyperbolic conservation laws. *J. Differential Equations*, 14:202–212, 1973.
- [45] C. M. Dafermos. *Hyperbolic conservation laws in continuum physics*, volume 325 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, fourth edition, 2016.
- [46] H. Darcy. *Les Fontaines Publiques de la Ville de Dijon*. 1856.
- [47] C. De Lellis and L. Székelyhidi, Jr. The Euler equations as a differential inclusion. *Ann. of Math. (2)*, 170(3):1417–1436, 2009.
- [48] C. De Lellis and L. Székelyhidi, Jr. On admissibility criteria for weak solutions of the Euler equations. *Arch. Ration. Mech. Anal.*, 195(1):225–260, 2010.
- [49] C. De Lellis and L. Székelyhidi, Jr. On turbulence and geometry: from Nash to Onsager. *Notices Amer. Math. Soc.*, 66(5):677–685, 2019.
- [50] C. De Lellis and L. Székelyhidi, Jr. Weak stability and closure in turbulence. *Philos. Trans. Roy. Soc. A*, 380(2218):Paper No. 20210091, 16, 2022.
- [51] F. Deng, Z. Lei, and F. Lin. On the two-dimensional Muskat problem with monotone large initial data. *Comm. Pure Appl. Math.*, 70(6):1115–1145, 2017.
- [52] H. Dong, F. Gancedo, and H. Q. Nguyen. Global well-posedness for the one-phase muskat problem. *Comm. Pure Appl. Math.*, 2023.
- [53] J. Escher and B.-V. Matioc. On the parabolicity of the Muskat problem: well-posedness, fingering, and stability results. *Z. Anal. Anwend.*, 30(2):193–218, 2011.
- [54] E. Feireisl. Maximal dissipation and well-posedness for the compressible Euler system. *J. Math. Fluid Mech.*, 16(3):447–461, 2014.
- [55] C. Förster and L. Székelyhidi, Jr. Piecewise constant subsolutions for the Muskat problem. *Comm. Math. Phys.*, 363(3):1051–1080, 2018.

- [56] F. Gancedo and O. Lazar. Global well-posedness for the three dimensional Muskat problem in the critical Sobolev space. *Arch. Ration. Mech. Anal.*, 246(1):141–207, 2022.
- [57] E. García-Juárez, J. Gómez-Serrano, H. Q. Nguyen, and B. Pausader. Self-similar solutions for the Muskat equation. *Adv. Math.*, 399:Paper No. 108294, 30, 2022.
- [58] E. García-Juárez, J. Gómez-Serrano, S. V. Haziot, and B. Pausader. Desingularization of small moving corners for the Muskat equation. *arXiv:2305.05046*.
- [59] B. Gebhard, J. Hirsch, and J. J. Kolombán. On a degenerate elliptic problem arising in the least action principle for Rayleigh-Taylor subsolutions. *arXiv:2208.14495*.
- [60] B. Gebhard and J. J. Kolombán. On bounded two-dimensional globally dissipative Euler flows. *SIAM J. Math. Anal.*, 54(3):3457–3479, 2022.
- [61] B. Gebhard and J. J. Kolombán. Relaxation of the Boussinesq system and applications to the Rayleigh-Taylor instability. *NoDEA Nonlinear Differential Equations Appl.*, 29(1):Paper No. 7, 38, 2022.
- [62] B. Gebhard, J. J. Kolombán, and L. Székelyhidi. A new approach to the Rayleigh-Taylor instability. *Arch. Ration. Mech. Anal.*, 241(3):1243–1280, 2021.
- [63] N. Gigli and F. Otto. Entropic Burgers’ equation via a minimizing movement scheme based on the Wasserstein metric. *Calc. Var. Partial Differential Equations*, 47(1-2):181–206, 2013.
- [64] R. Granero-Belinchón and O. Lazar. Growth in the Muskat problem. *Math. Model. Nat. Phenom.*, 15:Paper No. 7, 23, 2020.
- [65] L. Hitruhin and S. Lindberg. Lamination convex hull of stationary incompressible porous media equations. *SIAM J. Math. Anal.*, 53(1):491–508, 2021.
- [66] P. Isett and V. Vicol. Hölder continuous solutions of active scalar equations. *Ann. PDE*, 1(1):Art. 2, 77, 2015.
- [67] M. Jacobs, I. Kim, and A. R. Mészáros. Weak solutions to the Muskat problem with surface tension via optimal transport. *Arch. Ration. Mech. Anal.*, 239(1):389–430, 2021.
- [68] A. Keimer and L. Pflug. Chapter 6 - nonlocal balance laws – an overview over recent results. In *Numerical Control: Part B*, volume 24 of *Handbook of Numerical Analysis*, pages 183–216. Elsevier, 2023.

- [69] A. Kiselev and Y. Yao. Small scale formations in the incompressible porous media equation. *Arch. Ration. Mech. Anal.*, 247(1):Paper No. 1, 25, 2023.
- [70] S. N. Kružkov. First order quasilinear equations with several independent variables. *Mat. Sb. (N.S.)*, pages 228–255, 1970.
- [71] C. Marchioro and M. Pulvirenti. *Mathematical theory of incompressible non-viscous fluids*, volume 96 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1994.
- [72] B.-V. Matioc. The Muskat problem in two dimensions: equivalence of formulations, well-posedness, and regularity results. *Anal. PDE*, 12(2):281–332, 2019.
- [73] F. Mengual. H-principle for the 2-dimensional incompressible porous media equation with viscosity jump. *Anal. PDE*, 15(2):429–476, 2022.
- [74] F. Mengual and L. Székelyhidi, Jr. Dissipative Euler flows for vortex sheet initial data without distinguished sign. *Comm. Pure Appl. Math.*, 76(1):163–221, 2023.
- [75] G. Menon and F. Otto. Diffusive slowdown in miscible viscous fingering. *Commun. Math. Sci.*, 4(1):267–273, 2006.
- [76] M. Muskat. Two fluid systems in porous media. the encroachment of water into an oil sand. *Physics*, 5(9):250–264, 1934.
- [77] N. Muskhelishvili. *Singular Integral Equations*. Springer Dordrecht, 2011.
- [78] H. Q. Nguyen and B. Pausader. A paradifferential approach for well-posedness of the Muskat problem. *Arch. Ration. Mech. Anal.*, 237(1):35–100, 2020.
- [79] L. Nirenberg. An abstract form of the nonlinear Cauchy-Kowalewski theorem. *J. Differential Geometry*, 6:561–576, 1972.
- [80] T. Nishida. A note on a theorem of Nirenberg. *J. Differential Geometry*, 12(4):629–633, 1977.
- [81] F. Noisette and L. Székelyhidi, Jr. Mixing solutions for the Muskat problem with variable speed. *J. Evol. Equ.*, 21(3):3289–3312, 2021.
- [82] F. Otto. Personal communication.
- [83] F. Otto. Evolution of microstructure in unstable porous media flow: a relaxational approach. *Comm. Pure Appl. Math.*, 52(7):873–915, 1999.
- [84] M. Reissig. Ein abstraktes nichtlineares Cauchy-Kowalewskaja-Theorem mit singulären Koeffizienten. I. *Z. Anal. Anwendungen*, 6(1):35–41, 1987.

- [85] M. Reissig. Ein abstraktes nichtlineares Cauchy-Kowalewskaja-Theorem mit singulären Koeffizienten. II. *Z. Anal. Anwendungen*, 7(2):171–183, 1988.
- [86] P. G. Saffman and G. Taylor. The penetration of a fluid into a porous medium or Hele-Shaw cell containing a more viscous liquid. *Proc. Roy. Soc. London Ser. A*, 245:312–329. (2 plates), 1958.
- [87] E. Sanchez-Palencia. *Non-Homogeneous Media and Vibration Theory*. Springer Berlin Heidelberg, 1980.
- [88] J. Shi. Regularity of solutions to the Muskat equation. *Arch. Ration. Mech. Anal.*, 247(3):Paper No. 36, 46, 2023.
- [89] M. Siegel, R. E. Caflisch, and S. Howison. Global existence, singular solutions, and ill-posedness for the Muskat problem. *Comm. Pure Appl. Math.*, 57(10):1374–1411, 2004.
- [90] L. Székelyhidi. Weak solutions to the incompressible Euler equations with vortex sheet initial data. *C. R. Math. Acad. Sci. Paris*, 349(19-20):1063–1066, 2011.
- [91] L. Székelyhidi, Jr. Relaxation of the incompressible porous media equation. *Ann. Sci. Éc. Norm. Supér. (4)*, 45(3):491–509, 2012.
- [92] L. Tartar. Compensated compactness and applications to partial differential equations. In *Nonlinear analysis and mechanics: Heriot-Watt Symposium, Vol. IV*, volume 39 of *Res. Notes in Math.*, pages 136–212. Pitman, Boston, Mass.-London, 1979.
- [93] L. Tartar. Convergence of the homogenization process. *Appendix of Nonhomogeneous media and vibration theory, Lecture Notes in Phys. 127*, Springer-Verlag, New York, 1980.
- [94] S. Thalabard, J. Bec, and A. Mailybaev. From the butterfly effect to spontaneous stochasticity in singular shear flows. *Commun. Phys.*, 3(122), 2020.

Acknowledgements. A.C., D.F. and B.G. acknowledge financial support from the Severo Ochoa Programme for Centres of Excellence Grant CEX2019-000904-S funded by MCIN/AEI/10.13039/501100011033, A.C. from Grant PID2020-114703GB-I00 funded by MCIN/AEI/10.13039/501100011033 and D.F. and B.G. grant PI2021-124-195NB-C32 funded by MCIN/AEI/10.13039/501100011033. D.F. was also partially supported by CAM through the Line of excellence for University Teaching Staff between CM and UAM. D.F. and B.G. were also partially supported by the ERC Advanced Grant 834728. B.G. is supported by María Zambrano Grant CA6/RSUE/2022-00097. Finally A.C. and D.F. acknowledge financial support from Grants RED2022-134784-T and RED2018-102650-T funded by MCIN/AEI/10.13039/501100011033.

Ángel Castro

Instituto de Ciencias Matemáticas, CSIC-UAM-UC3M-UCM, 28049 Madrid, Spain
angel_castro@icmat.es

Daniel Faraco

Departamento de Matemáticas, Universidad Autónoma de Madrid, Instituto de
Ciencias Matemáticas, CSIC-UAM-UC3M-UCM, 28049 Madrid, Spain
daniel.faraco@uam.es

Björn Gebhard

Departamento de Matemáticas, Universidad Autónoma de Madrid, Instituto de
Ciencias Matemáticas, CSIC-UAM-UC3M-UCM, 28049 Madrid, Spain
gebhard.bjorn@uam.es