

# Simultaneous inference for generalized linear models with unmeasured confounders

Jin-Hong Du<sup>†‡</sup>      Larry Wasserman<sup>†‡</sup>      Kathryn Roeder<sup>†§</sup>

April 23, 2024

## Abstract

Tens of thousands of simultaneous hypothesis tests are routinely performed in genomic studies to identify differentially expressed genes. However, due to unmeasured confounders, many standard statistical approaches may be substantially biased. This paper investigates the large-scale hypothesis testing problem for multivariate generalized linear models in the presence of confounding effects. Under arbitrary confounding mechanisms, we propose a unified statistical estimation and inference framework that harnesses orthogonal structures and integrates linear projections into three key stages. It begins by disentangling marginal and uncorrelated confounding effects to recover the latent coefficients. Subsequently, latent factors and primary effects are jointly estimated through lasso-type optimization. Finally, we incorporate projected and weighted bias-correction steps for hypothesis testing. Theoretically, we establish the identification conditions of various effects and non-asymptotic error bounds. We show effective Type-I error control of asymptotic  $z$ -tests as sample and response sizes approach infinity. Numerical experiments demonstrate that the proposed method controls the false discovery rate by the Benjamini-Hochberg procedure and is more powerful than alternative methods. By comparing single-cell RNA-seq counts from two groups of samples, we demonstrate the suitability of adjusting confounding effects when significant covariates are absent from the model.

*Keywords:* Hidden variables; High-dimensional regression; Hypothesis testing; Multivariate response regression; Nuisance parameters; Surrogate variables analysis.

---

<sup>†</sup>Department of Statistics and Data Science, Carnegie Mellon University, Pittsburgh, PA 15213, USA.

<sup>‡</sup>Machine Learning Department, Carnegie Mellon University, Pittsburgh, PA 15213, USA.

<sup>§</sup>Computational Biology Department, Carnegie Mellon University, Pittsburgh, PA 15213, USA.

# 1 Introduction


To discover genes that are differentially expressed under different experimental conditions or across groups of samples, large numbers of simultaneous hypothesis tests must be performed. These tests are made more challenging by the presence of unmeasured covariates that bias the analyses. In 2008, [Leek and Storey \(2008\)](#) presented their pathbreaking “surrogate variable” approach to control for unmeasured confounding effects in differential expression (DE) studies using microarray data. These confounders go by various names in the literature, including batch effects, surrogate variables, latent effects, or simply unwanted variations ([Leek et al., 2010](#); [Gagnon-Bartsch and Speed, 2012](#); [Sun et al., 2012](#)). Adjusting for confounding effects is crucial because they may distort the correct null distribution of the test statistics, and consequently, standard statistical approaches can be substantially biased ([Wang et al., 2017a](#); [McKenna and Nicolae, 2019](#)). Due to burgeoning developments in the genomics field, DE testing has been dramatically expanded to include a variety of genomic readouts beyond microarray, in which the normality of the observed counts rarely holds. Inspired by modern-day omic studies, the concerns about confounding are more urgent than ever, and there is a pressing need to adapt statistical approaches to changing data types.

The problem of confounder adjustment has been an important topic in statistics in recent years. To characterize the confounding effects, the pioneering work in this field assumes a linear model  $\mathbf{Y} = \mathbf{X}\mathbf{B}^\top + \mathbf{Z}\mathbf{\Gamma}^\top + \mathbf{E}$ , where  $\mathbf{Y} \in \mathbb{R}^{n \times p}$  is the gene expression matrix,  $\mathbf{X} \in \mathbb{R}^{n \times d}$  is the measured covariate matrix,  $\mathbf{B} \in \mathbb{R}^{p \times d}$  is the direct effect to be estimated,  $\mathbf{Z} \in \mathbb{R}^{n \times r}$  is the latent factor matrix,  $\mathbf{\Gamma} \in \mathbb{R}^{p \times r}$  is the latent factor loading, and  $\mathbf{E} \in \mathbb{R}^{n \times p}$  is the additive noise. The early investigations study the statistical inference problem under this model by further imposing a linear relationship between  $\mathbf{X}$  and  $\mathbf{Z}$ , assuming either  $\mathbf{X}$  causes  $\mathbf{Z}$  as in Figure 1(a), i.e.,  $\mathbf{Z}$  is a hidden mediator ([Leek and Storey, 2008](#); [Wang et al., 2017a](#); [Gerard and Stephens, 2020](#)), or  $\mathbf{Z}$  causes  $\mathbf{X}$  as in Figure 1(b), i.e.,  $\mathbf{Z}$  is a hidden confounder ([Guo et al., 2022](#); [Sun et al., 2023](#)).

In the presence of hidden mediators, where the observed covariates are the cause of the hidden variables, [Wang et al. \(2017a\)](#) and [Gerard and Stephens \(2020\)](#) study the statistical inference problem for multiple outcomes ( $p > 1$ ) by assuming a linear relationship between the observed variables and the hidden variables. In the context of hidden confounders and a single outcome ( $p = 1$ ), [Guo et al. \(2022\)](#) propose a doubly debiased lasso estimator and establish asymptotic normality; [Ćevic et al. \(2020\)](#) propose a spectral de-confounding method; and [Sun et al. \(2023\)](#) analyze non-asymptotic and asymptotic false discovery control with high-dimensional covariates.

More recently, methods for estimating primary effects extend beyond linear dependence structures between covariates and confounders. For instance, [Jiang and Ning \(2022\)](#) model the interaction between the covariates and the confounders, and projection-based methods are employed to estimate the primary effects under arbitrary dependency ([Lee et al., 2017](#); [Bing et al., 2022](#); [McKenna and Nicolae, 2022](#)). For statistical inference, [McKenna and Nicolae \(2019\)](#) propose an estimator that is asymptotically equivalent to the ordinary least squares estimators obtained when every covariate is observed, and [Bing et al. \(2023\)](#) establish asymptotic normality, efficiency, and consistency.

The applicability of the aforementioned methods to the nonlinear model remains challenging. Limited research has been done to address adjustments for confounding effects



**Figure 1:** Causal diagrams on the generative models illustrating the relationship between the covariate  $\mathbf{X}$ , the latent variable  $\mathbf{Z}$ , and the response  $\mathbf{Y}$ . (a)  $\mathbf{Z}$  is a hidden mediator when  $\mathbf{X}$  causes  $\mathbf{Z}$ . (b) hidden confounder when  $\mathbf{Z}$  causes  $\mathbf{X}$ . Note that we do not require knowledge of the relationship between  $\mathbf{X}$  and  $\mathbf{Z}$  for the analysis in this paper.

under the setting of *arbitrary confounding mechanisms, nonlinear models, and multiple outcomes*. For empirical studies, [Salim et al. \(2022\)](#) propose a heuristic algorithm that utilizes a pseudo-replicate design matrix and negative control genes to remove unwanted variations. For theoretical analysis, to the best of our knowledge, the related literature that explores slightly broader settings is limited to [Feng \(2020\)](#), who studies nonlinear factor models concerning treatment effects with a single outcome by PCA-based matching, and [Ouyang et al. \(2023\)](#), who study the generalized linear models with a single outcome and linear hidden confounders. However, both of these works assume the covariates are some functions of the unobserved confounders.

Our work is inspired by the rapid developments in the field of genomics, particularly single-cell omics ([Editorial, 2023](#)). For example, CRISPR perturbations with single-cell sequencing readouts have promised extraordinary scientific insight ([Kampmann, 2020](#); [Hong and Iakoucheva, 2023](#); [Cheng et al., 2023](#)); due to the sparsity of outcomes and the nature of the molecular readout, these data are not suitable for analysis by linear models under Gaussianity assumptions ([Sarkar and Stephens, 2021](#); [Barry et al., 2023](#)). Hence, our development of generalized linear models for confounding is timely.

In this paper, we adopt the term “confounder” to encompass a broad category of latent variables, including both mediators and confounders, as defined in the context of causal inference literature. The purpose of this paper is to derive valid simultaneous inference for multivariate generalized linear models in the presence of unmeasured confounding effects. Existing methods in this domain typically focus on Gaussian linear models ([Wang et al., 2017a](#); [Bing et al., 2022, 2023](#)) or necessitate direct modeling of the relationship between covariates and confounders ([Feng, 2020](#); [Ouyang et al., 2023](#)). To the best of our knowledge, the proposed method is the first estimation and inference framework capable of (1) accommodating general relationships between observed covariates and unmeasured confounders, allowing for *arbitrary confounding mechanisms*; (2) utilizing generalized linear models, allowing for *nonlinear modeling*; and (3) incorporating information from *multiple outcomes*. Our approach leverages the orthogonal structures inherent in the problem, incorporating linear projection techniques into both estimation and inference processes to effectively mitigate confounding effects and elucidate primary effects. Notably, it exhibits significant utility in high-dimensional sparse count data, as demonstrated through the analysis of single-cell

datasets on systemic lupus erythematosus disease in Section 6.

Our proposed procedure GCATE (generalized confounder adjustment for testing and estimation) consists of three main steps. In the first step, we use joint maximum likelihood estimation (Chen et al., 2019; Chen and Li, 2022) to obtain the initial estimate of the marginal effects and uncorrelated latent components by projecting the latent factor  $\mathbf{Z}$  to the orthogonal space of  $\mathbf{X}$ , from which we recover the column space of  $\mathbf{\Gamma}$ . In the second step, we use a similar strategy to obtain the estimates of both  $\mathbf{Z}$  and primary effect  $\mathbf{B}$ , by constraining the latter to be orthogonal to the estimated latent coefficients  $\hat{\mathbf{\Gamma}}$  and using  $\ell_1$ -regularization to encourage sparsity. Lastly, the valid inference is guaranteed by a bias-corrected estimator of  $\hat{\mathbf{B}}$ , which innovates a link-specific weight function, similar to Javanmard and Montanari (2014); Cai et al. (2021), while incorporating projection-based score adjustments that combine the information from multivariate responses.

In our theoretical framework of confounded generalized linear models, we establish conditions for identifying the latent coefficients and direct effects. Furthermore, we provide non-asymptotic estimation error bounds for these estimated quantities, in the high-dimensional scenarios when both the sample size  $n$  and response size  $p$  tend to infinity. In particular, we derive element-wise  $\ell_2$ -norm and  $\ell_1$ -norm bounds for the estimation error of the primary effects by effectively controlling the column-wise estimation errors of the latent components. Lastly, we demonstrate the asymptotic normality of our proposed bias-corrected estimator and show the proper control of statistical errors, thereby enabling the construction of valid confidence intervals and hypothesis tests.

**Organization and Notation.** In Section 2, we set up our modeling framework, which extends existing results in the literature to the generalized linear model setting. In Section 3, we describe our strategy for estimation and establish bounds on the estimation error of the parameters of interest. In Section 4, we motivate and construct asymptotically valid confidence intervals and hypothesis tests. Finally, in Section 5 and Section 6, we study the empirical behavior of our estimators in realistic simulations and a study of gene expression in lupus patients. The code for reproducing experimental results in the paper can be found at <https://github.com/jaydu1/gcate/>.

Throughout our exposition, we will use the following notational conventions. For any matrix  $\mathbf{A} \in \mathbb{R}^{n \times p}$ , we use  $\mathbf{a}_i$ ,  $\mathbf{A}_j$ , and  $a_{ij}$  to denote its  $i$ th row,  $j$ th column, and  $(i, j)$ -th entry, respectively, for  $i = 1, \dots, n$ ,  $j = 1, \dots, p$ . For any matrix  $\mathbf{A} \in \mathbb{R}^{n \times p}$  with full column rank, let  $\mathcal{P}_{\mathbf{A}} = \mathbf{A}(\mathbf{A}^\top \mathbf{A})^{-1} \mathbf{A}^\top$  and  $\mathcal{P}_{\mathbf{A}}^\perp = \mathbf{I}_p - \mathcal{P}_{\mathbf{A}}$  be the orthogonal projection matrices on the  $\mathbf{A}$ 's column space and its orthogonal space, respectively. For any square matrix  $\mathbf{A} \in \mathbb{R}^{n \times n}$ ,  $\lambda_i(\mathbf{A})$  denotes its  $i$ th eigenvalue. The symbol " $\odot$ " denotes the Hadamard product. We use " $o$ " and " $\mathcal{O}$ " to denote the little-o and big-O notations and let " $o_{\mathbb{P}}$ " and " $\mathcal{O}_{\mathbb{P}}$ " be their probabilistic counterparts. For sequences  $\{a_n\}$  and  $\{b_n\}$ , we write  $a_n \ll b_n$  or  $b_n \gg a_n$  if  $a_n = o(b_n)$ ;  $a_n \lesssim b_n$  or  $b_n \gtrsim a_n$  if  $a_n = \mathcal{O}(b_n)$ ; and  $a_n \asymp b_n$  if  $a_n = \mathcal{O}(b_n)$  and  $b_n = \mathcal{O}(a_n)$ . Convergence in distribution and probability are denoted by " $\xrightarrow{d}$ " and " $\xrightarrow{\mathbb{P}}$ ".

## 2 Modeling Differential Expression

In the context of DE testing and related applied problems, the outcome variable can be a variety of measures, including gene expression, protein abundance, and open chromatin. For simplicity of exposition, we will describe our methods in the context of tests for differential gene expression. These tests aim to contrast outcomes from case versus control samples, wherein case and control observations may be derived from various study designs, spanning the spectrum from diseased versus healthy subjects to perturbed versus non-targeted cells.

### 2.1 Generalized linear model with hidden confounders

Suppose the gene expression  $\mathbf{y} \in \mathbb{R}^p$  is a  $p$ -dimensional random vector containing conditional independent entries from a one-dimensional exponential family with density:

$$p(y_j | \theta_j) = h(y_j) \exp(y_j \theta_j - A(\theta_j)),$$

where  $\theta_j \in \mathbb{R}$  is the *natural parameter*, and  $A(\cdot)$  and  $h(\cdot)$  are functions that depend on the member of the exponential family. We restrict ourselves to the regular families whose natural parameter space is a nonempty open set and  $A$  is continuously thrice differentiable, which is satisfied by most common exponential families as summarized in Table F1. Because the one-parameter exponential family is minimal, the natural parameter space is convex, and the *log-partition function*  $A$  is strictly convex. If we know the distribution of  $\mathbf{y}$ , then  $\boldsymbol{\theta} \in \mathbb{R}^p$  is a unique solution to the equation  $\mathbb{E}[\mathbf{y} | \boldsymbol{\theta}] = A'(\boldsymbol{\theta})$ , where  $A'$  is the first derivative of  $A$  and applied element-wise to  $\boldsymbol{\theta}$ ; equivalently,  $\boldsymbol{\theta} = A'^{-1}(\mathbb{E}[\mathbf{y} | \boldsymbol{\theta}])$ . In other words, we can recover  $\boldsymbol{\theta}$  based on the information of the first moment of  $\mathbf{y}$  and the log-partition function  $A$ .

To associate multiple outcomes with both covariates and hidden confounders, one can naturally consider the generalized linear model, where the natural parameters are linear functions of both the observed covariates  $\mathbf{x} \in \mathbb{R}^d$  and the unmeasured confounder  $\mathbf{z} \in \mathbb{R}^r$ :

$$\boldsymbol{\theta}_{p \times 1} = \mathbf{B}_{p \times d} \mathbf{x}_{d \times 1} + \boldsymbol{\Gamma}_{p \times r} \mathbf{z}_{r \times 1}.$$

Here,  $\mathbf{B}$  and  $\boldsymbol{\Gamma}$  are the linear coefficients. Denote  $\mathbf{D}\mathbf{x}$  the linear projection of  $\mathbf{z}$  onto  $\mathbf{x}$ , where  $\mathbf{D} := \mathbb{E}[\mathbf{z}\mathbf{x}^\top] \mathbb{E}[\mathbf{x}\mathbf{x}^\top]^{-1} \in \mathbb{R}^{r \times d}$  is the projection coefficient and  $\mathbf{w} = \mathbf{z} - \mathbf{D}\mathbf{x}$  is the residual uncorrelated with  $\mathbf{x}$ . To see how  $\mathbf{z}$  may affect the inference on  $\mathbf{B}$ , note that

$$\boldsymbol{\theta} = (\mathbf{B} + \boldsymbol{\Gamma}\mathbf{D})\mathbf{x} + \boldsymbol{\Gamma}\mathbf{w}. \tag{2.1}$$

When  $\mathbf{y}$  is normally distributed, the confounding effects occur even when regressing the mean response  $\boldsymbol{\theta} = \mathbb{E}[\mathbf{y} | \boldsymbol{\theta}]$  on  $\mathbf{x}$ , which yields the *confounded coefficient*  $\mathbf{B} + \boldsymbol{\Gamma}\mathbf{D}$  while the *direct effect* of interest is  $\mathbf{B}$ . When  $\mathbf{y}$  comes from general exponential families, the confounding effects are more intractable because all moments and cumulants of the response may be affected by the colinearity of  $\mathbf{x}$  and  $\mathbf{z}$ .

In the context of genomic analysis, the problem of confounding is more severe when the number of available covariates is limited. In particular, one typically encounters high-dimensional scenarios characterized by a substantial number of genes, often surpassing the available numbers of covariates and hidden confounding factors. In this paper, we also

consider such a challenging scenario where the number of genes is much larger than the numbers of the observed covariates and the unmeasured confounders, namely,  $p \gg d$  and  $p \gg r$ . Under such challenges, the first natural and essential question one may ask is whether there is any hope to disentangle the confounding effects and identify the direct effects.

The answer to this inquiry is affirmative. On the one hand, the column space of  $\mathbf{\Gamma}$  can be identified up to rotations if  $\text{Cov}(\mathbf{\Gamma}\mathbf{w}) = \mathbf{\Gamma}\mathbf{\Sigma}_w\mathbf{\Gamma}^\top$  has rank  $r$ , where  $\mathbf{\Sigma}_w = \text{Cov}(\mathbf{w})$  is the covariance of the uncorrelated latent factors. This fact originates from basic principles in linear algebra, frequently employed in factor analysis (Bai, 2003; Bai and Li, 2012). On the other hand, once the column space of  $\mathbf{\Gamma}$  is known, one can apply the orthogonal projections to remove the confounding effects based on (2.1):

$$\mathcal{P}_\Gamma^\perp \boldsymbol{\theta} = \mathcal{P}_\Gamma^\perp \mathbf{B}\mathbf{x},$$

where  $\mathcal{P}_\Gamma = \mathbf{\Gamma}(\mathbf{\Gamma}^\top\mathbf{\Gamma})^{-1}\mathbf{\Gamma}^\top$  and  $\mathcal{P}_\Gamma^\perp = \mathbf{I}_p - \mathcal{P}_\Gamma$  are the orthogonal projection matrices that project vectors on to the image of  $\mathbf{\Gamma}$  and the orthogonal complement of  $\mathbf{\Gamma}$ , respectively. By regressing  $\mathcal{P}_\Gamma^\perp \boldsymbol{\theta}$  on  $\mathbf{x}$ , we can further obtain the unconfounded primary effects  $\mathcal{P}_\Gamma^\perp \mathbf{B} = \mathcal{P}_\Gamma^\perp \mathbb{E}[\boldsymbol{\theta}\mathbf{x}^\top] \mathbb{E}[\mathbf{x}\mathbf{x}^\top]^{-1}$ . However, the other component,  $\mathcal{P}_\Gamma \mathbf{B}$ , often poses challenges in identifiability unless additional conditions are imposed. Typically, extra assumptions on the spectrum of  $\mathbf{\Gamma}$  and the sparsity of  $\mathbf{B}$  are necessary, to assert that  $\mathbf{\Gamma}$  and  $\mathbf{B}$  are asymptotically orthogonal, in the sense that  $\mathcal{P}_\Gamma \mathbf{B}$  is negligible (Lee et al., 2017; Wang et al., 2017a; Bing et al., 2023). In that case,  $\mathbf{B}$  can be well approximated by  $\mathcal{P}_\Gamma^\perp \mathbf{B}$ . Below, we give one sufficient identification condition.

**Proposition 1** (Identification of  $\mathbf{B}$ ). Suppose there exists a sequence  $\{\tau_p\}_{p \in \mathbb{N}}$  that is uniformly lower bounded away from zero such that the following conditions hold:

$$\lambda_r(\mathbf{\Gamma}\mathbf{\Sigma}_w\mathbf{\Gamma}^\top) \geq \tau_p, \quad \max_{1 \leq j \leq p} (\mathbf{\Gamma}\mathbf{\Sigma}_w\mathbf{\Gamma}^\top)_{jj} = \mathcal{O}(1), \quad \max_{1 \leq \ell \leq d} \|\mathbf{B}_\ell\|_1 = o(\tau_p). \quad (2.2)$$

Then as  $p$  tends to infinity, it follows that  $\mathbf{B} = \mathcal{P}_\Gamma^\perp \mathbf{B} + o(1)$  and  $\|\mathcal{P}_\Gamma \mathbf{B}\|_F \lesssim \sqrt{p} \|\mathbf{B}\|_{1,1} / \tau_p$ , where  $\|\cdot\|_{1,1}$  is the element-wise  $\ell_1$ -norm. Further,  $\mathcal{P}_\Gamma$  and  $\mathbf{B}$  can be identified from the first two moments of  $\mathbf{x}, \mathbf{y}$  asymptotically.

As hinted above, the lower bound condition of  $\mathbf{\Gamma}\mathbf{\Sigma}_w\mathbf{\Gamma}^\top$ 's spectrum in (2.2) ensures that the column space of  $\mathbf{\Gamma}$  can be identified up to rotations. The second condition guarantees that the diagonal entries of  $\mathbf{\Gamma}\mathbf{\Sigma}_w\mathbf{\Gamma}^\top$  are balanced. Finally, the last condition in (2.2) can hold when  $\mathbf{B}$  is sparse, and its entry is bounded. Compared to Bing et al. (2023, Theorem 1) where the response  $\mathbf{y}$  is normally distributed, the identifiability condition of Proposition 1 applies for exponential families, which is of much broader generality. Furthermore, the smallest eigenvalue of  $\mathbf{\Gamma}\mathbf{\Sigma}_w\mathbf{\Gamma}^\top$  can grow at a specific rate  $\tau_p$  in Proposition 1. When  $\tau_p = p$ , we can recover the result in Bing et al. (2023, Theorem 1). Lastly, we also provide a norm bound for the residual  $\mathcal{P}_\Gamma \mathbf{B}$ , which is helpful for later analysis of the estimation errors.

## 2.2 Random samples

While the preceding identification outcomes are applicable when population moments are known, practical scenarios involve the observation of independent and identically distributed

(i.i.d.) samples. Consequently, statistical estimation becomes imperative to disentangle direct effects from the confounding effects based on samples. Suppose  $(\mathbf{x}_i, \mathbf{y}_i)$  for  $i = 1, \dots, n$  are  $n$  i.i.d. samples coming from the same distribution as  $(\mathbf{x}, \mathbf{y})$ , and let  $\mathbf{X} \in \mathbb{R}^{n \times d}$  and  $\mathbf{Y} \in \mathbb{R}^{n \times p}$  denote the design matrix and the gene expression matrix, respectively. The expression  $y_{ij}$  of the  $i$ th observation and the  $j$ th gene has the density:

$$p(y_{ij} | \theta_{ij}) = h(y_{ij}) \exp(y_{ij}\theta_{ij} - A(\theta_{ij})),$$

where  $\theta_{ij}$  is the natural parameter. In matrix form, the natural parameters decompose as

$$\Theta = \mathbf{X}\mathbf{B}^\top + \mathbf{Z}\mathbf{\Gamma}^\top,$$

where  $\mathbf{B} \in \mathbb{R}^{p \times d}$ ,  $\mathbf{Z} \in \mathbb{R}^{n \times r}$ , and  $\mathbf{\Gamma} \in \mathbb{R}^{p \times r}$  are unknown. Note that  $y_{ij}$ 's are conditionally independent given the natural parameter  $\Theta$ .

One natural way to estimate the unknown variable  $\mathbf{Z}$  and parameters  $(\mathbf{B}, \mathbf{\Gamma})$  is to perform maximum likelihood estimation. Ignoring the constant terms, the negative log-likelihood function of  $\mathbf{Y}$  is given by

$$\mathcal{L}(\Theta) = \mathcal{L}(\mathbf{B}, \mathbf{Z}, \mathbf{\Gamma}) = -\frac{1}{n} \sum_{i=1}^n \sum_{j=1}^p (y_{ij}\theta_{ij} - A(\theta_{ij})). \quad (2.3)$$

The second notation  $\mathcal{L}(\mathbf{B}, \mathbf{Z}, \mathbf{\Gamma})$  reflects the dependence of  $\Theta$  on the model parameters  $\mathbf{B}$  and unknown quantities  $\mathbf{\Gamma}, \mathbf{Z}$ . To tackle the challenges raised by the nonconvexity and high-dimensionality, the strategic development of efficient algorithms for estimating these unknown quantities and conducting a thorough analysis of their statistical properties emerges as a critically significant endeavor.

To overcome the difficulty of estimation, one critical observation comes from the projection-based decomposition and Proposition 1:

$$\begin{aligned} \Theta &= \mathbf{X}\mathbf{B}^\top + \mathbf{Z}\mathbf{\Gamma}^\top \\ &= (\mathbf{X}\mathbf{B}^\top \mathcal{P}_\Gamma^\perp + \mathbf{X}\mathbf{B}^\top \mathcal{P}_\Gamma) + (\mathbf{X}\mathbf{D}^\top \mathbf{\Gamma}^\top + \mathbf{W}\mathbf{\Gamma}^\top) \\ &= \mathbf{X}\mathbf{B}^\top \mathcal{P}_\Gamma^\perp + \mathcal{P}_\mathbf{X} \mathbf{Z}\mathbf{\Gamma}^\top + \mathcal{P}_\mathbf{X}^\perp \mathbf{W}\mathbf{\Gamma}^\top + \mathbf{o}_\mathbb{P}(1), \end{aligned}$$

where we replace the best linear projection  $\mathbf{X}\mathbf{D}$  with its empirical counterpart  $\mathcal{P}_\mathbf{X} \mathbf{Z}$  in finite samples, which yield negligible terms that contribute to  $\mathbf{o}_\mathbb{P}(1)$ . It is worth noting that  $\mathbf{X}\mathbf{B}^\top \mathcal{P}_\Gamma^\perp$  and  $\mathcal{P}_\mathbf{X} \mathbf{Z}\mathbf{\Gamma}^\top + \mathbf{W}\mathbf{\Gamma}^\top$  have orthogonal columns, while  $\mathbf{X}\mathbf{B}^\top \mathcal{P}_\Gamma^\perp + \mathcal{P}_\mathbf{X} \mathbf{Z}\mathbf{\Gamma}^\top$  and  $\mathcal{P}_\mathbf{X}^\perp \mathbf{W}\mathbf{\Gamma}^\top$  have orthogonal rows. Our analysis will then take advantage of such two-way structural orthogonality to perform both estimation and inference for the parameters of interest, as detailed in the following sections.

### 3 Estimation

From now on, we will use an asterisk on the upper subscript to indicate the population parameters and the true latent factors. Specifically, we denote the underlying parameter as

$$\Theta^* = \mathbf{X}\mathbf{B}^{*\top} + \mathbf{Z}^*\mathbf{\Gamma}^{*\top} = \mathbf{X}(\mathbf{B}^* + \mathbf{\Gamma}^*\mathbf{D}^*)^\top + \mathbf{W}^*\mathbf{\Gamma}^{*\top}.$$

Let  $\mathcal{R} \subseteq \mathbb{R}$  be an open domain of  $\theta$  such that  $A(\theta) < \infty$  for all  $\theta \in \mathcal{R}$ . For a given  $C > 0$ , define  $\mathcal{R}_C = \mathcal{R} \cap [-C, C]$  for Gaussian, Binomial and Poisson distributions and  $\mathcal{R}_C = \mathcal{R} \cap [-C, -1/C]$  for Negative Binomial distributions. For our theoretical results, we assume the existence of constant  $C > 1$  such that the following common assumptions hold.

**Assumption 1** (Model parameters). Assume that  $\Theta^* \in \mathcal{R}_C^{n \times p}$  with probability  $\iota_n$  for some deterministic sequence  $\iota_n$  tending to one as  $n$  tends to infinity. The primary coefficient satisfies that  $\max_{1 \leq \ell \leq d} \|\mathbf{B}_\ell^*\|_0 \leq s$  for some  $1 \leq s \leq p$  and  $\max_{1 \leq j \leq p} \|\mathbf{b}_j^*\|_2 \leq C$ .

**Assumption 2** (Covariates). Assume that  $\mathbf{x}_1, \dots, \mathbf{x}_n$  are i.i.d.  $\nu$ -sub-Gaussian random vectors with second moment  $\Sigma_x := \mathbb{E}[\mathbf{x}_1\mathbf{x}_1^\top]$  such that  $C^{-1} \leq \lambda_p(\Sigma_x) \leq \lambda_1(\Sigma_x) \leq C$ .

**Assumption 3** (Latent vectors). Assume that the uncorrelated latent factors  $\mathbf{w}_1^*, \dots, \mathbf{w}_n^*$  are i.i.d.  $\nu$ -sub-Gaussian random vectors with zero means and covariance  $\Sigma_w$ , such that  $C^{-1} \leq \lambda_r(\Sigma_w) \leq \lambda_1(\Sigma_w) \leq C$ ; and the factor loadings  $\mathbf{\Gamma}^*$  satisfy that  $C^{-1} \leq \lambda_r(p^{-1}\mathbf{\Gamma}^*\mathbf{\Gamma}^{*\top}) \leq C$  and  $\max_{1 \leq j \leq p} \|\boldsymbol{\gamma}_j^*\|_2 \leq C$ .

Like all nonlinear (nonconvex) analyses, the rows of  $\mathbf{B}^*$  and  $\mathbf{\Gamma}^*$  are assumed to be in a bounded set, as in Assumptions 1 and 3. The boundedness of the natural parameter  $\Theta^*$  is required to control the tail probability of the response  $y$  conditional on observed covariates and latent factors. In Assumption 2, the sub-Gaussian assumptions admit the particular case when  $x_{i1} = 1$  for all  $1 \leq i \leq n$  so that the intercept can be incorporated into our model. In Assumption 3, the zero mean condition on  $\mathbf{W}^*$  is to simplify the theoretical analysis, which can be guaranteed if we include the intercept and project  $\mathbf{Z}^*$  onto the linear span of the columns of  $[\mathbf{1}_n, \mathbf{X}]$ . Finally, the sparsity and boundedness assumptions of  $\mathbf{B}^*$  in Assumption 1, and the bounded spectrum assumptions of  $\Sigma_w$  and  $\mathbf{\Gamma}^*$  in Assumption 3 imply the conditions of Proposition 1 with  $\tau_p = p$  therein. Finally, we note that the assumptions on the projection coefficient  $\mathbf{D}^*$  are less restrictive, as long as  $\Theta^*$  is within a bounded set with high probability.

**Remark 1** (The number of latent factors). For our theoretical results, we assume the number of latent factors  $r$  is known in advance. Note that the joint-likelihood-based information criterion (JIC) proposed by Chen and Li (2022) can be utilized to select the number of latent factors. The JIC value is the sum of deviance and a penalty on model complexity:

$$\text{JIC}(\widehat{\Theta}^{(r)}) = -2 \sum_{i \in [n], j \in [p]} \log p(y_{ij} \mid \widehat{\theta}_{ij}^{(r)}) + \nu(n, p, d + r), \quad (3.1)$$

where  $\widehat{\Theta}^{(r)}$  is the joint maximum likelihood estimator of the natural parameter matrix that minimizes (2.3) with  $r$  latent factors and  $d$  observed covariates, and  $\nu(n, p, r) =$



$c_{\text{JIC}} \cdot r \log(n \wedge p)(n \wedge p)^{-1}$  is the complexity measure with penalty level  $c_{\text{JIC}} > 0$ . As shown by [Chen and Li \(2022\)](#), minimizing the empirical JIC yields a consistent estimate for the number of factors in generalized linear factor models with an intercept parameter ( $d = 1$ ). The utility of this metric in our problem setting is also empirically examined for both the simulation in [Section 5](#) and the real data analysis in [Section 6](#).

As motivated in [Section 2.1](#), we consider the following optimization problem:

$$\begin{aligned} \widehat{\mathbf{B}}, \widehat{\mathbf{Z}}, \widehat{\mathbf{\Gamma}} = & \underset{\mathbf{B} \in \mathbb{R}^{p \times d}, \mathbf{Z} \in \mathbb{R}^{n \times r}, \mathbf{\Gamma} \in \mathbb{R}^{p \times r}}{\operatorname{argmin}} \quad \mathcal{L}(\mathbf{B}, \mathbf{Z}, \mathbf{\Gamma}) + \lambda \|\mathbf{B}\|_{1,1} \\ \text{s.t.} \quad & \mathbf{X}\mathbf{B}^\top + \mathbf{Z}\mathbf{\Gamma}^\top \in \mathcal{R}_C^{n \times p}, \quad \mathcal{P}_{\mathbf{\Gamma}}\mathbf{B} = \mathbf{0}. \end{aligned} \quad (3.2)$$

where the unregularized loss function  $\mathcal{L}(\mathbf{B}, \mathbf{Z}, \mathbf{\Gamma})$  is defined in [\(2.3\)](#) and  $\|\cdot\|_{1,1}$  denotes the element-wise  $\ell_1$ -norm. It is worth noting that for any feasible  $\widehat{\mathbf{\Gamma}}$  fixed, [\(3.2\)](#) reduces to

$$\begin{aligned} \widehat{\mathbf{B}}, \widehat{\mathbf{Z}} = & \underset{\mathbf{B} \in \mathbb{R}^{p \times d}, \mathbf{Z} \in \mathbb{R}^{n \times r}}{\operatorname{argmin}} \quad \mathcal{L}(\mathbf{B}, \mathbf{Z}, \widehat{\mathbf{\Gamma}}) + \lambda \|\mathbf{B}\|_{1,1} \\ \text{s.t.} \quad & \mathbf{X}\mathbf{B}^\top + \mathbf{Z}\widehat{\mathbf{\Gamma}}^\top \in \mathcal{R}_C^{n \times p}, \quad \mathcal{P}_{\widehat{\mathbf{\Gamma}}}\mathbf{B} = \mathbf{0}, \end{aligned} \quad (3.3)$$

which is a convex optimization problem in variables  $\mathbf{B}$  and  $\mathbf{Z}$ . This motivates us to solve optimization problem [\(3.2\)](#) in two steps: (1) firstly obtaining a good estimate of  $\widehat{\mathbf{\Gamma}}$  and (2) then based on  $\widehat{\mathbf{\Gamma}}$ , obtaining good estimates for  $\mathbf{B}^*$  and  $\mathbf{Z}^*$ . In [Algorithm 1](#), we incorporate the two-step procedure by solving two sub-problems [\(3.4\)](#) and [\(3.5\)](#) consecutively. We next analyze the statistical properties of estimators in each step of [Algorithm 1](#).

### 3.1 Estimation of uncorrelated latent components

To estimate the marginal effects  $\mathbf{F}^* = \mathbf{B}^* + \mathbf{\Gamma}^*\mathbf{D}^*$  and the uncorrelated latent components  $\mathbf{W}^*\mathbf{\Gamma}^{*\top}$ , we first solve optimization problem [\(3.4\)](#). This is also known as the joint maximum likelihood estimation ([Chen et al., 2019, 2020](#)), which is statistically optimal in the minimax sense when both the sample size  $n$  and the response dimension  $p$  grow to infinity. From optimization problem [\(3.4\)](#), we obtain the initial estimates of the natural parameter matrix  $\widehat{\mathbf{\Theta}}_0 = \mathbf{X}\widehat{\mathbf{F}}^\top + \widehat{\mathbf{W}}_0\widehat{\mathbf{\Gamma}}_0^\top$ . The following theorem characterizes the estimation error of the initial maximum likelihood estimate  $\widehat{\mathbf{\Theta}}_0$ .

**Theorem 2** (Estimation error of  $\widehat{\mathbf{\Theta}}_0$ ). Under [Assumptions 1–3](#), let  $\widehat{\mathbf{\Theta}}_0$  be any estimator such that  $\mathcal{L}(\widehat{\mathbf{\Theta}}_0) \leq \mathcal{L}(\mathbf{\Theta}^*)$ . For any constant  $\delta > 1$ , when  $np \geq 3$ , it holds with probability at least  $1 - (n + p)^{-\delta} - (np)^{-\delta} - \iota_n$  that

$$\|\widehat{\mathbf{\Theta}}_0 - \mathbf{\Theta}^*\|_{\text{F}} = \mathcal{O}\left(\sqrt{(d+r)((n \vee p) \vee \delta^3)}\right), \quad \max_{1 \leq j \leq p} \|(\widehat{\mathbf{\Theta}}_0)_j - \mathbf{\Theta}_j^*\|_2 = \mathcal{O}\left(\sqrt{(d+r)(n \vee \delta^3)}\right).$$

In our specific setting, where the dimensions represented by  $d$  and  $r$  are orders of magnitude smaller compared to  $n$  and  $p$ , the estimation error is primarily dominated by the scale of the larger dimensions  $n$  and  $p$ . Because the dimensions of natural parameters expand with both  $n$  and  $p$ , the bound implies that the error associated with each entry is approximately

---

**Algorithm 1** GCATE (generalized confounder adjustment for testing and estimation)
 

---

**Input:** A data matrix  $\mathbf{Y} \in \mathbb{R}^{n \times p}$ , a design matrix  $\mathbf{X} \in \mathbb{R}^{n \times d}$ , a natural number  $r \geq 1$  (the number of latent factors)

- 1: **Estimation of uncorrelated latent components  $\mathbf{W}\mathbf{\Gamma}^\top$ :** Solve optimization problem (3.4) to obtain  $\widehat{\mathbf{W}}_0\widehat{\mathbf{\Gamma}}_0^\top$  and the initial estimate  $\widehat{\mathbf{\Theta}}_0 = \mathbf{X}\widehat{\mathbf{F}}^\top + \widehat{\mathbf{W}}_0\widehat{\mathbf{\Gamma}}_0^\top$  by alternative maximization (Algorithm F.3) with initialization given in Appendix F.2:

$$\begin{aligned} \widehat{\mathbf{F}}, \widehat{\mathbf{W}}_0, \widehat{\mathbf{\Gamma}}_0 \in & \underset{\mathbf{F} \in \mathbb{R}^{p \times d}, \mathbf{W} \in \mathbb{R}^{n \times r}, \mathbf{\Gamma} \in \mathbb{R}^{p \times r}}{\operatorname{argmin}} \quad \mathcal{L}(\mathbf{X}\mathbf{F}^\top + \mathbf{W}\mathbf{\Gamma}^\top) \\ \text{subject to} \quad & \mathbf{X}\mathbf{F}^\top + \mathbf{W}\mathbf{\Gamma}^\top \in \mathcal{R}_C^{n \times p}, \quad \mathcal{P}_\mathbf{X}\mathbf{W} = \mathbf{0}. \end{aligned} \quad (3.4)$$

- 2: **Estimation of latent coefficients  $\mathbf{\Gamma}$ :** Set  $\widehat{\mathbf{W}} := \sqrt{n}\mathbf{U}\mathbf{\Sigma}^{1/2}$  and  $\widehat{\mathbf{\Gamma}} := \sqrt{p}\mathbf{V}\mathbf{\Sigma}^{1/2}$ , where  $\widehat{\mathbf{W}}_0\widehat{\mathbf{\Gamma}}_0^\top = \sqrt{np}\mathbf{U}\mathbf{\Sigma}\mathbf{V}^\top$  is the condensed SVD with  $\mathbf{U} \in \mathbb{R}^{n \times r}$ ,  $\mathbf{\Sigma} \in \mathbb{R}^{r \times r}$ ,  $\mathbf{V} \in \mathbb{R}^{p \times r}$ .
- 3: **Estimation of direct effects  $\mathbf{B}$  and latent factors  $\mathbf{Z}$ :** Solve optimization problem (3.5) to obtain  $(\widehat{\mathbf{B}}, \widehat{\mathbf{Z}})$  by Algorithm F.3 with initialization given in Appendix F.2:

$$\begin{aligned} \widehat{\mathbf{B}}, \widehat{\mathbf{Z}} = & \underset{\mathbf{B} \in \mathbb{R}^{p \times d}, \mathbf{Z} \in \mathbb{R}^{p \times r}}{\operatorname{argmin}} \quad \mathcal{L}(\mathbf{X}\mathbf{B}^\top + \mathbf{Z}\widehat{\mathbf{\Gamma}}^\top) + \lambda\|\mathbf{B}\|_{1,1} \\ \text{subject to} \quad & \mathbf{X}\mathbf{B}^\top + \mathbf{Z}\widehat{\mathbf{\Gamma}}^\top \in \mathcal{R}_C^{n \times p}, \quad \mathcal{P}_{\widehat{\mathbf{\Gamma}}}\mathbf{B} = \mathbf{0}. \end{aligned} \quad (3.5)$$

- 4: **Debiasing:** Construct the debiased estimate (4.1) and its estimated variance (4.6), based on  $(\widehat{\mathbf{B}}, \widehat{\mathbf{Z}}, \widehat{\mathbf{\Gamma}})$ . Compute  $p$ -values according to the asymptotic distribution (4.5).

**Output:** Return the  $p$ -values.

---

on the order of  $\sqrt{(n \vee p)/np}$ . As demonstrated in the upcoming subsection, these results empower us to attain robust estimates of the confounding effects.

### 3.2 Estimation of latent coefficients

From optimization problem (3.4), we obtain the initial estimates  $\widehat{\mathbf{E}} = \widehat{\mathbf{W}}_0\widehat{\mathbf{\Gamma}}_0^\top$  of the latent components that are uncorrelated to the observed covariates. Though  $\mathbf{W}^*$  and  $\mathbf{\Gamma}^*$  are only identified up to rotations, we can use the condensed singular value decomposition of the normalized latent components  $\widehat{\mathbf{E}}/\sqrt{np} = \mathbf{U}\mathbf{\Sigma}\mathbf{V}^\top$  to obtain the final estimates through the following optimization problem:

$$\begin{aligned} \widehat{\mathbf{W}}, \widehat{\mathbf{\Gamma}} \in & \underset{\mathbf{W} \in \mathbb{R}^{n \times r}, \mathbf{\Gamma} \in \mathbb{R}^{p \times r}}{\operatorname{argmin}} \quad \frac{1}{np} \|\widehat{\mathbf{E}} - \mathbf{W}\mathbf{\Gamma}^\top\|_F^2 \\ \text{subject to} \quad & \frac{1}{n}\mathbf{W}^\top\mathbf{W} = \frac{1}{p}\mathbf{\Gamma}^\top\mathbf{\Gamma} \text{ is diagonal.} \end{aligned} \quad (3.6)$$

A simple derivation yields the solution  $\widehat{\mathbf{W}} = \sqrt{n}\mathbf{U}\mathbf{\Sigma}^{1/2}$  and  $\widehat{\mathbf{\Gamma}} = \sqrt{p}\mathbf{V}\mathbf{\Sigma}^{1/2}$  to the above problem. The above procedure is also commonly used in the factor analysis literature for estimating factor loadings from regression residuals (Bai, 2003; Bing et al., 2023). The following theorem guarantees that the above estimate of the latent coefficients is provably

accurate in recovering the column space of the true coefficients.

**Theorem 3** (Estimation error of  $\mathcal{P}_{\hat{\Gamma}}$ ). Under Assumptions 1–3, as  $n, p \rightarrow \infty$ , it holds that

$$\|\mathcal{P}_{\hat{\Gamma}} - \mathcal{P}_{\Gamma^*}\|_{\text{op}} = \mathcal{O}_{\mathbb{P}}\left(\sqrt{\frac{1}{n \wedge p}}\right), \quad \max_{1 \leq i, j \leq p} |(\mathcal{P}_{\hat{\Gamma}} - \mathcal{P}_{\Gamma^*})_{ij}| = \mathcal{O}_{\mathbb{P}}\left(\sqrt{\frac{1}{p^2(n \wedge p)}}\right).$$

Theorem 3 implies that the image of  $\Gamma^*$  can be estimated well by  $\hat{\Gamma}$ . Furthermore, the column-wise error decays at a fast rate. The precise error control of each column individually enables us to disentangle the intricate relationships within the confounder-adorned high-dimensional dataset. However, these do not directly extend to error control for the latent factors  $\mathbf{Z}^*$  or the latent coefficients  $\Gamma^*$  themselves.

For instance, Bing et al. (2023, Theorem 4) shows the concentration of the latent coefficients  $\max_{1 \leq j \leq p} \|\gamma_j^* - \hat{\gamma}_j\|_2$  under Gaussian linear models. Their results rely on the special structure of the regression problem  $\mathbf{Y}_j = \mathbf{X}(\mathbf{b}_j^* + \mathbf{D}^{*\top} \gamma_j^*) + \epsilon_j$ , where the regression coefficient can be decomposed into a sparse component  $\mathbf{b}_j^*$  and a dense component  $\mathbf{D}^{*\top} \gamma_j^*$ . By using the lava estimator (Chernozhukov et al., 2017), they can derive the estimation error of the residual  $\epsilon_j$ , from whose covariance structure,  $\|\gamma_j^* - \hat{\gamma}_j\|_2$  can be further bounded. In the generalized linear model setting, we don't have the flexibility to utilize the additive noises' covariance structure. However, as we illustrate in the next section, it is still possible to estimate  $\mathbf{Z}^* \gamma_j^*$ , which suffices for recovering the direct effects  $\mathbf{B}^*$ .

### 3.3 Estimation of latent factors and direct effects

Once the column space of confounding effects becomes distinguishable, the subsequent phase entails retrieving direct effects by mitigating the influence of confounding variables and solving optimization problem (3.3). Through this, we can simultaneously obtain the estimates of latent factors  $\mathbf{Z}^*$  and direct effects  $\mathbf{B}^*$ . In the high-dimensional scenarios when  $p$  can be larger than  $n$ , one natural approach to estimate the sparse coefficient  $\mathbf{B}^*$  is via  $\ell_1$ -regularization, as employed in the optimization problem (3.5), which aims to obtain a sparse and consistent estimator  $\hat{\mathbf{B}}$  by  $\ell_1$ -regularization while simultaneously removing the unmeasured effects. Next, we analyze the properties of the two estimators  $\hat{\mathbf{Z}}$  and  $\hat{\mathbf{B}}$  in turn.

**Latent factors.** As previously alluded to, estimating latent factor  $\mathbf{Z}^*$  demands special consideration. To bypass the technical difficulty, we will use the estimation errors of  $\hat{\Theta}$  and  $\mathcal{P}_{\hat{\Gamma}}$  provided by Theorem 2 and Theorem 3, respectively, coupled with one extra identifiability condition for the latent factors  $\mathbf{Z}^*$  and their coefficients  $\Gamma^*$ . From Lin et al. (2021, Proposition 5.1), there exists an invertible matrix  $\mathbf{R} \in \mathbb{R}^{r \times r}$  with bounded operator norm, such that  $\text{Var}(\mathbf{R}\mathbf{z}_1^*)$  and  $\mathbf{R}^{-\top} \gamma_1^* \gamma_1^{*\top} \mathbf{R}^{-1}$  are the same diagonal matrix. The following assumption from Lin et al. (2021) restricts the spacing of  $\text{Var}(\mathbf{R}\mathbf{z}_1^*)$ 's eigenvalues.

**Assumption 4** (Identifiability of latent factors). Assume there exists positive numbers  $c_1 \leq c_2$  and  $1 < k_1 \leq k_2$  such that for all  $\ell \in \{1, \dots, r\}$ , the eigenvalues of  $\text{Var}(\mathbf{R}\mathbf{z}_1^*)$  satisfy  $c_1 \ell^{-k_1} \leq \lambda_\ell \leq c_2 \ell^{-k_1}$ , and  $\lambda_\ell - \lambda_{\ell+1} \geq c_1 \ell^{-k_2}$ , with the convention that  $\lambda_{r+1} = 0$ .

Intuitively, Assumption 4 guarantees that  $\mathbf{Z}^* \mathbf{R}^\top$  can be recovered up to sign from the matrix product  $\mathbf{Z}^* \Gamma^{*\top}$ . This implies that, if one can consistently estimate  $\mathbf{Z}^* \Gamma^{*\top}$  with  $\hat{\mathbf{Z}} \hat{\Gamma}^\top$ ,

then  $\mathbf{Z}^*$  and  $\mathbf{\Gamma}^*$  can also be consistently estimated by appropriate transformations of  $\widehat{\mathbf{Z}}$  and  $\widehat{\mathbf{\Gamma}}$ , respectively. A simple consequence from Lin et al. (2021, Proposition 5.2), coupled with Theorems 2 and 3, is the following error bound on the columns of latent components.

**Corollary 4** (Estimation of latent components). Let  $\widehat{\mathbf{\Gamma}}$  and  $\widehat{\mathbf{Z}}$  be solutions to optimization problems (3.6) and (3.5), respectively. Under Assumptions 1–4, suppose  $\min_{\{\mathbf{B} \in \mathbb{R}^{p \times d} \mid \mathcal{P}_{\widehat{\mathbf{\Gamma}}}\mathbf{B} = \mathbf{0}\}} \mathcal{L}(\mathbf{X}\mathbf{B}^\top + \widehat{\mathbf{Z}}\widehat{\mathbf{\Gamma}}^\top) \leq \mathcal{L}(\mathbf{\Theta}^*)$  with probability tending to one. Then, as  $n, p \rightarrow \infty$ , it holds that,

$$\max_{1 \leq j \leq p} \frac{1}{\sqrt{n}} \|\widehat{\mathbf{Z}}\widehat{\boldsymbol{\gamma}}_j - \mathbf{Z}^*\boldsymbol{\gamma}_j^*\|_2 = \mathcal{O}_{\mathbb{P}} \left( \sqrt{\frac{\log n}{n}} \vee \sqrt{\frac{r^{4k_2 - k_1 + 4} \log n}{n \wedge p}} \right).$$

In Corollary 4, we require the maximum likelihood based on the estimated latent components to be higher than the likelihood evaluated at the truth. It's important to note that, unlike the analysis for linear models in prior work (Bing et al., 2022, 2023) that projects the responses onto the orthogonal column space of  $\widehat{\mathbf{\Gamma}}$  and removes the effects of latent factors  $\mathbf{Z}$ , estimating  $\mathbf{Z}$  is unavoidable under generalized linear models. Therefore, we emphasize the need for the estimated latent components derived from (3.5) to exhibit stability. This stability ensures that the maximum likelihood with the estimated latent factors remains close to the joint maximum likelihood from (3.4). In the presence of nuisance parameters  $\mathbf{Z}^*$  and  $\mathbf{\Gamma}^*$ , the sharp control on estimation error of the column  $\mathbf{Z}^*\boldsymbol{\gamma}_j^*$  provided by Corollary 4 helps control the estimation error of  $\widehat{\mathbf{B}}$ .

**Direct effects.** In high-dimensional scenarios, controlling the estimation error of  $\mathcal{P}_{\mathbf{\Gamma}^*}\mathbf{B}^*$  requires the projection  $\mathcal{P}_{\mathbf{\Gamma}^*}$  does not excessively densify the primary effects  $\mathbf{B}^*$ . To this end, we require the ratio  $\|\mathcal{P}_{\mathbf{\Gamma}^*}\mathbf{B}^*\|_{1,1}/\|\mathcal{P}_{\mathbf{\Gamma}^*}\mathbf{B}^*\|_{\text{F}}$  to be of smaller order than  $\sqrt{p}$ . Coupled with the previous assumptions and results, the estimation error of  $\widehat{\mathbf{B}}$  returned by problem (3.5) can be controlled.

**Theorem 5** (Estimation error of  $\widehat{\mathbf{B}}$ ). Suppose the assumptions in Corollary 4 hold and  $\|\mathcal{P}_{\mathbf{\Gamma}^*}\mathbf{B}^*\|_{1,1} = \mathcal{O}(p^{k/2}\|\mathcal{P}_{\mathbf{\Gamma}^*}\mathbf{B}^*\|_{\text{F}})$  for some constant  $k \in [0, 1)$ . Then, as  $n, p \rightarrow \infty$  such that  $\sqrt{n}/\log(nd) = o(p)$  and  $\log(p) = o(n)$ , the estimate  $\widehat{\mathbf{B}}$  of optimization problem (3.5) with  $\lambda \asymp 8\nu^2 \log(nd)n^{-1/2}$  satisfies that

$$\|\widehat{\mathbf{B}} - \mathbf{B}^*\|_{\text{F}} = \mathcal{O}_{\mathbb{P}}(r_{n,p}), \quad \|\widehat{\mathbf{B}} - \mathbf{B}^*\|_{1,1} = \mathcal{O}_{\mathbb{P}}(r'_{n,p}),$$

where the sequences  $r_{n,p}$  and  $r'_{n,p}$  are defined as:

$$r_{n,p} := \sqrt{\frac{(sd \log^2(nd)) \vee \log(np)}{n}} + \sqrt{\frac{n^{1/2}}{(n \wedge p)^{3/2} \log(nd)}} + \sqrt{\frac{sd}{n \wedge p^{1-k}}},$$

$$r'_{n,p} := \sqrt{sd} r_{n,p} + \frac{\sqrt{n}}{(n \wedge p) \log(nd)}.$$

In Theorem 5, the parameter  $k$  captures the deviation of the projected  $\ell_1$ -norm  $\|\mathcal{P}_{\mathbf{\Gamma}^*}\mathbf{B}^*\|_{1,1}$  from  $\|\mathbf{B}^*\|_{1,1}$ . The smaller  $k$ , the more information of  $\mathbf{B}^*$  is retained after projection, and the signal-noise-ratio is larger. In the high-dimensional scenarios when  $n < p$ , the  $\ell_1$ -norm

and  $\ell_2$ -norm of the estimation error scale in  $\mathcal{O}_{\mathbb{P}}(n^{-1/2} \wedge p^{(k-1)/2})$  when ignoring lower order factors. The appearance of the response dimension  $p$  in the denominator reflects the blessing of dimensionality; namely, having more responses than the sample size is not detrimental to consistency, provided that  $p$  does not grow exponentially larger than  $n$ .

To prove Theorem 5, we establish the (approximate) optimal condition for  $(\widehat{\mathbf{B}}, \widehat{\mathbf{Z}}, \widehat{\mathbf{\Gamma}})$  from the two-step procedure to the joint optimization problem (3.2), as shown in Lemma D.1. This relies on the optimality condition of optimization problem (3.5) and the convergence rate of  $\mathcal{P}_{\widehat{\mathbf{\Gamma}}}^{\perp}$  provided by Theorem 3. It then allows us to establish the cone condition, obtain the upper and lower bounds of the first-order approximation error of the loss function, and derive the error rate in Appendix D. Compared to double machine learning in the presence of high-dimensional nuisance parameters (Chernozhukov et al., 2018,a), our estimation procedure does not require sample splitting. To establish consistency, we only need the convergence rate of  $\max_{1 \leq j \leq p} \|\mathbf{Z}^* \boldsymbol{\gamma}_j^* - \widehat{\mathbf{Z}} \widehat{\boldsymbol{\gamma}}_j\|_2$  to be the parametric rate  $(n \wedge p)^{-1/2}$ , as shown in the proof of Theorem 5; see also Remark 5 for discussion on the connection to Neyman orthogonality. However, to derive the asymptotic distribution for inference, one may need more stringent conditions or sample splitting, as illustrated next.

## 4 Inference

### 4.1 Projected and weighted bias correction

When evaluating uncertainty in high-dimensional inference, confidence intervals and statistical hypothesis tests are required. After obtaining the initial estimate  $\widehat{\mathbf{B}}$ , we need to remove the bias caused by  $\ell_1$ -regularization to have valid inferences on the estimated coefficients. Without loss of generality, we focus on testing the coefficients of the first covariate  $b_{j1}$  for  $j = 1, \dots, p$ . We consider the following debiased estimator for each of them:

$$\widehat{b}_{j1}^{\text{de}} = \widehat{b}_{j1} + \mathbf{u}^{\top} \frac{1}{n} \sum_{i=1}^n \mathbf{x}_i (\mathbf{y}_i - A'(\widehat{\boldsymbol{\theta}}_i))^{\top} \mathbf{v}_i, \quad (4.1)$$

where  $\widehat{\boldsymbol{\Theta}} = \mathbf{X} \widehat{\mathbf{B}}^{\top} + \widehat{\mathbf{Z}} \widehat{\mathbf{\Gamma}}^{\top}$  is the estimated natural parameter matrix, and  $\mathbf{u} \in \mathbb{R}^d$  and  $\mathbf{v}_i \in \mathbb{R}^p$  are projection vectors to be specified later, such that the correction term  $\mathbf{u}^{\top} n^{-1} \sum_{i=1}^n \mathbf{x}_i (\mathbf{y}_i - A'(\widehat{\boldsymbol{\theta}}_i))^{\top} \mathbf{v}_i$  is a reasonable estimate of the bias  $b_{j1}^* - \widehat{b}_{j1}$ .

By Taylor expansion of  $A'(\theta_{ij}^*)$  at  $\widehat{\theta}_{ij} := \mathbf{x}_i^{\top} \widehat{\mathbf{b}}_j + \widehat{\mathbf{z}}_i^{\top} \widehat{\boldsymbol{\gamma}}_j$ , we have

$$A'(\theta_{ij}^*) = A'(\widehat{\theta}_{ij}) + A''(\widehat{\theta}_{ij})(\theta_{ij}^* - \widehat{\theta}_{ij}) + \frac{1}{2} A'''(\psi_{ij})(\theta_{ij}^* - \widehat{\theta}_{ij})^2,$$

for some  $\psi_{ij}$  between  $\widehat{\theta}_{ij}$  and  $\theta_{ij}^*$ . Then, the residual of the  $i$ th sample can be decomposed into three sources of errors:

$$\mathbf{y}_i - A'(\widehat{\boldsymbol{\theta}}_i) = \underbrace{\boldsymbol{\epsilon}_i}_{\text{stochastic error}} + \underbrace{\mathbf{p}_i}_{\text{remaining bias}} + \underbrace{\mathbf{q}_i}_{\text{approximation error}} \quad (4.2)$$

where the three terms of errors are given by

$$\begin{aligned}\boldsymbol{\epsilon}_i &= \mathbf{y}_i - A'(\boldsymbol{\theta}_i^*) \\ \mathbf{p}_i &= A''(\widehat{\boldsymbol{\theta}}_i) \odot (\widehat{\boldsymbol{\theta}}_i - \boldsymbol{\theta}_i^*) \\ \mathbf{q}_i &= -\frac{1}{2}[A'''(\widehat{\boldsymbol{\theta}}_{ij} + t(\boldsymbol{\theta}_{ij}^* - \widehat{\boldsymbol{\theta}}_{ij}))(\boldsymbol{\theta}_{ij}^* - \widehat{\boldsymbol{\theta}}_{ij})^2]_{1 \leq j \leq p}.\end{aligned}$$

If we let  $\mathbf{v}_i = \omega_i \text{diag}(A''(\widehat{\boldsymbol{\theta}}_i))^{-1} \mathcal{P}_{\widehat{\Gamma}}^\perp \mathbf{e}_j$  where  $\omega_i$  is the sample-specific weight and  $\mathbf{e}_j \in \mathbb{R}^p$  is the unit vector with  $j$ th entry being one, then substituting (4.2) into (4.1) yields that

$$\begin{aligned}\widehat{b}_{j1}^{\text{de}} - b_{j1}^* &= (\widehat{b}_{j1} - b_{j1}^*) + \mathbf{u}^\top \frac{1}{n} \sum_{i=1}^n \mathbf{x}_i \boldsymbol{\epsilon}_i^\top \mathbf{v}_i + \mathbf{u}^\top \frac{1}{n} \sum_{i=1}^n \mathbf{x}_i \mathbf{p}_i^\top \mathbf{v}_i + \mathbf{u}^\top \frac{1}{n} \sum_{i=1}^n \mathbf{x}_i \mathbf{q}_i^\top \mathbf{v}_i \\ &= \mathbf{u}^\top \frac{1}{n} \sum_{i=1}^n \mathbf{x}_i \boldsymbol{\epsilon}_i^\top \mathbf{v}_i + \left( \mathbf{u}^\top \frac{1}{n} \sum_{i=1}^n \omega_i \mathbf{x}_i \mathbf{x}_i^\top - \mathbf{e}_1^\top \right) (\mathbf{b}_j^* - \widehat{\mathbf{b}}_j) + \text{Rem}.\end{aligned}\quad (4.3)$$

The estimation error rates provided by Theorems 3 and 5 guarantee that  $\|\mathbf{b}_j^* - \widehat{\mathbf{b}}_j\|_1 = \mathcal{O}_{\mathbb{P}}(r'_{n,p})$  and the remaining term is  $\text{Rem} = \mathcal{O}_{\mathbb{P}}(\max_{1 \leq i \leq n} |\mathbf{u}^\top \mathbf{x}_i|^3 r_{n,p}^2)$  for  $r_{n,p}$  and  $r'_{n,p}$  defined in Theorem 5. Based on (4.3), the idea of debiasing is to choose  $\mathbf{u}$  and  $\omega_i$ 's such that the second term and the remaining term is of order  $o_{\mathbb{P}}(n^{-1/2})$ , while enabling the convergence of the average of primary stochastic errors to a normal distribution by central limit theorem.

To facilitate our theoretical analysis, suppose we split the dataset into two parts  $\mathcal{D}_1 = \{(\mathbf{x}_i, \mathbf{y}_i), 1 \leq i \leq n\}$  and  $\mathcal{D}_2 = \{(\mathbf{x}_i, \mathbf{y}_i), n+1 \leq i \leq 2n\}$ , where  $\mathcal{D}_2$  is used to obtain the estimates  $\widehat{\mathbf{B}}$  and  $\widehat{\Gamma}$ , and  $\mathcal{D}_1$  is used to remove the bias for  $\widehat{\mathbf{B}}$  induced by  $\ell_1$ -regularization. There are also latent factors  $\{\mathbf{z}_i\}_{i=1}^n$  and  $\{\mathbf{z}_i\}_{i=n+1}^{2n}$  associated with  $\mathcal{D}_1$  and  $\mathcal{D}_2$ , respectively. Further, if  $\boldsymbol{\epsilon}_i$ 's are independent of the projection vectors  $\mathbf{u}$  and  $\mathbf{v}_i$  (or equivalently  $\omega_i$ ) conditional on  $(\mathbf{x}_i, \mathbf{z}_i)$ 's and  $\mathcal{D}_2$ , then we can approximate the scaled conditional variance of the stochastic errors as:

$$\begin{aligned}\sigma_j^2 &= \text{Var} \left( \mathbf{u}^\top \frac{1}{\sqrt{n}} \sum_{i=1}^n \mathbf{x}_i \boldsymbol{\epsilon}_i^\top \mathbf{v}_i \mid \{(\mathbf{x}_i, \mathbf{z}_i)\}_{i=1}^n, \mathcal{D}_2 \right) \\ &= \mathbf{u}^\top \frac{1}{n} \sum_{i=1}^n \omega_i^2 \mathbf{x}_i \mathbf{e}_j^\top \mathcal{P}_{\widehat{\Gamma}}^\perp \text{diag}(A''(\widehat{\boldsymbol{\theta}}_i))^{-1} \text{Cov}(\boldsymbol{\epsilon}_i \mid \boldsymbol{\theta}_i^*) \text{diag}(A''(\widehat{\boldsymbol{\theta}}_i))^{-1} \mathcal{P}_{\widehat{\Gamma}}^\perp \mathbf{e}_j \mathbf{x}_i \mathbf{u} \\ &= \mathbf{u}^\top \frac{1}{n} \sum_{i=1}^n \omega_i^2 (\mathbf{e}_j^\top \mathcal{P}_{\widehat{\Gamma}}^\perp \text{diag}(A''(\widehat{\boldsymbol{\theta}}_i))^{-1} \text{diag}(A''(\boldsymbol{\theta}_i^*)) \text{diag}(A''(\widehat{\boldsymbol{\theta}}_i))^{-1} \mathcal{P}_{\widehat{\Gamma}}^\perp \mathbf{e}_j) \mathbf{x}_i \mathbf{x}_i^\top \mathbf{u} \\ &\approx \mathbf{u}^\top \frac{1}{n} \sum_{i=1}^n \omega_i \mathbf{x}_i \mathbf{x}_i^\top \mathbf{u} =: \widehat{\sigma}_j^2,\end{aligned}$$

by using a proper data-dependent weight  $\omega_i = \widehat{\omega}_i$ . Then, the projection vector  $\widehat{\mathbf{u}}$  is constructed by minimizing the variance proxy while controlling the bias and remaining terms

in (4.3):

$$\begin{aligned}
\widehat{\mathbf{u}} &\in \underset{\mathbf{u} \in \mathbb{R}^d}{\operatorname{argmin}} \mathbf{u}^\top \frac{1}{n} \sum_{i=1}^n \widehat{\omega}_i \mathbf{x}_i \mathbf{x}_i^\top \mathbf{u} \\
\text{s.t.} &\quad \left\| \frac{1}{n} \sum_{i=1}^n \widehat{\omega}_i \mathbf{x}_i \mathbf{x}_i^\top \mathbf{u} - \mathbf{e}_1 \right\|_\infty \leq \lambda_n \\
&\quad \max_{1 \leq i \leq n} |\mathbf{x}_i^\top \mathbf{u}| \leq \tau_n,
\end{aligned} \tag{4.4}$$

where  $\lambda_n \asymp \sqrt{\log(nd)/n}$  and  $\tau_n \asymp \sqrt{\log n}$ . Based on  $\widehat{\omega}_i$  and  $\widehat{\mathbf{u}}$ , the resulting bias-corrected estimator (4.1) is similar to those used by van de Geer et al. (2014); Javanmard and Montanari (2014); Cai et al. (2021); however, we need to incorporate information from multiple responses with projection operator  $\mathcal{P}_{\widehat{\mathbf{\Gamma}}}^\perp$  to de-confound, in the spirit of proximal gradient descent. Under mild regularity conditions, the following theorem shows that the debiased estimator  $\widehat{b}_{j1}^{\text{de}}$  is asymptotically normal.

**Theorem 6** (Asymptotical normality of  $\widehat{\mathbf{B}}^{\text{de}}$ ). Under the same conditions in Theorem 5, for  $j = 1, \dots, p$ , additionally assume the following conditions hold: (i)  $n/\log(nd) = o(p^{3/2})$  and  $n = o(p^{2(1-k)})$ ; and (ii)  $\widehat{\omega}_i = \omega(\mathbf{x}_i, \mathbf{z}_i, \mathcal{D}_2)$  for some real-valued function  $\omega$  that is uniformly bounded away from 0 and  $\infty$ . Then as  $n, p \rightarrow \infty$ , it holds that

$$\sqrt{n} \frac{\widehat{b}_{j1}^{\text{de}} - b_{j1}^*}{\sigma_j} \xrightarrow{d} \mathcal{N}(0, 1). \tag{4.5}$$

With fewer assumptions on the correlation between the covariate  $\mathbf{X}$  and the confounder  $\mathbf{Z}^*$ , removing unmeasured confounders is only possible by utilizing multiple outcomes to disentangle the primary effect  $\mathbf{B}^*$  and the latent coefficient  $\mathbf{\Gamma}^*$ . In particular, because the estimation error rates of  $\mathbf{B}^*$  and  $\mathbf{\Gamma}^*$  are related to  $(n \wedge p)^{-1}$ , the number of outcomes  $p$  is expected to be larger than  $n$ , so that these errors are primarily affected by the sample size  $n$ .

In Theorem 6, condition (i) requires that the response dimension  $p$  grows faster than  $n^{2/3} \vee n^{1/(2(1-k))}$ , which ensures the remainder term Rem in (4.3) vanishes in the limit. Specifically, Rem a magnitude associated with the convergence rate of  $\|\widehat{\mathbf{B}} - \mathbf{B}^*\|_{\mathbb{F}}^2$ , as provided by Theorem 5. To derive the asymptotic normality,  $\text{Rem} = o_{\mathbb{P}}(n^{-1/2})$  is required; however, if  $n$  is too large compared to  $p$ , the convergence rate of  $\widehat{\mathbf{B}}$  from the first two steps of the proposed procedure is insufficient to establish the desired asymptotic normality. In this case, having a much larger sample size does not help. When  $k \leq 1/2$ , condition (i) is satisfied with  $n = o(p)$ , which is reasonable in most scientific scenarios of cohort-level differential expression analysis, as we shall see later from the real data example in Section 6.

In terms of condition (ii), a proper sample-specific and link-specific weight function is required. One can construct such weights  $\widehat{\omega}_i$  by sample splitting to fulfill this condition. For instance, using sample splitting procedure in Algorithm E.2, one valid choice is  $\widehat{\omega}_i = A''(\widehat{\theta}_{ij})$ . In Lemma E.3, we show that such a choice of  $\widehat{\omega}_i$  satisfies the condition (ii) in Theorem 6

with probability tending to one and the resulting variance estimator

$$\widehat{\sigma}_j^2 = \widehat{\mathbf{u}}^\top \frac{1}{n} \sum_{i=1}^n \widehat{\omega}_i \mathbf{x}_i \mathbf{x}_i^\top \widehat{\mathbf{u}}, \quad (4.6)$$

is also consistent with  $\sigma_j^2$ . Hence, Theorem 6 implies that  $t_j = \sqrt{n}(\widehat{b}_{j1}^{\text{de}} - b_{j1}^*)/\widehat{\sigma}_j \xrightarrow{d} \mathcal{N}(0, 1)$ . We reject the null hypothesis  $H_{0j} : b_{j1}^* = 0$  at level- $\alpha$  if  $|t_j| > z_{\alpha/2} := \Phi^{-1}(1 - \alpha/2)$ , where  $\Phi$  is the cumulative distribution function of the standard normal. Numerically, we show that the efficiency loss of sample splitting is negligible and similar to the result without sample splitting in Appendix G.1; and the proposed method performs well without sample splitting and is statistically more efficient than the alternative methods in Section 5.

**Remark 2** (Inference without unmeasured confounders). In the special case when there are no unmeasured confounders, the matrix  $\mathcal{P}_{\mathbf{F}}^\perp$  reduces to the identity matrix. Also, the projection vector  $\widehat{\mathbf{u}}$  is the  $j$ th column of  $(\mathbf{X}^\top \text{diag}(A''(\mathbf{X}\widehat{\mathbf{B}}_1))\mathbf{X})^{-1}$ , and  $\widehat{\sigma}_j$  is the asymptotic variance of  $\widehat{b}_{j1}$  under well-specified generalized linear models. In this case, (4.1) is simply a one-step adjustment based on the score function. For Bernoulli distributed binary outcomes without unmeasured confounders, the above choice of the weight function  $\omega(\theta) = A''(\theta)$  for optimization problem (4.4) coincide with  $f'(\theta)^2/(f(\theta)(1 - f(\theta)))$ , the one used in Cai et al. (2021) with  $f = A'$  being the link function.

**Remark 3** (Incorporate information from latent factors). In (4.1) and (4.4), we only use the covariate  $\mathbf{X}$  to adjust for the estimation bias. However, including the estimated latent factors  $\mathbf{Z}$  to construct a projection vector  $\mathbf{u}$  of dimension  $d + r$  is also feasible. The validity of this extension is also guaranteed by the sample splitting procedure in Algorithm E.2.

**Remark 4** (Estimation and inference with non-canonical links). Through Sections 3 and 4, we discuss the methodology to conduct inference on confounded generalized linear models (GLM) with canonical link functions, as outlined in Table F1. However, in practical scenarios, non-canonical link functions may also be employed. For instance, the log link function is commonly used with Negative Binomial GLMs. Fortunately, our method extends its applicability to GLMs with non-canonical link functions, as exemplified in the case of the Negative Binomial GLMs in Appendix F.4. Establishing theoretical guarantees for these scenarios may follow a similar framework with suitable assumptions to address the non-convexity of the objective functions, as elaborated in Appendix F.4.

## 4.2 Simultaneous inference

The asymptotic normality provided in Theorem 6 provides Type-I error controls for individual hypothesis tests  $H_{0j} : b_{j1}^* = 0$  for  $j = 1, \dots, p$ . The following proposition shows that we can also control the overall Type-I error and family-wise error rate (FWER) using the statistics  $t_j = \sqrt{n}(\widehat{b}_{j1}^{\text{de}} - b_{j1}^*)/\widehat{\sigma}_j$  for  $j = 1, \dots, p$ .



**Proposition 7** (Simultaneous inference). Let  $\mathcal{N}_p = \{j \mid b_{j1}^* = 0, j = 1, \dots, p\}$  be the true null hypotheses. Under the assumptions of Theorem 6, as  $n, p, |\mathcal{N}_p| \rightarrow \infty$ , it holds that

$$\frac{1}{|\mathcal{N}_p|} \sum_{j \in \mathcal{N}_p} \mathbb{1}\{|t_j| > z_{\frac{\alpha}{2}}\} \xrightarrow{P} \alpha, \quad \text{and} \quad \limsup \mathbb{P} \left( \sum_{j \in \mathcal{N}_p} \mathbb{1}\{|t_j| > z_{\frac{\alpha}{2p}}\} \geq 1 \right) \leq \alpha.$$

When  $p$  is large, controlling for the false discovery rate (FDR) is more desirable when performing simultaneous testing. In that regard, Cai et al. (2021, Section 2.3) provides insights on FDR controls using different techniques. From simulations in Section 5, we also show that FDR is usually well controlled by the Benjamini–Hochberg procedure empirically.


## 5 Numerical experiments

DE and related tests are frequently performed in two distinct settings in the genomic field. One relies on counts of gene expression to contrast the expression of each gene in case versus control observations. Typically, observations are either samples from RNA sequencing (RNA-seq) (Love et al., 2014) or pseudo-bulk cells obtained from single-cell sequencing by aggregating the expressions of single cells in the same homogeneous groups (Squair et al., 2021). Another setting is single-cell RNA-sequence (scRNA-seq) CRISPR screening (Dixit et al., 2016; Barry et al., 2023), where the fundamental task is to test for association between a designed genetic perturbation and gene expression (Dixit et al., 2016). In both settings, the measured gene expression is often assumed to approximately follow a Poisson or Negative Binomial (NB) distribution (Sarkar and Stephens, 2021). However, in the former, the mean expression per sample is much larger due to molecular design, and the distribution is often approximated by a normal distribution with an appropriate transformation. In the latter case, the observational unit is a single cell. Hence, the mean of the gene expression is near zero, and the data is not well approximated with a normal distribution.

Before we turn to the simulation details, we present a simulated bulk-cell dataset and a simulated single-cell dataset corresponding to the above two distinct scenarios, respectively (Figure 2). The Poisson distribution can often model the former scenario, while the NB distribution is a better option for the latter because the counts are sparser and typically exhibit strong overdispersion (Figure 2(a)-(b)). Furthermore, for single-cell data, the lower-expressed genes are typically more dispersed, and this feature is captured in our simulated data set (Figure 2(c)). In practice, both Poisson and NB models are available for analysis of either type of experiment; however, to simplify exposition, we use a Poisson distribution for bulk samples in Section 5.1 and a NB distribution for single-cell samples in Section 5.2. In the subsequent experiments, we adhere to the protocol described in Appendix F.3 for selecting both the hyperparameters and the number of factors pertinent to the proposed methods.

### 5.1 Well-specified simulated datasets


We simulate expression data  $\mathbf{Y}$  that consists of  $n \in \{100, 250\}$  cells and  $p = 3,000$  genes based on the Poisson likelihood with natural parameter  $\Theta$ . More specifically, we generate



**Figure 2:** Overview of the simulated data. **(a)** The first and second rows show the summary of one simulated dataset for bulk cells (Poisson) in Section 5.1 and single cells (Negative Binomial) by Splatter in Section 5.2, respectively. The first column shows the overall distribution of the generated counts; the second column shows the estimated dispersion parameters by methods of moments using the mean estimates from GLM with Poisson likelihood. **(b)** The proportions of zero and non-zero counts in the two datasets, colored in orange and blue, respectively. **(c)** The estimated dispersion parameter versus the estimated mean for the simulated single-cell dataset.

the covariate  $x_1$  to be a centered binary variable, i.e.,  $(x_1 + 1)/2 \sim \text{Bernoulli}(0.5)$ . We also include an intercept  $x_2 = 1$ , so that the covariate vector  $\mathbf{x} = [x_1, x_2]^\top$  has dimension  $d = 2$ . To allow for the most general confounding scenarios without assuming causal relationships as in Figure 1, we directly generate the latent factor matrix using  $\mathbf{Z} = \mathbf{X}\mathbf{D}^\top + \mathbf{W} \in \mathbb{R}^{n \times r}$  with the number of latent factors being  $r \in \{2, 10\}$ . Here, to generate  $\mathbf{D}$  and  $\mathbf{W}$ , we first sample their entries independently from  $\mathcal{N}(0, 1)$  and further modify the singular values to be  $s_1, \dots, s_r$  where  $s_k = a \cdot (2 - (k - 1)/(r - 1))$ , with  $a = n^{-3/2}$  for  $\mathbf{D}$  and  $a = (n/2)^{1/2}$  for  $\mathbf{W}$ . For the latent loading matrix  $\mathbf{\Gamma}$ , we follow Wang et al. (2017a) to take  $\mathbf{\Gamma} = \tilde{\mathbf{\Gamma}}\mathbf{\Lambda}$  where  $\tilde{\mathbf{\Gamma}}$  is a  $p \times r$  orthogonal matrix sampled uniformly from the set of all  $p \times r$  orthogonal matrix and  $\mathbf{\Lambda} = (p/2)^{1/2} \text{diag}(\lambda_1, \dots, \lambda_r)$  where  $\lambda_k = 2 - (k - 1)/(r - 1)$ . The primary effect of  $x_1$  on gene  $j$  is sampled from  $(b_{j1} + 0.2)/0.4 \sim \text{Bernoulli}(0.5)$  with probability 0.05 and set to be zero with probability 0.95. The coefficient for the intercept is set to be  $b_{j2} = 0.5$ .

Four methods are applied to the simulated datasets: (1) CATE (confounder adjustment for testing and estimation), which is a unified approach for surrogate variable analysis under linear models (Wang et al., 2017a) and operates on the log-normalized data. (2) GLM-naive, which fits generalized linear models with the Poisson likelihood method but only uses the measured covariates  $\mathbf{X}$  without adjusting for unmeasured confounders; (3) GLM-oracle, which fits generalized linear models with the Poisson likelihood method and uses both observed and unobserved covariates  $(\mathbf{X}, \mathbf{Z})$  for estimation and testing; (4) GCATE, our proposed method with the Poisson likelihood. For CATE, we use bi-cross-validation (BCV) (Owen and Wang, 2016) to select the number of factors, as suggested in their original paper (Wang et al.,




**Figure 3:** The Type-I errors and false discovery proportions (FDPs) of different methods on the simulated datasets over 100 runs, with varying numbers of samples  $n \in \{100, 250\}$  and numbers of latent factors  $r \in \{2, 10\}$ . For GLM, the maximum values of Type-I errors and FDPs are clipped at 0.1 and 0.5, respectively. The blue dashed lines indicate the desired cutoffs.


2017a). For GCATE, we use JIC described in Remark 1 to select the number of factors.

To evaluate different methods, we summarize the type-I error and FDP (false discovery proportion) after the Benjamini-Hochberg procedure in Figure 3, where the desirable thresholds for the two are set to be 5% and 20%, respectively. From Figure 3, we see that when the sample size  $n$  is small, or the latent factor dimension  $r$  is large, the performance of all methods gets slightly worse, especially those that are misspecified, which is expected. In all setups of  $(n, r)$ , because the multivariate-Gaussian assumptions of CATE are violated, it does not provide proper Type-I error control and FDP control. This suggests that CATE may inflate test statistics and cause anti-conservative inference. Similarly, GLM-naive also fails to control the FDPs because it cannot account for dependencies induced by the latent factors. On the other hand, GCATE performs as well as GLM-oracle that has knowledge of the latent factors  $\mathbf{Z}$ . This indicates that our modeling helps to accurately remove unwarranted sources of confounding effects. Note that variations of CATE may yield improved performance using empirical nulls or negative controls, but GCATE requires no such tuning.

We further inspect the FDP control of different methods with varying thresholds. In the ideal scenarios, FDP aligns closely with the specified  $\alpha$  cutoffs. From Figure 4, GLM-oracle has FDP aligning closely with the specified  $\alpha$  cutoffs and consistently performs admirably across different levels of confounding effects. Conversely, the GLM-naive approach struggles to control the FDP effectively, and this discrepancy becomes increasingly pronounced as the number of latent factors grows. However, in a commendable contrast to CATE, our method GCATE consistently outperforms in terms of FDP control at various alpha cutoffs. This superiority can be attributed to our method’s ability to model the data distribution




**Figure 4:** False discovery proportion at different  $\alpha$  levels for  $p$ -values adjusted by the Benjamini-Hochberg procedure on 100 simulated datasets when  $n = 250$ . The left and right panels show the results for different numbers of latent factors, **(a)**  $r = 2$  and **(b)**  $r = 10$ , respectively. When  $r = 10$ , the FDP of GLM-naive is above 0.15; hence it is not shown in the figure.



**Figure 5:** The powers and precisions of different methods on the simulated datasets over 100 runs, with varying numbers of samples  $n \in \{100, 250\}$  and numbers of latent factors  $r \in \{2, 10\}$ .

accurately and eliminate unwarranted variations.

Lastly, we also evaluate the statistical power and precision of different methods. Here, the power is evaluated when the Type-I error threshold is 5%. We anticipate that both CATE and the GLM-naive approach would yield higher power compared to other methods because they tend to allow more discoveries without adequately controlling the Type-I errors (Figure 3). In Figure 5, we observe that CATE exhibits the lowest power among the considered methods. In contrast, the GLM-naive approach concurrently registers the most insufficient precision among all the methods. As anticipated, the GLM-oracle approach boasts the highest power and



**Figure 6:** Simulation results on 100 simulated scRNA-seq datasets generated by Splatter with varying numbers of samples  $n \in \{100, 200\}$ . The four metrics are shown in four columns, respectively. The blue dashed lines indicate the desired cutoffs for the statistical errors.

precision because it operates in an ideal scenario without confounding effects. In contrast, our proposed method, GCATE, demonstrates a balanced and robust performance concerning power and precision. It achieves a competitive power level while maintaining a significantly higher precision than the GLM-naive method. Moreover, GCATE outperforms CATE regarding both power and precision. This suggests that correct modeling of confounding effects boosts the statistical power and precision in high-dimensional datasets.

## 5.2 Misspecified simulated datasets using scRNA simulators

To better evaluate the performance of various methods, we use the single-cell RNA sequencing data simulator Splatter (Zappia et al., 2017) to generate simulated count datasets. Splatter explicitly models the hierarchical Gamma-Poisson processes that give rise to data observed in scRNA-seq experiments and can model the multiple-faceted variability. Thus, the simulated datasets generated by Splatter are similar to real-world datasets and suitable for benchmarking differential expression testing methods.

Using Splatter,  $n$  cells are sampled from two groups with equal probability for  $n \in \{100, 200\}$ , containing  $p = 10,000$  genes. Because of the sparse nature of the simulated single-cell datasets, about 80% of the genes are only expressed in 10 cells. Hence, we exclude these lowly-expressed genes and evaluate the methods for the remaining genes. We include  $d = 3$  covariates for each cell: the intercept, the group indicator ( $\{\pm 1\}$ ), and the logarithm of the library sizes, which is the sum of expression across all genes. When simulating the datasets, we use Splatter to generate four batches, introducing three major confounders. Because the data is not generated from well-specified GLMs, the oracle model is unknown and hence not included. For GLM-naive and GCATE, we use the NB likelihood with log links

to directly model the count data, where the gene-level dispersion parameters are estimated by the method of moments based on the estimated mean returned by using the Poisson likelihood; see Appendix F.2 for more details. For CATE, we normalize the counts in each cell by its library size, then multiply them by a scale factor of  $10^4$  and shift them by one, and finally, apply the logarithm transform, following the standard preprocessing approach of single-cell data.

Compared to the previous bulk-cell simulation in Section 5.1, the simulated data from Splatter is sparser and more noisy. From Figure 6, both CATE and GLM fail to control the Type-I error at level 5% and have lower power than GCATE in this more challenging setting. The primary reason lies in the assumption underlying CATE is significantly violated, while the GLM approach fails to account for confounding effects. Though GLM may have reasonable control over the false discoveries, its power and precision are highly affected by the confounders. On the contrary, GCATE obtain valid Type-I error and FDP controls and higher power and precision with small sample sizes because of proper distributional modeling.

## 6 Lupus data example


### 6.1 The dataset

Systemic lupus erythematosus (SLE) is an autoimmune disease predominantly affecting women and individuals of Asian, African, and Hispanic descent. Perez et al. (2022) developed multiplexed single-cell RNA sequencing (mux-seq) to capture the complexity of immune cell populations and systematically profile the composition and transcriptional states of immune cells in a large multiethnic cohort. The dataset contains 1.2 million peripheral blood mononuclear cells from 8 major cell types and 261 individuals, including 162 SLE cases and 99 healthy controls of either Asian or European ancestry. The cell-type-specific DE analysis aims to provide insights into the diagnosis and treatment of SLE.

To remove the genes with small variations, we use the Python package `scanpy` (Wolf et al., 2018) to pre-process the single-cell data and select the top 2,000 highly variable genes (HVGs) within each cell type. For each cell type, we aggregate expression across cells from the same subject to obtain gene-level pseudo-bulk counts and then remove genes expressed in less than 10 subjects. The basic information of the preprocessed datasets is provided in Appendix G.2. For each subject, the recorded variables are SLE status (condition), the logarithm of the library size, sex, population, and processing cohorts (4 levels). The latter 3 variables, which account for  $r = 5$  degrees of freedom, are considered to be the measured confounders. To simulate unmeasured confounders, we exclude these variables in all our analyses, except the so-called GLM-oracle analysis.

### 6.2 Confounder adjustment

We conducted five analyses for each cell type. Three approaches are based on the NB GLM model: GLM-oracle (NB including the confounders), GLM-naive (NB ignoring the confounders) and GCATE. Two versions of CATE were applied: the basic CATE analysis and CATE-mad, which uses an estimated empirical null (Wang et al., 2017a) based on median



**Figure 7:** Results on the lupus datasets. **(a)** Histograms of lupus  $z$ -statistics of different methods on T4 cell type. The orange curves represent the density of the standard normal distribution. **(b)** The precision and specificity for four methods computed across 5 major cell types.

absolute deviation (MAD). Only 5 cell types (T4, cM, B cell, T8, NK) contain more than 50,000 single cells and have sufficient power to obtain significant findings using the GLM-oracle approach, so we restrict our comparisons to those types. In particular, we display our results for the largest T4 cell type in this section, and similar results for other cell types are included in Appendix G. To estimate the number of latent factors  $r$ , we analyze the JIC values according to Remark 1. As shown in Figure G4, the scree plot reveals a diminishing negative log-likelihood with increasing  $r$ , which plateaus for  $r = 4$  to  $r = 7$ , and the decrement becomes marginal beyond  $r = 7$ . Consequently, we recommend selecting  $r = 7$  for GCATE analysis.

The majority of the test statistics obtained for GLM-oracle are well approximated by a standard normal distribution, which suggests that the experiment conducted by Perez et al. (2022) was well controlled, and the impact of unmeasured confounders was negligible (Figure 7(a)). However, when we excluded the measured confounders, the GLM-naive statistics were poorly calibrated, indicating that controlling for these variables is essential to proper analysis, either directly or indirectly. The CATE statistics are even more poorly calibrated than GLM-naive, suggesting that these sparse data cannot be modeled using a linear model, though restricting the test to the top 250 HVGs yields test statistics closer to the expected distribution (Figure G7). With the empirical null adjustment, CATE-mad performed somewhat better, but this adaptation is insufficient, suggesting that CATE cannot remove the confounding effects when the data are unsuitable for a linear model. Finally, the performance of GCATE is ideal: the majority of the statistics are well approximated by the standard normal, and a few signals can be captured on the right tail. Similar results were obtained for each of the 5 biggest cell types, as shown in Figure G6.

For comparison, we label genes based on the GLM-oracle analysis with FDR control at cutoff 0.2 as “true positives”, resulting in 72 significant genes for the T4 cell type. With FDR control at cutoff 0.2, 15 of the 16 GCATE’s statistics overlap with the true positives,

indicating that the test loses power when the impact of confounders has to be removed using factor analysis. Still, the test appears to control the error rate. To illustrate the performance of the 4 competing analysis methods across all 5 large cell types, we calculate the precision and specificity using 0.2 as a cutoff for false discovery rate control. As shown in Figure 7(b), only GCATE achieves uniformly high precision and specificity.

To compare different methods in the biological significance of the discoveries, we conduct gene ontology over-representation analysis to identify the related biological processes. As shown in Figure G8, both GLM-oracle and GCATE discover genes that are pertinent to the immune-response-related pathways, which also appear in prior studies on lupus (Perez et al., 2022, Fig. 3). On the other hand, though hundreds of significant genes are claimed by CATEmad, they are not associated with meaningful biological pathways. The results indicate that GCATE identifies scientifically more relevant genes than CATE under unmeasured confounders.

Lastly, we inspect the sensitivity of GCATE to the number of latent factors  $r$ . By utilizing JIC (3.1), we have selected  $r = 7$ , which is close to the number of major covariates we drop. In Table G4, we examine the performance of GCATE for different values of  $r$ . Remarkably, the resulting distributions of  $z$ -statistics generated by GCATE, across varying numbers of factors  $r$ , are similar to the standard normal distribution when  $r \geq 4$  because the MAD is close to one. Thus, JIC can serve as a valuable criterion for determining the appropriate number of latent factors for GCATE. Furthermore, it is noteworthy that the number of discoveries remains consistent when  $r$  falls within a reasonable range. These observations collectively suggest the stability of GCATE’s inferential outcomes within this range of reasonable factor selections.

## 7 Discussion

We presented novel estimation and inference procedures for multivariate generalized linear models with unmeasured confounders in the high-dimensional scenarios when both the sample size  $n$  and response size  $p$  tend to infinity. Our approach comprises three main phases. In the first phase, we disentangle the marginal effects from the uncorrelated confounding effects, recovering the column space of latent coefficients  $\widehat{\Gamma}$  from the latter. We provide non-asymptotic estimation error bounds for both the estimated natural parameter matrix  $\widehat{\Theta}$  and the projection onto the column space of  $\widehat{\Gamma}$ . In the second phase, we estimate both latent factors  $\mathbf{Z}$  and primary effects  $\mathbf{B}$  by solving a constrained lasso-type problem that confines  $\mathbf{B}$  to the orthogonal space of  $\widehat{\Gamma}$ . From the column-wise estimation error of the latent components, we obtain the estimation error for the primary effects in the presence of nuisance parameters. In the third phase, we design an inferential procedure to correct the bias introduced by  $\ell_1$ -regularization and establish Type-I error and family-wise error rate controls.

Numerically, we demonstrate the usage of the proposed method with Poisson and Negative Binomial likelihoods for bulk-cell and single-cell simulations, respectively. Compared to alternative methods, the proposed method effectively controls the Type-I error and false discovery proportion while delivering enhanced statistical power and precision as the count data get sparser and more over-dispersed. Furthermore, our analysis of real single-cell datasets underscores the essential nature of accounting for confounding effects when major covariates



are unobserved. Notably, our proposed method consistently outperforms alternative techniques, demonstrating superior precision and specificity, thus establishing its suitability for high-dimensional sparse count data.

The present study, while offering valuable insights, is not without its limitations and opportunities for future exploration. Some of these include the development of hypothesis testing for confounding effects, the theoretical guarantee of the FDR, and more robust criteria for selecting the optimal number of latent factors. Recent works by [Dai et al. \(2023\)](#) and [Chen and Li \(2022\)](#) offer promising insights that may contribute to resolving some of these challenges. Although we have briefly touched upon the applicability of our proposed method under non-canonical link functions in [Appendix F.4](#), comprehensive theoretical guarantees remain an area deserving of further research and investigation.

## Acknowledgement

We extend our heartfelt appreciation to Kevin Lin for graciously sharing their meticulously prepared manuscript on cohort eSVD, which has proven to be an indispensable asset in the advancement of our current research endeavor. We are indebted to Pratik Patil, Catherine Wang, F. William Townes, and all other members of the CMU GenStats Lab Group for suggestions on the first draft and many insightful conversations. This work used the Bridges-2 system at the Pittsburgh Supercomputing Center (PSC) through allocations BIO220140 and MTH230023 from the Advanced Cyberinfrastructure Coordination Ecosystem: Services & Support (ACCESS) program. This project was funded by the National Institute of Mental Health (NIMH) grant R01MH123184.

## References

- Bai, J. (2003). Inferential theory for factor models of large dimensions. *Econometrica*, 71(1):135–171.
- Bai, J. and Li, K. (2012). Statistical analysis of factor models of high dimension. *The Annals of Statistics*, pages 436–465.
- Barry, T., Mason, K., Roeder, K., and Katsevich, E. (2023). Robust differential expression testing for single-cell crispr screens at low multiplicity of infection. *bioRxiv*.
- Bing, X., Cheng, W., Feng, H., and Ning, Y. (2023). Inference in high-dimensional multivariate response regression with hidden variables. *Journal of the American Statistical Association*, pages 1–12.
- Bing, X., Ning, Y., and Xu, Y. (2022). Adaptive estimation in multivariate response regression with hidden variables. *The Annals of Statistics*, 50(2):640–672.
- Cai, T. T., Guo, Z., and Ma, R. (2021). Statistical inference for high-dimensional generalized linear models with binary outcomes. *Journal of the American Statistical Association*, pages 1–14.

- Ćevid, D., Bühlmann, P., and Meinshausen, N. (2020). Spectral deconfounding via perturbed sparse linear models. *The Journal of Machine Learning Research*, 21(1):9442–9482.
- Chen, Y. and Li, X. (2022). Determining the number of factors in high-dimensional generalized latent factor models. *Biometrika*, 109(3):769–782.
- Chen, Y., Li, X., and Zhang, S. (2019). Joint maximum likelihood estimation for high-dimensional exploratory item factor analysis. *Psychometrika*, 84:124–146.
- Chen, Y., Li, X., and Zhang, S. (2020). Structured latent factor analysis for large-scale data: Identifiability, estimability, and their implications. *Journal of the American Statistical Association*, 115(532):1756–1770.
- Cheng, J., Lin, G., Wang, T., Wang, Y., Guo, W., Liao, J., Yang, P., Chen, J., Shao, X., Lu, X., Zhu, L., Wang, Y., and Fan, X. (2023). Massively parallel CRISPR-based genetic perturbation screening at single-cell resolution. *Adv Sci (Weinh)*, 10(4):e2204484.
- Chernozhukov, V., Chetverikov, D., Demirer, M., Duflo, E., Hansen, C., Newey, W., and Robins, J. (2018a). Double/debiased machine learning for treatment and structural parameters.
- Chernozhukov, V., Hansen, C., and Liao, Y. (2017). A lava attack on the recovery of sums of dense and sparse signals. *The Annals of Statistics*, 45(1):39–76.
- Chernozhukov, V., Nekipelov, D., Semenova, V., and Syrgkanis, V. (2018b). Plug-in regularized estimation of high-dimensional parameters in nonlinear semiparametric models. *arXiv preprint arXiv:1806.04823*.
- Dai, C., Lin, B., Xing, X., and Liu, J. S. (2023). A scale-free approach for false discovery rate control in generalized linear models. *Journal of the American Statistical Association*, pages 1–15.
- Dixit, A., Parnas, O., Li, B., Chen, J., Fulco, C. P., Jerby-Arnon, L., Marjanovic, N. D., Dionne, D., Burks, T., Raychowdhury, R., et al. (2016). Perturb-Seq: dissecting molecular circuits with scalable single-cell RNA profiling of pooled genetic screens. *cell*, 167(7):1853–1866.
- Editorial (2023). A focus on single-cell omics. *Nat Rev Genet*, 24(8):485.
- Feng, Y. (2020). Causal inference in possibly nonlinear factor models. *arXiv preprint arXiv:2008.13651*.
- Gagnon-Bartsch, J. A. and Speed, T. P. (2012). Using control genes to correct for unwanted variation in microarray data. *Biostatistics*, 13(3):539–52.
- Gerard, D. and Stephens, M. (2020). Empirical Bayes shrinkage and false discovery rate estimation, allowing for unwanted variation. *Biostatistics*, 21(1):15–32.
- Guo, Z., Ćevid, D., and Bühlmann, P. (2022). Doubly debiased lasso: High-dimensional inference under hidden confounding. *The Annals of Statistics*, 50(3):1320.

- Hong, D. and Iakoucheva, L. M. (2023). Therapeutic strategies for autism: targeting three levels of the central dogma of molecular biology. *Transl Psychiatry*, 13(1):58.
- Javanmard, A. and Montanari, A. (2014). Confidence intervals and hypothesis testing for high-dimensional regression. *The Journal of Machine Learning Research*, 15(1):2869–2909.
- Jiang, K. and Ning, Y. (2022). Treatment effect estimation with unobserved and heterogeneous confounding variables. *arXiv preprint arXiv:2207.14439*.
- Kampmann, M. (2020). Crispr-based functional genomics for neurological disease. *Nat Rev Neurol*, 16(9):465–480.
- Lee, S., Sun, W., Wright, F. A., and Zou, F. (2017). An improved and explicit surrogate variable analysis procedure by coefficient adjustment. *Biometrika*, 104(2):303–316.
- Leek, J. T., Scharpf, R. B., Bravo, H. C., Simcha, D., Langmead, B., Johnson, W. E., Geman, D., Baggerly, K., and Irizarry, R. A. (2010). Tackling the widespread and critical impact of batch effects in high-throughput data. *Nat Rev Genet*, 11(10):733–9.
- Leek, J. T. and Storey, J. D. (2008). A general framework for multiple testing dependence. *Proceedings of the National Academy of Sciences*, 105(48):18718–18723.
- Lin, K. Z., Lei, J., and Roeder, K. (2021). Exponential-family embedding with application to cell developmental trajectories for single-cell RNA-seq data. *Journal of the American Statistical Association*, 116(534):457–470.
- Love, M. I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, 15(12):550.
- McKenna, C. and Nicolae, D. (2019). Accounting for unobserved covariates with varying degrees of estimability in high-dimensional biological data. *Biometrika*, 106(4):823–840.
- McKenna, C. and Nicolae, D. (2022). Estimating and accounting for unobserved covariates in high-dimensional correlated data. *Journal of the American Statistical Association*, 117(537):225–236.
- Ouyang, J., Tan, K. M., and Xu, G. (2023). High-dimensional inference for generalized linear models with hidden confounding. *Journal of Machine Learning Research*, 24(296):1–61.
- Owen, A. B. and Wang, J. (2016). Bi-cross-validation for factor analysis. *Statistical Science*, pages 119–139.
- Perez, R. K., Gordon, M. G., Subramaniam, M., Kim, M. C., Hartoularos, G. C., Targ, S., Sun, Y., Ogorodnikov, A., Bueno, R., Lu, A., et al. (2022). Single-cell RNA-seq reveals cell type-specific molecular and genetic associations to lupus. *Science*, 376(6589):eabf1970.
- Salim, A., Molania, R., Wang, J., De Livera, A., Thijssen, R., and Speed, T. P. (2022). Ruv-iii-nb: Normalization of single cell rna-seq data. *Nucleic Acids Research*, 50(16):e96–e96.

- Sarkar, A. and Stephens, M. (2021). Separating measurement and expression models clarifies confusion in single-cell RNA sequencing analysis. *Nature genetics*, 53(6):770–777.
- Squair, J. W., Gautier, M., Kathe, C., Anderson, M. A., James, N. D., Hutson, T. H., Hudelle, R., Qaiser, T., Matson, K. J. E., Barraud, Q., Levine, A. J., La Manno, G., Skinnider, M. A., and Courtine, G. (2021). Confronting false discoveries in single-cell differential expression. *Nat Commun*, 12(1):5692.
- Sun, Y., Ma, L., and Xia, Y. (2023). A decorrelating and debiasing approach to simultaneous inference for high-dimensional confounded models. *Journal of the American Statistical Association*, pages 1–12.
- Sun, Y., Zhang, N. R., and Owen, A. B. (2012). Multiple hypothesis testing adjusted for latent variables, with an application to the agemap gene expression data. *The Annals of Applied Statistics*, pages 1664–1688.
- van de Geer, S., Bühlmann, P., Ritov, Y., and Dezeure, R. (2014). On asymptotically optimal confidence regions and tests for high-dimensional models. *The Annals of Statistics*, 42(3):1166–1202.
- Wang, J., Zhao, Q., Hastie, T., and Owen, A. B. (2017). Confounder adjustment in multiple hypothesis testing. *The Annals of Statistics*, 45(5):1863.
- Wolf, F. A., Angerer, P., and Theis, F. J. (2018). SCANPY: large-scale single-cell gene expression data analysis. *Genome biology*, 19:1–5.
- Zappia, L., Phipson, B., and Oshlack, A. (2017). Splatter: simulation of single-cell RNA sequencing data. *Genome biology*, 18(1):174.

# Appendix

The appendix includes the proof for all the theorems, computational details, and extra experiment results. The structure of the appendix is listed below:

Appendix	Content
Appendix A	Proof of Proposition 1
Appendix B	B.1 Proof of Theorem 2. B.2 Lemmas B.1–B.2, which are used in the proof of Theorem 2.
Appendix C	C.1 Preparatory definitions. C.2 Proof of Theorem 3. C.3 Lemmas C.1–C.4, which are used in the proof of Theorem 3.
Appendix D	D.1 Preparatory definitions. D.2 Proof of Corollary 4. D.3 Proof of Theorem 5 D.4 Lemmas D.1–D.6, which are used in the proof of Theorem 5.
Appendix E	E.1 Proof of Theorem 6. E.2 Proof of Proposition 7. E.3 Lemmas E.1–E.3, which are used in the proof of Theorem 6 and Proposition 7.
Appendix F	F.1 Summary of commonly used exponential families. F.2 Initialization procedure, alternative maximization algorithm, and the estimation of dispersion parameters. F.3 Choosing hyperparameters in practice. F.4 Discussion about non-canonical links.
Appendix G	G.1 Efficiency loss of sample splitting. G.2 Information about lupus data. G.3 Extra results on lupus datasets.

## A Proof of Proposition 1

*Proof of Proposition 1.* Because the one-parameter exponential family is minimal, the natural parameter space is convex, and the log-partition function  $A$  is strictly convex. Based on the information of the first moment of  $\mathbf{y}$  and the log-partition function  $A$ , we can identify  $\mathbf{B}\mathbf{x} + \mathbf{\Gamma}\mathbf{z} = \boldsymbol{\theta} = A'^{-1}(\mathbb{E}[\mathbf{y}])$ . Because  $\mathbf{\Gamma}\mathbf{w}$  has zero mean and is uncorrelated to  $\mathbf{x}$ ,  $\text{Cov}(\mathbf{\Gamma}\mathbf{w}) = \mathbf{\Gamma}\boldsymbol{\Sigma}_w\mathbf{\Gamma}^\top$  can be identified as the residual covariance of regression of  $\boldsymbol{\theta}$  on  $\mathbf{x}$ .

Because  $\lambda_r(\mathbf{\Gamma}\boldsymbol{\Sigma}_w\mathbf{\Gamma}^\top) \geq \tau_p$ ,  $\mathbf{\Gamma}$  and  $\boldsymbol{\Sigma}_w$  have full rank. Let  $\mathbf{U}_r\boldsymbol{\Lambda}_r\mathbf{U}_r^\top$  be the reduced eigenvalue decomposition of  $\mathbf{\Gamma}\boldsymbol{\Sigma}_w\mathbf{\Gamma}^\top$  where  $\mathbf{U}_r \in \mathbb{R}^{p \times r}$ . Note that

$$\begin{aligned} \mathcal{P}_\Gamma &= \mathbf{\Gamma}\boldsymbol{\Sigma}_w^{1/2}(\boldsymbol{\Sigma}_w^{1/2}\mathbf{\Gamma}^\top\mathbf{\Gamma}\boldsymbol{\Sigma}_w^{1/2})^{-1}\boldsymbol{\Sigma}_w^{1/2}\mathbf{\Gamma}^\top \\ &= \mathbf{U}_r\boldsymbol{\Lambda}_r^{\frac{1}{2}}(\boldsymbol{\Lambda}_r^{\frac{1}{2}}\mathbf{U}_r^\top\mathbf{U}_r\boldsymbol{\Lambda}_r^{\frac{1}{2}})^{-1}\boldsymbol{\Lambda}_r^{\frac{1}{2}}\mathbf{U}_r^\top \\ &= \mathbf{U}_r\boldsymbol{\Lambda}_r^{\frac{1}{2}}(\boldsymbol{\Lambda}_r^{\frac{1}{2}}\boldsymbol{\Lambda}_r^{\frac{1}{2}})^{-1}\boldsymbol{\Lambda}_r^{\frac{1}{2}}\mathbf{U}_r^\top \\ &= \mathbf{U}_r\mathbf{U}_r^\top. \end{aligned}$$

Thus,  $\mathcal{P}_\Gamma$  can be recovered.

By the orthogonal decomposition, we have  $\mathbf{B} = \mathcal{P}_\Gamma^\perp\mathbf{B} + \mathcal{P}_\Gamma\mathbf{B}$ . Let  $\mathbf{e}_{p,i} = (\delta_{i\ell})_{1 \leq \ell \leq p}$  and  $\mathbf{e}_{d,j} = (\delta_{j\ell})_{1 \leq \ell \leq d}$ . We consider the  $(i, j)$ -th entry of  $\mathcal{P}_\Gamma\mathbf{B}$ :

$$\begin{aligned} |e_{p,i}^\top \mathcal{P}_\Gamma \mathbf{B} e_{d,j}| &= |e_{p,i}^\top \mathbf{\Gamma} \boldsymbol{\Sigma}_w^{1/2} (\boldsymbol{\Sigma}_w^{1/2} \mathbf{\Gamma}^\top \mathbf{\Gamma} \boldsymbol{\Sigma}_w^{1/2})^{-1} \boldsymbol{\Sigma}_w^{1/2} \mathbf{\Gamma}^\top \mathbf{B} e_{d,j}| \\ &\leq \|\mathbf{\Gamma} \boldsymbol{\Sigma}_w^{1/2} (\boldsymbol{\Sigma}_w^{1/2} \mathbf{\Gamma}^\top \mathbf{\Gamma} \boldsymbol{\Sigma}_w^{1/2})^{-1} \boldsymbol{\Sigma}_w^{1/2} \mathbf{\Gamma}^\top e_{p,i}\|_\infty \cdot \|\mathbf{B} e_{d,j}\|_1 \\ &= \max_{\ell \in [p]} |e_\ell^\top \mathbf{\Gamma} \boldsymbol{\Sigma}_w^{1/2} (\boldsymbol{\Sigma}_w^{1/2} \mathbf{\Gamma}^\top \mathbf{\Gamma} \boldsymbol{\Sigma}_w^{1/2})^{-1} \boldsymbol{\Sigma}_w^{1/2} \mathbf{\Gamma}^\top e_{p,i}| \cdot \|\mathbf{B} e_{d,j}\|_1 \\ &\leq \max_{\ell \in [p]} \|\boldsymbol{\Sigma}_w^{1/2} \mathbf{\Gamma}^\top e_{p,\ell}\|_2 \cdot \|(\boldsymbol{\Sigma}_w^{1/2} \mathbf{\Gamma}^\top \mathbf{\Gamma} \boldsymbol{\Sigma}_w^{1/2})^{-1} \boldsymbol{\Sigma}_w^{1/2} \mathbf{\Gamma}^\top e_{p,i}\|_2 \cdot \|\mathbf{B} e_{d,j}\|_1 \\ &\leq \max_{\ell \in [p]} \|\boldsymbol{\Sigma}_w^{1/2} \mathbf{\Gamma}^\top e_{p,\ell}\|_2 \cdot \|(\boldsymbol{\Sigma}_w^{1/2} \mathbf{\Gamma}^\top \mathbf{\Gamma} \boldsymbol{\Sigma}_w^{1/2})^{-1}\|_{\text{op}} \cdot \|\boldsymbol{\Sigma}_w^{1/2} \mathbf{\Gamma}^\top e_{p,i}\|_2 \cdot \|\mathbf{B}_j\|_1 \\ &\leq \max_{\ell \in [p]} \|\boldsymbol{\Sigma}_w^{1/2} \mathbf{\Gamma}^\top e_{p,\ell}\|_2 \cdot \lambda_r(\mathbf{\Gamma} \boldsymbol{\Sigma}_w \mathbf{\Gamma}^\top)^{-1} \cdot \|\boldsymbol{\Sigma}_w^{1/2} \mathbf{\Gamma}^\top e_{p,i}\|_2 \cdot \|\mathbf{B}_j\|_1 \\ &= \mathcal{O}\left(\frac{\|\mathbf{B}_j\|_1}{\tau_p}\right) \\ &= o(1), \end{aligned} \tag{A.1}$$

where the first two inequalities are from Holder's inequality; the third inequality holds because of the sub-multiplicativity of the operator norm; and the last inequality holds because  $\boldsymbol{\Sigma}_w^{1/2} \mathbf{\Gamma}^\top \mathbf{\Gamma} \boldsymbol{\Sigma}_w^{1/2}$  and  $\mathbf{\Gamma} \boldsymbol{\Sigma}_w \mathbf{\Gamma}^\top$  have the same non-zero eigenvalues. Then we have

$$\|\mathcal{P}_\Gamma \mathbf{B}\|_F \leq \sqrt{p} \max_{1 \leq i \leq p} \|\mathbf{B}^\top \mathcal{P}_\Gamma e_{p,i}\|_2 \lesssim \frac{\sqrt{p} \|\mathbf{B}\|_{1,1}}{\tau_p}.$$

Thus, the conclusion follows.  $\square$

## B Estimation error of natural parameters by alternative maximization

In this section, we gather useful results to bound the estimation error for the natural parameter matrix. Let  $E_C = \{\Theta^* \in \mathcal{R}_C^{n \times p}\}$  be the event that all the natural parameters are bounded. From Assumption 1, we know that  $\mathbb{P}(E_C) = \iota_n \rightarrow 1$  as  $n \rightarrow \infty$ . Under event  $E_C$ , because  $A$  is strictly convex and trice continuously differentiable, we have that

$$\kappa_1 := \inf_{\theta \in \mathcal{R}_C} A''(\theta) > 0 \text{ and } \kappa_2 := \sup_{\theta \in \mathcal{R}_C} A''(\theta) < \infty. \quad (\text{B.1})$$

These facts enable us to derive Theorem 2, which will be used in Appendix C for proving Theorem 3 and in Appendix D for proving Theorem 5.

### B.1 Estimation error of natural parameters

*Proof of Theorem 2.* We split the proof into two parts under event  $E_C$ .

**Part (1) Bounding  $\|\widehat{\Theta} - \Theta^*\|_F$ .** From the assumption of Theorem 2, we have

$$\mathcal{L}(\Theta^*) - \mathcal{L}(\widehat{\Theta}) \geq 0,$$

which also holds when  $\widehat{\Theta}$  is the maximum likelihood estimator. From Lemma B.1 it further follows that

$$0 \leq \sqrt{2(d+r)} \|\mathbf{Y} - A'(\Theta^*)\|_{\text{op}} \|\widehat{\Theta} - \Theta^*\|_F - \frac{\kappa_2}{2} \|\widehat{\Theta} - \Theta^*\|_F^2.$$

Thus, we have

$$\|\widehat{\Theta} - \Theta^*\|_F \leq \frac{2\sqrt{2(d+r)}}{\kappa_2} \|\mathbf{Y} - A'(\Theta^*)\|_{\text{op}}.$$

Next, we bound the operator norm of  $\mathbf{Y} - A'(\Theta^*)$ . Conditional on  $\mathbf{X}$  and  $\mathbf{Z}$ , observe that  $z_{ij} = y_{ij} - A'(\theta_{ij}^*)$  ( $i \in [n]$  and  $j \in [p]$ ) are independent, zero-meaned, and sub-exponential with parameters  $\nu = \sqrt{\kappa_2}$  and  $\alpha = 1/C^2$ . To see this, note that its moment generating function is  $\mathbb{E}[\exp(tz_{ij})] = \exp(A(\theta_{ij}^* + t) - A(\theta_{ij}^*) - tA'(\theta_{ij}^*)) = \exp(A''(\theta_{ij}^* + t')t^2/2)$  for some  $|t'| < |t|$ . By Assumption 1, we have  $\mathbb{E}[\exp(tz_{ij})] \leq \kappa_2 t^2/2$  for all  $|t| < C^2$ , which shows that  $z_{ij}$  is sub-exponential. By Lemma B.2, for any  $\delta > 0$ , with probability at least  $1 - (n+p)^{-\delta} - (np)^{-\delta}$ , it follows that

$$\begin{aligned} \|\widehat{\Theta} - \Theta^*\|_F &\leq \frac{2\sqrt{2(d+r)}}{\kappa_2} (4\nu\sqrt{n \vee p} + 2\delta^{\frac{3}{2}}\sqrt{c}(\alpha \vee \nu) \log(np) \sqrt{\log(n+p)}) \\ &\lesssim \sqrt{(d+r)(n \vee p)}. \end{aligned}$$

**Part (2) Bounding**  $\max_{1 \leq j \leq p} \|\widehat{\Theta}_j - \Theta_j^*\|_2$ . Similarly to Part (1), by union bound, we have

$$\begin{aligned} \max_{1 \leq j \leq p} \|\widehat{\Theta}_j - \Theta_j^*\|_2 &\leq \max_{1 \leq j \leq p} \frac{2\sqrt{2(d+r)}}{\kappa_2} \|\mathbf{Y}_j - A'(\Theta_j^*)\|_2 \\ &\leq \frac{2\sqrt{2(d+r)}}{\kappa_2} (4\nu\sqrt{n} + 2(\delta+1)^{\frac{3}{2}}\sqrt{c}(\alpha \vee \nu) \log(n)\sqrt{\log(n+1)}), \end{aligned}$$

with probability at least  $1 - p(n+p)^{-\delta-1} - p(np)^{-\delta-1} \geq 1 - (n+p)^{-\delta} - (np)^{-\delta}$ , for any  $\delta > 0$ .

For  $\delta > 1$ , taking union bound over the above two events and  $E_C$  finishes the proof.  $\square$

## B.2 Technical lemmas

**Lemma B.1** (Upper bound of likelihood difference). Suppose that  $\Theta_1 \in \mathcal{R}_{r_1}, \Theta_2 \in \mathcal{R}_{r_2}$  with  $r_j = \text{rank}(\Theta_j)$  for  $j = 1, 2$ . Define  $\kappa_1 := \inf_{\theta \in \mathcal{R}} A''(\theta)$ . Then it holds that

$$\mathcal{L}(\Theta_2) - \mathcal{L}(\Theta_1) \leq \frac{\sqrt{r_1 + r_2}}{n} \|\mathbf{Y} - A'(\Theta_2)\|_{\text{op}} \|\Theta_1 - \Theta_2\|_{\text{F}} - \frac{\kappa_1}{2n} \|\Theta_1 - \Theta_2\|_{\text{F}}^2,$$

and

$$\mathcal{L}(\Theta_2) - \mathcal{L}(\Theta_1) \leq \frac{\sqrt{r_1 + r_2}}{n} \|\mathbf{Y} - A'(\Theta_1)\|_{\text{op}} \|\Theta_1 - \Theta_2\|_{\text{F}} + \frac{\kappa_2}{2n} \|\Theta_1 - \Theta_2\|_{\text{F}}^2.$$

*Proof of Lemma B.1.* Recall that  $\mathcal{L}(\Theta) = n^{-1}[-\text{tr}(\mathbf{Y}^\top \Theta) + \text{tr}(\mathbf{1}_{p \times n} A(\Theta))]$ . Then we have

$$\begin{aligned} \mathcal{L}(\Theta_2) - \mathcal{L}(\Theta_1) &= \frac{1}{n} \text{tr}((\mathbf{Y} - A'(\Theta_2))^\top (\Theta_1 - \Theta_2)) \\ &\quad - \frac{1}{n} \text{tr}(\mathbf{1}_{p \times n} (A(\Theta_1) - A(\Theta_2)) - A'(\Theta_2)^\top (\Theta_1 - \Theta_2)). \end{aligned} \quad (\text{B.2})$$

Next, we analyze the two terms separately.

For the first term, we have

$$\text{tr}((\mathbf{Y} - A'(\Theta_2))^\top (\Theta_1 - \Theta_2)) \quad (\text{B.3})$$

$$\begin{aligned} &\leq \sqrt{\text{rank}(\Theta_1 - \Theta_2)} \|\mathbf{Y} - A'(\Theta_2)\|_{\text{op}} \|\Theta_1 - \Theta_2\|_{\text{F}} \\ &\leq \sqrt{\text{rank}(\Theta_1) + \text{rank}(\Theta_2)} \|\mathbf{Y} - A'(\Theta_2)\|_{\text{op}} \|\Theta_1 - \Theta_2\|_{\text{F}}, \end{aligned} \quad (\text{B.4})$$

where the first inequality is from the matrix norm inequality  $|\text{tr}(\mathbf{A}^\top \mathbf{B})| \leq \sqrt{\text{rank}(\mathbf{B})} \|\mathbf{A}\|_{\text{op}} \|\mathbf{B}\|_{\text{F}}$  and the last inequality is due to the fact that  $\text{rank}(\mathbf{A} + \mathbf{B}) \leq \text{rank}(\mathbf{A}) + \text{rank}(\mathbf{B})$ .

For the second term, note that each entry inside the trace takes the form

$$\begin{aligned} A((\Theta_1)_{ij}) - A((\Theta_2)_{ij}) - A'((\Theta_2)_{ij})((\Theta_1)_{ij} - (\Theta_2)_{ij}) &= \frac{1}{2} A''(\theta) ((\Theta_1)_{ij} - (\Theta_2)_{ij})^2 \\ &\geq \frac{\kappa_1}{2} ((\Theta_1)_{ij} - (\Theta_2)_{ij})^2 \end{aligned}$$



where the first equality is from Taylor expansion with  $\theta$  lies between  $(\Theta_1)_{ij}$  and  $(\Theta_2)_{ij}$  and the definition of  $\kappa_1 := \inf_{\theta \in \mathcal{R}} A''(\theta) \geq 0$ . Thus we have

$$\text{tr}(\mathbf{1}_{p \times n}(A(\Theta_1) - A(\Theta_2)) - A'(\Theta_2)^\top(\Theta_1 - \Theta_2)) \geq \frac{\kappa_1}{2} \|\Theta_1 - \Theta_2\|_F^2. \quad (\text{B.5})$$

Combining (B.2), (B.4), and (B.5) finishes the proof of the first inequality. Similarly, we have

$$\begin{aligned} \mathcal{L}(\Theta_2) - \mathcal{L}(\Theta_1) &= \frac{1}{n} \text{tr}((\mathbf{Y} - A'(\Theta_1))^\top(\Theta_1 - \Theta_2)) \\ &\quad + \frac{1}{n} \text{tr}(\mathbf{1}_{p \times n}(A(\Theta_2) - A(\Theta_1)) - A'(\Theta_1)^\top(\Theta_2 - \Theta_1)) \\ &\leq \frac{\sqrt{r_1 + r_2}}{n} \|\mathbf{Y} - A'(\Theta_1)\|_{\text{op}} \|\Theta_1 - \Theta_2\|_F + \frac{\kappa_2}{2n} \|\Theta_1 - \Theta_2\|_F^2, \end{aligned}$$

which completes the proof of the second inequality.  $\square$

**Lemma B.2** (Operator norm of matrices with sub-exponential entries). Let  $\mathbf{X} = (x_{ij})_{i \in [n], j \in [p]}$  be a matrix with independent and centered entries such that  $x_{ij}$ 's are  $(\nu, \alpha)$ -sub-exponential random variables<sup>1</sup> with parameters  $\nu, \alpha > 0$ :

$$\mathbb{E}[\exp(tx_{ij})] \leq \exp(t^2\nu^2/2), \quad \forall |t| < \frac{1}{\alpha}.$$

Then for all  $\delta > 0$ , there exists a universal constant  $c > 0$  such that, with probability at least  $1 - (n+p)^{-\delta} - (np)^{-\delta}$ , when  $n, p$  are large enough, it holds that

$$\|\mathbf{X}\|_{\text{op}} \leq 4\nu\sqrt{n \vee p} + 2\delta^{3/2}\sqrt{c}(\alpha \vee \nu) \log(np) \sqrt{\log(n+p)}.$$

*Proof of Lemma B.2.* We define a symmetric matrix

$$\mathbf{Z} = (z_{ij}) = \begin{pmatrix} 0 & \widetilde{\mathbf{X}} \\ \widetilde{\mathbf{X}}^\top & 0 \end{pmatrix} \in \mathbb{R}^{(n+p) \times (n+p)},$$

where  $\widetilde{\mathbf{X}} := (\widetilde{x}_{ij}) = \mathbf{X} - \mathbf{X}'$  and  $\mathbf{X}' = (x'_{ij})$  is an independent copy of  $\mathbf{X}$ . Because  $\widetilde{x}_{ij}$ 's have symmetric distribution and are independent, it follows that  $z_{ij}$ 's are also independent and symmetric random variables, and  $\|\mathbf{Z}\|_{\text{op}} = \|\widetilde{\mathbf{X}}\|_{\text{op}}$ . Because the tail event  $\mathbb{1}\{\|\mathbf{X} - \mathbf{X}'\|_{\text{op}} \geq t\}$  is a convex function on  $\mathbf{X}'$ , by Jensen's inequality we have

$$\mathbb{1}\{\|\mathbf{X}\|_{\text{op}} \geq t\} = \mathbb{1}\{\|\mathbf{X} - \mathbb{E}_{\mathbf{X}'}[\mathbf{X}']\|_{\text{op}} \geq t\} \leq \mathbb{E}_{\mathbf{X}'}[\mathbb{1}\{\|\mathbf{X} - \mathbf{X}'\|_{\text{op}} \geq t\}] = \mathbb{E}_{\mathbf{X}'}[\mathbb{1}\{\|\widetilde{\mathbf{X}}\|_{\text{op}} \geq t\}].$$

By Fubini's theorem, it follows that

$$\mathbb{P}(\|\mathbf{X}\|_{\text{op}} \geq t) \leq \mathbb{E}_{\mathbf{X}, \mathbf{X}'}[\mathbb{1}\{\|\widetilde{\mathbf{X}}\|_{\text{op}} \geq t\}] = \mathbb{P}(\|\widetilde{\mathbf{X}}\|_{\text{op}} \geq t) = \mathbb{P}(\|\mathbf{Z}\|_{\text{op}} \geq t).$$

<sup>1</sup>Here we adopt the definition from [Wainwright \(2019, Definition 2.7\)](#). In some literature, the term 'sub-gamma' is used interchangeably with 'sub-exponential' to refer to this definition.

Then, it suffices to bound the tail probability of  $\|\mathbf{Z}\|_{\text{op}}$ .

We define a truncated random matrix  $\mathbf{Z}(\lambda)$  of  $\mathbf{Z}$ ,

$$\mathbf{Z}(\lambda) = (z_{ij}(\lambda))_{1 \leq i \leq n, 1 \leq j \leq p} = (z_{ij} \mathbf{1}(|z_{ij}| \leq \lambda))_{1 \leq i \leq n, 1 \leq j \leq p}$$

whose entries are independent, symmetric random variables bounded by  $\lambda$ . By [Bandeira and van Handel \(2016, Corollary 3.12\)](#), there exists a universal constant  $c > 0$  such that

$$\mathbb{P} \left( \|\mathbf{Z}(\lambda)\|_{\text{op}} \geq 2^{\frac{3}{2}} \max_{1 \leq i \leq n+p} \left( \sum_{j=1}^{n+p} \mathbb{E}[z_{ij}^2(\lambda)] \right)^{\frac{1}{2}} + t \right) \leq (n+p) \exp \left( -\frac{t^2}{c\lambda^2} \right).$$

Note that

$$\begin{aligned} \max_{1 \leq i \leq n+p} \left( \sum_{j=1}^{n+p} \mathbb{E}[z_{ij}^2(\lambda)] \right)^{\frac{1}{2}} &\leq \max_{1 \leq i \leq n+p} \left( \sum_{j=1}^{n+p} \mathbb{E}[z_{ij}^2] \right)^{1/2} \\ &= \max \left[ \max_{1 \leq i \leq n} \left( \sum_{j=1}^p \mathbb{E}[\tilde{x}_{ij}^2] \right)^{\frac{1}{2}}, \max_{1 \leq j \leq p} \left( \sum_{i=1}^n \mathbb{E}[\tilde{x}_{ij}^2] \right)^{\frac{1}{2}} \right] \\ &\leq \max\{\sqrt{p}, \sqrt{n}\} \max_{i,j} \mathbb{E}[\tilde{x}_{ij}^2]^{\frac{1}{2}} \\ &\leq \sqrt{2(n \vee p)} \max_{i,j} \mathbb{E}[x_{ij}^2]^{\frac{1}{2}} \\ &\leq \nu \sqrt{2(n \vee p)}, \end{aligned}$$

where the first inequality is from the definition of the truncated variable, the second and the third inequality is by Cauchy-Schwartz inequality, and the last inequality is because  $x_{ij}$  is  $(\nu, \alpha)$ -sub-exponential. Thus, the above two inequality yields that

$$\mathbb{P} (\|\mathbf{Z}(\lambda)\|_{\text{op}} \geq 4\nu\sqrt{n \vee p} + t) \leq (n+p) \exp \left( -\frac{t^2}{c\lambda^2} \right).$$

Then we have

$$\begin{aligned} \mathbb{P} (\|\mathbf{Z}\|_{\text{op}} \geq 4\nu\sqrt{n \vee p} + t) &\leq \mathbb{P} (\|\mathbf{Z}(\lambda)\|_2 \geq 4\nu\sqrt{n \vee p} + t) + \mathbb{P} \left( \max_{1 \leq i, j \leq n+p} |z_{ij}| > \lambda \right) \\ &\leq (n+p) \exp \left( -\frac{t^2}{c\lambda^2} \right) + \sum_{1 \leq i \leq n} \sum_{1 \leq j \leq p} \mathbb{P} (|\tilde{x}_{ij}| > \lambda) \\ &\leq (n+p) \exp \left( -\frac{t^2}{c\lambda^2} \right) + np \left( \exp \left( -\frac{\lambda^2}{4\nu^2} \right) \vee \exp \left( -\frac{\lambda}{2\alpha} \right) \right). \end{aligned}$$

where the last inequality follows because  $\tilde{x}_{ij} = x_{ij} - x'_{ij}$  is  $(2\nu^2, \alpha)$ -sub-exponential. For all  $\delta > 0$ , let  $\lambda = 2(\delta + 1)(\alpha \vee \nu) \log(np)$  and  $np \geq 3$ , the second term is bounded by  $(np)^{-\delta}$ . Let  $t = \lambda((\delta + 1)c \log(n+p))^{1/2}$ , the first term is bounded by  $(n+p)^{-\delta}$ . Combining these

yields that

$$\mathbb{P}\left(\|\mathbf{Z}\|_{\text{op}} \geq 4\nu\sqrt{n\vee p} + 2\delta^{3/2}\sqrt{c}(\alpha \vee \nu)\log(np)\sqrt{\log(n+p)}\right) \leq (n+p)^{-\delta} + (np)^{-\delta},$$

which completes the proof.  $\square$

## C Estimation of latent coefficients

### C.1 Preparatory definitions

Analogous to [Bing et al. \(2023\)](#), we define  $\mathbf{H}_0 := (np)^{-1}\mathbf{W}^{*\top}\mathbf{W}^*\mathbf{\Gamma}^{*\top}\widehat{\mathbf{\Gamma}}\mathbf{\Sigma}^{-3/2}$  and

$$\widetilde{\mathbf{\Gamma}} := \mathbf{\Gamma}^*\mathbf{H}_0 = (np)^{-1}\mathbf{\Gamma}^*\mathbf{W}^{*\top}\mathbf{W}^*\mathbf{\Gamma}^{*\top}\widehat{\mathbf{\Gamma}}\mathbf{\Sigma}^{-3/2}, \quad (\text{C.1})$$

which is identifiable because it depends on both the data  $\widehat{\mathbf{\Gamma}}\mathbf{\Sigma}^{-3/2}$  and the identifiable quantity  $\mathbf{\Gamma}^*\mathbf{W}^{*\top}\mathbf{W}^*\mathbf{\Gamma}^{*\top}$ . Note that

$$\begin{aligned} \mathcal{P}_{\widetilde{\mathbf{\Gamma}}} &= \widetilde{\mathbf{\Gamma}}(\widetilde{\mathbf{\Gamma}}^\top\widetilde{\mathbf{\Gamma}})^{-1}\widetilde{\mathbf{\Gamma}}^\top \\ &= \mathbf{\Gamma}^*(\mathbf{W}^{*\top}\mathbf{W}^*\mathbf{\Gamma}^{*\top}\mathbf{V})(\mathbf{V}^\top\mathbf{\Gamma}^*\mathbf{W}^{*\top}\mathbf{W}^*\mathbf{\Gamma}^{*\top}\mathbf{\Gamma}^*\mathbf{W}^{*\top}\mathbf{W}^*\mathbf{\Gamma}^{*\top}\mathbf{V})^{-1}(\mathbf{V}^\top\mathbf{\Gamma}^*\mathbf{W}^{*\top}\mathbf{W}^*)\mathbf{\Gamma}^{*\top} \\ &= \mathbf{\Gamma}^*(\mathbf{\Gamma}^{*\top}\mathbf{\Gamma}^*)^{-1}\mathbf{\Gamma}^{*\top} \\ &= \mathcal{P}_{\mathbf{\Gamma}^*} \end{aligned} \quad (\text{C.2})$$

because both  $\mathbf{\Gamma}^{*\top}\mathbf{\Gamma}^*$  and  $\mathbf{W}^{*\top}\mathbf{W}^*\mathbf{\Gamma}^{*\top}\mathbf{V} \in \mathbb{R}^{r \times r}$  have full rank. Thus, to quantify the error between  $\mathcal{P}_{\widehat{\mathbf{\Gamma}}}$  and  $\mathcal{P}_{\mathbf{\Gamma}^*}$ , we can first analyze the error between  $\widehat{\mathbf{\Gamma}}$  and  $\widetilde{\mathbf{\Gamma}}$ .

### C.2 Proof of Theorem 3

*Proof of Theorem 3.* We split the proof into three parts by bounding the operator norm, column-wise  $\ell_2$ -norm, and the sup norm consecutively.

**Part (1) Bounding the operator norm.** From (C.2), we have that  $\mathcal{P}_{\widetilde{\mathbf{\Gamma}}} = \mathcal{P}_{\mathbf{\Gamma}^*}$  for  $\widetilde{\mathbf{\Gamma}}$  defined in (C.1). Then we have

$$\begin{aligned} &\|\mathcal{P}_{\widehat{\mathbf{\Gamma}}} - \mathcal{P}_{\mathbf{\Gamma}^*}\|_{\text{op}} \\ &= \|\mathcal{P}_{\widehat{\mathbf{\Gamma}}} - \mathcal{P}_{\widetilde{\mathbf{\Gamma}}}\|_{\text{op}} \\ &= \|\widehat{\mathbf{\Gamma}}(\widehat{\mathbf{\Gamma}}^\top\widehat{\mathbf{\Gamma}})^{-1}\widehat{\mathbf{\Gamma}}^\top - \widetilde{\mathbf{\Gamma}}(\widetilde{\mathbf{\Gamma}}^\top\widetilde{\mathbf{\Gamma}})^{-1}\widetilde{\mathbf{\Gamma}}^\top\|_{\text{op}} \\ &\leq \|(\widehat{\mathbf{\Gamma}} - \widetilde{\mathbf{\Gamma}})(\widetilde{\mathbf{\Gamma}}^\top\widetilde{\mathbf{\Gamma}})^{-1}\widetilde{\mathbf{\Gamma}}^\top\|_{\text{op}} + \|\widehat{\mathbf{\Gamma}}((\widehat{\mathbf{\Gamma}}^\top\widehat{\mathbf{\Gamma}})^{-1} - (\widetilde{\mathbf{\Gamma}}^\top\widetilde{\mathbf{\Gamma}})^{-1})\widetilde{\mathbf{\Gamma}}^\top\|_{\text{op}} + \|\widehat{\mathbf{\Gamma}}(\widehat{\mathbf{\Gamma}}^\top\widehat{\mathbf{\Gamma}})^{-1}(\widehat{\mathbf{\Gamma}} - \widetilde{\mathbf{\Gamma}})^\top\|_{\text{op}}, \end{aligned}$$

where the last inequality is from the triangle inequality. Next, we bound the three terms separately. Recall  $\widetilde{\mathbf{\Gamma}}$  is defined in (C.1) with  $\|\widetilde{\mathbf{\Gamma}}\|_{\text{op}} \asymp \|\widehat{\mathbf{\Gamma}}\|_{\text{op}} \asymp \sqrt{p}$  by Assumption 3 and Lemma C.2. Then by Lemma C.1 and Assumption 3, for all  $\delta > 0$ , the first term in the

above display is bounded

$$\|(\widehat{\Gamma} - \widetilde{\Gamma})(\widetilde{\Gamma}^\top \widetilde{\Gamma})^{-1} \widetilde{\Gamma}^\top\|_{\text{op}} \leq \|\widehat{\Gamma} - \widetilde{\Gamma}\|_{\text{op}} \|(\widetilde{\Gamma}^\top \widetilde{\Gamma})^{-1}\|_{\text{op}} \|\widetilde{\Gamma}\|_{\text{op}} \leq C'(n \wedge p)^{-\frac{1}{2}},$$

with probability at least  $1 - 2(n+p)^{-\delta} - 2(np)^{-\delta} - \exp(-n)$ , for some constant  $C' > 0$  and  $\delta > 0$ . Similarly, the third term is also bounded by  $\mathcal{O}_{\mathbb{P}}((n \vee p)^{-1/2})$ . It remains to bound the second term:

$$\begin{aligned} & \|\widehat{\Gamma}((\widehat{\Gamma}^\top \widehat{\Gamma})^{-1} - (\widetilde{\Gamma}^\top \widetilde{\Gamma})^{-1}) \widetilde{\Gamma}^\top\|_{\text{op}} \\ &= \|\widehat{\Gamma}(\widehat{\Gamma}^\top \widehat{\Gamma})^{-1}(\widehat{\Gamma}^\top \widehat{\Gamma} - \widetilde{\Gamma}^\top \widetilde{\Gamma})(\widetilde{\Gamma}^\top \widetilde{\Gamma})^{-1} \widetilde{\Gamma}^\top\|_{\text{op}} \\ &\leq \|\widehat{\Gamma}(\widehat{\Gamma}^\top \widehat{\Gamma})^{-1}\|_{\text{op}} \cdot \|(\widehat{\Gamma}^\top \widehat{\Gamma} - \widetilde{\Gamma}^\top \widetilde{\Gamma})(\widetilde{\Gamma}^\top \widetilde{\Gamma})^{-1} \widetilde{\Gamma}^\top\|_{\text{op}} \\ &\lesssim \frac{1}{\sqrt{p}} \|(\widehat{\Gamma}^\top \widehat{\Gamma} - \widetilde{\Gamma}^\top \widetilde{\Gamma})(\widetilde{\Gamma}^\top \widetilde{\Gamma})^{-1} \widetilde{\Gamma}^\top\|_{\text{op}} \\ &\leq \frac{1}{\sqrt{p}} \|\widehat{\Gamma}^\top (\widehat{\Gamma} - \widetilde{\Gamma})(\widetilde{\Gamma}^\top \widetilde{\Gamma})^{-1} \widetilde{\Gamma}^\top\|_{\text{op}} + \frac{1}{\sqrt{p}} \|(\widehat{\Gamma} - \widetilde{\Gamma})^\top \widetilde{\Gamma}(\widetilde{\Gamma}^\top \widetilde{\Gamma})^{-1} \widetilde{\Gamma}^\top\|_{\text{op}} \\ &\leq \frac{1}{\sqrt{p}} \|\widehat{\Gamma}\|_{\text{op}} \|(\widehat{\Gamma} - \widetilde{\Gamma})(\widetilde{\Gamma}^\top \widetilde{\Gamma})^{-1} \widetilde{\Gamma}^\top\|_{\text{op}} + \frac{1}{\sqrt{p}} \|\widehat{\Gamma} - \widetilde{\Gamma}\|_{\text{op}} \|\mathcal{P}_{\widetilde{\Gamma}}\|_{\text{op}} \\ &\lesssim \mathcal{O}\left((n \wedge p)^{-\frac{1}{2}}\right), \end{aligned}$$

with probability at least  $1 - (n+p)^{-\delta} - (np)^{-\delta} - \exp(-n)$ . The proof for the operator norm is completed by combing the above inequality.

**Part (2) Bounding the column-wise  $\ell_2$ -norm.** Let  $\mathbf{e}_j \in \mathbb{R}^p$  be the unit vector such that its  $i$ -th entry is one if  $i = j$  and zero otherwise. Similar to Part (1), note that

$$\begin{aligned} & \|(\mathcal{P}_{\widehat{\Gamma}} - \mathcal{P}_{\Gamma^*})\mathbf{e}_j\|_2 \\ &\leq \|(\widehat{\Gamma} - \widetilde{\Gamma})(\widetilde{\Gamma}^\top \widetilde{\Gamma})^{-1} \widetilde{\Gamma}^\top \mathbf{e}_j\|_2 + \|\widehat{\Gamma}((\widehat{\Gamma}^\top \widehat{\Gamma})^{-1} - (\widetilde{\Gamma}^\top \widetilde{\Gamma})^{-1}) \widetilde{\Gamma}^\top \mathbf{e}_j\|_2 + \|\widehat{\Gamma}(\widehat{\Gamma}^\top \widehat{\Gamma})^{-1}(\widehat{\Gamma} - \widetilde{\Gamma})^\top \mathbf{e}_j\|_2. \end{aligned}$$

The first term can be bounded analogously as

$$\max_{1 \leq j \leq p} \|(\widehat{\Gamma} - \widetilde{\Gamma})(\widetilde{\Gamma}^\top \widetilde{\Gamma})^{-1} \widetilde{\Gamma}^\top \mathbf{e}_j\|_2 \leq \|\widehat{\Gamma} - \widetilde{\Gamma}\|_{\text{op}} \|(\widetilde{\Gamma}^\top \widetilde{\Gamma})^{-1}\|_{\text{op}} \max_{1 \leq j \leq p} \|\widetilde{\Gamma}^\top \mathbf{e}_j\|_2 \leq C'[p(n \wedge p)]^{-\frac{1}{2}},$$

for some constant  $C' > 0$ , by noting that  $\|\widetilde{\Gamma}^\top \mathbf{e}_j\|_2 = \mathcal{O}_{\mathbb{P}}(1)$ . The rest of the terms follow a similar argument as in Part (1), under the same probabilistic events therein.

**Part (3) Bounding the sup norm.** The sup norm  $\|\cdot\|_{\max}$  can be upper bounded analogously:

$$\begin{aligned} & \|\mathcal{P}_{\widehat{\Gamma}} - \mathcal{P}_{\Gamma^*}\|_{\max} \\ &= \max_{i,j \in [p]} |\mathbf{e}_i^\top (\mathcal{P}_{\widehat{\Gamma}} - \mathcal{P}_{\Gamma^*}) \mathbf{e}_j| \\ &\leq \max_{i,j \in [p]} (|\mathbf{e}_i^\top (\widehat{\Gamma} - \widetilde{\Gamma})(\widetilde{\Gamma}^\top \widetilde{\Gamma})^{-1} \widetilde{\Gamma}^\top \mathbf{e}_j| + |\mathbf{e}_i^\top \widehat{\Gamma}((\widehat{\Gamma}^\top \widehat{\Gamma})^{-1} - (\widetilde{\Gamma}^\top \widetilde{\Gamma})^{-1}) \widetilde{\Gamma}^\top \mathbf{e}_j| + |\mathbf{e}_i^\top \widehat{\Gamma}(\widehat{\Gamma}^\top \widehat{\Gamma})^{-1}(\widehat{\Gamma} - \widetilde{\Gamma})^\top \mathbf{e}_j|) \end{aligned}$$

The first term can be bounded as

$$\max_{1 \leq j \leq p} |\mathbf{e}_i^\top (\widehat{\Gamma} - \widetilde{\Gamma}) (\widetilde{\Gamma}^\top \widetilde{\Gamma})^{-1} \widetilde{\Gamma}^\top \mathbf{e}_j| \leq \|(\widehat{\Gamma} - \widetilde{\Gamma}) \mathbf{e}_i\|_{\text{op}} \|(\widetilde{\Gamma}^\top \widetilde{\Gamma})^{-1}\|_{\text{op}} \max_{1 \leq j \leq p} \|\widetilde{\Gamma}^\top \mathbf{e}_j\|_2 \leq C' [p^2(n \wedge p)]^{-\frac{1}{2}},$$

by involving both Part (1) and (2). The rest of the terms follow a similar argument as in Part (1), under the same probabilistic events therein. This completes the proof.  $\square$

### C.3 Technical lemmas

**Lemma C.1** (Estimation error of  $\widehat{\Gamma}$ ). Under Assumptions 1–3 and event  $E_C$ , for all  $\delta > 0$  and sufficiently large  $n$  and  $p$ , there exists a absolute constant  $C' > 0$  such that

$$\max_{1 \leq j \leq p} \|\widehat{\gamma}_j - \widetilde{\gamma}_j\|_2 \leq C', \quad \|\widehat{\Gamma} - \widetilde{\Gamma}\|_{\text{op}} \leq C' \sqrt{\frac{n \vee p}{n}},$$

with probability at least  $1 - 2(n+p)^{-\delta} - 2(np)^{-\delta} - \exp(-n)$ .

*Proof of Lemma C.1.* Define  $\mathbf{E} = \mathbf{W}^* \mathbf{\Gamma}^{*\top}$  and  $\widehat{\mathbf{E}} = \widehat{\mathbf{W}}_0 \widehat{\Gamma}^{\top 2}$ . Then we have  $\widehat{\mathbf{E}} = \mathbf{W}^* \mathbf{\Gamma}^{*\top} + \mathbf{\Delta}$  where  $\mathbf{\Delta} = (\widehat{\Theta} - \Theta) - (\mathbf{X} \widehat{\mathbf{F}}^\top - \mathbf{X} \mathbf{F}^{*\top})$ . By the definition of  $\widehat{\mathbf{E}}$  we have

$$\frac{1}{np} \widehat{\mathbf{E}}^\top \widehat{\mathbf{E}} = \mathbf{V} \mathbf{\Sigma}^2 \mathbf{V}^\top.$$

Note that  $\widehat{\Gamma} = \sqrt{p} \mathbf{V} \mathbf{\Sigma}^{1/2}$ , we further have

$$\frac{1}{np} \widehat{\mathbf{E}}^\top \widehat{\mathbf{E}} \widehat{\Gamma} = \widehat{\Gamma} \mathbf{\Sigma}^2$$

and

$$\frac{1}{n\sqrt{p}} \widehat{\mathbf{E}}^\top \widehat{\mathbf{E}} \mathbf{V} \mathbf{\Sigma}^{-3/2} = \widehat{\Gamma}.$$

It follows that

$$\widehat{\Gamma} - \widetilde{\Gamma} = \frac{1}{n\sqrt{p}} (\mathbf{E}^\top \mathbf{\Delta} + \mathbf{\Delta}^\top \mathbf{E} + \mathbf{\Delta}^\top \mathbf{\Delta}) \mathbf{V} \mathbf{\Sigma}^{-3/2}.$$

Because the operator norm is sub-multiplicative, the  $\ell_2$ -norm of the  $j$ th row of  $\widehat{\Gamma} - \widetilde{\Gamma}$  is bounded by

$$\begin{aligned} \|\widehat{\gamma}_j - \widetilde{\gamma}_j\|_2 &\leq \frac{1}{n\sqrt{p}} (\|\mathbf{E}^\top \mathbf{\Delta}_j\|_2 + \|\mathbf{\Delta}^\top \mathbf{E}_j\|_2 + \|\mathbf{\Delta}^\top \mathbf{\Delta}_j\|_2) \|\mathbf{V}\|_{\text{op}} \|\mathbf{\Sigma}^{-3/2}\|_{\text{op}} \\ &\leq \frac{1}{n\sqrt{p}} (\|\mathbf{E}\|_{\text{op}} \|\mathbf{\Delta}_j\|_2 + \|\mathbf{\Delta}\|_{\text{op}} \|\mathbf{E}_j\|_2 + \|\mathbf{\Delta}\|_{\text{op}} \|\mathbf{\Delta}_j\|_2) \|\mathbf{\Sigma}^{-3/2}\|_{\text{op}}, \end{aligned} \quad (\text{C.3})$$

---

<sup>2</sup>Throughout the manuscript, the notation  $\mathbf{e}_i$  is reserved for the unit vector and is not the  $i$ -th row of  $\mathbf{E}$ . We will only use the notations of  $\mathbf{E}$  and  $\mathbf{E}_j$  to denote matrix of latent components and its  $j$ -th column.

and

$$\|\widehat{\Gamma} - \widetilde{\Gamma}\|_{\text{op}} \leq \frac{1}{n\sqrt{p}}(2\|\mathbf{E}\|_{\text{op}}\|\Delta\|_{\text{op}} + \|\Delta\|_{\text{op}}^2)\|\Sigma^{-3/2}\|_{\text{op}}. \quad (\text{C.4})$$

To proceed, we split the proof into three parts.

**Part (1) Bounding operator norms of  $\Sigma$ ,  $\mathbf{E}$ , and  $\Delta$ .** Note that  $\mathbf{E} = \mathbf{W}^*\Gamma^{*\top}$  and  $\mathbf{w}_1, \dots, \mathbf{w}_n$  are mean-zero sub-Gaussian random vectors from Assumption 3. From Lemma C.2, for any  $\delta > 0$ , there exists  $C_\Sigma > 0$ , such that

$$\|\mathbf{E}\|_{\text{op}} \leq 2C_\Sigma\sqrt{np}, \quad \frac{1}{C_\Sigma} \leq \lambda_r(\Sigma) \leq \lambda_1(\Sigma) \leq C_\Sigma,$$

with probability at least  $1 - (n+p)^{-\delta} - (np)^{-\delta} - \exp(-n)$ .

Because

$$\Delta = \mathbf{E} - \widehat{\mathbf{E}} = \mathcal{P}_{\mathbf{X}}^\perp(\Theta - \widehat{\Theta}) + \mathcal{P}_{\mathbf{X}}\mathbf{E},$$

we have

$$\begin{aligned} \|\Delta\|_{\text{op}} &= \|\mathcal{P}_{\mathbf{X}}^\perp(\Theta - \widehat{\Theta}) + \mathcal{P}_{\mathbf{X}}\mathbf{E}\|_{\text{op}} \\ &\leq \|\mathcal{P}_{\mathbf{X}}^\perp\|_{\text{op}}\|\Theta - \widehat{\Theta}\|_{\text{op}} + \|\mathcal{P}_{\mathbf{X}}\mathbf{E}\|_{\text{op}} \\ &\leq \|\Theta - \widehat{\Theta}\|_{\text{op}} + \|\mathcal{P}_{\mathbf{X}}\mathbf{E}\|_{\text{op}}. \end{aligned}$$

On the one hand, from Theorem 2, it follows that when  $n, p$  are large enough,

$$\|\Theta - \widehat{\Theta}\|_{\text{op}} \leq \sqrt{c(d+r)(n \vee p)}$$

with probability at least  $1 - (n+p)^{-\delta} - (np)^{-\delta}$  for some constant  $c > 0$ . On the other hand, notice that

$$\mathcal{P}_{\mathbf{X}}\mathbf{E} = \mathcal{P}_{\mathbf{X}}\mathbf{W}^*\Gamma^{*\top} = \mathbf{X} \left( \frac{\mathbf{X}^\top \mathbf{X}}{n} \right)^{-1} \frac{\mathbf{X}^\top \mathbf{W}^*}{n} \Gamma^{*\top}$$

where  $\mathbf{X}^\top \mathbf{W}^* = \sum_{i=1}^n \mathbf{x}_i \mathbf{w}_i^{*\top}$  is the sum of  $n$  i.i.d. sub-exponential random matrices with zero means. By the matrix Bernstein's inequality,  $\|\mathbf{X}^\top \mathbf{W}^*/n\|_{\text{op}} \lesssim \sqrt{\log(nd)/n}$ . Thus, the second term can be bounded as

$$\|\mathcal{P}_{\mathbf{X}}\mathbf{E}\|_{\text{op}} \lesssim \sqrt{n} \cdot 1 \cdot \sqrt{\frac{\log(nd)}{n}} \cdot \sqrt{p}$$

with probability at least  $1 - n^{-\delta}$ . The above results suggest that

$$\|\Delta\|_{\text{op}} \leq \sqrt{c(d+r)(n \vee p)}$$

Below we condition on the two events above, which hold with probability at least  $1 - 2(n +$

$p)^{-\delta} - 2(np)^{-\delta} - \exp(-n)$  by union bound.

**Part (2) Bounding  $\|\mathbf{E}_j\|_2$  and  $\|\Delta_j\|_2$**  From Lemma C.3 and Assumption 3, we have

$$\max_{1 \leq j \leq p} \|\mathbf{E}_j\|_2 = \max_{1 \leq j \leq p} \|\mathbf{W}\boldsymbol{\gamma}_j\|_2 \leq \|\mathbf{W}\|_{\text{op}} \max_{1 \leq j \leq p} \|\boldsymbol{\gamma}_j\|_2 \leq 2C_\Sigma \sqrt{n}.$$

On the other hand, because  $\widehat{\mathbf{W}}^\top \widehat{\mathbf{W}} = n\boldsymbol{\Sigma}$  and  $\|\widehat{\boldsymbol{\gamma}}_j\|_2 \leq C^2 C_\Sigma^{1/2}$  for all  $j \in [p]$  from Lemma C.4, we also have

$$\max_{1 \leq j \leq p} \|\widehat{\mathbf{E}}_j\|_2 = \max_{1 \leq j \leq p} \|\widehat{\mathbf{W}}\widehat{\boldsymbol{\gamma}}_j\|_2 = \max_{1 \leq j \leq p} \sqrt{n}\|\widehat{\boldsymbol{\gamma}}_j\|_2 \lesssim C^2 C_\Sigma^{3/2} \sqrt{n}.$$

Thus, by triangle inequality, we have

$$\max_{1 \leq j \leq p} \|\Delta_j\|_2 \leq \max_{1 \leq j \leq p} (\|\mathbf{E}_j\|_2 + \|\widehat{\mathbf{E}}_j\|_2) \lesssim (2 + C^2)C_\Sigma \sqrt{n}.$$

**Part (3) Combining the previous results.** From (C.3), (C.4) and the previous two parts, we have

$$\begin{aligned} & \max_{1 \leq j \leq p} \|\widehat{\boldsymbol{\gamma}}_j - \widetilde{\boldsymbol{\gamma}}_j\|_2 \\ & \lesssim \frac{1}{n\sqrt{p}C_\Sigma} (2C_\Sigma \sqrt{np}(2 + C^2)C_\Sigma \sqrt{n} + \sqrt{c(d+r)(n \vee p)}C\sqrt{n} + \sqrt{c(d+r)(n \vee p)}(2C_\Sigma + C)\sqrt{n}) \\ & \lesssim 2(2 + C^2)C_\Sigma \end{aligned}$$

and

$$\|\widehat{\boldsymbol{\Gamma}} - \widetilde{\boldsymbol{\Gamma}}\|_2 \leq \frac{1}{n\sqrt{p}C_\Sigma} (4C_\Sigma \sqrt{np}\sqrt{c(d+r)(n \vee p)} + c(d+r)(n \vee p)) \lesssim 4\sqrt{c},$$

which finishes the proof.  $\square$

**Lemma C.2** (Spectrum of  $\boldsymbol{\Sigma}$ ). Under Assumptions 1–3 and event  $E_C$ , for all  $\delta > 0$  and sufficiently large  $n$  and  $p$ , there exists a absolute constant  $C_\Sigma > 1$  such that

$$\frac{1}{C_\Sigma} \leq \lambda_r(\boldsymbol{\Sigma}) \leq \lambda_1(\boldsymbol{\Sigma}) \leq C_\Sigma,$$

with probability at least  $1 - (n + p)^{-\delta} - (np)^{-\delta} - \exp(-n)$ .

*Proof of Lemma C.2.* In the following proof, for simplicity, we drop the superscript and use  $\mathbf{W}$ ,  $\boldsymbol{\Gamma}$ , and  $\boldsymbol{\Theta}$  to denote the true parameters. By Weyl's inequality, we have that for all  $k \in [r]$ ,

$$\begin{aligned} \left| \lambda_k(\boldsymbol{\Sigma}) - \frac{1}{\sqrt{np}} \lambda_k(\mathbf{W}\boldsymbol{\Gamma}^\top) \right| &= \frac{1}{\sqrt{np}} \left| \lambda_k(\widehat{\mathbf{W}}\widehat{\boldsymbol{\Gamma}}^\top) - \lambda_k(\mathbf{W}\boldsymbol{\Gamma}^\top) \right| \\ &\leq \frac{1}{\sqrt{np}} \|\widehat{\mathbf{W}}\widehat{\boldsymbol{\Gamma}}^\top - \mathbf{W}\boldsymbol{\Gamma}^\top\|_{\text{op}} \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{\sqrt{np}} \|\mathcal{P}_{\mathbf{X}}^\perp(\widehat{\Theta} - \Theta)\|_{\text{op}} \\
&\leq \frac{1}{\sqrt{np}} \|\widehat{\Theta} - \Theta\|_{\text{op}}.
\end{aligned} \tag{C.5}$$

We next bound  $\lambda_k(\mathbf{W}\mathbf{\Gamma}^\top)$  and  $\|\widehat{\Theta} - \Theta\|_{\text{op}}$  separately. Applying Lemma C.3 under Assumption 3 yields that

$$\frac{1}{C'_0} \leq \lambda_r\left(\frac{1}{n}\mathbf{W}^\top\mathbf{W}\right) \leq \lambda_1\left(\frac{1}{n}\mathbf{W}^\top\mathbf{W}\right) \leq C'_0.$$

with probability at least  $1 - \exp(-n)$  for some constant  $C'_0 > 1$ . From Assumption 3 we further have

$$\sqrt{\frac{1}{CC'_0}} \leq \frac{1}{\sqrt{np}} \lambda_r(\mathbf{W}\mathbf{\Gamma}^\top) \leq \frac{1}{\sqrt{np}} \lambda_1(\mathbf{W}\mathbf{\Gamma}^\top) \leq \sqrt{CC'_0}. \tag{C.6}$$

On the other hand, from Theorem 2 we have for all  $\delta > 0$ , there exists  $C > 0$  such that

$$\frac{1}{\sqrt{np}} \|\widehat{\Theta} - \Theta\|_{\text{op}} \leq \frac{1}{\sqrt{np}} \|\widehat{\Theta} - \Theta\|_{\text{F}} \leq C \sqrt{\frac{r(n \vee p)}{np}} =: C_{n,p}, \tag{C.7}$$

with probability at least  $1 - (n+p)^{-\delta} - (np)^{-\delta}$ .

Condition on the above two events, applying triangle inequality on (C.5) and combining (C.6) and (C.7), we have

$$\sqrt{\frac{1}{CC'_0}} - C_{n,p} \leq \lambda_r(\mathbf{\Sigma}) \leq \lambda_1(\mathbf{\Sigma}) \leq \sqrt{CC'_0} + C_{n,p}.$$

Note that  $C_{n,p} = o(1)$  as both  $n$  and  $p$  tend to infinity. When  $n$  and  $p$  is such that  $C_{n,p} < 1/2\sqrt{CC'_0}$ , setting  $C_\Sigma = 3\sqrt{CC'_0}/2$  gives the desired bound. By union bound, this holds with probability at least  $1 - (n+p)^{-\delta} - (np)^{-\delta} - \exp(-n)$ , which finishes the proof.  $\square$

**Lemma C.3** (Spectrum of  $\mathbf{W}$ ). Under Assumption 3, for sufficiently large  $n$ , there exists a absolute constant  $C'_0 > 0$  such that

$$\frac{1}{C'_0} \leq \lambda_r\left(\frac{1}{n}\mathbf{W}^\top\mathbf{W}\right) \leq \lambda_1\left(\frac{1}{n}\mathbf{W}^\top\mathbf{W}\right) \leq C'_0,$$

with probability at least  $1 - \exp(-n)$ .

*Proof of Lemma C.3.* Note that  $\mathbf{W} \in \mathbb{R}^{n \times r}$  is a random matrix whose rows  $\mathbf{w}_1, \dots, \mathbf{w}_n$  are i.i.d. sub-Gaussian random vectors. From the concentration inequality of the operator norm of the random matrices (Vershynin, 2018, Theorem 4.6.1, Exercise 4.7.3), for any  $u > 0$  we



have

$$\left\| \frac{1}{n} \mathbf{W}^\top \mathbf{W} - \boldsymbol{\Sigma}_w \right\|_{\text{op}} \leq CK^2 \left( \sqrt{\frac{r+u}{n}} + \frac{r+u}{n} \right) \|\boldsymbol{\Sigma}_w\|_{\text{op}},$$

with probability at least  $1 - 2\exp(-u)$ , where  $C > 0$  is some absolute constant and  $K = \max_{i \in [n]} \|\mathbf{w}_i\|_{\psi_2}$ . When  $n$  is sufficiently large such that  $2CK^2n^{-1/4} < 1$ , setting  $u = n^{1/2}$  and  $C' = 2CK^2n^{-1/4}C_0$  yields that

$$\left\| \frac{1}{n} \mathbf{W}^\top \mathbf{W} - \boldsymbol{\Sigma}_w \right\|_{\text{op}} \leq CK^2 \left( n^{-\frac{1}{4}} + n^{-\frac{1}{2}} \right) \|\boldsymbol{\Sigma}_w\|_{\text{op}} \leq C' < C_0.$$

By Weyl's inequality, we have that,

$$\max_{k \in [r]} \left| \lambda_k \left( \frac{1}{n} \mathbf{W}^\top \mathbf{W} \right) - \lambda_k(\boldsymbol{\Sigma}_w) \right| \leq \left\| \frac{1}{n} \mathbf{W}^\top \mathbf{W} - \boldsymbol{\Sigma}_w \right\|_{\text{op}} \leq C'.$$

By triangle inequality and the boundedness of  $\boldsymbol{\Sigma}_w$ 's spectrum from Assumption 3, it follows that

$$\frac{1}{C_0} - C' \leq \min_{k \in [r]} \lambda_k \left( \frac{1}{n} \mathbf{W}^\top \mathbf{W} \right) \leq \max_{k \in [r]} \lambda_k \left( \frac{1}{n} \mathbf{W}^\top \mathbf{W} \right) \leq C_0 + C',$$

with probability at least  $1 - \exp(-n)$ . Setting  $C'_0 = \min\{C_0 + C', (C_0^{-1} - C')^{-1}\}$  finishes the proof.  $\square$

**Lemma C.4** (Boundedness of latent factors and loadings). Under Assumptions 1–3 and event  $E_C$ , for all  $\delta > 0$  and sufficiently large  $n$  and  $p$ , there exists a absolute constant  $C_\Sigma > 1$  such that  $\|\widehat{\boldsymbol{\gamma}}\|_2 \leq C^2 C_\Sigma^{1/2}$ , with probability at least  $1 - (n+p)^{-\delta} - (np)^{-\delta} - \exp(-n)$ .

*Proof of Lemma C.4.* Recall that  $\widehat{\mathbf{W}}_0$  and  $\widehat{\boldsymbol{\Gamma}}_0$  are the solutions to the alternative maximization problems which satisfy that  $\|\widehat{\mathbf{w}}_{0,i}\|_2 \leq C$  and  $\|\widehat{\boldsymbol{\gamma}}_{0,j}\|_2 \leq C$  for  $i = 1, \dots, n$  and  $1, \dots, p$ . Let  $\widehat{\mathbf{W}}_0 \widehat{\boldsymbol{\Gamma}}_0^\top = \sqrt{np} \mathbf{U} \boldsymbol{\Sigma} \mathbf{V}^\top$  be the condensed SVD. Then  $\widehat{\boldsymbol{\Gamma}}$  is defined to be  $\sqrt{p} \mathbf{V} \boldsymbol{\Sigma}^{1/2} = \widehat{\boldsymbol{\Gamma}}_0 \widehat{\mathbf{W}}_0^\top \mathbf{U} \boldsymbol{\Sigma}^{1/2} / \sqrt{n}$ . From Lemma C.2, with probability at least  $1 - (n+p)^{-\delta} - (np)^{-\delta} - \exp(-n)$ , it holds that

$$\begin{aligned} \|\widehat{\boldsymbol{\gamma}}_j\|_2 &\leq \|\widehat{\mathbf{W}}_0 / \sqrt{n}\|_{\text{op}} \|\mathbf{U}\|_{\text{op}} \|\boldsymbol{\Sigma}^{1/2}\|_{\text{op}} \|\widehat{\boldsymbol{\gamma}}_{0,j}\|_2 \\ &\leq \max_{1 \leq i \leq n} \|\widehat{\mathbf{W}}_{0,i}\|_2 \cdot 1 \cdot C_\Sigma^{1/2} \cdot C \\ &\leq C_\Sigma^{1/2} C^2, \end{aligned}$$

which finishes the proof.  $\square$

## D Estimation of latent factors and direct effects

### D.1 Preparatory definitions

Towards proving Theorem 5, we first introduce the following notations. Recall that optimization problem (3.2) is the multivariate lasso with nuisance parameter. Define the response vector  $\tilde{\mathbf{y}} = \text{vec}(\mathbf{Y}) \in \mathbb{R}^{np}$ , the design matrices  $\tilde{\mathbf{X}} = (\mathbf{I}_p \otimes \mathbf{X}) \in \mathbb{R}^{np \times pd}$ ,  $\boldsymbol{\beta} = \text{vec}(\mathbf{B})$ ,  $\boldsymbol{\zeta} = \text{vec}(\mathbf{Z}\boldsymbol{\Gamma}^\top)$ , and the projection matrix  $\tilde{\mathcal{P}}_\Gamma = (\mathcal{P}_\Gamma \otimes \mathbf{I}_d) \in \mathbb{R}^{pd \times pd}$ . Here, symbol ‘ $\otimes$ ’ denotes the Kronecker product. With slight abuse of notations, we use  $\mathcal{L}(\boldsymbol{\beta}, \boldsymbol{\zeta})$  to denote the unregularized loss function of (3.2). Let  $\mathcal{F} = \{(\boldsymbol{\beta}, \boldsymbol{\zeta}) \in \mathbb{R}^{pd} \times \mathbb{R}^{np} \mid \boldsymbol{\beta} = \text{vec}(\mathbf{B}), \boldsymbol{\zeta} = \text{vec}(\mathbf{Z}\boldsymbol{\Gamma}^\top) \text{ for } \mathbf{B} \in \mathbb{R}^{p \times d}, \mathbf{Z} \in \mathbb{R}^{n \times r}, \boldsymbol{\Gamma} \in \mathbb{R}^{p \times r} \text{ such that } \mathcal{P}_\Gamma \mathbf{B} = \mathbf{0}, \tilde{\mathbf{X}}\boldsymbol{\beta} + \boldsymbol{\zeta} \in \mathcal{R}_C^{np}\}$  be the feasible set of  $\boldsymbol{\beta}$  and  $\boldsymbol{\zeta}$ . Then the joint optimization problem (3.2) is equivalent to

$$\hat{\boldsymbol{\beta}}, \hat{\boldsymbol{\zeta}} \in \underset{(\boldsymbol{\beta}, \boldsymbol{\zeta}) \in \mathcal{F}}{\text{argmin}} \mathcal{L}(\boldsymbol{\beta}, \boldsymbol{\zeta}) + \lambda \|\boldsymbol{\beta}\|_1. \quad (\text{D.1})$$

Let  $(\tilde{\boldsymbol{\beta}}^*, \tilde{\boldsymbol{\zeta}}^*) = (\text{vec}(\mathcal{P}_{\Gamma^*}^\perp \mathbf{B}^*), \text{vec}(\mathbf{X}\mathbf{B}^*\mathcal{P}_{\Gamma^*} + \mathbf{Z}^*\boldsymbol{\Gamma}^{*\top}))$ ,  $(\boldsymbol{\beta}^*, \boldsymbol{\zeta}^*) = (\text{vec}(\mathbf{B}^*), \text{vec}(\mathbf{Z}^*\boldsymbol{\Gamma}^{*\top}))$  denote the tuples of target coefficients and note that  $(\tilde{\boldsymbol{\beta}}^*, \tilde{\boldsymbol{\zeta}}^*) \in \mathcal{F}$ .

The organization of the following subsections is summarized as below:

- Appendix D.2 proves Corollary 4, which controls the column-wise  $\ell_2$ -norm of the estimation error of the latent component.
- Appendix D.3 proves Theorem 5. The proof of Theorem 5 consists of three main steps:
  - (1) Establish cone condition: We involve Lemma D.1 to show that the estimation from the sequential optimization problems (3.4)-(3.5) obtain approximately optimality condition to the joint optimization problem (3.2):

$$\mathcal{L}(\hat{\mathbf{B}}, \hat{\boldsymbol{\Gamma}}, \hat{\mathbf{Z}}) + \lambda \|\hat{\mathbf{B}}\|_{1,1} \leq \mathcal{L}(\mathbf{B}^*, \boldsymbol{\Gamma}^*, \mathbf{Z}^*) + \lambda \|\mathbf{B}^*\|_{1,1} + \tau_{n,p},$$

for some small order term  $\tau_{n,p}$  with high probability. This enables us to derive the cone condition.

- (2) Obtain upper and lower bound of the first-order approximation error: We involve Lemma D.2 to derive the upper bound, and Lemma D.3 to establish the locally strong convexity and hence the lower bound.
  - (3) Derive the estimation errors: We compute the  $\ell_2$ -norm and  $\ell_1$ -norm estimation error based on the previous two steps.
- Appendix D.4 gathers helper lemmas used in the current section.

### D.2 Proof of Corollary 4

*Proof of Corollary 4.* Define  $\hat{\mathbf{E}} = \hat{\mathbf{Z}}\hat{\boldsymbol{\Gamma}}^\top$  and  $\mathbf{E}^* = \mathbf{Z}^*\boldsymbol{\Gamma}^{*\top}$ . Let  $\hat{\mathbf{B}} = \underset{\{\mathbf{B} \in \mathbb{R}^{p \times d} \mid \mathcal{P}_\Gamma \mathbf{B} = \mathbf{0}\}}{\text{argmin}} \mathcal{L}(\mathbf{X}\mathbf{B}^\top + \hat{\mathbf{E}})$  and  $\hat{\boldsymbol{\Theta}} = \mathbf{X}\hat{\mathbf{B}}^\top + \hat{\mathbf{E}}$ . Since  $\mathcal{L}(\hat{\boldsymbol{\Theta}}) \leq \mathcal{L}(\boldsymbol{\Theta}^*)$  as assumed in Corollary 4, from Theorem 2

we have  $\|\widehat{\Theta} - \Theta^*\| \lesssim \mathcal{O}_{\mathbb{P}}(\sqrt{n \vee p})$ . From Theorem 3, we further have

$$\|\widehat{\mathbf{E}} - \mathbf{E}^*\|_{\text{F}} \leq \|(\widehat{\Theta} - \Theta^*)\mathcal{P}_{\widehat{\Gamma}}\|_{\text{F}} + \|\Theta^*(\mathcal{P}_{\widehat{\Gamma}} - \mathcal{P}_{\Gamma^*})\|_{\text{F}} \lesssim \mathcal{O}_{\mathbb{P}}(\sqrt{n \vee p}).$$

Then from Lin et al. (2021, Proposition 5.2) we have that, up to sign,

$$\frac{1}{n} \|\widehat{\mathbf{Z}} - \mathbf{Z}^* \mathbf{R}^{\top}\|_{\text{F}}^2 \lesssim \frac{r^{4k_2 - k_1 + 4}}{n \wedge p} =: \eta_n,$$

with probability at least  $1 - n^{-c} - p^{-c}$ .

Recall the invertible matrix  $\mathbf{R}$  with  $\|\mathbf{R}\|_{\text{op}} = \mathcal{O}_{\mathbb{P}}(1)$  from Assumption 4 and define the transformed parameters  $\widetilde{\mathbf{Z}}^* = \mathbf{Z}^* \mathbf{R}^{\top}$  and  $\widetilde{\Gamma}^* = \Gamma^* \mathbf{R}^{-1}$ . Because  $\mathbf{E}_j - \mathbf{E}_j^* = \widetilde{\mathbf{Z}}^*(\gamma_j - \widetilde{\gamma}_j^*) + (\mathbf{Z} - \widetilde{\mathbf{Z}}^*)\gamma_j$ , with probability tending to one, it follows that

$$\begin{aligned} \max_{1 \leq j \leq p} \frac{1}{\sqrt{n}} \|\mathbf{E}_j - \mathbf{E}_j^*\|_2 &= \max_{1 \leq j \leq p} \frac{1}{\sqrt{n}} \|\widetilde{\mathbf{Z}}^*(\gamma_j - \widetilde{\gamma}_j^*) + (\mathbf{Z} - \widetilde{\mathbf{Z}}^*)\gamma_j\|_2 \\ &\leq \max_{1 \leq j \leq p} \frac{1}{\sqrt{n}} \|\widetilde{\mathbf{Z}}^*(\gamma_j - \widetilde{\gamma}_j^*)\|_2 + \max_{1 \leq j \leq p} \frac{1}{\sqrt{n}} \|(\mathbf{Z} - \widetilde{\mathbf{Z}}^*)\gamma_j\|_2 \\ &\leq \max_{1 \leq j \leq p, 1 \leq i \leq n} \frac{1}{\sqrt{n}} \|\widetilde{\mathbf{z}}_i^*\|_{\infty} \|\gamma_j - \widetilde{\gamma}_j^*\|_1 + \max_{1 \leq j \leq p} \frac{1}{\sqrt{n}} \|\mathbf{Z} - \widetilde{\mathbf{Z}}^*\|_{\text{op}} \|\gamma_j\|_2 \\ &\lesssim \frac{\log n}{\sqrt{n}} + \sqrt{\eta_n} \\ &\lesssim \sqrt{n^{-1} \log n \vee \eta_n} \end{aligned}$$

where the second last inequality is because that  $\|\widetilde{\gamma}_j^*\|_2 \leq C$  from Assumption 3,  $\|\gamma_j - \widetilde{\gamma}_j^*\|_1 \leq \sqrt{r} \|\gamma_j - \widetilde{\gamma}_j^*\|_2 \leq 2\sqrt{r}C$  from Lemma C.1, and  $\widetilde{\mathbf{z}}_i^*$ 's are independent  $r$ -dimensional sub-Gaussian random vectors from Lemma D.5 so that  $\max_{1 \leq i \leq n} \|\widetilde{\mathbf{z}}_i^*\|_{\infty}$  scales in  $\log(nr)$ .  $\square$

### D.3 Proof of Theorem 5

*Proof of Theorem 5.* Define  $\Delta_{\beta} = \widehat{\beta} - \beta^*$ ,  $\Delta_{\zeta} = \widehat{\zeta} - \zeta^*$ , and  $\mathcal{S} = \text{supp}(\beta^*)$ . Let  $\mathbf{g}_{\beta} = \nabla_{\beta} \mathcal{L}(\widehat{\beta}, \widehat{\zeta})$ , and  $\mathbf{g}_{\zeta} = \nabla_{\zeta} \mathcal{L}(\widehat{\beta}, \widehat{\zeta})$  and analogously define  $\mathbf{g}_{\beta}^*, \mathbf{g}_{\zeta}^*$ .

(1) **Cone condition.** From Lemma D.1, we have the optimality condition

$$\mathcal{L}(\widehat{\beta}, \widehat{\zeta}) + \lambda \|\widehat{\beta}\|_1 \leq \mathcal{L}(\beta^*, \zeta^*) + \lambda \|\beta^*\|_1 + \tau_{n,p},$$

with  $\tau_{n,p}$  defined in Lemma D.1. Rearranging the above display, it follows that

$$\begin{aligned} \lambda \|\widehat{\beta}\|_1 &\leq \mathcal{L}(\beta^*, \zeta^*) - \mathcal{L}(\widehat{\beta}, \widehat{\zeta}) + \lambda \|\beta^*\|_1 + \tau_{n,p} \\ &\leq \Delta_{\beta}^{\top} \mathbf{g}_{\beta}^* + \Delta_{\zeta}^{\top} \mathbf{g}_{\zeta}^* + \lambda \|\beta^*\|_1 + \tau_{n,p} \\ &\leq \|\Delta_{\beta}\|_1 \|\mathbf{g}_{\beta}^*\|_{\infty} + \Delta_{\zeta}^{\top} \mathbf{g}_{\zeta}^* + \lambda \|\beta^*\|_1 + \tau_{n,p}, \end{aligned} \tag{D.2}$$

where the second inequality is from the convexity of  $\mathcal{L}$ , and the last is from Holder's inequality. The term involves nuisance parameters can be further bounded as

$$\mathbf{\Delta}_\zeta^\top \mathbf{g}_\zeta^* = \frac{1}{n} \sum_{\ell=1}^{np} [\tilde{y}_\ell - A'(\tilde{\mathbf{x}}_\ell^\top \boldsymbol{\beta}^* + \zeta_\ell^*)] \delta_{\zeta,\ell} \leq 2C \|\mathbf{g}_\zeta^*\|_\infty \leq \frac{4cC^2\alpha \log(2np)}{n}, \quad (\text{D.3})$$

where the last inequality is from Lemma D.2 (2), which holds with probability at least at least  $1 - (n+p)^{-c} - (np)^{-c} - \exp(-n)$  for some  $c > 0$ . On the other hand, the left hand side of (D.2) is lower bounded as

$$\begin{aligned} \|\widehat{\boldsymbol{\beta}}\|_1 &= \|\boldsymbol{\beta}^* + \mathbf{\Delta}_\beta\|_1 \\ &= \|\boldsymbol{\beta}_\mathcal{S}^* + \mathbf{\Delta}_{\beta,\mathcal{S}}\|_1 + \|\mathbf{\Delta}_{\beta,\mathcal{S}^c}\|_1 \\ &\geq \|\boldsymbol{\beta}_\mathcal{S}^*\|_1 - \|\mathbf{\Delta}_{\beta,\mathcal{S}}\|_1 + \|\mathbf{\Delta}_{\beta,\mathcal{S}^c}\|_1, \end{aligned} \quad (\text{D.4})$$

where  $\mathcal{S} = \{j \in [pd] \mid \beta_j^* \neq 0\}$  is the active set of the true coefficients. Combining (D.2), (D.3), and (D.4) yields that

$$(\lambda - \|\mathbf{g}_\beta^*\|_\infty) \|\mathbf{\Delta}_{\beta,\mathcal{S}^c}\|_1 \leq (\lambda + \|\mathbf{g}_\beta^*\|_\infty) \|\mathbf{\Delta}_{\beta,\mathcal{S}}\|_1 + \left( \frac{4cC^2\alpha \log(2np)}{n} + \tau_{n,p} \right).$$

Note that under the same probabilistic event above, from Lemma D.2 (1), we have

$$\|\mathbf{g}_\beta^*\|_\infty \leq 4\nu^2 \sqrt{c \log^2(2nd)/n}. \quad (\text{D.5})$$

When  $\lambda^* \asymp 8\nu^2 \sqrt{c \log^2(2nd)/n} \geq 2\|\mathbf{g}_\beta^*\|_\infty$ , this implies the approximate cone condition  $\mathbf{\Delta}_\beta \in \mathcal{C}(3, \mathcal{S})$ , where

$$\mathcal{C}(\xi, \mathcal{S}) := \left\{ \mathbf{\Delta}_\beta \in \mathbb{R}^{pd} \mid \|\mathbf{\Delta}_{\beta,\mathcal{S}^c}\|_1 \leq \xi \|\mathbf{\Delta}_{\beta,\mathcal{S}}\|_1 + \tau_{n,p}^* \right\}, \quad (\text{D.6})$$

and

$$\tau_{n,p}^* = \frac{C^2\alpha}{\nu^2} \sqrt{\frac{c \log^2(2np)}{n \log^2(2nd)}} + \frac{\sqrt{n}}{(n \wedge p) \log(2nd)} + \sqrt{\frac{(sd)^2}{n \wedge p^{1-k}}}.$$

From the cone condition (D.6), the  $\ell_1$ -norm bound follows by observing that

$$\begin{aligned} \|\mathbf{\Delta}_\beta\|_1 &\leq \|\mathbf{\Delta}_{\beta,\mathcal{S}}\|_1 + \|\mathbf{\Delta}_{\beta,\mathcal{S}^c}\|_1 \\ &\leq 4\|\mathbf{\Delta}_{\beta,\mathcal{S}}\|_1 + \tau_{n,p}^* \\ &\leq 4\sqrt{sd} \|\mathbf{\Delta}_{\beta,\mathcal{S}}\|_2 + \tau_{n,p}^* \\ &\leq 4\sqrt{sd} \|\mathbf{\Delta}_\beta\|_2 + \tau_{n,p}^*. \end{aligned} \quad (\text{D.7})$$

**(2) Upper and lower bound of the first-order approximation error.** To quantify the estimation of the coefficient, we next analyze the first-order approximation error of the normalized likelihood:  $\mathcal{E}(\hat{\beta}, \beta^*; \hat{\zeta}, \zeta^*) = \Delta_\beta^\top (\mathbf{g}_\beta - \mathbf{g}_\beta^*)$ . By the first-order optimality condition of convex optimization problem (3.3), we have

$$\begin{aligned} \mathcal{E}(\hat{\beta}, \beta^*; \hat{\zeta}, \zeta^*) &\leq -\Delta_\beta^\top \mathbf{g}_\beta^* \\ &\leq \|\Delta_\beta\|_1 \|\mathbf{g}_\beta^*\|_\infty \\ &\leq \|\Delta_\beta\|_2 16\nu^2 \sqrt{\frac{csd \log^2(2nd)}{n}} + 4C^2 \alpha \sqrt{c} \frac{\log(2np)}{n}, \end{aligned} \quad (\text{D.8})$$

where the second inequality is from Holder's inequality, and the last inequality is from (D.5) and (D.7). This establishes the upper bound for  $\mathcal{E}(\hat{\beta}, \beta^*; \hat{\zeta}, \zeta^*)$ .

On the other hand, Lemma D.3 implies that  $\mathcal{E}(\hat{\beta}, \beta^*; \hat{\zeta}, \zeta^*)$  is locally restricted strongly convex over an augmented cone  $\mathcal{C}_a(3, \mathcal{S}, \eta_n)$ , defined in (D.20):

$$\inf_{(\Delta_\beta, \Delta_\zeta) \in \mathcal{C}_a(3, \mathcal{S}, \eta_n)} \mathcal{E}(\beta^* + \Delta_\beta, \beta^*; \zeta^* + \Delta_\zeta, \zeta^*) \geq \frac{\kappa_1 C'}{2} \|\Delta_\beta\|_2^2 - C'' \sqrt{n^{-1} \vee \eta_n} \|\Delta_\beta\|_1, \quad (\text{D.9})$$

for some constant  $C' > 0$ , with probability at least  $1 - 2 \exp(-3 \log n)$ .

**(3) Estimation error.** From Part (2), (D.8) and (D.9) imply that

$$\frac{\kappa_1 C'}{2} \|\Delta_\beta\|_2^2 \leq \|\Delta_\beta\|_2 \left( 16\nu^2 \sqrt{\frac{csd \log^2(2nd)}{n}} + 4C'' \sqrt{\frac{sd}{n \wedge p}} \right) + 4C^2 \alpha \sqrt{c} \frac{\log(2np)}{n} + \frac{C'' r^{4k_2 - k_1 + 4}}{\sqrt{n \wedge p}} \tau_{n,p}^*.$$

over  $\mathcal{C}_a(3, \mathcal{S}, \eta_n)$ . This implies that, with probability at least  $1 - (n+p)^{-c'} - (np)^{-c'} - \exp(-n)$  for some  $c' > 0$ ,

$$\begin{aligned} \|\Delta_\beta\|_2 &\leq \frac{2}{\kappa_1 C'} \left( 16\nu^2 \sqrt{\frac{csd \log^2(2nd)}{n}} + C'' \sqrt{\frac{sd}{n \wedge p}} \right) + \sqrt{\frac{2}{\kappa_1 C'}} \sqrt{4C^2 \alpha \sqrt{c} \frac{\log(2np)}{n} + \frac{C'' r^{4k_2 - k_1 + 4} \tau_{n,p}^*}{\sqrt{n \wedge p}}} \\ &\lesssim \sqrt{\frac{(sd \log^2(nd)) \vee \log(np)}{n}} + \frac{n^{1/4}}{(n \wedge p)^{3/2} \log^{1/2}(nd)} + \sqrt{\frac{sd}{n \wedge p^{1-k}}}. \end{aligned} \quad (\text{D.10})$$

To establish the  $\ell_1$ -norm bound, from (D.7) and (D.10), we have

$$\begin{aligned} \|\Delta_\beta\|_1 &\leq 4\sqrt{sd} \|\Delta_\beta\|_2 + \tau_{n,p}^* \\ &\lesssim \sqrt{sd \frac{(sd \log^2(nd)) \vee \log(np)}{n}} + \sqrt{\frac{(sd)^2}{n \wedge p^{1-k}}} + \frac{\sqrt{n}}{(n \wedge p) \log(nd)}, \end{aligned} \quad (\text{D.11})$$

which completes the proof.  $\square$

## D.4 Technical lemmas

**Lemma D.1** (Sequential and joint optimization). Under the same conditions in Theorem 5, for any constant  $\delta > 0$  it holds with probability at least  $1 - (n + p)^{-\delta} - (np)^{-\delta} - \exp(-n)$  that

$$\mathcal{L}(\widehat{\boldsymbol{\beta}}, \widehat{\boldsymbol{\zeta}}) + \lambda \|\widehat{\boldsymbol{\beta}}\|_1 \leq \mathcal{L}(\boldsymbol{\beta}^*, \boldsymbol{\zeta}^*) + \lambda \|\boldsymbol{\beta}^*\|_1 + \tau_{n,p},$$

where

$$\tau_{n,p} = \mathcal{O} \left( \frac{1}{n \wedge p} + \lambda \sqrt{\frac{(sd)^2}{n \wedge p^{1-k}}} \right).$$

*Proof of Lemma D.1.* From Assumption 1, the entry of  $\mathbf{B}^*$  is bounded because  $\|\mathbf{B}^*\|_{\max} = \max_{1 \leq i \leq p, 1 \leq j \leq d} |b_{ij}| \leq \max_{1 \leq i \leq p} \|\mathbf{b}_i\|_{\infty} \leq \max_{1 \leq i \leq p} \|\mathbf{b}_i\|_2 \leq C$ . From Proposition 1 and Assumption 1, we further have that

$$\|\mathcal{P}_{\Gamma^*} \mathbf{B}^*\|_{\text{F}} \lesssim \sqrt{sd/p} \|\mathbf{B}^*\|_{\max} \lesssim \sqrt{sd/p}. \quad (\text{D.12})$$

Thus, from the assumption of Theorem 5, we have that

$$\|\mathcal{P}_{\Gamma^*} \mathbf{B}^*\|_{1,1} = \mathcal{O}(p^{k/2} \|\mathcal{P}_{\Gamma^*} \mathbf{B}^*\|_{\text{F}}) \lesssim \sqrt{sd} p^{(k-1)/2}. \quad (\text{D.13})$$

This ensures that  $\|\widetilde{\boldsymbol{\beta}}^*\|_{1,1}$  is close to  $\|\boldsymbol{\beta}^*\|_{1,1}$ .

From optimality condition of optimization problem (3.5), we have

$$\mathcal{L}(\widehat{\boldsymbol{\beta}}, \widehat{\boldsymbol{\zeta}}) + \lambda \|\widehat{\boldsymbol{\beta}}\|_1 \leq \mathcal{L}(\widetilde{\boldsymbol{\beta}}', \widetilde{\boldsymbol{\zeta}}') + \lambda \|\widetilde{\boldsymbol{\beta}}'\|_1,$$

where  $\widetilde{\boldsymbol{\beta}}' = \text{vec}(\mathcal{P}_{\widehat{\Gamma}}^{\perp} \mathbf{B}^*)$  and  $\widetilde{\boldsymbol{\zeta}}' = \text{vec}(\boldsymbol{\Theta}^* \mathcal{P}_{\widehat{\Gamma}})$ . It follows that

$$\mathcal{L}(\widehat{\boldsymbol{\beta}}, \widehat{\boldsymbol{\zeta}}) + \lambda \|\widehat{\boldsymbol{\beta}}\|_1 \leq \mathcal{L}(\boldsymbol{\Theta}^* - \mathbf{Z}^* \boldsymbol{\Gamma}^{*\top} \mathcal{P}_{\widehat{\Gamma}}^{\perp}) + \lambda \|\mathbf{B}^{*\top} \mathcal{P}_{\widehat{\Gamma}}^{\perp}\|_{1,1}. \quad (\text{D.14})$$

We next bound the first term in (D.14). From the proof of Lemma B.1, we have

$$\mathcal{L}(\boldsymbol{\Theta}^* - \mathbf{Z}^* \boldsymbol{\Gamma}^{*\top} \mathcal{P}_{\widehat{\Gamma}}^{\perp}) - \mathcal{L}(\boldsymbol{\Theta}^*) \leq \underbrace{\frac{1}{n} \text{tr}((\mathbf{Y} - \mathbf{A}'(\boldsymbol{\Theta}^*))^{\top} \mathbf{Z}^* \boldsymbol{\Gamma}^{*\top} \mathcal{P}_{\widehat{\Gamma}}^{\perp})}_{T_1} + \underbrace{\frac{\kappa_2}{2n} \|\mathbf{Z}^* \boldsymbol{\Gamma}^{*\top} \mathcal{P}_{\widehat{\Gamma}}^{\perp}\|_{\text{F}}^2}_{T_2}.$$

Let  $\mathbf{A} = n^{-1} \mathbf{Z}^* \boldsymbol{\Gamma}^{*\top} \mathcal{P}_{\widehat{\Gamma}}^{\perp}$ . From Lemmas D.4 and D.5, the second term  $T_2$  can be upper bounded as

$$T_2 = \frac{\kappa_2}{2} n \|\mathbf{A}\|_{\text{F}}^2 = \frac{\kappa_2}{2n} \|\mathbf{Z}^* \boldsymbol{\Gamma}^{*\top} \mathcal{P}_{\widehat{\Gamma}}^{\perp}\|_{\text{F}}^2 \leq \frac{\kappa_2}{2n} \|\mathbf{Z}^*\|_{\text{op}}^2 \|\boldsymbol{\Gamma}^{*\top} \mathcal{P}_{\widehat{\Gamma}}^{\perp}\|_{\text{F}}^2 = \mathcal{O} \left( \frac{r}{n \wedge p} \right),$$

with probability at least  $1 - (n + p)^{-\delta} - (np)^{-\delta} - \exp(-n)$ . In the equality, we use the fact that  $\mathbf{Z}^*$  is a matrix with independent sub-Gaussian rows to obtain similar results as Lemma C.2.

For the first term  $T_1$ , note that  $y_{ij} - A'(\theta_{ij}^*)$  is mean-zero  $(\nu, \alpha)$ -sub-exponential random variable when conditioned on  $(\mathbf{X}^*, \mathbf{Z}^*)$ , where  $\nu = \sqrt{\kappa_2}$  and  $\alpha = 1/C^2$ , as shown in the proof of Theorem 2. To bound the first term  $T_1$ , we apply Bernstein's inequality (Vershynin, 2018, Theorem 2.8.2) to obtain

$$\mathbb{P}(|T_1| \geq t \mid \mathbf{X}^*, \mathbf{Z}^*) \leq 2 \exp \left( - \min \left\{ \frac{t^2}{2\nu^2 \|\mathbf{A}\|_{\mathbb{F}}^2}, \frac{t}{2\alpha \|\mathbf{A}\|_{\max}} \right\} \right)$$

From the proof above, we have that  $\|\mathbf{A}\|_{\max} \leq \|\mathbf{A}\|_{\mathbb{F}} = \mathcal{O}_{\mathbb{P}}((nr(n \wedge p))^{-1/2})$ . Therefore, by choosing  $t \asymp (nr(n \wedge p))^{-1/2} \log(np)$ , we further have

$$|T_1| = \mathcal{O} \left( \sqrt{\frac{r}{n(n \wedge p)}} \right).$$

with probability at least  $1 - (np)^{-\delta}$ . The above results imply that

$$\mathcal{L}(\Theta^* - \mathbf{Z}^* \mathbf{\Gamma}^{*\top} \mathcal{P}_{\hat{\mathbf{F}}}^{\perp}) \leq \mathcal{L}(\Theta^*) + \mathcal{O} \left( \frac{r}{n \wedge p} \right), \quad (\text{D.15})$$

with probability at least  $1 - (n+p)^{-\delta} - (np)^{-\delta} - \exp(-n)$ .

Consider the second term of (D.14), from Theorem 3 we also have

$$\begin{aligned} \|\mathbf{B}^{*\top} \mathcal{P}_{\hat{\mathbf{F}}}^{\perp}\|_{1,1} - \|\mathbf{B}^{*\top} \mathcal{P}_{\mathbf{\Gamma}}^{*\perp}\|_{1,1} &\leq \|\mathbf{B}^{*\top} (\mathcal{P}_{\hat{\mathbf{F}}}^{\perp} - \mathcal{P}_{\mathbf{\Gamma}}^{*\perp})\|_{1,1} \\ &\leq \sum_{\ell=1}^d \|(\mathcal{P}_{\hat{\mathbf{F}}}^{\perp} - \mathcal{P}_{\mathbf{\Gamma}}^{*\perp}) \mathbf{B}_{\ell}^*\|_1 \\ &\leq dsC \max_{j \in [d]} \|(\mathcal{P}_{\hat{\mathbf{F}}}^{\perp} - \mathcal{P}_{\mathbf{\Gamma}}^{*\perp}) \mathbf{e}_j\|_1 \\ &= \mathcal{O} \left( \frac{ds}{\sqrt{p(n \wedge p)}} \right), \end{aligned} \quad (\text{D.16})$$

under the same probability event above.

Finally, combining (D.14)-(D.16), we have

$$\begin{aligned} \mathcal{L}(\hat{\boldsymbol{\beta}}, \hat{\boldsymbol{\zeta}}) + \lambda \|\hat{\boldsymbol{\beta}}\|_1 &\leq \mathcal{L}(\Theta^*) + \lambda \|\mathbf{B}^{*\top} \mathcal{P}_{\hat{\mathbf{F}}}^{\perp}\|_{1,1} + \mathcal{O}_{\mathbb{P}} \left( \frac{1}{n \wedge p} \right), \\ &= \mathcal{L}(\tilde{\boldsymbol{\beta}}^*, \tilde{\boldsymbol{\zeta}}^*) + \lambda \|\tilde{\boldsymbol{\beta}}^*\|_1 + \mathcal{O}_{\mathbb{P}} \left( \frac{1}{n \wedge p} \right) \\ &= \mathcal{L}(\boldsymbol{\beta}^*, \boldsymbol{\zeta}^*) + \lambda \|\boldsymbol{\beta}^*\|_1 + \mathcal{O}_{\mathbb{P}} \left( \frac{1}{n \wedge p} + \lambda \sqrt{\frac{(sd)^2}{n \wedge p^{1-k}}} \right), \end{aligned}$$

where the last inequality is from (D.13). This completes the proof.  $\square$

**Lemma D.2** (Infinity norm of the gradient). Under the same conditions in Theorem 5, for any constant  $c > 0$  it holds that

$$(1) \quad \|\nabla_{\beta} \mathcal{L}(\beta^*, \zeta^*)\|_{\infty} \leq 4\nu^2 \sqrt{\frac{c \log^2(2nd)}{n}},$$

$$(2) \quad \|\nabla_{\zeta} \mathcal{L}(\beta^*, \zeta^*)\|_{\infty} \leq \frac{2c\alpha \log(2np)}{n},$$

with probability at least  $1 - (2nd)^{-c} - (2np)^{-c}$ .

*Proof of Lemma D.2.* Recall that

$$\nabla_{\beta} \mathcal{L}(\beta^*, \zeta^*) = \frac{1}{n} \sum_{\ell=1}^{np} [\tilde{y}_{\ell} - A'(\tilde{\mathbf{x}}_{\ell}^{\top} \beta^* + \zeta_{\ell}^*)] \tilde{\mathbf{x}}_{\ell} \quad (\text{D.17})$$

$$\nabla_{\zeta_{\ell}} \mathcal{L}(\beta^*, \zeta^*) = \frac{1}{n} [\tilde{y}_{\ell} - A'(\tilde{\mathbf{x}}_{\ell}^{\top} \beta^* + \zeta_{\ell}^*)], \quad \ell \in [np], \quad (\text{D.18})$$

where  $\tilde{\mathbf{x}}_{\ell}$  is the  $\ell$ -th row of  $\tilde{\mathbf{X}}$ . We split the proof into different parts.

**Part (1).** Conditioned on  $\mathbf{X}$  and  $\mathbf{Z}^*$ , the term  $\tilde{y}_{\ell} - A'(\tilde{\mathbf{x}}_{\ell}^{\top} \beta^* + \zeta_{\ell}^*)$  is a zero-mean  $(\nu, \alpha)$ -sub-exponential random variable, where  $\nu = \sqrt{\kappa_2}$  and  $\alpha = 1/C^2$ , as shown in the proof of Theorem 2. Let  $C' = \max_{\ell \in [np]} \|\tilde{\mathbf{x}}_{\ell}^*\|_{\infty}$ . Because  $\tilde{\mathbf{x}}_{\ell}^*$  are sparse vectors with  $\|\tilde{\mathbf{x}}_{\ell}^*\|_0 = d$ , we have that  $[\tilde{y}_{\ell} - A'(\tilde{\mathbf{x}}_{\ell}^{\top} \beta^* + \zeta_{\ell}^*)] \tilde{x}_{\ell j}$  is a zero-mean  $(\nu, C'\alpha)$ -sub-exponential random variable for  $j = k, k+p, \dots, k+p(d-1)$  and zero otherwise, where  $k = \lfloor p/d \rfloor$ . By Bernstein's inequality, it follows that

$$\mathbb{P} \left( \frac{1}{n} \sum_{\ell=1}^{np} [\tilde{y}_{\ell} - A'(\tilde{\mathbf{x}}_{\ell}^{\top} \beta^* + \zeta_{\ell}^*)] \tilde{x}_{\ell j} \leq t \right) \geq 1 - 2 \exp \left( -\frac{n}{2} \min \left\{ \frac{t^2}{\nu^2}, \frac{t}{C'\alpha} \right\} \right).$$

Applying union bound over  $j = k, k+p, \dots, k+p(d-1)$  yields that

$$\mathbb{P} (\|\nabla_{\beta} \mathcal{L}(\beta^*, \zeta^*)\|_{\infty} \leq t) \geq 1 - 2d \exp \left( -n \min \left\{ \frac{t^2}{2\nu^2}, \frac{t}{2C'\alpha} \right\} \right).$$

By setting  $t = C' \sqrt{2\nu^2 c \log(2nd)/n}$  for some fixed constant  $c > 1$ , we have

$$\mathbb{P} \left( \|\nabla_{\beta} \mathcal{L}(\beta^*, \zeta^*)\|_{\infty} \leq C' \sqrt{\frac{2\nu^2 c \log(2nd)}{n}} \right) \geq 1 - (2nd)^{1-c},$$

when  $n$  is large enough such that  $t < \nu^2/(C'\alpha)$ . By Lemma D.6, we also have  $C' = \|\tilde{\mathbf{X}}\|_{\max} \leq 2\sqrt{2\nu} \sqrt{c \log(nd)}$  with probability at least  $1 - (2nd)^{1-c}$ . It follows that

$$\mathbb{P} \left( \|\nabla_{\beta} \mathcal{L}(\beta^*, \zeta^*)\|_{\infty} \leq 4\nu^2 c \sqrt{\frac{\log^2(2nd)}{n}} \right) \geq 1 - 2(2nd)^{1-c}, \quad (\text{D.19})$$

which finishes the proof for Part (1).



**Part (2).** Because  $\tilde{y}_\ell - A'(\tilde{\mathbf{x}}_\ell^\top \boldsymbol{\beta}^* + \zeta_\ell^*)$ 's are zero-mean  $(\nu, \alpha)$ -sub-exponential random variables, by union bound, we have

$$\mathbb{P}(n \|\nabla_\zeta \mathcal{L}(\boldsymbol{\beta}^*, \boldsymbol{\zeta}^*)\|_\infty \leq t) \geq 1 - 2np \exp\left(-\frac{1}{2} \min\left\{\frac{t^2}{\nu^2}, \frac{t}{\alpha}\right\}\right),$$

or equivalently,

$$\begin{aligned} \mathbb{P}(\|\nabla_\zeta \mathcal{L}(\boldsymbol{\beta}^*, \boldsymbol{\zeta}^*)\|_\infty \leq t) &\geq 1 - 2np \exp\left(-\frac{1}{2} \min\left\{\frac{(nt)^2}{\nu^2}, \frac{nt}{\alpha}\right\}\right) \\ &= 1 - 2np \exp\left(-\frac{nt}{2\alpha}\right) \end{aligned}$$

when  $t \geq \nu^2/(\alpha n)$  is sufficiently large. Choosing  $t = \max\{2c\alpha \log(2np), \nu^2/\alpha\}/n$  for any constant  $c > 1$  yields that

$$\mathbb{P}\left(\|\nabla_\zeta \mathcal{L}(\boldsymbol{\beta}^*, \boldsymbol{\zeta}^*)\|_\infty \leq \frac{2c\alpha \log(2np)}{n}\right) \leq 1 - (2np)^{1-c}.$$

Finally, we obtain the tail probability as the lemma states by taking the union bound over the above events.  $\square$

**Lemma D.3** (Locally restricted strongly convexity). Under the same conditions in Theorem 5, define the augmented cone

$$\begin{aligned} \mathcal{C}_a(\xi, \mathcal{S}, \eta_m) &:= \{(\boldsymbol{\Delta}_\beta, \boldsymbol{\Delta}_\zeta) \in \mathbb{R}^{pd} \times \mathbb{R}^{np} \mid \boldsymbol{\Delta}_\beta \in \mathcal{C}(\xi, \mathcal{S}), \frac{1}{n} \|\mathbf{Z} - \mathbf{Z}^* \mathbf{R}^\top\|_F^2 \leq \eta_m, \\ &\text{for } \mathbf{Z} \text{ such that } \boldsymbol{\zeta} = \text{vec}(\mathbf{Z} \boldsymbol{\Gamma}^\top)\}, \end{aligned} \quad (\text{D.20})$$

where the cone  $\mathcal{C}(\xi, \mathcal{S})$  defined in (D.6) and  $\eta_m = o(1)$ . Then, it holds that

$$\inf_{(\boldsymbol{\Delta}_\beta, \boldsymbol{\Delta}_\zeta) \in \mathcal{C}_a(\xi, \mathcal{S}, \eta_m)} \mathcal{E}(\boldsymbol{\beta}^* + \boldsymbol{\Delta}_\beta, \boldsymbol{\beta}^*; \boldsymbol{\zeta}^* + \boldsymbol{\Delta}_\zeta, \boldsymbol{\zeta}^*) \geq \frac{\kappa_1 C'}{2} \|\boldsymbol{\Delta}_\beta\|_2^2 - C'' \sqrt{n^{-1} \vee \eta_m} \|\boldsymbol{\Delta}_\beta\|_1,$$

for some constant  $C' > 0$ , with probability at least  $1 - 2 \exp(-\xi \log n)$ .

*Proof of Lemma D.3.* For all  $(\boldsymbol{\Delta}_\beta, \boldsymbol{\Delta}_\zeta) \in \mathcal{C}_a(\xi, \mathcal{S}, \eta_m)$ , let  $\boldsymbol{\beta} = \boldsymbol{\beta}^* + \boldsymbol{\Delta}_\beta$  and  $\boldsymbol{\zeta} = \boldsymbol{\zeta}^* + \boldsymbol{\Delta}_\zeta$ . Let  $\boldsymbol{\Delta}_{\beta_j} \in \mathbb{R}^d$  be the sub-vector containing the  $(j-1)d$ -th to  $(jd-1)$ -th entries of  $\boldsymbol{\Delta} \in \mathbb{R}^{pd}$ . It is also equivalent to  $\mathbf{b}_j - \mathbf{b}_j^*$ , the  $j$ -th row of  $\text{vec}^{-1}(\boldsymbol{\beta} - \boldsymbol{\beta}^*) \in \mathbb{R}^{p \times d}$ . Let  $\mathbf{E} = \text{vec}^{-1}(\boldsymbol{\zeta}) \in \mathbb{R}^{n \times p}$  and  $\mathbf{E}^* = \text{vec}^{-1}(\boldsymbol{\zeta}^*) \in \mathbb{R}^{n \times p}$ .

We begin by decomposing the error into different terms:

$$\mathcal{E}(\boldsymbol{\beta}, \boldsymbol{\beta}^*; \boldsymbol{\zeta}, \boldsymbol{\zeta}^*) = \sum_{j=1}^p \mathcal{E}_j(\boldsymbol{\beta}, \boldsymbol{\beta}^*; \boldsymbol{\zeta}, \boldsymbol{\zeta}^*),$$

where

$$\begin{aligned}
\mathcal{E}_j(\boldsymbol{\beta}, \boldsymbol{\beta}^*; \boldsymbol{\zeta}, \boldsymbol{\zeta}^*) &= \frac{1}{n} \sum_{i=1}^n [A'(\mathbf{x}_i^\top \mathbf{b}_j + \mathbf{z}_i^\top \boldsymbol{\gamma}_j) - A'(\mathbf{x}_i^\top \mathbf{b}_j^* + \mathbf{z}_i^{*\top} \boldsymbol{\gamma}_j^*)] \mathbf{x}_i^\top \boldsymbol{\Delta}_{\beta_j} \\
&= \frac{1}{n} \sum_{i=1}^n [A'(\mathbf{x}_i^\top \mathbf{b}_j + \mathbf{z}_i^\top \boldsymbol{\gamma}_j) - A'(\mathbf{x}_i^\top \mathbf{b}_j^* + \mathbf{z}_i^\top \boldsymbol{\gamma}_j)] \mathbf{x}_i^\top \boldsymbol{\Delta}_{\beta_j} \\
&\quad + \frac{1}{n} \sum_{i=1}^n [A'(\mathbf{x}_i^\top \mathbf{b}_j^* + \mathbf{z}_i^\top \boldsymbol{\gamma}_j) - A'(\mathbf{x}_i^\top \mathbf{b}_j^* + \mathbf{z}_i^{*\top} \boldsymbol{\gamma}_j^*)] \mathbf{x}_i^\top \boldsymbol{\Delta}_{\beta_j} \\
&= \frac{1}{n} \sum_{i=1}^n A''(\theta_{ij})(\mathbf{x}_i^\top \boldsymbol{\Delta}_{\beta_j})^2 + \frac{1}{n} \sum_{i=1}^n A''(\theta'_{ij})(\mathbf{z}_i^\top \boldsymbol{\gamma}_j - \mathbf{z}_i^{*\top} \boldsymbol{\gamma}_j^*) \mathbf{x}_i^\top \boldsymbol{\Delta}_{\beta_j} \\
&= T_{1j} + T_{2j}, \tag{D.21}
\end{aligned}$$

where  $\theta_{ij}$  is between  $\mathbf{x}_i^\top \mathbf{b}_j + \mathbf{z}_i^\top \boldsymbol{\gamma}_j$  and  $\mathbf{x}_i^\top \mathbf{b}_j^* + \mathbf{z}_i^\top \boldsymbol{\gamma}_j$ , and  $\theta'_{ij}$  is between  $\mathbf{x}_i^\top \mathbf{b}_j^* + \mathbf{z}_i^\top \boldsymbol{\gamma}_j$  and  $\mathbf{x}_i^\top \mathbf{b}_j^* + \mathbf{z}_i^{*\top} \boldsymbol{\gamma}_j^*$ .

For the first term, because by Assumption 1,  $\kappa_1 := \inf_{\theta \in \mathcal{R}} A''(\theta) > 0$ , we have

$$\begin{aligned}
T_{1j} &\geq \kappa_1 \frac{1}{n} \sum_{i=1}^n (\mathbf{x}_i^\top \boldsymbol{\Delta}_{\beta_j})^2 \\
&\geq \kappa_1 \lambda_p \left( \frac{1}{n} \mathbf{X}^\top \mathbf{X} \right) \|\boldsymbol{\Delta}_{\beta_j}\|_2^2 \\
&\geq \frac{\kappa_1}{C} \lambda_p \left( \frac{1}{n} \boldsymbol{\Sigma}_x^{-\frac{1}{2}} \mathbf{X}^\top \mathbf{X} \boldsymbol{\Sigma}_x^{-\frac{1}{2}} \right) \|\boldsymbol{\Delta}_{\beta_j}\|_2^2,
\end{aligned}$$

where the last inequality is from Assumption 2. From Vershynin (2018, Lemma 10.6.6), we further have that when  $n \gtrsim \xi \log d$

$$T_{1j} \geq \frac{\kappa_1 C'}{2} \|\boldsymbol{\Delta}_{\beta_j}\|_2^2 \tag{D.22}$$

over  $\mathcal{C}(\xi, \mathcal{S})$  (which contains  $\mathcal{C}_a(\xi, \mathcal{S}, C_0)$ ), with probability at least  $1 - 2 \exp(-\xi \log n)$  for some absolute constant  $C' > 0$ .

For the second term, by Holder's inequality, we have

$$\begin{aligned}
|T_{2j}| &= \left| \frac{1}{n} (\mathbf{E}_j - \mathbf{E}_j^*)^\top \text{diag}(A''(\boldsymbol{\Theta}'_j)) \mathbf{X} \boldsymbol{\Delta}_{\beta_j} \right| \\
&\leq \frac{1}{n} \|\mathbf{X}^\top \text{diag}(A''(\boldsymbol{\Theta}'_j)) (\mathbf{E}_j - \mathbf{E}_j^*)\|_\infty \cdot \|\boldsymbol{\Delta}_{\beta_j}\|_1 \\
&= \max_{1 \leq \ell \leq d} \frac{1}{n} |(\mathbf{E}_j - \mathbf{E}_j^*)^\top \text{diag}(A''(\boldsymbol{\Theta}'_j)) \mathbf{X}_\ell| \cdot \|\boldsymbol{\Delta}_{\beta_j}\|_1,
\end{aligned} \tag{D.23}$$

where  $\boldsymbol{\Theta}'_j = (\theta'_{1j}, \dots, \theta'_{nj})^\top$ . Recall the invertible matrix  $\mathbf{R}$  with  $\|\mathbf{R}\|_{\text{op}} = \mathcal{O}_{\mathbb{P}}(1)$  from Assumption 4 and define the transformed parameters  $\tilde{\mathbf{Z}}^* = \mathbf{Z}^* \mathbf{R}^\top$  and  $\tilde{\boldsymbol{\Gamma}}^* = \boldsymbol{\Gamma}^* \mathbf{R}^{-1}$ . Because

$\mathbf{E}_j - \mathbf{E}_j^* = \tilde{\mathbf{Z}}^*(\boldsymbol{\gamma}_j - \tilde{\boldsymbol{\gamma}}_j^*) + (\mathbf{Z} - \tilde{\mathbf{Z}}^*)\boldsymbol{\gamma}_j$ , it follows that

$$\begin{aligned}
|T_{2j}| &\leq \max_{1 \leq \ell \leq d} \frac{1}{n} |(\boldsymbol{\gamma}_j - \tilde{\boldsymbol{\gamma}}_j^*)^\top \tilde{\mathbf{Z}}^{*\top} \text{diag}(A''(\boldsymbol{\Theta}'_j)) \mathbf{X}_\ell| \cdot \|\Delta_{\beta_j}\|_1 \\
&\quad + \max_{1 \leq \ell \leq d} \frac{1}{n} |\tilde{\boldsymbol{\gamma}}_j^\top (\mathbf{Z} - \tilde{\mathbf{Z}}^*)^\top \text{diag}(A''(\boldsymbol{\Theta}'_j)) \mathbf{X}_\ell| \cdot \|\Delta_{\beta_j}\|_1 \\
&\leq \max_{1 \leq \ell \leq d, 1 \leq k \leq r} \frac{1}{\sqrt{n}} \|\boldsymbol{\gamma}_j - \tilde{\boldsymbol{\gamma}}_j^*\|_2 \cdot \frac{1}{\sqrt{n}} |\tilde{\mathbf{Z}}_k^{*\top} \text{diag}(A''(\boldsymbol{\Theta}'_j)) \mathbf{X}_\ell| \cdot \|\Delta_{\beta_j}\|_1 \\
&\quad + \kappa_2 \max_{1 \leq \ell \leq d} \|\boldsymbol{\gamma}_j\|_2 \frac{1}{\sqrt{n}} \|\mathbf{Z} - \tilde{\mathbf{Z}}^*\|_{\text{op}} \frac{1}{\sqrt{n}} \|\mathbf{X}\|_{\text{op}} \cdot \|\Delta_{\beta_j}\|_1 \\
&\lesssim \sqrt{n^{-1} \vee \eta_n} \|\Delta_{\beta_j}\|_1, \tag{D.24}
\end{aligned}$$

where the last inequality is because that  $\|\boldsymbol{\gamma}_j - \tilde{\boldsymbol{\gamma}}_j^*\|_2 \leq 2C$  and  $\tilde{\mathbf{Z}}_k^{*\top} \text{diag}(A''(\boldsymbol{\Theta}'_j)) \mathbf{X}_\ell$  scales in  $\sqrt{n}$  as the sum of  $n$  independent sub-exponential random variables, and  $\|\mathbf{X}\|_{\text{op}} \lesssim \sqrt{n}$ .

By combining (D.21), (D.22) and (D.24), we have that

$$\mathcal{E}_j(\boldsymbol{\beta}, \boldsymbol{\beta}^*; \boldsymbol{\zeta}, \boldsymbol{\zeta}^*) \geq \frac{\kappa_1 C'}{2} \|\Delta_{\beta_j}\|_2^2 - C'' \sqrt{n^{-1} \vee \eta_n} \|\Delta_{\beta_j}\|_1,$$

for some constant  $C'' > 0$ . Thus,

$$\mathcal{E}(\boldsymbol{\beta}, \boldsymbol{\beta}^*; \boldsymbol{\zeta}, \boldsymbol{\zeta}^*) \geq \frac{\kappa_1 C'}{2} \|\Delta_{\beta}\|_2^2 - C'' \sqrt{n^{-1} \vee \eta_n} \|\Delta_{\beta}\|_1,$$

over the cone  $\mathcal{C}_a(\xi, \mathcal{S}, \eta_n)$ . □

**Remark 5** (Neyman orthogonality). By coincidence, the proof above on  $T_{2j}$  actually verifies the uniform Neyman orthogonality of the empirical loss in semiparametric models (Chernozhukov et al., 2018; Foster and Syrgkanis, 2023). This relies on the estimation error rates for the nuisance parameters  $\mathbf{E}_j^*$  by using the consistent estimation for the latent factors  $\mathbf{Z}^*$  with rate  $\eta_n$  as assumed in the cone condition (D.21). To see this, recall that the pathwise derivative map of the gradient  $\nabla_{\mathbf{b}_j} \mathcal{L}(\boldsymbol{\beta}^*, \boldsymbol{\zeta}^*)$  evaluated at the true parameter  $\boldsymbol{\beta}^*$  and nuisance component value  $\boldsymbol{\zeta}^*$  (when  $t = 0$ ) is given by

$$\begin{aligned}
\left. \frac{\partial}{\partial t} \nabla_{\mathbf{b}_j} \mathcal{L}(\boldsymbol{\beta}^*, t(\boldsymbol{\zeta} - \boldsymbol{\zeta}^*) + \boldsymbol{\zeta}^*) \right|_{t=0} &= \frac{1}{n} \sum_{i \in [n]} A''(\mathbf{x}_i^\top \mathbf{b}_j^* + e_{ij}^*) (e_{ij} - e_{ij}^*) \mathbf{x}_i \\
&= \frac{1}{n} \mathbf{X}^\top \text{diag}(A''(\boldsymbol{\Theta}_j^*)) (\mathbf{E}_j - \mathbf{E}_j^*).
\end{aligned}$$

Compared to (D.23), up to a constant factor, (D.24) also suggest that the pathwise derivative's infinity norm vanishes with a rate of  $\sqrt{n^{-1} \vee \eta_n}$ . In other words, at the true parameter value, local perturbations of the nuisance component around its true value have a negligible effect on the gradient of the loss with respect to the primary parameter, with high probability; see (Chernozhukov et al., 2018; Foster and Syrgkanis, 2023) for more detailed discussions about the Neyman orthogonality.

**Lemma D.4** (Bounds related to projection). Under assumptions in Theorem 5, it holds that

$$\|\mathcal{P}_{\hat{\Gamma}}^\perp \Gamma^*\|_{\text{F}}^2 = \mathcal{O}\left(\frac{r}{n \wedge p}\right),$$

with probability at least  $1 - (n + p)^{-\delta} - (np)^{-\delta} - \exp(-n)$ .

*Proof of Lemma D.4.* From the proof of Theorem 3, the result follows by noting that

$$\begin{aligned} \|\mathcal{P}_{\hat{\Gamma}}^\perp \Gamma^*\|_{\text{F}}^2 &= \|(\mathcal{P}_{\hat{\Gamma}}^\perp - \mathcal{P}_{\Gamma^*}^\perp) \Gamma^*\|_{\text{F}}^2 \\ &= \sum_{\ell=1}^r \left\| \sum_{j=1}^p (\mathcal{P}_{\hat{\Gamma}}^\perp - \mathcal{P}_{\Gamma^*}^\perp) \mathbf{e}_j \cdot \gamma_{j,\ell}^* \right\|_2^2 \\ &\leq rpC^2 \max_{1 \leq \ell \leq p} \|(\mathcal{P}_{\hat{\Gamma}}^\perp - \mathcal{P}_{\Gamma^*}^\perp) \mathbf{e}_j\|_2^2 \\ &= \mathcal{O}(r(n \wedge p)^{-1}), \end{aligned}$$

with probability at least  $1 - (n + p)^{-\delta} - (np)^{-\delta} - \exp(-n)$ .  $\square$

**Lemma D.5** (Sub-Gaussianity of  $\mathbf{Z}$ ). Under Assumptions 1–3,  $\mathbf{z}_1, \dots, \mathbf{z}_n$  are independent and identically distributed sub-Gaussian random vectors.

*Proof of Lemma D.5.* For the proof below, we ignore the asterisks for the true parameters for simplicity. Recall that  $\mathbf{z}_i = \mathbf{D}\mathbf{x}_i + \mathbf{w}_i$  is the linear function of two independent sub-Gaussian random vectors. The mean is given by  $\mathbb{E}[\mathbf{z}_i] = \mathbf{D}\mathbb{E}[\mathbf{x}_i]$ . It suffices to bound the operator norm of  $\mathbf{D}$ . From  $\Theta = \mathbf{X}\mathbf{B}^\top + \mathbf{Z}\Gamma^\top$ , we have  $\mathbf{X}^\top \Theta = \mathbf{X}^\top \mathbf{X}\mathbf{B}^\top + \mathbf{X}^\top \mathbf{Z}\Gamma^\top$ . Taking expectation over  $\mathbf{X}$  and  $\mathbf{Z}$  yield that  $\mathbb{E}[\mathbf{X}^\top \Theta/n] = \Sigma_x(\mathbf{B}^\top + \mathbf{D}^\top \Gamma^\top)$ . Rearranging the formula yields that

$$\Gamma \mathbf{D} = \frac{1}{n} \mathbb{E}[\Theta^\top \mathbf{X}] \Sigma_x^{-1} - \mathbf{B}$$

and

$$\mathbf{D} = (\Gamma^\top \Gamma)^{-1} \Gamma^\top \left( \frac{1}{n} \mathbb{E}[\Theta^\top \mathbf{X}] \Sigma_x^{-1} - \mathbf{B} \right).$$

By the sub-multiplicative property of the operator norm, we have

$$\begin{aligned} \|\mathbf{D}\|_{\text{op}} &\leq \|(\Gamma^\top \Gamma)^{-1} \Gamma^\top\|_{\text{op}} \left( \frac{1}{n} \|\mathbb{E}[\Theta^\top \mathbf{X}]\|_{\text{op}} \|\Sigma_x^{-1}\|_{\text{op}} + \|\mathbf{B}\|_{\text{op}} \right) \\ &\lesssim \frac{1}{\sqrt{p}} \left( \frac{1}{n} \mathbb{E}[\|\Theta^\top\|_{\text{op}} \|\mathbf{X}\|_{\text{op}}] + \sqrt{p} \right) \\ &\lesssim \frac{1}{\sqrt{p}} \left( \frac{\sqrt{np} \sqrt{n}}{n} + \sqrt{p} \right) \\ &\lesssim 1, \end{aligned}$$

where in the second inequality, we use Jensen's inequality and the norm inequality  $\|\mathbf{B}\|_{\text{op}} \leq \sqrt{p}\|\mathbf{B}\|_{\text{max}}$ . This finishes the proof.  $\square$

**Lemma D.6** (Infinity norm of the covariates). Under Assumption 2, it holds that  $\|\widetilde{\mathbf{X}}\|_{\text{max}} \leq 2\sqrt{2\nu}\sqrt{c\log(2nd)}$ , with probability at least  $1 - (2nd)^{-c}$  for any fixed constant  $c > 0$ .

*Proof of Lemma D.6.* Let  $\text{vec}^{-1}(\mathbf{e}_\ell) = \mathbf{e}_{n,i}\mathbf{e}_{p,j}^\top$  where  $\mathbf{e}_{n,i}$  is the unit vector of dimension  $n$  and  $\mathbf{e}_{p,j}$  is defined analogously. Note that

$$\begin{aligned}\widetilde{\mathbf{x}}_\ell &= (\mathbf{I}_p \otimes \mathbf{X})^\top \mathbf{e}_\ell \\ &= \text{vec}(\mathbf{X}^\top \text{vec}^{-1}(\mathbf{e}_\ell) \mathbf{I}_p) \\ &= \text{vec}(\mathbf{X}^\top \mathbf{e}_{n,i} \mathbf{e}_{p,j}^\top) \\ &= \text{vec}(\mathbf{x}_i \mathbf{e}_{p,j}^\top).\end{aligned}$$

Because  $\mathbf{x}_i$ 's are  $\nu$ -sub-Gaussian random vectors, we have

$$\begin{aligned}\|\widetilde{\mathbf{X}}\|_{\text{max}} &= \max_{\ell \in [np]} \|\widetilde{\mathbf{x}}_\ell\|_\infty \\ &= \max_{i \in [n], j \in [p]} \|\mathbf{x}_i \mathbf{e}_{p,j}^\top\|_\infty \\ &\leq \max_{i \in [n]} \|\mathbf{x}_i\|_\infty \\ &\leq \max_{i \in [n], j \in [p]} |x_{ij}| \\ &\leq 2\sqrt{2\nu}\sqrt{c\log(2nd)}\end{aligned}$$

with probability at least  $1 - (2nd)^{-c}$  for any fixed constant  $c > 0$ .  $\square$

## E Asymptotic normality of the debiased estimator

### E.1 Proof of Theorem 6

*Proof of Theorem 6.* Condition on  $\mathcal{D}_2$ , Theorem 3 and Theorem 5 imply that the following event holds:

$$\mathcal{E}_1 = \left\{ \begin{aligned} \max_{1 \leq j \leq p} \|(\mathcal{P}_{\widehat{\Gamma}} - \mathcal{P}_{\Gamma^*})\mathbf{e}_j\|_2 &\lesssim (p(n \wedge p))^{-1/2}, \frac{1}{n} \|\widehat{\mathbf{Z}}\widehat{\boldsymbol{\gamma}}_j - \mathbf{Z}^*\boldsymbol{\gamma}_j^*\|_2^2 \lesssim r_{n,p}, \\ \|\widehat{\mathbf{B}} - \mathbf{B}^*\|_{1,1} &\lesssim \sqrt{sd}r_{n,p}, \|\widehat{\mathbf{B}} - \mathbf{B}^*\|_{\text{F}} \lesssim r_{n,p} \end{aligned} \right\}.$$

where  $r_{n,p}$  is defined in Theorem 5.

Recall that

$$\widehat{b}_{j1}^{\text{de}} = \widehat{b}_{j1} + \widehat{\mathbf{u}}^\top \frac{1}{n} \sum_{i=1}^n \widehat{\omega}_i \mathbf{x}_i (\mathbf{y}_i - A'(\widehat{\boldsymbol{\theta}}_i))^\top \mathbf{v}_i \quad (\text{E.1})$$

where  $\mathbf{v}_i = \text{diag}(A''(\widehat{\boldsymbol{\theta}}_i))^{-1} \mathcal{P}_{\widehat{\Gamma}}^\perp \mathbf{e}_j$ . By Taylor expansion of  $A'(\theta_{ij}^*)$  at  $\widehat{\theta}_{ij} := \mathbf{x}_i^\top \widehat{\mathbf{b}}_j + \widehat{\mathbf{z}}_i^\top \widehat{\boldsymbol{\gamma}}_j$ , we have

$$A'(\theta_{ij}^*) = A'(\widehat{\theta}_{ij}) + A''(\widehat{\theta}_{ij})(\theta_{ij}^* - \widehat{\theta}_{ij}) + \frac{1}{2}A'''(\psi_{ij})(\theta_{ij}^* - \widehat{\theta}_{ij})^2,$$

for some  $\psi_{ij}$  between  $\widehat{\theta}_{ij}$  and  $\theta_{ij}^*$ . Then, the residual of the  $i$ th sample can be decomposed into three sources of errors:

$$\mathbf{y}_i - A'(\widehat{\boldsymbol{\theta}}_i) = \underbrace{\boldsymbol{\epsilon}_i}_{\text{stochastic error}} + \underbrace{\mathbf{p}_i}_{\text{remaining bias}} + \underbrace{\mathbf{q}_i}_{\text{approximation error}} \quad (\text{E.2})$$

where the three terms of errors read that

$$\begin{aligned} \boldsymbol{\epsilon}_i &= \mathbf{y}_i - A'(\boldsymbol{\theta}_i^*) \\ \mathbf{p}_i &= A''(\widehat{\boldsymbol{\theta}}_i) \odot (\widetilde{\boldsymbol{\theta}}_i - \boldsymbol{\theta}_i^*) \\ \mathbf{q}_i &= -\frac{1}{2}[A'''(\widetilde{\theta}_{ij} + t(\theta_{ij}^* - \widetilde{\theta}_{ij}))(\theta_{ij}^* - \widetilde{\theta}_{ij})^2]_{j \in [p]}. \end{aligned}$$

Substituting (E.2) into (E.1) yields that

$$\begin{aligned} \widehat{b}_{j1}^{\text{de}} - b_{j1}^* &= (\widehat{b}_{j1} - b_{j1}^*) + \mathbf{u}^\top \frac{1}{n} \sum_{i=1}^n \omega_i \mathbf{x}_i \boldsymbol{\epsilon}_i^\top \mathbf{v}_i + \mathbf{u}^\top \frac{1}{n} \sum_{i=1}^n \omega_i \mathbf{x}_i \mathbf{p}_i^\top \mathbf{v}_i + \mathbf{u}^\top \frac{1}{n} \sum_{i=1}^n \omega_i \mathbf{x}_i \mathbf{q}_i^\top \mathbf{v}_i \\ &= \mathbf{u}^\top \frac{1}{n} \sum_{i=1}^n \omega_i \mathbf{x}_i \boldsymbol{\epsilon}_i^\top \mathbf{v}_i + \left( \mathbf{u}^\top \frac{1}{n} \sum_{i=1}^n \omega_i \mathbf{x}_i \mathbf{x}_i^\top - \mathbf{e}_1^\top \right) (\mathbf{b}_j^* - \widehat{\mathbf{b}}_j) \\ &\quad + \mathbf{u}^\top \frac{1}{n} \sum_{i=1}^n \mathbf{x}_i \mathbf{x}_i^\top \mathbf{B}^{*\top} ((\mathcal{P}_{\widehat{\Gamma}}^\perp - \mathcal{P}_{\Gamma^*}^\perp) \mathbf{e}_j - \mathcal{P}_{\Gamma^*} \mathbf{e}_j) \\ &\quad + \mathbf{u}^\top \frac{1}{n} \sum_{i=1}^n \mathbf{x}_i \mathbf{q}_i^\top \mathbf{v}_i, \end{aligned} \quad (\text{E.3})$$

In the second equality above, we use the properties that  $\widehat{\mathbf{B}}$  and  $\mathbf{B}^*$  are in the column spaces of the orthogonal projections  $\mathcal{P}_{\widehat{\Gamma}}^\perp$  and  $\mathcal{P}_{\Gamma^*}^\perp$ , respectively. Denote the four terms in the right-hand side of (E.3) by  $T_{1j}, T_{2j}, T_{3j}, T_{4j}$ , respectively. We will analyze each of them separately, conditioning on  $\mathcal{D}_2$  and  $\mathbf{X}$ . Then the randomness is from  $\boldsymbol{\epsilon}_i$ 's and  $\mathbf{u}$ . We will show that the  $T_1$  inherits  $\sqrt{n}$ -convergence rate and is asymptotically normally distributed, while the others have faster convergence rates.

Part (1)  $T_{1j}$ . From Lemma E.2, it follows that

$$\sqrt{n} \frac{T_{1j}}{\sigma_j} \xrightarrow{d} \mathcal{N}(0, 1).$$

Part (2)  $T_{2j}$ . By Holder's inequality and the constraint of optimization problem (4.4), it

follows that

$$|T_{2j}| \leq \lambda_n \|\mathbf{b}_j^* - \widehat{\mathbf{b}}_j\|_1 \lesssim \sqrt{\frac{\log(nd)}{n}} r'_{n,p},$$

with probability tending to one. Thus, we have  $\sqrt{n}|T_{2j}| \xrightarrow{P} 0$  as  $n, p \rightarrow \infty$ .

Part (3)  $T_{3j}$ . From Theorem 3 and (A.1) in the proof of Proposition 1, we have that

$$\begin{aligned} |T_{3j}| &\leq \|\mathbf{u}\|_2 \cdot \left\| \frac{1}{n} \sum_{i=1}^n \mathbf{x}_i \mathbf{x}_i^\top \right\|_{\text{op}} \left( \|\mathbf{B}^*\|_{\text{op}} \|(\mathcal{P}_{\widehat{\Gamma}}^\perp - \mathcal{P}_{\Gamma^*}^\perp) \mathbf{e}_j\|_2 + \|\mathbf{B}^{*\top} \mathcal{P}_{\Gamma^*} \mathbf{e}_j\|_2 \right) \\ &\lesssim \sqrt{\frac{d}{p(n \wedge p)}} + \frac{sd}{p}, \end{aligned}$$

with probability tending to one. Thus, if  $\sqrt{n}/p \rightarrow 0$ , we have  $\sqrt{n}|T_{3j}| \xrightarrow{P} 0$  as  $n, p \rightarrow \infty$ .

Part (4)  $T_{4j}$ . The higher-order term is bounded as below:

$$\begin{aligned} |T_{4j}| &\leq \frac{\kappa_2}{2\kappa_1} \max_{1 \leq i \leq n} |\mathbf{u}^\top \mathbf{x}_i| \cdot \frac{1}{n} \sum_{i=1}^n \sum_{j=1}^n |\mathbf{x}_i^\top (\widehat{\mathbf{b}}_j - \mathbf{b}_j^*)|^2 \cdot \|\mathcal{P}_{\widehat{\Gamma}}^\perp \mathbf{e}_j\|_2 \\ &\lesssim \max_{1 \leq i \leq n} |\mathbf{u}^\top \mathbf{x}_i|^3 \|\widehat{\mathbf{B}} - \mathbf{B}^*\|_{\text{F}}^2 \\ &\lesssim \tau_n^3 r_{n,p}^2, \end{aligned}$$

with probability tending to one. Thus, if  $n/(\log(nd)p^{3/2}) \rightarrow 0$  and  $\sqrt{n}/p^{1-k} \rightarrow 0$ , we have  $\sqrt{n}|T_{4j}| \xrightarrow{P} 0$  as  $n, p \rightarrow \infty$ .

We are now combining the above four terms. Because when  $n/\log(nd) = o(p^{3/2})$  and  $n = o(p^{2(1-k)})$ ,  $T_{2j}, \dots, T_{4j} = o_{\mathbb{P}}(1/\sqrt{n})$ , we have  $\sqrt{n}(\widehat{b}_{j1}^{\text{de}} - b_{j1}^*)/\sigma_j \xrightarrow{d} \mathcal{N}(0, 1)$ .  $\square$

## E.2 Proof of Proposition 7

*Proof of Proposition 7.* By the definition of  $t_j$  and (E.3) we have the decomposition

$$t_j = \vartheta_j + \varsigma_j,$$

where

$$\begin{aligned} \vartheta_j &= \sqrt{n} \frac{\widehat{\mathbf{u}}^\top \frac{1}{n} \sum_{i=1}^n \widehat{\omega}_i \mathbf{x}_i \epsilon_{ij} A''(\widehat{\theta}_{ij})^{-1}}{\widehat{\sigma}_j}, \\ \varsigma_j &= \sqrt{n} \frac{\widehat{\mathbf{u}}^\top \frac{1}{n} \sum_{i=1}^n \widehat{\omega}_i \mathbf{x}_i \epsilon_i^\top \text{diag}(A''(\widehat{\theta}_i))^{-1} (\mathcal{P}_{\widehat{\Gamma}}^\perp - \mathbf{I}_p) \mathbf{e}_j}{\widehat{\sigma}_j} + \sqrt{n} \frac{T_{2j} + T_{3j} + T_{4j}}{\widehat{\sigma}_j}. \end{aligned}$$

For the first component, note that  $\vartheta_j$  for  $j = 1, \dots, p$  are independent conditional on  $\{(\mathbf{x}_i, \mathbf{z}_i)\}_{i=1}^n$  and  $\mathcal{D}_2$ . Furthermore,  $\vartheta_j \xrightarrow{d} \mathcal{N}(0, 1)$  for  $j \in \mathcal{N}_p$  from Lemma E.2 by noting that  $\widehat{\sigma}_j$  is also consistent to the conditional variance of  $\vartheta_j$ . For the second component,

from the proof of Theorem 6 and Lemma E.3, we know that

$$\max_{1 \leq j \leq p} \sqrt{n} \frac{T_{2j} + T_{3j} + T_{4j}}{\hat{\sigma}_j} = o_{\mathbb{P}}(1).$$

On the other hand, from Theorem 3 and Assumption 3, we also have

$$\begin{aligned} & \max_{1 \leq j \leq p} \sqrt{n} \frac{\hat{\mathbf{u}}^\top \frac{1}{n} \sum_{i=1}^n \hat{\omega}_i \mathbf{x}_i \boldsymbol{\epsilon}_i^\top \text{diag}(A''(\hat{\boldsymbol{\theta}}_i))^{-1} (\mathcal{P}_{\hat{\Gamma}}^\perp - \mathbf{I}_p) \mathbf{e}_j}{\hat{\sigma}_j} \\ &= \max_{1 \leq j \leq p} \sqrt{n} \frac{\hat{\mathbf{u}}^\top \frac{1}{n} \sum_{i=1}^n \hat{\omega}_i \mathbf{x}_i \boldsymbol{\epsilon}_i^\top \text{diag}(A''(\hat{\boldsymbol{\theta}}_i))^{-1} (\mathcal{P}_{\hat{\Gamma}}^\perp - \mathcal{P}_{\Gamma^*}^\perp) \mathbf{e}_j}{\hat{\sigma}_j} \\ & \quad + \max_{1 \leq j \leq p} \sqrt{n} \frac{\hat{\mathbf{u}}^\top \frac{1}{n} \sum_{i=1}^n \hat{\omega}_i \mathbf{x}_i \boldsymbol{\epsilon}_i^\top \text{diag}(A''(\hat{\boldsymbol{\theta}}_i))^{-1} \mathcal{P}_{\Gamma^*} \mathbf{e}_j}{\hat{\sigma}_j} \\ &= \mathcal{O}_{\mathbb{P}} \left( \sqrt{p} \max_{1 \leq j \leq p} \|(\mathcal{P}_{\hat{\Gamma}}^\perp - \mathcal{P}_{\Gamma^*}^\perp) \mathbf{e}_j\|_2 \right) \\ & \quad + \max_{1 \leq j \leq p} \left\| \sqrt{n} \frac{\hat{\mathbf{u}}^\top \frac{1}{n} \sum_{i=1}^n \hat{\omega}_i \mathbf{x}_i \boldsymbol{\epsilon}_i^\top \text{diag}(A''(\hat{\boldsymbol{\theta}}_i))^{-1} \Gamma^*}{\hat{\sigma}_j} \right\|_2 \left\| (\Gamma^{*\top} \Gamma^*)^{-1} \right\|_{\text{op}} \|\boldsymbol{\gamma}_j^*\|_2 \\ &= \mathcal{O}_{\mathbb{P}}(p^{-\frac{1}{2}}) + \mathcal{O}_{\mathbb{P}}(\sqrt{rp} \cdot p^{-1} \cdot 1) \\ &= o_{\mathbb{P}}(1). \end{aligned}$$

where we use the subexponential concentration of  $\epsilon_{ij}$  conditional on  $\{(\mathbf{x}_i, \mathbf{z}_i)\}_{i=1}^n$  and  $\mathcal{D}_2$ . Therefore, we have that  $\max_{1 \leq j \leq p} |\varsigma_j| = o_{\mathbb{P}}(1)$ .

The rest of the proof follows similarly to the proof of Wang et al. (2017a, Theorem 3.4). We present here for completeness.

**Overall Type-I error control.** Let  $\varrho = |\mathcal{N}_p|^{-1} \sum_{j \in \mathcal{N}_p} \mathbf{1}(|t_j| > z_{\frac{\alpha}{2}})$ . To prove the overall Type-I error control, we will show the expectation of  $\varrho$  tends to  $\alpha$  and its variance tends to zero. For the expectation, for any  $\epsilon > 0$ , we have

$$\begin{aligned} \mathbb{E}[\varrho] &= \frac{1}{|\mathcal{N}_p|} \sum_{j \in \mathcal{N}_p} \mathbb{P}(|t_j| > z_{\frac{\alpha}{2}}) \\ &\leq \frac{1}{|\mathcal{N}_p|} \sum_{j \in \mathcal{N}_p} [\mathbb{P}(|\vartheta_j| > z_{\frac{\alpha}{2}} - \epsilon) + \mathbb{P}(|\varsigma_j| > \epsilon)] \\ &= \frac{1}{|\mathcal{N}_p|} \sum_{j \in \mathcal{N}_p} \mathbb{P}(|\vartheta_j| > z_{\frac{\alpha}{2}} - \epsilon) + \frac{1}{|\mathcal{N}_p|} \sum_{j \in \mathcal{N}_p} \mathbb{P}(|\varsigma_j| > \epsilon) \\ &\leq \frac{1}{|\mathcal{N}_p|} \sum_{j \in \mathcal{N}_p} \mathbb{P}(|\vartheta_j| > z_{\frac{\alpha}{2}} - \epsilon) + \mathbb{P}\left(\max_{1 \leq j \leq p} |\varsigma_j| > \epsilon\right) \\ &\rightarrow 2(1 - \Phi(z_{\frac{\alpha}{2}} - \epsilon)), \end{aligned}$$



where the last convergence is because the Cesaro mean converges to the same limit as  $\lim_{n,p} \mathbb{P}(|\vartheta_j| > z_{\frac{\alpha}{2}} - \epsilon) = 2(1 - \Phi(z_{\frac{\alpha}{2}} - \epsilon))$  and the second term  $\mathbb{P}(\max_{1 \leq j \leq p} |\varsigma_j| > \epsilon)$  vanishes. Similarly, we can also show that  $\liminf_{n,p \rightarrow \infty} \mathbb{E}[\varrho] \geq 2(1 - \Phi(z_{\frac{\alpha}{2}} - \epsilon))$  for all  $\epsilon > 0$ . This implies that  $\mathbb{E}[\varrho] \rightarrow \alpha$  as  $n, p \rightarrow \infty$ .

Next, we analyze the second moment. For any  $\epsilon > 0$ , the second moment can be upper bounded as

$$\begin{aligned}
\mathbb{E}[\varrho^2] &= \frac{1}{|\mathcal{N}_p|^2} \sum_{j,k \in \mathcal{N}_p} \mathbb{P}(|t_j| > z_{\frac{\alpha}{2}}, |t_k| > z_{\frac{\alpha}{2}}) \\
&= \frac{1}{|\mathcal{N}_p|^2} \sum_{j \in \mathcal{N}_p} \mathbb{P}(|t_j| > z_{\frac{\alpha}{2}}) + \frac{1}{|\mathcal{N}_p|^2} \sum_{j,k \in \mathcal{N}_p, j \neq k} \mathbb{P}(|t_j| > z_{\frac{\alpha}{2}}, |t_k| > z_{\frac{\alpha}{2}}) \\
&\leq \frac{1}{|\mathcal{N}_p|^2} \sum_{j \in \mathcal{N}_p} \mathbb{P}(|t_j| > z_{\frac{\alpha}{2}}) \\
&\quad + \frac{1}{|\mathcal{N}_p|^2} \sum_{j,k \in \mathcal{N}_p, j \neq k} \mathbb{P}(|\vartheta_j| > z_{\frac{\alpha}{2}} - \epsilon, |\vartheta_k| > z_{\frac{\alpha}{2}} - \epsilon) + \mathbb{P}(|\varsigma_j| > \epsilon) + \mathbb{P}(|\varsigma_k| > \epsilon) \\
&= \frac{1}{|\mathcal{N}_p|^2} \sum_{j,k \in \mathcal{N}_p, j \neq k} \mathbb{P}(|\vartheta_j| > z_{\frac{\alpha}{2}} - \epsilon, |\vartheta_k| > z_{\frac{\alpha}{2}} - \epsilon) + o(1) \\
&= \frac{1}{|\mathcal{N}_p|^2} \sum_{j,k \in \mathcal{N}_p, j \neq k} \mathbb{P}(|\vartheta_j| > z_{\frac{\alpha}{2}} - \epsilon) \mathbb{P}(|\vartheta_k| > z_{\frac{\alpha}{2}} - \epsilon) + o(1) \\
&\rightarrow |\mathcal{N}_p| [2(1 - \Phi(z_{\frac{\alpha}{2}} - \epsilon))]^2,
\end{aligned}$$

where the last equality is from the independence of  $\vartheta_j$  and  $\vartheta_k$ . We can similarly obtain the lower bound. This implies that  $\mathbb{E}[\varrho^2] \rightarrow \alpha^2$  and  $\text{Var}(\varrho) \rightarrow 0$  as  $n, p \rightarrow \infty$ . Combining the previous results yields that  $\varrho \xrightarrow{\mathbb{P}} \alpha$ .

**FWER control.** To prove the second statement, note that

$$\begin{aligned}
\mathbb{P}(|\mathcal{N}_p| \varrho \geq 1) &= \mathbb{P}\left(\max_{j \in \mathcal{N}_p} |t_j| > \Phi^{-1}(1 - \alpha/(2p))\right) \\
&= \mathbb{P}\left(\max_{j \in \mathcal{N}_p} |\vartheta_j + \varsigma_j| > \Phi^{-1}(1 - \alpha/(2p))\right) \\
&\leq \mathbb{P}\left(\max_{j \in \mathcal{N}_p} |\vartheta_j| > \Phi^{-1}(1 - \alpha/(2p)) - \max_{j \in \mathcal{N}_p} |\varsigma_j|\right) \\
&\leq \mathbb{P}\left(\max_{1 \leq j \leq p} |\vartheta_j| > \Phi^{-1}(1 - \alpha/(2p)) - \max_{j \in \mathcal{N}_p} |\varsigma_j|\right),
\end{aligned}$$

which is asymptotically upper bounded by  $\alpha$  because of the valid control of Bonferroni correction for i.i.d. normal random variables, the fact that  $\Phi^{-1}(1 - \alpha/(2p)) \rightarrow \infty$  as  $p \rightarrow \infty$ , and the result that  $\max_{j \in \mathcal{N}_p} |\varsigma_j| = o_{\mathbb{P}}(1)$ .  $\square$

### E.3 Technical lemmas

**Lemma E.1.** Under the same conditions as in Theorem 6, suppose event  $\mathcal{E}_1$  holds, then the solution to optimization problem (4.4) exists with probability at least  $1 - 2(nd)^{-c}$ .

*Proof of Lemma E.1.* Define the matrix  $\mathbf{S} = \mathbb{E}[\widehat{\omega}_i \mathbf{x}_i \mathbf{x}_i^\top]$ . We next show that (1)  $\mathbf{S}$  is invertible and (2) the  $j$ -th column  $\mathbf{u}^*$  of  $\mathbf{S}^{-1}$  is feasible for the constraints of the optimization problem (4.4) with high probability. We split the proof into two parts, as below.

**Part (1)** Because  $C^{-1} \leq \widehat{\omega}_i \leq C$ , we have  $C^{-1} \mathbb{E}[\mathbf{x}_1 \mathbf{x}_1^\top] \preceq \mathbf{S} \preceq C \mathbb{E}[\mathbf{x}_1 \mathbf{x}_1^\top]$ . On the other hand, note that  $\mathbb{E}[\mathbf{x}_1 \mathbf{x}_1^\top] = \boldsymbol{\Sigma}_x \succeq \lambda_{\min}(\boldsymbol{\Sigma}_x) \mathbf{I}_d$ . Thus, for any unit vector  $\mathbf{a} \in \mathbb{R}^d$ , we have  $\mathbf{a}^\top \mathbf{S} \mathbf{a} \geq C^{-1} \lambda_{\min}(\boldsymbol{\Sigma}_x) > 0$ . This establishes claim (1).

**Part (2)** Let  $\mathbf{u}^*$  be the  $j$ -th column of  $\mathbf{S}^{-1}$ . By definition, we have  $\mathbf{S} \mathbf{u}^* = \mathbf{e}_j$ . Conditional on  $\mathcal{D}_2$ , we have that  $\widehat{\omega}_i \mathbf{u}^{*\top} \mathbf{x}_i \mathbf{x}_i^\top \mathbf{e}_k$  for  $i = 1, \dots, n$  are independent random variables with mean  $\delta_{jk}$ . Because  $\widehat{\omega}_i$  is bounded, we further have that  $\widehat{\omega}_i \mathbf{u}^{*\top} \mathbf{x}_i \mathbf{x}_i^\top \mathbf{e}_k$ 's are independent sub-exponential random variables. Applying Bernstein's inequality as in the proof of Lemma D.2, we have with probability at least  $1 - (nd)^{-c}$ ,

$$\left\| \frac{1}{n} \sum_{i=1}^n \widehat{\omega}_i \mathbf{u}^{*\top} \mathbf{x}_i \mathbf{x}_i^\top - \mathbf{e}_j \right\|_\infty \leq \lambda_n,$$

where  $\lambda_n \asymp \sqrt{\log(nd)/n}$ . This also holds after taking account of the randomness of  $\mathcal{D}_2$ .

On the other hand, because  $\omega_i$  is bounded away from zero and infinity, and  $\|\boldsymbol{\Sigma}_x\|_{\text{op}} = \mathcal{O}(1)$ , it follows that  $\|\mathbf{S}\|_{\text{op}} = \mathcal{O}(1)$  and  $\|\mathbf{u}^*\|_2 = \mathcal{O}(1)$ . By the sub-Gaussianity of  $\mathbf{x}_i$ , we also have  $\|\mathbf{X} \mathbf{u}^*\|_\infty = \max_{1 \leq i \leq n} |\mathbf{x}_i^\top \mathbf{u}^*| \leq \tau_n$ , with probability at least  $1 - n^{-c}$ . The above shows that  $\mathbf{u}^*$  is feasible for optimization problem (4.4), which establishes the claim (2).

Finally, taking the union bound over the two probabilistic events finishes the proof.  $\square$

**Lemma E.2** (Asymptotic normality). Under the conditions in Theorem 6, it holds that

$$\sqrt{n} \sum_{i=1}^n \sigma_j^{-1} \widehat{\mathbf{u}}^\top \mathbf{x}_i \boldsymbol{\epsilon}_i \widehat{\mathbf{v}}_i \xrightarrow{d} \mathcal{N}(0, 1),$$

where  $\sigma_j^2 = n^{-1} \text{Var}(\mathbf{u}^\top \sum_{i=1}^n \mathbf{x}_i \boldsymbol{\epsilon}_i^\top \mathbf{v}_i \mid \{\mathbf{x}_i, \mathbf{z}_i\}_{i=1}^n, \mathcal{D}_2)$ .

*Proof of Lemma E.2.* Note that when conditioning on the natural parameters  $\boldsymbol{\Theta}^*$ ,  $\boldsymbol{\epsilon}_{ij}$ 's are independent  $(\nu, \alpha)$ -sub-exponential random variable as shown in the proof of Theorem 2. Define  $\xi_i := \sigma_j^{-1} \widehat{\mathbf{u}}^\top \mathbf{x}_i \boldsymbol{\epsilon}_i \widehat{\mathbf{v}}_i$  for  $i \in [n]$ . Then  $\xi_i$ 's are independent random variables with mean  $\mathbb{E}[\xi_i \mid \mathcal{D}_2, \mathbf{X}] = 0$  and variance  $\text{Var}(\xi_i \mid \mathcal{D}_2, \mathbf{X}) = 1$ . It suffices to check the bounded variance condition and Lindeberg's condition.

**Part (1) Boundedness of  $\sigma_j$ .** We first show the boundedness of the variance

$$\sigma_j^2 = \widehat{\mathbf{u}}^\top \frac{1}{n} \sum_{i=1}^n \widehat{\omega}_i^2 (\mathbf{e}_j^\top \mathcal{P}_{\widehat{\mathbf{F}}}^\perp \text{diag}(A''(\widehat{\boldsymbol{\theta}}_i))^{-1} \text{diag}(A''(\boldsymbol{\theta}_i^*)) \text{diag}(A''(\widehat{\boldsymbol{\theta}}_i))^{-1} \mathcal{P}_{\widehat{\mathbf{F}}}^\perp \mathbf{e}_j) \mathbf{x}_i \mathbf{x}_i^\top \widehat{\mathbf{u}}.$$

Because  $A''(\theta) \geq \kappa_1 > 0$  for all  $\theta \in \mathcal{R}$ , the quadratic term satisfies that

$$\mathbf{e}_j \mathcal{P}_{\hat{\mathbf{r}}}^\perp \text{diag}(A''(\hat{\boldsymbol{\theta}}_i))^{-1} \text{diag}(A''(\boldsymbol{\theta}_i^*)) \text{diag}(A''(\hat{\boldsymbol{\theta}}_i))^{-1} \mathcal{P}_{\hat{\mathbf{r}}}^\perp \mathbf{e}_j \geq 0,$$

with equality holds if and only if  $\mathcal{P}_{\hat{\mathbf{r}}}^\perp \mathbf{e}_j = \mathbf{0}_p$ . On the other hand, we have

$$\|\mathcal{P}_{\hat{\mathbf{r}}}^\perp \mathbf{e}_j\|_2 = \|\mathbf{e}_j - \mathcal{P}_{\hat{\mathbf{r}}} \mathbf{e}_j\|_2 \geq 1 - \|\mathcal{P}_{\hat{\mathbf{r}}} \mathbf{e}_j\|_2 \gtrsim \Omega(1 - p^{-1/2}),$$

where the last inequality is because

$$\|\mathcal{P}_{\hat{\mathbf{r}}} \mathbf{e}_j\|_2 \leq \|\mathcal{P}_{\mathbf{r}^*} \mathbf{e}_j\|_2 + \|(\mathcal{P}_{\mathbf{r}^*} - \mathcal{P}_{\hat{\mathbf{r}}}) \mathbf{e}_j\|_2 \lesssim \frac{1}{\sqrt{p}}$$

from Assumption 3 and Theorem 3. This implies that, when  $n, p$  are sufficiently large,

$$c^{-1} \leq \hat{\omega}_i (\mathbf{e}_j^\top \mathcal{P}_{\hat{\mathbf{r}}}^\perp \text{diag}(A''(\hat{\boldsymbol{\theta}}_i))^{-1} \text{diag}(A''(\boldsymbol{\theta}_i^*)) \text{diag}(A''(\hat{\boldsymbol{\theta}}_i))^{-1} \mathcal{P}_{\hat{\mathbf{r}}}^\perp \mathbf{e}_j) \leq c$$

for some constant  $c > 1$ , under event  $\mathcal{E}_1$ . Thus, it is equivalent to show the boundedness of  $\hat{\sigma}_j^2 = \hat{\mathbf{u}}^\top \hat{\mathbf{S}} \hat{\mathbf{u}}$ , where  $\hat{\mathbf{S}} = n^{-1} \sum_{i=1}^n \hat{\omega}_i \mathbf{x}_i \mathbf{x}_i^\top$ . From Lemma E.1, we know that  $\mathbf{S} = \mathbb{E}[\hat{\mathbf{S}}]$  has a bounded spectrum with high probability. The upper bound that  $\hat{\sigma}_j^2 \leq \mathbf{S}_{jj}$  with high probability then follows by the sub-exponential concentration results as in the proof of Lemma E.1.

Next, we proceed to show the lower bound. Because  $\hat{\mathbf{u}}$  satisfies the constraint  $|\mathbf{e}_j^\top \hat{\mathbf{S}} \hat{\mathbf{u}} - 1| \leq \lambda_n$ , we have that  $\sigma_j^2 \geq \hat{\mathbf{u}}^\top \hat{\mathbf{S}} \hat{\mathbf{u}} + t((1 - \lambda_n) - \mathbf{e}_j^\top \hat{\mathbf{S}} \hat{\mathbf{u}})$  for any  $t > 0$ . Note that  $\min_{\mathbf{v} \in \mathbb{R}^d} \mathbf{v}^\top \hat{\mathbf{S}} \mathbf{v} + t((1 - \lambda_n) - \mathbf{e}_j^\top \hat{\mathbf{S}} \mathbf{v}) = -t^2 \mathbf{e}_j^\top \hat{\mathbf{S}} \mathbf{e}_j / 4 + t(1 - \lambda_n)$  where the minimum is obtained when  $\hat{\mathbf{S}} \mathbf{v} = t \hat{\mathbf{S}} \mathbf{e}_j / 2$ . We further have  $\hat{\sigma}_j^2 \geq \max_{t \geq 0} -t^2 \mathbf{e}_j^\top \hat{\mathbf{S}} \mathbf{e}_j / 4 + t(1 - \lambda_n) \geq (1 - \lambda_n)^2 / (\mathbf{e}_j^\top \hat{\mathbf{S}} \mathbf{e}_j)$ . By the sub-Gaussianity of  $\mathbf{x}_i$ ,  $\mathbf{e}_j^\top \hat{\mathbf{S}} \mathbf{e}_j \geq \mathbf{S}_{jj} + o_{\mathbb{P}}(1)$ . We then have  $\hat{\sigma}_j^2 \geq 0.5 / \mathbf{S}_{jj}$  when  $n$  and  $p$  are large enough.

**Part (2) Lindeberg's condition.** On the other hand, because

$$\max_{1 \leq i \leq n} |\xi_i| \leq \max_{1 \leq i \leq n} |\hat{\mathbf{u}}^\top \mathbf{x}_i| \|\boldsymbol{\epsilon}_i\|_2 \sigma_j^{-1} \|\hat{\mathbf{v}}_i\|_2 \lesssim \sqrt{n},$$

with probability at least  $1 - 2(nd)^{-c}$ , the Lindeberg's condition holds that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\xi_i^2 \mathbb{1}\{|\xi_i| \geq \epsilon \sqrt{n}\}] = 0$$

for all  $\epsilon > 0$ . Applying Lindeberg's central limit theorem yields that

$$\sqrt{n} \sum_{i=1}^n \sigma_j^{-1} \hat{\mathbf{u}}^\top \mathbf{x}_i \boldsymbol{\epsilon}_i \hat{\mathbf{v}}_i \xrightarrow{d} \mathcal{N}(0, 1),$$

which finishes the proof.  $\square$

**Lemma E.3** (Consistent estimators of  $\sigma_j$ ). Under the conditions in Theorem 6,  $\widehat{\omega}_j = A''(\widehat{\boldsymbol{\theta}}_{ij})$  satisfies condition (ii) of Theorem 6. Furthermore, for the variance estimate defined in (4.6) using sample splitting procedure Algorithm E.2, it holds that  $\widehat{\sigma}_j \xrightarrow{P} \sigma_j$ .

*Proof of Lemma E.3.* The boundedness of  $\widehat{\omega}_j$  follows from (B.1).

By the sample splitting procedure Algorithm E.2, we know that for a given response  $j \in [p]$ ,  $\mathbf{Y}_j - A'(\boldsymbol{\Theta}_j^*) \in \mathbb{R}^n$  is independent of  $\widehat{\mathbf{u}}$ ,  $\widehat{\mathbf{B}}$ ,  $\widehat{\boldsymbol{\Gamma}}$ , and  $\widehat{\mathbf{Z}}$ , when conditioning on  $\mathbf{X}$ . However, noted that  $\widehat{\mathbf{B}}$ ,  $\widehat{\boldsymbol{\Gamma}}$ , and  $\widehat{\mathbf{Z}}$  may be specific to each  $j \in I$ . For the sake of simplicity, in the following proof, we will assume that  $\mathbf{Y} - A'(\boldsymbol{\Theta}^*)$  is independent of  $\widehat{\mathbf{u}}$ , and a common set of estimators  $\widehat{\mathbf{B}}$ ,  $\widehat{\boldsymbol{\Gamma}}$ , and  $\widehat{\mathbf{Z}}$  when conditioning on  $\mathbf{X}$ ; or equivalently  $I = [p]$ . Note that, however, the proof still works for the cases in Algorithm E.2, except the constructed debiased estimators only use responses in the index set  $I$ ; namely,  $\widehat{b}_{j1}^{\text{de}} = \widehat{b}_{j1} + \widehat{\mathbf{u}}^\top \frac{1}{n} \sum_{i=1}^n \widehat{\omega}_i \mathbf{x}_i ([\mathbf{y}_i]_I - [A'(\widehat{\boldsymbol{\theta}}_i)]_I)^\top \mathcal{P}_{\widehat{\boldsymbol{\Gamma}}_I}^\perp \mathbf{e}_j$ .

Recall that

$$\sigma_j^2 = \widehat{\mathbf{u}}^\top \frac{1}{n} \sum_{i=1}^n \widehat{\omega}_i^2 (\mathbf{e}_j^\top \mathcal{P}_{\widehat{\boldsymbol{\Gamma}}}^\perp \text{diag}(A''(\widehat{\boldsymbol{\theta}}_i))^{-1} \text{diag}(A''(\boldsymbol{\theta}_i^*)) \text{diag}(A''(\widehat{\boldsymbol{\theta}}_i))^{-1} \mathcal{P}_{\widehat{\boldsymbol{\Gamma}}}^\perp \mathbf{e}_j) \mathbf{x}_i \mathbf{x}_i^\top \widehat{\mathbf{u}},$$

and

$$\widehat{\sigma}_j^2 = \widehat{\mathbf{u}}^\top \frac{1}{n} \sum_{i=1}^n \widehat{\omega}_i \mathbf{x}_i \mathbf{x}_i^\top \widehat{\mathbf{u}}.$$

Let  $a_i = \widehat{\omega}_i^2 (\mathbf{e}_j^\top \mathcal{P}_{\widehat{\boldsymbol{\Gamma}}}^\perp \text{diag}(A''(\widehat{\boldsymbol{\theta}}_i))^{-1} \text{diag}(A''(\boldsymbol{\theta}_i^*)) \text{diag}(A''(\widehat{\boldsymbol{\theta}}_i))^{-1} \mathcal{P}_{\widehat{\boldsymbol{\Gamma}}}^\perp \mathbf{e}_j)$  and  $a'_i = \widehat{\omega}_i$ . We begin by bounding the difference between the two:

$$\begin{aligned} |\widehat{\sigma}_j^2 - \sigma_j^2| &= \left| \frac{1}{n} \sum_{i=1}^n (a'_i - a_i) \cdot (\widehat{\mathbf{u}} \mathbf{x}_i)^2 \right| \\ &\leq \sqrt{\frac{1}{n} \sum_{i=1}^n (\widehat{\mathbf{u}} \mathbf{x}_i)^4} \cdot \sqrt{\frac{1}{n} \sum_{i=1}^n (a'_i - a_i)^2} \\ &\lesssim \tau_n^2 \cdot \sqrt{\frac{1}{n} \sum_{i=1}^n (a'_i - a_i)^2}, \end{aligned}$$

where the first inequality is from Holder's inequality and the second inequality is due to the second constraint of the optimization problem (4.4). For the second factor above, note that

$$\begin{aligned} &\frac{1}{n} \sum_{i=1}^n (a'_i - a_i)^2 \\ &= \max_{i \in [n]} \widehat{\omega}_i^4 \cdot \frac{1}{n} \sum_{i=1}^n (\mathbf{e}_j^\top \mathcal{P}_{\widehat{\boldsymbol{\Gamma}}}^\perp \text{diag}(A''(\widehat{\boldsymbol{\theta}}_i))^{-1} \text{diag}(A''(\boldsymbol{\theta}_i^*)) \text{diag}(A''(\widehat{\boldsymbol{\theta}}_i))^{-1} \mathcal{P}_{\widehat{\boldsymbol{\Gamma}}}^\perp \mathbf{e}_j - A''(\widehat{\boldsymbol{\theta}}_i))^2. \quad (\text{E.4}) \end{aligned}$$

Each term inside the square can be decomposed into

$$\begin{aligned}
& \mathbf{e}_j^\top \mathcal{P}_{\widehat{\mathbf{F}}}^\perp \text{diag}(A''(\widehat{\boldsymbol{\theta}}_i))^{-1} \text{diag}(A''(\boldsymbol{\theta}_i^*)) \text{diag}(A''(\widehat{\boldsymbol{\theta}}_i))^{-1} \mathcal{P}_{\widehat{\mathbf{F}}}^\perp \mathbf{e}_j - A''(\widehat{\boldsymbol{\theta}}_{ij}) \\
&= \mathbf{e}_j^\top \mathcal{P}_{\widehat{\mathbf{F}}}^\perp \text{diag}(A''(\widehat{\boldsymbol{\theta}}_i))^{-1} \text{diag}(A''(\boldsymbol{\theta}_i^*) - A''(\widehat{\boldsymbol{\theta}}_i)) \text{diag}(A''(\widehat{\boldsymbol{\theta}}_i))^{-1} \mathcal{P}_{\widehat{\mathbf{F}}}^\perp \mathbf{e}_j \\
&\quad - [2\mathbf{e}_j^\top \mathcal{P}_{\widehat{\mathbf{F}}} A''(\widehat{\boldsymbol{\theta}}_i) \mathcal{P}_{\widehat{\mathbf{F}}}^\perp \mathbf{e}_j + \mathbf{e}_j^\top \mathcal{P}_{\widehat{\mathbf{F}}} A''(\widehat{\boldsymbol{\theta}}_i) \mathcal{P}_{\widehat{\mathbf{F}}} \mathbf{e}_j] \\
&=: T_1 + T_2.
\end{aligned} \tag{E.5}$$

Now note that

$$\begin{aligned}
\frac{1}{n} \sum_{i=1}^n T_1^2 &\lesssim \frac{1}{n} \sum_{i=1}^n (A''(\boldsymbol{\theta}_{ij}^*) - A''(\widehat{\boldsymbol{\theta}}_{ij}))^2 \\
&= \frac{1}{n} (\widehat{\boldsymbol{\Theta}}_j - \boldsymbol{\Theta}_j^*)^\top \text{diag}(A'''(\boldsymbol{\Theta}'_j)) (\widehat{\boldsymbol{\Theta}}_j - \boldsymbol{\Theta}_j^*) \\
&\lesssim \frac{1}{n} \|\widehat{\boldsymbol{\Theta}}_j - \boldsymbol{\Theta}_j^*\|_2^2,
\end{aligned} \tag{E.6}$$

where the first inequality is due to the boundedness of  $A''$  on  $\mathcal{R}$ , and the bounded spectral of the projection matrix  $\mathcal{P}_{\widehat{\mathbf{F}}}^\perp$ , and noting that  $\|\mathcal{P}_{\widehat{\mathbf{F}}} \mathbf{e}_j\|_2 \lesssim \mathcal{O}_{\mathbb{P}}(p^{-1/2})$ ; the second equality is from Taylor expansion with  $\boldsymbol{\Theta}'_j = (\theta'_{1j}, \dots, \theta'_{nj})$  and  $\theta'_{ij}$  being between  $\widehat{\theta}_{ij}$  and  $\theta_{ij}^*$  for  $i = 1, \dots, n$ ; and the second inequality is from the continuity and boundedness of  $A'''$  on  $\mathcal{R}_C$ .

On the other hand,

$$\frac{1}{n} \sum_{i=1}^n T_2^2 \lesssim p^{-1} \tag{E.7}$$

by noting that  $\|\mathcal{P}_{\widehat{\mathbf{F}}} \mathbf{e}_j\|_2 \lesssim \mathcal{O}_{\mathbb{P}}(p^{-1/2})$  again.

By applying triangle inequality on (E.4) and combining (E.5)-(E.7), we further have

$$\begin{aligned}
& \frac{1}{n} \sum_{i=1}^n (a'_i - a_i)^2 \\
&\lesssim \max_{i \in [n]} \widehat{\omega}_i^4 \cdot \left( \frac{1}{n} \sum_{i=1}^n T_1^2 + \frac{1}{n} \sum_{i=1}^n T_2^2 \right) \\
&\lesssim \frac{1}{n} \|\widehat{\boldsymbol{\Theta}}_j - \boldsymbol{\Theta}_j^*\|_2^2,
\end{aligned}$$

where in the last inequality, we also use the boundedness of  $\widehat{\omega}_i$ . This implies that

$$\begin{aligned}
|\widehat{\sigma}_j^2 - \sigma_j^2| &\lesssim \tau_n^2 \sqrt{\frac{1}{n} \|\widehat{\boldsymbol{\Theta}}_j - \boldsymbol{\Theta}_j^*\|_2^2} \\
&\lesssim \tau_n^2 \sqrt{\frac{1}{n} \|\mathbf{X}(\widehat{\mathbf{b}}_j - \mathbf{b}_j^*)\|_2^2 + \frac{1}{n} \|\widehat{\mathbf{E}}_j - \mathbf{E}_j^*\|_2^2} \\
&\lesssim \tau_n^2 r_{n,p} \\
&= o_{\mathbb{P}}(1),
\end{aligned}$$

where  $r_{n,p}$  is defined in Theorem 5. Here the concentration of  $\|\mathbf{X}(\widehat{\mathbf{b}}_j - \mathbf{b}_j^*)\|_2^2$  is from Theorem 5 and the one of  $\|\widehat{\mathbf{E}}_j - \mathbf{E}_j^*\|_2^2$  is from (D.24) as in the proof of Lemma D.3. This implies that  $\widehat{\sigma}_j^2 \xrightarrow{P} \sigma_j^2$ . The conclusion then follows by applying the continuous mapping theorem.  $\square$

---

**Algorithm E.2** Data splitting procedure.

---

**Input:** Data  $(\mathbf{x}_i, \mathbf{y}_i) \in \mathbb{R}^d \times \mathbb{R}^p$  for  $i = 1, \dots, 2n$ .

- 1: Split the full data into two disjoint datasets  $\mathcal{D}_1 = \{(\mathbf{x}_i, \mathbf{y}_i) \mid i = 1, \dots, n\}$  and  $\mathcal{D}_2 = \{(\mathbf{x}_i, \mathbf{y}_i) \mid i = n + 1, \dots, 2n\}$ .
- 2: Apply Algorithm 1 on  $\mathcal{D}_2$  to obtain the estimates  $\widehat{\mathbf{B}}$  and  $\widehat{\mathbf{\Gamma}}$ .
- 3: **for**  $j = 1, 2, \dots, p$  **do**
- 4:   Select a subset  $I \subseteq [p] \cap \{j\}$  and set  $I^c = [p] \setminus I$ .
- 5:   Based on  $\widehat{\mathbf{B}}$  and  $\widehat{\mathbf{\Gamma}}$ , use partial data  $(\mathbf{X}, \mathbf{Y}_{I^c})$  to estimate  $\widehat{\mathbf{Z}}$ , where  $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n]^\top$ ,  $\mathbf{Y} = [\mathbf{y}_1, \dots, \mathbf{y}_n]^\top$  and  $\mathbf{Y}_{I^c} = [\mathbf{Y}_\ell]_{\ell \in I^c}$ .  
(Alternatively, Step 2-5 can be combined such that  $\widehat{\mathbf{B}}$ ,  $\widehat{\mathbf{\Gamma}}$ , and  $\widehat{\mathbf{Z}}$  are estimated jointly for gene  $j$ .)
- 6:   Based on  $\widehat{\mathbf{B}}$ ,  $\widehat{\mathbf{\Gamma}}$ , and  $\widehat{\mathbf{Z}}$ , estimate  $\widehat{\omega}_i$ 's and  $\widehat{\mathbf{u}}$  on  $(\mathbf{X}, \mathbf{Y}_I)$ .
- 7:   Calculate the test statistics  $z_j$  for gene  $j$ .
- 8: **end for**

**Output:** A set of test statistics  $\{z_j \mid j = 1, \dots, p\}$ .

---

## F Computational aspects

### F.1 Exponential family

Some commonly used exponential families, the exact formulas of the log-partition functions and other statistics, are summarized in Table F1.

Distribution	Extra parameter	Base measure $h(y)$	Sufficient statistics $T(y)$	Domain $\text{dom}(A(\theta))$	Log-partition $A(\theta)$	Mean $\mu = A'(\theta)$	Variance $A''(\theta)$
Gaussian	variance $\sigma^2$	$\frac{e^{-\frac{x^2}{2\sigma^2}}}{\sqrt{2\pi}\sigma}$	$\frac{y}{\sigma}$	$\mathbb{R}$	$\frac{\theta^2}{2}$	$\theta$	1
Bernoulli		1	$y$	$\mathbb{R}$	$\log(1 + e^\theta)$	$\frac{1}{1 + e^{-\theta}}$	$\mu(1 - \mu)$
Binomial	number of trials $m$	$\binom{m}{y}$	$y$	$\mathbb{R}$	$m \log(1 + e^\theta)$	$\frac{m}{1 + e^{-\theta}}$	$\mu \left(1 - \frac{\mu}{m}\right)$
Poisson		$\frac{1}{y!}$	$y$	$\mathbb{R}$	$e^\theta$	$e^\theta$	$e^\theta$
Negative Binomial	number of failures $\phi$	$\binom{y + \phi - 1}{y}$	$y$	$\mathbb{R}_-$	$-\phi \log(1 - e^\theta)$	$\phi \frac{e^\theta}{1 - e^\theta}$	$\phi \frac{e^\theta}{(1 - e^\theta)^2}$

**Table F1:** Summary of exponential family in canonical form.

## F.2 Optimization details

**Initialization.** Our initialization procedure for optimization problem (3.4) is inspired by Lin et al. (2024).

- Initialize the marginal effects  $\mathbf{F}$  by solving a generalized linear model without considering the latent variables. When the fitting of GLM is numerically unstable, one can also add a small ridge penalty  $\lambda = 10^{-5}$ .
- Initialize  $\mathbf{W}$  and  $\mathbf{\Gamma}$  using the SVD of the matrix  $\log(\mathbf{Y} + 1) = \mathbf{U}_Y \mathbf{\Sigma}_Y \mathbf{V}_Y^\top$  for Poisson likelihood or Negative Binomial likelihood with log link. Let  $\mathbf{W} = (\mathcal{P}_{\mathbf{X}}^\perp \mathbf{U}_Y \mathbf{\Sigma}_Y^{1/2})_{1:r}$  and  $\mathbf{\Gamma} = (\mathbf{V}_Y \mathbf{\Sigma}_Y^{1/2})_{1:r}$  be the first  $r$  columns of the corresponding matrices. Here the projection  $\mathcal{P}_{\mathbf{X}}^\perp$  ensures that  $\mathbf{W}$  is uncorrelated with  $\mathbf{X}$ . In particular, when the intercept is included in the covariates, the initial value of  $\mathbf{W}$  also has zero means per column.

To initialize variables for optimization problem (3.5):

- Initialize the direct effects  $\mathbf{B}$  as  $\mathcal{P}_{\hat{\mathbf{F}}}^\perp \hat{\mathbf{F}}$ .
- Initialize  $\mathbf{Z}$  and  $\mathbf{\Gamma}$  using the SVD of the matrix  $\mathbf{X} \hat{\mathbf{F}}^\top \mathcal{P}_{\hat{\mathbf{F}}} + \hat{\mathbf{W}} \hat{\mathbf{\Gamma}}^\top = \mathbf{U}' \mathbf{\Sigma}' \mathbf{V}'^\top$ . Let  $\mathbf{Z} = (\mathbf{U}' \mathbf{\Sigma}'^{1/2})_{1:r}$  and  $\mathbf{\Gamma} = (\mathbf{V}' \mathbf{\Sigma}'^{1/2})_{1:r}$ . Because the latter has the same column space as  $\hat{\mathbf{\Gamma}}$ , we simply treat the latter as  $\hat{\mathbf{\Gamma}}$  in optimization problem (3.5) with a light abuse of notation.

**Alternative maximization** The alternative maximization Algorithm F.3 is used to perform nonconvex matrix factorization. In our setup where the objective function is convex in the natural parameter, each iteration of Algorithm F.3 is simply solving two convex optimization subproblems.

By default, we use the inexact line search algorithm with an initial step size of 0.1 and a shrinkage factor of 0.5 for each iteration. The search is stopped if the Armijo rule is satisfied with tolerance  $10^{-4}$  or the number of iterations reaches 20. We early stop the alternative maximization if the objective value does not increase more than a tolerance of  $10^{-4}$  for 20 iterations.

**Estimation of dispersion parameters** To estimate the dispersion parameter, we first fit GLMs on the data and obtain the estimated mean expression of gene  $j$ , denoted as  $\hat{\mu}_j$  for  $j = 1, \dots, p$ . Note that when  $y_{ij}$  comes from a Negative Binomial distribution, its variance is given by

$$\text{Var}(y_{ij} \mid \theta_{ij}) = \mu (1 + \alpha_j \mu)$$

where  $\mu = \mathbb{E}[y_{ij} \mid \theta_{ij}]$  is the conditional mean while  $\alpha_j$  is the dispersion parameter of the NB1 form. In the form of exponential family in Table F1 parameterized by the parameter  $\phi_j$ ,  $\alpha_j$  is the reciprocal of  $\phi_j$ , namely,  $\alpha_j = 1/\phi_j$ . By methods of moments, we can solve the

---

**Algorithm F.3** Joint maximum likelihood estimation by alternative maximization

---

**Input:** Data  $\mathbf{Y} \in \mathbb{R}^{n \times p}$  from exponential family with log-partition function  $A$ , the regularization parameter  $\lambda$ , and initial value  $\mathbf{l}_i^{(0)} \in \mathcal{D}_{l_i}$ ,  $\mathbf{r}_j^{(0)} \in \mathcal{D}_{r_j}$  for  $i \in [n]$  and  $j \in [p]$ .

1: Initialize the iteration number  $t = 0$ .

2: **while** not converged **do**

3:    $t \leftarrow t + 1$ .

4:   **for**  $i = 1, 2, \dots, n$  **do**

5:

$$\mathbf{l}_i^{(t)} \in \operatorname{argmax}_{\mathbf{l} \in \mathcal{D}_{l_i}} \frac{1}{p} \sum_{j=1}^p \left( y_{ij} \mathbf{l}^\top \mathbf{r}_j^{(t-1)} - A(\mathbf{l}^\top \mathbf{r}_j^{(t-1)}) \right)$$

6:   **end for**

7:   **for**  $j = 1, 2, \dots, p$  **do**

8:

$$\mathbf{r}_j^{(t)} \in \operatorname{argmax}_{\mathbf{r} \in \mathcal{D}_{r_j}} \frac{1}{n} \sum_{i=1}^n \left( y_{ij} \mathbf{l}_i^{(t)\top} \mathbf{r} - A(\mathbf{l}_i^{(t)\top} \mathbf{r}) \right) - \lambda \|\mathbf{r}\|_1$$

9:   **end for**

10: **end while**

**Output:**  $\mathbf{L} = [\mathbf{l}_i^{(t)}]_{i \in [n]}^\top$  and  $\mathbf{R} = [\mathbf{r}_j^{(t)}]_{j \in [p]}^\top$  with  $\mathbf{LR}^\top$  being the estimated natural parameters.

---

following equation to obtain an estimator  $\hat{\phi}_j$  for  $\phi_j$ :

$$\frac{1}{n} \sum_{i=1}^n (y_{ij} - \hat{\mu}_j)^2 = \hat{\mu}_j (1 + \alpha \hat{\mu}_j).$$

Finally, we clip  $\hat{\alpha}_j$  to be in  $[10^{-2}, 10^2]$  and set  $\hat{\phi}_j = 1/\hat{\alpha}_j$ .

### F.3 Choice of hyperparameters in practice

The main text provides theoretical results for the proposed method under certain assumptions. The proposed algorithm Algorithm 1 requires the choice of hyperparameters, such as the rank  $r$ , the regularization parameters  $\lambda$  in optimization problem (3.5), and  $(\tau_n, \lambda_n)$  in optimization problem (4.4). Although the theoretical orders of some parameters are provided for consistency and asymptotic normality, the choice of hyperparameters in practice is crucial for the performance of the proposed method. Below, we discuss the choice of hyperparameters in practice.

**Boundedness constant  $C$ .** The boundedness constant  $C$  is a reasonably large constant to ensure a finite solution to optimization problems exists. In our simulations, estimating the model parameters is not sensitive to the choice of boundedness constant as long as it is set to be sufficiently large; see also Chen and Li (2022, Appendix D) for detailed discussions. Therefore, in our implementation, instead of restricting the parameters to be bounded, we project the gradient at each step of the alternative maximization onto the  $L_2$ -norm ball with



radius  $2C'$  for some constant  $C'$ . A smaller value of  $C'$  equals decreasing the learning rates while improving the numerical stability. We set  $C'$  to be  $10^5$  and  $10^3$  for experiments with Poisson and Negative Binomial likelihoods, respectively.

**Lasso penalty  $\lambda$ .** For the lasso penalty  $\lambda = c_1 \sqrt{\log p/n}$ , one can use cross-validation to tune the lasso penalty for optimal log-likelihood. However, because the estimation results are insensitive to the choice of this penalty, we simply set  $c_1$  to be 0.02 and 0.01 for experiments with Poisson and Negative Binomial likelihoods, respectively.


**The number of factors  $r$ .** For the number of factors  $r$ , the joint-likelihood-based information criterion (JIC) proposed by [Chen and Li \(2022\)](#) can be utilized to select a proper number of latent factors. The JIC value is the sum of deviance and a penalty on model complexity:

$$\begin{aligned} \text{JIC}(\widehat{\Theta}^{(r)}) &= \text{deviance} + \nu(n, p, d + r) \\ &= -2 \sum_{i \in [n], j \in [p]} \log p(y_{ij} | \widehat{\theta}_{ij}^{(r)}) + c_{\text{JIC}} \cdot \frac{(d + r) \log(n \wedge p)}{n \wedge p}, \end{aligned}$$

where  $\widehat{\Theta}^{(r)}$  is the estimated natural parameter matrix with  $r$  latent factors and  $d$  observed covariates, and  $c_{\text{JIC}} > 0$  is a universal constant set to be one in all our simulations. As shown by [Chen and Li \(2022\)](#), minimizing the empirical JIC yields a consistent estimate for the number of factors in generalized linear factor models. As an illustration, we compute the values of JIC at different numbers of factors on simulated datasets and visualize them in [Figure F1](#). When the unmeasured confounding effects are strong, the default choice of  $c_{\text{JIC}} = 1$  gives reasonable estimates for the number of factors under both Poisson and Negative Binomial likelihoods. Because the complexity term is a linear function in  $r$ , one can also inspect the increment of log-likelihood compared to the increment of the complexity term as a function of  $r$ , as shown in the right panel of [Figure F1](#). For real-world datasets, this can help to adjust the penalty level  $c_{\text{JIC}}$  to select a suitable value of  $r$  to achieve a sufficient reduction of negative log-likelihood while avoiding overfitting.

**Debiasing parameters  $(\tau_n, \lambda_n)$ .** For inference, two parameters  $(\tau_n, \lambda_n)$  are to be specified. However, the parameter  $\tau_n$  is less important because as long as both the covariate  $\mathbf{x}_i$  and the projection direction  $\mathbf{u}$  are bounded in  $L_\infty$ -norm, the second constraint in Equation (4.4) will always be satisfied. For this reason, we can ignore the second constraint and solve the relaxed optimization problem, similar to the implementation of [Cai et al. \(2021\)](#). Therefore, one only needs to determine the parameter  $\lambda_n = c_2 \sqrt{\log(n)/n}$ .

To address this, [Cai et al. \(2021\)](#) only implemented a single value for  $c_2$ . However, we propose a more effective heuristic method to guide the selection of  $c_2$ . Specifically, we enumerate different values of  $c_2 \in \{0.001, 0.002, \dots, 0.01, 0.02, \dots, 0.1, 0.2, \dots, 1\}$  and compute the median and median absolute deviation (MAD) of the corresponding empirical  $z$ -statistics. We then generate the scree plot of the two summarized statistics. As shown in [Figure F2](#), as  $\lambda_n$  increases, both the median and the MAD of the empirical null distribution change. Specifically, as  $\lambda_n$  increases, the median decreases, while the MAD increases and



**Figure F1:** The left panel shows the deviance and the complexity penalty  $\nu$  at different numbers of factors  $r$ . The JIC is the sum of deviance and  $\nu$ . The right panel shows the decrement of the deviance and the complexity at different numbers of factors  $r$ . The values are computed from one simulation in Section 5 with  $n = 100$  and  $r^* = 2$  underlying factors.

then decreases. Therefore, when  $\lambda_n$  is too small, the empirical null distribution concentrates around 0, and the resulting tests will be conservative. On the other hand, when  $\lambda_n$  is too large, the tests will be anti-conservative. Therefore, a reasonable choice for  $\lambda_n$  is such that the absolute value of the median is not too large while the MAD of the corresponding test statistics is near one.

For simulations with Poisson likelihood, according to the scree plot, the adaptive choice of the value  $c_2$  would be the largest value that makes the median deviate from 0 by no more than a threshold of 0.1. Analogously, we set the median deviation threshold to be 0.025 for the Negative Binomial simulations. Note that any value below the selected  $\lambda_n$  also provides valid inference results but with lower power.

## F.4 Negative binomial likelihood with non-canonical link


While theoretically nice, the canonical link function for Negative Binomial distributions (NB-C) is not recommended in general because its natural parameter value is always negative, but linear predictors ought to be unbounded in general. Numerical instability may occur in the boundary of the natural parameters. Furthermore, the NB-C model is sensitive to the initial values and may converge to a local solution.

The common choice of a link function for generalized linear models with Negative Binomial likelihood is the log link (Agresti, 2015). Below, we show how to incorporate non-canonical link into our framework.

For Negative Binomial distribution, recall that  $\phi$  is a parameter that represents the number of failures as in Table F1. Define the Negative Binomial canonical link  $A'^{-1}$  and log link  $L^{-1}$ , such that  $A'(\theta) = \phi e^\theta / (1 - e^\theta)$  and  $L(\xi) = e^\xi$ .

Let  $\theta$  and  $\xi$  be the natural parameter and its representation under the log link; namely, the mean  $\mu$  can be obtained from them through the corresponding link functions:

$$\mu = A'(\theta) = L(\xi).$$



**Figure F2:** The median and MAD of the  $z$ -statistics as a function of the regularization parameter  $\lambda_n$  computed from one simulation in Section 5.1 with  $n = 100$  and  $r = 2$ . The shaded region indicates feasible values of  $\lambda_n$ , for which the absolute values of the medians of the corresponding test statistics are less than 0.1.

This gives rise to the transformation equations:

$$e^\xi = \phi \frac{e^\theta}{1 - e^\theta}, \quad e^\theta = \frac{e^\xi}{\phi + e^\xi},$$

and

$$\theta = A'^{-1}(L(\xi)) = \log \frac{e^\xi}{\phi + e^\xi}. \quad (\text{F.1})$$

Note that the negative log-likelihood is given by

$$\begin{aligned} l(\xi) &:= -y \cdot (A')^{-1}(L(\xi)) + A((A')^{-1}(L(\xi))) \\ &= -y \log \frac{e^\xi}{\phi + e^\xi} - \phi \log \frac{\phi}{\phi + e^\xi}, \end{aligned}$$

which has gradient and hessian:

$$\begin{aligned} \frac{\partial l}{\partial \xi} &= \frac{\partial l}{\partial \theta} \frac{\partial \theta}{\partial \xi} = -(y - A'(\theta)) \frac{\phi}{\phi + e^\xi} \\ \frac{\partial^2 l}{\partial \xi^2} &= \frac{\partial^2 l}{\partial \theta^2} \frac{\partial \theta}{\partial \xi} + \frac{\partial l}{\partial \theta} \frac{\partial^2 \theta}{\partial \xi^2} \\ &= A''(\theta) \left( \frac{\partial \theta}{\partial \xi} \right)^2 + \frac{\partial l}{\partial \theta} \frac{\partial^2 \theta}{\partial \xi^2} \end{aligned} \quad (\text{F.2})$$

$$\approx \frac{\phi e^\xi}{\phi + e^\xi} \quad (\text{F.3})$$

where the last line is because the conditional expectation on  $y - A'(\theta)$  given  $\theta$  is zero,  $\mathbb{E}[\partial l / \partial \theta \mid \theta] = 0$ , so that the second term is ignorable. The latter approximation approach is also used in classic GLM to derive the asymptotic variance of the estimates.

For  $n$  i.i.d. samples  $(\mathbf{x}_i, \mathbf{z}_i, \mathbf{y}_i)$ , the linear predictor reads that  $\Xi = \mathbf{X}\mathbf{B}^\top + \mathbf{Z}\mathbf{\Gamma}^\top$  when using the log link. Based on the relationship (F.1), we can perform estimation and inference for the log link function, as described below.

For estimation, the objective function (2.3) now becomes:

$$l(\Xi) = l(\mathbf{\Gamma}, \mathbf{Z}, \mathbf{B}) = -\frac{1}{n} \sum_{i \in [n]} \sum_{j \in [p]} (y_{ij} A'^{-1}(L(\xi_{ij})) - L(\xi_{ij})).$$

Even though the new objective is now nonconvex in the parameter  $\Xi$ , the alternative maximization algorithm Algorithm F.3 is still applicable to it, because the gradient can be computed based on (F.2). If we initialize  $\hat{\mathbf{F}}$  from GLM estimates and treat it as fixed, then solving optimization problem (3.2) reduces to a nonconvex matrix factorization problem. Under this setting, there is a rich literature on establishing the estimation error for  $\mathbf{W}^*$  and  $\mathbf{\Gamma}^*$  given that the initial value is close to the truth; see Wang et al. (2017b); Lin et al. (2024) among the others. In other words, we may also obtain error bounds on  $\|\Xi^* - \hat{\Xi}\|_{\text{F}}^2$  by imposing additional conditions.

For inference, we simply apply the chain rule and (F.2)-(F.3) to rewrite (4.1) as:

$$\hat{b}_{j1}^{\text{de}} = \hat{b}_{j1} + \mathbf{u}^\top \frac{1}{n} \sum_{i=1}^n \mathbf{x}_i (\mathbf{y}_i - A'(\hat{\boldsymbol{\theta}}_i))^\top \text{diag} \left( \left\{ \left. \frac{\partial \theta}{\partial \xi} \right|_{\xi = \hat{\xi}_{ij}} \right\}_{j \in [p]} \right) \mathbf{v}_i,$$

with

$$\begin{aligned} \mathbf{v}_i &= \omega_i \text{diag} \left( \left\{ \left. \frac{\partial \theta}{\partial \xi} \right|_{\xi = \hat{\xi}_{ij}} \right\}_{j \in [p]} \right)^{-2} \text{diag}(A''(\hat{\boldsymbol{\theta}}_i))^{-1} \mathcal{P}_{\hat{\mathbf{F}}}^\perp \mathbf{e}_j, \\ \omega_i &= \mathbb{E} \left[ \frac{\partial^2 l}{\partial \xi^2} \mid \xi = \hat{\xi}_{ij} \right] = A''(\hat{\theta}_{ij}) \left( \left. \frac{\partial \theta}{\partial \xi} \right|_{\xi = \hat{\xi}_{ij}} \right)^2 = \frac{\phi e^\xi}{\phi + e^\xi}. \end{aligned}$$

Because when  $\mathcal{R}_{\mathcal{C}}$  is bounded, the derivative function  $\partial \theta / \partial \xi$  is Lipschitz continuous, the estimation error of it:

$$\sum_{j=1}^p (\partial \theta / \partial \xi |_{\xi = \hat{\xi}_{ij}^*} - \partial \theta / \partial \xi |_{\xi = \hat{\xi}_{ij}})^2 \lesssim \|\boldsymbol{\xi}_i^* - \hat{\boldsymbol{\xi}}_i\|_2^2$$

can be bounded if  $\boldsymbol{\xi}_i^*$  can be well estimated. Similarly, the estimation error of  $\boldsymbol{\Theta}^*$  can be controlled because  $\theta_{ij}^*$  is a Lipschitz continuous function of  $\xi_{ij}^*$ . Thus, Theorem 6 also applies if

$$\|\Xi^* - \hat{\Xi}\|_{\text{F}}^2 \lesssim \sqrt{n \vee p} \quad (\text{F.4})$$

$$\max_{1 \leq j \leq p} \frac{1}{\sqrt{n}} \|\mathbf{Z}^* \boldsymbol{\Gamma}_j^{*\top} - \widehat{\mathbf{Z}} \widehat{\boldsymbol{\Gamma}}_j^\top\|_{\text{F}} \lesssim \frac{1}{\sqrt{n \wedge p}} \quad (\text{F.5})$$

$$\|\widehat{\mathbf{B}} - \mathbf{B}^*\|_{1,1} \lesssim \frac{\sqrt{sd}}{\sqrt{n \wedge p}} \quad (\text{F.6})$$

$$\|\widehat{\mathbf{B}} - \mathbf{B}^*\|_{\text{F}} \lesssim \frac{1}{\sqrt{n \wedge p}}, \quad (\text{F.7})$$

where (F.4) requires an analysis tool from nonconvex matrix factorization, (F.5) is a direct consequence from (F.4) similar to Corollary 4, and (F.6)-(F.7) requires non-asymptotic analysis as in the proof of Theorem 5 but for nonconvex objectives instead.

## G Extra experiment results

### G.1 Efficiency loss of sample splitting

To evaluate the efficiency loss caused by sample splitting described in Algorithm E.2, we conduct the experiments with different splitting proportions and compare their results. To apply Algorithm E.2, we split the  $p$  genes into 2 groups with equal sizes, so that  $I_1 = \{1, \dots, p/2\}$  and  $I_2 = \{p/2 + 1, \dots, p\}$ . For each of the groups  $I$ , the optimization is jointly conducted based on  $\mathbf{X}$ ,  $\mathbf{Y}_{I^c}$  and  $\mathcal{D}_2$ , and the inference is conducted for genes in  $I$ . As summarized in Table G2, the performance on Type-I error and FDP control is similar across different splitting ratios. However, the power and precision are affected when the ratio of observations reserved for inference is too small. This suggests that one should leave more observations to conduct the debias step. Lastly, we see similar performance even without sample splitting, suggesting that the validity of the inferential procedure could be true even without sample splitting.


ratio split	type-I error	FDP	power	precision
0.2	0.050	0.200	0.454	0.610
0.4	0.049	0.193	0.755	0.920
0.6	0.050	0.191	0.901	1.000
0.8	0.051	0.195	0.963	1.000
no splitting	0.051	0.219	0.987	1.000

**Table G2:** Performance with varying ratios of observations reserved for inference, under the same data setup in Section 5.1 with  $n = 250$  and  $r = 2$ . The reported values are medians summarized over 100 simulated datasets.

## G.2 Information about lupus data

Cell type	Number of samples $n$	Number of genes $p$	Proportion of non-zeros
T4	256	1255	0.398
cM	256	1208	0.434
B	254	1269	0.417
T8	256	1281	0.471
NK	256	1178	0.385

**Table G3:** Summary statistics of the preprocessed lupus datasets in each cell type. The last column represents the proportion of non-zero count in the gene expression matrix.




**Figure G3:** Histograms of expressions of 5 genes on the T4 cell type. The first row shows the raw pseudo-bulk counts and the second row shows the counts after library size normalization and  $\log_{1p}$  transformation, which is used for CATE. Due to the sparsity of the gene expressions, some genes are not distributed like normal after transformation.


### G.3 Extra results on lupus datasets

$r$	mean	median	mad	num_sig	deviance	JIC
1	0.348	-0.018	1.551	302	3.578	3.600
2	0.361	-0.040	1.539	313	3.565	3.592
3	0.141	-0.001	1.145	57	3.553	3.586
4	0.140	0.063	1.087	37	3.546	3.584
5	0.130	0.043	1.067	33	3.539	3.582
6	0.140	0.051	1.084	39	3.532	3.581
7	0.128	0.057	1.033	22	3.526	3.580
8	0.139	0.069	1.044	18	3.521	3.580
9	0.143	0.073	1.032	18	3.516	3.581
10	0.155	0.095	1.038	20	3.513	3.583

**Table G4:** The summary of the  $z$ -statistics and model fitness for varying number of latent factors  $r$ . The metrics include the mean, median, median absolute deviation (mad), and the total number of significant genes of  $q$ -value less than 0.2. The last two columns show the deviance (2 times the negative log-likelihood) and the JIC model selection criteria (3.1).




**Figure G4:** The right panel shows the deviance and the complexity penalty  $\nu$  at different numbers of factors  $r$ , computed on the T4 cell type of the Lupus dataset. The JIC is the sum of the deviance and  $\nu/4$ . The right panel shows the decrement of the deviance and the complexity at different numbers of factors  $r$ .




**Figure G5:** The median and MAD of the  $z$ -statistics as a function of the regularization parameter  $\lambda_n$  computed from the T4 cell type of the Lupus dataset. The shaded region indicates feasible values of  $\lambda_n$ , for which the MADs of the corresponding test statistics are less than 1.13.






**Figure G6:** Histograms of lupus  $z$ -statistics of different methods on cM, B, T8 and NK cell types.



**Figure G7:** Histograms of lupus  $z$ -statistics of CATE on T4, cM, B, and T8 cell types, when restricted to the top 250 highly variable genes. The preprocessing procedure is as described in Section 6, but with genes expressed less than 5 subjects excluded. The result on the NK cell type is not included because the fitting of CATE fails due to sparsity of the gene expressions.



**Figure G8:** The treemap plot produced by *rrvgo* (Sayols, 2023) of GO enrichment analysis results on **(a)** significant genes by the GLM-oracle method; **(b)** significant genes by both the GLM-oracle and GCATE methods; and **(c)** significant genes by the CATE-mad method but not the GLM-oracle method.

## References

- Agresti, A. (2015). *Foundations of linear and generalized linear models*. John Wiley & Sons.
- Bandeira, A. S. and van Handel, R. (2016). Sharp nonasymptotic bounds on the norm of random matrices with independent entries. *The Annals of Probability*, 44(4):2479–2506.
- Bing, X., Cheng, W., Feng, H., and Ning, Y. (2023). Inference in high-dimensional multivariate response regression with hidden variables. *Journal of the American Statistical Association*, pages 1–12.
- Cai, T. T., Guo, Z., and Ma, R. (2021). Statistical inference for high-dimensional generalized linear models with binary outcomes. *Journal of the American Statistical Association*, pages 1–14.
- Chen, Y. and Li, X. (2022). Determining the number of factors in high-dimensional generalized latent factor models. *Biometrika*, 109(3):769–782.
- Chernozhukov, V., Nekipelov, D., Semenova, V., and Syrgkanis, V. (2018). Plug-in regularized estimation of high-dimensional parameters in nonlinear semiparametric models. *arXiv preprint arXiv:1806.04823*.
- Foster, D. J. and Syrgkanis, V. (2023). Orthogonal statistical learning. *The Annals of Statistics*, 51(3):879–908.
- Lin, K. Z., Lei, J., and Roeder, K. (2021). Exponential-family embedding with application to cell developmental trajectories for single-cell RNA-seq data. *Journal of the American Statistical Association*, 116(534):457–470.
- Lin, K. Z., Qiu, Y., and Roeder, K. (2024). esvd-de: Cohort-wide differential expression in single-cell rna-seq data using exponential-family embeddings. *BMC bioinformatics*, 25(1):113.
- Sayols, S. (2023). rrvgo: a bioconductor package for interpreting lists of gene ontology terms. *Micropublication Biology*, 2023.
- Vershynin, R. (2018). *High-dimensional probability: An introduction with applications in data science*, volume 47. Cambridge university press.
- Wainwright, M. J. (2019). *High-dimensional statistics: A non-asymptotic viewpoint*, volume 48. Cambridge university press.
- Wang, J., Zhao, Q., Hastie, T., and Owen, A. B. (2017a). Confounder adjustment in multiple hypothesis testing. *The Annals of Statistics*, 45(5):1863.
- Wang, L., Zhang, X., and Gu, Q. (2017b). A unified computational and statistical framework for nonconvex low-rank matrix estimation. In *Artificial Intelligence and Statistics*, pages 981–990. PMLR.