

THE HEAT EQUATION WITH THE L^p PRIMITIVE INTEGRAL

ERIK TALVILA

ABSTRACT. For each $1 \leq p < \infty$ a Banach space of integrable Schwartz distributions is defined by taking the distributional derivative of all functions in $L^p(\mathbb{R})$. Such distributions can be integrated when multiplied by a function that is the integral of a function in $L^q(\mathbb{R})$, where q is the conjugate exponent of p . The heat equation on the real line is solved in this space of distributions. The initial data is taken to be the distributional derivative of an $L^p(\mathbb{R})$ function. The solutions are shown to be smooth functions. Initial conditions are taken on in norm. Sharp estimates of solutions are obtained and a uniqueness theorem is proved.

1. INTRODUCTION

In this paper we study solutions of the heat equation on the real line with initial value data that is the distributional derivative of an L^p function ($1 \leq p < \infty$). The solution is given by convolution of the Gauss–Weierstrass heat kernel with the initial data. This produces smooth solutions to the heat equation. We introduce a norm that is isometric to the L^p norm and show that solutions tend to the initial condition within this norm as $t \rightarrow 0^+$. Since the initial data is the distributional derivative of an L^p function it can fail to have pointwise values everywhere, can be zero almost everywhere, and can be the difference of translated Dirac distributions. And yet, the initial conditions are taken on in the strong (norm) sense rather than in the weak (distributional) sense.

For $u: \mathbb{R} \times (0, \infty) \rightarrow \mathbb{R}$ write $u_t(x) = u(x, t)$.

The classical problem of the heat equation on the real line is, given a function $F \in L^p$ for some $1 \leq p \leq \infty$, find a function $u: \mathbb{R} \times (0, \infty) \rightarrow \mathbb{R}$ such that $u_t \in C^2(\mathbb{R})$ for each $t > 0$, $u(x, \cdot) \in C^1((0, \infty))$ for each $x \in \mathbb{R}$ and

$$(1.1) \quad \frac{\partial^2 u(x, t)}{\partial x^2} - \frac{\partial u(x, t)}{\partial t} = 0 \text{ for each } (x, t) \in \mathbb{R} \times (0, \infty)$$

$$(1.2) \quad \lim_{t \rightarrow 0^+} \|u_t - F\|_p = 0.$$

If $p = \infty$ then F is also assumed to be continuous. Under suitable growth conditions on u the unique solution is given by $u(x, t) = f * \Theta_t(x)$, where the Gauss–Weierstrass heat kernel is $\Theta_t(x) = \exp(-x^2/(4t))/(2\sqrt{\pi t})$.

Instead of the initial data being an L^p function we take it to be the distributional derivative of such a function. A well-defined integration process for these functions and distributions appears in [16]. The space of derivatives of L^p functions is denoted L^p . It

Date: Preprint September 14, 2023. To appear in *Generalized integrals and applications* (World Scientific; Eds. T. P. Bercera, H. Kalita, A. Croitoru, B. Hazarika).

2020 Mathematics Subject Classification. Primary 35K05, 46F10, 46G12; Secondary 46E30.

Key words and phrases. Heat equation, Lebesgue space, Lebesgue space primitive integral, tempered distribution, Schwartz distribution, generalised function, Banach space, convolution, primitive, integral.

is a Banach space isometrically isomorphic to L^p . In L^p many properties of L^p functions continue to hold, such as convolution with functions in Lebesgue spaces, a type of Hölder inequality, continuity of translations, etc. Some of the distributions in L^p can be identified with signed measures, some can be identified with smooth functions and some have no pointwise values.

See [3] and [19] for classical results on the heat equation, including extensive bibliographies. For initial data in L^p spaces see [9] and [10]. Distributional solutions of the heat equation are considered in [4] and [14], where the initial data is taken on in the distributional sense. For initial data in the Sobolev space H^s , see [12]. Another paper studying the heat equation with distributions in Banach spaces is [17], in which the initial data is taken to be the distributional derivative of a continuous function.

For $1 \leq p < \infty$, the Lebesgue space on the real line is L^p , which is the set of measurable functions $f: \mathbb{R} \rightarrow \mathbb{R}$ such that $\int_{-\infty}^{\infty} |f(x)|^p dx < \infty$. To distinguish between other types of integrals introduced later, Lebesgue integrals will always explicitly show the integration variable and differential as above.

The Schwartz space (test functions), of rapidly decreasing smooth functions, is denoted \mathcal{S} and the tempered distributions \mathcal{S}' . See, for example, [7] or [8].

The outline of the paper is as follows.

The heat equation is considered in L^p in Section 2. With initial data $f \in L^p$ it is proved that convolution with f and the Gauss–Weierstrass heat kernel is a smooth function in L^s for each $p \leq s < \infty$ but need not be in L^s if $1 \leq s < p$ (Theorem 2.1). Sharp estimates are given in $\|\cdot\|'_r$, with $1/p + 1/q = 1 + 1/r$. This convolution gives the unique solution to the heat equation (Corollary 2.3). Existence is proven using a type of Hardy space. In Proposition 2.4 continuity is proved with respect to the initial conditions.

In Section 3 various sharp L^p and pointwise estimates are given. It is shown that these solutions cannot be of one sign, have limit 0 at infinity, and have a vanishing integral over the real line.

An appendix contains a lemma on pointwise differentiation of convolutions and proof of some sharp estimates for solutions of (1.1) and (1.2).

2. SOLUTIONS OF THE HEAT EQUATION IN L^p

For $1 \leq p < \infty$ let $L^p = \{f \in \mathcal{S}' \mid f = F'$ for some $F \in L^p\}$. As explained in [16], if $f \in L^p$ then it has a unique primitive in L^p . In this case, $\|f\|'_p = \|F\|_p$ and this makes L^p and L^p isometrically isomorphic Banach spaces. As in [16], define $I^p = \{G: \mathbb{R} \rightarrow \mathbb{R} \mid G(x) = \int_0^x g(t) dt \text{ for some } g \in L^p\}$

If $f \in L^p$ then $f * \Theta_t$ is a smooth solution of the heat equation. The initial value f is taken on in the $\|\cdot\|'_p$ norm and norm boundedness ensures uniqueness. Solutions are shown to be in L^r for each $p \leq r < \infty$ and sharp norm estimates are given using the results of Theorem 4.1 in the Appendix. If a solution v_t of the heat equation has bounded norms $\|v_t\|'_p$ for $t > 0$ then there exists a distribution $f \in L^p$ such that $v_t = \Theta_t * f$. This is proved using an analogue of a harmonic Hardy space. There is continuity with respect to initial conditions. The example of initial data that is the difference to two Dirac measures is considered.

Theorem 2.1. *Let $1 \leq p < \infty$. Let $f \in L^p$ with primitive $F \in L^p$. Let $t > 0$ and define $v(x, t) = v_t(x) = f * \Theta_t(x)$.*

(a) For each $x \in \mathbb{R}$ we have

$$v_t(x) = \int_{-\infty}^{\infty} f(y)\Theta_t(x-y) dy = F * \Theta'_t(x) = \Theta'_t * F(x) = (F * \Theta_t)'(x).$$

The function v_t is uniformly continuous on \mathbb{R} .

(b) For each $n \geq 1$ we have $v_t^{(n)}(x) = f * \Theta_t^{(n)}(x)$ and $\partial^n v(x, t)/\partial t^n = f * \partial^n \Theta(x, t)/\partial t^n$. Hence, $v \in C^\infty(\mathbb{R}) \times C^\infty((0, \infty))$.

(c) If $p \leq s < \infty$ then $f * \Theta_t \in L^s$.

(d) Let $q, r \in [1, \infty]$ such that $1/p + 1/q = 1 + 1/r$. There is a constant $K_{p,q}$ such that $\|f * \Theta_t\|'_r \leq K_{p,q} \|f\|'_p t^{-(1-1/q)/2}$ for all $t > 0$. The estimate is sharp in the sense that if $\psi : (0, \infty) \rightarrow (0, \infty)$ such that $\psi(t) = o(t^{-(1-1/q)/2})$ as $t \rightarrow 0^+$ or $t \rightarrow \infty$ then there is $g \in L^p$ such that $\|g * \Theta_t\|'_r / \psi(t)$ is not bounded. The constant $K_{p,q}$ cannot be replaced by any smaller number. In particular, $\|f * \Theta_t\|'_p \leq \|f\|'_p$ is a sharp estimate.

(e) If $1 \leq s < p$ then $f * \Theta_t$ need not be in L^s .

Note that if $1/p + 1/q = 1 + 1/r$ then, for suitable q , r can take on any of the values in $[p, \infty]$, with $r = p$ corresponding to $q = 1$ and $r = \infty$ corresponding to p and q being conjugates. And, for suitable r , q can take on any of the values in $[1, p/(p-1)]$, with $q = 1$ corresponding to $r = p$ and $q = p/(p-1)$ corresponding to $r = \infty$. If $p = 1$ then q can take on all values in $[1, \infty]$.

The constants $K_{p,q}$ in Theorem 4.1(b) and part (d) above are the same. When $r = p$ and $q = 1$ the inequality in part (d) reads $\|f * \Theta_t\|'_p \leq \|f\|'_p$.

Proof. (a) Since Θ_t and its derivatives are in each L^q space, the equalities follow by Theorem 5.1 in [16], Lemma 4.2 in the Appendix, and usual properties of convolution for Lebesgue integrals. To prove uniform continuity, note that if q is the conjugate of p then

$$|v_t(x) - v_t(y)| = |F * [\Theta'_t(x) - \Theta'_t(y)]| \leq \|F\|_p \|\Theta'_t(x - \cdot) - \Theta'_t(y - \cdot)\|_q \rightarrow 0 \text{ as } y \rightarrow x.$$

The last line uses the Hölder inequality and continuity in the L^q norm. (When $q = \infty$ this requires boundedness and uniform continuity of Θ'_t .)

(b) Differentiation in x follows from Lemma 4.2. For differentiation in t , use part (a) to write

$$v_t(x) = F * \Theta'_t(x) = -\frac{1}{4\sqrt{\pi} t^{3/2}} \int_{-\infty}^{\infty} F(x-y) y e^{-y^2/(4t)} dy.$$

Then

$$\frac{\partial v(x, t)}{\partial t} = \frac{3}{8\sqrt{\pi} t^{5/2}} \int_{-\infty}^{\infty} F(x-y) y e^{-y^2/(4t)} dy - \frac{1}{16\sqrt{\pi} t^{7/2}} \int_{-\infty}^{\infty} F(x-y) y^3 e^{-y^2/(4t)} dy.$$

Differentiation under the integral sign is justified by dominated convergence since, if $t \leq t_0$, then $|y^3 e^{-y^2/(4t)}| \leq |y|^3 e^{-y^2/(4t_0)}$.

(c), (d) From Theorems 3.6 and 5.1 in [16] we have

$$\|f * \Theta_t\|'_r = \|F * \Theta'_t\|'_r = \|(F * \Theta_t)'\|'_r = \|F * \Theta_t\|_r \leq C_{p,q} \|F\|_p \|\Theta_t\|_q.$$

The rest of the proof is identical to Theorem 4.1(b).

(e) Since $v_t = (F * \Theta_t)' \in L^s$ if and only if $F * \Theta_t \in L^s$, this result is equivalent to Theorem 4.1(c). \square

To prove uniqueness for the initial value problem we use a classical theorem, which follows from Theorem 6.1 in [19].

Theorem 2.2. *Let $u: \mathbb{R} \times (0, \infty) \rightarrow \mathbb{R}$ such that $u_t \in C^2(\mathbb{R})$ for each $t > 0$, $u(x, \cdot) \in C^1((0, \infty))$ for each $x \in \mathbb{R}$, $u \in C(\mathbb{R} \times [0, \infty))$, $\partial^2 u(x, t)/\partial x^2 - \partial u(x, t)/\partial t = 0$ for $(x, t) \in \mathbb{R} \times (0, \infty)$, u is bounded, $u(x, 0) = f(x)$ for a bounded continuous function $f: \mathbb{R} \rightarrow \mathbb{R}$. Then the unique solution is given by $u_t(x) = f * \Theta_t(x)$.*

Corollary 2.3. *Let $1 \leq p < \infty$.*

(a) *Let $f \in L^p$. The unique solution of the problem to find $v: \mathbb{R} \times (0, \infty) \rightarrow \mathbb{R}$ such that*

$$(2.1) \quad v_t \in L^p \text{ for each } t > 0$$

$$(2.2) \quad v_t \in C^2(\mathbb{R}) \text{ for each } t > 0, v(x, \cdot) \in C^1((0, \infty)) \text{ for each } x \in \mathbb{R}$$

$$(2.3) \quad \frac{\partial^2 v(x, t)}{\partial x^2} - \frac{\partial v(x, t)}{\partial t} = 0 \text{ for each } (x, t) \in \mathbb{R} \times (0, \infty)$$

$$(2.4) \quad \|v_t\|'_p \text{ is bounded}$$

$$(2.5) \quad \lim_{t \rightarrow 0^+} \|v_t - f\|'_p = 0$$

*is given by $v_t = f * \Theta_t$.*

(b) *If $f \in L^p$ and $v_t = f * \Theta_t$ then $\lim_{t \rightarrow 0^+} \|v_t\|'_p = \|f\|'_p$.*

(c) *Let $1 < p < \infty$ and let h^p be the set of functions satisfying (2.1), (2.2), (2.3) and (2.4). If $v \in h^p$ define $\|v\|_p^{h'} = \sup_{t > 0} \|v_t\|'_p$. Let $T: L^p \rightarrow h^p$ be given by $T[f] = \Theta_t * f$. Then T is a linear isometric isomorphism.*

(d) *Let $1 < p < \infty$. If (2.1), (2.2), (2.3) and (2.4) hold for some $v \in h^p$ then there is a unique distribution $f \in L^p$ such that $v_t = \Theta_t * f$ and (2.5) holds.*

The proof uses methods from [10, Theorem 9.2, p. 195] and [1, pp. 115-120]. Part (c) implies h^p is a Banach space.

Proof. (a) By part (b) of the theorem we can differentiate under the integral sign and this shows $v_t = f * \Theta_t$ is a smooth solution of the heat equation. By part (c) of the theorem, $v_t \in L^p$. To see that the initial conditions are taken on in the $\|\cdot\|'_p$ norm, note that $\|v_t - f\|'_p = \|F * \Theta_t - F\|_p$. Since $\int_{-\infty}^{\infty} \Theta_t(y) dy = 1$ we have

$$\|F * \Theta_t - F\|_p = \left(\int_{-\infty}^{\infty} \left| \int_{-\infty}^{\infty} [F(x-y) - F(x)] \Theta_t(y) dy \right|^p dx \right)^{1/p}.$$

By the Minkowski inequality for integrals, for example [7, p. 194],

$$\lim_{t \rightarrow 0^+} \|F * \Theta_t - F\|_p \leq \lim_{t \rightarrow 0^+} \int_{-\infty}^{\infty} \|F(\cdot - y) - F(\cdot)\|_p \Theta_t(y) dy = \lim_{y \rightarrow 0} \|F(\cdot - y) - F(\cdot)\|_p = 0.$$

The last line follows by the fact that Θ_t is an approximate identity (delta sequence) and by continuity in the L^p norm.

To prove the solution is unique, suppose there were two solutions v and w . Let $z = v - w$. Then z satisfies (2.1), (2.2), (2.3) and (2.4). And,

$$\|z_t\|'_p \leq \|v_t - f\|'_p + \|w_t - f\|'_p \rightarrow 0 \text{ as } t \rightarrow 0^+.$$

Now define $K : \mathbb{R} \rightarrow \mathbb{R}$ by $K(x) = 1 - |x|$ for $|x| \leq 1$ and $K(x) = 0$ for $|x| \geq 1$, and let $K_h(x) = (1/h)K(x/h)$ for $h > 0$. Then K_h is absolutely continuous with compact support. Note that $\int_{-\infty}^{\infty} K(y) dy = \int_{-\infty}^{\infty} K_h(y) dy = 1$. We have

$$K'_h(x) = \begin{cases} 0, & |x| > h \\ -\operatorname{sgn}(x)/h^2, & |x| < h. \end{cases}$$

Now define $\psi_h(x, t) = K_h * z_t(x) = \int_{-h}^h K_h(y) z_t(x - y) dy$. Since $\partial z_t(x)/\partial t$ and $z_t''(x)$ are continuous on compact intervals in $\mathbb{R} \times (0, \infty)$ we can use dominated convergence to differentiate under the integral sign and this shows ψ_h is a solution of the heat equation. By usual properties of convolutions, $\psi_h \in C^2(\mathbb{R}) \times C^1((0, \infty))$. From the Hölder inequality [16, Theorem 3.6], $|\psi_h(x, t)| \leq \|z_t\|'_p \|K'_h\|_q$, where q is the conjugate of p . And, $\|K'_h\|_q = 2^{1-1/p}/h^{1+1/p}$. It follows that ψ_h is bounded on $\mathbb{R} \times (0, \infty)$ and $\lim_{t \rightarrow 0^+} \psi_h(x, t) = 0$. Define $\psi_h(x, 0) = 0$. The above estimate now shows $\psi_h \in C(\mathbb{R} \times [0, \infty))$. By Theorem 2.2, $\psi_h = 0$. Since z_t is continuous, for $(x, t) \in \mathbb{R} \times (0, \infty)$ we get $\lim_{h \rightarrow 0^+} \psi_h(x, t) = 0 = \lim_{h \rightarrow 0^+} K_h * z_t(x) = z_t(x) \lim_{h \rightarrow 0^+} \int_{-\infty}^{\infty} K_h(y) dy = z_t(x)$.

(b) This follows from (2.5) and the triangle inequality.

(c) Clearly, T is linear.

Show T is an isometry. From Theorem 2.1(d), $\|T[f]\|_p^h \leq \sup_{t>0} \|f\|'_p = \|f\|'_p$. And, if $f \in L^p$ then

$$\|f\|'_p \leq \|f - f * \Theta_t\|'_p + \|f * \Theta_t\|'_p \leq \|f - f * \Theta_t\|'_p + \|T[f]\|_p^h.$$

Letting $t \rightarrow 0^+$ and using (2.5) shows $\|f\|'_p \leq \|T[f]\|_p^h$. Hence, T is an isometry and necessarily one-to-one.

Now show T is onto h'^p . Let $v \in h'^p$ and find $f \in L^p$ such that $T[f] = v$. For $h > 0$ use notation from the proof of (a) and define $w_h = K_h * v_t$. Let $\delta > 0$. Then the function $(x, t) \mapsto w_h(x, t + \delta)$ satisfies (2.1), (2.2), (2.3) and (2.4) and is continuous and bounded on $\mathbb{R} \times [0, \infty)$. By Theorem 2.2 we have $w_h(x, t + \delta) = \int_{-\infty}^{\infty} \Theta_t(x - y) w_h(y, \delta) dy$. And, $\|K_h * v_\delta - v_\delta\|'_p \rightarrow 0$ as $h \rightarrow 0^+$ by [16, Theorem 5.3(g)]. (The second sentence in this part of the theorem should read, Let $F \in L^1$.) If q is the conjugate of p ([16, Theorem 3.3(c)]) then $I^{q*} = L^p$ so, by weak* convergence, for each $G \in I^q$ we have $\int_{-\infty}^{\infty} (K_h * v_\delta - v_\delta) G \rightarrow 0$. In particular, take $G = \Theta_t$ and note Definition 3.5 in [16]. Then, also using the continuity of v ,

(2.6)

$$\lim_{h \rightarrow 0^+} w_h(x, t + \delta) = v_{t+\delta}(x) = \lim_{h \rightarrow 0^+} \int_{-\infty}^{\infty} \Theta_t(x - y) w_h(y, \delta) dy = \int_{-\infty}^{\infty} \Theta_t(x - y) v_\delta(y) dy.$$

By definition, $v \in h'^p$ is norm-bounded in $L^p = I^{q*}$ where q is the conjugate of p . Every norm-bounded sequence in a separable Banach space contains a weak* convergent subsequence. Since I^q is separable (since $q < \infty$) there is a sequence δ_i that decreases to 0 and a distribution $f \in L^p$ such that $\int_{-\infty}^{\infty} v_{\delta_i} G \rightarrow \int_{-\infty}^{\infty} f G$ for each $G \in I^q$. In particular, take $G = \Theta_t$. Then $\lim_{i \rightarrow \infty} \int_{-\infty}^{\infty} v_{\delta_i}(y) \Theta_t(x - y) dy = \int_{-\infty}^{\infty} f(y) \Theta_t(x - y) dy = f * \Theta_t(x)$. Replacing δ by δ_i in (2.6) and letting $i \rightarrow \infty$ now shows $v_t = T[f]$.

(d) This follows from (c). □

The following proposition shows continuity with respect to initial conditions.

Proposition 2.4. *Let $1 \leq p < \infty$. Let $f, g \in L^p$.*

(a) *Let $q, r \in [1, \infty]$ such that $1/p + 1/q = 1 + 1/r$. Let $K_{p,q}$ be the constant in the proof of Theorem 2.1. Then*

$$\|f * \Theta_t - g * \Theta_t\|'_r \leq K_{p,q} \|f - g\|'_p t^{-(1-1/q)/2}.$$

*When $r = p, q = 1$, this reads $\|f * \Theta_t - g * \Theta_t\|'_p \leq \|f - g\|'_p$.*

(b) *Let $v, w: \mathbb{R} \times (0, \infty) \rightarrow \mathbb{R}$ such that for each $t > 0$ we have v_t and w_t in L^p . Let $\epsilon > 0$. Suppose $\|v_t - f\|'_p \rightarrow 0$ and $\|w_t - g\|'_p \rightarrow 0$ as $t \rightarrow 0^+$. If $f, g \in L^p$ such that $\|f - g\|'_p < \epsilon$ then for small enough $t > 0$ we have $\|v_t - w_t\|'_p < 2\epsilon$.*

Proof. (a) This follows from Theorem 2.1(d) since $\|f * \Theta_t - g * \Theta_t\|'_r = \|(F - G) * \Theta_t\|_r$, where F and G are the respective primitives of f, g in L^p . (b) This follows from the triangle inequality. \square

We remark the initial conditions are also taken on in the weak (distributional) sense.

Remark 2.5. If $f \in L^p$ with primitive $F \in L^p$ for some $1 \leq p < \infty$ then we can define $v(x, t) = v_t(x) = f * \Theta_t(x)$. As shown in Corollary 2.3, v satisfies the heat equation and the initial condition is taken on in $\|\cdot\|'_p$ as $t \rightarrow 0^+$. Such a solution also takes on the initial condition in the weak distributional sense. To see this let $\phi \in \mathcal{S}$. Then

$$\langle v_t - f, \phi \rangle = \langle (F * \Theta_t - F)', \phi \rangle = -\langle F * \Theta_t - F, \phi' \rangle = -\int_{-\infty}^{\infty} [F * \Theta_t(x) - F(x)] \phi'(x) dx.$$

We know $\|F * \Theta_t - F\|_p \rightarrow 0$ so $\langle F * \Theta_t - F, \psi \rangle \rightarrow 0$ for each $\psi \in L^q$ where q is conjugate to p . In particular, this holds when $\psi = \phi'$. Therefore, the initial condition is taken on in the weak distributional sense.

Several distributions in L^p are given in [16], Section 4., for which (2.5) of course holds but (1.2) does not hold in any Lebesgue space. We now consider the example of the difference of two translated Dirac distributions.

Example 2.6. Denote the Dirac distribution supported at $a \in \mathbb{R}$ by δ_a . Let $F = \chi_{[a,b]}$ then F is in every L^p space. Therefore, $F' = \delta_a - \delta_b$ is in every L^p space. Hence, L^p is not a subset of any of the Lebesgue spaces. Since step functions are dense in L^p , linear combinations of such differences of Dirac distributions are dense in L^p .

Now let $f = \delta_{-a} - \delta_a$ for fixed $a > 0$. Then $v_t(x) = f * \Theta_t(x) = \Theta_t(x + a) - \Theta_t(x - a)$. Note that v is a solution of the heat equation in $\mathbb{R} \times (0, \infty)$. According to Corollary 2.3 the initial condition is taken on as $\|v_t - f\|'_p \rightarrow 0$ as $t \rightarrow 0^+$.

One might wonder in what other sense the initial conditions are realised. We've shown above that they are taken on in the weak distributional sense. For no $1 \leq r \leq \infty$ is $\|v_t - f\|_r$ defined so (1.2) is meaningless. As a signed Borel measure, we have $d\mu(x) = [\Theta_t(x+a) - \Theta_t(x-a)]dx - d\delta_{-a}(x) + d\delta_a(x)$. If the Jordan decomposition of μ is $\mu = \mu_+ - \mu_-$ then μ converges to 0 in variation if $V(\mu) = \mu_+(\mathbb{R}) + \mu_-(\mathbb{R}) \rightarrow 0$ as $t \rightarrow 0^+$. For any measurable set E we have

$$\begin{aligned} V(\mu) &\geq \mu_+(E) + \mu_-(E) \geq |\mu(E)| \\ &= \left| \frac{1}{2\sqrt{\pi t}} \int_E \left(e^{-(x+a)^2/(4t)} - e^{-(x-a)^2/(4t)} \right) dx - \chi_E(-a) + \chi_E(a) \right|. \end{aligned}$$

Take $E = (-\infty, -a)$. Then

$$\begin{aligned} V(\mu) &\geq \frac{1}{2\sqrt{\pi t}} \int_{-\infty}^{-a} \left(e^{-(x+a)^2/(4t)} - e^{-(x-a)^2/(4t)} \right) dx \\ &= \frac{1}{2\sqrt{\pi t}} \int_0^{2a} e^{-x^2/(4t)} dx = \frac{1}{\sqrt{\pi}} \int_0^{a/\sqrt{t}} e^{-y^2} dy \rightarrow \frac{1}{2} \text{ as } t \rightarrow 0^+. \end{aligned}$$

Hence, v_t does not converge to f in variation.

This also shows v_t does not converge to f in the Alexiewicz norm. See [15].

3. ESTIMATES

In the previous section we looked at estimates of $f * \Theta_t$ in L^p . Here we look at estimates in the spaces L^r where $1/p + 1/q = 1 + 1/r$. If $f \in L^p$ then $f * \Theta_t \in L^r$ and Young's inequality gives estimates in terms of $\|f\|'_p$. It is also shown that $f * \Theta_t(x)$ cannot be of one sign, has limit 0 as $|x| \rightarrow \infty$, and has a vanishing integral over the real line. Pointwise estimates are given when f has compact support.

Let $1 \leq q \leq \infty$. Then

$$(3.1) \quad \|\Theta'_t\|_q = \frac{\delta_q}{t^{(2-1/q)/2}} \text{ where } \delta_q = \begin{cases} \frac{1}{\sqrt{\pi}}, & q = 1 \\ \frac{\Gamma^{1/q}((q+1)/2)}{2^{1-1/q}\sqrt{\pi}q^{(1+1/q)/2}}, & 1 < q < \infty \\ \frac{1}{2^{3/2}\sqrt{\pi e}}, & q = \infty. \end{cases}$$

When $q = 1$ the integral is elementary. When $1 < q < \infty$ the calculation uses the definition of the gamma function, $\Gamma(s) = \int_0^\infty e^{-x} x^{s-1} dx$. When $q = \infty$ this is a calculus minimisation problem.

Theorem 3.1. *Let $1 \leq p < \infty$. Let $f \in L^p$ with primitive $F \in L^p$. Let $t > 0$ and define $v(x, t) = v_t(x) = f * \Theta_t(x)$.*

(a) *Let $q, r \in [1, \infty]$ such that $1/p + 1/q = 1 + 1/r$. There is a constant $L_{p,q}$ such that $\|f * \Theta_t\|_r \leq L_{p,q} \|f\|'_p t^{-(2-1/q)/2}$ for all $t > 0$. The estimate is sharp in the sense that if $\psi : (0, \infty) \rightarrow (0, \infty)$ such that $\psi(t) = o(t^{-(2-1/q)/2})$ as $t \rightarrow 0^+$ or $t \rightarrow \infty$ then there is $g \in L^p$ such that $\|g * \Theta_t\|_r / \psi(t)$ is not bounded as $t \rightarrow 0^+$ or $t \rightarrow \infty$.*

(b) *Fix $t > 0$. Then v_t cannot be of one sign.*

(c) *For each $t > 0$, $\int_{-\infty}^\infty v_t(x) dx = 0$.*

(d) *For each $t > 0$, $\lim_{|x| \rightarrow \infty} v_t(x) = 0$.*

(e) *Suppose f has compact support in $[-R, R]$. Then $v_t(x) = O(|x|^{1/p} e^{-(|x|-R)^2/(4t)})$ for fixed $t > 0$ as $|x| \rightarrow \infty$. Or,*

$$|v_t(x)| \leq \begin{cases} M_p \|f\|'_1 |x| e^{-(|x|-R)^2/(4t)} t^{-3/2}, & p = 1, \quad |x| \geq R + \sqrt{2t} \\ M_p \|f\|'_p |x|^{1/p} e^{-x^2/(16t)} t^{-(1/2+1/p)}, & 1 < p < \infty, \quad |x| \geq 2R, \end{cases}$$

where

$$M_p = \begin{cases} \frac{1}{4\sqrt{\pi}}, & p = 1 \\ \frac{3^{1/p}(p-1)^{1-1/p}}{2^{1+2/p}\sqrt{\pi}p^{1-1/p}}, & 1 < p < \infty. \end{cases}$$

In part (a) remarks about possible values of q and r are the same as those following Theorem 2.1.

When $r = p$ and $q = 1$ the inequality in part (a) reads $\|f * \Theta_t\|_p \leq \|F\|_p / \sqrt{\pi t}$. When $r = \infty$ then p and q are conjugates and the inequality in part (a) reads $\|f * \Theta_t\|_\infty \leq \delta_q \|F\|_p t^{-(1+1/p)/2}$. When $p = 1$ then $q = r$ and the sharp constant in (a) is $L_{p,q} = \delta_q$. This can be proved similarly to part (b) of Theorem 4.1 using $F = \Theta_t^\beta$ in the limit $\beta \rightarrow \infty$ (effectively, $f = \delta'$). It is not known in the other cases if the constant $L_{p,q}$ is sharp or not.

Part (c) appears for $f \in L^1$ as Exercise 8.1.5 in [5].

Positive solutions of the heat equation are known to be convolutions of the Gauss–Weierstrass heat kernel with an increasing function. See [19].

An example for part (d) is $F(x) = x^{-1/p} \sin(x)$ for $x > 1$ and $F(x) = 0$ for $x \leq 1$. Then $\int_{-\infty}^{\infty} F(y) dy$ exists. And, $F \in L^s$ if and only if $s > p$.

Proof. (a) By Young’s inequality,

$$\|v_t\|_r = \|F * \Theta'_t\|_r \leq C_{p,q} \|F\|_p \|\Theta'_t\|_q = \frac{C_{p,q} \|f\|_p' \delta_q}{t^{(2-1/q)/2}}.$$

The constant $C_{p,q}$ is given in the proof of Theorem 4.1. We then take $L_{p,q} = C_{p,q} \delta_q$.

As in the proof of Theorem 4.1, define $S_t : L^p \rightarrow L^r$ by $S_t[f](x) = f * \Theta_t(x) / \psi(t)$. The estimate $\|S_t[f]\|_r \leq C_{p,q} \delta_q \|f\|_p' t^{-(2-1/q)/2} / \psi(t)$ shows that, for each $t > 0$, S_t is a bounded linear operator. Let $f_t = \Theta'_t$. Then

$$\frac{\|S_t[f_t]\|_r}{\|f_t\|_p'} = \frac{\|\Theta'_{2t}\|_r}{\psi(t) \|\Theta_t\|_p} = \frac{\delta_r}{\alpha_p 2^{(2-1/r)/2} \psi(t) t^{(2-1/q)/2}}.$$

This is not bounded in the limit $t \rightarrow 0^+$. Hence, S_t is not uniformly bounded. By the Uniform Bounded Principle it is not pointwise bounded. Therefore, there is a distribution $g \in L^p$ such that $\|g * \Theta_t\|_r \neq O(\psi(t))$ as $t \rightarrow 0^+$. And, the growth estimate $\|v_t\|_r = O(t^{-(2-1/q)/2})$ as $t \rightarrow 0^+$ is sharp. Similarly for sharpness as $t \rightarrow \infty$.

(b) Note that $v_t(x) = f * \Theta_t(x) = (F * \Theta_t)'(x)$ for each $x \in \mathbb{R}$ by Theorem 2.1(a). If $v_t \geq 0$ then $F * \Theta_t$ is increasing. But $F * \Theta_t \in L^p$ by Theorem 4.1(a) so $F * \Theta_t = 0$ and hence $v_t = 0$.

(c) From Example 2.6 we have that linear combinations of differences of Dirac distributions are dense in L^p . Given $\epsilon > 0$ there are $-\infty < b_0 < b_1 < \dots < b_N < \infty$ and $a_n \in \mathbb{R}$ such that if $g = \sum_{n=1}^N a_n (\tau_{b_{n-1}} \delta - \tau_{b_n} \delta)$ then $\|f - g\|_p' < \epsilon$. Then, by Theorem 3.1(a),

$$\left| \int_{-\infty}^{\infty} f * \Theta_t(x) dx - \int_{-\infty}^{\infty} g * \Theta_t(x) dx \right| \leq \|f - g\|_p' t^{-(2-1/q)/2},$$

where q is conjugate to p . It suffices then to show $\int_{-\infty}^{\infty} g * \Theta_t(x) dx = 0$. We have

$$\begin{aligned} \int_{-\infty}^{\infty} g * \Theta_t(x) dx &= \int_{-\infty}^{\infty} \sum_{n=1}^N a_n (\tau_{b_{n-1}} \delta * \Theta_t(x) - \tau_{b_n} \delta * \Theta_t(x)) dx \\ &= \sum_{n=1}^N a_n \int_{-\infty}^{\infty} (\Theta_t(x - b_{n-1}) - \Theta_t(x - b_n)) dx = 0. \end{aligned}$$

(d) Let $R > 0$. Since $F \in L^1_{loc}$ and Θ'_t is bounded, dominated convergence shows

$$\lim_{|x| \rightarrow \infty} \int_{-R}^R F(y) \Theta'_t(x - y) dy = \int_{-R}^R F(y) \lim_{|x| \rightarrow \infty} \Theta'_t(x - y) dy = 0.$$

Given $\epsilon > 0$ now take $R > 0$ such that $\|F\chi_{(R,\infty)}\|_p < \epsilon$. By the Hölder inequality,

$$\left| \int_R^\infty F(y)\Theta'_t(x-y) dy \right| \leq \|F\chi_{(R,\infty)}\|_p \|\Theta'_t\|_q.$$

Similarly, for integration over $(-\infty, -R)$.

(d) From the Hölder inequality, $|v_t(x)| \leq \|F\|_p \|\chi_{(x-R, x+R)}\Theta'_t\|_q$. When $p = 1$ and $q = \infty$ we compute that if $|x| \geq R + \sqrt{2t}$ then the maximum of $|\Theta'_t|$ occurs at $|x| - R$. And, $\Theta'_t(|x| - R) = (|x| - R)e^{-(|x|-R)^2/(4t)}/[4\sqrt{\pi}t^{3/2}]$.

Now let $1 < p < \infty$ and q be its conjugate. Suppose $x > 2R$. Then

$$\begin{aligned} \|\chi_{(x-R, x+R)}\Theta'_t\|_q^q &= \frac{1}{(4\sqrt{\pi}t^{3/2})^q} \int_{x-R}^{x+R} y^q e^{-qy^2/(4t)} dy \\ &\leq \frac{(x+R)^{q-1}2t}{(4\sqrt{\pi}t^{3/2})^q q} \left[e^{-q(x-R)^2/(4t)} - e^{-q(x+R)^2/(4t)} \right] \\ &\leq \frac{3^{q-1}x^{q-1}te^{-qx^2/(16t)}}{2^{q-2}(4\sqrt{\pi}t^{3/2})^q q}. \end{aligned}$$

□

4. APPENDIX

The Appendix contains a proof of sharp L^p estimates for the heat equation and a lemma on differentiation of convolutions.

The solution $u(x, t)$ that is in $C^2(\mathbb{R})$ with respect to x and in $C^1((0, \infty))$ with respect to t of (1.1) and (1.2) is given by the convolution $u_t(x) = F * \Theta_t(x) = \int_{-\infty}^\infty F(x-y)\Theta_t(y) dy$ where the Gauss–Weierstrass heat kernel is $\Theta_t(x) = \exp(-x^2/(4t))/(2\sqrt{\pi t})$. For example, see [6].

The heat kernel has the following properties. Let $t > 0$ and let $s \neq 0$ such that $1/s + 1/t > 0$. Then

$$(4.1) \quad \Theta_t * \Theta_s = \Theta_{t+s}$$

$$(4.2) \quad \|\Theta_t\|_q = \frac{\alpha_q}{t^{(1-1/q)/2}} \text{ where } \alpha_q = \begin{cases} 1, & q = 1 \\ \frac{1}{(2\sqrt{\pi})^{1-1/q} q^{1/(2q)}}, & 1 < q < \infty \\ \frac{1}{2\sqrt{\pi}}, & q = \infty. \end{cases}$$

The last of these follows from the probability integral $\int_{-\infty}^\infty e^{-x^2} dx = \sqrt{\pi}$.

We now present some estimates for $F * \Theta_t$ when $F \in L^p$, as they are used in Theorem 2.1 with data in L^p . These estimates are well-known, for example, [9] when $r = \infty$. And, [11, Proposition 3.1]. However, we have not been able to find a proof in the literature that the estimates are sharp.

Theorem 4.1. *Let $1 \leq p \leq \infty$ and $F \in L^p$.*

(a) *If $p \leq s \leq \infty$ then $F * \Theta_t \in L^s$.*

(b) *Let $q, r \in [1, \infty]$ such that $1/p + 1/q = 1 + 1/r$. There is a constant $K_{p,q}$ such that $\|F * \Theta_t\|_r \leq K_{p,q} \|F\|_p t^{-(1-1/q)/2}$ for all $t > 0$. The estimate is sharp in the sense that if $\psi : (0, \infty) \rightarrow (0, \infty)$ such that $\psi(t) = o(t^{-(1-1/q)/2})$ as $t \rightarrow 0^+$ or $t \rightarrow \infty$ then there is $G \in L^p$ such that $\|G * \Theta_t\|_r / \psi(t)$ is not bounded as $t \rightarrow 0^+$ or $t \rightarrow \infty$. The constant*

$K_{p,q} = (c_p c_q / c_r)^{1/2} \alpha_q$, where $c_p = p^{1/p} / (p')^{1/p'}$ with p, p' being conjugate exponents. It cannot be replaced with any smaller number.

(c) If $1 \leq s < p$ then $F * \Theta_t$ need not be in L^s .

When $r = p$ and $q = 1$ the inequality in part (b) reads $\|F * \Theta_t\|_p \leq \|F\|_p$. When $r = \infty$ then p and q are conjugates and the inequality in part (b) reads $\|F * \Theta_t\|_\infty \leq \|F\|_p t^{-1/(2p)}$.

The condition for sharpness in Young's inequality is that both functions be Gaussians. This fact is exploited in the proof of part (b). See [13, p. 99], [2] and [18].

Proof. (a), (b) Young's inequality gives

$$(4.3) \quad \|F * \Theta_t\|_r \leq C_{p,q} \|F\|_p \|\Theta_t\|_q = \frac{C_{p,q} \|F\|_p \alpha_q}{t^{(1-1/q)/2}},$$

where α_q is given in (4.2). The sharp constant, given in [13, p. 99], is $C_{p,q} = (c_p c_q / c_r)^{1/2}$ where $c_p = p^{1/p} / (p')^{1/p'}$ with p, p' being conjugate exponents. Note that $c_1 = c_\infty = 1$. Also, $0 < C_{p,q} \leq 1$. We then take $K_{p,q} = C_{p,q} \alpha_q$.

To show the estimate $\|u_t\|_r = O(t^{-(1-1/q)/2})$ is sharp as $t \rightarrow 0^+$ and $t \rightarrow \infty$, let ψ be as in the statement of the theorem. Fix $p \leq r \leq \infty$. Define the family of linear operators $S_t: L^p \rightarrow L^r$ by $S_t[F](x) = F * \Theta_t(x) / \psi(t)$. The estimate $\|S_t[F]\|_r \leq K_{p,q} \|F\|_p t^{-(1-1/q)/2} / \psi(t)$ shows that, for each $t > 0$, S_t is a bounded linear operator. Let $F_t = \Theta_t$. Then, from (4.1) and (4.2),

$$\frac{\|S_t[F_t]\|_r}{\|F_t\|_p} = \frac{\|\Theta_t * \Theta_t\|_r}{\psi(t) \|\Theta_t\|_p} = \frac{\|\Theta_{2t}\|_r}{\psi(t) \|\Theta_t\|_p} = \frac{\alpha_r}{\alpha_p 2^{(1-1/r)/2} \psi(t) t^{(1-1/q)/2}}.$$

This is not bounded in the limit $t \rightarrow 0^+$. Hence, S_t is not uniformly bounded. By the Uniform Bounded Principle it is not pointwise bounded. Therefore, there is a function $F \in L^p$ such that $\|F * \Theta_t\|_r \neq O(\psi(t))$ as $t \rightarrow 0^+$. And, the growth estimate $\|F * \Theta_t\|_r = O(t^{-(1-1/q)/2})$ as $t \rightarrow 0^+$ is sharp. Similarly for sharpness as $t \rightarrow \infty$.

Now show the constant $K_{p,q}$ cannot be reduced. A calculation shows we have equality in (4.3) when $F = \Theta_t^\beta$ and β is given by the equation

$$(4.4) \quad \frac{\beta^{1-1/q}}{(\beta+1)^{1-1/r}} = \frac{c_p c_q}{c_r} \left(\frac{\alpha_p \alpha_q}{\alpha_r} \right)^2 = \left(1 - \frac{1}{p}\right)^{1-1/p} \left(1 - \frac{1}{q}\right)^{1-1/q} \left(1 - \frac{1}{r}\right)^{-(1-1/r)}.$$

First consider the case $p \neq 1$ and $q \neq 1$. Notice that $1 - 1/r = (1 - 1/q) + (1 - 1/p) > 1 - 1/q$. Let $g(x) = x^A (x+1)^{-B}$ with $B > A > 0$. Then g is strictly increasing on $(0, A/(B-A))$ and strictly decreasing for $x > A/(B-A)$ so there is a unique maximum for g at $A/(B-A)$. Put $A = 1 - 1/q$ and $B = 1 - 1/r$. Then

$$g\left(\frac{A}{B-A}\right) = \frac{\beta^{1-1/q}}{(\beta+1)^{1-1/r}} = \left(1 - \frac{1}{p}\right)^{1-1/p} \left(1 - \frac{1}{q}\right)^{1-1/q} \left(1 - \frac{1}{r}\right)^{-(1-1/r)}.$$

Hence, (4.4) has a unique positive solution for β given by $\beta = (1 - 1/q)/(1 - 1/p)$.

If $p = 1$ then $q = r$. In this case, (4.4) reduces to $(1 + 1/\beta)^{1-1/q} = 1$ and the solution is given in the limit $\beta \rightarrow \infty$. Sharpness of (4.3) is then given in this limit. It can also be seen that taking F to be the Dirac distribution gives equality.

If $q = 1$ then $p = r$. Now, (4.4) reduces to $(\beta+1)^{1-1/p} = 1$ and $\beta = 0$. There is equality in (4.3) when $F = 1$. This must be done in the limit $\beta \rightarrow 0^+$.

If $p = q = r = 1$ then there is equality in (4.3) for each $\beta > 0$.

Hence, the constant in (4.3) is sharp.

(c) Suppose $F \geq 0$ and F is decreasing on $[c, \infty)$ for some $c \in \mathbb{R}$. Let $x > c$. Then

$$\begin{aligned} F * \Theta_t(x) &\geq \int_c^x F(y) \Theta_t(x-y) dy \geq F(x) \int_c^x \Theta_t(x-y) dy \\ &= \frac{F(x)}{\sqrt{\pi}} \int_0^{(x-c)/(2\sqrt{t})} e^{-y^2} dy \sim F(x)/2 \quad \text{as } x \rightarrow \infty. \end{aligned}$$

Now put $F(x) = 1/[x^{1/p} \log^2(x)]$ for $x \geq e$ and $F(x) = 0$, otherwise. For $p = \infty$ replace $x^{1/p}$ by 1. \square

We prove a lemma on pointwise differentiation of L^p convolutions. A similar result for $p = 1$ and functions with bounded derivatives appears in [7, Proposition 8.10].

Lemma 4.2 (Differentiation of Convolutions). *Let $F \in L^p$ for some $1 \leq p \leq \infty$. Let q be the conjugate of p and suppose $\psi \in L^q$.*

(a) *Suppose ψ is a differentiable function such that ψ' is absolutely continuous and $\psi'' \in L^q$. Then $F * \psi$ is differentiable and $(F * \psi)'(x) = F * \psi'(x)$ for each $x \in \mathbb{R}$.*

(b) *Suppose ψ is a differentiable function such that ψ is absolutely continuous, ψ' is bounded on each compact interval and $\psi' \in L^q$. Then $F * \psi$ is differentiable and $(F * \psi)'(x) = F * \psi'(x)$ for each $x \in \mathbb{R}$.*

(c) *Suppose ψ is a C^∞ function such that $\psi^{(n)} \in L^q$ for each $n \geq 0$. Then $F * \psi \in C^\infty(\mathbb{R})$ and $(F * \psi)^{(n)}(x) = F * \psi^{(n)}(x)$ for each $n \geq 1$ and each $x \in \mathbb{R}$.*

Proof. Let $w = F * \psi$. (a) Without loss of generality, we can take $h > 0$ and consider

$$\frac{w(x+h) - w(x)}{h} = \int_{-\infty}^{\infty} F(y) \left[\frac{\psi(x+h-y) - \psi(x-y)}{h} \right] dy = F * \psi'(x) - R(h).$$

The remainder from Taylor's theorem is

$$R(h) = \frac{1}{h} \int_{-\infty}^{\infty} F(y) \int_{x-y}^{x+h-y} (z-x-h+y) \psi''(z) dz dy.$$

And, by the Hölder inequality, Jensen's inequality and the Fubini–Tonelli theorem,

$$\begin{aligned} |R(h)| &\leq \|F\|_p \left(\int_{-\infty}^{\infty} \int_{x-y}^{x+h-y} |z-x-h+y|^q |\psi''(z)|^q \frac{dz}{h} \right)^{1/q} dy \\ &= \|F\|_p \left(\int_{-\infty}^{\infty} |\psi''(z)|^q \int_{x-z}^{x+h-z} |z-x-h+y|^q \frac{dy}{h} dz \right)^{1/q} \\ &= \frac{\|F\|_p \|\psi''\|_q |h|}{(q+1)^{1/q}} \rightarrow 0 \text{ as } h \rightarrow 0. \end{aligned}$$

Minor changes are needed when $p = 1$. Therefore, $(F * \psi)'(x) = F * \psi'(x)$.

(b) Let $R > 0$. Write

$$F * \psi(x) = \int_{-R}^R F(y) \psi(x-y) dy + \int_{|y|>R} F(y) \psi(x-y) dy.$$

Since $F \in L^1_{loc}$, and ψ' is bounded on compact intervals, dominated convergence shows

$$\frac{d}{dx} \int_{-R}^R F(y)\psi(x-y) dy = \int_{-R}^R F(y)\psi'(x-y) dy.$$

Now, let $h > 0$ and consider

$$\left| \int_R^\infty F(y) \left[\frac{\psi(x+h-y) - \psi(x-y)}{h} \right] dy \right| \leq \|F\chi_{(R,\infty)}\|_p \left(\int_R^\infty \left| \frac{\psi(x+h-y) - \psi(x-y)}{h} \right|^q dy \right)^{1/q}.$$

Given $\epsilon > 0$ take R large enough so that $\|F\chi_{(R,\infty)}\|_p < \epsilon$. Using the fundamental theorem of calculus, Jensen's inequality and then the Fubini–Tonelli theorem, we have

$$\begin{aligned} \int_R^\infty \left| \frac{\psi(x+h-y) - \psi(x-y)}{h} \right|^q dy &= \int_R^\infty \left| \int_{x-y}^{x+h-y} \psi'(z) \frac{dz}{h} \right|^q dy \\ &\leq \int_R^\infty \int_{x-y}^{x+h-y} |\psi'(z)|^q \frac{dz}{h} dy \\ &= \int_{-\infty}^\infty |\psi'(z)|^q \int_{x-z}^{x+h-z} \frac{dy}{h} dz \\ &= \|\psi'\|_q^q. \end{aligned}$$

Therefore,

$$\left| \int_R^\infty F(y) \frac{\psi(x+h-y) - \psi(x-y)}{h} dy \right| \leq \epsilon \|\psi'\|_q.$$

The case $p = 1$ is similar.

(c) Follows from (a) or (b). □

REFERENCES

- [1] S. Axler, P. Bourdon and W. Ramey, *Harmonic function theory*, New York, Springer-Verlag, 2001.
- [2] W. Beckner, *Inequalities in Fourier analysis*, Ann. of Math. (2) **102**(1975), 159–182.
- [3] J.R. Cannon, *The one-dimensional heat equation*, Menlo Park, Addison–Wesley, 1984.
- [4] R. Dautray and J.-L. Lions, *Mathematical analysis and numerical methods for science and technology, vol. 5* (trans. A. Craig), Berlin, Springer-Verlag, 1992.
- [5] B. Epstein, *Partial differential equations*, New York, McGraw-Hill, 1962.
- [6] G.B. Folland, *Introduction to partial differential equations*, Princeton, Princeton University Press, 1995.
- [7] G.B. Folland, *Real analysis*, New York, Wiley, 1999.
- [8] F.G. Friedlander and M. Joshi, *Introduction to the theory of distributions*, Cambridge, Cambridge University Press, 1999.
- [9] K.E. Gustafson, *Introduction to partial differential equations and Hilbert space methods*, New York, Dover, 1999.
- [10] I.I. Hirschman and D.V. Widder, *The convolution transform*, Princeton, Princeton University Press, 1955.
- [11] T. Iwabuchi, T. Matsuyama and K. Taniguchi, *Boundedness of spectral multipliers for Schrödinger operators on open sets*, Rev. Mat. Iberoam. **34**(2018), 1277–1322.
- [12] R.J. Iorio, Jr. and V. de Magalhães Iorio, *Fourier analysis and partial differential equations*, New York, Cambridge University Press, 2001.

- [13] E.H. Lieb and M. Loss, *Analysis*, Providence, American Mathematical Society, 2001.
- [14] Z. Szmydt, *Fourier transforms and linear differential equations* (trans. M. Kuczma), Dordrecht, D. Reidel, 1977.
- [15] E. Talvila, *The regulated primitive integral*, Illinois J. Math. **53**(2009), 1187–1219.
- [16] E. Talvila, *The L^p primitive integral*, Math. Slovaca **64**(2014), 1497–1524.
- [17] E. Talvila, *The one-dimensional heat equation in the Alexiewicz norm*, Adv. Pure Appl. Math. **6**(2015), 13–37.
- [18] G. Toscani, *Heat equation and the sharp Young's inequality*, arXiv:1204.2086 (2012).
- [19] D.V. Widder, *The heat equation*, New York, Academic Press, 1975.

DEPARTMENT OF MATHEMATICS & STATISTICS, UNIVERSITY OF THE FRASER VALLEY, ABBOTSFORD, BC CANADA V2S 7M8

Email address: Erik.Talvila@ufv.ca