

On 3-generated axial algebras of Jordan type $1/2$

Ravil Bildanov, Ilya Gorshkov

June 25, 2024

Abstract: Axial algebras of Jordan type η are a special type of commutative non-associative algebras. They are generated by idempotents whose adjoint operators have the minimal polynomial dividing $(x - 1)x(x - \eta)$, where η is a fixed value that is not equal to 0 or 1. These algebras have restrictive multiplication rules that generalize the Peirce decomposition for idempotents in Jordan algebras. A universal 3-generated algebra of Jordan type $\frac{1}{2}$ as an algebra with 4 parameters was constructed by I. Gorshkov and A. Staroletov. Depending on the value of the parameter, the universal algebra may contain a non-trivial form radical. In this paper, we describe all semisimple 3-generated algebras of Jordan type $\frac{1}{2}$ over a quadratically closed field.

MSC 2020 code: 20D60; 17A30; 17C27

Keywords: axial algebras, Jordan type algebras.

Introduction

Axial algebras of Jordan type η were introduced by Hall, Rehren, and Shpectorov [2] within the framework of the general theory of axial algebras. These algebras are commutative non-associative algebras over a field \mathbb{F} , generated by special idempotents known as primitive axes. While Jordan algebras generated by primitive idempotents are an example of algebras of Jordan type $\frac{1}{2}$, not all algebras of this type are Jordan algebras. The Matsuo algebras, constructed from the group of 3-transpositions, are examples of such algebras. It was proved in [2] (with a correction in [3]) that for $\eta \neq 1/2$, algebras of Jordan type η are the Matsuo algebras or their quotient algebras. Therefore, the case $\eta = 1/2$ is special for algebras of Jordan type, and for this η , they are called algebras of Jordan type half. The class of Matsuo algebras was introduced by Matsuo [6] and later generalized in [2].

Algebras of Jordan type half are not exhausted by Matsuo algebras and their quotient algebras. Moreover, the quotient algebras of Matsuo algebras do not contain all Jordan algebras generated by primitive idempotents. For example, the 27-dimensional Albert algebra is generated by 4 primitive idempotents and hence it is an algebra of Jordan type half but not a Matsuo algebra [5].

A universal 3-generated algebra of Jordan type half $A(\alpha, \beta, \gamma, \psi)$ was constructed in [1]. It is proved there that if $(\alpha + \beta + \gamma - 2\psi - 1)(\alpha\beta\gamma - \psi^2) \neq 0$ and $\psi^2 - \alpha\beta\gamma$ is a square

in \mathbb{F} , then $A(\alpha, \beta, \gamma, \psi)$ is isomorphic to the algebra $M_3^+(\mathbb{F})$ of 3×3 matrices with Jordan multiplication. Otherwise, the algebra $A(\alpha, \beta, \gamma, \psi)$ is not simple.

A Frobenius form (\cdot, \cdot) on algebra A is a nonzero symmetric bilinear form that associates with multiplication in A , i.e., $\forall a, b, c \in A$, we have $(ab, c) = (ac, b)$ and $(a, a) \neq 0$ for each axis $a \in A$. Hall, Rehren, and Shpectorov [2] showed that for algebras of Jordan type, there exists a unique Frobenius form with the property $(a, a) = 1$ for every primitive axis a .

Let A be an algebra with a Frobenius form (\cdot, \cdot) . The radical of the form (\cdot, \cdot) is an ideal $R(A)$ generated by elements x such that $(x, a) = (a, x) = 0$ for every element $a \in A$.

The purpose of this article is to describe all 3-generated algebras of Jordan type half with trivial radical over a quadratically closed field.

The universal 3-generated algebra $A(\alpha, \beta, \gamma, \psi)$ from [1] is a Jordan algebra, so any quotient algebra is also Jordan. However, we use notation of universal algebra instead of notation of Jordan algebras. When we embedding a 3-generated algebra into some large algebra, its parameters remain unchanged, despite the fact that the large algebra may not be Jordan. For example, 2-generated subalgebras are in studied in [7, Section 3]. In this paper it is shown that isomorphic 2-generated subalgebras with different parameters of the Frobenius form have fundamentally different properties when embedded in a large algebra. This statement emphasizes the importance of studying semisimple 3-generated algebras of Jordan type as algebras with parameters.

1 Preliminary results

We consider commutative non-associative algebras over a ground field \mathbb{F} of characteristic not two. For definitions, we almost always follow [2] and [4].

By $L\langle X \rangle$ denote the linear span of the set X over \mathbb{F} , and by $\langle\langle X \rangle\rangle$ denote the algebra generated by the set X .

Notation 1. Given $a \in A$ and $\lambda \in \mathbb{F}$, consider the subspace $A_\lambda(a) = \{u \in A \mid au = \lambda u\}$.

Obviously, $A_\lambda(a)$ is the eigenspace of the operator $ad_a : x \rightarrow ax$, associated with $\lambda \in \mathbb{F}$.

Definition 1. An idempotent $a \in A$ is said to be primitive if $\dim(A_1(a)) = 1$.

Definition 2. An algebra A is an algebra of Jordan type half if A is generated by the set of primitive idempotents X such that for every $x \in X$, we have a decomposition $A = A_0(x) \oplus A_1(x) \oplus A_{1/2}(x)$ with the following fusion (multiplication) rules:

$$A_0(x)A_{1/2}(x) \subseteq A_{1/2}(x), A_1(x)A_{1/2}(x) \subseteq A_{1/2}(x), A_0(x)A_1(x) = 0,$$

$$(A_0(x))^2 \subseteq A_0(x), (A_1(x))^2 \subseteq A_1(x), (A_{1/2}(x))^2 \subseteq A_0(x) \oplus A_1(x).$$

Such idempotents are called axes. By an n -generated algebra we mean an algebra generated by n primitive axes.

Given an axial algebra of Jordan type half A and an axis $a \in A$, the map $\tau_a : A \rightarrow A$ which acts as follows, $x^{\tau_a} = (-1)^{2\lambda}x$ for $x \in A_\lambda(a)$, is an involution automorphism of A

called Miyamoto involution. It is clear that $y^{\tau_a} = y - 4ya + 4\phi_a(y)$ for each $y \in A$, where $\phi_a(y)$ is a projection of y on $A_1(a)$.

Let us introduce some classes of simple Jordan algebras.

Definition 3. Denote by $M_n^+(\mathbb{F})$ the matrix algebra $M_n(\mathbb{F})$ with Jordan product $A \circ B = \frac{1}{2}(AB + BA)$.

Definition 4. If j is an involutory automorphism of $M_n(\mathbb{F})$, then we define the Hermitian Jordan algebra $H(M_n(\mathbb{F}), j)$ as $\{A \in M_n^+(\mathbb{F}) \mid j(A) = A\}$.

Definition 5. Let V be an n -dimensional vector space over (\mathbb{F}) a symmetric bilinear form ϕ we define:

$$(a \oplus \mathbf{v}) \bullet (b \oplus \mathbf{w}) = (ab + \phi(\mathbf{v}, \mathbf{w})) \oplus (a\mathbf{w} + b\mathbf{v}), \text{ where } a, b \in \mathbb{F} \text{ and } \mathbf{v}, \mathbf{w} \in V.$$

Then the vector space $\mathbb{F} \oplus V$ with multiplication \oplus gives a Jordan algebra which will called Jordan algebra of symmetric bilinear form $JForm_n(\mathbb{F})$.

It is well known that $M_n^+(\mathbb{F})$, $H_n^+(\mathbb{F})$, $JForm_n(\mathbb{F})$ for $n \geq 2$ are simple Jordan algebras see [9, Theorem 23.1.2], so they are algebras of Jordan type half.

Lemma 1. [3, Theorem 4.1] Every algebra of Jordan type η admits a unique Frobenius form which satisfies the property $(a, a) = 1$ for all axes $a \in X$.

Lemma 2. [2, Proposition 2.7] The radical of Frobenius form $R(A)$ coincides with the largest ideal of A containing no axes from A .

Definition 6. An algebra of Jordan type η is semisimple if the radical of Frobenius form is trivial.

If A is a Jordan type η algebra, then $A/R(A)$ is a Jordan type η algebra too. It follows from Lemma 1 that $A/R(A)$ has a unique Frobenius form with the property $(\bar{a}, \bar{a}) = 1$ for every axis $\bar{a} \in A/R(A)$. In the next lemma we prove that when factorizing by the radical, the Frobenius form does not change.

Lemma 3. Let A be an algebra of Jordan type η . Then for all $a, b \in A$ and their images $\bar{a}, \bar{b} \in A/R(A)$, we have $(a, b) = (\bar{a}, \bar{b})$.

Proof. Let $a = \bar{a} + r_a, b = \bar{b} + r_b$, where $\bar{a}, \bar{b} \in A/R(A), r_a, r_b \in R(A)$. Then $(a, b) = (\bar{a} + r_a, \bar{b} + r_b) = (\bar{a}, \bar{b}) + (\bar{a}, r_b) + (\bar{b}, r_a) + (r_a, r_b) = (\bar{a}, \bar{b})$. \square

Lemma 4. [4, Lemma 2] Let A be a finitely generated algebra of Jordan type half, a, b are axes, let $\alpha := (a, b)$. Then we have the following equalities:

1. $(a_0(b))^2 = (1 - \alpha)a_0(b)$;
2. $(a_{1/2}(b))^2 = \alpha a_0(b) + (\alpha - \alpha^2)a$;
3. $a_0(b)a_{1/2}(b) = \frac{1}{2}(1 - \alpha)a_{1/2}(b)$.

From Lemma 1 it follows that on any algebra of Jordan type η A there is a unique Frobenius form with the property $(a, a) = 1$ for any axis $a \in A$.

Lemma 5. *Let $A = \langle\langle a, b \rangle\rangle$ be a 2-generated algebra of Jordan type half. Then one of the following holds:*

1. $\dim(A) = 1$, $(a, b) = 1$, $a = b$, $\dim(R(A)) = 0$;
2. $\dim(A) = 2$, $(a, b) = 0$, $A \cong \mathbb{F} \oplus \mathbb{F}$, $\dim(R(A)) = 0$;
3. $\dim(A) = 2$, $(a, b) = 1$, $\dim(R(A)) = 1$;
4. $\dim(A) = 3$, $(a, b) = 0$, $\dim(R(A)) = 1$, $A/R(A) \cong \mathbb{F} \oplus \mathbb{F}$;
5. $\dim(A) = 3$, $(a, b) = 1$, $\dim(R(A)) = 2$;
6. $\dim(A) = 3$, $(a, b) \neq 0, 1$, and A is a Matsuo algebra. In particular, it is a simple Jordan algebra isomorphic to $J\text{Form}_2(\mathbb{F})$.

Proof. The assertion of the lemma is a simple consequence of [2, Theorem 1.1]. □

Lemma 6. [4, Corollary 1] *Let A be a 2-generated algebra of Jordan type half with generating axes a and b . Let $\alpha := (a, b)$. Then we have*

1. $a(ab) = \frac{1}{2}(\alpha a + ab)$;
2. $(ab)b = \frac{1}{2}(\alpha b + ab)$;
3. $(ab)(ab) = \frac{\alpha}{4}(a + b + 2ab)$.

Lemma 7. [1, Theorem 1] *Let A be a 3-generated Jordan type half algebra. There exists a 3-generated 9-dimensional algebra $A(\alpha, \beta, \gamma, \psi)$ such that A is a quotient algebra of $A(\alpha, \beta, \gamma, \psi)$ for suitable values of parameters $\alpha, \beta, \gamma, \psi$.*

Let $A = \langle\langle a, b, c \rangle\rangle$, $\dim(A) = 9$, $\alpha = (a, b)$, $\beta = (b, c)$, $\gamma = (a, c)$, $\psi = (ab, c)$. In Table 1 below (that is similar to [1, Table 6] up to renumbering rows), we present all possible relations for $\alpha, \beta, \gamma, \psi$ for $A(\alpha, \beta, \gamma, \psi)$ to not be simple.

2 Main Results

In this section we assume that A is a 3-generated algebra of Jordan type half with a trivial radical over a quadratically closed field \mathbb{F} and denote with (\cdot, \cdot) the unique Frobenius form on A satisfying the property that $(a, a) = 1$ for every axis a of A .

Theorem. *Let A be a 3-generated algebra of Jordan type half with a trivial radical over a quadratically closed field \mathbb{F} with characteristic not equal to two or three. Then A is isomorphic to one of the following algebras:*

A_i	Relations	$\dim(A/R(A))$	Basis of the radical
A_1	$\psi = \alpha = \beta = \gamma = 1$	1	$b - a, c - a, ab - a, bc - a, ac - a,$ $a(bc) - a, b(ac) - a, c(ab) - a$
A_2	$\psi = \alpha = \beta = 0, \gamma = 1$	2	$c - a, ab, bc, ac - a,$ $a(bc), b(ac), c(ab)$
A_3	$\psi = \alpha = \beta = \gamma = 0$	3	$ab, bc, ac, a(bc), b(ac), c(ab)$
A_4	$\psi = \alpha = 0, \beta, \gamma \neq 0,$ $\beta + \gamma = 1$	3	$ab, \frac{1}{2}\gamma a - \frac{1}{2}\beta b - \frac{1}{2}c + bc,$ $-\frac{1}{2}\gamma a + \frac{1}{2}\beta b - \frac{1}{2}c + ac,$ $\frac{1}{4}\gamma a + \frac{1}{4}\beta b - \frac{1}{4}c + a(bc),$ $\frac{1}{4}\gamma a + \frac{1}{4}\beta b - \frac{1}{4}c + b(ac), c(ab)$
A_5	$\alpha\beta\gamma = \psi^2, \psi \neq 0, \alpha \neq 1,$ $\alpha + \beta + \gamma = 2\psi + 1$	3	$\alpha(\beta - 1)a + \alpha(\gamma - 1)b + \alpha(1 - \alpha)c + (2\alpha - 2\psi)ab,$ $(\alpha\beta - \alpha\psi)b + (\psi - \alpha\beta)ab + (\alpha^2 - \alpha)bc,$ $(\alpha\gamma - \alpha\psi)a + (\psi - \alpha\gamma)ab + (\alpha^2 - \alpha)ac,$ $(\alpha\psi - \alpha^2\beta)a + (\alpha + \psi - \alpha^2 - \alpha\gamma)ab + 2\alpha(\alpha - 1)a(bc),$ $\alpha(\psi - \alpha\gamma)b + (\alpha + \psi - \alpha^2 - \alpha\beta)ab + 2\alpha(\alpha - 1)b(ac),$ $(\psi - \alpha\beta)a + (\psi - \alpha\gamma)b + (1 - \alpha)ab + 2(\alpha - 1)c(ab)$
A_6	$\psi = \alpha = \beta = 0, \gamma \neq 0, 1$	4	$ab, bc, ac, a(bc), b(ac), c(ab)$
A_7	$\psi^2 \neq \alpha\beta\gamma,$ $\alpha + \beta + \gamma = 2\psi + 1,$ $\alpha \neq 1$	4	$\frac{1}{2}(\beta - 1)a + \frac{1}{2}(\beta - \alpha)b + \frac{1}{2}(1 - \alpha)c + (1 - \beta)ab + (\alpha - 1)bc,$ $\frac{1}{2}(\gamma - \alpha)a + \frac{1}{2}(\gamma - 1)b + \frac{1}{2}(1 - \alpha)c + (1 - \gamma)ab + (\alpha - 1)ac,$ $(2\psi - 2\alpha\beta + \beta - 1)a + (\gamma - 1)b + (1 - \alpha)c + (4 - 2\alpha - 2\gamma)ab + (4\alpha - 4)a(bc),$ $(\beta - 1)a + (2\psi - 2\alpha\gamma + \gamma - 1)b + (1 - \alpha)c + (4 - 2\alpha - 2\beta)ab + (4\alpha - 4)b(ac),$ $(\psi - \alpha)a + (\psi - \alpha)b + \alpha(1 - \alpha)c + (2 - \beta - \gamma)ab + (2\alpha - 2)c(ab)$
A_8	$\psi = \alpha = 0, \beta, \gamma \neq 0,$ $\beta + \gamma \neq 1$	6	$ab, b(ac) - a(bc), c(ab)$
A_9	$\alpha\beta\gamma = \psi^2, \psi \neq 0,$ $\alpha + \beta + \gamma \neq 2\psi + 1$	6	$-\beta\gamma ab - \alpha\beta ac + 2\psi a(bc),$ $-\beta\gamma ab - \alpha\gamma bc + 2\psi b(ac),$ $-\alpha\gamma bc - \alpha\beta ac + 2\psi c(ab)$

Table 1: Bases of the radical

1. $\mathbb{F}^n, n \in \{1, 2, 3\};$
2. $JForm_2(\mathbb{F});$
3. $\mathbb{F} \oplus JForm_2(\mathbb{F});$
4. $M_2^+(\mathbb{F});$
5. $H(M_3(\mathbb{F}), j)$ with $j(X) = X^T;$
6. $M_3^+(\mathbb{F}).$

It follows from Lemma 7, that any algebra of Jordan type half is isomorphic to the quotient algebra of $A(\alpha, \beta, \gamma, \psi)$ for some parameters $\alpha, \beta, \gamma, \psi$. From [9, Enlightenment Structure Theorem] we obtain that $A(\alpha, \beta, \gamma, \psi)/R(A(\alpha, \beta, \gamma, \psi))$ is a direct sum of simple algebras. Consequently, to describe 3-generated algebras of Jordan type half with a trivial radical, we need to describe for each choice of the parameters $\alpha, \beta, \gamma, \psi$ the quotient algebras of $A(\alpha, \beta, \gamma, \psi)$ by its radical.

We use the description of the algebra $A(\alpha, \beta, \gamma, \psi)$ from [1, Theorem 2]. Let us remind you that following [1], we use the following notation: $\alpha = (a, b), \beta = (b, c), \gamma = (a, c), \psi = (ab, c).$

In Table 1, one can find the dimensions and bases of the radicals of the algebra $A(\alpha, \beta, \gamma, \psi)$. Denote by A_i the universal 9-dimensional algebra $A(\alpha_i, \beta_i, \gamma_i, \psi_i)$ with parameters and numeration from Table 1, R_i the radical of this algebra and by S_i the quotient algebra A_i/R_i .

We begin with two trivial propositions for 1-dimensional and 2-dimensional algebras, which are not generated by three linearly independent axes.

Proposition 1. *If A is a 1-dimensional algebra of Jordan type half with a trivial radical, then $A \cong S_1$.*

Proof. It is easy to see that $S_1 \cong \mathbb{F}$. We have that A is 1-dimensional, so $\dim L\langle a, b, c \rangle = 1$ and $a = b = c$. Hence $A \cong \mathbb{F} \cong S_1$. \square

Proposition 2. *If A is a 2-dimensional 3-generated algebra of Jordan type half with a trivial radical, then $A \cong \mathbb{F} \oplus \mathbb{F} \cong S_2$.*

Proof. By Lemma 5, there is only one 2-dimensional algebra of Jordan type half with a trivial radical, so $A \cong \mathbb{F} \oplus \mathbb{F} \cong S_2$. \square

Lemma 8. *Algebras S_4 and S_5 are isomorphic.*

Proof. We will first show that $S_4 = \langle\langle a, c \rangle\rangle$. Put $S = \langle\langle a, c \rangle\rangle$. We have that $S = S_0(a) + S_1(a) + S_{1/2}(a)$ and $c = c_0(a) + \gamma a + c_{1/2}(a)$, where $c_0(a) \in S_0(a)$, $c_{1/2}(a) \in S_{1/2}(a)$. Assume that $c_{1/2}(a) = 0$. We have $c = c^2 = (c_0(a))^2 + (\gamma a)^2$, in particular $(c_0(a))^2 = c_0(a)$. Consequently $c_0(a)c = (c_0(a))^2 = c_0(a)$; which contradicts c being the primitive idempotent. Thus set $c_{1/2}(a) \neq 0$. Assume that $c_0(a) = 0$. Therefore, $\gamma a + c_{1/2}(a) = c = c^2 = \gamma^2 a + \gamma c_{1/2}(a) + (c_{1/2}(a))^2$. Hence $\gamma = 1$ and from definition of S_4 it follows that $\beta = 0$. In this case $(a - c, b) = (a, b) - (c, b) = 0$. It follows that $a - c \in R(S_4)$, which is a contradiction. Therefore, $\dim(S) = 3$. Thus, $S_4 = S$. Hence S_4 is generated by 2 axes. From Lemma 5 it follows that $S_4 \simeq JForm_2(\mathbb{F})$

Consider the algebra S_5 . By definition of S_5 , we have $(a, b) \notin \{0, 1\}$, where a, b are the axes from the generating set of the algebra S_5 . Lemma 5 implies that $\langle\langle a, b \rangle\rangle \simeq JForm_2(\mathbb{F})$. Since $\dim(\langle\langle a, b \rangle\rangle) = 3$, we obtain $\langle\langle a, b \rangle\rangle = S_5$, in particular $S_5 \simeq S_4$. \square

Proposition 3. *If A is a 3-dimensional 3-generated algebra of Jordan type half with a trivial radical, then A is isomorphic to either S_3 or S_5 .*

Proof. Assume that A is 2-generated and let a and b are the generating axes. From Lemma 5, it follows that there is only one 3-dimensional 2-generated algebra of Jordan type half with a trivial radical. In this case, we can choose any other axis of the algebra A . Let us call it c . Put $c = a^{\tau_b} = a - 4ab + 4ab$. We have $\beta = \alpha$, $\gamma = (1 - 2\alpha)^2$ and $\psi = \alpha(2\alpha - 1)$. Therefore $\alpha\beta\gamma = \psi^2$, $\psi \neq 0$, $\alpha \neq 1$, $\alpha + \beta + \gamma = 2\psi + 1$. So in this case $A \simeq S_5$.

Assume that A is not generated by 2 axes. Therefore, based on the dimension of A , we conclude that $A = L\langle a, b, c \rangle$.

Assume that $ab \notin L\langle a, b \rangle$. Hence $\dim(\langle\langle a, b \rangle\rangle) = 3$. Therefore $c \in L\langle a, b, ab \rangle = A$, which is a contradiction. Similarly, we can show that if $ac \in L\langle a, c \rangle$ and $bc \in L\langle b, c \rangle$. In particular,

we have $\dim(\langle\langle a, b \rangle\rangle) = \dim(\langle\langle a, c \rangle\rangle) = \dim(\langle\langle c, b \rangle\rangle) = 2$. From Lemma 5 it follows that $\{(a, b), (a, c), (b, c)\} \subseteq \{0, 1\}$. Moreover, if $(a, b) = 0$, then $\langle\langle a, b \rangle\rangle \simeq \mathbb{F} \oplus \mathbb{F}$. Therefore, if $(a, b) = (a, c) = (b, c) = 0$ then $A \simeq \mathbb{F} \oplus \mathbb{F} \oplus \mathbb{F}$ and $\psi = 0$. In this case the Gram matrix of the algebra A is the identity matrix and hence the radical of A is trivial. We conclude that in this case $A \simeq S_3$.

Assume that $(a, c) \neq 0$. We have $(a, c) = 1$. In this case, $R(\langle\langle a, c \rangle\rangle)$ is not trivial and contains the element $a - c$. Assume that $(a, b) = (b, c) = 0$. In this case we have $(a - c, b) = 0$. Consequently $a - c \in R(A)$, which is a contradiction. Therefore, without loss of generality, we can assume that $(b, c) = 1$. If $(a, b) = 1$ then $(a - c, b) = 0$ and consequently $a - c \in R(A)$, which is a contradiction. Therefore $(a, b) = 0$. From the description of 2-generated algebras of Jordan type half we have $ab = 0$, $a = c + a_h$, $b = c + b_h$ where $a_h, b_h \in A_{1/2}(c)$. Therefore $0 = ab = (c + a_h)(c + b_h) = c + 1/2(a_h + b_h) + a_h b_h$, where $c + a_h b_h \in A_0(c) \oplus A_1(c)$ and $a_h + b_h \in A_{1/2}(c)$. Therefore $a_h + b_h = 0$. In particular, $b = a^{r_c}$ and $\dim(A) = 2$.

Lemma 5 implies that in this case $A \simeq JForm_2(\mathbb{F})$. \square

Lemma 9. S_6 is isomorphic to $\mathbb{F} \oplus JForm_2(\mathbb{F})$.

Proof. Let $\langle\langle a, b, c \rangle\rangle \simeq S_6$. We have $(a, c) \notin \{0, 1\}$. Therefore $\langle\langle a, c \rangle\rangle$ is isomorphic to $JForm_2(\mathbb{F})$. From Table 1, it follows that the radical of $A(0, 0, \gamma, 0)$ contains ab and bc . Therefore $ab = bc = 0$ and $S_6 \simeq \langle\langle a, c \rangle\rangle \oplus \langle\langle b \rangle\rangle \simeq \mathbb{F} \oplus JForm_2(\mathbb{F})$. \square

Lemma 10. S_7 is isomorphic to $M_2^+(\mathbb{F})$.

Proof. Assume that $\alpha = \beta = \gamma = 0$. We have $(c^{r_b}, a) = 2\alpha\beta + \gamma - 4\psi = -2$ and $\langle\langle c^{r_b}, a, b \rangle\rangle = \langle\langle a, b, c \rangle\rangle = S_7$. Thus, up to redesignation of the generating elements, we can assume that $\gamma \neq 0$.

Let

$$A = \begin{pmatrix} 1 - \lambda_c & 1 \\ \lambda_c(1 - \lambda_c) & \lambda_c \end{pmatrix}, B = \begin{pmatrix} 1 & 0 \\ \lambda_b & 0 \end{pmatrix}, C = \begin{pmatrix} 1 & \lambda_a \\ 0 & 0 \end{pmatrix},$$

where $\lambda_a, \lambda_b, \lambda_c \in \mathbb{F} \setminus \{0\}$. Consider the following map $f : S_7 \rightarrow M_2^+(\mathbb{F})$, $f(a) = A$, $f(b) = B$, $f(c) = C$. It is easy to see that $\dim L\langle A, B, C, A \circ B \rangle = 4$, so $\langle\langle A, B, C \rangle\rangle = M_2^+(\mathbb{F})$.

A map $(\cdot, \cdot) : M_2^+(\mathbb{F})^2 \rightarrow \mathbb{F}$ such that $(X, Y) = tr(XY) = tr(X \circ Y)$, where $X, Y \in M_2^+(\mathbb{F})$, is a symmetric bilinear form on $M_2^+(\mathbb{F})$ (see [9, Chapter 1.6]). This form associates with the product \circ . Clearly, we have $tr(A \circ A) = tr(B \circ B) = tr(C \circ C) = 1$.

Furthermore, we see that $tr(A \circ B) = 1 - \lambda_a + \lambda_b = \alpha$, $tr(B \circ C) = 1 + \lambda_a \lambda_b = \beta$, $tr(A \circ C) = 1 - \lambda_c + \lambda_a \lambda_c(1 - \lambda_c) = \gamma$ and $tr(A \circ (B \circ C)) = tr(B \circ (A \circ C)) = tr(C \circ (A \circ B)) = \psi = \frac{1}{2}(1 - \alpha - \beta - \gamma)$. If $\alpha \neq 0$ then:

$$\lambda_a = \frac{1}{\alpha(\alpha - 1)}(\psi + \alpha\gamma \mp \sqrt{\psi^2 - \alpha\beta\gamma}), \lambda_b = \frac{1}{\gamma}(\psi + \alpha\gamma \pm \sqrt{\psi^2 - \alpha\beta\gamma}),$$

$$\lambda_c = \pm \frac{1}{\gamma}(\psi + \alpha + \sqrt{\psi^2 - \alpha\beta\gamma}),$$

as the solution of these equations.

*	a	b	c	ab
a	a	*	*	*
b	ab	b	*	*
c	$\frac{1}{2(\alpha-1)}((\gamma-\alpha)a + (\gamma-1)b + (1-\alpha)c + 2(-\gamma+1)ab)$	$\frac{1}{2(\alpha-1)}((\beta-1)a + (\beta-\alpha)b + (1-\alpha)c + 2(-\beta+1)ab)$	c	*
ab	$\frac{1}{2}(a\alpha + ab)$	$\frac{1}{2}(b\alpha + ab)$	$\frac{1}{2(\alpha-1)}((\psi-\alpha)a + (\psi-\alpha)b + (\alpha-\alpha^2)c + (2-\beta-\gamma)ab)$	$\frac{1}{4}\alpha(a + b + 2ab)$

Table 2: Multiplication table for S_7

If $\alpha = 0$ then:

$$\lambda_a = -\frac{\gamma^2 - \gamma}{-1 + \beta + \gamma}, \lambda_b = -\frac{-1 + \beta + \gamma}{\gamma}, \lambda_c = \frac{1 - \beta}{\gamma}$$

as the solution of these equations. Note that in this case $-1 + \beta + \gamma \neq 0$, otherwise $\psi = \alpha\beta\gamma = 0$.

Using computer calculations, we show that multiplication table for $f(\langle\langle a, b, c \rangle\rangle)$ ¹ coincides with multiplication table for S_7 . Hence, f is an isomorphism.

We also use computer calculations to check that $R(f(\langle\langle a, b, c \rangle\rangle)) = \{0\}$ and relations between $\alpha, \beta, \gamma, \psi$ hold². \square

Proposition 4. *If A is a 4-dimensional 3-generated algebra of Jordan type half with a trivial radical, then one of the following assertions holds:*

1. $A \simeq S_6 \simeq \mathbb{F} \oplus JForm_2(\mathbb{F})$;
2. $A \simeq S_7 \simeq M_2^+(\mathbb{F})$.

Proof. The algebra $M_2^+(\mathbb{F})$ is a simple Jordan algebra (see [9, Chapter 1.11]). Algebra $\mathbb{F} \oplus JForm_2(\mathbb{F})$ contains non-trivial ideals. Therefore $M_2^+(\mathbb{F}) \not\cong \mathbb{F} \oplus JForm_2(\mathbb{F})$. Hence, to prove this proposition, it suffices to show that $S_6 \simeq \mathbb{F} \oplus JForm_2(\mathbb{F})$ and $S_7 \simeq M_2^+(\mathbb{F})$. Thus the Proposition follows from Lemmas 8 and 9. \square

Lemma 11. $S_8 \cong H(M_3(\mathbb{F}), j)$.

¹Computer calculations for multiplication table in S_7 can be found in <https://github.com/RavilBildanov/3gen-axial-algebras/blob/main/S7multiplicationtable.nb>, see paragraph Tables.

²One can find our computer calculations here: [https://github.com/RavilBildanov/3gen-axial-algebras/blob/main/M2+\(S7\).nb](https://github.com/RavilBildanov/3gen-axial-algebras/blob/main/M2+(S7).nb)

Proof. Consider the following matrices in $H(M_3(\mathbb{F}), j)$ and map $f : S_8 \rightarrow H(M_3(\mathbb{F}), j)$, $f(a) = A, f(b) = B, f(c) = C$, $\lambda_a, \lambda_b, \lambda_c \in \mathbb{F} \setminus \{0\}$ are the invariant by θ parameters which are defined later from conditions to α, β, γ and ψ .

$$A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} B = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \frac{1+\sqrt{1-4\lambda_b^2}}{2} & \lambda_b \\ 0 & \lambda_b & \frac{1-\sqrt{1-4\lambda_b^2}}{2} \end{pmatrix} C = \begin{pmatrix} \frac{1+\sqrt{1-4\lambda_c^2}}{2} & 0 & \lambda_c \\ 0 & 0 & 0 \\ \lambda_c & 0 & \frac{1-\sqrt{1-4\lambda_c^2}}{2} \end{pmatrix}$$

Below we show that the mapping f is an isomorphism between the algebras S_8 and $H(M_3(\mathbb{F}), j)$.

It is easy to see that $A^2 = A, B^2 = B, C^2 = C$. We check that $f(\langle\langle a, b, c \rangle\rangle) = L\langle A, B, C, A \circ C, B \circ C, A \circ (B \circ C) \rangle$. Thus, $\dim L\langle A, B, C, A \circ C, B \circ C, A \circ (B \circ C) \rangle = 6$. Hence $\langle\langle A, B, C \rangle\rangle$ and $H(M_3(\mathbb{F}), j)$ are isomorphic as vector spaces.

A map $(\cdot, \cdot) : H(M_3(\mathbb{F}), j)^2 \rightarrow \mathbb{F}$ such that $(X, Y) = \text{tr}(XY) = \text{tr}(X \circ Y)$, where $X, Y \in H(M_3(\mathbb{F}), j)$ is a symmetric bilinear form on $H(M_3(\mathbb{F}), j)$ [9, Chapter 1.6]. This form associates with the product \circ . Clearly, we have $\text{tr}(A \circ A) = \text{tr}(B \circ B) = \text{tr}(C \circ C) = 1$. Furthermore, we see that $\text{tr}(A \circ B) = 0, \text{tr}(B \circ C) = \frac{1}{4}(1 - \sqrt{1 - 4\lambda_b^2})(1 - \sqrt{1 - 4\lambda_c^2}) = \beta, \text{tr}(A \circ C) = \frac{1+\sqrt{1-4\lambda_c^2}}{2} = \gamma$ and $\text{tr}(A \circ (B \circ C)) = \text{tr}(B \circ (A \circ C)) = \text{tr}(C \circ (A \circ B)) = 0$. So, we have conditions to λ_b, λ_c .

Take the basis $a, b, c, b \cdot c, a \cdot c, a \cdot (b \cdot c)$ for S_8 . Multiplication table for $f(\langle\langle a, b, c \rangle\rangle)$ ³ coincides with multiplication table for S_8 .

We also use computer calculations to check that $R(f(\langle\langle a, b, c \rangle\rangle)) = \{0\}$ and relations between $\alpha, \beta, \gamma, \psi$ hold⁴.

We have that $A, B, C, B \circ C, A \circ C, A \circ (B \circ C)$ is a basis of the algebra $H(M_3(\mathbb{F}), j)$ and hence the kernel of f is trivial. Thus f is an isomorphism of the algebras S_8 and $H(M_3(\mathbb{F}), j)$. \square

Lemma 12. *Algebras S_8 and S_9 are isomorphic.*

Proof. Assume that $\alpha = \beta = \gamma = 1$, but then $\psi = 1$ and we obtain a contradiction. Let $\alpha \neq 1$, take $d = x_a(b) = \frac{2ab - \alpha a - b}{\alpha - 1}$. It is known from [4], that d is a primitive idempotent in S_9 with $ad = 0$ and so d is an axis because S_9 is a Jordan algebra.

Assume that $(c, d) \neq 0$. It can be proved via computer calculations that $a, c, d, ac, cd, a(cd)$ is an additive basis of $B = \langle\langle a, c, d \rangle\rangle$. In particular $B = S_9$. Define a homomorphism f from S_9 to S_8 : $f(a) = \bar{a}, f(d) = \bar{b}, f(c) = \bar{c}$. We have $(\bar{a}, \bar{b}) = 0, (\bar{a}, \bar{b}\bar{c}) = 0, (\bar{a}, \bar{c}) = \bar{\beta} \neq 0, (\bar{b}, \bar{c}) = \bar{\gamma} \neq 0$. Relation $\bar{\beta} + \bar{\gamma} = \frac{2\psi - \beta - \alpha\gamma}{\alpha - 1} + \gamma \neq 1$ is equivalent to $\alpha + \beta + \gamma \neq 2\psi + 1$, so f is an isomorphism between S_8 and S_9 iff $(c, d) \neq 0$.

³Computer calculations for the multiplication table in S_8 can be found in <https://github.com/RavilBildanov/3gen-axial-algebras/blob/main/S8multiplicationtable.nb>, see paragraph Tables.

⁴Computer calculations for this proof can be found in [https://github.com/RavilBildanov/3gen-axial-algebras/blob/main/H3+\(S8\).nb](https://github.com/RavilBildanov/3gen-axial-algebras/blob/main/H3+(S8).nb)

*	a	b	c	bc	ac	$a(bc)$
a	a	*	*	*	*	*
b	0	b	*	*	*	*
c	ac	bc	c	*	*	*
bc	$a(bc)$	$\frac{1}{2}(b\beta + bc)$	$\frac{1}{2}(c\beta + bc)$	$\frac{\beta}{4}(b + c + 2bc)$	*	*
ac	$\frac{1}{2}(\gamma a + ac)$	$a(bc)$	$\frac{1}{2}(\gamma c + ac)$	$\frac{\gamma}{4}bc + \frac{\beta}{4}ac + \frac{1}{2}a(bc)$	$\frac{\gamma}{4}(a + c + 2ac)$	*
$a(bc)$	0	$\frac{1}{4}(\beta ac + 2a(bc))$	$\frac{1}{4}(\gamma bc + \beta ac)$	$\frac{\beta\gamma}{8}b + \frac{\beta}{8}ac + \frac{\beta}{4}a(bc)$	$\frac{\beta\gamma}{8}a + \frac{\gamma}{8}bc + \frac{\gamma}{4}a(bc)$	$\frac{\beta\gamma}{16}a + \frac{\beta\gamma}{16}b$

Table 3: Multiplication table for S_8

Now assume that $(c, d) = 0$. We will prove that $\gamma \neq 1$ in this case. We have that

$$\begin{aligned} 0 &= (c, (\alpha - 1)d) = (c, 2ab - \alpha a - b, c) = \\ &= (2ab - b, c) - \alpha = (2ab, c) - (b, c) - \alpha = 2\psi - \beta - \alpha \end{aligned}$$

If $\gamma = 1$, then $2\psi + 1 = \beta + \alpha + \gamma$, a contradiction. Therefore, $\gamma \neq 1$.

Put $d' = \frac{2ac - \gamma a - c}{\gamma - 1}$. Note that $(b, d') \neq 0$. Indeed, if $(b, d') = 0$, we have

$$0 = (b, 2ac - \gamma a - c) = (b, 2ac) - (b, \gamma a) - (b, c) = 2\psi - \gamma\alpha - \beta$$

Using these equalities, we obtain $\alpha = \gamma\alpha$ and so $\gamma = 1$ or $\alpha = 0$, a contradiction.

We then take a, b, d' as the new generating set of S_9 and, using the same computer calculations, prove that $a, b, x, ab, ax, b(ax)$ form an additive basis of $B' = \langle\langle a, b, d' \rangle\rangle$. Define a homomorphism f' from S_9 to S_8 : $f'(a) = \bar{a}', f'(d') = \bar{b}', f'(b) = \bar{c}'$. We have $(\bar{a}', \bar{b}') = 0, (\bar{a}', \bar{b}'\bar{c}') = 0, (\bar{a}', \bar{c}') = \bar{\beta}' \neq 0, (\bar{b}', \bar{c}') = \bar{\gamma}' \neq 0$. Relation $\bar{\beta}' + \bar{\gamma}' = \frac{2\psi - \beta - \alpha\gamma}{\gamma - 1} + \alpha \neq 1$ is equivalent to $\alpha + \beta + \gamma \neq 2\psi + 1$, so f' is an isomorphism between S_8 and S_9 iff $(b, d') \neq 0$.

We can also check that the multiplication table for $a, c, d, ac, cd, a(cd)$ coincides with the multiplication table for the standard basis $a, b, c, bc, ac, a(bc)$ of S_8 . This means that S_9 contains a 6-dimensional subalgebra isomorphic to S_8 ⁵. □

Proposition 5. *If A is a 6-dimensional 3-generated algebra of Jordan type half with a trivial radical, then $A \simeq S_8 \simeq S_9 \simeq H(M_3(\mathbb{F}), j)$, where $j(X) = X^T$.*

Proof. The proposition follows from Lemmas 10 and 11. □

⁵One can see our computer calculations here:

<https://github.com/RavilBildanov/3gen-axial-algebras/blob/main/3genaxialalgebra.nb>, see section "Isomorphism between S_8 and S_9 ".

3 Acknowledgements

The work was supported by the Theoretical Physics and Mathematics Advancement Foundation "BASIS". We are grateful to V. A. Afanasev, who read the draft of this paper and pointed out several errors.

References

- [1] *I. Gorshkov, A. Staroletov*, On primitive 3-generated axial algebras of Jordan type. *J. Algebra*, 563:74–99, 2020.
- [2] *J. I. Hall, F. Rehren, and S. Shpectorov*, Primitive axial algebras of Jordan type. *J. Algebra*, 437:79–115, 2015.
- [3] *J. I. Hall, Y. Segev, and S. Shpectorov*, On primitive axial algebras of Jordan type. *Bull. Inst. Math. Acad. Sin. (N.S.)*, 13(4):397–409, 2018.
- [4] *I. Gorshkov, V. Gubarev*, Quasi-definite axial algebras of Jordan type half, *Sib. Elektron. Mat. Izv.* 20(2):833–846, 2023.
- [5] *I. Gorshkov, A. Mamontov, A. Staroletov*, On Jordan algebras that are factors of Matsuo algebras. *arXiv:2305.10958*.
- [6] *M. Miyamoto*, Griess algebras and conformal vectors in vertex operator algebras, *J. Algebra* 179(2):523–548, 1996.
- [7] *I. Gorshkov, A. Staroletov, S. Shpectorov*, Solid subalgebras in algebras of Jordan type half. *arXiv:2401.16218*.
- [8] *S. Khasraw, J. McInroy, S. Shpectorov*, On the structure of axial algebras, *Trans. Amer. Math. Soc.* 373(3):2135–2156, 2020.
- [9] *K. McCrimmon*, *A Taste of Jordan Algebras*, Springer-Verlag, xxv+562, 2004.

Ravil Bildanov, SOBOLEV INSTITUTE OF MATHEMATICS, NOVOSIBIRSK, RUSSIA;
NOVOSIBIRSK STATE UNIVERSITY, NOVOSIBIRSK, RUSSIA;
E-mail address: `ravilbildanov@gmail.com`

Ilya Gorshkov, NOVOSIBIRSK STATE TECHNICAL UNIVERSITY, NOVOSIBIRSK, RUSSIA;
E-mail address: `ilygor8@gmail.com`