Generalized non-autonomous Cohen-Grossberg neural network model

Ahmed Elmwafy, José J. Oliveira, César M. Silva

Abstract

In the present paper, we investigate both the global exponential stability and the existence of a periodic solution of a general differential equation with unbounded distributed delays. The main stability criterion depends on the dominance of the non-delay terms over the delay terms. The criterion for the existence of a periodic solution is obtained with the application of the coincide degree theorem. We use the main results to get criteria for the existence and global exponential stability of periodic solutions of a generalized higher-order periodic Cohen-Grossberg neural network model with discrete-time varying delays and infinite distributed delays. Additionally, we provide a comparison with the results in the literature and a numerical simulation to illustrate the effectiveness of some of our results.

Keywords: Cohen-Grossberg neural network, Periodic solutions, Global exponential stability, Coincide degree theorem, discrete and distributed delays.

Mathematics Subject Classification System 2020: 34K20, 34K25, 34K60, 92B20.

1 Introduction

In the past decades, due to application in various sciences, delayed functional differential equations have attracted the attention of an increasing number of researchers. In many fields, such as population dynamics, ecology, epidemiology, disease evolution, and neural networks, differential equations with delay have served as models.

As a result of their widespread use in several fields including image and signal processing [20], pattern recognition [35], optimization [30], and content-addressable memory [34], delayed neural networks have had their dynamical behaviours extensively studied [39], [33], [14].

Obtaining results about the convergence characteristics of neural networks is crucial in these applications. To keep the entire network from acting chaotically, convergent dynamics are required. Significantly, the global convergent dynamics imply that every trajectory of the network can converge to some equilibrium state or invariable sets so that, when used as an associative memory, every state in the underlying space can serve as a key to recover certain stored memory. As an outcome, the state space is entirely covered by different basins of the stored memories. Furthermore, the globally convergent dynamics indicate that the neural network algorithm will ensure convergence to an optimal solution from each initial guess when used as an optimization solver [7].

The fact that the connectivity weights, the neuron charging time, and the external inputs change throughout time is another important consideration. Thus, it is relevant to introduce and investigate neural network models that incorporate the temporal structure of neural activities. Among the various neural network models that have been extensively investigated and applied, Cohen-Grossberg which was first introduced and investigated by Cohen and Grossberg [5] by the following system of ordinary differential equations,

$$\frac{dx_i(t)}{dt} = -a_i(x_i(t)) \Big[b_i(x_i(t)) - \sum_{j=1}^n c_{ij} f_j(x_j(t)) + I_i(t) \Big], \ t \ge 0, \ i = 1, \dots, n$$
(1.1)

where n is a natural number indicates the number of neurons, $x_i(t)$ is the *i*th neuron state at time t, $a_i(u)$ denote the amplification functions, $b_i(u)$ are the self-signal functions, $f_j(u)$ are the activation functions, c_{ij} represent the strengths of connectivity between neurons *i* and *j*, I_i denote the inputs from outside of the system.

Differential equations modelling neural networks should include time delays due to synaptic transmission time across neurons or, in artificial neural networks, communication time among amplifiers in order to be more realistic.

Since Cohen and Grossberg first proposed the CGNN model [5], the dynamical properties of CGNNs such as stability, instability, and periodic oscillation have been extensively studied for theoretical and application considerations. Some studies have already accomplished several positive results such as [1], [4], [19], [17], [6] and etc., most of the results in the literature require either the boundedness of the activation functions or the boundedness of delays. For example, [3] investigated the global exponential stability of the periodic solutions of delayed CGNNs but in the case of discontinuous activation functions. Besides that the existence, uniqueness and stability of almost periodic solutions for a class of NNs have been studied in [31]. Meanwhile, [24], [38], and [23] started studying the existence and exponential stability of high-order CGNNs depending on many techniques. For example, [24] and [38] used some differential inequality techniques, and [23] depended on using a proper Lyapunov function and the properties of M-matrix. Therefore, the present work is meaningful and the conclusion is novel.

Since as far as we know, there are few results on high-order CGNNs without using the Lyapunov technique, neither assuming the boundedness nor the discontinuity of the activation functions. Motivated by the proceeding studies, we consider a generalized high-order CGNN model with discrete time-varying and distributed delays to study the existence of periodic solutions and global exponential stability without using the Lyapunov technique nor the boundedness of the activation functions.

In this paper, we use the continuation theorem of coincidence degree theory to show the existence of a periodic solution of a generalized system of high-order CGNNs, and then we present sufficient conditions to guarantee the global exponential stability of that system. The remainder of this work is organized as follows. Section 2 is a preliminary section where we introduce our notations and our hypotheses. Section 4 introduces the global exponential stability of general neural network models. In Section 3, we investigate the existence and global exponential stability of the periodic solution of that generalized high-order CGNNs system under certain assumptions. In section 5, We show numerical simulations to demonstrate the efficacy of the results we have obtained.

2 Preliminaries and model description

In the present paper, for $n \in \mathbb{N}$, we consider the n-dimensional vector space \mathbb{R}^n equipped with the norm $|x| = \max\{|x_i|, i = 1, ..., n\}$.

For a positive real number ϵ , we consider the Banach space

$$UC_{\epsilon}^{n} = \left\{ \phi \in C((-\infty, 0]; \mathbb{R}^{n}) : \sup_{s \le 0} \frac{|\phi(s)|}{e^{-\epsilon s}} < +\infty, \frac{\phi(s)}{e^{-\epsilon s}} \text{ is uniformly continuous on } (-\infty, 0] \right\},$$

equipped with the norm $\|\phi\|_{\epsilon} = \sup_{s \le 0} \frac{|\phi(s)|}{e^{-\epsilon s}}.$

In [16], a basic theory about the existence, uniqueness, and continuation solutions is established for the general functional differential equation in the phase space UC_{ϵ}^{n}

$$x'(t) = f(t, x_t), \quad t \ge 0,$$
 (2.1)

where, for an open set $D \subseteq UC_{\epsilon}^n$, the function $f: [0, +\infty) \times D \to \mathbb{R}^n$ is continuous and x_t denotes the function $x_t: (-\infty, 0] \to \mathbb{R}^n$ defined by $x_t(s) = x(t+s)$ for $s \leq 0$.

We denote by $x(t, t_0, \phi)$ a solution of (2.1) with initial condition $x_{t_0} = \phi$ for $t_0 \ge 0$ and $\phi \in D$.

For $x \in \mathbb{R}^n$, we also use x to denote the constant function $\phi(s) = x$ in UC_{ϵ}^n . A vector $x \in \mathbb{R}^n$ is said to be positive if $x_i > 0$ for all $i = 1, \ldots, n$ and we denote it by x > 0.

Now, we introduce the Banach space BC of all continuous bounded functions $\phi : (-\infty, 0] \to \mathbb{R}^n$ equipped with the norm $\|\phi\| = \sup_{s \leq 0} |\phi(s)|$. It is clear that $BC \subseteq UC^n_{\epsilon}$ and we have $\|\phi\|_{\epsilon} \leq \|\phi\|$ for all $\phi \in BC$.

In the phase space UC_{ϵ}^n , for $n \in \mathbb{N}$ and $\epsilon > 0$, we consider the following general nonautonomous differential system with infinite delays,

$$x'_{i}(t) = a_{i}(t, x_{i}(t)) \Big[-b_{i}(t, x_{i}(t)) + f_{i}(t, x_{t}) \Big], \quad t \ge 0, \ i = 1, \dots, n,$$
(2.2)

where $a_i : [0, +\infty) \times \mathbb{R} \to (0, \infty), b_i : [0, +\infty) \times \mathbb{R} \to \mathbb{R}$, and $f_i : [0, +\infty) \times UC^n_{\epsilon} \to \mathbb{R}$ are continuous functions.

The goal is to apply the results to Cohen-Grossberg neural network-type models, thus we only consider bounded initial conditions. i.e.

$$x_{t_0} = \phi, \quad \text{for } \phi \in BC \text{ and } t_0 \ge 0.$$
 (2.3)

The continuity of a_i , b_i , and f_i functions assures that the initial value problem (2.2)-(2.3) has a solution (see [12, Theorem 2.1]).

As we always consider bounded initial conditions, in this paper we consider the following definition of global exponential stability.

Definition 2.1. The system (2.2) is said to be globally exponentially stable if there are $\delta > 0$ and $C \ge 1$ such that

$$|x(t, t_0, \phi) - x(t, t_0, \psi)| \le C e^{-\delta(t - t_0)} \|\phi - \psi\|, \quad \forall t_0 \ge 0, \, \forall t \ge t_0, \, \forall \phi, \psi \in BC.$$

It should be emphasized that the preceding definition of global exponential stability is the usually used one in the literature on neural networks ([40], [37], [37]).

3 Global exponential stability

In this section, we obtain sufficient conditions for the global exponential stability of (2.2). To do that in this section we assume the following hypotheses.

For each $i = 1, \ldots, n$:

(H1) there are $\underline{a}_i, \overline{a}_i > 0$ such that

$$\underline{a}_i < a_i(t, u) < \overline{a}_i, \quad \forall t \ge 0, \, \forall u \in \mathbb{R};$$

(H2) there exists a continuous function $D_i: [0, +\infty) \to \mathbb{R}$ such that

$$D_i(t)a_i^2(t,u) \le \frac{\partial a_i}{\partial t}(t,u), \quad \forall t > 0, \, \forall u \in \mathbb{R};$$

(H3) there exists a function $\beta_i : [0, +\infty) \to (0, +\infty)$ such that

$$\frac{b_i(t,u) - b_i(t,v)}{u - v} \ge \beta_i(t), \quad \forall t \ge 0, \, \forall u, v \in \mathbb{R}, \, u \neq v;$$

(H4) the function $f_i : [0, +\infty) \times UC^n_{\epsilon} \to \mathbb{R}$ is Lipschitz on its second variable i.e., there is a continuous function $\mathcal{L}_i : [0, +\infty) \to [0, +\infty)$ such that

$$|f_i(t,\phi) - f_i(t,\psi)| \le \mathcal{L}_i(t) ||\phi - \psi||_{\epsilon}, \quad \forall t \ge 0, \, \forall \phi, \psi \in UC_{\epsilon}^n;$$

(H5) for all $t \ge 0$,

$$\underline{a}_i \Big(\beta_i(t) + D_i(t) \Big) - \overline{a}_i \mathcal{L}_i(t) > \epsilon.$$
(3.1)

By the generalized Gronwall's inequality [13, Lemma 6.2] and the Continuation Theorem [12, Theorem 2.4], we can assure that the solutions of the initial value problem (2.2)-(2.3) are defined on \mathbb{R} .

Now, we are in a position the obtain the main stability criterion for system (2.2).

Theorem 3.1. If (H1)-(H5) hold, then the system (2.2) is globally exponentially stable.

Proof. Let $t_0 > 0$, $\phi = (\phi_1, \ldots, \phi_n) \in BC$, $\psi = (\psi_1, \ldots, \psi_n) \in BC$, and consider two solutions, $x(t) = x(t, t_0, \phi)$ and $y(t) = x(t, t_0, \psi)$, of (2.2).

For each $t \ge t_0$, define $V(t) = V(t, t_0, x(\cdot), y(\cdot)) = (V_i(t), \dots, V_n(t)) \in \mathbb{R}^n$ by

$$V_i(t) := e^{\epsilon(t-t_0)} sign\left(x_i(t) - y_i(t)\right) \int_{y_i(t)}^{x_i(t)} \frac{1}{a_i(t,u)} du, \quad i = 1, \dots, n.$$
(3.2)

From (H1), we conclude that

$$e^{-\epsilon(t-t_0)}\underline{a}_i V_i(t) \le |x_i(t) - y_i(t)| \le e^{-\epsilon(t-t_0)}\overline{a}_i V_i(t), \quad \forall t \ge t_0, \ i = 1, \dots, n.$$
(3.3)

Firstly, we show that

$$|V(t)| \le \max_{i} \{\underline{a}_{i}^{-1}\} \|\phi - \psi\|, \quad \forall t \ge t_{0}.$$
 (3.4)

Obviously, from (3.3), we have

$$|V(t_0)| \le \max_i \left\{ \underline{a}_i^{-1} |x_i(t_0) - y_i(t_0)| \right\} \le \max_i \{ \underline{a}_i^{-1} \} \| \phi - \psi \|.$$

Now, to obtain a contradiction, we assume that inequality (3.4) is false. Consequently, there exists $t_1 > t_0$ such that

$$|V(t_1)| > \max_i \{\underline{a}_i^{-1}\} \|\phi - \psi\|.$$

Define

$$T := \min\left\{t \in [t_0, t_1] : V(t) = \max_{s \in [t_0, t_1]} |V(s)|\right\}.$$

Choosing $i \in \{1, \ldots, n\}$ such that $V_i(T) = |V(T)|$, we have

$$V_i(T) > 0, \quad V'_i(T) \ge 0, \quad \text{and} \quad V_i(T) > |V(t)|, \ \forall t < T.$$
 (3.5)

From (2.2), and (H2), (H3), and (H4), we obtain

$$\begin{split} V_{i}'(T) &= \epsilon V_{i}(T) + \mathrm{e}^{\epsilon(T-t_{0})} sign \Big(x_{i}(T) - y_{i}(T) \Big) \left[\frac{1}{a_{i}(T, x_{i}(T))} x_{i}'(T) \\ &- \frac{1}{a_{i}(T, y_{i}(T))} y_{i}'(T) + \int_{y_{i}(T)}^{x_{i}(T)} - \frac{\frac{\partial a_{i}}{\partial t}(T, u)}{a_{i}^{2}(T, u)} du \right] \\ &= \epsilon V_{i}(T) + \mathrm{e}^{\epsilon(T-t_{0})} sign \Big(x_{i}(T) - y_{i}(T) \Big) \Big[b_{i}(T, y_{i}(T)) - b_{i}(T, x_{i}(T)) \\ &+ f_{i}(T, x_{T}) - f_{i}(T, y_{T}) + \int_{y_{i}(T)}^{x_{i}(T)} - \frac{\partial_{t}a_{i}(T, u)}{a_{i}^{2}(T, u)} du \Big] \\ &\leq \epsilon V_{i}(T) + \mathrm{e}^{\epsilon(T-t_{0})} \Big[- \beta_{i}(T) |x_{i}(T) - y_{i}(T))| + \mathcal{L}_{i}(T) ||x_{T} - y_{T}||_{\epsilon} \\ &- D_{i}(T) |x_{i}(T) - y_{i}(T)| \Big]. \end{split}$$

Hypothesis (H5) implies $\beta_i(T) + D_i(T) > 0$, and from (3.3), we obtain

$$V_{i}'(T) \leq \epsilon V_{i}(T) - \underline{a}_{i} \Big[\beta_{i}(T) + D_{i}(T) \Big] V_{i}(T) + e^{\epsilon(T-t_{0})} \mathcal{L}_{i}(T) \max \Big\{ \sup_{s \leq t_{0} - T} |x(T+s) - y(T+s)| e^{\epsilon s}, \sup_{t_{0} - T < s \leq 0} |x(T+s) - y(T+s)| e^{\epsilon s} \\ \leq \epsilon V_{i}(T) - \underline{a}_{i} \Big[\beta_{i}(T) + D_{i}(T) \Big] V_{i}(T) + e^{\epsilon(T-t_{0})} \mathcal{L}_{i}(T) \max \Big\{ \|\phi - \psi\| e^{\epsilon(t_{0} - T)}, \sup_{t_{0} - T < s \leq 0} |x(T+s) - y(T+s)| e^{\epsilon s} \Big\}.$$

By (3.3), we obtain

$$V_{i}'(T) \leq \epsilon V_{i}(T) - \underline{a}_{i} \Big[\beta_{i}(T) + D_{i}(T) \Big] V_{i}(T) + e^{\epsilon(T-t_{0})} \mathcal{L}_{i}(T) \max \Big\{ \| \phi - \psi \| e^{\epsilon(t_{0}-T)}, \sup_{t_{0}-T < s \leq 0} e^{-\epsilon(T+s-t_{0})+\epsilon s} \overline{a}_{i} V_{i}(T+s) \Big\} = \epsilon V_{i}(T) - \underline{a}_{i} \Big[\beta_{i}(T) + D_{i}(T) \Big] V_{i}(T) + \overline{a}_{i} \mathcal{L}_{i}(T) \max \Big\{ \frac{\| \phi - \psi \|}{\overline{a}_{i}}, \sup_{t_{0}-T < s \leq 0} V_{i}(T+s) \Big\}.$$

By (H1), the definition of T, and (3.5), we have

$$V_i'(T) \le \epsilon V_i(T) - \underline{a}_i \Big[\beta_i(T) + D_i(T) \Big] V_i(T) + \overline{a}_i \mathcal{L}_i(T) V_i(T).$$

From (3.5) and (H5), we conclude that

$$V_i'(T) \le \left[\epsilon - \underline{a}_i \left(\beta_i(T) + D_i(T)\right) + \overline{a}_i \mathcal{L}_i(T)\right] V_i(T) < 0,$$

which contradicts (3.5) and hence (3.4) holds.

From (3.3) and (3.4), we obtain

$$|x(t) - y(t)|e^{\epsilon(t-t_0)}\min\left\{\overline{a}_i^{-1}\right\} \le |V(t)| \le \max_i \{\underline{a}_i^{-1}\} \|\phi - \psi\|,$$

thus

$$|x(t) - y(t)| \le Ce^{-\epsilon(t-t_0)} \|\phi - \psi\|, \quad \forall t \ge t_0,$$

with $C = \frac{\max_i \{\underline{a}_i^{-1}\}}{\min_i \{\overline{a}_i^{-1}\}} = \frac{\max_i \{\overline{a}_i\}}{\min_i \{\underline{a}_i\}} \ge 1$, which shows that the system (2.2) is globally exponentially stable.

We remark that hypothesis (H2) trivially holds (with $D_i(t) = 0$ for all t > 0) in case of all functions a_i do not explicitly depend on time t, i.e. $a_i(t, u) = a_i(u)$ for all i = 1, ..., n and $u \in \mathbb{R}$. Thus, under the assumption

(h5) For all $t \ge 0$ and i = 1, ..., n, we have $\underline{a}_i \beta_i(t) - \overline{a}_i \mathcal{L}_i(t) > \epsilon$,

we have the following result for system

$$x'_{i}(t) = a_{i}(x_{i}(t)) \Big[-b_{i}(t, x_{i}(t)) + f_{i}(t, x_{t}) \Big], \quad t \ge 0, \ i = 1, \dots, n.$$
(3.6)

Corollary 3.2. Assume (H1), (H3), (H4), and (h5) hold. Then, system (3.6) is globally exponentially stable.

Proof. Hypothesis (H2) holds with D(t) = 0, thus the result comes from Theorem 3.1.

Now consider the model studied in [29]

$$x'(t) = a_i(t, x_i(t)) \left[-b_i(t, x_i(t)) + \sum_{k=1}^K \sum_{j=1}^n f_{ijk}(t, x_{j_t}) \right], \quad t \ge 0, \ i = 1, \dots, n,$$
(3.7)

where $n, K \in \mathbb{N}$, a_i and b_i are functions as in system (2.2) and $f_{ijk} : [0, +\infty) \times UC_{\epsilon}^1 \to \mathbb{R}$ are continuous functions for i, j = 1, ..., n and k = 1, ..., K.

We will also assume the following conditions:

(h4) for each i, j = 1, ..., n and k = 1, ..., K, there exists a continuous function \mathcal{F}_{ijk} : $[0, +\infty) \to [0, +\infty)$ such that

$$|f_{ijk}(t,\varphi) - f_{ijk}(t,\psi)| \le \mathcal{F}_{ijk}(t) \|\varphi - \psi\|_{\epsilon}, \quad \forall t \ge 0, \, \varphi, \phi \in UC^{1}_{\epsilon}.$$

(h5') for all $t \ge 0$ and $i = 1, \ldots, n$, we have

$$\underline{a}_i \Big(\beta_i(t) + D_i(t) \Big) - \overline{a}_i \sum_{k=1}^K \sum_{j=1}^n \mathcal{F}_{ijk}(t) > \epsilon.$$

As system (3.7) is a particular situation of (2.2), the following stability criterion holds.

Corollary 3.3. Assume that (H1), (H2), (H3), (h4) and (h5') hold. Then system (3.7) is globally exponentially stable.

Proof. System (3.7) is a particular situation of (2.2) with

$$f_i(t,\varphi) = \sum_{k=1}^K \sum_{j=1}^n f_{ijk}(t,\varphi_j), \quad \forall t \ge 0, \ \varphi = (\varphi_1, \dots, \varphi_n) \in UC_{\epsilon}^n$$

From (h4), we know that (H4) holds with

$$\mathcal{L}_i(t) = \sum_{k=1}^K \sum_{j=1}^n \mathcal{F}_{ijk}(t), \quad \forall t \ge 0, \ i = 1, \dots, n.$$

Moreover, (H5) reads as (h5'). Thus the results comes from Theorem 3.1.

Remark 3.4. We remark that the exponential stability of (3.7) was proved in [29] under the assumptions (H1), (H2), (H3), (h4), and a condition equivalent to

(h5") for all $t \ge 0$ and $i = 1, \ldots, n$, we have

$$\underline{a}_i \Big(\beta_i(t) + D_i(t) \Big) - \sum_{k=1}^K \sum_{j=1}^n \overline{a}_j \mathcal{F}_{ijk}(t) > \epsilon.$$
(3.8)

We emphasize that conditions (h5') and (3.8) are different, thus Corollary 3.3 presents a new exponential stability criterion for the system (3.7).

4 Existence of periodic solution

In this section, we assume that (2.2) is a periodic system and we establish sufficient conditions for the existence of a periodic solutions.

The existence of a periodic solutions will be proved through Mawhin's Continuation Theorem. Before stating the referred theorem, we need to recall some definitions and facts.

Definition 4.1. Let X and Z two Banach spaces.

A linear mapping L: Dom $L \subseteq X \to Z$ is called a Fredholm mapping of index zero if dim Ker_L = codim Im_L < ∞ and Im_L is closed in Z.

Given a Fredholm mapping of index zero, $L : \text{Dom } L \subseteq X \to Z$, it is well known that there are continuous projectors $P : X \to X$ and $Q : Z \to Z$ such that $\text{Im}_P = \text{Ker}_L$, $\text{Ker}_Q = \text{Im}_L = \text{Im}_{I-Q}$, $X = \text{Ker}_L \oplus \text{Ker}_P$ and $Z = \text{Im}_L \oplus \text{Im}_Q$. It follows that $L|_{\text{Dom }L \cap \text{ker}_P} : \text{Dom }L \cap \text{ker}_P \to \text{Im}_L$ is invertible. We denote the inverse of that map by K_P .

Definition 4.2. Let U be an open bounded subset of X. We say that a continuous mapping $N : \overline{U} \subseteq X \to Z$ is L-compact on \overline{U} if the set $QN(\overline{U})$ is bounded and the mapping $K_P(I-Q)N : \overline{U} \subseteq X \to X$ is compact.

Theorem 4.1 (Mawhin's Continuation Theorem). Let X be a Banach space and $\Omega \subseteq X$ an open bounded set. Suppose $L : \text{Dom } L \subset X \to X$ is a Fredholm operator with zero index and that $N : \overline{\Omega} \to X$ is L-compact on $\overline{\Omega}$. Moreover, assume that all the following conditions are satisfied:

- 1. $Lx \neq \lambda Nx$, $\forall x \in \partial \Omega \cap \text{Dom } L, \lambda \in (0, 1);$
- 2. $QNx \neq 0$, $\forall x \in \partial \Omega \cap \text{Ker } L$;
- 3. $\deg_B\{QN, \Omega \cap \text{Ker } L, 0\} \neq 0$, where \deg_B denotes the Brouwer degree.

Then, the equation Lx = Nx has at least one solution in $\overline{\Omega}$.

For studying the system (2.2) in case of being periodic, the following hypotheses will be considered:

(H1^{*}) For each i = 1, ..., n, there exist $\overline{a}_i, \underline{a}_i > 0$ such that

$$\underline{a}_i < a_i(t, u) < \overline{a}_i \text{ for all } t \ge 0, u \in \mathbb{R};$$

(H2*) There is $\omega > 0$ such that, for each $i = 1, \ldots, n$,

$$a_i(t,u) = a_i(t+\omega, u), \quad b_i(t,u) = b_i(t+\omega, u), \quad f_i(t,\phi) = f_i(t+\omega,\phi)$$

for all $t \ge 0$, $u \in \mathbb{R}$, and $\phi \in BC$;

(H3*) For each i = 1, ..., n, there exist ω -periodic continuous functions $\beta_i, \beta_i^* : [0, +\infty) \to (0, +\infty)$ such that

$$\beta_i(t) \le \frac{b_i(t, u) - b_i(t, v)}{u - v} \le \beta_i^*(t), \quad \forall t \in [0, \omega], \, \forall u, v \in \mathbb{R}, \, u \neq v;$$

(H4*) For each i = 1, ..., n, there exists a ω -periodic continuous function $\mathcal{L}_i : [0, +\infty) \to [0, +\infty)$ such that

$$|f_i(t,\phi) - f_i(t,\psi)| \le \mathcal{L}_i(t) \|\phi - \psi\|, \quad \forall t \in [0,\omega], \, \forall \phi, \psi \in BC;$$

(H5*) For each i = 1, ..., n,

$$\beta_i(t) > \mathcal{L}_i(t), \quad \forall t \in [0, \omega]$$

From (H2^{*}), we conclude that the continuous functions $t \mapsto b_i(t,0)$ and $t \mapsto f_i(t,0)$ are ω -periodic and therefore bounded. From (H3^{*}), we also conclude that β_i are bounded away from zero and β_i^* are bounded.

Defining

$$\underline{\beta}_i := \min_{t \in [0,\omega]} \beta_i(t), \quad \overline{\beta}_i^* := \max_{t \in [0,\omega]} \beta_i^*(t), \quad \overline{b}_i := \max_{t \in [0,\omega]} |b_i(t,0)|, \text{ and } \quad \overline{f}_i := \max_{t \in [0,\omega]} |f_i(t,0)|, \quad (4.1)$$

so that we have $0 < \underline{\beta}_i, \overline{\beta}_i^*$, and $0 \le \overline{b}_i, \overline{f}_i$.

We denote by X the Banach space

$$X = \left\{ \phi \in C(\mathbb{R} : \mathbb{R}^n) : \phi \text{ is } \omega - \text{periodic} \right\},\$$

with the norm $\|\phi\| = \sup_{t \in [0,\omega]} |\phi(t)|$, for $\phi \in X$.

For $\text{Dom}_L = \{\phi \in X : \phi' \in X\} \subseteq X$, define the linear operator $L : \text{Dom}_L \to X$ by

$$L\phi = \phi' \tag{4.2}$$

i.e., for all $t \in \mathbb{R}$ and $\phi(t) = (\phi_1(t), \dots, \phi_n(t)) \in \text{Dom}_L$, we have $(L\phi)(t) = (\phi'_1(t), \dots, \phi'_n(t))$. It is not difficult to show that $\text{Ker}_L \cong \mathbb{R}^n$ and

$$\operatorname{Im}_{L} = \left\{ \phi = (\phi_{1}, \dots, \phi_{n}) \in X : \int_{0}^{\omega} \phi_{1}(t) dt = \dots = \int_{0}^{\omega} \phi_{n}(t) dt = 0 \right\},$$
(4.3)

with Im_L closed in X and dim $\text{Ker}_L = \text{codim} \text{Im}_L = n$, thus L is a Fredhom operator with zero index.

Now, we consider the projection $P: X \to X$ defined by

$$P\phi = \frac{1}{\omega} \int_0^\omega \phi(t) dt = \frac{1}{\omega} \left(\int_0^\omega \phi_1(t) dt, \dots, \int_0^\omega \phi_n(t) dt \right), \quad \forall \phi = (\phi_1, \dots, \phi_n) \in X.$$
(4.4)

The projection P is continuous and, considering $Q\phi = P\phi$, we have $\operatorname{Im}_P = \operatorname{Ker}_L$, $\operatorname{Ker}_Q = \operatorname{Im}_L$, and the operator $L_{|\operatorname{Dom}_L \cap \operatorname{Ker}_P}$: $\operatorname{Dom}_L \cap \operatorname{Ker}_P \to \operatorname{Im}_L$ is invertible and we denote the inverse by K_P . By (4.2) and (4.3), we obtain that $K_p\phi = ((K_P\phi)_1, \cdots, (K_P\phi)_n)$ with

$$(K_P\phi)_i(t) = \int_0^t \phi_i(u) du - \frac{1}{\omega} \int_0^\omega \int_0^u \phi_i(s) ds du, \ \forall \phi = (\phi_1, \dots, \phi_n) \in \operatorname{Im}_L,$$
(4.5)

for i = 1, ..., n.

For a convenient bounded open set $\Omega \subseteq X$, define the function $N : \overline{\Omega} \to X$ by $N\phi = ((N\phi)_1, \ldots, (N\phi)_n)$, where

$$(N\phi)_{i}(t) = a_{i}(t,\phi_{i}(t)) \bigg[-b_{i}(t,\phi_{i}(t)) + f_{i}(t,\phi_{t}) \bigg],$$
(4.6)

for all $t \in \mathbb{R}$, $\phi = (\phi_1, \dots, \phi_n) \in X$, and $i = 1, \dots, n$.

We claim that, from the continuity of a_i , b_i , and f_i , (4.5) and (4.6), we can conclude that, for any $\alpha > 0$, the mapping N is L-compact in the set $\Omega = \{\phi \in X : \|\phi\| < \alpha\}$.

In fact, for any $t \in \mathbb{R}$ and any $x \in X$, we have $||QNx|| \leq \max_{i} \overline{a}_{i} [2\overline{\beta}_{i}^{*}\alpha + \overline{b}_{i} + \overline{f}_{i}]$, and we conclude that QN(X) is bounded, implying that $QN(\overline{\Omega})$ is bounded.

Additionally, we also need to show that the mapping $K_P(I-Q)N$ is compact. To achieve this, we show that for any bounded $V \subseteq \overline{\Omega}$, the set $\overline{K_P(I-Q)N(V)}$ is compact. It is easy to verify that, for any sequence, (ϕ_n) , with $\phi_n \in V$, $n \in \mathbb{N}$, such that $\phi_n \to \phi$, we have, for any $t, t_0 \in \mathbb{R}$,

$$\lim_{n \to +\infty} |K_P(I-Q)N(\phi_n)(t) - K_P(I-Q)N(\phi_n)(t_0)| \\\leq 3 \max_i [\overline{a}_i(2\overline{\beta}_i^*\alpha + \overline{b}_i + \overline{f}_i)] (t-t_0).$$

$$(4.7)$$

and

$$\lim_{n \to +\infty} \|K_P(I-Q)N(\phi_n)\| \le 3\omega \max_i [\overline{a}_i(2\overline{\beta}_i^*\alpha + \overline{b}_i + \overline{f}_i)].$$
(4.8)

Inequality (4.7) shows that the family of functions $\overline{K_P(I-Q)N(V)}$ is equicontinuous and inequality (4.8) shows that the norms of all the functions in the referred family of functions are bounded by the same constant. Ascoli-Arzela theorem allows us to conclude that the set $\overline{K_P(I-Q)N(V)}$ is compact. Thus the mapping $K_P(I-Q)N$ is compact and the claim is proved.

Notice that equation (4.8) only allows us to conclude that

$$\lim_{n \to +\infty} |K_P(I-Q)N(\phi_n)(t)| \le 3\omega \max_i [\overline{a}_i(\overline{b}_i + \overline{f}_i)], \text{ for any } t \in [0, \omega].$$

Thus we are not able to apply directly Ascoli-Arzela's theorem to functions in

$$\overline{K_P(I-Q)N(\overline{\Omega})}.$$

Instead, we must consider the space $\tilde{\Omega} = \{\phi \in C([0, \omega] : \mathbb{R}^n) : \|\phi\| < \alpha\}$ instead of Ω , with the norm defined in the same way. This is not a problem since once we show the compactness property for $\tilde{\Omega}$, the same property holds for Ω , because the functions on Ω are ω -periodic.

In view of (4.6) and (4.2), for $\lambda \in (0, 1)$ and $x(t) = (x_1(t), \dots, x_n(t)) \in X$, the operator equation $Lx = \lambda Nx$ is equivalent to the following equation:

$$x'_{i}(t) = \lambda a_{i}(t, x_{i}(t)) \bigg[-b_{i}(t, x_{i}(t)) + f_{i}(t, x_{t}) \bigg], \quad \forall \lambda \in (0, 1), \, i = 1, \dots, n.$$
(4.9)

Now we are in a position to prove the existence of a periodic solution of the general differential system (2.2).

Theorem 4.2. Suppose that (H1^{*}), (H2^{*}), (H3^{*}), (H4^{*}), and (H5^{*}) hold. Then, system (2.2) has at least one ω -periodic solution.

Proof. Our objective is to apply Theorem 4.1. To accomplish this, it is needed to define a bounded open set $\Omega \subseteq X$ for which the conditions 1., 2., and 3. in Theorem 4.1 hold.

Let $x = x(t) = (x_1(t), \dots, x_n(t))^T$ be an arbitrary ω -periodic solution of equation (4.9). The components $x_i(t)$ of x(t) are all continuously differentiable, thus, for each $i = 1, \dots, n$, there is $t_i \in [0, \omega]$ such that

$$|x_i(t_i)| = \max_{t \in [0,\omega]} |x_i(t)|.$$

Hence $x'_i(t_i) = 0$ for all $i = 1, \ldots, n$.

Choose $i \in \{1, \ldots, n\}$ such that $|x_i(t_i)| = \max_{t \in [0,\omega]} |x(t)|$. Consequently, from (4.9), we have

$$b_i(t_i, x_i(t_i)) = f_i(t_i, x_{t_i}),$$
(4.10)

thus

$$b_i(t_i, x_i(t_i)) - b_i(t_i, 0) + b_i(t_i, 0) = f_i(t_i, x_{t_i}) - f_i(t_i, 0) + f_i(t_i, 0).$$

By $(H3^*)$, $(H4^*)$, and (4.1) we obtain

$$\beta_i(t_i)|x_i(t_i)| - \overline{b}_i \leq \mathcal{L}_i(t_i)||x_{t_i}|| + \overline{f}_i,$$

and, as $||x_{t_i}|| = |x(t_i)| = |x_i(t_i)|$, we get

$$|x_i(t_i)| \left(1 - \frac{\mathcal{L}_i(t_i)}{\beta_i(t_i)}\right) \le \frac{\overline{f}_i + \overline{b}_i}{\beta_i(t_i)},$$

From $(H2^*)$, $(H5^*)$, and (4.1), we can define

$$\overline{\xi} = \max_{j,t} \left\{ \left(1 - \frac{\mathcal{L}_j(t)}{\beta_j(t)} \right)^{-1} \frac{\overline{f} + \overline{b}}{\underline{\beta}} \right\} + 1 > 0,$$
(4.11)

where $\overline{b} = \max_i \overline{b}_i$, $\overline{f} = \max_i \overline{f}_i$, and $\underline{\beta} = \min_i \underline{\beta}_i$, thus we conclude that

$$|x_i(t_i)| < \overline{\xi}. \tag{4.12}$$

Consequently, $||x|| < \overline{\xi}$, and taking

$$\Omega = \left\{ \phi \in X : \|\phi\| < \overline{\xi} \right\},\tag{4.13}$$

we conclude that the first condition of Theorem 4.1 is satisfied.

Now, we prove that the second condition of Theorem 4.1 holds.

Let $x = x(t) = (x_1(t), \ldots, x_n(t))^T \in \partial \Omega \cap \operatorname{Ker}_L$. As $\operatorname{Ker}_L \cong \mathbb{R}^n$, then x(t) is a constant vector in \mathbb{R}^n , i.e. $x(t) = (x_1, \ldots, x_n)$, and by (4.13), we conclude that there is $i \in \{1, \ldots, n\}$ such that $|x_i| = \overline{\xi}$. By (4.4) and (4.6), we have

$$(QNx)_i(t) = (QNx)_i = \frac{1}{\omega} \int_0^\omega a_i(u, x_i) \left[-b_i(u, x_i) + f_i(u, x) \right] du.$$

We claim that

$$|(QNx)_i| > 0. (4.14)$$

By contradiction, we assume that $|(QNx)_i| = 0$. Then there is $t_i^* \in [0, \omega]$ such that

$$b_i(t_i^*, x_i) = f_i(t_i^*, x).$$

Reproducing the same computations above (see how (4.10) implies (4.12)), we conclude that

$$\overline{\xi} = |x_i| < \overline{\xi},$$

which is a contradiction. Consequently, (4.14) holds and the second condition of Theorem 4.1 is proved.

In order to prove the last condition of Theorem 4.1, we consider the continuous function $\Psi : (\Omega \cap \operatorname{Ker}_L) \times [0,1] \to X$ defined by $\Psi(x,\mu) = (\Psi(x,\mu)_1, \ldots, \Psi(x,\mu)_n)$ with

$$\Psi(x,\mu)_i = -\mu \overline{a}_i \overline{\beta}_i^* x_i + (1-\mu)(QNx)_i,$$

for all $x = (x_1, \ldots, x_n) \in \Omega \cap \operatorname{Ker}_L \cong \Omega \cap \mathbb{R}^n$, $\mu \in (0, 1)$, and $i = 1, \ldots, n$. We claim that

$$|\Psi(x,\mu)| \neq 0, \quad \forall x \in (\partial\Omega) \cap \operatorname{Ker}_L, \ \mu \in [0,1].$$
 (4.15)

Consequently, defining $\Phi : \mathbb{R}^n \to \mathbb{R}^n$ by

$$\Phi x = \left(-\overline{a}_1 \overline{\beta}_1^* x_1, \dots, -\overline{a}_n \overline{\beta}_n^* x_n\right), \quad \forall x = (x_1, \dots, x_n) \in \mathbb{R}^n,$$

the homotopy invariance theorem [26] implies that

$$\deg_B \{QN, \Omega \cap \operatorname{Ker}_L, 0\} = \deg_B \{\Phi, \Omega \cap \operatorname{Ker}_L, 0\} \neq 0.$$

Now, it remains to prove that (4.15) holds to conclude the proof.

Let $x = (x_1, \ldots, x_n) \in (\partial \Omega) \cap \operatorname{Ker}_L$ and $\mu \in [0, 1]$. The function x is constant because Ker $\cong \mathbb{R}^n$ and, by (4.13), we conclude that there is $i \in \{1, \ldots, n\}$ such that $|x| = |x_i| = \overline{\xi}$. We claim that

$$|\Psi(x,\mu)_i| \neq 0.$$

By contradiction assume that

$$|\Psi(x,\mu)_i| = 0. \tag{4.16}$$

From (4.4), (4.6), and (4.16), we have

$$-\mu\overline{a}_i\overline{\beta}_i^*x_i + \frac{1-\mu}{\omega}\int_0^\omega a_i(t,x_i)\Big[-b_i(t,x_i) + f_i(t,x)\Big]dt = 0,$$

thus there exists $t_i^{**} \in [0, \omega]$ such that

$$-\mu \overline{a}_i \overline{\beta}_i^* x_i + (1-\mu) a_i(t_i^{**}, x_i) \Big[-b_i(t_i^{**}, x_i) + f_i(t_i^{**}, x) \Big] = 0.$$
(4.17)

Now, we assume that $|x| = x_i = \overline{\xi} > 0$ (the situation $|x| = -x_i = \overline{\xi}$ is analogous).

By condition $(H1^*)$ and $(H3^*)$, we have

$$a_{i}(t_{i}^{**}, x_{i})b_{i}(t_{i}^{**}, x_{i}) = a_{i}(t_{i}^{**}, x_{i})\left[b_{i}(t_{i}^{**}, x_{i}) - b_{i}(t_{i}^{**}, 0)\right] + a_{i}(t_{i}^{**}, x_{i})b_{i}(t_{i}^{**}, 0)$$
$$\leq \overline{a_{i}}\overline{\beta}_{i}^{*}x_{i} + a_{i}(t_{i}^{**}, x_{i})b_{i}(t_{i}^{**}, 0),$$

then

$$a_i(t_i^{**}, x_i)b_i(t_i^{**}, x_i) - \overline{a}_i\overline{\beta}_i^*x_i - a_i(t_i^{**}, x_i)b_i(t_i^{**}, 0) \le 0.$$

Consequently, from (4.17), we have

$$\begin{aligned} -a_{i}(t_{i}^{**}, x_{i})b_{i}(t_{i}^{**}, x_{i}) + (1 - \mu)a_{i}(t_{i}^{**}, x_{i})f_{i}(t_{i}^{**}, x) \\ &\geq \mu \left[a_{i}(t_{i}^{**}, x_{i})b_{i}(t_{i}^{**}, x_{i}) - \overline{a}_{i}\overline{\beta}_{i}^{*}x_{i} - a_{i}(t_{i}^{**}, x_{i})b_{i}(t_{i}^{**}, 0)\right] - a_{i}(t_{i}^{**}, x_{i})b_{i}(t_{i}^{**}, x_{i}) \\ &+ (1 - \mu)a_{i}(t_{i}^{**}, x_{i})f_{i}(t_{i}^{**}, x) \\ &= -\mu\overline{a}_{i}\overline{\beta}_{i}^{*}x_{i} + (1 - \mu)a_{i}(t_{i}^{**}, x_{i}) \left[-b_{i}(t_{i}^{**}, x_{i}) + f_{i}(t_{i}^{**}, x)\right] - \mu a_{i}(t_{i}^{**}, x_{i})b_{i}(t_{i}^{**}, 0) \\ &= -\mu a_{i}(t_{i}^{**}, x_{i})b_{i}(t_{i}^{**}, 0) \\ &\geq a_{i}(t_{i}^{**}, x_{i}) \min \left\{0, -b_{i}(t_{i}^{**}, 0)\right\} \end{aligned}$$

and by $(H1^*)$, we obtain

$$-b_i(t_i^{**}, x_i) + (1-\mu)f_i(t_i^{**}, x) \ge \min\left\{0, -b_i(t_i^{**}, 0)\right\}.$$

Consequently,

$$b_i(t_i^{**}, x_i) - b_i(t_i^{**}, 0) \le |f_i(t_i^{**}, x) - f_i(t_i^{**}, 0)| + \overline{b}_i + \overline{f}_i,$$

recalling that $x_i > 0$, and ||x|| = |x|, from (H3^{*}), (H4^{*}), and (4.1) we have

$$x_i \le \frac{\mathcal{L}_i(t_i^{**})}{\beta_i(t_i^{**})} |x| + \frac{\overline{b}_i + \overline{f}_i}{\underline{\beta}_i}.$$

As $|x| = x_i = \overline{\xi} > 0$, we obtain

$$\overline{\xi} = x_i \le \left(1 - \frac{\mathcal{L}_i(t_i^{**})}{\beta_i(t_i^{**})}\right)^{-1} \frac{\overline{b}_i + \overline{f}_i}{\underline{\beta}_i},$$

and by (4.11) we conclude that

$$\overline{\xi} = x_i \le \left(1 - \frac{\mathcal{L}_i(t_i^{**})}{\beta_i(t_i^{**})}\right)^{-1} \frac{\overline{b}_i + \overline{f}_i}{\underline{\beta}_i} < \overline{\xi},$$

which is a contradiction.

The case when $x_i < 0$ is very similar to the previous one and we present it briefly. From (H1^{*}), (H3^{*}), and (4.1), we obtain

$$a_i(t_i^{**}, x_i)b_i(t_i^{**}, x_i) - \overline{a}_i\overline{\beta}_i^* x_i - a_i(t_i^{**}, x_i)b_i(t_i^{**}, 0) \ge 0$$

and, from $(H1^*)$, $(H5^*)$, and (4.11), we obtain

$$-b_i(t_i^{**}, x_i) + (1-\mu)f_i(t_i^{**}, x) \le \max\left\{0, -b_i(t_i^{**}, 0)\right\}.$$

Therefore,

$$b_i(t_i^{**}, x_i) - b_i(t_i^{**}, 0) \ge -|f_i(t_i^{**}, x) - f_i(t_i^{**}, 0)| - \overline{b}_i - \overline{f}_i.$$

Since $x_i < 0$ and $||x||_{\epsilon} = |x|$, from (H3^{*}), (H4^{*}), (4.1) and taking into account that $|x| = -x_i = -\overline{\xi} < 0$, we obtain

$$x_i \ge \frac{\mathcal{L}_i(t_i^{**})}{\beta_i(t_i^{**})} x_i - \frac{\overline{b}_i + \overline{f}_i}{\underline{\beta}_i}.$$

Using this last equation and (4.11), we conclude that

$$-\overline{\xi} = x_i \ge -\left(1 - \frac{\mathcal{L}_i(t_i^{**})}{\beta_i(t_i^{**})}\right)^{-1} \frac{\overline{b}_i + \overline{f}_i}{\underline{\beta}_i} > -\overline{\xi},$$

and we obtain again a contradiction.

By the stability criteria established in the previous Section, now we are in a position to present the following results.

From Theorems 3.1 and 4.2, we have the following result.

Theorem 4.3. Assume (H1^{*}), (H2^{*}), (H2), (H3^{*}), (H4) with $\mathcal{L}_i \omega$ -periodic continuous functions, (H5^{*}), and (H5). Then the system (2.2) has an ω -periodic solution which is globally exponentially stable.

In the case of $D_i(t) \leq 0$, for all $t \geq 0$ and i = 1, ..., n, hypothesis (H5) implies (H5^{*}), thus the following result is an immediate consequence of Theorem 4.3.

Corollary 4.4. If (H1^{*}), (H2^{*}), (H2) with $D_i(t) \leq 0$, for all $t \geq 0$ and i = 1, ..., n, (H3^{*}), (H4) with $\mathcal{L}_i \omega$ -periodic continuous functions, and (H5) hold, then the system (2.2) has an ω -periodic solution which is globally exponentially stable.

In the particular case of functions a_i that do not explicitly depend on time t, from the Corollary 4.4, we have the following result.

Corollary 4.5. If (H1^{*}), (H3^{*}), (H4) with $\mathcal{L}_i \ \omega$ -periodic continuous functions, and (H5) hold, then system (3.6) has an ω -periodic solution which is globally exponentially stable.

Now, we assume that the system (3.7) is ω -periodic, i.e. the following hypothesis holds: (h1*) There is $\omega > 0$ such that, for each i, j = 1, ..., n and k = 1, ..., K,

$$a_i(t,u) = a_i(t+\omega, u), \quad b_i(t,u) = b_i(t+\omega, u), \quad f_{ijk}(t,\phi) = f_{ijk}(t+\omega,\phi),$$

for all $t \geq 0$, $u \in \mathbb{R}$, and $\phi \in BC$.

From Corollary 3.3, Remark 3.4, and Theorem 4.2, we obtain the next result.

Theorem 4.6. Assume (h1^{*}), (H1^{*}), (H2), (H3^{*}), (h4) with $\mathcal{F}_{ijk} \omega$ -periodic continuous functions, and

$$\beta_i(t) > \sum_{k=1}^K \sum_{j=1}^n \mathcal{F}_{ijk}(t), \quad \forall t \in [0, \omega], \ i = 1, \dots, n.$$

If one of the conditions (h5') or (h5") holds, then the system (3.7) has an ω -periodic solution which is globally exponentially stable.

5 Applications to Cohen-Grossberg neural network models

In this section, we apply the results in Sections 3 and 4 to Cohen-Grossberg type models. As we want to apply it to low-order and high-order models, we consider the following general Cohen-Grossberg model with discrete-time varying and distributed delays.

$$\begin{aligned} x_{i}'(t) &= a_{i}(t, x_{i}(t)) \bigg[-b_{i}(t, x_{i}(t)) + F_{i} \bigg(\sum_{p=1}^{P} \sum_{j,l=1}^{n} c_{ijlp}(t) h_{ijlp} \Big(x_{j}(t - \tau_{ijp}(t)), x_{l}(t - \tilde{\tau}_{ilp}(t)) \Big) \\ &+ G_{i} \bigg(\sum_{q=1}^{Q} \sum_{j,l=1}^{n} d_{ijlq}(t) f_{ijlq} \left(\int_{-\infty}^{0} g_{ijq}(x_{j}(t + s)) d\eta_{ijq}(s), \int_{-\infty}^{0} \tilde{g}_{ilq}(x_{l}(t + s)) d\tilde{\eta}_{ilq}(s) \right) \bigg) \\ &+ I_{i}(t) \bigg], \quad t \ge 0, \quad i = 1, \dots, n, \end{aligned}$$

$$(5.1)$$

where $n, P, Q \in \mathbb{N}$ and $a_i : [0, +\infty) \times \mathbb{R} \to (0, +\infty), b_i : [0, +\infty) \times \mathbb{R} \to \mathbb{R}, c_{ijlp}, d_{ijlq}, I_i : [0, +\infty) \to \mathbb{R}, \tau_{ijp}, \tilde{\tau}_{ilp} : [0, +\infty) \to [0, +\infty), h_{ijlp}, f_{ijlq} : \mathbb{R}^2 \to \mathbb{R}, F_i, G_i, g_{ijq}, \tilde{g}_{ilq} : \mathbb{R} \to \mathbb{R}$ are continuous functions, and $\eta_{ijq}, \tilde{\eta}_{ilq} : (-\infty, 0] \to \mathbb{R}$ are non-decreasing bounded functions such that $\eta_{ijq}(0) - \eta_{ijq}(-\infty) = 1$ and $\tilde{\eta}_{ilq}(0) - \tilde{\eta}_{ilq}(-\infty) = 1$, for each $i, j, l = 1, \ldots, n, p = 1, \ldots, P$, and $q = 1, \ldots, Q$.

Here, we assume the next Lipschitz conditions:

(H4**) For each i, j, l = 1, ..., n, p = 1, ..., P, and q = 1, ..., Q, there are positive numbers $\gamma_{ijlp}^{(1)}, \gamma_{ijlp}^{(2)}, \mu_{ijlq}^{(1)}, \mu_{ijlq}^{(2)}, \xi_{ijq}, \tilde{\xi}_{ilq}, \zeta_i$, and ζ_i such that

$$\begin{aligned} |h_{ijlp}(u_1, u_2) - h_{ijlp}(v_1, v_2)| &\leq \gamma_{ijlp}^{(1)} |u_1 - v_1| + \gamma_{ijlp}^{(2)} |u_2 - v_2| \\ |f_{ijlq}(u_1, u_2) - f_{ijlq}(v_1, v_2)| &\leq \mu_{ijlq}^{(1)} |u_1 - v_1| + \mu_{ijlq}^{(2)} |u_2 - v_2| \end{aligned}$$

for all $u_1, u_2, v_1, v_2 \in \mathbb{R}$, and

$$\begin{aligned} |g_{ijq}(u) - g_{ijq}(v)| &\leq \xi_{ijq} |u - v|, \quad |\tilde{g}_{ilq}(u) - \tilde{g}_{ilq}(v)| \leq \xi_{ilq} |u - v|, \\ |F_i(u) - F_i(v)| &\leq \zeta_i |u - v|, \quad |G_i(u) - G_i(v)| \leq \zeta_i |u - v|, \end{aligned}$$

for all $u, v \in \mathbb{R}$.

Now, we state our main stability criterion for model (5.1).

Theorem 5.1. Assume that (H1)-(H3), (H4**), the functions τ_{ijp} , $\tilde{\tau}_{ijp}$ are bounded, and there exists $\vartheta > 0$ such that

$$\int_{-\infty}^{0} e^{-\vartheta s} d\eta_{ijq}(s) < +\infty, \quad \int_{-\infty}^{0} e^{-\vartheta s} d\tilde{\eta}_{ilq}(s) < +\infty.$$
(5.2)

If there exist $\varepsilon > 0$ and $w = (w_1, \ldots, w_n) > 0$ such that for all $t \ge 0$, and $i = 1, \ldots, n$,

$$\underline{a}_{i}\left(\beta_{i}(t)+D_{i}(t)\right)-\overline{a}_{i}\sum_{j,l=1}^{n}\left[\sum_{p=1}^{P}\zeta_{i}|c_{ijlp}(t)|\left(\frac{w_{j}}{w_{i}}\gamma_{ijlp}^{(1)}+\frac{w_{l}}{w_{i}}\gamma_{ijlp}^{(2)}\right)\right.\\\left.+\sum_{q=1}^{Q}\varsigma_{i}|d_{ijlq}(t)|\left(\frac{w_{j}}{w_{i}}\mu_{ijlq}^{(1)}\xi_{ijq}+\frac{w_{l}}{w_{i}}\mu_{ijlq}^{(2)}\tilde{\xi}_{ilq}\right)\right]>\varepsilon,\quad(5.3)$$

then the model (5.1) is globally exponentially stable.

Proof. With the change of variables $y_i(t) = w_i^{-1} x_i(t)$, model (5.1) is transformed into

$$y_{i}'(t) = a_{i}(t, w_{i}y_{i}(t))w_{i}^{-1} \bigg[-b_{i}(t, w_{i}y_{i}(t)) + I_{i}(t) \\ +F_{i}\bigg(\sum_{p=1}^{P}\sum_{j,l=1}^{n}c_{ijlp}(t)h_{ijlp}\bigg(w_{j}y_{j}(t-\tau_{ijp}(t)), w_{l}y_{l}(t-\tilde{\tau}_{ilp}(t))\bigg)\bigg) + G_{i}\bigg(\sum_{q=1}^{Q}\sum_{j,l=1}^{n}d_{ijlq}(t)) \\ \cdot f_{ijlq}\bigg(\int_{-\infty}^{0}g_{ijq}(w_{j}y_{j}(t+s))d\eta_{ijq}(s), \int_{-\infty}^{0}\tilde{g}_{ilq}(w_{l}y_{l}(t+s))d\tilde{\eta}_{ilq}(s)\bigg)\bigg)\bigg],$$
(5.4)

for $t \ge 0$, and $i = 1, \ldots, n$.

From (5.3), there exists $\nu > 0$ such that

$$\underline{a}_{i}\left(\beta_{i}(t)+D_{i}(t)\right) - \overline{a}_{i}\sum_{j,l=1}^{n} \left[\sum_{p=1}^{P} \zeta_{i}|c_{ijlp}(t)| \left(\frac{w_{j}}{w_{i}}\gamma_{ijlp}^{(1)}+\frac{w_{l}}{w_{i}}\gamma_{ijlp}^{(2)}\right) + \sum_{q=1}^{Q} \zeta_{i}|d_{ijlq}(t)| \left(\frac{w_{j}}{w_{i}}\mu_{ijlq}^{(1)}\xi_{ijq}+\frac{w_{l}}{w_{i}}\mu_{ijlq}^{(2)}\tilde{\xi}_{ilq}\right)\right] (1+\nu) > \nu, \quad (5.5)$$

for all $t \ge 0$ and $i = 1, \ldots, n$.

As τ_{ijp} and $\tilde{\tau}_{ilp}$ are bounded functions, it is possible to define the non-negative real number

$$\tau := \max_{i,j,p} \left(\sup_{t \ge 0} \left\{ \tau_{ijp}(t), \tilde{\tau}_{ijp}(t) \right\} \right).$$

As in the proof of [8, Theorem 4.3], from (5.2), we can conclude that there exists $\alpha \in (0, \vartheta)$ such that

$$\int_{-\infty}^{0} e^{-\alpha s} d\eta_{ijq}(s) < 1 + \nu \quad \text{and} \quad \int_{-\infty}^{0} e^{-\alpha s} d\tilde{\eta}_{ijq}(s) < 1 + \nu, \tag{5.6}$$

for all i, j = 1, ..., n and q = 1, ..., Q.

Let $\epsilon := \min\{\nu, \alpha, \frac{\log(1+\nu)}{\tau+1}\}$ and consider the system (5.4) in the phase space UC_{ϵ}^{n} . Defining, for each $i = 1, \ldots, n$, $\tilde{a}_{i}(t, u) := a_{i}(t, w_{i}u)$, $\tilde{b}_{i}(t, u) = w_{i}^{-1}b_{i}(t, w_{i}u)$, and

$$\begin{split} \tilde{f}_{i}(t,\phi) &:= w_{i}^{-1} F_{i} \bigg(\sum_{p=1}^{P} \sum_{j,l=1}^{n} c_{ijlp}(t) h_{ijlp} \Big(w_{j} \phi_{j}(-\tau_{ijp}(t)), w_{l} \phi_{l}(-\tilde{\tau}_{ilp}(t)) \Big) \bigg) + w_{i}^{-1} I_{i}(t) \\ &+ w_{i}^{-1} G_{i} \bigg(\sum_{q=1}^{Q} \sum_{j,l=1}^{n} d_{ijlq}(t) f_{ijlq} \bigg(\int_{-\infty}^{0} g_{ijq}(w_{j} \phi_{j}(s)) d\eta_{ijq}(s), \int_{-\infty}^{0} \tilde{g}_{ilq}(w_{l} \phi_{l}(s)) d\tilde{\eta}_{ilq}(s) \bigg) \bigg) \end{split}$$

for all $u \in \mathbb{R}$ and $\phi = (\phi_1, \dots, \phi_n) \in UC^n_{\epsilon}$, model (5.4) has the form

$$y'_{i}(t) = \tilde{a}_{i}(t, y_{i}(t)) \Big[-\tilde{b}_{i}(t, y_{i}(t)) + \tilde{f}_{i}(t, y_{t}) \Big], \quad t \ge 0, \ i = 1, \dots, n.$$
(5.7)

For model (5.7), the hypotheses (H1), (H2), and (H3) hold with same constants $\underline{a}_i, \overline{a}_i$ and same functions $D_i(t), \beta_i(t)$.

From Theorem 3.1, the proof is concluded if hypotheses (H4) and (H5) hold.

For $\phi = (\phi_1, \ldots, \phi_n), \psi = (\psi_1, \ldots, \psi_n) \in UC^n_{\epsilon}, t \ge 0$, and $i = 1, \ldots, n$, from (H4^{**}) we have

$$\begin{split} |\tilde{f}_{i}(t,\phi) - \tilde{f}_{i}(t,\psi)| &\leq w_{i}^{-1} \sum_{j,l=1}^{n} \left[\zeta_{i} \sum_{p=1}^{P} |c_{ijlp}(t)| \\ &\cdot \left| h_{ijlp} \left(w_{j}\phi_{j}(-\tau_{ijp}(t)), w_{l}\phi_{l}(-\tilde{\tau}_{ilp}(t)) \right) - h_{ijlp} \left(w_{j}\psi_{j}(-\tau_{ijp}(t)), w_{l}\psi_{l}(-\tilde{\tau}_{ilp}(t)) \right) \right| \\ &+ \varsigma_{i} \sum_{q=1}^{Q} |d_{ijlq}(t)| \left| f_{ijlq} \left(\int_{-\infty}^{0} g_{ijq}(w_{j}\phi_{j}(s)) d\eta_{ijq}(s), \int_{-\infty}^{0} \tilde{g}_{ilq}(w_{l}\phi_{l}(s)) d\tilde{\eta}_{ilq}(s) \right) \right. \\ &- \left. f_{ijlq} \left(\int_{-\infty}^{0} g_{ijq}(w_{j}\psi_{j}(s)) d\eta_{ijq}(s), \int_{-\infty}^{0} \tilde{g}_{ilq}(w_{l}\psi_{l}(s)) d\tilde{\eta}_{ilq}(s) \right) \right| \right] \\ &\leq w_{i}^{-1} \sum_{j,l=1}^{n} \left[\zeta_{i} \sum_{p=1}^{P} |c_{ijlp}(t)| \\ &\cdot \left(\gamma_{ijlp}^{(1)} w_{j} \Big| \phi_{j}(-\tau_{ijp}(t)) - \psi_{j}(-\tau_{ijp}(t)) \Big| + \gamma_{ijlp}^{(2)} w_{l} \Big| \phi_{l}(-\tilde{\tau}_{ilp}(t)) - \psi_{l}(-\tilde{\tau}_{ilp}(t)) \Big| \right) \\ &+ \varsigma_{i} \sum_{q=1}^{Q} |d_{ijlq}(t)| \left(\mu_{ijlq}^{(1)} \Big| \int_{-\infty}^{0} g_{ijq}(w_{j}\phi_{j}(s)) - g_{ijq}(w_{j}\psi_{j}(s)) d\eta_{ijq}(s) \Big| \\ &+ \mu_{ijlq}^{(2)} \Big| \int_{-\infty}^{0} \tilde{g}_{ilq}(w_{l}\phi_{l}(s)) - \tilde{g}_{ilq}(w_{l}\psi_{l}(s)) d\tilde{\eta}_{ilq}(s) \Big| \right) \right]. \end{split}$$

Again from (H4**) and by the monotony of η_{ijq} and $\tilde{\eta}_{ijq}$ we obtain,

$$\begin{split} |\tilde{f}_{i}(t,\phi) - \tilde{f}_{i}(t,\psi)| &\leq \sum_{j,l=1}^{n} \left[\zeta_{i} \sum_{p=1}^{P} |c_{ijlp}(t)| \left(\gamma_{ijlp}^{(1)} \frac{w_{j}}{w_{i}} \Big| \phi_{j}(-\tau_{ijp}(t)) - \psi_{j}(-\tau_{ijp}(t)) \right| \right) \\ &+ \gamma_{ijlp}^{(2)} \frac{w_{l}}{w_{i}} \Big| \phi_{l}(-\tilde{\tau}_{ilp}(t)) - \psi_{l}(-\tilde{\tau}_{ilp}(t)) \Big| \right) \\ &+ \varsigma_{i} \sum_{q=1}^{Q} |d_{ijlq}(t)| \left(\mu_{ijlq}^{(1)} \int_{-\infty}^{0} \xi_{ijq} \frac{w_{j}}{w_{i}} |\phi_{j}(s) - \psi_{j}(s)| d\eta_{ijq}(s) \right) \\ &+ \mu_{ijlq}^{(2)} \int_{-\infty}^{0} \tilde{\xi}_{ilq} \frac{w_{l}}{w_{i}} |\phi_{l}(s) - \psi_{l}(s)| d\tilde{\eta}_{ilq}(s) \right) \end{split}$$
(5.8)

and consequently

$$\begin{split} |\tilde{f}_{i}(t,\phi) - \tilde{f}_{i}(t,\psi)| &\leq \sum_{j,l=1}^{n} \left[\zeta_{i} \sum_{p=1}^{P} |c_{ijlp}(t)| \left(\gamma_{ijlp}^{(1)} \frac{w_{j}}{w_{i}} \frac{|(\phi_{j} - \psi_{j})(-\tau_{ijp}(t))|}{\mathrm{e}^{-\epsilon(-\tau_{ijp}(t))}} \mathrm{e}^{\epsilon\tau_{ijp}(t)} \right) \\ &+ \gamma_{ijlp}^{(2)} \frac{w_{l}}{w_{i}} \frac{|(\phi_{l} - \psi_{l})(-\tilde{\tau}_{ilp}(t))|}{\mathrm{e}^{-\epsilon(-\tilde{\tau}_{iip}(t))}} \mathrm{e}^{\epsilon\tilde{\tau}_{ilp}(t)} \right) \\ &+ \varsigma_{i} \sum_{q=1}^{Q} |d_{ijlq}(t)| \left(\mu_{ijlq}^{(1)} \int_{-\infty}^{0} \xi_{ijq} \frac{w_{j}}{w_{i}} \frac{|(\phi_{j} - \psi_{j})(s)|}{\mathrm{e}^{-\epsilon s}} \mathrm{e}^{-\epsilon s} d\eta_{ijq}(s) \\ &+ \mu_{ijlq}^{(2)} \int_{-\infty}^{0} \tilde{\xi}_{ilq} \frac{w_{l}}{w_{i}} \frac{|(\phi_{l} - \psi_{l})(s)|}{\mathrm{e}^{-\epsilon s}} \mathrm{e}^{-\epsilon s} d\tilde{\eta}_{ilq}(s) \right) \Big] \\ &\leq \sum_{j,l=1}^{n} \left[\zeta_{i} \sum_{p=1}^{P} |c_{ijlp}(t)| \left(\gamma_{ijlp}^{(1)} \frac{w_{j}}{w_{i}} \| \phi - \psi \|_{\epsilon} \mathrm{e}^{\epsilon\tau_{ijp}(t)} + \gamma_{ijlp}^{(2)} \frac{w_{l}}{w_{i}} \| \phi - \psi \|_{\epsilon} \mathrm{e}^{\epsilon\tilde{\tau}_{ilp}(t)} \right) \\ &+ \varsigma_{i} \sum_{q=1}^{Q} |d_{ijlq}(t)| \left(\mu_{ijlq}^{(1)} \int_{-\infty}^{0} \xi_{ijq} \frac{w_{j}}{w_{i}} \| \phi - \psi \|_{\epsilon} \mathrm{e}^{-\epsilon s} d\eta_{ijq}(s) \\ &+ \mu_{ijlq}^{(2)} \int_{-\infty}^{0} \tilde{\xi}_{ilq} \frac{w_{l}}{w_{i}} \| \phi - \psi \|_{\epsilon} \mathrm{e}^{-\epsilon s} d\eta_{ijq}(s) \\ &+ \mu_{ijlq}^{(2)} \int_{-\infty}^{0} \tilde{\xi}_{ilq} \frac{w_{l}}{w_{i}} \| \phi - \psi \|_{\epsilon} \mathrm{e}^{-\epsilon s} d\tilde{\eta}_{ilq}(s) \Big) \Big] \\ &\leq \| \phi - \psi \|_{\epsilon} \sum_{j,l=1}^{n} \left[\zeta_{i} \sum_{p=1}^{P} |c_{ijlp}(t)| \left(\gamma_{ijlp}^{(1)} \frac{w_{j}}{w_{i}} + \gamma_{ijlp}^{(2)} \frac{w_{l}}{w_{i}} \right) \mathrm{e}^{\epsilon\tau} \\ &+ \varsigma_{i} \sum_{q=1}^{Q} |d_{ijlq}(t)| \left(\mu_{ijlq}^{(1)} \xi_{ijq} \frac{w_{j}}{w_{i}} \int_{-\infty}^{0} \mathrm{e}^{-\epsilon s} d\eta_{ijq}(s) + \mu_{ijlq}^{(2)} \tilde{\xi}_{ilq} \frac{w_{l}}{w_{i}} \int_{-\infty}^{0} \mathrm{e}^{-\epsilon s} d\tilde{\eta}_{ilq}(s) \right) \right] \end{aligned}$$

As $\epsilon \leq \alpha$ from (5.6) we have

$$\int_{-\infty}^{0} e^{-\epsilon s} d\eta_{ijq}(s) < 1 + \nu \quad \text{and} \quad \int_{-\infty}^{0} e^{-\epsilon s} d\tilde{\eta}_{ijq}(s) < 1 + \nu,$$

for all i, j = 1, ..., n. As $\epsilon \leq \frac{\log(1 + \nu)}{\tau + 1}$, then we also have

$$e^{\epsilon \tau} < e^{\epsilon(\tau+1)} \le 1 + \nu.$$

Consequently

$$\begin{split} |\tilde{f}_{i}(t,\phi) - \tilde{f}_{i}(t,\psi)| &\leq \left[\sum_{j,l=1}^{n} \left(\zeta_{i} \sum_{p=1}^{P} |c_{ijlp}(t)| \left(\gamma_{ijlp}^{(1)} \frac{w_{j}}{w_{i}} + \gamma_{ijlp}^{(2)} \frac{w_{l}}{w_{i}}\right) \right. \\ &+ \varsigma_{i} \sum_{q=1}^{Q} |d_{ijlq}(t)| \left(\mu_{ijlq}^{(1)} \xi_{ijq} \frac{w_{j}}{w_{i}} + \mu_{ijlq}^{(2)} \widetilde{\xi}_{ilq} \frac{w_{l}}{w_{i}}\right) \right) (1+\nu) \right] \|\phi - \psi\|_{\epsilon}, \end{split}$$

and hypothesis (H4) holds with

$$\mathcal{L}_{i}(t) = \sum_{j,l=1}^{n} \left(\zeta_{i} \sum_{p=1}^{P} |c_{ijlp}(t)| \left(\gamma_{ijlp}^{(1)} \frac{w_{j}}{w_{i}} + \gamma_{ijlp}^{(2)} \frac{w_{l}}{w_{i}} \right) + \varsigma_{i} \sum_{q=1}^{Q} |d_{ijlq}(t)| \left(\mu_{ijlq}^{(1)} \xi_{ijq} \frac{w_{j}}{w_{i}} + \mu_{ijlq}^{(2)} \tilde{\xi}_{ilq} \frac{w_{l}}{w_{i}} \right) \right) (1+\nu)$$
for all $i = 1$ or

for all $i = 1, \ldots, n$.

As $\epsilon \leq \nu$ and from (5.5), the hypothesis (H5) also holds and the proof is concluded. \Box Now, we assume that the model (5.1) is periodic, i.e. there is $\omega > 0$ such that (H2**) There is $\omega > 0$ such that, for each i, j, l = 1, ..., n, p = 1, ..., P, and q = 1, ..., Q,

$$a_i(t, u) = a_i(t + \omega, u), \quad c_{ijlp}(t) = c_{ijlp}(t + \omega), \quad \tau_{ijp}(t) = \tau_{ijp}(t + \omega),$$

$$b_i(t, u) = b_i(t + \omega, u), \quad d_{ijlq}(t) = d_{ijlq}(t + \omega), \quad \tilde{\tau}_{ijp}(t) = \tilde{\tau}_{ijp}(t + \omega), \text{ and}$$

$$I_i(t) = I_i(t + \omega)$$

for all $t \geq 0$ and $u \in \mathbb{R}$.

Theorem 5.2. Assume the hypotheses $(H2^{**})$, (H1), $(H3^{*})$, and $(H4^{**})$.

If there exists $w = (w_1, \ldots, w_n) > 0$ such that for all $t \in [0, \omega]$, and $i = 1, \ldots, n$,

$$\beta_{i}(t) > \sum_{j,l=1}^{n} \left[\sum_{p=1}^{P} \zeta_{i} |c_{ijlp}(t)| \left(\frac{w_{j}}{w_{i}} \gamma_{ijlp}^{(1)} + \frac{w_{l}}{w_{i}} \gamma_{ijlp}^{(2)} \right) + \sum_{q=1}^{Q} \zeta_{i} |d_{ijlq}(t)| \left(\frac{w_{j}}{w_{i}} \mu_{ijlq}^{(1)} \xi_{ijq} + \frac{w_{l}}{w_{i}} \mu_{ijlq}^{(2)} \widetilde{\xi}_{ilq} \right) \right],$$
(5.9)

then the model (5.1) has an ω -periodic solution.

Proof. As in previous proof, the change of variables $y_i(t) = w_i^{-1} x_i(t)$ transforms model (5.1) into (5.4).

Considering model (5.4) in the phase space UC_{ϵ}^{n} , for some $\epsilon > 0$, it has the form (5.7).

Proceeding as in the previous proof, functions \tilde{f}_i verify (5.8) for all $\phi = (\phi_1, \ldots, \phi_n), \psi = (\psi_1, \ldots, \psi_n) \in BC$ and $t \ge 0$. Consequently

$$\begin{split} |\tilde{f}_{i}(t,\phi) - \tilde{f}_{i}(t,\psi)| &\leq \|\phi - \psi\| \sum_{j,l=1}^{n} \left[\zeta_{i} \sum_{p=1}^{P} |c_{ijlp}(t)| \left(\gamma_{ijlp}^{(1)} \frac{w_{j}}{w_{i}} + \gamma_{ijlp}^{(2)} \frac{w_{l}}{w_{i}} \right) \right. \\ &+ \varsigma_{i} \sum_{q=1}^{Q} |d_{ijlq}(t)| \left(\mu_{ijlq}^{(1)} \xi_{ijq} \frac{w_{j}}{w_{i}} \int_{-\infty}^{0} d\eta_{ijq}(s) + \mu_{ijlq}^{(2)} \widetilde{\xi}_{ilq} \frac{w_{l}}{w_{i}} \int_{-\infty}^{0} d\widetilde{\eta}_{ilq}(s) \right) \right], \end{split}$$

and from the properties of η_{ijq} and $\tilde{\eta}_{ijq}$ we obtain

$$|\widetilde{f}_i(t,\phi) - \widetilde{f}_i(t,\psi)| \le \mathcal{L}_i(t) \|\phi - \psi\|,$$

with

$$\mathcal{L}_{i}(t) = \sum_{j,l=1}^{n} \left[\zeta_{i} \sum_{p=1}^{P} |c_{ijlp}(t)| \left(\gamma_{ijlp}^{(1)} \frac{w_{j}}{w_{i}} + \gamma_{ijlp}^{(2)} \frac{w_{l}}{w_{i}} \right) + \varsigma_{i} \sum_{q=1}^{Q} |d_{ijlq}(t)| \left(\mu_{ijlq}^{(1)} \xi_{ijq} \frac{w_{j}}{w_{i}} + \mu_{ijlq}^{(2)} \widetilde{\xi}_{ilq} \frac{w_{l}}{w_{i}} \right) \right],$$

thus $(H4^*)$ holds for model (5.7).

By hypothesis (5.9), (H5^{*}) also holds and the conclusion follows from Theorem 4.2. \Box

Immediately from Theorems 5.1 and 5.2, we have the following result.

Corollary 5.3. Assume (H1), (H2) with D_i an ω -periodic continuous function, (H2^{**}), (H3^{*}), (H4^{**}), and (5.2).

If there exists $w = (w_1, \ldots, w_n) > 0$ such that, for all $t \in [0, \omega]$ and $i = 1, \ldots, n$ inequality (5.9) holds and

$$\underline{a}_{i}(\beta_{i}(t) + D_{i}(t)) > \overline{a}_{i} \sum_{j,l=1}^{n} \left[\sum_{p=1}^{P} \zeta_{i} |c_{ijlp}(t)| \left(\frac{w_{j}}{w_{i}} \gamma_{ijlp}^{(1)} + \frac{w_{l}}{w_{i}} \gamma_{ijlp}^{(2)} \right) + \sum_{q=1}^{Q} \zeta_{i} |d_{ijlq}(t)| \left(\frac{w_{j}}{w_{i}} \mu_{ijlq}^{(1)} \xi_{ijq} + \frac{w_{l}}{w_{i}} \mu_{ijlq}^{(2)} \widetilde{\xi}_{ilq} \right) \right], \quad (5.10)$$

then the model (5.1) has an ω -periodic solution which is globally exponentially stable.

Proof. From (H2^{**}) functions τ_{ijp} , $\tilde{\tau}_{ijp}$ are bounded.

Moreover, from (H2^{**}) and (H3^{*}) we know that β_i , c_{ijlp} , and d_{ijlq} are ω -periodic functions. As D_i are also ω -periodic, then there is $\varepsilon > 0$ such that inequality (5.3) holds and the conclusion comes from Theorems 5.1 and 5.2.

Now, we consider model (5.1) with amplifications functions, a_i , do not explicitly depend on time t, i.e.

$$\begin{aligned} x_{i}'(t) &= a_{i}(x_{i}(t)) \bigg[-b_{i}(t, x_{i}(t)) + F_{i} \bigg(\sum_{p=1}^{P} \sum_{j,l=1}^{n} c_{ijlp}(t) h_{ijlp} \Big(x_{j}(t - \tau_{ijp}(t)), x_{l}(t - \tilde{\tau}_{ilp}(t)) \Big) \bigg) \\ &+ G_{i} \bigg(\sum_{q=1}^{Q} \sum_{j,l=1}^{n} d_{ijlq}(t) f_{ijlq} \left(\int_{-\infty}^{0} g_{ijq}(x_{j}(t + s)) d\eta_{ijq}(s), \int_{-\infty}^{0} \tilde{g}_{ilq}(x_{l}(t + s)) d\tilde{\eta}_{ilq}(s) \right) \bigg) \\ &+ I_{i}(t) \bigg], \quad t \ge 0, \quad i = 1, \dots, n, \end{aligned}$$

$$(5.11)$$

From Corollary 5.3 we have the following result.

Corollary 5.4. Assume (H1), (H2**), (H3*), (H4**), and (5.2).

If there exists $w = (w_1, \ldots, w_n) > 0$ such that, for all $t \in [0, \omega]$ and $i = 1, \ldots, n$,

$$\underline{a}_{i}\beta_{i}(t) > \overline{a}_{i}\sum_{j,l=1}^{n} \left[\sum_{p=1}^{P} \zeta_{i}|c_{ijlp}(t)| \left(\frac{w_{j}}{w_{i}}\gamma_{ijlp}^{(1)} + \frac{w_{l}}{w_{i}}\gamma_{ijlp}^{(2)}\right) + \sum_{q=1}^{Q} \zeta_{i}|d_{ijlq}(t)| \left(\frac{w_{j}}{w_{i}}\mu_{ijlq}^{(1)}\xi_{ijq} + \frac{w_{l}}{w_{i}}\mu_{ijlq}^{(2)}\tilde{\xi}_{ilq}\right)\right],$$
(5.12)

then the model (5.11) has an ω -periodic solution which is globally exponentially stable.

Proof. Noting that $a_i(t, u) = a_i(u)$ for all $t, u \in \mathbb{R}$ and $i = 1, \ldots, n$, the hypothesis (H2) trivially holds with $D_i(t) = 0$. Consequently inequality (5.12) implies (5.9) and the result comes from Corollary 5.3.

For model (5.11) under the hypotheses (H2^{**}), (H1), (H3^{*}), and (H4^{**}), consider the constants

$$\underline{\beta}_i := \min_{t \in [0,\omega]} \beta_i(t), \quad \overline{c}_{ijlp} := \max_{t \in [0,\omega]} c_{ijlp}(t), \quad \text{and} \quad \overline{d}_{ijlq} := \max_{t \in [0,\omega]} d_{ijlq}(t), \tag{5.13}$$

for each i, j = 1, ..., n, p = 1, ..., P, q = 1, ..., Q, and the square real matrix \mathcal{M} defined by

$$\mathcal{M} := diag(\underline{a}_{1}\underline{\beta}_{1}, \dots, \underline{a}_{n}\underline{\beta}_{n}) - [\mathfrak{m}_{ij}]_{i,j=1}^{n}, \qquad (5.14)$$

where, for each $i, j = 1, \ldots, n$,

$$\mathfrak{m}_{ij} := \overline{a}_i \sum_{l=1}^n \left(\zeta_i \sum_{p=1}^P \left(\overline{c}_{ijlp} \gamma_{ijlp}^{(1)} + \overline{c}_{iljp} \gamma_{iljp}^{(2)} \right) + \varsigma_i \sum_{q=1}^Q \left(\overline{d}_{ijlq} \mu_{ijlq}^{(1)} \xi_{ijq} + \overline{d}_{iljq} \mu_{iljq}^{(2)} \widetilde{\xi}_{ijq} \right) \right).$$

Corollary 5.5. Assume (H1), (H2**), (H3*), (H4**), and (5.2).

If \mathcal{M} is a non-singular M-matrix, then the model (5.11) has an ω -periodic solution which is globally exponentially stable.

Proof. As \mathcal{M} is a non-singular M-matrix, then (see [9]) there exists $w = (w_1, \ldots, w_n) > 0$ such that $\mathcal{M}w^T > 0$, i.e.,

$$\underline{a}_{i}\underline{\beta}_{i}w_{i} > \sum_{j=1}^{n} w_{j} \left[\overline{a}_{i} \sum_{l=1}^{n} \left(\zeta_{i} \sum_{p=1}^{P} \left(\overline{c}_{ijlp} \gamma_{ijlp}^{(1)} + \overline{c}_{iljp} \gamma_{iljp}^{(2)} \right) + \varsigma_{i} \sum_{q=1}^{Q} \left(\overline{d}_{ijlq} \mu_{ijlq}^{(1)} \xi_{ijq} + \overline{d}_{iljq} \mu_{iljq}^{(2)} \widetilde{\xi}_{ijq} \right) \right) \right],$$

for all $i = 1, \ldots, n$, which is equivalent to

$$\underline{a}_{i}\underline{\beta}_{i} > \overline{a}_{i}\sum_{j,l=1}^{n} \left[\zeta_{i}\sum_{p=1}^{P} \left(\overline{c}_{ijlp} \frac{w_{j}}{w_{i}} \gamma_{ijlp}^{(1)} + \overline{c}_{iljp} \gamma_{iljp}^{(2)} \frac{w_{j}}{w_{i}} \right) + \varsigma_{i}\sum_{q=1}^{Q} \left(\overline{d}_{ijlq} \frac{w_{j}}{w_{i}} \mu_{ijlq}^{(1)} \xi_{ijq} + \overline{d}_{iljq} \mu_{iljq}^{(2)} \widetilde{\xi}_{ijq} \frac{w_{j}}{w_{i}} \right)$$

and consequently

$$\underline{a}_{i}\underline{\beta}_{i} > \overline{a}_{i}\sum_{j,l=1}^{n} \left[\zeta_{i}\sum_{p=1}^{P} \left(\overline{c}_{ijlp} \frac{w_{j}}{w_{i}} \gamma_{ijlp}^{(1)} + \overline{c}_{ijlp} \gamma_{ijlp}^{(2)} \frac{w_{l}}{w_{i}} \right) + \varsigma_{i}\sum_{q=1}^{Q} \left(\overline{d}_{ijlq} \frac{w_{j}}{w_{i}} \mu_{ijlq}^{(1)} \xi_{ijq} + \overline{d}_{ijlq} \mu_{ijlq}^{(2)} \widetilde{\xi}_{ilq} \frac{w_{l}}{w_{i}} \right) \right].$$

$$(5.15)$$

From (5.13) and (5.15) we obtain (5.12). Now the result follows from Corollary 5.4. \Box

Example 5.1. Consider the following low-order Cohen-Grossberg neural network model

$$x'(t) = a_i(x_i(t)) \left[-b_i(t, x_i(t)) + G_i\left(\sum_{j=1}^n c_{ij}(t) \int_0^{+\infty} x_j(t-u) K_{ij}(u) du\right) \right], \quad (5.16)$$

for $t \ge 0$ and i = 1, ..., n, where $a_i : \mathbb{R} \to (0, +\infty), b_i : [0, +\infty) \times \mathbb{R} \to \mathbb{R}, c_{ij} : [0, +\infty) \to \mathbb{R}, G_i : \mathbb{R} \to \mathbb{R}$, and $K_{ij} : [0, +\infty) \to [0, +\infty)$ are continuous functions such that

$$\int_{0}^{+\infty} K_{ij}(u) du = 1, \qquad (5.17)$$

for all i, j = 1, ..., n.

The model (5.16) is a generalization of the following autonomous static neural network model

$$x'(t) = -x_i(t) + G_i\left(\sum_{j=1}^n c_{ij} \int_0^{+\infty} x_j(t-u) K_{ij}(u) du\right), \ t \ge 0, \ i = 1, \dots, n,$$
(5.18)

whose the existence and global asymptotic stability of an equilibrium point was studied in [27].

Defining, for each $i, j = 1, ..., n, \eta_{ij} : (-\infty, 0] \to \mathbb{R}$ by

$$\eta_{ij}(s) = \int_{-\infty}^{s} K_{ij}(-v)dv, \quad s \in (-\infty, 0]$$
(5.19)

we have η_{ij} non-decreasing and, from (5.17), $\eta_{ij}(0) - \eta_{ij}(-\infty) = 1$. Consequently, the model (5.16) can be written in the form

$$x'(t) = a_i(x_i(t)) \left[-b_i(t, x_i(t)) + G_i\left(\sum_{j=1}^n c_{ij}(t) \int_{-\infty}^0 x_j(t+s) d\eta_{ij}(s)\right) \right], \ t \ge 0, \ i = 1, \dots, n,$$

which is a particular situation of (5.11). Consequently, from Corollary 5.4, we obtain the following result.

Corollary 5.6. Assume (H1), (H3^{*}) and, for each i, j = 1, ..., n, the functions $t \mapsto b_i(t, u)$ and c_{ij} is ω -periodic, the function G_i is Lipschitz with Lipschitz constant $\varsigma_i > 0$, and there is $\alpha > 0$ such that

$$\int_0^{+\infty} K_{ij}(u) \mathrm{e}^{\alpha u} du < +\infty.$$
(5.20)

If there exists $w = (w_1, \ldots, w_n) > 0$ such that, for all $t \in [0, \omega]$ and $i = 1, \ldots, n$,

$$\underline{a}_{i}\beta_{i}(t)w_{i} > \overline{a}_{i}\sum_{j=1}^{n}\varsigma_{i}|c_{ij}(t)|w_{j}, \quad t \ge 0, \ i = 1, \dots, n,$$

$$(5.21)$$

then the model (5.16) has an ω -periodic solution which is globally exponentially stable. For model (5.18), condition (5.21) reads as

$$w_i > \sum_{j=1}^n \varsigma_i |c_{ij}| w_j, \quad t \ge 0, \ i = 1, \dots, n,$$

which is equivalent to matrix

$$\mathcal{N} = I_n - \left[\varsigma_i |c_{ij}|\right]_{i,j=1}^n,$$

where I_n denotes the identity matrix of *n*-dimension, being a non-singular M-matrix (see [Fidler]). Consequently, we also have the following result.

Corollary 5.7. For each i, j = 1, ..., n assume that the function G_i is Lipschitz with Lipschitz constant $\varsigma_i > 0$ and (5.20).

If \mathcal{N} is a non-singular M-matrix, then the model (5.18) has an equilibrium point which is globally exponentially stable.

Remark 5.8. We remark that in [27] the existence and global asymptotic stability of an equilibrium point of (5.18) was obtained assuming stronger conditions over G_i than being Lipschitz, \mathcal{N} be a non-singular M-matrix, and

$$\int_0^{+\infty} u K_{ij}(u) du < +\infty.$$

instead of (5.20).

Example 5.2. Consider the following low-order Cohen-Grossberg neural network model,

$$x'(t) = a_i(t, x_i(t)) \left[-b_i(t, x_i(t)) + \sum_{j=1}^n c_{ij1}(t) h_{ij1}(x_j(t)) + \sum_{j=1}^n c_{ij2}(t) h_{ij2}(x_j(t - \tau_{ij}(t))) + \sum_{j=1}^n d_{ij}(t) \int_0^{+\infty} g_{ij}(x_j(t - u)) K_{ij}(u) du + I_i(t) \right], t \ge 0, i = 1, \dots, n, \quad (5.22)$$

where, for each i, j = 1, ..., n, $a_i : [0, +\infty) \times \mathbb{R} \to (0, +\infty)$, $b_i : [0, +\infty) \times \mathbb{R} \to \mathbb{R}$, $c_{ij1}, c_{ij2}, d_{ij}, I_i : [0, +\infty) \to \mathbb{R}$, $\tau_{ij} : [0, +\infty) \to [0, +\infty)$, $h_{ij1}, h_{ij2}, g_{ij} : \mathbb{R} \to \mathbb{R}$, and $K_{ij} : [0, +\infty) \to [0, +\infty)$ are continuous functions such that K_{ij} verifies (5.17).

Sufficient conditions for the exponential stability of (5.22) were obtained in [22, 29].

The existence and global asymptotic stability of a periodic solution of (5.22), with finite delays, were studied in [15].

The following particular situation of (5.22)

$$x'(t) = a_i(x_i(t)) \left[-b_i(x_i(t)) + \sum_{j=1}^n c_{ij1}(t)h_{j1}(x_j(t)) + \sum_{j=1}^n c_{ij2}(t)h_{j2}(x_j(t-\tau_{ij}(t))) + \sum_{j=1}^n d_{ij}(t) \int_0^{+\infty} g_j(x_j(t-u))K_{ij}(u)du + I_i(t) \right], \ t \ge 0, \ i = 1, \dots, n, \quad (5.23)$$

was studied in [36], where conditions for the existence and global exponential stability of a pseudo almost automorphic solution were established.

Considering the definition of the bounded variation function η_{ij} as in (5.19), model (5.22) is a particular situation of (5.1), thus from Theorem 5.1 we obtain the following stability criterion

Corollary 5.9. Assume that (H1)-(H3), and, for each i, j = 1, ..., n, τ_{ij} is bounded, K_{ij} verifies (5.17) and (5.20), and h_{ij1}, h_{ij2}, g_{ij} are Lipschitz functions with Lipschitz constants $\gamma_{ij1}, \gamma_{ij2}, \xi_{ij} > 0$ respectively.

If there exist $\varepsilon > 0$ and $w = (w_1, \ldots, w_n) > 0$ such that, for all $t \ge 0$ and $i = 1, \ldots, n$,

$$\underline{a}_{i}\left(\beta_{i}(t)+D_{i}(t)\right)w_{i}-\overline{a}_{i}\sum_{j=1}^{n}\left[\left(|c_{ij1}(t)|\gamma_{ij1}+|c_{ij2}(t)|\gamma_{ij2}+|d_{ij}(t)|\xi_{ij}\right)w_{j}\right]>\varepsilon,\quad(5.24)$$

then the model (5.22) is globally exponentially stable.

Remark 5.10. In [29] the exponential stability of (5.22) was established assuming hypotheses in Corollary 5.9 with the condition

$$\underline{a}_{i}\left(\beta_{i}(t)+D_{i}(t)\right)-\sum_{j=1}^{n}\overline{a}_{j}\left[\left(|c_{ij1}(t)|\gamma_{ij1}+|c_{ij2}(t)|\gamma_{ij2}+|d_{ij}(t)|\xi_{ij}\right)\frac{w_{j}}{w_{i}}\right]>\varepsilon,\qquad(5.25)$$

instead of (5.24).

Model (5.23) is a particular situation of (5.11), from Corollaries 5.4 and 5.9 and Remark 5.10 we obtain the following result

Corollary 5.11. Assume (H1), (H3^{*}) with $\beta_i(t) \equiv \beta_i$, and, for each i, j = 1, ..., n, $c_{ij1}, c_{ij2}, \tau_{ij}, d_{ij}, I_i$ are ω -periodic for some $\omega > 0$, K_{ij} verifies (5.17) and (5.20), and h_{j1}, h_{j2}, g_j are Lipschitz functions with Lipschitz constants $\gamma_{j1}, \gamma_{j2}, \xi_j > 0$ respectively.

If there exists $w = (w_1, \ldots, w_n) > 0$ such that one of the following conditions

$$\underline{a}_{i}\beta_{i}w_{i} - \overline{a}_{i}\sum_{j=1}^{n} \left[\left(|c_{ij1}(t)|\gamma_{j1} + |c_{ij2}(t)|\gamma_{j2} + |d_{ij}(t)|\xi_{j} \right) w_{j} \right] > 0,$$
(5.26)

or

$$\underline{a}_{i}\beta_{i}w_{i} - \sum_{j=1}^{n} \overline{a}_{j} \left[\left(|c_{ij1}(t)|\gamma_{j1} + |c_{ij2}(t)|\gamma_{j2} + |d_{ij}(t)|\xi_{j} \right) w_{j} \right] > 0,$$
(5.27)

holds for all $t \in [0, \omega]$ and i = 1, ..., n, then the model (5.23) has an ω -periodic solution which is globally exponentially stable.

Remark 5.12. In [36] the existence of a unique pseudo almost automorphic solution of model (5.23) with $c_{ij1}, c_{ij2}, \tau_{ij}, d_{ij}$ being pseudo almost automorphic functions was obtained assuming (H1), (H3*) with $\beta_i(t) \equiv \beta_i$, K_{ij} verifying (5.17), (5.20), and h_{j1}, h_{j2}, g_j are Lipschitz functions with Lipschitz constants $\gamma_{j1}, \gamma_{j2}, \xi_j > 0$, and

$$\underline{a}_{i}\beta_{i}w_{i} - \sum_{j=1}^{n} \overline{a}_{j} \left[\left(\overline{c}_{ij1}\gamma_{j1} + \overline{c}_{ij2}\gamma_{j2} + \overline{d}_{ij}\xi_{j} \right) w_{j} \right] > 0, \qquad (5.28)$$

where $\overline{c}_{ijp} = \sup |c_{ij1}(t)|$, $\overline{d}_{ij} = \sup |d_{ij}(t)|$ for $i, j = 1, \ldots, n$ and p = 1, 2.

All periodic functions are pseudo almost automorphic functions. Thus Corollary 5.11 is not a generalization of [36, Theorem 3.1]. However, in case of (5.23) being a periodic model, the existence criterium in Corollary 5.11 is better than the corresponding criterium in [36, Theorem 3.1].

Example 5.3. Consider the following high-order Cohen-Grossberg neural network model,

$$x'(t) = a_i(x_i(t)) \left[-b_i(x_i(t)) + \sum_{j=1}^n c_{ij}(t) f_j(\rho_j x_j(t)) + \sum_{j=1}^n d_{ij11}(t) f_j\left(\rho_j \int_0^{+\infty} K_{ij}(u) x_j(t-u) du\right) + \sum_{j,l=1}^n d_{ijl2}(t) f_j\left(\rho_j \int_0^{+\infty} K_{ij}(u) x_j(t-u) du\right) f_l\left(\rho_l \int_0^{+\infty} K_{il}(u) x_l(t-u) du\right) + I_i(t) \right],$$
(5.29)

for $t \geq 0$, i = 1, ..., n, where, for each i, j, l = 1, ..., n and $q = 1, 2, \rho_i > 0, a_i : \mathbb{R} \to (0, +\infty), b_i : \mathbb{R} \to \mathbb{R}, c_{ij}, d_{ijlq}, I_i : [0, +\infty) \to \mathbb{R}, f_j : \mathbb{R} \to \mathbb{R}, \text{ and } K_{ij} : [0, +\infty) \to [0, +\infty)$ are continuous functions such that K_{ij} verifies (5.17).

The existence and global exponential stability of a periodic solution of (5.29) were studied in [23].

Considering the definition of the bounded variation functions η_{ij} as in (5.19), model (5.29) is a particular situation of (5.11), thus from Corollary 5.4 we obtain the following stability criterion.

Corollary 5.13. Assume (H1), (H3^{*}) with $\beta_i(t) \equiv \beta_i$, and, for each i, j, l = 1, ..., n and $q = 1, 2, c_{ij}, d_{ijlq}, I_i$ are ω -periodic for some $\omega > 0$, K_{ij} verifies (5.17) and (5.20), and there are $M_j > 0$ and $\mu_j > 0$ such that

$$|f_j(u) - f_j(v)| \le \mu_j |u - v|$$
 and $|f_j(u)| \le M_j$, $\forall u, v \in \mathbb{R}, j = 1, \dots, n$.

If there exists $w = (w_1, \ldots, w_n) > 0$ such that, for all $t \ge 0$ and $i = 1, \ldots, n$,

$$\underline{a}_{i}\beta_{i}w_{i} > \overline{a}_{i}\sum_{j=1}^{n} \left[|c_{ij}(t)|\mu_{j}\rho_{j}w_{j} + |d_{ij11}(t)|\rho_{j}\mu_{j}w_{j} + \sum_{l=1}^{n} |d_{ijl2}(t)| \left(w_{j}M_{j}\mu_{j}\rho_{j} + w_{l}M_{l}\mu_{l}\rho_{l}\right) \right], \quad (5.30)$$

then the model (5.22) has an ω -periodic solution which is globally exponentially stable.

Proof. If we take in model (5.11) P = 1, Q = 2, and, for each i, j, l = 1, ..., n, q = 1, 2, the functions $F_i(u) = G_i(u) = u$, $\tau_{ij1}(t) = \tilde{\tau}_{ij1}(t) = 0$, $h_{ijl1}(u_1, u_2) = f_j(\rho_j u_1)$, $c_{ij11}(t) = c_{ij}(t)$ $c_{ijl1}(t) = d_{ijl1}(t) = 0$ for $l \neq 1$, $f_{ijl1}(u_1, u_2) = f_j(\rho_j u_1)$, $f_{ijl2}(u_1, u_2) = f_j(\rho_j u_1)f_j(\rho_j u_2)$, $g_{ijq}(u) = \tilde{g}_{ijq}(u) = u$, and $\tilde{\eta}_{ijq}(s) = \eta_{ijq}(s)$ defined by (5.19) for all $u_1, u_2 \in \mathbb{R}$ and $s \leq 0$, then we obtain model (5.29).

For all $u_1, u_2, v_1, v_2 \in \mathbb{R}$, we have

$$\begin{aligned} |h_{ijl1}(u_1, u_2) - h_{ijl1}(v_1, v_2)| &= |f_j(\rho_j u_1) - f_j(\rho_j v_1)| \le \rho_j \mu_j |u_1 - v_1|, \\ |f_{ijl1}(u_1, u_2) - f_{ijl1}(v_1, v_2)| &= |f_j(\rho_j u_1) - f_j(\rho_j v_1)| \le \rho_j \mu_j |u_1 - v_1|, \end{aligned}$$

and

$$\begin{aligned} |f_{ijl2}(u_1, u_2) - f_{ijl2}(v_1, v_2)| &= |f_j(\rho_j u_1) f_j(\rho_j u_2) - f_j(\rho_j v_1) f_j(\rho_j u_2)| \\ &\leq M_j \rho_j \mu_j |u_1 - v_1| + M_j \rho_j \mu_j |u_2 - v_2|, \end{aligned}$$

for all i, j, l = 1, ..., n, thus hypothesis (H4^{**}) holds. Condition (5.2) follows from (5.20) and the inequality (5.12) reads as (5.30). Finally, the result follows from Corollary 5.4.

Remark 5.14. In [23], sufficient conditions for the existence and global exponential stability of a ω -periodic solution of (5.29) were presented. However, it is important to mention that the proof of the main result is not correct. Specifically, the way inequality [23, (3.10)] is obtained is problematic. In fact assuming the uniqueness of solution of (5.29) with initial condition $x_0 = \psi$ for $\psi \in BC$, denoting this solution by $x(t, 0, \psi)$, and defining $P: BC \to BC$ by $P(\psi) = x_{\omega}(\cdot, 0, \psi)$, we always have

$$||P^{N}(\psi_{1}) - P^{N}(\psi_{2})|| = ||x_{N\omega}(\cdot, 0, \psi_{1}) - x_{N\omega}(\cdot, 0, \psi_{2})||$$

=
$$\sup_{s \le 0} ||x(N\omega + s, 0, \psi_{1}) - x(N\omega + s, 0, \psi_{2})||$$

$$\ge ||\psi_{1} - \psi_{2}||$$

for all $\psi_1, \psi_2 \in BC$ and $N \in \mathbb{N}$, since model (5.29) is ω -periodic.

6 Numerical Example

Here, we present a numerical example to illustrate the applicability of some new results given in this work.

The system

$$x_{1}'(t) = \left(\frac{1}{48}\sin\left(x_{1}(t)\right) + \frac{7}{48}\right) \left[-(9 + \sin(t))x_{1}(t) + c\cos(t)\arctan\left(x_{1}(t - \sin(t))\right) \\ \cdot \arctan\left(x_{2}(t - \cos(t))\right) + d\sin(t)\int_{0}^{+\infty}e^{-u}x_{2}(t - u)du + \cos(t)\right]$$
(6.1)

$$\begin{aligned} x_2'(t) &= \left(2 + \cos\left(x_2(t)\right)\right) \left[-(2 + \cos(t))x_2(t) + \hat{c}\sin(t)\arctan\left(x_1(t - \cos(t))\right) \\ &+ \hat{d}\cos(t)\tanh\left(\int_0^{+\infty} e^{-u}x_1(t - u)du\right) \tanh\left(\int_0^{+\infty} e^{-u}x_2(t - u)du\right) + e^{\sin(t)}\right] \end{aligned}$$

for $t \ge 0$, where $c, d, \hat{c}, \hat{d} \in \mathbb{R}$, is a 2π -periodic example of a high-order Cohen-Grossberg neural network model.

Defining $\eta_{ij1}, \tilde{\eta}_{ij1} : (-\infty, 0] \to \mathbb{R}$ by

$$\eta_{ij1}(s) = \tilde{\eta}_{ij1}(s) = \int_{-\infty}^{s} e^{v} dv, \quad s \in (-\infty, 0],$$

system (6.1) is a particular situation of (5.11). However, (6.1) is not a particular case of (5.29), thus the model studied in [23] is not general enough to include (6.1) as a particular example.

Following the notations in (5.11) and (5.13), we have n = 2, P = Q = 1, $\underline{a}_1 = \frac{1}{8}$, $\overline{a}_1 = \frac{1}{6}$, $\underline{a}_2 = 1$, $\overline{a}_2 = 3$, $\underline{\beta}_1 = 8$, $\underline{\beta}_2 = 1$, $\zeta_i = \zeta_i = 1$, $\gamma_{1121}^{(1)} = \gamma_{1121}^{(2)} = \frac{\pi}{2}$, $\gamma_{2111}^{(1)} = 1$, $\mu_{1211}^{(2)} = \mu_{2121}^{(1)} = \mu_{2121}^{(2)} = 1$, $I_1(t) = \cos(t)$, $I_2(t) = e^{\sin(t)}$, $\gamma_{1121}^{(1)} = \gamma_{1121}^{(2)} = \frac{\pi}{2}$, $\gamma_{2111}^{(1)} = 1$, $\mu_{1211}^{(2)} = \mu_{2121}^{(1)} = 1$, $c_{1121}(t) = c\cos(t)$, $c_{2111}(t) = \hat{c}\sin(t)$, $d_{1211}(t) = d\sin(t)$, $d_{2121}(t) = \hat{d}\cos(t)$, and all other $c_{ijl1}(t) = d_{ijl1}(t) = 0$, for i, j, l = 1, 2.

Consequently, example (6.1) is 2π -periodic and the matrix \mathcal{M} , defined in (5.14), has the form

$$\mathcal{M} = \begin{bmatrix} 1 - \frac{\pi}{16} |c| & -\frac{1}{2} \left(\frac{\pi}{2} |c| + |d| \right) \\ -3 \left(\frac{\pi}{2} |\hat{c}| + |\hat{d}| \right) & 1 - 3 |\hat{d}| \end{bmatrix}.$$

Condition (5.2) trivially holds with $\vartheta \in (0, 1)$. Consequently, Corollary 5.5 assures the existence and exponential stability of a 2π -periodic solution of (6.1) in case \mathcal{M} being a non-singular M-matrix. For example, if we consider $c = \frac{1}{\pi}$, $d = \frac{1}{100}$, $\hat{c} = \frac{1}{30\pi}$, $\hat{d} = \frac{1}{30}$, we have

$$\mathcal{M} = \begin{bmatrix} \frac{15}{16} & -\frac{101}{200} \\ \\ -\frac{1}{5} & \frac{9}{10} \end{bmatrix},$$

which is a non-singular M-matrix.

Acknowledgments.

This work was partially supported by Fundação para a Ciência e a Tecnologia (Portugal) within the Projects UIDB/00013/2020, UIDP/00013/2020 of CMAT-UM (José J. Oliveira), and Project UIDB/00212/2020 of CMA-UBI (Ahmed Elmwafy and César M. Silva).

References

- [1] C. Aouiti and F. Dridi. New results on interval general cohen-grossberg bam neural networks. *Journal of Systems Science and Complexity*, 33:944–967, 2020.
- [2] S. Arik. Global asymptotic stability analysis of bidirectional associative memory neural networks with time delays. *IEEE Transactions on Neural Networks*, 16(3):580–586, 2005.

- [3] X. Chen and Q. Song. Global exponential stability of the periodic solution of delayed cohen–grossberg neural networks with discontinuous activations. *Neuro*computing, 73(16-18):3097–3104, 2010.
- [4] Y. Chen and S. Jia. Multiple stability and instability of cohen-grossberg neural network with unbounded time-varying delays. *Journal of Inequalities and Applications*, 2019(1):1–14, 2019.
- [5] M. A. Cohen and S. Grossberg. Absolute stability of global pattern formation and parallel memory storage by competitive neural networks. *IEEE transactions on* systems, man, and cybernetics, 5:815–826, 1983.
- [6] Z. Dong, X. Zhang, and X. Wang. Global exponential stability of discrete-time higher-order cohen-grossberg neural networks with time-varying delays, connection weights and impulses. *Journal of the Franklin Institute*, 358(11):5931–5950, 2021.
- [7] C. Eliasmith. A unified approach to building and controlling spiking attractor networks. *Neural computation*, 17(6):1276–1314, 2005.
- [8] T. Faria and J. J. Oliveira. General criteria for asymptotic and exponential stabilities of neural network models with unbounded delays. *Applied mathematics and computation*, 217(23):9646–9658, 2011.
- [9] M. Fiedler. Special matrices and their applications in numerical mathematics. Courier Corporation, 2008.
- [10] J. Haddock and W. Hornor. Precompactness and convergence in norm of positive orbits in a certain fading memory space. *Funkcial. Ekvac*, 31:349–361, 1988.
- [11] J. Haddock and J. Terjéki. On the location of positive limit sets for autonomous functional differential equations with infinite delay. *Journal of Differential Equations*, 86(1):1–32, 1990.
- [12] J. Hale and J. Kato. Phase space for retarded equations with infinite delay. funkcial. ekvac. 21. 1978.
- [13] J. K. Hale. Ordinary differential equations, robert e. Krieer, New York, 1980.
- [14] C. Huang, X. Long, and J. Cao. Stability of antiperiodic recurrent neural networks with multiproportional delays. *Mathematical Methods in the Applied Sciences*, 43(9):6093–6102, 2020.
- [15] L. Hien, T. Loan, B. H. Trang, and H. Trinh. Existence and global asymptotic stability of positive periodic solution of delayed cohen–grossberg neural networks. *Applied Mathematics and Computation*, 240:200–212, 2014.
- [16] Y. Hino, S. Murakami, and T. Naito. Functional differential equations with infinite delay. Springer, 2006.
- [17] H. Kang, X. Fu, and Z. Sun. Global exponential stability of periodic solutions for impulsive cohen–grossberg neural networks with delays. *Applied Mathematical Modelling*, 39(5-6):1526–1535, 2015.

- [18] M. M. Ketencigil, O. Faydasicok, and S. Arik. Novel criteria for robust stability of cohen-grossberg neural networks with multiple time delays. *Discrete and Continuous Dynamical Systems-S*, 15(11):3189–3203, 2022.
- [19] B. Li and Q. Song. Some new results on periodic solution of cohen-grossberg neural network with impulses. *Neurocomputing*, 177:401–408, 2016.
- [20] M. Liu, I. Dassios, and F. Milano. On the stability analysis of systems of neutral delay differential equations. *Circuits, Systems, and Signal Processing*, 38:1639– 1653, 2019.
- [21] Q. Liu and R. Xu. Periodic solutions of high-order cohen–grossberg neural networks with distributed delays. *Communications in Nonlinear Science and Numerical Simulation*, 16(7):2887–2893, 2011.
- [22] B. Liu. New convergence behavior of solutions to cohen–grossberg neural networks with delays and time-varying coefficients. *Physics Letters A*, 372(2):117–123, 2008.
- [23] Q. Liu and R. Xu. Periodic solutions of high-order cohen-grossberg neural networks with distributed delays. *Communications in Nonlinear Science and Numerical Simulation*, 16(7):2887–2893, 2011.
- [24] F. Long, Y. Wang, and S. Zhou. Existence and exponential stability of periodic solutions for a class of cohen–grossberg neural networks with bounded and unbounded delays. *Nonlinear Analysis: Real World Applications*, 8(3):797–810, 2007.
- [25] J. Mawhin. Periodic solutions of nonlinear functional differential equations. Journal of Differential Equations, 10(2):240–261, 1971.
- [26] J. Mawhin. Periodic solutions in the golden sixties: the birth of a continuation theorem. In 10 Mathematical Essays on Approximation in Analysis and Topology, pages 199–214. Elsevier, 2005.
- [27] I. Ncube. Existence, uniqueness, and global asymptotic stability of an equilibrium in a multiple unbounded distributed delay network. *Electronic Journal of Qualitative Theory of Differential Equations*, 2020(59):1–11, 2020.
- [28] J. J. Oliveira. Convergence of asymptotic systems of non-autonomous neural network models with infinite distributed delays. *Journal of Nonlinear Science*, 27:1463–1486, 2017.
- [29] J. J. Oliveira. Global exponential stability of nonautonomous neural network models with unbounded delays. *Neural Networks*, 96:71–79, 2017.
- [30] J. H. Park. Lmi optimization approach to asymptotic stability of certain neutral delay differential equation with time-varying coefficients. *Applied mathematics and computation*, 160(2):355–361, 2005.
- [31] S. Qin, X. Xue, and P. Wang. Global exponential stability of almost periodic solution of delayed neural networks with discontinuous activations. *Information Sciences*, 220:367–378, 2013.

- [32] F. A. Rihan, C. Tunc, S. Saker, S. Lakshmanan, and R. Rakkiyappan. Applications of delay differential equations in biological systems. *Complexity*, 2018:NA–NA, 2018.
- [33] J. Wu, S. A. Campbell, and J. Bélair. Time-delayed neural networks: Stability and oscillations. In *Encyclopedia of Computational Neuroscience*, pages 3434–3440. Springer, 2022.
- [34] R. Wu. Exponential convergence of bam neural networks with time-varying coefficients and distributed delays. Nonlinear Analysis: Real World Applications, 11(1):562–573, 2010.
- [35] Y. Yao and W. J. Freeman. Model of biological pattern recognition with spatially chaotic dynamics. *Neural networks*, 3(2):153–170, 1990.
- [36] H. Zhu, Q. Zhu, X. Sun, and H. Zhou. Existence and exponential stability of pseudo almost automorphic solutions for cohen-grossberg neural networks with mixed delays. Advances in Difference Equations, 2016(1):1–17, 2016.
- [37] W. Zhao. Dynamics of cohen–grossberg neural network with variable coefficients and time-varying delays. *Nonlinear Analysis: Real World Applications*, 9(3):1024– 1037, 2008.
- [38] F. Zhang, B. Liu, and L. Huang. Existence and exponential stability of periodic solutions for a class of cohen–grossberg neural networks with bounded and unbounded delays. *Computers & Mathematics with Applications*, 53(9):1325–1338, 2007.
- [39] J. Zhang and C. Huang. Dynamics analysis on a class of delayed neural networks involving inertial terms. Advances in Difference Equations, 2020(1):1–12, 2020.
- [40] Q. Zhang, X. Wei, and J. Xu. Global exponential stability for nonautonomous cellular neural networks with unbounded delays. *Chaos, Solitons & Fractals*, 39(3):1144–1151, 2009.