

SINGULAR EXTENSION OF CRITICAL SOBOLEV MAPPINGS UNDER AN EXPONENTIAL WEAK-TYPE ESTIMATE

BOHDAN BULANYI AND JEAN VAN SCHAFTINGEN

ABSTRACT. Given $m \in \mathbb{N} \setminus \{0\}$ and a compact Riemannian manifold \mathcal{N} , we construct for every map u in the critical Sobolev space $W^{m/(m+1), m+1}(\mathbb{S}^m, \mathcal{N})$, a map $U : \mathbb{B}^{m+1} \rightarrow \mathcal{N}$ whose trace is u and which satisfies an exponential weak-type Sobolev estimate. The result and its proof carry on to the extension to a half-space of maps on its boundary hyperplane and to the extension to the hyperbolic space of maps on its boundary sphere at infinity.

CONTENTS

1. Introduction	1
2. Mean oscillation on λ -adic skeletons	7
2.1. Truncated mean oscillation	7
2.2. λ -adic cubes and skeletons	8
2.3. Longitudinal estimate	8
2.4. Transversal estimate	9
2.5. Combining the estimates	11
3. Proofs of the singular extension theorems	11
3.1. Oscillation and gradient estimate on the skeleton	11
3.2. Sobolev and Sobolev–Marcinkiewicz extensions on cubes	13
3.3. Proofs of the theorems	14
References	19

1. INTRODUCTION

The classical trace theory in Sobolev spaces [8] states that for every $m \in \mathbb{N} \setminus \{0\}$ and $p \in (1, \infty)$, there is a well-defined surjective trace operator $\text{tr}_{\mathbb{S}^m} : W^{1,p}(\mathbb{B}_1^{m+1}, \mathbb{R}) \rightarrow W^{1-1/p,p}(\mathbb{S}^m, \mathbb{R})$ that coincides with the restriction on the subsets of continuous functions, where the first-order Sobolev space on the unit ball $\mathbb{B}_1^{m+1} \subseteq \mathbb{R}^{m+1}$ is defined as

$$W^{1,p}(\mathbb{B}_1^{m+1}, \mathbb{R}) = \left\{ U : \mathbb{B}_1^{m+1} \rightarrow \mathbb{R} \mid U \text{ is weakly differentiable and } \int_{\mathbb{B}_1^{m+1}} |DU|^p < \infty \right\}$$

2020 *Mathematics Subject Classification.* 58D15 (46E35).

Key words and phrases. Extension of traces in Sobolev spaces; trace theory; Sobolev embedding theorem; weak-type Marcinkiewicz spaces; Lorentz space.

The authors were supported by the Projet de Recherche T.0229.21 ‘‘Singular Harmonic Maps and Asymptotics of Ginzburg–Landau Relaxations’’ of the Fonds de la Recherche Scientifique–FNRS.

and the fractional Sobolev space $W^{s,p}(\mathbb{S}^m, \mathbb{R})$ on the unit sphere $\mathbb{S}^m := \partial\mathbb{B}_1^{m+1}$ is defined for $s \in (0, 1)$ as

$$(1) \quad W^{s,p}(\mathbb{S}^m, \mathbb{R}) = \left\{ u : \mathbb{S}^m \rightarrow \mathbb{R} \mid u \text{ is measurable and } \iint_{\mathbb{S}^m \times \mathbb{S}^m} \frac{|u(x) - u(y)|^p}{|x - y|^{m+sp}} dx dy < \infty \right\}.$$

Moreover, the trace operator $\text{tr}_{\mathbb{S}^m}$ has a continuous linear right inverse through a suitable linear operator, so that, in particular, for every function $u \in W^{1-1/p,p}(\mathbb{S}^m, \mathbb{R})$ there exists a function $U \in W^{1,p}(\mathbb{B}_1^{m+1}, \mathbb{R})$ such that $\text{tr}_{\mathbb{S}^m} U = u$ and

$$(2) \quad \int_{\mathbb{B}_1^{m+1}} |DU|^p \leq C \iint_{\mathbb{S}^m \times \mathbb{S}^m} \frac{|u(x) - u(y)|^p}{|x - y|^{m+p-1}} dx dy,$$

where the constant $C \in (0, \infty)$ in (2) depends only on m and p .

Given a compact Riemannian manifold \mathcal{N} , which we can assume without loss of generality to be linearly embedded into \mathbb{R}^ν by Nash's isometric embedding theorem [14], we can consider the Sobolev spaces of mappings

$$W^{1,p}(\mathbb{B}_1^{m+1}, \mathcal{N}) = \{U \in W^{1,p}(\mathbb{B}_1^{m+1}, \mathbb{R}^\nu) \mid U \in \mathcal{N} \text{ almost everywhere in } \mathbb{B}_1^{m+1}\}$$

and

$$W^{s,p}(\mathbb{S}^m, \mathcal{N}) = \{u \in W^{s,p}(\mathbb{S}^m, \mathbb{R}^\nu) \mid u \in \mathcal{N} \text{ almost everywhere in } \mathbb{S}^m\}.$$

Although it follows immediately from the classical trace theory that the trace operator is well defined and continuous from $W^{1,p}(\mathbb{B}_1^{m+1}, \mathcal{N})$ to $W^{1-1/p,p}(\mathbb{S}^m, \mathcal{N})$, the proof of the surjectivity fails to extend to this case: one can just prove that every mapping in $W^{1-1/p,p}(\mathbb{S}^m, \mathcal{N})$ can be extended to a function in $W^{1,p}(\mathbb{B}_1^{m+1}, \mathbb{R}^\nu)$; since the extension is constructed by averaging, there is no hope that the extension would satisfy the nonlinear manifold constraint in general.

In the case of subcritical dimension $m < p - 1$, maps in the nonlinear Sobolev spaces $W^{1-1/p,p}(\mathbb{S}^m, \mathcal{N})$ and $W^{1,p}(\mathbb{B}_1^{m+1}, \mathcal{N})$ can be assumed to be continuous by the Sobolev–Morrey embedding; it turns out that all maps in $W^{1-1/p,p}(\mathbb{S}^m, \mathcal{N})$ are traces of maps in $W^{1,p}(\mathbb{B}_1^{m+1}, \mathcal{N})$ if and only if all continuous maps from \mathbb{S}^m to \mathcal{N} are restrictions of continuous mappings, or equivalently, if and only if the m^{th} homotopy group $\pi_m(\mathcal{N})$ is trivial: $\pi_m(\mathcal{N}) \simeq \{0\}$ [2, Th. 1]. For the critical dimension $m = p - 1$, one gets similarly that all mappings in $W^{m/(m+1), m+1}(\mathbb{S}^m, \mathcal{N})$ are traces of maps in $W^{1, m+1}(\mathbb{B}_1^{m+1}, \mathcal{N})$ if and only if $\pi_m(\mathcal{N}) \simeq \{0\}$ [2, Th. 2]; this can be explained by the bounded mean oscillation property of maps in $W^{m/(m+1), m+1}(\mathbb{S}^m, \mathcal{N})$ and the possibility to extend homotopy and obstruction theories to such maps [6].

The situation is radically different in the supercritical dimension case $m > p - 1$, where several necessary or sufficient conditions, imposing either finiteness or triviality of some of the homotopy groups $\pi_j(\mathcal{N})$ for $j \in \{1, \dots, \lfloor p - 1 \rfloor\}$, have been proved [1, 2, 9, 13] and where the complete characterization of manifolds \mathcal{N} , for which the trace operator is surjective, still remains open.

Going back to the case of subcritical or critical dimension $m \leq p - 1$, one can wonder whether an estimate (2) can still hold for the extension of Sobolev mappings. It has been proved that every map $u \in W^{1-1/p,p}(\mathbb{S}^m, \mathcal{N})$ has an extension $U \in W^{1,p}(\mathbb{B}_1^{m+1}, \mathcal{N})$ with the estimate (2) when either $m = 1$ and $\pi_1(\mathcal{N}) \simeq \{0\}$ or $m \geq 2$, $\pi_1(\mathcal{N})$ is finite and $\pi_2(\mathcal{N}) \simeq \dots \simeq \pi_{\lfloor p \rfloor}(\mathcal{N}) \simeq \{0\}$ [9, 23]; moreover, if there is an extension satisfying (2), then the homotopy groups $\pi_1(\mathcal{N}), \dots, \pi_{\lfloor p-1 \rfloor}(\mathcal{N})$ have all to be finite [1, 13]; in the critical case $m = p - 1$, the additional condition that $\pi_m(\mathcal{N}) \simeq \{0\}$ is necessary for

a Sobolev estimate on the extension of continuous mappings that have a continuous extension [13].

In view of the obstructions to linear Sobolev estimates on the extension of the form (2), one can hope to get some nonlinear estimates instead. In the subcritical dimension case, a compactness argument [22, Th. 4] shows that given $m \in \mathbb{N} \setminus \{0\}$, $p > m + 1$ and a compact Riemannian manifold \mathcal{N} , there exists a function $\gamma \in C([0, \infty), [0, \infty))$ such that $\gamma(0) = 0$, and every map $u \in W^{1-1/p, p}(\mathbb{S}^m, \mathcal{N})$ that has a continuous extension has an extension $U \in W^{1, p}(\mathbb{B}_1^{m+1}, \mathcal{N})$ satisfying

$$(3) \quad \int_{\mathbb{B}_1^{m+1}} |\mathrm{D}U|^p \leq \gamma \left(\iint_{\mathbb{S}^m \times \mathbb{S}^m} \frac{d(u(x), u(y))^p}{|x - y|^{m+p-1}} \mathrm{d}x \mathrm{d}y \right),$$

where d is the geodesic distance on \mathcal{N} .

Because of bubbling phenomena, this estimate (3) does not go to the endpoint $p = m + 1$ when $\pi_m(\mathcal{N}) \not\cong \{0\}$. Indeed, any map in $W^{m/(m+1), m+1}(\mathbb{S}^m, \mathcal{N})$ is the weak limit of continuous maps in $W^{m/(m+1), m+1}(\mathbb{S}^m, \mathcal{N})$ that can be extended so that a weak compactness argument would imply that every map in $W^{m/(m+1), m+1}(\mathbb{S}^m, \mathcal{N})$ has an extension $W^{1, m+1}(\mathbb{B}_1^{m+1}, \mathcal{N})$, which cannot be the case when $\pi_m(\mathcal{N}) \not\cong \{0\}$ (see [21, Prop. 2.8]). In this situation, it has been proved in works by Petrace, Rivière and the second author that there is still an extension $U \in W^{1, (m+1, \infty)}(\mathbb{B}_1^{m+1}, \mathcal{N})$ satisfying a weak-type Marcinkiewicz–Sobolev estimate [21; 22, Th. 2]: for every $t \in (0, \infty)$,

$$(4) \quad t^{m+1} \mathcal{L}^{m+1}(\{x \in \mathbb{B}_1^{m+1} \mid |\mathrm{D}U(x)| > t\}) \leq \gamma \left(\iint_{\mathbb{S}^m \times \mathbb{S}^m} \frac{d(u(x), u(y))^p}{|x - y|^{2m}} \mathrm{d}x \mathrm{d}y \right).$$

The function γ , appearing in the estimate (4) for a general target manifold \mathcal{N} , is a wild *double exponential* function [22, discussion after Th. 2]. Petrace and Rivière get similar estimates with γ being a polynomial when $m = 2$ and $\mathcal{N} = \mathbb{S}^2$ [21, Th. C] (thanks to the Hopf fibration) and an exponential of a power when $m = 3$ and $\mathcal{N} = \mathbb{S}^3$ [21, Th. B].

In the present work, we construct an extension U that satisfies (4), where γ can be taken to be an exponential function.

Theorem 1.1. *Let $m \in \mathbb{N} \setminus \{0\}$ and let \mathcal{N} be a compact Riemannian manifold. There exist constants $A, B, \delta \in (0, \infty)$ such that for every $u \in W^{m/(m+1), m+1}(\mathbb{S}^m, \mathcal{N})$, there exists a mapping $U \in W_{\mathrm{loc}}^{1, 1}(\mathbb{B}_1^{m+1}, \mathcal{N})$ such that $\mathrm{tr}_{\mathbb{S}^m} U = u$ and for every $t \in (0, \infty)$,*

$$(5) \quad t^{m+1} \mathcal{L}^{m+1}(\{x \in \mathbb{B}_1^{m+1} \mid |\mathrm{D}U(x)| \geq t\}) \leq A \exp \left(B \iint_{\substack{(x, y) \in \mathbb{S}^m \times \mathbb{S}^m \\ d(u(x), u(y)) \geq \delta}} \frac{1}{|x - y|^{2m}} \mathrm{d}x \mathrm{d}y \right) \iint_{\mathbb{S}^m \times \mathbb{S}^m} \frac{d(u(x), u(y))^{m+1}}{|x - y|^{2m}} \mathrm{d}x \mathrm{d}y.$$

Moreover, one can take $U \in C(\mathbb{B}_1^{m+1} \setminus S, \mathcal{N})$, where the singular set $S \subset \mathbb{B}_1^{m+1}$ is a finite set whose cardinality is controlled by the right-hand side of (5).

The gap potential of the double integral appearing in the exponential in (5)

$$(6) \quad \iint_{\substack{(x, y) \in \mathbb{S}^m \times \mathbb{S}^m \\ d(u(x), u(y)) \geq \delta}} \frac{1}{|x - y|^{2m}} \mathrm{d}x \mathrm{d}y$$

first appeared in estimates by Bourgain, Brezis and Mironescu on the topological degree of maps from a sphere to itself [3, Open problem 2; 4, Th. 1.1; 17] (see also [20]) and in estimates on free homotopy decompositions of mappings [23], as well as in estimates on liftings [18, Th. 2; 24]; they characterize, in the limit $\delta \rightarrow 0$, first-order Sobolev spaces [15, 16, 19] and encompass a property stronger than VMO [5].

Theorem 1.1 implies, in particular, the weak-type estimates of [22]. The improvement of Theorem 1.1 is two-fold: the dependence of the weak-type Sobolev bound on the extension U is exponential in the Gagliardo energy of u , which is much more reasonable than the double exponential in [22] and the nonlinear part of the estimate, that is the exponential, relies on a gap potential (6) instead of a full fractional Gagliardo energy

$$\iint_{\mathbb{S}^m \times \mathbb{S}^m} \frac{d(u(x), u(y))^p}{|x - y|^{2m}} dx dy.$$

The latter controls the form by the immediate estimate

$$\iint_{\substack{(x,y) \in \mathbb{S}^m \times \mathbb{S}^m \\ d(u(x), u(y)) \geq \delta}} \frac{1}{|x - y|^{2m}} dx dy \leq \frac{1}{\delta^p} \iint_{\mathbb{S}^m \times \mathbb{S}^m} \frac{d(u(x), u(y))^p}{|x - y|^{2m}} dx dy.$$

We also have a counterpart of Theorem 1.1 for the extension on the half-space $\mathbb{R}_+^{m+1} := \mathbb{R}^m \times (0, \infty)$ of mappings defined on the hyperplane $\mathbb{R}^m \simeq \mathbb{R}^m \times \{0\}$.

Theorem 1.2. *Let $m \in \mathbb{N} \setminus \{0\}$ and let \mathcal{N} be a compact Riemannian manifold. There exist constants $A, B, \delta \in (0, \infty)$ such that for every $u \in W^{m/(m+1), m+1}(\mathbb{R}^m, \mathcal{N})$, there exists $U \in W_{\text{loc}}^{1,1}(\mathbb{R}_+^{m+1}, \mathcal{N})$ such that $\text{tr}_{\mathbb{R}^m} U = u$ and for every $t \in (0, \infty)$,*

$$(7) \quad t^{m+1} \mathcal{L}^{m+1}(\{x \in \mathbb{R}_+^{m+1} \mid |DU(x)| \geq t\}) \\ \leq A \exp\left(B \iint_{\substack{(x,y) \in \mathbb{R}^m \times \mathbb{R}^m \\ d(u(x), u(y)) \geq \delta}} \frac{1}{|x - y|^{2m}} dx dy \right) \iint_{\mathbb{R}^m \times \mathbb{R}^m} \frac{d(u(x), u(y))^{m+1}}{|x - y|^{2m}} dx dy.$$

Moreover, one can take $U \in C(\mathbb{R}_+^{m+1} \setminus S, \mathcal{N})$, where the singular set $S \subset \mathbb{R}_+^{m+1}$ is a finite set whose cardinality is controlled by the right-hand side of (7).

Finally, we consider the extension to the hyperbolic space \mathbb{H}^{m+1} of maps defined on its boundary sphere \mathbb{S}^m . Using the Poincaré ball model, the hyperbolic space \mathbb{H}^{m+1} is the ball \mathbb{B}_1^{m+1} , endowed with the metric g defined for $x \in \mathbb{B}_1^{m+1}$ in terms of the Euclidean metric g_{euc} by

$$g(x) = \frac{4g_{\text{euc}}(x)}{(1 - |x|^2)^2},$$

whose boundary \mathbb{S}^m is then considered to be the boundary sphere of \mathbb{H}^{m+1} . In this setting, we have a hyperbolic counterpart of Theorem 1.1 and Theorem 1.2.

Theorem 1.3. *Let $m \in \mathbb{N} \setminus \{0\}$ and let \mathcal{N} be a compact Riemannian manifold. There exist constants $A, B, \delta \in (0, \infty)$ such that for every $u \in W^{m/(m+1), m+1}(\mathbb{S}^m, \mathcal{N})$, there*

exists $U \in W_{\text{loc}}^{1,1}(\bar{\mathbb{B}}_1^{m+1}, \mathcal{N})$ such that $\text{tr}_{\mathbb{S}^m} U = u$ and for every $t \in (0, \infty)$,

$$(8) \quad \mathcal{H}^{m+1}(\{x \in \mathbb{H}^{m+1} \mid |DU(x)| \geq t\}) \leq \frac{A}{t^{m+1}} \exp\left(B \iint_{\substack{(x,y) \in \mathbb{S}^m \times \mathbb{S}^m \\ d(u(x), u(y)) \geq \delta}} \frac{1}{|x-y|^{2m}} dx dy \right) \iint_{\mathbb{S}^m \times \mathbb{S}^m} \frac{d(u(x), u(y))^{m+1}}{|x-y|^{2m}} dx dy.$$

Moreover, one can take $U \in C(\mathbb{H}^{m+1} \setminus S, \mathcal{N})$, where the singular set $S \subset \mathbb{H}^{m+1}$ is a finite set whose cardinality is controlled by the right-hand side of (8).

Although standard conformal transformations between \mathbb{B}_1^{m+1} , \mathbb{H}^{m+1} and \mathbb{R}_+^{m+1} preserve the fractional Gagliardo energy of the boundary values in the right-hand sides of (5), (7) and (8), the corresponding gap potentials and the strong-type quantities

$$\int_{\mathbb{B}_1^{m+1}} |DU|^{m+1}, \quad \int_{\mathbb{R}_+^{m+1}} |DU|^{m+1} \quad \text{and} \quad \int_{\mathbb{H}^{m+1}} |DU|^{m+1}$$

corresponding to the weak-type quantities in their left-hand side, the left-hand sides as such are not conformally invariant so that Theorems 1.1 to 1.3 are not equivalent to each other. In practice, this explains why we have the same construction of U in all three cases, but three different particular estimates on U .

At the core of the proof of Theorems 1.1 to 1.3 is a refined understanding of functions in the fractional critical Sobolev space $W^{s,p}(\mathbb{R}^m, \mathbb{R}^\nu)$. We start from the elegant and versatile approach through the vanishing mean oscillation property of critical Sobolev maps [6]: if $sp = m$, it follows from Lebesgue's dominated convergence theorem and from the definition in (1) that

$$(9) \quad \lim_{r \rightarrow 0} \sup_{x \in \mathbb{R}^m} \int_{\mathbb{B}_r^m(x)} \int_{\mathbb{B}_r^m(x)} |u(y) - u(z)|^p dy dz = 0.$$

If $u \in \mathcal{N}$ almost everywhere, then we have

$$(10) \quad \text{dist}\left(\int_{\mathbb{B}_r^m(x)} u, \mathcal{N}\right)^p \leq \int_{\mathbb{B}_r^m(x)} \int_{\mathbb{B}_r^m(x)} |u(y) - u(z)|^p dy dz.$$

Thus, by (9) and (10),

$$(11) \quad \lim_{r \rightarrow 0} \sup_{x \in \mathbb{R}^m} \text{dist}\left(\int_{\mathbb{B}_r^m(x)} u, \mathcal{N}\right)^p = 0.$$

Because of the lack of control on the rate of convergence of the limit in (11), we need to refine it to get the quantitative results we are looking for.

To give a glimpse of our core quantitative estimate in a simplified setting, let us replace (11) by the integral inequality

$$(12) \quad \begin{aligned} & \int_0^\infty \sup_{x \in \mathbb{R}^m} \int_{\mathbb{B}_r^m(x)} \int_{\mathbb{B}_r^m(x)} |u(y) - u(z)|^p dy dz \frac{dr}{r} \\ & \leq \frac{1}{\mathcal{L}^m(\mathbb{B}_1^m)^2} \int_0^\infty \iint_{\substack{(y,z) \in \mathbb{R}^m \times \mathbb{R}^m \\ |y-z| \leq 2r}} |u(y) - u(z)|^p dy dz \frac{dr}{r^{2m+1}} \\ & = \frac{4^m}{2m \mathcal{L}^m(\mathbb{B}_1^m)^2} \iint_{\mathbb{R}^m \times \mathbb{R}^m} \frac{|u(y) - u(z)|^p}{|y-z|^{2m}} dy dz. \end{aligned}$$

It follows from (12) and the Chebyshev inequality that for every $\lambda \in (1, \infty)$ there exists $\tau \in (1, \lambda)$ such that

$$\begin{aligned}
(13) \quad & \sum_{k \in \mathbb{Z}} \sup_{x \in \mathbb{R}^m} \text{dist} \left(\int_{\mathbb{B}_{\tau\lambda^{-k}}^m(x)} u, \mathcal{N} \right)^p \\
& \leq \sum_{k \in \mathbb{Z}} \sup_{x \in \mathbb{R}^m} \int_{\mathbb{B}_{\tau\lambda^{-k}}^m(x)} \int_{\mathbb{B}_{\tau\lambda^{-k}}^m(x)} |u(y) - u(z)|^p \, dy \, dz \\
& \leq \frac{4^m}{2m \mathcal{L}^m(\mathbb{B}_1^m)^2 \ln \lambda} \iint_{\mathbb{R}^m \times \mathbb{R}^m} \frac{|u(y) - u(z)|^p}{|y - z|^{2m}} \, dy \, dz.
\end{aligned}$$

If we take now an extension u by averaging on the half-space \mathbb{R}_+^{m+1} , (13) implies that when its right-hand side is small enough, that is, when

$$(14) \quad \ln \lambda \simeq \iint_{\mathbb{R}^m \times \mathbb{R}^m} \frac{|u(y) - u(z)|^p}{|y - z|^{2m}} \, dy \, dz,$$

there is a family of hyperplanes whose distance to the boundary is in geometric progression of ratio λ on which the extension is close to the manifold. In view of (14), the ratio λ is controlled by an exponential of the Gagliardo fractional energy.

In order to actually prove the results, we improve the estimate (12) in two directions. First, instead of considering parallel hyperplanes, we consider a decomposition of cubes on the boundary of which we control the mean oscillation. Second, we replace the mean oscillation by a truncated mean oscillation which can be controlled by the gap potential (6).

In comparison, the proof in [22] of (4) relies on the estimate (see [22, Prop. 4.2])

$$\begin{aligned}
& \int_0^R \sup_{\substack{x \in \mathbb{B}_1^{m+1} \\ |x|=r}} \text{dist} \left(\frac{(1-r^2)^m}{\mathcal{H}^m(\mathbb{S}^m)} \int_{\mathbb{S}^m} \frac{u(y)}{|x-y|^{2m}} \, dy, \mathcal{N} \right) \frac{dr}{1-r^2} \\
& \leq \int_0^R \frac{(1-|r|^2)^{2m-1}}{\mathcal{H}^m(\mathbb{S}^m)^2} \sup_{\substack{x \in \mathbb{B}_1^{m+1} \\ |x|=r}} \iint_{\mathbb{S}^m \times \mathbb{S}^m} \frac{|u(y) - u(z)|}{|x-y|^{2m} |x-z|^{2m}} \, dy \, dz \, dr \\
& \leq C \left(\ln \frac{2}{1-R} \right)^{1-\frac{1}{p}} \left(\iint_{\mathbb{S}^m \times \mathbb{S}^m} \frac{|u(y) - u(z)|^p}{|y-z|^{2m}} \, dy \, dz \right)^{\frac{1}{p}},
\end{aligned}$$

with $R \in (0, 1)$. This allows to find bad balls in the hyperbolic space in the Poincaré ball models outside of which the hyperharmonic extension to \mathbb{B}_1^{m+1} is close to the target manifold and whose radius is controlled by the exponential of the Gagliardo fractional energy. One cannot perform directly a homogeneous extension on the bad balls because they could intersect; through a Besicovitch-type covering argument, the construction is performed on collections of disjoint balls, with a number of collections bounded exponentially; combined with the exponential of the radius appearing in the equivalence between critical Marcinkiewicz–Sobolev quasinnorms, this explains the double exponential appearing in the final estimate. In the present work we avoid this pitfall by working directly with a decomposition into disjoint cubes.

2. MEAN OSCILLATION ON λ -ADIC SKELETONS

2.1. Truncated mean oscillation. Our analysis will rely on truncated mean oscillations, which have already been used in estimates on homotopy decompositions [23] and on liftings over noncompact Riemannian coverings [24].

Definition 2.1. Given $u : \mathbb{R}^m \rightarrow \mathcal{N}$, we define the function $\text{MO}_{\delta,p}u : \mathbb{R}_+^{m+1} \rightarrow [0, \infty]$ for every $x = (x', x_{m+1}) \in \mathbb{R}_+^{m+1}$ by

$$(15) \quad \text{MO}_{\delta,p}u(x) := \int_{\mathbb{B}_{x_{m+1}}^m(x')} \int_{\mathbb{B}_{x_{m+1}}^m(x')} (d(u(y), u(z)) - \delta)_+^p dy dz.$$

If $\delta_0, \delta_1 \in [0, \infty)$, one has

$$(16) \quad \text{MO}_{\delta_1,p}u(x)^{\frac{1}{p}} \leq \text{MO}_{\delta_0,p}u(x)^{\frac{1}{p}} + (\delta_0 - \delta_1)_+.$$

The truncated Gagliardo fractional energy can be written in terms of mean oscillations as

$$\iint_{\mathbb{R}^m \times \mathbb{R}^m} \frac{(d(u(y), u(z)) - \delta)_+^p}{|y - z|^{m+sp}} dy dz \simeq \int_{\mathbb{R}_+^{m+1}} \text{MO}_{\delta,p}u(x) \frac{dx}{x_{m+1}^{1-m+sp}}.$$

If $V : \mathbb{R}_+^{m+1} \rightarrow \mathbb{R}^\nu$ denotes an extension by averaging of $u : \mathbb{R}^m \rightarrow \mathbb{R}^\nu$, that is

$$(17) \quad V(x) := \frac{1}{x_{m+1}^m} \int_{\mathbb{R}^m} u(z) \varphi\left(\frac{x' - z}{x_{m+1}}\right) dz = \int_{\mathbb{R}^m} u(x' - x_{m+1}z) \varphi(z) dz,$$

with $\varphi \in C^\infty(\mathbb{R}^m, \mathbb{R})$ such that $\int_{\mathbb{R}^m} \varphi = 1$ and $\text{supp } \varphi \subseteq \mathbb{B}_1^m$, then one has for every $x = (x', x_{m+1}) \in \mathbb{R}_+^{m+1}$, if $u \in Y \subseteq \mathbb{R}^\nu$ almost everywhere on \mathbb{R}^m ,

$$(18) \quad \text{dist}(V(x), Y)^p \leq C_1(\text{MO}_{\delta,p}u(x) + \delta^p)$$

and

$$(19) \quad |DV(x)|^p \leq \frac{C_2}{x_{m+1}^p} (\text{MO}_{\delta,p}u(x) + \delta^p).$$

Remark 2.2. Replacing one of the averages in the (15) by the value of the mapping u , one gets the sharp truncated mean oscillation $\text{MO}_{\delta,p}^\sharp u : \mathbb{R}_+^{m+1} \rightarrow [0, \infty]$ defined for every $x = (x', x_{m+1}) \in \mathbb{R}_+^{m+1}$ by

$$(20) \quad \text{MO}_{\delta,p}^\sharp u(x) := \inf_{z \in \mathbb{R}^m} \int_{\mathbb{B}_{x_{m+1}}^m(x')} (d(u(y), u(z)) - \delta)_+^p dy;$$

it follows immediately by the triangle inequality that

$$(21) \quad \text{MO}_{\delta,p}u(x) \leq 2^p \text{MO}_{\delta/2,p}^\sharp u(x)$$

and by Fubini's theorem that

$$\int_{\mathbb{B}_{x_{m+1}}^m(x')} \text{MO}_{\delta,p}^\sharp u(z, x_{m+1}) dz \leq 2^{2m} \text{MO}_{\delta,p}u(x', 2x_m).$$

2.2. λ -adic cubes and skeletons. The domain of the linear extension is the half-space \mathbb{R}_+^{m+1} ; we will subdivide the latter in λ -adic cubes on which we will perform appropriate constructions.

Given $\lambda \in (1, \infty)$, we consider for every $k \in \mathbb{Z}$ the set of cubes of \mathbb{R}^m

$$\mathcal{Q}_{\lambda, \tau, k} := \{\tau\lambda^{-k}([0, 1]^m + j) \mid j \in \mathbb{Z}^m\}$$

and the corresponding set of cubes of \mathbb{R}_+^{m+1}

$$\begin{aligned} \mathcal{Q}_{\lambda, \tau, k}^+ &:= \{Q \times [\frac{\tau\lambda^{-k}}{\lambda-1}, \frac{\tau\lambda^{-(k-1)}}{\lambda-1}] \mid Q \in \mathcal{Q}_{\lambda, \tau, k}\} \\ &= \{\tau\lambda^{-k}([0, 1]^{m+1} + (j, (\lambda-1)^{-1})) \mid j \in \mathbb{Z}^m\}. \end{aligned}$$

In particular, we have the decompositions

$$\mathbb{R}^m = \bigcup_{Q \in \mathcal{Q}_{\lambda, \tau, k}} Q \quad \text{and} \quad \mathbb{R}_+^{m+1} = \bigcup_{Q \in \mathcal{Q}_{\lambda, \tau, k}^+} Q.$$

Moreover, we consider the part of the boundaries of the cubes in $\mathcal{Q}_{\lambda, \tau, k}^+$ that are parallel to the hyperplane $\mathbb{R}^m \times \{0\}$

$$(22) \quad \begin{aligned} \mathcal{Q}_{\lambda, \tau, k}^{\parallel} &:= \{Q \times \{\frac{\tau\lambda^{-k}}{\lambda-1}\} \mid Q \in \mathcal{Q}_{\lambda, \tau, k}\} \\ &= \{\tau\lambda^{-k}([0, 1]^m \times \{0\} + (j, (\lambda-1)^{-1})) \mid j \in \mathbb{Z}^m\} \end{aligned}$$

and those that are normal to the same hyperplane

$$(23) \quad \begin{aligned} \mathcal{Q}_{\lambda, \tau, k}^{\perp} &:= \{\partial Q \times [\frac{\tau\lambda^{-k}}{\lambda-1}, \frac{\tau\lambda^{-(k-1)}}{\lambda-1}] \mid Q \in \mathcal{Q}_{\lambda, \tau, k}\} \\ &= \{\tau\lambda^{-k}((\partial[0, 1]^m) \times [0, 1] + (j, (\lambda-1)^{-1})) \mid j \in \mathbb{Z}^m\}. \end{aligned}$$

Given $h \in \mathbb{R}^m$, we also define the corresponding translated sets

$$(24) \quad \begin{aligned} \mathcal{Q}_{\lambda, \tau, k, h} &:= \{Q + \tau\lambda^{-k}h \mid Q \in \mathcal{Q}_{\lambda, \tau, k}\}, \\ \mathcal{Q}_{\lambda, \tau, k, h}^+ &:= \{Q + \tau\lambda^{-k}(h, 0) \mid Q \in \mathcal{Q}_{\lambda, \tau, k}^+\}, \\ \mathcal{Q}_{\lambda, \tau, k, h}^{\parallel} &:= \{\Sigma + \tau\lambda^{-k}(h, 0) \mid \Sigma \in \mathcal{Q}_{\lambda, \tau, k}^{\parallel}\}, \\ \mathcal{Q}_{\lambda, \tau, k, h}^{\perp} &:= \{\Sigma + \tau\lambda^{-k}(h, 0) \mid \Sigma \in \mathcal{Q}_{\lambda, \tau, k}^{\perp}\}. \end{aligned}$$

2.3. Longitudinal estimate. We first have an estimate on the maximal mean oscillation on longitudinal faces of cubes of the λ -adic decomposition of \mathbb{R}_+^{m+1} .

Proposition 2.3. *For every $m \in \mathbb{N} \setminus \{0\}$, there exists a constant $C = C(m) \in (0, \infty)$ such that for every $p \in [1, \infty)$, for every measurable map $u : \mathbb{R}^m \rightarrow \mathcal{N}$, for every $\lambda \in [2, \infty)$ and for every $\delta \in [0, \infty)$, one has*

$$(25) \quad \int_1^\lambda \sum_{k \in \mathbb{Z}} \int_{[0, 1]^m} \sum_{\Sigma \in \mathcal{Q}_{\lambda, \tau, k, h}^{\parallel}} \sup_{x \in \Sigma} \text{MO}_{\delta, p} u(x) \, dh \frac{d\tau}{\tau} \leq C \iint_{\mathbb{R}^m \times \mathbb{R}^m} \frac{(d(u(y), u(z)) - \delta)_+^p}{|y - z|^{2m}} \, dy \, dz.$$

Proof. For any $\Sigma \in \mathcal{Q}_{\lambda, \tau, k, h}^{\parallel}$, we can write, in view of (22) and (24),

$$\Sigma = Q \times \{\frac{\tau\lambda^{-k}}{\lambda-1}\},$$

where $Q \in \mathcal{Q}_{\lambda, \tau, k, h}$. If $x \in \Sigma$ and $y \in \mathbb{R}^m$ satisfy $|x' - y| \leq \frac{\tau\lambda^{-k}}{\lambda-1}$, then $x' \in Q$ and we have immediately that

$$(26) \quad \text{dist}(y, Q) \leq |x' - y| \leq \frac{\tau\lambda^{-k}}{\lambda-1}.$$

Since Q is a cube of edge length $\tau\lambda^{-k}$, according to (26), we have

$$y \in (1 + \frac{2}{\lambda-1})Q = \frac{\lambda+1}{\lambda-1}Q$$

under the convention that $\frac{\lambda+1}{\lambda-1}Q$ is the cube with the same center that Q dilated by a factor $\frac{\lambda+1}{\lambda-1}$. Thus, $y \in 3Q$, since $\lambda \geq 2$. One has then, by monotonicity of the integral, for every $x \in \Sigma$,

$$(27) \quad \begin{aligned} \text{MO}_{\delta,p}u(x) &= \frac{1}{\mathcal{L}^m(\mathbb{B}_1^m)^2} \left(\frac{\lambda^k(\lambda-1)}{\tau} \right)^{2m} \iint_{\substack{(y,z) \in \mathbb{R}^m \times \mathbb{R}^m \\ |y-x'| \leq \frac{\tau\lambda^{-k}}{\lambda-1} \\ |z-x'| \leq \frac{\tau\lambda^{-k}}{\lambda-1}}} (d(u(y), u(z)) - \delta)_+^p \, dy \, dz \\ &\leq \frac{1}{\mathcal{L}^m(\mathbb{B}_1^m)^2} \left(\frac{\lambda^k(\lambda-1)}{\tau} \right)^{2m} \iint_{\substack{(y,z) \in 3Q \times 3Q \\ |y-z| \leq \frac{2\tau\lambda^{-k}}{\lambda-1}}} (d(u(y), u(z)) - \delta)_+^p \, dy \, dz. \end{aligned}$$

Summing (27) over the sets $\Sigma \in \mathcal{Q}_{\lambda,\tau,k,h}^{\parallel}$ and integrating it over the translations $h \in [0, 1]^m$, we get

$$(28) \quad \begin{aligned} \int_{[0,1]^m} \sum_{\Sigma \in \mathcal{Q}_{\lambda,\tau,k,h}^{\parallel}} \sup_{x \in \Sigma} \text{MO}_{\delta,p}u(x) \, dh \\ \leq \frac{5^m}{\mathcal{L}^m(\mathbb{B}_1^m)^2} \left(\frac{\lambda^k(\lambda-1)}{\tau} \right)^{2m} \iint_{\substack{(y,z) \in \mathbb{R}^m \times \mathbb{R}^m \\ |y-z| \leq \frac{2\tau\lambda^{-k}}{\lambda-1}}} (d(u(y), u(z)) - \delta)_+^p \, dy \, dz. \end{aligned}$$

Summing (28) over the scales $k \in \mathbb{Z}$ and integrating it over $\tau \in [1, \lambda]$, we get

$$\begin{aligned} &\int_1^\lambda \sum_{k \in \mathbb{Z}} \int_{[0,1]^m} \sum_{\Sigma \in \mathcal{Q}_{\lambda,\tau,k,h}^{\parallel}} \sup_{x \in \Sigma} \text{MO}_{\delta,p}u(x) \, dh \frac{d\tau}{\tau} \\ &\leq \frac{5^m}{\mathcal{L}^m(\mathbb{B}_1^m)^2} (\lambda-1)^{2m} \iint_{\mathbb{R}^m \times \mathbb{R}^m} \sum_{\substack{k \in \mathbb{Z} \\ \tau \in (1, \lambda) \\ \tau \geq (\lambda-1)\lambda^k |y-z|/2}} \int \frac{\lambda^{2km} (d(u(y), u(z)) - \delta)_+^p}{\tau^{2m+1}} \, d\tau \, dy \, dz \\ &= \frac{5^m}{\mathcal{L}^m(\mathbb{B}_1^m)^2} (\lambda-1)^{2m} \iint_{\mathbb{R}^m \times \mathbb{R}^m} \int_{\frac{(\lambda-1)|y-z|}{2}}^\infty \frac{(d(u(y), u(z)) - \delta)_+^p}{\theta^{2m+1}} \, d\theta \, dy \, dz \\ &= \frac{20^m}{2m\mathcal{L}^m(\mathbb{B}_1^m)^2} \iint_{\mathbb{R}^m \times \mathbb{R}^m} \frac{(d(u(y), u(z)) - \delta)_+^p}{|y-z|^{2m}} \, dy \, dz, \end{aligned}$$

which implies the announced conclusion (25) and completes our proof of Proposition 2.3. \square

2.4. Transversal estimate. We next prove a counterpart of Proposition 2.3, where we estimate the maximal mean oscillation on transversal faces of cubes of the λ -adic decomposition of \mathbb{R}_+^{m+1} instead of the longitudinal ones.

Proposition 2.4. *For every $m \in \mathbb{N} \setminus \{0\}$ and $p \in [1, \infty)$, there exists a constant $C = C(m, p) \in (0, \infty)$ such that for every measurable map $u : \mathbb{R}^m \rightarrow \mathcal{N}$, for every $\lambda \in [2, \infty)$ and for every $\delta \in [0, \infty)$, one has*

$$(29) \quad \int_1^\lambda \sum_{k \in \mathbb{Z}} \int_{[0,1]^m} \sum_{\Sigma \in \mathcal{Q}_{\lambda, \tau, k, h}^\perp} \sup_{x \in \Sigma} \text{MO}_{\delta, p} u(x) \, dh \frac{d\tau}{\tau} \leq C \iint_{\mathbb{R}^m \times \mathbb{R}^m} \frac{(d(u(y), u(z)) - \frac{\delta}{2})_+^p}{|y - z|^{2m}} \, dy \, dz.$$

Proof. We consider a set $\Sigma \in \mathcal{Q}_{\lambda, \tau, k, h}^\perp$, that we can write, in view of (23) and (24), as

$$\Sigma = \partial Q \times \left[\frac{\tau \lambda^{-k}}{\lambda - 1}, \frac{\tau \lambda^{-(k-1)}}{\lambda - 1} \right],$$

where $Q \in \mathcal{Q}_{\lambda, \tau, k, h}$. We first note that by (21),

$$(30) \quad \begin{aligned} \text{MO}_{\delta, p} u(x) &\leq 2^p \text{MO}_{\delta/2, p}^\sharp u(x) \\ &\leq 2^p \inf_{\substack{z \in \partial Q \\ |z - x'| \leq x_{m+1}}} \int_{\mathbb{B}_{x_{m+1}}^m(x')} (d(u(y), u(z)) - \frac{\delta}{2})_+^p \, dy \\ &\leq 2^p \int_{E_x} \int_{B_{x_{m+1}}(x')} (d(u(y), u(z)) - \frac{\delta}{2})_+^p \, dz \, dy, \end{aligned}$$

where

$$E_x := \{z \in \partial Q \mid |z - x'| \leq x_{m+1}\}.$$

If $x \in \Sigma$ and $y \in \mathbb{R}^m$ satisfy $|x' - y| \leq \frac{\tau \lambda^{-(k-1)}}{\lambda - 1}$, then $x' \in \partial Q \subset Q$ and

$$(31) \quad \text{dist}(y, Q) \leq \frac{\tau \lambda^{-(k-1)}}{\lambda - 1}.$$

Since Q is a cube of edge length $\tau \lambda^{-k}$ and $\lambda \geq 2$, using (31), we get

$$(32) \quad y \in \left(1 + \frac{2\lambda}{\lambda - 1}\right)Q = \frac{3\lambda - 1}{\lambda - 1}Q \subseteq 5Q.$$

One has then for every $x \in \Sigma$, by (30), (32) and by monotonicity of the integral,

$$(33) \quad \begin{aligned} \text{MO}_{\delta, p} u(x) &\leq \frac{C_1}{x_{m+1}^{2m-1}} \iint_{\substack{(y, z) \in \mathbb{R}^m \times \partial Q \\ |y - x'| \leq x_{m+1} \\ |z - x'| \leq x_{m+1}}} (d(u(y), u(z)) - \frac{\delta}{2})_+^p \, dy \, dz \\ &\leq C_2 \iint_{\substack{(y, z) \in 5Q \times \partial Q \\ |y - z| \leq \frac{2\tau \lambda^{-(k-1)}}{\lambda - 1}}} \frac{(d(u(y), u(z)) - \frac{\delta}{2})_+^p}{|y - z|^{2m-1}} \, dy \, dz, \end{aligned}$$

where the constant $C_2 > 0$ depends only on m and p . Summing (33) over $\Sigma \in \mathcal{Q}_{\lambda, \tau, k, h}^\perp$ and integrating it with respect to the translations h over $[0, 1]^m$, we have

$$(34) \quad \int_{[0,1]^m} \sum_{\Sigma \in \mathcal{Q}_{\lambda, \tau, k, h}^\perp} \sup_{x \in \Sigma} \text{MO}_{\delta, p} u(x) \, dh \leq \frac{C_2 \lambda^k}{\tau} \iint_{\substack{(y, z) \in \mathbb{R}^m \times \mathbb{R}^m \\ |y - z| \leq \frac{2\tau \lambda^{-(k-1)}}{\lambda - 1}}} \frac{(d(u(y), u(z)) - \frac{\delta}{2})_+^p}{|y - z|^{2m-1}} \, dy \, dz,$$

where we have also used Fubini's theorem. Thus, summing and integrating (34) over the scales, we get

$$\begin{aligned}
& \int_1^\lambda \sum_{k \in \mathbb{Z}} \int_{[0,1]^m} \sum_{\Sigma \in \mathcal{Q}_{\lambda, \tau, k, h}^\perp} \sup_{x \in \Sigma} \text{MO}_{\delta, p} u(x) \, dh \frac{d\tau}{\tau} \\
& \leq C_2 \iint_{\mathbb{R}^m \times \mathbb{R}^m} \sum_{k \in \mathbb{Z}} \int_{\substack{\tau \in (1, \lambda) \\ \tau \geq \lambda^k (\lambda-1) |y-z| / 2\lambda}} \frac{\lambda^k (d(u(y), u(z)) - \frac{\delta}{2})_+^p}{|y-z|^{2m-1} \tau^2} \, d\tau \, dy \, dz \\
(35) \quad & = C_2 \iint_{\mathbb{R}^m \times \mathbb{R}^m} \int_{\frac{(\lambda-1)|y-z|}{2\lambda}}^\infty \frac{(d(u(y), u(z)) - \frac{\delta}{2})_+^p}{|y-z|^{2m-1} \theta^2} \, d\theta \, dy \, dz \\
& = \frac{2\lambda}{\lambda-1} C_2 \iint_{\mathbb{R}^m \times \mathbb{R}^m} \frac{(d(u(y), u(z)) - \frac{\delta}{2})_+^p}{|y-z|^{2m}} \, dy \, dz,
\end{aligned}$$

so that the conclusion (29) follows. \square

2.5. Combining the estimates. We close this section by summarizing Propositions 2.3 and 2.4 in the following statement.

Proposition 2.5. *For every $m \in \mathbb{N} \setminus \{0\}$ and $p \in [1, \infty)$, there exists a constant $C = C(m, p) > 0$ such that for every measurable map $u : \mathbb{R}^m \rightarrow \mathcal{N}$, for every $\lambda \in [2, \infty)$ and for every $\delta \in [0, \infty)$, one has*

$$\int_1^\lambda \sum_{k \in \mathbb{Z}} \int_{[0,1]^m} \sum_{Q \in \mathcal{Q}_{\lambda, \tau, k, h}^+} \sup_{x \in \partial Q} \text{MO}_{\delta, p} u(x) \, dh \frac{d\tau}{\tau} \leq C \iint_{\mathbb{R}^m \times \mathbb{R}^m} \frac{(d(u(y), u(z)) - \frac{\delta}{2})_+^p}{|y-z|^{2m}} \, dy \, dz.$$

Proof. This follows immediately from Propositions 2.3 and 2.4. \square

3. PROOFS OF THE SINGULAR EXTENSION THEOREMS

3.1. Oscillation and gradient estimate on the skeleton. We first estimate the average number of cubes on which the extension V of u given by (17) is far away from the range of u .

Proposition 3.1. *Let $m \in \mathbb{N} \setminus \{0\}$. There exist constants $\eta \in (0, 1)$ and $C \in (0, \infty)$ depending only on m such that for every $\delta \in (0, \infty)$, for every $\lambda \in [2, \infty)$, for every measurable function $u : \mathbb{R}^m \rightarrow \mathbb{R}^\nu$ and every set $Y \subseteq \mathbb{R}^\nu$, if V is an extension by convolution given by (17) and if $u \in Y$ almost everywhere in \mathbb{R}^m , then*

$$\begin{aligned}
(36) \quad & \int_1^\lambda \sum_{k \in \mathbb{Z}} \int_{[0,1]^m} \#\{Q \in \mathcal{Q}_{\lambda, \tau, k, h}^+ \mid \sup_{x \in \partial Q} \text{dist}(V(x), Y) \geq \delta\} \, dh \frac{d\tau}{\tau} \\
& \leq \frac{C}{\delta^{m+1}} \iint_{\mathbb{R}^m \times \mathbb{R}^m} \frac{(d(u(y), u(z)) - \eta\delta)_+^{m+1}}{|y-z|^{2m}} \, dy \, dz.
\end{aligned}$$

Proof. Using (16), (18) and the elementary inequality $(a+b)^{m+1} \leq 2^m(a^{m+1} + b^{m+1})$, which is valid for all $a, b \in [0, \infty)$, we have for every $x \in \mathbb{R}_+^{m+1}$,

$$(37) \quad \text{dist}(V(x), Y)^{m+1} \leq C_1 \text{MO}_{0, m+1} u(x) \leq 2^m C_1 (\text{MO}_{\theta, m+1} u(x) + \theta^{m+1}).$$

Hence, taking

$$\eta := \frac{1}{2} \left(\frac{2}{2 + C_1} \right)^{\frac{1}{m+1}},$$

if

$$(38) \quad \text{dist}(V(x), Y) \geq \delta,$$

the inequality (37) with $\theta = 2\eta\delta$ implies that

$$(39) \quad (\eta\delta)^{m+1} \leq \text{MO}_{2\eta\delta, m+1} u(x).$$

We get, by (38), (39) and Proposition 2.5,

$$\begin{aligned} & \int_1^\lambda \sum_{k \in \mathbb{Z}} \int_{[0,1]^m} \#\{Q \in \mathcal{Q}_{\lambda, \tau, k, h}^+ \mid \sup_{x \in \partial Q} \text{dist}(V(x), Y) \geq \delta\} dh \frac{d\tau}{\tau} \\ & \leq \int_1^\lambda \sum_{k \in \mathbb{Z}} \int_{[0,1]^m} \#\{Q \in \mathcal{Q}_{\lambda, \tau, k, h}^+ \mid \sup_{x \in \partial Q} \text{MO}_{2\eta\delta, m+1} u(x) \geq (\eta\delta)^{m+1}\} dh \frac{d\tau}{\tau} \\ & \leq \frac{1}{(\eta\delta)^{m+1}} \int_1^\lambda \sum_{k \in \mathbb{Z}} \int_{[0,1]^m} \sum_{Q \in \mathcal{Q}_{\lambda, \tau, k, h}^+} \sup_{x \in \partial Q} \text{MO}_{2\eta\delta, m+1} u(x) dh \frac{d\tau}{\tau} \\ & \leq \frac{C_2}{\delta^{m+1}} \iint_{\mathbb{R}^m \times \mathbb{R}^m} \frac{(d(u(y), u(z)) - \eta\delta)_+^{m+1}}{|y - z|^{2m}} dy dz, \end{aligned}$$

where the constant $C_2 > 0$ depends only on m . This proves the estimate (36) and completes our proof of Proposition 3.1. \square

Next, we can prove an average uniform bound on the extension by convolution.

Proposition 3.2. *Let $m \in \mathbb{N} \setminus \{0\}$. There exists a constant $C = C(m) > 0$ such that for every $\lambda \in [2, \infty)$, for every measurable function $u : \mathbb{R}^m \rightarrow \mathbb{R}^\nu$, if V is an extension by convolution given by (17), then*

$$\int_1^\lambda \sum_{k \in \mathbb{Z}} \int_{[0,1]^m} \sum_{Q \in \mathcal{Q}_{\lambda, \tau, k, h}^+} \sup_{x \in \partial Q} x_{m+1}^{m+1} |DV(x)|^{m+1} dh \frac{d\tau}{\tau} \leq C \iint_{\mathbb{R}^m \times \mathbb{R}^m} \frac{d(u(y), u(z))^{m+1}}{|y - z|^{2m}} dy dz.$$

Proof. Similarly as in the proof of Proposition 3.1, since by (19)

$$x_{m+1}^{m+1} |DV(x)|^{m+1} \leq C_1 \text{MO}_{0, m+1} u(x),$$

the proof of Proposition 3.2 then follows from Proposition 2.5. Namely,

$$\begin{aligned} & \int_1^\lambda \sum_{k \in \mathbb{Z}} \int_{[0,1]^m} \sum_{Q \in \mathcal{Q}_{\lambda, \tau, k, h}^+} \sup_{x \in \partial Q} x_{m+1}^{m+1} |DV(x)|^{m+1} dh \frac{d\tau}{\tau} \\ & \leq C_1 \int_1^\lambda \sum_{k \in \mathbb{Z}} \int_{[0,1]^m} \sum_{Q \in \mathcal{Q}_{\lambda, \tau, k, h}^+} \sup_{x \in \partial Q} \text{MO}_{0, m+1} u(x) dh \frac{d\tau}{\tau} \\ & \leq C_2 \iint_{\mathbb{R}^m \times \mathbb{R}^m} \frac{d(u(y), u(z))^{m+1}}{|y - z|^{2m}} dy dz, \end{aligned}$$

where the constant $C_2 > 0$ only depends on m . \square

3.2. Sobolev and Sobolev–Marcinkiewicz extensions on cubes. The first construction we will perform is a classical extension of the boundary data on cubes with small energy. To simplify the presentation, the result is stated on a ball, which is bi-Lipschitz equivalent to a cube.

Lemma 3.3. *Let $m \in \mathbb{N} \setminus \{0\}$. There exists a constant $C = C(m) \in (0, \infty)$ such that for every $w \in W^{1,m+1}(\mathbb{S}_\rho^m, \mathbb{R}^\nu)$ there exists a function $W \in W^{1,m+1}(\mathbb{B}_\rho^{m+1}, \mathbb{R}^\nu) \cap C(\mathbb{B}_\rho^{m+1}, \mathbb{R}^\nu)$ such that $\text{tr}_{\mathbb{S}_\rho^m} W = w$,*

$$(40) \quad \int_{\mathbb{B}_\rho^{m+1}} |DW|^{m+1} \leq C\rho \int_{\mathbb{S}_\rho^m} |Dw|^{m+1},$$

$$(41) \quad \int_{\mathbb{B}_\rho^{m+1}} \int_{\mathbb{S}_\rho^m} |W(x) - w(y)|^{m+1} dx dy \leq C\rho^{2m+2} \int_{\mathbb{S}_\rho^m} |Dw|^{m+1}$$

and for every $x \in \mathbb{B}_\rho^{m+1}$ and almost every $y \in \mathbb{S}_\rho^m$,

$$(42) \quad |W(x) - w(y)|^{m+1} \leq C\rho \int_{\mathbb{S}_\rho^m} |Dw|^{m+1}.$$

Proof. Let

$$p := \frac{(m+1)^2}{m}.$$

We have, by the fractional Sobolev–Morrey embedding,

$$(43) \quad \iint_{\mathbb{S}_\rho^m \times \mathbb{S}_\rho^m} \frac{|w(y) - w(z)|^p}{|y - z|^{m+p-1}} dy dz \leq C_1 \left(\int_{\mathbb{S}_\rho^m} |Dw|^{m+1} \right)^{\frac{p}{m+1}}.$$

Assuming without loss of generality that $\int_{\mathbb{S}_\rho^m} w = 0$, Gagliardo’s classical trace theory [8] (see also [7, Prop. 17.1; 10, Section 6.9; 11, Th. 9.4; 12, Th. 10.1.1.1]) yields a function $W \in W^{1,p}(\mathbb{B}_\rho^{m+1}, \mathbb{R}^\nu)$ such that $\text{tr}_{\mathbb{S}_\rho^m} W = w$ and

$$(44) \quad \int_{\mathbb{B}_\rho^{m+1}} |DW|^p + \frac{|W|^p}{\rho^p} \leq C_2 \iint_{\mathbb{S}_\rho^m \times \mathbb{S}_\rho^m} \frac{|w(y) - w(z)|^p}{|y - z|^{m+p-1}} dy dz.$$

It follows from Hölder’s inequality, (44) and (43), that

$$(45) \quad \int_{\mathbb{B}_\rho^{m+1}} |DW|^{m+1} + \frac{|W|^{m+1}}{\rho^{m+1}} \leq C_3\rho \int_{\mathbb{S}_\rho^m} |Dw|^{m+1},$$

and (40) follows then from (45).

Since for every $x \in \mathbb{B}_\rho^{m+1}$ and $y \in \mathbb{S}_\rho^m$,

$$|W(x) - w(y)|^{m+1} \leq 2^m (|W(x)|^{m+1} + |w(y)|^{m+1}),$$

we have

$$(46) \quad \int_{\mathbb{B}_\rho^{m+1}} \int_{\mathbb{S}_\rho^m} |W(x) - w(y)|^{m+1} dx dy \leq C_4\rho^m \int_{\mathbb{B}_\rho^{m+1}} |W|^{m+1} + C_5\rho^{m+1} \int_{\mathbb{S}_\rho^m} |w|^{m+1}.$$

The estimate (41) follows from the estimates (46), (45) and the Poincaré inequality on the sphere.

Finally, by the Morrey–Sobolev inequality, (44) and (43), we have

$$\begin{aligned} |W(x) - w(y)| &\leq C_6 |x - y|^{1 - \frac{m+1}{p}} \left(\int_{\mathbb{B}_\rho^{m+1}} |DW|^p \right)^{\frac{1}{p}} \\ &\leq C_7 \rho^{1 - \frac{m}{m+1}} \left(\int_{\mathbb{S}_\rho^m} |Dw|^{m+1} \right)^{\frac{1}{m+1}}, \end{aligned}$$

and (42) follows. \square

When the oscillation is too large on the boundary of cubes, we will perform our construction of the controlled singular extension; the resulting map is quite wild but is sufficiently well controlled to provide an acceptable extension on those cubes.

Lemma 3.4. *Let $m \in \mathbb{N} \setminus \{0\}$. If $w \in W^{1,m+1}(\mathbb{S}_\rho^m, \mathbb{R}^\nu)$ and if we define $W : \mathbb{B}_\rho^{m+1} \rightarrow \mathbb{R}^\nu$ for each $x \in \mathbb{B}_\rho^{m+1} \setminus \{0\}$ by*

$$W(x) := w\left(\frac{\rho}{|x|}x\right),$$

then $W \in W^{1,1}(\mathbb{B}_\rho^{m+1}, \mathbb{R}^\nu)$ and for every $t \in (0, \infty)$,

$$(47) \quad \mathcal{L}^{m+1}(\{x \in \mathbb{B}_\rho^{m+1} \mid |DW(x)| > t\}) \leq \frac{\rho}{(m+1)t^{m+1}} \int_{\mathbb{S}_\rho^m} |Dw|^{m+1}.$$

Proof. It can be observed immediately that $W \in W^{1,m+1}(\mathbb{B}_\rho^{m+1} \setminus \mathbb{B}_\varepsilon^{m+1}, \mathbb{R}^\nu)$ for every $\varepsilon \in (0, \rho)$ and that for every $x \in \mathbb{B}_\rho^{m+1} \setminus \{0\}$, one has

$$|DW(x)| = \frac{\rho}{|x|} |Dw\left(\frac{\rho}{|x|}x\right)|.$$

Hence, using Fubini's theorem, we have

$$\begin{aligned} \mathcal{L}^{m+1}(\{x \in \mathbb{B}_\rho^{m+1} \mid |DW(x)| > t\}) &= \int_0^\rho \mathcal{H}^m(\{x \in \mathbb{S}_r^m \mid |Dw\left(\frac{\rho}{r}x\right)| \geq \frac{tr}{\rho}\}) \, dr \\ &= \int_0^\rho \mathcal{H}^m(\{y \in \mathbb{S}_\rho^m \mid |Dw(y)| \geq \frac{tr}{\rho}\}) \left(\frac{r}{\rho}\right)^m \, dr \\ &= \frac{\rho}{t^{m+1}} \int_0^t \mathcal{H}^m(\{y \in \mathbb{S}_\rho^m \mid |Dw(y)| \geq \tau\}) \tau^m \, d\tau \\ &\leq \frac{\rho}{t^{m+1}} \int_0^\infty \mathcal{H}^m(\{y \in \mathbb{S}_\rho^m \mid |Dw(y)| \geq \tau\}) \tau^m \, d\tau \\ &= \frac{\rho}{t^{m+1}} \int_0^\infty \mathcal{H}^m(\{y \in \mathbb{S}_\rho^m \mid |Dw(y)|^{m+1} \geq \tau\}) \frac{d\tau}{m+1} \\ &= \frac{\rho}{(m+1)t^{m+1}} \int_{\mathbb{S}_\rho^m} |Dw(x)|^{m+1} \, dx. \end{aligned}$$

This yields (47) and completes our proof of Lemma 3.4. \square

3.3. Proofs of the theorems. We first construct and estimate the singular extension on the half-space (Theorem 1.2).

Proof of Theorem 1.2. We fix $\delta_{\mathcal{N}} \in (0, \infty)$ so that the nearest-point retraction $\Pi_{\mathcal{N}} : \mathcal{N} + \mathbb{B}_{\delta_{\mathcal{N}}}^\nu \rightarrow \mathcal{N}$ is well defined and smooth up to the boundary. We take $V : \mathbb{R}_+^{m+1} \rightarrow \mathbb{R}^\nu$ to be an extension by convolution of u to \mathbb{R}_+^{m+1} as in (17).

Since by assumption $u \in \mathcal{N}$ almost everywhere on \mathbb{R}^m , by the averaged estimate on the distance to the target (see Proposition 3.1), we have

$$(48) \quad \int_1^\lambda \sum_{k \in \mathbb{Z}} \int_{[0,1]^m} \#\{Q \in \mathcal{Q}_{\lambda,\tau,k,h}^+ \mid \sup_{x \in \partial Q} \text{dist}(V(x), \mathcal{N}) \geq \delta_{\mathcal{N}}/2\} dh \frac{d\tau}{\tau} \\ \leq C_1 \iint_{\substack{(y,z) \in \mathbb{R}^m \times \mathbb{R}^m \\ d(u(y), u(z)) \geq \eta \delta_{\mathcal{N}}/2}} \frac{1}{|y-z|^{2m}} dy dz,$$

where $\eta > 0$ is the constant of Proposition 3.1 depending only on m , the constant $C_1 \in (0, \infty)$ depends only on m, \mathcal{N} , and we have also used that \mathcal{N} is compact, namely $\text{diam}(\mathcal{N}) < \infty$. We set $\delta := \eta \delta_{\mathcal{N}}/2$. Taking

$$(49) \quad \lambda := \exp\left(2C_1 \iint_{\substack{(y,z) \in \mathbb{R}^m \times \mathbb{R}^m \\ d(u(y), u(z)) \geq \delta}} \frac{1}{|y-z|^{2m}} dy dz\right)$$

and using (48), we have then

$$(50) \quad \frac{1}{\ln \lambda} \int_1^\lambda \sum_{k \in \mathbb{Z}} \int_{[0,1]^m} \#\{Q \in \mathcal{Q}_{\lambda,\tau,k,h}^+ \mid \sup_{x \in \partial Q} \text{dist}(V(x), \mathcal{N}) \geq \delta_{\mathcal{N}}/2\} dh \frac{d\tau}{\tau} \leq \frac{1}{2}.$$

Similarly, by Proposition 3.2,

$$(51) \quad \frac{1}{\ln \lambda} \int_1^\lambda \sum_{k \in \mathbb{Z}} \int_{[0,1]^m} \sum_{Q \in \mathcal{Q}_{\lambda,\tau,k,h}^+} \sup_{x \in \partial Q} x_{m+1}^{m+1} |DV(x)|^{m+1} dh \frac{d\tau}{\tau} \\ \leq \frac{C_2}{\ln \lambda} \iint_{\mathbb{R}^m \times \mathbb{R}^m} \frac{d(u(y), u(z))^{m+1}}{|y-z|^{2m}} dy dz.$$

In view of the estimates (50) and (51), there exists $\tau \in (1, \lambda)$ and, for every $k \in \mathbb{Z}$, $h_k \in [0,1]^m$ such that for every $k \in \mathbb{Z}$, every $Q \in \mathcal{Q}_{\lambda,\tau,k,h_k}^+$ and every $x \in \partial Q$, one has

$$(52) \quad \text{dist}(V(x), \mathcal{N}) \leq \frac{\delta_{\mathcal{N}}}{2}$$

and

$$(53) \quad \sum_{k \in \mathbb{Z}} \sum_{Q \in \mathcal{Q}_{\lambda,\tau,k,h_k}^+} \sup_{x \in \partial Q} x_{m+1}^{m+1} |DV(x)|^{m+1} \leq \frac{C_3}{\ln \lambda} \iint_{\mathbb{R}^m \times \mathbb{R}^m} \frac{d(u(y), u(z))^{m+1}}{|y-z|^{2m}} dy dz.$$

We define now the set of good cubes

$$(54) \quad \mathcal{G} := \left\{Q \in \mathcal{Q}_{\lambda,\tau,k,h_k}^+ \mid \sup_{x \in \partial Q} x_{m+1}^{m+1} |DV(x)|^{m+1} \leq \mu \text{ and } k \in \mathbb{Z}\right\}$$

and the set of bad cubes

$$(55) \quad \mathcal{B} := \left\{Q \in \mathcal{Q}_{\lambda,\tau,k,h_k}^+ \mid \sup_{x \in \partial Q} x_{m+1}^{m+1} |DV(x)|^{m+1} > \mu \text{ and } k \in \mathbb{Z}\right\}.$$

Clearly, any cube is either good or bad and thus

$$\bigcup_{Q \in \mathcal{G} \cup \mathcal{B}} Q = \mathbb{R}_+^{m+1}.$$

By (53) and (55),

$$\begin{aligned}
(56) \quad \#\mathcal{B} &\leq \frac{1}{\mu} \sum_{k \in \mathbb{Z}} \sum_{Q \in \mathcal{Q}_{\lambda, \tau, k, h_k}^+} \sup_{x \in \partial Q} x_{m+1}^{m+1} |\mathrm{DV}(x)|^{m+1} \\
&\leq \frac{C_3}{\mu \ln \lambda} \iint_{\mathbb{R}^m \times \mathbb{R}^m} \frac{d(u(y), u(z))^{m+1}}{|y-z|^{2m}} \, dy \, dz.
\end{aligned}$$

Notice also that for every $k \in \mathbb{Z}$ and every $Q \in \mathcal{Q}_{\lambda, \tau, k, h_k}^+$, we have

$$\begin{aligned}
(57) \quad \tau \lambda^{-k} \int_{\partial Q} |\mathrm{DV}|^{m+1} &\leq C_4 (\tau \lambda^{-k})^{m+1} \sup_{x \in \partial Q} |\mathrm{DV}(x)|^{m+1} \\
&\leq C_4 (\lambda - 1)^{m+1} \sup_{x \in \partial Q} x_{m+1}^{m+1} |\mathrm{DV}(x)|^{m+1},
\end{aligned}$$

since for every $x \in Q$, $x_{m+1} \geq \tau \lambda^{-k} / (\lambda - 1)$, in view of (24).

We are now going to define a map $W : \mathbb{R}_+^{m+1} \rightarrow \mathcal{N} + \mathbb{B}_{\delta_{\mathcal{N}}}^{\nu}$ separately on $\cup_{\mathcal{G}}$ and $\cup_{\mathcal{B}}$.

For every $Q \in \mathcal{G}$, we apply Lemma 3.3, up to a suitable bi-Lipschitz homeomorphism between a ball and a cube, to define the mapping W on Q as an extension of $V \upharpoonright_{\partial Q}$. We have, in view of (40) and of (57),

$$\begin{aligned}
(58) \quad \int_Q |\mathrm{DW}|^{m+1} &\leq C_5 \tau \lambda^{-k} \int_{\partial Q} |\mathrm{DV}|^{m+1} \\
&\leq C_6 (\lambda - 1)^{m+1} \sup_{x \in \partial Q} x_{m+1}^{m+1} |\mathrm{DV}(x)|^{m+1},
\end{aligned}$$

whereas by the triangle inequality, (52), (42), (57) and (54),

$$\mathrm{dist}(W(x), \mathcal{N}) \leq \frac{\delta_{\mathcal{N}}}{2} + C_7 \mu^{\frac{1}{m+1}} (\lambda - 1).$$

Hence if

$$(59) \quad \mu = \left(\frac{\delta_{\mathcal{N}}}{2C_7(\lambda - 1)} \right)^{m+1},$$

we have for every $x \in Q$, $W(x) \in \mathcal{N} + \mathbb{B}_{\delta_{\mathcal{N}}}^{\nu}$. Using the Chebyshev inequality and (58), we obtain

$$(60) \quad t^{m+1} \mathcal{L}^{m+1}(\{x \in Q \mid |\mathrm{DW}(x)| \geq t\}) \leq C_8 (\lambda - 1)^{m+1} \sup_{x \in \partial Q} x_{m+1}^{m+1} |\mathrm{DV}(x)|^{m+1}.$$

Next, we apply Lemma 3.4 on every bad cube $Q \in \mathcal{B}$ to define there W by homogeneous extension of $V \upharpoonright_{\partial Q}$. We compute then for such a cube, in view of (47) and (57), for every $t \in (0, \infty)$,

$$\begin{aligned}
(61) \quad \mathcal{L}^{m+1}(\{x \in Q \mid |\mathrm{DW}(x)| \geq t\}) &\leq \frac{C_9}{t^{m+1}} \tau \lambda^{-k} \int_{\partial Q} |\mathrm{DV}|^{m+1} \\
&\leq \frac{C_{10}}{t^{m+1}} (\lambda - 1)^{m+1} \sup_{x \in \partial Q} x_{m+1}^{m+1} |\mathrm{DV}(x)|^{m+1}.
\end{aligned}$$

Combining the estimates (60) and (61), we get

$$\begin{aligned}
& t^{m+1} \mathcal{L}^{m+1}(\{x \in \mathbb{R}_+^{m+1} \mid |DW(x)| \geq t\}) \\
& \leq C_{11}(\lambda - 1)^{m+1} \sum_{k \in \mathbb{Z}} \sum_{Q \in \mathcal{Q}_{\lambda, \tau, k, h_k}^+} \sup_{x \in \partial Q} x_{m+1}^{m+1} |DV(x)|^{m+1} \\
(62) \quad & \leq C_{12} \exp\left(C_{13} \iint_{\substack{(y,z) \in \mathbb{R}^m \times \mathbb{R}^m \\ d(u(y), u(z)) \geq \delta}} \frac{1}{|y-z|^{2m}} dy dz \right) \iint_{\mathbb{R}^m \times \mathbb{R}^m} \frac{d(u(y), u(z))^{m+1}}{|y-z|^{2m}} dy dz,
\end{aligned}$$

in view of (49) and (53), where C_{12} and C_{13} are positive constants depending only on m, \mathcal{N} . Moreover, if S denotes the set of the barycenters of the cubes $Q \in \mathcal{B}$, we have

$$(63) \quad \#S = \#\mathcal{B} \leq C_{14} \exp\left(C_{15} \iint_{\substack{(y,z) \in \mathbb{R}^m \times \mathbb{R}^m \\ d(u(y), u(z)) \geq \delta}} \frac{1}{|y-z|^{2m}} dy dz \right) \iint_{\mathbb{R}^m \times \mathbb{R}^m} \frac{d(u(y), u(z))^{m+1}}{|y-z|^{2m}} dy dz,$$

in view of (49), (56) and (59), where C_{14} and C_{15} are positive constants depending only on m, \mathcal{N} .

In order to check that u is the trace of W , we observe that on the one hand we have, since V is an extension by averaging, for every $Q \in \mathcal{G}$, by Poincaré's inequality, by (41) and by (57),

$$\begin{aligned}
(64) \quad & \int_Q |V - W|^{m+1} \\
& \leq \frac{C_{16}}{(\tau\lambda^{-k})^m} \left(\int_{\partial Q} \int_Q |V(x) - V(y)|^{m+1} dy dx + \int_{\partial Q} \int_Q |W(x) - V(y)|^{m+1} dy dx \right) \\
& \leq \frac{C_{16}}{(\tau\lambda^{-k})^m} \int_Q \int_{\partial Q} |V(x) - V(y)|^{m+1} dx dy + C_{17}(\tau\lambda^{-k})^{m+2} \int_{\partial Q} |DV|^{m+1} \\
& \leq C_{18}(\tau\lambda^{-k})^{m+1} \left(\int_Q |DV|^{m+1} + (\lambda - 1)^{m+1} \sup_{x \in \partial Q} x_{m+1}^{m+1} |DV(x)|^{m+1} \right).
\end{aligned}$$

By the classical theory of traces,

$$(65) \quad \int_{\mathbb{R}^{m+1}} |DV|^{m+1} < \infty.$$

It follows then from (53), (64) and (65) that

$$(66) \quad \sum_{k \in \mathbb{Z}} \sum_{Q \in \mathcal{G}} \frac{1}{(\lambda\tau^{-k})^{m+1}} \int_Q |V - W|^{m+1} < \infty.$$

Thus, since the set \mathcal{B} is finite, in view of (66), we have

$$\lim_{k \rightarrow \infty} \frac{1}{(\lambda\tau^{-k})^{m+1}} \sum_{Q \in \mathcal{Q}_{\lambda, \tau, k, h_k}^+} \int_Q |V - W|^{m+1} dx = 0.$$

This implies that W and V have the same trace, and hence u is the trace of W .

Finally, we define $U := \Pi_{\mathcal{N}} \circ W$. The map U also has u as the trace on $\mathbb{R}^m \times \{0\}$; the conclusion (7) follows from (62); $U \in C(\mathbb{R}_+^{m+1} \setminus S, \mathcal{N})$ with the cardinality of the singular set S being estimated by (63). This completes our proof of Theorem 1.2. \square

The proof of Theorem 1.2 can be adapted easily to the case of the hyperbolic space (Theorem 1.3).

Proof of Theorem 1.3. We proceed as in the proof of Theorem 1.2. Instead of (61), we have using the Poincaré metric on the half-space, if $Q \in \mathcal{Q}_{\lambda,\tau,k,h}^+$ (see (24)), then

$$\begin{aligned} \int_{\substack{x \in Q \\ |DV(x)|_{x_{m+1}} \geq t}} \frac{1}{x_{m+1}^{m+1}} dx &\leq \left(\frac{(\lambda-1)\lambda^k}{\tau} \right)^{m+1} \mathcal{L}^{m+1}(\{x \in Q \mid |DV(x)|_{x_{m+1}} \geq t\}) \\ &\leq \frac{\lambda^{m+1}}{t^{m+1}} (\tau\lambda^{-k})^{m+1} \sup_{x \in Q} |DV(x)|^{m+1} \\ &\leq \frac{\lambda^{m+1}(\lambda-1)^{m+1}}{t^{m+1}} \sup_{x \in Q} x_{m+1}^{m+1} |DV(x)|^{m+1}. \end{aligned}$$

The remainder of the proof is similar. \square

The case of the singular extension to the ball is slightly more complicated, as we will rely on the parameterization of \mathbb{B}_1^{m+1} by \mathbb{R}_+^{m+1} through a classical suitable conformal mapping.

Proof of Theorem 1.1. We recall that the map

$$\Psi(x) := \frac{4(x+e)}{|x+e|^2} - 2e,$$

where $e := (0, \dots, 0, 1) \in \mathbb{R}_+^{m+1}$, defines a diffeomorphism from \mathbb{B}_1^{m+1} to \mathbb{R}_+^{m+1} . Indeed, if $x \in \mathbb{B}_1^{m+1} \setminus \{e\}$, then

$$e \cdot \Psi(x) = \frac{2-2|x|^2}{|x+e|^2}.$$

Moreover, we have

$$\Psi^{-1}(x) = \frac{4(x+2e)}{|x+2e|^2} - e.$$

In particular, the ball \mathbb{B}_1^{m+1} is isometric to the half-space \mathbb{R}_+^{m+1} endowed with the metric g defined for $x \in \mathbb{R}_+^{m+1}$ and $v \in \mathbb{R}^{m+1}$ by

$$g(x)[v, v] = \frac{16|v|^2}{|x+2e|^4}.$$

Moreover, since Ψ is a conformal map and $u : \mathbb{R}^m \rightarrow \mathcal{N}$, one has

$$\iint_{\mathbb{S}^m \times \mathbb{S}^m} \frac{d(u(x), u(y))^p}{|x-y|^{2m}} dx dy = \iint_{\mathbb{R}^m \times \mathbb{R}^m} \frac{d(u(\Psi^{-1}(x)), u(\Psi^{-1}(y)))^p}{|\Psi^{-1}(x) - \Psi^{-1}(y)|^{2m}} dx dy$$

and

$$(67) \quad \iint_{\substack{x, y \in \mathbb{S}^m \\ d(u(x), u(y)) \geq \delta}} \frac{d(u(x), u(y))^p}{|x-y|^{2m}} dx dy = \iint_{\substack{x, y \in \mathbb{R}^m \\ d(u(\Psi^{-1}(x)), u(\Psi^{-1}(y))) \geq \delta}} \frac{d(u(\Psi^{-1}(x)), u(\Psi^{-1}(y)))^p}{|\Psi^{-1}(x) - \Psi^{-1}(y)|^{2m}} dx dy.$$

We proceed then as in the proof of Theorem 1.2, using $u \circ \Psi^{-1}$ instead of u . In order to replace the estimate (61), we proceed as follows.

Given $Q \in \mathcal{Q}_{\lambda,\tau,k,h}^+$ and setting

$$m_Q = \inf_{x \in Q} \frac{4}{|x+2e|^2} \quad \text{and} \quad M_Q = \sup_{x \in Q} \frac{4}{|x+2e|^2},$$

we have

$$(68) \quad \int_{\substack{x \in Q \\ |DV(x)| \geq 4t/|x+2e|^2}} \frac{4^{m+1}}{|x+2e|^{2m+2}} dx \leq M_Q^{m+1} \mathcal{L}^{m+1}(\{x \in Q \mid |DV(x)| \geq tm_Q\}) \\ \leq \frac{1}{t^{m+1}} \left(\frac{M_Q}{m_Q}\right)^{m+1} (\tau\lambda^{-k})^{m+1} \sup_{x \in Q} |DV(x)|^{m+1}.$$

We observe now that if $x, y \in Q$, then

$$(69) \quad \frac{|x+2e|}{|y+2e|} \leq 1 + \frac{|x-y|}{|x+2e|} \leq 1 + \frac{\tau\lambda^{-k}\sqrt{m+1}}{2 + \frac{\tau\lambda^{-k}}{\lambda-1}} \leq 1 + (\lambda-1)\sqrt{m+1},$$

and we have thus by (68) and (69),

$$\int_{\substack{x \in Q \\ |DV(x)| \geq 4t/|x+2e|^2}} \frac{4^{m+1}}{|x+2e|^{2m+2}} dx \leq \frac{(\lambda-1)^{m+1}}{t^{m+1}} \left(1 + (\lambda-1)\sqrt{m+1}\right)^{m+1} \sup_{x \in Q} x_{m+1}^{m+1} |DV(x)|^{m+1}.$$

The rest of the proof is similar to the proof of Theorem 1.2. \square

REFERENCES

- [1] F. Bethuel, *A new obstruction to the extension problem for Sobolev maps between manifolds*, J. Fixed Point Theory Appl. **15** (2014), no. 1, 155–183, doi:10.1007/s11784-014-0185-0. $\uparrow 2$
- [2] F. Bethuel and F. Demengel, *Extensions for Sobolev mappings between manifolds*, Calc. Var. Partial Differential Equations **3** (1995), no. 4, 475–491, doi:10.1007/BF01187897. $\uparrow 2$
- [3] J. Bourgain, H. Brezis, and P. Mironescu, *Lifting, degree, and distributional Jacobian revisited*, Comm. Pure Appl. Math. **58** (2005), no. 4, 529–551, doi:10.1002/cpa.20063. $\uparrow 4$
- [4] J. Bourgain, H. Brezis, and Nguyen H.-M., *A new estimate for the topological degree*, C. R. Math. Acad. Sci. Paris **340** (2005), no. 11, 787–791, doi:10.1016/j.crma.2005.04.007. $\uparrow 4$
- [5] H. Brezis and Nguyen H.-M., *On a new class of functions related to VMO*, C. R. Math. Acad. Sci. Paris **349** (2011), no. 3-4, 157–160, doi:10.1016/j.crma.2010.11.026. $\uparrow 4$
- [6] H. Brezis and L. Nirenberg, *Degree theory and BMO. I. Compact manifolds without boundaries*, Selecta Math. (N.S.) **1** (1995), no. 2, 197–263, doi:10.1007/BF01671566. $\uparrow 2, 5$
- [7] E. DiBenedetto, *Real analysis*, 2nd ed., Birkhäuser Advanced Texts: Basler Lehrbücher, Birkhäuser/Springer, New York, 2016, doi:10.1007/978-1-4939-4005-9. $\uparrow 13$
- [8] E. Gagliardo, *Caratterizzazioni delle tracce sulla frontiera relative ad alcune classi di funzioni in n variabili*, Rend. Sem. Mat. Univ. Padova **27** (1957), 284–305. $\uparrow 1, 13$
- [9] R. Hardt and F.-H. Lin, *Mappings minimizing the L^p norm of the gradient*, Comm. Pure Appl. Math. **40** (1987), no. 5, 555–588, doi:10.1002/cpa.3160400503. $\uparrow 2$
- [10] A. Kufner, O. John, and S. Fučík, *Function spaces*, Noordhoff, Leyden, 1977. $\uparrow 13$
- [11] G. Leoni, *A first course in fractional Sobolev spaces*, Graduate Studies in Mathematics, vol. 229, American Mathematical Society, Providence, R.I., 2023, doi:10.1090/gsm/229. $\uparrow 13$
- [12] V. Maz'ya, *Sobolev spaces with applications to elliptic partial differential equations*, Second, revised and augmented edition, Grundlehren der Mathematischen Wissenschaften, vol. 342, Springer, Heidelberg, 2011, doi:10.1007/978-3-642-15564-2. $\uparrow 13$
- [13] P. Mironescu and J. Van Schaftingen, *Trace theory for Sobolev mappings into a manifold*, Ann. Fac. Sci. Toulouse Math. (6) **30** (2021), no. 2, 281–299, doi:10.5802/afst.1675. $\uparrow 2, 3$
- [14] J. Nash, *The imbedding problem for Riemannian manifolds*, Ann. of Math. (2) **63** (1956), 20–63, doi:10.2307/1969989. $\uparrow 2$
- [15] Nguyen H.-M., *Some new characterizations of Sobolev spaces*, J. Funct. Anal. **237** (2006), no. 2, 689–720, doi:10.1016/j.jfa.2006.04.001. $\uparrow 4$
- [16] ———, *Γ -convergence and Sobolev norms*, C. R. Math. Acad. Sci. Paris **345** (2007), no. 12, 679–684, doi:10.1016/j.crma.2007.11.005. $\uparrow 4$
- [17] ———, *Optimal constant in a new estimate for the degree*, J. Anal. Math. **101** (2007), 367–395, doi:10.1007/s11854-007-0014-0. $\uparrow 4$

- [18] ———, *Inequalities related to liftings and applications*, C. R. Math. Acad. Sci. Paris **346** (2008), no. 17-18, 957–962, doi:10.1016/j.crma.2008.07.026. ↑4
- [19] ———, *Further characterizations of Sobolev spaces*, J. Eur. Math. Soc. (JEMS) **10** (2008), no. 1, 191–229, doi:10.4171/JEMS/108. ↑4
- [20] ———, *Estimates for the topological degree and related topics*, J. Fixed Point Theory Appl. **15** (2014), no. 1, 185–215, doi:10.1007/s11784-014-0182-3. ↑4
- [21] M. Petrache and T. Rivière, *Global gauges and global extensions in optimal spaces*, Anal. PDE **7** (2014), no. 8, 1851–1899, doi:10.2140/apde.2014.7.1851. ↑3
- [22] M. Petrache and J. Van Schaftingen, *Controlled singular extension of critical trace Sobolev maps from spheres to compact manifolds*, Int. Math. Res. Not. IMRN **12** (2017), 3647–3683, doi:10.1093/imrn/rnw109. ↑3, 4, 6
- [23] J. Van Schaftingen, *Estimates by gap potentials of free homotopy decompositions of critical Sobolev maps*, Adv. Nonlinear Anal. **9** (2020), no. 1, 1214–1250, doi:10.1515/anona-2020-0047. ↑2, 4, 7
- [24] ———, *Lifting of fractional Sobolev mappings to noncompact covering spaces*, Ann. Inst. H. Poincaré Anal. Non Linéaire, doi:10.4171/AIHPC/98. ↑4, 7

UNIVERSITÉ CATHOLIQUE DE LOUVAIN, INSTITUT DE RECHERCHE EN MATHÉMATIQUE ET PHYSIQUE,
CHEMIN DU CYCLOTRON 2 BTE L7.01.01, 1348 LOUVAIN-LA-NEUVE, BELGIUM

Email address: Bohdan.Bulanyi@UCLouvain.be

Email address: Jean.VanSchaftingen@UCLouvain.be