

ON τ -TILTING FINITE BOREL-SCHUR ALGEBRAS

QI WANG

On the occasion of Professor Susumu Ariki's retirement

ABSTRACT. We completely determine the τ -tilting finiteness of Borel-Schur algebras. To achieve this, we use two recently introduced techniques in silting theory: sign decomposition as introduced by Aoki, Higashitani, Iyama, Kase and Mizuno [AHIKM], and symmetry of silting quivers as investigated by Aihara and the author [AiW]. Besides, we explore some new properties for both Borel-Schur algebra and sign decomposition.

1. INTRODUCTION

Representation type plays a fundamental role in the representation theory of finite-dimensional algebras. One can classify finite-dimensional algebras into three classes based on the indecomposable modules the algebra admits. Let A be a finite-dimensional algebra over an algebraically closed field K . Then A is said to be *representation-finite* if it admits only finitely many indecomposable modules up to isomorphism; otherwise, A is said to be *representation-infinite*. We say that a representation-infinite algebra A is *tame* if all but finitely many d -dimensional indecomposable A -modules can be organized in a one-parameter family, for each dimension d . A representation-infinite algebra A is called *wild* if there exists a faithful exact K -linear functor from the module category of the free associative algebra $K\langle x, y \rangle$ to the module category of A . The famous Finite-Tame-Wild Trichotomy, established by Drozd [D], asserts that the representation type of any finite-dimensional algebra over K is exactly one of representation-finite, tame and wild.

Similar to the concept of representation-finiteness, Demonet, Iyama and Jasso [DIJ] introduced a modern notion called *τ -tilting finiteness*, which is inspired by the τ -tilting theory initially proposed by Adachi, Iyama and Reiten [AIR]. Here, τ denotes the Auslander-Reiten translation. Over the past decade, τ -tilting theory has arisen not only as a generalization of classical tilting theory but also as a field closely connected to various other objects in representation theory. These include torsion classes, wide subcategories, semibricks, two-term silting complexes, stability conditions, maximal green sequences, wall-and-chamber structures, cluster-tilted objects, fans and polytopes, and so on. We refer readers to some in-depth papers such as [AHIKM], [As], [BST], [DIRRT], [IR], [KT], [T], for more details. The study of τ -tilting finiteness and new results in τ -tilting theory may provide a deeper understanding of some classical problems, and then enhance the

Key words and phrases. Sign decomposition, Borel-Schur algebras, τ -tilting finiteness, silting objects.
 2020 *Mathematics Subject Classification.* 16G20, 16G60.

connection to the aforementioned objects. The methodologies developed during the process may lead to other applications that fall within their own research directions. These are the underlying motivations for studying this modern notion.

A finite-dimensional algebra over K is said to be τ -tilting finite if it has only a finite number of τ -tilting modules. It is easy to observe that a representation-finite algebra must be τ -tilting finite. Certain tame or wild algebras could also exhibit τ -tilting finiteness, e.g., Brauer graph algebras [AAC] and preprojective algebras of Dynkin type [Mi]. In fact, τ -tilting finiteness captures a *brick-finiteness*, as discussed in [DIJ], in both tame and wild algebras. Besides, τ -tilting finiteness has been determined for various classes of algebras, including radical square zero algebras [Ad], cycle finite algebras [MS], locally hereditary algebras [AHMW], (special) biserial algebras [STV, Mo], gentle algebras [P], some subclasses of two-point algebras [Wa1, Wa4], Hecke algebras of type A [ALS], Schur algebras [Wa3, AoW], simply connected algebras [Wa2] and their tensor product algebras [MW], cluster-tilted algebras [Z], and more.

In this paper, we pay particular attention to the bijection (as demonstrated in Theorem 2.6) between τ -tilting theory and silting theory. We are aiming to study additional properties of sign decomposition within the framework of silting theory. Sign decomposition was initially introduced by Aoki in [Ao] as a means to classify torsion classes in radical square zero algebras. Subsequently, this concept was extended and generalized by Aoki, Higashitani, Iyama, Kase and Mizuno in their work [AHIKM], where it was applied to the study of fans and polytopes in τ -tilting theory. We are also aiming to investigate the behavior of sign decomposition on some subclasses of *triangular algebras*, i.e., algebra without oriented cycles in its quiver. Indeed, the application of sign decomposition to certain Borel-Schur algebra yields interesting insights, see for example, Proposition 3.8.

Borel-Schur algebra $S^+(n, r)$ is a certain subalgebra of the Schur algebra $S(n, r)$ over an algebraically closed field K of any characteristic $p \geq 0$, see Subsection 2.2 for the definitions. Since the module category of $S(n, r)$ is equivalent to the category of r -th homogeneous polynomial representations over the general linear group $\mathrm{GL}_n(K)$ of degree n , the module category of $S^+(n, r)$ encodes the representation theory of r -th homogeneous polynomial representations over the group of upper triangular matrices in $\mathrm{GL}_n(K)$. Borel-Schur algebras also play an important role in the study of projective resolutions for Weyl modules associated with $S(n, r)$ or $\mathrm{GL}_n(K)$, see [SY, Wo1, Wo2].

Recently, Aoki and the author [Wa3, AoW] have completely determined the τ -tilting finiteness of $S(n, r)$. Although taking quotient and idempotent truncation preserves τ -tilting finiteness of algebras (see [DIRRT, DIJ]), taking subalgebra usually does not keep τ -tilting finiteness. Hence, one cannot expect to know the τ -tilting finiteness of $S^+(n, r)$ according to the results in [Wa3, AoW], by the fact that $S^+(n, r)$ is the upper part of the triangular decomposition of $S(n, r)$. In fact, $S(3, 2)$ is known to be τ -tilting finite (over any $p \geq 0$) but $S^+(3, 2)$ is τ -tilting infinite showing in this paper. Besides, the aforementioned fact enables $S^+(n, r)$ more nice properties compared with $S(n, r)$, e.g., $S^+(n, r)$ is a triangular algebra (Proposition 2.11), such that the $S^+(n, r)$'s we need to deal

with may have a large radical layer length. Since the support-rank reduction developed in [Wa3, AoW] is only applicable to algebras with a small radical layer length, we cannot use the same strategy as we did for $S(n, r)$. We have to introduce a new technique for $S^+(n, r)$, and this gives the reason why we consider sign decomposition in silting theory.

Based on the motivations for both sign decomposition and Borel-Schur algebras as mentioned above, we present in this paper a complete classification of Borel-Schur algebras in terms of τ -tilting finiteness. Erdmann, Santana and Yudin [ESY1, ESY2] have provided a complete classification for the representation type of Borel-Schur algebras. We first review their results as follows.

Theorem 1 ([ESY1, ESY2]). *The Borel-Schur algebra $S^+(n, r)$ is*

- *representation-finite if one of the following holds:*
 - $n = 2$ and $p = 0$, or $p = 2, r \leq 3$ or $p = 3, r \leq 4$ or $p \geq 5, r \leq p$;
 - $n \geq 3$ and $r = 1$.
- *tame if $n = 2, p = 3, r = 5$ or $n = 3, r = 2$.*

Otherwise, $S^+(n, r)$ is wild.

Representation-finite cases are naturally τ -tilting finite, but the converse is not valid in the class of Borel-Schur algebras. Our main result below implies that a tame case ($S^+(2, 5)$ over $p = 3$) and two wild cases ($S^+(2, 4)$ over $p = 2$, $S^+(2, 6)$ over $p = 5$) occur as τ -tilting finite algebras. These are the only cases that are representation-infinite but τ -tilting finite.

Theorem 2 (Theorem 4.6 and Theorem 5.5). *Let $S^+(n, r)$ be the Borel-Schur algebra. Then, it is τ -tilting finite if and only if one of the following holds:*

- $n = 2$ and $p = 0$, or $p = 2, r \leq 4$ or $p = 3, r \leq 5$ or $p = 5, r \leq 6$ or $p \geq 7, r \leq p$;
- $n \geq 3$ and $r = 1$.


In order to prove the above result, we first reduce the problem on $S^+(n, r)$ to cases involving small n and r via idempotent truncation, and then check the τ -tilting finiteness of these few cases via quotient methods and the sign decomposition in silting theory. The τ -tilting finiteness of all but three cases: $S^+(2, 4)$ over $p = 2$, $S^+(2, 5)$ over $p = 3$, $S^+(2, 6)$ over $p = 5$, can be obtained theoretically using reduction techniques in Section 2, while the exceptional three cases are handled by sign decomposition mentioned in Section 3. We then divide the proof into Section 4 and Section 5. In particular, the symmetry of silting quivers, tame concealed algebras and simply connected algebras, all of which play a crucial role in our proof.

An immediate observation is that a τ -tilting finite Borel-Schur algebra is representation-finite if $p = 0$ or $n = 2, p \geq 7$ or $n \geq 3$. This observation potentially introduces new examples of algebras for which τ -tilting finiteness is equivalent to representation-finiteness.

2. PRELIMINARIES

Any finite-dimensional algebra A over an algebraically closed field K can be presented as a bound quiver algebra KQ/I with a finite (connected) quiver Q and an admissible ideal I . Here, *admissible* stands for the condition $R_Q^m \subseteq I \subseteq R_Q^2$ with some integer $m \geq 2$, where R_Q is the arrow ideal of KQ .

In this paper, however, we do not always deal with admissible ideals. We will meet a quotient KQ/J of KQ , in which J satisfies $R_Q^m \subseteq J$ for some $m \geq 2$ and contains a relation $\alpha - \sum_{i=1}^k \lambda_i \omega_i$ for some arrow $\alpha \in Q$, paths $\omega_i \in KQ$, coefficients $\lambda_i \in K$ such that α does not occur in any of ω_i 's. It is obvious that J is not admissible. If this is the case, the quotient KQ/J is isomorphic to a bound quiver algebra KQ'/J' , where Q' is obtained by removing α from Q and J' is obtained by removing the relation $\alpha - \sum_{i=1}^k \lambda_i \omega_i$ from J as well as replacing all references to α by $\sum_{i=1}^k \lambda_i \omega_i$. For example,



Here, a dotted line stands for a commutativity relation in an obvious way. Throughout, we will use such a graph to indicate the bound quiver algebra presented by the quiver with commutativity relations.

We consider right A -modules in this paper. Let $(-)^* := \text{Hom}_A(-, A)$. We denote by $\text{rad } A$ the Jacobson radical of A and by A^{op} the opposite algebra of A . The category of finitely generated right A -modules is denoted by $\text{mod } A$ and the full subcategory of $\text{mod } A$ consisting of projective A -modules is denoted by $\text{proj } A$.

2.1. τ -tilting theory and sifting theory. We first review the fundamental definitions in τ -tilting theory. Then, we explain some details of sifting theory as well as its connection with τ -tilting theory.

Definition 2.1 ([AIR, Definition 0.1]). Let $M \in \text{mod } A$ and $|M|$ be the number of isomorphism classes of indecomposable direct summands of M .

- (1) M is called τ -rigid if $\text{Hom}_A(M, \tau M) = 0$.
- (2) M is called τ -tilting if it is τ -rigid and $|M| = |A|$.
- (3) M is called support τ -tilting if M is a τ -tilting (A/AeA) -module for an idempotent e of A . In this case, (M, P) with $P := eA$ is called a support τ -tilting pair.

We denote by $\tau\text{-tilt } A$ (resp., $s\tau\text{-tilt } A$) the set of isomorphism classes of basic (resp., support τ -tilting) τ -tilting A -modules. Obviously, $\tau\text{-tilt } A \subset s\tau\text{-tilt } A$. It is worth noting that support τ -tilting modules are also τ -rigid.

Definition 2.2 ([DIJ, Definition 1.1]). An algebra A is called τ -tilting finite if $\tau\text{-tilt } A$ is a finite set. Otherwise, A is said to be τ -tilting infinite.

It is shown by [AIR, Theorem 0.2] that any τ -rigid A -module is a direct summand of some τ -tilting A -module. Then, A is τ -tilting finite if and only if $s\tau\text{-tilt } A$ is finite, if and

only if A has only finitely many pairwise non-isomorphic indecomposable τ -rigid modules. The last condition yields the finiteness of bricks in $\mathbf{mod} A$, see [DIJ, Theorem 4.2].

Let $\mathcal{K}_A := \mathbf{K}^b(\mathbf{proj} A)$ be the homotopy category of bounded complexes of finitely generated projective A -modules. For any $T \in \mathcal{K}_A$, we denote by $\mathbf{thick} T$ the smallest thick subcategory of \mathcal{K}_A containing T . Let $\mathbf{add}(T)$ be the full subcategory of \mathcal{K}_A whose objects are direct summands of finite direct sums of copies of T .

Definition 2.3 ([AI, Definition 2.1]). Let $T \in \mathcal{K}_A$. Then,

- (1) T is presilting (resp., pretilting) if $\mathrm{Hom}_{\mathcal{K}_A}(T, T[i]) = 0$ for any $i > 0$ (resp., $i \neq 0$).
- (2) T is silting (resp., tilting) if T is presilting (resp., pretilting) and $\mathbf{thick} T = \mathcal{K}_A$.

We recall the construction of silting mutations, following from [AI, Definition 2.30]. Let $T = X \oplus Y \in \mathcal{K}_A$ be a basic silting complex with a direct summand X (not necessarily indecomposable). Take a minimal left $\mathbf{add}(Y)$ -approximation π of X and a triangle

$$X \xrightarrow{\pi} Z \longrightarrow X' \longrightarrow X[1],$$

where X' is the mapping cone of π . It is shown in [AI, Theorem 2.31] that $\mu_X^-(T) := X' \oplus Y$ is again a basic silting complex in \mathcal{K}_A . We call $\mu_X^-(T)$ the left (silting) mutation of T with respect to X . Dually, we can define the right (silting) mutation $\mu_X^+(T)$ of T with respect to X . If moreover, X is indecomposable, then $\mu_X^\pm(T)$ is said to be *irreducible*.

Let $\mathbf{silt} A$ be the set of isomorphism classes of basic silting complexes in \mathcal{K}_A . For any $T, S \in \mathbf{silt} A$, we define $T \geq S$ if $\mathrm{Hom}_{\mathcal{K}_A}(T, S[i]) = 0$ for all $i > 0$. This actually gives a partial order on the set $\mathbf{silt} A$. It is known from [AI, Theorem 2.35] that S is a left mutation of T if and only if T is a right mutation of S , if and only if, $T > S$.

We now restrict our attention to two-term silting complexes. A complex in \mathcal{K}_A is said to be *two-term* if it is homotopy equivalent to a complex T , which is concentrated in degrees 0 and -1 , that is,

$$T = (T^{-1} \xrightarrow{f} T^0) := \cdots \longrightarrow 0 \longrightarrow T^{-1} \xrightarrow{f} T^0 \longrightarrow 0 \longrightarrow \cdots.$$

Let $2\text{-silt} A$ be the set of two-term complexes in $\mathbf{silt} A$. Obviously, $2\text{-silt} A$ is again a poset under the partial order \geq on $\mathbf{silt} A$. We then denote by $\mathcal{H}(2\text{-silt} A)$ the Hasse quiver of $2\text{-silt} A$, which is compatible with the left/right mutation of two-term silting complexes.

Proposition 2.4 ([AI, Lemma 2.25, Theorem 2.27]). *Let $T = (T^{-1} \rightarrow T^0) \in 2\text{-silt} A$. Then, we have $\mathbf{add} A = \mathbf{add}(T^0 \oplus T^{-1})$ and $\mathbf{add}(T^0) \cap \mathbf{add}(T^{-1}) = \emptyset$.*

Let $\{P_1, P_2, \dots, P_n\}$ be a complete set of pairwise non-isomorphic indecomposable projective A -modules. We denote by $\underline{P}_1, \underline{P}_2, \dots, \underline{P}_n$ the isomorphism classes of indecomposable complexes concentrated in degree 0. Then, the classes $\underline{P}_1, \underline{P}_2, \dots, \underline{P}_n$ in \mathcal{K}_A induce a standard basis of the Grothendieck group $K_0(\mathcal{K}_A)$. If a two-term complex $T \in \mathcal{K}_A$ is written as

$$\left(\bigoplus_{i=1}^n P_i^{\oplus b_i} \longrightarrow \bigoplus_{i=1}^n P_i^{\oplus a_i} \right),$$

then the class \underline{T} can be identified by an integer vector

$$g(T) = (a_1 - b_1, a_2 - b_2, \dots, a_n - b_n) \in \mathbb{Z}^n,$$

which is called the g -vector of T . The above Proposition 2.4 says that a two-term silting complex $T \in 2\text{-silt } A$ must be of the form

$$(2.1) \quad \left(\bigoplus_{i \in I} P_i^{\oplus a_i} \longrightarrow \bigoplus_{j \in J} P_j^{\oplus a_j} \right)$$

with $I \cap J = \emptyset$ and $I \cup J = \{1, 2, \dots, n\}$. Then, each entry of $g(T)$ for $T \in 2\text{-silt } A$ is either positive or negative, and can not be zero. Without confusion, we sometimes display g -vectors of two-term silting complexes as the so-called g -matrices, in which the rows are given by g -vectors of indecomposable two-term presilting complexes.

The following is a critical statement.

Proposition 2.5 ([AIR, Theorem 5.5]). *Let T be a two-term silting complex in $2\text{-silt } A$. Then, the map $T \mapsto g(T)$ is an injection.*

We can explain the connection between τ -tilting theory and silting theory as follows. This will give the reason why we focus on two-term silting complexes.

Theorem 2.6 ([AIR, Theorem 3.2]). *There exists a poset isomorphism between $\text{s}\tau\text{-tilt } A$ and $2\text{-silt } A$. More precisely, the bijection is given by*

$$(M, P) \longmapsto (P_1 \oplus P \xrightarrow{\begin{pmatrix} f \\ 0 \end{pmatrix}} P_0),$$

where (M, P) is the support τ -tilting pair corresponding to M and $P_1 \xrightarrow{f} P_0 \longrightarrow M \longrightarrow 0$ is the minimal projective presentation of M .

We give an example to illustrate the above settings.

Example 2.7. Let A be the algebra presented by

$$\begin{array}{ccc} 1 & \xrightarrow{\alpha} & 2 \\ \beta \downarrow & \searrow & \downarrow \mu \\ 3 & \xrightarrow{\nu} & 4 \end{array}.$$

The indecomposable projective A -module P_i at vertex i can be displayed as

$$P_1 \simeq \begin{matrix} 1 \\ 2 \\ 4 \end{matrix} 3, \quad P_2 \simeq \begin{matrix} 2 \\ 4 \end{matrix}, \quad P_3 \simeq \begin{matrix} 3 \\ 4 \end{matrix}, \quad P_4 \simeq 4.$$

A complete list of support τ -tilting A -modules is given in [Wa2, Example A.3], and there are 46 support τ -tilting A -modules. We choose, e.g., $\begin{matrix} 1 \\ 2 \\ 4 \end{matrix} 3 \oplus \begin{matrix} 1 \\ 3 \end{matrix} \oplus \begin{matrix} 1 \\ 2 \end{matrix} \oplus \begin{matrix} 1 \\ 2 \\ 3 \end{matrix} \in \text{s}\tau\text{-tilt } A$, the aforementioned correspondences can be illustrated as

$$\left(\begin{matrix} 1 \\ 2 \\ 4 \end{matrix} 3 \oplus \begin{matrix} 1 \\ 3 \end{matrix} \oplus \begin{matrix} 1 \\ 2 \end{matrix} \oplus \begin{matrix} 1 \\ 2 \\ 3 \end{matrix}, 0 \right) \xrightarrow{\text{Th. 2.6}} \begin{bmatrix} 0 \longrightarrow P_1 \\ \oplus \\ P_2 \xrightarrow{\alpha} P_1 \\ \oplus \\ P_3 \xrightarrow{\beta} P_1 \\ \oplus \\ P_4 \xrightarrow{\alpha\mu} P_1 \end{bmatrix} \xrightarrow{\text{Prop. 2.5}} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & -1 & 0 & 0 \\ 1 & 0 & -1 & 0 \\ 1 & 0 & 0 & -1 \end{pmatrix} \xrightarrow{1:1} (4, -1, -1, -1).$$

We know from [AIR] that checking the τ -tilting finiteness of A is identical to finding either a finite connected component of $\mathcal{H}(2\text{-silt } A) \simeq \mathcal{H}(s\tau\text{-tilt } A)$ or an infinite left mutation chain in $\mathcal{H}(2\text{-silt } A) \simeq \mathcal{H}(s\tau\text{-tilt } A)$. This is the reason why we are able to solve certain small cases via direct calculation.

Proposition 2.8 ([AIR, Corollary 2.38]). *If the Hasse quiver $\mathcal{H}(2\text{-silt } A)$ contains a finite connected component \mathcal{C} , then $\mathcal{H}(2\text{-silt } A) = \mathcal{C}$.*

In order to identify a finite connected component of $\mathcal{H}(2\text{-silt } A)$, we usually calculate the left mutations starting from A because A is the maximal element in $2\text{-silt } A$. Although such a left mutation is always silting, it may not always be two-term. Hence, it becomes crucial to determine when such a left mutation is out of $2\text{-silt } A$. This leads us to the following essential statement.

Proposition 2.9 ([AIR, Corollary 3.8]). *If $S \in \mathcal{K}_A$ is a two-term presilting complex with $|S| = |A| - 1$, then S is a direct summand of exactly two basic two-term silting complexes.*

2.2. Borel-Schur algebras. Let n, r be two positive integers and K an algebraically closed field of characteristic $p \geq 0$. We take an n -dimensional vector space V over K with a basis $\{v_1, v_2, \dots, v_n\}$ and denote by $V^{\otimes r}$ the r -fold tensor product $V \otimes_K V \otimes_K \cdots \otimes_K V$. Then, the tensor product $V^{\otimes r}$ admits a K -basis given by

$$\{v_{i_1} \otimes v_{i_2} \otimes \cdots \otimes v_{i_r} \mid 1 \leq i_j \leq n \text{ for all } 1 \leq j \leq r\}.$$

Since the general linear group GL_n over K has a natural action on V , it acts on $V^{\otimes r}$ by

$$(v_{i_1} \otimes v_{i_2} \otimes \cdots \otimes v_{i_r}) \cdot g = gv_{i_1} \otimes gv_{i_2} \otimes \cdots \otimes gv_{i_r}$$

for any $g \in \text{GL}_n$. This is naturally extended to the group algebra $K\text{GL}_n$ of GL_n . We then obtain a homomorphism of algebras:

$$\rho : K\text{GL}_n \longrightarrow \text{End}_K(V^{\otimes r}).$$

The image of ρ , i.e., $\rho(K\text{GL}_n)$, is called the *Schur algebra*.

Definition 2.10. Let B^+ be the Borel subgroup of GL_n consisting of all upper triangular matrices. We call the subalgebra $\rho(B^+)$ of $\rho(K\text{GL}_n)$ the Borel-Schur algebra and denote it by $S_K^+(n, r)$, or simply by $S^+(n, r)$.

Borel-Schur algebra $S^+(n, r)$ admits many nice properties, for example, it is a basic algebra ([S]) and it has finite global dimension ([G]). We explore two more new properties of $S^+(n, r)$ in the following.

Let $\Lambda(n, r)$ be the set consisting of all compositions¹ of r into n parts, that is,

$$\Lambda(n, r) := \{(\lambda_1, \lambda_2, \dots, \lambda_n) \mid \lambda_1 + \lambda_2 + \cdots + \lambda_n = r, \lambda_i \geq 0, 1 \leq i \leq n\}.$$

¹This usually appears in the literature as *weak compositions*, and the *composition* of r into at most n parts in the literature is obtained by restricting $\lambda_i > 0$ in the definition of $\Lambda(n, r)$.

We may write a composition $(\lambda_1, \lambda_2, \dots, \lambda_n)$ as $\lambda_1\lambda_2\cdots\lambda_n$ for short. We define $\lambda < \mu \in \Lambda(n, r)$ in *lexicographic order* if $\lambda_j = \mu_j$ for $j < i$ and $\lambda_i < \mu_i$ for some $1 \leq i \leq n$. This gives a total order on $\Lambda(n, r)$, for instance,

$$003 < 012 < 021 < 030 < 102 < 111 < 120 < 201 < 210 < 300 \in \Lambda(3, 3).$$

We define an involution ι on $\Lambda(n, r)$ by $\iota(\lambda_1\lambda_2\cdots\lambda_n) := \lambda_n\cdots\lambda_2\lambda_1$.

The primitive orthogonal idempotents of $S^+(n, r)$ are labeled by elements of $\Lambda(n, r)$. The quiver Q of $S^+(n, r)$ has been determined in [S, Theorem 5.4] (see also [L, Chapter 4]). Since $S^+(n, r)$ is basic, we may identify the vertex set of Q by $\Lambda(n, r)$. If $p = 0$, there is an arrow from λ to μ in Q if and only if $\mu - \lambda = \gamma_i$, for some $1 \leq i \leq n$, where $\gamma_i = (0, \dots, 0, 1, -1, 0, \dots, 0)$ with 1 being the i -th entry. If $p > 0$, we have $\lambda \rightarrow \mu$ in Q if and only if $\mu - \lambda = p^d\gamma_i$, for some $d \geq 0$, $1 \leq i \leq n$.

We show that there is no oriented cycle in the quiver of $S^+(n, r)$.

Proposition 2.11. *The Borel-Schur algebra $S^+(n, r)$ is a triangular algebra.*

Proof. If there is an arrow $\lambda \rightarrow \mu$ in the quiver of $S^+(n, r)$, we obtain $\lambda_j = \mu_j$ for $j < i$, and $\lambda_i = \mu_i - 1$ if $p = 0$, $\lambda_i = \mu_i - p^d$ if $p > 0$, for some $d \geq 0$, $1 \leq i \leq n$. This gives us $\lambda < \mu$ in lexicographic order.

Suppose that there is an oriented cycle in $S^+(n, r)$, say, $\lambda^0 \rightarrow \lambda^1 \rightarrow \cdots \rightarrow \lambda^k \rightarrow \lambda^0$, for some $k \geq 0$. We then obtain both $\lambda^0 < \lambda^k$ and $\lambda^k < \lambda^0$, a contradiction. \square

Next, we show that the quiver Q of $S^+(n, r)$ admits an isomorphism $\zeta : Q \rightarrow Q$ sending $\lambda, \mu, \lambda \rightarrow \mu$ to $\zeta(\lambda), \zeta(\mu), \zeta(\mu) \rightarrow \zeta(\lambda)$ respectively, for any $\lambda, \mu \in \Lambda(n, r)$. It is possible that $\zeta(\lambda) = \mu$; in this case, $\lambda \rightarrow \mu$ is stable under the isomorphism ζ . Such an isomorphism of Q could give rise to an algebra anti-isomorphism of KQ .

Proposition 2.12. *Let Q be the quiver of $S^+(n, r)$. There is an isomorphism $\zeta : Q \rightarrow Q$ mapping λ to $\iota(\lambda)$, for $\lambda \in \Lambda(n, r)$, and ζ induces an algebra anti-isomorphism of KQ .*

Proof. The opposite quiver Q^{op} of Q consists of the same vertex set $\Lambda(n, r)$ and reversed arrows x^* for arrows x of Q , i.e., x^* is obtained by swapping the source and target of x . To keep symbol consistency, we write $\lambda \in \Lambda(n, r)$ as λ^* if it indicates a vertex in Q^{op} . We then conclude that $\lambda \rightarrow \mu$ in Q if and only if $\mu^* \rightarrow \lambda^*$ in Q^{op} , for any $\lambda, \mu \in \Lambda(n, r)$.

On the other hand, the condition $\mu - \lambda = \gamma_i$ in the above definition is equivalent to $\iota(\mu) - \iota(\lambda) = \bar{\gamma}_i$, where $\bar{\gamma}_i$ is obtained by rearranging the entries of γ_i in reverse order, and $-\bar{\gamma}_i = \gamma_j$ for some j . Since $\iota(\lambda), \iota(\mu) \in \Lambda(n, r)$, each arrow $\lambda \rightarrow \mu$ in Q gives an arrow $\iota(\mu) \rightarrow \iota(\lambda)$ in Q . We obtain an isomorphism $\sigma : Q^{\text{op}} \rightarrow Q$ of quivers by sending $\lambda^*, \mu^*, \mu^* \rightarrow \lambda^*$ to $\iota(\lambda), \iota(\mu), \iota(\mu) \rightarrow \iota(\lambda)$, respectively. This gives rise to an algebra isomorphism between KQ^{op} and KQ , which is also denoted as σ .

Hence, the composition $\zeta := \sigma \circ (-)^*$ gives the desired quiver isomorphism of Q and algebra anti-isomorphism of KQ . \square

There exists an explicit formula ([G]) for the multiplication in $S^+(n, r)$, which can be used to find the quiver presentation KQ/I of $S^+(n, r)$. For our purpose, however, we do

not need to explain more details of $S^+(n, r)$, but only constructions of $S^+(2, r)$. Since a composition (λ_1, λ_2) of r into 2 parts is uniquely determined by $\lambda_1 - \lambda_2$ and $\Lambda(2, r)$ contains $r + 1$ elements, we replace the indexed set $\Lambda(2, r)$ of Q by $\{0, 1, 2, \dots, r\}$.

Proposition 2.13 ([ESY1, Proposition 2.6 (a)]). *Let $p = 0$. Then $S^+(2, r)$ is isomorphic to the path algebra of the linear quiver $0 \longleftarrow 1 \longleftarrow 2 \longleftarrow \dots \longleftarrow r$.*

There are several different approaches to describe the quiver and relations of $S^+(2, r)$ over $p > 0$, see [S], [Y], [ESY1], etc. We follow the constructions in [L, Chapter 2] as follows. Given a positive integer s , it admits a unique p -adic decomposition $s = \sum_{i \geq 0} s_i p^i$ with $0 \leq s_i \leq p - 1$. Set $[s]_p := \sum_{i \geq 0} s_i$. We define a quiver Δ_r whose vertex set is labelled by $\{0, 1, 2, \dots, r\}$, and arrows are given by $\alpha_{s,t} : s \rightarrow t$ if $0 \leq t < s \leq r$ and $[s - t]_p = 1$. Then, we define a two-sided ideal I of $K\Delta_r$ generated by

$$\{\alpha_{s+p^{k+1}, s+(p-1)p^k} \alpha_{s+(p-1)p^k, s+(p-2)p^k} \cdots \alpha_{s+p^k, s} \mid s, k \geq 0, 0 \leq s + p^k \leq r\}$$

and

$$\{\alpha_{s+p^k+p^\ell, s+p^k} \alpha_{s+p^k, s} - \alpha_{s+p^k+p^\ell, s+p^\ell} \alpha_{s+p^\ell, s} \mid s, k \neq \ell \geq 0, 0 \leq s + p^k + p^\ell \leq r\}.$$

The following proposition is fully used in Section 4.

Proposition 2.14 ([L, Theorem 2.3.1]). *Let $p > 0$. Then, $S^+(2, r) \simeq K\Delta_r/I$.*

We find two new properties of $S^+(2, r)$ as follows. Recall that an algebra A is said to be *Schurian* ([S]) if $\dim_K eAe' \leq 1$ for any primitive orthogonal idempotents e, e' of A .

Proposition 2.15. *The Borel-Schur algebra $S^+(2, r)$ is a Schurian algebra.*

Proof. Let $\{e_i \mid 0 \leq i \leq r\}$ be the complete set of primitive orthogonal idempotents of $S^+(2, r)$. Since $S^+(2, r)$ is triangular, we have $\dim_K e_i S^+(2, r) e_i = 1$, for any $0 \leq i \leq r$. By the construction of the quiver of $S^+(2, r)$, there is no path from i to j for any $0 \leq i < j \leq r$, such that $\dim_K e_i S^+(2, r) e_j = 0$.

Suppose $0 \leq j < i \leq r$ and $p > 0$. If $i = j + 1$, there exists a unique path $\alpha_{j+1, j}$ from i to j such that $\dim_K e_i S^+(2, r) e_j = 1$. If $i = j + p^k$ for some $k \geq 1$, there are two paths $\alpha_{j+p^k, j}, \alpha_{j+p^k, j+(p-1)p^{k-1}} \cdots \alpha_{j+p^{k-1}, j}$ from i to j , but the latter one is zero by Proposition 2.14, hence, $\dim_K e_i S^+(2, r) e_j = 1$. If $i = j + p^k + p^\ell$ for some $k \neq \ell \geq 0$, there are two non-zero paths $\alpha_{s+p^k+p^\ell, s+p^k} \alpha_{s+p^k, s}, \alpha_{s+p^k+p^\ell, s+p^\ell} \alpha_{s+p^\ell, s}$ from i to j , but they are equal by Proposition 2.14, hence, $\dim_K e_i S^+(2, r) e_j = 1$. Otherwise, we may rewrite i as the form $j + p^{k_1} + \cdots + p^{k_m} + t$ with $0 \leq t \leq p - 1$; set $j' := j + p^{k_1} + \cdots + p^{k_m}$, and $\dim_K e_{j'} S^+(2, r) e_j = 1$ follows from the previous cases; we obtain $i = j' + t$ with $0 \leq t \leq p - 1$, there is a unique path $e_{j'}$ if $t = 0$, $\alpha_{i, j'+t-1} \cdots \alpha_{j'+1, j'}$ if $t \neq 0$, from i to j' , and hence $\dim_K e_i S^+(2, r) e_{j'} = 1$. \square

We denote by $\{S_i \mid 0 \leq i \leq r\}$ the set of all simple modules of $S^+(2, r)$. Then, the quiver isomorphism of $S^+(2, r)$ gives an anti-isomorphism of $S^+(2, r)$.

Proposition 2.16. *There exists an algebra anti-isomorphism $\zeta : S^+(2, r) \rightarrow S^+(2, r)$ by mapping S_i to S_{r-i} , for any $0 \leq i \leq r$.*

Proof. After replacing the indexed set $\Lambda(2, r)$ by $\{0, 1, \dots, r\}$, the involution ι on $\Lambda(2, r)$ is changed to the involution $\iota' : i \rightarrow r - i$ on $\{0, 1, \dots, r\}$. Suppose $S^+(2, r) \simeq KQ/I$. By Proposition 2.12, we have obtained an algebra isomorphism $\sigma : KQ^{\text{op}} \rightarrow KQ$ mapping $(S_i)^*$ to S_{r-i} . Moreover, reversing arrows produces the admissible ideal I^{op} of KQ^{op} such that $\sigma(I^{\text{op}}) = I$. We have $(KQ/I)^{\text{op}} \simeq KQ^{\text{op}}/I^{\text{op}} \xrightarrow{\sigma} KQ/I$, then $\zeta := \sigma \circ (-)^*$ gives the desired anti-isomorphism of $S^+(2, r)$. \square

2.3. Reduction on τ -tilting finiteness. We need several methods to check the τ -tilting finiteness of algebras.

Proposition 2.17 ([DIRRT, Theorem 5.12], [DIJ, Theorem 4.2]). *If A is τ -tilting finite,*

- (1) *the quotient algebra A/I is τ -tilting finite for any two-sided ideal I of A ,*
- (2) *the idempotent truncation eAe is τ -tilting finite for any idempotent e of A .*

Lemma 2.18. *Let $m \leq n$ and $s \leq r$. If $S^+(m, s)$ is τ -tilting infinite, then so is $S^+(n, r)$.*

Proof. It is shown in [ESY2, Proposition 3.2] that there exists an idempotent e of $S^+(n, r)$ satisfying $S^+(m, s) = eS^+(n, r)e$. Then, the statement follows from Proposition 2.17. \square

Let $A := K\Delta$ be the path algebra of an acyclic quiver Δ . Recall from [HR] that an A -module T is said to be *classical tilting* if $|T| = |A|$, $\text{Ext}_A^1(T, T) = 0$ and the projective dimension of T is at most one. If T is a tilting A -module, we call the endomorphism algebra $B := \text{End}_A T$ a *tilted algebra* of type Δ . If moreover, T is contained in a preprojective component of the Auslander-Reiten quiver of A , we call B a *concealed algebra* of type Δ . Tame concealed algebras are precisely the concealed algebras of type $\tilde{\mathbb{A}}_n, \tilde{\mathbb{D}}_n (n \geq 4), \tilde{\mathbb{E}}_6, \tilde{\mathbb{E}}_7$ and $\tilde{\mathbb{E}}_8$. The following result is well-known, for example, see [Mo, Remark 2.9].

Lemma 2.19. *A tame concealed algebra is τ -tilting infinite.*

It is worth mentioning that tame concealed algebras have been classified completely by quiver and relations in [HV] (see also [B]). We also mention (see [Wa2, Theorem 1.1]) that a simply connected algebra is τ -tilting finite if and only if it is representation-finite, if and only if it does not contain a concealed algebra of type $\tilde{\mathbb{D}}_n (n \geq 4), \tilde{\mathbb{E}}_6, \tilde{\mathbb{E}}_7$ or $\tilde{\mathbb{E}}_8$ as an idempotent truncation.

An algebra A is called *sincere* if there is an indecomposable A -module M such that all simple A -modules appear in M as composition factors. A complete list of representation-finite sincere simply connected algebras is given in [RT], in terms of quiver and relations.

3. SIGN DECOMPOSITION

This technical tool is first introduced in [Ao] to classify torsion classes for radical square zero algebras, and then generalized in [AHIKM] to study fans and polytopes in τ -tilting theory. Let $A = KQ/I$ be a bound quiver algebra whose vertex set is labeled by $[n] := \{1, 2, \dots, n\}$. We define

$$\mathfrak{s}_n := \{\epsilon = (\epsilon_1, \epsilon_2, \dots, \epsilon_n) : [n] \rightarrow \{\pm 1\}\}$$

to be the set of all maps from $[n]$ to $\{\pm 1\}$. In particular, \mathfrak{s}_n admits an involution mapping ϵ to $-\epsilon$ such that $(-\epsilon)_i = -\epsilon_i$. We sometimes write $\epsilon_i = +$ or $-$ instead of $\epsilon_i = +1$ or -1 without causing confusion.

Let \mathbb{Z}_ϵ^n be the area determined by the same sign with ϵ_i of $\epsilon \in \mathfrak{s}_n$, i.e.,

$$\mathbb{Z}_\epsilon^n := \{(x_1, x_2, \dots, x_n) \in \mathbb{Z}^n \mid \epsilon_i x_i > 0, \forall i \in [n]\}.$$

By comparing the signs of entries in each g -vector of two-term silting complexes, we may assign to each map $\epsilon \in \mathfrak{s}_n$ a subset of $2\text{-silt } A$, that is,

$$2\text{-silt}_\epsilon A := \{T \in 2\text{-silt } A \mid g(T) \in \mathbb{Z}_\epsilon^n\}.$$

It is obvious from the definition of \mathbb{Z}_ϵ^n that $2\text{-silt}_\epsilon A \cap 2\text{-silt}_{\epsilon'} A = \emptyset$ if $\epsilon \neq \epsilon'$. We have

$$(3.1) \quad 2\text{-silt } A = \bigsqcup_{\epsilon \in \mathfrak{s}_n} 2\text{-silt}_\epsilon A.$$

Let P_i be the indecomposable projective A -module at vertex i . In the definition of $\mu_{\underline{P}_i}^-(A)$, we take a minimal left $\text{add}(\underline{A}/\underline{P}_i)$ -approximation π of \underline{P}_i from the triangle

$$\underline{P}_i \xrightarrow{\pi} Z \longrightarrow X' \longrightarrow \underline{P}_i[1],$$

and it sends the degree 0 item of \underline{P}_i to the degree -1 part of X' . This means that the i -th entry of $g(\mu_{\underline{P}_i}^-(A))$ is negative. In particular, $2\text{-silt}_\epsilon A = \{A\}$ (resp., $\{A[1]\}$) if and only if $\epsilon = (+, \dots, +)$ (resp., $(-, \dots, -)$).

Fix $\epsilon \in \mathfrak{s}_n$. Suppose $[n] = I \cup J$ with $I = \{i \in [n] \mid \epsilon_i = -\}$ and $J = \{j \in [n] \mid \epsilon_j = +\}$. We define

$$P_I := \bigoplus_{i \in I} P_i \quad \text{and} \quad P_J := \bigoplus_{j \in J} P_j.$$

Proposition 3.1. *Suppose ϵ is neither $(+, \dots, +)$ nor $(-, \dots, -)$. We have*

$$2\text{-silt}_\epsilon A = \{T \in 2\text{-silt } A \mid \mu_{P_I}^-(A) \geq T \geq \mu_{P_J[1]}^+(A[1])\}.$$

Proof. As mentioned in (2.1), each entry in the g -vector $g(T)$ of $T \in 2\text{-silt } A$ is either positive or negative, and cannot be zero. For any $T \neq A$ in $2\text{-silt } A$, it is obtained by taking left mutations starting from A , hence, $\mu_{P_i}^-(A) \geq T$ for some $i \in [n]$. We conclude that $\mu_{P_i}^-(A) \geq T$ if and only if the i -th entry of $g(T)$ is negative, since the i -th entry of $g(\mu_{P_i}^-(A))$ is negative as mentioned above. By the definition of $2\text{-silt}_\epsilon A$, we find that $\epsilon_i = -$ if and only if $\mu_{P_i}^-(A) \geq T$. Then, the upper bound follows from the fact that $\epsilon_i = -$ for all $i \in I$. We may obtain the lower bound dually. \square

Note that $2\text{-silt}_\epsilon A$ could be a finite set even if $2\text{-silt } A$ is infinite. One also finds that $2\text{-silt}_\epsilon A$ is again a poset, and $\mathcal{H}(2\text{-silt}_\epsilon A)$ is a subquiver of $\mathcal{H}(2\text{-silt } A)$. Moreover, we have the following analog of Proposition 2.8.

Proposition 3.2. *Fix $\epsilon \in \mathfrak{s}_n$. If $2\text{-silt}_\epsilon A$ is finite, then $\mathcal{H}(2\text{-silt}_\epsilon A)$ is connected.*

Proof. It is shown in [AI, Proposition 2.36] that there exists an irreducible left mutation U of T satisfying $T > U \geq S$, for any $T > S$ in $\text{silt } A$. Dually, one may obtain a statement for irreducible right mutation. If both T and S are contained in $2\text{-silt}_\epsilon A$, then U also belongs to $2\text{-silt}_\epsilon A$ by Proposition 3.1. It suffices to say that $\mathcal{H}(2\text{-silt}_\epsilon A)$ is connected. \square

Proposition 3.3. *Fix $\epsilon \in \mathfrak{s}_n$. If $\mathcal{H}(2\text{-silt}_\epsilon A)$ contains a finite connected component \mathcal{C} , then \mathcal{C} exhausts all elements of $2\text{-silt}_\epsilon A$.*

Proof. Set $\Lambda = \mu_{P_I}^-(A)$ and $\Gamma = \mu_{P_J[1]}^+(A[1])$ in Proposition 3.1. For any $T \in 2\text{-silt}_\epsilon A$, we have $\Lambda \geq T$. Applying [AI, Proposition 2.36], we obtain an irreducible left mutation sequence $\Lambda = U_0 > U_1 > \cdots > U_i > \cdots > V_j > \cdots > V_1 > V_0 = \Gamma$ in \mathcal{C} such that $U_i \geq T \geq V_j$, for some i, j . Since \mathcal{C} is finite, we must have $T \in \mathcal{C}$. \square

Since we need to do direct calculations in some small cases and the calculation of left mutations in silting theory is very complicated, it is better to reformulate the above using the language of τ -tilting theory. We denote by $\mathcal{F} : 2\text{-silt } A \xrightarrow{\sim} \mathfrak{s}\tau\text{-tilt } A$ the poset isomorphism as mentioned in Theorem 2.6, and then, set

$$\mathfrak{s}\tau\text{-tilt}_\epsilon A := \{\mathcal{F}(T) \mid T \in 2\text{-silt}_\epsilon A\} \cap \mathfrak{s}\tau\text{-tilt } A.$$

We may define $\tau\text{-tilt}_\epsilon A$ similarly. Note that both $\tau\text{-tilt}_\epsilon A$ and $\mathfrak{s}\tau\text{-tilt}_\epsilon A$ are again posets.


Proposition 3.4. *Let $\epsilon \in \mathfrak{s}_n$. If the Hasse quiver $\mathcal{H}(\mathfrak{s}\tau\text{-tilt}_\epsilon A)$ (or $\mathcal{H}(\tau\text{-tilt}_\epsilon A)$) contains a finite connected component \mathcal{C} , then \mathcal{C} exhausts all elements of $\mathfrak{s}\tau\text{-tilt}_\epsilon A$ (or $\tau\text{-tilt}_\epsilon A$).*

Proof. The statement for $\mathfrak{s}\tau\text{-tilt}_\epsilon A$ is nothing but exactly a translation of Proposition 3.3. Since $\mathcal{H}(\tau\text{-tilt } A)$ is a connected subquiver of $\mathcal{H}(\mathfrak{s}\tau\text{-tilt } A)$ as well-known, we have the fact that a finite connected component of $\mathcal{H}(\tau\text{-tilt } A)$ exhausts all elements of $\tau\text{-tilt } A$ according to Proposition 2.8. Then, the statement for $\tau\text{-tilt}_\epsilon A$ is obtained by combining this fact and Proposition 3.3. \square

Example 3.5. Let A be the path algebra of the bipartite quiver: $1 \xrightarrow{\alpha_1} 2 \xleftarrow{\alpha_2} 3 \xrightarrow{\alpha_3} 4$. We take $I = \{2, 4\}$ and $J = \{1, 3\}$, i.e., $\epsilon = (+, -, +, -)$. Then,

$$\mu_{P_I}^-(A) = \begin{bmatrix} P_2 \xrightarrow{(\alpha_1, \alpha_2)^t} P_1 \oplus P_3 \\ \oplus \\ P_4 \xrightarrow{\alpha_3} P_3 \\ \oplus \\ 0 \longrightarrow P_1 \oplus P_3 \end{bmatrix} \quad \text{and} \quad \mu_{P_J[1]}^+(A[1]) = \begin{bmatrix} P_2 \oplus P_4 \xrightarrow{(\alpha_2, \alpha_3)} P_3 \\ \oplus \\ P_2 \xrightarrow{\alpha_1} P_1 \\ \oplus \\ P_2 \oplus P_4 \longrightarrow 0 \end{bmatrix}.$$

The Hasse quiver $\mathcal{H}(\mathfrak{s}\tau\text{-tilt}_\epsilon A)$ has the following connected component:



in which \ominus and $\boxed{-}$ indicate the elements in $\tau\text{-tilt}_\epsilon A$ and $\mathfrak{s}\tau\text{-tilt}_\epsilon A$, respectively. (Note that $\mathcal{H}(\mathfrak{s}\tau\text{-tilt}_\epsilon A)$ contains both the maximal and minimal elements, but $\mathcal{H}(\tau\text{-tilt}_\epsilon A)$ may only contain the maximal element.) The information on each vertex is given below.

- 1) $\mu_{P_1}^-(A) \mapsto \frac{1}{2} \oplus \frac{1}{2} \frac{3}{4} \oplus \frac{3}{4} \oplus \frac{3}{2} \mapsto ((1, 0, 0, 0), (1, -1, 1, 0), (0, 0, 1, 0), (0, 0, 1, -1)),$
- 2) $\frac{1}{2} \oplus \frac{1}{2} \frac{3}{4} \oplus \frac{1}{2} \frac{3}{2} \oplus \frac{3}{2} \mapsto ((1, 0, 0, 0), (1, -1, 1, 0), (1, -1, 1, -1), (0, 0, 1, -1)),$
- 3) $\frac{3}{4} \oplus \frac{1}{2} \frac{3}{4} \oplus \frac{3}{4} \oplus \frac{3}{2} \mapsto ((0, -1, 1, 0), (1, -1, 1, 0), (0, 0, 1, 0), (0, 0, 1, -1)),$
- 4) $\frac{1}{2} \oplus \frac{1}{2} \frac{3}{4} \oplus \frac{1}{2} \frac{3}{2} \oplus 1 \mapsto ((1, 0, 0, 0), (1, -1, 1, 0), (1, -1, 1, -1), (1, -1, 0, 0)),$
- 5) $(\frac{1}{2} \oplus \frac{1}{2} \frac{3}{2} \oplus \frac{3}{2}, 4) \mapsto ((1, 0, 0, 0), (1, -1, 1, -1), (0, 0, 1, -1), (0, 0, 0, -1)),$
- 6) $\frac{3}{4} \oplus \frac{1}{2} \frac{3}{4} \oplus \frac{1}{2} \frac{3}{2} \oplus \frac{3}{2} \mapsto ((0, -1, 1, 0), (1, -1, 1, 0), (1, -1, 1, -1), (0, 0, 1, -1)),$
- 7) $\frac{3}{4} \oplus 3 \oplus \frac{1}{2} \frac{3}{2} \oplus \frac{3}{2} \mapsto ((0, -1, 1, 0), (0, -1, 1, -1), (1, -1, 1, -1), (0, 0, 1, -1)),$
- 8) $\frac{3}{4} \oplus \frac{1}{2} \frac{3}{4} \oplus \frac{1}{2} \frac{3}{2} \oplus 1 \mapsto ((0, -1, 1, 0), (1, -1, 1, 0), (1, -1, 1, -1), (1, -1, 0, 0)),$
- 9) $(3 \oplus \frac{1}{2} \frac{3}{2} \oplus \frac{3}{2}, 4) \mapsto ((0, -1, 1, -1), (1, -1, 1, -1), (0, 0, 1, -1), (0, 0, 0, -1)),$
- 10) $\frac{3}{4} \oplus 3 \oplus \frac{1}{2} \frac{3}{2} \oplus 1 \mapsto ((0, -1, 1, 0), (0, -1, 1, -1), (1, -1, 1, -1), (1, -1, 0, 0)),$
- 11) $(\frac{1}{2} \oplus \frac{1}{2} \frac{3}{2} \oplus 1, 4) \mapsto ((1, 0, 0, 0), (1, -1, 1, -1), (1, -1, 0, 0), (0, 0, 0, -1)),$
- 12) $(3 \oplus \frac{1}{2} \frac{3}{2} \oplus 1, 4) \mapsto ((0, -1, 1, -1), (1, -1, 1, -1), (1, -1, 0, 0), (0, 0, 0, -1)),$
- 13) $(\frac{3}{4} \oplus 3 \oplus 1, 2) \mapsto ((0, -1, 1, 0), (0, -1, 1, -1), (1, -1, 0, 0), (0, -1, 0, 0)),$
- 14) $\mu_{P_J[1]}^+(A[1]) \mapsto (3 \oplus 1, 2 \oplus 4) \mapsto ((0, -1, 1, -1), (1, -1, 0, 0), (0, -1, 0, 0), (0, 0, 0, -1)).$

We next are aiming to characterize the elements in $2\text{-silt}_\epsilon A$ via a simpler algebra A_ϵ . Let e_i be the primitive idempotent of A associated with the vertex i . For each $\epsilon \in \mathfrak{s}_n$, set

$$e_{\epsilon,+} := \sum_{i \in [n], \epsilon_i = +} e_i \quad \text{and} \quad e_{\epsilon,-} := \sum_{i \in [n], \epsilon_i = -} e_i.$$

Then, let $J_{\epsilon,+}$ be the two-sided ideal of $e_{\epsilon,+} A e_{\epsilon,+}$ generated by all $x \in \text{rad}(e_{\epsilon,+} A e_{\epsilon,+})$ satisfying $xy = 0$ for any $y \in e_{\epsilon,+} A e_{\epsilon,-}$. Similarly, let $J_{\epsilon,-}$ be the two-sided ideal of $e_{\epsilon,-} A e_{\epsilon,-}$ generated by all $x \in \text{rad}(e_{\epsilon,-} A e_{\epsilon,-})$ satisfying $yx = 0$ for any $y \in e_{\epsilon,+} A e_{\epsilon,-}$.

Definition 3.6 ([AoW, Definition 2.17]). We define an upper triangular matrix algebra:

$$A_\epsilon := \begin{pmatrix} e_{\epsilon,+} A e_{\epsilon,+} / J_{\epsilon,+} & e_{\epsilon,+} A e_{\epsilon,-} \\ 0 & e_{\epsilon,-} A e_{\epsilon,-} / J_{\epsilon,-} \end{pmatrix}.$$

It is better to give an example to illustrate the structure of A_ϵ .

Example 3.7. Let A be the path algebra of the linear quiver²: $1 \xrightarrow{\alpha_1} 2 \xrightarrow{\alpha_2} 3 \xrightarrow{\alpha_3} 4$.

- If $\epsilon = (+, +, +, +)$, then $e_{\epsilon,+} A e_{\epsilon,-} = 0$ and $\alpha_i \in J_{\epsilon,+}$. Thus, $A_\epsilon = K \oplus K \oplus K \oplus K$.
- If $\epsilon = (+, -, +, +)$, then $e_{\epsilon,+} A e_{\epsilon,-} = \{\alpha_1\}$, $\alpha_1 \alpha_2, \alpha_1 \alpha_2 \alpha_3, \alpha_3 \in J_{\epsilon,+}$, and $\alpha_2, \alpha_2 \alpha_3$ are removed. Thus, $A_\epsilon = K(1 \xrightarrow{\alpha_1} 2) \oplus K \oplus K$.
- If $\epsilon = (+, -, +, -)$, then $e_{\epsilon,+} A e_{\epsilon,-} = \{\alpha_1, \alpha_3, \alpha_1 \alpha_2 \alpha_3\}$, $e_{\epsilon,+} A e_{\epsilon,+} = \{e_1, e_3, \alpha_1 \alpha_2\}$, $e_{\epsilon,-} A e_{\epsilon,-} = \{e_2, e_4, \alpha_2 \alpha_3\}$, $J_{\epsilon,+} = J_{\epsilon,-} = 0$ and α_2 is removed. Thus, A_ϵ is exactly the algebra given in Example 2.7.

It turns out that A_ϵ shares the same vertex set $\{1, 2, \dots, n\}$ with A . Furthermore,

²Since we consider right modules in this paper, the paths here are read from left to right.

Proposition 3.8. *Let $\epsilon = (\epsilon_1, \epsilon_2, \dots, \epsilon_n) \in \mathfrak{s}_n$ and i a vertex in the quiver of A .*

- (1) *If i is a source and $\epsilon_i = -$, then i is an isolated vertex in the quiver of A_ϵ .*
- (2) *If i is a sink and $\epsilon_i = +$, then i is an isolated vertex in the quiver of A_ϵ .*

Proof. This is obvious from the definition of A_ϵ . □

The above Proposition 3.8 is useful when one deals with one-point extension algebras. Recall that the *one-point extension* $A[M]$ of A by an A -module M is defined as

$$A[M] := \begin{bmatrix} A & 0 \\ M & K \end{bmatrix}.$$

The quiver of $A[M]$ then contains the quiver of A as a full subquiver with exactly one additional extension vertex which is always a source. Hence, $(A[M])_\epsilon = (K \oplus A)_\epsilon$ if the sign of the extension vertex is $-$.

It is shown in [ESY2] that $S^+(2, r)$ is an one-point extension of $S^+(2, r-1)$. More precisely, let P_r be the indecomposable projective module of $S^+(2, r)$ at vertex r , then $\text{rad } P_r$ is an indecomposable module of $S^+(2, r-1)$ and

$$S^+(2, r) = \begin{bmatrix} S^+(2, r-1) & 0 \\ \text{rad } P_r & K \end{bmatrix}.$$

One finds that the vertex corresponding to $\text{rad } P_r$ in the quiver of $S^+(2, r)$ is a source. Recall that we use $\{0, 1, 2, \dots, r\}$ to label the vertices in the quiver of $S^+(2, r)$. Then, 0 is a sink and r is a source. We have the following reduction result for Borel-Schur algebras.

Proposition 3.9. *Let $\epsilon = (\epsilon_0, \epsilon_1, \dots, \epsilon_r)$. If $\epsilon_r = -$, then $S^+(2, r)_\epsilon = (S^+(2, r-1) \oplus K)_\epsilon$.*

One of the most important properties of sign decomposition is stated as follows.

Proposition 3.10 ([AHIKM, Theorem 4.23 and Proposition 4.25]). *For any $\epsilon \in \mathfrak{s}_n$, there is an isomorphism*

$$2\text{-silt}_\epsilon A \xrightarrow{\sim} 2\text{-silt}_\epsilon A_\epsilon,$$

which preserves g -vectors of two-term silting complexes.

As a corollary of Proposition 3.10 together with (3.1), it follows that A is τ -tilting finite if $2\text{-silt}_\epsilon A_\epsilon$ is a finite set for all $\epsilon \in \mathfrak{s}_n$. Although A has a complicated structure, the finiteness of $2\text{-silt}_\epsilon A_\epsilon$ can be easily verified for most maps in \mathfrak{s}_n , with only a few cases remaining. Moreover, these remaining cases may be resolved by the tilting mutation process as mentioned below. (This approach has been applied to the block algebras of Schur algebras, see [AoW].)

Let $\Phi := \{\epsilon \in \mathfrak{s}_n \mid \epsilon_i = -\}$ and $-\Phi := \{-\epsilon \mid \epsilon \in \Phi\}$. We define

$$2\text{-silt}_\Phi A := \bigsqcup_{\epsilon \in \Phi} 2\text{-silt}_\epsilon A.$$

Proposition 3.11 ([AoW, Proposition 2.20]). *Let $B := \text{End } \mu_{P_i}^-(A)$ be the endomorphism algebra of the left mutation $\mu_{P_i}^-(A)$ of A with respect to the indecomposable projective module P_i . If $\mu_{P_i}^-(A)$ is tilting, then there is a poset isomorphism*

$$(3.2) \quad 2\text{-silt}_{\Phi} A \xrightarrow{\sim} 2\text{-silt}_{-\Phi} B.$$

Remark 3.12. The poset isomorphism between $2\text{-silt}_{\Phi} A$ and $2\text{-silt}_{-\Phi} B$ first appeared in [AMN, Lemma 4.6], in which the authors deal with Brauer tree algebras. Then, a similar proof for arbitrary algebras is given in [AoW, Proposition 2.20]. See also [AHIKM, Theorem 4.28] for a proof in terms of g -fans.

It is worth mentioning that the poset isomorphism in Proposition 3.11 does not imply an involution on the sign of g -vectors, i.e., $2\text{-silt}_{\epsilon} A \not\cong 2\text{-silt}_{-\epsilon} B$ for $\epsilon \in \Phi$. However, we may describe the relation precisely as follows. For any $T \in \text{silt } A$, we define $\mathcal{G}(T)$ to be the g -matrix of T .

Proposition 3.13. *Suppose $\mu_{P_i}^-(A)$ is a tilting complex. For any $T \in 2\text{-silt}_{\Phi} A$, the poset isomorphism in (3.2) is induced by $g(T) \mapsto g(T) \cdot \mathcal{G}(\mu_{P_i}^-(A))$.*

Proof. It is known from Rickard [Ric] that there is a triangle equivalence from $\text{D}^b(\text{mod } A)$ to $\text{D}^b(\text{mod } B)$ given by $\mu_{P_i}^-(A) \mapsto B$. Here, $\text{D}^b(\text{mod } A)$ is the derived category of $\text{mod } A$. In this way, the indecomposable direct summand S_i of $\mu_{P_i}^-(A)$ is mapped to the indecomposable projective module Q_i of B . The triangle equivalence naturally induces an isomorphism $K_0(\mathcal{K}_A) \rightarrow K_0(\mathcal{K}_B)$ of Grothendieck groups by $[S_i] \mapsto [Q_i]$, which sends a g -vector $g(T)$ to a g -vector $g(T) \cdot \mathcal{G}(\mu_{P_i}^-(A))$. By Proposition 2.5, each two-term silting complex T in $2\text{-silt } A$ is uniquely determined by its g -vector $g(T)$, and hence, $g(T) \cdot \mathcal{G}(\mu_{P_i}^-(A))$ uniquely gives a two-term silting complex in $2\text{-silt } B$. \square

We will provide an example in Section 5 to explain how to apply Proposition 3.13.

At the end of this section, we reformulate the anti-isomorphism between $2\text{-silt } A$ and $2\text{-silt } A^{\text{op}}$ and the anti-automorphism of $2\text{-silt } A$ as follows. Let $(-)^* := \text{Hom}_A(-, A)$. For a given $T \in 2\text{-silt } A$, we have

$$T = \left(0 \longrightarrow \bigoplus_{i \in I} P_i^{\oplus a_i} \xrightarrow{f} \bigoplus_{j \in J} P_j^{\oplus a_j} \longrightarrow 0 \right)$$

with $I \cap J = 0$ and $I \cup J = [n]$ (see (2.1)). Then,

$$T^* = \left(0 \longrightarrow 0 \longrightarrow \bigoplus_{j \in J} (P_j^*)^{\oplus a_j} \xrightarrow{f^*} \bigoplus_{i \in I} (P_i^*)^{\oplus a_i} \right).$$

If there exists an algebra isomorphism $\sigma : A^{\text{op}} \rightarrow A$, then σ induces a permutation on $\{1, 2, \dots, n\}$ by $\sigma(e_i^*) = e_j$. Here, $P_i^* = e_i^* A$. We then obtain

$$\sigma(T^*) = \left(0 \longrightarrow 0 \longrightarrow \bigoplus_{j \in J} (P_{\sigma(j)})^{\oplus a_{\sigma(j)}} \xrightarrow{\sigma(f^*)} \bigoplus_{i \in I} (P_{\sigma(i)})^{\oplus a_{\sigma(i)}} \right),$$

which is again a silting complex in \mathcal{K}_A . Set $S_{\sigma} := [1] \circ \sigma \circ (-)^*$.

Proposition 3.14 ([AIR, Theorem 2.14]). *For any $\epsilon \in \mathfrak{s}_n$, the duality $(-)^*$ induces a bijection $2\text{-silt}_\epsilon A \xrightarrow{1:1} 2\text{-silt}_{-\epsilon} A^{\text{op}}$.*

Proposition 3.15 ([AiW, Theorem 1.2]). *Let $\epsilon = (\epsilon_1, \epsilon_2, \dots, \epsilon_n) \in \mathfrak{s}_n$ and $\sigma : A^{\text{op}} \rightarrow A$ an algebra isomorphism. Then, the functor S_σ gives a bijection $2\text{-silt}_\epsilon A \xrightarrow{1:1} 2\text{-silt}_{-\sigma(\epsilon)} A$, where $\sigma(\epsilon) = (\epsilon_{\sigma(1)}, \epsilon_{\sigma(2)}, \dots, \epsilon_{\sigma(n)})$.*

Corollary 3.16. *Let $\epsilon = (\epsilon_1, \epsilon_2, \dots, \epsilon_n) \in \mathfrak{s}_n$. If there is an algebra isomorphism $\sigma : A^{\text{op}} \rightarrow A$, then we have $A_\epsilon \simeq (A_{-\sigma(\epsilon)})^{\text{op}}$.*

4. τ -TILTING INFINITE BOREL-SCHUR ALGEBRAS

In this section, we show that most representation-infinite Borel-Schur algebras are τ -tilting infinite. We will frequently use Proposition 2.17, Lemma 2.18 and Lemma 2.19 without repeated citations.

Proposition 4.1. *Let $p \geq 0$ and $n \geq 3$. Then $S^+(n, r)$ is τ -tilting infinite for any $r \geq 2$.*

Proof. By Lemma 2.18, it suffices to consider $S^+(3, 2)$. By [ESY2, Section 5], $S^+(3, 2)$ contains a tame concealed algebra of type $\tilde{\mathbb{A}}_3$, i.e., the path algebra of the quiver:

$$\begin{array}{ccc} \circ & \longrightarrow & \circ \\ \downarrow & & \downarrow \\ \circ & \longrightarrow & \circ \end{array},$$

as an idempotent truncation. Based on Proposition 2.17 and Lemma 2.19, we conclude that $S^+(3, 2)$ is τ -tilting infinite. \square

Proposition 4.2. *Let $p = 2$. Then $S^+(2, r)$ is τ -tilting infinite for any $r \geq 5$.*

Proof. We show that $S^+(2, 5)$ is τ -tilting infinite. By Proposition 2.14, $S^+(2, 5)$ over $p = 2$ is isomorphic to KQ/I with

$$Q : \begin{array}{ccccccccc} & & & \gamma_1 & & & & & \\ & & & \curvearrowright & & & & & \\ & & & \beta_2 & & & & & \\ & & & \curvearrowright & & & & & \\ & & & \beta_0 & & & & & \\ & & & \curvearrowright & & & & & \\ 5 & \xrightarrow{\alpha_4} & 4 & \xrightarrow{\alpha_3} & 3 & \xrightarrow{\alpha_2} & 2 & \xrightarrow{\alpha_1} & 1 & \xrightarrow{\alpha_0} & 0 \\ & & & \curvearrowleft & & & & & & & \\ & & & \beta_3 & & & & & & & \\ & & & \curvearrowleft & & & & & & & \\ & & & \beta_1 & & & & & & & \\ & & & \curvearrowleft & & & & & & & \\ & & & \gamma_0 & & & & & & & \end{array}$$

and

$$I : \left\langle \begin{array}{l} \alpha_4\alpha_3, \alpha_3\alpha_2, \alpha_2\alpha_1, \alpha_1\alpha_0, \beta_3\beta_1, \beta_2\beta_0, \gamma_1\alpha_0 - \alpha_4\gamma_0 \\ \alpha_4\beta_2 - \beta_3\alpha_2, \alpha_3\beta_1 - \beta_2\alpha_1, \alpha_2\beta_0 - \beta_1\alpha_0 \end{array} \right\rangle.$$


It is obvious that $S^+(2, 5)$ over $p = 2$ contains a tame concealed algebra of type $\tilde{\mathbb{A}}_5$, i.e., the path algebra of the bipartite quiver:

$$\begin{array}{ccc} 0 & \longleftarrow & 2 & \longrightarrow & 1 \\ \uparrow & & & & \uparrow \\ 4 & \longrightarrow & 3 & \longleftarrow & 5 \end{array},$$

as a quotient algebra. Then, it is τ -tilting infinite. \square

Proposition 4.3. *Let $p = 3$. Then $S^+(2, r)$ is τ -tilting infinite for any $r \geq 6$.*


Proof. By Proposition 2.14, $S^+(2, 6)$ over $p = 3$ is isomorphic to KQ/I with



and

$$I : \langle \alpha_5\alpha_4\alpha_3, \alpha_4\alpha_3\alpha_2, \alpha_3\alpha_2\alpha_1, \alpha_2\alpha_1\alpha_0, \alpha_5\beta_2 - \beta_3\alpha_2, \alpha_4\beta_1 - \beta_2\alpha_1, \alpha_3\beta_0 - \beta_1\alpha_0 \rangle.$$


It turns out that $S^+(2, 6)$ over $p = 3$ contains a tame concealed algebra of type $\widetilde{\mathbb{D}}_6$, i.e.,



as a quotient algebra, and it is τ -tilting infinite. \square

Proposition 4.4. *Let $p = 5$. Then $S^+(2, r)$ is τ -tilting infinite for any $r \geq 7$.*


Proof. By Proposition 2.14, $S^+(2, 7)$ over $p = 5$ is isomorphic to KQ/I with



and

$$I : \langle \alpha_6\alpha_5\alpha_4\alpha_3\alpha_2, \alpha_5\alpha_4\alpha_3\alpha_2\alpha_1, \alpha_4\alpha_3\alpha_2\alpha_1\alpha_0, \alpha_6\beta_1 - \beta_2\alpha_1, \alpha_5\beta_0 - \beta_1\alpha_0 \rangle.$$


It turns out that $S^+(2, 7)$ over $p = 5$ contains a tame concealed algebra of type $\widetilde{\mathbb{E}}_7$, i.e.,



as a quotient algebra, and it is τ -tilting infinite. \square

Proposition 4.5. *Let $p \geq 7$. Then $S^+(2, r)$ is τ -tilting infinite for any $r \geq p + 1$.*

Proof. It is shown in [ESY1, Theorem 6.6] that $S^+(2, p+1)$ over $p \geq 7$ contains a concealed algebra of type $\widetilde{\mathbb{E}}_7$, i.e.,



as an idempotent truncation, and it is τ -tilting infinite. \square

In summary, we have the following result.

Theorem 4.6. *Let $p \geq 0$. Then, $S^+(n, r)$ is τ -tilting infinite if*

- $n \geq 3$ and $r \geq 2$; or
- $n = 2$ and $p = 2, r \geq 5$ or $p = 3, r \geq 6$ or $p = 5, r \geq 7$ or $p \geq 7, r \geq p + 1$.

5. τ -TILTING FINITE BOREL-SCHUR ALGEBRAS

Representation-finite Borel-Schur algebras are naturally τ -tilting finite. By combining this fact with the τ -tilting infinite cases established in the previous section, we are left with the remaining cases: $S^+(2, 4)$ over $p = 2$, $S^+(2, 5)$ over $p = 3$ and $S^+(2, 6)$ over $p = 5$. In this section, we show that all these cases are τ -tilting finite.

We introduce some technical methods for later use.

- Let w be a maximal path of $A = KQ/I$, i.e., a path w such that $w \notin I$ but $\alpha w, w\alpha \in I$ for all arrows $\alpha \in Q$. If $w = \alpha_1\alpha_2 \cdots \alpha_k \in e_{\epsilon,+}Ae_{\epsilon,-}$ and α_i is removed in A_ϵ for some i , then we should replace the minimal subpath containing α_i by a new arrow β in A_ϵ . If $w \in e_{\epsilon,+}Ae_{\epsilon,+}$ (or $e_{\epsilon,-}Ae_{\epsilon,-}$), then $w \in J_{\epsilon,+}$ (or $J_{\epsilon,-}$) by the definition.
- Let A be the path algebra of $1 \rightarrow 2 \rightarrow 3 \rightarrow 4$ and B the algebra presented by

$$\begin{array}{ccc} 1 & \rightarrow & 3 \\ \downarrow & \searrow & \downarrow \\ 2 & \rightarrow & 4 \end{array}.$$

Then, $A_{(+,-,+,-)} \simeq B_{(+,-,+,-)}$ as explained in Example 3.7. If the quiver of A or B appears as a subquiver in the quiver of $S^+(n, r)$, then we may exchange $A_{(+,-,+,-)}$ and $B_{(+,-,+,-)}$ (in an appropriate way) to reduce the complexity.

- Let $\sigma : A^{\text{op}} \rightarrow A$ be an algebra isomorphism. We use $\epsilon \sim_\sigma \epsilon'$ to indicate $\epsilon' = -\sigma(\epsilon)$, and then, $A_\epsilon \simeq (A_{\epsilon'})^{\text{op}}$ by Corollary 3.16.
- We use the *String Applet* to check the representation type of gentle algebras and special biserial algebras.

Proposition 5.1. *Let $p = 2$. Then $S^+(2, 4)$ is τ -tilting finite.*

Proof. By Proposition 2.14, $S^+(2, 4)$ over $p = 2$ is isomorphic to $A := KQ/I$ with

$$Q : \begin{array}{ccccccc} & & \beta_2 & & \beta_0 & & \\ & & \curvearrowright & & \curvearrowright & & \\ 4 & \xrightarrow{\alpha_3} & 3 & \xrightarrow{\alpha_2} & 2 & \xrightarrow{\alpha_1} & 1 & \xrightarrow{\alpha_0} & 0 \\ & & \curvearrowleft & & \curvearrowleft & & & & \\ & & & & \beta_1 & & & & \\ & & & & \curvearrowright & & & & \\ & & & & \gamma_0 & & & & \end{array} \quad \text{and} \quad I : \left\langle \begin{array}{l} \alpha_3\alpha_2, \alpha_2\alpha_1, \alpha_1\alpha_0, \beta_2\beta_0, \\ \alpha_3\beta_1 - \beta_2\alpha_1, \alpha_2\beta_0 - \beta_1\alpha_0 \end{array} \right\rangle.$$

We check the τ -tilting finiteness of A_ϵ for $\epsilon = (\epsilon_0, \epsilon_1, \dots, \epsilon_4) \in \mathfrak{s}_5$ case by case.

If $\epsilon_0 = +$ or $\epsilon_4 = -$, we have $A_\epsilon \simeq (K \oplus S^+(2, 3))_\epsilon$ by Proposition 3.8 and Proposition 3.9. Then, A_ϵ is τ -tilting finite since $S^+(2, 3)$ over $p = 2$ is representation-finite.

Suppose $\epsilon_0 = -$ and $\epsilon_4 = +$. Since there is an algebra isomorphism $\sigma : A^{\text{op}} \rightarrow A$ sending e_i^* to e_{4-i} , it suffices to consider the following 4 cases by Corollary 3.16.

$$(-, +, +, +, +) \sim_{\sigma} (-, -, -, -, +), \quad (-, +, +, -, +) \sim_{\sigma} (-, +, -, -, +),$$

$$(-, +, -, +, +) \sim_{\sigma} (-, -, +, -, +), \quad (-, -, -, +, +) \sim_{\sigma} (-, -, +, +, +).$$

Let P_i be the indecomposable projective A -module at vertex i . Then, $\text{Hom}_A(P_i, P_j)$ is given in the following table.

Hom	0	1	2	3	4
0	e_0	α_0	β_0	$\alpha_2\beta_0$	γ_0
1	0	e_1	α_1	β_1	$\alpha_3\beta_1$
2	0	0	e_2	α_2	β_2
3	0	0	0	e_3	α_3
4	0	0	0	0	e_4

Some A_{ϵ} 's can be easily found as follows:

$$\begin{array}{ccc}
\begin{array}{c} 3 \longrightarrow 2 \\ \downarrow \quad \searrow \quad \downarrow \\ 1 \longrightarrow 0 \longleftarrow 4 \end{array} & \begin{array}{c} 1 \longleftarrow 2 \\ \uparrow \quad \searrow \quad \uparrow \\ 3 \longleftarrow 4 \longrightarrow 0 \end{array} & \begin{array}{c} 2 \\ \downarrow \\ 1 \longrightarrow 0 \longleftarrow 4 \longrightarrow 3 \end{array} & \begin{array}{c} 2 \\ \uparrow \\ 3 \longleftarrow 4 \longrightarrow 0 \longleftarrow 1 \end{array} \\
A_{(-,+,+,+,+)} & A_{(-,-,-,-,+)} & A_{(-,+,+,-,+)} & A_{(-,+,-,-,+)}
\end{array}$$

i.e., the first two are representation-finite sincere simply connected algebras and the last two are path algebras of type \mathbb{D}_5 . The reader may use these to verify Corollary 3.16. We give the details as follows.

- Set $\epsilon = (-, +, +, +, +)$. Then, $\alpha_1, \alpha_3, \beta_2, \alpha_3\beta_1 \in J_{\epsilon,+}$ and all others are survived.
- Set $\epsilon = (-, +, +, -, +)$. Then, $\alpha_1, \beta_2, \alpha_3\beta_1 \in J_{\epsilon,+}$, $\alpha_2\beta_0 \in J_{\epsilon,-}$, α_2, β_1 are removed and all others are survived.
- Set $\epsilon = (-, +, -, +, +)$. Then, $\alpha_3, \alpha_3\beta_1 \in J_{\epsilon,+}$, α_1 is removed and all others are survived. We obtain that

$$A_{\epsilon} = K \left(\begin{array}{c} 3 \xrightarrow{\alpha_2} 2 \xleftarrow{\beta_2} \\ \beta_1 \downarrow \quad \searrow \quad \downarrow \beta_0 \\ 1 \xrightarrow{\alpha_0} 0 \xleftarrow{\gamma_0} 4 \end{array} \right) / \langle \alpha_2\beta_0 - \beta_1\alpha_0, \beta_2\beta_0 \rangle.$$

We define

$$B := K \left(\begin{array}{c} 3 \xrightarrow{\alpha_2} 2 \xrightarrow{x} 1 \xrightarrow{\alpha_0} 0 \\ \beta_2 \searrow \quad \quad \nearrow \gamma_0 \\ 4 \end{array} \right) / \langle \beta_2x \rangle,$$

i.e., we replace a commutativity square subquiver in A_{ϵ} with a linear subquiver without relation in B . It is easy to check that $A_{\epsilon} \simeq B_{\epsilon}$. Since B is a representation-finite gentle algebra, we deduce that $2\text{-silt}_{\epsilon} A_{\epsilon} \simeq 2\text{-silt}_{\epsilon} B_{\epsilon}$ is a finite set.

- Set $\epsilon = (-, -, -, +, +)$. This case is slightly more complicated because of $A_{\epsilon} \simeq A$. We apply Proposition 3.1 and Proposition 3.4 to this case. Set $P_I = P_0 \oplus P_1 \oplus P_2$,

$$\mu_{P_I}^-(A) = \left[\begin{array}{ccc} P_1 & \xrightarrow{\beta_1} & P_3 \\ & \oplus & \\ P_0 & \xrightarrow{(\alpha_2\beta_0, \gamma_0)^t} & P_3 \oplus P_4 \\ & \oplus & \\ P_2 & \xrightarrow{(\alpha_2, \beta_2)^t} & P_3 \oplus P_4 \\ & \oplus & \\ 0 & \longrightarrow & P_3 \oplus P_4 \end{array} \right] \in 2\text{-silt}_\epsilon A.$$

Then we have $g(\mu_{P_I}^-(A)) = (-1, -1, -1, 4, 3)$ and

$$\mathcal{F}(\mu_{P_I}^-(A)) \simeq \begin{array}{c} 3 \\ \diagdown \quad \diagup \\ 2 \end{array} \oplus \begin{array}{c} 3 \quad 4 \\ \diagdown \quad \diagup \\ 1 \quad 3 \\ \diagdown \quad \diagup \\ 0 \quad 1 \end{array} 2 \oplus \begin{array}{c} 3 \quad 4 \\ \diagdown \quad \diagup \\ 1 \quad 2 \quad 3 \quad 0 \end{array} \oplus \begin{array}{c} 3 \\ \diagdown \quad \diagup \\ 2 \quad 1 \\ \diagdown \quad \diagup \\ 0 \end{array} \oplus \begin{array}{c} 4 \\ \diagdown \quad \diagup \\ 0 \quad 3 \\ \diagdown \quad \diagup \\ 1 \quad 2 \end{array}.$$

All elements in $\tau\text{-tilt}_\epsilon A$ can be obtained by left mutations starting from $\mathcal{F}(\mu_{P_I}^-(A))$ if $\mathcal{H}(\tau\text{-tilt}_\epsilon A)$ admits a finite connected component. By direct calculation, we find that $\#\tau\text{-tilt}_\epsilon A = 34$. A complete list of g -matrices of τ -tilting modules in $\tau\text{-tilt}_\epsilon A$ is given below.

- 1) $((0, -1, 0, 1, 0), (0, 0, -1, 1, 1), (-1, 0, 0, 1, 1), (0, 0, 0, 1, 0), (0, 0, 0, 0, 1))$,
- 2) $((0, -1, 0, 1, 0), (0, 0, -1, 1, 1), (-1, 0, 0, 1, 1), (0, 0, 0, 1, 0), (-1, 0, -1, 2, 1))$,
- 3) $((0, -1, 0, 1, 0), (0, 0, -1, 1, 1), (-1, 0, 0, 1, 1), (0, 0, 0, 0, 1), (-1, 0, 0, 0, 1))$,
- 4) $((0, -1, 0, 1, 0), (0, 0, -1, 1, 1), (-1, 0, 0, 0, 1), (0, -1, -1, 1, 1), (0, 0, 0, 0, 1))$,
- 5) $((0, -1, 0, 1, 0), (0, 0, -1, 1, 1), (-1, 0, 0, 0, 1), (0, -1, -1, 1, 1), (-1, 0, -1, 1, 1))$,
- 6) $((0, -1, 0, 1, 0), (0, 0, -1, 1, 1), (-1, 0, 0, 0, 1), (-1, 0, -1, 2, 1), (-1, 0, -1, 1, 1))$,
- 7) $((0, -1, 0, 1, 0), (0, 0, -1, 1, 1), (-1, 0, 0, 0, 1), (-1, 0, -1, 2, 1), (-1, 0, 0, 1, 1))$,
- 8) $((0, -1, 0, 1, 0), (0, 0, -1, 1, 1), (0, 0, -1, 1, 0), (-1, 0, -1, 2, 1), (0, 0, 0, 1, 0))$,
- 9) $((0, -1, 0, 1, 0), (0, 0, -1, 1, 1), (0, 0, -1, 1, 0), (-1, 0, -1, 2, 1), (-1, 0, -1, 1, 1))$,
- 10) $((0, -1, 0, 1, 0), (0, 0, -1, 1, 1), (0, 0, -1, 1, 0), (0, -1, -1, 1, 1), (-1, 0, -1, 1, 1))$,
- 11) $((0, -1, 0, 1, 0), (0, -1, -1, 1, 1), (-1, 0, 0, 0, 1), (0, -1, 0, 0, 1), (0, 0, 0, 0, 1))$,
- 12) $((0, -1, 0, 1, 0), (0, -1, -1, 1, 1), (-1, 0, 0, 0, 1), (0, -1, 0, 0, 1), (-1, -1, 0, 0, 1))$,
- 13) $((0, -1, 0, 1, 0), (0, -1, -1, 1, 1), (-1, 0, 0, 0, 1), (-1, 0, -1, 1, 1), (-1, -1, -1, 1, 1))$,
- 14) $((0, -1, 0, 1, 0), (0, -1, -1, 1, 1), (-1, 0, 0, 0, 1), (-1, -1, 0, 0, 1), (-1, -1, -1, 1, 1))$,
- 15) $((0, -1, 0, 1, 0), (0, -1, -1, 1, 1), (0, 0, -1, 1, 0), (-1, 0, -1, 1, 1), (-1, -1, -1, 1, 1))$,
- 16) $((0, -1, 0, 1, 0), (0, -1, -1, 1, 1), (0, 0, -1, 1, 0), (0, -1, -1, 1, 0), (-1, -1, -1, 1, 1))$,
- 17) $((0, -1, 0, 1, 0), (-1, 0, 0, 1, 0), (-1, 0, -1, 2, 1), (0, 0, -1, 1, 0), (0, 0, 0, 1, 0))$,
- 18) $((0, -1, 0, 1, 0), (-1, 0, 0, 1, 0), (-1, 0, -1, 2, 1), (-1, 0, 0, 1, 1), (0, 0, 0, 1, 0))$,
- 19) $((0, -1, 0, 1, 0), (-1, 0, 0, 1, 0), (-1, 0, -1, 2, 1), (-1, 0, 0, 1, 1), (-1, 0, 0, 0, 1))$,
- 20) $((0, -1, 0, 1, 0), (-1, 0, 0, 1, 0), (-1, 0, -1, 2, 1), (-1, 0, -1, 1, 1), (-1, 0, 0, 0, 1))$,
- 21) $((0, -1, 0, 1, 0), (-1, 0, 0, 1, 0), (-1, 0, -1, 2, 1), (-1, 0, -1, 1, 1), (0, 0, -1, 1, 0))$,
- 22) $((0, 0, -1, 0, 1), (0, -1, -1, 1, 1), (-1, 0, 0, 0, 1), (0, 0, -1, 1, 1), (0, 0, 0, 0, 1))$,
- 23) $((0, 0, -1, 0, 1), (0, -1, -1, 1, 1), (-1, 0, 0, 0, 1), (0, 0, -1, 1, 1), (-1, 0, -1, 1, 1))$,
- 24) $((0, 0, -1, 0, 1), (0, -1, -1, 1, 1), (-1, 0, 0, 0, 1), (0, -1, 0, 0, 1), (0, 0, 0, 0, 1))$,
- 25) $((0, 0, -1, 0, 1), (0, -1, -1, 1, 1), (-1, 0, 0, 0, 1), (0, -1, 0, 0, 1), (-1, -1, 0, 0, 1))$,
- 26) $((0, 0, -1, 0, 1), (0, -1, -1, 1, 1), (-1, 0, 0, 0, 1), (-1, 0, -1, 1, 1), (-1, -1, -1, 1, 1))$,
- 27) $((0, 0, -1, 0, 1), (0, -1, -1, 1, 1), (0, 0, -1, 1, 0), (-1, 0, -1, 1, 1), (-1, -1, -1, 1, 1))$,
- 28) $((0, 0, -1, 0, 1), (0, -1, -1, 1, 1), (0, 0, -1, 1, 0), (-1, 0, -1, 1, 1), (0, 0, -1, 1, 1))$,
- 29) $((0, 0, -1, 0, 1), (0, -1, -1, 1, 1), (0, 0, -1, 1, 0), (0, -1, -1, 1, 0), (-1, -1, -1, 1, 1))$,
- 30) $((0, 0, -1, 0, 1), (-1, -1, -1, 1, 1), (-1, 0, 0, 0, 1), (-1, 0, -1, 1, 1), (-1, 0, -1, 0, 1))$,
- 31) $((0, 0, -1, 0, 1), (-1, -1, -1, 1, 1), (-1, 0, 0, 0, 1), (-1, 0, -1, 0, 1), (-1, -1, 0, 0, 1))$,
- 32) $((0, 0, -1, 0, 1), (-1, -1, -1, 1, 1), (-1, 0, 0, 0, 1), (0, -1, -1, 1, 1), (-1, -1, 0, 0, 1))$,
- 33) $((0, 0, -1, 0, 1), (-1, -1, -1, 1, 1), (0, 0, -1, 1, 0), (-1, 0, -1, 0, 1), (0, -1, -1, 1, 0))$,
- 34) $((0, 0, -1, 0, 1), (-1, -1, -1, 1, 1), (0, 0, -1, 1, 0), (-1, 0, -1, 0, 1), (-1, 0, -1, 1, 1))$.

To verify the list, a useful fact from Proposition 2.9 is that an irreducible left mutation of a two-term silting complex T changes exactly one row of the g -matrix $\mathcal{G}(T)$. By comparing the rows of g -matrices, it is not difficult to find the finite connected component of $\mathcal{H}(\tau\text{-tilt}_\epsilon A)$. However, we leave the details to the readers, as it is hard to visualize the Hasse quiver on a plane. In conclusion, $2\text{-silt}_\epsilon A_\epsilon \simeq 2\text{-silt}_\epsilon A$ is a finite set.

We have proved that $2\text{-silt}_\epsilon A_\epsilon$ is finite for any $\epsilon \in \mathfrak{s}_5$. \square

Proposition 5.2. *Let $p = 3$. Then $S^+(2, 5)$ is τ -tilting finite.*

Proof. By Proposition 2.14, $S^+(2, 5)$ over $p = 3$ is isomorphic to $A := KQ/I$ with

$$Q : 5 \xrightarrow{\alpha_4} 4 \xrightarrow{\alpha_3} 3 \xrightarrow{\alpha_2} 2 \xrightarrow{\alpha_1} 1 \xrightarrow{\alpha_0} 0 \quad \text{and} \quad I : \left\langle \begin{array}{l} \alpha_4\alpha_3\alpha_2, \alpha_3\alpha_2\alpha_1, \alpha_2\alpha_1\alpha_0, \\ \alpha_4\beta_1 - \beta_2\alpha_1, \alpha_3\beta_0 - \beta_1\alpha_0 \end{array} \right\rangle.$$

Let $\epsilon = (\epsilon_0, \epsilon_1, \dots, \epsilon_5) \in \mathfrak{s}_6$. If $\epsilon_0 = +$ or $\epsilon_5 = -$, we have $A_\epsilon \simeq (K \oplus S^+(2, 4))_\epsilon$ by Proposition 3.8 and Proposition 3.9, and then A_ϵ is τ -tilting finite since $S^+(2, 4)$ over $p = 3$ is representation-finite.

Suppose $\epsilon_0 = -$ and $\epsilon_5 = +$. Since there is an algebra isomorphism $\sigma : A^{\text{op}} \rightarrow A$ sending e_i^* to e_{5-i} , it suffices to consider the following 10 cases by Corollary 3.16.

$$\begin{aligned} (-, +, +, -, +, +) &\sim_\sigma (-, -, +, -, -, +), & (-, -, -, -, -, +) &\sim_\sigma (-, +, +, +, +, +), \\ (-, -, +, -, +, +), & & (-, +, +, -, -, +), & & (-, +, +, +, -, +) &\sim_\sigma (-, +, -, -, -, +), \\ (-, +, -, +, -, +), & & (-, -, +, +, -, +) &\sim_\sigma (-, +, -, -, +, +), & & (-, -, -, +, +, +), \\ (-, -, -, -, +, +) &\sim_\sigma (-, -, +, +, +, +), & & & (-, -, -, +, -, +) &\sim_\sigma (-, +, -, +, +, +). \end{aligned}$$

Let P_i be the indecomposable projective A -module at vertex i . Then, $\text{Hom}_A(P_i, P_j)$ is given in the following table.

Hom	0	1	2	3	4	5
0	e_0	α_0	$\alpha_1\alpha_0$	β_0	$\alpha_3\beta_0$	$\alpha_4\alpha_3\beta_0$
1	0	e_1	α_1	$\alpha_2\alpha_1$	β_1	$\alpha_4\beta_1$
2	0	0	e_2	α_2	$\alpha_3\alpha_2$	β_2
3	0	0	0	e_3	α_3	$\alpha_4\alpha_3$
4	0	0	0	0	e_4	α_4
5	0	0	0	0	0	e_5

- If $\epsilon = (-, +, +, -, +, +)$, then $\alpha_3\alpha_2 \in J_{\epsilon,+}$, $\alpha_2, \alpha_2\alpha_1$ are removed and all others are survived; If $\epsilon = (-, -, -, -, -, +)$, then $\alpha_2, \alpha_3\alpha_2, \alpha_2\alpha_1 \in J_{\epsilon,-}$; If $\epsilon = (-, -, +, -, +, +)$, then $\alpha_3\alpha_2 \in J_{\epsilon,+}$, $\alpha_2\alpha_1 \in J_{\epsilon,-}$, α_2 is removed and all others are survived. In all these cases, A_ϵ is presented as

$$\begin{array}{ccccc} 5 & \longrightarrow & 4 & \longrightarrow & 3 \\ \downarrow & \searrow & \downarrow & \searrow & \downarrow \\ 2 & \longrightarrow & 1 & \longrightarrow & 0 \end{array},$$

which is a representation-finite sincere simply connected algebra.

- If $\epsilon = (-, +, +, -, -, +)$, then $\alpha_2, \beta_1, \alpha_3\alpha_2, \alpha_2\alpha_1$ are removed and all others are survived, and A_ϵ is presented as

$$\begin{array}{ccccc} 5 & \longrightarrow & 4 & \longrightarrow & 3 \\ \downarrow & \searrow & \downarrow & \searrow & \downarrow \\ 2 & \longrightarrow & 1 & \longrightarrow & 0 \end{array};$$

If $\epsilon = (-, +, +, +, -, +)$, then $\alpha_2, \alpha_2\alpha_1 \in J_{\epsilon,+}$, $\beta_1, \alpha_3, \alpha_3\alpha_2$ are removed and all others are survived, and A_ϵ is presented as

$$\begin{array}{ccccc} 5 & \longrightarrow & 4 & \longrightarrow & 3 \\ \downarrow & \searrow & \downarrow & \searrow & \downarrow \\ 2 & \longrightarrow & 1 & \longrightarrow & 0 \end{array}.$$

In both two cases, we have $A_\epsilon \simeq B_\epsilon$ and B is presented as

$$\begin{array}{ccccc} & & 4 & \longrightarrow & 3 \\ & \nearrow & & & \searrow \\ 5 & \longrightarrow & & & 0 \\ & \searrow & & & \nearrow \\ & & 2 & \longrightarrow & 1 \end{array},$$

which is a representation-finite sincere simply connected algebra.

- If $\epsilon = (-, +, -, +, -, +)$, then $\alpha_2\alpha_1 \in J_{\epsilon,+}$, $\alpha_3\alpha_2 \in J_{\epsilon,-}$, $\alpha_1, \alpha_3, \beta_1$ are removed and all others are survived. In this case, $A_\epsilon = KQ/I$ is given by

$$Q : \begin{array}{ccccc} & & 1 & & \\ & \nearrow & & \searrow & \\ & & 2 & & \\ & \nearrow & & \searrow & \\ 5 & \longrightarrow & & & 0 \\ & \searrow & & & \nearrow \\ & & 3 & & \\ & \nearrow & & \searrow & \\ & & 4 & & \end{array} \quad \text{and} \quad I : \langle y\alpha_0 - \beta_2w, \beta_2w - z\beta_0, z\beta_0 - \alpha_4x, z\alpha_2, \alpha_2w \rangle.$$

We replace two commutativity square subquivers, and it gives $B := KQ/I$ with

$$Q : \begin{array}{ccccc} & & 4 & \xrightarrow{s} & 3 \\ & \nearrow & & \searrow & \\ & & 2 & & \\ & \nearrow & & \searrow & \\ 5 & \longrightarrow & & & 0 \\ & \searrow & & & \nearrow \\ & & 2 & \xrightarrow{t} & 1 \end{array} \quad \text{and} \quad I : \langle \alpha_4s\beta_0 - \beta_2t\alpha_0, s\alpha_2, \alpha_2t \rangle.$$

It is easy to check that $A_\epsilon \simeq B_\epsilon$. Since B is a representation-finite special biserial algebra, we conclude that $2\text{-silt}_\epsilon A_\epsilon$ is a finite set.

- If $\epsilon = (-, -, +, +, -, +)$, then $\alpha_3, \alpha_3\alpha_2$ are removed and all others are survived. In this case, $A_\epsilon = KQ/I$ is presented by

$$Q : \begin{array}{ccccc} & & y & & \\ & & \curvearrowright & & \\ & 5 & \xrightarrow{\beta_2} & 2 & \xleftarrow{\alpha_2} & 3 \\ & \alpha_4 \downarrow & \dashrightarrow & \downarrow \alpha_1 & \downarrow \beta_0 & \\ & 4 & \xrightarrow{\beta_1} & 1 & \xrightarrow{\alpha_0} & 0 \end{array} \quad \text{and} \quad I : \left\langle \begin{array}{c} y\alpha_2, \alpha_2\alpha_1\alpha_0, \beta_2\alpha_1 - \alpha_4\beta_1, \\ y\beta_0 - \alpha_4\beta_1\alpha_0 \end{array} \right\rangle.$$

We replace the commutativity square subquiver consisting of 1, 2, 4, 5 with a linear subquiver, say, $B := KQ/I$ with

$$Q : \begin{array}{ccccccc} 5 & \xrightarrow{\alpha_4} & 4 & \xrightarrow{z} & 2 & \xrightarrow{\alpha_1} & 1 & \xrightarrow{\alpha_0} & 0 \\ & & \searrow y & & \alpha_2 \uparrow & & & & \\ & & & & 3 & & \xrightarrow{\beta_0} & & \end{array} \quad \text{and} \quad I : \left\langle \begin{array}{c} y\alpha_2, \alpha_2\alpha_1\alpha_0, \\ y\beta_0 - \alpha_4z\alpha_1\alpha_0 \end{array} \right\rangle.$$


Then, $A_\epsilon \simeq B_\epsilon$. We apply Proposition 3.1 and Proposition 3.4 to this case. Let R_i be the indecomposable projective B -module at vertex i . Then,

$$\mu_{R_0 \oplus R_1 \oplus R_4}^-(B) = \left[\begin{array}{c} R_0 \xrightarrow{(\alpha_1\alpha_0, \beta_0)^t} R_2 \oplus R_3 \\ \oplus \\ R_1 \xrightarrow{\alpha_1} R_2 \\ \oplus \\ R_4 \xrightarrow{\alpha_4} R_5 \\ \oplus \\ 0 \longrightarrow R_2 \oplus R_3 \oplus R_5 \end{array} \right] \in 2\text{-silt}_\epsilon B$$

and

$$\mathcal{F}(\mu_{R_0 \oplus R_1 \oplus R_4}^-(B)) \simeq \begin{array}{c} 2 \\ | \\ 1 \\ | \\ 0 \end{array} \begin{array}{c} 3 \\ / \quad \backslash \\ 0 \quad 2 \\ | \\ 1 \end{array} \oplus 2 \oplus \begin{array}{c} 5 \\ | \\ 3 \end{array} \oplus \begin{array}{c} 2 \\ | \\ 1 \\ | \\ 0 \end{array} \oplus \begin{array}{c} 3 \\ / \quad \backslash \\ 0 \quad 2 \\ | \\ 1 \end{array} \oplus \begin{array}{c} 5 \\ / \quad \backslash \\ 3 \quad 4 \\ | \quad | \\ 0 \quad 1 \end{array}.$$

By direct calculation of left mutations starting from $\mathcal{F}(\mu_{R_0 \oplus R_1 \oplus R_4}^-(B))$, we find that $\mathcal{H}(\tau\text{-tilt}_\epsilon B)$ admits a finite connected component:



where the g -matrix of each vertex is listed below:

- 1) $((0, 0, 1, 0, 0, 0), (0, 0, 0, 1, 0, 0), (0, 0, 0, 0, 0, 1), (-1, 0, 1, 1, 0, 0), (0, -1, 1, 0, 0, 0), (0, 0, 0, 0, -1, 1))$,
- 2) $((0, 0, 1, 0, 0, 0), (0, 0, 0, 0, 0, 1), (-1, 0, 1, 1, 0, 0), (-1, 0, 1, 0, 0, 0), (0, -1, 1, 0, 0, 0), (0, 0, 0, 0, -1, 1))$,
- 3) $((0, 0, 0, 1, 0, 0), (0, 0, 0, 0, 0, 1), (-1, 0, 1, 1, 0, 0), (-1, 0, 0, 1, 0, 0), (0, -1, 1, 0, 0, 0), (0, 0, 0, 0, -1, 1))$,
- 4) $((0, 0, 0, 1, 0, 0), (0, 0, 0, 0, 0, 1), (0, -1, 0, 1, 0, 1), (-1, 0, 0, 1, 0, 0), (0, -1, 1, 0, 0, 0), (0, 0, 0, 0, -1, 1))$,
- 5) $((0, 0, 0, 0, 0, 1), (-1, 0, 1, 1, 0, 0), (-1, 0, 1, 0, 0, 0), (-1, 0, 0, 1, 0, 0), (0, -1, 1, 0, 0, 0), (0, 0, 0, 0, -1, 1))$,
- 6) $((0, 0, 0, 0, 0, 1), (0, -1, 0, 1, 0, 1), (-1, 0, 0, 1, 0, 0), (-1, -1, 0, 1, 0, 1), (0, -1, 1, 0, 0, 0), (0, 0, 0, 0, -1, 1))$,
- 7) $((0, 0, 0, 1, 0, 0), (0, -1, 0, 1, 0, 1), (-1, 0, 0, 1, 0, 0), (0, -1, 0, 1, 0, 0), (0, -1, 1, 0, 0, 0), (0, 0, 0, 0, -1, 1))$,
- 8) $((0, 0, 0, 0, 0, 1), (-1, 0, 1, 0, 0, 0), (-1, 0, 0, 1, 0, 0), (-1, 0, 0, 0, 0, 1), (0, -1, 1, 0, 0, 0), (0, 0, 0, 0, -1, 1))$,
- 9) $((0, 0, 0, 0, 0, 1), (0, -1, 0, 1, 0, 1), (0, -1, 0, 0, 0, 1), (-1, -1, 0, 1, 0, 1), (0, -1, 1, 0, 0, 0), (0, 0, 0, 0, -1, 1))$,
- 10) $((0, -1, 0, 1, 0, 1), (-1, 0, 0, 1, 0, 0), (-1, -1, 0, 1, 0, 1), (0, -1, 0, 1, 0, 0), (0, -1, 1, 0, 0, 0), (0, 0, 0, 0, -1, 1))$,
- 11) $((0, 0, 0, 0, 0, 1), (-1, 0, 0, 1, 0, 0), (-1, -1, 0, 1, 0, 1), (-1, 0, 0, 0, 0, 1), (0, -1, 1, 0, 0, 0), (0, 0, 0, 0, -1, 1))$,
- 12) $((0, -1, 0, 1, 0, 1), (0, -1, 0, 0, 0, 1), (-1, -1, 0, 1, 0, 1), (0, -1, 0, 1, 0, 0), (0, -1, 1, 0, 0, 0), (0, 0, 0, 0, -1, 1))$,

- 13) $((0, 0, 0, 0, 0, 1), (0, -1, 0, 0, 0, 1), (-1, -1, 0, 1, 0, 1), (-1, 0, 0, 0, 0, 1), (0, -1, 1, 0, 0, 0), (0, 0, 0, 0, -1, 1)),$
 14) $((-1, 0, 0, 1, 0, 0), (-1, -1, 0, 1, 0, 1), (0, -1, 0, 1, 0, 0), (-1, -1, 0, 1, 0, 0), (0, -1, 1, 0, 0, 0), (0, 0, 0, 0, -1, 1)),$
 15) $((0, -1, 0, 0, 0, 1), (-1, -1, 0, 1, 0, 1), (0, -1, 0, 1, 0, 0), (-1, -1, 0, 1, 0, 0), (0, -1, 1, 0, 0, 0), (0, 0, 0, 0, -1, 1)).$

If the reader wishes to verify the above quiver, Proposition 2.9 serves as a useful tool, as mentioned in the proof of Proposition 5.1. Another helpful observation is that the second and third direct summands of $\mathcal{F}(\mu_{R_0 \oplus R_1 \oplus R_4}^-(B))$ are immutable in $\tau\text{-tilt}_\epsilon B$. Consequently, we conclude that $2\text{-silt}_\epsilon A_\epsilon \simeq 2\text{-silt}_\epsilon B_\epsilon$ is a finite set.

- If $\epsilon = (-, -, -, +, +, +)$, then $A_\epsilon \simeq A$. In this case, we have

$$\mu_{P_0 \oplus P_1 \oplus P_2}^-(A) = \left[\begin{array}{c} P_0 \xrightarrow{\beta_0} P_3 \\ \oplus \\ P_1 \xrightarrow{(\alpha_2 \alpha_1, \beta_1)^t} P_3 \oplus P_4 \\ \oplus \\ P_2 \xrightarrow{(\alpha_2, \beta_2)^t} P_3 \oplus P_5 \\ \oplus \\ 0 \longrightarrow P_3 \oplus P_4 \oplus P_5 \end{array} \right] \in 2\text{-silt}_\epsilon A$$

and

$$\mathcal{F}(\mu_{P_0 \oplus P_1 \oplus P_2}^-(A)) \simeq \begin{array}{c} 3 \\ \vdots \\ 2 \\ \vdots \\ 1 \end{array} \oplus \begin{array}{c} 3 \\ \swarrow 1 \quad \searrow 2 \\ 1 \quad 4 \\ \swarrow 1 \quad \searrow 3 \\ 2 \end{array} \oplus \begin{array}{c} 3 \\ \swarrow 1 \quad \searrow 2 \\ 0 \quad 5 \\ \swarrow 1 \quad \searrow 4 \\ 1 \quad 3 \end{array} \oplus \begin{array}{c} 3 \\ \swarrow 1 \quad \searrow 2 \\ 0 \quad 2 \\ \swarrow 1 \quad \searrow 1 \end{array} \oplus \begin{array}{c} 4 \\ \swarrow 1 \quad \searrow 3 \\ 1 \quad 3 \\ \swarrow 0 \quad \searrow 2 \end{array} \oplus \begin{array}{c} 5 \\ \swarrow 2 \quad \searrow 4 \\ 1 \quad 4 \\ \swarrow 1 \quad \searrow 3 \\ 0 \end{array}.$$

Similar to the previous case, we find $\#\tau\text{-tilt}_\epsilon A = 157$ using Proposition 3.1 and Proposition 3.4. Here is nothing new but only more calculations of left mutations, we thus omit the details. It turns out that $2\text{-silt}_\epsilon A_\epsilon$ is a finite set.


- Set $\epsilon = (-, -, -, -, +, +)$. We denote $P_I = P_0 \oplus P_1 \oplus P_2 \oplus P_3$. We have

$$\mu_{P_I}^-(A) = \left[\begin{array}{c} P_0 \xrightarrow{\alpha_3 \beta_0} P_4 \\ \oplus \\ P_1 \xrightarrow{\beta_1} P_4 \\ \oplus \\ P_3 \xrightarrow{\alpha_3} P_4 \\ \oplus \\ P_2 \xrightarrow{(\alpha_3 \alpha_2, \beta_2)^t} P_4 \oplus P_5 \\ \oplus \\ 0 \longrightarrow P_4 \oplus P_5 \end{array} \right] \in 2\text{-silt}_\epsilon A$$

and

$$\mathcal{F}(\mu_{P_I}^-(A)) \simeq \begin{array}{c} 4 \\ \swarrow 1 \quad \searrow 3 \\ 1 \quad 3 \\ \swarrow 1 \quad \searrow 2 \\ 2 \end{array} \oplus \begin{array}{c} 4 \\ \swarrow 1 \quad \searrow 2 \\ 4 \\ \swarrow 1 \quad \searrow 1 \end{array} \oplus \begin{array}{c} 4 \\ \swarrow 1 \quad \searrow 2 \\ 4 \\ \swarrow 1 \quad \searrow 1 \end{array} \oplus \begin{array}{c} 4 \\ \swarrow 1 \quad \searrow 3 \\ 1 \quad 3 \\ \swarrow 0 \quad \searrow 2 \\ 2 \quad 5 \\ \swarrow 1 \quad \searrow 4 \\ 4 \quad 3 \end{array} \oplus \begin{array}{c} 4 \\ \swarrow 1 \quad \searrow 3 \\ 1 \quad 3 \\ \swarrow 0 \quad \searrow 2 \end{array} \oplus \begin{array}{c} 5 \\ \swarrow 2 \quad \searrow 4 \\ 1 \quad 4 \\ \swarrow 1 \quad \searrow 3 \\ 0 \end{array}.$$

The Hasse quiver $\mathcal{H}(\tau\text{-tilt}_\epsilon A)$ has the following connected component:



where the g -matrix of each vertex is given below:

- 1) $((0, 0, 0, 0, 1, 0), (0, 0, 0, 0, 0, 1), (0, 0, 0, -1, 1, 0), (0, 0, -1, 0, 1, 1), (0, -1, 0, 0, 1, 0), (-1, 0, 0, 0, 1, 0)),$
- 2) $((0, 0, 0, 0, 1, 0), (0, 0, 0, -1, 1, 0), (0, 0, -1, 0, 1, 1), (0, 0, -1, 0, 1, 0), (0, -1, 0, 0, 1, 0), (-1, 0, 0, 0, 1, 0)),$
- 3) $((0, 0, 0, 0, 0, 1), (0, 0, 0, -1, 1, 0), (0, 0, -1, 0, 1, 1), (0, -1, 0, 0, 1, 0), (-1, 0, 0, 0, 1, 0), (-1, 0, -1, 0, 1, 1)),$
- 4) $((0, 0, 0, 0, 0, 1), (0, 0, 0, -1, 1, 0), (0, 0, -1, 0, 1, 1), (0, -1, 0, 0, 1, 0), (0, -1, -1, 0, 1, 1), (-1, 0, -1, 0, 1, 1)),$
- 5) $((0, 0, 0, -1, 1, 0), (0, 0, -1, 0, 1, 1), (0, 0, -1, 0, 1, 0), (0, -1, 0, 0, 1, 0), (-1, 0, 0, 0, 1, 0), (-1, 0, -1, 0, 1, 1)),$
- 6) $((0, 0, 0, 0, 0, 1), (0, 0, 0, -1, 1, 0), (0, -1, 0, 0, 1, 0), (-1, 0, 0, 0, 1, 0), (-1, 0, -1, 0, 1, 1), (-1, 0, 0, 0, 0, 1)),$
- 7) $((0, 0, 0, -1, 1, 0), (0, 0, -1, 0, 1, 0), (0, -1, 0, 0, 1, 0), (-1, 0, 0, 0, 1, 0), (-1, 0, -1, 0, 1, 1), (-1, 0, -1, 0, 1, 0)),$
- 8) $((0, 0, 0, 0, 0, 1), (0, 0, 0, -1, 1, 0), (0, 0, -1, 0, 1, 1), (0, 0, -1, 0, 0, 1), (0, -1, -1, 0, 1, 1), (-1, 0, -1, 0, 1, 1)),$
- 9) $((0, 0, 0, -1, 1, 0), (0, 0, -1, 0, 1, 1), (0, 0, -1, 0, 1, 0), (0, -1, 0, 0, 1, 0), (0, -1, -1, 0, 1, 1), (-1, 0, -1, 0, 1, 1)),$
- 10) $((0, 0, 0, 0, 0, 1), (0, 0, 0, -1, 1, 0), (0, -1, 0, 0, 1, 0), (0, -1, -1, 0, 1, 1), (-1, 0, -1, 0, 1, 1), (-1, 0, 0, 0, 0, 1)),$
- 11) $((0, 0, 0, 0, 0, 1), (0, 0, 0, -1, 1, 0), (0, 0, -1, 0, 0, 1), (0, -1, -1, 0, 1, 1), (-1, 0, -1, 0, 1, 1), (-1, 0, 0, 0, 0, 1)),$
- 12) $((0, 0, 0, -1, 1, 0), (0, 0, -1, 0, 1, 1), (0, 0, -1, 0, 0, 1), (0, 0, -1, 0, 1, 0), (0, -1, -1, 0, 1, 1), (-1, 0, -1, 0, 1, 1)),$
- 13) $((0, 0, 0, -1, 1, 0), (0, 0, -1, 0, 1, 0), (0, -1, 0, 0, 1, 0), (0, -1, -1, 0, 1, 1), (-1, 0, -1, 0, 1, 1), (-1, 0, -1, 0, 1, 0)),$
- 14) $((0, 0, 0, 0, 0, 1), (0, 0, 0, -1, 1, 0), (0, -1, 0, 0, 1, 0), (0, -1, -1, 0, 1, 1), (0, -1, 0, -1, 1, 1), (-1, 0, 0, 0, 0, 1)),$
- 15) $((0, 0, 0, 0, 0, 1), (0, -1, 0, 0, 1, 0), (0, -1, -1, 0, 1, 1), (0, -1, 0, 0, 0, 1), (0, -1, 0, -1, 1, 1), (-1, 0, 0, 0, 0, 1)),$
- 16) $((0, 0, 0, -1, 1, 0), (0, 0, -1, 0, 0, 1), (0, 0, -1, 0, 1, 0), (0, -1, -1, 0, 1, 1), (-1, 0, -1, 0, 1, 1), (-1, 0, -1, 0, 1, 0)),$
- 17) $((0, 0, 0, -1, 1, 0), (0, 0, -1, 0, 1, 0), (0, -1, 0, 0, 1, 0), (0, -1, -1, 0, 1, 1), (0, -1, -1, 0, 1, 0), (-1, 0, -1, 0, 1, 0)),$
- 18) $((0, 0, 0, 0, 0, 1), (0, 0, 0, -1, 1, 0), (0, 0, -1, 0, 0, 1), (0, -1, -1, 0, 1, 1), (0, -1, 0, -1, 1, 1), (-1, 0, 0, 0, 0, 1)),$
- 19) $((0, 0, 0, 0, 0, 1), (0, 0, -1, 0, 0, 1), (0, -1, -1, 0, 1, 1), (0, -1, 0, 0, 0, 1), (0, -1, 0, -1, 1, 1), (-1, 0, 0, 0, 0, 1)),$
- 20) $((0, 0, 0, -1, 1, 0), (0, 0, -1, 0, 0, 1), (0, 0, -1, 0, 1, 0), (0, -1, -1, 0, 1, 1), (0, -1, -1, 0, 1, 0), (-1, 0, -1, 0, 1, 0)),$
- 21) $((0, 0, 0, 0, 0, 1), (0, 0, 0, -1, 1, 0), (0, 0, 0, -1, 0, 1), (0, 0, -1, 0, 0, 1), (0, -1, 0, -1, 1, 1), (-1, 0, 0, 0, 0, 1)),$
- 22) $((0, 0, 0, 0, 0, 1), (0, 0, 0, -1, 0, 1), (0, 0, -1, 0, 0, 1), (0, -1, 0, -1, 1, 1), (-1, 0, 0, 0, 0, 1), (0, -1, 0, 0, 0, 1)).$

We obtain $\#\tau\text{-tilt}_e A = 22$ and thus, $2\text{-silt}_e A$ is finite by Proposition 3.4.

- Set $\epsilon = (-, -, -, +, -, +)$. We have $P_I = P_0 \oplus P_1 \oplus P_2 \oplus P_4$. Then,

$$\mu_{P_I}^-(A) = \left[\begin{array}{c} P_0 \xrightarrow{\beta_0} P_3 \\ \oplus \\ P_4 \xrightarrow{\alpha_4} P_5 \\ \oplus \\ P_1 \xrightarrow{(\alpha_2 \alpha_1, \alpha_4 \beta_1)^t} P_3 \oplus P_5 \\ \oplus \\ P_2 \xrightarrow{(\alpha_2, \beta_2)^t} P_3 \oplus P_5 \\ \oplus \\ 0 \longrightarrow P_3 \oplus P_5 \end{array} \right] \in 2\text{-silt}_\epsilon A$$

and

$$\mathcal{F}(\mu_{P_I}^-(A)) \simeq \begin{array}{c} 3 \\ \vdots \\ 2 \\ \vdots \\ 1 \end{array} \oplus \begin{array}{c} 5 \\ \vdots \\ 2 \end{array} \oplus \begin{array}{c} 3 \\ \vdots \\ 0 \end{array} \begin{array}{c} 5 \\ \vdots \\ 2 \\ \vdots \\ 1 \end{array} \oplus \begin{array}{c} 3 \\ \vdots \\ 0 \end{array} \begin{array}{c} 5 \\ \vdots \\ 2 \\ \vdots \\ 1 \end{array} \oplus \begin{array}{c} 3 \\ \vdots \\ 0 \end{array} \begin{array}{c} 5 \\ \vdots \\ 2 \\ \vdots \\ 1 \end{array} \oplus \begin{array}{c} 5 \\ \vdots \\ 2 \\ \vdots \\ 1 \end{array} \begin{array}{c} 3 \\ \vdots \\ 0 \end{array}.$$

We find $\#\tau\text{-tilt}_\epsilon A = 60$ by similar calculations in the previous case.

We have proved that $2\text{-silt}_\epsilon A \simeq 2\text{-silt}_\epsilon A_\epsilon$ is finite for any $\epsilon \in \mathfrak{s}_6$. \square

Proposition 5.3. *Let $p = 5$. Then $S^+(2, 6)$ is τ -tilting finite.*

Proof. By Proposition 2.14, $S^+(2, 6)$ over $p = 5$ is isomorphic to $A := KQ/I$ with

$$Q : 6 \xrightarrow{\alpha_5} 5 \xrightarrow{\alpha_4} 4 \xrightarrow{\alpha_3} 3 \xrightarrow{\alpha_2} 2 \xrightarrow{\alpha_1} 1 \xrightarrow{\alpha_0} 0$$

and

$$I : \langle \alpha_5 \alpha_4 \alpha_3 \alpha_2 \alpha_1, \alpha_4 \alpha_3 \alpha_2 \alpha_1 \alpha_0, \alpha_5 \beta_0 - \beta_1 \alpha_0 \rangle.$$

Let $\epsilon = (\epsilon_0, \epsilon_1, \dots, \epsilon_6) \in \mathfrak{s}_7$. If $\epsilon_0 = +$ or $\epsilon_6 = -$, we have $A_\epsilon \simeq (K \oplus S^+(2, 5))_\epsilon$ by Proposition 3.8 and Proposition 3.9, and then A_ϵ is τ -tilting finite since $S^+(2, 5)$ over $p = 5$ is representation-finite.

Suppose $\epsilon_0 = -$ and $\epsilon_6 = +$. Since there is an algebra isomorphism $\sigma : A^{\text{op}} \rightarrow A$ sending e_i^* to e_{6-i} , it suffices to consider the following 16 cases by Corollary 3.16.

$$\begin{aligned} (-, +, +, +, +, +, +) &\sim_\sigma (-, -, -, -, -, -, +), & (-, +, +, +, +, -, +) &\sim_\sigma (-, +, -, -, -, -, +), \\ (-, -, +, +, +, -, +) &\sim_\sigma (-, +, -, -, -, +, +), & (-, +, +, +, -, +, +) &\sim_\sigma (-, -, +, -, -, -, +), \\ (-, +, +, +, -, -, +) &\sim_\sigma (-, +, +, -, -, -, +), & (-, +, +, -, -, +, +) &\sim_\sigma (-, -, +, +, -, -, +), \\ (-, +, +, -, +, -, +) &\sim_\sigma (-, +, -, +, -, -, +), & (-, +, -, +, +, -, +) &\sim_\sigma (-, +, -, -, +, -, +), \\ (-, -, -, -, +, -, +) &\sim_\sigma (-, +, -, +, +, +, +), & (-, -, -, +, -, -, +) &\sim_\sigma (-, +, +, -, +, +, +), \\ (-, -, -, +, +, -, +) &\sim_\sigma (-, +, -, -, +, +, +), & (-, -, +, -, +, -, +) &\sim_\sigma (-, +, -, +, -, +, +), \\ (-, -, -, -, +, +, +) &\sim_\sigma (-, -, -, +, +, +, +), & (-, -, -, -, -, +, +) &\sim_\sigma (-, -, +, +, +, +, +), \\ (-, -, -, +, -, +, +) &\sim_\sigma (-, -, +, -, +, +, +), & (-, -, +, -, -, +, +) &\sim_\sigma (-, -, +, +, -, +, +). \end{aligned}$$

Let P_i be the indecomposable projective A -module at vertex i . Then, $\text{Hom}_A(P_i, P_j)$ is given in the following table.

Hom	0	1	2	3	4	5	6
0	e_0	α_0	$\alpha_1 \alpha_0$	$\alpha_2 \alpha_1 \alpha_0$	$\alpha_3 \alpha_2 \alpha_1 \alpha_0$	β_0	$\alpha_5 \beta_0$
1	0	e_1	α_1	$\alpha_2 \alpha_1$	$\alpha_3 \alpha_2 \alpha_1$	$\alpha_4 \alpha_3 \alpha_2 \alpha_1$	β_1
2	0	0	e_2	α_2	$\alpha_3 \alpha_2$	$\alpha_4 \alpha_3 \alpha_2$	$\alpha_5 \alpha_4 \alpha_3 \alpha_2$
3	0	0	0	e_3	α_3	$\alpha_4 \alpha_3$	$\alpha_5 \alpha_4 \alpha_3$
4	0	0	0	0	e_4	α_4	$\alpha_5 \alpha_4$
5	0	0	0	0	0	e_5	α_5
6	0	0	0	0	0	0	e_6

- If $\epsilon = (-, +, +, +, +, +, +)$, then we have $\alpha_4, \alpha_4 \alpha_3, \alpha_4 \alpha_3 \alpha_2, \alpha_4 \alpha_3 \alpha_2 \alpha_1, \alpha_5 \alpha_4, \alpha_5 \alpha_4 \alpha_3, \alpha_5 \alpha_4 \alpha_3 \alpha_2 \in J_{\epsilon, +}$, and all others are survived; If $\epsilon = (-, +, +, +, +, -, +)$, then $\alpha_5 \alpha_4, \alpha_5 \alpha_4 \alpha_3, \alpha_5 \alpha_4 \alpha_3 \alpha_2 \in J_{\epsilon, +}$, $\alpha_4, \alpha_4 \alpha_3, \alpha_4 \alpha_3 \alpha_2, \alpha_4 \alpha_3 \alpha_2 \alpha_1$ are removed

and all others are survived; If $\epsilon = (-, -, +, +, +, -, +)$, then $\alpha_5\alpha_4$, $\alpha_5\alpha_4\alpha_3$, $\alpha_5\alpha_4\alpha_3\alpha_2 \in J_{\epsilon,+}$, $\alpha_4\alpha_3\alpha_2\alpha_1 \in J_{\epsilon,-}$, α_4 , $\alpha_4\alpha_3$, $\alpha_4\alpha_3\alpha_2$ are removed and all others are survived. In all these cases, A_ϵ is presented as

$$\begin{array}{ccccccc} & & 5 & \longleftarrow & 6 & & \\ & & \downarrow & \nearrow & \downarrow & & \\ & & 0 & \longleftarrow & 1 & \longleftarrow & 2 \longleftarrow 3 \longleftarrow 4 \end{array},$$

which is a representation-finite sincere simply connected algebra.

- If $\epsilon = (-, +, +, +, -, +, +)$, then α_4 , $\alpha_4\alpha_3$, $\alpha_4\alpha_3\alpha_2$, $\alpha_4\alpha_3\alpha_2\alpha_1$, $\alpha_5\alpha_4\alpha_3$, $\alpha_5\alpha_4\alpha_3\alpha_2 \in J_{\epsilon,+}$, $\alpha_3\alpha_2\alpha_1\alpha_0 \in J_{\epsilon,-}$, α_3 , $\alpha_3\alpha_2$, $\alpha_3\alpha_2\alpha_1$ are removed and all others are survived; If $\epsilon = (-, +, +, +, -, -, +)$, then $\alpha_5\alpha_4\alpha_3$, $\alpha_5\alpha_4\alpha_3\alpha_2 \in J_{\epsilon,+}$, $\alpha_3\alpha_2\alpha_1\alpha_0 \in J_{\epsilon,-}$, α_3 , $\alpha_3\alpha_2$, $\alpha_3\alpha_2\alpha_1$, $\alpha_4\alpha_3$, $\alpha_4\alpha_3\alpha_2$, $\alpha_4\alpha_3\alpha_2\alpha_1$ are removed and all others are survived. In both cases, A_ϵ is presented as

$$\begin{array}{ccccccc} & & & & 6 & \longrightarrow & 5 \longrightarrow 4 \\ & & & & \downarrow & \nearrow & \downarrow \\ & & & & 1 & \longrightarrow & 0 \\ 3 & \longrightarrow & 2 & \longrightarrow & 1 & \longrightarrow & 0 \end{array},$$

which is a representation-finite sincere simply connected algebra.

- If $\epsilon = (-, +, +, -, -, +, +)$, then $\alpha_4\alpha_3\alpha_2$, $\alpha_4\alpha_3\alpha_2\alpha_1$, $\alpha_5\alpha_4\alpha_3\alpha_2 \in J_{\epsilon,+}$, $\alpha_2\alpha_1\alpha_0$, $\alpha_3\alpha_2\alpha_1\alpha_0 \in J_{\epsilon,-}$, α_2 , $\alpha_2\alpha_1$, $\alpha_3\alpha_2$, $\alpha_3\alpha_2\alpha_1$ are removed and all others are survived. In this case, A_ϵ is presented as


$$\begin{array}{ccccccc} & & & & 6 & \longrightarrow & 5 \longrightarrow 4 \longrightarrow 3 \\ & & & & \downarrow & \nearrow & \downarrow \\ & & & & 1 & \longrightarrow & 0 \\ 2 & \longrightarrow & 1 & \longrightarrow & 0 & & \end{array},$$

which is again representation-finite.

- If $\epsilon = (-, +, +, -, +, -, +)$, then $\alpha_5\alpha_4\alpha_3\alpha_2 \in J_{\epsilon,+}$, α_2 , α_4 , $\alpha_2\alpha_1$, $\alpha_4\alpha_3\alpha_2$, $\alpha_4\alpha_3\alpha_2\alpha_1$ are removed and all others are survived. One may check that $A_\epsilon \simeq B_\epsilon$ and $B := KQ/I$ is given by

$$Q : \begin{array}{ccccccc} 6 & \xrightarrow{\alpha_5} & 5 & \xrightarrow{x} & 1 & \xrightarrow{\alpha_0} & 0 \\ & & \downarrow \alpha_4 & & \uparrow \alpha_1 & & \\ & & 4 & \xrightarrow{\alpha_3} & 3 & \xrightarrow{\alpha_2} & 2 \end{array} \quad \text{and} \quad I : \langle \alpha_4\alpha_3\alpha_2 \rangle.$$

By using Proposition 3.1 and Proposition 3.4, we find that $\mathcal{H}(\tau\text{-tilt}_\epsilon B)$ has the following connected component:



where the g -matrices are given as follows:


- 1) $((0, 1, 0, 0, 0, 0, 0), (0, 0, 1, 0, 0, 0, 0), (0, 0, 0, 0, 1, 0, 0), (0, 0, 0, 0, 0, 0, 1), (0, 0, 0, -1, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0))$,
- 2) $((0, 0, 1, 0, 0, 0, 0), (0, 0, 0, 0, 1, 0, 0), (0, 0, 0, 0, 0, 0, 1), (0, 0, 0, -1, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 1, 0, 0, 0, 1))$,
- 3) $((0, 1, 0, 0, 0, 0, 0), (0, 0, 1, 0, 0, 0, 0), (0, 0, 0, 0, 0, 0, 1), (0, 0, 0, -1, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (0, 0, 0, -1, 0, 0, 1))$,
- 4) $((0, 0, 1, 0, 0, 0, 0), (0, 0, 0, 0, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 1, 0, 0, 0, 1), (-1, 0, 1, 0, 0, 0, 0), (0, 0, 0, -1, 1, 0, 0))$,
- 5) $((0, 0, 0, 0, 1, 0, 0), (0, 0, 0, 0, 0, 0, 1), (0, 0, 0, -1, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 1, 0, 0, 0, 1), (-1, 0, 0, 0, 1, 0, 1))$,
- 6) $((0, 0, 1, 0, 0, 0, 0), (0, 0, 0, 0, 0, 0, 1), (0, 0, 0, -1, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 1, 0, 0, 0, 1), (0, 0, 0, -1, 0, 0, 1))$,
- 7) $((0, 0, 0, 0, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 1, 0, 0, 0, 1), (-1, 0, 1, 0, 0, 0, 0), (0, 0, 0, -1, 1, 0, 0), (-1, 0, 0, 0, 1, 0, 1))$,
- 8) $((0, 0, 1, 0, 0, 0, 0), (0, 0, 0, -1, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 1, 0, 0, 0, 1), (0, 0, 0, -1, 0, 0, 1), (-1, 0, 1, -1, 0, 0, 1))$,
- 9) $((0, 0, 0, 0, 0, 0, 1), (0, 0, 0, -1, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 1, 0, 0, 0, 1), (-1, 0, 0, 0, 1, 0, 1), (-1, 0, 0, 0, 0, 0, 1))$,
- 10) $((0, 0, 0, 0, 0, 0, 1), (0, 0, 0, -1, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 1, 0, 0, 0, 1), (0, 0, 0, -1, 0, 0, 1), (-1, 0, 0, 0, 0, 0, 1))$,
- 11) $((0, 0, 1, 0, 0, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 1, 0, 0, 0, 1), (-1, 0, 1, 0, 0, 0, 0), (0, 0, 0, -1, 1, 0, 0), (-1, 0, 1, -1, 0, 0, 1))$,
- 12) $((0, 0, 0, 0, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 1, 0, 0, 0, 0), (0, 0, 0, -1, 1, 0, 0), (-1, 0, 0, 0, 1, 0, 1), (-1, 0, 0, 0, 1, 0, 0))$,
- 13) $((0, 0, 0, -1, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 1, 0, 0, 0, 1), (-1, 0, 0, 0, 1, 0, 1), (-1, 0, 0, 0, 0, 0, 1), (-1, 0, 1, 0, 0, 0, 0))$,
- 14) $((0, 0, 0, -1, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 1, 0, 0, 0, 1), (0, 0, 0, -1, 0, 0, 1), (-1, 0, 0, 0, 0, 0, 1), (-1, 0, 1, -1, 0, 0, 1))$,
- 15) $((0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 1, 0, 0, 0, 1), (-1, 0, 1, 0, 0, 0, 0), (0, 0, 0, -1, 1, 0, 0), (-1, 0, 1, -1, 0, 0, 1), (-1, 0, 0, 0, 0, 0, 1))$,
- 16) $((0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 1, 0, 0, 0, 0), (0, 0, 0, -1, 1, 0, 0), (-1, 0, 0, 0, 1, 0, 1), (-1, 0, 0, 0, 1, 0, 0), (-1, 0, 0, 0, 0, 0, 1))$,
- 17) $((0, 0, 0, -1, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (0, 0, 0, -1, 0, 0, 1), (-1, 0, 0, 0, 0, 0, 1), (-1, 0, 1, -1, 0, 0, 1), (-1, 0, 0, -1, 0, 0, 1))$,
- 18) $((0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 1, 0, 0, 0, 0), (0, 0, 0, -1, 1, 0, 0), (-1, 0, 1, -1, 0, 0, 1), (-1, 0, 0, 0, 0, 0, 1), (-1, 0, 0, -1, 0, 0, 1))$.

This implies that $\#\tau\text{-tilt}_\epsilon B = 18$. Hence, $2\text{-silt}_\epsilon A_\epsilon \simeq 2\text{-silt}_\epsilon B_\epsilon$ is finite.

- If $\epsilon = (-, +, -, +, +, -, +)$, then $\alpha_1, \alpha_4, \alpha_4\alpha_3, \alpha_4\alpha_3\alpha_2\alpha_1$ are removed and all others are survived. We find that $A_\epsilon \simeq B_\epsilon$ and $B := KQ/I$ is given by

$$Q : \begin{array}{ccccccc} 6 & \xrightarrow{\alpha_5} & 5 & \xrightarrow{x} & 1 & \xrightarrow{\alpha_0} & 0 \\ & & \downarrow \alpha_4 & & \uparrow \alpha_1 & & \\ & & 4 & \xrightarrow{\alpha_3} & 3 & \xrightarrow{\alpha_2} & 2 \end{array} \quad \text{and} \quad I : \langle \alpha_4\alpha_3\alpha_2\alpha_1 \rangle.$$

We then find that $\mathcal{H}(\tau\text{-tilt}_\epsilon B)$ has the following structure:




where the g -matrices are given as follows:

- 1) $((0, 1, 0, 0, 0, 0, 0), (0, 0, 0, 1, 0, 0, 0), (0, 0, 0, 0, 1, 0, 0), (0, 0, 0, 0, 0, 0, 1), (0, 0, -1, 1, 0, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0))$,
- 2) $((0, 1, 0, 0, 0, 0, 0), (0, 0, 0, 0, 1, 0, 0), (0, 0, 0, 0, 0, 0, 1), (0, 0, -1, 1, 0, 0, 0), (0, 0, -1, 0, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0))$,
- 3) $((0, 0, 0, 1, 0, 0, 0), (0, 0, 0, 0, 1, 0, 0), (0, 0, 0, 0, 0, 0, 1), (0, 0, -1, 1, 0, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 0, 1, 0, 0, 1))$,
- 4) $((0, 0, 0, 0, 1, 0, 0), (0, 0, 0, 0, 0, 0, 1), (0, 0, -1, 1, 0, 0, 0), (0, 0, -1, 0, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 0, 0, 1, 0, 1))$,
- 5) $((0, 1, 0, 0, 0, 0, 0), (0, 0, 0, 0, 0, 0, 1), (0, 0, -1, 1, 0, 0, 0), (0, 0, -1, 0, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (0, 0, -1, 0, 0, 0, 1))$,
- 6) $((0, 0, 0, 0, 1, 0, 0), (0, 0, 0, 0, 0, 0, 1), (0, 0, -1, 1, 0, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 0, 1, 0, 0, 1), (-1, 0, 0, 0, 1, 0, 1))$,
- 7) $((0, 0, 0, 1, 0, 0, 0), (0, 0, 0, 0, 1, 0, 0), (0, 0, -1, 1, 0, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 0, 1, 0, 0, 1), (-1, 0, 0, 1, 0, 0, 0))$,
- 8) $((0, 0, 0, 0, 0, 0, 1), (0, 0, -1, 1, 0, 0, 0), (0, 0, -1, 0, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 0, 0, 1, 0, 1), (-1, 0, 0, 0, 0, 0, 1))$,
- 9) $((0, 0, 0, 0, 1, 0, 0), (0, 0, -1, 1, 0, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 0, 1, 0, 0, 1), (-1, 0, 0, 0, 1, 0, 1), (-1, 0, 0, 1, 0, 0, 0))$,
- 10) $((0, 0, 0, 0, 0, 0, 1), (0, 0, -1, 1, 0, 0, 0), (0, 0, -1, 0, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (0, 0, -1, 0, 0, 0, 1), (-1, 0, 0, 0, 0, 0, 1))$,
- 11) $((0, 0, 0, 0, 0, 0, 1), (0, 0, -1, 1, 0, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 0, 1, 0, 0, 1), (-1, 0, 0, 0, 1, 0, 1), (-1, 0, 0, 0, 0, 0, 1))$,
- 12) $((0, 0, 0, 0, 1, 0, 0), (0, 0, -1, 1, 0, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 0, 1, 0, 0, 1), (-1, 0, 0, 1, 0, 0, 0), (-1, 0, 0, 0, 1, 0, 0))$,
- 13) $((0, 0, -1, 1, 0, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 0, 1, 0, 0, 1), (-1, 0, 0, 0, 1, 0, 1), (-1, 0, 0, 0, 0, 0, 1), (-1, 0, 0, 1, 0, 0, 0))$,
- 14) $((0, 0, -1, 1, 0, 0, 0), (0, 0, -1, 0, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (0, 0, -1, 0, 0, 0, 1), (-1, 0, 0, 0, 0, 0, 1), (-1, 0, -1, 0, 0, 0, 1))$,
- 15) $((0, 0, -1, 1, 0, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 0, 0, 1, 0, 1), (-1, 0, 0, 0, 0, 0, 1), (-1, 0, 0, 1, 0, 0, 0), (-1, 0, 0, 0, 1, 0, 0))$,
- 16) $((0, 0, 0, 0, 1, 0, 0), (0, 0, -1, 1, 0, 0, 0), (0, 0, -1, 0, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 0, 0, 1, 0, 1), (-1, 0, 0, 0, 1, 0, 0))$,
- 17) $((0, 0, -1, 1, 0, 0, 0), (0, 0, -1, 0, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (-1, 1, 0, 0, 0, 0, 0), (-1, 0, 0, 0, 1, 0, 1), (-1, 0, 0, 0, 0, 0, 1), (-1, 0, 0, 0, 1, 0, 0))$.

Then, $\#\tau\text{-tilt}_\epsilon B = 17$ and $2\text{-silt}_\epsilon A_\epsilon \simeq 2\text{-silt}_\epsilon B_\epsilon$ is a finite set by Proposition 3.4.

- If $\epsilon = (-, -, -, -, +, -, +)$, we find that $\mathcal{H}(\tau\text{-tilt}_\epsilon A)$ has the following structure:



where the g -matrices are given as follows:

- 1) $((0, 0, 0, 0, 1, 0, 0), (0, 0, 0, 0, 0, 0, 1), (0, 0, 0, -1, 1, 0, 0), (0, 0, -1, 0, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (0, -1, 0, 0, 1, 0, 1), (-1, 0, 0, 0, 1, 0, 1))$,
- 2) $((0, 0, 0, 0, 1, 0, 0), (0, 0, 0, -1, 1, 0, 0), (0, 0, -1, 0, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (0, -1, 0, 0, 1, 0, 1), (-1, 0, 0, 0, 1, 0, 1), (-1, 0, 0, 0, 1, 0, 0))$,
- 3) $((0, 0, 0, 0, 0, 0, 1), (0, 0, 0, -1, 1, 0, 0), (0, 0, -1, 0, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (0, -1, 0, 0, 1, 0, 1), (-1, 0, 0, 0, 1, 0, 1), (-1, 0, 0, 0, 0, 0, 1))$,
- 4) $((0, 0, 0, -1, 1, 0, 0), (0, 0, -1, 0, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (0, -1, 0, 0, 1, 0, 1), (-1, 0, 0, 0, 1, 0, 1), (-1, 0, 0, 0, 0, 0, 1), (-1, 0, 0, 0, 1, 0, 0))$,
- 5) $((0, 0, 0, 0, 0, 0, 1), (0, 0, 0, -1, 1, 0, 0), (0, 0, -1, 0, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (0, -1, 0, 0, 1, 0, 1), (0, -1, 0, 0, 0, 0, 1), (-1, 0, 0, 0, 0, 0, 1))$,

- 6) $((0, 0, 0, 0, 1, 0, 0), (0, 0, 0, -1, 1, 0, 0), (0, 0, -1, 0, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (0, -1, 0, 0, 1, 0, 1), (0, -1, 0, 0, 1, -1, 1), (-1, 0, 0, 0, 1, 0, 0))$,
- 7) $((0, 0, 0, -1, 1, 0, 0), (0, 0, -1, 0, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (0, -1, 0, 0, 1, 0, 1), (-1, 0, 0, 0, 0, 0, 1), (-1, 0, 0, 0, 1, 0, 0), (-1, -1, 0, 0, 1, 0, 1))$,
- 8) $((0, 0, 0, 0, 0, 0, 1), (0, 0, 0, -1, 1, 0, 0), (0, 0, -1, 0, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (0, 0, -1, 0, 0, 0, 1), (0, -1, 0, 0, 0, 0, 1), (-1, 0, 0, 0, 0, 0, 1))$,
- 9) $((0, 0, 0, -1, 1, 0, 0), (0, 0, -1, 0, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (0, -1, 0, 0, 1, 0, 1), (0, -1, 0, 0, 0, 0, 1), (-1, 0, 0, 0, 0, 0, 1), (-1, -1, 0, 0, 1, 0, 1))$,
- 10) $((0, 0, 0, -1, 1, 0, 0), (0, 0, -1, 0, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (0, -1, 0, 0, 1, 0, 1), (0, -1, 0, 0, 1, -1, 1), (-1, 0, 0, 0, 1, 0, 0), (-1, -1, 0, 0, 1, 0, 1))$,
- 11) $((0, 0, 0, 0, 1, 0, 0), (0, 0, 0, -1, 1, 0, 0), (0, 0, -1, 0, 1, 0, 0), (0, -1, 0, 0, 1, 0, 1), (0, -1, 0, 0, 1, 0, 0), (0, -1, 0, 0, 1, -1, 1), (-1, 0, 0, 0, 1, 0, 0))$,
- 12) $((0, 0, 0, -1, 1, 0, 0), (0, 0, -1, 0, 1, 0, 0), (0, -1, 0, 0, 1, 0, 1), (0, -1, 0, 0, 1, 0, 0), (0, -1, 0, 0, 1, -1, 1), (-1, 0, 0, 0, 1, 0, 0), (-1, -1, 0, 0, 1, 0, 1))$,
- 13) $((0, 0, 0, -1, 1, 0, 0), (0, 0, -1, 0, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (0, -1, 0, 0, 1, 0, 1), (0, -1, 0, 0, 0, 0, 1), (0, -1, 0, 0, 1, -1, 1), (-1, -1, 0, 0, 1, 0, 1))$,
- 14) $((0, 0, 0, -1, 1, 0, 0), (0, 0, -1, 0, 1, 0, 0), (0, -1, 0, 0, 1, 0, 1), (0, -1, 0, 0, 1, 0, 0), (0, -1, 0, 0, 1, -1, 1), (0, -1, 0, 0, 0, 0, 1), (-1, -1, 0, 0, 1, 0, 1))$,
- 15) $((0, 0, 0, -1, 1, 0, 0), (0, 0, -1, 0, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (0, 0, -1, 0, 0, 0, 1), (0, -1, 0, 0, 0, 0, 1), (-1, 0, 0, 0, 0, 0, 1), (-1, 0, -1, 0, 0, 0, 1))$,
- 16) $((0, 0, 0, 0, 0, 0, 1), (0, 0, 0, -1, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (0, 0, 0, -1, 0, 0, 1), (0, 0, -1, 0, 0, 0, 1), (0, -1, 0, 0, 0, 0, 1), (-1, 0, 0, 0, 0, 0, 1))$,
- 17) $((0, 0, 0, -1, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (0, 0, 0, -1, 0, 0, 1), (0, 0, -1, 0, 0, 0, 1), (0, -1, 0, 0, 0, 0, 1), (-1, 0, 0, 0, 0, 0, 1), (-1, 0, -1, 0, 0, 0, 1))$,
- 18) $((0, 0, 0, -1, 1, 0, 0), (0, 0, -1, 0, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (0, 0, -1, 0, 0, 0, 1), (0, -1, 0, 0, 0, 0, 1), (0, -1, -1, 0, 0, 0, 1), (-1, 0, -1, 0, 0, 0, 1))$,
- 19) $((0, 0, 0, -1, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (0, 0, 0, -1, 0, 0, 1), (0, 0, -1, 0, 0, 0, 1), (0, -1, 0, 0, 0, 0, 1), (0, -1, -1, 0, 0, 0, 1), (-1, 0, -1, 0, 0, 0, 1))$,
- 20) $((0, 0, 0, -1, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (0, 0, 0, -1, 0, 0, 1), (0, -1, 0, 0, 0, 0, 1), (-1, 0, 0, 0, 0, 0, 1), (-1, 0, -1, 0, 0, 0, 1), (-1, 0, 0, -1, 0, 0, 1))$,
- 21) $((0, 0, 0, -1, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (0, 0, 0, -1, 0, 0, 1), (0, -1, 0, 0, 0, 0, 1), (0, -1, -1, 0, 0, 0, 1), (-1, 0, -1, 0, 0, 0, 1), (-1, 0, 0, -1, 0, 0, 1))$,
- 22) $((0, 0, 0, -1, 1, 0, 0), (0, 0, 0, 0, 0, -1, 1), (0, 0, 0, -1, 0, 0, 1), (0, -1, 0, 0, 0, 0, 1), (0, -1, -1, 0, 0, 0, 1), (0, -1, 0, -1, 0, 0, 1), (-1, 0, 0, -1, 0, 0, 1))$.

It turns out that $\#\tau\text{-tilt}_\epsilon A = 22$ and $2\text{-silt}_\epsilon A$ is finite.

We address the remaining 7 cases in a manner similar to the approach outlined above. More precisely, we first find the left mutation $\mu_{P_l}^-(A)$ of A corresponding to ϵ , and then calculate the left mutation sequences starting from $\mathcal{F}(\mu_{P_l}^-(A))$ until we find a finite connected component in $\mathcal{H}(\tau\text{-tilt}_\epsilon A)$. The utilization of Aoki's GAP-QPA program enables us to reduce this heavy process to a computer. We have

ϵ	$\#\tau\text{-tilt}_\epsilon A$
$(-, -, -, +, -, -, +)$	51
$(-, -, -, +, +, -, +)$	73
$(-, -, +, -, +, -, +)$	102
$(-, -, -, -, +, +, +)$	115
$(-, -, -, -, -, +, +)$	142
$(-, -, -, +, -, +, +)$	242
$(-, -, +, -, -, +, +)$	1067

By Proposition 3.4, the poset $2\text{-silt}_\epsilon A$ is finite for each of these 7 cases.

We have proved that $2\text{-silt}_\epsilon A$ is finite for any $\epsilon \in \mathfrak{s}_7$. □

Remark 5.4. We give an application of Proposition 3.13. Let $A = S^+(2, 6)$ over $p = 5$. Then, we have

$$\mu_{P_3}^-(A) = \left[\begin{array}{ccc} 0 & \longrightarrow & P_0 \oplus P_1 \oplus P_2 \\ & & \oplus \\ & & P_3 \xrightarrow{\alpha_3} P_4 \\ & & \oplus \\ 0 & \longrightarrow & P_4 \oplus P_5 \oplus P_6 \end{array} \right] \quad \text{and} \quad \mathcal{G}(\mu_{P_3}^-(A)) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

It is easy to check that $\mu_{P_3}^-(A)$ is a tilting complex. Set $B = \text{End } \mu_{P_3}^-(A)$. Then, B is isomorphic to KQ/I with

$$Q : \begin{array}{ccccccc} & & 6 & \xrightarrow{\alpha_5} & 5 & \xrightarrow{\alpha_4} & 4 & \xleftarrow{x} & 3 \\ & & \beta_1 \downarrow & \swarrow & \downarrow \beta_0 & & \downarrow y & & \\ & & 1 & \xrightarrow{\alpha_0} & 0 & & 2 & & \\ & & & \searrow & \alpha_1 & & & & \end{array} \quad \text{and} \quad I : \langle \alpha_5 \beta_0 - \beta_1 \alpha_0, \alpha_5 \alpha_4 y, y \alpha_1 \alpha_0 \rangle.$$

We take, for example, $\epsilon = (-, +, +, -, -, +, +)$, and it is sent to $\epsilon' = (-, +, +, +, -, +, +)$ under the action of $\mathcal{G}(\mu_{P_3}^-(A))$. We then obtain $2\text{-silt}_\epsilon A \simeq 2\text{-silt}_{\epsilon'} B$.

In summary, we have the following result.

Theorem 5.5. *A representation-infinite Borel-Schur algebra $S^+(n, r)$ is τ -tilting finite if and only if $n = 2$ and $p = 2, r = 4$, or $p = 3, r = 5$, or $p = 5, r = 6$.*

ACKNOWLEDGEMENTS

The author is grateful to Toshitaka Aoki for many useful discussions on silting theory. The author is partially supported by National Key Research and Development Program of China (Grant No. 2020YFA0713000) and China Postdoctoral Science Foundation (Grant No. 315251 and No. 2023M731988).

DATA AVAILABILITY STATEMENTS

This is a research in pure mathematics. All data generated or analyzed during this study are included in this manuscript.

ETHICAL STATEMENT

Not applicable.

CONFLICT OF INTEREST STATEMENT

Not applicable.

REFERENCES

- [Ad] T. Adachi, Characterizing τ -tilting finite algebras with radical square zero. *Proc. Amer. Math. Soc.* **144** (2016), no. 11, 4673–4685.
- [AIR] T. Adachi, O. Iyama and I. Reiten, τ -tilting theory. *Compos. Math.* **150** (2014), no. 3, 415–452.
- [AAC] T. Adachi, T. Aihara and A. Chan, Classification of two-term tilting complexes over Brauer graph algebras. *Math. Z.* **290** (2018), no. 1–2, 1–36.
- [AI] T. Aihara and O. Iyama, Silting mutation in triangulated categories. *J. Lond. Math. Soc.* **85** (2012), no. 3, 633–668.
- [AiW] T. Aihara and Q. Wang, A symmetry of silting quivers. *Glasg. Math. J.* **65** (2023), no. 3, 687–696.
- [AHMW] T. Aihara, T. Honma, K. Miyamoto and Q. Wang, Report on the finiteness of silting objects. *Proc. Edinb. Math. Soc. (2)* **64** (2021), no. 2, 217–233.
- [Ao] T. Aoki, Classifying torsion classes for algebras with radical square zero via sign decomposition. *J. Algebra* **610** (2022), 167–198.
- [AHIKM] T. Aoki, A. Higashitani, O. Iyama, R. Kase and Y. Mizuno, Fans and polytopes in tilting theory. Preprint (2022), arXiv: 2203.15213.
- [AoW] T. Aoki and Q. Wang, On τ -tilting finiteness of blocks of Schur algebras. Preprint (2021), arXiv: 2110.02000.
- [ALS] S. Ariki, S. Lyle and L. Speyer, Schurian-finiteness of blocks of type A Hecke algebras. *J. Lond. Math. Soc. (2)* **108** (2023), no.6, 2333–2376.
- [As] S. Asai, Semibricks. *Int. Math. Res. Not.* (2020), no. 16, 4993–5054.
- [AMN] H. Asashiba, Y. Mizuno and K. Nakashima, Simplicial complexes and tilting theory for Brauer tree algebras. *J. Algebra* **557** (2020), 119–153.
- [B] K. Bongartz, Critical simply connected algebras. *Manuscripta Math.* **46** (1984), no. 1-3, 117–136.
- [BST] T. Brüstle, D. Smith and H. Treffinger, Stability conditions, τ -tilting theory and maximal green sequences. Preprint (2017), arXiv:1705.08227.
- [DIJ] L. Demonet, O. Iyama and G. Jasso, τ -tilting finite algebras, bricks, and g -vectors. *Int. Math. Res. Not.* (2019), no. 3, 852–892.
- [DIRRT] L. Demonet, O. Iyama, N. Reading, I. Reiten and H. Thomas, Lattice theory of torsion classes: beyond τ -tilting theory. *Trans. Amer. Math. Soc. Ser. B* **10** (2023), 542–612.
- [D] Yu. A. Drozd, Tame and wild matrix problems, in: Representation Theory II, Lecture Notes in Math., vol. 832. *Springer Verlag*, (1980), pp. 242–258.
- [ESY1] K. Erdmann, A. P. Santana and I. Yudin, On Auslander-Reiten sequences for Borel-Schur algebras. *J. Algebra Appl.* **17** (2018), no. 2, 1850028.
- [ESY2] K. Erdmann, A. P. Santana and I. Yudin, Representation type of Borel-Schur algebras. *Algebr. Represent. Theory* **24** (2021), no. 6, 1387–1413.
- [G] J. A. Green, On certain subalgebras of the Schur algebra. *J. Algebra* **131** (1) (1990), 265–280.
- [HR] D. Happel and C. M. Ringel, Tilted algebras. *Trans. Amer. Math. Soc.* **274** (1982), no. 2, 399–443.
- [HV] D. Happel and D. Vossieck, Minimal algebras of infinite representation type with preprojective component. *Manuscripta Math.* **42** (1983), no. 2-3, 221–243.
- [IR] O. Iyama and I. Reiten, Introduction to τ -tilting theory. *Proc. Natl. Acad. Sci.* **111** (2014), no. 27, 9704–9711.
- [KT] M. Kaipel and H. Treffinger, Wall-and-chamber structures for finite-dimensional algebras and τ -tilting theory. Preprint (2023), arXiv: 2302.12699.

- [L] D. G. Liang, On the quiver and relations of the Borel Schur algebras. *Ph.D. Thesis*, 2007.
- [MS] P. Malicki and A. Skowroński, Cycle-finite algebras with finitely many τ -rigid indecomposable modules. *Comm. Algebra* **44** (2016), no. 5, 2048–2057.
- [MW] K. Miyamoto and Q. Wang, On τ -tilting finiteness of tensor products between simply connected algebras. Preprint (2021), arXiv: 2106.06423.
- [Mi] Y. Mizuno, Classifying τ -tilting modules over preprojective algebras of Dynkin type. *Math. Z.* **277** (2014), no. 3-4, 665–690.
- [Mo] K. Mousavand, τ -tilting finiteness of biserial algebras. Preprint (2019), arXiv: 1904.11514.
- [P] P. G. Plamondon, τ -tilting finite gentle algebras are representation-finite. *Pacific J. Math.* **302** (2019), no. 2, 709–716.
- [Ric] J. Rickard, Morita theory for derived categories. *J. London Math. Soc.* **39** (1989), no. 3, 436–456.
- [RT] A. Rogat and T. Tesche, The Gabriel quivers of the sincere simply connected algebras. *Er-ganzungsreihe*, Bielefeld (1993).
- [S] A. P. Santana, The Schur algebra $S(B^+)$ and projective resolutions of Weyl modules. *J. Algebra* **161** (1993), no. 2, 480–504.
- [SY] A. P. Santana and I. Yudin, Characteristic-free resolutions of Weyl and Specht modules. *Adv. Math.* **229** (2012), no. 4, 2578–2601.
- [STV] S. Schroll, H. Treffinger and Y. Valdivieso, On band modules and τ -tilting finiteness. *Math. Z.* **299** (2021), no. 3-4, 2405–2417.
- [T] H. Treffinger, τ -tilting theory - An introduction. Preprint (2021), arXiv: 2106.00426.
- [Wa1] Q. Wang, τ -tilting finiteness of two-point algebras I. *Math. J. Okayama Univ.* **64** (2022), 117–141.
- [Wa2] Q. Wang, On τ -tilting finite simply connected algebras. *Tsukuba J. Math.* **46** (1) (2022), 1–37.
- [Wa3] Q. Wang, On τ -tilting finiteness of the Schur algebra. *J. Pure Appl. Algebra* **226** (1) (2022), 106818.
- [Wa4] Q. Wang, τ -tilting finiteness of two-point algebras II. *To appear in Journal of Algebra and its Applications*, (2022), arXiv: 2206.00239.
- [Wo1] D. Woodcock, Schur algebras and global bases: new proofs of old vanishing theorems. *J. Algebra* **191** (1997), no.1, 331–370.
- [Wo2] D. Woodcock, Borel Schur algebras. *Comm. Algebra* **22** (1994), no.5, 1703–1721.
- [Y] I. Yudin, On projective resolutions of simple modules over the Borel subalgebra $S^+(n, r)$ of the Schur algebra $S(n, r)$ for $n \leq 3$. *J. Algebra* **319** (2008), no.5, 1870–1902.
- [Z] S. Zito, τ -tilting finite cluster-tilted algebras. *Proc. Edinb. Math. Soc.* (2) **63** (2020), no. 4, 950–955.

YAU MATHEMATICAL SCIENCES CENTER, TSINGHUA UNIVERSITY, BEIJING 100084, CHINA.
 Email address: infinite-wang@outlook.com & infinite-wang@tsinghua.edu.cn