A note on Gauss operators and quantizations of Yang-Mills theories

Bruno Bucciotti[∗](#page-0-0)

Scuola Normale Superiore, Piazza dei Cavalieri 7, I-56126 Pisa, Italy INFN, Sezione di Pisa, Largo B. Pontecorvo, 3, I-56127 Pisa, Italy

Franco Strocchi[†](#page-0-1)

Dipartimento di Fisica, Università di Pisa, Largo B. Pontecorvo, 3, I-56127 Pisa, Italy

(Dated: 02 Oct 2023)

The quantization of Yang-Mills field theories requires the introduction of a gauge fixing which leads to a violation of the Local Gauss Law described by the so-called Gauss operator. We discuss the local quantizations of Yang-Mills theories in terms of the possible Gauss operators, which are argued to have a more physical meaning than the gauge fixings.

We focus the attention on the local quantizations which leave the global gauge group and a subgroup of local gauge transformations unbroken, as the Feynman quantization of quantum Electrodynamics, and show that in the non-abelian case such properties cannot be realized together with Lorentz covariance; thus, quite generally, one cannot have the structural properties of the Feynman quantization of Quantum Electrodynamics.

By relaxing the condition of Lorentz covariance, we obtain a classification of Gauss operators, which satisfy gauge covariant conservation laws and generate non-trivial residual subgroups of local gauge transformations.

I. INTRODUCTION

Given a Lie *(global)* gauge group G , the correspondent local gauge group $\mathcal G$ is conventionally obtained by replacing the constant group parameters of G by infinitely differentiable functions, which is convenient to choose of compact support in space and of slow (i.e. at most poly- $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ increase in time¹.

A classical gauge field theory is defined by a Lagrangian \mathcal{L}_{inv} invariant under a local gauge group $\mathcal{G},$ with the fields (which enter in the definition of \mathcal{L}_{inv}) transforming as representations of \mathcal{G} . In order to have a well posed Cauchy problem, i.e. a deterministic time evolution of the fields, one must add a gauge fixing \mathcal{L}_{GF} , which need not break $\mathcal G$ to the identity (as implied by the functional integral argument), but only to the extent of allowing a deterministic field dynamics [\[1](#page-6-0)] (for example, the Feynman gauge in electrodynamics leaves the unbroken residual group with parameters satisfying $\Box \varepsilon = 0$.

The introduction of a gauge fixing is crucial for a quantization of the field algebra since, according to the principles of quantum mechanics, the dynamics is described by a unitary operator $U(t) = e^{-itH}$, and therefore in order to have quantum fields the dynamics of the field operators must be deterministic.

Thus, a quantization of a gauge field theory is defined by the choice of a gauge fixing, different gauge fixings identifying different quantum field algebras F , even if the resulting representations of the observable field subalgebra \mathcal{F}_{obs} are all equivalent; one of the main roles of the field algebra $\mathcal F$ is to allow, through its vacuum representation, the construction of the representations of \mathcal{F}_{obs} beyond its vacuum sector [\[1](#page-6-0)], namely the expectations of the observables on the states with non-trivial G charges.

We consider quantizations of Yang-Mills field theories (with G a compact Lie group) defined by vacuum representations of local field algebras $\mathcal F$, in a vector space $V = \mathcal{F}\Psi_0$, obtained by gauge fixings which do not break the global gauge group G, satisfy locality and leave a residual local gauge subgroup $\mathcal{G}_r \subset \mathcal{G}$ unbroken.

We briefly mention the motivations for such a choice.

Gauge fixings which break the global gauge group G down to the identity (like the unitary gauge and the ξ gauges) involve a mean field parameter which is difficult to justify beyond the perturbative approach and it is at most allowed for the discussion of the phenomenon of gauge symmetry breaking, as in the electroweak theory.

A residual local gauge group \mathcal{G}_r is very useful for exploiting the related Ward identities, a crucial tool for the proof of renormalization, as displayed by the prototypical case of Feynman gauge in Quantum Electrodynamics.

[∗] bruno.bucciotti@sns.it

[†] franco.strocchi@sns.it

¹ Regularity conditions on the gauge functions are needed, since the quantum fields are operator valued distributions, so that the transformations induced by the action of $\mathcal G$ should have the property of multipliers of the related test functions. The requirement of compact support in space guarantees the very useful property of space localization of the gauge transformations and it also provides the needed clear cut distinction between the global and the local gauge group. For the time dependence it is enough a slow increase in time; the condition of compact support in time would exclude the local gauge transformations in the temporal gauge and it is not required. Technically, the group parameters of G are required to belong to $\mathcal{D}(\mathbf{R}^3) \times \mathcal{O}_M(\mathbf{R})$). Very problematic is the generalization of the local gauge transformations with the gauge functions replaced by field operators, since it involves the singular (actually divergent) multiplication of operator valued distributions at the same point. For more detailed motivations and useful consequences of such choices see [\[1\]](#page-6-0).

Furthermore, the unbroken residual group allows us to exploit its topology for codifying the interplay between gauge invariance and other symmetries, in particular chiral symmetries. Indeed, the representation of the topology of the residual group yields a non-perturbative solution of the $U(1)$ problem in Quantum Chromodynamics, without relying on the semi-classical instanton approximation, which is not free of mathematical problems [\[1](#page-6-0)].

The consideration of local field algebras, i.e. of gauge fixings which do not violate locality, is motivated by the fact most of what is known on quantum field theories at the non-perturbative level relies on locality [\[2\]](#page-6-1), which is also shared by the (perturbatively) renormalizable gauges.

Another ingredient of our analysis is the distinguished role of the Gauss operators, for the classification of the possible quantizations of Yang-Mills theories, since they have a more physical status than the gauge fixings, as argued below.

In fact, the presence of the gauge fixing invalidates the conclusion of the second Noether theorem, namely the validity of Local Gauss Law

$$
J^a_\mu = \partial^\nu F^a_{\nu\,\mu},\tag{1}
$$

satisfied by the currents J^a_μ , $(a = 1, ..., n, n =$ the dimension of G), which generate the infinitesimal transformations of G on the field algebra \mathcal{F} :

$$
\delta^a F = \lim_{R \to \infty} i \left[Q_R^a, F \right], \quad \forall F \in \mathcal{F}, \tag{2}
$$

where Q_R^a is a suitably regularized integral of J_0^a over a sphere of radius R.

Hence, instead of eq. [\(1\)](#page-1-0) one rather gets

$$
J^a_\mu = \partial^\nu F^a_{\nu\,\mu} + G^a_\mu,\tag{3}
$$

where G^a_μ , henceforth called the *Gauss operator*, depends on the choice of the gauge fixing.

The unbroken global group G implies that $\partial^{\mu} J_{\mu}^{a} = 0$ and therefore eq. [\(3\)](#page-1-1) and the antisymmetry of $F_{\nu\mu}$ yield

$$
\partial^{\mu}G^{a}_{\mu}=0,\t\t(4)
$$

i.e. the Gauss operator satisfies a continuity equation. Thus, it plays the role of an "external" conserved current, (a meaning not shared by the gauge fixing).

Moreover, for any physical state Ψ (typically obtained by a non-local construction through the vacuum representation of $\mathcal F$, as for example in the Gupta-Bleuer quantization of QED), the Local Gauss Law holds (for more details and proofs, see [\[1\]](#page-6-0)):

$$
(\Psi, (J_{\mu}^{a} - \partial^{\nu} F_{\nu \mu}^{a}) \Psi) = (\Psi, G_{\mu}^{a} \Psi) = 0.
$$
 (5)

Actually, such a property qualifies the physical states (in Quantum Electrodynamics amounts to the validity of the Maxwell equations on the physical states), so that, by eq. [\(5\)](#page-1-2), the Gauss operator provides the characterization of the physical states.

In addition, by the locality of the field algebra it has been shown [\[3\]](#page-6-2) that in the Feynman quantization of QED and in general in the temporal gauge of Yang-Mills theories the Gauss operator generates the infinitesimal transformations of the residual local gauge group in the given representation of F , so that it belongs to the commutant of the \mathcal{F}_{obs} and therefore it classifies the representations of \mathcal{F}_{obs} contained in the representation space of \mathcal{F} . In the abelian case, the Gauss operator even belongs to \mathcal{F}_{obs} and therefore to its center.

The role and meaning of the Gauss operator in such a construction of the physical states become more evident in connection with locality.

In fact, the Gauss law, eq. [\(1\)](#page-1-0), implies that the flux of F_{i0}^{a} at infinity, briefly called *Gauss charge* and denoted by Q_G^a , coincides with the total charge, a suitable integral of J_0^a , called the *current charge* and denoted by Q_J^a . Then, since by locality the Gauss charge has vanishing expectations on local states, so does the charge Q_J^a and, as a consequence, the global gauge group G acts trivially in the representation space of a local field algebra \mathcal{F} .

Hence, the role of the Gauss operator is to screen/compensate the contribution of the current charge to the flux of F_{i0}^a at infinity allowing for local states with non-zero charge and, consequently, for a non-trivial action of G in the representation space of \mathcal{F} .

Furthermore, a clever choice of the Gauss operator may make more explicit the hyperbolic and elliptic contents of eq. [\(3\)](#page-1-1), which in the abelian case is equivalent to:

$$
\Box F_{\mu\nu} = \partial_{\mu}J_{\nu} - \partial_{\nu}J_{\mu} - \partial_{\mu}G_{\nu} + \partial_{\nu}G_{\mu}, \quad \partial^{i}F_{i\,0} = J_{0} - G_{0}.
$$
\n(6)

In fact, in the abelian case the choice of the temporal gauge $G_i = 0$, which implies $\partial_0 G_0 = 0$, leads to a standard hyperbolic equation for $F_{i,j}$,

$$
\Box F_{i\,j} = \partial_i J_j - \partial_j J_i,\tag{7}
$$

without involving the Gauss operator, which only enters in the "elliptic" constraint involving F_{i0} and J_0 , consistent with the time evolution of $F_{i\,0}$

$$
\partial^0 F_{0i} = \partial^j F_{i\,j} + J_i. \tag{8}
$$

Thus, the role of G_0 is that of a screening *static charge* density which is seen by the local states but not by the non-local physical charged states.

Furthermore, in the temporal gauge the Gauss operator satisfies a (gauge) covariant conservation law, namely $D^{\mu}G_{\mu}=0$, in agreement with its interpretation as the current of a (fictitious) charge distribution.

A similar role and meaning of the Gauss operator occurs also in the temporal gauge of the non-abelian case.

For these reasons, for the characterization of the possible (local) quantizations of a gauge field theory, the Gauss operator is better qualified than the gauge fixing, which does not have such properties.

The aim of this note is to discuss the constraints on the Gauss operators in the non-abelian case (and therefore on the possible quantizations) and point out the general obstruction for a quantization with the properties similar to that of the Feynman gauge in Quantum Electrodynamics.

We show that it is impossible to have a quantization of non-abelian Yang-Mills field theories which satisfies locality, Lorentz covariance and leaves the global gauge group G and a subgroup $\mathcal{G}_r \subset \mathcal{G}$ of local gauge transformations unbroken. This provides a general argument for the recourse to BRST quantization in the case of non-abelian Yang-Mills theories.

In addition, we obtain a classification of the Gauss operators which satisfy a (gauge) covariant conservation law. This property yields that the gauge fixing and the corresponding Gauss operator minimally affect the gauge group geometry of the Yang-Mills equations, being required in particular if the current which generates the global group has to be left unchanged by the gauge fixing.

As a consequence, one has the desirable property that the action on the matter fields by the Gauss operator and by the topological operators (namely, the operators which describe the topology of the residual group) is the same as that of the current in the absence of gauge fixing.

This pattern does not seem to be available in the BRST quantization [\[4](#page-6-3)]: the invariance group of the Lagrangian is no longer the group $\mathcal G$ of local gauge transformations but rather the BRST group which involves field transformations with parameters which are field operators. As discussed above, the mathematical control of such transformations, which involve singular multiplication of distributions at the same point, is rather problematic, beyond the perturbative approach.

The BRST gauge fixing does not leave a subgroup \mathcal{G}_r of local gauge transformations unbroken and the singular (distributional) properties of the fields which parametrize the BRST transformations preclude the possibility of considering their topological properties.

From this point of view, the quantization of the temporal gauge proved more effective for a non-perturbative analysis of non-abelian Yang-Mills theory, at the only expense of losing manifest Lorentz covariance. In exchange, the vacuum representation of the field algebra of the temporal gauge satisfies positivity and does not involve ghosts (so that the fields may be defined as operators in a Hilbert space), whereas the vacuum representation of the BRST field algebra is indefinite and, in particular, the ghosts are not operators in a Hilbert space. Needless to say, the identification or construction of the Hilbert space of physical states is much easier if the vacuum representation of the field algebra has a Hilbert space structure.

II. GAUSS OPERATORS IN YANG-MILLS GAUGE THEORIES

For the reasons discussed above, in the following we consider the possible realization of a Gauss operators G_u which has the following properties:

- i) it leaves the global gauge group G unbroken,
- ii) it leaves a residual group \mathcal{G}_r of local gauge transformations unbroken; this allows for the exploitation of the corresponding Ward identities, and in the nonabelian case the effect of the topology of \mathcal{G}_r on the properties of the physical states,
- iii) it defines a *local field*, in the representation space V of the local field algebra,
- iv) it allows a simple characterization of the physical states in terms of the subspace of the representation space of F , or of a suitable completion of it, with vanishing expectation of G_{μ} ^{[2](#page-2-0)}

A. Group theoretical constraints on Gauss operators

Since by ii) the Gauss operators under our consideration are related to residual subgroups of local gauge transformations, the characterization of such subgroups provides group theoretical constraints on the analysis of possible Gauss operators.

As mentioned above, the local gauge group $\mathcal G$ is parametrized by the set of infinitely differentiable functions of compact support in space, $\varepsilon^a(x)$, $a = 1,...n$, which form not only a vector space V with the sum defined by

$$
(\varepsilon_1 + \varepsilon_2)^a(x) \equiv \varepsilon_1^a(x) + \varepsilon_2^a(x),
$$

but also an algebra $\mathcal E$ with product

$$
(\varepsilon_1 \wedge \varepsilon_2)^a(x) \equiv i f_{bc}^a \varepsilon_1^b(x) \varepsilon_2^c(x).
$$

The abelian subgroups \mathcal{G}_i of \mathcal{G}_i , stable under G by i), are identified by subspaces V_i of V , whereas the non-abelian subgroups are identified by sub-algebras \mathcal{E}_i of \mathcal{E}_i , since in the non-abelian case the group composition law requires stability under the above product (see the Zassenhaus formula below).

The stability of \mathcal{G}_i under the global group G requires that if $\varepsilon^a(x)T^a$ belongs to the Lie algebra of \mathcal{G}_i so does $\lbrack \varepsilon^b T^b, \varepsilon^a(x)T^a \rbrack, \forall b, \text{ i.e. if } \varepsilon^a \text{ belongs to the parameter }$ space of \mathcal{G}_i so does $\varepsilon^c(x) = f^c_{a b} \varepsilon^a(x)$, $\forall b$.

² For examples which realize such a structure see [\[1](#page-6-0)].

The above properties must be satisfied by the local gauge subgroups related to the Gauss operators and therefore provide a first classification of them.

Another restriction comes from the condition that the Gauss operators admit a derivation from a Lagrangian.

The invariance of the gauge fixing under the residual group \mathcal{G}_r , with gauge parameters ε_r is codified by the following equations in terms of the gauge fixing \mathcal{L}_{GF} :

$$
L_a^{\mu} \, \partial_{\mu} \varepsilon_r^a \equiv \tag{9}
$$

$$
\left\{\frac{\delta \mathcal{L}_{GF}}{\delta A_{\mu}^{a}} + iT_{bc}^{a} \left[\frac{\delta \mathcal{L}_{GF}}{\delta \partial_{\mu} A_{\nu}^{b}} A_{\nu}^{c} + \frac{\delta \mathcal{L}_{GF}}{\delta \partial_{\mu} B^{b}} B^{c}\right]\right\} \partial_{\mu} \varepsilon_{r}^{a} = 0,
$$

$$
L_a^{\mu\nu} \partial_\mu \partial_\nu \varepsilon_r^a \equiv \frac{\delta \mathcal{L}_{GF}}{\delta \partial_\mu A_\nu^a} \partial_\mu \partial_\nu \varepsilon_r^a = 0, \tag{10}
$$

where for simplicity, we omitted the presence of matter fields and considered gauge fixings \mathcal{L}_{GF} which are functions of A^a_μ and of fields B^a , of the Nakanishi-Lautrup type, which transform according to irreducible adjoint representations of G, taken to be a simple group.

In conclusion, in the case of an abelian global group G the possible residual groups, equivalently the possible Gauss operators, are identified by G invariant subspaces of V with gauge parameters $\varepsilon_r(x)$ satisfying eqs. [\(9\)](#page-3-0) and [\(10\)](#page-3-1).

For non-abelian G the characterization is provided by G invariant sub-algebras of $\mathcal E$ with gauge parameters $\varepsilon_r(x)$ satisfying eqs. [\(9\)](#page-3-0) and [\(10\)](#page-3-1).

Explicit cases are discussed below.

B. Feynman Gauss operator

A distinguished choice of the Gauss operator is that of the Feynman gauge in Quantum Electrodynamics, which preserves locality and covariance: $G_{\mu} = \partial_{\mu}L$, with L a massless longitudinal field.

In addition to the properties i)-iv), the Feynman Gauss operator in QED satisfies

- v) it is Lorentz covariant, allowing for a Lorentz covariant field algebra,
- vi) it provides a simple subsidiary condition, which linearizes the constraint eq. [\(5\)](#page-1-2) for the identification of the physical states,
- vii) it does not require ghost fields, since it involves only the vector potential and possibly the Nakanishi-Lautrup field B.

The technical advantages of such a choice are well known, in particular for the control of renormalization, the explicit construction of the physical charged states, etc. (see e.g. [\[1\]](#page-6-0)). It is then natural to ask whether a Gauss operator with such properties may be obtained also in the non-abelian case.

A negative answer has been presented in the literature and it is part of the standard textbook presentation of the need for BRST quantization. However, such a negative conclusion has been reached for the choice of a gauge fixing which is a literal transcription of the Feynman choice, namely for the gauge fixing $\frac{1}{2}\xi(\partial^{\mu}A_{\mu}^{a})^{2}$ or for its Nakanishi-Lautrup variations, e.g. $B^a \partial^{\mu} A^a_{\mu} + \frac{1}{2} \xi (\partial^{\mu} A^a_{\mu})^2$.

One may reasonably ask whether in the non-abelian case a more clever choice of the gauge fixing may work. Quite generally this leads to investigating the possible Lorentz covariant solutions of the equation $\partial^{\mu}G_{\mu}=0$.

For simplicity, we discuss the problem in the absence of matter fields and consider gauge fixings \mathcal{L}_{GF} which are functions of A^a_μ and of fields B^a , of the Nakanishi-Lautrup type, which transform according to irreducible adjoint representations of G , for simplicity taken to be a simple group. Then, the invariance of the gauge fixing \mathcal{L}_{GF} under the global gauge group G requires

$$
T_{bc}^a \left[\frac{\delta \mathcal{L}_{GF}}{\delta A_\mu^b} A_\mu^c + \frac{\delta \mathcal{L}_{GF}}{\delta B^b} B^c + \right] \tag{11}
$$

$$
+\frac{\delta \mathcal{L}_{GF}}{\delta \partial_{\nu} A_{\mu}^{b}} \partial_{\nu} A_{\mu}^{c} + \frac{\delta \mathcal{L}_{GF}}{\delta \partial_{\nu} B^{b}} \partial_{\nu} B^{c} \bigg] = 0 \tag{12}
$$

The additional restriction of Lorentz covariance plays a crucial role, requiring the Lorentz covariance of the subspaces of gauge parameters in the abelian case and of the sub-algebras of $\mathcal E$ in the non-abelian case. Then, the group parameters of \mathcal{G}_r should define a vector space stable under G and under Lorentz transformations. Now, since G does not have invariant subgroups, eq. (9) cannot provide restrictions on the index a, and similarly there is no linear subspace invariant under the Lorentz transformations to be spanned by the four vectors $\partial_{\mu} \varepsilon^{a}$.

Then, in the absence of restrictions on the group parameters, in order to satisfy eq. [\(9\)](#page-3-0) the expression in curly brackets should vanish

$$
\left\{\frac{\delta \mathcal{L}_{GF}}{\delta A_{\mu}^{a}} + iT_{bc}^{a} \left[\frac{\delta \mathcal{L}_{GF}}{\delta \partial_{\mu} A_{\nu}^{b}} A_{\nu}^{c} + \frac{\delta \mathcal{L}_{GF}}{\delta \partial_{\mu} B^{b}} B^{c}\right]\right\} = 0. \quad (13)
$$

It is instructive to check that eq. [\(9\)](#page-3-0) provides restrictions on the gauge parameter ε_r if Lorentz covariance is not required, as it happens for the temporal gauge. In fact, in this case the term in square brackets and $\delta\mathcal{L}_{GF}/\delta A_i^a$, $i = 1, 2, 3$ vanish, (see below), and eq. [\(9\)](#page-3-0) gives the restriction $\partial_0 \varepsilon_r = 0$, i.e. the residual group of the temporal gauge.

It remains to exploit eq. [\(10\)](#page-3-1), which (assuming Lorentz invariance) yields the following characterization

$$
\frac{\delta \mathcal{L}_{GF}}{\delta \partial_{\mu} A_{\nu}^{a}} + \frac{\delta \mathcal{L}_{GF}}{\delta \partial_{\nu} A_{\mu}^{a}} = g_{\mu \nu} L^{a}, \qquad \Box \varepsilon_{r}^{a}(x) = 0, \qquad (14)
$$

with $L^a \neq 0$, a scalar function of the fields. Clearly, if $L^a = 0$, the condition $\Box \varepsilon^a = 0$ is not required, but then the absence of conditions on the ε^a would imply $\mathcal{G}_r = \mathcal{G}$, i.e. no effective gauge fixing.

Hence, the possible residual subgroup should be characterized by gauge parameters $\varepsilon^a(x)$ satisfying the (free) wave equation. Indeed, as we shall see, this works in the abelian case, but not in the non-abelian case.

In fact, in the non-abelian case there cannot be a residual group \mathcal{G}_r characterized by gauge parameters ε_r satisfying $\Box \varepsilon_r = 0$, since this condition keeps being satisfied by linear combination of such functions (vector space structure), but not by their products in conflict with the group composition rule.

This is displayed by the Zassenhaus formula, [\[5,](#page-6-4) [6\]](#page-6-5), which is known to converge for small values of the parameters; in fact, it gives

$$
e^{i\varepsilon_1^a(x)t^a}\ e^{i\varepsilon_2^b(x)t^b}=e^{i\varepsilon_1^a(x)t^a+i\varepsilon_2^b(x)t^b}\ e^{-\frac{1}{2}\varepsilon_1^a(x)\varepsilon_2^b(x)[t^a,t^b]}...
$$

The first term on the right hand side is the exponential of an element of the Lie algebra, and therefore an element of the group, but the other factors, which do not vanish in the non-abelian case, do not define elements of a gauge group characterized by the above gauge parameters, because the product of gauge functions which satisfy the wave equation does not satisfy the wave equation.

In conclusion, quite generally in the non-abelian case it is impossible to have a covariant Gauss operator satisfying i)-ii), as a function of A^a_μ and possible Nakanishi-Lautrup fields; hence the properties of the Feynman Gauss operator are in general excluded. Lorentz covariance can be rescued by relaxing the condition of a nontrivial residual gauge group, as in BRST quantization.

C. Gauss operator and covariant conservation law

By the defining eq. [\(3\)](#page-1-1), $\partial^{\mu}G_{\mu} = 0$, where J_{μ}^{a} is the current which generates the unbroken global gauge group G thanks to Noether theorem.

Now, in the absence of a gauge fixing, the Gauss law may also be written as

$$
D^{\nu}F_{\nu\mu}^{a} = j_{\mu}^{a}(\psi),\tag{15}
$$

where D^{μ} denotes the covariant derivative and $j^a_{\mu}(\psi)$ is the contribution to the conserved current by the matter fields (the so-called matter current). Both terms of this equation satisfy a covariant conservation law

$$
D^{\mu}D^{\nu}F_{\nu\mu}^{a} = D^{\mu}j_{\mu}^{a}(\psi) = 0.
$$
 (16)

One may then ask whether such covariant conservation laws, strictly related to the gauge group geometry, keep holding also in the presence of a gauge fixing. Thus we look for Gauss operators which satisfy a covariant conservation law (besides the ordinary conservation law), i.e.

$$
D^{\mu}G_{\mu}^{a}=0.\t\t(17)
$$

We allow for gauge fixings which are functions of A^a_μ and of auxiliary fields of the Nakanishi-Lautrup type B_i^a , $i = 1, \dots k$, which transform as the adjoint representation of G , and possibly G invariant fields B_i .

We shall see that if

a) the gauge fixing does not depend on the derivatives of A^a_μ , and

b) the current

$$
j^a_\mu(B) = \frac{\delta \mathcal{L}}{\delta \partial_\mu B_i^b} \delta^a B_i^b = \frac{\delta \mathcal{L}_{GF}}{\delta \partial_\mu B_i^b} \delta^a B_i^b
$$

satisfies a covariant conservation law (in particular $j^a_\mu(B) = 0$ if \mathcal{L}_{GF} does not depend on the derivatives of the B_i^a)

then, the corresponding Gauss operator satisfies a covariant conservation law. Indeed we shall see that, contrary to what implicitly argued in the literature, there are solutions of eq. [\(17\)](#page-4-0) and of the continuity equation.

We denote by \mathcal{L}_0 the gauge invariant Lagrangian, so that $\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_{GF}$. By Noether theorem, the invariance of $\mathcal L$ under the global gauge group G determines the conserved current J^a_μ which generates the transformations of G:

$$
J_{\mu}^{a} = i T_{bc}^{a} \left[\frac{\delta \mathcal{L}}{\delta \partial_{\mu} A_{\nu}^{b}} A_{\nu}^{c} + \frac{\delta \mathcal{L}}{\delta \partial_{\mu} B_{i}^{b}} B_{i}^{c} \right] + j_{\mu}^{a}(\psi) = \qquad (18)
$$

$$
= i T^a_{bc} \left[\frac{\delta \mathcal{L}_0}{\delta \partial_\mu A^b_\nu} A^c_\nu + \frac{\delta \mathcal{L}_{GF}}{\delta \partial_\mu B^b_i} B^c_i \right] + j^a_\mu(\psi) = j^a_\mu(A) + j^a_\mu(B_i) + j^a_\mu(\psi), \tag{19}
$$

where $j^a_\mu(\psi)$ denotes the contribution of the matter fields, the so-called matter current, and $j^a_\mu(A)$ = $-iT^a_{bc}(A^b)^{\nu}F^c_{\nu\mu}.$

The second Noether theorem applied to $\mathcal{L} - \mathcal{L}_{GF}$ provides information on the effect of \mathcal{L}_{GF} in yielding eq. [\(3\)](#page-1-1).

The coefficient of $\partial_{\mu}\varepsilon$ in the infinitesimal variation of

 $\mathcal{L}_0 = \mathcal{L} - \mathcal{L}_{GF}$ must vanish and one gets

$$
J_{\mu}^{a} + \frac{\delta \mathcal{L}}{\delta A_{\mu}^{a}} - \frac{\delta \mathcal{L}_{GF}}{\delta A_{\mu}^{a}} - j_{\mu}^{a}(B_{i}) =
$$

$$
J_{\mu}^{a} + \partial^{\nu} \frac{\delta \mathcal{L}_{0}}{\delta \partial_{\nu} A_{\mu}^{a}} - \frac{\delta \mathcal{L}_{GF}}{\delta A_{\mu}^{a}} - j_{\mu}^{a}(B_{i}) =
$$
(20)

$$
=J_{\mu}^{a} + \partial^{\nu} F_{\mu\nu}^{a} - \frac{\delta \mathcal{L}_{GF}}{\delta A_{\mu}^{a}} - j_{\mu}^{a}(B_{i}) = 0, \qquad (21)
$$

where we have used that

1) \mathcal{L}_{GF} does not depend on the derivatives of A^a_μ ,

2) \mathcal{L}_0 does not depend on the auxiliary fields $B_i^{a'}$, so that the derivatives of \mathcal{L}_{GF} with respect to the fields B_i^a and to their derivatives may be replaced by the corresponding derivatives of \mathcal{L} , allowing to use the equations of motion for the fields B_i^a .

Hence, the Gauss operator is given by

$$
G_{\mu}^{a} = \frac{\delta \mathcal{L}_{GF}}{\delta A_{\mu}^{a}} + j_{\mu}^{a}(B_{i}).
$$
\n(22)

and eq. [\(21\)](#page-4-1) gives

$$
D^{\nu}F^{a}_{\nu\mu} = j^{a}_{\mu}(\psi) + j^{a}_{\mu}(B) - G^{a}_{\mu}.
$$
 (23)

Now, the matter current satisfies a covariant conservation law and, by assumption so does also $j^a_\mu(B_i)$:

$$
D^{\mu}j_{\mu}^{a}(B_{i})=0.
$$

Then, since $D^{\nu}D^{\mu}F_{\nu\mu}^{a}=0$, eq. [\(21\)](#page-4-1) gives

$$
D^{\mu}G_{\mu}^{a}=0.\t\t(24)
$$

D. Solutions of the equations $\partial^{\mu}G_{\mu}^{a}=0$, $D^{\mu}G_{\mu}^{a}=0$

Given a solution G^a_μ of the conservation law $\partial^\mu G^a_\mu = 0$, the additional covariant conservation law requires

$$
T_{bc}^a (A^b)^{\mu} G_{\mu}^c = 0.
$$
 (25)

The vanishing of the left hand side may be due either to the summation over the Lorentz indices or to the summation over the G -group indices b, c (for each Lorentz index μ).

We have already shown that in the non-abelian case the equation $\partial^{\mu} G_{\mu}^{a} = 0$ does not have Lorentz covariant solutions fulfilling our requirements. One can see that in the non-abelian case also eq. [\(25\)](#page-5-0) does not admit Lorentz covariant solutions.

In fact, the invariance of the solutions under the corresponding residual group requires

$$
T^a_{bc} \,\partial^\mu \varepsilon^b_r(x) \, G^c_\mu = 0.
$$

Since the parameters $\partial^{\mu} \varepsilon_r(x)$ span a Lorentz covariant vector subspace and no such subspace exits except the whole Lorentz covariant vector space,

$$
(v^b)^\mu T^a_{bc} G^c_\mu = 0, \tag{26}
$$

for any Lorentz vector v^b_μ . Therefore $G^c_\mu = 0$ and $\mathcal{G}_r = \mathbb{1}$.

Thus, one must look for solutions which violate Lorentz covariance, still keeping the conditions i)-iv).

Now, the covariance of eq. [\(3\)](#page-1-1) under the unbroken global group requires that the Gauss operator must transform according to the adjoint representation.

Therefore, from a group theoretical point of view, in terms of the (irreducible) adjoint representation of $SU(N)$, eq. [\(24\)](#page-5-1) may be written in the form of scalar product

$$
(G,T^aA)=0,
$$

where A and G are vectors of the (irreducible) adjoint representation.

Now, in an irreducible representation given any vector by applying the generators T^a and the identity one reaches any other vector; therefore the above equation requires $G \sim A$.

Thus, the solutions of eq. [\(24\)](#page-5-1) are $G^a_\mu = A^a_\mu B$, with B an operator invariant under G transformations. This is easily obtained in the case $G = SU(2)$, since then eq. [\(24\)](#page-5-1) requires the vanishing of the vector product $\mathbf{A}_{\mu} \wedge \mathbf{G}_{\mu}$.

A very relevant example is provided by the Gauss operator of the temporal gauge, characterized by G_i^a = $0, \, \partial_0 G_0^a = 0.$

It is uniquely selected by the condition of rotational covariance of G^a_μ . In this case if $v^b_\mu = (v^b_i, v^b_0)$ is an allowed vector, so are the space-rotated ones $Rv_{\mu}^{b} = ((Rv^{b})_{i}, v_{0}^{b})$ as well as $Rv_{\mu}^{b}-v_{\mu}^{b}=((Rv^{b})_{i}-v_{i}^{b},0)$ for any rotation R. The set of such vectors spans the whole three-dimensional space and therefore eq. [\(26\)](#page-5-2) implies $G_i^c = 0$.

Its explicit form is obtainable by the gauge fixing term $\mathcal{L}_{GF}=A_0^a\,B^a,$ with B^a a Nakanishi-Lautrup field which plays the role of a Lagrange multiplier leading to $A_0^a = 0$. Then, one has

$$
G_{\mu}^{a} = \delta \mathcal{L}_{GF} / \delta A_{\mu}^{a} = \delta_{\mu,0} B^{a}, \qquad (27)
$$

and eq. [\(24\)](#page-5-1) is obviously satisfied since $G_i^a = 0$ and $A_0^a =$ 0.

In this case, the residual local covariance group of eq. [\(3\)](#page-1-1) is given by time independent gauge parameters, which characterize the *temporal gauge*. Indeed, the con-straint of eq. [\(9\)](#page-3-0) is satisfied by unrestricted $\partial_i \varepsilon^a$, which requires $\delta \mathcal{L}_{GF} / \delta A_i^a = 0$, and by $\partial_0 \varepsilon^a = 0$, which allows for $\delta \mathcal{L}_{GF}/\delta A_0^a \neq 0$.

This realization of the temporal gauge is particularly convenient because it directly provides the violation of the Local Gauss Law in terms of the time independent Nakanishi-Lautrup local field B^a . It is one of the basic fields which generate the local field algebra and it does not arise as a composite field with the related problem of point splitting regularization; thus, its algebraic properties are derivable from canonical quantization.

A similar pattern is realized by the axial gauge or by variations of it, like e.g. the $n^{\mu} A_{\mu}^{a} = 0$ gauges, with n_{μ} timelike, spacelike or lightlike. In this case, rotational covariance is replaced by the covariance under the stability group of n_{μ} .

III. CONCLUSION

We have shown that it is impossible to quantize Yang-Mills gauge theories with the properties of the Feynman quantization of Quantum Electrodynamics, namely with a Gauss operator which is Lorentz covariant and preserves a residual subgroup of local gauge transformations. If the covariance condition is relaxed, one may obtain a classification of the Gauss operators, which satisfy (gauge) covariant conservation laws and generate nontrivial subgroups of local gauge transformations.

It is worthwhile to remark that most of the nonperturbative results on gauge field theories (see the discussion in [\[1\]](#page-6-0)). have been obtained by using such a class of Gauss operators, thanks to their properties of a) preserving locality, b) leaving a residual local gauge group unbroken (helpful for renormalization), whose topology plays a crucial role in solving the $U(1)$ problem and c) satisfying a covariant conservation law, which allows for a simple control of its action on the matter fields. Moreover, the vacuum representation of the correspondent field algebra satisfies positivity and therefore has a Hilbert space structure.

Thus, on one side our results provide a general argument for justifying the recourse to Becchi-Rouet-Stora-Tyutin (BRST) quantization in non-abelian Yang-Mills theories in order to have manifest Lorentz covariance and locality, even if the residual group is not a subgroup of local gauge transformations and involves transformations with ghost fields as group parameters, violating positivity. On the other side it adds further support for the distinguished role of the temporal gauge, which proved very useful for a non-perturbative approach and results in non-abelian gauge theories.

IV. ACKNOWLEDGMENTS

B.B. is partially supported by INFN Iniziativa Specifica TPPC.

- [1] F. Strocchi, An Introduction to Non-Perturbative Foundations of Quantum Field Theory, Oxford University Press 2013, 2016.
- [2] R. Haag, Local Quantum Physics, Springer, Berlin-Heidelberg, 2nd ed., 1996.
- [3] F. Strocchi, Local Gauss law and local gauge symmetries in QFT, [arXiv:2301.06543v](http://arxiv.org/abs/2301.06543)2 [hep-th]; published in Trails in Theoretical and Mathematical Physics, edited by A. Cintio and A. Michelangeli eds., Springer 2024.
- [4] S. Weinberg, The Quantum Theory of Fields, Vol. II, Cambridge University Press 1996.
- [5] W. Magnus, Commun. Pure Appl. Math. 7, 649, (1954).
- [6] R.M. Wilcox, J. Math. Phys., 8, 962 (1967).