Thermal Bekenstein-Hawking entropy from the worldsheet

Indranil Halder and Daniel L. Jafferis

Jefferson Physical Laboratory, Harvard University, Cambridge, MA 02138, USA

ABSTRACT: We define and compute the leading sphere diagram contribution to the entropy of the BTZ black hole supported by Kalb-Ramond flux in bosonic string theory. In a winding condensate description, integrating exactly over the constant mode for the radial direction of AdS_3 reduces the problem to one of the correlation functions of winding operators in the free theory. The volume of the residual $PSL(2,\mathbb{C})$ gauge group of the sphere is canceled by the action of conformal transformations on the winding interaction insertions. We formulate a precise version of the replica trick in terms of (infinitesimally) non-integer winding condensates to produce the entropy of the BTZ black hole. The resulting entropy can be calculated from the one-point function of a non-local operator on the worldsheet.

¹ihalder@g.harvard.edu

²jafferis@g.harvard.edu

Contents

1	Introduction		1
2	Review of the duality		4
	2.1	Sigma model description	4
	2.2	Winding condensate description	5
3	Thermal free energy		7
	3.1	Non-zero modes on AdS_3	8
		3.1.1 Radial constant mode	8
		3.1.2 Wick contractions	8
	3.2	The divergent factor	11
	3.3	The finite factor	12
	3.4	Thermal entropy	14
4	Stringy replica trick		15
	4.1	Understanding the zero	15
	4.2	Getting rid of the zero	16
	4.3	The area operator	20
5	5 Summary and future directions		20
\mathbf{A}	A Three point function at zero temperature		23

1 Introduction

Entropies in gravitational theories have a deep significance as their leading order value in weakly gravitating system is encoded in classical geometry via the Bekenstein-Hawking [1, 2] and Hubeny-Rangamani-Ryu-Takayanagi [3, 4] area formulas. These can be derived in the long-distance effective theory, explaining their universality, using thermodynamic relations such as the Euclidean free energy $[5, 6]^1$ and the replica trick [10] ²calculations. It is therefore of great interest to find the string theory

¹Alternatively, the Bekenstein-Hawking entropy can be calculated using the method of orbifolds [7, 8]. Further gravitational loop corrections were found Sen [9].

²In this paper we need only the leading order in small g_s version of Lewkowycz-Maldacena [10]. The first quantum correction was found by Faulkner-Lewkowycz-Maldacena [11]. For further sub-leading order corrections and current status see the work of Engelhardt-Wall [12].

generalization of the area-entropy relation. This is the question we will address for the BTZ black hole - what is the α' exact analog of the area operator on the worldsheet?³

There are two conceptual hurdles to overcome in attempting to perform the Gibbons-Hawking-York Euclidean black hole calculation of the entropy in string theory. First, our interest is in the leading term in the (logarithm of the) partition function, of order g_s^{-2} , which comes from the genus zero diagram with no insertions. This corresponds to the on-shell action of the target space theory. This sphere diagram is notoriously difficult to define, due to the unfixed PSL(2, \mathbb{C}) gauge group⁴ (more so because there is no natural regularization that assigns a finite volume to PSL(2, \mathbb{C})⁵) and the infinite volume of the zero mode integration over the target space.⁶ Furthermore, string theory, even in its formulation as string field theory, is defined with respect to a background, so the only well-posed question appears to be the difference in the on-shell action of one background with respect to another.

We resolve these issues by using an exact dual description of the BTZ worldsheet theory in terms of an Euclidean time winding condensate on a free field background [20-22]. This description is strongly coupled in the large k limit, however, we can perform the calculations exactly. Note that these backgrounds have the same asymptotics, so they differ by a finite normalizable deformation (assuming the length of AdS in the strings unit is large⁷). The free linear dilaton background provides the reference point for the genus zero calculation, which can be performed using the standard technique of analytic continuation away values with integer numbers of insertions of the winding interaction vertex operators. The residual conformal transformations can now be fixed by localizing three of those insertions.

The second conceptual point is that the result of the above computation in

⁵The partition function on the disc is also plagued by similar issues of unfixed gauge modes. In this case, Polchinski showed how to get a sensible finite answer for the volume of the residual unfixed gauge group using a suitable regularization [15]. For a re-derivation of D-brane tension [16] using this result see [17].

⁶The problem of unfixed gauge degrees of freedom also appears in on-shell two or lower-point amplitudes of string theory. In the context of the two-point amplitude in flat-space string theory, the authors of [18] showed that the divergence from the unfixed gauge modes is canceled by the IR divergence of zero modes in target space to produce a sensible result (a similar conceptual point was speculated in [19] for the one-point function of the central charge operator in AdS₃. A non-trivial $l_{AdS}/l_s = \sqrt{k-2}$ dependent map between the IR cut-off in the target space and the regulator used to define the volume of PSL(2, \mathbb{C}) is proposed).

⁷More precisely we are considering k > 3.

³More precisely we are looking at string compactifications of the type BTZ×M, where M is an compact internal manifold. To have a tractable description of the worldsheet in the NSR formalism we will consider the bosonic string analog of the setup in [13] with pure NS-NS flux.

⁴In a beautiful recent work [14], the zero temperature partition function in AdS₃ is analyzed. In their approach, a new parameter μ is introduced in the worldsheet CFT. The triple derivative of the partition function with respect to μ is studied, thereby avoiding the issue of unfixed gauge modes.

fact vanishes. This corresponds to the fact that the bulk on-shell action of string field theory is zero [23] by the dilaton theorem [24] (considerations of a boundary in spacetime is much more subtle [8, 25, 26], see [27] for a review), and the our worldsheet calculation does not add any boundary terms (like the Gibbons-Hawking-York term, these boundary terms are responsible for the nonzero result in spacetime gravity, the Einstein-Hilbert bulk term can be written (up to total derivatives) in terms of the β functions of the worldsheet sigma model which vanishes on-shell to all orders in α' [28–31]).

Therefore, we turn to the Lewkowycz-Maldacena method of computing the entropy from analytic extrapolation of the Z_n quotient of the n^{th} Renyi geometries, which only requires knowledge of the bulk action. In semi-classical gravity, the result is that the entropy can be extracted from the change in the bulk action upon the addition of an infinitesimal conical defect in the interior [32–36] (see also [37]), keeping fixed the asymptotic geometry. The string theory analog is an analytic continuation of the worldsheet theories corresponding to a Z_n orbifold in the interior and the same Euclidean black hole asymptotics.

This is somewhat related to the proposal of Susskind-Uglum [34], utilizing the off-shell worldsheet formalism of [28, 29]. Here one is instructed to take a derivative of the off-shell worldsheet non-linear sigma model partition function with respect to the renormalization group UV cut-off to produce the string theoretic partition function (for a recent discussion of it see [38, 39]). The method relies on the details of the perturbative renormalization group flow and the simplest way of performing the path integral produces the bulk term in the target space, order by order in α' expansion. In this paper, we will produce exact in α' results without dealing with renormalization group flow.

The conical deformation we will use corresponds, in the winding condensate description, to a background obtained by adding a non-integer winding operator to the free theory. Such operators are not mutually local with the properly quantized Matsubara frequency vertex operators, but they are local amongst themselves, so the sphere diagram remains well-defined. Moreover, this conical defect-inducing operator is weight (1, 1) in the free worldsheet theory.

One might worry that such a (1, 1) deformation would correspond to the inclusion of a physical source for the curvature of the conical defect, which would result in a cancelation of its contribution to the action (at the linear order relevant for the entropy). However, the non-integer winding deformation that we use precisely does not include the reflected wave component in the radial direction. Including the reflected wave would result in a deformation that was not normalizable at the AdS boundary, and not including it corresponds to taking a singular conical bulk geometry without a compensating brane, as desired.

The worldsheet sphere amplitude associated with this deformed background is non-vanishing, and we compute it with the same free field methodology. From this, we find the g_s^{-2} leading order contribution to the BTZ entropy in string theory, exact in α' , which in $\alpha' \to 0$ limit reproduces the expected result. The derivative of the winding operator with respect to the non-integer replica number gives a non-local worldsheet operator that is the stringy generalization of the area.

Beyond the leading order in string coupling the thermal entropy gets a contribution both from the one-point function of the stringy area operator mentioned above and the entanglement entropy of the graviton and the other bulk fields outside the horizon. A systematic discussion of the finite value of entanglement entropy of the bulk fields at one loop including all stringy corrections can be obtained by the 'extrapolated' (order of the orbifold is fractional) Dabholkar-Witten orbifold method in [40, 41] (see also [42]).

The paper is organized as follows. In section 2 we review the sigma model and the winding condensate description of the worldsheet CFT. Then in section 3 we calculate the sphere path integral for BTZ carefully separating the divergent and finite parts. In section 4 we formulate the replica trick on the worldsheet. We conclude by discussing several open questions in section 5.

2 Review of the duality

In this paper, we will consider bosonic strings on $AdS_3 \times M$, M is a compact manifold, supported purely by Kalb-Ramond flux (the left and right central charges of the holographic dual CFT₂ are taken equal to each other). We will focus our discussion on the worldsheet CFT for the AdS₃ part. In subsection 2.1 we review the standard description of the worldsheet CFT in terms of WZW model. In subsection 2.2 we review the winding condensate description developed in [20, 21]. Throughout this section, we will follow the notation and the conventions used in [21].

2.1 Sigma model description

The usual sigma model description of the worldsheet CFT in Poincare patch⁸ takes the following form

$$S_{\text{sigma}} = \frac{k}{2\pi} \int 2d^2\sigma \left(\partial\hat{\phi}\bar{\partial}\hat{\phi} + \bar{\partial}\hat{\Gamma}\partial\bar{\hat{\Gamma}}e^{2\hat{\phi}} \right)$$
(2.1)

The Kalb-Ramond flux k determines the radius of AdS in string units

$$l_{AdS} = \sqrt{k - 2l_s} \tag{2.2}$$

For the rest of the paper, unless otherwise noted, we choose to work with the units set by $l_s = 1$. The sigma model description in (2.1) is exact at leading order in

⁸The metric in our convention is given by $ds^2 = l_{AdS}^2 (d\hat{\phi}^2 + e^{2\hat{\phi}} d\hat{\Gamma} d\hat{\bar{\Gamma}}).$

the large k limit. It is well known that for finite values of k, the (Euclidean target) worldsheet CFT is given by the $SL(2,\mathbb{C})_k/SU(2)$ WZW model.

The most convenient way to represent the simplest set of vertex operators in the large k limit is the following:⁹

$$V_{j,m,\bar{m}}(z,\bar{z}) = \int d^2x \ x^{j+m} \bar{x}^{j+\bar{m}} \Phi_{-1-j}(x,\bar{x}|z,\bar{z})$$

$$\Phi_j(x,\bar{x}|z,\bar{z}) = -\frac{2j+1}{\pi} \left(|\hat{\Gamma}(z,\bar{z}) - x|^2 e^{\hat{\phi}(z,\bar{z})} + e^{-\hat{\phi}(z,\bar{z})} \right)^{2j}$$
(2.3)

The function $\Phi_j(x, \bar{x})$ satisfies the massive scalar Laplace equation in the target space

$$(\nabla^2 - M^2)\Phi_j(x, \bar{x}|z, \bar{z}) = 0, \quad M^2 l_{AdS}^2 = (2j+2)(2j)$$
 (2.4)

with the following boundary condition

$$\lim_{\hat{\phi} \to \infty} e^{\hat{\phi}(2j+2)} \Phi_j(x, \bar{x}|z, \bar{z}) = \delta(\hat{\Gamma} - x) \delta(\hat{\bar{\Gamma}} - \bar{x})$$
(2.5)

These properties dictate Φ_j represent a field of mass M [44, 45] (see Appendix A for important details on proper normalization of vertex operators).

For the purpose of this work will not need to look into more general vertex operators, see [43-52] and [53-56] for more details.

2.2 Winding condensate description

Thermal AdS_3 has a dual description obtained by deformation of a free theory by a winding condensate on the spatial circle [20, 21]

$$S_{\text{TAdS}} = \frac{1}{2\pi} \int 2d^2 \sigma (\partial \varphi \bar{\partial} \varphi + \frac{1}{4b'} \varphi R + \beta \bar{\partial} \gamma + \bar{\beta} \partial \bar{\gamma} + \frac{\pi \mu}{2b'^2} (V^+ + V^-))$$
(2.6a)

$$V^{\pm} := e^{\pm \frac{k}{4}(\gamma + \bar{\gamma})} e^{\mp (\int_{(0,0)}^{(z,\bar{z})} \beta dz' + \int_{(0,0)}^{(z,\bar{z})} \bar{\beta} d\bar{z}')} e^{b'\varphi}, \quad \mu := 2b'^4 \left(\frac{\mu'\gamma(b'^2)}{\pi}\right)^{1/2}, \qquad (2.6b)$$

$$b' := \frac{1}{b''}, \quad \pi \mu' \gamma(b'^2) := (\pi \mu'' \gamma(b''^2))^{1/(b''^2)}, \quad b'' = \frac{1}{\sqrt{k-2}}, \quad \mu'' = \frac{b''^2}{\pi^2}$$
(2.6c)

The geometric meaning of the fields becomes apparent in large k limit: $\gamma = \hat{\xi} + i\hat{\theta}$, $\varphi = -\sqrt{k}\hat{r}$ (β is an auxiliary field that keeps track of the amount of winding in space circle $\hat{\theta}$), where $\hat{r}, \hat{\xi}, \hat{\theta}$ are standard global co-ordinates. At leading order in large k the metric and the Kalb-Ramond field are given by

$$ds^{2} = l_{AdS}^{2}(\sinh^{2}(\hat{r})d\hat{\theta}^{2} + \cosh^{2}(\hat{r})d\hat{\xi}^{2} + d\hat{r}^{2})$$

$$B = i l_{AdS}^{2}\sinh^{2}(\hat{r}) \ (d\hat{\theta} \otimes d\hat{\xi} - d\hat{\xi} \otimes d\hat{\theta})$$
(2.7)

⁹There is a sign different between the convention here and in [43]: $\Phi_{-1-j}(\text{here}) = -\Phi_j(\text{there}).$

Putting this background at temperature T, as seen by the spacetime CFT_2 living on the boundary of thermal AdS_3 , amounts to the following identification of coordinates

$$\hat{\xi} \sim \hat{\xi} + (Tl_{AdS})^{-1}, \quad \hat{\theta} \sim \hat{\theta} + 2\pi$$
 (2.8)

Vertex operators of relevance are

$$\tilde{V}_{\alpha,a,\bar{a}} = \frac{1}{\pi} e^{a\gamma + \bar{a}\gamma} e^{2\alpha\varphi}, \qquad (2.9)$$

They are dual to the vertex operators in (2.3)

$$\tilde{V}_{\alpha,a,\bar{a}} \leftrightarrow V_{j,m,\bar{m}}, \quad \alpha = -\frac{j}{2b}, \quad a = m - \frac{k\nu}{4}, \quad \bar{a} = \bar{m} - \frac{k\nu}{4}$$
(2.10)

These vertex operators carry no winding in either the space or time circle.

The focus of this paper is string theory around the Euclidean BTZ black hole. It can be described by the background given in (2.7) with a different identification¹⁰

$$\hat{\xi} \sim \hat{\xi} + 4\pi^2 (T l_{AdS}), \quad \hat{\theta} \sim \hat{\theta} + 2\pi$$
 (2.11)

In EBTZ background the winding around the temporal circle $\hat{\xi}$ is not conserved. This is not apparent in this presentation. This feature becomes visible in an equivalent description of EBTZ at temperature T obtained by exchanging the roles of $\hat{\xi}, \hat{\theta}$ and switching the holomorphic and anti-holomorphic coordinates on the worldsheet (needed to keep the same B field)

$$ds^{2} = l_{AdS}^{2}(\cosh^{2}(\hat{r})d\hat{\theta}^{2} + \sinh^{2}(\hat{r})d\hat{\xi}^{2} + d\hat{r}^{2})$$

$$B = -il_{AdS}^{2}\sinh^{2}(\hat{r}) \ (d\hat{\theta} \otimes d\hat{\xi} - d\hat{\xi} \otimes d\hat{\theta})$$

$$\hat{\theta} \sim \hat{\theta} + 4\pi^{2}(Tl_{AdS}), \quad \hat{\xi} \sim \hat{\xi} + 2\pi$$
(2.12)

The worldsheet theory in this background is given by¹¹

$$S_{\text{EBTZ}} = \frac{1}{2\pi} \int 2d^2 \sigma (\partial \varphi \bar{\partial} \varphi + \frac{1}{4b'} \varphi R + \beta \bar{\partial} \gamma + \bar{\beta} \partial \bar{\gamma} + \frac{\pi \mu}{2b'^2} (W^+ + W^-)) \qquad (2.13a)$$
$$W^{\pm} - e^{\pm i \frac{k}{4} (\gamma - \bar{\gamma})} e^{\pm i (\int_{(0,0)}^{(z,\bar{z})} \beta dz' - \int_{(0,0)}^{(z,\bar{z})} \bar{\beta} d\bar{z}')} e^{b'\varphi} \qquad (2.13b)$$

w =
$$e^{-4(0.6)}$$
 $e^{-4(0.6)}$ $e^{-4(0.6)}$ $e^{-4(0.6)}$ (2.13b)
upling constant μ is still given by its expression in (2.6). Geometrically we

The coupling constant μ is still given by its expression in (2.6). Geometrically we still identify $\gamma = \hat{\xi} + i\hat{\theta}, \hat{\theta} \sim \hat{\theta} + 4\pi^2 (Tl_{AdS}), \hat{\xi} \sim \hat{\xi} + 2\pi$ as a result W^{\pm} above carries non-trivial winding on the temporal circle $\hat{\xi}$ as opposed to V^{\pm} which carries non-zero winding on the spatial circle $\hat{\theta}$.

¹⁰Note that trying to convert this identification to the one in (2.8) while keeping the background metric and B field fixed through field redefinition is a non-trivial exercise because in the process one generates a non-zero B field on the boundary torus.

¹¹This is obtained from (2.6) by the replacement $\gamma \to i\bar{\gamma}, \beta \to -i\hat{\beta}, z \to \bar{z}$ along with their complex conjugates. While presenting (2.12) we dropped hats on variables.

3 Thermal free energy

The goal of this section is to calculate the leading order in g_s partition function of string theory in the EBTZ background staying entirely within the framework of on-shell string theory.

We will proceed to evaluate the partition function of the worldsheet CFT on unit sphere. This includes AdS_3 , compact manifold M and ghosts, such that the total central charge vanishes. This ensures that the partition function of the worldsheet CFT is well-defined and independent of the radius of the sphere [57].

$$Z_{CFT}((Tl_{AdS})^{-1}, l_{AdS}/l_s) = Z^M Z^{AdS_3} Z_c$$
(3.1)

The sphere partition function of the worldsheet CFT with target space M is given by Z^M . We will not discuss the evaluation of Z^M , except the assumption that the integration of compact zero modes on M is normalized to give a factor of volume V_M in string units, and contribution of non-zero modes are sub-leading in large k limit, more precisely

$$Z^{M} = Z_{0}^{M} C_{S^{2}}^{M}, \quad Z_{0}^{M} = V_{M}, \quad C_{S^{2}}^{M} = 1 + \mathcal{O}\left(\frac{1}{k}\right)$$
(3.2)

The AdS₃ part of the partition function is defined with the measure as in [21] - the normalization of the measure on Liouville-like field φ is the same as in [58], and the measure of the non-zero modes of the $\beta\gamma$ system is same as in [59]. To understand the zero mode structure of the $\beta\gamma$ system we note that the free $\beta\gamma$ system is a type of 'toric' sigma model whose Hilbert is analyzed in section 5 of [60]. After we integrate out the radial constant mode we will be left with the free $\beta\gamma$ system (see next subsection for explanation). The zero mode contribution from translation in $\hat{\xi}, \hat{\theta}$ gives

$$Z_0^{AdS_3} = 8\pi^3 (Tl_{AdS}) \tag{3.3}$$

The non-zero mode $\gamma', \bar{\gamma'}$ contributes as

$$C_{S^2}^{AdS_3} = \int d\varphi d\beta' d\gamma' d\bar{\beta'} d\bar{\gamma'} e^{-S_{\text{EBTZ}}}$$
(3.4)

These two factors $C_{S^2}^{AdS_3}$, $Z_0^{AdS_3}$ together give Z^{AdS_3} . The remaining k, β independent factor Z_c , taking care of the contribution of b, c ghosts (omitting the ghost zero modes).

The same factors Z^M , Z_c appear in the calculation of the effective three-dimensional Newton's constant. Therefore when the entropy of the BTZ black hole is expressed in terms of the three-dimensional Newton's constant these factors disappear. For this reason we will not be determine them in this paper explicitly.

3.1 Non-zero modes on AdS₃

3.1.1 Radial constant mode

First we focus on the radial constant mode φ_c defined by

$$\varphi(z,\bar{z}) = \varphi_c + \varphi'(z,\bar{z}) \tag{3.5}$$

where φ_c denotes the kernel of the scalar Laplacian and $\varphi'(z, \bar{z})$ is the space of functions orthogonal to the kernel, using the following identity [58, 61–63]

$$\int_{-\infty}^{+\infty} d\varphi_c \ e^{2a\varphi_c - \alpha e^{2b\varphi_c}} = \frac{1}{2b} \Gamma\left(\frac{a}{b}\right) \alpha^{-\frac{a}{b}} \tag{3.6}$$

We can integrate out φ_c exactly to get

$$C_{S^{2}}^{AdS_{3}} = \frac{2\pi}{b'} \Gamma\left(-2s'\right) \left(\frac{\mu}{2b'^{2}}\right)^{2s'} \int d\varphi' d\beta' d\gamma' d\bar{\beta}' d\bar{\gamma'} e^{-S_{\text{EBTZ}}|_{\mu=0}} \left(\int d^{2}z \left(W^{+}(z,\bar{z}) + W^{-}(z,\bar{z})\right)\right)^{2s'} (3.7)$$

Here we have defined

$$s' = \frac{1}{b'^2}$$
(3.8)

3.1.2 Wick contractions

After integrating out the radial constant mode the leftover path integration is very simple due to winding and momentum conservation - see [21] for more details. Essentially only a very specific expression of the winding condensate contributes to the path integral when s' is a positive integer.¹² For this special case, we perform the Wick contractions on (3.7), to write ¹³

$$C_{S^{2}}^{AdS_{3}} = \frac{2\pi}{b'} \Gamma(-2s') \left(\frac{-\mu}{2b'^{2}}\right)^{2s'} \frac{\Gamma(2s'+1)}{\Gamma(s'+1)^{2}} \int \prod_{i=1}^{s'} d^{2}z_{i} \int \prod_{i'=1}^{s'} d^{2}z_{i'} \prod_{i'< j'} \left[(z_{i'j'})(\bar{z}_{i'j'}) \right] \prod_{i< j} \left[(z_{ij})(\bar{z}_{ij}) \right] \prod_{i,j'} \left[(z_{ij'})^{1-k}(\bar{z}_{ij'})^{1-k} \right]$$

$$(3.9)$$

 12 We remind the reader that

$$k = 2\left(1 + \frac{1}{s}\right)$$

Therefore the calculation done for the residue is valid for small k. This is no surprise given that the winding condensate description is best suited for calculation in this range. This in particular implies k is not an integer. This is compatible even with compact internal manifolds, for example arising from the compactification of M theory on a Calabi-Yau fourfold singularity (see for instance [64]).

¹³The skeptical reader might ask for the justification of not including a factor corresponding to the partition function of the linear dilaton and free $\beta\gamma$ (and complex conjugate) system at this step. The contribution of the non-zero modes of $\beta\gamma$ system is independent of k, β , so it is assumed to be absorbed in Z_c . For the linear dilaton note that the curvature coupling on unit radius is only through the zero mode as a result the contribution is again independent of k. Here $z_{ij} = z_i - z_j$, $z_{ij'} = z_i - z'_{j'}$, $z_{i'j'} = z'_{i'} - z'_{j'}$. Before going forward we notice that if we change variables as

$$z_i \to \lambda z_i, \quad \bar{z}_i \to \lambda \bar{z}_i, \quad z'_i \to \lambda z'_i, \quad \bar{z}'_i \to \lambda z'_i,$$

$$(3.10)$$

Then all the factors of λ cancel out since s'(k-2) = 1

$$\lambda^{2s' \times 2 + s'(s'-1)/2 \times 4 + 2(1-k)s'^2} = \lambda^{4s'+2s'(s'-1)-2s'^2(k-1)}$$
(3.11)

Thus we have effective scale invariance even after integrating out the radial constant mode.

We single out $z_a, a = 1, 2, z'_{a'}, a = 1$ (this process works only for s' = 2, 3, 4, ...we focus on this particular case) to write the above expression as

$$C_{S^{2}}^{AdS_{3}} = \frac{2\pi}{b'} \Gamma(-2s') \left(\frac{-\mu}{2b'^{2}}\right)^{2s'} \frac{\Gamma(2s'+1)}{\Gamma(s'+1)^{2}} \int \prod_{a=1}^{2} d^{2}z_{a} \prod_{I=3}^{s'} d^{2}z_{I} \int \prod_{a'=1} d^{2}z'_{a'} \int \prod_{I'=2}^{s'} d^{2}z'_{I'} |z_{12}|^{2} |z_{11'}|^{2(1-k)} |z_{21'}|^{2(1-k)} \prod_{a} \left[(\prod_{I} |z_{aI}|^{2}) (\prod_{I'} |z_{aI'}|^{2(1-k)}) \right] \prod_{a'} \left[(\prod_{I} |z_{a'I}|^{2(1-k)}) (\prod_{I'} |z_{a'I'}|^{2}) \right] \prod_{I'(3.12)$$

We make the following transformations (and their conjugates) in the given order below

$$z_{I} \to z_{I} + z_{1}, z'_{I'} \to z'_{I'} + z_{1}$$

$$z_{I} \to z_{21} z_{I}, z'_{I'} \to z_{21} z'_{I'}$$
(3.13)

Due to the scale invariance discussed above

$$C_{S^{2}}^{AdS_{3}} = \frac{2\pi}{b'} \Gamma(-2s') \left(\frac{-\mu}{2b'^{2}}\right)^{2s'} \frac{\Gamma(2s'+1)}{\Gamma(s'+1)^{2}}$$

$$\int \prod_{a=1}^{2} d^{2}z_{a} \int \prod_{a'=1} d^{2}z'_{a'} \prod_{I=3}^{s'} d^{2}z_{I} \int \prod_{I'=2}^{s'} d^{2}z'_{I'} \frac{|z_{12}|^{2}|z_{11'}|^{2(1-k)}|z_{21'}|^{2(1-k)}}{|z_{12}|^{6+2+2(1-k)+2(1-k)}}$$

$$\left[(\prod_{I} |z_{I}|^{2}|z_{I}-1|^{2}|z_{I}-x|^{2(1-k)}) (\prod_{I'} |z'_{I'}|^{2(1-k)}|z'_{I'}-1|^{2(1-k)}|z'_{I'}-x|^{2}) \right]$$

$$\prod_{I'< J'} [|z_{I'J'}|^{2}] \prod_{I< J} [|z_{IJ}|^{2}] \prod_{I< J} [|z_{IJ}|^{2}] \qquad (3.14)$$

Here we have defined $x = \frac{z_{1'1}}{z_{21}}$. Next we do the following change of co-ordinates for all I, i'

$$z_I = \frac{\tilde{z}_I x}{\tilde{z}_I + x - 1}, z'_{I'} = \frac{\tilde{z}'_{I'} x}{\tilde{z}'_{I'} + x - 1},$$
(3.15)

This transformation has a non-trivial Jacobian associated with it

$$dz_i = \frac{x(x-1)}{(z_i + x - 1)^2} d\tilde{z}_i$$
(3.16)

This transformation has the following properties

$$z_{I} - 1 = (\tilde{z}_{I} - 1) \frac{(x - 1)}{\tilde{z}_{I} + x - 1}, \quad z_{I} - x = \frac{-x(x - 1)}{\tilde{z}_{I} + x - 1}$$

$$z_{I} - z_{J} = (\tilde{z}_{I} - \tilde{z}_{J}) \frac{x(x - 1)}{(\tilde{z}_{I} + x - 1)(\tilde{z}_{J} + x - 1)}$$

$$z_{I} - z_{I'} = (\tilde{z}_{I} - \tilde{z}_{I'}) \frac{x(x - 1)}{(\tilde{z}_{I} + x - 1)(\tilde{z}_{I'} + x - 1)}$$
(3.17)

The change of variables becomes simple because the factor

$$\lambda = \frac{1}{z_I + x - 1} \tag{3.18}$$

for a given I drops out of the integration over z_I

$$\lambda^{4+2\times 2+2(1-k)+2(s'-3)+2(1-k)(s'-1)} = \lambda^{4+6-2k+2s'-6+2s'(1-k)-2(1-k)} = \lambda^{2+2s'(2-k))} = 1$$
(3.19)

(3.19) Similarly for each I' integration this factor drops out.¹⁴ As a result the integrations over $z_I, z'_{I'}$ does not involve any factor of $z_a, z'_{a'}$. We can present the result of the manipulation as

$$C_{S^2}^{AdS_3} = z_{div} \ z \tag{3.21}$$

where

$$z_{div} = \int \prod_{a=1}^{2} d^{2}z_{a} \int d^{2}z_{1'}' \frac{|z_{12}|^{2}|z_{11'}|^{2(1-k)}|z_{21'}|^{2(1-k)}}{|z_{12}|^{6+2+2(1-k)+2(1-k)}} \frac{1}{|x|^{2(2-k)}|1-x|^{2(2-k)}}$$

$$= \int \prod_{a=1}^{2} d^{2}z_{a} \int d^{2}z_{1'}' \frac{1}{|z_{12}|^{2}|z_{11'}|^{2}|z_{21'}|^{2}}$$
(3.22)

and

$$z = \frac{2\pi}{b'} \Gamma(-2s') \left(\frac{-\mu}{2b'^2}\right)^{2s'} \frac{\Gamma(2s'+1)}{\Gamma(s'+1)^2} \prod_{I=3}^{s'} d^2 z_I \int \prod_{I'=2}^{s'} d^2 z'_{I'} \\ \left[(\prod_I |z_I|^2 |z_I - 1|^2) (\prod_{I'} |z'_{I'}|^{2(1-k)} |z'_{I'} - 1|^{2(1-k)}) \right] \\ \prod_{I'(3.23)$$

¹⁴Essentially due to

$$\lambda^{4+2(1-k)\times 2+2+2(s'-2)+2(1-k)(s'-2)} = \lambda^{4+4(1-k)+2+2s'-4+2s'(1-k)-4(1-k)} = 1$$
(3.20)

In the next two subsections, we will show show how z_{div} gets cancelled against the volume of unfixed gauge group and simplify the expression of z in terms of Liouville theory.

3.2 The divergent factor

The divergent factor z_{div} can be identified with the volume of the unfixed residual gauge group PSL(2,C) (after fixing the conformal gauge). To show this we follow the footsteps of [18] and write zero temperature three-point functions in two different ways ¹⁵

$$C_{S^{2}}\langle\Phi_{j_{1}}(0|0)\Phi_{j_{2}}(1|1)\Phi_{j_{3}}(\infty|\infty)\rangle = \frac{C_{S^{2}}}{z_{PSL(2,C)}}\int d^{2}z_{1}d^{2}z_{2}d^{2}z_{3}\langle\Phi_{j_{1}}(0|z_{1},\bar{z}_{1})\Phi_{j_{2}}(1|z_{2},\bar{z}_{2})\Phi_{j_{3}}(\infty|z_{3},\bar{z}_{3})\rangle$$

$$(3.24)$$

Here C_{S^2} is the normalization factor for the sphere diagram at zero temperature given in (A.7). The expectation value on both sides is taken in AdS₃ sigma model, i.e., with the measure (note that the zero mode path integration is included)

$$\langle \dots \rangle = \frac{\int d\varphi d\beta d\gamma d\bar{\beta} d\bar{\gamma} e^{-S_{\rm EBTZ}} \dots}{\int d\varphi d\beta' d\gamma' d\bar{\beta}' d\bar{\gamma}' e^{-S_{\rm EBTZ}}}$$
(3.25)

On the LHS of (3.24) we fixed the position of three operators, thus we have inserted three factors of c, \bar{c} ghosts at those locations in the complete worldsheet path integral (see section 5.3 of [59] for more explanation). The three-point function of ghosts cancels the z_i, \bar{z}_i dependence of the matter three-point function just as in flat space and we are left with the three-point function of the matter vertex operators at three fixed points with the conformal factor involving z_i, \bar{z}_i stripped off. On the RHS of (3.24) we have not fixed the location of any vertex operator as a result there is no insertion of the ghost operator (and we are supposed to ignore the ghost zero modes in RHS¹⁶), however in this case we have divided by the explicit factor $z_{SL(2,C)}$ of the volume of the residual gauge group. This gives

$$z_{PSL(2,C)} = z_{div} \tag{3.26}$$

This is similar to the cancellation discussed in the context of Liouville theory in [65, 66].

¹⁵The actual on-shell vertex operators in the BRST cohomology of string theory are (1, 1) on the worldsheet made up of Φ_j multiplied with an operator on the internal manifold M. Once this is taken into account the final conclusion made through arguments presented here remains unchanged.

¹⁶Another equivalent way of saying it is as follows: we could include the ghost zero modes in RHS of (3.24) and then the RHS of (3.26) includes ghost zero mode contribution contribution. In this process we also include ghost zero modes in RHS of (3.4) and in effect the final result is unchanged.

3.3 The finite factor

Now we turn to simplify the finite part in terms of Liouville theory correlator. Note that in effect the finite factor is obtained by fixing location of three winding operator insertions after radial zero mode integration (just like ordinary string theoretic three point function, a careful consideration including b, c ghosts shows in effect the ghost zero modes are taken into account).

$$z = \frac{2\pi}{b'} \Gamma(-2s') \left(\frac{-\mu}{2b'^2}\right)^{2s'} \frac{\Gamma(2s'+1)}{\Gamma(s'+1)^2} \prod_{I=3}^{s'} d^2 z_I \int \prod_{I'=2}^{s'} d^2 z'_{I'} \\ \left[(\prod_I |z_I|^2 |z_I - 1|^2) (\prod_{I'} |z'_{I'}|^{2(1-k)} |z'_{I'} - 1|^{2(1-k)}) \right] \\ \prod_{I'(3.27)$$

We integrate out $z'_{I'}$ using the integral identity in appendix B of [21] with

$$n = s' - 1, \quad m = 0$$

$$t_j = z_j, \quad j = 1, \dots, s' - 2, \quad t_{s'-1} = 0, \quad t_{s'} = 1$$

$$p_j = 1 - k, \quad p_{s'-1} = (1 - k), \quad p_{s'} = 1 - k$$
(3.28)

We remind the reader that the expression for the finite factor here is only valid when s' = 2, 3, 4... (for s' = 1 we do not have enough winding operators to fix), resulting in the following formula for the residue

$$\operatorname{Res}_{s=2s'\to 2\mathbb{Z}} z = \frac{2\pi}{b'} \left(\frac{-\mu}{2b'^2}\right)^{2s'} \frac{1}{\Gamma(s'+1)} \frac{\pi^{s'-1}\Gamma(s')\gamma(2-k)^{s'}}{\gamma(s'(2-k))} \int \prod_{I=3}^{s'} d^2 z_I \prod_{I

$$(3.29)$$$$

We have written down the residue using Liouville theory three-point function (see appendix A for conventions)

$$C_{(b',\mu')}(\alpha_1,\alpha_2,\alpha_3) = \left[\pi\mu'\gamma(b'^2)b'^{2-2b'^2}\right]^{\frac{Q'-\sum_k\alpha_k}{b'}} \frac{\Upsilon_{b'}(0)\prod_{k=1}^3\Upsilon_{b'}(2\alpha_k)}{\Upsilon_{b'}(\sum_k\alpha_k-Q')\prod_{k=1}^3\Upsilon_{b'}(\sum_k\alpha_k-2\alpha_k)}$$
(3.30)

The $\Upsilon_{b'}(x)$ functions have zeros at

$$x = -mb' - nb'^{-1}, \quad x = Q' + mb' + nb'^{-1}, \quad m, n = 0, 1, 2, \dots$$
 (3.31)

$$\operatorname{Res}_{\sum_{i} \alpha_{i} = Q' - (s'-2)b'} C(\alpha_{1}, \alpha_{2}, \alpha_{3}) = b' \operatorname{Res}_{s' = 2 + \frac{Q' - \sum_{i} \alpha_{i}}{b'}} C(\alpha_{1}, \alpha_{2}, \alpha_{3})$$

$$= b' \operatorname{Res}_{s' = 2 + \frac{Q' - \sum_{i} \alpha_{i}}{b'}} \left[\pi \mu' \gamma(b'^{2}) b'^{2 - 2b'^{2}} \right]^{\frac{Q - \sum_{i} \alpha_{i}}{b'}} \frac{\Upsilon'_{b'}(0) \Upsilon_{b'}(2b')^{3}}{\Upsilon_{b'}(-(s'-2)b') \Upsilon_{b'}(b')^{3}}$$
(3.32)

Here we get a pole when s' = 2, 3, 4, ... from $\Upsilon_{b'}(-(s'-2)b')$. As discussed above there are precisely the values of s' where we also get a pole in z. We generalize this result for all values of s' as follows ¹⁷

$$z = (-1)^{2ls'} \left(\frac{-\mu}{2b'^2}\right)^{2s'} \frac{\pi}{(s'-1)(s')^2} \frac{\pi^{s'-1}\gamma(2-k)^{s'}}{\gamma(s'(2-k))} (-\mu')^{2-s'} \\ \left[\pi\mu'\gamma(b'^2)b'^{2-2b'^2}\right]^{\frac{Q-\sum_i\alpha_i}{b'}} \frac{\Upsilon'_{b'}(0)\Upsilon_{b'}(2b')^3}{\Upsilon_{b'}(-(s'-2)b')\Upsilon_{b'}(b')^3} \\ = (-1)^{2ls'} \left(\frac{-\mu}{2b'^2}\right)^{2s'} \frac{\pi}{(s'-1)(s')^2} \frac{\pi^{s'-1}\gamma(2-k)^{s'}}{\gamma(s'(2-k))} (-\mu')^{2-s'} \\ \left[\pi\mu'\gamma(b'^2)b'^{2-2b'^2}\right]^{s'-2} \frac{\Upsilon'_{b'}(0)\Upsilon_{b'}(2b')^3}{\Upsilon_{b'}(-(s'-2)b')\Upsilon_{b'}(b')^3} \\ = (-1)^{2ls'} (-1)^{s'} \left(\frac{\mu}{2b'^2}\right)^{2s'} \frac{\pi}{(s'-1)(s')^2} \frac{\pi^{2s'-3}\gamma(2-k)^{s'}}{\gamma(s'(2-k))} \\ \left[\gamma(b'^2)b'^{2-2b'^2}\right]^{s'-2} \frac{\Upsilon'_{b'}(0)\Upsilon_{b'}(2b')^3}{\Upsilon_{b'}(-(s'-2)b')\Upsilon_{b'}(b')^3} \\ (3.34)$$

Here l is an unfixed integer, since we are analytically continuing from integer values of s', l is not determined at this stage. Later by demanding reality of the resulting formula will determine l.

We can actually simplify the above expression a lot owing to the special argument of $\Upsilon_{b'}$. We use the following identities [67]

$$\Upsilon_{b'}(Q'-x) = \Upsilon_{b'}(x), \quad \Upsilon_{b'}(x) = \frac{\Upsilon_{b'}(x+b')}{\gamma(b'x)b^{1-2b'x}} = \frac{\Upsilon_{b'}(x+\frac{1}{b'})}{\gamma(\frac{x}{b'})\left(\frac{1}{b'}\right)^{1-\frac{2x}{b'}}}$$
(3.35)

First factor in (3.34) simplifies to (see (3.35))

$$\frac{\Upsilon_{b'}(2b')^3}{\Upsilon_{b'}(b')^3} = \left(\gamma(b'^2)b'^{1-2b'^2}\right)^3 \tag{3.36}$$

The other factor can be manipulated as follows

$$\Upsilon_{b'}(-(s'-2)b') = \Upsilon_{b'}(-s'b'+2b') = \Upsilon_{b'}\left(2b'-\frac{1}{b'}\right) = \gamma(b'^2-1)b'^{3-2b'^2}\Upsilon_{b'}\left(b'-\frac{1}{b'}\right)$$
$$= \gamma(b'^2-1)b'^{3-2b'^2}b'^3\gamma(-1)\Upsilon_{b'}\left(-\frac{1}{b'}\right)$$
(3.37)

 17 We used

$$\operatorname{Res}_{2s'\to 2\mathbb{Z}} z = 2\operatorname{Res}_{s'\to \mathbb{Z}} z \tag{3.33}$$

The last two factors are divergent and zero respectively. A proper limiting process can be obtained by using $s' = (k-2)^{-1} + \epsilon$

$$\gamma(-1) \to \gamma(-1 - \epsilon b'^2) = \frac{1}{b'^2 \epsilon}$$

$$\Upsilon_{b'} \left(-\frac{1}{b'}\right) \to \Upsilon_{b'}(-\frac{1}{b'} - \epsilon b') = \frac{\Upsilon_{b'}(-\epsilon b')}{\gamma(-b'^{-2})b'^{2b'^{-1}(-b'^{-1})-1}} = -\epsilon b' \Upsilon_{b'}'(0) \frac{b'^{\frac{2}{b'^2}+1}}{\gamma(-\frac{1}{b'^2})}$$
(3.38)

So in summary, we have

$$\Upsilon_{b'}(-(s'-2)b') = -\gamma(b'^2-1)b'^{6-2b'^2}\frac{1}{b'^2}b'\Upsilon'_{b'}(0)\frac{b'^{\frac{2}{b'^2}+1}}{\gamma(-\frac{1}{b'^2})} = -b'^{6-2b'^2+\frac{2}{b'^2}}\gamma(b'^2-1)\frac{1}{\gamma(-\frac{1}{b'^2})}\Upsilon'_{b'}(0)$$
(3.39)

Putting it back into (3.34) we get

$$z = \pi (-1)^{(2l+1)s'} \left(\frac{\mu}{2b'^2}\right)^{2s'} \frac{1}{(s'-1)(s')^2} \frac{\pi^{2s'-3}\gamma(2-k)^{s'}}{\gamma(s'(2-k))} \\ \left[\gamma(b'^2)b'^{2-2b'^2}\right]^{s'-2} \frac{\left(\gamma(b'^2)b'^{1-2b'^2}\right)^3}{-b'^{6-2b'^2+\frac{2}{b'^2}}\gamma(b'^2-1)\frac{1}{\gamma(-\frac{1}{b'^2})}} \\ = \frac{(-1)^{\frac{2l}{k-2}}}{\Gamma(-1)} \mu^{\frac{2}{k-2}} \frac{(-1)^{\frac{k}{k-2}}4^{\frac{1}{2-k}}(k-3)(k-2)^{\frac{3k+2}{4-2k}}\pi^{\frac{2}{k-2}-2}\Gamma\left(\frac{1}{2-k}\right)}{\Gamma\left(1+\frac{1}{k-2}\right)}$$
(3.40)

Now we substitute the value of cosmological constant from (2.6)

$$\mu = 2(k-2)^2 \pi^{-\frac{k}{2}} \left(-\frac{\Gamma\left(\frac{1}{k-2}\right)}{\Gamma\left(\frac{1}{2-k}\right)} \right)^{\frac{k-2}{2}}$$
(3.41)

Plugging it back into (3.40), and setting l = -1 to get rid of the complex phase we get

$$z = \frac{1}{\Gamma(-1)\pi^3} \left(\sqrt{k-2} - \frac{1}{\sqrt{k-2}} \right) = \frac{1}{\Gamma(-1)\pi^3} \sqrt{k} \left(1 - \frac{2}{k} + \mathcal{O}\left(\frac{1}{k^2}\right) \right) \quad (3.42)$$

Curiously, the dependence of z on k is exactly given by the slope of the linear dilaton of the dual conformal field theory on the boundary of AdS [68].

3.4 Thermal entropy

The string theoretic thermal free energy of BTZ black hole gets contributions from connected worldsheets only [69], thus it is determined by combining results from (3.3), (3.26), (3.42) to be

$$\log Z_{BTZ}(1/(Tl_{AdS}), l_{AdS}/l_s) = \frac{1}{g_s^2} \frac{Z_{CFT}(1/(Tl_{AdS}), l_{AdS}/l_s)}{z_{PSL(2,C)}}$$
(3.43)

Here g_s is the string coupling at the tip of the Euclidean cigar.¹⁸ To the leading order in large k we find

$$\log Z_{BTZ}(1/(Tl_{AdS}), l_{AdS}/l_s) = \frac{8Z_c}{\Gamma(-1)} \frac{V_M}{g_s^2} \frac{\sqrt{k}}{(Tl_{AdS})^{-1}} = \frac{8Z_c}{\Gamma(-1)} \frac{V_M}{g_s^2 l_s} l_{AdS}(Tl_{AdS})$$
(3.44)

This is precisely the formula for the entropy for BTZ black hole found from target space gravity analysis [13] (up to constant factors). We remind the reader that the radius of the horizon is related to the inverse temperature measured in AdS units $(Tl_{AdS})^{-1}$ according to

$$\frac{r_H}{l_{AdS}} = \frac{2\pi}{(Tl_{AdS})^{-1}} \tag{3.45}$$

The factor $\Gamma(-1)^{-1}$ is 0, and tells us that our method is insensitive to the boundary terms in the target space effective action. In the next section, we will carefully discuss the conceptual origin of the overall factor $\Gamma(-1)^{-1}$ in terms of the replica trick.

4 Stringy replica trick

In the previous section, we performed the calculation of the thermal entropy of the BTZ black hole using the on-shell worldsheet description and noticed that the final result is zero. In section 4.1 we will show that the technical reason for the factor of zero is a brunch cut becoming transparent. As a remedy in section 4.2 we will develop a precise version of the replica trick [10-12] in the language of non-integer winding condensates.

4.1 Understanding the zero

The integral identity in appendix B of [21] was used to obtain (3.29) that generated the factor of $\gamma(s'(2-k))^{-1}$. We turn to understand this factor of zero for the simplest case of the identity¹⁹

$$G(a,b) = \int d^2 z |z|^{2a} |1-z|^{2b} = \pi \frac{\gamma(1+a)\gamma(1+b)}{\gamma(2+a+b)}$$
(4.2)

 19 In the notation of [21], this corresponds to

$$n = 1, \ m = 0, \ t_1 = 0, t_2 = 1, p_1 = a, p_2 = b$$
 (4.1)

¹⁸In the BTZ geometry given by (2.12) the angular circle remains of the finite size at the tip, whereas the Euclidean time circle shrinks. When one takes into account the perturbative α' corrections to it and the effect of winding tachyon in near horizon geometry, the physical meaning of the tip might be more subtle [70].

We write z = x + iy, $x, y \in \mathbb{R}$ and analytically continue the integration over y as follows

$$y \to iye^{-2i\epsilon}, \ \epsilon \to 0+$$
 (4.3)

This gives

$$G(a,b) = -\frac{i}{2} \int_{-\infty}^{+\infty} dz_{+} (z_{+} - i\epsilon(z_{+} - z_{-}))^{a} (z_{+} - 1 - i\epsilon(z_{+} - z_{-}))^{b} \int_{-\infty}^{+\infty} dz_{-} (z_{-} + i\epsilon(z_{+} - z_{-}))^{a} (z_{-} - 1 + i\epsilon(z_{+} - z_{-}))^{b}$$

$$(4.4)$$

Here we defined $z_{\pm} = x \pm y$. Say $z_{+} \in (-\infty, 0)$, the contour of integration over z_{-} runs below the singularity at $z_{-} = 0, 1$. Provided the integration contour can be deformed away near infinity on the lower half-plane, the contribution is zero. A similar argument holds for $z_{+} \in (1, \infty)$ (in this case we can deform the contour in the upper half plane). For $z_{+} \in (0, 1)$, the contour of integration over z_{-} runs above the singularity at $z_{-} = 0$ and below at $z_{-} = 1$. We can deform the contour to run around $z_{-} \in (1, \infty)$ encircling the point at $z_{-} = 1$. If the integration over the circle around $z_{-} = 1$ does not contribute anything, we get

$$G(a,b) = -\sin(\pi b) \int_0^1 dz_+(z_+)^a (1-z_+)^b \int_1^\infty dz_-(z_-)^a (z_--1)^b$$

= $-\sin(\pi b) \int_1^\infty dz_+(z_+)^{-a-b-2} (z_+-1)^b \int_1^\infty dz_-(z_-)^a (z_--1)^b$ (4.5)

The factor of $\sin(\pi b)$ comes from the fact that the integration around the branch cuts starting at $z_{-} = 1$ has the opposite orientation. To go from the first to the second line we changed variables from $z_{+} \rightarrow 1/z_{+}$. It is easy to check that this gives the same answer as in (4.2) by using

$$\int_0^1 dz_+(z_+)^a (1-z_+)^b = \frac{\Gamma(1+a)\Gamma(1+b)}{\Gamma(2+a+b)}, \quad \Gamma(z)\Gamma(1-z) = \frac{\pi}{\sin(\pi z)}$$
(4.6)

The first factor in the second line of (4.5) shows when

$$2 + a + b = 0, -1, -2, \dots$$
(4.7)

we have a transparent branch cut that makes the integral vanish. This explains the origin of the overall factor of $\Gamma(-1)^{-1}$ we had in our calculation in the previous section. A branch cut becoming transparent is highly suggestive of the smooth limit of a replica geometry. We make this precise in the next sub-section.

4.2 Getting rid of the zero

In this sub-section, we will propose a precise version of the replica trick on the worldsheet that is valid for all orders in α' . In order to do that we need to first

discuss operators that create a conical singularity in the target space. Given that this is a subtle topic we proceed for notational convenience by first discussing it in the thermal AdS_3 background made out of a spatial winding condensate, and then proceed to Euclidean BTZ.

A generic vertex operator in TAdS_3 carrying spatial winding ν takes the following form (see (see [21] for more details)

$$\tilde{V}^{\nu}_{\alpha,a,\bar{a}} = N^{\nu}_{\alpha,a,\bar{a}} \ e^{a\gamma + \bar{a}\gamma} \ e^{n(\int_{(0,0)}^{(z,\bar{z})} \beta dz' + \int_{(0,0)}^{(z,\bar{z})} \bar{\beta} d\bar{z}')} \ e^{2\alpha\varphi}$$
(4.8)

with

$$\alpha := -\frac{j}{b'}, \quad a := m - \frac{k\nu}{4}, \quad n := -\nu$$
(4.9)

We start with a vertex operator with no spatial winding, i.e., $\nu = 0$ and with angular momentum

$$m = \bar{m} = \frac{k}{2} \tag{4.10}$$

At this stage the radial momentum j is arbitrary. We proceed by turning on $\nu = 1$ unit of spatial winding. The resulting operator carries no momentum in any circle.

$$m' = m - \frac{k}{2}\nu = 0, \quad \bar{m}' = \bar{m} - \frac{k}{2}\nu = 0$$
 (4.11)

Then j is fixed by requiring the resulting operator to be on-shell (and decaying at the AdS boundary $\varphi \to -\infty$) in the free theory

$$j = 1 - \frac{k}{2} \tag{4.12}$$

This gives us V^+ .

The off-shell background for the replica trick with $2\pi(1-\delta)$ opening angle is obtained as follows. We start with a vertex operator carrying no winding and momentum close to (4.10).

$$m = \bar{m} = \frac{k}{2}(1 - c\delta) \tag{4.13}$$

Here c is an order one constant. After turning on winding

$$\nu = 1 - \delta \tag{4.14}$$

we require that the resulting operator conserves momentum on both circles which fixes^{20}

$$c = 1 \tag{4.15}$$

Now we fix the radial momentum by requiring the operator to be (1,1) to get

$$V_{\delta}^{\pm} := e^{\pm \frac{k}{4}(1-\delta)(\gamma+\bar{\gamma})} e^{\mp (1-\delta)(\int_{(0,0)}^{(z,\bar{z})} \beta dz' + \int_{(0,0)}^{(z,\bar{z})} \bar{\beta} d\bar{z}')} e^{\left(1-\frac{1}{1-\frac{3}{k}}\delta\right)\sqrt{k-2}\varphi}$$
(4.16)

²⁰Here we are assuming the expression of the currents in terms of free fields remain unchanged as we introduce the conical singularity.

This operator is not on-shell in the interacting theory because it does not have integer quantized winding (so it is not even a local operator). Consider the theory obtained by replacing V^{\pm} with V_{δ}^{\pm} in the action of TAdS₃ (2.6). The operator V_{δ}^{\pm} has a nonzero reflection coefficient of order δ . Therefore to have a smooth interior we would have to add the reflected wave as well in the action (moreover, the reflected wave would change the AdS asymptotics for k > 3). Our proposal is that we just add V_{δ}^{\pm} to the action and do not add any reflected wave associated with it. This keeps the AdS asymptotics unchanged but creates a conical singularity in the $\hat{r}, \hat{\theta}$ plane.

This discussion is easily generalized to Euclidean BTZ and we propose the replica geometry required for the evaluation of the BTZ black hole entropy is given by replacing W^{\pm} by²¹

$$W_{\delta}^{\pm} = e^{\pm i\frac{k}{4}(1-\delta)(\gamma-\bar{\gamma})} e^{\pm i(1-\delta)(\int^{(z,\bar{z})}\beta dz' - \int^{(z,\bar{z})}\bar{\beta} d\bar{z}')} e^{\left(1-\frac{1}{1-\frac{3}{k}}\delta\right)\sqrt{k-2}\varphi}$$
(4.17)

in (2.13) (with no reflected wave added).

For the purpose of calculating the thermal entropy, we need to understand the variation of the free energy of this theory to first order in δ . We need to evaluate the free energy with the insertion of winding operators W_{δ}^{\pm} . However, since evaluation of entropy requires only the knowledge of the first derivative with respect to δ we can set all W_{δ}^{\pm} to $W_{\delta=0}^{\pm}$ except one. In the context of the calculation in the previous section, we fix the $W_{\delta\neq0}^{\pm}$ to be at the origin.

After carefully taking into account the symmetry factor (2s'), the regulated expression is given by (all the terms of order δ^2 or higher are thrown away)

$$z = \frac{2\pi}{b'} \Gamma(-2s') \left(\frac{-\mu}{2b'^2}\right)^{2s'} \frac{\Gamma(2s'+1)}{\Gamma(s'+1)^2} \prod_{I=3}^{s'} d^2 z_I \int \prod_{I'=2}^{s'} d^2 z'_{I'} \\ \left[(\prod_{I} |z_I|^{2(1+\frac{k}{k-3}s'\delta)} |z_I-1|^2) (\prod_{I'} |z'_{I'}|^{2(1-k+(2k+\frac{k}{k-3})s'\delta)} |z'_{I'}-1|^{2(1-k)}) \right]$$

$$\prod_{I'

$$(4.18)$$$$

Now we turn to simplify this expression in terms of Liouville theory correlators. We integrate out $z_{I'}$ with

$$n = s' - 1, \quad m = 0$$

$$t_j = z_j, \quad j = 1, \dots, s' - 2, \quad t_{s'-1} = 0, \quad t_{s'} = 1$$

$$p_j = 1 - k, \quad p_{s'-1} = 1 - k + a(k)\delta, \quad p_{s'} = 1 - k$$
(4.19)

²¹Note that W_{δ}^{\pm} remains weight (1, 1) if we change the coordinate periodicities to $\hat{\theta} \sim \hat{\theta} + 4\pi^2 (Tl_{AdS}), \hat{\xi} \sim \hat{\xi} + 2\pi (1-\delta)$, which corresponds to a background with asymptotic temperature $T(1+\delta)$. Then it would have been a conventional on-shell vertex operator.

Interestingly, we have a k dependent factor in the regulator

$$a(k) = \left(2 + \frac{1}{k-3}\right) \left(\frac{k}{k-2}\right) \tag{4.20}$$

We remind the reader that the expression for the finite factor here is only valid when s' = 2, 3, 4... (for s' = 1 we do not have enough winding operators to fix). This gives the following formula for the residue at those values

In the second line of (4.21), we have written down the residue using the Liouville theory three-point function. To go from the second to the third line of (4.21) we have kept only terms of leading order in $\delta \to 0$ limit. This is precisely the expression in section 3.3 up to the overall factor of the regulator $-a(k)\delta$.

In this language, the entropy is obtained from

$$S_{BTZ} = \lim_{\delta \to 0} \left(-\partial_n (\log Z_{BTZ}(n) - n \log Z_{BTZ}(1)) \Big|_{n=1+\delta} \right)$$
(4.22)

We get the following exact in l_{AdS}/l_s expression for the entropy of the BTZ black hole

$$S_{BTZ} = 16Z_c \left(\frac{Z^M}{g_s^2 \sqrt{k-2}}\right) k \left(1 - \frac{1}{2(k-2)}\right) (Tl_{AdS})$$

$$= \left(\frac{1}{g_s^2} Z_c Z^M Tl_{AdS}\right) 16\sqrt{k} \left(1 + \frac{1}{2k} + \mathcal{O}\left(\frac{1}{k^2}\right)\right)$$

$$(4.23)$$

On the other hand, from the target space gravity analysis the entropy of the BTZ black hole is given by

$$S_{BTZ} = \frac{2\pi V_M}{g_s^2} \left(2\pi T l_{AdS}\right) \left(2\pi \sqrt{k}\right) \left(1 + \mathcal{O}\left(\frac{1}{k}\right)\right) \tag{4.24}$$

The numerical coefficient matches precisely if the path integral is normalized according to

$$Z_c = \frac{\pi^3}{2} \tag{4.25}$$

4.3 The area operator

The calculation performed in this note makes it clear that for the purpose of evaluating thermal entropy, we need to only go away from the on-shell background by an infinitesimal amount, in particular, this leads to the fact that entropy comes entirely from one point function of the stringy area operator

$$A = \frac{\mu}{b^{\prime 4}} \int d^2 \sigma \left. \frac{\partial}{\partial \delta} (W_{\delta}^+ + W_{\delta}^-) \right|_{\delta=0}$$
(4.26)

This operator is not a local operator, in the sense that it has a logarithmic branch cut attached to it. This can be seen from the following OPE

$$W_{\delta}^{+}(z,\bar{z})\tilde{V}_{\alpha,a,\bar{a}}^{0}(0,0) \sim (z)^{-\alpha b'(1-\frac{1}{1-\frac{3}{k}}\delta)-im(1-\delta)}(\bar{z})^{-\alpha b'(1-\frac{1}{1-\frac{3}{k}}\delta)+i\bar{m}(1-\delta)}(1+\ldots)$$

$$W_{\delta}^{-}(z,\bar{z})\tilde{V}_{\alpha,a,\bar{a}}^{0}(0,0) \sim (z)^{-\alpha b'(1-\frac{1}{1-\frac{3}{k}}\delta)+im(1-\delta)}(\bar{z})^{-\alpha b'(1-\frac{1}{1-\frac{3}{k}}\delta)-i\bar{m}(1-\delta)}(1+\ldots)$$
(4.27)

Because of these branch cuts the area operator A given in (4.26) is not a local operator and therefore evaluation of its one-point function is tricky (see a similar discussion for non-normalizable operators in [71]).

5 Summary and future directions

In this paper, we have calculated the thermal Bekenstein-Hawking entropy of the BTZ black hole in bosonic string theory on $AdS_3 \times M$ from a first principles analysis using the winding condensate description of the worldsheet theory. Here, we want to emphasize to the reader the significance of the derivation and its underlying logic. From the point of view of the worldsheet, there are two dimensionless quantities

$$\frac{l_{AdS}}{l_s}, \quad Tl_{AdS} \tag{5.1}$$

The free energy is order g_s^{-2} on a spherical worldsheet due to the usual contribution of the string dilaton. Integration over the angular zero modes of the worldsheet fields in the BTZ background produces a factor of Tl_{AdS} in free energy. At this point, the dependence on l_{AdS}/l_s is completely unknown (the curious reader might demand here that we use the additional input from target space physics that entropy must be order G_N^{-1} , and since G_N has a scale it will fix the l_{AdS} dependence of the answer when l_{AdS}/l_s is scaled to infinity). The central point of this paper is to avoid any such inputs from the target space physics and derive the result intrinsically. Naively one might expect a very simple answer for the coefficient of the linear in temperature dependence of the entropy of the BTZ blackhole,²² based on AdS₃/CFT₂, in terms

²²As an observation we note that the explicit k dependence in (4.23) is the same as the product of the central charge of the SL(2,R) WZW model and the maximum allowed Liouville-like momentum of an operator in the boundary conformal field theory corresponding to the discrete representations [68].

of central charge of the dual conformal field theory (although there are potential subtleties in the present context due to long strings rendering the dual theory not a standard CFT_2 , as the spectrum is not discrete, and our setting in the bosonic string which has a closed string tachyon, so it is not truly a valid background). However, it is not clear that the states which contribute to the Cardy formula of the boundary CFT all come from black holes as we enter the highly stringy regime.²³ Moreover, the nature of BTZ as a quotient makes it clear that the worldsheet genus zero free energy is exactly linear in T, since the temperature therefore only enters through the volume of zero modes, and there are no worldsheet instantons corresponding to maps of S² into the angular directions of the target space which is contractible to S¹. In this work, we determined the exact dependence on k using the following ingredients –

- 1. We performed an honest path integration of the AdS_3 part of worldsheet CFT, however, we did not attempt to consider the path integration over the worldsheet CFT on the compact manifold M in detail. The functional dependence of the entropy on the compact manifold M is only through an overall factor that also appears in the three-point function of vertex operators contained entirely in AdS_3 (such vertex operators always exist in compactifications of the type $AdS_3 \times M$). Therefore, the dependence on M is absorbed in the definition of effective lower dimensional Planck constant. Keeping this in mind, it will be very interesting to compare our results against higher derivative corrections in the target space [72–74]. We leave this to future work.
- 2. We proposed a prescription for the replica trick based on the expectation that non-integer winding introduces a conical singularity at the origin. This procedure is uniquely fixed by the symmetries. In summary, the calculation presented here is correct up to the leading order in small g_s , and we have managed to identify the α' exact 'area' operator whose one-point function produces the leading order entropy of the BTZ black hole. Given the explicit form of the area operator (4.26), we can extend the calculation of its one-point function to higher orders in g_s , say when the worldsheet is a torus. From the point of view of the replica trick in the bulk effective field theory, the one-loop contribution to the thermal entropy comes from two pieces - the entanglement of the bulk fields (including gravitons) outside the horizon of the classical black hole geometry, and the first derivative of the log of the partition function of the replica geometry, having a conical singularity in the interior, with respect to (the inverse of) the opening angle of the conical singularity. Therefore it is

 $^{^{23}}$ Arbitrary higher derivative corrections to the entropy of non-rotating BTZ black holes can be put in a suggestive form of a Cardy formula, i.e, the entropy is proportional to the temperature (we are considering a non-chiral theory), however it is not clear if the coefficient of proportionality is given by the central charge of the dual CFT₂ in general.

likely that the one-point function of the area operator is related to the latter and presumably the former corresponds to the correction to the entropy obtained through the use of the orbifold method developed in [40, 41]. It is a fascinating open problem to explore these questions further possibly keeping in mind any potential connection to logarithmic conformal field theories [75] and algebra of observables in the target space [76-79].²⁴

Any consistent quantum theory of gravity must provide a microscopic explanation for the thermal Bekenstein-Hawking entropy [1, 2] of a black hole. In the context of string theory, Strominger and Vafa showed the microstates of a certain black hole at zero temperature belong to the Hilbert space of the theory living on the world volume of the constituent D-branes [81] (see also Sen's work on black holes built out of elementary strings [82, 83]). Later, Gokakumar and Vafa formulated a different approach to microstate counting in M theory exploiting its connection with topological strings [84, 85].²⁵ Over the years these techniques greatly developed [87– 91], however, the reliance on supersymmetry²⁶ implied that the explanation of the thermal part of the entropy remained elusive [98]. For instance, in the example of black D3 branes, the temperature dependence of the entropy can be obtained using extensivity and conformal properties of the theory on the brane [99], however, dependence on the 't Hooft coupling remains unknown. The similar scaling of entropy with temperature in D0 brane quantum mechanics is determined in [100] based on the added assumption of the existence of certain scaling behavior of the effective theory of the 'moduli' $[101, 102]^{27}$. Conceptually, thermal physics is inherently much more complicated due to the presence of chaotic dynamics.²⁸ Nevertheless, in this paper, we have presented a *well-controlled* worldsheet exact calculation for the thermal Bekenstein-Hawkin entropy of the BTZ black hole.²⁹. A state counting picture of our calculation is still lacking. It would be fantastic to interpret the calculation in terms of the state counting in the Hilbert space of the worldsheet in angular quantization [116] following the proposal in [20] for the Lorentzian meaning of time winding operators, or through the effective 'mini-superspace' approximation of [117, 118].

The major success of the target space version of the replica trick is to produce systematic quantum corrections to the Hubeny-Rangamani-Ryu-Takayanagi formula

 $^{^{24}}$ Also see the prospectives in [80].

²⁵For the successful evaluation of BPS entropy along these lines see [86].

 $^{^{26}}$ For more discussion on the contribution of complex thermal geometries to supersymmetric index see [91–95] (see also [96, 97]).

 $^{^{27}}$ In the context of disorder averaged Sachdev-Ye-Kitaev (SYK) model and Jackiw–Teitelboim (JT) gravity, the leading order thermal entropy of a black hole (further quantum corrections to thermal entropy in JT gravity is calculated in [103, 104]) has been evaluated on the field theory side by Maldacena and Stanford [105].

 $^{^{28}}$ At strong coupling, black holes act as the maximally chaotic quantum systems [106–111].

 $^{^{29}}$ For a discussion of absorption coefficients in AdS₃/CFT₂ based on thermal two-point functions see [112–115].

[3, 4] for the entanglement entropy of a region on the spacetime CFT and its crucial role in the discussion of the black hole information paradox (consult [119, 120] and references therein for a review). It would be very interesting to understand this generalization from the thermal entropy to entanglement entropy of a sub-region of the spacetime CFT directly on the worldsheet.³⁰ Perhaps a good starting point for discussion involves understanding the area operator here in terms of metric deformation using the map of the tachyon vertex operator to the graviton vertex operator as given in [126–128].

Acknowledgments

We thank Yiming Chen for collaboration in the initial stage of this project. We thank David Kolchmeyer, Prahar Mitra, David Tong, Aron Wall, Xi Yin, and especially Nicholas Agia, Amr Ahmadain, Juan Maldacena and Douglas Stanford for insightful discussions and comments. IH is supported by the Harvard Quantum Initiative Fellowship. The work of DLJ and IH is supported in part by DOE grants DE-SC0007870 and DE-SC0021013.

A Three point function at zero temperature

In this Appendix, we will compare the three-point function as calculated from the worldsheet with that of the gravity in target space for large values of l_{AdS}/l_s . The process is not completely fixed in principle because of the unknown normalization of vertex operators (in flatspace string theory, the normalization of vertex operators is unambiguously fixed by demanding the unitarity of spacetime S matrix). In AdS, there is no notion of S-matrix. One might hope to take advantage of the unitarity of the spacetime CFT living on the boundary of AdS. This procedure is tricky in the AdS₃ background we are talking about. This is because of the issues related to the normalization of spacetime stress tensor [129], which in turn related to the issues of normalization of the vacuum [130]. We will fix the normalization of vertex operators by looking at the example of AdS₅ × S⁵ up to an order one constant factor.

It will be easier to work in the Poincare patch in $l_{AdS} = 1$ unit for this purpose:

$$ds^2 = d\hat{\phi}^2 + e^{2\hat{\phi}}d\hat{\Gamma}d\hat{\bar{\Gamma}} = \frac{du^2 + d\vec{x}^2}{u^2}$$
(A.1)

Here we used the following change of variables

$$u = e^{-\hat{\phi}}, \quad d\vec{x}^2 = d\hat{\Gamma}d\hat{\bar{\Gamma}} \tag{A.2}$$

 $^{^{30}}$ See [121–125] for some of the related discussions.

By the complete analogy with the discussion of $AdS_5 \times S^5$ in [131], we demand the following effective action for a set of three scalar fields ϕ_i of mass M_i

$$S_{g} = \frac{1}{2} \int d^{2}\vec{x} du \sqrt{g} \left(g^{\mu\nu} \partial_{\mu} \phi \partial_{\nu} \phi + M^{2} \phi^{2} + \sqrt{l_{3}} \phi_{1} \phi_{2} \phi_{3} \right), \quad \frac{1}{l_{3}} = \frac{V_{M}}{g_{s}^{2} l_{s}}$$
(A.3)

In direct quantization, the bulk-to-boundary propagator $K_{\Delta_i}(u, \vec{x}; \vec{y})$ from (u, \vec{x}) to \vec{y} satisfies

$$(\nabla^2 - M_i^2) K_{\Delta_i}(u, \vec{x}; \vec{y}) = 0, \quad \Delta_i = 1 + \sqrt{1 + M_i^2}$$
 (A.4)

with the following boundary condition

$$\lim_{u \to 0} u^{\Delta_i - 2} K_{\Delta_i}(u, \vec{x}; \vec{y}) = \delta^{(2)}(\vec{x} - \vec{y})$$
(A.5)

Comparing (A.4),(A.5) with (2.4),(2.5) (note the former is calculated in $l_{AdS} = 1$ units, whereas later is calculated in $l_s = 1$ unit) suggests that Φ_j represents a scalar field in the target space with the map $\Delta(j) = -2j$. To make this claim sharp we calculate the three-point function using the Witten diagram. Up to a constant factor it is given by [132]

$$C_{123} = \sqrt{l_3} \Gamma\left(\frac{\Delta}{2} - 1\right) \prod_{i=1}^3 \frac{\Gamma\left(\frac{\Delta - 2\Delta_i}{2}\right)}{\Gamma\left(\Delta_i - 1\right)}, \quad \Delta = \sum_{i=1}^3 \Delta_i \tag{A.6}$$

Now we turn to calculate the string theoretic three-point function from the point of view of the worldsheet. We have to multiply the three-point function of Φ_{j_i} in SL(2,R) WZW model with the normalization of path integration chosen in this paper with the sphere partition function of the worldsheet CFT. The sphere partition function is already determined in the main text for non-zero temperature. The only difference with the zero temperature calculation considered here is due to the zero mode integral in $\gamma + \bar{\gamma}$. Here the zero mode ends up giving a conservation law for the momentum of the external operators (see section 6.2 of [59] for more explanation), as a result, the normalization is given by

$$C_{S^2} = \frac{1}{g_s^2} C_{S^2}^{AdS_3} Z_0^M C_{S^2}^M \frac{Z_c}{z_{SL(2,\mathbb{C})}} \sim \frac{1}{l_3}$$
(A.7)

Here we omitted any order one constant. The three-point function in WZW model is given by [21, 43] (we stripped off the conformal factor involving x, \bar{x})

$$\begin{split} \langle \Phi_{j_1}(0|0)\Phi_{j_2}(1|1)\Phi_{j_3}(\infty|\infty)\rangle &= \frac{1}{2\pi^3 b''} \left[\frac{\gamma(b''^2)b''^{2-2b''^2}}{\pi}\right]^{-2-\sum_k j_k} \\ &\frac{\Upsilon_{b''}(0)\prod_{k=1}^3\Upsilon_{b''}(-b''(2j_k+1))}{\Upsilon_{b''}(-b''(\sum_i j_i+1))\prod_{k=1}^3\Upsilon_{b''}(-b''(\sum_i j_i-2j_k))} \end{split}$$
(A.8)

For the purpose of obtaining the large k limit of the formula, we use [133]

$$\Upsilon_b(\sigma b) = \frac{b^{-\sigma}}{\Gamma(\sigma)} F_1 \sqrt{b} e^{-\frac{1}{4b^2} \log b + \frac{F_0}{b^2} + \mathcal{O}(b^2 \log b)}$$

$$\implies \Upsilon'_b(0)) = \frac{F_1}{\sqrt{b}} e^{-\frac{1}{4b^2} \log b + \frac{F_0}{b^2} + \mathcal{O}(b^2 \log b)}$$
(A.9)

Here $F_{0,1}$ are two constants whose value will not be important for our purpose. Using these formulae we get

$$\lim_{b''\to 0} \frac{1}{b''} \frac{\Upsilon'_{b''}(0) \prod_{k=1}^{3} \Upsilon_{b''}(-b''(2j_k+1))}{\Upsilon_{b''}(-b''(\sum_k j_k+1)) \prod_{k=1}^{3} \Upsilon_{b''}(-b''(\sum_i j_i-2j_k))} = \Gamma\left(-(\sum_i j_i+1)\right) \frac{\prod_{k=1}^{3} \Gamma(-(\sum_i j_i-2j_k))}{\prod_{k=1}^{3} \Gamma(-(2j_k+1))} \tag{A.10}$$

$$\lim_{b''\to 0} \gamma(b''^2) b''^{2-2b''^2} = 1$$

This shows, the string theoretic three-point function of operators Φ_i^G defined as³¹

$$\Phi_j^G = \sqrt{l_3} \ \pi^{-j_i} \ \Phi_j \leftrightarrow \phi \text{ with } \Delta(j) = -2j \tag{A.11}$$

evaluated using (A.8) multiplied with (A.7) in large k limit matches precisely (up to order one factor) with the result of Witten diagram in (A.6).

To compare our result against the one in [134] (see also [68]), one must carefully take into account the relation between g_s, Z^M and the order of the symmetric product on the spacetime CFT (N in the notation of [134]).³² At present these relations are not known precisely because of the issues related to the normalization of spacetime stress tensor [129], which in turn related to the issues of normalization of the vacuum [130] and we leave these questions to the future.

References

- [1] J. D. Bekenstein, Black holes and entropy, Phys. Rev. D 7 (Apr, 1973) 2333-2346.
- [2] S. W. Hawking, Particle Creation by Black Holes, Commun. Math. Phys. 43 (1975) 199–220. [Erratum: Commun.Math.Phys. 46, 206 (1976)].
- [3] S. Ryu and T. Takayanagi, Holographic derivation of entanglement entropy from AdS/CFT, Phys. Rev. Lett. **96** (2006) 181602, [hep-th/0603001].

³¹There overall factor of $\sqrt{l_3}$ is obtained by keeping in mind the large central charge expansion of the spacetime CFT (two-point function in the space-time CFT is normalized to be order one). Here we introduced the factor of π arbitrarily.

 $^{^{32}}$ For a recent attempt see [14].

- [4] V. E. Hubeny, M. Rangamani, and T. Takayanagi, A Covariant holographic entanglement entropy proposal, JHEP 07 (2007) 062, [arXiv:0705.0016].
- [5] J. W. York, Role of conformal three-geometry in the dynamics of gravitation, Phys. Rev. Lett. 28 (Apr, 1972) 1082–1085.
- [6] G. W. Gibbons and S. W. Hawking, Action integrals and partition functions in quantum gravity, Phys. Rev. D 15 (May, 1977) 2752–2756.
- [7] A. Dabholkar, Strings on a cone and black hole entropy, Nucl. Phys. B 439 (1995) 650-664, [hep-th/9408098].
- [8] A. Dabholkar, Tachyon condensation and black hole entropy, Phys. Rev. Lett. 88 (2002) 091301, [hep-th/0111004].
- [9] A. Sen, Quantum Entropy Function from AdS(2)/CFT(1) Correspondence, Int. J. Mod. Phys. A 24 (2009) 4225-4244, [arXiv:0809.3304].
- [10] A. Lewkowycz and J. Maldacena, Generalized gravitational entropy, JHEP 08 (2013) 090, [arXiv:1304.4926].
- [11] T. Faulkner, A. Lewkowycz, and J. Maldacena, Quantum corrections to holographic entanglement entropy, JHEP 11 (2013) 074, [arXiv:1307.2892].
- [12] N. Engelhardt and A. C. Wall, Quantum Extremal Surfaces: Holographic Entanglement Entropy beyond the Classical Regime, JHEP 01 (2015) 073, [arXiv:1408.3203].
- [13] J. M. Maldacena and A. Strominger, AdS(3) black holes and a stringy exclusion principle, JHEP 12 (1998) 005, [hep-th/9804085].
- [14] L. Eberhardt and S. Pal, Holographic Weyl anomaly in string theory, [arXiv:2307.03000].
- [15] J. Liu and J. Polchinski, Renormalization of the Mobius Volume, Phys. Lett. B 203 (1988) 39–43.
- [16] J. Polchinski, Dirichlet Branes and Ramond-Ramond charges, Phys. Rev. Lett. 75 (1995) 4724-4727, [hep-th/9510017].
- [17] L. Eberhardt and S. Pal, The disk partition function in string theory, JHEP 08 (2021) 026, [arXiv:2105.08726].
- [18] H. Erbin, J. Maldacena, and D. Skliros, Two-Point String Amplitudes, JHEP 07 (2019) 139, [arXiv:1906.06051].
- [19] J. Troost, The AdS_3 central charge in string theory, Phys. Lett. B 705 (2011) 260–263, [arXiv:1109.1923].
- [20] D. L. Jafferis and E. Schneider, Stringy ER=EPR, [arXiv:2104.07233].
- [21] I. Halder, D. L. Jafferis, and D. Kolchmeyer, A duality in string theory on AdS₃, [arXiv:2208.00016].
- [22] I. Halder and D. L. Jafferis, Double winding condensate CFT, [arXiv:2308.11702].

- [23] T. Erler, The closed string field theory action vanishes, JHEP 10 (2022) 055, [arXiv:2204.12863].
- [24] O. Bergman and B. Zwiebach, The Dilaton theorem and closed string backgrounds, Nucl. Phys. B 441 (1995) 76-118, [hep-th/9411047].
- [25] A. Adams, J. Polchinski, and E. Silverstein, Don't panic! Closed string tachyons in ALE space-times, JHEP 10 (2001) 029, [hep-th/0108075].
- [26] Y. Okawa and B. Zwiebach, Twisted tachyon condensation in closed string field theory, JHEP 03 (2004) 056, [hep-th/0403051].
- [27] M. Headrick, S. Minwalla, and T. Takayanagi, Closed string tachyon condensation: An Overview, Class. Quant. Grav. 21 (2004) S1539–S1565, [hep-th/0405064].
- [28] A. A. Tseytlin, Mobius Infinity Subtraction and Effective Action in σ Model Approach to Closed String Theory, Phys. Lett. B **208** (1988) 221–227.
- [29] A. A. Tseytlin, SIGMA MODEL APPROACH TO STRING THEORY, Int. J. Mod. Phys. A 4 (1989) 1257.
- [30] H. Osborn, General Bosonic σ Models and String Effective Actions, Annals Phys. **200** (1990) 1.
- [31] H. Osborn, Weyl consistency conditions and a local renormalization group equation for general renormalizable field theories, Nucl. Phys. B 363 (1991) 486–526.
- [32] S. Carlip and C. Teitelboim, The Off-shell black hole, Class. Quant. Grav. 12 (1995) 1699–1704, [gr-qc/9312002].
- [33] M. Banados, C. Teitelboim, and J. Zanelli, Black hole entropy and the dimensional continuation of the Gauss-Bonnet theorem, Phys. Rev. Lett. 72 (1994) 957–960, [gr-qc/9309026].
- [34] L. Susskind and J. Uglum, Black hole entropy in canonical quantum gravity and superstring theory, Phys. Rev. D 50 (1994) 2700–2711, [hep-th/9401070].
- [35] D. V. Fursaev, Proof of the holographic formula for entanglement entropy, JHEP 09 (2006) 018, [hep-th/0606184].
- [36] M. Headrick, Entanglement Renyi entropies in holographic theories, Phys. Rev. D 82 (2010) 126010, [arXiv:1006.0047].
- [37] R. Brustein and Y. Zigdon, Black hole entropy sourced by string winding condensate, JHEP 10 (2021) 219, [arXiv:2107.09001].
- [38] A. Ahmadain and A. C. Wall, Off-Shell Strings I: S-matrix and Action, [arXiv:2211.08607].
- [39] A. Ahmadain and A. C. Wall, Off-Shell Strings II: Black Hole Entropy, [arXiv:2211.16448].
- [40] E. Witten, Open Strings On The Rindler Horizon, JHEP 01 (2019) 126, [arXiv:1810.11912].

- [41] A. Dabholkar, Quantum Entanglement in String Theory, [arXiv: 2207.03624].
- [42] S. He, T. Numasawa, T. Takayanagi, and K. Watanabe, Notes on Entanglement Entropy in String Theory, JHEP 05 (2015) 106, [arXiv:1412.5606].
- [43] S. Ribault and J. Teschner, H+(3)-WZNW correlators from Liouville theory, JHEP 06 (2005) 014, [hep-th/0502048].
- [44] A. Giveon, D. Kutasov, and N. Seiberg, Comments on string theory on AdS(3), Adv. Theor. Math. Phys. 2 (1998) 733-782, [hep-th/9806194].
- [45] D. Kutasov and N. Seiberg, More comments on string theory on AdS(3), JHEP 04 (1999) 008, [hep-th/9903219].
- [46] J. Teschner, On structure constants and fusion rules in the SL(2,C) / SU(2) WZNW model, Nucl. Phys. B 546 (1999) 390-422, [hep-th/9712256].
- [47] J. Teschner, Operator product expansion and factorization in the H+(3) WZNW model, Nucl. Phys. B 571 (2000) 555-582, [hep-th/9906215].
- [48] J. M. Maldacena and H. Ooguri, Strings in AdS(3) and SL(2,R) WZW model 1.: The Spectrum, J. Math. Phys. 42 (2001) 2929–2960, [hep-th/0001053].
- [49] J. M. Maldacena, H. Ooguri, and J. Son, Strings in AdS(3) and the SL(2,R) WZW model. Part 2. Euclidean black hole, J. Math. Phys. 42 (2001) 2961–2977, [hep-th/0005183].
- [50] J. M. Maldacena and H. Ooguri, Strings in AdS(3) and the SL(2,R) WZW model. Part 3. Correlation functions, Phys. Rev. D 65 (2002) 106006, [hep-th/0111180].
- [51] A. Dei and L. Eberhardt, String correlators on AdS₃: four-point functions, JHEP 09 (2021) 209, [arXiv:2107.01481].
- [52] A. Dei and L. Eberhardt, String correlators on AdS₃: three-point functions, JHEP 08 (2021) 025, [arXiv:2105.12130].
- [53] M. R. Gaberdiel and R. Gopakumar, Tensionless string spectra on AdS₃, JHEP 05 (2018) 085, [arXiv:1803.04423].
- [54] L. Eberhardt, M. R. Gaberdiel, and R. Gopakumar, The Worldsheet Dual of the Symmetric Product CFT, JHEP 04 (2019) 103, [arXiv:1812.01007].
- [55] L. Eberhardt, M. R. Gaberdiel, and R. Gopakumar, Deriving the AdS₃/CFT₂ correspondence, JHEP 02 (2020) 136, [arXiv:1911.00378].
- [56] A. Dei, M. R. Gaberdiel, R. Gopakumar, and B. Knighton, Free field world-sheet correlators for AdS₃, JHEP 02 (2021) 081, [arXiv:2009.11306].
- [57] A. Zamolodchikov, Scaling Lee-Yang model on a sphere. 1. Partition function, JHEP 07 (2002) 029, [hep-th/0109078].
- [58] A. B. Zamolodchikov and A. B. Zamolodchikov, Structure constants and conformal bootstrap in Liouville field theory, Nucl. Phys. B 477 (1996) 577–605, [hep-th/9506136].

- [59] J. Polchinski, String theory. Vol. 1: An introduction to the bosonic string. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 12, 2007.
- [60] E. Frenkel and A. Losev, Mirror symmetry in two steps: A-I-B, Commun. Math. Phys. 269 (2006) 39-86, [hep-th/0505131].
- [61] M. Goulian and M. Li, Correlation functions in liouville theory, Phys. Rev. Lett. 66 (Apr, 1991) 2051–2055.
- [62] H. Dorn and H. J. Otto, Two and three point functions in Liouville theory, Nucl. Phys. B 429 (1994) 375–388, [hep-th/9403141].
- [63] J. Teschner, Liouville theory revisited, Class. Quant. Grav. 18 (2001) R153–R222, [hep-th/0104158].
- [64] B. Balthazar, A. Giveon, D. Kutasov, and E. J. Martinec, Asymptotically free AdS₃/CFT₂, JHEP **01** (2022) 008, [arXiv:2109.00065].
- [65] R. Mahajan, D. Stanford, and C. Yan, Sphere and disk partition functions in Liouville and in matrix integrals, JHEP 07 (2022) 132, [arXiv:2107.01172].
- [66] G. Giribet and M. Leoni, A Twisted FZZ-like dual for the 2D black hole, Rept. Math. Phys. 61 (2008) 151–162, [arXiv:0706.0036].
- [67] V. A. Fateev and A. V. Litvinov, Multipoint correlation functions in Liouville field theory and minimal Liouville gravity, Theor. Math. Phys. 154 (2008) 454–472, [arXiv:0707.1664].
- [68] L. Eberhardt, A perturbative CFT dual for pure NS-NS AdS₃ strings, J. Phys. A 55 (2022), no. 6 064001, [arXiv:2110.07535].
- [69] J. Polchinski, Evaluation of the One Loop String Path Integral, Commun. Math. Phys. 104 (1986) 37.
- [70] I. Halder and D. L. Jafferis, Stretched horizon, replica trick and off-shell winding condensate, and all that, [arXiv:2402.00932].
- [71] P. Kraus, A. Ryzhov, and M. Shigemori, Strings in noncompact space-times: Boundary terms and conserved charges, Phys. Rev. D 66 (2002) 106001, [hep-th/0206080].
- [72] P. Kraus and F. Larsen, Microscopic black hole entropy in theories with higher derivatives, JHEP 09 (2005) 034, [hep-th/0506176].
- [73] P. Kraus and F. Larsen, Holographic gravitational anomalies, JHEP 01 (2006) 022, [hep-th/0508218].
- [74] J. R. David, B. Sahoo, and A. Sen, AdS(3), black holes and higher derivative corrections, JHEP 07 (2007) 058, [arXiv:0705.0735].
- [75] T. Creutzig and D. Ridout, Logarithmic Conformal Field Theory: Beyond an Introduction, J. Phys. A 46 (2013) 4006, [arXiv:1303.0847].

- [76] S. Leutheusser and H. Liu, Causal connectability between quantum systems and the black hole interior in holographic duality, [arXiv:2110.05497].
- [77] S. Leutheusser and H. Liu, *Emergent times in holographic duality*, [arXiv:2112.12156].
- [78] E. Witten, Gravity and the crossed product, JHEP 10 (2022) 008, [arXiv:2112.12828].
- [79] V. Chandrasekaran, G. Penington, and E. Witten, Large N algebras and generalized entropy, [arXiv:2209.10454].
- [80] H. Verlinde, ER = EPR revisited: On the Entropy of an Einstein-Rosen Bridge, [arXiv:2003.13117].
- [81] A. Strominger and C. Vafa, Microscopic origin of the Bekenstein-Hawking entropy, Phys. Lett. B 379 (1996) 99–104, [hep-th/9601029].
- [82] A. Sen, Extremal black holes and elementary string states, Mod. Phys. Lett. A 10 (1995) 2081–2094, [hep-th/9504147].
- [83] A. Sen, How does a fundamental string stretch its horizon?, JHEP 05 (2005) 059, [hep-th/0411255].
- [84] R. Gopakumar and C. Vafa, M theory and topological strings. 1., [hep-th/9809187].
- [85] R. Gopakumar and C. Vafa, M theory and topological strings. 2., [hep-th/9812127].
- [86] I. Halder and Y.-H. Lin, *Blackhole/blackring transition*, [arXiv:2307.13735].
- [87] J. Kinney, J. M. Maldacena, S. Minwalla, and S. Raju, An Index for 4 dimensional super conformal theories, Commun. Math. Phys. 275 (2007) 209-254, [hep-th/0510251].
- [88] J. Bhattacharya, S. Bhattacharyya, S. Minwalla, and S. Raju, Indices for Superconformal Field Theories in 3,5 and 6 Dimensions, JHEP 02 (2008) 064, [arXiv:0801.1435].
- [89] H. Ooguri, A. Strominger, and C. Vafa, Black hole attractors and the topological string, Phys. Rev. D 70 (2004) 106007, [hep-th/0405146].
- [90] F. Denef and G. W. Moore, Split states, entropy enigmas, holes and halos, JHEP 11 (2011) 129, [hep-th/0702146].
- [91] F. Benini, K. Hristov, and A. Zaffaroni, Black hole microstates in AdS₄ from supersymmetric localization, JHEP 05 (2016) 054, [arXiv:1511.04085].
- [92] A. Cabo-Bizet, D. Cassani, D. Martelli, and S. Murthy, Microscopic origin of the Bekenstein-Hawking entropy of supersymmetric AdS₅ black holes, JHEP 10 (2019) 062, [arXiv:1810.11442].
- [93] S. Choi, J. Kim, S. Kim, and J. Nahmgoong, Large AdS black holes from QFT, [arXiv:1810.12067].

- [94] L. V. Iliesiu, M. Kologlu, and G. J. Turiaci, Supersymmetric indices factorize, JHEP 05 (2023) 032, [arXiv:2107.09062].
- [95] A. A. H., P. V. Athira, C. Chowdhury, and A. Sen, Logarithmic Correction to BPS Black Hole Entropy from Supersymmetric Index at Finite Temperature, [arXiv:2306.07322].
- [96] Y. Chen and G. J. Turiaci, Spin-Statistics for Black Hole Microstates, [arXiv:2309.03478].
- [97] Y. Chen, M. Heydeman, Y. Wang, and M. Zhang, Probing Supersymmetric Black Holes with Surface Defects, [arXiv:2306.05463].
- [98] G. T. Horowitz, J. M. Maldacena, and A. Strominger, Nonextremal black hole microstates and U duality, Phys. Lett. B 383 (1996) 151–159, [hep-th/9603109].
- [99] S. S. Gubser, I. R. Klebanov, and A. W. Peet, Entropy and temperature of black 3-branes, Phys. Rev. D 54 (1996) 3915–3919, [hep-th/9602135].
- [100] A. Biggs and J. Maldacena, Scaling similarities and quasinormal modes of D0 black hole solutions, [arXiv:2303.09974].
- [101] S. Paban, S. Sethi, and M. Stern, Constraints from extended supersymmetry in quantum mechanics, Nucl. Phys. B 534 (1998) 137–154, [hep-th/9805018].
- [102] S. Paban, S. Sethi, and M. Stern, Supersymmetry and higher derivative terms in the effective action of Yang-Mills theories, JHEP 06 (1998) 012, [hep-th/9806028].
- [103] M. Heydeman, L. V. Iliesiu, G. J. Turiaci, and W. Zhao, The statistical mechanics of near-BPS black holes, J. Phys. A 55 (2022), no. 1 014004, [arXiv:2011.01953].
- [104] L. V. Iliesiu and G. J. Turiaci, The statistical mechanics of near-extremal black holes, JHEP 05 (2021) 145, [arXiv:2003.02860].
- [105] J. Maldacena and D. Stanford, Remarks on the Sachdev-Ye-Kitaev model, Phys. Rev. D 94 (2016), no. 10 106002, [arXiv:1604.07818].
- [106] J. Maldacena, S. H. Shenker, and D. Stanford, A bound on chaos, JHEP 08 (2016) 106, [arXiv:1503.01409].
- [107] I. Halder, Global symmetry and maximal chaos, [arXiv:1908.05281].
- [108] X. O. Camanho, J. D. Edelstein, J. Maldacena, and A. Zhiboedov, Causality constraints on corrections to the graviton three-point coupling, JHEP 02 (2016) 020, [arXiv:1407.5597].
- [109] S. D. Chowdhury, A. Gadde, T. Gopalka, I. Halder, L. Janagal, and S. Minwalla, Classifying and constraining local four photon and four graviton S-matrices, JHEP 02 (2020) 114, [arXiv:1910.14392].
- [110] D. Chandorkar, S. D. Chowdhury, S. Kundu, and S. Minwalla, Bounds on Regge growth of flat space scattering from bounds on chaos, JHEP 05 (2021) 143, [arXiv:2102.03122].

- [111] M. Blake and H. Liu, On systems of maximal quantum chaos, JHEP 05 (2021) 229, [arXiv:2102.11294].
- [112] S. R. Das and S. D. Mathur, Comparing decay rates for black holes and D-branes, Nucl. Phys. B 478 (1996) 561-576, [hep-th/9606185].
- [113] S. R. Das and S. D. Mathur, Interactions involving D-branes, Nucl. Phys. B 482 (1996) 153-172, [hep-th/9607149].
- [114] A. Dhar, G. Mandal, and S. R. Wadia, Absorption versus decay of black holes in string theory and T symmetry, Phys. Lett. B 388 (1996) 51–59, [hep-th/9605234].
- [115] J. M. Maldacena and A. Strominger, Universal low-energy dynamics for rotating black holes, Phys. Rev. D 56 (1997) 4975-4983, [hep-th/9702015].
- [116] N. Agia and D. L. Jafferis, Angular Quantization in CFT, [arXiv:2204.11872].
- [117] R. Dijkgraaf, H. L. Verlinde, and E. P. Verlinde, String propagation in a black hole geometry, Nucl. Phys. B 371 (1992) 269–314.
- [118] T. G. Mertens, H. Verschelde, and V. I. Zakharov, Random Walks in Rindler Spacetime and String Theory at the Tip of the Cigar, JHEP 03 (2014) 086, [arXiv:1307.3491].
- [119] A. Almheiri, T. Hartman, J. Maldacena, E. Shaghoulian, and A. Tajdini, The entropy of Hawking radiation, Rev. Mod. Phys. 93 (2021), no. 3 035002, [arXiv:2006.06872].
- [120] S. Raju, Lessons from the information paradox, Phys. Rept. 943 (2022) 1–80, [arXiv:2012.05770].
- [121] S. A. Hartnoll and E. Mazenc, Entanglement entropy in two dimensional string theory, Phys. Rev. Lett. 115 (2015), no. 12 121602, [arXiv:1504.07985].
- [122] S. R. Das, A. Kaushal, G. Mandal, and S. P. Trivedi, Bulk Entanglement Entropy and Matrices, J. Phys. A 53 (2020), no. 44 444002, [arXiv:2004.00613].
- [123] S. R. Das, A. Kaushal, S. Liu, G. Mandal, and S. P. Trivedi, Gauge invariant target space entanglement in D-brane holography, JHEP 04 (2021) 225, [arXiv:2011.13857].
- [124] S. R. Das, A. Kaushal, G. Mandal, K. K. Nanda, M. H. Radwan, and S. P. Trivedi, Entanglement entropy in internal spaces and Ryu-Takayanagi surfaces, JHEP 04 (2023) 141, [arXiv:2212.11640].
- [125] V. Balasubramanian and O. Parrikar, Remarks on entanglement entropy in string theory, Phys. Rev. D 97 (2018), no. 6 066025, [arXiv:1801.03517].
- [126] A. Giveon, N. Itzhaki, and D. Kutasov, Stringy Horizons II, JHEP 10 (2016) 157, [arXiv:1603.05822].
- [127] E. J. Martinec, S. Massai, and D. Turton, *Stringy Structure at the BPS Bound*, *JHEP* **12** (2020) 135, [arXiv:2005.12344].

- [128] E. J. Martinec, AdS3 Orbifolds, BTZ Black Holes, and Holography, [arXiv:2307.02559].
- [129] H. Bertle, A. Dei, and M. R. Gaberdiel, Stress-energy tensor correlators from the world-sheet, JHEP 03 (2021) 036, [arXiv:2012.08486].
- [130] J. Kim and M. Porrati, On the central charge of spacetime current algebras and correlators in string theory on AdS₃, JHEP 05 (2015) 076, [arXiv:1503.07186].
- [131] S. Lee, S. Minwalla, M. Rangamani, and N. Seiberg, Three point functions of chiral operators in D = 4, N=4 SYM at large N, Adv. Theor. Math. Phys. 2 (1998) 697-718, [hep-th/9806074].
- [132] D. Z. Freedman, S. D. Mathur, A. Matusis, and L. Rastelli, Correlation functions in the CFT(d) / AdS(d+1) correspondence, Nucl. Phys. B 546 (1999) 96-118, [hep-th/9804058].
- [133] D. Harlow, J. Maltz, and E. Witten, Analytic Continuation of Liouville Theory, JHEP 12 (2011) 071, [arXiv:1108.4417].
- [134] A. Dei and L. Eberhardt, String correlators on AdS₃: Analytic structure and dual CFT, SciPost Phys. 13 (2022), no. 3 053, [arXiv:2203.13264].