

Ergodic and mixing quantum channels: From two-qubit to many-body quantum systems

S. Aravinda,^{1,*} Shilpak Banerjee,^{2,†} and Ranjan Modak^{1,‡}

¹*Department of Physics, Indian Institute of Technology Tirupati, Tirupati, India 517619*

²*Department of Mathematics and Statistics, Indian Institute of Technology Tirupati, Tirupati, India 517619*

The development of classical ergodic theory has had a significant impact in the areas of mathematics, physics, and, in general, applied sciences. The quantum ergodic theory of Hamiltonian dynamics has its motivations to understand thermodynamics and statistical mechanics. Quantum channel, a completely positive trace-preserving map, represents a most general representation of quantum dynamics and is an essential aspect of quantum information theory and quantum computation. In this work, we study the ergodic theory of quantum channels by characterizing different levels of ergodic hierarchy from integrable to mixing. The quantum channels on single systems are constructed from the unitary operators acting on bipartite states and tracing out the environment. The interaction strength of these unitary operators measured in terms of operator entanglement provides sufficient conditions for the channel to be mixing. By using block diagonal unitary operators, we construct a set of non-ergodic channels. By using canonical form of two-qubit unitary operator, we analytically construct the channels on single qubit ranging from integrable to mixing. Moreover, we also study interacting many-body quantum systems that include the famous Sachdev-Ye-Kitaev (SYK) model and show that they display mixing within the framework of the quantum channel.

I. INTRODUCTION

In classical ergodic theory, we study the statistical properties of a probability-measure-preserving transformation, and the five most widely studied properties in the community are ergodicity, weak-mixing, (strong) mixing, K-property, and Bernoulli. It turns out that these properties can be stacked in a nice hierarchy, often referred to as the ergodic hierarchy, with the strongest one being Bernoulli. More precisely: Bernoulli $\implies$ K-property $\implies$ (strong) mixing $\implies$ weak mixing $\implies$ ergodicity [1]. The reverse implications are not true in general and counterexamples exist. Predictable, the set of transformations becomes smaller when one requires a stronger property in the hierarchy, and in fact, invertible Bernoulli transformations are completely classified [2] while invertible ergodic transformations cannot be classified [3] answering the isomorphism question laid down in von Neumann's seminal paper [4] in the negative.

In quantum theory, early works on quantum ergodic theory in Hamiltonian unitary dynamics are by von Neumann which he named it as quantum ergodic theorem and H theorem [5, 6]. These theorems met with criticisms [7–9] and counter justifications [10]. Peres defined and studied ergodicity and mixing based on his definition of quantum chaos [8, 9]. Many others continued to study ergodic and mixing behaviour in quantum Hamiltonian dynamics, maps, and its relation to chaos and various other related quantities [11–

17]. In the quantum many-body systems, these concepts link with that of the thermalization and has produced a host of interesting results (see review [18] and reference therein). Remarkably, certain spacetime dual quantum many-body circuit provides an analytically solvable minimal model for the correlation function between two spatiotemporal observables [19–22], and thereby facilitates the characterization of the entire ergodic hierarchy [20, 23]. The question of how an isolated many-body quantum system thermalizes while going through unitary dynamics also brought lots of interest in the last two decades thanks to extraordinary advancements in cold atom experiments [24, 25]. It is now well-established that if the many-body energy level spacing of a system are of the Wigner-Dyson type, one expects thermalization to occur [26] i.e. the system acts as its own bath, obey the eigenstate thermalization Hypothesis [27–30].

Quantum channels, a linear map that maps quantum states, is a generalization of unitary quantum dynamics [31, 32]. Mathematically, it is a linear, completely positive trace preserving (CPTP) map [33, 34]. At this juncture, we would like to map the results of ergodic theory from classical to quantum. Here, the quantum channel acts as a transformation on the set of quantum states, and we are interested in studying the iterative application of quantum channels. Ergodic theory of quantum channels has been studied in the past [35, 36] using the algebraic framework. With the emergence of quantum information theory and computation, the ergodic theory of quantum channels in finite-dimensional systems took the forefront as it has been the topic of many recent works [37–41]. The theoretical results of iterative application of quantum channels has been used in various tasks in quantum information theory. For

* aravinda@iittp.ac.in

† shilpak@iittp.ac.in

‡ ranjan@iittp.ac.in




Figure 1. Sketch of the iterative process studied in this work (see text Sec. II for the description).

example, in the studies of structure-preserving maps, which have led to a better understanding of quantum error correction [42–44], the convergence rates of quantum channels are related to critical exponents in the tensor representation of many-body systems [45], to construct any measurement using sequential generalized measurements [46, 47], and asymptotic results in quantum channels [48, 49]. Mixing of quantum channels is related to information-theoretic channel capacities [50].

II. SUMMARY OF MAIN RESULTS

In this section, we first briefly describe our setup and then summarize our main findings. We prepare a system that is described by the Hamiltonian H_s and state ρ_s . Then, we connect the system with a bath and let the entire system (system+bath) evolve. ρ_b and H_b are referred to as the state and the Hamiltonian of the bath, respectively, and H_{sb} stands for the interaction between the system and the bath. After the evolution, the state of the entire system becomes, $\rho_{sb} = U(\rho \otimes \sigma)U^\dagger$, where $U = e^{-i(H_s + H_b + H_{sb})}$. We trace out the bath degrees of freedom from ρ_{sb} and obtain the reduced density matrix for the system $\rho_s^1 = \text{Tr}_b \rho_{sb}$. Again, we connect the system with the bath and repeat it iteratively n times. Schematically, this is explained in Fig. 1.

In a broader theoretical perspective, every quantum channel acting on the single quantum system has a representation in terms of a unitary operator acting on a bipartite system and then tracing out one of the systems [51, 52]. By using the notions of matrix realignment [51, 53, 54] used extensively in entanglement theory [55, 56] and recently in the many-body systems literature [57, 58], we provide the representation of the channel in terms of the unitary operator and the environment, and is explained in Sec. (III A). Our study focuses largely on the spectral properties of channels, and the details are presented in Ref. [31, 32]. For completeness and continuity, we present the spectral results of the channel in detail in Sec. (III B), and we define the various equivalent definitions of ergodic and mixing in Sec. (III C). The main results of our paper at the formalism level and application to many-body physics are presented in Sec. (IV) and Sec. (V), respectively. The sufficiency condition on mixing of quantum channels in terms of the strength of unitary operator measured as operator entanglement [59, 60] is proved in Sec. (IV A), the stability of sufficiency condition under any local unitary operators is presented in Sec. (IV B), ergodic and non-ergodic channels are constructed by using block diagonal unitary operators in Sec. (IV C), and bounds on the convergence of any initial state to the fixed state in Sec. (IV D). Finally, we construct various single qubit channels using two-qubit unitary operators as an illustration in Sec. (IV E) and also investigate many-body quantum Hamiltonians (e.g. the Sachdev-Ye-Kitaev (SYK) model [61, 62] and experimentally realizable short-ranged model of interacting fermions in the presence of quasi-periodic potential [63] in Sec. (V). Strikingly, we find that mixing is the most common phenomenon in such systems. Moreover, we find the mixing time increases with the increase of the Hilbert space dimension, and in parallel, it also becomes harder for the system to meet the sufficiency condition of the mixing, which was derived in Sec. (IV A) in terms of the operator entanglement. Finally, we also qualitatively predict the scrambling time for such systems using the generalized spectral form factor (SFF).

III. QUANTUM CHANNELS

In this Section, we provide a brief introduction to the quantum channels and their spectral properties. In Sec. (III A), various representations of quantum channels and their interrelations are explained by explicitly expressing the linear operator acting on a single system in terms of the bipartite unitary operator. In Sec. (III B), we reproduce the spectral results for the quantum channel, which is required for our work. In Sec. (III C), we explain the definition of ergodic and mixing quantum

channels.

A. Representation of quantum channels

Let $\mathcal{H}^d$ is the Hilbert space of dimension d . The quantum states are represented by the density matrix ρ such that $\rho \geq 0$ and $\text{Tr}(\rho) = 1$, and the state space $\mathcal{S}(\mathcal{H})$ is defined over the set of density matrices. The map $\mathcal{E}$ is a linear, completely positive trace preserving (CPTP), defined over the state space $\mathcal{S}(\mathcal{H})$, and is called a quantum channel [32]. The action of $\mathcal{E}$ on the state ρ is

$$\rho' = \mathcal{E}(\rho), \quad (1)$$

and its corresponding linear operator is represented as $\mathcal{L}$ [52]

$$|\rho'\rangle = \mathcal{L}|\rho\rangle \quad (2)$$

where the vectorization is defined as $\langle i|\rho|j\rangle = \langle ij|\rho\rangle$. The linear operator $\mathcal{L}$ is a $d^2 \times d^2$ matrix, and we use bipartite index notation to represent the matrix, $\langle i\alpha|\mathcal{L}|j\beta\rangle \equiv \mathcal{L}_{i\alpha,j\beta}$. Note that the operator $\mathcal{L}$ is not Hermitian. The channel can be represented using an Hermitian operator, *dynamical matrix* [51, 54] $\mathcal{D}$, defined as

$$\mathcal{D} = \mathcal{L}^{R_2}, \quad (3)$$

where R_2 is a matrix reshaping operation defined as $\langle i\alpha|A|j\beta\rangle = \langle ij|A^{R_2}|\alpha\beta\rangle$. Expressing $\mathcal{D}$ in its eigenbasis as

$$\mathcal{D} = \sum_i |A_i\rangle\langle A_i|, \quad (4)$$

where we rescaled the basis as $|A_i\rangle = \sqrt{g_i}|\gamma_i\rangle$, with g_i and $|\gamma_i\rangle$ are eigenvalues and eigenstates of $\mathcal{D}$ respectively. The operator $\mathcal{L}$ we can be written as

$$\mathcal{L} = \sum_i g_i \gamma_i \otimes \gamma_i^* = \sum_i A_i \otimes A_i^*. \quad (5)$$

Here we used the identity $(A \otimes B)^{R_2} = |A\rangle\langle B^*|$. This provides a canonical form of Kraus representation with $\sum_i A_i^\dagger A_i = I$,

$$\mathcal{E}(\rho) = \sum_i A_i \rho A_i^\dagger. \quad (6)$$

The operator $\mathcal{D}$ is also called as *Choi matrix* [64] and represented as

$$\mathcal{D} = (\mathcal{E} \otimes I) |I\rangle\langle I|, \quad (7)$$

with the action of the channel can be recovered as

$$\mathcal{E}(\rho) = \text{Tr}_2[(I \otimes \rho^T)\mathcal{D}], \quad (8)$$

where T is the matrix transposition operation, I is the identity operator, and $|I\rangle = \sum_i |ii\rangle$.

Every channel $\mathcal{E}$ has its unitary representation in a larger dimension. The channel $\mathcal{E}$ acting on a single system in $\mathcal{H}^d$ is obtained from the unitary operator acting on the bipartite system in $\mathcal{H}^d \otimes \mathcal{H}^d$ by tracing the second system,

$$\mathcal{E}(\rho) = \text{Tr}_2[U(\rho \otimes \sigma)U^\dagger]. \quad (9)$$

This represents the physical scenario in which the joint system evolves under the unitary operator U , in which the second system can be treated as an environment (or bath). We are interested in the dynamics of the system alone. Let $\sigma = \sum_{ij} \lambda_{ij} |i\rangle\langle j|$ and for the channel $\mathcal{E}(\rho)$ in Eqn. (9), the matrix representation is

$$\begin{aligned} \langle \alpha | \mathcal{E}(\rho) | \beta \rangle &= \sum_\gamma \langle \alpha \gamma | U(\rho \otimes \sigma) U^\dagger | \beta \gamma \rangle \\ &= \sum_{\gamma i j m k} \lambda_{ij} \langle \alpha \gamma | U | m i \rangle \langle k j | U^\dagger | \beta \gamma \rangle \langle m | \rho | k \rangle \\ &= \sum_{\gamma i j m k} \lambda_{ij} \langle \alpha m | U^{R_2} | \gamma i \rangle \langle \gamma j | U^{R_2^\dagger} | \beta k \rangle \langle m k | \rho \rangle \\ &= \sum_{m k} \langle \alpha \beta | \left[U^{R_2} (I \otimes \sigma) U^{R_2^\dagger} \right]^{R_2} | m k \rangle \langle m k | \rho \rangle \end{aligned} \quad (10)$$

By referring Eq. (1) and Eq. (2), the matrix representation of $\mathcal{L}$ in terms of the unitary operator U is

$$\mathcal{L} = \left[U^{R_2} (I \otimes \sigma) U^{R_2^\dagger} \right]^{R_2}. \quad (11)$$

B. Spectral properties of channels

The ergodic properties of a channel depend largely on the spectral properties of quantum channels. For completeness, we comprehensively explain these results here [31, 32].

The eigenvalue equation of the channel $\mathcal{E}$ with eigenvalue λ and an eigenoperator X is

$$\mathcal{E}(X) = \lambda X. \quad (12)$$

The eigenvalues and eigenoperators are the solution of the characteristic equation $\det(\lambda I - \sum_i A_i \otimes A_i^*) = 0$. Denote $\text{spec}(\mathcal{E})$ as the spectrum of the channel $\mathcal{E}$. However, note that the channel $\mathcal{L}$ need not be diagonalizable.

Theorem 1. *The channel $\mathcal{E}$ has a density matrix ρ_* as a fixed point.*

Proof. The proof follows from the fact that the channel $\mathcal{E}$ is linear continuous mapping on the set of density matrices. Since the set of density matrices are convex and compact, due to Markov-Kakutani theorem [V.10.6 of Ref. [65]], the theorem follows [66]. $\square$

Corollary 1. If λ_0 is an eigenvalue of the channel with the density matrix as its eigenstate say ρ_* , i.e., $\mathcal{E}(\rho_*) = \lambda_0 \rho_*$, then $\lambda_0 = 1$

Proof. Since $\text{Tr}\{\rho_*\} = 1$, the proof follows from the trace-preserving property of the channel. $\square$

Theorem 2. λ is an eigenvalue of the channel $\mathcal{E}$ with an eigenvector X iff λ^* is also an eigenvalue with eigenvector $X^\dagger$.

Proof. Using the Kraus representation, we can express the action of the channel as $\mathcal{E}(\rho) = \sum_i A_i \rho A_i^\dagger$ where $\sum_i A_i^\dagger A_i = I$. So, starting with the eigenvalue equation,

$$\begin{aligned} \mathcal{E}(X) &= \lambda X \\ \iff \sum_i A_i X A_i^\dagger &= \lambda X \\ \iff \left(\sum_i A_i X A_i^\dagger \right)^\dagger &= (\lambda X)^\dagger \\ \iff \sum_i A_i X^\dagger A_i^\dagger &= \lambda^* X^\dagger \\ \iff \mathcal{E}(X^\dagger) &= \lambda^* X^\dagger \end{aligned} \quad (13)$$

$\square$

If $\mathcal{E}$ is a linear map, the adjoint map $\mathcal{E}^\dagger$ is defined as $\mathcal{E}^\dagger(X) = \sum_i A_i^\dagger X A_i$. The adjoint map $\mathcal{E}^\dagger$ of a completely positive map is also completely positive [32]. The adjoint of a CPTP map, a quantum channel, is an unital map. If the channel $\mathcal{E}$ is unital, then its adjoint $\mathcal{E}^\dagger$ is trace-preserving and thereby forms an unital channel. The spectrum of $\mathcal{E}$ and $\mathcal{E}^\dagger$ is related as follows: $\text{spec}(\mathcal{E}^\dagger) = \text{spec}(\mathcal{E})^*$, due to which a channel $\mathcal{E}$ and its adjoint $\mathcal{E}^\dagger$ has same spectrum.

Theorem 3. The eigenvalues of the channel $\mathcal{E}$ are bounded as:

$$|\lambda| \leq 1. \quad (14)$$

Proof. First we switch from the Schrödinger picture for $\mathcal{E}$ to the Heisenberg picture for $\mathcal{E}^\dagger$ induced by

$$\text{Tr}[\mathcal{E}(X)Y] = \text{Tr}[X\mathcal{E}^\dagger(Y)] \quad (15)$$

We also recall that linearity and complete positivity of $\mathcal{E}$ translates to linearity and complete positivity of $\mathcal{E}^\dagger$ whereas the trace-preserving property of $\mathcal{E}$ becomes unitality ($\mathcal{E}^\dagger(I) = I$) of $\mathcal{E}^\dagger$, and moreover, both $\mathcal{E}$ and $\mathcal{E}^\dagger$ have the same spectrum. So, proving the statement for $\mathcal{E}^\dagger$ will suffice.

Since for any $X \in \mathcal{H}^d$, $X/\|X\|_\infty$ is a contraction, a consequence of the theorem of Russo and Dye [67] can be stated as follows [68]:

$$\|\mathcal{E}^\dagger(X)\|_\infty \leq \|\mathcal{E}^\dagger(I)\|_\infty \|X\|_\infty. \quad (16)$$

Here $\|\cdot\|_\infty$ is an operator norm and for matrix it is defined as the maximum absolute row sum of the matrix. In the particular case where X is an eigenvector of $\mathcal{E}^\dagger$ with eigenvalue λ , we have

$$\begin{aligned} |\lambda| \|X\|_\infty &\leq \|\mathcal{E}^\dagger(I)\|_\infty \|X\|_\infty \\ \implies |\lambda| &\leq \|\mathcal{E}^\dagger(I)\|_\infty = \|I\|_\infty = 1 \end{aligned} \quad (17)$$

$\square$

The ergodic properties of quantum channels largely depend on their spectrum. Arrange the spectrum of $\mathcal{E}$ in the increasing order of magnitude as

$$\lambda_0 = 1 \geq |\lambda_1| \geq |\lambda_2| \cdots |\lambda_{d^2-1}|, \quad (18)$$

which will be used to define the various properties of quantum channels.

C. Definitions: Ergodic and mixing quantum channels

Ergodic theory is a mathematically well-characterized field [1]. All definitions related to the ergodic hierarchy are explained in various sources [69]. Here, we define ergodic and mixing of quantum channels [38, 41]. The essential aspect of the dynamical system is the measure space equipped with transformation. In the ergodic theory of quantum systems [70], the set of events is represented by the positive operator with the measure given by trace rule due to Gleason's theorem [71–73]. We focus on quantum channels as the transformations.

The composition of the channel $\mathcal{E}_1$ and $\mathcal{E}_2$ is denoted as $(\mathcal{E}_1 \circ \mathcal{E}_2)(\rho) = \mathcal{E}_1(\mathcal{E}_2(\rho))$. Note that if $\mathcal{E}$ is CPTP map, then any iterative application of the map

$$\mathcal{E}^n := \mathcal{E} \circ \cdots \circ \mathcal{E} \quad (19)$$

is also a CPTP map [32]. The following quantity, similar to taking an average, is also a quantum channel,

$$\Lambda_N^\mathcal{E}(\rho) := \frac{1}{N+1} \sum_{n=0}^N \mathcal{E}^n(\rho), \quad (20)$$

where $\mathcal{E}^0$, an identity channel.

Definition 1 (Ergodic [38]). A quantum channel is ergodic if there exists a unique state ρ_* such that

$$\mathcal{E}(\rho_*) = \rho_*. \quad (21)$$

Here ρ_* is the unique fixed point. In terms of the spectrum, the channel $\mathcal{E}$ is ergodic if and only if it has $\lambda_0 = 1$ and $\lambda_i \neq 1, \forall i \neq 0$. The ergodicity of the channel can be defined through the convergence of $\Lambda_N^\mathcal{E}$. The channel $\mathcal{E}$ is ergodic if and only if

$$\lim_{N \rightarrow \infty} \|\Lambda_N^\mathcal{E}(\rho) - \rho_*\|_1 = 0, \quad (22)$$

where $\|A\|_1 := \text{Tr}[\sqrt{A^\dagger A}]$ is the trace norm. It instructs us to see the convergence in terms of the function of operators. Let $\mathcal{M}_d$ is the set of $d \times d$ complex matrices. For any $A, B \in \mathcal{M}_d$, the quantum channel $\mathcal{E}$ is ergodic if and only if the following equivalent conditions holds [41],

$$\begin{aligned} \lim_{N \rightarrow \infty} \Lambda_N^\mathcal{E}(A) &= \text{Tr}(A)\rho_* \\ \lim_{N \rightarrow \infty} \frac{1}{N+1} \sum_{n=0}^N \text{Tr}(\mathcal{E}^n(A)B) &= \text{Tr}(A) \text{Tr}(B\rho_*). \end{aligned} \quad (23)$$

Definition 2 (Mixing [38]). $\mathcal{E}$ is mixing if there exists a unique state ρ_* such that

$$\lim_{n \rightarrow \infty} \|\mathcal{E}^n(\rho) - \rho_*\|_1 = 0 \quad \forall \rho \quad (24)$$

In terms of the spectrum, the channel $\mathcal{E}$ is mixing if and only if its spectrum has the property that $|\lambda_i| \neq 1, \forall i \neq 0$. Similar to the ergodic property, for any $A, B \in \mathcal{M}_d$, the equivalent definition of mixing holds if and only if the following equivalent conditions are satisfied,

$$\begin{aligned} \lim_{n \rightarrow \infty} \mathcal{E}^n(A) &= \text{Tr}(A)\rho_* \\ \lim_{n \rightarrow \infty} \text{Tr}(\mathcal{E}^n(A)B) &= \text{Tr}(A) \text{Tr}(B\rho_*). \end{aligned} \quad (25)$$

Mixing implies ergodicity but not the converse. If the spectrum is such that there exist λ_k for which $\lambda_k \neq 1$ but $|\lambda_k| = 1$. Such channels are ergodic but not mixing. In classical ergodic theory, further classifications of mixing channels like weak mixing and strong mixing are defined. It has been shown recently that in the case of quantum channels, weak mixing and strong mixing is equivalent to mixing [41], and we focus on mixing in this work.

IV. ERGODICITY, MIXING AND OPERATOR ENTANGLEMENT

In this Section, we study the ergodic and mixing nature of quantum channels defined in Eq. 9, by considering the spectrum of the quantum channel with its representation in terms of the unitary operator as in Eq. 11.

A. Sufficient condition for mixing

The Hilbert-Schmidt norm or Frobenius norm of an operator A is defined as $\|A\| = \sqrt{\text{Tr}(AA^\dagger)}$. By using an easily provable identity $\text{Tr}(X^{R_2} X^{R_2\dagger}) = \text{Tr}(XX^\dagger)$ for any square matrix X , we can see

$$\|\mathcal{L}\|^2 = \text{Tr}[(U^{R_2}(I \otimes \sigma)U^{R_2\dagger})^2]. \quad (26)$$

We are more interested in the case in which $\sigma = I/d$, and for this we have

$$\|\mathcal{L}\|^2 = \frac{1}{d^2} \text{Tr}[(U^{R_2}U^{R_2\dagger})^2]. \quad (27)$$

In order to understand the mixing of quantum channel $\mathcal{L}$ as a function of the unitary operator U , it is required to understand the entangling nature of the unitary operator. Physically, it encodes the interaction that the bipartite unitary operator U induces between the system and environment. Consider an operator Schmidt decomposition [74] of a unitary operator U

$$U = \sum_i \sqrt{\mu_i} A_i \otimes B_i, \quad (28)$$

where $\mu_i \geq 0$ and $\text{Tr}(A_i A_j^\dagger) = \text{Tr}(B_i B_j^\dagger) = \delta_{ij}$. The unitarity of U implies that $\sum_{i=0}^{d^2-1} \mu_i = d^2$, and hence μ_i/d^2 can be treated as probabilities and related entropies can be defined. By using linear entropy, the normalized operator entanglement $E(U)$ is defined as [59],

$$E(U) = \left(\frac{d^2}{d^2-1} \right) \left(1 - \frac{1}{d^4} \text{Tr}(U^{R_2}U^{R_2\dagger})^2 \right). \quad (29)$$

As we are interested in nontrivial eigenvalues $\lambda_{i \neq 0}$, let's express $\tilde{\mathcal{L}} = \mathcal{L} - |I\rangle\langle I|$ and the spectrum of $\tilde{\mathcal{L}}$ represents eigenvalues of $\mathcal{L}$ excluding $\lambda_0 = 1$,

$$\|\tilde{\mathcal{L}}\|^2 = \|\mathcal{L}\|^2 - 1 = (d^2-1)(1-E(U)). \quad (30)$$

Consider the Schur decomposition of $\tilde{\mathcal{L}} = V_1 T V_2$, where V_1 and V_2 are unitary operators and T is an upper triangular matrix with diagonal entries λ_i . The Hilbert-Schmidt norm is unitarily invariant which implies that $\|\mathcal{L}\| = \|T\|$, and since T is upper triangular matrix, it follows that

$$\sum_{i=1}^{d^2-1} |\lambda_i|^2 \leq (d^2-1)(1-E(U)). \quad (31)$$

Interesting results follow from the inequality (31). $|\lambda_{d^2-1}|^2(d^2-1) \leq \sum_{i=1}^{d^2-1} |\lambda_i|^2 \leq (d^2-1)(1-E(U))$ and we see that

$$|\lambda_{d^2-1}| \leq \sqrt{1-E(U)}. \quad (32)$$

This captures that whenever $E(U) \neq 0$, at least there exists one mode in which $|\lambda_{d^2-1}| < 1$.

The upper bound for $|\lambda_1|$ can be obtained similarly as $|\lambda_1|^2 \leq \sum_{i=1}^{d^2-1} |\lambda_i|^2 \leq (d^2-1)(1-E(U))$,

$$|\lambda_1| \leq \sqrt{(d^2-1)(1-E(U))}. \quad (33)$$

From this, it follows that whenever

$$E(U) > E_* = \frac{d^2-2}{d^2-1}, \quad (34)$$

guarantees $|\lambda_1| < 1$, which imply mixing. Thus, this provides a sufficient condition for the quantum channel to be mixed. A similar condition for discrete Hamiltonian dynamics holds for mixing in the many-body spacetime dual circuits in terms of entangling power [23].

For general σ , the spectrum is bounded as

$$\sum_{i=1}^{d^2-1} |\lambda_i|^2 \leq \text{Tr} \left[(U^{R_2} (I \otimes \sigma) U^{R_2^\dagger})^2 \right] - 1, \quad (35)$$

which provides the sufficiency condition for mixing $|\lambda_1| < 1$ if

$$\text{Tr} \left[(U^{R_2} (I \otimes \sigma) U^{R_2^\dagger})^2 \right] < 2. \quad (36)$$

If U is *dual-unitary*, $U^{R_2} U^{R_2^\dagger} = I$, then the purity of the state σ determines mixing nature, $|\lambda_1| < 1$ if

$$\text{Tr}(\sigma^2) < \frac{2}{d}. \quad (37)$$

For $d = 2$, any mixed state guarantees the mixing, and for general d , the bound (37) provides the sufficiency condition for mixing.

B. Local unitary stability of mixing condition

Local unitary operators act as a disorder. Two unitary operators U and U' are said to be local unitarily (LU) equivalent if U' can be written as

$$U' = (u_1 \otimes u_2) U (u_3 \otimes u_4) \quad (38)$$

where $u_i \in \mathcal{H}^d$ are any unitary operators. The spectrum of the quantum channel $\mathcal{E}(\rho)$ with U and U' with $\mathcal{E}'(\rho) = \text{Tr}_2[U'(\rho \otimes \sigma)U'^\dagger]$ are not identical. The linear operator $\mathcal{L}'$ for the channel $\mathcal{E}'$ is

$$\begin{aligned} \mathcal{L}' &= \left[U'^{R_2} (I \otimes \sigma) U'^{R_2^\dagger} \right]^{R_2} \\ &= \left[(u_1 \otimes u_3^T) \left[U^{R_2} (I \otimes u_4 \sigma u_4^\dagger) U^{R_2^\dagger} \right] (u_1^\dagger \otimes u_3^*) \right]^{R_2} \\ &= (u_1 \otimes u_1^*) \left[U^{R_2} (I \otimes \sigma') U^{R_2^\dagger} \right] (u_3 \otimes u_3^*) \end{aligned} \quad (39)$$

where $\sigma' = u_4 \sigma u_4^\dagger$. Here we have used the identity

$$[(u_1 \otimes u_2) A (u_3 \otimes u_4)]^{R_2} = (u_1 \otimes u_3^T) A^{R_2} (u_2^T \otimes u_4). \quad (40)$$

However, it can be seen that $\|\mathcal{L}\|^2 = \|\mathcal{L}'\|^2$ and hence the sufficient condition for mixing remains invariant under any local unitary operations of the form Eq. (38). For $\sigma = I/d$, the eigenvalue $|\lambda_1|$ is upper bounded by the operator entanglement $E(U)$ and $E(U)$ is LU equivalent quantity. Even for general σ it can be seen the sufficient condition is invariant with local unitary operators. The bound provides stable mixing criteria that are invariant under any local unitary operators.

C. Block diagonal construction of quantum channels

We further study the ergodic hierarchy of channels using block diagonal unitary operators. Define block diagonal unitary operator U_D as $U_D = \bigoplus_{i=1}^d u_i$ in which u_i are $d \times d$ unitary operators. For $\sigma = I/d$ the linear operator $\mathcal{L}[U_D]$,

$$\mathcal{L}[U_D] = \frac{1}{d} [U_D^{R_2} U_D^{R_2^\dagger}]^{R_2}. \quad (41)$$

The $U_D^{R_2}$ of block diagonal matrix contains d rows and the inner product of these rows forms the d^2 elements of $U_D^{R_2} U_D^{R_2^\dagger}$, and hence $\mathcal{L}[U_D]$ is diagonal with diagonal entries

$$\lambda_{ij} = \frac{1}{d} \text{Tr}(u_i u_j^\dagger). \quad (42)$$

There are d eigenvalues $\lambda_{ii} = 1$, and hence the channel is non-ergodic. If we choose an orthonormal unitary operator set u_j , then there are d eigenvalue $\lambda_{ii} = 1$, and the remaining eigenvalues vanish. This is an interesting case as the channel is non-ergodic as a whole but there are d eigenmodes which is highly mixing. If $u_j = u$, all the eigenvalues $\lambda_{ij} = 1$ and an example of integrable channel.

D. Bound on the convergence

The convergence of any state ρ to the fixed state ρ_* under the action of map $\mathcal{E}$ can be bounded as [66]

$$\|\mathcal{E}^n(\rho) - \rho_*\|_1 \leq C_d \text{poly}(n) |\lambda_1|^n, \quad (43)$$

where C_d is some constant depending on the local dimensions d . Due to the upper bound on $|\lambda_1|$ in Eq. (33), we can write

$$\begin{aligned} \|\mathcal{E}^n(\rho) - \rho_*\|_1 &\leq C_d \text{poly}(n) \left((d^2 - 1)(1 - E(U)) \right)^{\frac{n}{2}} \\ &\leq C(d, n) (1 - E(U))^{\frac{n}{2}}. \end{aligned} \quad (44)$$

The operator entanglement $E(U)$ thus provides an upper bound to the convergence of ρ to ρ_* for a fixed iteration n .




Figure 2. Set of all two-qubit unitary LU equivalent unitary operators represented in the tetrahedron called Weyl chamber. Here the coordinates represent the parameters of unitary operators in Eq. (45).

E. Two-qubit unitary operators and single qubit quantum channels

For simplest scenerio of $d = 2$, any two qubit unitary operator U is unitarily equivalent to U_{nl} [75],

$$U_{nl} = \exp[-i(x\sigma_x \otimes \sigma_x + y\sigma_y \otimes \sigma_y + z\sigma_z \otimes \sigma_z)]$$

$$= \begin{pmatrix} e^{-iz}c_- & 0 & 0 & -ie^{-iz}s_- \\ 0 & e^{iz}c_+ & -ie^{iz}s_+ & 0 \\ 0 & -ie^{iz}s_+ & e^{iz}c_+ & 0 \\ -ie^{-iz}s_- & 0 & 0 & e^{-iz}c_- \end{pmatrix}, \quad (45)$$

where $c_{\pm} = \cos(x \pm y)$, $s_{\pm} = \sin(x \pm y)$ and $0 \leq |c_z| \leq c_y \leq c_x \leq \frac{\pi}{4}$. For such U_{nl} with $\sigma = I/d$, $\mathcal{L}$ can be explicitly diagonalized as

$$\mathcal{L} = \sum_{i=0}^3 \tilde{\lambda}_i |\phi_i\rangle\langle\phi_i|, \quad (46)$$

where

$$|\phi_0\rangle = \frac{|00\rangle + |11\rangle}{\sqrt{2}} \quad |\phi_1\rangle = \frac{|00\rangle - |11\rangle}{\sqrt{2}}$$

$$|\phi_2\rangle = \frac{|01\rangle + |10\rangle}{\sqrt{2}} \quad |\phi_3\rangle = \frac{|01\rangle - |10\rangle}{\sqrt{2}} \quad (47)$$

The eigenvalues with trivial $\lambda_0 = \tilde{\lambda}_0 = 1$ and unordered eigenvalues are

$$\tilde{\lambda}_1 = \frac{1}{2} [\cos(2(x+y)) + \cos(2(x-y))] \quad (48)$$

$$\tilde{\lambda}_2 = \cos(2y) \cos(2z), \quad \tilde{\lambda}_3 = \cos(2x) \cos(2z).$$




Figure 3. Variation of the real part vs. the imaginary part of the eigenvalue spectrum of the matrix representation of quantum channel $\mathcal{L}_H$ for the SYK model and the Hamiltonian H_{SR} .

The variables (x, y, z) in Eq. (45) uniquely represent a set of LU equivalent unitary operators [75]. It forms a tetrahedron called the Weyl chamber, as shown in Fig. (2). The ergodic properties of qubit quantum channel (46) can be characterized using these variables as :

1. **Case 1:** for the line 'LOCAL' to 'CNOT' :: for which (x, y, z) varies from $(0, 0, 0)$ to $(\pi/4, 0, 0)$. The ordered eigenvalues can be then $\lambda_1 = \tilde{\lambda}_2 = 1$ and $\lambda_2 = \lambda_3 = \tilde{\lambda}_1 = \tilde{\lambda}_3 = \cos 2x$. The entire line represents a non-ergodic channel. For $x = 0$, local, $\lambda_2 = \lambda_3 = 1$ represents integrable channel.
2. **Case 2:** for the line 'CNOT' to 'DCNOT' :: for which (x, y, z) varies from $(\pi/4, 0, 0)$ to $(\pi/4, \pi/4, 0)$. We have $\lambda_1 = \tilde{\lambda}_2 = \cos 2y$ and $\lambda_2 = \lambda_3 = \tilde{\lambda}_1 = \tilde{\lambda}_3 = 0$. Except $y = \pi/4$ i.e., CNOT gate, all other unitaries with will have $\lambda_1 = \cos(2y) < 1$. Single qubit channels with these unitaries forms mixing channels.
3. **Case 3:** The qubit channel corresponding to the unitary operators from line from SWAP to DCNOT for which (x, y, z) varies from $(\pi/4, \pi/4, 0)$ to $(\pi/4, \pi/4, \pi/4)$, represents completely depolarizing channel with $\lambda_i = 0 \quad \forall i \neq 0$.
4. **Case 4:** For the line LOCAL to DCNOT and from LOCAL to SWAP all represent mixing vertices.

V. ERGODIC AND MIXING CHANNELS WITH MANY-BODY HAMILTONIANS

In the previous section, we study a two-qubit system, i.e., the Hilbert space dimension of the system




Figure 4. Variation of the 1st two eigenvalues of $\mathcal{L}_H$, λ_0 and $|\lambda_1|$ (after arranging in descending order in magnitudes) with h for the Hamiltonian H_{SR} . Inset shows the approach of $|\lambda_1|$ to 1 with increasing h in semi-log scale.




Figure 5. Variation of $\Delta_n = \|\mathcal{L}_H^n \rho - \rho_*\|_1$ with n for the SYK model and the Hamiltonian H_{SR} for the initial Neel state. Inset shows the variation of the operator entanglement with h for the Hamiltonian H_{SR} .

$d = 2$ (the Hilbert space dimension of the system+ bath $d^2 = 4$). In this section, we focus on the many-body quantum systems where $d \geq 2$. We study two types of many-body Hamiltonian of interacting fermions on a one-dimensional lattice of length L . The first one is an experimentally realizable [63, 76] short-range (SR) model consisting of nearest-neighbor hopping and nearest-neighbor interaction in the presence of quasi-periodic potential. The other one is a model with random all-to-all interactions, famously known as the SYK model. The Hamiltonians for these two models are




Figure 6. (Left panel) Variation of operator entanglement $E(U)$ with h and for the Hamiltonian H_{SR} for different values of d . The dotted lines correspond to E_* . (Right panel) Variation of the spectral gap $1 - |\lambda_1|$ with h for the Hamiltonian H_{SR} and for different values of d .

given by:

$$H_{SR} = - \sum_{i=1}^{L-1} (\hat{c}_i^\dagger \hat{c}_{i+1} + \text{H.c.}) + V \sum_{i=1}^{L-1} \hat{n}_i \hat{n}_{i+1} + h \sum_{i=1}^L \cos(2\pi\alpha i) \hat{n}_i, \quad (49)$$

and

$$H_{SYK} = \sum_{i,j,k,l}^L J_{ij,kl} \hat{c}_i^\dagger \hat{c}_j^\dagger \hat{c}_k \hat{c}_l, \quad (50)$$

Where, $\hat{c}_i^\dagger$ ($\hat{c}_i$) is the fermionic creation (annihilation) operator at site i , $\hat{n}_i = \hat{c}_i^\dagger \hat{c}_i$ is the number operator. $\hat{c}_i^\dagger$ and $\hat{c}_i$ obey following anti-commutation relations i.e. $\{\hat{c}_i, \hat{c}_j^\dagger\} = \delta_{ij}$, $\{\hat{c}_i, \hat{c}_j\} = 0$, and $\{\hat{c}_i^\dagger, \hat{c}_j^\dagger\} = 0$. In the case of the Hamiltonian H_{SR} , α is an irrational number which we choose to be $\alpha = (\sqrt{5} - 1)/2$. We also set the interaction parameter $V = 1$ for all our calculations. In the absence of the interaction, i.e., $V = 0$, H_{SR} undergoes a delocalization-localization transition as one tunes the strength of the quasi-periodic potential, and $h = 2$ turns out to be the transition point [77]. In the presence of interaction, it is believed that within an isolated quantum dynamics framework, changing the strength of the quasi-periodic potential h , one can drive the system from ergodic to many-body localized (MBL) phase [78], though, in recent days, the stability of MBL transition in the thermodynamic limit has been questioned [79].

On the other, the SYK Hamiltonian H_{SYK} in the large L limit is exactly solvable [62, 80]. The study of this model has uncovered a remarkable direct connection

to quantum gravity with a black hole in AdS_2 [81–83]. As in the case of a black hole, the SYK model gives rise to thermalization and many-body quantum chaos with a Lyapunov exponent that saturates the quantum limit [84]. The random four-fermion coupling J_{ijkl} drawn from a Gaussian distribution with zero mean and variance $|J_{ijkl}|^2 = J^2/L^3$; J_{ijkl} are properly antisymmetrized, i.e., $J_{ijkl} = -J_{jikl} = -J_{ijlk}$ and $J_{ijkl} = J_{klji}^*$ [61]. It is important to point out that though both of these systems are fermionic in nature, using Jordan-Wigner (JW) transformation, they can be mapped into models of spin-1/2 particles (qubit). Interestingly, even though JW transformation is non-local, the Hamiltonian H_{SR} becomes the spin-1/2 XXZ model in the presence of the incommensurate magnetic field, which involves spin-exchange term only between nearest sites. On the other hand, the SYK model has all-to-all interactions. Recently, both the SYK model and the H_{SR} model have been realized in quantum computing platforms using proper Qubitization and Trotterization [85–87].

We consider the first $L/2$ sites as the system and the rest of the $L/2$ sites as the bath. Hence, the Hilbert space dimension of the sub-system and the bath are $d = 2^{L/2}$. To understand the ergodic hierarchy of these systems according to the formalism described in Sec. III, we need to investigate the matrix $\mathcal{L}$ (see Eq. 11). Here, for all the calculations, we consider an infinite temperature bath; hence, we replace σ in Eq. 11 by I_d/d (where I_d is a d dimensional identity matrix) and the Unitary operator $U = e^{iH} = U_H$ (where $H = H_{SR}$ or $H = H_{SYK}$). Eq. 11 reads as,

$$\mathcal{L}_H = \frac{1}{d} \left[U_H^{R_2} U_H^{R_2^\dagger} \right]^{R_2}. \quad (51)$$

First, we investigate the eigenvalue spectrum of the matrix $\mathcal{L}_H$. Figure 3 shows the eigenvalue spectrum of $\mathcal{L}_H$ in the complex plane, where we plot the real part vs. the imaginary part of the eigenvalues λ_i . We find that for the Hamiltonian H_{SR} , as we increase h , the effective distance from the center of most of the data points increases, indicating the absolute value of a large fraction of eigenvalues $|\lambda_i|$ approaches toward 1. On the other hand, for the SYK model, most of the data points are in the neighborhood close to the center.

Moreover, we arrange the spectrum in descending order of magnitude (as mentioned in Eq. (18)), plot λ_0 (which is real) and $|\lambda_1|$ as a function of h in Fig. 4. While $\lambda_0 = 1$, $|\lambda_1|$ approaches towards 1 with increasing h . However, the inset of Fig. 4 ensures that for a finite value of h , $|\lambda_1|$ does not reach 1, though $1 - |\lambda_1|$ approaches to zero with increasing h . It is important to emphasize that in the $h \rightarrow \infty$ limit, U_H will be completely diagonal. Hence, it is a non-ergodic channel (see Sec. (IVC)). $|\lambda_1|$ for the SYK model for $L = 8$ turns

out to be 0.5275, which is much smaller than $|\lambda_1|$ obtained for H_{SR} . Intuitively, one expects that given the SYK model is maximally chaotic. This confirms that the quantum channel representations of the Hamiltonian H_{SR} and H_{SYK} show mixing.

To strengthen the claim, we also plot $\Delta_n = \|\mathcal{L}_H^n \rho - \rho_*\|_1$ vs n . For a mixing channel $\Delta_n \rightarrow 0$ as $n \rightarrow \infty$, $\forall \rho$ and where $\mathcal{L}_H \rho_* = \rho_*$. The results shown in Fig. 5 are for Neel state $|\psi\rangle = \prod_{i=1}^{L/2} \hat{c}_{2i-1}^\dagger |0\rangle$ [88–90]; however, we check our results for a few other states as well. We find that, indeed, Δ_n approaches zero as n increases. As expected, with increasing h in H_{SR} , the fall-off of Δ_n becomes slower. On the other hand, the decay of Δ_n is extremely fast for the SYK model. Moreover, we also study the variation of the operator entanglement $E(U)$ (see Eq. (29)) with h for the Hamiltonian H_{SR} in the inset of Fig. 5 and find that it decreases with increasing h , and always remains below E_* (see Eq. (34)). On the other hand, for the SYK model (for $L = 8$), $E(U) \simeq 0.9925$ is very close to E_* .

Next, we also investigate the effect of the Hilbert space dimension of the system d in the ergodic hierarchy. In Fig. 6 (right panel), we show the variation of the spectral gap $1 - |\lambda_1|$ (a measure of the mixing time) with h of the short-range Hamiltonian H_{SR} for different values of d . The spectral gap appears to decrease quickly with increasing d . It says that as d increases, the mixing time decreases for the short-range model. Similar features can also be observed in the variation of the operator entanglement vs. h plot in Fig. 6 (left panel). Interestingly, we find that only for $d = 2$ and for a certain parameter range ($h \in [0, 3]$), $E(U)$ satisfies the sufficient condition of the mixing, i.e., $E(U) > E_*$. For $d > 2$, $E(U)$ always appears to be less than E_* . Intuitively, these results can be understood in the following way. In the case of the H_{SR} , the coupling term H_{sb} (according to schematic Fig. 1) between the system and the bath involves only two adjacent sites, does not depend on the system size L or the Hilbert space dimension of the system d . Hence, one would expect that if L (or d) increases, the bath will take more time to thermalize the system; hence, the mixing time is expected to go down with increasing d . On the contrary, for the all-to-all SYK model, this is not the case; hence, there is not much of a dependence of the mixing time on d .

Further, we also make a quantitative connection between thermalization (measured in terms of scrambling [91]) and the ergodic hierarchy using the spectral form factor (SFF) [92–94], a statistical measure of quantum chaos with that of the ergodic hierarchy. The generalized SFF for a quantum channel is defined as [91],

$$K := \frac{1}{d^2} \sum_i |\text{Tr } A_i|^2 \quad (52)$$

where A_i are the Kraus operator of the dynamical equa-




Figure 7. Variation of generalized SFF $K(n)$ with n for the Hamiltonian H_{SR} . Inset shows the variation of the scrambling time n_s (iteration) with h . The dashed horizontal line corresponds to $1/d$.

tion (see Eq. (6)). By using a simple identity $|X|^2 = \text{Tr}(X \otimes X^*)$, we can write generalized SFF as

$$K(n) = \frac{1}{d^2} \text{Tr} \mathcal{L}_H^n, \quad (53)$$

where d stands for the Hilbert space dimension of the system. It has been argued that the necessary condition for scrambling in many-body systems is $K(n) < 1/d$ [91]. In terms of the eigenvalues of $\mathcal{L}$, the generalized SFF

$$K(n) = \frac{1}{d^2} \left(1 + \sum_{i=1}^{d^2-1} \lambda_i^n \right). \quad (54)$$

It is very interesting to note that in the block diagonal construction of quantum channels constructed in Sec. (IV C), there exists a model for which d eigenvalues equals $\lambda_i = 1$ and remaining eigenvalues vanish giving rise to a non-thermalizing model. From the ergodic theory perspective, the model itself is non-ergodic, and it matches with that of the many-body perspectives.

Given the necessary condition for scrambling has been argued as $K(n) < 1/d$ [91], here, we roughly estimate the scrambling time by identifying $n_s = \min(n)$, such as the $K(n_s) \leq 1/d$. We have considered $H = H_{SR}$, computed $K(n)$ as a function of n . Previously, we have shown in Fig. 3 and Fig. 4 that for any finite value h , the channel $\mathcal{L}_H$ is mixing. However, as we increase the h , the spectral gap $1 - |\lambda_1|$ (which is also a measure of mixing time) decreases. One would naively expect that mixing time and the scrambling time should be related to each other. If one observes the decrease in mixing time, similar features should also be noticed in the context of scrambling. Figure. 7 validates our hypothesis; we show the variation of $K(n)$ vs. n for different values of h . As expected, with increasing h , the fall-off of

$K(n)$ with n is slower. Moreover, we also identify the scrambling time (iteration) n_s and find that, indeed, it increases with increasing h .

VI. CONCLUSION AND DISCUSSIONS

In this work, we study the ergodic hierarchy of quantum channels by proving the necessary condition for the channel to be mixing in terms of operator entanglement of the unitary operator, and we use these tools to study the mixing in quantum many-body systems. We also prove the stability of the sufficiency of the mixing condition under any single particle local unitary operator. We construct various analytical models using block diagonal unitary operators to explore the ergodic hierarchy of quantum channels. As an illustration, by using the canonical two-qubit unitary operator, we construct various quantum channels of a single qubit that show different ergodic behaviors. Then, we explore unitary operators constructed from quantum many-body Hamiltonians, especially focused on two models: 1) short-range interacting models in the presence of quasiperiodic potential and 2) all-to-all interacting SYK model. We find both the models show mixing, while SYK, being maximally chaotic, almost satisfies the sufficiency conditions for mixing; the other model fails to do so. Also, studying different spectral measures of quantum channels and the convergence criteria, it was apparent that the mixing rate for the SYK model is much higher compared to the short-ranged model. Intuitively, it fits well with our notion that the SYK is maximally chaotic in the thermodynamic limit. However, both in the two-qubit case and for the many-body systems, we could not find an example where the channel is ergodic but not mixing. It indicates that maybe in typical quantum systems mixing is more common, while ergodicity is a rare phenomena. We also study the quantitative relation between mixing and scrambling using spectral form factor.

In summary, based on our results, we make a few conjectures for generic quantum many-body systems, and they are 1) In the context of quantum many-body systems, mixing is much more common than other ergodic hierarchies, 2) The mixing time decreases with increasing d for the short-range model, 3) As d increases, the sufficient conditions of mixing in terms of the operator entanglement become harder to achieve. 4) The mixing and scrambling times are typically expected to behave similarly for a generic quantum system. However, we must emphasize one can possibly cook up an extremely fine-tuned model where some of our above conjectures may get violated; it will be interesting to search for such models in future studies.

In the last century, the mathematics of ergodic theory

has provided many interesting results and opened up many applications [95]. It would be intriguing to explore how far one can draw parallels from the world of classical ergodic theory to the world of quantum channels. In particular, one might be able to characterize total ergodicity (all powers of the given transformation are ergodic) [96] and prove that it sits distinctively in between ergodicity and weak mixing, but on the other hand, the notion of mild mixing [96] defined in any conceivable way will be equivalent to mixing as it would sit between mixing and weak mixing. It would be interesting to see the relation between the ergodic theory of

quantum channels and its information-theoretic capabilities.

VII. ACKNOWLEDGEMENTS

RM acknowledges the DST-Inspire fellowship by the Department of Science and Technology, Government of India, SERB start-up grant (SRG/2021/002152). SA acknowledges the start-up research grant from SERB, Department of Science and Technology, Govt. of India (SRG/2022/000467).

-
- [1] I. P. Cornfeld, S. V. Fomin, and Ya. G. Sinai, *Ergodic Theory*, Vol. 245 (Springer, 1982).
 - [2] Donald Ornstein, "Bernoulli shifts with the same entropy are isomorphic," *Advances in Mathematics* **4**, 337–352 (1970).
 - [3] Matthew Foreman, Daniel J. Rudolph, and Benjamin Weiss, "The conjugacy problem in ergodic theory," *Annals of Mathematics* **173**, 1529–1586 (2011).
 - [4] John von Neumann, "Zur operatorenmethode in der klassischen mechanik," *Annals of Mathematics* **33**, 587–642 (1932).
 - [5] John von Neumann, "Proof of the ergodic theorem and the h-theorem in quantum mechanics: Translation of: Beweis des ergodensatzes und des h-theorems in der neuen mechanik," *The European Physical Journal H* **35**, 201–237 (2010).
 - [6] J v Neumann, "Beweis des ergodensatzes und des h-theorems in der neuen mechanik," *Zeitschrift für Physik* **57**, 30–70 (1929).
 - [7] George M Zaslavsky, "Stochasticity in quantum systems," *Physics Reports* **80**, 157–250 (1981).
 - [8] Asher Peres, "Ergodicity and mixing in quantum theory. i," *Phys. Rev. A* **30**, 504–508 (1984).
 - [9] Mario Feingold, Nimrod Moiseyev, and Asher Peres, "Ergodicity and mixing in quantum theory. ii," *Phys. Rev. A* **30**, 509–511 (1984).
 - [10] Sheldon Goldstein, Joel L Lebowitz, Roderich Tumulka, and Nino Zanghi, "Long-time behavior of macroscopic quantum systems: Commentary accompanying the english translation of john von neumann's 1929 article on the quantum ergodic theorem," *The European Physical Journal H* **35**, 173–200 (2010).
 - [11] Tomaž Prosen, "Time evolution of a quantum many-body system: Transition from integrability to ergodicity in the thermodynamic limit," *Physical review letters* **80**, 1808 (1998).
 - [12] Roberto Artuso, Giulio Casati, and Italo Guarneri, "Numerical study on ergodic properties of triangular billiards," *Physical Review E* **55**, 6384 (1997).
 - [13] Giulio Casati and Tomaž Prosen, "Mixing property of triangular billiards," *Physical Review Letters* **83**, 4729 (1999).
 - [14] Tomaž Prosen, "Ergodic properties of a generic nonintegrable quantum many-body system in the thermodynamic limit," *Physical Review E* **60**, 3949 (1999).
 - [15] Tomaž Prosen, "General relation between quantum ergodicity and fidelity of quantum dynamics," *Physical Review E* **65**, 036208 (2002).
 - [16] Martin Horvat, Mirko Degli Esposti, Stefano Isola, Tomaž Prosen, and Leonid Bunimovich, "On ergodic and mixing properties of the triangle map," *Physica D: Nonlinear Phenomena* **238**, 395–415 (2009).
 - [17] Dongliang Zhang, HT Quan, Biao Wu, *et al.*, "Ergodicity and mixing in quantum dynamics," *Physical Review E* **94**, 022150 (2016).
 - [18] Luca D'Alessio, Yariv Kafri, Anatoli Polkovnikov, and Marcos Rigol, "From quantum chaos and eigenstate thermalization to statistical mechanics and thermodynamics," *Advances in Physics* **65**, 239–362 (2016), <https://doi.org/10.1080/00018732.2016.1198134>.
 - [19] M Akila, D Waltner, B Gutkin, and T Guhr, "Particle-time duality in the kicked Ising spin chain," *Journal of Physics A: Mathematical and Theoretical* **49**, 375101 (2016).
 - [20] Bruno Bertini, Pavel Kos, and Tomaž Prosen, "Exact correlation functions for dual-unitary lattice models in 1 + 1 dimensions," *Phys. Rev. Lett.* **123**, 210601 (2019).
 - [21] Boris Gutkin, Petr Braun, Maram Akila, Daniel Waltner, and Thomas Guhr, "Exact local correlations in kicked chains," *Phys. Rev. B* **102**, 174307 (2020).
 - [22] Petr Braun, Daniel Waltner, Maram Akila, Boris Gutkin, and Thomas Guhr, "Transition from quantum chaos to localization in spin chains," *Phys. Rev. E* **101**, 052201 (2020).
 - [23] S. Aravinda, Suhail Ahmad Rather, and Arul Lakshminarayan, "From dual-unitary to quantum Bernoulli circuits: Role of the entangling power in constructing a quantum ergodic hierarchy," *Phys. Rev. Research* **3**, 043034 (2021).
 - [24] Immanuel Bloch, Jean Dalibard, and Wilhelm Zwerger, "Many-body physics with ultracold gases," *Rev. Mod. Phys.* **80**, 885–964 (2008).
 - [25] M. A. Cazalilla, R. Citro, T. Giamarchi, E. Orignac, and M. Rigol, "One dimensional bosons: From condensed matter systems to ultracold gases," *Rev. Mod. Phys.* **83**, 1405–1466 (2011).
 - [26] Lea F. Santos and Marcos Rigol, "Onset of quantum chaos in one-dimensional bosonic and fermionic sys-

- tems and its relation to thermalization," *Phys. Rev. E* **81**, 036206 (2010).
- [27] Marcos Rigol, Vanja Dunjko, Vladimir Yurovsky, and Maxim Olshanii, "Relaxation in a completely integrable many-body quantum system: An ab initio study of the dynamics of the highly excited states of 1d lattice hard-core bosons," *Phys. Rev. Lett.* **98**, 050405 (2007).
- [28] Marcos Rigol and Mark Srednicki, "Alternatives to eigenstate thermalization," *Phys. Rev. Lett.* **108**, 110601 (2012).
- [29] Mark Srednicki, "Chaos and quantum thermalization," *Phys. Rev. E* **50**, 888–901 (1994).
- [30] J. M. Deutsch, "Quantum statistical mechanics in a closed system," *Phys. Rev. A* **43**, 2046–2049 (1991).
- [31] Michael M Wolf, "Quantum channels & operations: Guided tour," Lecture notes available at <http://www-m5.ma.tum.de/foswiki/pub/M5,13> (2012).
- [32] Mark M Wilde, *Quantum information theory* (Cambridge university press, 2013).
- [33] Vern Paulsen, *Completely bounded maps and operator algebras*, 78 (Cambridge University Press, 2002).
- [34] Rajendra Bhatia, "Positive definite matrices," in *Positive Definite Matrices* (Princeton university press, 2009).
- [35] Robert Alicki and Karl Lendi, *Quantum dynamical semigroups and applications*, Vol. 717 (Springer, 2007).
- [36] Alexander S Holevo, *Statistical structure of quantum theory*, Vol. 67 (Springer Science & Business Media, 2003).
- [37] Daniel Burgarth and Vittorio Giovannetti, "The generalized iyapunov theorem and its application to quantum channels," *New Journal of Physics* **9**, 150 (2007).
- [38] Daniel Burgarth, Giulio Chiribella, Vittorio Giovannetti, Paolo Perinotti, and Kazuya Yuasa, "Ergodic and mixing quantum channels in finite dimensions," *New Journal of Physics* **15**, 073045 (2013).
- [39] Ramis Movassagh and Jeffrey Schenker, "An ergodic theorem for homogeneously distributed quantum channels with applications to matrix product states," in *American Physical Society (March Meeting)* (2020).
- [40] Ramis Movassagh and Jeffrey Schenker, "Theory of ergodic quantum processes," *Physical Review X* **11**, 041001 (2021).
- [41] Satvik Singh, Nilanjana Datta, and Ion Nechita, "Ergodic theory of diagonal orthogonal covariant quantum channels," arXiv preprint arXiv:2206.01145 (2022).
- [42] Maxim Raginsky, "Strictly contractive quantum channels and physically realizable quantum computers," *Physical Review A* **65**, 032306 (2002).
- [43] Robin Blume-Kohout, Hui Khoon Ng, David Poulin, and Lorenza Viola, "Characterizing the structure of preserved information in quantum processes," *Physical review letters* **100**, 030501 (2008).
- [44] Robin Blume-Kohout, Hui Khoon Ng, David Poulin, and Lorenza Viola, "Information-preserving structures: A general framework for quantum zero-error information," *Physical Review A* **82**, 062306 (2010).
- [45] Vittorio Giovannetti, Simone Montangero, and Rosario Fazio, "Quantum multiscale entanglement renormalization ansatz channels," *Physical review letters* **101**, 180503 (2008).
- [46] Wen-Long Ma, Shu-Shen Li, and Ren-Bao Liu, "Sequential generalized measurements: Asymptotics, typicality, and emergent projective measurements," *Physical Review A* **107**, 012217 (2023).
- [47] Noah Linden and Paul Skrzypczyk, "How to use arbitrary measuring devices to perform almost perfect measurements," arXiv preprint arXiv:2203.02593 (2022).
- [48] Victor V Albert, "Asymptotics of quantum channels: conserved quantities, an adiabatic limit, and matrix product states," *Quantum* **3**, 151 (2019).
- [49] Daniele Amato, Paolo Facchi, and Arturo Konderak, "Asymptotics of quantum channels," *Journal of Physics A: Mathematical and Theoretical* **56**, 265304 (2023).
- [50] Satvik Singh, Mizanur Rahaman, and Nilanjana Datta, "Zero-error communication, scrambling, and ergodicity," arXiv preprint arXiv:2402.18703 (2024).
- [51] Karol Życzkowski and Ingemar Bengtsson, "On duality between quantum maps and quantum states," *Open Systems & Information Dynamics* **11**, 3–42 (2004).
- [52] Ingemar Bengtsson and Karol Życzkowski, *Geometry of quantum states: an introduction to quantum entanglement* (Cambridge University Press, 2007).
- [53] Clinton J Oxenrider and Richard D Hill, "On the matrix reorderings γ and ψ ," *Linear algebra and its applications* **69**, 205–212 (1985).
- [54] ECG Sudarshan, PM Mathews, and Jayaseetha Rau, "Stochastic dynamics of quantum-mechanical systems," *Physical Review* **121**, 920 (1961).
- [55] Kai Chen and Ling-An Wu, "A matrix realignment method for recognizing entanglement," arXiv preprint quant-ph/0205017 (2002).
- [56] Oliver Rudolph, "Some properties of the computable cross-norm criterion for separability," *Physical Review A* **67**, 032312 (2003).
- [57] Suhail Ahmad Rather, S. Aravinda, and Arul Lakshminarayan, "Creating ensembles of dual unitary and maximally entangling quantum evolutions," *Phys. Rev. Lett.* **125**, 070501 (2020).
- [58] Suhail Ahmad Rather, S Aravinda, and Arul Lakshminarayan, "Construction and local equivalence of dual-unitary operators: from dynamical maps to quantum combinatorial designs," *PRX Quantum* **3**, 040331 (2022).
- [59] Paolo Zanardi, "Entanglement of quantum evolutions," *Phys. Rev. A* **63**, 040304 (2001).
- [60] Paolo Zanardi, Christof Zalka, and Lara Faoro, "Entangling power of quantum evolutions," *Phys. Rev. A* **62**, 030301 (2000).
- [61] Subir Sachdev, "Bekenstein-hawking entropy and strange metals," *Phys. Rev. X* **5**, 041025 (2015).
- [62] Subir Sachdev and Jinwu Ye, "Gapless spin-fluid ground state in a random quantum heisenberg magnet," *Phys. Rev. Lett.* **70**, 3339–3342 (1993).
- [63] Michael Schreiber, Sean S Hodgman, Pranjal Bordia, Henrik P Lüschen, Mark H Fischer, Ronen Vosk, Ehud Altman, Ulrich Schneider, and Immanuel Bloch, "Observation of many-body localization of interacting fermions in a quasirandom optical lattice," *Science* **349**, 842–845 (2015).
- [64] Man-Duen Choi, "Completely positive linear maps on complex matrices," *Linear algebra and its applications* **10**, 285–290 (1975).
- [65] Nelson Dunford and Jacob T Schwartz, *Linear operators, part 1: general theory*, Vol. 10 (John Wiley & Sons, 1988).

- [66] Barbara M Terhal and David P DiVincenzo, "Problem of equilibration and the computation of correlation functions on a quantum computer," *Physical Review A* **61**, 022301 (2000).
- [67] B. Russo and H. A. Dye, "A note on unitary operators in C^* -algebras," *Duke Math. J.* **33**, 413–416 (1966).
- [68] Jean-Christophe Bourin and Eun-Young Lee, "On the russo-dye theorem for positive linear maps," *Linear Algebra and its Applications* **571**, 92–102 (2019).
- [69] Joseph Berkovitz, Roman Frigg, and Fred Kronz, "The ergodic hierarchy, randomness and hamiltonian chaos," *Studies in History and Philosophy of Science Part B: Studies in History and Philosophy of Modern Physics* **37**, 661–691 (2006).
- [70] Robert Alicki and Mark Fannes, "Quantum dynamical systems," (2001).
- [71] Andrew M Gleason, "Measures on the closed subspaces of a hilbert space," in *The Logico-Algebraic Approach to Quantum Mechanics: Volume I: Historical Evolution* (Springer, 1975) pp. 123–133.
- [72] Paul Busch, "Quantum states and generalized observables: a simple proof of gleason's theorem," *Physical Review Letters* **91**, 120403 (2003).
- [73] Stephen M Barnett, James D Cresser, John Jeffers, and David T Pegg, "Quantum probability rule: a generalization of the theorems of gleason and busch," *New Journal of Physics* **16**, 043025 (2014).
- [74] Michael A Nielsen, Christopher M Dawson, Jennifer L Dodd, Alexei Gilchrist, Duncan Mortimer, Tobias J Osborne, Michael J Bremner, Aram W Harrow, and Andrew Hines, "Quantum dynamics as a physical resource," *Physical Review A* **67**, 052301 (2003).
- [75] Jun Zhang, Jiri Vala, Shankar Sastry, and K. Birgitta Whaley, "Geometric theory of nonlocal two-qubit operations," *Phys. Rev. A* **67**, 042313 (2003).
- [76] Dmitry A Abanin, Ehud Altman, Immanuel Bloch, and Maksym Serbyn, "Colloquium: Many-body localization, thermalization, and entanglement," *Reviews of Modern Physics* **91**, 021001 (2019).
- [77] Serge Aubry and Gilles André, "Analyticity breaking and anderson localization in incommensurate lattices," *Ann. Israel Phys. Soc* **3**, 18 (1980).
- [78] Shankar Iyer, Vadim Oganesyan, Gil Refael, and David A. Huse, "Many-body localization in a quasiperiodic system," *Phys. Rev. B* **87**, 134202 (2013).
- [79] Jan Šuntajs, Janez Bonča, Tomaž Prosen, and Lev Vidmar, "Quantum chaos challenges many-body localization," *Phys. Rev. E* **102**, 062144 (2020).
- [80] Olivier Parcollet and Antoine Georges, "Transition from overscreening to underscreening in the multichannel kondo model: Exact solution at large N ," *Phys. Rev. Lett.* **79**, 4665–4668 (1997).
- [81] Subir Sachdev, "Holographic metals and the fractionalized fermi liquid," *Phys. Rev. Lett.* **105**, 151602 (2010).
- [82] Subir Sachdev, "Bekenstein-hawking entropy and strange metals," *Phys. Rev. X* **5**, 041025 (2015).
- [83] Juan Maldacena and Douglas Stanford, "Remarks on the sachdev-ye-kitaev model," *Phys. Rev. D* **94**, 106002 (2016).
- [84] Juan Maldacena, Stephen H Shenker, and Douglas Stanford, "A bound on chaos," *Journal of High Energy Physics* **2016**, 1–17 (2016).
- [85] Adam Smith, MS Kim, Frank Pollmann, and Johannes Knolle, "Simulating quantum many-body dynamics on a current digital quantum computer," *npj Quantum Information* **5**, 106 (2019).
- [86] L. García-Álvarez, I. L. Egusquiza, L. Lamata, A. del Campo, J. Sonner, and E. Solano, "Digital quantum simulation of minimal AdS/CFT," *Phys. Rev. Lett.* **119**, 040501 (2017).
- [87] Muhammad Asaduzzaman, Raghav G Jha, and Bharath Sambasivam, "A model of quantum gravity on a noisy quantum computer," *arXiv preprint arXiv:2311.17991* (2023).
- [88] Vincenzo Alba and Pasquale Calabrese, "Entanglement dynamics after quantum quenches in generic integrable systems," *SciPost Physics* **4**, 017 (2018).
- [89] S. Aravinda and Ranjan Modak, "Complexity growth for one-dimensional free-fermionic lattice models," *Phys. Rev. B* **108**, 064309 (2023).
- [90] Ranjan Modak, Vincenzo Alba, and Pasquale Calabrese, "Entanglement revivals as a probe of scrambling in finite quantum systems," *Journal of Statistical Mechanics: Theory and Experiment* **2020**, 083110 (2020).
- [91] Amit Vikram and Victor Galitski, "Exact universal bounds on quantum dynamics and fast scrambling," *Phys. Rev. Lett.* **132**, 040402 (2024).
- [92] Fritz Haake, *Quantum signatures of chaos* (Springer, 1991).
- [93] Sebastian Müller, Stefan Heusler, Petr Braun, Fritz Haake, and Alexander Altland, "Semiclassical foundation of universality in quantum chaos," *Physical review letters* **93**, 014103 (2004).
- [94] Amit Vikram and Victor Galitski, "Dynamical quantum ergodicity from energy level statistics," *Physical Review Research* **5**, 033126 (2023).
- [95] Anatole Katok, AB Katok, and Boris Hasselblatt, *Introduction to the modern theory of dynamical systems*, 54 (Cambridge university press, 1995).
- [96] Tanja Eisner, Bálint Farkas, Markus Haase, and Rainer Nagel, *Operator Theoretic Aspects of Ergodic Theory*, Vol. 272 (Springer, 2015).