

## Hyperelliptic families and 4d $\mathcal{N} = 2$ SCFT

---

Dan Xie<sup>a</sup>, Zekai Yu<sup>b</sup>

<sup>a</sup>*Department of Mathematics, Tsinghua University, Beijing, 100084, China*

<sup>b</sup>*Qiuzhen College, Tsinghua University, Beijing, 100084, China*

ABSTRACT: We classify four dimensional  $\mathcal{N} = 2$  SCFTs whose Seiberg-Witten (SW) geometries can be written as hyperelliptic families. By using special Kähler condition of SW geometry, we reduce the problem to one parameter quasi-homogeneous hyperelliptic families  $y^2 = f(x, t)$ . The classification is given by further demanding that the complex algebraic surface defined by  $y^2 = f(x, t)$  has an isolated singularity. We then write down the full SW geometry by looking at mini-versal deformations of the one parameter family, and the SW differential is also written down. The detailed physical data for these theories are found by matching the theory with other known construction. Our solutions recover the known rank one and rank two results, and give some infinite sequences valid at arbitrary ranks. We also studied  $Z_2$  quotient of above hyperelliptic families which give rise to  $B$  type and  $D$  type conformal gauge theory, and further generalizations.

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Hyperelliptic families and SCFTs</b>	<b>2</b>
2.1	Detailed data of SCFTs: type I-IV	6
2.2	Detailed data of SCFTs: type V-VIII	7
<b>3</b>	<b><math>\mathbb{Z}_2</math> quotient</b>	<b>10</b>
<b>4</b>	<b>SW differential and other possibilities</b>	<b>11</b>
<b>5</b>	<b>Conclusion</b>	<b>13</b>

---

## 1 Introduction

The Coulomb branch solution of a general  $\mathcal{N} = 2$  theory can be described by finding a family of algebraic varieties  $F(z_i, u_i, \lambda_i, m_i) = 0$  fibered over the space of generalized Coulomb branch parameters involving expectation value  $u_i$  of Coulomb branch operators, mass parameters  $m_i$ , and coupling constants  $\lambda_i$  (including relevant couplings and exactly marginal deformations). One also need to find a SW differential  $\lambda_{sw}$  to describe the central charge of BPS particles. The combination of varieties  $F$  and  $\lambda_{sw}$  are called SW geometry, see [1, 2] for the SW curves<sup>1</sup> of  $SU(2)$  gauge theory.

The SW geometry is usually found through string/M theory-motivated methods [3, 4] and the connection of SW solution to integrable system [5, 6]. Most SW geometries were discovered by following two methods: a) The spectral curve of an integral system, and the most well studied ones are the so-called Hitchin system [5, 7–11]; b) The mini-versal deformation of an isolated three dimensional canonical singularities [12, 13].

While a large number of SW geometries can be found using the methods above, one should not forget that the resulting geometries are often written in a complicated way which makes it hard to perform a detailed study of the low energy physics. This is in particular true for some SW geometries engineered using Hitchin system. Often an equivalent but simpler SW geometry would make the physical exploration much more viable.

Holding this in mind, here we try to classify SW geometry which can be expressed in terms of the simplest algebraic curves, i.e. the hyperelliptic curves:

$$y^2 = f(x, u_i, \lambda_i, m_i);$$

---

<sup>1</sup>The SW geometries for some 4d  $\mathcal{N} = 2$  theories must be described by a high dimensional algebraic varieties.

Here  $f$  is a degree  $2g + 1$  or  $2g + 2$  polynomial in  $x$ , and  $u_i, i = 1, \dots, g$  denotes the expectation value of Coulomb branch operators. We focus on the SW geometry for the SCFT, and perform the classification through the following steps:

1. In the first step, we turn off the mass parameters and coupling constants, so that the geometry  $y^2 = f(x, u_i)$  have a  $\mathbb{C}^*$  action to reflect the  $U(1)_R$  symmetry of a SCFT.
2. We then impose the special Kähler condition which implies that the derivative of SW differential  $\lambda_{sw}$  should be in the space of holomorphic differential of the hyperelliptic curve [14].

$$\frac{\partial \lambda_{sw}}{\partial u_i} \in H^{1,0}.$$

The simplest solution would indicate that the SW curve and SW differential are written in the form 2.5 and 2.6.

3. The above two considerations reduce the classification into finding out possible one-parameter quasi-homogeneous families  $y^2 = f(x, t)$ . We impose a further condition, namely the surface singularity defined by the equation is isolated, and so a complete classification of one-parameter geometries is possible, see 2.8 for the resulting 8 types of geometries.
4. We then find the full SW geometry by looking at the versal deformations of one parameter geometry, and then replace the parameter  $t$  by a polynomial in  $x$  which involves VEV of Coulomb branch operators and possibly other coupling constants.

The detailed studies of those theories are listed in table. [1,2,3,4,5,6,7,8,9]. Our list recovers the rank one [15, 16] and rank two results [17], and generalizes them to arbitrary rank.

Two further generalizations are studied: first we study the  $\mathbb{Z}_2$  quotient of the hyperelliptic families and get three classes of theories, see Table. 10 and 11. These can be thought of as the arbitrary rank-generalization of the rank one  $I_4$  series studied in [18], and include the  $B$  type conformal gauge theory. We then also study the case where the square root of the holomorphic differential is used, which would describe  $D$  type conformal gauge theory, and generalizations.

This paper is organized as follows: section 2 gave a detailed illustration of our classification strategy and also the detailed physical properties of those theories, by matching them with known constructions;  $\mathbb{Z}_2$  quotients of the curves are studied in section 3; section 4 discusses the SW differential and the  $D$  type gauge theory; A conclusion is given in section 5.

## 2 Hyperelliptic families and SCFTs

First we assume that the SW curve of a 4d  $\mathcal{N} = 2$  SCFT is given by a family of genus  $g$  hyperelliptic curves:

$$y^2 = f(x, u_i) \tag{2.1}$$

where  $f(x, u_i)$  is a polynomial in  $x$  of degree  $2g + 1$  or  $2g + 2$ . Here  $u_i$ ,  $i = 1, \dots, g$  denote the Coulomb branch parameters. To get the SW geometry of a SCFT, the above curve needs to have a  $\mathbb{C}^*$  action acting on  $(x, y, u_i)$  which is the incarnation of the (complexified)  $U(1)_R$  symmetry of the SCFT.

The SW differential is given by  $\lambda_{sw} = R(x, y, u_i)dx$  with  $R$  a rational function of  $(x, y, u_i)$ , and the derivative of  $\lambda_{sw}$  with respect to Coulomb branch operator is [19]:

$$\frac{\partial}{\partial u_i} \int_{\lambda} \lambda_{sw} = \frac{\partial}{\partial u_i} \int_{\lambda} R dx = \int \left( \frac{\partial R}{\partial u_i} - \frac{\partial R}{\partial y} \left( \frac{\partial F}{\partial y} \right)^{-1} \frac{\partial F}{\partial u_i} \right) dx; \quad (2.2)$$

Here  $\lambda$  is the cycle on Riemann surface, and  $F$  is the equation  $F = y^2 - f(x, u_i) = 0$ . The special Kähler condition requires that [14]:

$$\frac{\partial \lambda_{sw}}{\partial u_i} \in H^{1,0}(C); \quad (2.3)$$

Here  $H^{1,0}(C)$  denotes the space of holomorphic differential of the hyperelliptic curve  $C$ , which has a canonical basis

$$\omega_i = \frac{x^i dx}{y}, \quad i = 0, \dots, g - 1.$$

Let  $t$  denote the Coulomb branch operator with maximal scaling dimension, and assume that the derivative of SW differential with respect of  $t$  is proportional to  $\frac{dx}{y}$ , i.e.

$$\frac{\partial \lambda_{sw}}{\partial t} = \left( \frac{\partial R}{\partial t} - \frac{\partial R}{\partial y} \left( \frac{\partial F}{\partial y} \right)^{-1} \frac{\partial F}{\partial t} \right) dx = c \frac{dx}{y}; \quad (2.4)$$

Here  $c$  is a nonzero constant. Now if the SW curve takes the following special form

$$\boxed{y^2 = f(x, u_{g-1}x^{g-1} + u_{g-2}x^{g-2} + \dots + u_1x + t).} \quad (2.5)$$

and similarly the SW differential takes the form

$$\boxed{\lambda_{sw} = R(x, y, u_{g-1}x^{g-1} + u_{g-2}x^{g-2} + \dots + u_1x + t)dx.} \quad (2.6)$$

Then we have

$$\frac{\partial \lambda_{sw}}{\partial u_i} = \left( \frac{\partial R}{\partial u_i} - \frac{\partial R}{\partial y} \left( \frac{\partial F}{\partial y} \right)^{-1} \frac{\partial F}{\partial u_i} \right) dx = x^i \frac{\partial \lambda_{sw}}{\partial t} = c \frac{x^i dx}{y}; \quad (2.7)$$

Here we used the fact  $\frac{\partial R}{\partial u_i} = x^i \frac{\partial R}{\partial t}$ , and  $\frac{\partial F}{\partial u_i} = x^i \frac{\partial F}{\partial t}$ . So the special Kähler condition is satisfied automatically for the SW curve 2.5 and SW differential 2.6, as long as the equation 2.4 has a solution (we will discuss this issue in section 4).

**One parameter curve:** By looking at the curve 2.5 and 2.6, one can see that the SW family is determined by the curve with just one parameter, which corresponds to the Coulomb branch operator of maximal scaling dimension. The classification of theories (or SW geometries) is now reduced to the classification of the one parameter hyperelliptic curve  $y^2 = f(x, t)$  with a  $\mathbb{C}^*$  action. We impose a further constraint on the one parameter

family: There is an isolated singularity (of the complex surface defined by  $y^2 = f(x, t)$ ) at the origin  $x = y = t = 0$ . Such one parameter hyperelliptic families can be easily classified as follows:

$$\begin{aligned}
I : y^2 &= x^{2g+1} + t^a, \\
II : y^2 &= x^{2g+2} + t^a, \\
III : y^2 &= x(x^{2g} + t^a), \\
IV : y^2 &= x(x^{2g+1} + t^a), \\
V : y^2 &= t(x^{g+2} + t^a), \\
VI : y^2 &= t(x^{g+3} + t^a), \\
VII : y^2 &= xt(x^{g+1} + t^a), \\
VIII : y^2 &= xt(x^{g+2} + t^a).
\end{aligned} \tag{2.8}$$

**Proof:** The isolated quasi-homogeneous singularity of two variables  $(x, t)$  is classified as  $x^a + t^b, x^a + xt^b, x^at + y^bt$  [20], so one has the above 8 types.

**Coulomb branch spectrum:** Now one can proceed to the calculation of Coulomb branch spectrum of the theory. The basic idea is that the SW differential  $\lambda_{sw}$  gives the mass of BPS particle and so it has scaling dimension one. The equation 2.7 gives a relation for the scaling dimension

$$1 - [u_i] = i[x] - [y];$$

We find that

$$[u_{i-1}] - [u_i] = [x];$$

So if the scaling dimension of  $x$  is positive, the maximal scaling dimension of Coulomb branch operators is  $[t]$  and the gap of the Coulomb branch spectrum is given by  $[x]$ . The  $\mathbb{C}^*$  action of the curve  $y^2 = f(x, t)$  yields two equations relating the scaling dimensions, which would help us to completely determine the spectrum.

*Example:* Let's consider one parameter curve  $y^2 = x^{2g+1} + t^2$ . We have the following three equations

$$\begin{aligned}
2[y] &= (2g + 1)[x], \\
2[y] &= 2[t], \\
1 - [t] &= [x] - [y].
\end{aligned}$$

and so we find  $[t] = \frac{g+1}{2}$ ,  $[x] = 1$ , so the Coulomb branch spectrum is  $\frac{3}{2}, \frac{5}{2}, \dots, \frac{2g+1}{2}$ .

**Full SW curve:** One can turn on exactly marginal deformations, relevant deformations, and mass deformations without lifting the Coulomb branch. These are encoded in the deformations of above SW curves. They can be derived as follows:

1. To get the SW curve with general Coulomb branch deformations, we replace the parameter  $t$  in the one parameter curve by a degree  $g - 1$  polynomial in  $x$ . More generally, substitution by a polynomial in higher degree in  $x$  is possible, as long as

the scaling dimensions of the parameters that come with the higher order terms are non-negative:

$$t \rightarrow \dots + \lambda_2 x^{g+1} + \lambda_1 x^g + u_{g-1} x^{g-1} + \dots + t.$$

The parameters  $\lambda_1, \lambda_2, \dots$  includes mass parameters, relevant deformations, and exactly marginal deformations. Extra care must be taken to ensure that the above replacement do not affect the genus of the curve.

2. Mass deformations can be found by finding the mini-versal deformation of the one parameter curve. The mini-versal deformation of an isolated singularity of our type is defined as follows

$$y^2 = f(x, t) + \sum_{\alpha} \lambda_{\alpha} \phi_{\alpha}; \quad (2.9)$$

Here  $\phi_{\alpha}$  is the monomial basis of the Jacobi algebra defined as follows

$$J_f = \frac{C[x, t]}{\left\{ \frac{\partial f}{\partial x}, \frac{\partial f}{\partial t} \right\}}.$$

The scaling dimension of  $\lambda_{\alpha}$  could be larger than or equal to one. Parameters with scaling dimension integral and larger than one should be interpreted as the Casimiers of the non-abelian flavor symmetry.

*Example:* Let's consider the one parameter curve  $y^2 = x(x^{2g} + t^2)$ , and the scaling dimensions are  $[x] = 2, [t] = 2g$ . The basis of Jacobi algebra  $J_f$  is

$$J_f = \{t, x^{2g}, x^{2g-1}, \dots, 1\};$$

and so there are a total of  $2g + 2$  mass parameters. The deformed one parameter curve is

$$y^2 = x^{2g+1} + xt^2 + [T_2 x^{2g} + T_4 x^{2g-1} + \dots + T_{4g+2}] + T'_{2g+2} t.$$

Now the replacement of  $t$  (with maximal number of deformations) is

$$t \rightarrow \tau x^g + u_{g-1} x^{g-1} + \dots + t;$$

We see that there is an exactly marginal deformation represented by . The full SW curve is

$$y^2 = x^{2g+1} + x(\tau x^g + u_g x^{g-1} + \dots + t)^2 + [T_2 x^{2g} + T_4 x^{2g-1} + \dots + T_{4g+2}] + T'_{2g+2}(\tau x^g + u_{g-1} x^{g-1} + \dots + t).$$

and the scaling dimension of the mass deformations  $(T_2, T_4, \dots, T_{4g+2}, T'_{2g+2})$  is  $(2, 4, \dots, 4g+2, 2g+2)$ , so the flavor symmetry is  $SO(4g+4)$ .

## 2.1 Detailed data of SCFTs: type I-IV

Now we study those SCFTs whose SW curve is given by the form 2.5. The one parameter family 2.8 is *I, II, III, IV*. These families share the property that the one parameter family already has genus  $g$ .

We list the Coulomb branch spectrum, the central charge, flavor symmetry, and the number of  $A_1$  singularities for the generic deformations. The corresponding periodic map [21]. Factorization of the related mapping class group element is also listed (see [22]), which would be important to understand the low energy physics [23]. The  $(A_1, G)$  sequence was first found in [24], and the class  $S$  construction was first found in [4].

**Table 1:** Type I:  $y^2 = x^{2g+1} + t^a$ .

Theory	$(A_1, A_{2g})$	$D_2(SU(2g+1))$	
curve	$y^2 = x^{2g+1} + t$	$y^2 = x^{2g+1} + t^2$	$y^2 = x^{2g+1} + t^a, a < \frac{4g+2}{2g-1}$
CB spectrum	$\frac{2g+4}{2g+3}, \frac{2g+6}{2g+3}, \dots, \frac{4g+2}{2g+3}$	$\frac{3}{2}, \frac{5}{2}, \dots, \frac{2g+1}{2}$	$a \leq 5, g = 1$
periodic map	$\frac{1}{2(2g+1)} + \frac{1}{2} + \frac{g}{2g+1}$	$\frac{g}{2g+1} + \frac{g}{2g+1} + \frac{1}{2g+1}$	$a \leq 3, g = 2$
factorization	$\phi_1 = \tau_1 \tau_2 \dots \tau_{2g}$	$\phi_1^2$	$a \leq 2, g \geq 3$
central charges	$a = \frac{g(24g+19)}{24(2g+3)}, c = \frac{g(6g+5)}{6(2g+3)}$	$a = \frac{7(g^2+g)}{24}, c = \frac{g^2+g}{3}$	
flavor symmetry	None	$su(2g+1)$	
# $A_1$ singularity	$2g$	$4g$	
Full SW curve $a = 1$	$y^2 = x^{2g+1} + (\lambda_1 x^{2g-1} + \lambda_2 x^{2g-2} + \dots + u_{g-1} x^{g-1} + \dots + t)$		
Full SW curve $a = 2$	$y^2 = x^{2g+1} + (T_2 x^{2g-1} + \dots + T_{2g+1}) + (\lambda_1 x^g + u_{g-1} x^{g-1} + \dots + t)^2$		
Full SW curve $a = 3, g = 2$ flavor $e_8$	$y^2 = x^5 + (u_4 x + t)^3 + (T_{12} x^3 + T_{18} x^2 + T_{24} x + T_{30}) + (T_2 x^3 + T_8 x^2 + T_{14} x + T_{20})(u_4 x + t)$		

**Table 2:** Type II:  $y^2 = x^{2g+2} + t^a$ .

Theory	$(A_1, A_{2g+1})$	$SU(g+1) - (2g+2)$	
curve	$y^2 = x^{2g+2} + t$	$y^2 = x^{2g+2} + t^2$	$y^2 = x^{2g+2} + t^a$
CB spectrum	$\frac{g+3}{g+2}, \dots, \frac{2g+2}{g+2}$	$2, 3, \dots, g+1$	$a < \frac{2g+2}{g}$
periodic map	$\frac{1}{2g+2} + \frac{1}{2g+2} + \frac{g}{g+1}$	$\frac{1}{g+1} + \frac{1}{g+1} + \frac{g}{g+1} + \frac{g}{g+1}$	$a \leq 3, g = 1$
factorization	$\phi_2 = \tau_1 \tau_2 \dots \tau_{2g} \tau_{2g}$	$\phi_2^2$	$a \leq 2, g \geq 2$
central charges	$a = \frac{12g^2+19g+2}{24(g+2)}, c = \frac{3g^2+5g+1}{6(g+2)}$	$a = \frac{7g^2+14g+2}{24}, c = \frac{2g^2+4g+1}{6}$	
flavor symmetry	$u(1)$	$su(2g+2) \times u(1)$	
# $A_1$ singularity	$2g+1$	$4g+2$	
Full SW curve $a = 1$	$y^2 = x^{2g+2} + (\lambda_1 x^{2g} + \lambda_2 x^{2g-1} + \dots + m x^g + u_{g-1} x^{g-1} + \dots + t)$		
Full SW curve $a = 2$	$y^2 = x^{2g+2} + (T_2 x^{2g} + \dots + T_{2g+2}) + (\tau x^{g+1} + m x^g + u_{g-1} x^{g-1} + \dots + t)^2$		

**Table 3:** Type III:  $y^2 = x(x^{2g} + t^a)$ .

Theory	$(A_1, D_{2g+1})$	$USp(2g) - (2g + 2)$	
curve	$y^2 = x(x^{2g} + t)$	$y^2 = x(x^{2g} + t^2)$	$y^2 = x(x^{2g} + t^a), a < \frac{4g}{2g-1}$
CB spectrum	$\frac{2g+2}{2g+1}, \dots, \frac{4g}{2g+1}$	$2, 4, \dots, 2g$	$a \leq 3, g = 1$
periodic map	$\frac{1}{4g} + \frac{2g-1}{4g} + \frac{1}{2}$	$\frac{1}{2} + \frac{1}{2} + \frac{1}{2g} + \frac{2g-1}{2g}$	$a \leq 2, g \geq 2$
factorization	$\phi_3 = \tau_1 \tau_2 \dots \tau_{2g} \tau_{2g+1}$	$\phi_3^2$	$\phi_3^a$
central charges	$a = \frac{g(8g+3)}{8(2g+1)}, c = \frac{g}{2}$	$a = \frac{14g^2+9g}{24}, c = \frac{4g^2+3g}{6}$	
flavor symmetry	$su(2)$	$so(4g + 4)$	
# $A_1$ singularity	$2g + 1$	$4g + 2$	
Full SW curve $a = 1$	$y^2 = x^{2g+1} + x(\lambda_1 x^{2g-1} + \dots + u_{g-1} x^{g-1} + \dots + t) + T_2$		
Full SW curve $a = 2$	$y^2 = x^{2g+1} + x(\tau x^g + u_{g-1} x^{g-1} + \dots + t)^2 + [T_2 x^{2g} + T_4 x^{2g-1} + \dots + T_{4g+2}] + T'_{2g+2}(\tau x^g + u_{g-1} x^{g-1} + \dots + t)$		

**Table 4:** Type IV:  $y^2 = x(x^{2g+1} + t^a)$ .

Theory	$(A_1, D_{2g+2})$	Class S	
curve	$y^2 = x(x^{2g+1} + t)$	$y^2 = x(x^{2g+1} + t^2)$	$y^2 = x(x^{2g+1} + t^a), a < \frac{2g+1}{g}$
CB spectrum	$\frac{g+2}{g+1}, \dots, \frac{2g+1}{g+1}$	$3, 5, \dots, 2g + 1$	$a \leq 2$
periodic map	$\frac{1}{2g+1} + \frac{1}{2g+1} + \frac{2g-1}{2g+1}$	$\frac{2g}{2g+1} + \frac{g+1}{2g+1} + \frac{g+1}{2g+1}$	
factorization	$\phi_4$	$\phi_4^2$	$\phi_4^a$
central charges	$a = \frac{g}{2} + \frac{1}{12}, c = \frac{g}{2} + \frac{1}{6}$	$a = \frac{14g^2+23g+4}{24}, c = \frac{4g^2+7g+2}{6}$	
flavor symmetry	$su(2) \times u(1)$	$so(4g + 6) \times u(1) (E_6 \text{ if } g = 1)$	
# $A_1$ singularity	$2g + 2$	$4g + 4$	
Full SW curve $a = 1$	$y^2 = x^{2g+2} + x(\lambda_1 x^{2g} + \dots + m x^g + u_{g-1} x^{g-1} + \dots + t) + T_2$		
Full SW curve $a = 2$	$y^2 = x^{2g+2} + x(m x^g + u_{g-1} x^{g-1} + \dots + t)^2 + [T_2 x^{2g+1} + T_4 x^{2g-1} + \dots + T_{4g+4}] + T'_{2g+3}(m x^g + u_{g-1} x^{g-1} + \dots + t)$		

## 2.2 Detailed data of SCFTs: type V-VIII

The one parameter families of those theories have genus less than  $g$ , so that substituting  $t$  by a degree  $g-1$  polynomial in  $x$  would give us a genus  $g$  hyperelliptic family. The periodic map is for the one parameter curve.

**Table 5:** Type V:  $y^2 = t(x^{g+2} + t^a)$ .

Theory		
curve	$y^2 = t(x^{g+2} + t)$	$y^2 = t(x^{g+2} + t^a), a < \frac{g+2}{g}$
CB spectrum	$3, 4, \dots, g+2$	$a \leq 1$
Class S	$([g, 1^2], [1^{g+2}], [1^{g+2}])$	
Periodic map	$\frac{g+1}{g+2} + \frac{g+1}{g+2} + \frac{2}{g+2} g \text{ odd}$ $\frac{g+1}{g+2} + \frac{g+1}{g+2} + \frac{1}{(g+2)/2} g \text{ even}$	
central charges	$a = \frac{7g^2+28g+6}{24}, c = \frac{2g^2+8g+3}{6}$	
flavor	$su(2g+4) \times su(2)(E_6 \text{ if } g=1)$	
# $A_1$ singularities	$4g+4$	
Full SW curve $a=1$	$y^2 = (T'_2 x^g + u_{g-1} x^{g-1} + \dots + t)[x^{g+2} + (T'_2 x^g + u_{g-1} x^{g-1} + \dots + t)]$ $+ [T_2 x^{2g+2} + T_3 x^{2g+1} + \dots + T_{2g+4}]$	

**Table 6:** Type VI:  $y^2 = t(x^{g+3} + t^a)$ .

Theory		
curve	$y^2 = t(x^{g+3} + t)$	$y^2 = t(x^{g+3} + t^a), a < \frac{g+3}{g+1}$
CB spectrum	$4, 5, \dots, g+3$	$a \leq 1$
Class S	$[(g+1), 2], [1^{g+3}]$	
Periodic map	$\frac{g+2}{g+3} + \frac{g+2}{g+3} + \frac{2}{g+3} g \text{ even}$ $\frac{g+2}{g+3} + \frac{g+2}{g+3} + \frac{1}{(g+3)/2} g \text{ odd}$	
central charges	$a = \frac{7g^2+42g+10}{24}, c = \frac{2g^2+12g+5}{6}$	
flavor	$su(2g+6)$	
# $A_1$ singularities	$4g+5$	
Full SW curve $a=1$	$y^2 = (u_{g-1} x^{g-1} + \dots + t)[x^{g+3} + (u_{g-1} x^{g-1} + \dots + t)]$ $+ [T_2 x^{2g+4} + T_3 x^{2g-1} + \dots + T_{2g+6}]$	

**Table 7:** Type VII:  $y^2 = xt(x^{g+1} + t^a)$ .

Theory		
curve	$y^2 = xt(x^{g+1} + t)$	$y^2 = xt(x^{g+1} + t^a), a < \frac{g+1}{g}$
CB spectrum	$4, 6, \dots, 2g + 2$	$a \leq 1$
Class S	$[(g+1)^2], [g^2, 1^2], [1^{2g+2}]$	
Periodic map	$\frac{2g+1}{2g+2} + \frac{g+2}{2g+2} + \frac{1}{2} g \text{ odd}$ $\frac{2g+1}{2g+2} + \frac{(g+2)/2}{g+1} + \frac{1}{2} g \text{ even}$	
central charges	$a = \frac{14g^2+37g+8}{24}, c = \frac{4g^2+11g+4}{6}$	
flavor	$so(4g+8) \times su(2)$	
# $A_1$ singularities	$4g+5$	
Full SW curve $a = 1$	$y^2 = x(T'_2x^g + u_{g-1}x^{g-1} + \dots + t)[x^{g+1} + (T'_2x^g + u_{g-1}x^{g-1} + \dots + t)]$ $+ [T_2x^{2g+2} + T_4x^{2g+1} + \dots + T_{4g+6}] + T'_{2g+4}(T'_2x^g + u_{g-1}x^{g-1} + \dots + t)$	

**Table 8:** Type VIII:  $y^2 = xt(x^{g+2} + t^a)$ .

Theory		
curve	$y^2 = xt(x^{g+2} + t)$	$y^2 = xt(x^{g+2} + t^a), a < \frac{g+2}{g+1}$
CB spectrum	$6, 8, \dots, 2g + 4$	$a \leq 1$
Class S	$[(g+2)^2], [(g+1)^2, 2], [1^{2g+4}]$	
Periodic map	$\frac{2g+3}{2g+4} + \frac{g+3}{2g+4} + \frac{1}{2} g \text{ even}$ $\frac{2g+3}{2g+4} + \frac{(g+3)/2}{g+2} + \frac{1}{2} g \text{ odd}$	
central charges	$a = \frac{14g^2+65g+16}{24}, c = \frac{4g^2+19g+8}{6}$	
flavor	$so(4g+12)$	
# $A_1$ singularities	$4g+6$	
Full SW curve $a = 1$	$y^2 = x(u_{g-1}x^{g-1} + \dots + t)[x^{g+2} + (u_{g-1}x^{g-1} + \dots + t)] +$ $[T_2x^{2g+4} + T_4x^{2g+3} + \dots + T_{4g+10}] + T'_{2g+10}(u_{g-1}x^{g-1} + \dots + t)$	

**Table 9:** The case when  $[x] = 0$ .

Theory	$2 - \underbrace{SU(2) - \dots - SU(2)}_g - 2$	
curve	$y^2 = t(x^{g+3} + \tau_1 x^{g+2} + \dots + \tau_{g+2})$	
CB spectrum	$2, \dots, 2$	
Class S	$[1^2] \times (g+3)$	
central charges	$a = \frac{19g+4}{24}, c = \frac{5g+2}{6}$	
Periodic map	$(\frac{1}{2})^{g+4}$	
flavor	$su(2)^{g+3}$	
# $A_1$ singularities	$4g+2$	
Full SW curve	$y^2 = (u_{g-1}x^{g-1} + \dots + t)[x^{g+3} + \tau_1 x^{g+2} + \dots + \tau_{g+2}]$	

### 3 $\mathbb{Z}_2$ quotient

The Coulomb branch solution is given by the fibration of abelian varieties over the generalized Coulomb branch [5]. In above studies, the abelian varieties are given by the Jacobian of a hyperelliptic curves. Some abelian varieties are given by the finite quotient of the Jacobian variety, and are called Prym variety. Given above hyperelliptic families, we would like to study the finite quotient to get the SW geometries for other SCFTs.

Let's consider the curves which admit a  $\mathbb{Z}_2$  action  $x \rightarrow -x$ , then the anti-invariant holomorphic differentials are

$$\frac{x^{2i} dx}{y}, \quad i = 0, 1, \dots,$$

So the number would be  $\frac{g}{2}$  for even  $g$ , and  $\frac{g+1}{2}$  for  $g$  odd. Now the special Kähler condition is

$$\frac{\partial \lambda_{sw}}{\partial t} = c \frac{dx}{y};$$

So  $\lambda_{sw}$  is required to be odd under  $x \rightarrow -x$  if we'd like to preserve the  $t$  in the spectrum. Once we keep  $t$  in the spectrum, the operator  $u_i$  with  $i$  even would be preserved. The Dirac quantization would put further constraint on the  $t$  transformation of the curve (we would like to discuss this subtle issue elsewhere).

*Example:* Let's now consider the curve defined as  $y^2 = x^{2g+2} + t^2$ , with  $g$  odd. The full SW curve would be

$$y^2 = x^{2g+2} + (\tau x^{g+1} + u_{g-1} x^{g-1} + u_{g-3} x^{g-3} + \dots + t)^2 + (T_2 x^{2g} + T_4 x^{2g-2} + \dots + T_{2g} x^2).$$

Terms of odd powers of  $x$  are eliminated in the last bracket since the curve is required to be invariant under  $x \rightarrow -x$ . This curve should describe the conformal gauge theory  $SO(g+2)$  coupled with  $2g$  half-hypermultiplets, so the flavor symmetry is  $usp(2g)$ . Notice that the constant deformation term is not allowed (see [25]).

We search all such theories and list them in table. 10 and 11. Notice the rank one theories are the  $I_4$  series found in [18].

**Table 10:**  $\mathbb{Z}_2$  quotient:  $y^2 = x^{2g+2} + t^2$ .

Theory	$SO(g+2) - 2g(\text{half} - \text{hyper})$	
curve	$y^2 = x^{2g+2} + t^2, g \text{ odd}$	$y^2 = x^{2g+2} + t^2, g \text{ even}$
CB spectrum	$2, 4, \dots, g+1$	$3, 5, \dots, g+1$
central charges	$a = \frac{1}{48}(g+2)(7g+5)$ $c = \frac{1}{12}(g+2)(2g+1)$	$a = \frac{1}{48}(g(7g+19)+2)$ $c = \frac{1}{12}(g(2g+5)+1)$
Singularities	$I_4 I_1^{2g}$	$I_4 I_1^{2g}$
free hyper on CB	$g$	$g$
flavor	$usp(2g)$	$usp(2g) \times u(1)$
Full SW curve $g \text{ odd}$	$y^2 = x^{2g+2} + (\tau x^{g+1} + u_{g-1} x^{g-1} + u_{g-3} x^{g-3} + \dots + t)^2 + (T_2 x^{2g} + T_4 x^{2g-2} + \dots + T_{2g} x^2)$	
Full SW curve $g \text{ even}$	$y^2 = x^{2g+2} + (m x^g + u_{g-2} x^{g-2} + u_{g-4} x^{g-4} + \dots + t)^2 + (T_2 x^{2g} + T_4 x^{2g-2} + \dots + T_{2g} x^2)$	

**Table 11:**  $\mathbb{Z}_2$  quotient:  $y^2 = t(x^{g+2} + t)$ .

Theory	
curve	$y^2 = t(x^{g+2} + t), g \text{ even}$
CB spectrum	$4, 6, \dots, g+2$
central charges	$a = \frac{1}{48}g(7g+36)$ $c = \frac{1}{24}g(4g+21)$
Singularities	$I_4 I_1^{2g+1}$
free hyper	$\frac{7g}{2} - 4$
flavor	$usp(2g+2) \times su(2)$
Full SW curve $g \text{ even}$	$y^2 = (T_2' x^g + u_{g-2} x^{g-2} + \dots + t)[x^{g+2} + (T_2' x^g + u_{g-2} x^{g-2} + \dots + t)] + [T_2 x^{2g+2} + T_4 x^{2g} + \dots + T_{2g+2} x^2]$

#### 4 SW differential and other possibilities

In our previous studies, we have assumed that the derivative of  $\lambda_{sw}$  with respect to  $t$  is given as  $\frac{\partial \lambda_{sw}}{\partial t} = c \frac{dx}{y}$ . We would like to solve  $\lambda_{sw}$  explicitly. Let's assume the SW curve and SW differential take the following form

$$y^2 = f(x, P(x, u_i)), \quad \lambda_{sw} = R(x, y, \sum_i P(x, u_i));$$

and the polynomial which contains the Coulomb branch operators could take the following general form

$$P(x, u_i) = \sum_i u_i x^i.$$

Now let's consider the constraints on SW differential  $\lambda_{sw} = R(P, x, y) \frac{dx}{y}$ , and the derivative reads

$$\frac{\partial \lambda_{sw}}{\partial u_i} = \left[ \frac{\partial R}{\partial u_i} + \frac{R}{y^2} \frac{\partial f}{\partial P} x^i \right] \frac{dx}{y}. \quad (4.1)$$

We'd like to write it in the form of  $\frac{\partial \lambda_{sw}}{\partial u_i} = a \frac{x^i dx}{y} + bd \left( \frac{x^{i+1}}{y} \right)$ . Using

$$d \left( \frac{x^{i+1}}{y} \right) = \frac{(i+1)x^i dx}{y} - \frac{1}{y^2} x^{i+1} \left( \frac{\partial f}{\partial P} P' + \frac{\partial f}{\partial x} \right) \frac{dx}{y}. \quad (4.2)$$

So if we'd like to derivative is given by a holomorphic differential, one has

$$\boxed{-x \left( \frac{\partial f}{\partial P} P' + \frac{\partial f}{\partial x} \right) = R \frac{\partial f}{\partial P} + cf.} \quad (4.3)$$

Here  $c$  is a constant. The final result would be

$$\frac{\partial \lambda_{sw}}{\partial u_i} = \frac{\partial R}{\partial u_i} dx + [c - (i+1)] \frac{x^i dx}{y} + d \left( \frac{x^{i+1}}{y} \right).$$

and the constant in equation 4.3 is  $c = -k$ .

*Example 1:* If  $f$  is linear in  $P$  (Type I and type II), the solution of SW differential is simple:  $R = y^2$  an  $\lambda_{sw} = y dx$ . A total derivative term is not needed. Similarly for type III and type IV, one may need to take  $R = \frac{y^2}{x}$ , and  $\lambda_{sw} = y \frac{dx}{x}$ .

*Example 2:* If  $f$  takes the form  $f = P^2 + x^k$ , the solution would be

$$R = \frac{k}{2} P - x P'.$$

There exists an interesting possibility that involves the  $D_n$  type conformal gauge theory. In this case, the polynomial encoding the Coulomb branch operator is written as the form

$$P(x, u_i) = \sum u_i x^i + t^2.$$

and the derivative of  $\lambda_{sw}$  with respect to  $t$  is now proportional the "square root" of the differential  $\frac{dx}{y}$ .

*Example:* The curve is  $y^2 = f(x, t^2) = x^{4g} + (t^2)^2$ , We impose a  $\mathbb{Z}_2$  invariance  $x \rightarrow -x$ , and so the full SW curve is written as

$$y^2 = x^{4g} + (\tau x^{2g} + u_2 x^{2g-2} + u_{2g-2} x^2 + t^2)^2 + T_2 x^{4g-2} + \dots + T_{4g-4} x^4.$$

This gives the SW curve of  $SO(2g)$  coupled with  $2g - 2$  fundamental hypermultiplets. Notice that the mass deformations from  $x^2$  term and constant are not allowed [26]. We might generalized above construction to the following curve ( $g$  is odd)

$$y^2 = x^{2g} + (m x^{g-1} + \dots + u_{g-2} x^2 + t^2)^2 + T_2 x^{2g-2} + \dots + T_{2g-4} x^4.$$

The Coulomb branch spectrum is  $(3, 5, \dots, g - 2, \frac{g}{2})$ , and the flavor symmetry is  $usp(2g - 4) \times U(1)$ . When  $g = 7$ , a theory with same Coulomb branch spectrum and flavor symmetry was indeed found in [27].

## 5 Conclusion

We classify 4d  $\mathcal{N} = 2$  SCFTs whose SW geometry can be written as hyperelliptic family. The classification is made possible by assuming 2.5 and 2.6, and reduce the problem to one parameter family. It would be interesting to try to find other solutions to the special Kähler conditions.

Once the full SW family is given, one can further study the low energy physics of any vacuum [14] by using the period integral. Geometrically, one can also compute the flavor symmetry by computing the Mordell-Weil lattice [28]. One might also compute the torsion part of Mordell-Weil lattice which could be identified as higher form symmetry.

We focus on the SCFT in this paper, and the method can be applied to non conformal theory. The SW geometry can still be written as the form 2.5 and 2.6, and the only difference is that there is no  $\mathbb{C}^*$  action acting on the one parameter family.

Another interesting generalization is to consider other algebraic curves such as order three algebraic curves  $x^3 = f(y, u_i)$ . We hope to report the progress in this direction in the near future.

## References

- [1] N. Seiberg and E. Witten, *Electric - magnetic duality, monopole condensation, and confinement in  $N=2$  supersymmetric Yang-Mills theory*, *Nucl. Phys. B* **426** (1994) 19–52, [[hep-th/9407087](#)]. [Erratum: *Nucl.Phys.B* 430, 485–486 (1994)].
- [2] N. Seiberg and E. Witten, *Monopoles, duality and chiral symmetry breaking in  $N=2$  supersymmetric QCD*, *Nucl. Phys. B* **431** (1994) 484–550, [[hep-th/9408099](#)].
- [3] E. Witten, *Solutions of four-dimensional field theories via M theory*, *Nucl. Phys. B* **500** (1997) 3–42, [[hep-th/9703166](#)].
- [4] D. Gaiotto,  *$N=2$  dualities*, *JHEP* **08** (2012) 034, [[arXiv:0904.2715](#)].
- [5] R. Donagi and E. Witten, *Supersymmetric yang-mills theory and integrable systems*, *Nuclear Physics B* **460** (feb, 1996) 299–334.
- [6] E. J. Martinec and N. P. Warner, *Integrable systems and supersymmetric gauge theory*, *Nucl. Phys. B* **459** (1996) 97–112, [[hep-th/9509161](#)].
- [7] N. J. Hitchin, *Stable bundles and integrable systems*, *Duke Math. J.* **54** (1987) 91–114.
- [8] D. Nanopoulos and D. Xie, *Hitchin Equation, Singularity, and  $N=2$  Superconformal Field Theories*, *JHEP* **03** (2010) 043, [[arXiv:0911.1990](#)].
- [9] D. Xie, *General Argyres-Douglas Theory*, *JHEP* **01** (2013) 100, [[arXiv:1204.2270](#)].
- [10] Y. Wang and D. Xie, *Classification of Argyres-Douglas theories from M5 branes*, *Phys. Rev. D* **94** (2016), no. 6 065012, [[arXiv:1509.00847](#)].
- [11] Y. Wang and D. Xie, *Codimension-two defects and Argyres-Douglas theories from outer-automorphism twist in 6d (2, 0) theories*, *Phys. Rev. D* **100** (2019), no. 2 025001, [[arXiv:1805.08839](#)].
- [12] D. Xie and S.-T. Yau, *4d  $n=2$  scft and singularity theory part i: Classification*, 2015.

- [13] Y. Wang, D. Xie, S. S. T. Yau, and S.-T. Yau, *4d  $\mathcal{N} = 2$  SCFT from complete intersection singularity*, *Adv. Theor. Math. Phys.* **21** (2017) 801–855, [[arXiv:1606.06306](#)].
- [14] D. Xie and D. Zhang, *Mixed Hodge structure and  $\mathcal{N} = 2$  Coulomb branch solution*, [arXiv:2107.11180](#).
- [15] P. C. Argyres, M. R. Plesser, N. Seiberg, and E. Witten, *New  $N=2$  superconformal field theories in four-dimensions*, *Nucl. Phys. B* **461** (1996) 71–84, [[hep-th/9511154](#)].
- [16] J. A. Minahan and D. Nemeschansky, *Superconformal fixed points with  $E(n)$  global symmetry*, *Nucl. Phys. B* **489** (1997) 24–46, [[hep-th/9610076](#)].
- [17] P. C. Argyres and M. Martone, *The rank 2 classification problem I: scale invariant geometries*, [arXiv:2209.09248](#).
- [18] P. C. Argyres, M. Lotito, Y. Lü, and M. Martone, *Geometric constraints on the space of  $\mathcal{N} = 2$  SCFTs. Part II: construction of special Kähler geometries and RG flows*, *JHEP* **02** (2018) 002, [[arXiv:1601.00011](#)].
- [19] E. Brieskorn and H. Knörrer, *Plane Algebraic Curves: Translated by John Stillwell*. Springer Science & Business Media, 2012.
- [20] V. I. Arnold, V. V. Goryunov, O. Lyashko, and V. A. Vasilev, *Singularity theory I*, vol. 1. Springer Science & Business Media, 1998.
- [21] D. Xie, *Pseudo-periodic map and classification of theories with eight supercharges*, 2023.
- [22] M. ISHIZAKA, *Presentation of hyperelliptic periodic monodromies and splitting families*, *Revista Matemática Complutense*, 2007 **20** (09, 2007).
- [23] D. Xie, *On rank two theories with eight supercharges part II: Lefschetz pencils*, *To appear*, .
- [24] T. Eguchi, K. Hori, K. Ito, and S.-K. Yang, *Study of  $N=2$  superconformal field theories in four-dimensions*, *Nucl. Phys. B* **471** (1996) 430–444, [[hep-th/9603002](#)].
- [25] U. H. Danielsson and B. Sundborg, *The Moduli space and monodromies of  $N=2$  supersymmetric  $SO(2r+1)$  Yang-Mills theory*, *Phys. Lett. B* **358** (1995) 273–280, [[hep-th/9504102](#)].
- [26] A. Brandhuber and K. Landsteiner, *On the monodromies of  $N=2$  supersymmetric Yang-Mills theory with gauge group  $SO(2n)$* , *Phys. Lett. B* **358** (1995) 73–80, [[hep-th/9507008](#)].
- [27] B. Li, D. Xie, and W. Yan, *On low rank 4d  $\mathcal{N} = 2$  SCFTs*, *JHEP* **05** (2023) 132, [[arXiv:2212.03089](#)].
- [28] T. Shioda, *Mordell-weil lattices for higher genus fibration over a curve*, *London Mathematical Society Lecture Note Series* (1999) 359–374.