

Biorthogonal Majorana zero modes, extended waves in continuum of bound states and non-Hermitian Toda soliton-fermion duality

Harold Blas

Instituto de Física

Universidade Federal de Mato Grosso

Av. Fernando Correa, N^o 2367

Bairro Boa Esperança, Cep 78060-900, Cuiabá - MT - Brazil

harold.achic@ufmt.br

We study the non-Hermitian (NH) Toda model coupled to fermions through soliton theory techniques and the realizations of the pseudo-chiral and pseudo-Hermitian symmetries. The interplay of non-Hermiticity, integrability, nonlinearity, and topology significantly influence the formation and behavior of a continuum of bound state modes (CBM) and extended waves in the localized continuum (ELC). The non-Hermitian soliton-fermion duality, the complex scalar field topological charges and winding numbers in the spectral topology are uncovered. The Hermitian bound states/solitons lie on the unit circle $|z| = 1$ defined by the uniformization parameter $z \in \mathbb{C} \setminus \{0\}$ related to the complex energy eigenvalue, whereas the non-Hermitian bound states/solitons lie on the complex plane such that $|z| \neq 1$. The biorthogonal Majorana zero modes, dual to the NH Toda solitons with topological charges ± 1 , appear at the complex-energy point gap and are pinned at zero energy. The regions of $\mathbb{C} \setminus \{0\}$ with real eigenvalues are uncovered, and these come in real pairs $\pm \lambda_1$ ($\lambda_1 \in \mathbb{R}$) preserving the pseudo-chiral symmetry. Our findings improve the understanding of exotic quantum states, but also paves the way for future research in harnessing non-Hermitian phenomena for topological quantum computation, as well as the exploration of integrability and NH solitons in the theory of topological phases of matter.

1 Introduction

A common assumption underlying the physics of isolated systems is the Hermiticity property of the Hamiltonian. However, non-Hermitian (NH) Hamiltonians have attracted considerable interest in recent years since they are quite ubiquitous in nature and remarkable theoretical and experimental results have been uncovered (see e.g. [1, 2, 3]). They have been studied in three different approaches, a generic NH Hamiltonians with complex eigenvalues [4, 5], (anti-)pseudo-Hermitian Hamiltonians [6, 7, 8], and \mathcal{PT} symmetric Hamiltonians with real eigenvalues [9, 10, 11]. For related quasi/pseudo Hermitian systems see the classical paper [12]. Theoretical aspects and some experimental validations of nonlinear wave dynamics in \mathcal{PT} symmetric systems have been reviewed in [13].

In physical systems with Hermitian and NH Hamiltonians a considerable amount of work on topological phases has emerged as a rapidly growing research area relevant to a variety of fields. An important concept in topological physics is the bulk-boundary correspondence in topological phases, such that the topological invariant related to the bulk states counts the number of gapless boundary states. So, topological phenomena can be related to boundary phenomena. For a review on non-Hermitian lattice systems in condensed matter physics, focusing on the boundary phenomena, see e.g. [14]. Originally found in condensed matter physics, robust edge states are currently found in practically all wave dynamics originating from a variety of physical models. For example, remarkable similarities have recently been found between the shallow water equations in their linearized form and topological phases of matter [15]. A duality relationship between the bound state waves and the topological solitons has been uncovered for non-linear shallow water waves [16].

The study of topological edge modes has initially been related to Hermitian linear systems concepts, such as orthogonal eigenvectors and real energy spectra. Actually, many physical systems exhibiting these phenomena naturally present a non-linear behavior. This motivated recently a growing interest in the confluence of nonlinearity and topology with applications to a wide range of phenomena (see e.g. [17] and references therein). From the quantum field theory perspective, it has been shown the presence of edge and bulk solitons related to the problem of fractionally charged solitons and electron fractionalization in $(1 + 1)$ -dimensional systems [18, 19, 20]. In various ways the concepts involved in the $(1 + 1)$ -dimensional systems can be extended to higher dimensions.

In this context, the exploration of the interplay of NH solitons and fermionic systems emerges as an intriguing subject in theoretical and mathematical physics. In this paper, we delve into the interplay between non-Hermiticity, nonlinearity, integrability and topology in the framework of the non-Hermitian Toda soliton coupled to Dirac fermions; so, extending the Hermitian soliton-fermion duality reported in [21, 22, 23]. The exploration of non-Hermitian systems has emerged as a fascinating and rapidly growing field, yielding intriguing insights into the fundamental nature of quantum mechanics. Non-Hermitian matrices exhibit unconventional properties compared with Hermitian ones: eigenstates are, in general, nonorthogonal, instead it has been introduced the concept of biorthogonality and a complex-energy spectrum can possess exceptional

points (EPs) giving rise to spectral topology. Non-Hermiticity drastically changes the properties of topological boundary states. Non-Hermiticity amplifies the edge states, which enables novel topologically protected states. Recently a modified biorthogonal bulk-boundary correspondence has been established that works in the presence of non-Hermiticity [24].

The non-Hermitian Toda model coupled to matter exhibits an extraordinary confluence of non-Hermiticity, nonlinearity, integrability and topological solitons. Central to our investigations are the concepts of pseudo-chiral and pseudo-Hermitian symmetries, which shed lights to disentangle the intricate interplay between the mentioned properties. Remarkably, a relationship between the complex energy, number wave and mass parameters in the NH soliton sector of the model can be defined on the whole complex plane through a uniformization parameter z , which allows us to discover that while Hermitian bound state/soliton states lie on the unit circle $|z| = 1$, the non-Hermitian generalizations of that correspondence can be established on the continuous region $|z| \neq 1$ of the complex plane. In this scenario we show the emergence of biorthogonal Majorana zero modes as dual to the non-Hermitian topological solitons with topological charges ± 1 , which manifest at the complex-energy point gap and pinned at zero energy. These zero modes, we find, hold the key to unlocking the topologically protected bulk-boundary correspondence from the integrable field theory perspective.

An essential distinction between Hermitian and non-Hermitian systems is the degrees of freedom that we have access to; nonunitary operations forbidden in Hermitian systems can be performed in non-Hermitian systems. In other words, a change in the spectrum from real to complex increases the number of parameters that describe the system. Since topology crucially depends on the underlying manifold, non-Hermiticity is expected to alter the topological classification of the physical systems such as insulators and superconductors. The Hermitian Toda model coupled to matter has been found in a variety of models, such as the continuum limit of s -wave superconductor [25], effective theory describing Majorana bound states [26], a model for high T_c superconductivity [27], low-energy effective Lagrangian in QCD₂ [28], etc; so, our study not only becomes of academic interest, but physically motivated. As pointed out above, pseudo-Hermiticity is a constraint unique to non-Hermitian systems and may provide novel topological features. In view of the rapid theoretical and experimental advances in non-Hermitian physics, there has been a great interest and an urgent need for comprehensive topological classifications that provide reference points for experiments and predict novel non-Hermitian topological phases.

Our exploration has led us to unveil remarkable phenomena: the existence of continuum spectra of bound states and extended waves embedded within this localized continuum (ELC). These phenomena are profoundly shaped by the interplay of four key characteristics: non-Hermiticity, integrability, nonlinearity, and topology. The continuum of biorthogonal bound states emerge coupled to a continuous family of topological complex solitons, such that their topological charges depend on the energy eigenvalues. This type of continuum of bound states in a non-Hermitian Hamiltonian have recently been uncovered also in [29]. In the Hermitian Toda system coupled to fermion, which has recently been presented in [21], the in-gap and the

bound states in the continuum (BIC) have quantized energies, whereas the free states form a continuum. On the other hand, the NH extended wave states embedded into the NH localized continuum can be regarded as some kind of inverted BIC states due to the non-Hermiticity of the model. Recently, these novel types of states in a NH system have been dubbed as extended states in a localized continuum (ELC) [30].

So, the NH models exhibit behaviors that challenge conventional understanding. Within the NH Toda theory coupled to matter integrable field theory we clarify the concept of the non-Hermitian soliton-fermion duality, revealing a profound connection between these entities which survive non-Hermiticity effects, and shed lights on the appearance of a pair of topological charges, such that the second charge arises purely as the result of the interplay between the non-Hermiticity, topology, and nonlinear effects. We believe that beyond the pure theoretical inquiry, our discoveries might have important implications for the realm of practical quantum technologies. The harnessing of non-Hermitian phenomena in the context of topological quantum computation promise new frontiers in the development of quantum computing paradigms. Furthermore, our work establishes a bridge between quantum field theory, condensed matter physics and integrable systems, inviting further exploration at the crossroads of these three fields.

The paper is organized as follows. In the next section we present the integrable NH Toda model coupled to fermions and the relevant eigenvalue equations for the right and left eigenvectors. In the section 3 the pseudo-chirality symmetry is discussed, as well as the concept of biorthogonality of the left and right eigenvectors. The section 4 discusses the anti-pseudo-Hermiticity symmetry. Some theorems and properties of pseudo-Hermitian Hamiltonians are discussed in the presence of \mathcal{P} , \mathcal{T} and \mathcal{PT} transformations. In the section 5 we uncover the topological complex solitons and the biorthogonal spinor bound states through the tau function formalism. The complex energy manifold is defined, identifying the regions of real and complex eigenvalues and the points of eigenvector degeneracies. Moreover, the spectral topology and the associated winding number are discussed. In section 6 a classification of these solutions are discussed. The continuum of bound states/solitons and the extended waves embedded into this localized continuum are presented. The biorthogonality relationships between the left and right eigenvectors are computed. In section 7 the biorthogonal Majorana zero mode-topological soliton duality is studied by providing a mapping between the spinor and the scalar fields. Section 8 presents the conclusions and discussions. In the appendix A we present the static equations of motion in components.

2 The model

There is a variety of ways to introduce non-Hermiticity in the integrable models. Earlier versions of non-Hermitian affine Toda theories consider a purely imaginary coupling constant which provide soliton solutions with real classical masses [31]. Some recent works introduce the non-Hermiticity by making the field of the model a complex field, as in the complex KdV, mKdV and sine-Gordon equations which posses PT-symmetric soliton solutions with real energies [32]. Likewise, non-Hermiticity has been induced by many

ways in field theory models, partially motivated by the search for another viewpoint to understand the topological properties of non-Hermitian systems. In particular, non-Hermiticity in the Dirac-type equations has been introduced through Lorentz symmetry violation and the generalization of some parameters from real to complex, e.g. the complex mass parameter [33, 34]. A generic one-dimensional Dirac model with both real and imaginary masses has been considered in optics [35] and the authors argued that those results can be tested experimentally.

Here we consider two fermion fields chirally coupled to a complex scalar field defined by the following non-Hermitian Lagrangian[36, 22, 23]¹

$$\mathcal{L} = \frac{1}{2}\partial_\mu\Phi\partial^\mu\Phi + i\bar{\psi}\gamma^\mu\partial_\mu\psi - M(\bar{\psi}\psi\cos(\beta\Phi) + i\bar{\psi}\gamma_5\psi\sin(\beta\Phi)), \quad (2.1)$$

with Φ being a complex field and $\tilde{\psi}$ and ψ two Dirac spinors. Below we will consider a complex mass parameter $M \in \mathbb{C}$ and a real β as the coupling strength between the fermions and the complex scalar. We define the model (2.1) as the *non-Hermitian Toda model coupled to Dirac fields* (NHTD). The general model (2.1), with two Dirac spinors ψ and $\tilde{\psi}$ and a complex field $\Phi \in \mathbb{C}$, belongs to a wide class of integrable soliton theories, the so-called $\hat{sl}(2)$ affine Toda systems coupled to matter fields [36]. The integrability properties, the construction of the general solution including the solitonic ones, the soliton-fermion duality properties, as well as its symplectic structures were discussed in [22, 23, 37]. We will consider the non-Hermitian model (2.1) and show that the dynamics of the system exhibits a continuum of kink type states and its associated biorthogonal spinor bound states. In addition, it will be shown the appearance of extended wave states embedded into the continuum of bound states. In this context we will realize the various symmetry properties of the model such as the pseudo-chirality and its associated biorthogonal bound states, such that the generalized non-Hermitian particle-hole (NHPH) symmetry holds.

The non-Hermiticity in the model (2.1) can be traced to the presence of at least one of the next conditions. First, $\Phi \neq \Phi^*$, where \star stands for complex conjugation. Second, $\tilde{\psi} \neq \psi^*$. Third, in clear contrast to the Hermitian case one has the complex mass parameter M . The presence of any of these conditions would make the Lagrangian non-Hermitian. As we will see below, the interplay between these conditions gives rise to interesting new results related to non-Hermitian systems. Among the new results of the NHTD model (2.1), beyond the known results in the literature for the Hermitian case, one has the emergence of new type of topological solitons associated to the complex sector of Φ and their related spinor bound states, as well as the appearance of extended wave states in the continuum of bound states supported by the complex scalar and spinor fields. So, one can argue that the novel solitons and extended waves associated to the imaginary sector $\Im(\Phi)$ will be generated due to the interplay between the non-Hermiticity, topological properties and nonlinearity of the model, as compared to the Hermitian counterpart of the model [21] which possesses kink solutions of the sine-Gordon type and BIC states associated to $\Re(\Phi)$.

¹Notation: $\partial^2 = \partial_t^2 - \partial_x^2$. $\gamma_0 = \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}$, $\gamma_1 = \begin{pmatrix} 0 & -i \\ -i & 0 \end{pmatrix}$, $\gamma_5 = \gamma_0\gamma_1 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, and $\bar{\psi} = \tilde{\psi}^T\gamma_0$; $\bar{\chi} = \tilde{\chi}^T\gamma_0$.

Regarding the physical motivation to introduce the complex parameter M one can justify in hindsight by some physical and mathematical concepts. As we will show below, the complex values $\pm M$ become the branch point singularities of the multivalued complex energy function with the dispersion relation $\lambda(k) = \pm\sqrt{M^2 - k^2}$, with k a complex wave number, in the enlarged complex parameter space manifold. The zero modes are defined by $\lambda = 0$, and so, in this case one has $k = \pm M$. Taking the complex number notation $z \equiv z_1 + iz_2$, one notices that for wave solutions with the x -dependence of type e^{kx} the real part of the wave number $k_1 = \pm M_1$ could be associated to the localized bound states and the imaginary part $k_2 = \pm M_2$ to the extended wave states. On the other hand, for large M and small wave number k one has $\lambda_2 = \pm M_2$, e.g. this will happen for bound states with sufficiently large width or extended waves with large wave-length. We will see below that some solutions for the spinors exhibit the dissipation/amplification factors of type $e^{\lambda_2 t}$. This behavior calls for an alternative interpretation of the physical origin of the non-Hermiticity in the realm of a topological non-Hermitian system without resorting to the gain/loss concept, for example following the recent alternative approach introduced by means of the skin effect of reflected waves [38]. As a special case in the soliton sector of the non-Hermitian model (2.1) we will uncover a parameter subspace related to the scalar and spinor solutions with real eigenvalues $\lambda \in \mathbb{R}$ which come in pairs $\{\pm\lambda_1, \lambda_1 \in \mathbb{R}\}$ for an arbitrary parameter $(\frac{M_2}{M_1})$, such that the pseudo-chiral symmetry is maintained.

Related to the points discussed above one can define a topological charge associated to the scalar field as $\mathcal{Q}_{top} \equiv \frac{\beta}{2\pi}(\Phi(x = +\infty) - \Phi(x = -\infty))$. For certain field configurations the both real and imaginary components of Φ will develop non-vanishing boundary values, and so, this could imply a non-trivial complex charge \mathcal{Q} . This charge will be regarded as a complex topological charge. In fact, by introducing the so-called uniformization complex variable z through $\lambda = -\frac{M}{2}(z + z^{-1})$, we will show below a profound connection between this charge and the topological properties of the complex manifold in parameter space expressed by the remarkable relationship $\mathcal{Q}_{top} = \pm\frac{2}{\pi i} \log z$. So, one can argue that the appearance of this complex topological charge is due to the interplay between the non-Hermitian, topological and nonlinear properties of the model (2.1).

The Hermitian version of the model (2.1) has been studied in the literature. In this case one has a real scalar field, the fermions satisfy the reality condition $\tilde{\psi} = \psi^*$ and M is a real mass parameter. In that case the model has been defined by a real Lagrangian and associated Hermitian hamiltonian [22, 37, 23, 28, 21]. The Hermitian model has been shown to describe the low-energy effective Lagrangian of QCD₂ with one flavor and N colors [28] and the BCS coupling in spinless fermions in a two dimensional model of high T superconductivity in which the solitons play the role of the Cooper pairs [27]. Moreover, the model has earlier been studied as a model for fermion confinement in a chiral invariant theory [39] and the mechanism of fermion mass generation without spontaneously chiral symmetry breaking in two-dimensions [40]. Recently, through analytical and numerical methods the Hermitian version of the model (2.1) plus a scalar potential has been shown to exhibit Majorana zero mode-soliton duality and in-gap and BIC bound states, such that the Noether, topological and a novel nonlocal charge densities satisfy a formula of the Atiyah-Patodi-Singer-type

[21].

The equations of motion become

$$\partial^2 \Phi - M\beta\tilde{\psi}\psi \sin(\beta\Phi) + iM\beta\tilde{\psi}\gamma_5\psi \cos(\beta\Phi) = 0 \quad (2.2)$$

$$i\gamma_\mu \partial^\mu \psi - M(\psi \cos(\beta\Phi) + i\gamma_5\psi \sin(\beta\Phi)) = 0 \quad (2.3)$$

$$i\gamma_\mu \partial^\mu \tilde{\psi} - M(\tilde{\psi} \cos(\beta\Phi) + i\gamma_5\tilde{\psi} \sin(\beta\Phi)) = 0 \quad (2.4)$$

Let us define

$$\psi \equiv e^{-i\lambda t}\xi, \quad \tilde{\psi} \equiv e^{i\lambda t}\eta \quad (2.5)$$

$$\xi = \begin{pmatrix} \xi_A \\ \xi_B \end{pmatrix}, \quad \eta = \begin{pmatrix} \eta_A \\ \eta_B \end{pmatrix}, \quad (2.6)$$

with λ being a complex parameter. The equations (2.3)-(2.4) can be written in matrix form as

$$H\xi = \lambda\xi, \quad (2.7)$$

$$\tilde{H}\eta = \lambda\eta \quad (2.8)$$

where

$$H = \begin{pmatrix} i\frac{d}{dx} & iMe^{-i\beta\Phi} \\ -iMe^{i\beta\Phi} & -i\frac{d}{dx} \end{pmatrix}, \quad \tilde{H} = \begin{pmatrix} -i\frac{d}{dx} & -iMe^{i\beta\Phi} \\ iMe^{-i\beta\Phi} & i\frac{d}{dx} \end{pmatrix}. \quad (2.9)$$

Notice that the equations above define a pair of eigenvalue equations for the non-Hermitian operators H and \tilde{H} with right eigenvectors ξ and η , respectively. Recently, the Hermitian version of the model (2.7) has been considered as the continuum limit of a one-dimensional system of spin-orbit coupled Dirac Hamiltonian and a s -wave superconducting pairing, such that the s -wave pairing is represented by $-iMe^{i\beta\Phi}$ ($\Phi \in \mathbb{R}$) [25]. Moreover, the system (2.7) has been considered in [26] as an effective theory describing Majorana bound states such that the function $iMe^{-i\beta\Phi}$ is related to the superconducting order parameter.

One can show that the NH operators H and \tilde{H} are related by

$$\Gamma H \Gamma = \tilde{H}, \quad \Gamma = i\gamma_1. \quad (2.10)$$

Due to the last relationship (2.10), given an eigenvector η of \tilde{H} with eigenvalue λ one can get an eigenvector $\Gamma\eta$ of H with the same eigenvalue λ . In fact, from (2.8) and taking into account (2.10) one can get

$$\Gamma H \Gamma \eta = \lambda \eta \rightarrow H(\Gamma\eta) = \lambda(\Gamma\eta) \quad (2.11)$$

So, comparing (2.11) with (2.7) one can write

$$\xi = \Gamma\eta \Rightarrow \xi_A = \eta_B, \quad \xi_B = \eta_A. \quad (2.12)$$

From (2.2) and taking into account the Ansatz (2.5) one can get a static version of the complex scalar field equation of motion

$$-\partial_x^2 \Phi + M\beta[e^{i\beta\Phi}\eta_B\xi_A + e^{-i\beta\Phi}\eta_A\xi_B] = 0. \quad (2.13)$$

We will examine the static solutions satisfying the relationship

$$\partial_x \Phi = -\frac{\beta}{2}[\eta_A \xi_A + \eta_B \xi_B], \quad (2.14)$$

$$= -\frac{\beta}{2}[\eta_3 \xi_3 - \eta_4 \xi_4 + \eta_1 \xi_1 - \eta_2 \xi_2 + i(\eta_3 \xi_4 + \eta_4 \xi_3 + \eta_1 \xi_2 + \eta_2 \xi_1)], \quad (2.15)$$

where in the last equation we have written the real and imaginary parts of the bilinear current terms using the component of ξ and η defined in (A.1) and (A.2).

Remarkably, the first order equation (2.14) together with the first order equations of motion for the Dirac spinors (2.7)-(2.8) imply the second order equation for Φ in (2.13). Therefore, a given set of solutions of the system of static first order differential equations (2.7)-(2.8) and (2.14) will also be solutions of the second order equation for Φ . The equation (2.14) suggests a certain kind of equivalence relationship between the x -derivative of the scalar and the fermion bilinear in the NH model. In fact, this is a NH analog of the equivalence between the $U(1)$ Noether and topological soliton currents considered in the literature in the Hermitian case of the model [21]. In both condensed matter (e.g. Bogoliubov-de Gennes (BdG) and gap equations) and high energy physics (e.g. Gross-Neveu (GN) and Nambu-Jona-Lasinio models) there have been studied analogous relationships to (2.14) by self-consistency requirements with (2.7)-(2.8) which amounts to write e.g. $iM e^{i\beta\Phi} \sim [\xi_A^* \xi_B + \eta_A^* \eta_B]$ [41]. Actually, for a real scalar and one Dirac fermion these relationships resemble to the bosonization rules which map some fermion bilinears to a topological current and exponentials of the scalar [42].

One can notice that the imaginary component of the bilinear current in (2.15) vanishes provided that

$$\eta = \pm \xi^* \Rightarrow \left\{ \begin{array}{l} \text{Case I : } \eta_3 = \xi_3, \quad \eta_4 = -\xi_4, \quad \eta_1 = \xi_1, \quad \eta_2 = -\xi_2, \\ \text{Case II : } \eta_3 = -\xi_3, \quad \eta_4 = \xi_4, \quad \eta_1 = -\xi_1, \quad \eta_2 = \xi_2. \end{array} \right\} \quad (2.16)$$

Then, for the cases (2.16) the scalar field must be real, i.e. $\Re(\Phi) = \Phi_1$ ($\Im(\Phi) = 0$). So, this implies $\tilde{H} = H^*$ (for real M) and one recovers a Hermitian model with

$$H^\dagger = H, \quad \tilde{H}^\dagger = \tilde{H}, \quad (2.17)$$

and real eigenvalues $\lambda \in \mathbb{R}$. In the both cases (2.16) the imaginary part of the second order equation (2.13) for the scalar field identically vanishes for $\Phi_2 = 0$ and $M_2 = 0$. The bilinear terms of the spinor components are presented in (A.14).

However, for $M_2 \neq 0$, $\eta \neq \pm \xi^*$ and $\Phi_2 = 0$ the eq. (2.15) provides the general constraint $(\eta_3 \xi_4 + \eta_4 \xi_3 + \eta_1 \xi_2 + \eta_2 \xi_1) = 0$ and the second order eq. (A.14) will provide an additional constraint (A.17).

In the case of NH Hamiltonians it is convenient to define the so-called left eigenvectors, since in general they are not the same as the relevant Hermitian conjugated right eigenvectors. So, the left eigenvectors $\bar{\xi}_j^T$ and $\bar{\eta}_j^T$ are defined as

$$\bar{\xi}_j^T H = \lambda_j \bar{\xi}_j^T, \quad \bar{\eta}_j^T \tilde{H} = \lambda_j \bar{\eta}_j^T \quad (2.18)$$

Next, we will consider certain solutions satisfying

$$\partial_x \bar{\Phi} = -\frac{\beta}{2}[\bar{\eta}_A \bar{\xi}_A + \bar{\eta}_B \bar{\xi}_B], \quad (2.19)$$

where we have introduced a complex scalar field $\bar{\Phi}$, and $\bar{\eta}^T \equiv (\bar{\eta}_A, \bar{\eta}_B)$ and $\bar{\xi}^T \equiv (\bar{\xi}_A, \bar{\xi}_B)$. In general one has that the left and right eigenvectors are different, i.e.

$$\bar{\eta} \neq \eta, \quad \bar{\xi} \neq \xi, \quad (2.20)$$

and, so, the scalar field $\bar{\Phi}$ can be considered to be in general different from Φ in (2.14). However, we will see below that the left and right eigenvectors satisfy some symmetry relationships, which imply a relationship between the scalars Φ and $\bar{\Phi}$ provided that the equation (2.19) holds. So, the second order differential equation for the scalar $\bar{\Phi}$ becomes

$$\partial_x^2 \bar{\Phi} + M\beta[e^{i\beta\bar{\Phi}} \bar{\eta}_A \bar{\xi}_B + e^{-i\beta\bar{\Phi}} \bar{\eta}_B \bar{\xi}_A] = 0. \quad (2.21)$$

Notice that the equations (2.13) and (2.21) have different origins since the first one has been derived from the Lagrangian, whereas the second one is a consequence of the equivalence relationship (2.19) which has been assumed to hold between a topological current associated to $\bar{\Phi}$ and the left-eigenvector current density, respectively. However, we will see below that for λ and $-\lambda$ eigenvalues the relevant right and left eigenvectors will be related by symmetry arguments, and so, those equations will be related by the so-called pseudo-chirality symmetry. So, we are establishing some equivalences between topological currents and biorthogonal fermion currents represented by the correspondence relationships (2.14) and (2.19), respectively.

3 Pseudo-chirality and biorthogonality of the eigenvectors

For a generic complex scalar field Φ the operators H and \tilde{H} are non-Hermitian with a spectra of complex eigenvalues λ . Recently, there have been various attempts to realize particle-hole symmetry in non-Hermitian (NH) systems. However, due to the fact that these effective particles and holes have complex energies in a NH system, they do not need to fulfill the Fermi-Dirac statistics. Non-Hermitian Hamiltonian matrices exhibit more types of symmetries than the Hermitian counterparts [14]. So, in order to examine certain extensions of the Noether's theorem, it has been proposed distinct ways to define the inner product in NH systems. An inner product, without the complex conjugation, has been proposed in [5] which focuses on asymmetric non-Hermitian Hamiltonians. So, let us write the next inner product

$$(\xi_i | \xi_j) \equiv \xi_i^T \xi_j. \quad (3.1)$$

With this inner product and using an alternative form of the Ehrenfest's theorem for the time evolution of $(\psi | \mathcal{G} | \psi)$ one can show that the operator \mathcal{G} is a constant of motion provided that H satisfies [5]

$$\mathcal{G}H\mathcal{G}^{-1} = -H^T, \quad (3.2)$$

where H^T stands for matrix transpose. For the NH Hamiltonian H in (2.9) one has that

$$\mathcal{G} = -i\gamma_0. \quad (3.3)$$

In [5] the symmetry (3.2) has been defined as pseudo-chirality, and it guaranties, as we will see below, a symmetric spectrum about the origin of the complex energy plane, i.e. for each λ there exists $-\lambda$, similar to the consequence of particle-hole symmetry in the hermitian counterpart of the model (2.1) studied recently in [21].

Below we will follow the biorthogonal approach to describe the orthogonality properties of eigenvectors and eigenstates [2, 5]. So, let us define the right eigenvectors ξ_j and the relevant left eigenvectors $\bar{\xi}_j^T$ as

$$H\xi_i = \lambda_i \xi_i, \quad \bar{\xi}_i^T H = \lambda_i \bar{\xi}_i^T. \quad (3.4)$$

The eigenvectors ξ_i and $\bar{\xi}_i$ are different in general but share the same eigenvalue λ_i . Moreover, they satisfy the biorthogonality relationship [5]

$$(\bar{\xi}_j|\xi_i) \equiv \bar{\xi}_j^T \xi_i = \delta_{ji}, \quad (3.5)$$

which holds away from a zero mode ($\lambda_j = 0$). In fact, considering the both eigenvalue equations in (3.4) for ξ_i and $\bar{\xi}_j^T$, respectively, one can write the identity

$$\bar{\xi}_j^T H \xi_i = \lambda_i \bar{\xi}_j^T \xi_i, \quad \bar{\xi}_j^T H \xi_i = \lambda_j \bar{\xi}_j^T \xi_i \quad \Rightarrow \quad 0 = (\lambda_i - \lambda_j) \bar{\xi}_j^T \xi_i. \quad (3.6)$$

Then, from this equation one can see that the biorthogonality relationship (3.5) is satisfied for $\lambda_i \neq \lambda_j$. The normalization $(\bar{\xi}_i|\xi_i) = 1$ ($\lambda_i \neq 0$) will be considered in the sections below in the context of the soliton type solutions of the model.

Next, for the Hamiltonian H in (2.9) which possesses the property (3.2), one can show the following property

$$H^T[\mathcal{G}\xi_{\lambda_j}] = -\mathcal{G}H\xi_{\lambda_j} = -\lambda_j[\mathcal{G}\xi_{\lambda_j}]. \quad (3.7)$$

Since the second eq. in (3.4) can be written as $H^T\bar{\xi}_{-\lambda_j} = -\lambda_j\bar{\xi}_{-\lambda_j}$, then from (3.7) one can write

$$\bar{\xi}_{-\lambda_j} = \mathcal{G}\xi_{\lambda_j}. \quad (3.8)$$

This relation suggests that for each right eigenstate ξ_{λ_j} with eigenvalue λ_j , one can find a left eigenstate $\bar{\xi}_{-\lambda_j}$ for the relevant eigenvalue $-\lambda_j$, such that in components one has the next relationships

$$\bar{\xi}_{-\lambda_j A} = \xi_{\lambda_j B}, \quad \bar{\xi}_{-\lambda_j B} = -\xi_{\lambda_j A}. \quad (3.9)$$

In terms of the Dirac field ψ_{λ_j} defined in (2.5), and defining $\bar{\psi}_{-\lambda_j} \equiv e^{-i\lambda_j t} \bar{\xi}_{-\lambda_j}$, one can write

$$\bar{\psi}_{-\lambda_j} = \mathcal{G}\psi_{\lambda_j}. \quad (3.10)$$

So, the operator \mathcal{G} maps a right Dirac eigenstate with eigenvalue λ_j to one of the left eigenstates with corresponding eigenvalue $-\lambda_j$. This relationship exhibits the so-called NH particle-hole symmetry of the system and then one expects that the ‘particles’ and ‘holes’ will have identical spectra, since the spectra of H is symmetric about the origin of the complex eigenvalue plane. So, for certain zero modes one can formulate the NH analog of the Majorana fermions for a system of this kind since there exists a biorthogonal ‘particle-hole’ symmetry.

A Majorana fermion in a Hermitian model is defined by considering the particle and hole with the same energy as a single particle. Here we generalize the Majorana zero modes to NH model which exhibits topological solitons, and show a general but quite different properties from their Hermitian counterparts. Then, following analogous reasoning, from the relationship (3.10) and for zero-mode solutions ($\lambda_j = 0$), one can write

$$\bar{\psi}_0 = \mathcal{G}\psi_0. \quad (3.11)$$

The biorthogonal states obeying the condition (3.11) will be regarded as NH Majorana zero modes. These states are well known to lead to interesting phenomena in condensed matter and we will search for them as bound states for the biorthogonal states.

Notice that the biorthogonal relationship (3.5) is different from the NH inner product (3.1). For example, the product (3.5) vanishes at the zero mode $\lambda_j = 0$, since taking into account (3.11), one has $\bar{\xi}_{0A} = \xi_{0B}$, $\bar{\xi}_{0B} = -\xi_{0A}$, which implies

$$\begin{aligned} (\bar{\xi}_0|\xi_0) &\equiv \bar{\xi}_0^T \xi_0 \\ &= 0. \end{aligned} \quad (3.12)$$

On the other hand, the product (3.1) does not vanish at the zero mode $\lambda_j = 0$, i.e. it becomes

$$(\xi_0|\xi_0) \equiv \xi_0^T \xi_0 \quad (3.13)$$

$$= (\xi_{0A})^2 + (\xi_{0B})^2, \quad (3.14)$$

which is a complex number in general. This is in contradistinction to the Hermitian case in which an inner product has been defined to provide a real value for the Hermitian Majorana zero modes [21].

Similar constructions as above can be performed for the second Dirac field $\tilde{\psi}$. Since the pseudo-chirality property (3.2) also holds for the second operator \tilde{H} , i.e.

$$\mathcal{G}\tilde{H}\mathcal{G}^{-1} = -\tilde{H}^T, \quad (3.15)$$

then one can follow throughout all the discussions above for the right eigenvectors η and the relevant left eigenvectors $\bar{\eta}^T$

$$\tilde{H}\eta_i = \lambda_i \eta_i, \quad \bar{\eta}_i^T \tilde{H} = \lambda_i \bar{\eta}_i^T. \quad (3.16)$$

So one can show

$$\bar{\eta}_{-\lambda_j} = \mathcal{G}\eta_{\lambda_j}. \quad (3.17)$$

So, for each right eigenstate η_{λ_j} , one can find a left eigenstate $\bar{\eta}_{-\lambda_j}$, which in components become

$$\bar{\eta}_{-\lambda_j A} = \eta_{\lambda_j B}, \quad \bar{\eta}_{-\lambda_j B} = -\eta_{\lambda_j A}. \quad (3.18)$$

For the Dirac field $\tilde{\psi}$ as defined in (2.5) ($\tilde{\psi} = e^{i\lambda t}\eta$) and taking into account (3.17) one can written

$$\bar{\tilde{\psi}}_{-\lambda_j} = \mathcal{G}\tilde{\psi}_{\lambda_j}. \quad (3.19)$$

So, one can argue that this relationship also exhibits the NH particle-hole symmetry for the second Dirac field $\tilde{\psi}$ and then allows us to discuss the Majorana zero modes. Following analogous reasoning from the relationship (3.19) and for zero-mode solutions ($\lambda_j = 0$) one can write

$$\bar{\tilde{\psi}}_0 = \mathcal{G}\tilde{\psi}_0. \quad (3.20)$$

The biorthogonal states obeying the condition (3.20) will be regarded as the second NH Majorana zero mode of the system. The analog of the product (3.5) becomes

$$(\bar{\eta}_i|\eta_j) = \bar{\eta}_i^T \eta_j = \delta_{ij}. \quad (3.21)$$

This product vanishes at the zero mode point $\lambda_j = 0$, since taking into account (3.20), one has $\bar{\eta}_{0A} = \eta_{0B}$, $\bar{\eta}_{0B} = -\eta_{0A}$, which implies

$$\begin{aligned} (\bar{\eta}_0|\eta_0) &\equiv \bar{\eta}_0^T \eta_0 \\ &= 0. \end{aligned} \quad (3.22)$$

At the zero mode one has $\lambda = 0$, at this point the bi-orthogonal conditions (3.5) and (3.21) break down, and the set of states $\{\bar{\xi}_0, \xi_0\}$ and $\{\bar{\eta}_0, \eta_0\}$ coalesce, i.e. they become orthogonal, and the Hamiltonian becomes non-diagonalizable, see e.g. [43, 44]. The results (3.12) and (3.22) will be verified below for spinor bound state eigenvectors. Notice that at the point $\lambda = 0$ we will have an energy degeneracy for the both types of eigenvector spaces, left and right. In fact, the set of right and left eigenvectors $\{\xi_0, \eta_0\}$ and $\{\bar{\xi}_0, \bar{\eta}_0\}$, respectively, correspond to the vanishing eigenvalue $\lambda = 0$.

At this point, let us examine the relationships between the currents equivalence equations (2.14) and (2.19) in view of the pseudo-chirality symmetry discussed above. So, taking into account the relationships (3.9) and (3.18) one can write

$$\bar{\eta}_{-\lambda,A} \bar{\xi}_{-\lambda,A} + \bar{\eta}_{-\lambda,B} \bar{\xi}_{-\lambda,B} = \eta_{\lambda,A} \xi_{\lambda,A} + \eta_{\lambda,B} \xi_{\lambda,B}. \quad (3.23)$$

Then, due to the identity (3.23) one can argue that the both scalars Φ and $\bar{\Phi}$ satisfy the identity

$$\bar{\Phi} = \Phi, \quad (3.24)$$

provided that they satisfy the equations (2.14) and (2.19) for the right-eigenvectors with λ eigenvalue and the left-eigenvectors with $-\lambda$ eigenvalue, respectively. Likewise, the second order differential equations (2.13) and (2.21) can be mapped to each other using the pseudo-chiral symmetry relationships (3.9) and (3.18) of the right and left eigenvectors for eigenvalues λ and $-\lambda$, respectively, and the relationship (3.24) for the scalars which holds for field configurations satisfying the currents equivalence equations (2.14) and (2.19).

4 Anti-pseudo-Hermiticity, \mathcal{P} , \mathcal{T} and \mathcal{PT} transformations

Breaking the Hermiticity property leads to a larger number of symmetries since the non-Hermitian Hamiltonian (2.9) exhibits the properties $H \neq H^\dagger$ and $H^* \neq H^T$. In that scenario the time-reversal (TR) and particle-hole symmetries constitute two inequivalent representations, i.e. depending on whether the symmetry considers conjugation or transposition of H , respectively (see e.g. [45]). Notice that the symmetry property considering the transposition of H has already been discussed as the pseudo-chiral symmetry in the last section. Next, we consider the other inequivalent representation by taking the conjugation of H .

4.1 Anti-pseudo-Hermiticity

Let us consider a TR operator \mathcal{T} such that $\mathcal{T}z\mathcal{T}^{-1} = z^*$ and $\mathcal{T}^2 = \pm 1$ and so, \mathcal{T} is an anti-linear operator. Next, apply this operator on the both hand sides of the pseudo-chirality relationship (3.2) for a generic complex scalar field Φ and complex parameter M . So, one can write

$$\sigma_- H \sigma_-^{-1} = -H^\dagger, \quad (4.1)$$

$$\sigma_- \equiv \mathcal{GT}, \quad (4.2)$$

where \mathcal{G} is given in (3.3). Notice that σ_- is an antilinear operator. Following the notations and conventions in the current literature one has that the equations (4.1)-(4.2) define H to be an anti-pseudo-(anti)-Hermitian Hamiltonian² [44, 7]. A similar property can be discussed for the companion Hamiltonian \tilde{H} , since it also satisfies the pseudo-chirality relationship (3.15); so, \tilde{H} is also an anti-pseudo-(anti)-Hermitian Hamiltonian

$$\sigma_- \tilde{H} \sigma_-^{-1} = -\tilde{H}^\dagger. \quad (4.3)$$

Next, let us define the right and left eigenvector equations for the operator H^*

$$H^* \chi_j = \lambda_j \chi_j, \quad \hat{\chi}_i^T H^* = \lambda_i \hat{\chi}_i^T. \quad (4.4)$$

Taking the transpose of the left eigenvalue equation in (4.4) and subsequently considering the pseudo-anti-Hermiticity relationship (4.1) one can write

$$H^\dagger \hat{\chi}_j = \lambda_j \hat{\chi}_j, \quad (4.5)$$

²We denote σ_\pm for pseudo-, and pseudo-(anti)-Hermitian cases, respectively, such that $\sigma_\pm H \sigma_\pm^{-1} = \pm H^\dagger$. Further generalizations are possible, σ_\pm operators can be either linear or anti-linear. In our case σ_- is an anti-linear operator. So, H can be dubbed as an anti-pseudo-(anti)-Hermitian Hamiltonian

$$-\sigma_- H \sigma_-^{-1} \hat{\chi}_j = \lambda_j \hat{\chi}_j, \quad (4.6)$$

$$H^*(\mathcal{G}^{-1} \hat{\chi}_j) = -\lambda_j (\mathcal{G}^{-1} \hat{\chi}_j), \quad (4.7)$$

where (4.6) follows upon using (4.1) into (4.5). In order to get (4.7) we have used $\mathcal{T}H\mathcal{T}^{-1} = H^*$ and multiplied on the left by \mathcal{G}^{-1} on the b.h.s. of (4.6). So, comparing the last relationship (4.7) and the right eigenvalue equation in (4.4) one gets

$$\chi_{\bar{j}} = \mathcal{G}^{-1} \hat{\chi}_j, \quad (4.8)$$

where $\chi_{\bar{j}}$ denotes a right eigenvector with eigenvalue $(-\lambda_j)$.

Notice that one can also relate the above left eigenvector $\hat{\chi}_j^T$ to the left eigenvector of type $\bar{\xi}^T$ in (3.4) for the same eigenvalue λ_j . In fact, by taking the complex conjugate of (3.4) one has

$$\bar{\xi}_j^\dagger H^* = \lambda_j^* \bar{\xi}_j^\dagger, \quad (4.9)$$

which upon comparing to the relevant equation in (4.4) provides the left eigenvector of H^*

$$\hat{\chi}_j^T = \bar{\xi}_j^\dagger \quad (4.10)$$

with eigenvalue λ_j^* . Moreover, using the symmetry (4.1) and the right eigenvector equation (3.4) one can write the following left eigenvector equations

$$(\xi^\dagger \sigma_-) H = -\lambda^* (\xi^\dagger \sigma_-), \quad (4.11)$$

$$(\xi^\dagger \mathcal{G}) H^* = -\lambda^* (\xi^\dagger \mathcal{G}), \quad (4.12)$$

where in order to get the last relationship we have used the form of σ_- in (4.2). So, by comparing (4.12) with the left-eigenvector equation in (4.4) one can conclude that for each left eigenvector $\hat{\chi}_j^T$ with eigenvalue λ_j one must have another eigenvector $(\xi^\dagger \mathcal{G})$ with eigenvalue $-\lambda_j^*$, i.e.

$$\hat{\chi}_{-\lambda_j^*}^T = (\xi_{\lambda_j}^\dagger \mathcal{G}), \quad (4.13)$$

with ξ_j being the eigenvector of H with eigenvalue λ_j defined in (3.4).

Analogous construction can be performed for the companion Hamiltonian \tilde{H} . So, let us define the right and left eigenvector equations for \tilde{H}^*

$$\tilde{H}^* \phi_j = \lambda_j \phi_j, \quad \hat{\phi}_j^T \tilde{H}^* = \lambda_j \hat{\phi}_j^T. \quad (4.14)$$

So, taking the transpose of the left eigenvalue equation in (4.14) and subsequently considering the pseudo-anti-Hermiticity relationship (4.3) one can show the next relationship

$$\phi_{\bar{j}} = \mathcal{G}^{-1} \hat{\phi}_j, \quad (4.15)$$

where $\phi_{\bar{j}}$ denotes a right eigenvector with eigenvalue $(-\lambda_j)$.

Similarly, one can relate the above left eigenvector $\hat{\phi}^T$ to the left eigenvector of type $\bar{\eta}^T$ in (3.16) for the conjugate eigenvalue λ_j^*

$$\hat{\phi}_j^T = \bar{\eta}_j^\dagger. \quad (4.16)$$

Moreover, using the symmetry (4.3) and the right eigenvector equation (3.16) one can write the following left eigenvector equations

$$(\eta^\dagger \sigma_-) \tilde{H} = -\lambda^* (\eta^\dagger \sigma_-), \quad (4.17)$$

$$(\eta^\dagger \mathcal{G}) \tilde{H}^* = -\lambda^* (\eta^\dagger \mathcal{G}). \quad (4.18)$$

So, by comparing (4.18) with the left-eigenvector equation in (4.14) one can conclude that for each left eigenvector $\hat{\phi}_j^T$ with eigenvalue λ_j one must have another eigenvector $(\eta^\dagger \mathcal{G})$ with eigenvalue $-\lambda_j^*$, i.e.

$$\hat{\phi}_{-\lambda_j^*}^T = (\eta_{\lambda_j}^\dagger \mathcal{G}), \quad (4.19)$$

with η_j being the right eigenvector of \tilde{H} with eigenvalue λ_j defined in (3.16).

So, in contradistinction to the pseudo-chiral symmetry (3.2) which ensures that for each energy level λ there exists the opposite $-\lambda$, i.e. the pairs $(\lambda, -\lambda)$, the symmetries (4.1) and (4.3) ensure the energy levels in pairs with identical imaginary part and opposite real part, i.e. the pairs $(\lambda, -\lambda^*)$. Remarkably, in our constructions of the next section, we will provide analytic realizations of these eigenvalue and eigenvector symmetries for a continuum of spinor bound states and extended wave states in the localized continuum.

We write the biorthogonality relationship in this case. So, from (4.4) one can get

$$\hat{\chi}_i^T H^* \chi_j = \lambda_j \hat{\chi}_i^T \chi_j, \quad \hat{\chi}_i^T H^* \chi_j = \lambda_i \hat{\chi}_i^T \chi_j, \quad (4.20)$$

which implies

$$0 = (\lambda_i - \lambda_j) \hat{\chi}_i^T \chi_j \Rightarrow \hat{\chi}_i^T \chi_j = \delta_{ij}. \quad (4.21)$$

Similar construction from (4.14) allows one to write

$$0 = (\lambda_i - \lambda_j) \hat{\phi}_i^T \phi_j \Rightarrow \hat{\phi}_i^T \phi_j = \delta_{ij}. \quad (4.22)$$

The Table 1 summarizes the main results related to the pseudo-chiral and (σ_-) -anti-pseudo-Hermitian symmetries and their relevant representations, respectively, for the set of Hamiltonians $\{H, \tilde{H}\}$ and $\{H^*, \tilde{H}^*\}$. So, we have organized, for each type of symmetry, the corresponding eigenvectors and their mutual relationships, as well as the allowed values of the spectral parameter λ permitted by the relevant symmetry. The corresponding biorthogonality relationships and the appearance of the Majorana zero modes at $\lambda = 0$ and the coalescence of the eigenvectors are also included.

Table 1: Pseudo-chiral and anti-pseudo-(anti-)Hermitian symmetries and representations

	Pseudo-chiral	Anti-pseudo-Hermitian
Symmetry	$\mathcal{G}X\mathcal{G}^{-1} = -X^T, X \equiv \{H, \tilde{H}\}$	$\sigma_- X \sigma_-^{-1} = -X^\dagger, X \equiv \{H, \tilde{H}\}$
Right-eigenvectors	$H\xi = \lambda\xi, \tilde{H}\eta = \lambda\eta$	$H^*\chi = \lambda\chi, \tilde{H}^*\phi = \lambda\phi$
Left-eigenvectors	$\Gamma H \Gamma^{-1} = \tilde{H} \rightarrow \eta = \Gamma\xi, \bar{\eta} = \Gamma\bar{\xi}$	$\Gamma H^* \Gamma^{-1} = \tilde{H}^* \rightarrow \phi = \Gamma\chi, \hat{\phi} = \Gamma\hat{\chi}$
	$\bar{\xi}^T H = \lambda\bar{\xi}^T, \bar{\eta}^T \tilde{H} = \lambda\bar{\eta}^T$	$\hat{\chi}^T H^* = \lambda\hat{\chi}^T, \hat{\phi}^T \tilde{H}^* = \lambda\hat{\phi}^T$
Eigenvalues	$\{\lambda, -\lambda\}$	$\{\lambda, \lambda^*, -\lambda^*\}$
R/L-eigenvectors, $(\lambda, -\lambda)$	$\bar{\xi}_{-\lambda} = \mathcal{G}\xi_\lambda, \bar{\eta}_{-\lambda} = \mathcal{G}\eta_\lambda$	$\chi_{-\lambda} = \mathcal{G}^{-1}\hat{\chi}_\lambda, \phi_{-\lambda} = \mathcal{G}^{-1}\hat{\phi}_\lambda$
R/L-eigenvectors, $(\lambda, \pm\lambda^*)$	-	$\hat{\chi}_{-\lambda^*} = -\mathcal{G}\xi_\lambda^*, \hat{\phi}_{-\lambda^*} = -\mathcal{G}\eta_\lambda^*$
	-	$\hat{\chi}_{\lambda^*} = -\bar{\xi}_\lambda^*, \hat{\phi}_{\lambda^*} = -\bar{\eta}_\lambda^*$
Biorthogonality	$\bar{\xi}_i^T \xi_j = \delta_{ij}, \bar{\eta}_i^T \eta_j = \delta_{ij}$	$\hat{\chi}_i^T \chi_j = \delta_{ij}, \hat{\phi}_i^T \phi_j = \delta_{ij}$
Majorana modes ($\lambda = 0$)	$\bar{\xi}_0^T \xi_0 = 0, \bar{\eta}_0^T \eta_0 = 0$	$\hat{\chi}_0^T \chi_0 = 0, \hat{\phi}_0^T \phi_0 = 0$

4.2 Some properties of pseudo-Hermitian Hamiltonians

Let us discuss some theorems and properties of pseudo-Hermiticity and anti-pseudo-Hermiticity for NH Hamiltonians following the approach in [6, 7, 8, 46], which are relevant to our model. A NH Hamiltonian H is said to be a ζ -pseudo-Hermitian Hamiltonian if one has

$$\zeta H \zeta^{-1} = H^\dagger, \quad (4.23)$$

for ζ being a linear invertible Hermitian operator. Notice that for $\zeta = \mathbf{1}$ one recovers a Hermitian Hamiltonian, i.e. $H^\dagger = H$. On the other hand, a σ -anti-pseudo-Hermitian Hamiltonian is defined as [7, 46]

$$\sigma H \sigma^{-1} = H^\dagger, \quad (4.24)$$

for σ being an anti-linear Hermitian automorphism. These types of anti-linear operators appearing in the definition of pseudo-Hermiticity have further been classified into σ_\pm , Hermitian [46] and anti-Hermitian ones [7], respectively. The anti-linear operator σ_- in (4.1) belongs to the anti-Hermitian type in this classification; so, H in (2.9) which satisfies (4.1), can be dubbed as (σ_-) -anti-pseudo-(anti-)Hermitian Hamiltonian. An equivalence between a ζ -pseudo-Hermiticity and a σ -anti-pseudo-Hermiticity has been established using spectral decomposition techniques in [7, 46].

Next, we summarize the main properties of pseudo-Hermitian operators [6, 8] to be useful for our discussions of the spectra of the pseudo-Hermitian Hamiltonian (2.9). Let $H : \mathcal{H} \rightarrow \mathcal{H}$ be a diagonalizable linear operator acting on a Hilbert space \mathcal{H} and $\zeta : \mathcal{H} \rightarrow \mathcal{H}$ be a linear Hermitian automorphism. Then

- 1). H is pseudo-Hermitian if and only if its eigenvalues are real or come in complex-conjugate pairs.

2). if H is pseudo-Hermitian with respect to two linear Hermitian automorphisms ζ_1 and ζ_2 , then $\zeta_1^{-1}\zeta_2$ generates a symmetry of H , i.e.

$$[\zeta_1^{-1}\zeta_2, H] = 0. \quad (4.25)$$

Moreover, in [7] it has been shown that every diagonalizable Hamiltonian possessing an antilinear symmetry is necessarily pseudo-hermitian. This applies, in particular, to diagonalizable PT-symmetric Hamiltonians. The proof of the equivalence of pseudo-hermiticity and the presence of antilinear symmetries has been performed provided that the Hamiltonian is diagonalizable and has a discrete spectrum. It used the fact that such Hamiltonian H is anti-pseudo-hermitian, i.e. there is an antilinear, Hermitian, invertible operator as σ in (4.24). The analogous properties relevant to field theory models, to our knowledge, have not yet been proved; however, it is expected that some of them must hold, with relevant modifications, for the NH Hamiltonians in the field theory context. In fact, we will see below that the eigenvalues of the field theory NH model described by the (σ_-) -anti-pseudo-Hermitian Hamiltonian H in (2.9) come in complex-conjugate pairs and some of the Hamiltonian symmetries (4.25) can be realized provided that the complex scalar Φ components satisfy some definite parities under space inversion.

4.3 \mathcal{P} , \mathcal{T} and \mathcal{PT} symmetries

So, in order to examine additional symmetries we consider the \mathcal{P} , \mathcal{T} and \mathcal{PT} symmetry transformations of the NH Hamiltonian (2.9). Let us define the next symmetry transformations for the scalar field components Φ_a , the complex parameter z and the space coordinate x

$$\Phi \equiv \Phi_1 + i\Phi_2, \quad (4.26)$$

$$\mathcal{P}\Phi_a\mathcal{P} = \nu_a\Phi_a, \quad \nu_a = \pm 1, \quad a = 1, 2. \quad (4.27)$$

$$\mathcal{P}x\mathcal{P} = -x, \quad \mathcal{T}z\mathcal{T}^{-1} = z^*. \quad (4.28)$$

Next, we will assume the scalar field components Φ_a ($a = 1, 2$) to possess definite parities $\nu_a = \pm 1$ as in (4.27). So, let us consider the following cases: **Case A.** $\nu_1 = +1$, $\nu_2 = +1$. **Case B.** $\nu_1 = +1$, $\nu_2 = -1$. **Case C.** $\nu_1 = -1$, $\nu_2 = +1$. And **Case D.** $\nu_1 = -1$, $\nu_2 = -1$.

One can show the next relationships for the Hamiltonian H in (2.9) upon making the relevant transformations in its matrix representation.

Case A. $\nu_1 = +1$, $\nu_2 = +1$. In this case one has

$$\mathcal{P}\gamma_5 H \gamma_5 \mathcal{P}^{-1} = -H. \quad (4.29)$$

This implies the anti-symmetry $\{\mathcal{P}\gamma_5, H\} = 0$.

Case B. $\nu_1 = +1$, $\nu_2 = -1$. In this case one has for M real

$$\mathcal{P}H\mathcal{P}^{-1} = H^\dagger. \quad (4.30)$$

Making use of the anti-pseudo-Hermiticity relationship (4.1) together with (4.30) one can get the anti-symmetry $\{\sigma_-^{-1}\mathcal{P}, H\} = 0$.

Case C. $\nu_1 = -1, \nu_2 = +1$.

$$\mathcal{PT}\gamma_5 H [\mathcal{PT}\gamma_5]^{-1} = H. \quad (4.31)$$

Then, in this case the NH Hamiltonian H possesses the antilinear symmetry $\mathcal{PT}\gamma_5$, i.e. $[\mathcal{PT}\gamma_5, H] = 0$. So, the antilinear \mathcal{PT} -like symmetry (4.31) must be taken into account together with the pseudo-chirality and anti-pseudo-Hermiticity of the model in order to examine its spectral properties in this case.

Case D. $\nu_1 = -1, \nu_2 = -1$.

In this case one has the relationship

$$\bar{\sigma}_- H \bar{\sigma}_-^{-1} = -H^\dagger, \quad (4.32)$$

$$\bar{\sigma}_- \equiv \mathcal{PT}. \quad (4.33)$$

Notice that $\bar{\sigma}_-$ is an antilinear operator. So, this Hamiltonian is a (anti)-pseudo-anti-Hermitian Hamiltonian as in (4.1), however in this particular case under the \mathcal{PT} operator $\bar{\sigma}_-$. Considering the both pseudo-anti-Hermiticity relationships (4.32) and (4.1) one can show that

$$\sigma_-^{-1} \bar{\sigma}_- H (\sigma_-^{-1} \bar{\sigma}_-)^{-1} = H \quad (4.34)$$

$$(\mathcal{PG})H(\mathcal{PG})^{-1} = H, \quad (4.35)$$

where the product $\sigma_-^{-1} \bar{\sigma}_- = \mathcal{PG}$ has been inserted in the eq. (4.35). So, (4.35) implies the symmetry

$$[H, \mathcal{PG}] = 0. \quad (4.36)$$

Then, (4.36) establishes a symmetry of type $\sigma_-^{-1} \bar{\sigma}_-$ for a Hamiltonian which is pseudo-anti-Hermitian with respect to two different anti-linear operators σ_- and $\bar{\sigma}_-$. So, it is an analogous symmetry to the one in (4.25) which establishes a type of symmetry $(\zeta_1^{-1} \zeta_2)$ for pseudo-Hermitian Hamiltonians, i.e. with pseudo-Hermiticity property with respect to two different linear Hermitian operators ζ_1 and ζ_2 .

Below, in the context of the tau-function approach, we will perform the construction of the right and left eigenvectors satisfying the pseudo-chirality and pseudo-Hermitian properties discussed above, and for a special solution we will show a NH Majorana zero mode-soliton duality description of the model, such that the static scalar field is a self-consistent solution of the equation (2.14) and (2.19). In fact, we will realize the theorem 1) above by showing that the both NH Hamiltonians in (2.9) possess complex-conjugate pairs of eigenvalues.

In the Table 2 we summarize the main results of the cases A, B, C and D discussed above. We present the interplay of the \mathcal{P} , \mathcal{T} and \mathcal{PT} symmetry transformations together with the pseudo-chiral and anti-pseudo-Hermiticity properties, which imply a (anti)-symmetry of the Hamiltonian (last column of the Table 2) for each set of parities (ν_1, ν_2) of the scalar fields Φ_1 and Φ_2 , respectively.

Table 2: \mathcal{P} , \mathcal{T} and \mathcal{PT} symmetries

Case	Parity:(ν_1, ν_2)	H' 's property	(Anti-)symmetry
A	(+ +)	Pseudo-chiral+Anti-pseudo-Hermitian	$\{\gamma_5 \mathcal{P}, H\} = 0$
B	(+ -)	Anti-pseudo-Hermitian	$\{\sigma^{-1} \mathcal{P}, H\} = 0$ ($M \in \mathbb{R}$)
C	(- +)	Pseudo-chiral+Anti-pseudo-Hermitian	$[\gamma_5 \mathcal{PT}, H] = 0$
D	(- -)	Anti-pseudo-Hermitian	$[\mathcal{PG}, H] = 0$

5 NH topological solitons coupled to the fermions

In this section we find exact analytical solutions of the system of eqs. (2.7)-(2.8) and (2.13) associated to the NH model (2.1) and search for nontrivial topological configurations for the complex field Φ interacting with the bound states formed by the two fermion fields ξ and η . The Hermitian case has been considered before by defining a submodel for Φ being a real scalar and by imposing the reality condition $\tilde{\psi} \equiv \psi^*$ to the model (2.1); so, the reduced model defined a Dirac field interacting with real Toda solitons [36, 40, 37, 23, 28, 22, 21].

5.1 Topological solitons and spinor bound states

The soliton solutions, as well as the bound states of the Hermitian version of this model has been considered in [21] through the tau function formalism. Here we perform the construction of the general complex 1-solitons and spinor bound state configurations of the non-Hermitian model (2.1). In this context, for a suitable subspace in parameter space, it will emerge also some extended wave states in the localized continuum.

Since the pair of first order differential equations (2.7)-(2.8) for the spinors plus the first order equation (2.14) for Φ imply the second order differential eq. for the complex scalar (2.13), each set of solutions of the first order system of equations will be solutions of the second order differential equation (2.13). So, it is sufficient to consider the system of equations (2.7)-(2.8) and (2.14) in order to find the complex solitons and related bound state solutions.

In the next steps we rewrite the equations of motion of the model in terms of the tau functions $\tau_{0,1}$, $\tau_{A,B}$, $\tilde{\tau}_{A,B}$ as

$$e^{i\beta\Phi} = \rho \left(\frac{\tau_0}{\tau_1} \right)^2, \quad (5.1)$$

$$\xi_A = \frac{\tau_A}{\tau_0}, \quad \xi_B = \frac{\tau_B}{\tau_1}, \quad \eta_A = \frac{\tilde{\tau}_A}{\tau_1}, \quad \eta_B = \frac{\tilde{\tau}_B}{\tau_0}, \quad (5.2)$$

with ρ being a constant complex parameter. Next, substituting the field components (5.1)-(5.2) into the system of first order equations (2.7)-(2.8) and (2.14) one has, respectively

$$i\lambda\tau_1\tau_B - M\rho\tau_0\tau_A + \tau_B\tau_1' - \tau_1\tau_B' = 0 \quad (5.3)$$

$$i\lambda\tau_0\tau_A + M\rho^{-1}\tau_1\tau_B - \tau_A\tau_0' + \tau_0\tau_A' = 0 \quad (5.4)$$

$$i\lambda\tau_0\tilde{\tau}_B + M\rho^{-1}\tau_1\tilde{\tau}_A - \tilde{\tau}_B\tau_0' + \tau_0\tilde{\tau}_B' = 0 \quad (5.5)$$

$$i\lambda\tau_1\tilde{\tau}_A - M\rho\tau_0\tilde{\tau}_B + \tilde{\tau}_A\tau_1' - \tau_1\tilde{\tau}_A' = 0. \quad (5.6)$$

and

$$2i(\tau_0'\tau_1 - \tau_0\tau_1') + M\beta(\tau_A\tilde{\tau}_A + \tau_B\tilde{\tau}_B) = 0. \quad (5.7)$$

Next, let us assume the following tau functions

$$\tau_1 = 1 + ae^{2kx}, \quad \tau_0 = 1 + be^{2kx}, \quad (5.8)$$

$$\tau_A = ce^{kx}, \quad \tau_B = de^{kx}, \quad (5.9)$$

$$\tilde{\tau}_A = \tilde{c}e^{kx}, \quad \tilde{\tau}_B = \tilde{d}e^{kx}, \quad (5.10)$$

with the complex parameters $a, b, c, d, \tilde{c}, \tilde{d}$ and k . So, the above tau functions satisfy the system of equations (5.3)-(5.6) and (5.7) provided that

$$a = \frac{\tilde{c}c}{2M} \frac{\beta^2 z^3}{(z^2 - 1)^2}, \quad \tilde{d} = \frac{\tilde{c}c}{d}, \quad \rho = -i\frac{d}{c}z, \quad (5.11)$$

$$b = \frac{\tilde{c}c}{2M} \frac{\beta^2 z}{(z^2 - 1)^2}, \quad z \equiv -\frac{\lambda}{M} \pm \sqrt{\frac{\lambda^2}{M^2} - 1} \quad (5.12)$$

$$\lambda = -\frac{M}{2}(z + z^{-1}), \quad k = \frac{iM}{2}(z - z^{-1}). \quad (5.13)$$

From the above equations one can write

$$\lambda^2 + k^2 = M^2. \quad (5.14)$$

The equation (5.14) can be regarded as a dispersion relationship between the complex energy λ and the complex wave number k of the quasi-particles for a fixed value of the complex mass parameter M . Notice that this equation is invariant under the transformations

$$\lambda \rightarrow -\lambda \quad (5.15)$$

$$k \rightarrow -k. \quad (5.16)$$

So, the eigenvalues come in pairs of type $(\lambda, -\lambda)$. This is in accordance to the pseudo-chirality symmetry property discussed in section 3. Then, one can argue that the eigenvectors corresponding to this pair of eigenvalues realize the pseudo-chirality symmetry (3.2) and the left $\bar{\xi}_{-\lambda}$ and right eigenvectors ξ_λ can be related by (3.8).

Next, one can take the complex conjugate of the dispersion relationship (5.14) and get

$$(\lambda^*)^2 + (k^*)^2 = (M^*)^2. \quad (5.17)$$

Since this equation defines a new eigenvalue λ^* , one has that the eigenvalues come in complex-conjugate pairs. Notice that (5.17) is invariant under $\lambda^* \rightarrow -\lambda^*$, $k^* \rightarrow -k^*$. So, from (5.14) and (5.17) and their

relevant symmetries under inversion $\lambda \rightarrow -\lambda$ and $\lambda^* \rightarrow -\lambda^*$, one can argue that the solutions above describe the corresponding left and right eigenvectors with the set of eigenvalues $\{\lambda, -\lambda, \lambda^*, -\lambda^*\}$.

One can see that the complex dispersion relationship (5.17) arises from the tau function representation of the eigenvalue equation in (4.4) for the right eigenvector χ and the operator H^* . In fact, by taking the complex conjugate of $H\xi = \lambda\xi$ in (3.4) and the relevant tau function components in (5.1)-(5.2) and subsequent substitution into the corresponding tau function equations analogous to (5.3)-(5.6) one can get (5.17).

In fact, these solutions become the realizations of the pseudo-chiral and pseudo-Hermitian symmetries of the model discussed in the sections 3 and 4, respectively. Remarkably, the relevant spectral properties associated to the symmetries (3.2) and (4.1) are realized above by providing the set of eigenvectors with eigenvalues $\{\lambda, -\lambda\}$ and $\{\lambda, -\lambda^*\}$, respectively. Then, the eigenvalues come in complex conjugate pairs and their reversions. This is in accordance to the first theorem 1) which must hold for pseudo-anti-Hermitian Hamiltonians, once the equivalence between the pseudo-Hermiticity and the anti-pseudo-Hermiticity properties is assumed [7, 46].

The energy λ and wave number k in (5.13) are defined in terms of the uniformization complex variable z . The Fig.1 shows some special points where $\lambda(z)$ exhibits a pole at the origin $z = 0$ and the zero modes at $z = \pm i$. The points $z = \pm 1$ correspond to the edge states $k = 0 \rightarrow \lambda = \pm M$. Next, we define the complex parameters such that

$$M \equiv M_1 + i M_2, \quad (5.18)$$

$$\lambda \equiv \lambda_1 + i \lambda_2, \quad (5.19)$$

$$k \equiv k_1 + i k_2, \quad (5.20)$$

and

$$a = |a|e^{i\alpha_1}, b = |b|e^{i\alpha_2}, \rho = |\rho|e^{i\delta}, c = |c|e^{i\zeta_1}, d = |d|e^{i\zeta_2}. \quad (5.21)$$

So, from the eqs. (5.11)-(5.12) the complex variable z can be written as

$$z = \sqrt{|a|/|b|} e^{i\sigma}, \quad \sigma \equiv \frac{\alpha_1 - \alpha_2}{2}. \quad (5.22)$$

Next, taking into account (5.18)-(5.20) and (5.11)-(5.13) one can write

$$z = \sqrt{\frac{(\lambda_1 - k_2)^2 + (\lambda_2 + k_1)^2}{|M|^2}} e^{i\sigma}, \quad \sigma \equiv \arctan \left[\frac{M_2(k_2 - \lambda_1) + M_1(k_1 + \lambda_2)}{M_2(k_1 + \lambda_2) - M_1(k_2 - \lambda_1)} \right]. \quad (5.23)$$

Notice that from (5.14) one can consider a complex function $\lambda = \pm\sqrt{M^2 - k^2}$ defined on a Riemann surface consisting of two sheets, with $\pm M$ being the branch points, which are necessary to make the square-root function single valued. However, the convenient parametrization in the complex z variable defined in (5.13) avoids the introduction of this two-sheet surface, see below more about this construction.




Figure 1: The complex plane defined by the uniformization variable z . The complex energy $\lambda(z)$ (5.13) exhibits a pole at the origin $z = 0$ and the zeros at $z = \pm i$. The points $z = \pm 1$ correspond to the edge states $k = 0$ ($\lambda = \pm M$).

The scalar field component solutions from (5.1) can be written as

$$\Phi_1 = \frac{2}{\beta} \arctan \left[i \frac{e^{-i\frac{\delta}{2}} \tau_0^* \tau_1 - e^{i\frac{\delta}{2}} \tau_0 \tau_1^*}{e^{i\frac{\delta}{2}} \tau_0 \tau_1^* + e^{-i\frac{\delta}{2}} \tau_0^* \tau_1} \right] \quad (5.24)$$

$$\Phi_2 = -\frac{1}{\beta} \log \left[|\rho| \frac{\tau_0^* \tau_0}{\tau_1^* \tau_1} \right]. \quad (5.25)$$

Next, taking into account the tau functions (5.8)-(5.10) one can write

$$\Phi_1 = \frac{2}{\beta} \arctan \left[\frac{\sin \frac{\delta}{2} + (|b| \sin(\Xi_2(x)) - |a| \sin(\Xi_1(x))) e^{2k_1 x} + |a||b| \sin(\Theta) e^{4k_1 x}}{\cos \frac{\delta}{2} + (|b| \cos(\Xi_2(x)) + |a| \cos(\Xi_1(x))) e^{2k_1 x} + |a||b| \cos(\Theta) e^{4k_1 x}} \right] \quad (5.26)$$

$$\Xi_1(x) \equiv 2k_2 x + \alpha_1 - \frac{\delta}{2}, \quad \Xi_2(x) \equiv 2k_2 x + \alpha_2 - \delta/2, \quad \Theta \equiv \alpha_2 - \alpha_1 + \frac{\delta}{2}$$

and

$$\Phi_2 = -\frac{1}{\beta} \log \left[|\rho| \frac{1 + 2|b| \cos(2k_2 x + \alpha_2) e^{2k_1 x} + |b|^2 e^{4k_1 x}}{1 + 2|a| \cos(2k_2 x + \alpha_1) e^{2k_1 x} + |a|^2 e^{4k_1 x}} \right]. \quad (5.27)$$

For the wave number with real component different from zero, i.e. $k_1 \neq 0$, one can define the next topological invariants for each component of the complex scalar field as

$$\Phi_1(+\infty) - \Phi_1(-\infty) = \pm \frac{2}{\beta} (\alpha_1 - \alpha_2), \quad (5.28)$$

$$= \pm \frac{4}{\beta} \sigma, \quad (5.29)$$

$$= \pm \frac{4}{\beta} \arg z \quad (5.30)$$

$$\Phi_2(+\infty) - \Phi_2(-\infty) = \pm \frac{2}{\beta} \log \frac{|b|}{|a|}, \quad (5.31)$$

$$= \pm \frac{4}{\beta} \log |z| \quad (5.32)$$

where σ ($\sigma = \arg z$) and $|z|$ are the argument and modulus of the complex parameter z , respectively, presented in (5.22)-(5.23). Remarkably, from the results (5.30) and (5.32) one can write the next suggestive

formula

$$e^{i\beta(\Phi(+\infty)-\Phi(-\infty))} = e^{i\beta(\Phi_1(+\infty)-\Phi_1(-\infty))} e^{-\beta(\Phi_2(+\infty)-\Phi_2(-\infty))} \quad (5.33)$$

$$= z^{\pm 4}. \quad (5.34)$$

This implies that

$$\Phi(+\infty) - \Phi(-\infty) = \pm \frac{4}{i\beta} \log z. \quad (5.35)$$

Next, we will regard the first order differential equation (2.14) as an equivalence relationship between the topological charge density associated to the complex scalar and the spinor current density defined by a sum of the spinor bilinear terms. So, integrating (2.14) in the whole real line, one can define the next topological charges

$$[\Phi_1 + i\Phi_2] \Big|_{x=-\infty}^{x=+\infty} = -\frac{\beta}{2} \int_{-\infty}^{+\infty} (\eta_A \xi_A + \eta_B \xi_B) dx, \quad (5.36)$$

$$\mathcal{Q}_1 \equiv \frac{\beta}{2\pi} [\Phi_1(+\infty) - \Phi_1(-\infty)] \quad (5.37)$$

$$\mathcal{Q}_2 \equiv \frac{\beta}{2\pi} [\Phi_2(+\infty) - \Phi_2(-\infty)]. \quad (5.38)$$

Then, one can write

$$\mathcal{Q}_1 \equiv \text{Re} \left[-\beta^2 \int_{-\infty}^{+\infty} \xi_A \xi_B dx \right] \quad (5.39)$$

$$\mathcal{Q}_2 \equiv \text{Im} \left[-\beta^2 \int_{-\infty}^{+\infty} \xi_A \xi_B dx \right] \quad (5.40)$$

where the relationship (2.11) between the ξ and η fields has been used.

In some cases the solutions $\Phi_{1,2}$ exhibit vanishing topological charges. In fact, for $\alpha_2 = \alpha_1$ ($\arg z = 0$) one has $\Phi_1(+\infty) = \Phi_1(-\infty)$, while for $|b| = |a|$ ($|z| = 1$) one gets $\Phi_2(+\infty) = \Phi_2(-\infty)$. These would imply $\mathcal{Q}_1 = 0$ or $\mathcal{Q}_2 = 0$ in (5.37)-(5.38), respectively.

Next, let us discuss the analytical properties of the multi-valued function λ associated to the complex plane defined by the variable z . The so-called uniformization variable z has been introduced for the energy λ and wave number k in (5.13). This parametrization avoids the introduction of the Riemann surface consisting of two sheets with relevant branching points as mentioned above. Notice that the relationship (5.14) will hold for any arbitrary complex number z in the entire complex plane excluding the origin, i.e. $z \in \mathbb{C} \setminus \{0\}$.

From (5.13) one can show the next useful identity

$$\lambda = -\frac{M}{2}(z + z^{-1}) \text{ and } k = \frac{iM}{2}(z - z^{-1}) \Rightarrow z^2 = \frac{\lambda + ik}{\lambda - ik}, \quad (M \neq 0). \quad (5.41)$$

Notice that the phase and modulus of the uniformization complex variable z encode the topological charges of the two types of solitons as shown in (5.30) and (5.32). So, we write the complex variable z in polar form such that the modulus $|z|$ and $\arg z$ depend on the real and complex parts of λ and k as follow

$$z = \left(\frac{|\lambda + ik|}{|\lambda - ik|} \right)^{1/2} e^{i\sigma}, \quad \sigma = \arctan \left\{ \frac{|k|^2 - |\lambda|^2 + |\lambda - ik||\lambda + ik|}{\lambda k^* + \lambda^* k} \right\}. \quad (5.42)$$




Figure 2: A representative of the continuum of bound states. Topological complex soliton $\Phi = \Phi_1 + i\Phi_2$ and bound state $\xi^T = (\xi_3 + i\xi_4, \xi_1 + i\xi_2)$ for $|a| = 2, |b| = 1.5, |c| = 1, |d| = 1.3, \zeta_1 = 1.2, \zeta_2 = 1.5, \alpha_1 = 1.1, \alpha_2 = 1.7, k_1 = 1, k_2 = 1, \beta = 1, |\rho| = 1.5, M_1 = 4.04, M_2 = 1.49$.

Then, using the results (5.30) and (5.32) together with the polar decomposition (5.42) into the formulas (5.37) and (5.38) for the topological charges one has

$$\mathcal{Q}_1 = \pm \frac{2}{\beta\pi} \arctan \left\{ \frac{|k|^2 - |\lambda|^2 + |\lambda - ik| |\lambda + ik|}{\lambda k^* + \lambda^* k} \right\}, \quad (5.43)$$

$$\mathcal{Q}_2 = \pm \frac{1}{\beta\pi} \log \frac{|\lambda + ik|}{|\lambda - ik|}. \quad (5.44)$$

Therefore, one has a family of localized continuum states with complex energy λ and wave number k defining two types of solitons with topological charges provided in (5.43)-(5.44). The second topological soliton with charge \mathcal{Q}_2 owns its existence purely to the non-Hermiticity and non-linearity of the model. In fact for λ and k real one has a trivial topological charge, i.e. from (5.44) one has that $\mathcal{Q}_2 = 0$. So, the appearance of the second soliton shows an interesting phenomena in which the interplay between non-linearity and non-Hermiticity of the model gives rise to a new state provided by the composite topological soliton Φ with components (5.26) and (5.27) and the next spinor bound states

$$\xi_A = |c| e^{-i(k_2 x + \alpha_2 - \zeta_1)} \frac{e^{k_1 x} (|b| e^{2k_1 x} + e^{i(2k_2 x + \alpha_2)})}{(1 + |b| e^{2k_1 x})^2 - 2|b| e^{2k_1 x} (1 - \cos(2k_2 x + \alpha_2))}, \quad (5.45)$$

$$\xi_B = |d| e^{-i(k_2 x + \alpha_1 - \zeta_2)} \frac{e^{k_1 x} (|a| e^{2k_1 x} + e^{i(2k_2 x + \alpha_1)})}{(1 + |a| e^{2k_1 x})^2 - 2|a| e^{2k_1 x} (1 - \cos(2k_2 x + \alpha_1))}, \quad (5.46)$$

where the tau functions (5.8)-(5.10) have been inserted into (5.2) in order to write the spinor solutions (5.45)-(5.46). The components η_A and η_B can be shown to satisfy the relationships (2.12), so from (5.45)-(5.46) one can get $\eta_A = \xi_B$ and $\eta_B = \xi_A$.

The Fig. 2 shows a representative plot of the above type of localized states. In this figure the spinor bound states become localized inside the corresponding topological complex solitons supported by the both scalar field components. Qualitatively inspecting the Fig. 2 one can argue that for a general set of soliton parameters the solution does not present any of the \mathcal{PT} -like symmetries discussed in the Table 2, since the scalar fields do not have definite parities.

Moreover, the x -integrated current density in (2.14) can be written as

$$\int_{-\infty}^{+\infty} [\eta_A \xi_A + \eta_B \xi_B] dx = \frac{|c||d|}{|b|} \left\{ \frac{(\log \frac{|a|}{|b|})^2 + (\alpha_1 - \alpha_2)^2}{(k_1^2 + k_2^2) \left[(\frac{|a|}{|b|})^2 + 1 - 2 \frac{|a|}{|b|} \cos(\alpha_1 - \alpha_2) \right]} \right\}^{1/2} e^{i\Delta} \quad (5.47)$$

$$\Delta \equiv \zeta_1 + \zeta_2 + \delta_o + \alpha_o + \nu, \quad \nu = -\arctan \left[\frac{k_2}{k_1} \right], \quad (5.48)$$

$$\delta_o = \arctan \left\{ \frac{\alpha_1 - \alpha_2}{\log |a| - \log |b|} \right\}, \quad \alpha_o = \arctan \left\{ \frac{|b| \sin \alpha_2 - |a| \sin \alpha_1}{|a| \cos \alpha_1 - |b| \cos \alpha_2} \right\}. \quad (5.49)$$

So, the above integration (5.47) provides a complex number such that the real part will be proportional to \mathcal{Q}_1 and the imaginary part to \mathcal{Q}_2 as in (5.39)-(5.40), respectively. The equations (5.43)-(5.44) together with (5.47) will provide two relationships for the parameters. Additional relationships which must be satisfied by the parameters come from the biorthogonality relationship $(\bar{\xi}_\lambda | \xi_\lambda)$ in (3.5) which will be computed below for any complex eigenvalue λ .

Notice that for the Hermitian case one has $|a| = |b|$, $k_2 = 0$ and since $\tau_0 = \tau_1^*$ in (5.8) one can take $\alpha_1 = \frac{\pi}{2}$, $\alpha_2 = \frac{3\pi}{2}$, and then $\nu = 0$, $\delta_o = \frac{\pi}{2}$, $\alpha_o = \frac{\pi}{2}$. So, it must be $\zeta_1 + \zeta_2 = n\pi$ in order to fix $e^{i\Delta} = \pm 1$, such that the charge \mathcal{Q}_2 vanishes.

So, the right-eigenvector fermion and the complex topological soliton Φ currents, which satisfy the equivalence relationship (2.14), as well as the relevant equivalence of the left-eigenvector fermion current to the scalar field $\bar{\Phi}$ topological current in (2.19), can be regarded as the analogs of the biorthogonal generalized bulk-boundary correspondence in non-Hermitian condensed matter systems [24].

One can see that an essential distinction between the Hermitian Toda model coupled to fermion and its non-Hermitian counterpart is the degrees of freedom that one can access to; i.e. nonunitary transformations forbidden in the Hermitian system can be realized in its non-Hermitian version. In fact, the change in the spectrum from real parameters (λ_1, k_1, M_1) to the complex ones (λ, k, M) increases the number of parameters that describe the system. Since topology mainly depends on the underlying manifold, non-Hermiticity is expected to alter the topological classification of the solitons related to the scalar field components Φ_1 and Φ_2 . So, in order to classify the topological sectors of the NH model, the pseudo-chiral and anti-pseudo-Hermitian symmetries will play an important role.

Next let us discuss the physical origin of the non-Hermiticity. Some physical effects which introduce non-Hermiticity into Hermitian systems are, gain and loss, dissipation of energy in open quantum systems, finite life-time of resonances, etc. In this context, it is usual to ascribe some physical meaning to the real and imaginary parts of a complex eigenvalue, i.e. the real part codifies the excitation energy, whereas the imaginary part encrypts the decay or amplification rate. However, in our constructions above, one can see that the modulus and phase of the uniformization complex parameter z in (5.41) acquire some physical meaning. In fact, as mentioned above, the phase, $\arg z$, and modulus, $|z|$, encode the topological charges of the scalar fields Φ_1 and Φ_2 in (5.30) and (5.32), respectively. Remarkably, the circle $|z| = 1$ will correspond to the trivial topological charge associated to the field Φ_2 . As we will discuss below, the bound states in the

Hermitian system (for $\Phi_2 = 0$) lie on the unit circle $|z| = 1$. So, one can argue that the topological soliton associated to Φ_2 is purely of NH origin. An alternative interpretation into the realm of non-Hermitian physics, which involves neither gain nor loss, has recently been introduced in [38], in which the complex eigenvalues emerge from the amplitudes and phase differences of waves backscattered from the boundary of insulators. So, we believe that the physical origin of the nonhermiticity of our model can be understood in the framework of the inverse scattering transform formalism associated to the model, which deserves a careful consideration in future.

In summary, the tau-function Ansatz (5.1)-(5.2) together with (5.8)-(5.10) provide the solution $\{\Phi, \xi, \eta\}$ ($\eta = \Gamma\xi$) which realizes the pseudo-chiral and anti-pseudo-Hermitian symmetries of the NH system defined by the Hamiltonians H and \tilde{H} . In fact, in order to realize the pseudo-chirality one can see in the Table 1 that once the right-eigenvectors $\{\xi_\lambda, \eta_\lambda\}$ are known one can construct the left-eigenvectors as $\bar{\xi}_\lambda = \mathcal{G}\xi_{-\lambda}$ and $\bar{\eta}_\lambda = \mathcal{G}\eta_{-\lambda}$. Likewise, the anti-pseudo-Hermitian symmetry is realized by constructing the right and left eigenvectors of H^* and \tilde{H}^* , respectively, as $\chi_\lambda = -\xi_{-\lambda}^*$ and $\hat{\chi}_\lambda = -\mathcal{G}\xi_{-\lambda}^*$, and supplemented by $\phi = \Gamma\chi_\lambda$ and $\hat{\phi} = \Gamma\hat{\chi}_\lambda$ for the relevant eigenvectors of \tilde{H}^* .

5.2 Real eigenvalues and pseudo-chiral symmetry

The model exhibits a \mathcal{PT} -like symmetry under special circumstances; for example in the case C of Table 2, in which the scalar field components exhibit the parities $(-, +)$. So, in this case one could resort to the balanced gain and loss concept, as in the so-called ‘ \mathcal{PT} -symmetric’ systems, in order to determine the real eigenvalue sector of the model. However, as we have seen above the model presents a rich variety of topological solitons and related bound states without presenting definite parities for the scalar field components. For example, inspecting qualitatively the Fig. 2 one can see that the scalar solitons do not exhibit definite parities. So, some questions arise regarding the conditions under which the NH model (2.1) would present real eigenvalues and the existence of points for pseudo-Hermiticity breaking. Here, we examine the first issue by considering the dispersion relation in the complex plane defined by the uniformization parameter z . So, taking $M = |M|e^{i\delta}$ and $z = |z|e^{i\sigma}$ in (5.13) one can write

$$\lambda = -\frac{1}{2}|M|e^{i(\delta+\theta)}\sqrt{|z|^2 + \frac{1}{|z|^2} + 2\cos(2\sigma)}; \quad \theta \equiv \arctan\left[\left(\frac{|z|^2 - 1}{|z|^2 + 1}\right)\tan\sigma\right], \quad \delta \equiv \arctan\left(\frac{M_2}{M_1}\right). \quad (5.50)$$

From the last equation one can conclude that the eigenvalues λ will be complex numbers provided that $(\delta + \theta) \neq n\pi$ ($n \in \mathbb{Z}$). Therefore, from (5.50) it is clear that the eigenvalue λ will be a real number if $(\delta + \theta) = n\pi$ ($n \in \mathbb{Z}$). This last condition implies

$$\lambda_1(\sigma) = \pm|M|\sqrt{\frac{(1 + (\frac{M_2}{M_1})^2)(\sin\sigma)^2}{(\tan\sigma)^2 - (\frac{M_2}{M_1})^2}}, \quad \lambda_2 = 0, \quad (5.51)$$

$$|z|^2 = \frac{\tan\sigma - \frac{M_2}{M_1}}{\tan\sigma + \frac{M_2}{M_1}}, \quad \sigma \in (-\pi + |\delta|, -|\delta|) \cup (|\delta|, \pi - |\delta|). \quad (5.52)$$

Then, the NH model (2.1) supports real eigenvalues for the parameter subspace defined by the relationships (5.51)-(5.52). The Fig. 3 shows a family of curves in the complex plane z defined by (5.52) and parametrized by $\frac{M_2}{M_1}$. Notice that the family of curves intersect at the singular point $z = 0$, and at the points of the complex plane z corresponding to the zero mode $\lambda(z = \pm i) = 0$. Remarkably, one has $|z| \in [0, +\infty)$ for a fixed value of $\frac{M_2}{M_1}$. However, some regions are prohibited for the phase σ in the complex plane, as $\sigma \notin [-|\delta|, |\delta|]$ and $\sigma \notin \{[-\pi, -\pi + |\delta|] \cup [\pi - |\delta|, \pi]\}$. In the limit $\frac{M_2}{M_1} \rightarrow 0$ one has a unit circle at the center, and then $|z| = 1$, which implies $\lambda_1 = \pm|M| \cos \sigma$ ($\sigma \in [-\pi, +\pi]$). The Fig. 4 shows λ_1 vs σ for $\frac{M_2}{M_1} = 0.05$ (left) and for $\frac{M_2}{M_1} \rightarrow 0$ (right) defined by (5.51). As mentioned above, some intervals for σ are missing since λ_1 is not defined on these regions (Fig.4 left). Notice that the pseudo-chiral symmetry is maintained since the real eigenvalues come in pairs $\pm\lambda_1$.

So, our findings add new results to the current exploration whether the non-Hermitian topological models can exhibit real energy spectra. The studies in the literature along this line have used mainly the \mathcal{PT} symmetry in order to preserve the real spectra when non-Hermiticity is relatively weak. Recently, it has been put forward a systematic way to construct non-Hermitian topological systems exhibiting energy spectra that are always real, regardless of weak or strong non-Hermiticity [47]. The classification of the eigenspace of the anti-pseudo-Hermitian Hamiltonians H and \tilde{H} , such as the regions of real and complex eigenvalues and the points of eigenvector degeneracies, would be important in order to identify the points of pseudo-Hermiticity breaking. Certainly, a systematic examination of the necessary and sufficient conditions for symmetry breaking to occur in the anti-pseudo-Hermitian matrix H constitutes a future direction of research, e.g. in the lines recently developed in [48].

5.3 Spectral topology and winding number

One can analyze the spectral properties of the model by examining the behavior of the curves on the complex plane defined by the uniformization variable z and the energy λ and wave number k expressions in (5.41)-(5.42). Notice that for the bound states in the Hermitian case one has the real parameters $M = M_1$, $\lambda = \lambda_1$. So, when $k = k_1$ ($k_1 \in \mathbb{R}$) one has the unit circle $|z| = 1$ (see Fig.1). This implies that the bound states lie on this unit circle ($\lambda_1 = \pm\sqrt{M_1^2 - k_1^2}$). However, for purely imaginary $k = ik_2$ one notices that the scattering states are allowed for $z \in \mathbb{R}$ ($\lambda_1 = \pm\sqrt{M_1^2 + k_2^2}$). The so called half-bound states at the edges of the continuous spectrum are given at $\lambda_1 = \pm M_1 \rightarrow z = \pm 1$. So, the thresholds of the energy gap of the free massive fermions are at $\lambda_1 = \pm M_1$.

In the non-Hermitian case one has a complex eigenvalue spectrum, which must be studied on the complex plane, and so leads to the emergence of eigenvalue topology, or spectral topology. So, one has a continuum of bound states, such that the energy λ is defined as an analytical function of the complex variable z . This additional input of spectral topological concepts is a unique feature in non-Hermitian systems. In this context, the zero-energy points at $z = \pm i$ in (5.41) can arise as branch points in the k -space eigenvalue




Figure 3: Family of curves in the complex plane z for $M_1 = 1$ defined by the real energy condition (5.52). For these points λ becomes real $\{\lambda_1 \in \mathbb{R}, \lambda_2 = 0\}$. They are parametrized by $M_2 = 0, \pm 0.05, \pm 0.5, \pm 1, \pm 2$. The unit circle at the center (dashed line) emerges in the limit $\frac{M_2}{M_1} \rightarrow 0$, and so, in this case $|\sigma| \leq \pi$.




Figure 4: The plot λ_1 vs σ for $\frac{M_2}{M_1} = 0.05$ (left) and for $\frac{M_2}{M_1} \rightarrow 0$ (right) defined by (5.51). λ_1 is not defined on the intervals $|\sigma| \leq \delta$ and $\{-\pi \leq \sigma \leq -\pi + \delta\} \cup \{\pi - \delta \leq \sigma \leq \pi\}$ ($\delta \approx 0.05$) of the left figure.

manifold as can be written from (5.14)

$$\lambda = \pm\sqrt{M^2 - k^2}. \quad (5.53)$$

As we will see below, the branch points ($k = \pm M$) exhibit exotic topological phenomena associated with the winding of eigenvalues and eigenvectors. The complex plane defined by the uniformization variable z turns out to be convenient for performing the relevant calculations in order to study the analytical properties of the energy $\lambda(z)$ and wave number $k(z)$ functions, i.e. in order to examine the zero modes and the points corresponding to the edges of the model, which we will discuss below. The analytic function $\lambda(z)$ in (5.41) has two zeros and one pole, located on the imaginary axis at the points $z = \pm i \rightarrow k = \pm M$, each one of first order, and the pole at the origin $z = 0$. Next, let us define the winding number

$$W = \frac{1}{2\pi i} \oint_{\Gamma} \frac{d\lambda}{\lambda} \quad (5.54)$$

$$= Z - P. \quad (5.55)$$

In the second line above we have used the Cauchy's argument principle, such that Z is the number of zeros and P is the number of poles of the function λ inside the contour Γ (see Fig. 1). The Z and P must be counted with their multiplicities and order, respectively. It is interesting to compute the winding number of $\lambda(z)$ in (5.13) as it traverses the contour Γ encircling the zero modes $\lambda = 0$ located at the points $z = \pm i$ and the pole at $z = 0$ in the complex plane z . So, for the function $\lambda(z)$ in (5.41) and using (5.54) one has

$$\begin{aligned} W &= \frac{1}{2\pi i} \oint_{\Gamma} \frac{z^2 - 1}{z(z^2 + 1)} dz \\ &= 1. \end{aligned} \quad (5.56)$$

Then, one can see that the eq. (5.55) is also verified, i.e. $W = 1$, since $Z = 2$ and $P = 1$ are the number of zeros and poles of the function $\lambda(z)$ inside Γ .

6 Continuum of bound states/solitons and extended ELC modes

In this section we discuss the general solution presented above and classify it into two types of solutions: The dual continuum of bound states/solitons and the extended modes embedded into the localized states (ELC). Moreover, we also compute analytically the biorthogonality relationships between the relevant left and right eigenvectors.

6.1 Continuum of bound state modes/solitons

In the Hermitian Toda system coupled to fermion, which has recently been presented in [21], the in-gap and BIC bound states have quantized energies, whereas the free states form a continuum. We have shown above how this principle fails for the non-Hermitian Toda system coupled to fermions, by analyzing the

non-Hermitian Hamiltonian with a complex Toda field and the properties of pseudo-chirality and pseudo-Hermiticity. In fact, the biorthogonal eigenstates, which we may call “continuum bound state modes” (CBM), have spatial envelopes and form a continuum filling the complex energy plane. We have shown that their strong coupling sector host a continuum complex Toda field solitons, such that the fermion eigenstates are localized inside the topological solitons with topological invariant quantities (5.47)-(5.48). In the strong coupling sector the family of complex solitons can be dubbed as continuum of topological solitons (CTS). These type of novel features of a non-Hermitian Hamiltonian, i.e. the spatially localized eigenstates at every value of the energy in the complex energy plane λ , have recently been uncovered also in [29].

The spinor bound states requires the condition $k_1 \neq 0$ in the next relationships

$$k_1 = \pm \frac{\sqrt{2}}{2} \sqrt{\Delta + (\widetilde{M} - \widetilde{\lambda})} \quad (6.1)$$

$$k_2 = \pm \frac{\sqrt{2}}{2} \sqrt{\Delta - (\widetilde{M} - \widetilde{\lambda})} \quad (6.2)$$

$$\Delta \equiv \sqrt{|M - \lambda|^2 |M + \lambda|^2} \quad (6.3)$$

$$\widetilde{M} \equiv M_1^2 - M_2^2, \quad (6.4)$$

$$\widetilde{\lambda} \equiv \lambda_1^2 - \lambda_2^2, \quad (6.5)$$

Notice that the parameters defined above satisfy $\Delta > |\widetilde{M} - \widetilde{\lambda}|$ without any further restriction on the complex plane $k = k_1 + ik_2$ for a fixed value of the complex mass parameter $M = M_1 + iM_2$. So, for each complex eigenvalue $\lambda \in \mathbb{C}$ one has a continuum of complex scalar solutions with wave numbers k_1 and k_2 given in (6.1)-(6.2). As discussed above, a continuum of bound states emerge coupled to a continuous family of topological solitons (see Fig 4.) such that their topological charges depend on the energy eigenvalues as in (5.28)-(5.32). In addition, the zero mode $\lambda = 0$ implies $k = \pm M$ in (6.1)-(6.2).

On the other hand, the complex energy components become

$$\lambda_1 = \pm \frac{\sqrt{2}}{2} \sqrt{\Delta + (\widetilde{M} - \widetilde{k})} \quad (6.6)$$

$$\lambda_2 = \pm \frac{\sqrt{2}}{2} \sqrt{\Delta - (\widetilde{M} - \widetilde{k})} \quad (6.7)$$

$$\Delta \equiv \sqrt{|M - k|^2 |M + k|^2} \quad (6.8)$$

$$= \sqrt{[(k_1 - M_1)^2 + (k_2 - M_2)^2][(k_1 + M_1)^2 + (k_2 + M_2)^2]} \quad (6.9)$$

$$\widetilde{M} \equiv M_1^2 - M_2^2, \quad (6.10)$$

$$\widetilde{k} \equiv k_1^2 - k_2^2. \quad (6.11)$$

From the above equations (6.6)-(6.11) one has the following $M = \pm k \Rightarrow \Delta = 0$ and $\widetilde{M} - \widetilde{k} = 0$, which imply $\lambda = 0$. This is just the zero mode $\lambda = 0$. In the Fig. 5 the left and right plots show the energy surfaces $\lambda_1(k_1, k_2)$ and $\lambda_2(k_1, k_2)$ for $M_1 = 2, M_2 = -2$ in the eqs. (6.6)-(6.7). The middle figure shows the curves for extended wave states in the localized continuum (ELC) states (6.16)-(6.19)(line colors distinguish the left(right) figure surface each one belongs to). The points of the left and right manifolds (excluding




Figure 5: The left and right plots show the energy surfaces $\lambda(k_1, k_2)$ for $M_1 = 2$, $M_2 = -2$. The middle figure shows the curves $\lambda(k_1 = 0, k_2)$ (line colors distinguish the left(right) figure surface each one belongs to) which correspond to the extended waves in the localized continuum (ELC) states (6.16)-(6.19). The points of the left and right manifolds (excluding the curves $\lambda(k_1 = 0, k_2)$ of the middle plot) represent the localized continuum states (5.26)-(5.27) and (5.45)-(5.46). Notice the presence of the points $k = \pm M$ (left and right plots) which correspond to the zero modes $\lambda = 0$.

the curves $\lambda(k_1 = 0, k_2)$ of the middle plot) represent the localized continuum states (5.26)-(5.27) and (5.45)-(5.46). The points $k = \pm M$ (left and right plots) correspond to the zero modes $\lambda = 0$.

Moreover, from the complex dispersion relationship (5.14) one can get the next two equations, considering the real and imaginary parts, respectively

$$k_1^2 - k_2^2 - M_1^2 + M_2^2 + \lambda_1^2 - \lambda_2^2 = 0 \quad (6.12)$$

$$k_1 k_2 - M_1 M_2 + \lambda_1 \lambda_2 = 0. \quad (6.13)$$

These two relationships are equivalent to the next system of equations

$$k_1^2(k_1^2 - M_1^2 + M_2^2 + \lambda_1^2 - \lambda_2^2) - (M_1 M_2 - \lambda_1 \lambda_2)^2 = 0, \quad k_1 \neq 0, \quad (6.14)$$

$$k_2^2(k_2^2 + M_1^2 - M_2^2 - \lambda_1^2 + \lambda_2^2) - (M_1 M_2 - \lambda_1 \lambda_2)^2 = 0, \quad k_2 \neq 0. \quad (6.15)$$

Once the values of M_1 and M_2 are fixed one has that the equations (6.14)-(6.15) define a couple of families of curves in the plane (λ_1, λ_2) parametrized by the variables k_1 and k_2 , respectively. The Fig. 6 shows the spectral curves filling some regions of the complex plane corresponding to the family of curves (6.14)-(6.15). Notice the overlap regions around the points $(M_1, M_2) = (\pm 2, \pm 2)$ (left Fig.) and $(M_1, M_2) = (\pm 2, \mp 2)$ (right Fig.). In the Fig. 7 we plot the family of curves (λ_1, λ_2) , some of which intersect to each other at the origin $\lambda = 0$. So, the point $\lambda = 0$ constitutes a coalescence point in which the zero mode eigenvectors become biorthogonal.




Figure 6: Spectral curves on the plane $\lambda = (\lambda_1, \lambda_2)$ for $(M_1, M_2) = (\pm 2, \pm 2)$ (left) and $(M_1, M_2) = (\pm 2, \mp 2)$ (right). The k_1 and k_2 families correspond to (6.14) and (6.15), respectively. The figures show the curves for the parameter ranges $|k_a| < 1.5$ ($a = 1, 2$). The zero modes are at $\lambda = 0$ ($k_1 = M_1, k_2 = M_2$). The points $\lambda = (M_1, M_2)$ correspond to the threshold points $k = 0$.

The parabolas $\lambda_1 \lambda_2 = M_1 M_2 = \text{const.}$ in the plane (λ_1, λ_2) can be obtained when one of the components k_1 or k_2 vanishes, and correspond to the extended wave states in the localized continuum for $k_1 = 0$, or to a subset of the continuum bound states for $k_2 = 0$.

So, the confluence of non-Hermiticity, nonlinearity and spectral topology generate new phenomena such as the non-Hermitian currents equivalence (2.14), the complex valued topological charges ($\mathcal{Q} \equiv \mathcal{Q}_1 + i\mathcal{Q}_2$) and the presence of the winding numbers (5.56) related to the spectral topology. The model exhibits the three phenomena unique to non-Hermitian systems: the non-vanishing fermion current, nonzero winding number of energy and the presence of the pair of topological solitons in the model. One can argue that the validity of any one of the three properties is the sufficient and necessary condition for the validity of the other two in this non-Hermitian system.

In summary, we have shown that the non-Hermitian Hamiltonian H can allow an uncountably infinite set of bound states, contrary to the results in [21] in which the in-gap and BIC bound states appear with quantized energies. The biorthogonal continuum bound states possess localized envelopes and are related to a continuum of topological complex Toda solitons. Regarding the zero modes, one can also argue that the biorthogonal Majorana zero modes are embedded into a family of continuum of bound states, with properties different from those studied in their Hermitian counterparts [21].

6.2 Extended wave states embedded in the continuum of bound states

Next, we construct the extended wave states localized in the continuum of bound states (ELC). These states can be obtained by setting $k_1 = 0$ into the tau functions (5.8)-(5.10). So, from (5.1) the scalar components




Figure 7: The family of curves (λ_1, λ_2) for $M_1 = -M_2 = 1$ (left) and $M_1 = M_2 = 1$ (right) in the parameter ranges $|k_{1,2}| < 1.5$. Notice the coalescence of some states at $\lambda = 0$.

become

$$\Phi_1 = -\frac{2}{\beta} \arctan \left[\frac{|a| \sin(2k_2x + \alpha_1 - \frac{\delta}{2}) - |b| \sin(2k_2x + \alpha_2 + \frac{\delta}{2}) + |a||b| \sin(\alpha_1 - \alpha_2 - \frac{\delta}{2}) - \sin \frac{\delta}{2}}{|a| \cos(2k_2x + \alpha_1 - \frac{\delta}{2}) + |b| \cos(2k_2x + \alpha_2 + \frac{\delta}{2}) + |a||b| \cos(\alpha_1 - \alpha_2 - \frac{\delta}{2}) + \cos \frac{\delta}{2}} \right], \quad (6.16)$$

$$\Phi_2 = -\frac{1}{\beta} \log \left[|\rho| \frac{1 + |b|^2 + 2|b| \cos(2k_2x + \alpha_2)}{1 + |a|^2 + 2|a| \cos(2k_2x + \alpha_1)} \right]. \quad (6.17)$$

Similarly, inserting (5.8)-(5.10) into (5.2) for $k_1 = 0$ we have the spinor components

$$\xi_A = |c| \frac{|b| \cos(k_2x + \alpha_2 - \zeta_1) + \cos(k_2x + \zeta_1)}{1 + |b|^2 + 2|b| \cos(2k_2x + \alpha_2)} - i|c| \frac{|b| \sin(k_2x + \alpha_2 - \zeta_1) - \sin(k_2x + \zeta_1)}{1 + |b|^2 + 2|b| \cos(2k_2x + \alpha_2)}, \quad (6.18)$$

$$\xi_B = |d| \frac{|a| \cos(k_2x + \alpha_1 - \zeta_2) + \cos(k_2x + \zeta_2)}{1 + |a|^2 + 2|a| \cos(2k_2x + \alpha_1)} - i|d| \frac{|a| \sin(k_2x + \alpha_1 - \zeta_2) - \sin(k_2x + \zeta_2)}{1 + |a|^2 + 2|a| \cos(2k_2x + \alpha_1)}. \quad (6.19)$$

Using the relationships (2.12) from (6.18)-(6.19) one can get $\eta_A = \xi_B$ and $\eta_B = \xi_A$ with the $\xi_{A,B}$'s given above.

The Fig. 8 shows a representative plot of this type of extended states. Notice that the spinor waves exhibit a period $\frac{2\pi}{k_2}$, whereas the scalar waves present a periodic profile with period $\frac{\pi}{k_2}$. The middle plot in Fig. 5 shows the curves $\lambda(k_1 = 0, k_2)$ which correspond to the ELC extended wave states (6.16)-(6.19).

The family of spectral curves for $k_1 = 0$ in (6.14)-(6.15) can be written as

$$k_2^2 M_1^2 - (\lambda_1^2 - M_1^2)(\lambda_2^2 + M_2^2) = 0. \quad (6.20)$$

The Fig. 9 shows this family of curves parametrized by k_2 for $M_1^2 = M_2^2 = 2$. Notice that these curves lie inside the continuum of bound states in the Fig. 6. So, these states constitute the extended waves in the




Figure 8: A representative of the extended wave states in the continuum of bound states. The components of $\Phi = \Phi_1 + i\Phi_2$ and $\xi^T = (\xi_3 + i\xi_4, \xi_1 + i\xi_2)$ for $|a| = 1.5, |b| = 1.5, |c| = 1, |d| = 1.3, \zeta_1 = 1.2, \zeta_2 = 1.5, \alpha_1 = \pi/3, \alpha_2 = \pi/2, k_1 = 0, k_2 = 1, \beta = 1, |\rho| = 1.5, M_1 = -0.05, M_2 = 3.9$. Notice the periods 2π and π for the spinors and scalars, respectively.




Figure 9: The family of curves (λ_1, λ_2) for $k_1 = 0$ and $M_1^2 = M_2^2 = 2$ in the parameter range $|k_2| < 1.5$ which correspond to the extended waves in the localized continuum (ELC) states parametrized by k_2 (6.20). Notice the ELC zero modes $\lambda = 0$ ($k_2 = \pm M_2$).

localized continuum (ELC), in contradistinction to the well known Hermitian bound states in the continuum (BIC), which appear embedded into the continuum of scattering states.

The bound states in the continuum (BIC) are localized states such that spatially and spectrally overlap with a continuum of free states. However, those states do not hybridize with any of the extended or radiating states. It has recently been established that those bound states remain localized even though they coexist with the continuous spectrum of radiating waves that can carry energy away, i.e. for energies above or below the threshold of the Hermitian Toda coupled to fermion [21]. On the other hand, the extended wave states on the localized continuum defined in (6.18)-(6.19) can be regarded as some kind of inverted BIC states due to the non-Hermiticity of the model (2.1). Recently, these novel type of states in a NH system have been dubbed as extended states in a localized continuum (ELC) [30].

6.3 Left and right eigenvectors and biorthogonality relationships

In the NH case there are some types of time-independent inner products [43, 44]. Thus, for the model above one can write: $\eta^T \xi$, $\xi^T \Gamma \xi$, $\eta^T \Gamma \eta$, $\bar{\xi}^T \xi$, $\xi^T \mathcal{G} \xi$, $\bar{\eta}^T \eta$, $\eta^T \mathcal{G} \eta$, $\bar{\xi}^T \mathcal{G}^{-1} \bar{\xi}$, $\bar{\eta}^T \mathcal{G}^{-1} \bar{\eta}$, and the analog products associated to the χ , ϕ , $\hat{\chi}$, $\hat{\phi}$ eigenvectors. In the following, before verifying the biorthogonality property (3.5), let us compute the NH inner product (3.1) between two right spinors of types η and ξ at the point $\lambda = 0$. So, let us consider the inner product

$$(\eta_{-\lambda} | \xi_\lambda) = \int_{-\infty}^{+\infty} [\eta_{-\lambda, A} \xi_{\lambda, A} + \eta_{-\lambda, B} \xi_{\lambda, B}] dx. \quad (6.21)$$

The soliton parameters in (5.11)-(5.13) can be mapped as $\lambda \rightarrow -\lambda$, $k \rightarrow k$ by making the complex variable transformation $z \rightarrow -\frac{1}{z}$; in addition, through this transformation one can also relate the parameters of $\eta_{-\lambda}$ to that of ξ_λ . So, from (6.21) one can write

$$(\eta_{-\lambda} | \xi_\lambda) = \frac{1}{2k} \left(\frac{\log(a'/b)}{a' - b} + \frac{\log(a/b')}{a - b'} \right), \quad (6.22)$$

$$= \pm \frac{M}{k\beta^2} \frac{(1 - z^2)^2}{\tilde{c}cz^2 + \tilde{c}'c'} \frac{1 + z^2}{z} \log \left(\frac{\tilde{c}c}{\tilde{c}'c'} z^2 \right). \quad (6.23)$$

where the parameters a' , b' , c' , \tilde{c}' and a , b , c , \tilde{c} correspond to the spinors $\eta_{-\lambda}$ and ξ_λ , respectively, according to the notation in (5.11)-(5.13). For the zero modes, i.e. $\lambda = 0$ ($z = \pm i$), from (6.23) one has

$$(\eta_0 | \xi_0) = 0. \quad (6.24)$$

In the next steps we will compute the biorthogonality relationship (3.5) for some special cases. So, let us find the left eigenvectors satisfying the equations (2.18)-(2.19) for the relevant eigenvalue $\bar{\lambda}$. Following similar steps as above we consider the tau functions $\bar{\tau}_{0,1}$, $\bar{\tau}_{A,B}$, $\bar{\tau}'_{A,B}$ such that

$$e^{i\beta\Phi} = \bar{\rho} \left(\frac{\bar{\tau}_0}{\bar{\tau}_1} \right)^2, \quad (6.25)$$

$$\bar{\xi}_A = \frac{\bar{\tau}_A}{\bar{\tau}_1}, \quad \bar{\xi}_B = \frac{\bar{\tau}_B}{\bar{\tau}_0}, \quad \bar{\eta}_A = \frac{\bar{\tau}'_A}{\bar{\tau}_0}, \quad \bar{\eta}_B = \frac{\bar{\tau}'_B}{\bar{\tau}_1}, \quad (6.26)$$

with $\bar{\rho}$ being a constant complex parameter.

So, let us assume the following tau functions

$$\bar{\tau}_1 = 1 + \bar{a} e^{2\bar{k}x}, \quad \bar{\tau}_0 = 1 + \bar{b} e^{2\bar{k}x}, \quad (6.27)$$

$$\bar{\tau}_A = \bar{c} e^{\bar{k}x}, \quad \bar{\tau}_B = \bar{d} e^{\bar{k}x}, \quad (6.28)$$

$$\bar{\tau}'_A = \bar{c}' e^{\bar{k}x}, \quad \bar{\tau}'_B = \bar{d}' e^{\bar{k}x}, \quad (6.29)$$

with the complex parameters \bar{a} , \bar{b} , \bar{c} , \bar{d} , \bar{c}' , \bar{d}' and \bar{k} . Next, making use of these tau functions and substituting the expressions (6.25)-(6.26) into the eigenvalue equations for the left eigenvectors (2.18) with eigenvalue λ , one can show that the equations are satisfied provided that the following parameter relationships hold

$$\bar{a} = \frac{\bar{c}'\bar{c}}{2M} \frac{\beta^2 w^3}{(w^2 - 1)^2}, \quad \bar{d}' = \frac{\bar{c}'\bar{c}}{\bar{d}}, \quad \bar{\rho} = i \frac{\bar{c}}{\bar{d}} w, \quad (6.30)$$

$$\bar{b} = \frac{\bar{c}'\bar{c}}{2M} \frac{\beta^2 w}{(w^2 - 1)^2}, \quad w \equiv -\frac{\bar{\lambda}}{M} \pm \sqrt{\frac{\bar{\lambda}^2}{M^2} - 1} \quad (6.31)$$

$$\bar{\lambda} = -\frac{M}{2}(w + w^{-1}), \quad \bar{k} = \frac{iM}{2}(w - w^{-1}). \quad (6.32)$$

Notice that the form of the relationship (5.14) also holds in this case, i.e. $\bar{\lambda}^2 + \bar{k} = M^2$, and the $\bar{\lambda}$ and \bar{k} parameters depend on the uniformization parameter w in the same form as in the right eigenvalue case (5.13).

Next, let us compute $(\bar{\xi}_{\bar{\lambda}}|\xi_{\lambda})$ for $\bar{\lambda} = \pm\lambda$ by using the expressions (5.2) and (6.26) and the relevant tau functions (5.8)-(5.10) and (6.27)-(6.29), respectively. First, let us consider $\bar{\lambda} = \lambda (\lambda \neq 0)$. The transformation $w = \frac{1}{z}$ maps $\bar{\lambda}(w) \rightarrow \lambda(z)$ and $\bar{k} \rightarrow -k$. So, let us compute the next integral in x for $\bar{k} = -k$

$$\begin{aligned} (\bar{\xi}_{\lambda}|\xi_{\lambda}) &= \int_{-\infty}^{\infty} [\bar{\xi}_{\lambda,A}\xi_{\lambda,A} + \bar{\xi}_{\lambda,B}\xi_{\lambda,B}] dx \\ &= \int_{-\infty}^{\infty} [\eta_{-\lambda,A}\xi_{\lambda,A} - \eta_{-\lambda,B}\xi_{\lambda,B}] dx \\ &= \frac{1}{2k} \left(\frac{\bar{c}c \log(\bar{a}b)}{\bar{a}b - 1} + \frac{\bar{d}d \log(\bar{b}a)}{\bar{a}b - 1} \right) \\ &= 1. \end{aligned} \quad (6.33)$$

Then, in order to satisfy the biorthogonality relationship (3.5) the parameters must satisfy the normalization condition (6.33).

Next, let us compute (3.5) for the special case $(\bar{\xi}_{-\lambda}|\xi_{\lambda}) = 0$. Notice that the transformation $w = -\frac{1}{z}$ maps $\bar{\lambda}(w) \rightarrow -\lambda(z)$ and $\bar{k} \rightarrow k$. So, one has $\bar{\lambda} = -\lambda$. Let us compute the integral in x for $\bar{k} = k$

$$(\bar{\xi}_{-\lambda}|\xi_{\lambda}) = \int_{-\infty}^{\infty} [\bar{\xi}_{-\lambda,A}\xi_{\lambda,A} + \bar{\xi}_{-\lambda,B}\xi_{\lambda,B}] dx \quad (6.34)$$

$$= \frac{1}{2k} \left(\frac{\bar{c}c \log(\bar{a}/b)}{\bar{a} - b} + \frac{\bar{d}d \log(\bar{b}/a)}{\bar{b} - a} \right) \quad (6.35)$$

In order to show the vanishing of the expression above let us impose the pseudo-chirality relationship (3.9) to the left and right eigenvectors, i.e. $\bar{\xi}_{-\lambda,A} = \xi_{\lambda,B}$, $\bar{\xi}_{-\lambda,B} = -\xi_{\lambda,A}$. So, from the the expressions (5.2) and (6.26) and the relevant tau functions (5.8)-(5.10) and (6.27)-(6.29), respectively, one has the relationships

$$\bar{c} = d, \quad \bar{a} = a, \quad \bar{d} = -c, \quad \bar{b} = b. \quad (6.36)$$

Therefore, making use of the relationships (6.36) into (6.35) one can show that

$$(\bar{\xi}_{-\lambda}|\xi_{\lambda}) = 0. \quad (6.37)$$

The zero mode biorthogonality equation (3.12) follows by setting $\lambda = 0$ in the last equation (6.37), i.e. $(\bar{\xi}_0|\xi_0) = 0$. Notice that for the general case of arbitrary states with λ_i and λ_j ($\lambda_i \neq \lambda_j \rightarrow k_i \neq k_j$), the x -integration $(\bar{\xi}_{\lambda_j}|\xi_{\lambda_j}) = \int_{-\infty}^{+\infty} \bar{\xi}_{\lambda_j}^T \xi_{\lambda_j} dx$, in general, will not provide an exact integral, and one could perform the x -integration using numerical methods. A similar procedure as above can be performed in the case of η and $\bar{\eta}$ eigenvectors and show $(\bar{\eta}_{\lambda_i}|\eta_{\lambda_j}) = \delta_{ij}$, except at $\lambda = 0$.

7 Biorthogonal Majorana zero mode-topological soliton duality

Firstly, we find the zero mode *right* eigenvectors. So, let us set $\lambda = 0$ into the system of equations (5.3)-(5.6) or equivalently one can set $z = \pm i$ into the set of parameter equations (5.11)-(5.12). So, the tau functions (5.8)-(5.10) satisfy the equations provided that

$$k = \pm M, \quad b = -a, \quad \rho = \pm \frac{\tilde{c}}{d}, \quad \tilde{d} = \frac{c\tilde{c}}{d}. \quad (7.1)$$

and the charge densities equivalence equation (2.14) for

$$a = \pm i \frac{c\tilde{c}\beta^2}{8M}, \quad (7.2)$$

without any further restriction. Therefore, the one-soliton solution satisfy the condition (2.14) for the equivalence between the spinor bilinear $\eta^T \xi$ and the x -derivative of the complex field Φ . An analog relationship in the one-soliton sector has been defined as the equivalence between Noether and topological currents in the Hermitian version of the model [22, 37, 21].

So, from (5.37)-(5.38) and using (5.30) and (5.32), one can compute the relevant topological charges associated to the complex scalar by setting $z = \pm i$ in the next equations

$$\mathcal{Q}_1 \equiv \frac{\beta}{2\pi} [\Phi_1(+\infty) - \Phi_1(-\infty)] \quad (7.3)$$

$$= \frac{2}{\pi} \arg z \quad (7.4)$$

$$= \pm 1 \quad (7.5)$$

$$\mathcal{Q}_2 \equiv \frac{1}{2\pi} [\Phi_2(+\infty) - \Phi_2(-\infty)]. \quad (7.6)$$

$$= \frac{2}{\pi} \log |z| \quad (7.7)$$

$$= 0. \quad (7.8)$$

The above soliton and bound state solutions of the model, provided the parameters (7.1)-(7.2) relationships are satisfied, allow us to construct the following mappings between the spinor components and the complex scalar field

$$\xi_{0A} = \pm \frac{c}{2\sqrt{a}} e^{-i\beta\Phi/2} (\rho - e^{i\beta\Phi})^{1/2} \quad (7.9)$$

$$\xi_{0B} = \frac{d}{2\sqrt{a}\rho} (\rho - e^{i\beta\Phi})^{1/2} \quad (7.10)$$

$$\eta_{0A} = \frac{d}{2\sqrt{a}\rho} (\rho - e^{i\beta\Phi})^{1/2} \quad (7.11)$$

$$\eta_{0B} = \pm \frac{c}{2\sqrt{a}} e^{-i\beta\Phi/2} (\rho - e^{i\beta\Phi})^{1/2}. \quad (7.12)$$

Notice that these fields satisfy the relationships (2.12). In the Fig. 10 we show the plots of ξ_0 Majorana zero mode components and the associated scalar fields. The field Φ_1 is a kink-type soliton with topological charge $\mathcal{Q}_1 = 1$, whereas the excitation Φ_2 develops a trivial topological charge $\mathcal{Q}_2 = 0$. The second spinor




Figure 10: Majorana zero mode bound state ξ_0 and the associated scalar fields for $|a| = |b| = 1$, $|c| = |d| = 1$, $\zeta_1 = -\zeta_2 = 1.2$, $\alpha_1 = \pi + 1$, $\alpha_2 = 1$, $k_1 = 1$, $k_2 = -1$, $\beta = 1$. The field Φ_1 is a soliton with $\mathcal{Q}_1 = 1$, whereas Φ_2 possesses zero topological charge.

η_0 can be plotted using $\eta_{0A} = \xi_{0B}$, $\eta_{0B} = \xi_{0A}$. One can see that this non-Hermitian Majorana zero mode real components ξ_j , $j = 1, 2, 3, 4$ are independent from each other as compared to its Hermitian counterpart reported in [21], in which the four real components reduce to two independent fields due to the particle-hole symmetry, or, a charge conjugation symmetry which holds in the relativistic Hermitian Toda model coupled to one Dirac fermion.

Next, using the mapping (7.9)-(7.12) into the equations of motion (2.13) and defining $\rho \equiv e^{\rho_o} e^{i\alpha_o}$ with ρ_o and α_o being real parameters, one can write the next equation for the complex scalar

$$-\partial_x^2 \Phi + \frac{4M^2}{\beta} \sin(\beta\Phi - \alpha_o + i\rho_o) = 0. \quad (7.13)$$

Likewise, using the mapping (7.9)-(7.12) into the system of equations of motion (2.7)-(2.8) with $\lambda = 0$ one can get the following equations for the zero mode components

$$i\xi'_{0A} + i\frac{M}{\rho} \xi_{0B} + \frac{1}{2}\beta^2 \xi_{0A} \xi_{0B} \eta_{0B} = 0, \quad (7.14)$$

$$-i\xi'_{0B} - iM\rho \xi_{0A} + \frac{1}{2}\beta^2 \xi_{0A} \xi_{0B} \eta_{0A} = 0, \quad (7.15)$$

$$-i\eta'_{0A} - iM\rho \eta_{0B} + \frac{1}{2}\beta^2 \eta_{0A} \eta_{0B} \xi_{0B} = 0, \quad (7.16)$$

$$i\eta'_{0B} + i\frac{M}{\rho} \eta_{0A} + \frac{1}{2}\beta^2 \eta_{0A} \eta_{0B} \xi_{0A} = 0. \quad (7.17)$$

The model (7.13) can be considered as a NH sine-Gordon model describing the dynamics of the complex scalar field Φ , whereas the system (7.14)-(7.17) as the NH Thirring model describing the biorthogonal Majorana fermions. Notice that the coupling parameter β^2 appears in the three-fermion coupling (or four-fermion term in the Lagrangian description) of the system (7.14)-(7.17), whereas in the Lagrangian description of the NHSG model (7.13) it would appear as $\frac{1}{\beta^2}$ factor of the $\cos(\beta\Phi - \alpha_o + i\rho_o)$ term of the relevant potential. So, following the ideas of [21], the mapping (7.9)-(7.12) can be regarded as the strong \rightleftharpoons weak coupling mappings of the dual sectors of the NH Toda model coupled to Majorana fermions. The Hermitian version of the model (7.14)-(7.17) has recently been considered for two independent spinors and shown to be integrable in the bosonic and Grassmanian versions, respectively [49, 50]. A Hermitian (generalized) massive

Thirring/sine-Gordon duality has also been discussed in [51, 52]. Non-Hermitian \mathcal{PT} -symmetric versions of the massive Thirring model and the sine-Gordon model have recently been considered, and some aspects of their duality relationships in the NH context were examined [53, 54].

Next, for completeness we provide the zero mode *left* eigenvectors. So, one can set $\lambda = 0$ into the system of equations (2.18)-(2.19). Since we have already found above the left eigenvectors for the general λ eigenvalue following the tau functions formalism (6.25)-(6.26) and (6.27)-(6.29), we simply set $w = \pm i$ ($\lambda = 0$) for the zero modes into the relevant parameter relationships (6.30)-(6.32). So, one has

$$\bar{k} = \pm M, \quad \bar{b} = -\bar{a}, \quad \bar{\rho} = \pm \frac{\bar{c}'}{d'}, \quad \bar{d}' = \frac{\bar{c}\bar{c}'}{d}, \quad (7.18)$$

$$\bar{a}' = \pm i \frac{\bar{c}'\bar{c}\beta^2}{8M}. \quad (7.19)$$

Topological superconductors can host Majorana zero modes and are currently intensively pursued as a promising candidate for topological quantum computation, but their experimental realization has so far been inconclusive. It has recently been shown, using non-Hermitian exceptional points, that non-Hermiticity can strongly enhance topological superconductivity, implying more robust Majorana zero modes [55].

8 Discussions and conclusions

We have studied the NH Toda model coupled to Dirac fields (2.1) by examining the pseudo-chiral and (anti)-pseudo-Hermitian symmetries of the model and their relevant representations. The pseudo-chiral symmetry has recently been introduced as a manifestation of Noether's theorem in NH systems, such that a new inner product was defined different from the one in standard quantum mechanics [5]. We have discussed the properties of the right and left eigenvectors and the related eigenvalues, as well as the role played by the complex scalar field components. Among the key characteristics of the model are the biorthogonality relationship between the left and right eigenvectors, and the fact that the complex eigenvalues come in pairs of type $\{\lambda, -\lambda\}$ and $\{\lambda, -\lambda^*\}$. The Table 1 summarizes the main results related to these symmetries and their relevant representations, respectively, for the set of Hamiltonians $\{H, \tilde{H}\}$ and $\{H^*, \tilde{H}^*\}$. This Table describes each type of symmetry and the corresponding eigenvectors and their mutual relationships, as well as the spectral parameter λ properties allowed by the relevant symmetry. The biorthogonality relationships and the presence of the Majorana zero modes at $\lambda = 0$ and the coalescence of the eigenvectors are also indicated. Moreover, we present the interplay of the \mathcal{P} , \mathcal{T} and \mathcal{PT} symmetry transformations with the pseudo-chiral and anti-pseudo-Hermiticity properties, which imply a symmetry of the Hamiltonian for each set of parities (ν_1, ν_2) which can take the scalar fields Φ_1 and Φ_2 , respectively. In Table 2 we summarize the main results of the cases *A*, *B*, *C* and *D* discussed such that the last column of this Table indicates their relevant symmetries.

Some relationships between the complex scalar fields Φ and $\bar{\Phi}$ and the biorthogonal spinor sectors $\{\xi, \eta\}$ and $\{\bar{\xi}, \bar{\eta}\}$ of the model are furnished by the correspondences between the spinors and topological currents

densities in (2.14) and (2.19), respectively, together with the equations (3.23) and (3.24) provided by the pseudo-chirality symmetry. The emergence of the fields $\bar{\Phi}$ and $\{\bar{\xi}, \bar{\eta}\}$ is a feature which characterizes the non-Hermitian Toda model coupled to Dirac spinors. We have shown that the solutions (5.1) and (5.2) with the tau functions in (5.8)-(5.10) satisfy the system of equations (2.7)-(2.8) which coupled the spinors and the scalar field and the currents equivalence equation (2.14). In the soliton sector, the x -integral in the whole real line of the current density (5.47) provides a complex number, which are proportional to the topological charges \mathcal{Q}_1 and \mathcal{Q}_2 , respectively. In fact, the real part of (5.47) is proportional to \mathcal{Q}_1 , whereas the imaginary part to \mathcal{Q}_2 as in (5.39)-(5.40), respectively. So, the above NH feature is the analog of the equivalence between the $U(1)$ Noether and real topological currents in the Hermitian version of the model [21].

The construction of the analytical biorthogonal spinor bound states and their associated complex scalar topological solitons were performed. Analytical solutions of the static version of the system were obtained by using the tau function approach. The complex eigenvalue λ , in each topological sector, satisfies a second order algebraic equation as the dispersion relation (5.14), $\lambda^2 = M^2 - k^2$, plus its complex conjugate. So, a related uniformization parameter z defined in (5.13) has played a key role in understanding the asymptotic behavior of the scalar field. The solutions exhibit some topological charges associated to the asymptotic values of the complex scalar field (5.35), i.e. $\Phi(+\infty) - \Phi(-\infty) = \pm \frac{4}{i\beta} \log z$, with $z \in \mathbb{C} \setminus \{0\}$ being the uniformization parameter. The biorthogonal spinor bound states are found for each topological configurations of the background scalar fields. In these developments the solitons shapes depend crucially on the spinor bound state complex energy and wave number parameters, λ and k , i.e. the soliton shape depends on the fermionic state to which it is coupled. So, it is worth mentioning that the back-reactions of the spinors on the solitons are described nonperturbatively and exactly through our analytical methods.

The energy λ has been defined as an analytical function of the complex variable z (5.13). The spectral topology concept is a unique feature in non-Hermitian systems. In this scenario, the zero-energy points at $z = \pm i$ in (5.41) arise as branch points in the k -space eigenvalue manifold (see the points at $k = \pm M$ in the left and right plots of the Fig. 5). The winding number of $\lambda(z)$ has been computed as it traverses the contour Γ encircling the zero modes $\lambda = 0$ located at the points $z = \pm i$ and the pole at $z = 0$ in the complex plane z (see Fig. 1). The contour integral of the function $\lambda(z)$ using (5.54) provides the winding number $W = 1$ in (5.56), and so, it is in accordance with the Cauchy's argument principle (5.55).

The general solution (5.1) and (5.2) with the tau functions in (5.8)-(5.10) can be classified into two types of solutions:

1. The continuum of bound states/solitons (5.45)-(5.46) and (5.26)-(5.27), respectively. For the wave number with non-vanishing real component, i.e. $k_1 \neq 0$, we have defined the topological invariants for each component of the scalar field (5.28)-(5.31). So, the spinor solutions (5.45)-(5.46) become spatially localized eigenstates at every value of the energy in the complex plane λ which satisfies the dispersion relation (5.14) provided that $\Re(k) \neq 0$. So, this feature violates the general wisdom that bound states should be quantized. In section 6 we have discussed in more detail this important class of solutions of the non-Hermitian Toda

model coupled to fermions, and presented the Fig. 2 as a representative plot. Recently, this phenomenon has also been reported in the context of NH lattice models [29]. The charges $\Re(Q_{topol}) = \pm 1$, $\Im(Q_{topol}) = 0$ are special in the sense that the relevant solitons host the Majorana zero-modes, as we will discuss below.

2. Another class of solutions are the extended waves embedded into the localized states (ELC). These states hold provided that one sets $k_1 = 0$ into the tau functions (5.8)-(5.10). The scalar fields take the form (6.16)-(6.17) and the extended wave spinors become (6.18)-(6.19). The Fig. 8 shows a representative plot of this type of state. The middle plot in Fig. 5 shows the curves $\lambda(k_1 = 0, k_2)$ corresponding to the ELC states. These states can be regarded as a generic NH phenomenon representing some kind of inverted BIC states due to the non-Hermiticity of the model (2.1). Recently, these novel type of ELC states in a NH system have been reported in the context of an active mechanical lattice [30].

The above type of solutions may further be classified into subsets according to the \mathcal{P} , \mathcal{T} and \mathcal{PT} symmetries of the Hamiltonians H and \tilde{H} for each set of parities (ν_1, ν_2) of the scalar field components Φ_1 and Φ_2 , respectively. In this context, the representations of the symmetries in the last column of the Table 2 presented in section 4.3 deserve to be performed. We are currently investigating these and other important issues through analytical and numerical methods.

In the subsection 5.2 we have shown that the non-Hermitian model (2.1) exhibits real energy spectra in a special parameter subspace. The regions of real and complex eigenvalues, and the points of eigenvector degeneracies, have been identified. It has also been shown that the real eigenvalues come in pairs $\pm\lambda_1$, so preserving the pseudo-chiral symmetry. These issues deserve careful examinations in order to see the anti-pseudo-Hermitian symmetry breaking; e.g. as recently developed in [48].

In order to understand the NH soliton-particle duality we have examined the zero mode bound states and corresponding scalar field solutions in section 7. In contradistinction to the Hermitian case in [21] the NH Majorana zero mode components ξ_j , $j = 1, 2, 3, 4$ are independent from each other, and they are hosted by the complex scalar field supporting a topological soliton in its real component and a wave with vanishing topological charge in its imaginary component (see Fig. 10). Then, one can argue that these non-Hermitian zero modes are stable even in the presence of the imaginary component scalar wave with vanishing topological charge. We uncovered the mapping (7.9)-(7.12) which can be regarded as a strong \rightleftharpoons weak duality mappings of the dual sectors of the NH Toda model coupled to Majorana fermions. In this picture the strong coupling sector is described by a NH sine-Gordon model (7.13), whereas the weak coupling sector by the NH Thirring model (7.14)-(7.17) of biorthogonal Majorana fermions. In future works it would be interesting to consider the NH (generalized) massive Thirring/sine-Gordon duality extension of the relevant Hermitian models [51, 52].

Some analytical and numerical extensions of this work deserve to be considered, such as the NH kink-antikink scattering in order to understand the behavior of the NH fermions and hosted bound states, as well as the extensions for more than one species of chiral fermions and scalars in the scalar-fermion interaction term. In particular, a scalar potential term can be introduced into the Lagrangian in order to study the NH

version of the model studied in [21]. So, our work presented the pseudo-chiral and (anti)-pseudo-Hermitian symmetries and their representations from the field theory perspective which were mainly reported before in the context of NH lattice models. We hope our approach will be relevant to the understanding of the physical origins of non-Hermiticity, the finite lifetimes of resonances, back reaction of fermions on solitons and the interplay of nonlinearity, topology, and integrability in the context of NH integrable field theory models.

Finally, it would be interesting to study the dynamics of kinks and bound state solutions of the NH model (2.1) in the context of quasi-integrability, and the relevant quasi-conserved quantities in the so-called (quasi-)integrable approach [56]. So far, to our knowledge, a NH model possessing a strong-weak duality property had not been studied in the quasi-integrability context. Besides, our findings could be relevant in the context of one-dimensional topological superconductors in order to understand how the non-hermiticity influence the shape and the lifetime of the bound states, in particular the Majorana zero modes. It will be an interesting issue to study the robustness against decoherence and disorder of our non-Hermitian Majorana zero modes, as compared to their Hermitian counterparts. These type of non-Hermitian modes, being more stable in the presence of imperfections, make them appealing for quantum information processing and fault-tolerant quantum computing. These and other extensions will be reported elsewhere.

Acknowledgements

The author acknowledges the anonymous referee for relevant comments and suggestions in order to improve the manuscript. He is also grateful to J.J. Monsalve and J. R. V. Pereira for useful discussions.

A Equations in components

Taking $\lambda = \lambda_1 + i\lambda_2$, $\Phi = \Phi_1 + i\Phi_2$, $M = M_1 + iM_2$ and

$$\psi = e^{-i\lambda t}\xi, \quad \xi = \begin{pmatrix} \xi_3 + i\xi_4 \\ \xi_1 + i\xi_2 \end{pmatrix}, \quad (\text{A.1})$$

$$\chi = e^{i\lambda t}\eta, \quad \eta = \begin{pmatrix} \eta_3 + i\eta_4 \\ \eta_1 + i\eta_2 \end{pmatrix}, \quad (\text{A.2})$$

one has the next equations for the spinor components

$$\xi'_1 + (\lambda_1\xi_2 + \lambda_2\xi_1) + e^{-\beta\Phi_2}[(M_1\xi_3 - M_2\xi_4) \cos(\beta\Phi_1) - (M_2\xi_3 + M_1\xi_4) \sin(\beta\Phi_1)] = 0, \quad (\text{A.3})$$

$$\xi'_2 - (\lambda_1\xi_1 - \lambda_2\xi_2) + e^{-\beta\Phi_2}[(M_2\xi_3 + M_1\xi_4) \cos(\beta\Phi_1) + (M_1\xi_3 - M_2\xi_4) \sin(\beta\Phi_1)] = 0, \quad (\text{A.4})$$

$$\xi'_3 - (\lambda_1\xi_4 + \lambda_2\xi_3) + e^{\beta\Phi_2}[(M_1\xi_1 - M_2\xi_2) \cos(\beta\Phi_1) + (M_2\xi_1 + M_1\xi_2) \sin(\beta\Phi_1)] = 0, \quad (\text{A.5})$$

$$\xi'_4 + (\lambda_1\xi_3 - \lambda_2\xi_4) + e^{\beta\Phi_2}[(M_2\xi_1 + M_1\xi_2) \cos(\beta\Phi_1) + (M_2\xi_2 - M_1\xi_1) \sin(\beta\Phi_1)] = 0, \quad (\text{A.6})$$

and the equations for the η'_a 's ($a = 1, 2, 3, 4$) can be obtained from (A.3)-(A.6) by replacing

$$\xi_3 \rightarrow \eta_1, \quad \xi_4 \rightarrow \eta_2, \quad (\text{A.7})$$

$$\xi_1 \rightarrow \eta_3, \quad \xi_2 \rightarrow \eta_4, \quad (\text{A.8})$$

in accordance to the relationship $\xi = \Gamma\eta$ in (2.11). So, let us write

$$\eta'_3 + (\lambda_1\eta_4 + \lambda_2\eta_3) + e^{-\beta\Phi_2}[(M_1\eta_1 - M_2\eta_2) \cos(\beta\Phi_1) - (M_2\eta_1 + M_1\eta_2) \sin(\beta\Phi_1)] = 0, \quad (\text{A.9})$$

$$\eta'_4 - (\lambda_1\eta_3 - \lambda_2\eta_4) + e^{-\beta\Phi_2}[(M_2\eta_1 + M_1\eta_2) \cos(\beta\Phi_1) + (M_1\eta_1 - M_2\eta_2) \sin(\beta\Phi_1)] = 0, \quad (\text{A.10})$$

$$\eta'_1 - (\lambda_1\eta_2 + \lambda_2\eta_1) + e^{\beta\Phi_2}[(M_1\eta_3 - M_2\eta_4) \cos(\beta\Phi_1) + (M_2\eta_3 + M_1\eta_4) \sin(\beta\Phi_1)] = 0, \quad (\text{A.11})$$

$$\eta'_2 + (\lambda_1\eta_1 - \lambda_2\eta_2) + e^{\beta\Phi_2}[(M_2\eta_3 + M_1\eta_4) \cos(\beta\Phi_1) + (M_2\eta_4 - M_1\eta_3) \sin(\beta\Phi_1)] = 0, \quad (\text{A.12})$$

For the scalar field components one has

$$\begin{aligned} & \partial^2\Phi_1 + \beta \cos(\beta\Phi_1) \left[-e^{\beta\Phi_2}(\eta_4(M_2\xi_1 + M_1\xi_2) - \eta_3(M_1\xi_1 - M_2\xi_2)) + \right. \\ & \left. e^{-\beta\Phi_2}(\eta_1(M_1\xi_3 - M_2\xi_4) - \eta_2(M_1\xi_4 + M_2\xi_3)) \right] + \\ & \beta \sin(\beta\Phi_1) \left[e^{\beta\Phi_2}(\eta_3(M_2\xi_1 + M_1\xi_2) + \eta_4(M_1\xi_1 - M_2\xi_2)) - \right. \\ & \left. e^{-\beta\Phi_2}(\eta_1(M_2\xi_3 - M_1\xi_4) + \eta_2(M_2\xi_4 - M_1\xi_3)) \right] \end{aligned} \quad (\text{A.13})$$

and

$$\begin{aligned} & \partial^2\Phi_2 + \beta \cos(\beta\Phi_1) \left[e^{\beta\Phi_2}(\eta_3(M_2\xi_1 + M_1\xi_2) + \eta_4(M_1\xi_1 - M_2\xi_2)) + \right. \\ & \left. e^{-\beta\Phi_2}(\eta_1(M_2\xi_3 + M_1\xi_4) - \eta_2(M_2\xi_4 - M_1\xi_3)) \right] + \\ & \beta \sin(\beta\Phi_1) \left[e^{\beta\Phi_2}(\eta_4(M_2\xi_1 + M_1\xi_2) - \eta_3(M_1\xi_1 - M_2\xi_2)) + \right. \\ & \left. e^{-\beta\Phi_2}(\eta_1(M_1\xi_3 - M_2\xi_4) - \eta_2(M_2\xi_3 + M_1\xi_4)) \right] \end{aligned} \quad (\text{A.14})$$

The relationship (2.14) becomes

$$\Phi'_1 = -\frac{\beta}{2}[\eta_1\xi_1 - \eta_2\xi_2 + \eta_3\xi_3 - \eta_4\xi_4], \quad (\text{A.15})$$

$$\Phi'_2 = -\frac{\beta}{2}[\eta_2\xi_1 + \eta_1\xi_2 + \eta_4\xi_3 + \eta_3\xi_4]. \quad (\text{A.16})$$

For $M_2 \neq 0$, $\Phi_2 = 0$ and $\eta \neq \pm\xi^*$ one has that the eq. (A.13) provides the constraint

$$\tan(\beta\Phi_1) = \frac{\eta_3(M_2\xi_1 + M_1\xi_2) + \eta_4(M_1\xi_1 - M_2\xi_2) + \eta_1(M_2\xi_3 + M_1\xi_4) - \eta_2(M_2\xi_4 - M_1\xi_3)}{\eta_4(M_2\xi_1 + M_1\xi_2) - \eta_3(M_1\xi_1 - M_2\xi_2) + \eta_1(M_1\xi_3 - M_2\xi_4) - \eta_2(M_2\xi_3 + M_1\xi_4)}. \quad (\text{A.17})$$

References

- [1] Y. Ashida, Z. Gong and M. Ueda, *Advances in Physics* **69** (2020) 249.

- [2] E. J. Bergholtz, J. C. Budich and F. K. Kunst, *Rev. Mod. Phys.* **93** (2021) 015005-1.
- [3] K. Kawabata, K. Shiozaki, M. Ueda and M. Sato, *Physical Review X* **9** (2019) 041015.
- [4] E.A. Ivanov and A. V. Smilga, *JHEP* **07** (2007) 036.
- [5] J. D. H. Rivero and L. Ge, *Phys. Rev. Lett.* **125** (2020) 083902.
- [6] A. Mostafazadeh, *J. Math. Phys.* **43** (2002) 205.
- [7] A. Mostafazadeh, *J. Math. Phys.* **43** (2002) 3944.
- [8] A. Mostafazadeh, *J. Math. Phys.* **43** (2002) 2814.
- [9] C. M. Bender, and S. Boettcher, *Phys. Rev. Lett.* **80** (1998) 5243.
- [10] C.M. Bender, *Rep. Prog. Phys.* **70** (2007) 947.
- [11] C. Bender, R. Tateo, P. Dorey, T. Dunning, G. Levai, S. Kuzhel, H. Jones, A. Fring, and D. Hook, *PT-Symmetry in Quantum And Classical Physics* (World Scientific Publishing Company, 2018).
- [12] F. G. Scholtz, H.B. Geyer, and F. J. W. Hahne, *Ann. of Phys.* **213** (1992) 74.
- [13] V. V. Konotop, J. Yang and D. A. Zezyulin, *Rev. Mod. Phys.* **88** (2016) 035002-1.
- [14] N. Okuma and M. Sato, *Annu. Rev. Condens. Matter Phys.* **14** (2023) 83.
- [15] P. Delplace, J.B. Marston and A. Venaille, *Science* **358** (2017) 1075.
- [16] H. Blas, R. A. DeLaCruz-Araujo, N. I. Reynaldo Jr., N. Santos, S. Tech and H.E.P. Cardoso, *Zero mode-soliton duality and pKdV kinks in Boussinesq system for non-linear shallow water waves*, arXiv:2305.04037[hep-th].
- [17] L. Jezequel and P. Delplace, *Phys. Rev.* **B105** (2022) 035410-1.
- [18] R. Jackiw and C. Rebbi, *Phys. Rev.* **D13** (1976) 3398.
- [19] J. Goldstone and F. Wilzeck, *Phys. Rev. Lett.* **47** (1981) 986.
- [20] R. MacKenzie and F. Wilczek, *Phys. Rev.* **D30** (1984) 2194.
- [21] H. Blas, J.J. Monsalve, R. Quicaño and J. R. V. Pereira, *JHEP* **09** (2022) 082.
- [22] H. Blas and L.A. Ferreira, *Nucl. Phys.* **B571** (2000) 607.
- [23] H. Blas, *Nucl. Phys.* **B596** (2001) 471.
- [24] F. K. Kunst, E. Edvardsson, J. C. Budich and E. J. Bergholtz, *Phys. Rev. Lett.* **121** (2018) 026808.

- [25] A. Udupa, A. Banerjee, K. Sengupta and D. Sen, *J. Phys.: Condens. Matter* **33** (2021) 145301.
- [26] C. Spanslatt, E. Ardonne, J. C. Budich and T. H. Hansson, *J. Phys.: Condens. Matter* **27** (2015) 405701.
- [27] R. Eneias and A. Ferraz, *New J. Phys.* **17** (2015) 123006.
- [28] H. Blas, *Phys. Rev.* **D66** (2002) 127701.
- [29] Q. Wang, C. Zhu, X. Zheng, H. Xue, B. Zhang and Y. D. Chong, *Phys. Rev. Lett.* **130** (2023) 103602.
- [30] W. Wang, X. Wang and G. Ma, *Phys. Rev. Lett.* **129** (2023) 264301.
- [31] T. Hollowood, *Nucl. Phys.* **B384** (1992) 523.
- [32] J. Cen, and A. Fring, *J. Physics* **A49** (2016) 365202.
- [33] Z. Y. Ge , Y.R. Zhang, T. Liu, S.-W. Li, H. Fan, and F. Nori, *Phys. Rev.* **B100** (2019) 054105.
- [34] C.-G. Oh, S.-H Han, and S. Cheon, *Sci Rep* **11** (2021) 21652.
- [35] D. Sticlet, B. Dóra , and C.P. Moca, *Phys. Rev. Lett.* **128** (2022) 016802
- [36] L.A. Ferreira, J-L. Gervais, J.Sánchez Guillen and M.V.Saveliev, *Nucl. Phys.* **B470** (1996) 236-288.
- [37] H. Blas and B.M. Pimentel. *Ann. of Phys.* **282** (2000) 67.
- [38] S. Franca, V. Konye, F. Hassler, J. van den Brink, and C. Fulga, *Phys. Rev. Lett.* **129** (2022) 086601.
- [39] S.-J. Chang, S.D. Ellis and B.W. Lee, *Phys. Rev.* **D11** (1975) 3572.
- [40] E. Witten, *Nucl. Phys.* **B145** (1978) 110.
- [41] D. A. Takahashi and M. Nitta, *Phys. Rev. Lett.* **110** (2013) 131601.
- [42] M. Stone (Ed.) *Bosonization*. (1994). Singapore: World Scientific.
- [43] M. Sato, K. Hasebe, K. Esaki and M. Kohmoto, *Prog. Theor. Phys.* **127** (2012) 937.
- [44] A. Ghatak and T. Das, *J. Phys.: Condens. Matter* **31** (2019) 263001.
- [45] Z.-H. Wang, F. Xu, L. Li, D.-H. Xu, W.-Q. Chen and B. Wang, *Phys. Rev.* **B103** (2021) 134507
- [46] A. Mostafazadeh, *J. Math. Phys.* **44** (2003) 974.
- [47] Y. Long , H. Xue , and B. Zhang, *Phys. Rev.* **B105** (2022) L100102.
- [48] A. Melkani, *Physical Review Research* **5** (2023) 023035.

- [49] B. Basu-Mallick, D. Sinha, *Integrable coupled bosonic massive Thirring model and its nonlocal reductions*, arXiv:2307.00351 [nlin.SI].
- [50] B. Basu-Mallick, F. Finkel, A. González-López and D. Sinha, *JHEP* **11** (2023) 018.
- [51] H. Blas, *JHEP* **11** (2003) 054.
- [52] H. Blas and H.L. Carrion. *JHEP* **0701** (2007) 027.
- [53] B. Liégeois, C. Ramasubramanian, and N. Defenu, *Tunable tachyon mass in the PT-broken massive Thirring model*, arXiv:2212.08110 [hep-th].
- [54] Y. Ashida, S. Furukawa and M. Ueda, *Nat Commun* **8** (2017) 15791.
- [55] R. Arouca, J. Cayao and A. M. Black-Schaffer, *Phys. Rev.* **B108** (2023) L060506.
- [56] H. Blas, H. F. Callisaya and J.P.R. Campos, *Nucl. Phys.* **B950** (2020) 114852.