# S-move matrix in the NS sector of N = 1 super Liouville field theory

Elena Apresyan<sup>1,2\*</sup> and Gor Sarkissian<sup>1,2</sup>  $\dagger$ 

<sup>1</sup>Yerevan Physics Institute, Alikhanian Br. 2, 0036 Yerevan, Armenia

and

<sup>2</sup> Bogoliubov Laboratory of Theoretical Physics, JINR, Dubna, Moscow region, 141980 Russia

#### Abstract

In this paper we calculate matrix of modular transformations of the onepoint toric conformal blocks in the Neveu-Schwarz sector of N = 1 super Liouville field theory. For this purpose we use explicit expression for this matrix as integral of product of certain elements of fusion matrix. This integral is computed using the chain of integral identities for supersymmetric hyperbolic gamma functions derived by the degeneration of the integrals of parafermionic elliptic gamma functions.

elena-apresyan@mail.ru

<sup>&</sup>lt;sup>†</sup>sarkissn@theor.jinr.ru, gor.sargsyan@yerphi.am

# Contents

1	Introduction	3	
<b>2</b>	Basics on $N = 1$ super Liouville field theory	8	
3	Elements of S-matrix for three NS primaries	12	
4	Elements of S-matrix for two NS and one $\tilde{N}S$ primaries	16	
5	Conclusion	19	
$\mathbf{A}$	Properties of supersymmetric hyperbolic and double gamma	functions	21
в	Parafermionic integral identities I	22	
$\mathbf{C}$	Parafermionic integral identities II	<b>24</b>	
D	Parafermionic integral identities III	25	
E	Paraformionic integral identities IV	26	

#### 1 Introduction

One of the basic objects in two-dimensional conformal field theory is the *S*-matrix of modular transformation of one-point toric conformal blocks.

One can write the one-point correlation function on a torus with modular parameter  $\tau$  in the following form:

$$\langle \phi_{\lambda,\bar{\lambda}} \rangle = \sum_{(\gamma,\bar{\gamma})} \mathcal{F}^{\lambda}_{\gamma}(q) \, \bar{\mathcal{F}}^{\bar{\lambda}}_{\bar{\gamma}}(\bar{q}) \, C^{\lambda,\bar{\lambda}}_{\gamma,\bar{\gamma}} \,\,, \tag{1}$$

where  $q = e^{2\pi i \tau}$ ,  $\mathcal{F}^{\lambda}_{\gamma}(q)$  are one-point conformal toric blocks and  $C^{\lambda,\bar{\lambda}}_{\gamma,\bar{\gamma}}$  are structure constants.

The S-matrix of modular transformation of the one-point toric blocks on a torus is defined by relation

$$\mathcal{F}_{\beta_1}^{\beta_3}(q(\tau)) = (-i\tau)^{-\Delta_{\beta_3}} \sum_{\beta_2} S_{\beta_1\beta_2}(\beta_3) \mathcal{F}_{\beta_2}^{\beta_3}\left(q\left(-\frac{1}{\tau}\right)\right) . \tag{2}$$

The S-matrix is one of the most important notions of 2D conformal field theories. The S-matrix, together with fusion F-matrix, braiding B-matrix and T-matrix, representing the second generator of the modular group  $\tau \rightarrow \tau + 1$ , form the Moore-Seiberg groupoid [34]. These generators satisfy a set of relations called Moore-Seiberg's equations [34]. One of them will be discussed below in detail. The Moore-Seiberg groupoid gives in a sense the full number of operations on the space of conformal block. More abstractly one can consider the space of conformal blocks as a representation of Moore-Seiberg groupoid. For better understanding of the Moore-Seiberg's equations we recall that the space of conformal blocks in rational conformal field theory also arises as Hilbert space of states of 3D topological field theory [22, 54]. It was observed that the Moore-Seiberg's equations are equivalent to three-dimensional topological invariance [33, 54]. In fact, it was proved for rational theories that given a representation of the Moore-Seiberg groupoid, namely a solution to the Moore-Seiberg's equations, one can reconstruct the 3D topological field theory [19].

The Moore-Seiberg groupoid proved to be useful also for non-rational theories.

It was proposed in [53] that the space of conformal block of Liouville theory with its mapping class group representation is isomorphic to the Hilbert space of states of quantized Teichmuller space. Later this proposal was refined in [50,51] and it was shown that quantized Teichmuller space carries unitary projective representation of Moore-Seiberg grupoid equivalent to that of Liouville theory. Using these data it was built in [17,18], in full analogy with rational case, topological field theory based on conformal blocks of the Liouville theory, called Virasoro topological quantum field theory. In these papers this Virasoro topological quantum field theory was related to 3D quantum gravity with negative cosmological constant.

The S-matrix  $S_{\beta_1\beta_2}(\beta_3)$  satisfies the following Moore-Seiberg duality identity [33,34]:

$$S_{\beta_{1}\beta_{2}}(\beta_{3})\sum_{\beta_{4}}F_{\beta_{3},\beta_{4}}\begin{bmatrix}\beta_{2}&\alpha_{1}\\\beta_{2}&\alpha_{2}\end{bmatrix}e^{-2\pi i(\Delta_{\beta_{4}}-\Delta_{\beta_{2}})}F_{\beta_{4},\beta_{5}}\begin{bmatrix}\alpha_{2}&\alpha_{1}\\\beta_{2}&\beta_{2}\end{bmatrix}$$
(3)  
$$=\sum_{\beta_{6}}F_{\beta_{3},\beta_{6}}\begin{bmatrix}\beta_{1}&\alpha_{1}\\\beta_{1}&\alpha_{2}\end{bmatrix}e^{-\pi i(\Delta_{\alpha_{1}}+\Delta_{\alpha_{2}}-\Delta_{\beta_{5}})}F_{\beta_{1},\beta_{5}}\begin{bmatrix}\alpha_{1}&\alpha_{2}\\\beta_{6}&\beta_{6}\end{bmatrix}}S_{\beta_{6}\beta_{2}}(\beta_{5}).$$

Here  $F_{\sigma_5,\sigma_6}\begin{bmatrix} \sigma_2 & \sigma_1 \\ \sigma_3 & \sigma_4 \end{bmatrix}$  is fusion matrix and  $\Delta_{\sigma}$  is conformal weight. Setting in eq. (3)  $\beta_1 = \beta_3 = 0$  one obtains an explicit expression of the *S*-matrix in terms of the fusion matrix and conformal weights [6,33]:

$$S_{0\beta_{2}}\sum_{\beta_{4}}F_{0,\beta_{4}}\begin{bmatrix}\beta_{2}&\alpha\\\beta_{2}&\alpha\end{bmatrix}e^{-2\pi i\Delta_{\beta_{4}}}F_{\beta_{4},\beta_{5}}\begin{bmatrix}\alpha&\alpha\\\beta_{2}&\beta_{2}\end{bmatrix}$$

$$=e^{-\pi i(2\Delta_{\alpha}-\Delta_{\beta_{5}}+2\Delta_{\beta_{2}})}F_{0,\beta_{5}}\begin{bmatrix}\alpha&\alpha\\\alpha&\alpha\end{bmatrix}S_{\alpha\beta_{2}}(\beta_{5})$$
(4)

or this can be written also in the form [6]

$$S_{0\beta_{2}}\sum_{\beta_{4}}F_{0,\beta_{4}}\begin{bmatrix}\beta_{2}&\alpha\\\beta_{2}&\alpha\end{bmatrix}e^{2\pi i\Delta_{\beta_{4}}}F_{\beta_{4},\beta_{5}}\begin{bmatrix}\alpha&\alpha\\\beta_{2}&\beta_{2}\end{bmatrix}$$

$$=e^{\pi i(2\Delta_{\alpha}+2\Delta_{\beta_{2}})}F_{0,\beta_{5}}\begin{bmatrix}\alpha&\alpha\\\alpha&\alpha\end{bmatrix}S_{\alpha\beta_{2}}(\beta_{5}).$$
(5)

Here we used that  $S_{\alpha\beta_2}(\beta_5)$  in the limit  $\beta_5 \to 0$  becomes matrix  $S_{\alpha\beta_2}$  of the modular transformation of characters and  $S_{0\beta_2}$  in (5) is the corresponding element of this matrix. The formula (5) was used in [52] to calculate  $S_{\beta_6\beta_2}(\beta_5)$  in the Liouville field theory. The calculations are very involved and use sequence of integral identities for various products of hyperbolic gamma functions. Important point here is that all the necessary identities can be derived by some chains of the reductions from the elliptic beta integral [45] and symmetry relations of so called elliptic V-function [46, 47]. This becomes possible since as shown in [38, 39] the hyperbolic gamma function is an asymptotic form of the elliptic gamma function.

As a result it was obtained that the  $S^{L}_{\alpha\beta_{2}}(\beta_{5})$ -matrix in Liouville theory is given by the expression [50, 52]:

$$S_{\alpha\beta_2}^L(\beta_5) = \frac{\mathcal{N}(\alpha, \beta_5, \alpha)}{\mathcal{N}(\beta_2, \beta_5, \beta_2)} S_{0\beta_2}^L e^{-\frac{i\pi}{2}\Delta_{\beta_5}^L} S_{\alpha\beta_2}^T(\beta_5) , \qquad (6)$$

where  $\mathcal{N}(\alpha_1, \alpha_2, \alpha_3)$  are some normalization coefficients of chiral vertex operators,  $\Delta_{\alpha}^L = \alpha(Q - \alpha)$  is conformal weight, Q = b + 1/b,

$$S_{0\alpha}^{L} = \frac{1}{\sqrt{2}S_{b}(\pm 2(\alpha - Q/2))} = -2\sqrt{2}\sin\frac{\pi(2\alpha - Q)}{b}\sin\pi(2\alpha - Q)b \quad (7)$$

is the corresponding component of the matrix of modular transformation of characters, and

$$S_{\alpha\beta_{2}}^{T}(\beta_{5}) = \frac{1}{S(\beta_{5})} \int_{-i\infty}^{i\infty} S_{b}(y - Q/2 + \beta_{2} + \beta_{5}/2) S_{b}(y + Q/2 - \beta_{2} + \beta_{5}/2) \quad (8)$$
  
  $\times S_{b}(-y - Q/2 + \beta_{2} + \beta_{5}/2) S_{b}(-y + Q/2 - \beta_{2} + \beta_{5}/2) e^{2i\pi y(2\alpha - Q)} dy.$ 

Here  $S_b(x)$  is hyperbolic gamma function defined in appendix A. For the physical states all the entering parameters belong to the set  $Q/2 + i\mathbb{R}$ .

The function (8) gives the bulk-boundary coupling in LFT as well [29], since the sewing constraints imply that the bulk-boundary coupling proportional to  $S^{L}_{\alpha\beta_{2}}(\beta_{5})$  [6,41].

Using identities (108) and (140) in appendices B and E correspondingly for r = 1 one can show that this function enjoys the following properties

$$S_{\alpha\beta_2}^T(\beta_5) = S_{\beta_2\alpha}^T(Q - \beta_5) \tag{9}$$

and

$$S_{\alpha\beta_{2}}^{T}(Q-\beta_{5}) = S_{b}(-Q+2\alpha+\beta_{5})S_{b}(Q-2\alpha+\beta_{5})$$
(10)  
 
$$\times S_{b}(2Q-\beta_{5}-2\beta_{2})S_{b}(-\beta_{5}+2\beta_{2})S_{\alpha\beta_{2}}^{T}(\beta_{5}).$$

Note that function (8) is nothing else as partition function of mass-deformed N = 4 SQED with two electron hypermultiplets, called T[SU(2)], on squashed three-sphere  $S_b^3$  and eq.(9) expresses its mirror symmetry [27,28]. This coincidence is not accidental but consequence of AGT dualty [3] between  $4D \ N = 2 SU(2)$  supersymmetryc gauge theory on  $S^4$  and Liouville theory in the presence of domain walls/defects [20,28]. It was proposed in [20] that partition function of 3D S-duality domain wall inside  $4D \ N = 2^*$  theory, which is the mentioned mass-deformed T[SU(2)] theory, should coincide with the kernel of S-move transformation of Liouville theory. The actual coincidence was established in [28].

It is possible to show that the invariance of the one-point correlation function (1) under the modular transformation requires S-matrix to satisfy the following Moore-Seiberg equation (see for example [17, 52]):

$$\int d\beta_2 S^L_{\beta_1 \beta_2}(\beta_5) S^L_{\beta_2 \beta_3}(\beta_5) = \delta(\beta_1 - \beta_3) e^{-i\pi \Delta^L_{\beta_5}} \,. \tag{11}$$

It is easy to check that using the relations (9) and (10) the eq.(11) can be written as

$$\int d\beta_2 S_{\beta_2\beta_1}^T(\beta_5) S_{\beta_2\beta_3}^T(\beta_5) S_{0\beta_2}^L S_b(-Q+2\beta_2+\beta_5) S_b(Q-2\beta_2+\beta_5) (12)$$
  
=  $\frac{1}{S_{0\beta_1}^L} \delta(\beta_1-\beta_3) S_b(-Q+2\beta_1+\beta_5) S_b(Q-2\beta_1+\beta_5) .$ 

This property is closely related to the fact that the function  $S_{\alpha\beta_2}^T(\beta_5)$  satisfies the following eigenvalue equation [24,35]

$$H(S_{\alpha\beta_{2}}^{T}(\beta_{5})) = 2\cos \pi b(Q - 2\beta_{2})S_{\alpha\beta_{2}}^{T}(\beta_{5}), \qquad (13)$$

where H is the finite-difference operator:

$$H = \frac{\sin \pi b (2\alpha - Q - \beta_5)}{\sin \pi b (2\alpha - Q)} e^{-\frac{b}{2}\partial_{\alpha}} + \frac{\sin \pi b (2\alpha - Q + \beta_5)}{\sin \pi b (2\alpha - Q)} e^{\frac{b}{2}\partial_{\alpha}}.$$
 (14)

This equation can be derived in the numerous ways. First one can obtain it evaluating the Moore-Seiberg equation (3) for degenerate primaries [35]. It is possible directly to check [24] that  $S_{\alpha\beta_2}^T(\beta_5)$  satisfies (13). And finally one can derive it by certain limiting procedure from the difference equations discussed in [4]. The operator (14) is well known Hamiltonian of the relativistic Calogero-Sutherland model [40], sometimes also called Ruijsenaars-Schneider model, and was extensively studied recently is series of works [11–13]. Now presence of the  $\delta$ -function in the r.h.s. of (12) is easily understood from the observation that the Hamiltonian (14) is hermitian w.r.t. the scalar product given by the l.h.s. of (12). The integral in l.h.s. of (12) is computed in [11–13] and indeed coincides with r.h.s. of (12). We see that the function (8) is on the junction of three theories: Liouviile field theory, T[SU(2)] theory and relativistic Calogero-Sutherland model, and various its properties have different interpretation in each of them.

In this paper we use the formula (5) to calculate S-matrix of modular transformation of one-point toric conformal block in N = 1 super Liouville field theory (SLFT). For this purpose we should elaborate "supersymmetric counterpart" of the mentioned integral identities. At this point we would like to point out that if the essential part of the fusion matrix in the Liouville field theory is integral of product of hyperbolic gamma functions  $S_b(x)$ , defined in appendix A, then the essential part of the fusion matrix in N = 1 super Liouville field theory is sum of integrals of product of so called Neveu-Schwarz and Ramond hyperbolic gamma functions:

$$S_1(y) \equiv S_{\rm NS}(y) \equiv S_b\left(\frac{y}{2}\right) S_b\left(\frac{y}{2} + \frac{Q}{2}\right),\tag{15}$$

$$S_0(y) \equiv S_{\rm R}(y) \equiv S_b\left(\frac{y}{2} + \frac{b}{2}\right) S_b\left(\frac{y}{2} + \frac{b^{-1}}{2}\right) ,$$
 (16)

where Q = b + 1/b. To derive the necessary integral formulas for them first we note that the functions (15) and (16) are particular case of so called parafermionic hyperbolic gamma functions [30, 36, 42], defined for any  $r \in \mathbb{N}$ :

$$\Lambda(y,m) = \prod_{k=0}^{m-1} S_b\left(\frac{y}{r} + b\left(1 - \frac{m}{r}\right) + Q\frac{k}{r}\right)$$
$$\times \prod_{k=0}^{r-m-1} S_b\left(\frac{y}{r} + \frac{m}{br} + Q\frac{k}{r}\right), \qquad (17)$$

where  $0 \le m \le r$ . Obviously, for r = 2 one obtains the functions (15) and (16):

$$\Lambda(y,0) = S_1(y),\tag{18}$$

$$\Lambda(y,1) = S_0(y) \,. \tag{19}$$

In paper [49], rarefied elliptic gamma functions are constructed and corresponding generalizations of the elliptic beta integral and elliptic V-function are studied. It was checked also in [42] that the functions (17) are asymptotic forms of the rarefied elliptic gamma functions. Therefore we can get the necessary identities using the same chains of the reductions as in the case r = 1. Some of the necessary integral identities were derived in this way in papers [4, 42]. Others are obtained in appendices B-E. Our main results are given by formulae (63) and (80). In fact these integrals describe also bulk-boundary coupling in super Liouville theory as we explained above.

In super Liouville theory we find as well that integrals describing matrix of modular transformation reside on the junction of different theories.

First one can see that integral (63) is zero-holonomy value of the partition function of the above mentioned mass-deformed T[SU(2)] theory on squashed lens space  $S_b^3/\mathbb{Z}_2$  found in [36]. This is in agreement with AGT correspondence between N = 2 SU(2) supersymmetric gauge theory on  $S^4/\mathbb{Z}_2$  and super Liouville theory [8–10, 15, 16]. Thus we can consider our results as an evidence for generalization of the AGT correspondence between gauge theory on  $S^4/\mathbb{Z}_2$ and super Liouville theory in the presence of the domain walls. It is important further to study correspondence between holonomies entering in partition function of T[SU(2)] theory on squashed lens space  $S_b^3/\mathbb{Z}_2$  and types of primaries entering in matrix of modular transformation.

The paper is organized as follows. In section 2, we review some basic facts on N = 1 Super Liouville field theory. In section 3, we calculate  $S_{\beta_1\beta_2}(\beta_3)$  in the case when all  $\beta_i$ , i = 1, 2, 3 are NS primaries. The corresponding result is given by formulae (63) and (64). In section 4, we calculate  $S_{\beta_1\beta_2}(\beta_3)$  in the case when  $\beta_i$ , i = 1, 2 are NS primaries and  $\beta_3$  is  $\tilde{N}S$  primary. It is given by (80) and (81). In section 5, we comment our results and outline possible applications. In appendices A-E the necessary special functions, their properties and integral identities are reviewed.

#### **2** Basics on N = 1 super Liouville field theory

N = 1 super Liouville field theory is defined on a two-dimensional surface with metric  $g_{ab}$  by the Lagrangian density:

$$\mathcal{L} = \frac{1}{2\pi} g_{ab} \partial_a \varphi \partial_b \varphi + \frac{1}{2\pi} (\psi \bar{\partial} \psi + \bar{\psi} \partial \bar{\psi}) + 2i\mu b^2 \bar{\psi} \psi e^{b\varphi} + 2\pi \mu^2 b^2 e^{2b\varphi} \,. \tag{20}$$

The energy-momentum tensor and the superconformal current are

$$T = -\frac{1}{2} (\partial \varphi \partial \varphi - Q \partial^2 \varphi + \psi \partial \psi), \qquad (21)$$

$$G = i(\psi \partial \varphi - Q \partial \psi), \qquad (22)$$

where  $Q = b + \frac{1}{b}$ . The superconformal algebra is

$$[L_m, L_n] = (m-n)L_{m+n} + \frac{c_{SL}}{12}m(m^2 - 1)\delta_{m+n}, \qquad (23)$$

$$[L_m, G_k] = \frac{m - 2k}{2} G_{m+k} , \qquad (24)$$

$$\{G_k, G_l\} = 2L_{l+k} + \frac{c}{3}\left(k^2 - \frac{1}{4}\right)\delta_{k+l}, \qquad (25)$$

with the central charge

$$c_{SL} = \frac{3}{2} + 3Q^2 \,. \tag{26}$$

Here k and l take integer values for the Ramond algebra and half-integer values for the Neveu-Schwarz algebra. Since in the Neveu-Schwarz sector of N = 1SLFT one has besides the Virasoro generators  $L_m$ , also supercurrent generators  $G_k$  with half-integer k, descendant fields are broken into two sets of integer and half-integer levels. Thus we will work with primary fields associated with the vertex operators  $N_{\alpha} = e^{\alpha\varphi}$ , which we call NS field, and its supercurrent descendant  $\tilde{N}_{\alpha} = G_{-1/2}N_{\alpha}$ , which we call  $\tilde{N}S$  field [7]. The  $N_{\alpha}$  primary field has conformal dimension

$$\Delta_{\alpha} = \frac{1}{2}\alpha(Q - \alpha).$$
(27)

The physical states have  $\alpha = \frac{Q}{2} + iP$ . The  $\tilde{N}_{\alpha}$  has the conformal dimension

$$\Delta_{\tilde{\alpha}} = \frac{1}{2}\alpha(Q - \alpha) + \frac{1}{2}.$$
(28)

The degenerate states are given by the momenta:

$$\alpha_{m,n} = \frac{1}{2b}(1-m) + \frac{b}{2}(1-n)$$
(29)

with even m-n in the NS sector and odd m-n in the R sector. For future use we need matrix of modular transformation of the NS characters. The corresponding characters for generic P which have no null-states are

$$\chi_P^{NS}(\tau) = \sqrt{\frac{\theta_3(q)}{\eta(q)}} \frac{q^{P^2/2}}{\eta(\tau)}, \qquad (30)$$

where  $q = \exp(2\pi i \tau)$  and

$$\eta(\tau) = q^{1/24} \prod_{n=1}^{\infty} (1 - q^n).$$
(31)

Modular transformation of characters (30) is well-known:

$$\chi_P^{NS}(\tau) = \int \chi_{P'}^{NS}(-1/\tau) e^{-2i\pi PP'} dP' \,. \tag{32}$$

For the degenerate representations, the characters are given by those of the corresponding Verma modules subtracted by those of null submodules:

$$\chi_{m,n}^{NS} = \chi_{\frac{1}{2}(nb+mb^{-1})}^{NS} - \chi_{\frac{1}{2}(nb-mb^{-1})}^{NS}$$
(33)

Modular transformations of (33) is

$$\chi_{m,n}^{NS}(\tau) = \int \chi_P^{NS}(-1/\tau) 2\sinh(\pi m P/b)\sinh(\pi n b P)dP.$$
(34)

Given that the identity field is specified by (m, n) = (1, 1) one finds the vacuum component of the matrix of modular transformation:

$$S_{0\alpha} = \sin \pi b^{-1} (\alpha - Q/2) \sin \pi b (\alpha - Q/2) \,. \tag{35}$$

For brevity the  $N_{\alpha}$  and  $\tilde{N}_{\alpha}$  primary fields will be denoted by  $\alpha$  and  $\tilde{\alpha}$  correspondingly. Therefore, if for example we take in the relation (5)  $\alpha$ ,  $\beta_2$  and  $\beta_5$  of NS type, it takes the form

$$S_{0\beta_{2}}\left[\int \frac{d\beta_{4}}{i}F_{0,\beta_{4}}\left[\begin{array}{cc}\beta_{2} & \alpha\\\beta_{2} & \alpha\end{array}\right]e^{2\pi i\Delta_{\beta_{4}}}F_{\beta_{4},\beta_{5}}\left[\begin{array}{cc}\alpha & \alpha\\\beta_{2} & \beta_{2}\end{array}\right]$$
(36)  
$$+\int \frac{d\beta_{4}}{i}F_{0,\tilde{\beta}_{4}}\left[\begin{array}{cc}\beta_{2} & \alpha\\\beta_{2} & \alpha\end{array}\right]e^{2\pi i\Delta_{\tilde{\beta}_{4}}}F_{\tilde{\beta}_{4},\beta_{5}}\left[\begin{array}{cc}\alpha & \alpha\\\beta_{2} & \beta_{2}\end{array}\right]\right] =$$
$$e^{\pi i(2\Delta_{\alpha}+2\Delta_{\beta_{2}})}F_{0,\beta_{5}}\left[\begin{array}{cc}\alpha & \alpha\\\alpha & \alpha\end{array}\right]S_{\alpha\beta_{2}}(\beta_{5}).$$

As we mentioned before  $S_{\alpha\beta_2}(\beta_5)$  in the limit  $\beta_5 \to 0$  should coincide with the matrix of modular transformations of the characters, and  $S_{0\beta_2}$  is given by (35).

The elements of the fusion matrix in the Neveu-Schwarz sector of N = 1SLFT are given by the formulae [25, 26]:

$$F_{\alpha_s,\alpha_t} \begin{bmatrix} \alpha_3 & \alpha_2 \\ \alpha_4 & \alpha_1 \end{bmatrix} = \frac{\mathcal{N}_{NS}(\alpha_s, \alpha_2, \alpha_1)\mathcal{N}_{NS}(\alpha_4, \alpha_3, \alpha_s)}{\mathcal{N}_{NS}(\alpha_t, \alpha_3, \alpha_2)\mathcal{N}_{NS}(\alpha_4, \alpha_t, \alpha_1)} \Big\{ \begin{array}{c} \alpha_1 & \alpha_3 \\ \alpha_2 & \alpha_4 \end{array} \Big| \begin{array}{c} \alpha_s \\ \alpha_t \end{array} \Big\}_{11}^{11}, \quad (37)$$

$$F_{\alpha_s,\tilde{\alpha_t}} \begin{bmatrix} \alpha_3 & \alpha_2 \\ \alpha_4 & \alpha_1 \end{bmatrix} = \frac{\mathcal{N}_{NS}(\alpha_s, \alpha_2, \alpha_1)\mathcal{N}_{NS}(\alpha_4, \alpha_3, \alpha_s)}{\mathcal{N}_R(\alpha_t, \alpha_3, \alpha_2)\mathcal{N}_R(\alpha_4, \alpha_t, \alpha_1)} \Big\{ \begin{array}{c} \alpha_1 & \alpha_3 \\ \alpha_2 & \alpha_4 \end{array} \big| \begin{array}{c} \alpha_s \\ \alpha_t \end{array} \Big\}_{00}^{11}, \quad (38)$$

$$F_{\tilde{\alpha_s},\alpha_t} \begin{bmatrix} \alpha_3 & \alpha_2 \\ \alpha_4 & \alpha_1 \end{bmatrix} = \frac{\mathcal{N}_R(\alpha_s,\alpha_2,\alpha_1)\mathcal{N}_R(\alpha_4,\alpha_3,\alpha_s)}{\mathcal{N}_{NS}(\alpha_t,\alpha_3,\alpha_2)\mathcal{N}_{NS}(\alpha_4,\alpha_t,\alpha_1)} \Big\{ \begin{smallmatrix} \alpha_1 & \alpha_3 \\ \alpha_2 & \alpha_4 \end{smallmatrix} \Big| \begin{smallmatrix} \alpha_s \\ \alpha_t \end{smallmatrix} \Big\}_{11}^{00}, \quad (39)$$

$$F_{\tilde{\alpha_s},\tilde{\alpha_t}} \begin{bmatrix} \alpha_3 & \alpha_2 \\ \alpha_4 & \alpha_1 \end{bmatrix} = \frac{\mathcal{N}_R(\alpha_s, \alpha_2, \alpha_1)\mathcal{N}_R(\alpha_4, \alpha_3, \alpha_s)}{\mathcal{N}_R(\alpha_t, \alpha_3, \alpha_2)\mathcal{N}_R(\alpha_4, \alpha_t, \alpha_1)} \Big\{ \begin{smallmatrix} \alpha_1 & \alpha_3 \\ \alpha_2 & \alpha_4 \end{smallmatrix} \Big| \begin{smallmatrix} \alpha_s \\ \alpha_t \end{smallmatrix} \Big\}_{00}^{00}, \qquad (40)$$

where normalization coefficients are described by the relations:

$$\mathcal{N}_{NS}(\alpha_3, \alpha_2, \alpha_1) = \tag{41}$$

$$\frac{\Gamma_{NS}(2\alpha_1)\Gamma_{NS}(2\alpha_2)\Gamma_{NS}(2Q - 2\alpha_3)}{\Gamma_{NS}(2Q - \alpha_1 - \alpha_2 - \alpha_3)\Gamma_{NS}(Q - \alpha_1 - \alpha_2 + \alpha_3)\Gamma_{NS}(\alpha_2 + \alpha_3 - \alpha_1)\Gamma_{NS}(\alpha_3 + \alpha_1 - \alpha_2)},$$

$$\mathcal{N}_{T}(\alpha_1, \alpha_2, \alpha_3) = \tag{42}$$

$$\frac{\Gamma_{NS}(2\alpha_1)\Gamma_{NS}(2\alpha_2)\Gamma_{NS}(2Q-2\alpha_3)}{\Gamma_R(2Q-\alpha_1-\alpha_2-\alpha_3)\Gamma_R(Q-\alpha_1-\alpha_2+\alpha_3)\Gamma_R(\alpha_2+\alpha_3-\alpha_1)\Gamma_R(\alpha_3+\alpha_1-\alpha_2)},$$
(42)

where the functions  $\Gamma_{NS}(x)$ ,  $\Gamma_R(x)$  are described in appendix A, and 6*j*-symbols of the quantum algebra  $U_q(osp(1|2))$  are [26]:

$$\left\{ \begin{array}{l} \alpha_{1} \alpha_{2} \\ \alpha_{3} \overline{\alpha_{4}} \\ \alpha_{4} \end{array} \right\}_{\nu_{1}\nu_{2}}^{\nu_{3}\nu_{4}} = |S_{1}(2\alpha_{t})|^{2} \frac{S_{\nu_{4}}(\alpha_{s} + \alpha_{2} - \alpha_{1})S_{\nu_{1}}(\alpha_{1} + \alpha_{t} - \alpha_{4})}{S_{\nu_{2}}(\alpha_{t} + \alpha_{2} - \alpha_{3})S_{\nu_{3}}(\alpha_{3} + \alpha_{s} - \alpha_{4})} I_{\alpha_{s},\alpha_{t}} \left[ \begin{array}{c} \alpha_{3} & \alpha_{2} \\ \alpha_{4} & \alpha_{1} \end{array} \right]_{\nu_{1}\nu_{2}}^{\nu_{3}\nu_{4}},$$

$$(43)$$

where  $\overline{\alpha}_4 = Q - \alpha_4, \nu_i = 0, 1, i = 1, 2, 3, 4, \sum_{i=1}^4 \nu_i = 0 \mod 2$ , and

$$I_{\alpha_{s},\alpha_{t}} \begin{bmatrix} \alpha_{3} & \alpha_{2} \\ \alpha_{4} & \alpha_{1} \end{bmatrix}_{\nu_{1}\nu_{2}}^{\nu_{3}\nu_{4}} = (-1)^{\nu_{3}\nu_{2}+\nu_{4}} \int_{-i\infty}^{i\infty} \sum_{\nu=0}^{1} (-1)^{\nu(\nu_{2}+\nu_{4})} S_{1+\nu+\nu_{3}}(y+\gamma_{1}^{\circ}) \qquad (44)$$

$$\times S_{1+\nu+\nu_{4}}(y+\gamma_{2}^{\circ})S_{1+\nu+\nu_{3}}(y+\gamma_{3}^{\circ})S_{1+\nu+\nu_{4}}(y+\gamma_{4}^{\circ})S_{\nu}(-y+\beta_{1}^{\circ})$$

$$\times S_{\nu+\nu_{2}+\nu_{3}}(-y+\beta_{2}^{\circ})S_{\nu+\nu_{2}+\nu_{3}}(-y+\beta_{3}^{\circ})S_{\nu}(-y+\beta_{4}^{\circ})\frac{dy}{2i}$$

$$= \int_{-i\infty}^{i\infty} \sum_{\nu=0}^{1} \Lambda(y+\gamma_{1}^{\circ},\nu_{3}+\nu)\Lambda(y+\gamma_{2}^{\circ},\nu+\nu_{4})\Lambda(y+\gamma_{3}^{\circ},\nu+\nu_{3})$$

$$\times \Lambda(y+\gamma_{4}^{\circ},\nu-\nu_{4})\Lambda(-y+\beta_{1}^{\circ},-\nu-1)\Lambda(-y+\beta_{2}^{\circ},-\nu+1+\nu_{2}-\nu_{3})$$

$$\times \Lambda(-y+\beta_{3}^{\circ},-\nu+1-\nu_{2}-\nu_{3})\Lambda(-y+\beta_{4}^{\circ},-\nu-1)\frac{dy}{2i},$$

where

$$\gamma_{1}^{\circ} = -Q/2 + \alpha_{3} - \alpha_{4}, \qquad \beta_{1}^{\circ} = Q/2 + \alpha_{s}, 
\gamma_{2}^{\circ} = -Q/2 + \alpha_{1} - \alpha_{2}, \qquad \beta_{2}^{\circ} = Q/2 - \alpha_{t} + \alpha_{4} + \alpha_{2}, 
\gamma_{3}^{\circ} = Q/2 - \alpha_{3} - \alpha_{4}, \qquad \beta_{3}^{\circ} = -Q/2 + \alpha_{t} + \alpha_{4} + \alpha_{2}, 
\gamma_{4}^{\circ} = Q/2 - \alpha_{1} - \alpha_{2}, \qquad \beta_{4}^{\circ} = 3Q/2 - \alpha_{s}.$$
(45)

The second equality in (44) was established in [4]. Let us explain briefly how it arises. First recall the following periodicity and reflection properties of  $\Lambda(y,\nu)$  [42, 43]:

$$\Lambda(y, m+2k) = (-1)^{mk+r\frac{k(k-1)}{2}}\Lambda(y, m), \qquad (46)$$

$$\Lambda(Q - x, -\nu)\Lambda(x, \nu) = (-1)^{\nu}.$$
(47)

For r = 2 the relation (46) implies

$$\Lambda(y, m+2k) = (-1)^{mk} \Lambda(y, m).$$
(48)

The functions  $S_{\nu}(x)$  are supposed to be periodic in the discrete variable:  $S_{\nu}(x) = S_{\nu+2}(x)$ . This together with (48) implies

$$\Lambda(y,2m) = S_1(y), \quad \Lambda(y,2m+1) = (-1)^m S_0(y).$$
(49)

Repeatedly using (49) one can arrive to the second equality in (44). One can check that the expressions (37)-(40) are invariant under the reflection  $\alpha \to Q-\alpha$  of any of six variables entering in the fusion matrix. Inserting (37)-(40) in (36) we derive:

$$S_{\alpha\beta_2}^{SL}(\beta_5) = \frac{\mathcal{N}_{NS}(\alpha, \beta_5, \alpha)}{\mathcal{N}_{NS}(\beta_2, \beta_5, \beta_2)} S_{\alpha\beta_2}(\beta_5)$$
(50)

where  $S_{\alpha\beta_2}(\beta_5)$  is computed by the same formula (36) where inserted only corresponding elements of 6j-symbols of the quantum algebra  $U_q(osp(1|2))$  appearing in the l.h.s of (37)-(40):

$$S_{0\beta_2}I_{\alpha\beta_2}(\beta_5) = e^{\pi i (2\Delta_{\alpha} + 2\Delta_{\beta_2})} \left\{ \begin{smallmatrix} \alpha & \alpha \\ \alpha & \alpha \end{smallmatrix} \middle| \begin{smallmatrix} 0 \\ \beta_5 \end{smallmatrix} \right\}_{11}^{11} S_{\alpha\beta_2}(\beta_5) , \qquad (51)$$

where

$$I_{\alpha\beta_{2}}(\beta_{5}) = \left[ \int \frac{d\beta_{4}}{i} \left\{ \begin{smallmatrix} \alpha & \beta_{2} & | & 0 \\ \alpha & \beta_{2} & | & \beta_{4} \end{smallmatrix} \right\}_{11}^{11} e^{2\pi i \Delta_{\beta_{4}}} \left\{ \begin{smallmatrix} \beta_{2} & \alpha & | & \beta_{4} \\ \alpha & \beta_{2} & | & \beta_{5} \end{smallmatrix} \right\}_{11}^{11}$$
(52)  
+ 
$$\int \frac{d\beta_{4}}{i} \left\{ \begin{smallmatrix} \alpha & \beta_{2} & | & 0 \\ \alpha & \beta_{2} & | & \beta_{4} \end{smallmatrix} \right\}_{00}^{10} e^{2\pi i \Delta_{\beta_{4}}} \left\{ \begin{smallmatrix} \beta_{2} & \alpha & | & \beta_{4} \\ \alpha & \beta_{2} & | & \beta_{5} \end{smallmatrix} \right\}_{11}^{00} \right].$$

#### 3 Elements of S-matrix for three NS primaries

Now we turn to the calculation of the matrix  $S_{\alpha\beta_2}(\beta_5)$  in the case when all the entering primaries of the NS type. For this purpose we use the formula (51). Start to calculate the left hand side integral (52). The first term in (52) is the following element of the 6*j*-symbol:  $\left\{ \begin{array}{c} \alpha \ \beta_2 \\ \alpha \ \beta_2 \end{array} \middle|_{\beta_4}^0 \right\}_{11}^{11}$ . Setting in (43)-(44)  $\nu_1 = \nu_2 = \nu_3 = \nu_4 = 1$ , and  $\alpha_1 = \alpha_2 = \alpha$ ,  $\alpha_3 = \beta_2$ ,

 $\alpha_4 = Q - \beta_2, \ \alpha_t = \beta_4, \ \alpha_s = \delta \to 0$  we obtain

$$\begin{cases} \alpha \beta_{2} \mid \beta_{4} \end{cases}_{11}^{11} = |S_{1}(2\beta_{4})|^{2} \frac{S_{1}(\delta)S_{1}(\alpha + \beta_{4} + \beta_{2} - Q)}{S_{1}(\beta_{4} + \alpha - \beta_{2})S_{1}(2\beta_{2} - Q)}$$

$$\times \int_{-i\infty}^{i\infty} \sum_{\nu=0}^{1} (-1)^{\nu+1} \Lambda(y - 3Q/2 + 2\beta_{2}, 1 + \nu)\Lambda(y - Q/2, \nu + 1)\Lambda(y + Q/2 - 2\alpha, \nu - 1)$$

$$\times \Lambda(-y + Q/2 + \delta, -\nu - 1)\Lambda(-y + 3Q/2 - \beta_{4} - \beta_{2} + \alpha, -\nu + 1)$$

$$\times \Lambda(-y + Q/2 + \beta_{4} - \beta_{2} + \alpha, -\nu - 1)\frac{dy}{2i}$$

$$= |S_{1}(2\beta_{4})|^{2}S_{1}(\delta)S_{1}(\alpha + \beta_{4} + \beta_{2} - Q)\Lambda(\delta, 0)\Lambda(Q - \beta_{2} - \beta_{4} + \alpha, 2)\Lambda(-2\alpha, -2) .$$
(53)

To compute integral here we used formula (106) in appendix B for r = 2.

Now we should compute the second term in (52):  $\begin{cases} \beta_2 & \alpha \\ \alpha & \beta_2 \end{cases} \begin{vmatrix} \beta_4 \\ \beta_5 \end{vmatrix}_{11}^{11}$ . Setting in (43)  $\nu_1 = \nu_2 = \nu_3 = \nu_4 = 1$ , and  $\alpha_1 = \beta_2$ ,  $\alpha_2 = \alpha$ ,  $\alpha_3 = \alpha$ ,  $\alpha_4 = Q - \beta_2$ ,  $\alpha_t = \beta_5, \ \alpha_s = \beta_4$  we have

$$\begin{cases} \beta_{2} \alpha \\ \alpha \beta_{2} \ | \beta_{5} \\ \beta_{5} \\ \beta_{11} \\ = |S_{1}(2\beta_{5})|^{2} \frac{S_{1}(\alpha + \beta_{4} - \beta_{2})S_{1}(2\beta_{2} + \beta_{5} - Q)}{S_{1}(\beta_{4} + \alpha + \beta_{2} - Q)S_{1}(\beta_{5})} \\ \times \int_{-i\infty}^{i\infty} \sum_{\nu=0}^{1} \Lambda(-y + Q/2 + \beta_{4}, -\nu - 1)\Lambda(-y + 3Q/2 - \beta_{4}, -\nu - 1)F(y, \nu + 1)(-1)^{\nu - 1} \frac{dy}{2i}, \end{cases}$$
(54)

where

$$F(y,\nu) = \Lambda(y - 3Q/2 + \beta_2 + \alpha, \nu)\Lambda(y - Q/2 + \beta_2 - \alpha, \nu)$$
(55)  
 
$$\times \Lambda(y - Q/2 - \alpha + \beta_2, \nu)\Lambda(y + Q/2 - \alpha - \beta_2, \nu)$$
  
 
$$\times \Lambda(-y + 3Q/2 - \beta_5 - \beta_2 + \alpha, -\nu + 2)\Lambda(-y + Q/2 + \beta_5 - \beta_2 + \alpha, -\nu).$$

Writing in this way we separated in integral appearing in (54)  $\beta_4$ -dependent terms and collected in  $F(y, \nu)$  all  $\beta_4$ -independent terms. The next element of the fusion matrix to be computed is  $\left\{ \begin{array}{c} \alpha & \beta_2 \\ \alpha & \beta_2 \end{array} | \begin{array}{c} \delta \\ \beta_4 \end{array} \right\}_{00}^{11}$ . Putting in (43)  $\nu_1 = \nu_2 = 0$ ,  $\nu_3 = \nu_4 = 1$ , and  $\alpha_1 = \alpha_2 = \alpha$ ,  $\alpha_3 = \beta_2$ ,  $\alpha_4 = Q - \beta_2$ ,  $\alpha_t = \beta_4$ ,  $\alpha_s = \delta \rightarrow 0$  one has

z

$$\begin{cases} \alpha \beta_{2} | \delta_{4} \rangle_{00}^{11} = |S_{1}(2\beta_{4})|^{2} \frac{S_{1}(\delta)S_{0}(\alpha + \beta_{4} + \beta_{2} - Q)}{S_{0}(\beta_{4} + \alpha - \beta_{2})S_{1}(2\beta_{2} - Q)}$$

$$\times \int_{-i\infty}^{i\infty} \sum_{\nu=0}^{1} (-1)^{\nu+1} \Lambda(y - 3Q/2 + 2\beta_{2}, 1 + \nu)\Lambda(y - Q/2, \nu + 1)\Lambda(y + Q/2 - 2\alpha, \nu - 1)$$

$$\times \Lambda(-y + Q/2 + \delta, -\nu - 1)\Lambda(-y + 3Q/2 - \beta_{4} - \beta_{2} + \alpha, -\nu)\Lambda(-y + Q/2 + \beta_{4} - \beta_{2} + \alpha, -\nu) \frac{dy}{2i}$$

$$= |S_{1}(2\beta_{4})|^{2}S_{1}(\delta)S_{0}(\alpha + \beta_{4} + \beta_{2} - Q)\Lambda(\delta, 0)\Lambda(Q - \beta_{2} - \beta_{4} + \alpha, 1)\Lambda(-2\alpha, -2).$$
(56)

The last element of the fusion matrix which we need to compute is  $\begin{cases} \beta_2 & \alpha \\ \alpha & \beta_2 \end{cases} \begin{vmatrix} \beta_4 \\ \beta_5 \end{vmatrix}_{11}^{00}$ . Now setting in (43)  $\nu_1 = \nu_2 = 1$ ,  $\nu_3 = \nu_4 = 0$ , and  $\alpha_1 = \beta_2$ ,  $\alpha_2 = \alpha$ ,  $\alpha_3 = \alpha$ ,  $\alpha_4 = Q - \beta_2$ ,  $\alpha_t = \beta_5$ ,  $\alpha_s = \beta_4$  one gets:

$$\begin{cases} \beta_{2} \alpha \\ \alpha \beta_{2} \end{vmatrix} \beta_{4}^{\beta_{2}} \beta_{11}^{00} = |S_{1}(2\beta_{5})|^{2} \frac{S_{0}(\alpha + \beta_{4} - \beta_{2})S_{1}(2\beta_{2} + \beta_{5} - Q)}{S_{0}(\beta_{4} + \alpha + \beta_{2} - Q)S_{1}(\beta_{5})}$$
(57)  
 
$$\times \int_{-i\infty}^{i\infty} \sum_{\nu=0}^{1} \Lambda(-y + Q/2 + \beta_{4}, -\nu - 1)\Lambda(-y + 3Q/2 - \beta_{4}, -\nu - 1)F(y, \nu)\frac{dy}{2i},$$

were the function  $F(y, \nu)$  was defined in (55).

Since  $\beta_4$  has the form  $\beta_4 = Q/2 + iP$  for some real P, one can write

$$|S_1(2\beta_4)|^2 = S_1(2\beta_4)S_1(2Q - 2\beta_4) = \frac{1}{S_1(Q - 2\beta_4)S_1(-Q + 2\beta_4)}$$
$$= \frac{1}{S_1(2(Q/2 - \beta_4))S_1(2(-Q/2 + \beta_4))} = \frac{1}{S_1(2z)S_1(-2z)},$$

where  $z = -Q/2 + \beta_4$ . Recalling also formulae for conformal dimensions (27) and (28) we obtain

$$I_{\alpha\beta_{2}}(\beta_{5})$$

$$= C \int_{-i\infty}^{i\infty} \frac{dy}{i} F(y,0) \int_{0}^{i\infty} \frac{dz}{2i} \sum_{m=0}^{1} \frac{\Lambda(Q/2 + \alpha - \beta_{2} \pm z, \pm m)\Lambda(-y + Q \pm z, \pm m)}{\Lambda(\pm 2z, \pm 2m)} e^{-i\pi z^{2}} (-1)^{m}$$

$$- C \int_{-i\infty}^{i\infty} \frac{dy}{i} F(y,1) \int_{0}^{i\infty} \frac{dz}{2i} \sum_{m=0}^{1} \frac{\Lambda(Q/2 + \alpha - \beta_{2} \pm z, 1 \pm m)\Lambda(-y + Q \pm z, \pm m)}{\Lambda(\pm 2z, \pm 2m)} e^{-i\pi z^{2}} (-1)^{m} ,$$

$$= \beta_{4} - \frac{Q}{2} , \qquad C = 2S_{1}(\delta)\Lambda(\delta, 0)\Lambda(-2\alpha, -2) \frac{|S_{1}(2\beta_{5})|^{2}S_{1}(2\beta_{2} + \beta_{5} - Q)}{S_{1}(\beta_{5})} e^{Q^{2}\pi i/4} .$$

$$(58)$$

$$(58)$$

Here we used the shorthand notation  $\Lambda(y \pm x, m \pm n) \equiv \Lambda(y + x, m + n)\Lambda(y - x, m - n)$ . The integrals appearing in the third and the fourth lines of (58)

are particular cases of the integral (114) in appendix B for r = 2. Hence using (114) in appendix B for r = 2 one has

$$\begin{split} I_{\alpha\beta_{2}}(\beta_{5}) &= C \int_{-i\infty}^{i\infty} \frac{dy}{i} F(y,0) \Lambda(3Q/2 + \alpha - \beta_{2} - y,0) e^{-\frac{i\pi}{4}(-Q/2 + \alpha - \beta_{2} + y)^{2}} e^{\frac{i\pi}{4}(3Q/2 + \alpha - \beta_{2} - y)Q} \ (60) \\ &- C \int_{-i\infty}^{i\infty} \frac{dy}{i} F(y,1) \Lambda(3Q/2 + \alpha - \beta_{2} - y,1) e^{-\frac{i\pi}{4}(-Q/2 + \alpha - \beta_{2} + y)^{2}} e^{\frac{i\pi}{4}(3Q/2 + \alpha - \beta_{2} - y)Q} e^{-\frac{i\pi}{4}} e^{\frac{i\pi}{2}} \\ &= C' \int_{-i\infty}^{i\infty} \frac{dy}{i} \sum_{\nu=0}^{1} \Lambda(y - 3Q/2 + \beta_{2} + \alpha, \nu) \Lambda(y - Q/2 - \alpha + \beta_{2}, \nu) \Lambda(y + Q/2 - \alpha - \beta_{2}, \nu) \\ &\times \Lambda(-y + 3Q/2 - \beta_{5} - \beta_{2} + \alpha, -\nu + 2) \Lambda(-y + Q/2 + \beta_{5} - \beta_{2} + \alpha, -\nu) \\ &\times e^{-\frac{i\pi}{4}y^{2}} e^{-\frac{i\pi}{2}y(\alpha - \beta_{2})} e^{-\frac{i\pi\nu}{4}} e^{\frac{3i\pi\nu}{2}}, \end{split}$$

where

$$C' = C e^{-\frac{i\pi}{4}(-Q/2 + \alpha - \beta_2)^2} e^{\frac{i\pi}{4}(3Q/2 + \alpha - \beta_2)Q}.$$
(61)

Changing sign of y and using that for  $\nu = 0, 1$  hold the relations:  $\Lambda(y, -\nu) = (-1)^{\nu} \Lambda(y, \nu)$  and  $\Lambda(y, 2 - \nu) = \Lambda(y, \nu)$  we can write the last integral in (60) in the form

$$I_{\alpha\beta_{2}}(\beta_{5}) = C' \int_{-i\infty}^{i\infty} \frac{dy}{i} \sum_{\nu=0}^{1} \Lambda(-y - 3Q/2 + \beta_{2} + \alpha, -\nu) \Lambda(-y - Q/2 - \alpha + \beta_{2}, -\nu) \\ \times \Lambda(-y + Q/2 - \alpha - \beta_{2}, -\nu) \Lambda(y + 3Q/2 - \beta_{5} - \beta_{2} + \alpha, \nu) \Lambda(y + Q/2 + \beta_{5} - \beta_{2} + \alpha, \nu) \\ \times e^{-\frac{i\pi}{4}y^{2}} e^{\frac{i\pi}{2}y(\alpha - \beta_{2})} e^{-\frac{i\pi\nu}{4}} e^{\frac{3i\pi\nu}{2}}.$$

It is easy to see that this integral has the same structure as the integral on the left hand side of relation (128) in appendix D for r = 2, therefore we can write for him taking as  $\gamma_1 = Q/2 + \beta_5 - \beta_2 + \alpha$  and  $\gamma_2 = 3Q/2 - \beta_5 - \beta_2 + \alpha$ 

$$I_{\alpha\beta_{2}}(\beta_{5}) = C'e^{\frac{i\pi}{4}((\alpha-\beta_{2})^{2}+Q^{2}/4-Q(Q/2+\beta_{5}+2\alpha-2\beta_{2}))}$$
(62)  
  $\times \Lambda(2\alpha-\beta_{5},0)\Lambda(Q-\beta_{5},0)\Lambda(2Q-2\beta_{2}-\beta_{5},0)$   
  $\times \int_{-i\infty}^{i\infty} \sum_{m=0}^{1} \frac{\Lambda(\pm y-Q+\beta_{2}+\alpha+\beta_{5}/2,\pm m)\Lambda(\pm y-\alpha+\beta_{2}+\beta_{5}/2,\pm m)}{\Lambda(2y,2m)\Lambda(-2y,-(2m))}$   
  $\times \Lambda(\pm y+Q-\alpha-\beta_{2}+\beta_{5}/2,\pm m)\Lambda(\pm y+\alpha-\beta_{2}+\beta_{5}/2,\pm m)e^{-i\pi y^{2}}e^{-i\pi m^{2}}\frac{dy}{2i}.$ 

Note that integral in the r.h.s. of (62) coincides with the integral in the r.h.s. of (124) in appendix C for r = 2. Thus we can write for the latter

$$\begin{split} &\int_{-i\infty}^{i\infty} \sum_{m=0}^{1} \Lambda(\pm y - Q/2 + \beta_2 + \beta_5/2, \pm m) \Lambda(\pm y + Q/2 - \beta_2 + \beta_5/2, \pm m) e^{2i\pi y(-Q/2+\alpha)} \frac{dy}{i} = \\ &\Lambda(2\alpha - \beta_5, 0) \Lambda(-2\alpha - \beta_5 + 2Q, 0) e^{-i\pi \left(\frac{1}{4}\beta_5^2 - (-Q/2+\alpha)^2 - (Q/2-\beta_2)^2\right)} \\ &\times \int_{-i\infty}^{i\infty} \sum_{m=0}^{1} \frac{\Lambda(\pm y - Q + \beta_2 + \alpha + \beta_5/2, \pm m) \Lambda(\pm y - \alpha + \beta_2 + \beta_5/2, \pm m)}{\Lambda(2y, 2m) \Lambda(-2y, -(2m))} \\ &\times \Lambda(\pm y + Q - \alpha - \beta_2 + \beta_5/2, \pm m) \Lambda(\pm y + \alpha - \beta_2 + \beta_5/2, \pm m) e^{-i\pi y^2} e^{-i\pi m^2} \frac{dy}{2i} \,. \end{split}$$

Collecting all we obtain:

$$S_{\alpha\beta_{2}}(\beta_{5}) = S_{0\beta_{2}} \frac{e^{-\frac{i\pi}{2}\Delta_{\beta_{5}}}}{S_{1}(\beta_{5})}$$

$$\times \int_{-i\infty}^{i\infty} \sum_{m=0}^{1} \Lambda(\pm y - Q/2 + \beta_{2} + \beta_{5}/2, \pm m)\Lambda(\pm y + Q/2 - \beta_{2} + \beta_{5}/2, \pm m)e^{2i\pi y(-Q/2 + \alpha)}\frac{dy}{i}$$
(63)

or using (49):

$$S_{\alpha\beta_{2}}(\beta_{5}) = S_{0\beta_{2}} \frac{e^{-\frac{i\pi}{2}\Delta_{\beta_{5}}}}{S_{\rm NS}(\beta_{5})}$$

$$\times \int_{-i\infty}^{i\infty} \left[ S_{\rm NS}(\pm y - Q/2 + \beta_{2} + \beta_{5}/2) S_{\rm NS}(\pm y + Q/2 - \beta_{2} + \beta_{5}/2) + S_{\rm R}(\pm y - Q/2 + \beta_{2} + \beta_{5}/2) S_{\rm R}(\pm y + Q/2 - \beta_{2} + \beta_{5}/2) \right] e^{2i\pi y(-Q/2 + \alpha)} \frac{dy}{i}.$$
(64)

This expression has number of properties analogous to that of in usual Liouville field theory [52]. First we obtain easily

$$S_{\alpha\beta_2}(\beta_5) = S_{Q-\alpha,\beta_2}(\beta_5) = S_{\alpha,Q-\beta_2}(\beta_5).$$
(65)

Using formula (140) in appendix E for r = 2 one can establish

$$S_{\alpha\beta_{2}}(Q - \beta_{5}) = S_{\rm NS}(-Q + 2\alpha + \beta_{5})S_{\rm NS}(Q - 2\alpha + \beta_{5})$$
(66)  
 
$$\times S_{\rm NS}(2Q - \beta_{5} - 2\beta_{2})S_{\rm NS}(-\beta_{5} + 2\beta_{2})S_{\alpha\beta_{2}}(\beta_{5}).$$

Combining (65) and (66) with definition of the normalization factors (41) and using formulae (103) in appendix A we obtain that

$$S_{\alpha\beta_2}^{SL}(\beta_5) = \frac{\mathcal{N}_{NS}(\alpha, \beta_5, \alpha)}{\mathcal{N}_{NS}(\beta_2, \beta_5, \beta_2)} S_{\alpha\beta_2}(\beta_5)$$
(67)

is invariant under reflection of all three variables.

Now let us check that (64) is self-consistent in the limit  $\beta_5 \to 0$ . In this limit using formulae in appendix A one has that  $\frac{1}{S_{\rm NS}(\beta_5)} \sim \pi\beta_5 \to 0$ , and

thus we should pick up only diverging part of integral. This diverging part originates due to pinching of integration contour in this limit between poles of  $S_{\rm NS}(y-Q/2+\beta_2+\beta_5/2)$  and  $S_{\rm NS}(-y+Q/2-\beta_2+\beta_5/2)$  and also between poles of  $S_{\rm NS}(-y-Q/2+\beta_2+\beta_5/2)$  and  $S_{\rm NS}(y+Q/2-\beta_2+\beta_5/2)$ . Analyzing poles of  $S_R(x)$  reviewed in appendix A we see that the second term in (64) has no diverging part. The diverging part caused by pinching is pole with coefficient given by the sum of residues:

$$\begin{aligned} &\frac{2\pi i}{i\pi^2\beta_5}S_{\rm NS}(-Q+2\beta_2)S_{\rm NS}(Q-2\beta_2)(e^{2i\pi(Q/2-\beta_2)(-Q/2+\alpha)}+e^{-2i\pi(Q/2-\beta_2)(-Q/2+\alpha)})\\ &=-\frac{1}{\pi\beta_5}\frac{\cos\pi(Q/2-\beta_2)(-Q/2+\alpha)}{\sin\frac{\pi}{b}(Q/2-\beta_2)\sin\pi b(Q/2-\beta_2)},\end{aligned}$$

where the product  $S_{\rm NS}(-Q+2\beta_2)S_{\rm NS}(Q-2\beta_2)$  was computed using formulae (97) and (98) in appendix A. Recalling (35) and (32) we obtain that indeed

$$\lim_{\beta_5 \to 0} S^{SL}_{\alpha\beta_2}(\beta_5) = S_{\alpha\beta_2} \,. \tag{68}$$

# 4 Elements of S-matrix for two NS and one NS primaries

Here we consider the case when  $\alpha, \beta_2$  are NS primaries and  $\beta_5$  is NS primary. In this case the relation (36) takes the form

$$S_{\alpha\beta_2}^{SL}(\tilde{\beta_5}) = \frac{\mathcal{N}_R(\alpha, \beta_5, \alpha)}{\mathcal{N}_R(\beta_2, \beta_5, \beta_2)} S_{\alpha\beta_2}(\tilde{\beta_5}), \qquad (69)$$

where  $S_{\alpha\beta_2}(\tilde{\beta}_5)$  is given by the relation:

$$S_{0\beta_2}I_{\alpha\beta_2}(\tilde{\beta}_5) = e^{\pi i (2\Delta_\alpha + 2\Delta_{\beta_2})} \left\{ \begin{smallmatrix} \alpha & \alpha \\ \alpha & \alpha \end{smallmatrix} \middle| \begin{smallmatrix} \delta \\ \beta_5 \end{smallmatrix} \right\}_{00}^{11} S_{\alpha\beta_2}(\tilde{\beta}_5)$$
(70)

and

$$I_{\alpha\beta_{2}}(\tilde{\beta}_{5}) = \left[ \int \frac{d\beta_{4}}{i} \left\{ \begin{array}{c} \alpha \ \beta_{2} \ \alpha \ \beta_{2} \ \beta_{4} \end{array} \right\}_{11}^{11} e^{2\pi i \Delta_{\beta_{4}}} \left\{ \begin{array}{c} \beta_{2} \ \alpha \ \beta_{2} \ \beta_{4} \end{array} \right\}_{00}^{11} \right.$$
(71)
$$+ \int \frac{d\beta_{4}}{i} \left\{ \begin{array}{c} \alpha \ \beta_{2} \ \beta_{4} \end{array} \right\}_{00}^{11} e^{2\pi i \Delta_{\tilde{\beta}_{4}}} \left\{ \begin{array}{c} \beta_{2} \ \alpha \ \beta_{2} \ \beta_{4} \end{array} \right\}_{00}^{00} \right].$$

Here  $\delta$  should be taken to 0. The first terms in the first and second lines  $\left\{ \begin{array}{c} \alpha \ \beta_2 \\ \alpha \ \beta_2 \end{array} \middle|_{\beta_4}^{\delta} \right\}_{11}^{11}$  and  $\left\{ \begin{array}{c} \alpha \ \beta_2 \\ \alpha \ \beta_2 \end{array} \middle|_{\beta_4}^{\delta} \right\}_{00}^{11}$  are the same as before and computed in (53) and (56) respectively.

To compute the second fusion matrix element in the first line we set in (43)  $\nu_1 = \nu_2 = 0$ ,  $\nu_3 = \nu_4 = 1$ , and  $\alpha_1 = \beta_2$ ,  $\alpha_2 = \alpha$ ,  $\alpha_3 = \alpha$ ,  $\alpha_4 = Q - \beta_2$ ,  $\alpha_t = \beta_5$ ,

 $\alpha_s = \beta_4$  and receive

$$\begin{cases} \beta_{2} \alpha \\ \alpha \beta_{2} \\ \beta_{5} \end{cases}^{11}_{00} = |S_{1}(2\beta_{5})|^{2} \frac{S_{1}(\alpha + \beta_{4} - \beta_{2})S_{0}(2\beta_{2} + \beta_{5} - Q)}{S_{1}(\beta_{4} + \alpha + \beta_{2} - Q)S_{0}(\beta_{5})}$$

$$\times \int_{-i\infty}^{i\infty} \sum_{\nu=0}^{1} \Lambda(-y + Q/2 + \beta_{4}, -\nu - 1)\Lambda(-y + 3Q/2 - \beta_{4}, -\nu - 1)F_{2}(y, \nu + 1)(-1)^{\nu - 1} \frac{dy}{2i},$$
(72)

where we defined

$$F_{2}(y,\nu) = \Lambda(y - 3Q/2 + \beta_{2} + \alpha,\nu)\Lambda(y - Q/2 + \beta_{2} - \alpha,\nu)$$
(73)  
  $\times \Lambda(y - Q/2 - \alpha + \beta_{2},\nu)\Lambda(y + Q/2 - \alpha - \beta_{2},\nu)$   
  $\times \Lambda(-y + 3Q/2 - \beta_{5} - \beta_{2} + \alpha, -\nu + 1)\Lambda(-y + Q/2 + \beta_{5} - \beta_{2} + \alpha, -\nu + 1).$ 

To derive the second fusion matrix element in the second line we set in (43):  $\nu_1 = \nu_2 = 0, \ \nu_3 = \nu_4 = 0, \ \text{and} \ \alpha_1 = \beta_2, \ \alpha_2 = \alpha, \ \alpha_3 = \alpha, \ \alpha_4 = Q - \beta_2, \ \alpha_t = \beta_5, \ \alpha_s = \beta_4 \ \text{and obtain}$ 

$$\begin{cases} \beta_{2} \alpha \\ \alpha \beta_{2} \\ \beta_{5} \end{cases}_{00}^{00} = |S_{1}(2\beta_{5})|^{2} \frac{S_{0}(\alpha + \beta_{4} - \beta_{2})S_{0}(2\beta_{2} + \beta_{5} - Q)}{S_{0}(\beta_{4} + \alpha + \beta_{2} - Q)S_{0}(\beta_{5})}$$
(74)  
 
$$\times \int_{-i\infty}^{i\infty} \sum_{\nu=0}^{1} \Lambda(-y + Q/2 + \beta_{4}, -\nu - 1)\Lambda(-y + 3Q/2 - \beta_{4}, -\nu - 1)F_{2}(y, \nu)\frac{dy}{2i}.$$

Collecting all, and repeating the same steps as in the first case and using the integral (114) in appendix B, we obtain

$$I_{\alpha\beta_{2}}(\tilde{\beta}_{5}) = C'' \int_{-i\infty}^{i\infty} \frac{dy}{i} \sum_{\nu=0}^{1} \Lambda(-y - 3Q/2 + \beta_{2} + \alpha, -\nu) \Lambda(-y - Q/2 - \alpha + \beta_{2}, -\nu)$$
(75)  
  $\times \Lambda(-y + Q/2 - \alpha - \beta_{2}, -\nu) \Lambda(y + 3Q/2 - \beta_{5} - \beta_{2} + \alpha, \nu + 1)$   
  $\times \Lambda(y + Q/2 + \beta_{5} - \beta_{2} + \alpha, \nu + 1) e^{-\frac{i\pi}{4}y^{2}} e^{\frac{i\pi}{2}y(\alpha - \beta_{2})} e^{-\frac{i\pi\nu}{4}} e^{\frac{i\pi\nu}{2}},$ 

where

$$C'' = S_1(\delta)\Lambda(\delta,0)\Lambda(-2\alpha,-2)\frac{|S_1(2\beta_5)|^2 S_0(2\beta_2+\beta_5-Q)}{S_0(\beta_5)}$$
(76)  
 
$$\times e^{-\frac{i\pi}{4}(-Q/2+\alpha-\beta_2)^2} e^{\frac{i\pi}{4}(3Q/2+\alpha-\beta_2)Q} e^{Q^2\pi i/4}.$$

The integral (75) is of the form (128) in appendix D, which yields

$$I_{\alpha\beta_{2}}(\tilde{\beta}_{5}) = C'' e^{\frac{i\pi}{4}((\alpha-\beta_{2})^{2}+Q^{2}/4-Q(Q/2+\beta_{5}+2\alpha-2\beta_{2}))} e^{\frac{i\pi}{4}}$$
(77)  
  $\times \Lambda(2\alpha-\beta_{5},1)\Lambda(Q-\beta_{5},1)\Lambda(2Q-2\beta_{2}-\beta_{5},1)\int_{-i\infty}^{i\infty}\sum_{m=0}^{1} \frac{\Lambda(\pm y-Q+\beta_{2}+\alpha+\beta_{5}/2,\pm m-\theta(\mp 1))\Lambda(\pm y-\alpha+\beta_{2}+\beta_{5}/2,\pm m-\theta(\mp 1))}{\Lambda(2y,2m+1)\Lambda(-2y,-2m-1)} \times \Lambda(\pm y+Q-\alpha-\beta_{2}+\beta_{5}/2,\pm m-\theta(\mp 1))\Lambda(\pm y+\alpha-\beta_{2}+\beta_{5}/2,\pm m+1+\theta(\pm 1))) \times e^{-i\pi y^{2}}\frac{dy}{2i}.$ 

Here  $\theta(x)$  is Heaviside step function:  $\theta(-1) = 0$  and  $\theta(1) = 1$ . As before the functions with double sign in arguments are used as shorthand for their product where the first term is taken with upper signs everywhere and the second term is taken with lower signs everywhere. For example the first term in integrand reads

$$\Lambda(\pm y - Q + \beta_2 + \alpha + \beta_5/2, \pm m - \theta(\mp 1)) \equiv$$
(78)  
$$\Lambda(y - Q + \beta_2 + \alpha + \beta_5/2, m)\Lambda(-y - Q + \beta_2 + \alpha + \beta_5/2, -m - 1).$$

And finally we express the r.h.s. of (77) via (124) in appendix C

$$\int_{-i\infty}^{i\infty} \sum_{m=0}^{1} \Lambda(y - Q/2 + \beta_2 + \beta_5/2, m - 1)\Lambda(y + Q/2 - \beta_2 + \beta_5/2, m + 1)$$
(79)  
$$\Lambda(-y - Q/2 + \beta_2 + \beta_5/2, -m)\Lambda(-y + Q/2 - \beta_2 + \beta_5/2, -m)e^{2i\pi y(-Q/2 + \alpha)}e^{i\pi m}\frac{dy}{i} =$$
  
$$\Lambda(2\alpha - \beta_5, 1)\Lambda(-2\alpha - \beta_5 + 2Q, -1)e^{-i\pi(\frac{1}{4}\beta_5^2 - (\alpha - Q/2)^2 - (Q/2 - \beta_2)^2)}e^{\frac{i\pi}{2}}\int_{-i\infty}^{i\infty}\sum_{m=0}^{1}$$
  
$$\frac{\Lambda(\pm y - Q + \beta_2 + \alpha + \beta_5/2, \pm m - \theta(\mp 1))\Lambda(\pm y - \alpha + \beta_2 + \beta_5/2, \pm m - \theta(\mp 1))}{\Lambda(2y, 2m + 1)\Lambda(-2y, -2m - 1)}$$
  
$$\times\Lambda(\pm y + Q - \alpha - \beta_2 + \beta_5/2, \pm m - \theta(\mp 1))\Lambda(\pm y + \alpha - \beta_2 + \beta_5/2, \pm m + 1 + \theta(\pm 1))$$
  
$$e^{-i\pi y^2}\frac{dy}{2i}.$$

Collecting all we derive

$$S_{\alpha\beta_{2}}(\tilde{\beta}_{5}) = -S_{0\beta_{2}} \frac{e^{-\frac{i\pi}{2}(\Delta_{\beta_{5}}+1/2)}}{S_{0}(\beta_{5})}$$

$$\times \int_{-i\infty}^{i\infty} \sum_{m=0}^{1} \Lambda(y - Q/2 + \beta_{2} + \beta_{5}/2, m - 1)\Lambda(y + Q/2 - \beta_{2} + \beta_{5}/2, m + 1)$$

$$\times \Lambda(-y - Q/2 + \beta_{2} + \beta_{5}/2, -m)\Lambda(-y + Q/2 - \beta_{2} + \beta_{5}/2, -m)e^{2i\pi y(-Q/2 + \alpha)}e^{i\pi m}\frac{dy}{i},$$
or using (49):

$$S_{\alpha\beta_{2}}(\tilde{\beta}_{5}) = S_{0\beta_{2}} \frac{e^{-\frac{i\pi}{2}\Delta_{\beta_{5}}}}{S_{\mathrm{R}}(\beta_{5})}$$

$$\times \int_{-i\infty}^{i\infty} \left[ S_{\mathrm{NS}}(y - Q/2 + \beta_{2} + \beta_{5}/2) S_{\mathrm{NS}}(y + Q/2 - \beta_{2} + \beta_{5}/2) \right] \times S_{\mathrm{R}}(-y - Q/2 + \beta_{2} + \beta_{5}/2) S_{\mathrm{R}}(-y + Q/2 - \beta_{2} + \beta_{5}/2) + S_{\mathrm{R}}(y - Q/2 + \beta_{2} + \beta_{5}/2) S_{\mathrm{R}}(y + Q/2 - \beta_{2} + \beta_{5}/2) \times S_{\mathrm{NS}}(-y - Q/2 + \beta_{2} + \beta_{5}/2) S_{\mathrm{NS}}(-y + Q/2 - \beta_{2} + \beta_{5}/2) \right] e^{2i\pi y(-Q/2 + \alpha)} \frac{dy}{i} .$$

$$(81)$$

Note the following properties of this expression. First we obtain easily

$$S_{\alpha\beta_2}(\tilde{\beta}_5) = S_{Q-\alpha,\beta_2}(\tilde{\beta}_5) = S_{\alpha,Q-\beta_2}(\tilde{\beta}_5).$$
(82)

Using formula (140) in appendix E one can establish

$$S_{\alpha\beta_2}(\widetilde{Q-\beta_5}) = S_{\mathrm{R}}(-Q+2\alpha+\beta_5)S_{\mathrm{R}}(Q-2\alpha+\beta_5)$$

$$\times S_{\mathrm{R}}(2Q-\beta_5-2\beta_2)S_{\mathrm{R}}(-\beta_5+2\beta_2)S_{\alpha\beta_2}(\tilde{\beta}_5).$$
(83)

Combining (82) and (83) with definition of the normalization factors (42) and using formulae (103) in apendix A we obtain that

$$S_{\alpha\beta_2}^{SL}(\tilde{\beta}_5) = \frac{\mathcal{N}_R(\alpha, \beta_5, \alpha)}{\mathcal{N}_R(\beta_2, \beta_5, \beta_2)} S_{\alpha\beta_2}(\tilde{\beta}_5)$$
(84)

is invariant under the reflection of all three variables.

## 5 Conclusion

In this paper we calculated the matrix  $S_{\alpha\beta_2}(\beta_5)$  in two cases. These results allow us to conjecture that in all cases the integral part of the matrix  $S_{\alpha\beta_2}(\beta_5)$  will be given by the following function with the various choices of discrete variables  $k_a, t_a, a = 1, 2$  and N:

$$F_{\alpha,\beta_2}(\beta_5)(\underline{k},\underline{t},N) = \int_{-i\infty}^{i\infty} \sum_{m=0}^{1} \Lambda(y-Q/2+\beta_2+\beta_5/2,k_1+m)$$
(85)  
  $\times \Lambda(y+Q/2-\beta_2+\beta_5/2,k_2+m)\Lambda(-y-Q/2+\beta_2+\beta_5/2,t_1-m)$   
  $\times \Lambda(-y+Q/2-\beta_2+\beta_5/2,t_2-m)e^{2i\pi y(-Q/2+\alpha)}e^{i\pi mN}dy.$ 

Applying some limiting procedures to the finite-difference equation derived in [4] we obtained that for the parameters  $t_1, t_2, k_1, k_2$  obeying the condition

$$k_1 - k_2 = t_1 - t_2 + 2l, (86)$$

the function (85) satisfies the eigenvalue equation:

$$e^{\frac{i\pi(L-l)}{2}} \frac{\sin\frac{\pi b}{2}(2\alpha - Q - \beta_5 - (N-M)/b)}{\sin\frac{b\pi}{2}(2\alpha - Q - (N+l)/b)} F_{\alpha-b/2,\beta_2}(\beta_5)(\underline{k},\underline{t},N+1) \quad (87)$$

$$+e^{-\frac{i\pi(L+l)}{2}} \frac{\sin\frac{\pi b}{2}(2\alpha - Q + \beta_5 - (N+M)/b)}{\sin\frac{b\pi}{2}(2\alpha - Q - (N+l)/b)} F_{\alpha+b/2,\beta_2}(\beta_5)(\underline{k},\underline{t},N-1)$$

$$= 2\cos\frac{\pi b}{2}(Q - 2\beta_2 - (t_2 - t_1)/b)F_{\alpha,\beta_2}(\beta_5)(\underline{k},\underline{t},N) ,$$

where

$$M = \frac{1}{2} \sum_{1,2} (t_a + k_a), \quad L = \frac{1}{2} \sum_{1,2} (t_a - k_a).$$
(88)

As in the case of Liouville theory one might hope that the correspondence of the elements of the S-move matrix in N = 1 super Liouville theory with the eigenvalues of the l.h.s. operator could help us to prove the Moore-Seiberg condition of orthogonality (11). It is intriguing to study the relation of equation (87) with the supersymmetric Ruijsenaars-Sutherland problem [14].

There is also another application of S-move matrix in the AGT correspondence. Remember that logarithm of one-point toric block in semi-classical limit yields Seiberg-Witten prepotential of  $\mathcal{N} = 2^*$  theory [21, 23, 32]. Therefore asymptotic form of the S-move matrix of Liouville field theory implements electromagnetic duality of SW prepotential. For Liouville field theory this was studied in [23, 31, 32]. We are planning to pursue these issues for S-move matrix in N = 1 SLFT using relation between the Nekrasov partition function of  $\mathcal{N} = 2^*$  theory on  $S^4/\mathbb{Z}_2$  and N = 1 SLFT one-point toric blocks found in [15, 16].

We expect that the S-move matrix of super Liouville theory also should play important role in quantization of the super Teichmuller space [1,2] and building supersymmetric analogue of the Virasoro topological field theory studied in [17,18].

Acknowledgements. The work of Elena Apresyan was supported by Armenian SCS grants 21AG-1C024 and 20TTAT-QTa009. We would like also to thank V. P. Spiridonov for many useful discussions.

# A Properties of supersymmetric hyperbolic and double gamma functions

The function  $S_b(y)$  has the integral representation

$$S_b(y) = \exp\left(-\int_0^\infty \left(\frac{\sinh(2y-b-b^{-1})x}{2\sinh(bx)\sinh(b^{-1}x)} - \frac{2y-b-b^{-1}}{2x}\right)\right)\frac{dx}{x}, \quad (89)$$

and obeys the equations:

$$\frac{S_b(y+b)}{S_b(y)} = 2\sin\pi by, \quad \frac{S_b(y+b^{-1})}{S_b(y)} = 2\sin\frac{\pi y}{b}.$$
(90)

Let us mention for further use the following of properties of the the function  $\Lambda(y,m)$ . The function  $\Lambda(y,m)$  has the asymptotics [36,42]:

$$\lim_{y \to \infty} \Lambda(y, m) = e^{\mp \frac{i\pi}{2} \left( \frac{1}{r} B_{2,2}(y, b, b^{-1}) + \frac{m^2}{r} - m + \frac{r}{6} - \frac{1}{6r} \right)}, \quad \text{for} \quad \Im \mathfrak{m}(y) \to \pm \infty \,, \quad (91)$$

where  $B_{2,2}(y, b, b^{-1})$  is the second order Bernoulli polynomial

$$B_{2,2}(y;b,b^{-1}) = \left(y - \frac{Q}{2}\right)^2 - \frac{b^2 + b^{-2}}{12}.$$
(92)

The poles and zeros of the function  $\Lambda(y, m)$  are given by the relations [36, 43]:

$$u_{\text{poles}}^{(m)} \in \{-n_1b - n_2b^{-1}\}, \ u_{\text{zeros}}^{(m)} \in \{(n_1+1)b + (n_2+1)b^{-1}\},$$
 (93)

where  $n_1 - n_2 = m \mod r, \ n_1, n_2 \in \mathbb{Z}_{\geq 0}$ .

The function  $\Lambda(y, m)$  satisfies the relation:

$$\frac{\Lambda(y+b,m-1)}{\Lambda(y,m)} = -2\sin\frac{\pi b(y-mb^{-1})}{r}, \ \frac{\Lambda(y+b^{-1},m+1)}{\Lambda(y,m)} = 2\sin\frac{\pi(y+mb)}{rb}.$$
(94)

Specializing these properties to the case r = 2 we derive, that poles and zeros of  $S_{NS}(x)$  are given by the set

$$u_{\text{poles}}^{(0)} \in \{-n_1b - n_2b^{-1}\}, \ u_{\text{zeros}}^{(0)} \in \{(n_1+1)b + (n_2+1)b^{-1}\},$$
 (95)

where  $n_1 - n_2 = 0 \mod 2$ ,  $n_1, n_2 \in \mathbb{Z}_{\geq 0}$ , and poles and zeros of  $S_R(x)$  are given by the set

$$u_{\text{poles}}^{(1)} \in \{-n_1b - n_2b^{-1}\}, \ u_{\text{zeros}}^{(1)} \in \{(n_1+1)b + (n_2+1)b^{-1}\},$$
 (96)

where  $n_1 - n_2 = 1 \mod 2$ ,  $n_1, n_2 \in \mathbb{Z}_{\geq 0}$ .

The relations (94) for the case r = 2 take the form:

$$\frac{S_{NS}(y+b)}{S_R(y)} = -2\sin\frac{\pi b(y-b^{-1})}{2}, \quad \frac{S_{NS}(y+b^{-1})}{S_R(y)} = 2\sin\frac{\pi(y+b)}{2b}$$
(97)

and

$$\frac{S_R(y+b)}{S_{NS}(y)} = 2\sin\frac{\pi by}{2}, \quad \frac{S_R(y+b^{-1})}{S_{NS}(y)} = 2\sin\frac{\pi y}{2b}.$$
(98)

We need also relations of sypersymmetric hyperbolic gamma functions with the supersymmetric double gamma functions [26, 37].

Start with definition of  $\Gamma_b(x)$  function.

The function  $\Gamma_b(x)$  is a close relative of the double Gamma function studied in [5,44]. It can be defined by means of the integral representation

$$\log \Gamma_b(x) = \int_0^\infty \frac{dt}{t} \left( \frac{e^{-xt} - e^{-Qt/2}}{(1 - e^{-bt})(1 - e^{-t/b})} - \frac{(Q - 2x)^2}{8e^t} - \frac{Q - 2x}{t} \right).$$
(99)

Important properties of  $\Gamma_b(x)$  are

- 1. Functional equation:  $\Gamma_b(x+b) = \sqrt{2\pi} b^{bx-\frac{1}{2}} \Gamma^{-1}(bx) \Gamma_b(x).$
- 2. Analyticity:  $\Gamma_b(x)$  is meromorphic, poles:  $x = -nb mb^{-1}, n, m \in \mathbb{Z}^{\geq 0}$ .
- 3. Self-duality:  $\Gamma_b(x) = \Gamma_{1/b}(x)$ .

The function  $S_b(x)$  may be defined in terms of  $\Gamma_b(x)$  as follows

$$S_b(x) = \frac{\Gamma_b(x)}{\Gamma_b(Q-x)}.$$
(100)

Similar role in the super Liouville theory play the functions:

$$\Gamma_1(x) \equiv \Gamma_{\rm NS}(x) = \Gamma_b\left(\frac{x}{2}\right)\Gamma_b\left(\frac{x+Q}{2}\right) \,, \tag{101}$$

$$\Gamma_0(x) \equiv \Gamma_{\rm R}(x) = \Gamma_b\left(\frac{x+b}{2}\right)\Gamma_b\left(\frac{x+b^{-1}}{2}\right).$$
(102)

The sypersymmetric hyperbolic gamma functions are related to them by the formulae similar to (100)

$$S_1(x) \equiv S_{\rm NS}(x) = \frac{\Gamma_{\rm NS}(x)}{\Gamma_{\rm NS}(Q-x)}, \quad S_0(x) \equiv S_{\rm R}(x) = \frac{\Gamma_{\rm R}(x)}{\Gamma_{\rm R}(Q-x)}.$$
 (103)

#### **B** Parafermionic integral identities I

In paper [42] the rarefied hyperbolic beta integral has been evaluated<sup> $\ddagger$ </sup>:

$$\int_{-i\infty}^{i\infty} \sum_{m=0}^{r-1} \frac{\prod_{a=1}^{6} \Lambda(s_a \pm y, n_a \pm m)}{\Lambda(\pm 2y, \pm 2m)} \frac{dy}{i} = 2r \prod_{1 \le a < b \le 6} \Lambda(s_a + s_b, n_a + n_b) (104)$$

<sup>&</sup>lt;sup>‡</sup>We put periods  $\omega_1$  and  $\omega_2$  there equal  $b^{-1}$  and b correspondingly and also  $\epsilon = 0$ .

where

$$\sum_{a=1}^{6} n_a = 0, \quad \sum_{a=1}^{6} s_a = Q.$$
(105)

Here we used the shorthand notation  $\Lambda(y \pm x, m \pm n) \equiv \Lambda(y + x, m + n)\Lambda(y - x, m - n)$ .

It is shown in [42], that in certain limit (104) yields

$$\int_{-i\infty}^{i\infty} \sum_{m=0}^{r-1} (-1)^m \prod_{a=1}^3 \Lambda(y+f_a, n_a+m) \Lambda(-y+g_a, l_a-m) \frac{dy}{i}$$
(106)  
=  $(-1)^{n_1+n_2+n_3} r \prod_{a,b=1}^3 \Lambda(f_a+g_b, n_a+l_b),$ 

where

$$\sum_{a=1}^{3} (n_a + l_a) = 0, \quad \sum_{a=1}^{3} (f_a + g_a) = Q.$$
 (107)

In certain limit (106) leads to another useful identity:

$$\int_{-i\infty}^{i\infty} \sum_{m=0}^{r-1} (-)^m e^{\frac{2i\pi Nm}{r}} e^{\frac{2i\pi}{r}yt} \Lambda(f \pm y, n \pm m) \frac{dy}{i}$$
(108)  
=  $(-1)^{N-n} r \Lambda(2f, 2n) \Lambda(Q/2 - f \pm t, -n \pm N).$ 

We can show that taking the following limit in (104):

$$s_5 \to i\infty, \qquad s_6 = Q - s_5 - \sum_{j=1}^4 s_j \to -i\infty$$
 (109)

and using the asymptotics (91) one derives

$$\int_{-i\infty}^{i\infty} \sum_{m=0}^{r-1} \frac{\prod_{a=1}^{4} \Lambda(s_a \pm y, n_a \pm m)}{\Lambda(\pm 2y, \pm 2m)} \frac{dy}{i} = 2r \frac{\prod_{1 \le a < b \le 4} \Lambda(s_a + s_b, n_a + n_b)}{\Lambda(\sum_{i=1}^{4} s_i, \sum_{i=1}^{4} n_i)}$$
(110)

Taking in (110) further limit

$$s_4 \to i\infty$$
 (111)

and again using the asymptotics (91) one gets

$$\int_{-i\infty}^{i\infty} \sum_{m=0}^{r-1} \frac{\prod_{a=1}^{3} \Lambda(s_a \pm y, n_a \pm m)}{\Lambda(\pm 2y, \pm 2m)} e^{-\frac{i\pi}{r}y^2} e^{-\frac{i\pi}{r}m^2} \frac{dy}{i} = 2re^{\frac{i\pi}{r}(s_1s_2+s_2s_3+s_1s_3)} \quad (112)$$
$$\times e^{\frac{i\pi}{r}(n_1n_2+n_2n_3+n_1n_3)} \Lambda(s_1+s_2, n_1+n_2) \Lambda(s_2+s_3, n_2+n_3) \Lambda(s_1+s_3, n_1+n_3).$$

And finally taking the limit

$$s_3 \to i\infty$$
 (113)

in (112) and again using the asymptotics (91) we obtain

$$\int_{-i\infty}^{i\infty} \sum_{m=0}^{r-1} \frac{\prod_{a=1}^{2} \Lambda(s_a \pm y, n_a \pm m)}{\Lambda(\pm 2y, \pm 2m)} e^{-\frac{2i\pi}{r}y^2} e^{-\frac{2i\pi}{r}m^2} \frac{dy}{i}$$
(114)  
=  $2re^{-\frac{i\pi}{2r}(s_1 - s_2)^2} e^{\frac{i\pi}{2r}(s_1 + s_2)Q} e^{-\frac{i\pi}{2r}(n_1 - n_2)^2} e^{\frac{i\pi}{2}(n_1 + n_2)} \Lambda(s_1 + s_2, n_1 + n_2).$ 

## C Parafermionic integral identities II

In paper [4] some parafermionic hypergeometric functions were defined. The first is the parafermionic generalization of the Ruijsenaars function [40]:

$$J(\underline{\beta},\underline{l};\underline{\gamma},\underline{t}) = \int_{-i\infty}^{i\infty} \sum_{m=0}^{r-1} \prod_{a=1}^{4} \Lambda(y+\gamma_a,t_a+m)\Lambda(-y+\beta_a,l_a-m)\frac{dy}{ir} .$$
(115)

Here  $t_a, l_a \in \mathbb{Z}$ . Parameters  $\gamma_j, \beta_j$  and  $l_j, t_j$  satisfy the constraints:

$$\sum_{j=1}^{4} (\gamma_j + \beta_j) = 2Q, \quad \sum_{j=1}^{4} (l_j + t_j) = 0.$$
 (116)

The second function is :

$$E_{\delta}(s_a, n_a) = \int_{-i\infty}^{i\infty} \sum_{m=0}^{r-1} \frac{\prod_{a=1}^{6} \Lambda(y + s_a, n_a + m + \delta) \Lambda(-y + s_a, n_a - m)}{\Lambda(2y, 2m + \delta) \Lambda(-2y, -(2m + \delta))} \frac{dy}{2ir} .$$
(117)

We proved in [4] the following relation between (115) and (117):

$$J(\underline{\beta}, \underline{l}; \underline{\gamma}, \underline{t}) = e^{i\pi(t_4 + l_4)} \prod_{i=1}^{3} \Lambda(\beta_i + \gamma_4, l_i + t_4) \Lambda(\gamma_i + \beta_4, t_i + l_4) \quad (118)$$
  
 
$$\times E_{\delta}(\beta_1 + \xi, l_1 - L_1; \beta_2 + \xi, l_2 - L_1; \beta_3 + \xi, l_3 - L_1;$$
  
 
$$\gamma_1 - \xi, t_1 - L_2; \gamma_2 - \xi, t_2 - L_2; \gamma_3 - \xi, t_3 - L_2), \quad (119)$$

where

$$L_1 = \frac{1}{2}(l_1 + l_2 + l_3 + t_4 + \delta), \qquad (120)$$

$$L_2 = \frac{1}{2}(t_1 + t_2 + t_3 + l_4 + \delta), \qquad (121)$$

$$\xi = \frac{1}{2}(Q - \gamma_4 - \sum_{j=1}^3 \beta_j).$$
(122)

The parameter  $\delta = 0, 1$  should be determined from the requirement that  $L_1$  and  $L_2$  are integer numbers. Let us now paparemeterize parameters  $\underline{\beta}, \underline{\gamma}$  in (118) in the following way:

$$\begin{aligned} \gamma_1 &= A - \alpha_1 + Q/2 , & \gamma_4 &= -A - \alpha_2 + \alpha_1 , \\ \gamma_2 &= \beta_2 &= \alpha_2/2 - \alpha_3 + Q/4 , & \beta_1 &= A + \alpha_1 + Q/2 , \\ \gamma_3 &= \beta_3 &= \alpha_2/2 + \alpha_3 + Q/4 , & \beta_4 &= -A - \alpha_2 - \alpha_1 . \end{aligned}$$
(123)

Obviously parametrized in this way  $\underline{\beta}$  and  $\underline{\gamma}$  satisfy (116).

Now putting (123) in (118) and taking the limit  $A \to \infty$ , denoting  $l_1 + t_4 = T$ , and at the end renaming indices  $2, 3 \to 1, 2$  for all variables, what we can do since variables with indices 1, 4 disappear, we obtain the identity

$$\begin{split} &\int_{-i\infty}^{i\infty} \sum_{m=0}^{r-1} \prod_{a=1}^{2} \Lambda(y+\gamma_{a},t_{a}+m)\Lambda(-y+\gamma_{a},l_{a}-m)e^{\frac{4i}{r}\pi y\alpha_{1}}e^{\frac{i\pi m}{r}(2T+\sum_{a=1}^{2}(t_{a}+l_{a}))}dy = (124) \\ &\Lambda(2\alpha_{1}-\gamma_{1}-\gamma_{2}+Q,T)\Lambda(-2\alpha_{1}-\gamma_{1}-\gamma_{2}+Q,-T-\sum_{a=1}^{2}(t_{a}+l_{a}))e^{-i\pi\left[\sum_{a=1}^{2}(t_{a}+l_{a})+\frac{2}{r}\left(\gamma_{1}\gamma_{2}-\alpha_{1}^{2}\right)\right]} \\ &\times \int_{-i\infty}^{i\infty} \sum_{m=0}^{r-1} \frac{\prod_{a=1}^{2}\Lambda(y+\gamma_{a}-\alpha_{1},l_{a}-M_{1}+m+\delta)\Lambda(-y+\gamma_{a}-\alpha_{1},l_{a}-M_{1}-m)}{\Lambda(2y,2m+\delta)\Lambda(-2y,-(2m+\delta))} \\ &\times \prod_{a=1}^{2}\Lambda(y+\gamma_{a}+\alpha_{1},t_{a}-M_{2}+m+\delta)\Lambda(-y+\gamma_{a}+\alpha_{1},t_{a}-M_{2}-m)e^{-\frac{2i\pi}{r}y^{2}}e^{-\frac{2i\pi}{r}(m^{2}+m\delta)}\frac{dy}{2} \\ &\times e^{\frac{i\pi}{2r}\left[l_{1}^{2}+l_{2}^{2}+t_{1}^{2}+t_{2}^{2}-2M_{1}^{2}-2M_{2}^{2}+2(T+l_{1}+l_{2})^{2}-(l_{1}+l_{2})^{2}\right]}, \end{split}$$

where

$$M_1 = \frac{1}{2}(T + l_1 + l_2 + \delta), \qquad (125)$$

$$M_2 = \delta - M_1 \,, \tag{126}$$

and  $\delta = 0, 1$  is determined from the requirement that  $M_1$  and therefore also  $M_2$  are integer. This formula for r = 1 case can be found in [48,52].

#### D Parafermionic integral identities III

Now let us set in (118)

$$\beta_4 = -\frac{\lambda}{3} - 2A + Q/2, \quad \gamma_1 = -\frac{\lambda}{3} + A + Q/2, \quad \gamma_2 = -\frac{\lambda}{3} + A + Q/2. \quad (127)$$

Taking the limit  $A \to \infty$ , we obtain, after renaming at the end the variables  $\gamma_3, \gamma_4, t_3, t_4 \to \gamma_1, \gamma_2, t_1, t_2$ , the identity

$$\begin{split} &\int_{-i\infty}^{i\infty} \sum_{m=0}^{r-1} \prod_{a=1}^{2} \Lambda(y+\gamma_{a},t_{a}+m) \prod_{a=1}^{3} \Lambda(-y+\beta_{a},l_{a}-m) e^{\frac{i\pi}{r}(\lambda y-\frac{y^{2}}{2})} e^{\frac{i\pi}{r} \left[m(t_{1}+t_{2}+\sum_{i=1}^{3}l_{i}+\frac{3r}{2})-\frac{m^{2}}{2}\right]} dy \quad (128) \\ &= \int_{-i\infty}^{i\infty} \sum_{m=0}^{r-1} \frac{\prod_{a=1}^{3} \Lambda(y+\beta_{a}+\eta,l_{a}-L_{1}+m+\delta)\Lambda(-y+\beta_{a}+\eta,l_{a}-L_{1}-m)}{\Lambda(2y,2m+\delta)\Lambda(-2y,-(2m+\delta))} \\ &\times \Lambda(y+\gamma_{1}-\eta,t_{1}-L_{2}+m+\delta)\Lambda(-y+\gamma_{1}-\eta,t_{1}-L_{2}-m) e^{-\frac{2i\pi}{r}y^{2}} e^{-\frac{2i\pi}{r}(m^{2}+m\delta)} \frac{dy}{2} \\ &\times e^{\frac{i\pi}{2r}(\lambda^{2}+Q^{2}/4-Q(\lambda+\gamma_{1}))} e^{\frac{i\pi}{2r} \left[4L_{2}^{2}-4L_{2}(t_{1}+\delta)+2\delta t_{1}+t_{1}^{2}\right]} e^{\frac{i\pi}{2} \left[-t_{1}+2(l_{1}+l_{2}+l_{3})\right]} \prod_{a=1}^{3} \Lambda(\beta_{a}+\gamma_{2};l_{a}+t_{2}) \,, \end{split}$$

where

$$\sum_{i=1}^{2} \gamma_i + \sum_{i=1}^{3} \beta_i = Q/2 + \lambda, \qquad (129)$$

and

$$2\eta = Q - \gamma_2 - \sum_{i=1}^{3} \beta_i = Q/2 - \lambda + \gamma_1, \qquad (130)$$

$$L_1 = \frac{1}{2}(l_1 + l_2 + l_3 + t_2 + \delta), \qquad (131)$$

$$L_2 = \delta - L_1 \,, \tag{132}$$

and  $\delta = 0, 1$  is determined from the requirement that  $L_1$  and therefore also  $L_2$  are integer. For r = 1 case this formula can be found in [52].

## **E** Parafermionic integral identities IV

We proved also in [4] that (115) satisfies the transformation property

$$J(\underline{\beta}, \underline{l}; \underline{\gamma}, \underline{t}) = e^{i\pi[t_3 + t_4 + l_1 + l_2]} J(\underline{\tilde{\beta}}, \underline{\tilde{l}}; \underline{\tilde{\gamma}}, \underline{\tilde{t}}) \times$$

$$\prod_{j,k=1}^{2} \Lambda(\gamma_j + \beta_k, t_j + l_k) \prod_{j,k=3}^{4} \Lambda(\gamma_j + \beta_k, t_j + l_k),$$
(133)

$$\begin{split} \tilde{\beta}_{a} &= \beta_{a} + \Theta(a)\eta \,, \quad \tilde{l}_{a} = l_{a} - (\Theta(a) + 1)K \,, \quad a = 1, 2, 3, 4 \,, \quad (134) \\ \tilde{\gamma}_{a} &= \gamma_{a} + \Theta(a)\eta \,, \quad \tilde{t}_{a} = t_{a} - (\Theta(a) - 1)K \,, \quad a = 1, 2, 3, 4 \,, \end{split}$$

where  $\Theta(a)$  is the sign function taking the values

$$\Theta(a) = 1, \quad a = 1, 2 \quad \text{and} \quad \Theta(a) = -1, \quad a = 3, 4,$$
 (135)

and

$$K = \frac{1}{2}(t_1 + t_2 + l_1 + l_2) = -\frac{1}{2}(t_3 + t_4 + l_3 + l_4), \qquad (136)$$

$$\eta = \frac{1}{2}(Q - \gamma_1 - \gamma_2 - \beta_1 - \beta_2).$$
(137)

Setting in (133):

$$\gamma_1 = A - \alpha_1 + Q/2, \quad \beta_1 = A + \alpha_1 + Q/2, \quad (138)$$
  
$$\gamma_2 = -A - \alpha_2 + \alpha_1 + Q/2, \quad \beta_2 = -A - \alpha_1 - \alpha_2 + Q/2,$$

$$\gamma_3 = \beta_3 = \alpha_2/2 - \alpha_3, \quad \gamma_4 = \beta_4 = \alpha_2/2 + \alpha_3,$$

and denoting

$$N = \frac{1}{2}(t_2 + l_1 - t_1 - l_2), \qquad (139)$$

we obtain in the limit  $A \to \infty$ 

$$\sum_{m \in \mathbb{Z}_{r}} \int_{-i\infty}^{i\infty} \Lambda(y + \alpha_{2}/2 + \alpha_{3}, t_{4} + m)\Lambda(-y + \alpha_{2}/2 + \alpha_{3}, l_{4} - m)$$
(140)  
  $\times \Lambda(y + \alpha_{2}/2 - \alpha_{3}, t_{3} + m)\Lambda(-y + \alpha_{2}/2 - \alpha_{3}, l_{3} - m)e^{\frac{4i\pi}{r}y\alpha_{1}}e^{\frac{2miN\pi}{r}}\frac{dy}{i} = e^{\frac{2iNK\pi}{r}}e^{i\pi(t_{3}+t_{4}-N)}\Lambda(Q - 2\alpha_{1} - \alpha_{2}, K - N)\Lambda(Q + 2\alpha_{1} - \alpha_{2}, K + N)$   
  $\times \Lambda(\alpha_{2} - 2\alpha_{3}, t_{3} + l_{3})\Lambda(\alpha_{2} + 2\alpha_{3}, t_{4} + l_{4})\Lambda(\alpha_{2}, t_{3} + l_{4})\Lambda(\alpha_{2}, t_{4} + l_{3})$   
  $\times \sum_{m \in \mathbb{Z}_{r}}\int_{-i\infty}^{i\infty}\Lambda(y + (Q - \alpha_{2})/2 + \alpha_{3}, t_{4} + 2K + m)\Lambda(-y + (Q - \alpha_{2})/2 + \alpha_{3}, l_{4} - m)$   
  $\times \Lambda(y + (Q - \alpha_{2})/2 - \alpha_{3}, t_{3} + 2K + m)\Lambda(-y + (Q - \alpha_{2})/2 - \alpha_{3}, l_{3} - m)e^{\frac{4i\pi}{r}y\alpha_{1}}e^{\frac{2miN\pi}{r}}\frac{dy}{i},$ 

where  $K + N \in \mathbb{Z}$  and  $K - N \in \mathbb{Z}$ .

#### References

- N. Aghaei, M. Pawelkiewicz and J. Teschner, "Quantisation of super Teichmüller theory," Commun. Math. Phys. 353 (2017) no.2, 597-631
- [2] N. Aghaei, M. K. Pawelkiewicz and M. Yamazaki, "Towards Super Teichmüller Spin TQFT," Adv. Theor. Math. Phys. 26 (2022) no.2, 245-294
- [3] L. F. Alday, D. Gaiotto and Y. Tachikawa, "Liouville Correlation Functions from Four-dimensional Gauge Theories," Lett. Math. Phys. 91 (2010), 167-197
- [4] E. Apresyan, G. Sarkissian and V. P. Spiridonov, "A parafermionic hypergeometric function and supersymmetric 6j-symbols," Nucl. Phys. B 990 (2023), 116170.
- [5] E. W. Barnes, "Theory of the double gamma function", Phil. Trans. Roy. Soc A196 (1901) 265-388
- [6] R. E. Behrend, P. A. Pearce, V. B. Petkova and J. B. Zuber, "Boundary conditions in rational conformal field theories," Nucl. Phys. B 570 (2000) 525 [Nucl. Phys. B 579 (2000) 707] [arXiv:hep-th/9908036].
- [7] A. Belavin, V. Belavin, A. Neveu and A. Zamolodchikov, "Bootstrap in Supersymmetric Liouville Field Theory. I. NS Sector," Nucl. Phys. B 784 (2007), 202-233
- [8] V. Belavin and B. Feigin, "Super Liouville conformal blocks from N=2 SU(2) quiver gauge theories," JHEP 07 (2011), 079

- [9] A. Belavin and B. Mukhametzhanov, N=1 superconformal blocks with Ramond fields from AGT correspondence, JHEP 01 (2013) 178.
- [10] A. Belavin, V. Belavin, and M. Bershtein, Instantons and 2d Superconformal field theory, JHEP 09 (2011) 117.
- [11] N. Belousov, S. Derkachov, S. Kharchev and S. Khoroshkin, "Baxter operators in Ruijsenaars hyperbolic system I. Commutativity of Q-operators," [arXiv:2303.06383 [math-ph]].
- [12] N. Belousov, S. Derkachov, S. Kharchev and S. Khoroshkin, "Baxter Qoperators in Ruijsenaars–Sutherland hyperbolic systems: one- and twoparticle cases," Zap. Nauchn. Semin. **520** (2023) no.0, 50-123
- [13] N. Belousov, S. Derkachov, S. Kharchev and S. Khoroshkin, "Baxter operators in Ruijsenaars hyperbolic system III. Orthogonality and completeness of wave functions," [arXiv:2307.16817 [math-ph]].
- [14] O. Blondeau-Fournier, P. Desrosiers and P. Mathieu, "Supersymmetric Ruijsenaars-Schneider Model," Phys. Rev. Lett. **114** (2015), 121602
- [15] G. Bonelli, K. Maruyoshi and A. Tanzini, "Instantons on ALE spaces and Super Liouville Conformal Field Theories," JHEP **1108** (2011) 056
- [16] G. Bonelli, K. Maruyoshi and A. Tanzini, "Gauge Theories on ALE Space and Super Liouville Correlation Functions," Lett. Math. Phys. 101 (2012) 103
- [17] S. Collier, L. Eberhardt and M. Zhang, "Solving 3d gravity with Virasoro TQFT," SciPost Phys. 15 (2023) no.4, 151
- [18] S. Collier, L. Eberhardt and M. Zhang, "3d gravity from Virasoro TQFT: Holography, wormholes and knots," [arXiv:2401.13900 [hep-th]].
- [19] P. Degiovanni, "Moore and Seiberg's equations and 3D topological field theory", Comm. Math. Phys. 145(3), 459-505 (1992).
- [20] N. Drukker, D. Gaiotto and J. Gomis, "The Virtue of Defects in 4D Gauge Theories and 2D CFTs," JHEP 06 (2011), 025
- [21] V. A. Fateev and A. V. Litvinov, "On AGT conjecture," JHEP 02 (2010), 014
- [22] S. Elitzur, G. W. Moore, A. Schwimmer and N. Seiberg, "Remarks on the Canonical Quantization of the Chern-Simons-Witten Theory," Nucl. Phys. B 326 (1989), 108-134

- [23] D. Galakhov, A. Mironov and A. Morozov, "S-duality as a beta-deformed Fourier transform," JHEP 08 (2012), 067
- [24] D. Galakhov, A. Mironov and A. Morozov, "S-Duality and Modular Transformation as a non-perturbative deformation of the ordinary pq-duality," JHEP 06 (2014), 050
- [25] L. Hadasz, "On the fusion matrix of the N=1 Neveu-Schwarz blocks," JHEP 0712 (2007) 071.
- [26] L. Hadasz, M. Pawelkiewicz and V. Schomerus, "Self-dual Continuous Series of Representations for  $U_q(sl(2))$  and  $U_q(osp(1|2))$ ," JHEP **1410** (2014) 91.
- [27] N. Hama, K. Hosomichi and S. Lee, "Notes on SUSY Gauge Theories on Three-Sphere," JHEP 03 (2011), 127
- [28] K. Hosomichi, S. Lee and J. Park, "AGT on the S-duality Wall," JHEP 12 (2010), 079
- [29] K. Hosomichi, "Bulk boundary propagator in Liouville theory on a disc," JHEP 11 (2001), 044
- [30] Y. Imamura and D. Yokoyama, " $S^3/Z_n$  partition function and dualities," JHEP **11** (2012), 122
- [31] A. K. Kashani-Poor and J. Troost, "Quantum geometry from the toroidal block," JHEP 08 (2014), 117
- [32] A. Marshakov, A. Mironov and A. Morozov, "On AGT Relations with Surface Operator Insertion and Stationary Limit of Beta-Ensembles," J. Geom. Phys. 61 (2011), 1203-1222
- [33] G. W. Moore and N. Seiberg, "Lectures on RCFT," Published in Trieste Superstrings 1989:1-129. Also in Banff NATO ASI 1989:263-362.
- [34] G. W. Moore and N. Seiberg, "Classical and Quantum Conformal Field Theory," Commun. Math. Phys. 123 (1989) 177.
- [35] N. Nemkov, "On modular transformations of toric conformal blocks," JHEP 10 (2015), 039.
- [36] F. Nieri and S. Pasquetti, Factorisation and holomorphic blocks in 4d, JHEP 1511 (2015) 155

- [37] H. Poghosyan and G. Sarkissian, "Comments on fusion matrix in N=1 super Liouville field theory," Nucl. Phys. B 909 (2016) 458.
- [38] E. M. Rains, Limits of elliptic hypergeometric integrals, Ramanujan J. 18:3 (2009), 257–306, arXiv:math/0607093 [math.CA].
- [39] S. N. M. Ruijsenaars, First order analytic difference equations and integrable quantum systems, J. Math. Phys. 38 (1997), 1069–1146.
- [40] S. N. M. Ruijsenaars, Systems of Calogero-Moser type, in: Proceedings of the 1994 Banff summer school "Particles and fields", CRM series in mathematical physics, (G. Semenoff, L. Vinet, Eds.), pp. 251–352, Springer, New York, 1999.
- [41] I. Runkel, "Boundary structure constants for the A series Virasoro minimal models," Nucl. Phys. B 549 (1999), 563-578
- [42] G. Sarkissian and V. P. Spiridonov, "From rarefied elliptic beta integral to parafermionic star-triangle relation," JHEP 10 (2018), 097.
- [43] G. Sarkissian and V. P. Spiridonov, "General modular quantum dilogarithm and beta integrals," Proc. Steklov Inst. Math. 309, 251–270 (2020).
- [44] T. Shintani, "On a Kronecker limit formula for real quadratic fields", J. Fac. Sci. Univ. Tokyo Sect. 1A Math. 24 (1977) 167-199
- [45] V. P. Spiridonov, On the elliptic beta function, Uspekhi Mat. Nauk 56:1 (2001), 181–182 (Russ. Math. Surveys 56:1 (2001), 185–186).
- [46] V. P. Spiridonov, Theta hypergeometric integrals, St. Petersburg Math. J. 15 (2004), 929–967, arXiv:math.CA/0303205.
- [47] V. P. Spiridonov, Essays on the theory of elliptic hypergeometric functions, Russian Math. Surveys 63 (2008), 405–472, arXiv:0805.3135.
- [48] V. P. Spiridonov and G. S. Vartanov, "Elliptic hypergeometry of supersymmetric dualities II. Orthogonal groups, knots, and vortices," Commun. Math. Phys. **325** (2014), 421-486
- [49] V. P. Spiridonov, "Rarefied elliptic hypergeometric functions," Adv. Math. 331 (2018) 830.
- [50] J. Teschner, "From Liouville theory to the quantum geometry of Riemann surfaces," in 14th International Congress on Mathematical Physics, 8, 2003, [arXiv:hep-th/0308031 [hep-th]].

- [51] J. Teschner, "An Analog of a modular functor from quantized teichmuller theory," [arXiv:math/0510174 [math.QA]].
- [52] J. Teschner and G. S. Vartanov, "Supersymmetric gauge theories, quantization of *M*<sub>flat</sub>, and conformal field theory," Adv. Theor. Math. Phys. **19** (2015) 1.
- [53] H. L. Verlinde, "Conformal Field Theory, 2-D Quantum Gravity and Quantization of Teichmuller Space," Nucl. Phys. B 337 (1990), 652-680
- [54] E. Witten, "Quantum Field Theory and the Jones Polynomial," Commun. Math. Phys. **121** (1989), 351-399