

Gravitational radiations of Kerr black hole from warped symmetries

Jianfei Xu

Shing-Tung Yau Center and School of Mathematics, Southeast University, Nanjing, 210000, China

E-mail: jfxu@seu.edu.cn

ABSTRACT: The warped conformal symmetries have been found in the covariant phase space of a set of non-trivial diffeomorphism near the Kerr black hole horizon. In this paper, we consider the retarded Green's function and the absorption probability for the scalar and higher spin perturbations on a generic non-extreme Kerr black hole background, and perform their holographic calculations in terms of the warped CFT. This provide further evidence on the conjecture that the warped CFT is relevant to the microstate description of the Kerr black hole.

Contents

1	Introduction	1
2	Review of Kerr spacetime and hidden conformal symmetry	3
3	Scalar radiations in Kerr spacetime	5
3.1	Near region	5
3.2	Far region	6
3.3	Near-far matching	7
3.4	Dual calculations in WCFT	8
4	Higher spin radiations in Kerr spacetime	12
4.1	Near region	13
4.2	Far region	14
4.3	Near-far matching	14
4.4	Dual explanations in WCFT	15
5	Summary and discussion	16

1 Introduction

In exploring the gravitational phenomena in the vicinity of a black hole, the conformal field theory (CFT) plays an essential role. This was not only due to the holographic duality (anti de-Sitter (AdS))/CFT, but also due to the large amount of studying go beyond it. One of the typical example is the Kerr/CFT [1] which aim to understand the Kerr black hole in the universe in terms of a two dimensional (2D) CFT. For a extreme Kerr black hole, there is a near horizon scaling region, know as the NHEK (near horizon extreme Kerr) geometry [2], where the asymptotic symmetry analysis under specific boundary conditions shows that the dual field theory should be a chiral left-moving 2D CFT [1]. Other boundary conditions that support the extreme Kerr black hole dual to a chiral 2D CFT can be found in [3, 4]. Evidence of the Kerr/CFT come from the facts that the Cardy formula of a 2D CFT reproduces the black hole entropy [1, 3] and the matching of the CFT correlators with the scattering amplitudes of the near-extreme Kerr black hole [5].

For non-extreme Kerr black holes, the NHEK geometry disappears and the near horizon geometry is just Rindler space without conformal symmetry. However, as proposed in [6], a geometry with conformal symmetry is not a necessary condition for the interactions to exhibit conformal invariance. The conformal symmetries may hide in the solution space and still govern the dynamics of the fields. In the low frequency limit $\omega \ll 1/M$, where ω is the frequency of the propagating field and M is the mass of the Kerr black hole,

the spacetime can be divided into the near region $r \ll 1/\omega$ and far region $r \gg M$ which have a large overlap. The hidden conformal symmetry acting on the near region wave equation. Intuitively, this is highly because of the independence of choosing a matching surface in getting the full wave solution by near-far matching [6]. This hidden conformal symmetry can help to fix the black hole entropy and scattering amplitudes in term of Cardy formula and correlators in 2D CFT [6]. Also the covariant phase space formalism [7] and gravitational wave analysis [8] support the validity of the hidden conformal symmetry.

In this paper, we will phenomenologically determine the scattering amplitudes of a non-extreme Kerr black hole by incident perturbative scalar and higher spin waves originating in the asymptotically flat region, and see whether or not the black hole will react like a warped CFT (WCFT) other than 2D CFT. This will generalise the work [5] to the non-extreme case. The WCFT is a different kind of field theory which also allows holographic interpretations. To be specific, it is a two dimensional nonrelativistic quantum field theory with warped conformal symmetry. The symmetry algebra contains one Virasoro algebra and one $U(1)$ Kac-Moody algebra [9, 10]. The global warped conformal symmetry is $SL(2, R) \times U(1)$. Concrete examples of WCFTs include chiral Liouville gravity [11], Weyl fermion models [12, 13], free scalars [14], and the complex Sachdev-Ye-Kitaev models as a symmetry-broken phase [15]. Much like its CFT cousin, the two and three point correlation functions of the WCFT can be fixed by the global warped conformal symmetry up to normalizations and OPE coefficients [16]. The entanglement entropy and the outer-of-time-order correlators of the WCFT have been studied in [17, 18].

From a holographic perspective, WCFT was initiated from the studying of the holography of a large class of geometries with $SL(2, R) \times U(1)$ isometry. Typical examples are the warped AdS_3 (WAdS) [19–25] and the NHEK geometry [2]. By imposing Dirichlet-Neumann boundary conditions on AdS_3 in Einstein gravity, WCFT can be viewed as an alternative holographic dual to the AdS_3 [26]. The corresponding conjectures WAdS/WCFT [10] and AdS_3 /WCFT [26] are possible holographic dualities beyond the standard AdS /CFT. Evidence of the above mentioned dualities are gathered from the following results. A Cardy like formula induced from the modular properties of WCFT partition functions recovers the thermal entropy of the WAdS black holes [10]. The WCFT thermal retarded correlators matches the scattering amplitudes in the thermal WAdS [16]. The holographic entanglement entropy [27–32], the mutual information [33], and the reflected entropy [34] of WCFT were realised in the AdS_3 or the WAdS. A holographic calculation of the asymptotic growth of three point coefficient in WCFT was provided on a BTZ black hole background [35]. See [17, 36] for another kind of holographic realization of WCFT in the lower spin Chern-Simons theory.

When the Kerr black hole is in or near extreme, the near horizon geometry with $SL(2, R) \times U(1)$ isometry at fixed polar angle can be viewed as WAdS [1, 5]. As mentioned above, the WAdS is conjectured to be dual to the WCFT under Dirichlet-Neumann boundary conditions [21]. So it is natural to believe that other than the Kerr/CFT, the WCFT could be relevant to the microstate description of the Kerr black hole. When the Kerr black hole is away from extreme, the warped symmetry is not manifest in the geometry. However, in the covariant phase space, by a specific choice of vector fields, the gravitational

charges associate to these vector fields can be shown to satisfy the warped conformal algebra with non-trivial central extensions [37]. This result will support the WCFT as a possible dual field theory to the Kerr black hole in the non-extreme case. Along this line, we will further investigate the scattering amplitudes in a generic Kerr spacetime, and explain the corresponding results in terms of the WCFT thermal correlators.

This paper is organized as follows. Section 2 is a review of the properties of the Kerr black hole as well as the corresponding hidden conformal symmetry. In section 3, the massless scalar field solutions are discussed in the near and far regions respectively. By imposing the boundary conditions and the near-far matching, the Kerr black hole absorption probability of the scalar flux is calculated at infinity. The dual calculations of the absorption probability as well as the retarded Green's function are further performed in the WCFT. In section 4, we generalise the analysis to the higher spin perturbations on the Kerr black hole background. Section 5 is for summary and discussion.

2 Review of Kerr spacetime and hidden conformal symmetry

In the Boyer-Lindquist coordinates, the Kerr black hole metric can be written as

$$ds^2 = -\frac{\Delta}{\rho^2}(dt - a \sin^2 \theta d\phi)^2 + \frac{\sin^2 \theta}{\rho^2} ((r^2 + a^2)d\phi - a dt)^2 + \frac{\rho^2}{\Delta} dr^2 + \rho^2 d\theta^2, \quad (2.1)$$

where

$$\Delta = r^2 - 2Mr + a^2, \quad \rho^2 = r^2 + a^2 \cos^2 \theta, \quad (2.2)$$

M is the black hole mass, and a is a constant with $J = Ma$ being the angular momentum of the black hole. The outer and the inner horizons are located at

$$r_{\pm} = M \pm \sqrt{M^2 - a^2}. \quad (2.3)$$

Consider a massless scalar field Φ on the Kerr black hole background. The Klein-Gordon equation is

$$\frac{1}{\sqrt{-g}} \partial_{\mu} (\sqrt{-g} g^{\mu\nu} \partial_{\nu} \Phi) = 0. \quad (2.4)$$

Note that both ∂_t and ∂_{ϕ} are Killing vectors, the scalar field can be expanded by the following eigenmodes

$$\Phi(t, r, \theta, \phi) = e^{-i\omega t + im\phi} S(\theta) R(r). \quad (2.5)$$

Substituting into the eigenmodes, the Klein-Gordon equation splits into

$$\left[\frac{1}{\sin \theta} \partial_{\theta} (\sin \theta \partial_{\theta}) - \frac{m^2}{\sin^2 \theta} + \omega^2 a^2 \cos^2 \theta \right] S(\theta) = -K_{\ell} S(\theta), \quad (2.6)$$

and

$$\left[\frac{d}{dr} \Delta \frac{d}{dr} + \frac{(2Mr_+ \omega - am)^2}{(r - r_+)(r_+ - r_-)} - \frac{(2Mr_- \omega - am)^2}{(r - r_-)(r_+ - r_-)} + (r^2 + 2M(r + 2M))\omega^2 \right] R(r) = K_{\ell} R(r). \quad (2.7)$$

The above equations can be solved by Heun functions with K_ℓ be the separation constant which are the eigenvalues on a sphere.

We will consider the case in which the wave frequency is much less than the inverse of the black hole mass

$$\omega \ll \frac{1}{M}. \quad (2.8)$$

The region $r \ll \frac{1}{\omega}$ is called the near region, and the region $r \gg M$ is called the far region. When the frequency ω is small, the near region can extend to the far region with a large overlap. The region $M \ll r \ll \frac{1}{\omega}$ is called the matching region. The hidden conformal symmetry shows up in the solution space of the wave equation for the propagating field in the near region [6].

In the near region, the angular equation with $\omega a < \omega M \ll 1$ reduce to the standard Laplacian on the 2-sphere

$$\left[\frac{1}{\sin \theta} \partial_\theta (\sin \theta \partial_\theta) - \frac{m^2}{\sin^2 \theta} \right] S(\theta) = -\ell(\ell - 1)S(\theta), \quad \ell = -m + 1, \dots, m + 1. \quad (2.9)$$

while the radial wave equation is

$$\left[\frac{d}{dr} \Delta \frac{d}{dr} + \frac{(2Mr_+ \omega - am)^2}{(r - r_+)(r_+ - r_-)} - \frac{(2Mr_- \omega - am)^2}{(r - r_-)(r_+ - r_-)} \right] R(r) = \ell(\ell - 1)R(r). \quad (2.10)$$

To see the structure of the wave equation more clearly, it is convenient to define the ‘‘conformal’’ coordinates [6]

$$\begin{aligned} \omega^+ &= \sqrt{\frac{r - r_+}{r - r_-}} e^{2\pi T_R \phi}, \\ \omega^- &= \sqrt{\frac{r - r_+}{r - r_-}} e^{2\pi T_L \phi - \frac{t}{2M}}, \\ y &= \sqrt{\frac{r_+ - r_-}{r - r_-}} e^{\pi(T_L + T_R)\phi - \frac{t}{4M}}, \end{aligned} \quad (2.11)$$

where

$$T_R = \frac{r_+ - r_-}{4\pi a}, \quad T_L = \frac{r_+ + r_-}{4\pi a}. \quad (2.12)$$

Define the local vector fields

$$\begin{aligned} H_1 &= i\partial_+, \\ H_0 &= i(\omega^+ \partial_+ + \frac{1}{2}y\partial_y), \\ H_{-1} &= i(\omega^{+2} \partial_+ + \omega^+ y \partial_y - y^2 \partial_-), \end{aligned} \quad (2.13)$$

and

$$\begin{aligned} \bar{H}_1 &= i\partial_-, \\ \bar{H}_0 &= i(\omega^- \partial_- + \frac{1}{2}y\partial_y), \\ \bar{H}_{-1} &= i(\omega^{-2} \partial_- + \omega^- y \partial_y - y^2 \partial_+). \end{aligned} \quad (2.14)$$

The above vector fields form the $SL(2, R)$ Lie algebra

$$[H_0, H_{\pm 1}] = \mp i H_{\pm 1}, \quad [H_{-1}, H_1] = -2i H_0. \quad (2.15)$$

A similar relation holds for $(\bar{H}_0, \bar{H}_{\pm 1})$. The corresponding $SL(2, R)$ quadratic Casimir operator is

$$\mathcal{H}^2 = -H_0^2 + \frac{1}{2}(H_1 H_{-1} + H_{-1} H_1). \quad (2.16)$$

One can check that

$$\mathcal{H}^2 = \partial_r \Delta \partial_r - \frac{(2Mr_+ \partial_t + a \partial_\phi)^2}{(r - r_+)(r_+ - r_-)} + \frac{(2Mr_- \partial_t + a \partial_\phi)^2}{(r - r_-)(r_+ - r_-)}. \quad (2.17)$$

So in the near region, the scalar field with fixed θ coordinate satisfies the following equation

$$\mathcal{H}^2 \Phi = \ell(\ell - 1) \Phi, \quad (2.18)$$

which indicate that the Φ is within the global conformal representation of the $SL(2, R)$ with conformal weight being ℓ . This equation have both right and left copy from $(H_0, H_{\pm 1})$ and $(\bar{H}_0, \bar{H}_{\pm 1})$. The $SL(2, R) \times SL(2, R)$ symmetry is spontaneously broken by the angular identification of the ϕ coordinate, this requires the corresponding 2D CFT at temperature

$$T_R = \frac{r_+ - r_-}{4\pi a}, \quad T_L = \frac{r_+ + r_-}{4\pi a}. \quad (2.19)$$

The above statement is supported by the black hole entropy, scattering amplitudes [6], the covariant phase space formalism [7] and the gravitational radiations [8].

3 Scalar radiations in Kerr spacetime

3.1 Near region

As mentioned in [37], the potential term in the near region radial wave equation (2.7) can be approximately written with additional $4M^2\omega^2$ term due to the $r\omega$ terms have negligible contributions in the frequency space. Take this into account, the near region radial wave equation becomes

$$\left[\frac{d}{dr} \Delta \frac{d}{dr} + \frac{(2Mr_+ \omega - am)^2}{(r - r_+)(r_+ - r_-)} - \frac{(2Mr_- \omega - am)^2}{(r - r_-)(r_+ - r_-)} \right] R(r) = \ell'(\ell' - 1)R(r), \quad (3.1)$$

where

$$\ell' = \frac{1}{2} + \frac{1}{2} \sqrt{(2\ell - 1)^2 - 16M^2\omega^2}. \quad (3.2)$$

The $SL(2, R)$ Casimir equation still holds

$$\mathcal{H}^2 \Phi = \ell'(\ell' - 1) \Phi, \quad (3.3)$$

but now with a frequency dependent conformal weight ℓ' . This also happens in the story of the WAdS [16], where the bulk Green's function is interpreted by the WCFT two point function. In the present paper, we are going to phenomenologically determine the scattering

amplitudes in the Kerr spacetime in terms of WCFT correlators. We will keep using ℓ' instead of ℓ in the following discussion.

In the near region, the radial wave equation (3.1) can be solved by the hypergeometric functions. Imposing an ingoing boundary condition at the horizon, the solution can be written as

$$R(r) = \left(\frac{r-r_+}{r-r_-} \right)^{-i \frac{2Mr_+\omega-am}{r_+-r_-}} (r-r_-)^{-\ell'} \times {}_2F_1 \left(\ell' - i \frac{4M^2\omega - 2am}{r_+-r_-}, \ell' - i2M\omega; 1 - i \frac{4Mr_+\omega - 2am}{r_+-r_-}; \frac{r-r_+}{r-r_-} \right), \quad (3.4)$$

where ${}_2F_1(a, b; c; z)$ is the hypergeometric function. When the frequency ω is very small, the near region $r \ll \frac{1}{\omega}$ extends to the far region where the asymptotic behavior of the scalar field can be analysed. The near region radial wave solution in the far region behaves as

$$R(r \gg M) \sim A_{near} r^{-\ell'} + B_{near} (r_+ - r_-)^{-2\ell'+1} r^{\ell'-1}, \quad (3.5)$$

where

$$A_{near} = \frac{\Gamma \left(1 - i \frac{4Mr_+\omega - 2am}{r_+-r_-} \right) \Gamma(-2\ell' + 1)}{\Gamma(1 - \ell' - i2M\omega) \Gamma \left(1 - \ell' - i \frac{4M^2\omega - 2am}{r_+-r_-} \right)}, \quad (3.6)$$

$$B_{near} = \frac{\Gamma \left(1 - i \frac{4Mr_+\omega - 2am}{r_+-r_-} \right) \Gamma(2\ell' - 1)}{\Gamma(\ell' - i2M\omega) \Gamma \left(\ell' - i \frac{4M^2\omega - 2am}{r_+-r_-} \right)}. \quad (3.7)$$

The retarded Green's function for the scalar field Φ in the frequency space can be read off from the asymptotic expansion

$$G_R^{bulk} \sim \frac{A_{near}}{B_{near}} = \frac{\Gamma(-2\ell' + 1)}{\Gamma(2\ell' - 1)} \frac{\Gamma(\ell' - i2M\omega)}{\Gamma(1 - \ell' - i2M\omega)} \frac{\Gamma \left(\ell' - i \frac{4M^2\omega - 2am}{r_+-r_-} \right)}{\Gamma \left(1 - \ell' - i \frac{4M^2\omega - 2am}{r_+-r_-} \right)}, \quad (3.8)$$

here \sim stands for equaling up to a factor independent of ω and m .

3.2 Far region

In the far region, the radial wave equation reduces to

$$\frac{d}{dr} \left(r^2 \frac{dR(r)}{dr} \right) + r^2 \omega^2 R(r) = \ell'(\ell' - 1)R(r). \quad (3.9)$$

This is the spherical Bessel's equation. There are two linearly independent solutions [8]

$$R(r) = A_{far} \frac{1}{\sqrt{\omega r}} J_{-\ell'+\frac{1}{2}}(\omega r) + B_{far} \frac{1}{\sqrt{\omega r}} J_{\ell'-\frac{1}{2}}(\omega r), \quad (3.10)$$

where A_{far} and B_{far} are constants. These solutions have the following asymptotic behaviors

$$\frac{1}{\sqrt{\omega r}} J_{-\ell'+\frac{1}{2}}(\omega r) \rightarrow \omega^{-\ell'} r^{-\ell'} \frac{2^{\ell'-\frac{1}{2}}}{\Gamma(-\ell'+\frac{3}{2})}, \quad r \rightarrow 0, \quad (3.11)$$

$$\rightarrow \sqrt{\frac{2}{\pi}} \frac{1}{\omega r} \sin(\omega r + \pi\ell'/2), \quad r \rightarrow \infty, \quad (3.12)$$

$$\frac{1}{\sqrt{\omega r}} J_{\ell'-\frac{1}{2}}(\omega r) \rightarrow \omega^{\ell'-1} r^{\ell'-1} \frac{2^{-\ell'+\frac{1}{2}}}{\Gamma(\ell'+\frac{1}{2})}, \quad r \rightarrow 0, \quad (3.13)$$

$$\rightarrow \sqrt{\frac{2}{\pi}} \frac{1}{\omega r} \cos(\omega r - \pi\ell'/2), \quad r \rightarrow \infty. \quad (3.14)$$

3.3 Near-far matching

At the outer boundary of the far region, each of these two solutions has outgoing and ingoing modes. One can separate the pure outgoing and pure ingoing part in the solution with the following recombination

$$R(r) = Z_{out} \left(i e^{i\pi\ell'} \frac{1}{\sqrt{\omega r}} J_{-\ell'+\frac{1}{2}}(\omega r) + \frac{1}{\sqrt{\omega r}} J_{\ell'-\frac{1}{2}}(\omega r) \right) + Z_{in} \left(-i e^{-i\pi\ell'} \frac{1}{\sqrt{\omega r}} J_{-\ell'+\frac{1}{2}}(\omega r) + \frac{1}{\sqrt{\omega r}} J_{\ell'-\frac{1}{2}}(\omega r) \right), \quad (3.15)$$

where

$$Z_{out} = \frac{1}{2 \cos(\pi\ell')} (-i A_{far} + e^{-i\pi\ell'} B_{far}), \quad (3.16)$$

$$Z_{in} = \frac{1}{2 \cos(\pi\ell')} (i A_{far} + e^{i\pi\ell'} B_{far}), \quad (3.17)$$

are the amplitudes of the pure outgoing and ingoing part respectively. With the asymptotic behaviors given in (3.12) and (3.14), one can show that the scalar fields at the outer boundary of the far region take the forms

$$\Phi_{out}(r \rightarrow \infty) = \sum_{\ell, m} e^{-i\omega t + im\phi} S_\ell(\theta) Z_{out} \left(\sqrt{\frac{2}{\pi}} \frac{e^{i\omega r + i\pi\ell'/2}}{\omega r} \cos(\pi\ell') \right), \quad (3.18)$$

$$\Phi_{in}(r \rightarrow \infty) = \sum_{\ell, m} e^{-i\omega t + im\phi} S_\ell(\theta) Z_{in} \left(\sqrt{\frac{2}{\pi}} \frac{e^{-i\omega r - i\pi\ell'/2}}{\omega r} \cos(\pi\ell') \right). \quad (3.19)$$

The Klein-Gordon particle number flux is defined as

$$\mathcal{F} = \int \sqrt{-g} J^r d\theta d\phi, \quad \text{with } J^\mu = \frac{i}{8\pi} (\Phi^* \nabla^\mu \Phi - \Phi \nabla^\mu \Phi^*). \quad (3.20)$$

At infinity, the fluxes of outgoing and ingoing particle are

$$\mathcal{F}_{out} \propto |Z_{out}|^2, \quad \mathcal{F}_{in} \propto |Z_{in}|^2, \quad (3.21)$$

with the same proportional constants. The absorption probability, which is the ratio of the net incoming flux to total incoming flux, can be expressed as

$$\sigma_{abs}^{bulk} = \frac{\mathcal{F}_{in} - \mathcal{F}_{out}}{\mathcal{F}_{in}} = 1 - \frac{|Z_{out}|^2}{|Z_{in}|^2} = \frac{2i \cos(\pi \ell') (A_{far} B_{far}^* - B_{far} A_{far}^*)}{|i A_{far} + e^{i\pi \ell'} B_{far}|^2}. \quad (3.22)$$

The radial solutions at far and near region should match with each other at the matching region. The far region $R(r)$ at $r \rightarrow 0$ should have the same coefficients with the near region $R(r)$ at $r \gg M$. By comparing with (3.10), (3.11), (3.13) and (3.5), these coefficients are related through

$$A_{far} = 2^{-\ell' + \frac{1}{2}} \Gamma\left(-\ell' + \frac{3}{2}\right) \omega^{\ell'} A_{near}, \quad (3.23)$$

$$B_{far} = 2^{\ell' - \frac{1}{2}} \Gamma\left(\ell' + \frac{1}{2}\right) (r_+ - r_-)^{-2\ell' + 1} \omega^{-\ell' + 1} B_{near}. \quad (3.24)$$

Using (3.6) and (3.7), and the above relations, we find

$$2i \cos(\pi \ell') (A_{far} B_{far}^* - B_{far} A_{far}^*) = \pi \omega (r_+ - r_-)^{-2\ell'} (4Mr_+ \omega - 2am). \quad (3.25)$$

Note that we assume $\omega M \ll 1$, and $r_+ - r_-$ is at order of M , the coefficient B_{far} will be much larger than A_{far} when $\text{Re}(\ell') > 1$. So the absorption probability in this case can be written as

$$\begin{aligned} \sigma_{abs}^{bulk} &= \frac{2i \cos(\pi \ell') (A_{far} B_{far}^* - B_{far} A_{far}^*)}{|B_{far}|^2} \\ &= \frac{(\omega/2)^{2\ell' - 1} (r_+ - r_-)^{2\ell' - 1}}{\Gamma(\ell' + \frac{1}{2})^2 \Gamma(2\ell' - 1)^2} \sinh\left(\pi \frac{4Mr_+ \omega - 2am}{r_+ - r_-}\right) \\ &\quad \times |\Gamma(\ell' - i2M\omega)|^2 \left| \Gamma\left(\ell' - i \frac{4M^2 \omega - 2am}{r_+ - r_-}\right) \right|^2. \end{aligned} \quad (3.26)$$

It is clear that when the modes parameters ω and m satisfying

$$0 < \omega < m \frac{a}{2Mr_+}, \quad (3.27)$$

the absorption probability is negative, which means that the outgoing flux is more energetic than the incoming one. This corresponds to the superradiant scattering case [38, 39].

In the next section, we will calculate the thermal retarded correlator and the absorption cross section of the WCFT, and show their matchings with the bulk quantities (3.8) and (3.26), respectively.

3.4 Dual calculations in WCFT

For a WCFT with coordinates (x, y) , the general warped conformal symmetry transformation can be written as

$$x' = f(x), \quad y' = y + g(x), \quad (3.28)$$

where $f(x)$ and $g(x)$ are two arbitrary functions. First consider the zero temperature case in which the WCFT is defined on a plane. Infinitesimally, the warped conformal symmetries are generated by a set of vector fields

$$L_n = -x^{n+1}\partial_x, \quad P_n = ix^n\partial_y. \quad (3.29)$$

These vector fields form the Virasoro-Kac-Moody algebra

$$\begin{aligned} [L_n, L_m] &= (n-m)L_{n+m}, \\ [L_n, P_m] &= -mP_{n+m}, \\ [P_n, P_m] &= 0. \end{aligned} \quad (3.30)$$

The global symmetries are generated by $(L_{\pm 1}, L_0, P_0)$ which form a $SL(2, R) \times U(1)$ sub-algebra. At charge level, there are infinitely many conserved charges associated with translations along x and y , with corresponding Noether currents $T(x)$ and $P(x)$, respectively

$$\mathcal{L}_n = -\frac{i}{2\pi} \int dx x^{n+1} T(x), \quad \mathcal{P}_n = -\frac{1}{2\pi} \int dx x^n P(x). \quad (3.31)$$

The commutation relations for the charges form a warped conformal algebra consists of one Virasoro algebra and one Kac-Moody algebra with central extension [10]

$$\begin{aligned} [\mathcal{L}_n, \mathcal{L}_m] &= (n-m)\mathcal{L}_{n+m} + \frac{c}{12}n(n^2-1)\delta_{n,-m}, \\ [\mathcal{L}_n, \mathcal{P}_m] &= -m\mathcal{P}_{n+m}, \\ [\mathcal{P}_n, \mathcal{P}_m] &= k\frac{n}{2}\delta_{n,-m}, \end{aligned} \quad (3.32)$$

where c is the central charge and k is the Kac-Moody level. Similar to the CFT story, one can define the primary operators of the WCFT through the following transformation law [16]

$$\mathcal{O}'(x', y') = \left(\frac{\partial x'}{\partial x} \right)^{-\delta} \mathcal{O}(x, y), \quad (3.33)$$

where δ is the scaling dimension of the primary operator $\mathcal{O}(x, y)$. The infinitesimal version can be expressed as

$$[\mathcal{L}_n, \mathcal{O}(x, y)] = [x^{n+1}\partial_x + (n+1)x^n\delta]\mathcal{O}(x, y), \quad (3.34)$$

$$[\mathcal{P}_n, \mathcal{O}(x, y)] = ix^n\partial_y\mathcal{O}(x, y) = -x^n Q\mathcal{O}(x, y), \quad (3.35)$$

where the last equation is because we are choosing a basis with definite \mathcal{P}_0 charge Q . The quadratic Casimir operator acting on $\mathcal{O}(x, y)$ as an eigenoperator with eigenvalue $\delta(\delta-1)$

$$L^2\mathcal{O} = \left(L_0 - \frac{1}{2}(L_1L_{-1} + L_{-1}L_1) \right) \mathcal{O} = \delta(\delta-1)\mathcal{O}. \quad (3.36)$$

The finite temperature case can be obtained by applying a warped conformal transformation [16]

$$x = e^{\frac{2\pi X}{\beta}}, \quad y = Y + \bar{q}X. \quad (3.37)$$

(X, Y) stands for the finite temperature coordinates with the following thermal identification

$$(X, Y) \sim (X + i\beta, Y - i\bar{\beta}), \quad (3.38)$$

where β is the inverse temperature along X and $\bar{\beta} = \bar{q}\beta + \alpha$ is the inverse temperature along Y . \bar{q} is the spectral flow parameter. In terms of (X, Y) , the warped conformal generators can be written as

$$L_n = -\frac{\beta}{2\pi} e^{\frac{2\pi n X}{\beta}} (\partial_X - \bar{q}\partial_Y), \quad P_n = i e^{\frac{2\pi n X}{\beta}} \partial_Y. \quad (3.39)$$

We relate the finite temperature spectral flowed WCFT coordinates (X, Y) to the Kerr coordinates (t, ϕ) in the following way

$$X = 2\pi T_R \phi, \quad Y = \frac{1}{2M} t - 2\pi(T_L + \bar{q}T_R)\phi. \quad (3.40)$$

The angular identification $\phi \sim \phi + 2\pi$ can be expressed as

$$(X, Y) \sim (X + 4\pi^2 T_R, Y - 4\pi^2(T_L + \bar{q}T_R)). \quad (3.41)$$

Further more, due to the facts that the the Kerr black hole has temperature and the corresponding coordinates have the following thermal identification

$$(t, \phi) \sim (t + iT_H^{-1}, \phi + iT_H^{-1}\Omega_H), \quad (3.42)$$

where T_H and Ω_H are the Hawking temperature and the angular velocity of the outer horizon

$$T_H = \frac{r_+ - r_-}{8\pi M r_+}, \quad \Omega_H = \frac{a}{2M r_+}. \quad (3.43)$$

Then the WCFT coordinates inherit the following thermal identification

$$(X, Y) \sim (X + 2\pi i, Y - 2\pi i(\bar{q} - 1)), \quad (3.44)$$

with $\beta = 2\pi$ and $\bar{\beta} = 2\pi(\bar{q} - 1)$. So the finite temperature WCFT is defined on a torus with both spatial and thermal identifications

$$(X, Y) \sim (X + 4\pi^2 T_R, Y - 4\pi^2(T_L + \bar{q}T_R)) \sim (X + 2\pi i, Y - 2\pi i(\bar{q} - 1)). \quad (3.45)$$

At low frequency $\omega \ll 1$, the near region $r \ll \frac{1}{\omega}$ extends to very far and satisfying $r \gg M$. The WCFT lives on this fixed large r surface. In this place, the conformal local vector fields $(H_{\pm 1}, H_0, \bar{H}_0)$ which form $SL(2, R) \times U(1)$ sub-algebra can be identified as the global part of the warped conformal generators $(L_{\pm 1}, L_0, P_0)$ in (3.39) by using (3.40), through

$$L_{\pm 1} = iH_{\mp 1}, \quad L_0 = iH_0, \quad P_0 = -\bar{H}_0. \quad (3.46)$$

With the above relation, the quadratic Casimir operator L^2 of the $SL(2, R)$ in the warped conformal algebra

$$L^2 = L_0^2 - \frac{1}{2}(L_1 L_{-1} + L_{-1} L_1) = \mathcal{H}^2, \quad (3.47)$$

is relate to the conformal Casimir operator \mathcal{H}^2 in the scalar wave equation (2.17). Thus we can relate the scaling dimension δ of the WCFT primary operator to the $SL(2, R)$ conformal weight of the scalar field Φ

$$\delta = \ell' = \frac{1}{2} + \frac{1}{2} \sqrt{(2\ell - 1)^2 - 16M^2\omega^2}. \quad (3.48)$$

The action of the deformed WCFT can be written as

$$S = S_{WCFT} + \sum_{\delta} \int dX dY J_{\delta, Q}(X, Y) \mathcal{O}_{\delta, Q}(X, Y), \quad (3.49)$$

where S_{WCFT} is the original WCFT action, and $\mathcal{O}_{\delta, Q}(X, Y)$ are the WCFT operator with scaling dimension δ and fix \mathcal{P}_0 charge Q . The source term, which comes from the unnormalised part of the asymptotic scalar field in the Kerr spacetime, can be expanded in terms of the periodic ϕ modes

$$J_{\delta, Q}(X, Y) = \sum_m J_{\delta m} e^{-i\Omega X + iQY}, \quad (3.50)$$

where $J_{\delta m} \propto B_{near}$, and

$$\Omega = \frac{4M^2\omega - 2am}{r_+ - r_-} + 2M\omega\bar{q}, \quad Q = -2M\omega. \quad (3.51)$$

Here we view Q as fixed constant and Ω is the frequency along X .

Inspired by the CFT's result, the Fermi's Golden rule gives the transition rate out of the thermal state [40, 41]

$$\mathcal{R} = 2\pi \sum_{\delta m} |J_{\delta m}|^2 \int dX dY e^{i\Omega X - iQY} G(X, Y), \quad (3.52)$$

where $G(X, Y)$ is the finite temperature two point function of the WCFT [16]

$$G(X, Y) = \langle \mathcal{O}_{\delta, Q}(X, Y) \mathcal{O}_{\delta, -Q}(0, 0) \rangle \sim \mathcal{C}_{\mathcal{O}} (-1)^{\delta} e^{iQ(Y + \bar{q}X)} \left(\frac{\beta}{\pi} \sinh \frac{\pi X}{\beta} \right)^{-2\delta}, \quad (3.53)$$

with $\beta = 2\pi$. We introduce an operator normalization factor $\mathcal{C}_{\mathcal{O}}$ in (3.53) with the following form for later convenience

$$\mathcal{C}_{\mathcal{O}} = e^{\Theta(\text{Im}(Y))\beta Q}, \quad (3.54)$$

where Θ is the step function. From the transition rate (3.52), one can write down the thermal retarded correlator and the absorption cross section in the following form [5]

$$G_R \sim \int dX dY e^{i\Omega X - iQY} G(X - i\epsilon, Y - i\epsilon), \quad (3.55)$$

$$\sigma_{abs} \sim \int dX dY e^{i\Omega X - iQY} [G(X - i\epsilon, Y - i\epsilon) - G(X + i\epsilon, Y + i\epsilon)], \quad (3.56)$$

where $-i\epsilon$ and $+i\epsilon$ correspond to absorption and emission. Using (3.53) in (3.55) and further notice that the Euclidean frequency at finite temperature takes discrete values of the Matsubara frequencies, the thermal retarded correlator then takes the form [16]

$$G_R \sim (2\pi)^{1-2\delta} \frac{e^{\pi Q\bar{q}}}{\sin(\pi(\delta + iQ\bar{q}))} \frac{\Gamma(\delta - i(\Omega + Q\bar{q}))}{\Gamma(1 - \delta - i(\Omega + Q\bar{q}))}, \quad (3.57)$$

where we have set $\beta = 2\pi$. Invoking the relations (3.48) and (3.51), one can show that the retarded Green's function for the scalar field in the Kerr spacetime (3.8) matches the WCFT thermal retarded correlator (3.57) in the Ω dependence.

Taking into account the normalization factor (3.54) in the two point function, the two terms in (3.56) then have additional relative factors in terms of Q . The absorption cross section takes the form

$$\sigma_{abs} \sim \frac{e^{\pi Q}}{\Gamma(2\delta)} \sinh(\pi(\Omega + Q\bar{q} - Q)) |\Gamma(\delta - i(\Omega - Q\bar{q}))|^2. \quad (3.58)$$

Given the relations (3.48) and (3.51), the absorption probability of the scalar flux in the Kerr spacetime (3.26) matches the WCFT absorption cross section (3.58) in the Ω dependence.

4 Higher spin radiations in Kerr spacetime

In this section, we generalise the previous discussion on the scalar to the higher spin field case, for instance the gravitational perturbation on the Kerr background. To explore the higher spin case, it is convenient to apply the Newman-Penrose formalism, by introducing the NP null tetrad of the Kerr metric in components (t, r, θ, ϕ)

$$\begin{aligned} \ell^\mu &= \left(\frac{r^2 + a^2}{\Delta}, 1, 0, \frac{a}{\Delta} \right), \\ n^\mu &= \frac{1}{2(r^2 + a^2 \cos^2 \theta)} (r^2 + a^2, -\Delta, 0, a), \\ m^\mu &= \frac{1}{\sqrt{2}(r + ia \cos \theta)} \left(ia \sin \theta, 0, 1, \frac{i}{\sin \theta} \right). \end{aligned} \quad (4.1)$$

The perturbations of the following gauge invariant Weyl scalars describe the gravitational radiations

$$\Psi_{s=-2} = \Psi_4^{(1)} = C_{\mu\nu\rho\sigma} n^\mu m^{*\nu} n^\rho m^{*\sigma}, \quad (4.2)$$

$$\Psi_{s=2} = \Psi_0^{(1)} = C_{\mu\nu\rho\sigma} \ell^\mu m^\nu \ell^\rho m^\sigma, \quad (4.3)$$

where $\Psi_{s=-2}$ is for outgoing radiation and $\Psi_{s=2}$ is for ingoing. These perturbations satisfy the following master equation on the Kerr background [42, 43]

$$\begin{aligned} &\left(\frac{(r^2 + a^2)^2}{\Delta} - a^2 \sin^2 \theta \right) \frac{\partial^2 \Psi_s}{\partial t^2} + \frac{4Mar}{\Delta} \frac{\partial^2 \Psi_s}{\partial t \partial \phi} + \left(\frac{a^2}{\Delta} - \frac{1}{\sin \theta} \right) \frac{\partial^2 \Psi_s}{\partial \phi^2} \\ &- \Delta^{-s} \frac{\partial}{\partial r} \left(\Delta^{s+1} \frac{\partial \Psi_s}{\partial r} \right) - \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial \Psi_s}{\partial \theta} \right) - 2s \left(\frac{a(r-M)}{\Delta} + \frac{i \cos \theta}{\sin^2 \theta} \right) \frac{\partial \Psi_s}{\partial \phi} \\ &- 2s \left(\frac{M(r^2 - a^2)}{\Delta} - r - ia \cos \theta \right) \frac{\partial \Psi_s}{\partial t} + (s^2 \cot^2 \theta - s) \Psi_s = 0. \end{aligned} \quad (4.4)$$

This equation is valid for a generic spin-weight parameter s and we have set the source term vanishing on the right hand side. The master equation (4.4) is separable by writing

$$\Psi_s = e^{-i\omega t + im\phi} S^s(\theta) R^s(r), \quad (4.5)$$

which leads to the following angular and radial equations

$$\begin{aligned} \frac{1}{\sin \theta} \frac{d}{d\theta} \left(\sin \theta \frac{dS^s(\theta)}{d\theta} \right) - \left(\frac{m^2}{\sin^2 \theta} + \frac{2ms \cos \theta}{\sin^2 \theta} + s^2 \cot^2 \theta - s \right) S^s(\theta) \\ - 2\omega a s \cos \theta S^s(\theta) + \omega^2 a^2 \cos^2 \theta S^s(\theta) = -K_\ell^s S^s(\theta), \end{aligned} \quad (4.6)$$

$$\begin{aligned} \Delta^{-s} \frac{d}{dr} \left(\Delta^{s+1} \frac{dR^s(r)}{dr} \right) + \left(\frac{(2Mr_+\omega - am)^2}{(r-r_+)(r_+-r_-)} - \frac{(2Mr_-\omega - am)^2}{(r-r_+)(r_+-r_-)} \right) R^s(r) \\ - \left(is \frac{2Mr_+\omega - am}{r-r_+} + is \frac{2Mr_-\omega - am}{r-r_-} \right) R^s(r) + 2is\omega(r-M)R^s(r) \\ + (r^2 + 2M(r+2M))\omega^2 R^s(r) = K_\ell^s R^s(r), \end{aligned} \quad (4.7)$$

with K_ℓ^s being the separation constant.

4.1 Near region

In the near region and low frequency limit, the angular equation becomes

$$\frac{1}{\sin \theta} \frac{d}{d\theta} \left(\sin \theta \frac{dS^s(\theta)}{d\theta} \right) - \left(\frac{m^2}{\sin^2 \theta} + \frac{2ms \cos \theta}{\sin^2 \theta} + s^2 \cot^2 \theta - s \right) S^s(\theta) = -K_\ell^s S^s(\theta), \quad (4.8)$$

which is the angular part of spin weighted spherical harmonic equation with $K_\ell^s = (\ell + s)(\ell - s - 1)$ [44]. And the radial equation reduces to

$$\begin{aligned} \Delta^{-s} \frac{d}{dr} \left(\Delta^{s+1} \frac{dR^s(r)}{dr} \right) + \left(\frac{(2Mr_+\omega - am)^2}{(r-r_+)(r_+-r_-)} - \frac{(2Mr_-\omega - am)^2}{(r-r_+)(r_+-r_-)} \right) R^s(r) \\ - \left(is \frac{2Mr_+\omega - am}{r-r_+} + is \frac{2Mr_-\omega - am}{r-r_-} \right) R^s(r) = K_{\ell'}^s R^s(r). \end{aligned} \quad (4.9)$$

Here we preserve the additional $4M^2\omega^2$ term in the potential and keep using the ω deformed ℓ' as in (3.2) in stead of ℓ in the radial equation. The solution to the radial equation in the near region can be found by the Green function method [8]. Imposing an ingoing boundary condition at the horizon, the radial solution can be expressed as

$$\begin{aligned} R^s(r) = \left(\frac{r-r_+}{r-r_-} \right)^{-i \frac{2Mr_+\omega - am}{r_+-r_-}} (r-r_-)^{-\ell'-s} \\ \times {}_2F_1 \left(\ell' - i \frac{4M^2\omega - 2am}{r_+-r_-}, \ell' - s - i2M\omega; 1 - s - i \frac{4Mr_+\omega - 2am}{r_+-r_-}; \frac{r-r_+}{r-r_-} \right). \end{aligned} \quad (4.10)$$

When the ω is very small, the near region extends to the far region. At the outer boundary of the near region, the above solution behaves as

$$R^s(r \gg M) \sim A_{near}^s r^{-\ell'-s} + B_{near}^s (r_+-r_-)^{-2\ell'+1} r^{\ell'-s-1}, \quad (4.11)$$

where

$$A_{near}^s = \frac{\Gamma \left(1 - s - i \frac{4Mr_+\omega - 2am}{r_+-r_-} \right) \Gamma(-2\ell' + 1)}{\Gamma(1 - \ell' - s - i2M\omega) \Gamma \left(1 - \ell' - i \frac{4M^2\omega - 2am}{r_+-r_-} \right)}, \quad (4.12)$$

$$B_{near}^s = \frac{\Gamma \left(1 - s - i \frac{4Mr_+\omega - 2am}{r_+-r_-} \right) \Gamma(2\ell' - 1)}{\Gamma(\ell' - s - i2M\omega) \Gamma \left(\ell' - i \frac{4M^2\omega - 2am}{r_+-r_-} \right)}. \quad (4.13)$$

The retarded Green's function for the spin- s field Ψ_s in the frequency space then proportional to the ratio

$$G_R^{s,bulk} \sim \frac{A_{near}^s}{B_{near}^s} = \frac{\Gamma(-2\ell' + 1)}{\Gamma(2\ell' - 1)} \frac{\Gamma(\ell' - s - i2M\omega)}{\Gamma(1 - \ell' - s - i2M\omega)} \frac{\Gamma\left(\ell' - i\frac{4M^2\omega - 2am}{r_+ - r_-}\right)}{\Gamma\left(1 - \ell' - i\frac{4M^2\omega - 2am}{r_+ - r_-}\right)}. \quad (4.14)$$

4.2 Far region

In the far region, the radial equation reduces to

$$r^{-2s} \frac{d}{dr} \left(r^{2s+2} \frac{dR^s(r)}{dr} \right) + (r^2\omega^2 + 2isr\omega)R^s(r) = (\ell' + s)(\ell' - s - 1)R^s(r). \quad (4.15)$$

The solutions of the above equation are the confluent hypergeometric functions [8]. The generic solution can be expressed as

$$R^s(r) = A_{far}^s R_1^s + B_{far}^s R_2^s, \quad (4.16)$$

where A_{far}^s and B_{far}^s are constants, and

$$R_1^s = r^{-\ell' - s} e_1^{-i\omega r} F_1(-\ell' - s + 1, -2\ell' + 2; 2i\omega r), \quad (4.17)$$

$$R_2^s = r^{\ell' - s - 1} e_1^{-i\omega r} F_1(\ell' - s, 2\ell'; 2i\omega r). \quad (4.18)$$

The asymptotic behaviors of R_1^s and R_2^s are

$$R_1^s \rightarrow r^{-\ell' - s}, \quad r \rightarrow 0, \quad (4.19)$$

$$\rightarrow \alpha_{in}^s r^{-1} e^{-i\omega r} + \alpha_{out}^s r^{-2s-1} e^{i\omega r}, \quad r \rightarrow \infty, \quad (4.20)$$

$$R_2^s \rightarrow r^{\ell' - s - 1}, \quad r \rightarrow 0, \quad (4.21)$$

$$\rightarrow \beta_{in}^s r^{-1} e^{-i\omega r} + \beta_{out}^s r^{-2s-1} e^{i\omega r}, \quad r \rightarrow \infty, \quad (4.22)$$

where

$$\begin{aligned} \alpha_{in}^s &= \frac{(2)^{\ell' + s - 1} (-i\omega)^{\ell' + s - 1} \Gamma(2 - 2\ell')}{\Gamma(-\ell' + s + 1)}, & \alpha_{out}^s &= \frac{(2i\omega)^{\ell' - s - 1} \Gamma(2 - 2\ell')}{\Gamma(-\ell' - s + 1)}, \\ \beta_{in}^s &= \frac{(2)^{-\ell' + s} (-i\omega)^{-\ell' + s} \Gamma(2\ell')}{\Gamma(\ell' + s)}, & \beta_{out}^s &= \frac{(2i\omega)^{-\ell' - s} \Gamma(2\ell')}{\Gamma(\ell' - s)}. \end{aligned} \quad (4.23)$$

4.3 Near-far matching

Matching the far region solution at $r \rightarrow 0$ to the near region solution at $r \gg M$ with ingoing boundary condition at the horizon (4.11), one find the relation

$$A_{far}^s = A_{near}^s, \quad B_{far}^s = (r_+ - r_-)^{-2\ell' + 1} B_{near}^s. \quad (4.24)$$

For the gravitational perturbation, the $s = 2$ case is for the ingoing radiative part while the $s = -2$ is for outgoing part. At infinity, the radial solutions have both outgoing and

ingoing parts. One can extract the outgoing and ingoing amplitudes by the following linear combination

$$Z_{out}^s \propto A_{far}^s \alpha_{out}^{-s} + B_{far}^s \beta_{out}^{-s}, \quad (4.25)$$

$$Z_{in}^s \propto A_{far}^s \alpha_{in}^s + B_{far}^s \beta_{in}^s, \quad (4.26)$$

where we flip the sign of s in the outgoing asymptotic coefficients since the spin components are defined according to the ingoing direction in (4.10). The corresponding perturbation of the outgoing and ingoing Weyl scalars at infinity then take the form

$$\Psi_{out}(r \rightarrow \infty) = \sum_{\ell, m} e^{-i\omega t + im\phi} S_\ell^s(\theta) Z_{out}^s r^{-1+2s} e^{i\omega r}, \quad (4.27)$$

$$\Psi_{in}(r \rightarrow \infty) = \sum_{\ell, m} e^{-i\omega t + im\phi} S_\ell^s(\theta) Z_{in}^s r^{-1} e^{i\omega r}. \quad (4.28)$$

The absorption probability is the net incoming flux to total incoming flux ratio. In the gravitational case, the energy flux at infinity is proportional to $|\Psi^s|^2$ [43]. In this sense, the absorption probability of the gravitational radiation can be written as

$$\begin{aligned} \sigma_{abs}^{s,bulk} &= 1 - \frac{|Z_{out}^s|^2}{|Z_{in}^s|^2} = 1 - \frac{|A_{far}^s \alpha_{out}^{-s} + B_{far}^s \beta_{out}^{-s}|^2}{|A_{far}^s \alpha_{in}^s + B_{far}^s \beta_{in}^s|^2} \\ &= \frac{2(A_{far}^s B_{far}^{*s} - B_{far}^s A_{far}^{*s}) \left(\frac{\alpha_{in}^s}{\beta_{in}^s} \right)}{|B_{far}^s|^2}, \end{aligned} \quad (4.29)$$

where in the last line, we invoke equations (4.23) and use the fact that in the low frequency limit, the coefficient $B_{far}^s \beta_{in}^s$ is much larger than $A_{far}^s \alpha_{in}^s$ when $\text{Re}(\ell') > 1$. Substituting equations (4.24), (4.12), (4.13) and (4.23) into the above formula, and take $s = 2$ for the gravitational case, we have

$$\begin{aligned} \sigma_{abs}^{s,bulk} &= (2\omega)^{2\ell'-1} (r_+ - r_-)^{2\ell'-1} \frac{\Gamma(\ell' + 2)\Gamma(-2\ell' + 2)(\ell' - 2)(\ell' - 1)\ell'(\ell' + 1)}{\Gamma(-\ell' + 3)\Gamma(2\ell')\Gamma(2\ell' - 1)^2\Gamma(\ell' + 1/2)\Gamma(-\ell' + 3/2)} \\ &\times \sinh\left(\pi \frac{4Mr_+ \omega - 2am}{r_+ - r_-}\right) |\Gamma(\ell' - s - i2M\omega)|^2 \left| \Gamma\left(\ell' - i \frac{4M^2\omega - 2am}{r_+ - r_-}\right) \right|^2 \\ &+ \mathcal{O}(\omega^{2\ell'-1}). \end{aligned} \quad (4.30)$$

4.4 Dual explanations in WCFT

For the higher spin case, the form of the frequency dependence in the bulk retarded Green's function and the absorption probability are expressed in (4.14) and (4.30), where there are possible different normalization factors for different spin weights. Note that in the WCFT, the charge Q is held fixed, invoking the relations (3.48) and (3.51), these scattering quantities (4.14) and (4.30) match the WCFT thermal retarded correlator (3.57) and the absorption cross section (3.58) in the Ω dependence, respectively.

However, the purely ω dependence and the spin weight parameters in the bulk scattering quantities are missing in the WCFT calculations. This is partially due to the non-locality of

the WCFT operator correlations in the Y coordinate, which exhibit a plane wave form [16], so the charge dependence in the frequency space is a delta function with fixed value. The WCFT is a non-relativistic field theory, a similar notion of spinning operator as in CFT is absent as well. Therefore, if the WCFT features the holographic dual of the Kerr black hole, one needs to incorporate charge-dependent normalization factors of the corresponding operators to resemble the local interactions in a 2D CFT.

5 Summary and discussion

In this paper, we consider the scalar and higher spin radiations on a generic non-extreme Kerr black hole background. We calculate the retarded Green's function and the absorption probability for the radiations, and find their matching to the thermal retarded correlator and the absorption cross section in the dual WCFT, respectively.

Unlike CFT, the WCFT has one Virasoro and one Kac-Moody algebra determining the underlying symmetry. This makes it a good candidate as the holographic dual for a class of geometries with $SL(2, R)$ and $U(1)$ isometry. The extreme Kerr black hole enjoys such isometry in its near horizon scaling region. For non-extreme Kerr black hole, the warped conformal symmetry is not manifest in the geometry, but it can be obtained in the covariant phase space. In this sense, the WCFT could also be relevant to the holographic description of the Kerr black hole, and the present work provides further evidence on it. In the WCFT, the primary operators' momenta are translated by the energy eigenmode parameters of the bulk fields, which are inspired by the identifications between the hidden conformal generators and the global warped conformal generators. The frequency dependent $SL(2, R)$ conformal weight of the bulk scalar field is set equal to the scaling dimension of the primary operator. This leads to a charge dependent scaling dimension. However, the $U(1)$ symmetry requires the charge to be fixed on the field theory side, so the bulk fields correspondingly have given fixed frequency. The matching between the bulk scattering amplitudes and the WCFT thermal correlators are then realized in the angular momentum space given a charge dependent normalization factor. For the higher spin case, the matching in the angular momentum space exists, but the spin weight parameters are missing on the WCFT side. How to introduce additional quantum numbers in the WCFT to acquire such interpretations is an interesting next step.

Acknowledgement

We are grateful to Jianxin Lu, Jun Nian, and Wei Song for helpful discussions. This work is supported by the NSFC Grant No. 12105045.

References

- [1] M. Guica, T. Hartman, W. Song and A. Strominger, "The Kerr/CFT Correspondence," *Phys. Rev. D* **80**, 124008 (2009) [arXiv:0809.4266 [hep-th]].
- [2] J. M. Bardeen and G. T. Horowitz, "The Extreme Kerr throat geometry: A Vacuum analog of $AdS(2) \times S^2$," *Phys. Rev. D* **60**, 104030 (1999) [arXiv:hep-th/9905099 [hep-th]].

- [3] Y. Matsuo, T. Tsukioka and C. M. Yoo, “Another Realization of Kerr/CFT Correspondence,” Nucl. Phys. B **825**, 231-241 (2010) [arXiv:0907.0303 [hep-th]].
- [4] B. Chen, B. Ning and J. J. Zhang, “Boundary Conditions for NHEK through Effective Action Approach,” Chin. Phys. Lett. **29**, 041101 (2012) [arXiv:1105.2878 [hep-th]].
- [5] I. Bredberg, T. Hartman, W. Song and A. Strominger, “Black Hole Superradiance From Kerr/CFT,” JHEP **04**, 019 (2010) [arXiv:0907.3477 [hep-th]].
- [6] A. Castro, A. Maloney and A. Strominger, “Hidden Conformal Symmetry of the Kerr Black Hole,” Phys. Rev. D **82**, 024008 (2010) [arXiv:1004.0996 [hep-th]].
- [7] S. Haco, S. W. Hawking, M. J. Perry and A. Strominger, “Black Hole Entropy and Soft Hair,” JHEP **12**, 098 (2018) [arXiv:1810.01847 [hep-th]].
- [8] J. Nian and W. Tian, “Gravitational Waves of Non-Extremal Kerr Black Holes from Conformal Symmetry,” [arXiv:2308.03577 [hep-th]].
- [9] D. M. Hofman and A. Strominger, “Chiral Scale and Conformal Invariance in 2D Quantum Field Theory,” Phys. Rev. Lett. **107**, 161601 (2011) [arXiv:1107.2917 [hep-th]].
- [10] S. Detournay, T. Hartman and D. M. Hofman, “Warped Conformal Field Theory,” Phys. Rev. D **86**, 124018 (2012) [arXiv:1210.0539 [hep-th]].
- [11] G. Compère, W. Song and A. Strominger, “Chiral Liouville Gravity,” JHEP **05**, 154 (2013) [arXiv:1303.2660 [hep-th]].
- [12] D. M. Hofman and B. Rollier, “Warped Conformal Field Theory as Lower Spin Gravity,” Nucl. Phys. B **897**, 1-38 (2015) [arXiv:1411.0672 [hep-th]].
- [13] A. Castro, D. M. Hofman and G. Sárosi, “Warped Weyl fermion partition functions,” JHEP **11**, 129 (2015) [arXiv:1508.06302 [hep-th]].
- [14] K. Jensen, “Locality and anomalies in warped conformal field theory,” JHEP **12**, 111 (2017) [arXiv:1710.11626 [hep-th]].
- [15] P. Chaturvedi, Y. Gu, W. Song and B. Yu, “A note on the complex SYK model and warped CFTs,” JHEP **12**, 101 (2018) [arXiv:1808.08062 [hep-th]].
- [16] W. Song and J. Xu, “Correlation Functions of Warped CFT,” JHEP **04**, 067 (2018) [arXiv:1706.07621 [hep-th]].
- [17] A. Castro, D. M. Hofman and N. Iqbal, “Entanglement Entropy in Warped Conformal Field Theories,” JHEP **02**, 033 (2016) [arXiv:1511.00707 [hep-th]].
- [18] L. Apolo, S. He, W. Song, J. Xu and J. Zheng, “Entanglement and chaos in warped conformal field theories,” JHEP **04**, 009 (2019) [arXiv:1812.10456 [hep-th]].
- [19] D. Anninos, W. Li, M. Padi, W. Song and A. Strominger, “Warped AdS(3) Black Holes,” JHEP **03**, 130 (2009) [arXiv:0807.3040 [hep-th]].
- [20] G. Compère and S. Detournay, “Semi-classical central charge in topologically massive gravity,” Class. Quant. Grav. **26**, 012001 (2009) [erratum: Class. Quant. Grav. **26**, 139801 (2009)] [arXiv:0808.1911 [hep-th]].
- [21] G. Compère and S. Detournay, “Boundary conditions for spacelike and timelike warped AdS_3 spaces in topologically massive gravity,” JHEP **08**, 092 (2009) [arXiv:0906.1243 [hep-th]].
- [22] M. Blagojevic and B. Cvetkovic, “Asymptotic structure of topologically massive gravity in spacelike stretched AdS sector,” JHEP **09**, 006 (2009) [arXiv:0907.0950 [gr-qc]].

- [23] D. Anninos, G. Compère, S. de Buyl, S. Detournay and M. Guica, “The Curious Case of Null Warped Space,” JHEP **11**, 119 (2010) [arXiv:1005.4072 [hep-th]].
- [24] D. Anninos, S. de Buyl and S. Detournay, “Holography For a De Sitter-Esque Geometry,” JHEP **05**, 003 (2011) [arXiv:1102.3178 [hep-th]].
- [25] M. Henneaux, C. Martinez and R. Troncoso, “Asymptotically warped anti-de Sitter spacetimes in topologically massive gravity,” Phys. Rev. D **84**, 124016 (2011) [arXiv:1108.2841 [hep-th]].
- [26] G. Compère, W. Song and A. Strominger, “New Boundary Conditions for AdS₃,” JHEP **05**, 152 (2013) [arXiv:1303.2662 [hep-th]].
- [27] W. Song, Q. Wen and J. Xu, “Modifications to Holographic Entanglement Entropy in Warped CFT,” JHEP **02**, 067 (2017) [arXiv:1610.00727 [hep-th]].
- [28] Q. Wen, “Towards the generalized gravitational entropy for spacetimes with non-Lorentz invariant duals,” JHEP **01**, 220 (2019) [arXiv:1810.11756 [hep-th]].
- [29] B. Gao and J. Xu, “Holographic entanglement entropy in AdS₃/WCFT,” Phys. Lett. B **822**, 136647 (2021) [arXiv:1912.00562 [hep-th]].
- [30] L. Apolo, H. Jiang, W. Song and Y. Zhong, “Swing surfaces and holographic entanglement beyond AdS/CFT,” JHEP **12**, 064 (2020) [arXiv:2006.10740 [hep-th]].
- [31] L. Apolo, H. Jiang, W. Song and Y. Zhong, “Modular Hamiltonians in flat holography and (W)AdS/WCFT,” JHEP **09**, 033 (2020) [arXiv:2006.10741 [hep-th]].
- [32] S. Detournay, D. Grumiller, M. Riegler and Q. Vandermiers, “Uniformization of Entanglement Entropy in Holographic Warped Conformal Field Theories,” [arXiv:2006.16167 [hep-th]].
- [33] B. Chen, P. X. Hao and W. Song, “Rényi mutual information in holographic warped CFTs,” JHEP **10**, 037 (2019) [arXiv:1904.01876 [hep-th]].
- [34] B. Chen, Y. Liu and B. Yu, “Reflected entropy in AdS₃/WCFT,” JHEP **12**, 008 (2022) [arXiv:2205.05582 [hep-th]].
- [35] W. Song and J. Xu, “Structure Constants from Modularity in Warped CFT,” JHEP **10**, 211 (2019) [arXiv:1903.01346 [hep-th]].
- [36] T. Azeanagi, S. Detournay and M. Riegler, “Warped Black Holes in Lower-Spin Gravity,” Phys. Rev. D **99**, no.2, 026013 (2019) [arXiv:1801.07263 [hep-th]].
- [37] A. Aggarwal, A. Castro and S. Detournay, “Warped Symmetries of the Kerr Black Hole,” JHEP **01**, 016 (2020) [arXiv:1909.03137 [hep-th]].
- [38] Y. B. Zel’dovich, Pis’ma Zh. Eksp. Teor. Fiz. **14**, 270 (1971) [JETP Lett. **14**, 180 (1971)].
- [39] Y. B. Zel’dovich, Zh. Eksp. Teor. Fiz. **62**, 2076 (1972) [Sov. Phys. JETP **35**, 1085 (1972)].
- [40] J. M. Maldacena and A. Strominger, “Universal low-energy dynamics for rotating black holes,” Phys. Rev. D **56**, 4975-4983 (1997) [arXiv:hep-th/9702015 [hep-th]].
- [41] S. S. Gubser, “Absorption of photons and fermions by black holes in four-dimensions,” Phys. Rev. D **56**, 7854-7868 (1997) [arXiv:hep-th/9706100 [hep-th]].
- [42] S. A. Teukolsky, “Rotating black holes - separable wave equations for gravitational and electromagnetic perturbations,” Phys. Rev. Lett. **29**, 1114-1118 (1972)

- [43] S. A. Teukolsky, "Perturbations of a rotating black hole. 1. Fundamental equations for gravitational electromagnetic and neutrino field perturbations," *Astrophys. J.* **185**, 635-647 (1973)
- [44] J. N. Goldberg, A. J. Macfarlane, E. T. Newman, F. Rohrlich, E. C. G. Sudarshan, "Spin-s Spherical Harmonics and edth," *J. Math. Phys.* **8**, 2155-2161 (1967)