GEOMETRIC MODEL FOR WEIGHTED PROJECTIVE LINES OF TYPE (p,q)

JIANMIN CHEN, SHIQUAN RUAN AND HONGXIA ZHANG*

ABSTRACT. We give a geometric model for the category of coherent sheaves over the weighted projective line of type (p,q) in terms of an annulus with marked points on its boundary. We establish a bijection between indecomposable sheaves over the weighted projective line and certain homotopy classes of oriented curves in the annulus, and prove that the dimension of extension group between indecomposable sheaves equals to the positive intersection number between the corresponding curves.

By using the geometric model, we provide a combinatorial description for the titling graph of tilting bundles, which is composed by quadrilaterals (or degenerated to a line). Moreover, we obtain that the automorphism group of the coherent sheaf category is isomorphic to the mapping class group of the marked annulus, and show the compatibility of their actions on the tilting graph of coherent sheaves and on the triangulation of the geometric model respectively. A geometric description of the perpendicular category with respect to an exceptional sheaf is presented at the end of the paper.

1. INTRODUCTION

Geometric models for categories have been studied by various authors in recent years. For instance, a geometric construction of cluster categories of type A was given by Caldero-Chapton-Schiffler in [12], of type D was given by Schiffler in [43] and of type A_{∞} was given by Holm-Jørgensen in [29]. There is also some progress for geometric models of abelian categories. Baur-Marsh in [7] provided a geometric model for tube categories. In [48], Warkentin established a bijection between string modules over a quiver of affine type A and certain oriented curves in a marked annulus. By these geometric realizations, many algebraic properties (e.g. the extension dimensions, Auslander-Reiten triangles, Auslander-Reiten sequences) of these categories can be studied in geometric terms. We refer to [5,6,8, 9,17,21,28,43,46] for related topics.

Weighted projective lines and their coherent sheaves categories were introduced by Geigle and Lenzing in [23], in order to give a geometric realization of canonical algebras in the sense of Ringel [41]. The study of weighted projective lines has a high contact with many mathematical branches, such as Lie theory [14–16,44], representation theory [13,22,26,40] and singularity theory [19,30,31,33–35], which in particular including the aspects of Arnold's strange duality [19,20] and homological mirror symmetry [18,42].

By Geigle-Lenzing [23], the coherent sheaf category coh- $\mathbb{X}(p,q)$ of type (p,q) is derived equivalent to the finitely generated module category mod $\tilde{A}_{p,q}$ of the canonical algebra $\tilde{A}_{p,q}$ (affine type A). Inspired by [9, 48], we hope to give a geometric model for the category coh- $\mathbb{X}(p,q)$ in terms of a marked annulus.

²⁰²⁰ Mathematics Subject Classification. 05C10, 05E10, 16D90, 16G70, 57K20.

Key words and phrases. Weighted projective line; geometric model; tilting graph; mapping class group.

^{*} the corresponding author.

Let $A_{p,q}$ be an annulus with p marked points on the inner boundary and q marked points on the outer boundary. We establish a bijection between indecomposable sheaves over $\operatorname{coh-X}(p,q)$ and certain homotopy classes of oriented curves in the annulus $A_{p,q}$ (see Theorem 3.4). Under this correspondence, Auslander-Reiten sequences in $\operatorname{coh-X}(p,q)$ can be realized via elementary moves in $A_{p,q}$ (see Proposition 3.6), and the dimension of extension space between two indecomposable coherent sheaves equals to the positive intersection number of the correspondence curves (see Theorem 3.10). Moreover, we obtain that the tilting sheaves in $\operatorname{coh-X}(p,q)$ are in natural bijection with the triangulations of $A_{p,q}$ (see Theorem 4.3), and the flip of an arc is compatible with the tilting mutation (see Proposition 4.7).

We point out here that the geometric model given in Theorem 3.4 has intuitive difference from the geometric realization of $\operatorname{mod} \tilde{A}_{p,q}$ in [9, 48]. More precisely, the indecomposable modules in the postprojective (*resp.* preinjective) components of $\operatorname{mod} \tilde{A}_{p,q}$ correspond to the bridging curves whose orientation is from the outer boundary to the inner boundary (*resp.* from the inner boundary to the outer boundary). However, all line bundles in coh- $\mathbb{X}(p,q)$ correspond to the bridging curves whose orientations are from the outer boundary to the inner boundary.

There is another interesting point, since the dimensions of the Hom-space and Ext-space between two indecomposable sheaves have explicit formulas due to the structure of the category coh- $\mathbb{X}(p,q)$, Theorem 3.10 makes the positive intersection number of the correspondence curves in $A_{p,q}$ easily calculated. Therefore, it seems that the category of coherent sheaves over $\mathbb{X}(p,q)$ provides a nice categorification model for the annulus $A_{p,q}$.

The geometric model has applications on the automorphism group of $\operatorname{coh-X}(p,q)$ and the tilting graph of coherent sheaves. Denote by

$$\mathcal{T}_{A_{p,q}} := \{ \text{Triangulations of } A_{p,q} \} \text{ and } \mathcal{T}_{\mathbb{X}} := \{ \text{Tilting sheaves in coh-} \mathbb{X}(p,q) \}$$

Then there is a bijection $\phi : \mathcal{T}_{A_{p,q}} \to \mathcal{T}_{\mathbb{X}}$, see (3.1).

There is an isomorphism ψ between the mapping class group $\mathcal{MG}(A_{p,q})$ of the marked annulus $A_{p,q}$ and the automorphism group $\operatorname{Aut}(\operatorname{coh}-\mathbb{X}(p,q))$ of $\operatorname{coh}-\mathbb{X}(p,q)$, see (7.4). Any automorphism of $\operatorname{coh}-\mathbb{X}(p,q)$ preserves tilting sheaves. Hence there is a natural group action of $\operatorname{Aut}(\operatorname{coh}-\mathbb{X}(p,q))$ on $\mathcal{T}_{\mathbb{X}}$. On the other hand, the mapping class group $\mathcal{MG}(A_{p,q})$ naturally acts on the set of triangulations of $A_{p,q}$. It turns out that these two actions are compatible. That is, we have the following commutative diagram, where the commutativity is in the sense of (1.1).



Theorem 1.1. For any $f \in \mathcal{MG}(A_{p,q})$ and any triangulation Γ of $A_{p,q}$, we have

$$\phi(f(\Gamma)) = \psi(f)(\phi(\Gamma)) \tag{1.1}$$

The tilting graph $\mathcal{G}(\mathcal{T}_{\mathbb{X}})$ of coh- $\mathbb{X}(p,q)$ has as vertices the isomorphism classes of tilting sheaves in coh- $\mathbb{X}(p,q)$, while two vertices are connected by an edge if and only if the associated

tilting sheaves differ by precisely one indecomposable direct summand. The full subgraph of $\mathcal{G}(\mathcal{T}_{\mathbb{X}})$ consisting of tilting bundles will be denoted by $\mathcal{G}(\mathcal{T}_{\mathbb{X}}^{\nu})$.

The connectedness of titling graph for weighted projective lines has been investigated widely in the literature through category aspect, see for example [3,22,25,27]. However, the explicit shape of the tilting graph is still unknown. By using the above geometric model, we provide a combinatorial description of $\mathcal{G}(\mathcal{T}_{\mathbb{X}}^{\nu})$.

Let $\Lambda_{(p,q)}$ be a graph with vertices

$$\Lambda^0_{(p,q)} = \{(c_1,\cdots,c_p) \in \mathbb{Z}^p | c_1 \leq \cdots \leq c_p \leq c_1 + q\},\$$

and there exists an edge between two vertex (c_1, \dots, c_p) and (d_1, \dots, d_p) if and only if

$$\sum_{i=1}^{p} |c_i - d_i| = 1.$$

Theorem 1.2. The tilting graph $\mathcal{G}(\mathcal{T}^{\nu}_{\mathbb{X}})$ coincides with the graph $\Lambda_{(p,q)}$.

Denote by η the bijection from $\Lambda_{(p,q)}$ to $\mathcal{G}(\mathcal{T}^{\nu}_{\mathbb{X}})$ obtained in Theorem 1.2. Let $H_{p,q} = \langle r_1, r_2 | r_1 r_2 = r_2 r_1, r_1^p = r_2^q \rangle$ and

$$\widetilde{H}_{p,q} = \begin{cases} H_{p,q}, & p \neq q; \\ H_{p,q} \times \mathbb{Z}_2, & p = q. \end{cases}$$

Then $\widetilde{H}_{p,q}$ coincides with the mapping class group $\mathcal{MG}(A_{p,q})$, hence there exists a group isomorphism from $\widetilde{H}_{p,q}$ to Aut(coh- $\mathbb{X}(p,q)$), still denoted by ψ . We construct an unexpected group action of $\widetilde{H}_{p,q}$ on the graph $\Lambda_{(p,q)}$ (c.f. Proposition 7.6), which is compatible with the group action of Aut(coh- $\mathbb{X}(p,q)$) on $\mathcal{G}(\mathcal{T}^{\nu}_{\mathbb{X}})$ in the following sense.

Theorem 1.3. For any $f \in \widetilde{H}_{p,q}$ and any vertex ν in $\Lambda_{(p,q)}$, we have

$$\eta(f(\nu)) = \psi(f)(\eta(\nu)). \tag{1.2}$$

Consequently, we have the following commutative diagram



The paper is organized as follows. In Section 2, we recall some basic facts on weighted projective lines of type (p,q). In Section 3, we show that the marked annulus $A_{p,q}$ gives a geometric model of the category $\operatorname{coh-X}(p,q)$. In Section 4, we establish a bijection between tilting sheaves in $\operatorname{coh-X}(p,q)$ and triangulations of $A_{p,q}$, and show that the flip of an arc is compatible with the tilting mutation of an indecomposable sheaf. Sections 5 and 6 focus on tilting mutation and tilting graphs, with the aim to prove Theorem 1.2. We give a geometric interpretation of the automorphism group of $\operatorname{coh-X}(p,q)$ in Section 7, and prove Theorem 1.1 and Theorem 1.3. In the final Section 8, we present a geometric description of the perpendicular category of an exceptional sheaf in $\operatorname{coh-X}(p,q)$.

2. WEIGHTED PROJECTIVE LINES OF TYPE (P,Q)

In this section, we recall from [23,36] some basic facts about the weighted projective lines of type (p,q) for $p,q \in \mathbb{Z}_{>1}$.

Let $\mathbb{L}(p,q)$ be the abelian group on generators \vec{x}_1, \vec{x}_2 with relations

$$p\vec{x}_1 = q\vec{x}_2 := \bar{c}$$

Then each $\vec{x} \in \mathbb{L}(p,q)$ can be uniquely written in normal form

$$\vec{x} = l_1 \vec{x}_1 + l_2 \vec{x}_2 + l\vec{c}$$
, where $0 \le l_1 \le p - 1, \ 0 \le l_2 \le q - 1$ and $l \in \mathbb{Z}$.

 $\mathbb{L}(p,q)$ is an ordered group whose cone of positive elements is $\mathbb{N}\vec{x}_1 + \mathbb{N}\vec{x}_2$. We equip $\mathbb{L}(p,q)$ with the structure of a partially ordered set: $\vec{x} \leq \vec{y}$ if and only if $\vec{y} - \vec{x} \in \mathbb{N}\vec{x}_1 + \mathbb{N}\vec{x}_2$.

Let **k** be an algebraically closed field and $\lambda = (\lambda_1, \lambda_2)$ be a sequence of pairwise distinct closed points on the projective line $\mathbb{P}^1_{\mathbf{k}}$. A weighted projective line $\mathbb{X}(p,q)$ of weight type (p,q)and parameter sequence λ is obtained from the projective line $\mathbb{P}^1_{\mathbf{k}}$ by attaching the weight p, q to λ_1, λ_2 , respectively. The parameter sequence can be normalized as $\lambda_1 = \infty$, $\lambda_2 = 0$.

The homogeneous coordinate algebra S(p,q) of the weighted projective line $\mathbb{X}(p,q)$ is given by $\mathbf{k}[x_1, x_2]$, which is $\mathbb{L}(p,q)$ -graded by means of deg $x_i = \vec{x}_i$ for i = 1, 2. That is, $S(p,q) = \bigoplus_{\vec{x} \in \mathbb{L}(p,q)} S(p,q)_{\vec{x}}$, where $S(p,q)_{\vec{x}}$ is the homogeneous component of degree \vec{x} . In particular, if we write $\vec{x} = l_1\vec{x}_1 + l_2\vec{x}_2 + l\vec{c}$ in its normal form, then $S(p,q)_{\vec{x}} \neq 0$ if and only if $l \geq 0$. Moreover, $\{x_1^{l_1+pa}x_2^{l_2+qb} \mid a+b=l, a, b \geq 0\}$ form a **k**-basis of $S(p,q)_{\vec{x}}$; see [23, Proposition 1.3].

We recall the definition of the category coh- $\mathbb{X}(p,q)$ of coherent sheaves over $\mathbb{X}(p,q)$ by a convenient description via graded S(p,q)-modules. Let $\mathrm{mod}^{\mathbb{L}(p,q)}$ -S(p,q) be the abelian category of finitely generated $\mathbb{L}(p,q)$ -graded S(p,q)-modules, and $\mathrm{mod}_{0}^{\mathbb{L}(p,q)}$ -S(p,q) be its Serre subcategory formed by finite dimensional modules. Denote by $\mathrm{qmod}^{\mathbb{L}(p,q)}$ -S(p,q) := $\mathrm{mod}^{\mathbb{L}(p,q)}$ - $S(p,q)/\mathrm{mod}_{0}^{\mathbb{L}(p,q)}$ -S(p,q) the quotient abelian category. By [23, Theorem 1.8] the sheafification functor yields an equivalence

$$\operatorname{qmod}^{\mathbb{L}(p,q)}$$
- $S(p,q) \xrightarrow{\sim} \operatorname{coh-} \mathbb{X}(p,q).$

From now on we will identify these two categories.

Notice that in this paper, we will abbreviate $\operatorname{Hom}_{\operatorname{coh}-\mathbb{X}(p,q)}(-,-)$ and $\operatorname{Ext}^{1}_{\operatorname{coh}-\mathbb{X}(p,q)}(-,-)$ as $\operatorname{Hom}(-,-)$ and $\operatorname{Ext}^{1}(-,-)$ respectively.

Proposition 2.1 ([23,36]). The category coh-X(p,q) is connected, Hom-finite and k-linear with the following properties:

- (1) coh- $\mathbb{X}(p,q)$ satisfies Serre duality in the form $\operatorname{Ext}^1(X,Y) \cong \operatorname{DHom}(Y,\tau(X))$, where the **k**-equivalence τ : coh- $\mathbb{X}(p,q) \longrightarrow$ coh- $\mathbb{X}(p,q)$ is the shift $X \mapsto X(\vec{\omega})$ with the dualizing element $\vec{\omega} = -(\vec{x}_1 + \vec{x}_2) \in \mathbb{L}(p,q)$.
- (2) coh-X(p,q) = vec-X(p,q) ∨ coh₀-X(p,q), where vec-X(p,q) denotes the full subcategory of coh-X(p,q) consisting of vector bundles over X(p,q), and coh₀-X(p,q) denotes the full subcategory of coh-X(p,q) consisting of all objects of finite length, ∨ means that each indecomposable object of coh-X(p,q) is either in vec-X(p,q) or in coh₀-X(p,q), and there are no non-zero morphism from coh₀-X(p,q) to vec-X(p,q).
- (3) Each indecomposable bundle over X(p,q) is a line bundle.
- (4) Each line bundle L is exceptional, i.e., $\operatorname{Hom}(L, L) \cong \mathbf{k}$ and $\operatorname{Ext}^{1}(L, L) = 0$.

(5) For any $\vec{x}, \vec{y} \in \mathbb{L}(p,q)$, there has

 $\operatorname{Hom}(\mathcal{O}(\vec{x}), \mathcal{O}(\vec{y})) \cong S(p, q)_{\vec{y} - \vec{x}}.$

In particular, $\operatorname{Hom}(\mathcal{O}(\vec{x}), \mathcal{O}(\vec{y})) \neq 0$ if and only if $\vec{x} \leq \vec{y}$.

(6) Denote by $K_0(\operatorname{coh}-\mathbb{X}(p,q))$ the Grothendieck group of the category $\operatorname{coh}-\mathbb{X}(p,q)$. Then the classes $[\mathcal{O}(\vec{x})]$ $(0 \le \vec{x} \le \vec{c})$ form a \mathbb{Z} -basis of $K_0(\operatorname{coh}-\mathbb{X}(p,q))$. Moreover, the rank of Grothendieck group $K_0(\operatorname{coh}-\mathbb{X}(p,q))$ equals p+q.

By [38, Proposition 1.1], the subcategory $\operatorname{coh}_0 X(p, q)$ decomposes into a coproduct $\coprod_{\lambda \in \mathbb{P}^1_k} \mathcal{U}_{\lambda}$ of connected uniserial length categories, whose associated Auslander-Reiten quiver is a stable tube $\mathbb{Z}A_{\infty}/(\tau^r)$ for some $r \in \mathbb{Z}_{\geq 1}$ (see e.g. [45]). Here, r is called the rank of the stable tube $\mathbb{Z}A_{\infty}/(\tau^r)$, which depends on λ . Precisely, r = p, q for $\lambda = \infty$, 0, respectively and r = 1 for $\lambda \in \mathbf{k}^* = \mathbf{k} \setminus \{0\}$. Objects that lie at the bottom of the stable tube are all simple objects of coh- $\mathbb{X}(p, q)$. A stable tube of r = 1 is called a *homogeneous stable tube*. Each $\lambda \in \mathbf{k}^*$ is associated with a unique simple sheaf S_{λ} , called *ordinary simple*; while $\lambda = \infty$ (*resp.* $\lambda = 0$) is associated with p (*resp.* q) simple sheaves $S_{\infty,i}$ ($i \in \mathbb{Z}/p\mathbb{Z}$) (*resp.* $S_{0,i}$ ($i \in \mathbb{Z}/q\mathbb{Z}$)) called *exceptional simples*.

Now we recall some important short exact sequences in coh- $\mathbb{X}(p,q)$. For each ordinary simple sheaf S_{λ} , there is an exact sequence

$$0 \longrightarrow \mathcal{O} \xrightarrow{u_{\lambda}} \mathcal{O}(\vec{c}) \longrightarrow S_{\lambda} \longrightarrow 0,$$

where the homomorphism $u_{\lambda} : \mathcal{O} \longrightarrow \mathcal{O}(\vec{c})$ is given by multiplication with $x_2^q - \lambda x_1^p$. By contrast, if $\lambda \in \{\infty, 0\}$, there are exact sequences

$$0 \longrightarrow \mathcal{O}((i-1)\vec{x}_1) \xrightarrow{u_{\infty}} \mathcal{O}(i\vec{x}_1) \longrightarrow S_{\infty,i} \longrightarrow 0, \quad i \in \mathbb{Z}/p\mathbb{Z};$$
(2.1)

$$0 \longrightarrow \mathcal{O}((i-1)\vec{x}_2) \xrightarrow{u_0} \mathcal{O}(i\vec{x}_2) \longrightarrow S_{0,i} \longrightarrow 0, \quad i \in \mathbb{Z}/q\mathbb{Z};$$
(2.2)

where u_{∞} (resp. u_0) is given by multiplication with x_1 (resp. x_2).

For each $\vec{x} = l_1 \vec{x}_1 + l_2 \vec{x}_2 \in \mathbb{L}(p, q)$, one has

$$S_{\lambda}(\vec{x}) = S_{\lambda} \text{ for } \lambda \in \mathbf{k}^*;$$

and

$$S_{\infty,i}(\vec{x}) = S_{\infty,i+l_1}, \ S_{0,i}(\vec{x}) = S_{0,i+l_2}.$$

For $\lambda \in \{\infty, 0\}$ and each $j \in \mathbb{Z}_{\geq 1}$, let $S_{\lambda,i}^{(j)}$ denote the indecomposable object in \mathcal{U}_{λ} of length j with top $S_{\lambda,i}$. More precisely, the composition series of $S_{\lambda,i}^{(j)}$ has the following form:

$$S_{\lambda,i-j+1} \hookrightarrow S_{\lambda,i-j+2}^{(2)} \hookrightarrow \cdots \hookrightarrow S_{\lambda,i-2}^{(j-2)} \hookrightarrow S_{\lambda,i-1}^{(j-1)} \hookrightarrow S_{\lambda,i}^{(j)}$$

with $S_{\lambda,i-k}^{(j-k)}/S_{\lambda,i-k-1}^{(j-k-1)} \cong S_{\lambda,i-k}$ for $0 \le k \le j-2$.

3. Geometric model for the category $\operatorname{coh-}\mathbb{X}(p,q)$

In this section, we aim to give a geometric model for $\operatorname{coh-X}(p,q)$ in terms of an annulus with p marked points on the inner boundary and q marked points on the outer boundary. We will establish a correspondence between indecomposable coherent sheaves and homotopy classes of oriented curves in the marked annulus and then study algebraic properties (e.g. Auslander-Reiten sequences, the dimension of extension space of degree 1, exact sequences) of $\operatorname{coh-X}(p,q)$ in geometric aspects. 3.1. The universal cover of $A_{p,q}$. Let $A_{p,q}$ be an annulus with p marked points on the inner boundary and q marked points on the outer boundary. Without loss of generality, assume $1 \le p \le q$. In this subsection, we recall from [9] for the universal cover of $A_{p,q}$.

Assume the marked points are distributed in equidistance on the two boundaries of $A_{p,q}$. We label the marked points on the inner boundary with 0_{∂} , $(\frac{1}{p})_{\partial}$, \cdots , $(\frac{p-1}{p})_{\partial}$ in an anticlockwise direction, and label the marked points on the outer boundary with $0_{\partial'}$, $(\frac{q-1}{q})_{\partial'}$, \cdots , $(\frac{1}{q})_{\partial'}$ in a clockwise direction; c.f. Figure 1.



FIGURE 1. Marked annulus $A_{p,q}$

Let $\operatorname{Cyl}_{p,q}$ be a rectangle of height 1 and width 1 in \mathbb{R}^2 with p marked points on the upper boundary ∂ and q marked points on the lower boundary ∂' , identifying its two vertical sides. Denote the points on the upper boundary of $\operatorname{Cyl}_{p,q}$ by i_{∂} , for $0 \leq i \leq \frac{p-1}{p}$:

$$0_{\partial} := (0,1), \ (\frac{1}{p})_{\partial} := (\frac{1}{p},1), \ \cdots, \ (\frac{p-1}{p})_{\partial} := (\frac{p-1}{p},1),$$

and denote the points on the lower boundary of $\operatorname{Cyl}_{p,q}$ by $j_{\partial'}$, for $0 \leq j \leq \frac{q-1}{q}$:

$$0_{\partial'} := (0,0), \ (\frac{1}{q})_{\partial'} := (\frac{1}{q},0), \ \cdots, \ (\frac{q-1}{q})_{\partial'} := (\frac{q-1}{q},0).$$

Then $A_{p,q}$ can be identified with $\operatorname{Cyl}_{p,q}$ in the sense of viewing the upper (*resp.* lower) boundary of $\operatorname{Cyl}_{p,q}$ as the inner (*resp.* outer) boundary of $A_{p,q}$, where the marked points of $\operatorname{Cyl}_{p,q}$ are oriented from left to right on the upper boundary ∂ and from right to left on the lower boundary ∂' , as in Figure 2.



FIGURE 2. Annulus via rectangle Cyl_{p,q}

It will be most convenient to work with in the universal cover $(\mathbb{U}, \pi_{p,q})$ of $\operatorname{Cyl}_{p,q}$, where $\mathbb{U} = \{(x, y) \in \mathbb{R}^2 | 0 \le y \le 1\}$ inherits the orientation of $\operatorname{Cyl}_{p,q}$, and the covering map

$$\pi := \pi_{p,q} : \mathbb{U} \to \operatorname{Cyl}_{p,q}, \ (x,y) \mapsto (x - \lfloor x \rfloor, y)$$

Here, $\lfloor x \rfloor$ denotes the integer part of x. Naturally, π is also a covering map of $A_{p,q}$.

Denote the marked point (i, 1) on the upper boundary ∂ of \mathbb{U} by i_{∂} and the marked point (j, 0) on the lower boundary ∂' of \mathbb{U} by $j_{\partial'}$, for $i \in \frac{\mathbb{Z}}{n}$ and $j \in \frac{\mathbb{Z}}{n}$, see Figure 3.



FIGURE 3. Universal cover \mathbb{U} of $A_{p,q}$

3.2. Arcs in $A_{p,q}$ and \mathbb{U} . In this subsection, we recall some definitions of curves and arcs in $A_{p,q}$ and \mathbb{U} . In this paper, we consider curves and arcs up to homotopy. The following definition refers to [11, 47].

Definition 3.1. A curve in \mathbb{U} is a continuous function $c : [0, 1] \longrightarrow \mathbb{U}$ such that $c(t) \in \mathbb{U}^0 := \mathbb{U} \setminus \{\partial, \partial'\}$ for any $t \in (0, 1)$. An arc in \mathbb{U} is a curve whose endpoints are marked points of \mathbb{U} , satisfying that it does not intersect itself in the interior of \mathbb{U} , the interior of the arc is disjoint from the boundary of \mathbb{U} , and it does not cut out a monogon or digon.

If an arc in \mathbb{U} starts at a marked point x_{b_1} and ends at a marked point y_{b_2} with $x, y \in \frac{\mathbb{Z}}{p}$ or $\frac{\mathbb{Z}}{q}$, $b_i \in \{\partial, \partial'\}$ for i = 1, 2, it will be denoted by $[x_{b_1}, y_{b_2}]$. The following definition refers to [9].

Definition 3.2. Let $\alpha = [x_{b_1}, y_{b_2}]$ be an arc in U. If $b_1 \neq b_2$, α is called a *bridging arc*. If $b_1 = b_2 = \partial$ and $y - x \ge \frac{2}{p}$, or $b_1 = b_2 = \partial'$ and $y - x \ge \frac{2}{q}$, then α is called a *peripheral arc*.

In order to give a geometric model for line bundles in coh- $\mathbb{X}(p,q)$, we only need to consider the bridging arcs that oriented from the boundary ∂' to the boundary ∂ . Such an arc $[x_{\partial'}, y_{\partial}]$ is called a *positive bridging arc*.

Similarly, one can define curves and arcs in the marked annulus $A_{p,q}$. Note that the image of an arc in \mathbb{U} under $\pi : \mathbb{U} \to A_{p,q}$ maybe intersect itself in the interior of $A_{p,q}$. We define bridging (resp. peripheral) curve in $A_{p,q}$ as the image $\pi(\alpha)$ for a bridging (resp. peripheral) arc α in \mathbb{U} . If $\pi(\alpha)$ is an arc additionally, then it will be called a bridging (resp. peripheral) arc in $A_{p,q}$.

In order to give a geometric model for homogeneous torsion sheaves in the subcategory $\coprod_{\lambda \in \mathbf{k}^*} \mathcal{U}_{\lambda}$ of coh- $\mathbb{X}(p,q)$, we introduce the notion of \mathbf{k}^* -parameterized *n*-cycles as follows.

Definition 3.3. For $n \in \mathbb{Z}_{\geq 1}$, an *n*-cycle in $A_{p,q}$ is a curve $\pi(c)$, where *c* is a curve in \mathbb{U}^0 with c(1) - c(0) = (n, 0). In particular, the 1-cycle will be called a *loop*. For the notion of \mathbf{k}^* -parameterized *n*-cycles we refer to the set $\{(\lambda, L^n) \mid \lambda \in \mathbf{k}^*\}$, where *L* is a loop in $A_{p,q}$ with the orientation in an anti-clockwise direction.

Recall from Proposition 2.1 that the category $\operatorname{coh-X}(p,q)$ contains two parts: vector bundles and torsion sheaves. Each indecomposable vector bundle is a line bundle. The torsion sheaves consists of two stable tubes of rank p and q respectively, and a family of homogeneous tubes indexed by \mathbf{k}^* .

Let $\operatorname{ind}(\operatorname{coh}-\mathbb{X}(p,q))$ denote the full subcategory of $\operatorname{coh}-\mathbb{X}(p,q)$ consisting of all indecomposable objects. Then the objects of $\operatorname{ind}(\operatorname{coh}-\mathbb{X}(p,q))$ are line bundles $\mathcal{O}(\vec{x})$ ($\vec{x} \in \mathbb{L}(p,q)$), and torsion sheaves $S_{\infty,i}^{(j)}$ ($i \in \mathbb{Z}/p\mathbb{Z}$), $S_{0,i}^{(j)}$ ($i \in \mathbb{Z}/q\mathbb{Z}$) and $S_{\lambda}^{(j)}$ ($\lambda \in \mathbf{k}^*$) for $j \in \mathbb{Z}_{\geq 1}$. Now we can state the main result of this subsection.

Theorem 3.4. A parametrization of the isoclasses of ind(coh-X(p,q)) is given as follows:

- (1) the indecomposable vector bundles are indexed by the homotopy classes of positive bridging curves $\pi([(\frac{j}{q})_{\partial'}, (\frac{i}{p})_{\partial}])$ in $A_{p,q}$;
- (2) the objects in the stable tube of rank p (resp. q) are indexed by the homotopy classes of peripheral curves $\pi([(\frac{i}{p})_{\partial}, (\frac{j}{p})_{\partial}])$ (resp. $\pi([(\frac{i}{q})_{\partial'}, (\frac{j}{q})_{\partial'}]))$ in $A_{p,q}$;
- (3) the objects in homogeneous tubes are indexed by \mathbf{k}^* -parameterized n-cycles in $A_{p,q}$.

Proof. Recall that $\pi : \mathbb{U} \to A_{p,q}$ is the covering map. Let \mathcal{C} be the set consisting of the following elements with $i, j \in \mathbb{Z}$ and $n \geq 1$:

- positive bridging curves: $\pi([(\frac{i}{q})_{\partial'}, (\frac{i}{p})_{\partial}]);$
- peripheral curves: $\pi([(\frac{i}{p})_{\partial}, (\frac{j}{p})_{\partial}])$ and $\pi([(\frac{i}{q})_{\partial'}, (\frac{j}{q})_{\partial'}])$, with $j i \ge 2$;
- **k***-parameterized *n*-cycles: $\{(\lambda, L^n) | \lambda \in \mathbf{k}^*\}$.

Then the following assignments:

$$\pi([(\frac{j}{q})_{\partial'}, (\frac{i}{p})_{\partial}]) \mapsto \mathcal{O}(i\vec{x}_1 - j\vec{x}_2), \quad \pi([(\frac{i-j-1}{p})_{\partial}, (\frac{i}{p})_{\partial}]) \mapsto S_{\infty,i}^{(j)},$$
(3.1)

$$\pi([(\frac{-i}{q})_{\partial'}, \ (\frac{j-i+1}{q})_{\partial'}]) \mapsto S_{0,i}^{(j)}, \quad (\lambda, L^j) \mapsto S_{\lambda}^{(j)},$$

define a bijective map $\phi : \mathcal{C} \longrightarrow \operatorname{ind}(\operatorname{coh-}\mathbb{X}(p,q))$. This proves Theorem 3.4.

3.3. Elementary moves. Basing on the bijection given in Theorem 3.4, we consider the geometric interpretation of Auslander-Reiten sequences in coh- $\mathbb{X}(p,q)$. Firstly, we introduce the notion of elementary moves of positive bridging curves and peripheral curves in $A_{p,q}$. The following definition refers to [9, 11].

Definition 3.5. For any positive bridging curve or peripheral curve γ in $A_{p,q}$, denote by ${}_{s}\gamma$ the elementary move of γ on its starting point, meaning that the curve ${}_{s}\gamma$ is obtained from γ moving its starting point to the next marked point on the same boundary, such that ${}_{s}\gamma$ is rotated in clockwise direction around the ending point of γ . Similarly, denote by γ_e the elementary move of γ on its ending point. Iterated elementary moves are denoted ${}_{s}\gamma_e = {}_{s}(\gamma_e) = ({}_{s}\gamma)_{e}, {}_{s}{}_{2}\gamma = {}_{s}({}_{s}\gamma)$ and $\gamma_{e^2} = (\gamma_e)_e$, respectively.

For the follow-up discussion, we recall from [9] for the intuitional expression of elementary moves of positive bridging curves and peripheral curves in $A_{p,q}$.

(1) If γ is a positive bridging curve in $A_{p,q}$, then $\gamma = \pi([(\frac{i}{q})_{\partial'}, (\frac{j}{p})_{\partial}])$ for some $i, j \in \mathbb{Z}$, and

$${}_{s}\gamma = \pi([(\frac{i-1}{q})_{\partial'}, (\frac{j}{p})_{\partial}]), \quad \gamma_{e} = \pi([(\frac{i}{q})_{\partial'}, (\frac{j+1}{p})_{\partial}]).$$

The corresponding picture in \mathbb{U} is:



(2) If γ is a peripheral curve in $A_{p,q}$ with endpoints on the inner boundary, then $\gamma = \pi([(\frac{i}{p})_{\partial}, (\frac{j}{p})_{\partial}])$ for some $i, j \in \mathbb{Z}, j-i \geq 2$, and

$${}_s\gamma = \pi([(\frac{i+1}{p})_{\partial}, (\frac{j}{p})_{\partial}]) \ (j-i>2), \qquad \gamma_e = \pi([(\frac{i}{p})_{\partial}, (\frac{j+1}{p})_{\partial}]).$$

In \mathbb{U} , they look as follows:



(3) If γ is a peripheral curve in $A_{p,q}$ with endpoints on the outer boundary, then $\gamma = \pi([(\frac{i}{q})_{\partial'}, (\frac{j}{q})_{\partial'}])$ for some $i, j \in \mathbb{Z}, j - i \geq 2$, and

$${}_{s}\gamma = \pi([(\frac{i-1}{q})_{\partial'}, (\frac{j}{q})_{\partial'}]), \quad \gamma_{e} = \pi([(\frac{i}{q})_{\partial'}, (\frac{j-1}{q})_{\partial'}]) \ (j-i>2).$$

In \mathbb{U} , they are described as follows:



For the sake of simplicity, from now on, we denote by

$$D_{\frac{j}{q}}^{\frac{i}{p}} := [(\frac{j}{q})_{\partial'}, (\frac{i}{p})_{\partial}], \quad D^{\frac{i}{p}, \frac{j}{p}} := [(\frac{i}{p})_{\partial}, (\frac{j}{p})_{\partial}] \quad \text{and} \quad D_{\frac{i}{q}, \frac{j}{q}} := [(\frac{i}{q})_{\partial'}, (\frac{j}{q})_{\partial'}]$$

for positive bridging arcs and peripheral arcs in \mathbb{U} , and denote their image under $\pi : \mathbb{U} \to A_{p,q}$ as $[D_{\frac{i}{q}}^{\frac{i}{p}}], [D^{\frac{i}{p},\frac{j}{p}}]$ and $[D_{\frac{i}{q},\frac{j}{q}}]$ respectively. It follows that for any $k \in \mathbb{Z}$,

$$[D_{\frac{i}{p}+k}^{\frac{i}{p}+k}] = [D_{\frac{j}{q}}^{\frac{i}{p}}], \quad [D^{\frac{i}{p}+k,\frac{j}{p}+k}] = [D^{\frac{i}{p},\frac{j}{p}}] \quad \text{and} \quad [D_{\frac{i}{q}+k,\frac{j}{q}+k}] = [D_{\frac{i}{q},\frac{j}{q}}]. \tag{3.2}$$

Proposition 3.6. Let $X \in ind(coh-\mathbb{X}(p,q))$ be a line bundle or a torsion sheaf supported at an exceptional point. Assume $\phi^{-1}(X) = \gamma$. Then the irreducible morphisms in coh- $\mathbb{X}(p,q)$ starting at X are obtained by elementary moves on endpoints of γ . Moreover, the Auslander-Reiten sequence starting at X has the form

$$0 \longrightarrow \phi(\gamma) \longrightarrow \phi(s\gamma) \oplus \phi(\gamma_e) \longrightarrow \phi(s\gamma_e) \longrightarrow 0$$

Proof. If X is a line bundle, then $X = \mathcal{O}(i\vec{x}_1 - j\vec{x}_2)$ for some $i, j \in \mathbb{Z}$, and then $\gamma = [D_{\frac{j}{q}}^{\frac{1}{p}}]$. Notice that the irreducible morphisms in coh- $\mathbb{X}(p,q)$ starting at $\mathcal{O}(i\vec{x}_1 - j\vec{x}_2)$ are given by

$$\mathcal{O}(i\vec{x}_1 - j\vec{x}_2) \longrightarrow \mathcal{O}((i+1)\vec{x}_1 - j\vec{x}_2) = \phi(\gamma_e)$$

and

$$\mathcal{O}(i\vec{x}_1 - j\vec{x}_2) \longrightarrow \mathcal{O}(i\vec{x}_1 - (j-1)\vec{x}_2) = \phi(s\gamma),$$

and

$$\tau^{-1}(\mathcal{O}(i\vec{x}_1 - j\vec{x}_2)) = (\mathcal{O}(i\vec{x}_1 - j\vec{x}_2))(\vec{x}_1 + \vec{x}_2) = \mathcal{O}((i+1)\vec{x}_1 - (j-1)\vec{x}_2) = \phi(_s\gamma_e).$$

Then the result follows.

If X is a torsion sheaf supported at ∞ , then $X = S_{\infty,i}^{(j)}$ for some $i \in \mathbb{Z}/p\mathbb{Z}$ and $j \in \mathbb{Z}_{\geq 1}$, and then $\gamma = [D^{\frac{i-j-1}{p},\frac{i}{p}}]$. If j = 1, we have $\phi(_{s}\gamma) = 0$ and there is only one irreducible morphism starting at $S_{\infty,i}^{(1)}$, that is, $S_{\infty,i}^{(1)} \longrightarrow S_{\infty,i+1}^{(2)} = \phi(\gamma_e)$. If $j \geq 2$, there are two irreducible morphisms starting at $S_{\infty,i}^{(j)}$, given by $S_{\infty,i}^{(j)} \longrightarrow S_{\infty,i+1}^{(j+1)} = \phi(\gamma_e)$ and $S_{\infty,i}^{(j)} \longrightarrow S_{\infty,i}^{(j-1)} = \phi(_{s}\gamma)$. Moreover, $\tau^{-1}(S_{\infty,i}^{(j)}) = S_{\infty,i+1}^{(j)} = \phi(_{s}\gamma_e)$. Similarly, if X is a torsion sheaf supported at the exceptional point 0, we can also obtain the result. We are done.

Remark 3.7. For any indecomposable torsion sheaf X supported at an ordinary point $\lambda \in \mathbf{k}^*$, we know that $X = S_{\lambda}^{(j)}$ for some $j \in \mathbb{Z}_{\geq 1}$. By Theorem 3.4, we have $\phi^{-1}(S_{\lambda}^{(j)}) = (\lambda, L^j)$. If we denote by ${}_{s}(\lambda, L^k) = (\lambda, L^{k-1}), \ (\lambda, L^k)_e = (\lambda, L^{k+1}) \text{ and } {}_{s}(\lambda, L^k)_e = {}_{s}((\lambda, L^k)_e) = ({}_{s}(\lambda, L^k))_e = (\lambda, L^k)$, then Proposition 3.6 holds for any $X \in \operatorname{ind}(\operatorname{coh-X}(p,q))$.

3.4. Translation quiver and equivalence between categories. In this subsection, we define a translation quiver $(\Gamma_{A_{p,q}}, \tau')$ associated to \mathcal{C} . The vertices of the quiver are the oriented curves in \mathcal{C} . There is an arrow $\gamma \to \gamma'$ if and only if $\gamma' = {}_{s}\gamma$ or γ_{e} . For the positive bridging curves and peripheral curves in \mathcal{C} , let the translation τ' be induced by the map $(\frac{i}{p})_{\partial} \mapsto (\frac{i-1}{p})_{\partial}$ and $(\frac{j}{q})_{\partial'} \mapsto (\frac{j+1}{q})_{\partial'}$ for $i, j \in \mathbb{Z}$, that is,

$$\tau'([D_{\frac{j}{q}}^{\frac{i}{p}}]) = [D_{\frac{j+1}{q}}^{\frac{i-1}{p}}], \ \tau'([D^{\frac{i}{p},\frac{j}{p}}]) = [D^{\frac{i-1}{p},\frac{j-1}{p}}], \ \tau'([D_{\frac{i}{q},\frac{j}{q}}]) = [D_{\frac{i+1}{q},\frac{j+1}{q}}].$$

For the **k**^{*}-parameterized k-cycles (λ, L^k) in \mathcal{C} , let $\tau'((\lambda, L^k)) = (\lambda, L^k)$.

Let $\mathcal{C}_{A_{p,q}}$ be the mesh category of the translation quiver $(\Gamma_{A_{p,q}}, \tau')$. Let $F : \mathcal{C}_{A_{p,q}} \longrightarrow$ ind(coh- $\mathbb{X}(p,q)$) be the functor between mesh categories defined as follows. On objects, let $F(\gamma) = \phi(\gamma)$. To define F on morphisms, it suffices to define it on the elementary moves. Define $F(\gamma \to {}_{s}\gamma)$ (resp. $F(\gamma \to \gamma_{e})$) to be the irreducible morphism $F(\gamma) \to F({}_{s}\gamma)$ (resp. $F(\gamma) \to F(\gamma_{e})$).

Let $(\Gamma_{\operatorname{coh}-\mathbb{X}(p,q)}, \tau)$ denote the Auslander-Reiten quiver of the category coh- $\mathbb{X}(p,q)$.

Theorem 3.8. The functor $F : \mathcal{C}_{A_{p,q}} \longrightarrow \operatorname{ind}(\operatorname{coh-}\mathbb{X}(p,q))$ is an equivalence of categories. In particular,

- (1) F induces an isomorphism of translation quivers $(\Gamma_{A_{p,q}}, \tau') \longrightarrow (\Gamma_{\operatorname{coh-}\mathbb{X}(p,q)}, \tau).$
- (2) τ' corresponds to the Auslander-Reiten translation τ in the following sense

$$F \circ \tau' = \tau \circ F.$$

(3) F is an exact functor with respect to the induced abelian structure on $\mathcal{C}_{A_{n,a}}$.

Proof. By Theorem 3.4 and Proposition 3.6, we get the statement (1). For the statement (2), we only consider the case $F \circ \tau'([D_{\frac{i}{q}}^{\frac{i}{p}}]) = \tau \circ F([D_{\frac{i}{q}}^{\frac{i}{p}}])$, the others being similar. In fact,

$$F \circ \tau'([D_{\frac{j}{q}}^{\frac{i}{p}}]) = F([D_{\frac{j+1}{q}}^{\frac{i-1}{p}}]) = \mathcal{O}((i-1)\vec{x}_1 - (j+1)\vec{x}_2)$$
$$= \tau(\mathcal{O}(i\vec{x}_1 - j\vec{x}_2))$$
$$= \tau \circ F([D_{\frac{j}{q}}^{\frac{i}{p}}]).$$

Therefore, the statement (2) holds.

Since both categories $C_{A_{p,q}}$ and $\operatorname{ind}(\operatorname{coh-X}(p,q))$ are the mesh categories of their translation quivers $(\Gamma_{A_{p,q}}, \tau')$ and $(\Gamma_{\operatorname{coh-X}(p,q)}, \tau)$ respectively, therefore, statement (1) implies that F is an equivalence of categories. In particular, this equivalence induces the structure of an abelian category on $C_{A_{p,q}}$. With respect to this structure, F is an exact functor since every equivalence between abelian categories is exact.

3.5. **Positive intersection.** In this subsection, we want to give a geometric explanation for the dimension of extension group $\text{Ext}^{i}(X, Y)$ for $X, Y \in \text{ind}(\text{coh-}\mathbb{X}(p, q))$. Since $\text{coh-}\mathbb{X}(p, q)$ is hereditary, the extension group is zero for $i \geq 2$. Hence we only need to consider the case i = 1.

Note that if X, Y are supported at distinct points, then $\operatorname{Ext}^1(X, Y) = 0$. If they are supported at the same ordinary point, then $X = S_{\lambda}^{(j_1)}, Y = S_{\lambda}^{(j_2)}$ for some $\lambda \in \mathbf{k}^*$ and $j_1, j_2 \in \mathbb{Z}_{\geq 1}$. In this case, $\dim_{\mathbf{k}}\operatorname{Ext}^1(X, Y) = \min\{j_1, j_2\}$. Hence, we only consider the remaining cases in the following. We will show that $\dim_{\mathbf{k}}\operatorname{Ext}^1(X, Y)$ can be interpreted as the positive intersection number between the corresponding oriented curves in the marked annulus $A_{p,q}$.

Definition 3.9 ([48]). A point of intersection of two oriented curves α and β in $A_{p,q}$ is called *positive intersection* of α and β , if β intersects α from the right, i.e., the picture looks like as:



FIGURE 4. Positive intersection

Denoted by

$$I^{+}(\alpha,\beta) = \min_{\alpha' \sim \alpha, \beta' \sim \beta} |\alpha' \cap^{+} \beta'|,$$

called the *positive intersection number* of α, β , where $\alpha' \cap^+ \beta'$ is the set of the positive intersections of α' and β' , excluding their endpoints, and the sign $a \sim b$ means that the two curves are homotopy. It follows that $I^+(\alpha, \beta) = 0$ or 1 if α, β are oriented arcs in \mathbb{U} .

Theorem 3.10. Let X, Y be two indecomposable objects in coh-X(p,q). Assume X, Y are not supported at ordinary points. Then

$$\dim_{\mathbf{k}} \operatorname{Ext}^{1}(X, Y) = I^{+}(\phi^{-1}(X), \phi^{-1}(Y)).$$
(3.3)

Proof. If $X, Y \in \mathcal{U}_{\infty} \coprod \mathcal{U}_0$, then the result follows from [7, Theorem 3.8]. If $X \in \text{vec-X}(p,q)$ and $Y \in \mathcal{U}_{\infty} \coprod \mathcal{U}_0$, then according to Proposition 2.1 (2), we have $\text{Ext}^1(X,Y) = 0$, and by Definition 3.9, we also have $I^+(\phi^{-1}(X), \phi^{-1}(Y)) = 0$. So we only need to consider the following two cases:

(1) $X, Y \in \text{vec-}\mathbb{X}(p, q).$

By Proposition 2.1 (3), we know that each indecomposable bundle over $\mathbb{X}(p,q)$ is a line bundle, thus $X = \mathcal{O}(\vec{x})$ and $Y = \mathcal{O}(\vec{y})$ for some $\vec{x}, \vec{y} \in \mathbb{L}(p,q)$. Write $\vec{x} = l_1\vec{x}_1 + l_2\vec{x}_2 + l\vec{c}$ and $\vec{y} = k_1\vec{x}_1 + k_2\vec{x}_2 + k\vec{c}$ in normal forms. According to Proposition 2.1 (1) and (5), we have

$$\operatorname{Ext}^{1}(\mathcal{O}(\vec{x}), \mathcal{O}(\vec{y})) \cong D\operatorname{Hom}(\mathcal{O}(\vec{y}), \mathcal{O}(\vec{x} - \vec{x}_{1} - \vec{x}_{2})) \cong S_{\vec{x} - \vec{x}_{1} - \vec{x}_{2} - \vec{y}}.$$

Observe that $\vec{x} - \vec{x}_1 - \vec{x}_2 - \vec{y} = (l_1 - k_1 - 1)\vec{x}_1 + (l_2 - k_2 - 1)\vec{x}_2 + (l - k)\vec{c}$, and $-p \le l_1 - k_1 - 1 \le p - 2$, $-q \le l_2 - k_2 - 1 \le q - 2$. Denote $h_1 := l_1 - k_1 - 1$ and $h_2 := l_2 - k_2 - 1$, we get

$$\dim_{\mathbf{k}} \operatorname{Ext}^{1}(X,Y) = \begin{cases} l-k+1 & \text{if } 0 \leq h_{1} \leq p-2, \ 0 \leq h_{2} \leq q-2, \ k \leq l; \\ l-k & \text{if } -p \leq h_{1} \leq -1, \ 0 \leq h_{2} \leq q-2, \ k \leq l; \\ l-k & \text{if } 0 \leq h_{1} \leq p-2, \ -q \leq h_{2} \leq -1, \ k \leq l; \\ l-k-1 & \text{if } -p \leq h_{1} \leq -1, \ -q \leq h_{2} \leq -1, \ k+1 \leq l; \\ 0 & \text{otherwise.} \end{cases}$$

On the other hand, we consider $I^+(\phi^{-1}(X), \phi^{-1}(Y))$. Since $\phi^{-1}(X) = [D_{-\frac{l_2}{q}-l}^{\frac{l_1}{p}}]$ and $\phi^{-1}(Y) = [D_{-\frac{l_2}{q}-k}^{\frac{k_1}{p}}]$. By similar arguments as in [7], a point of intersection of $\phi^{-1}(X)$ and $\phi^{-1}(Y)$ is positive if and only if there exists $m \in \mathbb{Z}$ such that $\frac{l_1}{p} + m > \frac{k_1}{p}$ and $-\frac{l_2}{q} - l + m < -\frac{k_2}{q} - k$ (c.f. Figure 5).



FIGURE 5.

These two inequalities can be combined into

$$\frac{k_1 - l_1}{p} < m < \frac{q(l-k) + l_2 - k_2}{q}$$

Therefore,

$$I^{+}(\phi^{-1}(X),\phi^{-1}(Y)) = |\{m \in \mathbb{Z} | \frac{k_{1}-l_{1}}{p} < m < \frac{q(l-k)+l_{2}-k_{2}}{q} \}|.$$

In order to prove that $\dim_{\mathbf{k}} \operatorname{Ext}^{1}(X, Y) = I^{+}(\phi^{-1}(X), \phi^{-1}(Y))$, we only consider the case: $0 \leq h_{1} \leq p-2, \ 0 \leq h_{2} \leq q-2, \ k \leq l$; the others being similar. In this case, we have

$$I^{+}(\phi^{-1}(X),\phi^{-1}(Y)) = |\{m \in \mathbb{Z} | 0 \le m \le l-k\}| = l-k+1 = \dim_{\mathbf{k}} \operatorname{Ext}^{1}(X,Y).$$

We are done.

(2) $X \in \mathcal{U}_{\infty} \coprod \mathcal{U}_0$ and $Y \in \text{vec-}\mathbb{X}(p,q)$.

We only consider the case $X \in \mathcal{U}_{\infty}$, that is, $X = S_{\infty,i}^{(j)}$ for $i \in \mathbb{Z}/p\mathbb{Z}$, $j \in \mathbb{Z}_{\geq 1}$, the case $X \in \mathcal{U}_0$ being similar.

By Proposition 2.1 (3), $Y = \mathcal{O}(\vec{x})$ for some $\vec{x} \in \mathbb{L}(p,q)$, write $\vec{x} = l_1\vec{x}_1 + l_2\vec{x}_2 + l\vec{c}$ in normal form. By exact sequence (2.1), we get $\dim_{\mathbf{k}} \operatorname{Hom}(\mathcal{O}(\vec{x} + \vec{x}_1 + \vec{x}_2), S_{\infty,l_1+1}) = 1$. Notice that $S_{\infty,i}^{(j)}$ has a composition series of the form

$$S_{\infty,i-j+1} \hookrightarrow S_{\infty,i-j+2}^{(2)} \hookrightarrow \cdots \hookrightarrow S_{\infty,i-2}^{(j-2)} \hookrightarrow S_{\infty,i-1}^{(j-1)} \hookrightarrow S_{\infty,i}^{(j)}$$

with $S_{\infty,i-k}^{(j-k)}/S_{\infty,i-k-1}^{(j-k-1)} \cong S_{\infty,i-k}$ for $0 \le k \le j-2$. Write j = np+r with $0 \le r \le p-1$. We get

$$\dim_{\mathbf{k}} \operatorname{Ext}^{1}(S_{\infty,i}^{(j)}, \mathcal{O}(\vec{x})) = \begin{cases} n & \text{if } p \nmid i - l_{1} - k \text{ for each } 1 \leq k \leq r, \\ n+1 & \text{otherwise.} \end{cases}$$

On the other hand, we consider $I^+(\phi^{-1}(S^{(j)}_{\infty,i}), \phi^{-1}(\mathcal{O}(\vec{x})))$. Since

$$\phi^{-1}(S_{\infty,i}^{(j)}) = [D^{\frac{i-j-1}{p},\frac{i}{p}}], \qquad \phi^{-1}(\mathcal{O}(\vec{x})) = [D^{\frac{i_1}{p}}_{-\frac{l_2}{q}-l}].$$

By similar arguments as in [7], a point of intersection of $\phi^{-1}(S_{\infty,i}^{(j)})$ and $\phi^{-1}(\mathcal{O}(\vec{x}))$ is positive if and only if there exists $m \in \mathbb{Z}$ such that $\frac{i-j-1}{p} < \frac{l_1}{p} + m < \frac{i}{p}$, (c.f. Figure 6).



FIGURE 6.

Hence

$$I^{+}(\phi^{-1}(S_{\infty,i}^{(j)}),\phi^{-1}(\mathcal{O}(\vec{x}))) = |\{m \in \mathbb{Z} | i - j - l_{1} \le pm \le i - l_{1} - 1\}|.$$

Observe that

$$\lfloor \frac{(i-l_1-1)-(i-j-l_1)+1}{p} \rfloor = \lfloor \frac{j}{p} \rfloor = \lfloor \frac{np+r}{p} \rfloor = n + \lfloor \frac{r}{p} \rfloor = n.$$

It follows that

$$I^+(\phi^{-1}(S^{(j)}_{\infty,i}),\phi^{-1}(\mathcal{O}(\vec{x}))) = \begin{cases} n & \text{if } p \nmid i - l_1 - k \text{ for each } 1 \le k \le r, \\ n+1 & \text{otherwise.} \end{cases}$$

Hence $\dim_{\mathbf{k}} \operatorname{Ext}^{1}(X, Y) = I^{+}(\phi^{-1}(X), \phi^{-1}(Y))$. This finishes the proof.

Remark 3.11. If one of X, Y is supported at an ordinary point, then (3.3) still holds.

Proposition 3.12. Let α, β be positive bridging arcs or peripheral arcs in \mathbb{U} with $I^+(\alpha, \beta) =$ 1. Then there is a natural short exact sequence in $\operatorname{coh-X}(p,q)$ associated to the intersection in geometric terms. More precisely, consider the following figure of the chosen intersection,



FIGURE 7. Short exact sequence

we have a short exact sequence

$$0 \longrightarrow \phi(\pi(\beta)) \longrightarrow \phi(\pi(\gamma_1)) \oplus \phi(\pi(\gamma_2)) \longrightarrow \phi(\pi(\alpha)) \longrightarrow 0.$$
(3.4)

Proof. We discuss case by case according to the types of α and β . If α , β are peripheral arcs, then the result follows from [4, Section 6] and [48, Remark 4.25]. Hence we only consider the following two cases.

(1) α , β are positive bridging arcs.

Then $\alpha = D_{\frac{j}{q}}^{\frac{i}{p}}$ and $\beta = D_{\frac{l}{q}}^{\frac{k}{p}}$ for some integers k < i and j < l. Hence $\gamma_1 = D_{\frac{j}{q}}^{\frac{k}{p}}$ and $\gamma_2 = D_{\frac{l}{q}}^{\frac{i}{p}}$. Note that

 $\phi(\pi(\alpha)) = \mathcal{O}(i\vec{x}_1 - j\vec{x}_2), \quad \phi(\pi(\beta)) = \mathcal{O}(k\vec{x}_1 - l\vec{x}_2).$

And we have the following short exact sequence in $\operatorname{coh-}\mathbb{X}(p,q)$:

$$0 \longrightarrow \mathcal{O}(k\vec{x}_1 - l\vec{x}_2) \longrightarrow \mathcal{O}(k\vec{x}_1 - j\vec{x}_2) \oplus \mathcal{O}(i\vec{x}_1 - l\vec{x}_2) \longrightarrow \mathcal{O}(i\vec{x}_1 - j\vec{x}_2) \longrightarrow 0.$$

Then (3.4) holds since

$$\phi(\pi(\gamma_1)) = \mathcal{O}(k\vec{x}_1 - j\vec{x}_2), \quad \phi(\pi(\gamma_2)) = \mathcal{O}(i\vec{x}_1 - l\vec{x}_2).$$

(2) α is a peripheral arc and β is a positive bridging arc.

We only consider the case that both the endpoints of α are in the upper boundary of \mathbb{U} , the other being similar.

By assumption, $\alpha = D^{\frac{i-j-1}{p}, \frac{i}{p}}$ and $\beta = D^{\frac{k}{p}}_{\frac{l}{q}}$ for some $i, j, k, l \in \mathbb{Z}$ with i - j - 1 < k < i. Then we have

$$\phi(\pi(\alpha)) = S_{\infty,i}^{(j)}, \ \phi(\pi(\beta)) = \mathcal{O}(k\vec{x}_1 - l\vec{x}_2).$$

In coh- $\mathbb{X}(p,q)$, there is a short exact sequence:

$$0 \longrightarrow \mathcal{O}(k\vec{x}_1 - l\vec{x}_2) \longrightarrow \mathcal{O}(i\vec{x}_1 - l\vec{x}_2) \oplus S^{(k+j-i)}_{\infty,k} \longrightarrow S^{(j)}_{\infty,i} \longrightarrow 0.$$

Therefore, the short exact sequence (3.4) holds.

4. The geometric interpretation of tilting sheaf

In this section, we investigate the correspondence between tilting sheaves in the category $\operatorname{coh-X}(p,q)$ and triangulations in the marked annulus $A_{p,q}$, and then study the relationship between the flip of an arc in a triangulation and the mutation of indecomposable direct summand of the corresponding tilting sheaf.

4.1. Tilting sheaves and triangulations. First let us recall from [23,25] for the definition of tilting sheaves and tilting bundles in coh-X(p,q) as follows.

Definition 4.1. A sheaf T in $\operatorname{coh-} \mathbb{X}(p,q)$ is called a *tilting sheaf* if

- (1) T is rigid, that is, $\text{Ext}^1(T,T) = 0$.
- (2) For any object $X \in \operatorname{coh-X}(p,q)$, the condition $\operatorname{Hom}(T,X) = 0 = \operatorname{Ext}^1(T,X)$ implies that X = 0.

Moreover, if a tilting sheaf T is given by a vector bundle, i.e., T has no direct summand of finite length, then T is called a *tilting bundle*.

Proposition 4.2 ([26]). Assume that \mathcal{H} is a hereditary abelian category with a tilting object. Then T is a tilting object in \mathcal{H} if and only if $\operatorname{Ext}^1(T,T) = 0$ and the number of non-isomorphic indecomposable direct summands of T equals the rank of the Grothendieck group $K_0(\mathcal{H})$.

Geigle-Lenzing [23] constructed a canonical tilting sheaf for any weighted projective line. As a consequence, the rank of the Grothendieck group $K_0(\operatorname{coh-X}(p,q))$ equals to p+q, and any tilting sheaf in $\operatorname{coh-X}(p,q)$ contains p+q indecomposable direct summands. A rigid sheaf in $\operatorname{coh-X}(p,q)$ with p+q-1-many (pairwise non-isomorphic) indecomposable direct summands is called an *almost complete tilting sheaf*. Let \overline{T} be an almost complete tilting sheaf in $\operatorname{coh-X}(p,q)$. If there exists an indecomposable sheaf E, such that $\overline{T} \oplus E$ is a tilting sheaf, then E is called a *complement* of \overline{T} .

Recall from [1] that a triangulation Γ of $A_{p,q}$ is a maximal collection of arcs that do not intersect in the interior of $A_{p,q}$. According to Proposition [1, Proposition 2.1], any triangulation Γ of $A_{p,q}$ consists of p+q-many arcs. In our model, we always fix the orientations for the arcs in Γ , namely, we take the arcs from the set \mathcal{C} appeared in Theorem 3.4. For example, all the bridging arcs are oriented from the outer boundary to the inner boundary.

Theorem 4.3. The tilting sheaves in the category $\operatorname{coh-X}(p,q)$ are in natural bijection with the triangulations of $A_{p,q}$.

Proof. Let Γ be a triangulation of $A_{p,q}$ and $\gamma_1, \gamma_2, \cdots, \gamma_{p+q}$ be the set of arcs in Γ . Let

$$T = \bigoplus_{i=1}^{p+q} \phi(\gamma_i).$$

According to Theorem 3.10 and the definition of triangulation, we have

$$\dim_{\mathbf{k}} \operatorname{Ext}^{1}(\phi(\gamma_{i}), \phi(\gamma_{j})) = I^{+}(\gamma_{i}, \gamma_{j}) = 0$$

for $1 \le i, j \le p + q$. It follows that $\dim_{\mathbf{k}} \operatorname{Ext}^{1}(T, T) = 0$. Combining with Proposition 4.2 and Propositions 2.1 (6), we get that $T = \bigoplus_{i=1}^{p+q} \phi(\gamma_{i})$ is a tilting sheaf in coh- $\mathbb{X}(p,q)$.

On the other hand, if $T = \bigoplus_{i=1}^{p+q} T_i$ is a tilting sheaf in coh- $\mathbb{X}(p,q)$, then we have $\operatorname{Ext}^1(T,T) = 0$. By Theorem 3.10, $\{\phi^{-1}(T_1), \phi^{-1}(T_2), \cdots, \phi^{-1}(T_{p+q})\}$ is a collection of arcs that do not intersect in the interior of $A_{p,q}$, which provides a triangulation of $A_{p,q}$ by Proposition [1, Proposition 2.1]. Therefore, we complete the proof of Theorem.

Remark 4.4. The following general result has been stated in [31] for weighted projective lines of arbitrary type. We give a short proof for weight type (p, q) by using the correspondence in Theorem 4.3.

Corollary 4.5. Let \overline{T} be an almost complete tilting sheaf in the category coh- $\mathbb{X}(p,q)$. Then there exist precisely two complements of \overline{T} .

Proof. By the proof of Theorem 4.3, we see that \overline{T} can be represented by p + q - 1 nonintersecting arcs in \mathcal{C} . To get a triangulation of $A_{p,q}$, we should add just one further arc, which must be a diagonal in a quadrilateral. Since there are exactly two diagonals in a quadrilateral, we get two arcs belong to \mathcal{C} by attaching suitable orientations and represent the two complements of \overline{T} .

4.2. Flips and mutations. In this subsection, we study the compatibility of the flip of an arc in a triangulation and the mutation of indecomposable direct summand of the corresponding tilting sheaf.

Definition 4.6. ([25]) Let $T = \overline{T} \oplus X$ and $T' = \overline{T} \oplus X'$ be two tilting sheaves in coh- $\mathbb{X}(p,q)$, where X and X' are two non-isomorphic indecomposable objects in coh- $\mathbb{X}(p,q)$. Then T' is called the *mutation* of T at X and denoted by $T' = \mu_X(T)$.

In this paper, we call X' the *mutation* of X w.r.t T and denote by $\mu_T(X) = X'$.

Let γ be an arc of a triangulation Γ of $A_{p,q}$. Then γ is one diagonal of the quadrilateral formed by the two triangles of Γ that contain γ . Recall from [1] that the *flip* of γ replaces

the arc γ by the other diagonal γ' of the same quadrilateral (c.f. Figure 8). Keeping all other arcs unchanged, one obtains a new triangulation $\mu_{\gamma}(\Gamma)$. For convenient, we call γ' the *flip* of γ w.r.t Γ , and denote by $\gamma' = \mu_{\Gamma}(\gamma)$.



FIGURE 8. Flip of an arc γ

For a triangulation Γ of $A_{p,q}$, by choosing suitable orientation for bridging arcs, we can always assume $\mu_{\gamma}(\Gamma)$ is also a triangulation of $A_{p,q}$. By Theorem 4.3, we know that a triangulation Γ corresponds to a tilting sheaf T in coh- $\mathbb{X}(p,q)$, and $\phi(\gamma)$ is an indecomposable direct summand of T. The following proposition shows that the flip of γ and the mutation of $\phi(\gamma)$ are compatible.

Proposition 4.7. Let Γ be a triangulation of $A_{p,q}$ and γ be an arc in Γ . Then

$$\phi(\mu_{\Gamma}(\gamma)) = \mu_T(\phi(\gamma)).$$

Proof. Let γ , γ_1 , γ_2 , \cdots , γ_{p+q-1} be the set of arcs in the triangulation Γ . Therefore, $\mu_{\gamma}(\Gamma)$ is a triangulation of $A_{p,q}$ with arcs γ' , γ_1 , γ_2 , \cdots , γ_{p+q-1} . By Theorem 4.3, we get two tilting sheaves

$$T = \phi(\gamma) \oplus \phi(\gamma_1) \oplus \phi(\gamma_2) \oplus \dots \oplus \phi(\gamma_{p+q-1})$$

and

$$T' = \phi(\gamma') \oplus \phi(\gamma_1) \oplus \phi(\gamma_2) \oplus \dots \oplus \phi(\gamma_{p+q-1})$$

in coh- $\mathbb{X}(p,q)$. It follows that $\phi(\mu_{\Gamma}(\gamma)) = \phi(\gamma') = \mu_T(\phi(\gamma))$.

5. Tilting bundles and bundle-mutations

In this section, we use the geometric model to investigate tilting bundles in the category $\operatorname{coh-X}(p,q)$, basing on Theorem 4.3 and Proposition 4.7. Denote by

 $\mathcal{T}^{\nu}_{\mathbb{X}} := \{ \text{Tilting bundles in coh-} \mathbb{X}(p,q) \}.$

5.1. A combinatorial description of $\mathcal{T}_{\mathbb{X}}^{\nu}$. Let $T = \bigoplus_{i=1}^{p+q} T_i$ be a tilting bundle in coh- $\mathbb{X}(p,q)$.

Under Theorem 3.4, we can view indecomposable objects in the category $\operatorname{coh-X}(p,q)$ as oriented curves in the marked annulus $A_{p,q}$. Hence each T_i corresponds to a positive bridging arc $[D_{b_i}^{a_i}]$, where $a_i \in \frac{\mathbb{Z}}{p}$ and $b_i \in \frac{\mathbb{Z}}{q}$ for $1 \leq i \leq p+q$. Moreover, by Theorem 4.3, T corresponds to a triangulation Γ with arcs $[D_{b_i}^{a_i}]$, $1 \leq i \leq p+q$. Therefore, $\{D_{b_i}^{a_i}, 1 \leq i \leq p+q\}$ is a triangulation of a parallelogram in \mathbb{U} . Thanks to (3.2), without loss of generality we may assume T has the following normal form:

$$0 = a_1 < a_2 \le \dots \le a_{p+q} = 1, \quad b_1 \le b_2 \le \dots \le b_{p+q}.$$
 (5.1)

Convention 1. From now on, for a tilting bundle $T = \bigoplus_{i=1}^{p+q} T_i$ in coh- $\mathbb{X}(p,q)$, we always assume $T = \bigoplus_{i=1}^{p+q} [D_{b_i}^{a_i}]$ satisfying (5.1), and denote by $\mu_i(T)$ for the mutation of T at T_i , where *i* is taken modulo p + q. In particular, $\mu_{i-1}(T) = \mu_{p+q}(T)$ for i = 1, and $\mu_{i+1}(T) = \mu_1(T)$ for i = p + q.

We will give a combinatorial description for the set of tilting bundles $\mathcal{T}_{\mathbb{X}}^{\nu}$ in coh- $\mathbb{X}(p,q)$. For this, we consider the following set

$$\Lambda^{0}_{(p,q)} := \{ (c_1, c_2, \cdots, c_p) \in \mathbb{Z}^p | c_1 \le c_2 \le \cdots \le c_p \le c_1 + q \}.$$

Theorem 5.1. There exists a bijection between $\mathcal{T}^{\nu}_{\mathbb{X}}$ and $\Lambda^{0}_{(p,q)}$.

Proof. For any tilting bundle T, by Theorem 4.3, T corresponds to a triangulation Γ of (a parallelogram in) U. Then for any $1 \le i \le p$, the segment $[(\frac{i-1}{p})_{\partial}, (\frac{i}{p})_{\partial}]$ belongs to a unique triangle in Γ :



FIGURE 9. *p*-triangles

Therefore, we obtain a sequence $(c_1, c_2, \dots, c_p) \in \mathbb{Z}^p$. Since Γ is a triangulation, we obtain that $\frac{c_1}{q} \leq \frac{c_2}{q} \cdots \leq \frac{c_p}{q} \leq \frac{c_1}{q} + 1$, i.e., $c_1 \leq c_2 \cdots \leq c_p \leq c_1 + q$. That is, $(c_1, c_2, \dots, c_p) \in \Lambda^0_{(p,q)}$. This defines a map

$$\varphi: \mathcal{T}^{\nu}_{\mathbb{X}} \longrightarrow \Lambda^{0}_{(p,q)}; \quad T \mapsto (c_{1}, c_{2}, \cdots, c_{p}).$$
(5.2)

By Theorem 4.3, we know that φ is injective. Now we show that it is surjective.

For any $(c_1, c_2, \dots, c_p) \in \Lambda^0_{(p,q)}$, we can obtain *p*-triangles as in Figure 9. For any marked point $c_{\partial'}$ satisfying $\frac{c_i}{q} < c < \frac{c_{i+1}}{q}$, there exists a unique bridging arc connecting to c which does not intersect with the above *p*-triangles, namely, the arc $D_c^{\frac{1}{p}}$ (see the following picture):



Therefore, the p-triangles in Figure 9 can be extended to a unique triangulation Γ of a parallelogram in U. Then by the bijection between tilting bundles and triangulations, we obtain a tilting bundle T, which satisfies $\varphi(T) = (c_1, c_2, \cdots, c_p)$ by the construction. Hence φ is surjective. We are done. 5.2. Bundle-mutations. Let T and T' be two tilting sheaves in coh- $\mathbb{X}(p,q)$ such that T' is a mutation of T. If T, T' are both tilting bundles, then such a mutation is called a *bundle-mutation*, cf. [25].

Proposition 5.2. Let $T = \bigoplus_{i=1}^{p+q} T_i$ be a tilting bundle in coh- $\mathbb{X}(p,q)$.

(1) $\mu_i(T)$ is a tilting bundle if and only if the positions of the arcs in \mathbb{U} corresponding to T_{i-1}, T_i, T_{i+1} exactly satisfy one of the following two conditions:



(2) $\mu_i(T)$ is not a tilting bundle if and only if the positions of the arcs in \mathbb{U} corresponding to T_{i-1}, T_i, T_{i+1} exactly satisfy one of the following two conditions:



Proof. The necessity is obvious. We only prove the sufficiency. By Proposition 4.7, we have T_i and $\mu_T(T_i)$ are two diagonals of the same quadrilateral. If $\mu_i(T)$ is a tilting bundle, then $\mu_T(T_i)$ is a line bundle, it follows that $\mu_T(T_i)$ corresponds to a positive bridging arc of $A_{p,q}$ by Theorem 3.4. Thus the statement (1) holds. If $\mu_i(T)$ is not a tilting bundle, then $\mu_T(T_i)$ corresponds to a peripheral arc of $A_{p,q}$ by Theorem 3.4. Therefore, the statement (2) holds.

Proposition 5.3. Assume $T = \bigoplus_{i=1}^{p+q} T_i$ is a tilting bundle in coh- $\mathbb{X}(p,q)$, and $\mu_i(T)$ is also a tilting bundle for some $1 \le i \le p+q$. Then

- (1) $\mu_{i-1}(T)$ is a tilting bundle if and only if $\mu_{i-1}(\mu_i(T))$ is not a tilting bundle;
- (2) $\mu_{i+1}(T)$ is a tilting bundle if and only if $\mu_{i+1}(\mu_i(T))$ is not a tilting bundle.

Proof. We only prove the statement (1), the statement (2) is similar.

If $\mu_{i-1}(T)$ is a tilting bundle, by Proposition 5.2, the position of the arcs in \mathbb{U} corresponding to $T_{i-2}, T_{i-1}, T_i, T_{i+1}$ exactly satisfies one of the following two conditions:



It is easy to see that $\mu_{i-1}(\mu_i(T))$ is not a tilting bundle in both situation.

If $\mu_{i-1}(\mu_i(T))$ is not a tilting bundle, we claim that $\mu_{i-1}(T)$ is a tilting bundle. For contradiction, we assume $\mu_{i-1}(T)$ is not a tilting bundle, by Proposition 5.2, the position of

the arcs in \mathbb{U} corresponding to $T_{i-2}, T_{i-1}, T_i, T_{i+1}$ exactly satisfies one of the following two conditions:



It follows that $\mu_{i-1}(\mu_i(T))$ is a tilting bundle in both cases, a contradiction. We are done. \Box

5.3. Bundle-mutation index n(T). For any tilting bundle $T = \bigoplus_{i=1}^{p+q} T_i$ in coh- $\mathbb{X}(p,q)$, let

 $I(T) = \{i | \mu_i(T) \text{ is a tilting bundle}\},\$

and define the bundle-mutation index of T as n(T) = |I(T)|.

Under the following bijection from (5.2),

$$\varphi: \mathcal{T}^{\nu}_{\mathbb{X}} \longrightarrow \Lambda^{0}_{(p,q)}; \quad T \mapsto (c_1, c_2, \cdots, c_p),$$

we define

$$J(T) = \{i | c_i = c_{i+1} \text{ for } 1 \le i \le p-1\}$$

and denote by r(T) = |J(T)|.

Proposition 5.4. For any tilting bundle T in coh-X(p,q), we have n(T) = 2(p - r(T)). In particular, $2 \le n(T) \le 2p$.

Proof. For any tilting bundle T, there exists a unique $(c_1, c_2, \dots, c_p) \in \Lambda^0_{(p,q)}$ associated to T by Theorem 5.1, moreover, T corresponds to the following triangulation of (a parallelogram in) U:



We denote by $m_T(T_i) = 1$ if $\mu_i(T)$ is a tilting bundle and $m_T(T_i) = 0$ otherwise. Combining the above figure and Proposition 5.2, we get

$$m_T([D_a^b]) = \begin{cases} 1, & \text{if } a = \frac{c_i}{q}, \ b = \frac{i-1}{p} \text{ or } b = \frac{i}{p} \text{ for some } i; \\ 0, & \text{otherwise.} \end{cases}$$

It follows that n(T) = 2(p - r(T)) and then $2 \le n(T) \le 2p$.

By Proposition 5.4, we have the following corollary.

Corollary 5.5. Keep notations as above. Let T be a tilting bundle in $\operatorname{coh-X}(p,q)$. Then

- (1) n(T) = 2 if and only if $c_1 = c_2 = \cdots = c_p$.
- (2) n(T) = 2p if and only if $c_1 < c_2 < \cdots < c_p$.

Remark 5.6. Let T be a tilting bundle, which corresponds to a triangulation Γ of $A_{p,q}$ by Theorem 4.3. Let Q_{Γ} be the quiver associated to the triangulation Γ . Then n(T) = 2 if and only if $\mathbf{k}Q_{\Gamma}$ is a canonical algebra of type (p,q).

For any $a \in \frac{\mathbb{Z}}{p}$, $b \in \frac{\mathbb{Z}}{q}$, denote by

$$T_{b}^{a} = [D_{b}^{a}] \oplus [D_{b}^{a+\frac{1}{p}}] \oplus [D_{b+\frac{1}{q}}^{a+\frac{1}{p}}] \oplus [D_{b+\frac{1}{q}}^{a+\frac{2}{p}}] \oplus \dots \oplus [D_{b+\frac{p-1}{q}}^{a+1}] \oplus [D_{b+\frac{p}{q}}^{a+1}] \oplus \dots \oplus [D_{b+\frac{q-1}{q}}^{a+1}].$$

Intuitively, T_b^a corresponds to the following triangulation of a parallelogram in \mathbb{U} up to shift:



We have the following observation.

Proposition 5.7. For any $a \in \frac{\mathbb{Z}}{p}$, $b \in \frac{\mathbb{Z}}{q}$, T_b^a is a tilting bundle with $n(T_b^a) = 2p$. Moreover,

$$\mu_{2p-1} \cdot \mu_{2p-3} \cdots \mu_1(T_b^a) = T_{b-\frac{1}{q}}^a \text{ and } \mu_{2p} \cdot \mu_{2p-2} \cdots \mu_2(T_b^a) = T_{b+\frac{1}{q}}^a$$

Proof. The first result follows from Theorem 4.3 and Proposition 5.4 immediately. Note that the triangulation $\mu_{2p-1} \cdot \mu_{2p-3} \cdots \mu_1(T_b^a)$ can be obtained from that of T_b^a (black arcs and green arcs) by iterated mutations, and each mutation μ_{2i-1} for $1 \le i \le p$ replaces one green arc by a red arc as below:



More precisely,

$$\begin{split} &\mu_{2p-1} \cdot \mu_{2p-3} \cdots \mu_1(T_b^a) \\ &= [D_{b+\frac{q-1}{q}}^{a+\frac{p+1}{p}}] \oplus [D_b^{a+\frac{1}{p}}] \oplus [D_b^{a+\frac{2}{p}}] \oplus \cdots \oplus [D_{b+\frac{p-2}{q}}^{a+1}] \oplus \cdots \oplus [D_{b+\frac{q-1}{q}}^{a+1}] \\ &= [D_{b-\frac{1}{q}}^a] \oplus [D_{b-\frac{1}{q}}^{a+\frac{1}{p}}] \oplus [D_b^{a+\frac{1}{p}}] \oplus [D_b^{a+\frac{2}{p}}] \oplus \cdots \oplus [D_{b+\frac{p-2}{q}}^{a+1}] \oplus \cdots \oplus [D_{b+\frac{q-2}{q}}^{a+1}] \\ &= T_{b-\frac{1}{q}}^a. \end{split}$$

Similarly, one can obtain the other equation. We are done.

The tilting bundle T_b^a plays an important role in the next subsection for the connectedness of the tilting graph of vector bundles category.

5.4. Connectedness of the tilting graph $\mathcal{G}(\mathcal{T}_{\mathbb{X}}^{\nu})$. Recall from [3,25] that the *tilting graph* $\mathcal{G}(\mathcal{T}_{\mathbb{X}})$ of coh- $\mathbb{X}(p,q)$ has as vertices the isomorphism classes of tilting sheaves in coh- $\mathbb{X}(p,q)$, while two vertices are connected by an edge if and only if the associated tilting sheaves differ by precisely one indecomposable direct summand. The *tilting graph* $\mathcal{G}(\mathcal{T}_{\mathbb{X}}^{\nu})$ of vec- $\mathbb{X}(p,q)$ is the full subgraph of $\mathcal{G}(\mathcal{T}_{\mathbb{X}})$ consisting of tilting bundles.

The connectedness of titling graph for weighted projective lines has been investigated widely in the literature through category aspect, see for example [3, 22, 25, 27]. In this subsection, we investigate the connectedness of $\mathcal{G}(\mathcal{T}_{\mathbb{X}}^{\nu})$ by using our geometric model.

Let T and T' be tilting bundles in coh- $\mathbb{X}(p,q)$. We say T is bundle-mutations to T' if there is a sequence of tilting bundles $T = T^0, T^1, \dots, T^{n-1}, T^n = T'$ such that T^i is a mutation of T^{i-1} for any $1 \leq i \leq n$. Under the bijection Theorem 3.4, we call a triangulation Γ is bundle-flips to Γ' if the associated tilting bundles T and T' satisfy that T is bundle-mutations to T'.

By Convention 1, we know that the direct summands of any tilting bundle T can be arranged in a unique order. Hence we can define a map ι from the set of tilting bundles in $\operatorname{coh-X}(p,q)$ to \mathbb{Z}_2^{p+q} , in the sense that $\iota(T)_i = 1$ if $\mu_i(T)$ is a tilting bundle, and $\iota(T)_i = 0$ otherwise. For example, in the case p = 2 and q = 2, if $\mu_1(T), \mu_2(T)$ are tilting bundles and $\mu_3(T), \mu_4(T)$ are not tilting bundles, we have $\iota(T) = (1, 1, 0, 0)$.

Lemma 5.8. Let T be a tilting bundle in $\operatorname{coh-X}(p,q)$ and assume

$$\iota(T) = (\underbrace{1, \cdots, 1}_{s_1 \ times}, \underbrace{0, \cdots, 0}_{t_1 \ times}, \underbrace{1, \cdots, 1}_{s_2 \ times}, \underbrace{0, \cdots, 0}_{t_2 \ times}).$$

Then T is bundle-mutations to some tilting bundle T' satisfying

$$\iota(T') = (\underbrace{1, \cdots, 1}_{m_1 \ times}, \underbrace{0, \cdots, 0}_{n_1 \ times}, \underbrace{1, \cdots, 1}_{m_2 \ times})$$

and $m_1 + m_2 \ge s_1 + s_2$.

Proof. Note that $p + q = s_1 + t_1 + s_2 + t_2$. We divide the proof into two cases by considering s_2 is even or not.

If s_2 is even, then $s_2 = 2s$ for some $s \in \mathbb{Z}_{\geq 1}$. Denote by

$$\mu_{[a,b]} = \mu_a \cdot \mu_{a-1} \cdots \mu_b, \quad b \le a.$$

Let

$$\nu_k = \mu_{[p+q+1-2k, \ p+q-t_2+2-2k]}, \ 1 \le k \le s.$$

By iterative use of Proposition 5.3, we obtain

$$\iota(\nu_s \cdot \nu_{s-1} \cdots \nu_1(T)) = (\underbrace{1, \cdots, 1}_{s_1 \text{ times}}, \underbrace{0, \cdots, 0}_{t_1+t_2 \text{ times}}, \underbrace{1, \cdots, 1}_{s_2 \text{ times}}),$$

and $\nu_s \cdot \nu_{s-1} \cdots \nu_1(T)$ is a tilting bundle.

If s_2 is odd, then $s_2 = 2s + 1$ for some $s \in \mathbb{Z}_{\geq 0}$. We have

$$\iota(\nu_s \cdot \nu_{s-1} \cdots \nu_1(T)) = (\underbrace{1, \cdots, 1}_{s_1 \text{ times}}, \underbrace{0, \cdots, 0}_{t_1 \text{ times}}, 1, \underbrace{0, \cdots, 0}_{t_2 \text{ times}}, \underbrace{1, \cdots, 1}_{2s \text{ times}}).$$

If $t_2 \geq t_1$, let

$$\delta_k = \mu_{[p+q-2s-2k, p+q-t_2-2s+1-k]}, \quad 1 \le k \le t_1.$$

$$\iota(\delta_{t_1} \cdot \delta_{t_1-1} \cdots \delta_1 \cdot \nu_s \cdot \nu_{s-1} \cdots \nu_1(T)) = (\underbrace{1, \cdots, 1}_{s_1+1 \text{ times}}, \underbrace{0, \cdots, 0}_{t_2-t_1 \text{ times}}, \underbrace{1, \cdots, 1}_{2(s+t_1) \text{ times}}),$$

and $s_1 + 1 + 2(s + t_1) = s_1 + s_2 + t_1 \ge s_1 + s_2$. If $t_2 < t_1$, let

$$\delta_k = \mu_{s_1+2k} \cdot \mu_{s_1+2k+1} \cdots \mu_{s_1+t_1+k}, \ 1 \le k \le t_2.$$

Then

$$\iota(\delta_{t_2} \cdot \delta_{t_2-1} \cdots \delta_1 \cdot \nu_s \cdot \nu_{s-1} \cdots \nu_1(T)) = (\underbrace{1, \cdots, 1}_{s_1+2t_2 \text{ times}}, \underbrace{0, \cdots, 0}_{t_1-t_2 \text{ times}}, \underbrace{1, \cdots, 1}_{s_2 \text{ times}}),$$

and $s_1 + 2t_2 + s_2 \ge s_1 + s_2$. We are done.

Now we can state the main result of this subsection.

Proposition 5.9. Any tilting bundle T in coh- $\mathbb{X}(p,q)$ is bundle-mutations to T_b^a for some $a \in \frac{\mathbb{Z}}{p}, b \in \frac{\mathbb{Z}}{q}$. Consequently, the tilting graph $\mathcal{G}(\mathcal{T}_{\mathbb{X}}^{\nu})$ is connected.

Proof. According to equations (3.2) and Proposition 5.4, we can always assume $\mu_1(T)$ is a tilting bundle. Hence $\iota(T)$ must be one of the following two forms by considering $\mu_{p+q}(T)$ is a tilting bundle or not:

$$\iota(T) = (\underbrace{1, \cdots, 1}_{s_1 \text{ times}}, \underbrace{0, \cdots, 0}_{t_1 \text{ times}}, \underbrace{1, \cdots, 1}_{s_2 \text{ times}}, \underbrace{0, \cdots, 0}_{t_2 \text{ times}}, \underbrace{1, \cdots, 1}_{s_3 \text{ times}}, \underbrace{1, \cdots, 1}_{s_m \text{ times}}),$$

or

$$\iota(T) = (\underbrace{1, \cdots, 1}_{s_1 \text{ times}}, \underbrace{0, \cdots, 0}_{t_1 \text{ times}}, \underbrace{1, \cdots, 1}_{s_2 \text{ times}}, \underbrace{0, \cdots, 0}_{t_2 \text{ times}}, \underbrace{1, \cdots, 1}_{s_3 \text{ times}}, \underbrace{1, \cdots, 1}_{s_m \text{ times}}, \underbrace{0, \cdots, 0}_{t_m \text{ times}}).$$

Using Lemma 5.8, we can reduce $\iota(T)$ to the following shapes:

$$\underbrace{(\underbrace{1,\cdots,1}_{m'_1 \text{ times}},\underbrace{0,\cdots,0}_{n'_1 \text{ times}},\underbrace{1,\cdots,1}_{m'_2+s_3 \text{ times}},\underbrace{0,\cdots,0}_{t_3 \text{ times}},\underbrace{1,\cdots,1}_{s_m \text{ times}})}_{s_m \text{ times}}$$

or

$$\underbrace{(\underbrace{1,\cdots,1}_{m_1' \text{ times}},\underbrace{0,\cdots,0}_{n_1' \text{ times}},\underbrace{1,\cdots,1}_{m_2'+s_3 \text{ times}},\underbrace{0,\cdots,0}_{t_3 \text{ times}},\cdots,\underbrace{1,\cdots,1}_{s_m \text{ times}},\underbrace{0,\cdots,0}_{t_m \text{ times}})$$

with $m'_1 + m'_2 \ge s_1 + s_2$ respectively.

By iterating the above process, one finally obtains that T is bundle-mutations to a tilting bundle T' such that

$$\iota(T') = (\underbrace{1, \cdots, 1}_{m_1 \text{ times}}, \underbrace{0, \cdots, 0}_{n_1 \text{ times}}, \underbrace{1, \cdots, 1}_{m_2 \text{ times}})$$

and $m_1 + m_2 \ge s_1 + s_2 + \dots + s_m$. Assume $T' = \bigoplus_{i=1}^{p+q} [D_{b_i}^{a_i}]$ with $0 = a_1 < a_2 \le \dots \le a_{p+q} = 1, \ b_1 \le b_2 \le \dots \le b_{p+q}$. According to (3.2), we have

$$T' = \left(\bigoplus_{i=1}^{m_1+n_1} [D_{b_i}^{a_i}]\right) \oplus \left(\bigoplus_{i=m_1+n_1+1}^{p+q} [D_{b_i}^{a_i}]\right) = \left(\bigoplus_{i=m_1+n_1+1}^{p+q} [D_{b_i-1}^{a_i-1}]\right) \oplus \left(\bigoplus_{i=1}^{m_1+n_1} [D_{b_i}^{a_i}]\right),$$

which implies that

$$\iota(T') = (\underbrace{1, \cdots, 1}_{m_1 + m_2 \text{ times}}, \underbrace{0, \cdots, 0}_{n_1 \text{ times}}).$$

By the proof of Proposition 5.4, we get $T' = T_b^a$ for some $a \in \frac{\mathbb{Z}}{p}$, $b \in \frac{\mathbb{Z}}{q}$. Combining with Proposition 5.7, we obtain the connectedness of $\mathcal{G}(\mathcal{T}_{\mathbb{X}}^{\nu})$. This finishes the proof. \Box

6. TILTING GRAPH

In this section, we will give more explicit structure for the tilting graph $\mathcal{G}(\mathcal{T}_{\mathbb{X}})$ of coh- $\mathbb{X}(p,q)$ and the tilting graph $\mathcal{G}(\mathcal{T}_{\mathbb{X}}^{\nu})$ of vec- $\mathbb{X}(p,q)$ respectively by using the geometric model.

6.1. Local shape of the tilting graphs. In this subsection, we consider the local shape of the tilting graphs $\mathcal{G}(\mathcal{T}_{\mathbb{X}})$ and $\mathcal{G}(\mathcal{T}_{\mathbb{X}})$ respectively.

For the case p = q = 1, i.e., for the classical projective line $\mathbb{P}^1_{\mathbf{k}}$ case, it is well known that any tilting sheaf has the form $T_n := \mathcal{O}(n) \oplus \mathcal{O}(n+1)$ for some $n \in \mathbb{Z}$. Moreover, there exists a tilting mutation between T_n and T_m if and only if $m = n \pm 1$. Therefore, the tilting subgraph $\mathcal{G}(\mathcal{T}^{\nu}_{\mathbb{X}})$ coincides with the tilting graph $\mathcal{G}(\mathcal{T}_{\mathbb{X}})$, which has the form:

FIGURE 10. Tilting graph of coh- $\mathbb{P}^1_{\mathbf{k}}$

From now on we focus on the weighted projective line $\mathbb{X}(p,q)$ with $(p,q) \neq (1,1)$.

. . .

Proposition 6.1. Assume $(p,q) \neq (1,1)$. Let $T = \bigoplus_{i=1}^{p+q} T_i$ be a tilting sheaf in coh- $\mathbb{X}(p,q)$ and Γ be the associated triangulation of $A_{p,q}$. Let γ_i be the arc in Γ corresponding to T_i for $1 \leq i \leq p+q$. For any $1 \leq i \neq j \leq p+q$, one of the following holds:

(1) if two arcs γ_i and γ_j are in the same triangle, then $\mu_i \mu_j \mu_i(T) = \mu_i \mu_j(T)$;

(2) if two arcs γ_i and γ_j are not in the same triangle, then $\mu_i \mu_j(T) = \mu_j \mu_i(T)$.

Proof. For any arc γ_i in Γ , there exists a quadrilateral formed by the two triangles of Γ containing γ_i , drawn as follows:



(1) If two arcs γ_i and γ_j are in the same triangle, without loss of generality, we may assume that the position of γ_i and γ_j is drawn as below:



By applying the flips μ_i and μ_j on the corresponding arcs, and noting that $(p,q) \neq (1,1)$, we have the following commutative diagram:



Therefore $\mu_i \mu_j \mu_i(T) = \mu_i \mu_j(T)$.

(2) If two arcs γ_i and γ_j are not in the same triangle, then the mutations $\mu_T(T_i)$ and $\mu_T(T_j)$ do not affect each other. Therefore, we have $\mu_i \mu_j(T) = \mu_j \mu_i(T)$.

As an immediate consequence of the above proposition, we obtain the following result, c.f. [21, Theorem 3.10]).

Corollary 6.2. Assume $(p,q) \neq (1,1)$, then the tilting graph $\mathcal{G}(\mathcal{T}_{\mathbb{X}})$ is composed of quadrilaterals and pentagons.

By considering the tilting subgraph $\mathcal{G}(\mathcal{T}^{\nu}_{\mathbb{X}})$ of tilting bundles, we have the following result.

Proposition 6.3. (1) The tilting graph $\mathcal{G}(\mathcal{T}^{\nu}_{\mathbb{X}})$ of vec- $\mathbb{X}(1,q)$ is a line.

(2) For $2 \leq p \leq q$, the tilting graph $\mathcal{G}(\mathcal{T}^{\nu}_{\mathbb{X}})$ of vec- $\mathbb{X}(p,q)$ is composed of quadrilaterals.

Proof. (1) If X has weight type (1,q), then by Theorem 4.3 $T = T_b^0$ for some $b \in \frac{\mathbb{Z}}{q}$, and the bundle-mutation index n(T) = 2 by Proposition 5.4. Combining with Proposition 5.7, we obtain that the tilting graph $\mathcal{G}(\mathcal{T}_{\mathbb{X}}^{\nu})$ of vec-X(1,q) is a line.

(2) Assume X has weight type (p,q) with $2 \leq p \leq q$. Let $T = \bigoplus_{i=1}^{p+q} T_i$ be a tilting bundle. dle. Assume $\mu_i(T)$ and $\mu_j(T)$ are both tilting bundles for some i, j, let γ_i and γ_j be the corresponding arcs of T_i and T_j respectively. If $\mu_j\mu_i(T)$ is also a tilting bundle, then γ_i and γ_j are not in the same triangle by Proposition 5.3. It follows that $\mu_i\mu_j(T) = \mu_j\mu_i(T)$ by Proposition 6.1 (2). Hence, the tilting graph $\mathcal{G}(\mathcal{T}^{\nu}_{\mathbb{X}})$ of vector bundle category is composed of quadrilaterals.

6.2. Proof of Theorem 1.2. Recall that $\Lambda_{(p,q)}$ is the graph with vertex set

$$\Lambda^{0}_{(p,q)} = \{ (c_1, \cdots, c_p) \in \mathbb{Z}^p | c_1 \le \cdots \le c_p \le c_1 + q \},\$$

and there exists an edge between two vertices (c_1, \dots, c_p) and (d_1, \dots, d_p) if and only if $\sum_{i=1}^{p} |c_i - d_i| = 1$, or equivalently, there exists a unique j such that

$$d_i = \begin{cases} c_i \pm 1 & i = j; \\ c_i & i \neq j. \end{cases}$$

By Theorem 5.1, the vertices of $\mathcal{G}(\mathcal{T}^{\nu}_{\mathbb{X}})$ and $\Lambda^{0}_{(p,q)}$ coincide. Let T be a tilting bundle, recall from (5.2) that we can assume $\varphi(T) = (c_1, \cdots, c_{i-1}, c_i, c_{i+1}, \cdots, c_p)$, and we define

$$\mu_i^+(T) := (T \setminus [D_{\frac{c_i}{q}}^{\frac{i}{p}}]) \oplus [D_{\frac{c_i+1}{q}}^{\frac{i-1}{p}}]$$

and

$$\mu_i^-(T) := (T \setminus [D_{\frac{c_i}{q}}^{\frac{i-1}{p}}]) \oplus [D_{\frac{c_i-1}{q}}^{\frac{i}{p}}].$$

Then $\mu_i^+(T)$ and $\mu_i^-(T)$ are both tilting bundles with

$$\varphi(\mu_i^{\pm}(T)) = (c_1, \cdots, c_{i-1}, c_i \pm 1, c_{i+1}, \cdots, c_p) = \varphi(T) \pm \epsilon_i,$$

here, $\epsilon_i = (0, \dots, 0, 1, 0, \dots, 0)$ is *i*-th canonical row vector in \mathbb{Z}^p . Therefore, if there exists an edge in the tilting graph $\mathcal{G}(\mathcal{T}^{\nu}_{\mathbb{X}})$, then there exists a corresponding edge in $\Lambda_{(p,q)}$. Moreover, according to Proposition 5.4, there are n(T) = 2(p-r(T))-many edges attached to the tilting bundle T in the graph $\mathcal{G}(\mathcal{T}^{\nu}_{\mathbb{X}})$.

On the other hand, there is an edge between $\alpha = (c_1, c_2, \dots, c_p)$ and $\beta = (d_1, d_2, \dots, d_p)$ if and only if there exists a unique j such that

$$d_i = \begin{cases} c_i \pm 1 & i = j \\ c_i & i \neq j; \end{cases}$$

if and only if $\beta = \alpha \pm \epsilon_i$. Observe that $\beta = \alpha + \epsilon_i \in \Lambda^0_{(p,q)}$ if and only if

$$c_1 \leq c_2 \leq \cdots \leq c_i < c_{i+1} \leq c_{i+2} \leq \cdots \leq c_p,$$

and $\beta = \alpha - \epsilon_i \in \Lambda^0_{(p,q)}$ if and only if

$$c_1 \le c_2 \le \cdots \le c_{i-1} < c_i \le c_{i+1} \cdots \le c_p$$

Therefore, $c_i = c_{i+1}$ if and only if $\alpha + \epsilon_i \notin \Lambda^0_{(p,q)}$, $\alpha - \epsilon_{i+1} \notin \Lambda^0_{(p,q)}$. Assume $T = \varphi^{-1}(\alpha)$, then there are 2(p - r(T)) = n(T)-many edges attached to α in $\Lambda_{(p,q)}$. This finishes the proof.

6.3. Typical examples of tilting graphs.

Example 6.4. For the case of p = 1 and q = 2, recall that

$$T^0_{\frac{a}{2}} := \mathcal{O}(-a\vec{x}_2) \oplus \mathcal{O}(-a\vec{x}_2 + \vec{x}_2) \oplus \mathcal{O}(-a\vec{x}_2 + \vec{c}), \ a \in \mathbb{Z}$$

is a tilting bundle. We denote by

$$T_{\frac{a}{2}} = \mathcal{O}(-a\vec{x}_2) \oplus \mathcal{O}(-a\vec{x}_2 + \vec{c}) \oplus S_{0,1},$$

hence $T_{\frac{a}{2}}$ is a tilting sheaf but not a tilting bundle.

By using Proposition 6.1, the tilting graph $\mathcal{G}(\mathcal{T}_{\mathbb{X}})$ of coh- $\mathbb{X}(1,2)$ can be drawn as below (see also [48] Figure 8.1), where the second line is the tilting graph $\mathcal{G}(\mathcal{T}_{\mathbb{X}}^{\nu})$ of vec- $\mathbb{X}(1,2)$.



FIGURE 11. The tilting graph $\mathcal{G}(\mathcal{T}_{\mathbb{X}})$ of coh- $\mathbb{X}(1,2)$

Example 6.5. For the case of p = 2 and q = 2, by Theorem 1.2, we obtain that the tilting graph $\mathcal{G}(\mathcal{T}^{\nu}_{\mathbb{X}})$ of vec- $\mathbb{X}(2,2)$ can be described by the graph $\Lambda_{(2,2)}$, it follows that the shape of $\mathcal{G}(\mathcal{T}^{\nu}_{\mathbb{X}})$ can be drawn as below:



FIGURE 12. The tilting graph $\mathcal{G}(\mathcal{T}^{\nu}_{\mathbb{X}})$ of vec- $\mathbb{X}(2,2)$

Combining with Proposition 6.1, the tilting graph $\mathcal{G}(\mathcal{T}_{\mathbb{X}})$ of coh- $\mathbb{X}(2,2)$ can be drawn as below (see also [48] Figure 7.2):



FIGURE 13. The tilting graph $\mathcal{G}(\mathcal{T}_{\mathbb{X}})$ of coh- $\mathbb{X}(2,2)$

Example 6.6. Consider the case of p = 2 and q = 3, by using Theorem 1.2, the shape of the tilting graph $\mathcal{G}(\mathcal{T}_{\mathbb{X}}^{\nu})$ of vec- $\mathbb{X}(2,3)$ is drawn as below:



FIGURE 14. The tilting graph $\mathcal{G}(\mathcal{T}^{\nu}_{\mathbb{X}})$ of vec- $\mathbb{X}(2,3)$

It can be imaginable that the shape of the tilting graph $\mathcal{G}(\mathcal{T}_{\mathbb{X}})$ of coh- $\mathbb{X}(2,3)$ is very difficult, here we omit the description of the tilting graph of coh- $\mathbb{X}(2,3)$.

7. Geometric interpretation of automorphism group of $coh-\mathbb{X}(p,q)$

This section is devoted to giving a geometric interpretation for the automorphism group $\operatorname{Aut}(\operatorname{coh}-\mathbb{X}(p,q))$ of the category $\operatorname{coh}-\mathbb{X}(p,q)$. We show that $\operatorname{Aut}(\operatorname{coh}-\mathbb{X}(p,q))$ is isomorphic to the mapping class group of $A_{p,q}$.

7.1. Mapping class group. Let's recall from [2] for the mapping class group of the marked annulus $A_{p,q}$ first. Write $A_{p,q} = (S, M)$, where S is an annulus and M is the set of all the marked points in $A_{p,q}$.

Recall that two homeomorphisms f, g of S are *isotopic* if there is a continuous function $H : S \times [0,1] \longrightarrow S$ such that H(x,0) = f, H(x,1) = g and $H(x,t) : S \longrightarrow S$ is a homeomorphism for each $t \in [0,1]$.

Denote by Homeo⁺(S, M) the group of orientation preserving homeomorphisms from S to S which map M to M. Note that there do not require that the points in M are fixed, neither that the points on the boundary of S are fixed nor that each boundary component is mapped to itself. However, if a boundary component is mapped to another component, then p = q.

Definition 7.1. A homeomorphism $f: S \longrightarrow S$ is *isotopic to the identity relative to* M, if f is isotopic to the identity via an isotopy H that fixes M pointwise, i.e., H(x,t) = x for all $x \in M$ and $t \in [0, 1]$.

Let $Homeo_0(S, M)$ be the set

 $\{f \in \text{Homeo}^+(S, M) | f \text{ is isotopic to the identity relative to } M\}.$

The mapping class group $\mathcal{MG}(S, M)$ of (S, M) is defined to be the quotient

 $\mathcal{MG}(S, M) = \operatorname{Homeo}^+(S, M) / \operatorname{Homeo}_0(S, M).$

Denote by $H_{p,q} = \langle r_1, r_2 | r_1 r_2 = r_2 r_1, r_1^p = r_2^q \rangle$ and

$$\widetilde{H}_{p,q} = \begin{cases} H_{p,q}, & p \neq q; \\ H_{p,q} \times \mathbb{Z}_2, & p = q. \end{cases}$$
(7.1)

It is well known that the mapping class group $\mathcal{MG}(A_{p,q})$ of the marked annulus $A_{p,q}$ is isomorphism to $\widetilde{H}_{p,q}$, c.f. [2]. We identify them in the following.

Remark 7.2. In our paper, let r_1 (resp. r_2) be the rotation in anti-clockwise direction (resp. in clockwise direction) sending each marked point in ∂ (resp. ∂') to the next marked point in ∂ (resp. ∂'), i.e., for any $a, b \in \mathbb{Z}$,

$$r_1((\frac{a}{p})_{\partial}) = (\frac{a+1}{p})_{\partial}, \qquad r_1((\frac{b}{q})_{\partial'}) = (\frac{b}{q})_{\partial'};$$

$$r_2((\frac{a}{p})_{\partial}) = (\frac{a}{p})_{\partial}, \qquad r_2((\frac{b}{q})_{\partial'}) = (\frac{b-1}{q})_{\partial'}.$$

For the case of p = q, there exists an involution σ of $\mathcal{MG}(A_{p,q})$ maps ∂ to ∂' , satisfying that $\sigma((\frac{a}{p})_{\partial}) = (-\frac{a}{p})_{\partial'}$ for any $a \in \mathbb{Z}$, in particular, $\sigma(0_{\partial}) = 0_{\partial'}$.

Let $\operatorname{Aut}(\operatorname{coh}-\mathbb{X}(p,q))$ be the automorphism group of the category $\operatorname{coh}-\mathbb{X}(p,q)$ consists of isomorphism classes of **k**-linear self-equivalences on $\operatorname{coh}-\mathbb{X}(p,q)$. Then we have

Proposition 7.3. Aut(coh- $\mathbb{X}(p,q)$) $\cong \mathcal{MG}(A_{p,q})$.

Proof. By [37, Theorem 3.4], we have

$$\operatorname{Aut}(\operatorname{coh-}\mathbb{X}(p,q)) \cong \begin{cases} \mathbb{L}(p,q), & p \neq q; \\ \mathbb{L}(p,q) \times \mathbb{Z}_2, & p = q. \end{cases}$$
(7.2)

Observe that $\mathbb{L}(p,q)$ is generated by \vec{x}_1, \vec{x}_2 subject to the relation $p\vec{x}_1 = q\vec{x}_2$. By comparing (7.1) and (7.2), we see that the map $\vec{x}_1 \mapsto r_1$, $\vec{x}_2 \mapsto r_2$ gives a group isomorphism from $\mathbb{L}(p,q)$ to $H_{p,q}$, which induces a group isomorphism from $\operatorname{Aut}(\operatorname{coh-X}(p,q))$ to $\mathcal{MG}(A_{p,q})$. \Box

Remark 7.4. In the case p = q, recall from [37] that there exists a geometric automorphism of $\mathbb{X}(p,q)$ by exchanging the exceptional points ∞ and 0, which induces isomorphism

$$\sigma_{1,2}: \operatorname{coh-X}(p,q) \longrightarrow \operatorname{coh-X}(p,q), \tag{7.3}$$

by exchanging $S_{\infty,i}^{(j)}$ with $S_{0,i}^{(j)}$ for any $0 \le i \le p-1$ and $j \in \mathbb{Z}_{\ge 1}$, and sending line bundles $\mathcal{O}(l_1\vec{x}_1 + l_2\vec{x}_2 + l\vec{c})$ to $\mathcal{O}(l_1\vec{x}_2 + l_2\vec{x}_1 + l\vec{c})$ for any $0 \le l_i \le p-1$ with i = 1, 2 and $l \in \mathbb{Z}$.

For any $\vec{x} \in \mathbb{L}(p,q)$, denote by $(\vec{x}) := - \otimes \mathcal{O}(\vec{x})$, which is the auto-equivalence functor in Aut(coh- $\mathbb{X}(p,q)$) given by the grading twist with \vec{x} . Then we have an isomorphism of groups

$$\psi : \mathcal{MG}(A_{p,q}) \longrightarrow \operatorname{Aut}(\operatorname{coh-X}(p,q)),$$
(7.4)

which sends

$$r_1 \mapsto (\vec{x}_1); \quad r_2 \mapsto (\vec{x}_2); \text{ and } \sigma \mapsto \sigma_{1,2} \text{ (when } p = q).$$

7.2. **Proof of Theorem 1.1.** For any triangulation Γ of the marked annulus $A_{p,q}$, we assume Γ consists of the following positive bridging arcs and peripheral arcs:

$$\left[D_{\frac{b_i}{q}}^{\frac{a_i}{p}}\right] (1 \le i \le n_1), \ \left[D^{\frac{i_k - j_k - 1}{p}, \frac{i_k}{p}}\right] (1 \le k \le n_2), \ \left[D_{\frac{-i_k}{q}, \frac{j_k - i_k + 1}{q}}\right] (1 \le k \le n_3)$$

According to the map (3.1), we obtain a tilting sheaf

$$\phi(\Gamma) = \bigoplus_{i=1}^{n_1} \mathcal{O}(a_i \vec{x}_1 - b_i \vec{x}_2) \oplus (\bigoplus_{k=1}^{n_2} S_{\infty, i_k}^{(j_k)}) \oplus (\bigoplus_{k=1}^{n_3} S_{0, i_k}^{(j_k)}).$$

Recall from (7.4) that ψ is a group isomorphism, we only need to prove that (1.1) holds for any generator $f \in \mathcal{MG}(A_{p,q})$. First assume $f = r_1$, then we have $\psi(f) = (\vec{x}_1)$. Recall that

$$r_1((\frac{a}{p})_{\partial}) = (\frac{a+1}{p})_{\partial}, \ r_1((\frac{b}{q})_{\partial'}) = (\frac{b}{q})_{\partial'}, \ a, b \in \mathbb{Z}.$$

It follows that the triangulation $f(\Gamma)$ consists of the following arcs:

$$\left[D_{\frac{b_i}{q}}^{\frac{a_i+1}{p}}\right] (1 \le i \le n_1), \ \left[D^{\frac{i_k-j_k}{p},\frac{i_k+1}{p}}\right] (1 \le k \le n_2), \ \left[D_{\frac{-i_k}{q},\frac{j_k-i_k+1}{q}}\right] (1 \le k \le n_3).$$

Then

$$\phi(f(\Gamma)) = \bigoplus_{i=1}^{n_1} \mathcal{O}((a_i+1)\vec{x}_1 - b_i\vec{x}_2) \oplus (\bigoplus_{k=1}^{n_2} S_{\infty,i_k+1}^{(j_k)}) \oplus (\bigoplus_{k=1}^{n_3} S_{0,i_k}^{(j_k)})$$

= $\phi(\Gamma)(\vec{x}_1)$
= $\psi(f)(\phi(\Gamma)).$

Hence (1.1) holds in this case. For $f = (r_2)$, the proof is similar.

If p = q, then $\mathcal{MG}(A_{p,q})$ has another generator (involution) $f = \sigma$ such that $\sigma((\frac{a}{p})_{\partial}) = (-\frac{a}{p})_{\partial'}$. It follows that the triangulation $f(\Gamma)$ consists of the following arcs:

$$\left[D_{-\frac{a_i}{p}}^{-\frac{b_i}{p}}\right] (1 \le i \le n_1), \ \left[D_{-\frac{i_k}{p}, \frac{j_k - i_k + 1}{p}}\right] (1 \le k \le n_2), \ \left[D^{\frac{i_k - j_k - 1}{p}, \frac{i_k}{p}}\right] (1 \le k \le n_3).$$

Then

$$\phi(f(\Gamma)) = \bigoplus_{i=1}^{n_1} \mathcal{O}(a_i \vec{x}_2 - b_i \vec{x}_1) \oplus (\bigoplus_{k=1}^{n_2} S_{0,i_k}^{(j_k)}) \oplus (\bigoplus_{k=1}^{n_3} S_{\infty,i_k}^{(j_k)}) = \psi(f)(\phi(\Gamma)).$$

Hence (1.1) holds. This finishes the proof.

Recall from [10] that the exchange graph $EG(A_{p,q})$ of $A_{p,q}$ has as vertices the triangulations of $A_{p,q}$, with an edge between two vertices Γ and Γ' whenever Γ' is obtained from Γ by the flip of an arc. By Theorem 1.1, we have the following corollary.

Corollary 7.5. For any $f \in \mathcal{MG}(A_{p,q})$, we have $\phi(f(EG(A_{p,q})) = \psi(f)(\phi(EG(A_{p,q})))$.

7.3. An involution on $\Lambda_{(p,p)}$. Recall from (5.2) that there exists a bijective map

$$\varphi: \mathcal{T}^{\nu}_{\mathbb{X}} \longrightarrow \Lambda^{0}_{(p,p)}; \quad T \mapsto (c_1, c_2, \cdots, c_p)$$

where $\Lambda^0_{(p,p)} = \{(c_1, \dots, c_p) \in \mathbb{Z}^p | c_1 \leq \dots \leq c_p \leq c_1 + p\}.$ In case p = q, by (7.3) there exists an involution $\sigma_{1,2}$ on coh- $\mathbb{X}(p,p)$, which induces an automorphism on $\mathcal{G}(\mathcal{T}^{\nu}_{\mathbb{X}})$, still denoted by $\sigma_{1,2}$.

Proposition 7.6. There exists an involution $\rho : \Lambda_{(p,p)} \longrightarrow \Lambda_{(p,p)}$ such that the following diagram commutes



Proof. Let T be a tilting bundle in coh- $\mathbb{X}(p, p)$, recall from (5.2) that we can assume

$$\varphi(T) = (c_1, \cdots, c_{i-1}, c_i, c_{i+1}, \cdots, c_p),$$

that is, T corresponds to a triangulation Γ of (a parallelogram in) U as follows:



According to Remark 7.2, we obtain that $\sigma(\Gamma)$ is a triangulation of the following form:



which corresponds to the tilting bundle $\sigma_{1,2}(T)$; see (7.3). By using (3.2), the above triangulation can be converted to its normal form (5.1) as follows:



This induces a well-defined map:

$$\rho: \Lambda^0_{(p,p)} \longrightarrow \Lambda^0_{(p,p)}, \quad (c_1, c_2, \cdots, c_p) \mapsto (d_1, d_2, \cdots, d_p).$$

By construction we have $\varphi(\sigma_{1,2}(T)) = \rho(\varphi(T))$. According to the proof of Theorem 1.2, we see that φ is an isomorphism between the graphs $\mathcal{G}(\mathcal{T}_{\mathbb{X}}^{\nu})$ and $\Lambda_{(p,p)}$. This proves the commutativity of the diagram. Moreover, since $\sigma_{1,2}$ is an involution, it follows that ρ is also an involution.

Example 7.7. For the case of p = q = 4, let T be a tilting bundle and $\varphi(T) = (0, 0, 1, 4)$, that is, T corresponds to a triangulation Γ of (a parallelogram in) U as follows:



Then $\sigma_{1,2}(T)$ corresponds to a triangulation $\sigma(\Gamma)$ of (a parallelogram in) U as follows:



By using (3.2), we can change the above triangulation to its normal form (red arcs):



Then $\rho(0, 0, 1, 4) = (1, 1, 1, 2)$, and $\varphi(\sigma_{1,2}(T)) = \rho(\varphi(T))$.

7.4. **Proof of Theorem 1.3.** Assume $\nu = (c_1, c_2, \dots, c_p)$. Let *T* be a tilting bundle in coh- $\mathbb{X}(p,q)$ corresponding to $\nu = (c_1, c_2, \dots, c_p)$, that is, $\varphi(T) = (c_1, c_2, \dots, c_p)$; see (5.2). According to the proof of Theorem 5.1, we obtain

$$\varphi(T(\vec{x}_1)) = (c_p - q, c_1, c_2, \cdots, c_{p-1}), \quad \varphi(T(\vec{x}_2)) = (c_1 - 1, c_2 - 1, \cdots, c_p - 1).$$

The following assignments define bijective maps

$$\rho_1: \Lambda^0_{(p,q)} \longrightarrow \Lambda^0_{(p,q)}; \quad (c_1, c_2, \cdots, c_{p-1}, c_p) \mapsto (c_p - q, c_1, c_2, \cdots, c_{p-1});$$

and

$$\rho_2: \Lambda^0_{(p,q)} \longrightarrow \Lambda^0_{(p,q)}; \quad (c_1, c_2, \cdots, c_{p-1}, c_p) \mapsto (c_1 - 1, c_2 - 1, \cdots, c_p - 1).$$

Observe that the grading shift by (\vec{x}_i) on coh- $\mathbb{X}(p,q)$ induces an automorphism on $\mathcal{G}(\mathcal{T}^{\nu}_{\mathbb{X}})$, still denoted by (\vec{x}_i) . Similar as the proof of Proposition 7.6, we have the following commutative diagrams for i = 1, 2



It is easy to check that $\rho_1\rho_2 = \rho_2\rho_1$ and $\rho_1^p = \rho_2^q$. Hence, the assignments $r_1 \mapsto \rho_1, r_2 \mapsto \rho_2$ defines a group homomorphism from $\widetilde{H}_{p,q}$ to Aut $(\Lambda_{(p,q)})$. Combining with Proposition 7.6, we obtain a group action of $\widetilde{H}_{p,q}$ on $\Lambda_{(p,q)}$. Then (1.2) follows from Theorem 1.2, and hence we finish the proof of Theorem 1.3.

8. Geometric interpretation of perpendicular category

In this section, we focus on the geometric interpretation of the perpendicular category of an exceptional object in $\operatorname{coh-X}(p,q)$.

Recall that a coherent sheaf E is called *exceptional* if $\text{Hom}(E, E) = \mathbf{k}$ and $\text{Ext}^1(E, E) = 0$. For an exceptional sheaf $E \in \text{coh-}\mathbb{X}(p, q)$, define the *right perpendicular category*

$$E^{\perp} := \{ X \in \operatorname{coh-} \mathbb{X}(p,q) | \operatorname{Hom}(E,X) = 0 = \operatorname{Ext}^{1}(E,X) \};$$

and the *left perpendicular category*

$${}^{\perp}E := \{ X \in \operatorname{coh-} \mathbb{X}(p,q) | \operatorname{Hom}(X,E) = 0 = \operatorname{Ext}^{1}(X,E) \}.$$

By Proposition 2.1 (1), we obtain that ${}^{\perp}E = (E(-\vec{\omega}))^{\perp}$, where $\vec{\omega} = -(\vec{x}_1 + \vec{x}_2)$ is the dualizing element in $\mathbb{L}(p,q)$.

Lemma 8.1. Let E be an indecomposable sheaf in coh-X(p,q), then E is an exceptional object if and only if $\phi^{-1}(E)$ is an arc in $A_{p,q}$.

Proof. By [40, Lemma 3.2.3], E is exceptional in coh- $\mathbb{X}(p,q)$ if and only if $\operatorname{Ext}^1(E, E) = 0$. Combining with Theorem 3.10, we get the result.

J. CHEN, S. RUAN AND H. ZHANG

We recall some basic facts on cutting marked annulus $A_{p,q}$ along an arc, cf. [39].

Let α be an arc in $A_{p,q}$. Denote by $A_{p,q}/\alpha$ the new marked surface obtained from $A_{p,q}$ by cutting along the arc α and then removing components which are homeomorphic to a triangle. Up to homeomorphism, it does not depend on the choice of representative of α .

Theorem 8.2. Let E be an exceptional object in coh- $\mathbb{X}(p,q)$, and assume the oriented arc corresponding to E is α . Then the marked surface $A_{p,q}/\alpha$ gives a geometric model for the category E^{\perp} .

Proof. To prove the theorem, we consider the following two cases:

(1) $E \in \text{vec-}\mathbb{X}(p,q).$

In this case, E is a line bundle, hence $E^{\perp} \cong \text{mod}(A_{p+q-1})$ by [36, Proposition 2.14]. On the other hand, the marked surface $A_{p,q}/\alpha$ is a disk with p+q+2 marked points on its boundary. According to [8], this gives a geometric model for the category $\text{mod}(A_{p+q-1}) \cong E^{\perp}$.

(2) $E \in \operatorname{coh}_0 - \mathbb{X}(p, q)$.

In this case, $E = S_{\infty,i}^{(j)}$, where $i \in \mathbb{Z}/p\mathbb{Z}$ and $1 \leq j \leq p-1$; or $E = S_{0,i}^{(j)}$, where $i \in \mathbb{Z}/q\mathbb{Z}$ and $1 \leq j \leq q-1$. We only consider the second case, the other one being similar.

Note that the composition factors of $S_{0,i}^{(j)}$ are given by $S_{0,i}$, $S_{0,i-1}$, \cdots , $S_{0,i-j+1}$. According to [24], $(S_{0,i}^{(j)})^{\perp}$ contains two disjoint components, one is $\{S_{0,i}, S_{0,i-1}, \cdots, S_{0,i-j+1}\}^{\perp}$, which equivalent to the weighted projective line of weight type (p, q - j); the other one is $\langle S_{0,i-1}, S_{0,i-2}, \cdots, S_{0,i-j+1} \rangle$, which is equivalent to the module category of type A_{j-1} . Therefore,

$$(S_{0,i}^{(j)})^{\perp} \cong \operatorname{coh-X}(p,q-j) \coprod \operatorname{mod}(A_{j-1}).$$

On the other hand, the marked surface $A_{p,q}/\alpha$ has two connected components: an annulus with p marked points on the inner boundary and q-j marked points on the outer boundary, and a disk with j + 2 marked points on its boundary. Combining with [8], this gives a geometric model for the category $(S_{0,i}^{(j)})^{\perp}$. We are done.

Acknowledgements. Jianmin Chen and Hongxia Zhang were partially supported by the National Natural Science Foundation of China (Grant Nos. 12371040, 11971398, 12131018 and 12161141001). Shiquan Ruan was partially supported by the Natural Science Foundation of Xiamen (No. 3502Z20227184), the Natural Science Foundation of Fujian Province (No. 2022J01034), the National Natural Science Foundation of China (Nos. 12271448), and the Fundamental Research Funds for Central Universities of China (No. 20720220043).

References

- I. Assem, T. Brüstle, G. Charbonneau-Jodoin, and P. G. Plamondon. Gentle algebras arising from surface triangulations. *Algebra Number Theory* 4 (2010), no. 2, 201–229.
- [2] I. Assem, R. Schiffler, and V. Shramchenko. Cluster automorphisms. Proc. Lond. Math. Soc. (3) 104 (2012), no. 6, 1271–1302.
- [3] M. Barot, D. Kussin, and H. Lenzing. The cluster category of a canonical algebra. Trans. Amer. Math. Soc. 362 (2010), no. 8, 4313–4330.
- [4] K. Baur, A. B. Buan, and R. J. Marsh. Torsion pairs and rigid objects in tubes. Algebr. Represent. Theory 17 (2014), no. 2, 565–591.
- [5] K. Baur, and R. J. Marsh. A geometric description of the *m*-cluster categories of type D_n . Int. Math. Res. Not. IMRN (2007), no.4.

- [6] K. Baur, and R. J. Marsh. A geometric description of m-cluster categories. Trans. Amer. Math. Soc. 360 (2008), no.11, 5789–5803.
- [7] K. Baur, and R. J. Marsh. A geometric model of tube categories. J. Algebra 362 (2012), 178–191.
- [8] K. Baur, and R. C. Simões. A geometric model for the module category of a gentle algebra. Int. Math. Res. Not. IMRN 2021, no. 15, 11357–11392.
- [9] K. Baur, and H. A. Torkildsen. A geometric interpretation of categories of type A and of morphisms in the infinite radical. *Algebr. Represent. Theory* 23 (2020), no. 3, 657–692.
- [10] T. Brüstle, and Y. Qiu. Tagged mapping class groups: Auslander-Reiten translation. Math. Z. 279 (2015), no. 3-4, 1103–1120.
- [11] T. Brüstle, and J. Zhang. On the cluster category of a marked surface without punctures. Algebra Number Theory 5 (2011), no. 4, 529–566.
- [12] P. Caldero, F. Chapoton, and R. Schiffler. Quivers with relations arising from clusters (A_n case). *Trans. Amer. Math. Soc.* 358 (2006), no. 3, 1347–1364.
- [13] J. Chen, Y. Lin, P. Liu, and S. Ruan. Tilting objects on tubular weighted projective lines: a cluster tilting approach. *Sci. China Math.* 64 (2021), no. 4, 691–710.
- [14] W. Crawley-Boevey. Kac's theorem for weighted projective lines. J. Eur. Math. Soc. (JEMS) 12 (2010), no. 6, 1331–1345.
- [15] B. Deng, S. Ruan, and J. Xiao. Applications of mutations in the derived categories of weighted projective lines to Lie and quantum algebras. Int. Math. Res. Not. IMRN 2020, no. 19, 5814–5871.
- [16] R. Dou, Y. Jiang, and J. Xiao. The Hall algebra approach to Drinfeld's presentation of quantum loop algebras. Adv. Math. 231 (2012), no. 5, 2593–2625.
- [17] B. Duan, L. Lamberti, and J. R. Li. Combinatorial model for *m*-cluster categories in type *E*, arXiv:1911.12042, 2019.
- [18] W. Ebeling. The Poincaré series of some special quasihomogeneous surface singularities. Publ. Res. Inst. Math. Sci. 39 (2003), no. 2, 393–413.
- [19] W. Ebeling, and D. Ploog. McKay correspondence for the Poincaré series of Kleinian and Fuchsian singularities. Math. Ann. 347 (2010), no. 3, 689–702.
- [20] W. Ebeling, and A. Takahashi. Mirror symmetry between orbifold curves and cusp singularities with group action. Int. Math. Res. Not. IMRN 2013, no. 10, 2240–2270.
- [21] S. Fomin, M. Shapiro, and D. Thurston. Cluster algebras and triangulated surfaces. Part I: Cluster complexes. Acta Math. 201 (2008), no. 1, 83–146.
- [22] C. Fu, and S. Geng. On cluster-tilting graphs for hereditary categories. Adv. Math. 383 (2021).
- [23] W. Geigle, and H. Lenzing. A class of weighted projective curves arising in representation theory of finite-dimensional algebras. Singularities, representation of algebras, and vector bundles (Lambrecht, 1985), 265–297, Lecture Notes in Math., 1273, Springer, Berlin, 1987.
- [24] W. Geigle, and H. Lenzing. Perpendicular categories with applications to representations and sheaves. J. Algebra 144 (1991), no. 2, 273–343.
- [25] S. Geng. Mutation of tilting bundles of tubular type. J. Algebra 550 (2020), 186–209.
- [26] D. Happel, and I. Reiten. Hereditary categories with tilting object. Math. Z. 232 (1999), no. 3, 559– 588.
- [27] D. Happel, and L. Unger. On the set of tilting objects in hereditary categories. Fields Inst. Commun., 45, Amer. Math. Soc., Providence, RI, 2005.
- [28] P. He, Y. Zhou, and B. Zhu. A geometric model for the module category of a skew-gentle algebra. Math. Z. 304 (2023), no.1, Paper No. 18, 41.
- [29] T. Holm, and P. Jørgensen. On a cluster category of infinite Dynkin type, and the relation to triangulations of the infinity-gon. Math. Z. 270 (2012), no. 1-2, 277–295.
- [30] T. Hübner. Classification of Indecomposable Vector Bundles on Weighted Curves. *Diplomarbeit, Universität Paderborn*, 1989.
- [31] T. Hübner. Exzeptionelle Vektorbündel und Reflektionen an Kippgarben über projektiven gewichteten Kurven. Dissertation, Universität Paderborn, 1996.
- [32] L. Lamberti. Combinatorial model for the cluster categories of type E. J. Algebraic Combin. 41 (2015), no. 4, 1023–1054.

- [33] H. Lenzing. Wild canonical algebras and rings of automorphic forms. Finite-dimensional algebras and related topics (Ottawa, ON, 1992), 191–212, NATO Adv. Sci. Inst. Ser. C: Math. Phys. Sci., 424, Kluwer Acad. Publ., Dordrecht, 1994.
- [34] H. Lenzing. Representations of finite-dimensional algebras and singularity theory. Trends in ring theory (Miskolc, 1996), 71–97, CMS Conf. Proc., 22, Amer. Math. Soc., Providence, RI, 1998.
- [35] H. Lenzing. Rings of singularities. Bull. Iranian Math. Soc. 37 (2011), no. 2, 235–271.
- [36] H. Lenzing. Weighted projective lines and applications. Representations of algebras and related topics, 153–187, EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2011.
- [37] H. Lenzing, and H. Meltzer. The automorphism group of the derived category for a weighted projective line. Comm. Algebra 28 (2000), no. 4, 1685–1700.
- [38] H. Lenzing, and I. Reiten. Hereditary Noetherian categories of positive Euler characteristic. Math. Z. 254 (2006), no. 1, 133–171
- [39] R. B. Marsh, and Y. Palu. Coloured quivers for rigid objects and partial triangulations: the unpunctured case. Proc. Lond. Math. Soc. (3) 108 (2014), no. 2, 411–440.
- [40] H. Meltzer. Exceptional vector bundles, tilting sheaves and tilting complexes for weighted projective lines. Mem. Amer. Math. Soc. 171 (2004), no. 808.
- [41] C. M. Ringel. Tame algebras and integral quadratic forms. Lecture Notes in Mathematics, 1099. Springer-Verlag, Berlin, 1984.
- [42] I. G. Sčerbak. Algebras of automorphic forms with three generators. Funkcional. Anal. i Priložen. 12 (1978), no. 2, 93–94.
- [43] R. Schiffler. A geometric model for cluster categories of type D_n . J. Algebraic Combin. 27 (2008), no. 1, 1–21.
- [44] O. Schiffmann. Noncommutative projective curves and quantum loop algebras. Duke Math. J. 121 (2004), no. 1, 113–168.
- [45] D. Simson, and A. Skowroński. Elements of the representation theory of associative algebras. Vol.
 2. Tubes and concealed algebras of Euclidean type. London Mathematical Society Student Texts, 71. Cambridge University Press, Cambridge, 2007.
- [46] H. A. Torkildsen. A geometric realization of the m-cluster category of affine type A. Comm. Algebra 43 (2015), no.6, 2541–2567.
- [47] H. Vogel. Asymptotic triangulations and Coxeter transformations of the annulus. Glasg. Math. J. 60 (2018), no. 1, 63–96.
- [48] M. Warkentin. Fadenmoduln über A_n und Cluster-Kombinatorik (string modules over A_n and cluster combinatorics). Diploma Thesis, University of Bonn, December 2008.

Jianmin Chen, Shiquan Ruan and Hongxia Zhang

School of Mathematical Sciences,

Xiamen University, Xiamen, 361005, Fujian, PR China.

E-mails: chenjianmin@xmu.edu.cn, sqruan@xmu.edu.cn, hxzhangxmu@163.com