Gromov-Witten Generating Series of Elliptic Curves and Iterated Integrals of Eisenstein-Kronecker Forms

Jie Zhou

Abstract

We compute various types of iterated integrals of Eisenstein-Kronecker forms that are constructed from the Kronecker theta function. Furthermore, we relate the generating series of Gromov-Witten invariants of elliptic curves to these iterated integrals.

Contents

1 Introduction

The Gromov-Witten (GW) theory of elliptic curves has remained an interesting and active research topic in modern enumerative geometry. Explicit formulas for the generating series of the corresponding GW invariants in terms of Jacobi theta functions have been derived by Okounkov-Pandharipande [\[OP06\]](#page-29-0) and Bloch-Okounkov [\[BO00\]](#page-28-0), and turn out to be quasi-elliptic functions [\[EZ85\]](#page-29-1) that are interesting in the studies of modular forms. These formulas have even since received a lot of attention due to their concreteness and effectiveness in studying the interaction between enumerative geometry, modular forms, and mathematical physics, see e.g., [\[Dij95,](#page-28-1) [KZ95,](#page-29-2) [Zag16\]](#page-30-0).

Recently, it is shown in [\[Zho23a\]](#page-30-1) that these purely enumerative generating series are given by configuration space integrals of cohomological classes constructed from sections of Poincaré bundles. Furthermore, they admit new sum-over-partitions formulas that are much simpler than the original ones given in [\[BO00,](#page-28-0) [OP06\]](#page-29-0).

We now review and rephrase these results briefly. For any point τ on the upper half-plane **H**, let

$$
E = \mathbb{C}/\Lambda_{\tau}, \quad \Lambda_{\tau} := \mathbb{Z} \oplus \mathbb{Z}\tau \tag{1.1}
$$

be the corresponding elliptic curve. We fix once and for all a linear holomorphic coordinate *z* on the universal cover $\widetilde{E} = \mathbb{C} \to E$ and the coordinate $u = e^{2\pi i z}$ on the cover $\mathbb{C}^* = \mathbb{C}/2\pi i \mathbb{Z} \to E$. The origin on *E* is taken to be the image of $0 \in \mathbb{C}$ under the universal covering map $\mathbb{C} \to E$. We also fix a weak Torelli marking $\{A, B\}$, namely a symplectic frame for $H_1(E, \mathbb{Z})$. The class A is taken to be the image of the segment connecting τ and $\tau + 1$ under the map $\mathbb{C} \to E$.

For any $n \geq 1$, denote $[n] := (1, 2, \dots, n)$ and let Conf_[*n*] $(E) \subseteq E^n$ be the configuration space of *n* ordered points on *E*. Let the linear coordinates on the *n* components of \tilde{E}^n , E^n in $\tilde{E}^n \times E^n$ be w_1, \cdots, w_n and z_1, \cdots, z_n respectively, and $q = e^{2\pi i \tau}$.

Theorem 1.1 (=Theorem [4.1\)](#page-20-2)**.** *Assume the following notation.*

• *Let*

$$
\omega_n = \frac{\theta(\sum_{i=1}^n w_i)}{\prod_{i=1}^n \theta(w_i)} \cdot \frac{\prod_{i < j} \theta(z_i + w_i - z_j - w_j) \theta(z_i - z_j)}{\prod_{i < j} \theta(z_i + w_i - z_j) \theta(z_i - w_j - z_j)} \cdot \bigwedge_{i=1}^n \frac{\mathbf{i} \, dz_i \wedge d\bar{z}_i}{2 \operatorname{im} \tau},\tag{1.2}
$$

where $\theta(w)$ *is the odd Jacobi theta function normalized such that* $(\partial_w \theta)(0) = 1$ *.*

• Let $\widehat{B}_n(w)$ be the complete Bell polynomial in the variables $\widehat{\mathcal{E}}_m^*(w)$, $m\geq 1$ which are the Eisenstein-*Kronecker series*[1](#page-1-0)

$$
\widehat{\mathcal{E}}_m^*(w) := (-1)^{m-1} (m-1)! \left(\frac{1}{w^m} + \sum_{\lambda \in \Lambda_\tau - \{(0,0)\}} \left(\frac{1}{(w+\lambda)^m} + (-1)^{m-1} \frac{1}{\lambda^m} \right) + \delta_{m,1} \frac{-\pi}{\mathrm{im } \,\tau} (\overline{w} - w) \right).
$$

• *Denote* $\Pi_{[n]\setminus\{j\}}$ to be the set of partitions of $[n]\setminus\{j\}$ consisting of elements of the form π = $\{\pi_1, \cdots, \pi_\ell\}.$

Then the regularized integral [\[LZ21\]](#page-29-3) $\int_{E^n} \omega_n$ *satisfies*

$$
\oint_{E^n} \omega_n = \sum_{j \in [n]} \sum_{\pi \in \Pi_{[n] \setminus \{j\}}} \frac{\widehat{B}_{\pi}}{|\pi|}, \quad \frac{\widehat{B}_{\pi}}{|\pi|} := \prod_{k=1}^{\ell} \frac{\widehat{B}_{|\pi_k|}(\sum_{i \in \pi_k} w_i)}{|\pi_k|}.
$$
\n(1.3)

The significance of this result lies in the following. Let $F_n(w_1, \dots, w_n; q)$ be the *n*-point function of disconnected stationary GW invariants of the elliptic curve \mathcal{E} [\[OP06\]](#page-29-0). Consider its scaled version

$$
T_n(w_1, \cdots, w_n; q) := \theta(\sum_{i=1}^n w_i) \cdot \prod_{k=1}^{+\infty} (1 - q^k)^{-1} \cdot F_n(w_1, \cdots, w_n; q).
$$
 (1.4)

¹A quick definition of the complete Bell polynomial in the variables x_1, \dots, x_m, \dots is given by $B_m(x_1, \dots, x_m)$ = $m! \cdot [t^m] \exp(\sum_{k=1}^{\infty} \frac{t^k}{k!} x_k)$. The quantity $\widehat{\mathcal{E}}_m^*$ can be alternatively written as $\partial_w^m(\ln \theta) + 2G_m + \delta_{m,1} \frac{-\pi}{\ln \tau} (\overline{w} - w)$, where $2G_m$ is the ordinary Eisenstein series defined as the summation of $(m-1)! \cdot \lambda^{-m}$ over $\lambda \in \Lambda_{\tau} - \{(0,0)\}.$

This quantity admits very explicit expressions in terms of the Jacobi theta function *θ* and turns out to be a quasi-elliptic function, due to Okounkov-Pandharipande [\[OP06\]](#page-29-0) and Bloch-Okounkov [\[BO00\]](#page-28-0). Let $\hat{T}_n(w_1, \dots, w_n; q)$ be the elliptic and modular completion [\[EZ85\]](#page-29-1) of $T_n(w_1, \dots, w_n; q)$. Then one has the following relation [\[Zho23a\]](#page-30-1) between this enumerative-geometric quantity and the regularized integral

$$
\widehat{T}_n(w_1,\cdots,w_n;q)=\int_{E^n}\omega_n\,.
$$
\n(1.5)

The results [\(1.3\)](#page-1-1), [\(1.5\)](#page-2-0) are proved [\[Zho23a,](#page-30-1) Theorem 1.1, Theorem 1.2] in their holomorphic limit versions, which correspond to an iterated A-cycle integral on the configuration space Conf_[*n*](*E*). The results are then promoted [\[Zho23a,](#page-30-1) Remark 4.13] to the present version using the machinery of quasi-elliptic functions, and the relation between iterated *A*-cycle integrals and regularized integrals established in [\[LZ21,](#page-29-3) [Zho23a\]](#page-30-1).

The main goal of this work is to offer a direct and much simpler proof of Theorem [1.1](#page-1-2) using the tool of regularized integrals [\[LZ21,](#page-29-3) [LZ23,](#page-29-4) [Zho23b\]](#page-30-2) and properties of the Eisenstein-Kronecker forms (reviewed in Section [2](#page-3-0) below). In particular, this work serves as a demonstration of the usefulness of the notion of regularized integrals which seems to have provided an ideal tool in regularizing divergent integrals arising from 2d chiral conformal field theories (see [\[LZ23\]](#page-29-4) for discussions on this). In the course, we compute various types of iterated integrals of Eisenstein-Kronecker forms which are of independent interest in understanding the geometry of configuration spaces of elliptic curves. We also reveal a relation between regularized integrals of Eisenstein-Kronecker forms and Chen's iterated path integrals [\[Che77\]](#page-28-2), and obtain some combinatorial results for the generating series of ${\{\hat{T}_n\}}_{n\geq 1}.$

Structure of the paper

In Section [2,](#page-3-0) we recall the basics of quasi-elliptic and almost-elliptic functions, including Jacobi theta functions, Kronecker theta function, and Eisenstein-Kronecker series.

In Section [3,](#page-9-0) after a brief review on the notion of regularized integrals on configuration spaces of elliptic curves, we compute regularized integrals of Eisenstein-Kronecker forms and discuss their connections to Chen's iterated path integrals.

In Section [4,](#page-20-0) we prove Theorem [1.1](#page-1-2) based on the results obtained in Section [3.](#page-9-0) We also discuss some combinatorial properties of the generating series of $\{\widehat{T}_n\}_{n\geq 1}$.

Acknowledgement

The author would like to thank Si Li for fruitful collaborations on related topics, and Jin Cao, Xinxing Tang and Zijun Zhou for helpful conversations.

This work was supported by the national key research and development program of China (No. 2022YFA1007100) and the Young overseas high-level talents introduction plan of China.

Notation and conventions

• Let Ω^p_X $\frac{p}{X}$ be the sheaf of holomorphic *p* forms on *X* and $H^0(X, \Omega_X^p)$ $\binom{p}{X}$ be the corresponding space of global sections over *X*.

- Let $A_X^{p,q}$ $\chi^{\mathit{p,q}}$ (resp. $A_X^{\mathit{p,q}}$ $\chi^{\mathcal{P},q}(\star D)$) be the space of (p,q) forms on *X* (resp. the space of (p,q) forms that are smooth on *X* except for holomorphic poles along the effective divisor *D*). Let $A^{n}(X) = A_{X}^{n} = \bigoplus_{p+q=n} A_{X}^{p,q}$ $P_{X}^{p,q}$, $A_{X}^{n}(\star D) = \bigoplus_{p+q=n} A_{X}^{p,q}$ $X^{p,q}(\star D).$
- The notation $[t^k]$ *f* represents the degree t^k coefficient of the formal Laurent series f in t .
- With the map exp : **C** → **C**[∗] understood, we use the additive coordinate *z* on **C** and the multiplicative coordinate $u = e^{2\pi i z}$ on \mathbb{C}^* interchangeably.
- Throughout this work, when we need to keep track of the variables in a construction such as a function f_n that are taken from some finite set *J* or sequence *K* of variables with cardinality *n*, we use the notation f_I or f_K . Correspondingly, we shall use the notation *S* abusively for both the set *S* and its cardinality |*S*|.
- We denote collectively the points P_i , $1 \le i \le n$ and Q_j , $1 \le j \le n$ by P , Q , respectively. Similarly we apply the same convention to denote a collection of coordinates, e.g., $z = (z_1, \dots, z_n)$.
- For notational simplicity, we shall often suppress the arguments of a function whenever they are clear from the surrounding texts.

2 Preliminaries

We now collect a few results in the literature about the space of almost-elliptic functions, which includes in particular the subspace $A_E^0({\star}D)$ that we are mostly interested in.

Throughout this work, constructions on the elliptic curve $E_\tau = C/\Lambda_\tau$ (and on the corresponding universal family $\mathcal{E}_H \to H$) will be identified with their lifts to the universal cover. For example, meromorphic functions on *E* are identified with those on **C** that are periodic/elliptic under the translation action by Λ*τ*.

2.1 Jacobi theta functions and Eisenstein-Kronecker series

We first recall some standard facts about Jacobi theta functions and Eisenstein series, mainly following [\[EZ85,](#page-29-1) [Sil09\]](#page-30-3). Define θ to be the normalized Jacobi theta function $\vartheta_{(\frac{1}{2},\frac{1}{2})}$:

$$
\theta(z) = \frac{\vartheta_{(\frac{1}{2},\frac{1}{2})}(z)}{\vartheta'_{(\frac{1}{2},\frac{1}{2})}(0)} = \frac{\vartheta_{(\frac{1}{2},\frac{1}{2})}(z)}{-2\pi i\eta^3},\tag{2.1}
$$

where *η* is the Dedekind eta function. For a positive integer *k*, define the Eisenstein series

$$
G_k = \frac{1}{2} \sum_{\lambda \in \Lambda_{\tau} - \{(0,0)\}}^e \frac{1}{\lambda^k},\tag{2.2}
$$

When *k* is odd, it is defined to be zero by convention. When $k = 2$, one uses the Eisenstein summation prescription \sum^e (summing over the 1 direction first then τ direction in Λ_τ) to deal with the non-absolute convergence issue.

Let ζ , \wp be the Weierstrass ζ -function and \wp -function, with $\wp = -\partial_z \zeta = -\zeta'$. Hereafter the superscript ' indicates derivative in z. The following relation between Eisenstein series and semiperiods is classical

$$
\int_{A} \wp \, dz = -\eta_1 \,, \quad \eta_1 := 2G_2 \,. \tag{2.3}
$$

One also has the following expansions (see e.g., [\[Sil09\]](#page-30-3))

$$
\theta(z) = z \prod_{\lambda \in \Lambda_{\tau} - \{(0,0)\}} (1 - \frac{z}{\lambda}) e^{\frac{z}{\lambda} + \frac{1}{2}(\frac{z}{\lambda})^2} = z \exp\left(\sum_{k \ge 1} -2G_{2k} \frac{z^{2k}}{2k}\right),
$$

$$
\zeta(z) = \frac{1}{z} + \sum_{\lambda \in \Lambda_{\tau} - \{(0,0)\}} \left(\frac{1}{z + \lambda} - \frac{1}{\lambda} + \frac{z}{\lambda^2}\right) = \frac{1}{z} - \sum_{k \ge 2} 2G_{2k} z^{2k-1}.
$$
 (2.4)

For later use, we introduce the following notation.

Definition 2.1. Define

$$
\mathbf{Y} = -\frac{\pi}{\text{im }\tau}, \quad \mathbf{A} = \mathbf{Y}(\bar{z} - z) = 2\pi \mathbf{i} \frac{\bar{z} - z}{\bar{\tau} - \tau}.
$$

Introduce

$$
\hat{\eta}_1 := \eta_1 + \Upsilon,
$$
\n
$$
2\hat{G}_k := 2G_k + \delta_{k,2}\Upsilon,
$$
\n
$$
\hat{\theta}(z) := \theta(z) \cdot \exp(-2\pi \frac{(\text{im } z)^2}{\text{im } \tau}),
$$
\n
$$
Z(z) := \partial_z \ln \theta = \zeta(z) - z\eta_1,
$$
\n
$$
\hat{Z}(z) := \partial_z \ln \hat{\theta} = \zeta(z) - z\hat{\eta}_1 + \Upsilon \bar{z}.
$$
\n(2.6)

Hereafter we suppress the notation for the variable \bar{z} that is linearly independent of z . However, when considering properties such as parity, a notation *f*(−*z*) stands for the quantity obtained from replacing both *z*, \bar{z} in f by $-z$, $-\bar{z}$.

We also recall the Eisenstein-Kronecker series [\[Wei76\]](#page-30-4).

Definition 2.2. The Eisenstein-Kronecker series are defined to be

$$
\mathcal{E}_m(z) = \sum_{\lambda \in \Lambda_{\tau}}^e \frac{1}{(z + \lambda)^m}, \quad m \ge 1.
$$
 (2.7)

Introduce the normalized versions

$$
\mathcal{E}_m^*(z) = (-1)^{m-1} (m-1)! \mathcal{E}_m(z) + (m-1)! \cdot 2G_m ,
$$

$$
\widehat{\mathcal{E}}_m^*(z) = \mathcal{E}_m^*(z) + \delta_{m,1} A(z) .
$$
 (2.8)

Definition [2.2](#page-4-0) and the relation [\(2.4\)](#page-4-1) give

$$
\partial_z \mathcal{E}_m = -m \mathcal{E}_{m+1}, \quad \mathcal{E}_m = \frac{(-1)^{m-1}}{(m-1)!} (\ln \theta)^{(m)}, \quad m \ge 1.
$$
 (2.9)

2.2 Laurent coefficients of the Kronecker theta function

Of crucial importance in this work are the Kronecker theta function *Sc*(*z*) [\[Wei76\]](#page-30-4) or Szegö kernel, and their Laurent coefficients which will be used to define the Eisenstein-Kronecker forms in Section [3.1](#page-10-0) below.

Definition 2.3. Set

$$
S_c(z) := \frac{\theta(z+c)}{\theta(z)\theta(c)}, \quad \widehat{S}_c(z) := e^{cA(z)} \frac{\theta(z+c)}{\theta(z)\theta(c)}.
$$
 (2.10)

These functions are meromorphic in *c*. Denote their Laurent expansions in *c* by

$$
S_c(z) = \frac{1}{c} \sum_{m \ge 0} c^m \frac{B_m(z)}{m!}, \quad \widehat{S}_c(z) = \frac{1}{c} \sum_{m \ge 0} c^m \frac{\widehat{B}_m(z)}{m!}.
$$
 (2.11)

Remark 2.4. The quantity $S_c(z)$ appears as the two-point function in chiral free fermions [\[Rai89\]](#page-29-5), and $\hat{S}_c(z)$ is its elliptic completion. The above Laurent coefficients enter the elliptic polylogarithms [\[BL11,](#page-28-3) Section 3.5] (see also [\[BKT07,](#page-28-4) [BKT10\]](#page-28-5)), and the GW generating series of elliptic curves [\[Zho23a,](#page-30-1) Section 4.3]. See [\[Spr19,](#page-30-5) [Spr20\]](#page-30-6) for a possible unified viewpoint via the Poincaré sheaf, and [\[FM20,](#page-29-6) [FM23\]](#page-29-7) for a purely algebraic description of these real-analytic objects.

2.2.1 Ring of almost-elliptic functions

The Laurent coefficients of the Kronecker theta function $S_c(z)$ can be computed concretely in terms of the Eisenstein-Kronecker series $\mathcal{E}_m(z)$, $m \geq 1$ as follows. One has from [\(2.9\)](#page-4-2) that

$$
\mathcal{E}_1(z+c) = \sum_{m\geq 0} \partial_z^m \mathcal{E}_1(z) \frac{c^m}{m!} = \sum_{m\geq 0} (-1)^m m! \, \mathcal{E}_{m+1}(z) \frac{c^m}{m!},
$$

and thus

$$
\frac{\theta(z+c)}{\theta(z)} = \exp\left(\sum_{m\geq 0} (-1)^m m! \, \mathcal{E}_{m+1}(z) \frac{c^{m+1}}{(m+1)!}\right) \,. \tag{2.12}
$$

Combining [\(2.4\)](#page-4-1) and [\(2.8\)](#page-4-3), this gives

$$
S_c(z) = \frac{1}{c} \exp\left(\sum_{m\geq 1} \frac{c^m}{m!} \mathcal{E}_m^*(z)\right).
$$
 (2.13)

By the set partition version of the Faà di Bruno formula, one then sees that *Bm*(*z*) is the complete Bell polynomial $B_m(\mathcal{E}_1^*, \mathcal{E}_2^*, \cdots, \mathcal{E}_m^*)$ in the variables $\mathcal{E}_k^*, k \ge 1$. Similarly, $\widehat{B}_m(z)$ is given by the complete Bell polynomial in the variables $\hat{\mathcal{E}}_k^*, k \geq 1$. More details regarding various expansions and formulas for $S_c(z)$, $\hat{S}_c(z)$ can be found in [\[Zag91\]](#page-30-7) (see also [\[BL11,](#page-28-3) [GM20,](#page-29-8) [Zho23a\]](#page-30-1)).

Definition 2.5. Let the notation be as above. Set

$$
e_m(z) := \frac{\mathcal{B}_m(z)}{m!}, \quad \widehat{e}_m(z) := \frac{\widehat{\mathcal{B}}_m(z)}{m!}, \quad m \ge 0,
$$
 (2.14)

and use the convention $e_m(z) = 0 = \hat{e}_m(z)$ if $m \le -1$. Define the rings of *quasi-elliptic* and *almost-elliptic* functions to be the graded polynomial rings^{[2](#page-5-1)}

$$
\widetilde{\mathfrak{F}} = \mathbb{C}\big[\{e_m\}_{m \ge 1}\big], \quad \widehat{\mathfrak{F}} = \mathbb{C}\big[\{\widehat{e}_m\}_{m \ge 1}\big], \tag{2.15}
$$

²The quantities $\{e_m(z)\}_{m\geq 1}$, $\{\hat{e}_m(z)\}_{m\geq 1}$ in fact satisfy relations such as ones induced by the Weierstrass relations. These relation however do not play an important role in this work.

respectively. Here for $m \geq 1$, the gradings of e_m , \hat{e}_m are m and the polynomial degrees are 1.

Using [\(2.2\)](#page-3-2), it is direct to see that the first few terms are given by

$$
\hat{e}_0(z) = 1
$$
, $\hat{e}_1(z) = \hat{Z}(z)$, $\hat{e}_2(z) = \frac{1}{2}(-\wp(z) + \hat{Z}^2(z))$. (2.16)

From the binomial identity, we also have

$$
\widehat{e}_m(z) = \sum_{k=0}^m e_k(z) \frac{A^{m-k}(z)}{(m-k)!}.
$$
\n(2.17)

For example, from [\(2.6\)](#page-4-4) and the automorphy

$$
\theta(z+1) = -\theta(z), \quad \theta(z+\tau) = -e^{-\pi i \tau} e^{-2\pi i z} \theta(z), \tag{2.18}
$$

one has the quasi-ellipticity and almost-ellipticity behavior

$$
e_1(z+1) = e_1(z), e_1(z+\tau) = e_1(z) - 2\pi i,
$$

\n $\hat{e}_1(z+1) = \hat{e}_1(z), \hat{e}_1(z+\tau) = \hat{e}_1(z).$ (2.19)

The generators $\{e_m(z)\}_{m>1}$, $\{\hat{e}_m(z)\}_{m>1}$ are actually quasi-Jacobi and almost-Jacobi forms of index zero with the weight given by the grading. See [\[EZ85,](#page-29-1) [Lib09,](#page-29-9) [GM20\]](#page-29-8) for more details.

We shall frequently make use of the following relation without explicit mentioning

$$
\widehat{\mathfrak{F}}\subseteq \widetilde{\mathfrak{F}}\otimes \mathbb{C}[A].
$$

The map $\hat{\mathfrak{F}} \to \tilde{\mathfrak{F}}$, obtained by taking the degree zero term in the polynomial expansion in *A*, offers an isomorphism of these two rings. We call this map the "holomorphic limit" and denote it by $\lim_{A=0}$; its inverse map is called the "elliptic completion".

Remark 2.6. Since ${B_m}_{m\geq0}$, ${\widehat{B}_m}_{m\geq0}$ are the complete Bell polynomials in the generators ${\{\mathcal{E}_m^*\}_{m\geq1}}$, $\{\widehat{\mathcal{E}}_m^*\}_{m\geq 1}$ *, the above rings can alternatively defined by taking the generators to be* $\{\mathcal{E}_m^*\}_{m\geq 1}$ *,* $\{\widehat{\mathcal{E}}_m^*\}_{m\geq 1}$ *,* respectively. While the latter sets are more convenient in analyzing properties relate to nonmeromorphicity (such as holomorphic anomaly [\[LZ23\]](#page-29-4)) due to the fact that {E[∗] *^m*}*m*≥² are all meromorphic, the former sets can sometimes be more convenient for actual computations due to the simple pole structure of them (see Lemma [2.7](#page-6-0) below), as we shall see in Section [3.](#page-9-0)

2.2.2 Singularities and differential relations of Laurent coefficients

We recall some simple facts about the Laurent coefficients $\hat{e}_m(z)$, $m \ge 1$, mostly from [\[BL11\]](#page-28-3). For self-containedness, we also give their short proofs.

Due to [\(2.19\)](#page-6-1) and [\[Zho23a,](#page-30-1) Equation A.9] (cf. [\[BKT10,](#page-28-5) Proposition 1.25]), *em*, *m* ≥ 2 is smooth at *z* ∈ **Z**, while e_1 = *Z* has a simple pole with residue 1 at *z* ∈ **Z**. Note that since the e_m 's are not almost-elliptic, these functions do not desend to functions on *E*.

We have the following more precise statement regarding the singularities of \hat{e}_m , $m \geq 1$.

Lemma 2.7. *Let the notation be as above.*

- *i*). *The functions* $\widehat{e}_m(z)$ *are elliptic in z, satisfying* $\widehat{e}_m(-z) = (-1)^m \widehat{e}_m(z)$ *.*
- *ii). The function* $\hat{e}_m(z)$, $m \ge 1$ *is smooth everywhere except for a simple pole along* $z \in \Lambda_\tau$ *. More precisely, the polar part of* $\hat{e}_m(z)$ *at* $z = 0$ *is*

$$
\frac{1}{z}\frac{(\mathbf{\Upsilon}\bar{z})^{m-1}}{(m-1)!}.
$$

- *Proof.* i). The ellipticity follows from that of their generating series $\hat{S}_c(z)$ which can be proved directly using the automorphy of θ . The parity follows from the parity of $A(z)$, $\theta(z)$ which gives $\widehat{S}_c(-z) = -\widehat{S}_{-c}(z)$.
- ii). The claim that \hat{e}_m is smooth and has at worst simple poles along $z \in \Lambda_\tau$ on *E* follows from the fact that the generating series $\hat{S}_c(z)$ of the \hat{e}_m 's is so. The claim about its polar part follows from the following formula for the generating series of the polar parts of all \hat{e}_m 's

$$
\frac{1}{z} \cdot \left(\frac{\theta(z+c)}{\theta(c)} e^{cA} \right) |_{z=0} = \frac{1}{z} \cdot e^{cY\overline{z}},
$$

where we have used the fact that z , \bar{z} are independent complex quantities.

 \Box

We next study some differential and quadratic relations between the $\hat{e}_m(z)$'s.

Definition 2.8. Suppose c_i , $i \in [N]$ are constants which are not necessarily distinct. Denote a formal sum ⊕ on the space of sequences

$$
\mathbf{m}=(m_1,\cdots,m_i,\cdots,m_N)\in\mathbb{N}^N.
$$

For two sequences $m \in \mathbb{N}^N$, $n \in \mathbb{N}^N$, define

$$
\widehat{f}_{\mathbf{m}} = \prod_{i=1}^N \widehat{e}_{m_i}(z+c_i), \quad \widehat{f}_{\mathbf{m}\oplus\mathbf{n}} = \widehat{f}_{\mathbf{m}} + \widehat{f}_{\mathbf{n}}.
$$

For $m \in \mathbb{N}^N$, define $|m| = \sum_{i=1}^N m_i$. For $\ell \in \mathbb{N}$, define

$$
m \ominus 1 := \bigoplus_{i=1}^N (m_1, \cdots, m_i - 1, \cdots, m_N), \quad m \ominus \ell := m \underbrace{\ominus 1 \ominus 1 \cdots \ominus 1}_{\ell \text{ times}}.
$$

Lemma 2.9. *Let the notation be as above.*

i). One has

$$
\partial_{\bar{z}}\widehat{e}_m(z)=\Upsilon\widehat{e}_{m-1}(z),\quad m\geq 1.
$$

ii). Suppose $m_i \geq 1$ *,* $i \in [N]$ *and* c_i *,* $i \in [N]$ *are constants which are not necessarily distinct. Denote* ${\bf r}$ = (m_1 , ⋅ ⋅ ⋅ , m_{N-1}) ∈ ${\mathbb N}^{N-1}$ *. Then*

$$
\begin{array}{rcl}\n\hat{Y}_{\text{fm}} & = & \partial_{\bar{z}} \left(\sum_{\ell \geq 0} (-1)^{\ell} \hat{f}_{\text{r} \ominus \ell} \cdot \hat{e}_{m_{N} + \ell + 1} \right) \\
& = & \partial_{\bar{z}} \left(\sum_{k_{1}, \dots, k_{N-1}} (-1)^{\sum_{i=1}^{N-1} k_{i}} \prod_{i=1}^{N-1} \hat{e}_{m_{i} - k_{i}}(z + c_{i}) \cdot \hat{e}_{m_{N} + \sum_{i=1}^{N-1} k_{i} + 1}(z + c_{N}) \right).\n\end{array}
$$

iii). Assume $x \neq y$ modulo Λ_{τ} , then one has

$$
\begin{array}{rcl}\n\widehat{e_i}(x)\widehat{e_j}(y) & = & \sum\limits_{b=0}^{i+j} \left(\sum\limits_{\ell=0}^i (-1)^\ell \binom{b}{j+\ell} \right) \widehat{e_{i+j-b}}(x-y)\widehat{e_b}(y) \\
& + & \sum\limits_{a=0}^{i+j} (-1)^{i+j-a} \left(\sum\limits_{\ell=0}^i (-1)^\ell \binom{a}{i-\ell-1} \right) \widehat{e_{i+j-a}}(x-y)\widehat{e_a}(x)\,.\n\end{array}
$$

In particular, one has for $m \geq 1$

$$
\widehat{e}_1(x)\widehat{e}_m(y)=\widehat{e}_1(x-y)\widehat{e}_m(y)+\widehat{e}_{m+1}(y)+\sum_{k+\ell=1+m}\widehat{e}_k(x)\widehat{e}_\ell(y-x).
$$

Proof. i). The first part of the claim is essentially [\[BL11,](#page-28-3) Equation 3.6]. It follows by comparing the Laurent coefficients of the following identity [\[BL11,](#page-28-3) Lemma 6] proved by a direct computation

$$
\partial_{\bar{z}}(\widehat{S}_c(z)) = c \partial_{\bar{z}} A \widehat{S}_c(z) = c \mathbf{Y} \widehat{S}_c(z). \tag{2.20}
$$

ii). From (i) and the product rule, we have

$$
\partial_{\bar{z}}\widehat{f}_{\bf r}=\widehat{f}_{{\bf r}\ominus 1}.
$$

Integration by parts gives

$$
\mathbf{Y}\widehat{f}_{\mathbf{r}}\widehat{e}_{m_N}=\widehat{f}_{\mathbf{r}}\cdot\partial_{\bar{z}}\widehat{e}_{m_N+1}=\partial_{\bar{z}}(\widehat{f}_{\mathbf{r}}\widehat{e}_{m_N+1})-\mathbf{Y}\widehat{f}_{\mathbf{r}\ominus 1}\widehat{e}_{m_N+1}.
$$

Applying this identity iteratively to the part involving *Y* and noting that $f_{r\ominus\ell} = 0$ for $\ell > |r|$ (since $\hat{e}_m = 0$ for $m < 0$), we obtain the first identity. The second identity follows directly.

iii). This is part of the quadratic Arnold-type relations given in [\[BL11,](#page-28-3) Equation 3.7] which are derived by using Fay's trisecant identity. To be more precise, Fay's trisecant identity gives

$$
S_a(x)S_b(y) = S_a(x - y)S_{a+b}(y) + S_{a+b}(x)S_b(y - x),
$$

and thus the identity for the \hat{S} version. Multiplying the resulting identity throughout by $ab(a + b)$ and comparing the Taylor coefficients in a, b , one obtains

$$
\widehat{e}_i(x)\widehat{e}_j(y)=-\widehat{e}_{i-1}(x)\widehat{e}_j(y)+\sum_{k+\ell=i+j} {\ell \choose j}\widehat{e}_k(x-y)\widehat{e}_\ell(y)+\sum_{k+\ell=i+j} {k \choose i-1}\widehat{e}_k(x)\widehat{e}_\ell(y-x).
$$

Iterating, one has

$$
\begin{array}{rcl}\n\widehat{e_i}(x)\widehat{e_j}(y) & = & \sum\limits_{\ell=0}^i (-1)^\ell \sum\limits_{a+b=i+j} \left(\binom{b}{j+\ell} \widehat{e_a}(x-y) \widehat{e_b}(y) + \binom{a}{i-\ell-1} \widehat{e_a}(x) \widehat{e_b}(y-x) \right) \\
& = & \sum\limits_{a+b=i+j} \widehat{e_a}(x-y) \widehat{e_b}(y) \sum\limits_{\ell=0}^i (-1)^\ell \binom{b}{j+\ell} \\
& & + & \sum\limits_{a+b=i+j} \widehat{e_a}(x) \widehat{e_b}(y-x) \sum\limits_{\ell=0}^i (-1)^\ell \binom{a}{i-\ell-1}.\n\end{array}
$$

Simplifying, we obtain the desired claim.

 \Box

Remark 2.10.

i). The generating series technique also gives rise to formulas expressing $∂_zê_m$ in terms of elements $\hat{\mathbf{m}} = \mathbb{C}[\{\hat{\epsilon}_m\}_{m\geq1}] = \mathbb{C}[\{\hat{\epsilon}_m^*\}_{m\geq1}]$ that are parallel to [\(2.9\)](#page-4-2). To be more explicit, from [\(2.12\)](#page-5-2) and [\(2.8\)](#page-4-3), [\(2.9\)](#page-4-2) we obtain

$$
\partial_z \widehat{S}_c(z) = \widehat{S}_c(z) \cdot \left(-c\mathbf{Y} + \sum_{m \geq 1} (-1)^m \mathcal{E}_{m+1} c^m \right)
$$

=
$$
\frac{1}{c} \sum_{m \geq 0} c^m \widehat{e}_m \cdot \left(\sum_{m \geq 1} \frac{c^m}{m!} (\widehat{\mathcal{E}}_{m+1}^* - m! \cdot 2 \widehat{G}_{m+1} - \delta_{m,0} \mathbf{A}(z)) \right).
$$

Comparing Laurent coefficients in *c* gives a desired relation

$$
\partial_z \widehat{e}_m(z) = \sum_{\substack{a+b=m\\b\geq 1}} \widehat{e}_a(z) \cdot \frac{1}{b!} (\widehat{\mathcal{E}}_{b+1}^* - b! \cdot 2\widehat{G}_{b+1}). \tag{2.21}
$$

This lands in the ring $M \otimes \tilde{\mathfrak{F}}$, where $M = \mathbb{C}[\{\tilde{G}_k\}_{k \geq 1}]$ is the ring of almost-holomorphic modular forms [\[KZ95\]](#page-29-2) with the generators $\{G_k\}_{k\geq 1}$ introduced in Definition [2.1.](#page-4-5) The ring \hat{M} is isomorphic to $\mathbb{C}[G_2, G_4, G_6]$ after quotienting out the relations among the generators.

ii). Setting $m = 1$ in Lemma [2.9](#page-7-0) (iii) and using Definition [2.2](#page-4-0) and [\(2.16\)](#page-6-2), one arrives at the familiar addition formula

$$
(\zeta(x) + \zeta(y) + \zeta(-x - y))^2 = \wp(x) + \wp(y) + \wp(-x - y).
$$

3 Iterated integrals of Eisenstein-Kronecker forms

In this part we compute iterated regularized integrals of Eisenstein-Kronecker forms and discuss their connections to Chen's iterated path integrals.

We first construct some differential forms in terms of the linear coordinate *z* on the universal cover **C** of *E*. A holomorphic volume form on *E* is given by *dz*, and a volume form vol satisfying \int_E vol = 1 is

$$
\text{vol} := \frac{\mathbf{i}}{2 \operatorname{im} \tau} dz \wedge d\bar{z} \,. \tag{3.1}
$$

We also denote

$$
\beta = \frac{d\overline{z} - dz}{\overline{\tau} - \tau} \,. \tag{3.2}
$$

Note that the cohomology class of *β* in *H*¹ (*E*, **C**) is the Poincaré dual of the *A*-cycle class in the weak Torelli marking {*A*, *B*}.

Recall that $A_E^1(\star D)$ is the space of 1-forms that are smooth everywhere on *E* except for possible holomorphic poles along an effective divisor *D* on *E*. One can introduce a notion [\[FK17\]](#page-29-10) (see also [\[LZ21\]](#page-29-3)) of holomorphic residue res*[∂]* that acts on *A* 1 *E* (⋆*D*). Concretely, this is given by

$$
res_{\partial} = \frac{1}{2\pi i} \lim_{\varepsilon \to 0} \int_{\partial B_{\varepsilon}(D)} , \qquad (3.3)
$$

where $B_{\varepsilon}(D)$ is a disk bundle of radius ε over the divisor *D*. The result is independent of the choice of the local defining equation for *D* and the Hermitian metric that is compatible with the complex structure. In particular, from the description of the right hand side of [\(3.3\)](#page-9-1) one has [\[LZ21,](#page-29-3) Lemma 2.2]

$$
res_{\partial}(\omega) = 0, \quad \forall \omega \in A_E^{0,1}(\star D). \tag{3.4}
$$

An analytic notion of regularized integral

$$
\oint_{E} : A_{E}^{1}(\star D) \to \mathbb{C}
$$
\n(3.5)

is introduced, extended, and further developed in [\[LZ21,](#page-29-3) [LZ23\]](#page-29-4). It extends the ordinary notion of integral on smooth forms and provides a nice regularization scheme that enjoys many nice properties such as the Fubini theorem. A purely cohomological formulation of this notion is provided in [\[Zho23b\]](#page-30-2).

For a function $f \in A^0_E(\star D)$, when no confusion might arise we adapt short-hand notation such as

$$
\oint_{E} f := \oint_{E} f \text{ vol}, \quad \int_{A} f := \int_{A} f dz.
$$
\n(3.6)

Similarly, we write res*∂*, *^z*=*^p* (*f*) for the residue res*∂*, *^z*=*^p* (*f dz*) of the corresponding differential *f dz* at *z* = *p*, and write res*[∂]* (*f*) for the summation of all local residues.

Set $[n] := (1, 2, \dots, n)$, denote $E_{[n]} := E_n \times E_{n-1} \times \dots \times E_1$, where E_k represents a copy of elliptic curve equipped with the corresponding linear coordinate z_k , $k \in [n]$. For a function $f \in A^0_{E_{[n]}} (\star D)$, we use the following short-hand notation for the iterated residues, iterated *A*-cycle integrals, and iterated regularized integrals

$$
\begin{array}{rcl}\n\operatorname{res}_{\partial}(f) & := & \operatorname{res}_{\partial} dz_1 \cdots \operatorname{res}_{\partial} dz_2 \operatorname{res}_{\partial} (f dz_1), \\
\int_{A_{\sigma([n])}} f & := & \int_{A_{\sigma(n)}} dz_{\sigma(n)} \cdots \int_{A_{\sigma(2)}} dz_{\sigma(2)} \int_{A_{\sigma(1)}} f dz_{\sigma(1)}, \\
\int_{E_{\sigma([n])}} f & := & \int_{E_{\sigma(n)}} \operatorname{vol}_{\sigma(n)} \cdots \int_{E_{\sigma(2)}} \operatorname{vol}_{\sigma(2)} \int_{E_{\sigma(1)}} f \operatorname{vol}_{\sigma(1)}, \quad \sigma \in \mathfrak{S}([n]).\n\end{array} \tag{3.7}
$$

Here the integration domain $A_{\sigma([n])}$ is constructed such that it avoids the singularities of *f*, with

$$
A_i = \{\varepsilon_i \tau + \tau + t \mid t \in [0,1]\}, \quad \varepsilon_{\sigma(n)} < \dots < \varepsilon_{\sigma(2)} < \varepsilon_{\sigma(1)}
$$

for sufficiently small positive real numbers ε_i , $i \in [n]$. The integration domain in $f_{E_{\sigma([n])}}$ can also be constructed explicitly [\[LZ21,](#page-29-3) Section 4.3]. Note that unlike the iterated residues and iterated *A*-cycle integrals, the iterated regularized integral in [\(3.7\)](#page-10-1) is independent of the ordering of the iteration [\[LZ21\]](#page-29-3) and is therefore simply called the regularized integral.

3.1 Eisenstein-Kronecker forms

Since by Lemma [2.7](#page-6-0) one has $\hat{e}_m(z) \in A_E^0(*0)$, one can then consider their holomorphic residues and regularized integrals, as defined in [\[LZ21\]](#page-29-3) and further developed in [\[LZ23,](#page-29-4) [Zho23b\]](#page-30-2).

We first establish the following simple result which indicates that the set of generators $\{\widehat{e}_m(z)\}_{m>1}$, which have at worst simple poles, is an ideal one (in contrast to e.g., $\{\widehat{\mathcal{E}}_m^*(z)\}_{m\geq 1}$) for exact computations.

Definition 3.1. Let the notation and conventions be as above. Consider the following subvector space of $\hat{\mathfrak{F}} = \mathbb{C}\big[\{\widehat{e}_m(z)\}_{m \geq 1}\big]$ consisting of the linear span of degree 0 and 1 elements

$$
\mathcal{V} = \bigoplus_{m \ge 0} \mathbb{C} \,\widehat{e}_m(z) \,. \tag{3.8}
$$

We call Vdz , Vvol the space of Eisenstein-Kronecker forms.

Lemma 3.2. *Let the notation be as above. Then one has*

$$
\operatorname{res}_{\partial}(\widehat{e}_m(z)dz)=\delta_{m,1},\quad \int_E \widehat{e}_m(z)\operatorname{vol}=\delta_{m,0}.
$$

Proof. We prove the statements by computing the holomorphic residue res*[∂]* and regularized integral f_E of the generating series $\hat{S}_c(z)$ of $\hat{e}_m(z)$, $m \ge 0$. The stament on the holomorphic residue follows from Lemma [2.7.](#page-6-0) For the regularized integral, from [\(2.20\)](#page-8-0) we have

$$
\bar{\partial}(\widehat{S}_c(z)dz) = c \cdot 2\pi i \beta \wedge \widehat{S}_c(z)dz = -c \cdot 2\pi i \widehat{S}_c(z)dz \wedge \psi = -c \cdot 2\pi i \widehat{S}_c(z) \text{vol},
$$

and thus

$$
\oint_E \widehat{S}_c(z) \text{vol} = -\frac{1}{2\pi i c} \oint_E \bar{\partial}(\widehat{S}_c(z) dz).
$$

According to the Cauchy integral formula [\[LZ21\]](#page-29-3) for the regularized integral (see also [\[Zho23b,](#page-30-2) Section 2.2.2] for a review), this gives

$$
\oint_{E} \widehat{S}_c(z) \text{vol} = \frac{1}{c} \text{res}_{\partial} (\widehat{S}_c(z) dz) = \frac{1}{c} \text{res}_{\partial} (\frac{1}{z} e^{cA} dz).
$$

The desired result then follows from the fact

$$
\operatorname{res}_{\partial}\left(\frac{\bar{z}^k}{z^n}dz\right) = 0, \quad k \ge 1,\tag{3.9}
$$

 \Box

proved using [\(3.3\)](#page-9-1).

Remark 3.3. Note that a smooth function with holomorphic poles can be even under the action $(z,\bar{z}) \mapsto (-z,-\bar{z})$ and have a simple pole at $z = 0$. For example, from [\(2.16\)](#page-6-2) and [\(2.4\)](#page-4-1), one can see that locally near $z = 0$ one has

$$
\widehat{e}_1^2(z)=\widehat{Z}^2(z)=\frac{1}{z^2}+2\Upsilon\frac{\bar{z}}{z}+\cdots, \quad \widehat{e}_2(z)=\frac{1}{2}\cdot(2\Upsilon\frac{\bar{z}}{z}+\cdots).
$$

Both of them have trivial holomorphic residues under res*[∂]* .

In fact, according to [\(3.9\)](#page-11-0), as long as the holomorphic residue operation res*[∂]* is concerned, monomials in the $\hat{e}_m(z)$'s behave as if they were ordinary meromorphic functions. In particular, from Lemma [2.7](#page-6-0) and [\(2.4\)](#page-4-1) we have

$$
\operatorname{res}_{\partial}(\hat{e}_{m}^{N}dz) = 0, \quad m \ge 2, \qquad \operatorname{res}_{\partial}(\hat{e}_{1}^{N}dz) = \sum_{\substack{(k_{1}, \cdots, k_{N}) \\ \sum_{i=1}^{N} (2k_{i}-1) = -1}} \prod_{i=1}^{N} (-2\hat{G}_{2k_{i}}), \tag{3.10}
$$

where we have set the convention $-2G_0 = -1$.

We shall also study iterated residues and iterated regularized integrals of almost-elliptic functions of the following type.

Definition 3.4. Let the notation be as above. Fix a set of generic values $c_{k,\ell}, k, \ell \in [n]$. Consider the smooth hypersurface arrangement D in $E_{[n]}$ given by (the image under the quotient $\mathbb{C}_{[n]} \to E_{[n]}$)

$$
D = \bigcup_{\substack{k,\ell \in [n] \\ k \neq \ell}} \{s_{k,\ell} = 0\}, \quad s_{k,\ell} := c_{k,\ell} + z_k - z_\ell.
$$
 (3.11)

We also regard $s_{k,l}$ as defining a morphism $E_{[n]} \to E$. Define the following spaces

$$
\mathcal{F}_{\mathsf{D}} = \bigotimes_{k,\ell \in [n]} s_{k,\ell}^* \widehat{\mathfrak{F}}, \quad \mathcal{V}_{\mathsf{D}} = \bigotimes_{k \in [n]} \left(\bigoplus_{\ell \in [n]} s_{k,\ell}^* \mathcal{V} \right), \quad \mathcal{V}_{\mathsf{D}}^{\circ} = \bigotimes_{k \in [n]} s_{k,k+1}^* \mathcal{V}, \tag{3.12}
$$

with the conventions $k \neq \ell$, $z_{n+1} = z_1$, $c_{n,n+1} = c_{n,1}$. In this work we call the latter two spaces of Eisenstein-Kronecker forms associated to $D \subseteq E_{[n]}$, with convention [\(3.7\)](#page-10-1) understood.

One has the following bases of the vector spaces \mathcal{F}_{D} , \mathcal{V}_{D} , \mathcal{V}_{D}° given in Table [1.](#page-12-0)

vector space	basis: $k, \ell \in [n]$, $k \neq \ell$
\mathcal{F}_{D}	$\prod_{k=1}^n \prod_{\ell=1}^n \hat{e}_{m_{k,\ell}}^{r_{k,\ell}}(s_{k,\ell}), \quad r_{k,\ell} \geq 0$
\mathcal{V}_{D}	$\left \prod_{k=1}^n \prod_{\ell=1}^n \hat{c}_{m_{k,\ell}}^{r_{k,\ell}}(s_{k,\ell}), \right r_{k,\ell} \in \{0,1\}, \quad \sum_{\ell=1}^n r_{k,\ell} \leq 1$
$\mathcal{V}_\mathsf{D}^\circ$	$\prod_{k=1}^n \hat{e}_{m_{k,k+1}}^{r_{k,k+1}}(s_{k,k+1}), \quad r_{k,\ell} \in \{0,1\}$

Table 1: Bases for the vector spaces \mathcal{F}_{D} , \mathcal{V}_{D} , \mathcal{V}_{D}° .

Many constructions and results regarding these spaces are most easily formulated and visualized by the so-called *indicating graphs* which we now introduce.

Definition 3.5. For any monomial $f = \prod_{k,\ell=1}^n \hat{e}_{m_{k,\ell}}^{r_{k,\ell}}(s_{k,\ell}) \in \mathcal{F}_D$, its indicating graph $\Gamma(f)$ is defined to be the following graph:

- it has *n* vertices labelled by $k \in [n]$,
- for each factor $\hat{e}_{m_{k,\ell}}^{r_{k,\ell}}(s_{k,\ell})$ with $k \neq \ell$, assign $r_{k,\ell}$ oriented edges starting at *k* and ending at ℓ if $m_{k,\ell} \geq 1$ $m_{k,\ell} \geq 1$, and no edges if $m_{k,\ell} = 0$. See Figure 1 below for an illustration.

For each vertex k , let $v_{k,in}$, $v_{k,out}$ be the inner and outer valencies at k . Since each edge contributes one inner valency and one outer valency, we have

$$
\sum_{k=1}^{n} v_{k,in} = \sum_{k=1}^{n} v_{k,out}.
$$
\n(3.13)

The geometric data of the monomial *f* is conveniently encoded in the indicating graph Γ(*f*). For example:

Figure 1: *Indicating graphs that are chain, loop, tree, respectively. Here the labellings of the vertices are omitted.*

- the number of edges, given by $\delta_{m_k,\ell \geq 1} \cdot r_{k,\ell}$, corresponds to the degree of the monomial $\hat{e}_{m_k,\ell}^{r_{k,\ell}}(s_{k,\ell})$. This is also the *multiplicity* of the irreducible component $(s_{k,\ell} = 0)$ of the polar divisor of $\hat{e}_{m_{k,\ell}}^{r_{k,\ell}}(s_{k,\ell})$. In particular, $\otimes_{k,\ell \in [n]} s_{k,\ell}^* \mathcal{V}$ is the subspace of \mathcal{F}_{D} that consists of elements whose corresponding polar divisors are reduced. This space is of great interest in that it underlies a model for the cohomology of $E_{[n]}$ – D, as shown in [\[BL11\]](#page-28-3).
- the outer valency at the vertex *k* is given by $v_{\ell, \text{out}} = \sum_{k \neq \ell} \delta_{m_{k,\ell} \geq 1} r_{k,\ell}$. In particular,

$$
f \in \mathcal{V}_{\mathsf{D}} \quad \Rightarrow \quad v_{\ell, \text{out}} \le 1 \, \text{ for } \ell \in [n]. \tag{3.14}
$$

• decomposition of Γ(*f*) into connected components corresponds to factorization of *f* and thus also those of the operations res_∂, $f_{E_{[n]}}$ on *f*. In particular, the existence of a trivial connected component (i.e., one with only one vertex *b* and no edges) represents the fact that *f* is independent of z_b and thus the f_{E_b} part in $f_{E_{[n]}}$ gives the identity operator.^{[3](#page-13-2)}

3.2 Iterated residues and iterated regularized integrals

We next study iterated residues and regularized integrals for elements in V_D . The more general case \mathcal{F}_{D} will not be discussed here since the combinatorics is more involved (and can be described using rooted labelled forests as exhibited in [\[LZ23\]](#page-29-4)).

3.2.1 Reduction from \mathcal{V}_D to $\mathcal{V}_\mathsf{D}^\circ$

We first have the following simple fact based on combinatorial properties of the indicating graphs.

Lemma 3.6. *Let the notation be as above.*

i.) Let the notation be as above. Let $f = \prod_{k=1}^n \prod_{\ell=1}^n \hat{\epsilon}_{m_{k\ell}}^{r_{k\ell}}(s_{k\ell}) \in \mathcal{F}_{\mathsf{D}}$. If there exists some vertex b in $\Gamma(f)$ *such that*

$$
v_{b,in}+v_{b,out}=1,
$$

then $f_{E_{[n]}} f = 0$ *.*

ii.) Let $f = \prod_{k=1}^n \prod_{\ell=1}^n \hat{e}_{m_{k\ell}}^{r_{k\ell}}(s_{k\ell}) \in V_D$. Then

$$
\int_{E_{[n]}} f \neq 0
$$

³Thanks to the fact that iterated regularized integral $f_{E_{[n]}}f$ is independent of the ordering for the iteration [\[LZ21\]](#page-29-3).

only if any connected component of the indicating graph Γ(*f*) *is trivial (i.e., has no edges) or a loop.*

Proof. i.) Denote the other vertex of the unique edge at *b* by *c*, then the function *f* must have the form

$$
f = \hat{e}_m(s_{bc})g
$$
 or $\hat{e}_m(s_{cb})g$,

for some $m \geq 1$ and function g that is independent of z_b . Applying Lemma [3.2](#page-11-1) we see that

$$
\int_{E_b} f \text{vol}_b = 0.
$$

This gives the desired claim.

ii.) By the factorization property of the iterated regularized integration, one can assume that Γ(*f*) is already connected. There is nothing to prove if it has no edges. Otherwise, from [\(3.13\)](#page-12-1) and [\(3.14\)](#page-13-3) we have

$$
\sum_{k=1}^{n} (v_{k,in} + v_{k,out}) = 2 \sum_{k=1}^{n} v_{k,out} \le 2n.
$$

If there exists a vertex, say *a*, with

 $v_{a,in} + v_{a,out} > 2$,

then there must exist another vertex *b* such that

$$
v_{b,\text{in}} + v_{b,\text{out}} \leq 1.
$$

Since $\Gamma(f)$ is connected, one can not have $v_{b,in} + v_{b,out} = 0$. Thus $v_{b,in} + v_{b,out} = 1$ and by (i) the regularized integral $f_{E_{[n]}}f$ is zero. Therefore, $f_{E_{[n]}}f$ is nonzero only if

$$
v_{a,\text{in}} + v_{a,\text{out}} = 2, \quad a \in [n].
$$

This also tells that the number of edges is $2n/2 = n$. It is now easy to see (for example by computing the Euler characteristic in two ways) that Γ(*f*) is a loop.

 \Box

Lemma [3.6](#page-13-4) allows to reduce the study of regularized integral for elements in V_D to that for elements in \mathcal{V}_{D}° . However, this simplification does not apply directly to iterated residues since the latter depend on the ordering of iteration.

3.2.2 Evaluation of iterated residues and iterated regularized integrals

For later simplification, we introduce the following short-hand notation.

Definition 3.7. Let the notation be as above. Define

$$
\widehat{e}_{m} = \prod_{k=1}^{n} \widehat{e}_{m_k}(s_{k,k+1}), \quad m = (m_1, m_2, \cdots, m_n) \in \mathbb{N}^n.
$$

Then

$$
\mathcal{V}_{\mathsf{D}}^{\circ} = \bigoplus_{\mathsf{m} \in \mathbb{N}^n} \mathbb{C} \,\widehat{e}_{\mathsf{m}}\,. \tag{3.15}
$$

The number of nonzero entries in the sequence m is called the length and denoted by $\ell(m)$, while the quantity $|m| := \sum_{k=1}^m m_k$ is called the size of m. They correspond to the degree and grading of $\widehat{e}_{m} \in \mathcal{F}_{D}$, respectively.

Following the methods developed in [\[LZ21,](#page-29-3) [LZ23\]](#page-29-4), we prove the following results.

Proposition 3.8. Let the notation and conventions be as above. Let $n \geq 2$ and z_0 be a constant. Define

$$
\zeta_{\text{chain}} = \prod_{k=1}^{n-1} \widehat{e}_{m_k}(s_{k,k+1}) \cdot \widehat{e}_{m_n}(z_n - z_0), \quad \zeta_{\text{loop}} = \widehat{e}_{\text{m}}.
$$

i). One has

$$
\operatorname{res}_{\partial}(\zeta_{\mathrm{chain}} \cdot \wedge_{k=1}^{n} dz_{k}) = \prod_{k=1}^{n} \delta_{m_{k},1}, \quad \operatorname{res}_{\partial}(\zeta_{\mathrm{loop}} \cdot \wedge_{k=1}^{n} dz_{k}) = 0.
$$

ii). One has

$$
\int_{E_{[n]}} \xi_{\text{chain}} \cdot \wedge_{k=1}^{n} \text{vol}_{k} = \prod_{k=1}^{n} \delta_{m_{k},0},
$$
\n
$$
\int_{E_{[n]}} \xi_{\text{loop}} \cdot \wedge_{k=1}^{n} \text{vol}_{k} = \hat{e}_{|m|} \left(\sum_{k=1}^{n} s_{k,k+1} \right) \cdot \left(\prod_{k=1}^{n} \delta_{m_{k},0} - \prod_{k=1}^{n} (\delta_{m_{k},0} - 1) \right).
$$

The nomenclature of *ξ*chain, *ξ*loop originates from the types of the indicating graphs as shown in Figure [1.](#page-13-1)

Proof. i). The first relation follows by direct computations using Lemma [3.2.](#page-11-1)

For the second relation, consider $C_m = \text{res}_{\partial}(\hat{e}_m \cdot \wedge_{k=1}^n dz_k)$. From Lemma [3.2](#page-11-1) and the expression for $t_{k,l}$ given in Definition [\(3.4\)](#page-12-2), we see that

$$
\text{res}_{\partial} \left(\hat{e}_{m_1}(s_{12}) \hat{e}_{m_n}(s_{n1}) dz_1 \right) = \delta_{m_1,1} \cdot \hat{e}_{m_n}(s_{n1}) \big|_{s_{12}=0} + (-1)^{m_n} \delta_{m_n,1} \cdot \hat{e}_{m_1}(s_{12}) \big|_{s_{n1}=0}
$$
\n
$$
= \delta_{m_1,1} \cdot \hat{e}_{m_n}(s_{n1} + s_{12}) - \delta_{m_n,1} \cdot \hat{e}_{m_1}(s_{n1} + s_{12}). \tag{3.16}
$$

.

Then using [\(3.16\)](#page-15-0) we have the recursion

$$
C_{(m_1,m_2,\cdots,m_n)}=\delta_{m_1,1}C_{(m_2,\cdots,m_{n-1},m_n)}-\delta_{m_n,1}C_{(m_2,\cdots,m_{n-1},m_1)}
$$

The values for the initial case $n = 2$ are

$$
C_{(m_1,m_2)} = \operatorname{res}_{\partial} dz_2 \left(\delta_{m_1,1} \cdot \hat{e}_{m_n}(s_{n1} + s_{12}) - \delta_{m_n,1} \cdot \hat{e}_{m_n}(s_{21} + s_{12}) \right) = 0,
$$

since $s_{21} + s_{12}$ is a constant independent of z_2 . Iterating, one proves the desired claim.

ii). The first relation follows from Lemma [3.6.](#page-13-4)

For the second relation, using Lemma [2.7](#page-6-0) (i), Lemma [2.9](#page-7-0) (ii), and the Cauchy integral formula [\[LZ21\]](#page-29-3) for the regularized integral as in Lemma [3.2,](#page-11-1) we have

$$
\int_{E_1} \hat{e}_{m_1}(s_{12}) \hat{e}_{m_n}(s_{n1}) \text{vol}_1 = (-1)^{m_n} \int_{E_1} \hat{e}_{m_1}(s_{12}) \hat{e}_{m_n}(-s_{n1}) \text{vol}_1
$$
\n
$$
= (-1)^{m_n} \cdot \frac{-1}{2\pi \mathbf{i}} \int_{E_1} \bar{\partial} \left(\sum_{k=0}^{m_1} (-1)^k \hat{e}_{m_1-k}(s_{12}) \hat{e}_{m_n+1+k}(-s_{n1}) dz \right)
$$
\n
$$
= (-1)^{m_n} \text{res}_{\partial} \left(\sum_{k=0}^{m_1} (-1)^k \hat{e}_{m_1-k}(s_{12}) \hat{e}_{m_n+1+k}(-s_{n1}) dz \right)
$$
\n
$$
= -\text{res}_{\partial} \left(\sum_{k=0}^{m_1} \hat{e}_{m_1-k}(s_{12}) \hat{e}_{m_n+1+k}(s_{n1}) dz_1 \right).
$$

Using [\(3.16\)](#page-15-0), this gives

$$
\oint_{E_1} \widehat{e}_{m_1}(s_{12}) \widehat{e}_{m_n}(s_{n1}) \text{vol}_1 = -\delta_{m_1 \ge 1} \cdot \widehat{e}_{m_n + m_1}(s_{n1} + s_{12}) + \delta_{m_n,0} \cdot \widehat{e}_{m_n + m_1}(s_{n1} + s_{12}). \tag{3.17}
$$

Now set

$$
C_{\mathbf{m}} = \frac{1}{\widehat{e}_{|\mathbf{m}|}(\sum_{k=1}^{n} s_{k,k+1})} \int_{E_{[n]}} \widehat{e}_{\mathbf{m}} \cdot \wedge_{k=1}^{n} \mathrm{vol}_{k}.
$$

Then using [\(3.17\)](#page-16-0) one has the recursion

$$
C_{(m_1,m_2,\cdots,m_n)}=(\delta_{m_n,0}-\delta_{m_1\geq 1})C_{(m_2,\cdots,m_n+m_1)},
$$

with the values for the initial case $n = 2$ determined from

$$
\oint_{E_{[2]}} \widehat{e}_m \cdot \wedge_{k=1}^2 \mathrm{vol}_k = \oint_{E_2} (\delta_{m_2,0} - \delta_{m_1 \ge 1}) \widehat{e}_{m_2+m_1}(s_{21}+s_{12}) \mathrm{vol}_2 = (\delta_{m_2,0} - \delta_{m_1 \ge 1}) \widehat{e}_{m_2+m_1}(s_{21}+s_{12}).
$$

If $m_n = 0$, then ξ_{loop} reduces to the chain type one ξ_{chain} discussed above and the result is $\prod_{k=1}^{n-1} \delta_{m_k,0}$. Otherwise, we have

$$
C_m = \prod_{k=1}^{n-1} (-\delta_{m_k \ge 1}) = \prod_{k=1}^{n-1} (\delta_{m_k,0} - 1).
$$

Combining these results and noting that $\hat{e}_0(z) = 1$, this proves the desired claim.

- \Box
- **Remark 3.9.** i). Strictly speaking, partial regularized integration such as the one in [\(3.17\)](#page-16-0) does not preserve the space \mathcal{F}_{D} : for example the polar divisor $s_{n1} + s_{12} = 0$ does not necessarily lie in D. Thus one needs to enlarge the divisor D so that it includes all divisors defined by any linear combination of the *sij*'s. This does not cause trouble since all existing results apply to the enlarged divisor straightforwardly.
- ii). The relation [\(3.17\)](#page-16-0) can be alternatively derived by reducing an iterated regularized integral of loop type to one of chain type as follows. From Lemma [2.9](#page-7-0) (iii), we see that 4

$$
\begin{array}{lll}\n\widehat{e}_{m_1}(s_{12})\widehat{e}_{m_n}(s_{n1}) & = & (-1)^{m_n}\widehat{e}_{m_1+m_n}(s_{n1}+s_{12})\delta_{m_n,0} - \widehat{e}_{m_1+m_n}(s_{n1}+s_{12}) \\
& & + \sum_{b=1}^{m_1+m_n} (-1)^{m_n-b} C_+(m_1;m_n,b)\widehat{e}_{m_1+m_n-b}(s_{n1}+s_{12})\widehat{e}_b(s_{n1}) \\
& & + (-1)^{m_n} \sum_{a=1}^{m_1+m_n} C_-(m_1;m_n,a)\widehat{e}_{m_1+m_n-a}(s_{n1}+s_{12})\widehat{e}_a(s_{12})\,,\n\end{array}
$$

where *C*+, *C*− are some combinatorial constants. The second and third lines on the right hand side above give rise to a linear combination of forms, each of which factorizes into

non-constant form of chain type \times form of loop type.

The resulting regularized integral is then zero due to the factorization of integrals and the vanishing result on chain type forms. The regularized integral of the first line gives [\(3.17\)](#page-16-0) as desired.

⁴The same identity here can also be used to find the residue of the differential, but now one picks up the \hat{e}_1 terms instead of \hat{e}_0 terms due to Lemma [3.2.](#page-11-1)

In the calculations in Proposition [3.8,](#page-15-1) we need to use the iterated residue formula [\[LZ23\]](#page-29-4) for the iterated regularized integral. This requires computing the ¯*∂*-primitive[5](#page-17-1) of the integrand lying in the space V_D . Here we have done this using Lemma [2.9](#page-7-0) (ii) under the generators $\{\hat{e}_m\}_{m>1}$ for the ring $\hat{\mathfrak{F}}$. One can alternatively use the set of generators $\{\widehat{\mathcal{E}}_m^*\}_{m\geq 1}$, as is done in studying properties related to non-meromorphicity (such as holomorphic anomaly in [\[LZ23\]](#page-29-4)) in which the addition formula for the only non-holomorphic generator $\hat{\mathcal{E}}_1^*$ plays the role of Lemma [2.9](#page-7-0) (iii) here. The combinatorics involved in finding the ¯*∂*-primitive for these two sets of generators are essentially the same, as exhibited in Lemma [2.9](#page-7-0) (ii). However, for the latter set $\{\widehat{\mathcal{E}}_m^*\}_{m\geq 1}$, monomials in generators can have much more complicated residues due to the higher order pole structure present in the generators.

One encounters the same difficulty when considering iterated residues and iterated regularized integrals of elements in \mathcal{F}_D .

We demonstrate this through the following example.

Example 3.10. Consider $f = \hat{e}_1^r(z)$, $r \ge 1$. Using the $\bar{\partial}$ -primitive of integrands provided in Lemma [2.9](#page-7-0) (ii), one has

$$
\oint_E \tilde{e}_1^r(z) \text{vol} = \text{res}_{\partial} \left(\sum_{\ell \ge 0} (-1)^{\ell} \tilde{e}_1^{r-1-\ell}(z) \hat{e}_{\ell+2}(z) dz \right).
$$

This is not convenient for actual computations. We instead use the alternative ¯*∂*-primitive as in [\[LZ23\]](#page-29-4)

$$
\bar{\partial}\left(\frac{1}{r+1}\hat{e}_{1}^{r+1}(z)dz\right) = \bar{\partial}\hat{e}_{1}(z) \wedge \hat{e}_{1}^{r}(z)dz = 2\pi i d\bar{z} \wedge \hat{e}_{1}^{r}(z)dz = -2\pi i \hat{e}_{1}^{r}(z) \text{vol}
$$

to obtain

$$
\oint_E \tilde{e}_1^r(z) \text{vol} = \text{res}_{\partial} \left(\frac{1}{r+1} \tilde{e}_1^{r+1}(z) dz \right).
$$

Combining [\(3.10\)](#page-11-2), this then leads to a formula for the desired regularized integral. The result is zero when *r* is odd, as it should be the case by the parity reason explained in Remark [3.3.](#page-11-3)

 \triangle

3.3 Iterated *A***-cycle integrals and Chen's iterated path integrals**

We now discuss iterated A-cycle integrals for elements in V_D and their relations to Chen's iterated path integrals [\[Che77\]](#page-28-2).

3.3.1 Relation between iterated *A***-cycle integrals and iterated regularized integrals**

We first recall a few results obtained in [\[LZ21,](#page-29-3) [LZ23,](#page-29-4) [Zho23a\]](#page-30-1) on the relation between iterated *A*-cycle integrals and iterated regularized integrals.

First, by the same reasoning in [\[Zho23a,](#page-30-1) Lemma 4.12], when acting on elements in V_D , one has from the simple pole structure that

$$
\lim_{A=0} \int_{E_{[n]}} = \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}([n])} \int_{A_{\sigma([n])}} . \tag{3.18}
$$

 5 The result for the regularized integral is independent of the choice of the primitive by the global residue theorem, as can be seen from the residue formula for the regularized integral.

Hence $f_{E_{[n]}}$ can be obtained as the elliptic completion of iterated *A*-cycle integrals averaged over the orderings.

Conversely, denote

$$
R_0^i = \int_A dz_i, \quad R_j^i = \text{res}_{z_i = z_j} dz_i,
$$
 (3.19)

Let R be the vector space consisting of the operators R_0^i , $i \in [N]$, and Lie (R) be the corresponding free Lie algebra, satisfying

$$
[R_0^i, R_0^j] = R_0^j R_j^i. \tag{3.20}
$$

By the Poincaré-Birkhoof-Witt theorem, one has

$$
T^{\otimes} \mathcal{R} \cong \mathcal{U}(\text{Lie}(\mathcal{R})) \cong \text{Sym}^{\otimes} \text{Lie}(\mathcal{R}). \tag{3.21}
$$

This tells that [\[LZ21\]](#page-29-3) an iterated *A*-cycle integral can be expressed in terms of averaged iterated *A*-cycle integrals of residues. Applying [\(3.18\)](#page-17-2), it can in turn be expressed in terms of holomorphic limits of iterated regularized integrals of residues.

3.3.2 Chen's iterated path integrals and KZB connection

Following [\[CEE09\]](#page-28-6), let $t_{1,n}$ be the graded Lie algebra with generators $x_i, y_i, i \in [n]$ of degree 1 and t_{ii} , $i \neq j \in [n]$, subject to the relations

$$
t_{ij} = t_{ji}, \quad [t_{ij}, t_{ik} + t_{jk}] = 0, \quad [t_{ij}, t_{k\ell}] = 0,
$$

\n
$$
[x_i, y_j] = t_{ij}, \quad [x_i, x_j] = [y_i, y_j] = 0, \quad [x_i, y_i] = -\sum_{k \neq i} t_{ik},
$$

\n
$$
[x_i, t_{jk}] = [y_i, t_{jk}] = 0, \quad [x_i + x_j, t_{ij}] = [y_i + y_j, t_{ij}] = 0,
$$

where i, j, k, ℓ are distinct. The elements $\sum_{i=1}^{n} x_i$, $\sum_{i=1}^{n} y_i$ lie in the center of $t_{1,n}$ and one denotes by $\bar{t}_{1,n}$ the quotient of $t_{1,n}$ by the subspace generated by them.

Denote $ad_x = [x, -] \in Der(t_{1,n})$. The Knizhnik-Zamolodchikov-Bernard (KZB) connection is given [\[CEE09\]](#page-28-6) by $\nabla = d - \omega_{KZB}$, with

$$
\omega_{\text{KZB}} = \sum_{i=1}^{n} dz_i \otimes \left(-y_i + \sum_{j \neq i} (S_c(z_{ij}) - \frac{1}{c})|_{c = \text{ad}_{x_i}}(t_{ij}) \right), \quad z_{ij} := z_i - z_j
$$

=
$$
- \sum_{i=1}^{n} dz_i \otimes \sum_{i=1}^{n} y_i + \sum_{i=1}^{n} dz_i \otimes \left(\sum_{j \neq i} cS_c(z_{ij})|_{c = \text{ad}_{x_i}}(y_j) \right) \in A^1(\text{Conf}_n(E)) \otimes t_{1,n}.
$$
 (3.22)

This gives a flat connection on the principal bundle $P_{1,n}$ on Conf_n(*E*) whose Lie algebra is t_{1,*n*}. In fact, one has $d\omega_{KZB} = \omega_{KZB} \wedge \omega_{KZB} = 0$. This connection descends to flat connections on the $\text{principal bundle } \overline{\mathcal{P}}_{1,n} \to \text{Conf}_n(E) \text{ and } \overline{\mathcal{P}}_{1,n} \to \text{Conf}_n(E)/E \cong \text{Conf}_{n-1}(E^\star)$, see [\[CEE09\]](#page-28-6).

Chen's *π*¹ de Rham theorem tells that homotopy invariant integrals on the path spaces of *E* and *E* [⋆] are given by holonomies under the above connections, that is, by iterated path integrals of the corresponding connections. See [\[Che77,](#page-28-2) [LR07,](#page-29-11) [BL11\]](#page-28-3) for more details on this.

For simplicity, we consider the fiber F_n of the relative configuration space $Conf_{n+1}(E) \to Conf_n(E)$ which is isomorphic to $E - \{n \text{ points}\}\$. Let z_{n+1} be the coordinate along the fiber, and $z_j, j \in [n]$

coordinates on the base. The KZB connection then induces [\[CEE09\]](#page-28-6) a connection $d - \omega_{F_n}$ over a bundle \mathcal{P}_{F_n} on the fiber space F_n , with

$$
\omega_{F_n} = -dz_{n+1} \otimes \sum_{i=1}^{n+1} y_i + dz_{n+1} \otimes \sum_{j=1}^{n} c S_c (z_{n+1} - z_j)|_{c = ad_{x_{n+1}}} (y_j)
$$

\n
$$
= -dz_{n+1} \otimes \sum_{i=1}^{n+1} y_i + dz_{n+1} \otimes c S_c (z_{n+1}))|_{c = ad_{x_{n+1}}} (\sum_{i=1}^{n+1} y_i)
$$

\n
$$
-dz_{n+1} \otimes c S_c (z_{n+1})|_{c = ad_{x_{n+1}}} y_{n+1} + dz_{n+1} \otimes \sum_{j=1}^{n} (c S_c (z_{n+1} - z_j) - c S_c (z_{n+1}))|_{c = ad_{x_{n+1}}} (y_j).
$$

Applying the gauge transformation $e^{Ax_{n+1}}$ changes ω_{F_n} to

$$
\hat{\omega}_{F_n} = 2\pi i \beta_{n+1} \otimes x_{n+1} + e^{A \Delta x_{n+1}} \omega_{F_n}
$$
\n
$$
= -dz_{n+1} \otimes e^{A \Delta x_{n+1}} \left(\sum_{i=1}^{n+1} y_j \right) + 2\pi i \beta_{n+1} \otimes x_{n+1} + dz_{n+1} \otimes \sum_{j=1}^n c \widehat{S}_c (z_{n+1} - z_j)|_{c = \Delta x_{n+1}} (y_j).
$$
\n(3.23)

Note the induced flat connection valued in $\bar{t}_{1,n}$ is exactly the one given in [\[BL11\]](#page-28-3).

Consider now Chen's iterated path integrals of *^ω*b*Fⁿ* which then give rise to homotopy invariant integrals of the path space of F_n . Recall that for a piecewise smooth path $\gamma : [0,1] \to M$ and 1-forms ω_k , $k = 1, 2, \cdots, N$, the iterated path integral is defined by [\[Che77\]](#page-28-2)

$$
\int_{\gamma} \omega_N \cdots \omega_1 := \int_{0 \le t_1 \le t_2 \le \cdots \le t_N \le 1} \gamma^* \omega_N(t_N) \cdots \gamma^* \omega_1(t_1).
$$
\n(3.24)

We now take $M = F_n$ and $\omega_k = \hat{\omega}_{F_n}, k = 1, 2, \cdots, N$. This leads to

$$
\int_{\gamma} \widehat{\omega}_{F_n} \cdots \widehat{\omega}_{F_n} = \int_{0 \le t_1 \le t_2 \le \cdots \le t_N \le 1} \gamma^* \widehat{\omega}_{F_n}(t_N) \cdots \gamma^* \widehat{\omega}_{F_n}(t_1) \in F^N \mathcal{U}(\mathfrak{k}_{1,n}), \tag{3.25}
$$

where F^{\bullet} is the increasing filtration on $\mathcal{U}(\bar{t}_{1,n})$ defined by the length of words. Using [\(3.23\)](#page-19-0) and Definition [2.3,](#page-5-3) we write

$$
\gamma^*\widehat{\omega}_{F_n}(t)=2\pi i\beta(t)\otimes x_{n+1}+\sum_{j=1}^n\sum_{m\geq 0}\widehat{e}_m(t-z_j)dt\otimes \mathrm{ad}^m_{x_{n+1}}(y_j)\in A^1(F_n)\otimes \overline{\mathfrak{t}}_{1,n}.
$$

The iterated path integral [\(3.25\)](#page-19-1) then gets simplified into

$$
\int_{\gamma} \widehat{\omega}_{F_n} \cdots \widehat{\omega}_{F_n} = \sum_{m_k \geq 0} \sum_{j_k \geq 0} \int_{\gamma} \prod_{k=1}^N \widehat{e}_{m_k} (t_k - z_{j_k}) dt_k \bigotimes \otimes_{k=1}^N \mathrm{ad}_{x_{n+1}}^{m_k} (y_{j_k}) \in \mathcal{U}(\mathfrak{k}_{1,n}), \tag{3.26}
$$

where we have used the convention that $\hat{e}_{m_k}(t_k - z_{j_k})dt = \delta_{m_k,0}2\pi i\beta$, $ad_{x_{n+1}}^{m_k}(y_{j_k}) = \delta_{m_k,0}x_{n+1}$ if $j_k = 0$. It follows from Chen's π_1 de Rham theorem that the coefficients of various words in the expansion [\(3.26\)](#page-19-2) are homotopy invariant in *γ*.

We have the following result regarding these coefficients.

Lemma 3.11. *Let the notation be as above. Take γ to be a (non-unique) A-cycle class on E avoiding the* pole divisor of $\widehat{\omega}_{F_n}$ and consider the image of $\int_{\gamma} \widehat{\omega}_{F_n} \cdots \widehat{\omega}_{F_n}$ in $gr_F^N U(\overline{t}_{1,n})$. Then the coefficient of the image of $\otimes_{k=1}^N$ ad $^{m_k}_{x_{n+1}}(y_{j_k})$ is given by the iterated A-cycle integral $\int_{A_{[N]}}\prod_{k=1}^N\widehat{e}_{m_k}(t_k-z_{j_k})dt_k$. Furthermore, the *result is independent of the ordering in the iterated A-cycle integral, that is,*

$$
\int_{A_{[N]}} \prod_{k=1}^N \widehat{e}_{m_k}(t_k-z_{j_k})dt_k = \int_{A_{\sigma([N])}} \prod_{k=1}^N \widehat{e}_{m_k}(t_k-z_{j_k})dt_k, \quad \sigma \in \mathfrak{S}_{[N]}.
$$

Proof. A cycle γ on *E* avoiding the pole divisor of $\hat{\omega}_{F_n}$ gives rise to a cycle on F_n . According to [\(3.26\)](#page-19-2), if we take the path γ be the *A*-cycle, integrals involving $j_k = 0$ terms are eliminated since $\gamma^* \beta = 0$.

Let us focus on the image of $\int_{\gamma} \hat{\omega}_{F_n} \cdots \hat{\omega}_{F_n}$ in $gr_F^N \mathcal{U}(\bar{t}_{1,n})$, we can pretend that $XY = YX$ for $X, Y \in \bar{t}_{1,n}$ since $XY - YX = 0 \in \operatorname{gr}^N_F \mathcal{U}(\bar{t}_{1,n})$. The coefficient of the image of $\otimes_{k=1}^N \operatorname{ad}^{m_k}_{x_{n+1}}(y_{j_k})$ in $gr_F^N\mathcal{U}(\bar{\mathfrak{t}}_{1,n})$ then becomes

$$
\sum_{\sigma \in \mathfrak{S}_N} \int_{0 \le t_1 \le t_2 \le \dots \le t_N \le 1} \prod_{k=1}^N \widehat{e}_{m_{\sigma(k)}}(t_k - z_{j_{\sigma(k)}}) dt_k
$$
\n
$$
= \int_{[0,1]^N} \prod_{k=1}^N \widehat{e}_{m_k}(t_k - z_{j_k}) dt_k
$$
\n
$$
= \int_{A_{[N]}} \prod_{k=1}^N \widehat{e}_{m_k}(t_k - z_{j_k}) dt_k.
$$

That the result is independent of the ordering of integration in $\int_{A_{[N]}}$ follows from [\(3.20\)](#page-18-0) and [\(3.2\)](#page-11-1). \Box

Applying the discussions in Section [3.3.1,](#page-17-3) the iterated *A*-cycle integrals in Lemma [3.11](#page-19-3) above can be further expressed in terms of iterated regularized integrals computed in Section [3.2.](#page-13-0)

We can also take the path *γ* to be the *B*-cycle, the same discussions lead to iterated *B*-cycle integrals. They are related to iterated *A*-cycle integrals in a nice way. Indeed, by the reciprocity law applied to $\hat{e}_m(z)dz, dz$, one has (using Lemma [3.2\)](#page-11-1)

$$
\int_B \widehat{e}_m(z)dz = \tau \int_A \widehat{e}_m(z)dz + \operatorname{res}_{\partial}(z\widehat{e}_m(z)dz) = \tau \int_A \widehat{e}_m(z)dz.
$$

It might be interesting to find the image in $gr_F^{\bullet}U(\bar{t}_{1,n})$ besides the top component above. It would also be interesting to find realizations of more general iterated *A*-cycle integrals considered in Section [3.2](#page-13-0) in terms of Chen's iterated path integrals, up to a reparametrization of the form $t_k \mapsto t_k + z_k$, $k \in [N]$. This would then imply the GW generating series of elliptic curves, discussed in Section [4](#page-20-0) below, are in fact linear combinations of Chen's iterated path integrals.

4 Gromov-Witten generating series of elliptic curves

4.1 Proof of the main Theorem

Using the notation and conventions in earlier sections (e.g., Definition [2.5,](#page-5-4) Convention [\(3.7\)](#page-10-1)), Theorem [1.1](#page-1-2) is recalled as follows.

Theorem 4.1. *Let*

$$
\varpi_n = \frac{\theta(\sum_{i=1}^n w_i)}{\prod_{i=1}^n \theta(w_i)} \cdot \frac{\prod_{i < j} \theta(z_i + w_i - z_j - w_j) \theta(z_i - z_j)}{\prod_{i < j} \theta(z_i + w_i - z_j) \theta(z_i - w_j - z_j)} \,. \tag{4.1}
$$

Then the regularized integral $\int_{E_{[n]}} \varpi_n$ *satisfies*

$$
\oint_{E_{[n]}} \varpi_n = \sum_{j \in [n]} \sum_{\pi \in \Pi_{[n] \setminus \{j\}} \atop k=1} \prod_{k=1}^\ell (|\pi_k| - 1)! \cdot \widehat{e}_{|\pi_k|} (\sum_{i \in \pi_k} w_i).
$$
\n(4.2)

From [\(2.18\)](#page-6-3), one can check that the function ω_n is indeed elliptic in both z_i , w_i , where the z_i 's are regarded as variables on *E*[*n*] , while the *wⁱ* 's as nonzero parameters. As before we use the additive coordinates z_i , w_i and the corresponding multiplicative coordinates $u_i = e^{2\pi i w_i}$, $v_i = e^{2\pi i z_i}$ interchangeably.

Denote the zero and polar divisor of ϖ_n by D_0 , D_∞ respectively. For generic values of the parameters w_i 's, the divisor D_∞ is a smooth hypersurface arrangement divisor (in fact a simple normal crossing divisor). In particular, this fits in the setting of cohomological regularized integral in [\[Zho23b\]](#page-30-2).

In what follows, we shall evaluate $\int_{E_{[n]}} \varpi_n$ using the results established in Section [3.](#page-9-0) To reduce the difficulty caused by the complexity of the polar divisor D_{∞} , we first need to split ω_n into a sum of forms with simpler polar divisors in a way similar to the one provided by Lemma [2.9](#page-7-0) (iii). This is possible thanks to the following version of Fay's multi-secant identity.

Lemma 4.2. *Let the notation be as above. Then one has*

$$
\varpi_n = \det \begin{pmatrix} 0 & 1 & \cdots & 1 \\ -1 & \frac{\partial}{\partial} \left(\frac{u_1 v_1}{v_1} \right) & \cdots & \frac{\partial}{\partial} \left(\frac{u_1 v_1}{v_n} \right) \\ \vdots & \vdots & \frac{\partial}{\partial} \left(\frac{u_i v_i}{v_j} \right) & \vdots \\ -1 & \frac{\partial}{\partial} \left(\frac{u_n v_n}{v_1} \right) & \cdots & \frac{\partial}{\partial} \left(\frac{u_n v_n}{v_n} \right) \end{pmatrix}
$$
(4.3)

Proof. The Frobenius-Stickelberger formula/Fay's multi-secant identity gives (see e.g., [\[Zho23a,](#page-30-1) Lemma 3.3])

$$
\varpi_n = \det \begin{pmatrix} 0 & 1 & \cdots & 1 \\ -1 & \frac{\theta'}{\theta}(\frac{u_1v_1}{v_1}) & \cdots & \frac{\theta'}{\theta}(\frac{u_1v_1}{v_n}) \\ \vdots & \vdots & \cdots & \vdots \\ -1 & \frac{\theta'}{\theta}(\frac{u_nv_n}{v_1}) & \cdots & \frac{\theta'}{\theta}(\frac{u_nv_n}{v_n}) \end{pmatrix}
$$
(4.4)

The right hand side in [\(4.3\)](#page-21-0) is almost-elliptic in z_i , w_i , whose holomorphic limit gives the right hand side of [\(4.4\)](#page-21-1) which is already almost-elliptic in z_i , w_i since ω_n is so.^{[6](#page-21-2)} Therefore, they are identical.

 \Box

Note that although the entire determinant in [\(4.4\)](#page-21-1) above have trivial automorphy in the z_i , w_i 's, the individual summands in its expansion are not so. This is why we passed from $e_1 = \theta' / \theta$ to its elliptic completion $\hat{e}_1 = \hat{\theta}'/\hat{\theta}$, and later on work with the expansion of [\(4.3\)](#page-21-0). The entry $\hat{\theta}'(\frac{u_i v_i}{v_j})$ $\frac{u_iv_i}{v_j}$)/ $\widehat{\theta}(\frac{u_iv_i}{v_j})$ $\frac{\partial_i v_i}{\partial_j})$ is recognized as giving the angular form [\[Zho23b\]](#page-30-2)

$$
\xi_{ij} = \partial_{z_i} \log |\theta(s_{ij})|^2 \cdot dz_i = \hat{e}_1(s_{ij}) dz_i
$$

associated to the divisor $(s_{ij} = 0)$, now with

$$
s_{ij} := w_i + z_i - z_j. \tag{4.5}
$$

Here $| - |^2$ is the standard Hermitian metric on the line bundle $\mathcal{O}_E([0])$, with $|\theta|^2 = \hat{\theta}\hat{\theta}$.

 6 One can also check the almost-ellipticity of the right hand side of [\(4.4\)](#page-21-1) directly, using the relations in Definition [2.1,](#page-4-5) the automorphy behavior [\(2.19\)](#page-6-1), and the linearity of determinant in row and column vectors.

Proof of Theorem [4.1.](#page-20-2) Denote the matrix in Lemma [4.2](#page-21-3) by Z whose rows and columns we label by {0} ∪ [*n*]. By the expansion of determinant, one has

$$
\varpi_n = \det \mathcal{Z} = \sum_{\rho \in \mathfrak{S}_{\{0\} \cup [n]}} (-1)^{\text{sign}(\rho)} \prod_{k \in \{0\} \cup [n]} \mathcal{Z}_{k\rho(k)}.
$$
\n(4.6)

Denote

$$
\xi_{\rho} = (-1)^{\text{sign}(\rho)} \prod_{k \in \{0\} \cup [n]} \mathcal{Z}_{k\rho(k)} \cdot \wedge_{k=1}^{n} \text{vol}_{k} . \tag{4.7}
$$

Representing any permutation *ρ* by its standard cycle decomposition, this leads to factorization of the corresponding iterated regularized integral $f_{E_{[n]}}$ *ξ*_{*ρ*}. Focus on the cycle that contains 0. It is either the trivial one (0), or one of the form $(0j_0j_1 \cdots j_{r-1}j_r)$ with length $r + 2, r \ge 0$. For the former, the contribution to $f_{E_{[n]}}\,\zeta_\rho$ gives zero since $\mathcal{Z}_{00}=0$ in [\(4.7\)](#page-22-1). For the latter, it gives

$$
(-1)^{r+2-1}\mathcal{Z}_{0j_0}\mathcal{Z}_{j_0j_1}\cdots\mathcal{Z}_{j_{r-1}j_r}\mathcal{Z}_{j_r,j_{r+1}=0}=(-1)^{r+1-1}\mathcal{Z}_{j_0j_1}\cdots\mathcal{Z}_{j_{r-1}j_r}.
$$
\n(4.8)

If $j_r \neq j_0$, then the above cycle $(0j_0 j_1 \cdots j_{r-1} j_r)$ gives a form of the type ξ_{chain} and thus the corresponding regularized integral − R *E*[*n*] *ξ^ρ* is zero by Proposition [3.8](#page-15-1) (ii). It follows that the nontrivial contributions $f_{E_{[n]}}$ *ξ_ρ* to the regularized integral $f_{E_{[n]}}$ *ω*_{*n*} arise from those *ρ* whose cycle containing 0 in the standard cycle decomposition takes the form $(0j_0j_1 \cdots j_{r-1}j_0)$. That is, one has

$$
\oint_{E_{[n]}} \varpi_n = \oint_{E_{[n]}} \sum_{j_0=1}^n \sum_{\sigma \in \mathfrak{S}_{[n]-\{j_0\}} (-1)^{\sigma} \mathcal{Z}_{1\sigma_1} \mathcal{Z}_{2\sigma_2} \cdots \check{\mathcal{Z}}_{j_0j_0} \cdots \mathcal{Z}_{n\sigma_n} \cdot \wedge_{k=1}^n \mathrm{vol}_k,
$$
\n(4.9)

where $\check{\mathcal{Z}}_{j_0j_0}$ stands for the omission of $\mathcal{Z}_{j_0j_0}$ due to [\(4.8\)](#page-22-2).

To compute [\(4.9\)](#page-22-3), again we use the standard cycle decomposition $\sigma = \sigma_1 \cdots \sigma_\ell$ for any permutation *σ* ∈ S[*n*]−{*j*0} . Observe that the iterated regularized integral corresponding to a cycle *σ^k* only depends on its underlying set π_k , by Proposition [3.8](#page-15-1) (ii) and the structure of the polar divisor given in [\(4.5\)](#page-21-4), with

$$
\sum_{i \in \pi_k} s_{i,\sigma_i} = \sum_{i \in \pi_k} w_i \,. \tag{4.10}
$$

 \Box

It follows that the sum in [\(4.9\)](#page-22-3) becomes a double sum, with the outer sum being a sum over partitions $\pi = \{\pi_1, \cdots, \pi_\ell\} \in \Pi_{[n] - \{j_0\}}$, and the inner sum over permutations within the blocks of the partitions. The number of permutations $\sigma \in \mathfrak{S}_{[n]-\{j_0\}}$ whose standard cycle decompositions share the same underlying partition $\pi \in \Pi_{[n]-\{j_0\}}$ is $\prod_{k=1}^{\ell}(|\pi_k|-1)!$. Since the sign contributing to $(-1)^{\sigma}$ of each block π_k is $(-1)^{|\pi_k|-1}$, applying Proposition [3.8](#page-15-1) (ii) to each such block simplifies [\(4.9\)](#page-22-3) into

$$
\int_{E_{[n]}} \varpi_n = \sum_{j_0=1}^n \sum_{\pi \in \Pi_{[n] - \{j_0\}}}\prod_{k=1}^\ell (-1)^{|\pi_k|-1} \cdot (|\pi_k|-1)! \cdot \widehat{e}_{|\pi_k|} (\sum_{i \in \pi_k} s_{i, \sigma_i}) \cdot \left(\prod_{i \in \pi_k} \delta_{1,0} - \prod_{i \in \pi_k} (\delta_{1,0}-1)\right).
$$

Simplifying further using [\(4.10\)](#page-22-4), one obtains the desired claim.

4.2 Generating series of *n***-point functions**

The results we have established in previous sections, such as Lemma [4.2,](#page-21-3) provide new persepectives in studying finer combinatorial properties of the GW generating series. In the rest of the work,

we initiate these studies by working with the generating series of $\{\widehat{T}_n\}_{n\geq 1}$. We hope to connect our results to constructions on infinite Grassmannians and integrable hierarchies along these lines in further investigations.

We first introduce some notation.

Definition 4.3. Let the notation be as before. Denote

$$
\widehat{Z}_n = \begin{pmatrix}\n\frac{\widehat{\theta}'}{\widehat{\theta}}\left(\frac{u_1v_1}{v_1}\right) & \cdots & \frac{\widehat{\theta}'}{\widehat{\theta}}\left(\frac{u_1v_1}{v_n}\right) \\
\vdots & \ddots & \vdots \\
\frac{\widehat{\theta}'}{\widehat{\theta}}\left(\frac{u_nv_n}{v_1}\right) & \cdots & \frac{\widehat{\theta}'}{\widehat{\theta}}\left(\frac{u_nv_n}{v_n}\right)\n\end{pmatrix}, \quad \delta_n = \begin{pmatrix}\n\varepsilon_1 \text{vol}_1 & \cdots & 0 \\
\vdots & \ddots & \vdots \\
0 & \cdots & \varepsilon_n \text{vol}_n\n\end{pmatrix},
$$
\n(4.11)\n
\n
$$
\widehat{Z}_n = \begin{pmatrix}\n0 & 1 & \cdots & 1 \\
-1 & \frac{\widehat{\theta}}{\widehat{\theta}}\left(\frac{u_1v_1}{v_1}\right) & \cdots & \frac{\widehat{\theta}}{\widehat{\theta}}\left(\frac{u_1v_1}{v_n}\right) \\
\vdots & \vdots & \ddots & \vdots \\
-1 & \frac{\widehat{\theta}}{\widehat{\theta}}\left(\frac{u_nv_n}{v_1}\right) & \cdots & \frac{\widehat{\theta}}{\widehat{\theta}}\left(\frac{u_nv_n}{v_n}\right)\n\end{pmatrix}, \quad \Delta_n = \begin{pmatrix}\n\varepsilon_0 & 0 & \cdots & 0 \\
0 & \varepsilon_1 \text{vol}_1 & \cdots & 0 \\
\vdots & \vdots & \ddots & \vdots \\
0 & 0 & \cdots & \varepsilon_n \text{vol}_n\n\end{pmatrix},
$$

where ε_0 , ε_k , $1 \leq k \leq n$ are indeterminants. If needed, we could add in an auxiliary curve E_0 and replace ε_0 by ε_0 vol₀ correspondingly. Similarly, we define Z_n , Z_n to be the corresponding holomorphic limit versions obtained by replacing $\hat{\theta}$ by θ everywhere.

By Lemma [4.2](#page-21-3) and results in Section [4.1,](#page-20-1) we have

$$
\det \mathcal{Z}_n = \det \widehat{\mathcal{Z}}_n, \quad \prod_{k=0}^n \varepsilon_k \cdot \widehat{T}_{[n]} = \int_{E_{[n]}} \det \left(\Delta_n \widehat{\mathcal{Z}}_n \right). \tag{4.12}
$$

Constructions using the matrix Z_n lead to results parallel to those obtained using \hat{Z}_n . In fact, many properties are easier to understand by putting them together.

Before proceeding, we need more definitions.

Definition 4.4. Let the notation and conventions be as above. Define

$$
\prod_{k=1}^{n} \varepsilon_k \cdot \widehat{G}_n := \int_{E_{[n]}} \det \left(\delta_n \widehat{Z}_n \right).
$$
\n(4.13)

We extend the definition of \widehat{T}_n , \widehat{G}_n to any set $S \subseteq \mathbb{N}_+$, and set by convention

$$
\widehat{T}_{\emptyset} = \widehat{T}_{0} = 1, \quad \widehat{G}_{\emptyset} = \widehat{G}_{0} = 1.
$$
\n(4.14)

Lemma 4.5. *Let the notation and conventions be as above. Then one has*

$$
\widehat{G}_{[n]} = \sum_{\pi \in \Pi_{[n]}} \frac{\widehat{B}_{\pi}}{|\pi|},\tag{4.15}
$$

where the right hand side is defined as in [\(1.3\)](#page-1-1) *of Theorem [1.1.](#page-1-2)*

Proof. This follows by repeating the proof of Theorem [4.1.](#page-20-2)

Note that unlike [\(4.12\)](#page-23-0), one has det $Z_n \neq$ det \hat{Z}_n , and only the regularized integral of det \hat{Z}_n makes sense since det Z_n does not define a function on $E_{[n]}$ due to its quasi-ellipticity.

 \Box

Definition 4.6. Define

$$
\mathcal{F}_{\mathsf{D}}\left[\left[\varepsilon_0,\varepsilon_1,\cdots,\varepsilon_n,\cdots\right]\right]=\lim_{\leftarrow}\mathcal{F}_{\mathsf{D}}\left[\left[\varepsilon_0,\varepsilon_1,\cdots,\varepsilon_n\right]\right],\tag{4.16}
$$

where the morphisms in the projective system are given by

$$
\mathcal{F}_{\mathsf{D}}[[\varepsilon_0,\varepsilon_1,\cdots,\varepsilon_n,\varepsilon_{n+1}]]\to \mathcal{F}_{\mathsf{D}}[[\varepsilon_0,\varepsilon_1,\cdots,\varepsilon_n]],\quad \varepsilon_k\mapsto\varepsilon_k, k\leq n, \varepsilon_{n+1}\mapsto 0.
$$

It follows that

$$
\mathcal{F}_{\mathsf{D}}[[\varepsilon_0,\varepsilon_1,\cdots,\varepsilon_n,\cdots]] = \lim_{\leftarrow} \left(\mathcal{F}_{\mathsf{D}}[\varepsilon_0,\varepsilon_1,\cdots,\varepsilon_n]/(\varepsilon_0,\varepsilon_1,\cdots,\varepsilon_n)^{n+1} \right).
$$

Let $\mathcal J$ be the ideal of $\mathcal F_{\mathsf{D}}[[\varepsilon_0,\varepsilon_1,\cdots,\varepsilon_n,\cdots]]$ generated by $\varepsilon_1^2,\cdots,\varepsilon_n^2,\cdots$.

Definition 4.7. Let $\mathbb{1}_{\infty}$, $\hat{\mathbb{Z}}_{\infty}$, $\hat{\mathbb{Z}}_{\infty}$, Δ_{∞} , δ_{∞} be the limit matrices of $\{\mathrm{id}_n\}_{n\geq 1}$, $\{\hat{\mathbb{Z}}_n\}_{n\geq 1}$, $\{\hat{\mathbb{Z}}_n\}_{n\geq 1}$, $\{\Delta_n\}_{n\geq 1}$, ${\{\delta_n\}_{n\geq 1}}$, respectively. Define

$$
\det\left(\mathbb{1}_{\infty} + \Delta_{\infty}\widehat{\mathcal{Z}}_{\infty}\right), \quad \det\left(\mathbb{1}_{\infty} + \delta_{\infty}\widehat{Z}_{\infty}\right) \in \mathcal{F}_{\mathsf{D}}\left[\left[\varepsilon_{0},\varepsilon_{1},\cdots,\varepsilon_{n},\cdots\right]\right]
$$
(4.17)

to be the summation of the principal minors of $\Delta_{\infty}\widehat{\mathcal{Z}}_{\infty}$, $\delta_{\infty}\widehat{Z}_{\infty}$, respectively.

Let $\varepsilon_S := \prod_{k \in S} \varepsilon_k$. Define

$$
\widehat{\mathcal{T}}_{[n]}(\varepsilon) = \sum_{S \subseteq [n]} \varepsilon_S \widehat{T}_S = 1 + \varepsilon_0 (\varepsilon_1 + \varepsilon_2 + \dots + \varepsilon_n) + \dots, \quad \widehat{\mathcal{G}}_{[n]}(\varepsilon) = \sum_{S \subseteq [n]} \varepsilon_S \widehat{G}_S = 1 + \dots, \quad (4.18)
$$

and correspondingly the following elements in the ring $\mathcal{F}_{D}[[\varepsilon_{0}, \varepsilon_{1}, \cdots, \varepsilon_{n}, \cdots]]$

$$
\widehat{\mathcal{T}}(\varepsilon) = \sum_{S \subseteq \mathbb{N}_+} \varepsilon_S \widehat{T}_S = 1 + \varepsilon_0 (\varepsilon_1 + \varepsilon_2 + \dots) + \dots, \quad \widehat{\mathcal{G}}(\varepsilon) = \sum_{S \subseteq \mathbb{N}_+} \varepsilon_S \widehat{G}_S = 1 + \dots \tag{4.19}
$$

Proposition 4.8. *Let the notation and conventions be as above.*

i). One has

$$
\widehat{\mathcal{G}}(\varepsilon) = \int_{E_{\infty}} \det \left(\mathbb{1}_{\infty} + \delta_{\infty} \widehat{Z}_{\infty} \right),
$$

 ν here \int_{E_∞} is understood to be the identity operator when acting on a constant form.

ii). One has

$$
\widehat{\mathcal{T}}(\varepsilon)-1-\varepsilon_0(\varepsilon_1+\varepsilon_2+\cdots)=(\widehat{\mathcal{T}}(\varepsilon)-1)(\widehat{\mathcal{G}}(\varepsilon)-1) \mod \mathcal{J}.
$$

iii). Define

$$
\widehat{\mathcal{H}}(\varepsilon) = \int_{E_{\infty}} \det \left(\mathbb{1}_{\infty} + \Delta_{\infty} \widehat{\mathcal{Z}}_{\infty} \right).
$$

Then one has

$$
\widehat{\mathcal{H}}(\varepsilon)=\varepsilon_0\cdot(\widehat{\mathcal{T}}(\varepsilon)-1)+\widehat{\mathcal{G}}(\varepsilon).
$$

Proof. i). This follows immediately from Definition [4.7](#page-24-0) and the proof of Theorem [4.1.](#page-20-2)

ii). Consider a square matrix *M* whose rows and columns are labelled by $S \subseteq \mathbb{N}$. For any subset $I \subseteq S$, let M_I be the submatrix of M formed by collecting the rows and columns with labels from *I*. For a subset $I \subseteq S$, set $I^c = S \setminus I$.

By the proof of Theorem [4.1,](#page-20-2) we have

$$
\int_{E_{[n]}} \det(\Delta_n \widehat{Z}_n) = \int_{E_{[n]} \oslash \subsetneq I \subsetneq [n]} \det(\Delta_I \widehat{Z}_I) \cdot \det(\delta_{I^c} \widehat{Z}_{I^c})
$$
\n
$$
= \sum_{\oslash \subsetneq I \subsetneq [n]} \oint_{E_I} \det(\Delta_I \widehat{Z}_I) \cdot \oint_{E_{I^c}} \det(\delta_{I^c} \widehat{Z}_{I^c}), \quad n \ge 2.
$$

Summing over *n* leads to the desired claim, where the need of modulo J arises from the relation $I \cap I^c = \emptyset$.

iii). By Definition [4.7,](#page-24-0) we have

$$
\epsilon_0^1 \cdot [\epsilon_0^1] \widehat{\mathcal{H}}(\epsilon) = \widehat{\mathcal{T}}(\epsilon) - 1, \quad [\epsilon_0^0] \widehat{\mathcal{H}}(\epsilon) = \int_{E_{\infty}} \det \left(\mathbb{1}_{\infty} + \delta_{\infty} \widehat{Z}_{\infty} \right) = \widehat{\mathcal{G}}(\epsilon).
$$

Combining these, one obtains the desired claim.

 \Box

4.3 Deformation in Grassmannian

Proposition [4.8](#page-24-1) above tells that any of $\hat{\tau}$, \hat{G} , $\hat{\mathcal{H}}$ determine the rest two. Therefore, it suffices to study the properties of one of them. In this work we prefer to work with \hat{G} as [\(4.15\)](#page-23-1) in Lemma [4.5](#page-23-2) seems to be simpler than [\(1.3\)](#page-1-1) in Theorem [1.1.](#page-1-2) Furthermore, \hat{G} admits a natural geometric formulation similar to the *τ*-function in the study of KP hierarchy [\[DJKM81,](#page-29-12) [DJKM83,](#page-29-13) [SW85\]](#page-30-8), as we now explain.

Let *J* be the Jacobian of *E* and Pic⁽⁰⁾(*J*) \cong *J* be its Picard variety. Denote by *m* the addition map

$$
m: J \times Pic^{(0)}(J) \to J.
$$

The Poincaré bundle on $J \times \text{Pic}^{(0)}(J)$ satisfies

$$
\mathcal{P} \cong \mathcal{O}_{J \times \text{Pic}^{(0)}(J)}(-[J \times 0] - [0 \times \text{Pic}^{(0)}(J)] + [\text{ker } m]). \tag{4.20}
$$

We consider constructions on $\mathcal{P}(J \times 0)|_{2(I \times 0)}$ which fits in the decomposition sequence

$$
0 \to \mathcal{P}|_{J\times 0} \to \mathcal{P}(J\times 0)|_{2(J\times 0)} \to \mathcal{P}(J\times 0)|_{J\times 0} \to 0,
$$
\n(4.21)

where

$$
\mathcal{P}|_{J\times 0}\cong \check{N}_{J\times 0}\cong K_{J\times 0},\quad \mathcal{P}(J\times 0)|_{J\times 0}\cong \mathcal{O}_{J\times 0}.
$$
\n(4.22)

Lemma 4.9. *One has*

$$
H^{0}(J \times 0, \mathcal{P}(J \times 0)|_{2(J \times 0)}) = \text{span}\left\{f_{0}, f_{-1} + \frac{\partial_{z}\theta(z)}{\theta(z)}f_{0}\right\},\tag{4.23}
$$

*where f*₀, *f*_{−1} are certain trivializations (detailed below) of $\mathcal{P}|_{J\times 0}$, $\mathcal{P}(J\times 0)|_{J\times 0}$, respectively.

Proof. The long exact sequence in cohomology associated to [\(4.21\)](#page-25-1) gives

$$
0 \to H^0(J \times 0, \mathcal{P}|_{J \times 0}) \to H^0(J \times 0, \mathcal{P}(J \times 0)|_{2(J \times 0)}) \to H^0(J \times 0, \mathcal{P}(J \times 0)|_{J \times 0}) \to \cdots
$$

From [\(4.22\)](#page-25-2), one has

$$
H^{0}(J \times 0, \mathcal{P}|_{J \times 0}) \subseteq H^{0}(J \times 0, \mathcal{P}(J \times 0)|_{2(J \times 0)}), \quad \dim H^{0}(J \times 0, \mathcal{P}(J \times 0)|_{2(J \times 0)}) \leq 2. \tag{4.24}
$$

Recall from [\[Rai89\]](#page-29-5) (see also [\[Zho23a\]](#page-30-1) in the current notation) that when $c \neq 0$ mod Λ_{τ} , one has

$$
H0(J \times Pic(0)(J), P(J \times 0)) = \text{span}\left\{S_c(z)\right\}.
$$
 (4.25)

Under the restriction map $\mathcal{P}(J\times 0)\to \mathcal{P}(J\times 0)|_{2(J\times 0)}$, the image of $S_c(z)$ in $H^0(J\times 0, \mathcal{P}(J\times 0)|_{2(J\times 0)})$ is represented by the first two terms of its the Laurent expansion in *c*

$$
c^{-1} + c^0 \frac{\partial_z \theta(z)}{\theta(z)},
$$
\n(4.26)

where c^{-1} , c^0 are trivializations of $\mathcal{P}(J \times 0)|_{J \times 0} \cong \mathcal{O}_{J \times 0}$, $\mathcal{P}|_{J \times 0} \cong K_{J \times 0}$ respectively. Setting $f_{-1} =$ c^{-1} , $f_0 = c^0$, the desired relation [\(4.23\)](#page-25-3) then follows from [\(4.24\)](#page-26-0) and [\(4.26\)](#page-26-1).

 \Box

Note that the c^{-1} -term of $S_c(z)$ is given by the residue along the map $\mathcal{P}(J \times 0) \to \mathcal{P}(J \times 0)|_{J \times 0}$ and is thus independent of the choice of local defining equation for $J \times 0$ inside $J \times Pic^{(0)}(J)$. The quasi-ellipticity of *θ* ′/*θ* given in [\(2.19\)](#page-6-1) and [\(4.26\)](#page-26-1) yield the automorphy behavior

$$
\left(f_{-1} \quad f_0\right) \mapsto \left(f_{-1} \quad f_0\right) \begin{pmatrix} 1 & 0 \\ 2\pi i & 1 \end{pmatrix} . \tag{4.27}
$$

From this we see the non-splitness of the decomposition sequence [\(4.21\)](#page-25-1). Furthermore, from [\(4.27\)](#page-26-2) it follows that the real-analytic trivializations

$$
\widehat{f}_{-1} := f_{-1} - A(z)f_0, \quad \widehat{f}_0 := f_0 \tag{4.28}
$$

are almost-elliptic, with the sections in [\(4.23\)](#page-25-3) now expressed as

$$
\widehat{f}_0, \quad \widehat{f}_{-1} + \frac{\widehat{\theta}'}{\widehat{\theta}}(z) \cdot \widehat{f}_0. \tag{4.29}
$$

For convenience, in what follows we replace $\mathcal{P}(J\times 0)$ by $\mathcal{L}(0\times \mathsf{Pic}^{(0)}(J))$, with

$$
\mathcal{L} := \mathcal{O}_{J \times \text{Pic}^{(0)}(J)}([\ker m] - [0 \times \text{Pic}^{(0)}(J)]). \tag{4.30}
$$

Working with the section $\frac{\theta(z+c)}{\theta(z)}$ of $\mathcal{L}(0 \times Pic^{(0)}(J))$, the frame [\(4.23\)](#page-25-3) then changes accordingly to

$$
c^1, \quad c^0 + \frac{c^1 \partial_z \theta(z)}{\theta(z)}, \tag{4.31}
$$

where c^0 , c^1 are trivializations of $\mathcal{L}(0 \times \text{Pic}^{(0)}(J))|_{J \times 0} \cong \mathcal{O}_{J \times 0}(0 \times 0)$, $\mathcal{L}(0 \times \text{Pic}^{(0)}(J))|_{J \times 0} \otimes \text{N}|_{J \times 0}$, respectively.

In what follows we also make the following identifications without explicit mentioning

$$
J \times 0 \cong J, \quad \mathcal{O}_{J \times 0} \cong \mathcal{O}_J \cong K_J, \quad \check{N}|_{J \times 0} \cong K_J. \tag{4.32}
$$

We now consider the map

$$
s_{ij}: E_i \times E_{\bar{j}} \to J_{i\bar{j}} \times E_{\bar{j}}, \quad (w_i, z_j) \mapsto (w_i + z_i - z_j, z_j), \tag{4.33}
$$

where $J_{i\bar{j}}$ is a copy of *J* labelled by i , \bar{j} . We then apply the same discussions as above, with $J \times \text{Pic}^{(0)}(J)$ replaced by $J_{i\bar{j}} \times E_{\bar{j}}$. Then we have

$$
\prod_{i=1}^{n} \frac{\theta(w_i + z_i - z_j + c)}{\theta(w_i + z_i - z_j)} \in H^0(E_{[n]} \times E_{\bar{j}'} \otimes_{i=1}^{n} s_{ij}^* \mathcal{L}(0 \times \text{Pic}^{(0)}(J)))
$$

and thus

$$
s_j := f_0^{(j)} + f_1^{(j)} \sum_{i=1}^n \frac{\theta'(w_i + z_i - z_j)}{\theta(w_i + z_i - z_j)} \in H^0(E_{[n]} \times E_{\bar{j}}, \otimes_{i=1}^n s_{ij}^* \mathcal{L}(0 \times \text{Pic}^{(0)}(J))|_{2(J \times 0)}), \tag{4.34}
$$

where $f_0^{(j)}$ $f_0^{(j)}$, $f_1^{(j)}$ $\mathcal{L}(0\times \mathsf{Pic}^{(0)}(J))|_{J_{i\bar{j}}\times 0}$, $\mathcal{L}(0\times \mathsf{Pic}^{(0)}(J))|_{J_{i\bar{j}}\times 0}\otimes K_{J_{i\bar{j}'}}$ respectively.

Denote the trivializations $(f_0^{(j)})$ 0)1≤*j*≤*n*,(*f* (*j*) $\int_1^{(1)}$)_{1 $\leq j \leq n$} by f_0 , f_1 collectively, and similarly the frame $(s_1, \dots, s_j, \dots, s_n)$ by *s*. Then [\(4.34\)](#page-27-0) can be written compactly as

$$
s = \begin{pmatrix} f_0 & f_1 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ Z_n \end{pmatrix} . \tag{4.35}
$$

Passing to the real-analytic trivializations $\widehat{f}_0^{(j)}$ $\widehat{f}_1^{(j)}$, $\widehat{f}_1^{(j)}$ $\int_1^{(j)}$, $j \ge 1$ constructed similar to that in [\(4.29\)](#page-26-3), this gives

$$
\mathbf{s} = \begin{pmatrix} \hat{f}_0 & \hat{f}_1 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ \hat{Z}_n \end{pmatrix} . \tag{4.36}
$$

This can be regarded as describing a point in the Grassmannian

$$
Gr(n, V_n), V_n := \bigoplus_{j=1}^n \otimes_{i=1}^n s_{ij}^* H^0 \left(J \times 0, \mathcal{L}(0 \times \text{Pic}^{(0)}(J))|_{2(J \times 0)} \right), \tag{4.37}
$$

with det \widehat{Z}_n being one of the affine coordinates in the corresponding affine patch. To be more precise, using the frame $s = (s_j)_{1 \leq j \leq n}$, the Plücker embedding is given by

$$
\begin{array}{rcl}\n\text{Gr}\,(n, V_n) & \to & \mathbb{P}(\wedge^n V_n) \\
\text{span}\,(s_j)_{1 \leq j \leq n} & \mapsto & [\wedge^n s] \,, \qquad \wedge^n s = s_1 \wedge s_2 \wedge \cdots \wedge s_n \,. \end{array}\n\tag{4.38}
$$

From [\(4.36\)](#page-27-1), we obtain the Plücker coordinates of \wedge^n s:

$$
\wedge^n s = \sum_{\emptyset \subseteq I \subseteq [n]} \wedge^{\text{top}} \widehat{f}_I \cdot \det \widehat{Z}_I, \quad \wedge^{\text{top}} \widehat{f}_I := \wedge_{j \in I^c} \widehat{f}_0^{(j)} \wedge \wedge_{j \in I} \widehat{f}_1^{(j)}.
$$
\n(4.39)

Introducing a pairing $\langle -, - \rangle$ on $\wedge^n V_n$ by declaring $(\widehat{f}_I)_I$ to be an orthonormal basis, then

 $\langle \wedge^{top} \hat{f}_I, \wedge^n s \rangle = det \hat{Z}_I.$

Due to [\(4.32\)](#page-27-2), we can identify s with

$$
s(\varepsilon) := \begin{pmatrix} \widehat{f}_0 & \widehat{f}_1 \end{pmatrix} \cdot \begin{pmatrix} vol & 0 \\ 0 & \delta_n \end{pmatrix} \cdot \begin{pmatrix} 1 \\ \widehat{Z}_n \end{pmatrix}, \quad vol = \begin{pmatrix} vol_1 & \cdots & 0 \\ \vdots & vol_k & \vdots \\ 0 & \cdots & vol_n \end{pmatrix} .
$$
 (4.40)

It follows that

$$
\wedge^{n} s(\varepsilon) = \sum_{\emptyset \subseteq I \subseteq [n]} \wedge^{\text{top}} \widehat{f}_I \cdot \varepsilon_I \cdot \det \widehat{Z}_I \cdot \wedge_{k=1}^{n} \text{vol}_k. \tag{4.41}
$$

The iterated regularized integral then gives

$$
\oint_{E_{[n]}} \wedge^n s(\varepsilon) = \sum_{\emptyset \subseteq I \subseteq [n]} \wedge^{\text{top}} \hat{f}_I \cdot \varepsilon_I \hat{G}_I.
$$
\n(4.42)

In particular, we have

$$
\int_{E_{[n]}} \langle \wedge^{\text{top}} \widehat{f}_{[n]}, \wedge^n s(\varepsilon) \rangle = \varepsilon_{[n]} \cdot \widehat{G}_{[n]}.
$$

The relation [\(4.42\)](#page-28-7) tells that the generating series $\mathcal{G}_{[n]}(\varepsilon)$ given in Definition [4.7](#page-24-0) can be regarded as the iterated regularized integral of $\wedge_{j=1}^{n} s_j$, which is a section of the determinant line bundle over the Grassmannian Gr (n, V_n) . It is the truncation, induced by the map $\mathcal{L}|_{\infty(J\times 0)}\to \mathcal{L}|_{2(J\times 0)}$, of the corresponding construction using the formal neighborhood ∞ (*J* × 0). Concretely, the latter is realized by replacing the Taylor polynomial [\(4.31\)](#page-26-4) by the full Taylor series.

The construction using $\mathcal{L}|_{2(J\times 0)}$ is also the deformation of the construction with $\varepsilon = 0$, which corresponds to the restriction of ${\cal L}$ to the 0th order neighborhood ${\cal L}|_{(J\times0)}$ instead of the restriction to the 1st order neighborhood $\mathcal{L}|_{2\left(J\times 0 \right)} .$

References

- [BKT07] K. Bannai, S. Kobayashi, and T. Tsuji, *Realizations of the Elliptic Polylogarithm for CM elliptic curves*, RIMS Kôkyûroku Bessatsu, **B12** (2009), 33–50.
- [BKT10] K. Bannai, S. Kobayashi, and T. Tsuji, *On the de Rham and p-adic realizations of the elliptic polylogarithm for CM elliptic curves*, Ann. Sci. Ec. Norm. Super. (4) **43** (2010), no. 2, 185–234.
- [BL11] F. Brown and A. Levin, *Multiple Elliptic Polylogarithms*, [arXiv:1110.6917](http://arxiv.org/abs/1110.6917) [math.NT].
- [BO00] S. Bloch and A. Okounkov, *The Character of the Infinite Wedge Representation*, Advances in Mathematics **149** (2000), 1–60.
- [CEE09] D. Calaque, B. Enriquez and P. I. Etingof, *Universal KZB equations: the elliptic case*, in Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. I, 165–266, Progr. Math., **269**, Birkhäuser Boston, Boston, MA, 2009.
- [Che77] K. T. Chen, *Iterated path integrals*, Bulletin of the American Mathematical Society, Volume **83**, Number 5, September 1977.
- [Dij95] R. Dijkgraaf, *Mirror symmetry and elliptic curves*, The moduli space of curves (Texel Island, 1994), Progr. Math., vol. **129**, Birkhäuser Boston, Boston, MA, 1995, pp. 149–163.
- [DJKM81] E. Date, M. Jimbo, M. Kashiwara, and T. Miwa, *Transformation groups for soliton equations*, Proc. Japan Acad. **57A** (1981), 342–347, 387–392; J. Phys. Soc. Japan **50** (1981), 3806–3818; Physica **4D** (1982), 343–365; Publ. RIMS, Kyoto Univ. **18** (1982), 1077–1119.
- [DJKM83] E. Date, M. Jimbo, M. Kashiwara, and T. Miwa, *Transformation groups for soliton equations*, in Jimbo and Miwa (ed), Non-linear Integrable Systems - Classical Theory and Quantum Theory (Kyoto, 1981), Proc. R.I.M.S. Symposium, World Scientific, Singapore, 1983, pp. 39–119.
- [Dup15] C. Dupont, *The Orlik-Solomon model for hypersurface arrangements*. Annales de l'Institut Fourier, Volume **65** (2015) no. 6, pp. 2507–2545.
- [EZ85] M. Eicher and D. Zagier, *The theory of Jacobi forms*, Progress in Math. **55**, Birkhäuser Boston, Boston, MA, 1985.
- [FK17] G. Felder and D. Kazhdan, *Divergent Integrals, Residues of Dolbeault Forms, and Asymptotic Riemann Mappings*, International Mathematics Research Notices, Vol. **2017**, No. 19, pp. 5897–5918.
- [FM20] T. Fonseca and N. Matthes, *Towards algebraic iterated integrals on elliptic curves via the universal vectorial extension*, RIMS Kokyuroku, no. **2160** (2020), 114–125.
- [FM23] T. Fonseca and N. Matthes, *Elliptic KZB connections via universal vector extensions*, [arXiv:2301.12560](http://arxiv.org/abs/2301.12560) [math.AG].
- [GM20] E. Goujard and M. Möller, *Counting Feynman-like graphs: Quasimodularity and Siegel-Veech weight*, J. Eur. Math. Soc., **22** (2020), No 2, pp. 365–412.
- [KZ95] M. Kaneko and D. Zagier, *A generalized Jacobi theta function and quasimodular forms*, The moduli space of curves (Texel Island, 1994), Progr. Math., vol. **129**, Birkhäuser Boston, Boston, MA, 1995, 165–172.
- [Lib09] A. Libgober, *Elliptic genera, real algebraic varieties and quasi-Jacobi forms*, [arXiv:0904.1026](http://arxiv.org/abs/0904.1026) [math.AG].
- [LR07] A. Levin and G. Racinet, *Towards multiple elliptic polylogarithms*, [arXiv:math/0703237](http://arxiv.org/abs/math/0703237) [math.NT].
- [LZ21] S. Li and J. Zhou, *Regularized Integrals on Riemann Surfaces and Modular Forms*, Commun. Math. Phys. **388** (2021), 1403–1474.
- [LZ23] S. Li and J. Zhou, *Regularized Integrals on Elliptic Curves and Holomorphic Anomaly Equations*, Commun. Math. Phys. **401** (2023), 613–645.
- [OP06] A. Okounkov and R. Pandharipande, *Gromov-Witten theory, Hurwitz theory, and completed cycles*, Ann. of Math. (2) **163** (2006), no. 2, 517–560.
- [Rai89] A. Raina, *Fay's Trisecant Identity and Conformal Field Theory*, Commun. Math. Phys. **122**, 625-641 (1989).
- [Sil09] J. Silverman, *The arithmetic of elliptic curves*, vol. **106**, Springer Science & Business Media, 2009.
- [Spr19] J. Sprang, *Eisenstein-Kronecker series via the Poincaré bundle*. Forum Math. Sigma **7** (2019), Paper No. e34, 59 pp.
- [Spr20] J. Sprang, *The algebraic de Rham realization of the elliptic polylogarithm via the Poincaré bundle*. Algebra Number Theory **14** (3) 545–585, 2020.
- [SW85] G. Segal and G. Wilson, Loop groups and equations of KdV-type, Publ. Math. I.H.E.S, **63** (1985), 1–64.
- [Wei76] A. Weil, *Elliptic functions according to Eisenstein and Kronecker*, Ergebnisse der Mathematik bind **88**, Springer (1976).
- [Zag91] D. Zagier, *Periods of modular forms and Jacobi theta functions*, Inventiones mathematicae **104** (1991), no. 1, 449–465.
- [Zag16] , *Partitions, quasimodular forms, and the Bloch–Okounkov theorem*, The Ramanujan Journal **41** (2016), no. 1-3, 345–368.
- [Zho23a] J. Zhou, *Gromov-Witten Generating Series of Elliptic Curves as Configuration Space Integrals*, [arXiv:2304.03912](http://arxiv.org/abs/2304.03912) [math.AG].
- [Zho23b] , *Regularized Integrals on Configuration Spaces of Riemann Surfaces and Cohomological Pairings*, [arXiv:2305.12362](http://arxiv.org/abs/2305.12362) [math.AG].

Yau Mathematical Sciences Center, Tsinghua University, Beijing 100084, P. R. China Email: jzhou2018@mail.tsinghua.edu.cn