

Generalized GM-MDS: Polynomial Codes are Higher Order MDS

Joshua Brakensiek*

Manik Dhar[†]

Sivakanth Gopi[‡]

Abstract

The GM-MDS theorem, conjectured by Dau-Song-Dong-Yuen and proved by Lovett and Yildiz-Hassibi, shows that the generator matrices of Reed-Solomon codes can attain every possible configuration of zeros for an MDS code. The recently emerging theory of higher order MDS codes has connected the GM-MDS theorem to other important properties of Reed-Solomon codes, including showing that Reed-Solomon codes can achieve list decoding capacity, even over fields of size linear in the message length.

A few works have extended the GM-MDS theorem to other families of codes, including Gabidulin and skew polynomial codes. In this paper, we generalize all these previous results by showing that the GM-MDS theorem applies to any *polynomial code*, i.e., a code where the columns of the generator matrix are obtained by evaluating linearly independent polynomials at different points. We also show that the GM-MDS theorem applies to dual codes of such polynomial codes, which is non-trivial since the dual of a polynomial code may not be a polynomial code. More generally, we show that GM-MDS theorem also holds for *algebraic codes* (and their duals) where columns of the generator matrix are chosen to be points on some irreducible variety which is not contained in a hyperplane through the origin. Our generalization has applications to constructing capacity-achieving list-decodable codes as shown in a follow-up work [BDGZ24], where it is proved that randomly punctured algebraic-geometric (AG) codes achieve list-decoding capacity over constant-sized fields.

*University of California, Berkeley. Email: josh.brakensiek@berkeley.edu. Research supported in part by a Microsoft Research PhD fellowship while a student at Stanford University.

[†]Department of Mathematics, Massachusetts Institute of Technology. Email: dmanik@mit.edu. Part of this work was done while this author was a graduate student at the Department of Computer Science, Princeton University where his research was supported by NSF grant DMS-1953807.

[‡]Microsoft Research. Email: sigopi@microsoft.com

Contents

1	Introduction	3
1.1	Main Results	5
1.2	Technical Overview	6
1.2.1	Generalized GM-MDS Theorem for Monomial Codes	6
1.2.2	Generalized GM-MDS Theorem for Duals of Monomial Codes	8
1.2.3	Extension to Polynomial Codes and Irreducible Varieties	8
1.3	Open Questions	9
2	Preliminaries	9
2.1	Notation	9
2.2	Properties of Higher Order MDS Codes	10
3	Higher Order MDS Codes with a Basis	11
3.1	Definition	11
3.2	Comparison to $\text{MDS}(\ell)$ Codes	11
3.3	Properties of $\text{MDSb}(\ell)$ Codes	13
3.4	Inductive Framework	15
4	Algebraic GM-MDS for Monomial Codes	17
5	A GM-MDS Theorem for Dual Monomial Codes	19
5.1	Reduction	20
5.2	Proof of Theorem 5.2	22
6	Extension to Irreducible Varieties	24
6.1	Algebraic Preliminaries	24
6.2	Algebraic GM-MDS	25
6.3	Reduction to Power Series	26
6.4	Reduction of Power Series to Univariate Monomials	27
A	Omitted Proofs	32
A.1	Proof of Lemma 3.7	32
A.2	Proof of Lemma 6.2	33

1 Introduction

The optimal rate distance tradeoff of an (n, k) -code (over large alphabets) is given by the Singleton bound [Sin64] which states that the minimum distance d is at most $n - k + 1$. A code which achieves this bound is called an MDS (Maximum Distance Separable) code. MDS codes are some of the most important codes because of their optimality, for example they are extensively used in distributed storage. Reed-Solomon codes [RS60] are the most well-known family of MDS codes and one of the first polynomial codes to be discovered. An (n, k) Reed-Solomon code is obtained by evaluating degree $< k$ polynomials over \mathbb{F}_q at n distinct points from \mathbb{F}_q . These codes achieve the Singleton bound over any field of size $q \geq n$. In this paper, we will only consider linear codes defined over finite fields. It is not hard to show that a linear (n, k) -code C with generator matrix $G_{k \times n}$ is MDS iff for every $k \times k$ minor of G is nonzero. The generator matrix of a Reed-Solomon code is of the form

$$G_{k \times n} = \begin{bmatrix} 1 & 1 & \dots & 1 \\ \alpha_1 & \alpha_2 & \dots & \alpha_n \\ \alpha_1^2 & \alpha_2^2 & \dots & \alpha_n^2 \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_1^{k-1} & \alpha_2^{k-1} & \dots & \alpha_n^{k-1} \end{bmatrix} \quad (1)$$

where $\alpha_1, \alpha_2, \dots, \alpha_n \in \mathbb{F}_q$ are distinct. There are several natural generalizations of MDS codes and recently many of them have been unified into a single concept called *higher order MDS codes* [BGM22, BGM23]. Higher order MDS codes tie together the GM-MDS theorem, optimally list-decodable codes and MR tensor codes. We will now briefly discuss these connections.

GM-MDS theorem. In some coding theory applications, we need MDS codes whose generator matrices have constrained support, i.e., with a specific pattern of zeros. See [DSY14, HHYD14, YS13, DSDY13] for applications of such codes to multiple access networks and secure data exchange. Let $\mathcal{S} = (S_1, S_2, \dots, S_n)$ where $S_1, \dots, S_k \subset [n]$, we call such an \mathcal{S} a zero pattern for $k \times n$ matrices. If there are ℓ distinct non-empty sets among S_1, \dots, S_k , we say that \mathcal{S} is an order- ℓ zero pattern. We say that an (n, k) -code C with some generator matrix $G_{k \times n}$ attains the zero pattern \mathcal{S} if there exists an invertible matrix $M_{k \times k}$ such that $\tilde{G} = MG$ has zeros in the positions specified by \mathcal{S} (i.e., in positions $\cup_{i=1}^k \{i\} \times S_i$). Since \tilde{G} is also a generator matrix of C , equivalently, we say C attains the zero pattern \mathcal{S} if there exists a generator matrix of C which has zeros in positions of \mathcal{S} . For an MDS code to attain a zero pattern \mathcal{S} , a simple necessary condition is that the zero pattern should satisfy a Hall type condition:

$$\forall I \subset [k], \quad |\cap_{i \in I} S_i| \leq k - |I|. \quad (2)$$

It is easy to see why (2) is a necessary condition, if (2) is not satisfied, then there exists a $k \times k$ submatrix of G which has too many zeros and hence cannot be invertible. On the other hand, if (2) is satisfied, then by Hall's matching theorem every $k \times k$ submatrix has some matching of non-zeros entries and hence can be made invertible by a suitable (or generic) choice of the non-zero entries [DSY14]. Therefore, over large enough alphabet, one can always find an MDS matrix with a given zero pattern \mathcal{S} if (2) is satisfied. A pattern which satisfies condition (2) is called a *generic zero pattern* [BGM23].

Dau et al. [DSY14] conjectured that given any generic zero pattern, there exists a Reed-Solomon code over any field of size $q \geq n + k - 1$ which attains the zero pattern \mathcal{S} . This conjecture which came to be known as the *GM-MDS conjecture*¹ was eventually proved independently by Lovett [Lov21] and Yildiz and Hassibi [YH19]. The heart of the proof (using a reduction shown by [DSY14]) is to show that generic Reed-Solomon codes can achieve the zero pattern \mathcal{S} which itself reduces to showing that a symbolic matrix has non-zero determinant. By observing that the degree of each variable in the determinant polynomial is at most $n + k - 2$, one can immediately get an upper bound on the field size needed to achieve the zero pattern \mathcal{S} [DSY14]. So the GM-MDS theorem can equivalently be stated as follows.

¹It is unclear what GM stands as [DSY14] don't explain it in the paper where they introduced the conjecture. Our best guess is that it stands for 'Generator Matrix' given that we are looking for MDS codes with constrained generator matrices.

Theorem 1.1 (GM-MDS Theorem [DSY14, Lov21, YH19]). *Let $\mathcal{S} = (S_1, S_2, \dots, S_n)$ be any generic zero pattern. Then for any finite field \mathbb{F}_q , there exists some $\alpha_1, \dots, \alpha_n \in \overline{\mathbb{F}_q}$ (algebraic closure of \mathbb{F}_q) such that the Reed-Solomon code with generator matrix G as in (1) attains the zero pattern \mathcal{S} .*

An MDS code which achieves all generic zero patterns of order ℓ is called a GZP(ℓ) code [BGM23]. In other words, the GM-MDS theorem states that Reed-Solomon codes are GZP(ℓ) for all ℓ . GM-MDS type theorems have been proved for Gabidulin codes [Gab21] in [YH20] and for linearized Reed-Solomon codes (or skew polynomial codes) in [LWWS23]. Are these codes somehow very special or is there a common generalization of these results?

Question 1.2. *Is there a common generalization of the GM-MDS theorem which implies the existing results for Reed-Solomon codes, Gabidulin codes and linearized Reed-Solomon codes?*

Note that all these codes are polynomial codes. We will see shortly that we can in fact prove a GM-MDS theorem for all polynomial codes!

Optimally list-decodable codes. A code with minimum distance d can be uniquely decoded up to half the minimum distance. *List-decoding* introduced by [Woz58, Eli57] allows us to decode from beyond half the minimum distance, by instead outputting a small list of possible codewords. An (n, k) -code is (ρ, L) -list decodable if any Hamming ball of radius ρn contains at most L codewords. *Average-radius list-decoding* is a slight strengthening of list-decoding defined as follows.

Definition 1.1. *Let $\text{wt}(x)$ denote the hamming weight (i.e., number of nonzeros) of a vector x . We say that an (n, k) -code is (ρ, L) -average-radius list-decodable if there doesn't exist $L + 1$ codewords $c_0, c_1, \dots, c_L \in C$ and $y \in \mathbb{F}^n$ such that*

$$\frac{1}{L+1} \sum_{i=0}^L \text{wt}(c_i - y) \leq \rho n.$$

A code which is (ρ, L) -average-radius list-decodable is also (ρ, L) -list-decodable. A generalized Singleton bound for list-decoding was proved recently [ST23, Rot22, GST24] which states that a (ρ, L) -list-decodable (and hence also average-radius list-decodable) code should satisfy

$$\rho \leq \frac{L}{L+1}(n-k). \quad (3)$$

A code which meets this bound exactly for all list sizes $\leq L$ is called an LD-MDS($\leq L$) (list-decoding MDS) code.

MR tensor codes. Give an $(m, m-a)$ -code C_{col} and an $(n, n-b)$ -code C_{row} , the tensor code $C_{col} \otimes C_{row}$ consists of all $m \times n$ matrices where each row belongs to the row code C_{row} and each column belongs to the column code C_{col} . Tensor codes are used for erasure coding in data centers [MLR+14]. A Maximally Recoverable (MR) tensor code are ‘optimal’ tensor codes which can recover from any erasure pattern as long as it is information theoretically possible [GHK+17]. [GHK+17] also give an explicit combinatorial condition on which erasure patterns are information theoretically recoverable which called ‘regularity’.

Higher order MDS codes. In [BGM22], Brakensiek, Gopi and Makam introduced a generalization of MDS codes called *higher order MDS codes*.

Definition 1.2 ([BGM22]). *Let C be an (n, k) -code with generator matrix G . For $\ell \geq 1$, we say C is MDS(ℓ) if for any subsets $A_1, A_2, \dots, A_\ell \subset [n]$, we have that*

$$\dim(G_{A_1} \cap G_{A_2} \cap \dots \cap G_{A_\ell}) = \dim(W_{A_1} \cap W_{A_2} \cap \dots \cap W_{A_\ell})$$

where $G|_{A_i}$ is the submatrix of G with columns indexed by A_i , G_{A_i} is the column span of $G|_{A_i}$, and $W_{k \times n}$ is a generic matrix (over fields of the same characteristic).²

²Actually, the results of [BGM23] show that the RHS is independent of the characteristic of the field.

In their follow up work, [BGM23] showed a remarkable equivalence between various natural generalizations of MDS codes by showing that all of them are equivalent to higher order MDS codes.

Theorem 1.3 ([BGM23]). *The following are equivalent for any (n, k) -code C and $\ell \geq 2$.*

1. C is $\text{MDS}(\ell)$.
2. C is $\text{GZP}(\ell)$.
3. $C_{\text{parity}} \otimes C$ is an MR tensor code where C_{parity} is the $(\ell, \ell - 1)$ parity check code.
4. C^\perp is $\text{LD-MDS}(\leq \ell - 1)$ where C^\perp is the dual code of C .

As a result of this theorem, an equivalent way to state the GM-MDS theorem is to state that generic Reed-Solomon codes are higher order MDS codes, i.e., $\text{MDS}(\ell)$ for all ℓ . In particular, this implies that Reed-Solomon codes over sufficiently large alphabet are optimally list decodable, i.e., they achieve the list-decoding singleton bound (3) for all list sizes, which was originally conjectured by Shangguan and Tamo [ST23]. In subsequent work, Guo and Zhang [GZ23] showed that Reed-Solomon codes over fields of size $O_\epsilon(nk)$ can achieve list-decoding capacity, i.e., (ρ, L) list decodable with $\rho = 1 - R - \epsilon$ and $L = O(1/\epsilon)$ where $R = k/n$ is the rate of the code. Note that this is weaker than being optimally list-decodable, but still very useful for applications and crucially now requires only $O_\epsilon(nk)$ field size. In fact this relaxation is necessary, [BDG23] showed that achieving $\text{LD-MDS}(\leq 2)$ (i.e., optimally list decodable with for list sizes $L \leq 2$) already requires exponential field sizes. Shortly after, Alrabiah, Guruswami and Li [AGL24] further improved this to show that Reed-Solomon codes over $O_\epsilon(n)$ fields can achieve list-decoding capacity.

Given the rich set of connections of higher order MDS codes, answering Question 1.2 or equivalently understanding what codes have the higher order MDS property is a fundamental question which will likely lead to progress in several areas of coding theory such as list-decoding, MR tensor codes etc..

1.1 Main Results

We now discuss the main contributions of this paper. We satisfactorily answer Question 1.2 by showing that there is nothing very special about Reed-Solomon, Gabidulin or Linearized Reed-Solomon codes. We prove that GM-MDS theorem holds for all polynomial codes.

Definition 1.3 (Polynomial codes). *Let $f = (f_1, f_2, \dots, f_k) \in (\mathbb{F}[x_1, \dots, x_r])^k$ be a k -dimensional vector of polynomials such that f_1, \dots, f_k are linearly independent over \mathbb{F} . Let $\alpha_1, \alpha_2, \dots, \alpha_k \in \mathbb{F}^r$ be distinct. The polynomial code C_f is defined as the linear code with generator matrix $G_{k \times n}$ where*

$$G = \begin{bmatrix} f_1(\alpha_1) & f_1(\alpha_2) & \cdots & f_1(\alpha_k) \\ f_2(\alpha_1) & f_2(\alpha_2) & \cdots & f_2(\alpha_k) \\ \vdots & \vdots & \ddots & \vdots \\ f_k(\alpha_1) & f_k(\alpha_2) & \cdots & f_k(\alpha_k) \end{bmatrix}. \quad (4)$$

We say that C_f is generic if $\alpha_1, \dots, \alpha_k \in \overline{\mathbb{F}}^r$ are chosen to be algebraically independent over \mathbb{F} .

Theorem 1.4 ((Informal) Generalized GM-MDS Theorem for Polynomial Codes). *Generic polynomial codes are higher order MDS, i.e., is $\text{MDS}(\ell)$ for all ℓ .*

Example 1.1. *We can recover all the known examples of GM-MDS theorem using Theorem 1.4.*

1. Reed-Solomon codes are polynomial codes with $f(x) = (1, x, x^2, \dots, x^{k-1})$.
2. Gabidulin codes are polynomial codes with $f(x) = (x, x^q, x^{q^2}, \dots, x^{q^{k-1}})$ where $|\mathbb{F}| = q^m$ for some m .

3. *Linearized Reed-Solomon or Skew polynomial codes are polynomial codes with*

$$f(x, y) = (x, x^q y, x^{q^2} y^{1+q}, \dots, x^{q^{k-1}} y^{1+q+\dots+q^{k-2}})$$

where $|\mathbb{F}| = q^m$ for some m . Note that in general, linearized Reed-Solomon codes can be MDS while sharing some of the x, y among various columns. But the GM-MDS theorem in [LWWS23] is only proved when the x, y in each column are different. So the result of [LWWS23] is indeed a special case of our result.

If a polynomial code $f = (f_1, \dots, f_k)$ is composed of degree at most d polynomials then it is not hard to show that any particular order ℓ pattern can be achieved by $[n, k]$ -code over a field of size $d(\ell - 1) + 1$ (this does not include the condition that the code is MDS). It simply follows from showing the individual degree of each variable in the determinant of Lemma 2.3 is at most $d(\ell - 1)$. As mentioned earlier, for Reed-Solomon codes an MDS code which satisfies a particular zero pattern can be constructed over a field of size $q \geq n + k - 1$ (this is because one could simplify the determinant in Lemma 2.3 to get better individual degree bounds). In a follow up paper, starting from a generalization of these theorems we will show how to construct codes which satisfy ‘relaxed’ generic zero patterns over a field of constant size (see end of this section).

All known examples of codes satisfying GM-MDS theorem such as Reed-Solomon, Gabidulin, Linearized Reed-Solomon are self-duality property, i.e., their dual codes are also of the same kind. For example, the dual of a Reed-Solomon code is again a (generalized) Reed-Solomon code. However, this is not true for most polynomial codes. And since a code C being LD-MDS($\leq L$) (i.e., optimally list decodable) is equivalent to the dual code C^\perp being MDS($L + 1$), it is important to know whether duals of generic polynomials codes are also higher order MDS and hence optimally list-decodable.

Theorem 1.5 ((Informal) Generalized GM-MDS Theorem for Dual Polynomial Codes). *Duals of generic polynomial codes are higher order MDS, i.e., is MDS(ℓ) for all ℓ . As a result, generic polynomial codes and their duals are MDS(ℓ) for all ℓ and LD-MDS($\leq L$) for all L .*

We can generalize Theorems 1.4 and 1.5 even further. Instead of specifying an explicit parameterization for f , we instead consider the columns of our code come from the solutions to an *implicit* set of polynomial equations (or, more formally an *irreducible variety*).

Theorem 1.6 ((Informal) Generalized GM-MDS for irreducible varieties). *Let $X \subset \overline{\mathbb{F}}^k$ be an irreducible variety which is not contained in any hyperplane. Let v_1, v_2, \dots, v_n be generic points from X . Then the (n, k) -code C with generator matrix $G = [v_1, v_2, \dots, v_k]$ as well as the dual code C^\perp are MDS(ℓ) for all ℓ .*

Theorems 1.4 and 1.5 are a special case of this theorem where the variety is $X = \{f(x) : x \in \overline{\mathbb{F}}^r\}$. Note that any variety which can be parametrized is irreducible and the linear Independence of f_1, f_2, \dots, f_k implies that X is not contained in any hyperplane.

In a follow up work [BDGZ24], Theorem 1.6 is crucially used to prove that randomly punctured Algebraic-Geometric codes achieve list decoding capacity over constant field sizes.

1.2 Technical Overview

We now outline how each of our main results are proved. First, it is possible to reduce the GM-MDS theorem for arbitrary irreducible varieties to the case of monomial codes. A monomial code is a polynomial code as in Definition 1.3 where each f_i is a monomial. We first describe an overview of the proof for monomial codes and then discuss the reduction.

1.2.1 Generalized GM-MDS Theorem for Monomial Codes

We prove Theorem 1.4 by an inductive approach similar to how the original GM-MDS theorem was proved [Lov21, YH19]. We first recall how the GM-MDS theorem is proved.

The work of [DSY14] showed that the GM-MDS theorem is equivalent to the following fact. Let $A_1, \dots, A_k \subseteq [n]$ be sets of size $k-1$, such that for any $J \subseteq [k]$ nonempty, we have that $|\bigcap_{j \in J} A_j| \leq k - |J|$. It suffices to show for any sufficiently large field \mathbb{F} , (i.e., $|\mathbb{F}| \geq n + k - 1$), there exists distinct $\alpha_1, \dots, \alpha_n \in \mathbb{F}$ such that the polynomials $g_i(x) := \prod_{j \in A_i} (x - \alpha_j)$ are linearly independent over \mathbb{F} .

The works [Lov21, YH19] prove this linear independence by inductively proving a generalized statement where they allow each polynomial g_i to have 0 as a repeated root.³ In particular, for each $i \in [k]$, they keep track of a set $A_i \subseteq [n]$ and $\sigma_i \geq 0$ such that $\sigma_i + |A_i| = k - 1$. They inductively show that the polynomials $g_i(x) := x^{\sigma_i} \prod_{j \in A_i} (x - \alpha_j)$ are linearly dependent over \mathbb{F} if and only if for all $J \subseteq [k]$ nonempty, we have that

$$\min_{j \in J} \sigma_j + \left| \bigcap_{j \in J} A_j \right| \leq k - |J|. \quad (5)$$

Outside of simple base cases, the inductive proof splits into three cases (per the convention used by [YH19]):

Case 1. There is $J \subseteq [k]$ with $2 \leq |J| \leq k-1$ for which (5) is tight. Define $g_J(x) := x^{\min_{j \in J} \sigma_j} \prod_{i \in \bigcap_{j \in J} A_j} (x - \alpha_i)$. The authors use the inductive hypothesis twice: first they argue that $\text{span}\{g_i : i \in J\} = \text{span}\{x^i g_J(x) : i \in \{0, 1, \dots, |J| - 1\}\}$. They then apply a second use of the induction hypothesis is to argue that $\{x^i g_J(x) : i \in \{0, 1, \dots, |J| - 1\}\} \cup \{g_j(x) : j \in [k] \setminus J\}$ is linearly independent.

Case 2. There is exactly one $i \in [\ell]$ for which $\sigma_i = 0$. Note that at least one $\sigma_i = 0$ or else (5) fails for $J = [k]$. Since every $g_j(x)$ with $j \neq i$ is a multiple of x , it suffices to argue that $\{g_j(x)/x : j \in [k] \setminus i\}$ are linearly independent over \mathbb{F} , which follows from the inductive hypothesis.

Case 3. Any other situation. In this case the authors identify an index $j \in [n]$ such that after assigning $\alpha_j \rightarrow 0$ —that is, if $j \in A_i$, remove j from A_i and increase σ_i by 1—we have that (5) still holds. After applying the inductive hypothesis, they show algebraically that undoing this operation preserves the linear independence of the g_i .

For Theorem 1.4, the connection between arbitrarily polynomial codes with the roots of polynomials is less clear. As such we generalize this technique of “repeated roots” within the higher order MDS framework of [BGM22, Rot22, BGM23]. Theorem 1.3 of [BGM23] shows that proving a GM-MDS theorem is equivalent to the following. Let C be our polynomial code with generator matrix G as in (4), we seek to show for any $A_1, \dots, A_k \subseteq [n]$ of size $k-1$, such that for any $J \subseteq [k]$, we have that $|\bigcap_{j \in J} A_j| \leq k - |J|$, we have that $G_{A_1} \cap \dots \cap G_{A_k} = 0$.

Higher Order MDS Codes with a Basis. The combinatorial structure of the induction is the same [Lov21, YH19], but our key novelty is how we adapt the “repeated root” trick. A first idea is that when one seeks to substitute α_j with 0, we replace each instance of G_j with a fixed vector v . However, after few substitutions, one quickly runs into a situation in which one wants to make the same substitution for multiple elements in the same A_i . To get around this, we have ordered set of vectors $u_1, \dots, u_k \in \mathbb{F}^k$ such that if s vectors in A_i are “set to zero,” then we replace those vectors with u_1, \dots, u_s . More formally, we call this a “higher order MDS code with a basis.”

Definition 1.4. Consider $n, k, b \geq 1$ with $b \leq k$. Let $U \in \mathbb{F}^{k \times b}$ and $V \in \mathbb{F}^{k \times n}$ be matrices. We say that (U, V) is a (b, n, k) -order- ℓ higher order MDS code with a basis, or (b, n, k) -MDSb(ℓ), if for all configurations

³The natural question of allowing each g_i to have multiple repeated roots is in general open, see [GS19]. One important observation is that the answer to the problem depends on the characteristic of the field, so it cannot have as simple of an answer as the GM-MDS theorem.

$(\sigma_1, A_1), \dots, (\sigma_\ell, A_\ell)$, we have that

$$\dim \left(\bigcap_{i=1}^{\ell} (U_{[\sigma_i]} + V_{A_i}) \right) = \dim \left(\bigcap_{i=1}^{\ell} (W_{[\sigma_i]} + W'_{A_i}) \right), \quad (6)$$

where $W \in \mathbb{F}^{k \times b}$ and $W' \in \mathbb{F}^{k \times n}$ are generic matrices.

The generalized statement we prove is that for any code C with a generator matrix like (4), there is a matrix $U \in \mathbb{F}^{k \times k}$ such that (U, G) is $\text{MDSb}(\ell)$ for all $\ell \geq 2$. Note by invoking (6) in the case that each $\sigma_i = 0$, we get that G is $\text{MDS}(\ell)$. Once a suitable basis is identified for our code C , the strategy of [Lov21, YH19], can be followed; although it takes some effort to show that conditions like (6) behave analogous to results on repeated roots of polynomials—this groundwork is laid in Section 3.

In the case of Reed-Solomon codes, we can set U to be the identity matrix; that is, the appropriate choice of basis is just the standard basis vectors $e_1, \dots, e_k \in \mathbb{F}^k$. In general for polynomial codes, the choice of basis depends on the degrees of the nonzero coefficients of f_1, \dots, f_k . However, for “monomial codes”, where each $f_i(x) = x^{d_i}$, for some integers $d_1 < d_2 < \dots < d_k$, we can again use the standard basis. In Section 4 we give all the details of the proof for monomial codes, and in Section 6, we give a reduction from polynomial codes to monomial codes (see also Section 1.2.3).

1.2.2 Generalized GM-MDS Theorem for Duals of Monomial Codes

As explained earlier, except in special cases, a generic polynomial code C does not necessarily have that the property that its dual code C^\perp is a generic polynomial code. Therefore, we cannot use the (exact) same approach to prove that C^\perp is $\text{MDS}(\ell)$ (and thus C is $\text{LD-MDS}(\leq \ell - 1)$) for any generic polynomial code C .

An approach which one might take is to directly compute a parity check matrix H of C in terms of the entries of the generator matrix G from (4). However, the entries of H depend on the values of many of the α_i 's and thus make it nearly impossible to directly show that H is $\text{MDS}(\ell)$, even with a suitable basis.

Instead, we show a surprising connection between higher order MDS codes with a basis and dual codes. In particular, we show that if (G^T, I_n) is $\text{MDSb}(\ell)$, then C^\perp is $\text{MDS}(\ell)$ (see Theorem 5.4). In other words, if we think of the rows of G as a (partial) basis of \mathbb{F}^n , it suffices to show that the criteria of (6) hold when the “main” part of the code is the standard basis of \mathbb{F}^n .

Like in the case of Theorem 1.4, we can assume for the proof of Theorem 1.5 that the generator matrix G of the code C consists of monomials. In this case, we can prove that (G^T, I_n) is $\text{MDSb}(\ell)$ using a GM-MDS-style induction (see Theorem 5.2). A crucial detail in the proof is that since the algebraic complexity of analyzing statements like (6) comes from G^T , the inductive proof removes “repeated roots” rather than adding them. In particular, the base case in which each $\sigma_i = 0$ is easy to analyze due to I_n being the standard basis. Once the order of induction is properly chosen, the remainder of the proof of Theorem 1.5 is similar to that of Theorem 1.4.

1.2.3 Extension to Polynomial Codes and Irreducible Varieties

So far, we have only explained the proofs of Theorem 1.4 and Theorem 1.5 for monomial codes, where each polynomial of (4) has a single nonzero coefficient. To extend this result to general polynomials, we observe that the higher order MDS conditions are equivalent to certain determinants based on the columns of the generator matrix G being nonzero (see Lemma 2.3). We show that once a suitable change of basis is performed to G , a leading monomial in the determinant expansion corresponds to the analogous determinant where each polynomial of G is replaced by its leading monomial. This only works when the column of the generator matrix comes from the same polynomial family. As such, the GM-MDS theorems we proved for monomial codes also apply to general polynomial codes.

We also extend these results to general irreducible varieties. In this case, we use the fact that any irreducible variety can be (generically) parameterized by a *power series*, essentially a polynomial where there are infinitely many terms. These power series can be analyzed essentially like polynomials, in particular they

can be reduced to analyzing a suitable monomial code. As such, Theorem 1.6 follows from the proofs of Theorem 1.4 and Theorem 1.5.

1.3 Open Questions

We conclude the paper with some open questions.

- Can we generalize Theorems 1.4 and 1.5 to the setting where different columns are evaluations of different polynomial maps? More generally, can we generalize 1.6 to the setting where different columns are generic points from different irreducible varieties?
- Can we generalize our GM-MDS theorems to the setting where the columns are evaluations of f at points which share some of the variables across columns. For example, linearized Reed-Solomon codes are such an example of MDS codes. See Example 1.1.
- One very broad generalization of the GM-MDS theorem which immediately implies all known GM-MDS type theorems (including ours) is as follows.

Conjecture 1.7 (Ultimate GM-MDS conjecture). *Let C_0 be any (N, k) -code which is MDS. Let C be an (n, k) -code obtained by a random puncturing of C_0 (i.e., its generator is formed by randomly selecting n columns from the generator of C_0). Then as with probability $1 - o_N(1)$, C is MDS(ℓ) for all ℓ .*

If Conjecture 1.7 is true, it says that GM-MDS is not really a special property of polynomial codes, but rather a generic property of MDS codes.

- The study of higher order MDS codes was motivated by the study of MR Tensor Codes. Two MDS codes $C_1 \subseteq \mathbb{F}^m$ and $C_2 \subseteq \mathbb{F}^n$ form an MR tensor code if for every subset $E \subseteq [m] \times [n]$, $(C_1 \otimes C_2)|_E$ achieves the dimension it should if C_1 and C_2 are generically chosen. The appropriate generalization of Theorem 1.4 in this setting is to show that the tensor of two generic polynomial codes is MR, although this question is open just in the case that C_1 and C_2 are generic Reed-Solomon codes.

Organization

In Section 2, we define our notation and state some known facts about higher order MDS codes. In Section 3, we formally define a Higher Order MDS code with a basis and use this to build an inductive framework for proving GM-MDS-type theorems. In Section 4, we prove our GM-MDS theorem for monomial codes. In Section 5, we prove our GM-MDS theorem for duals of monomial codes. In Section 6, we show the results of Sections 4 and 5 extend to polynomial codes and irreducible varieties. In Appendix A, we include proofs omitted from the main paper.

2 Preliminaries

In this section, we go over some technical background material not explained in the introduction.

2.1 Notation

In this paper we let \mathbb{F} denote an arbitrary field. Although applications to coding theory require \mathbb{F} to be finite, our results generally apply to any field (of any characteristic).

A *linear code* is a subspace $C \subseteq \mathbb{F}^n$, where n is number of symbols of the code. If k is the dimension of n , then we say that C is a (n, k) -code. We call points $x \in C$ *codewords* of C . The *Hamming weight* of a codeword $c \in C$ is the number of nonzero symbols which we denote by $\text{wt}(c)$. A *generator matrix* $G \in \mathbb{F}^{k \times n}$ of a linear code $C \subseteq \mathbb{F}^n$ is any matrix whose rows form a basis of C .

The *dual code* of C is the linear code $C^\perp := \{y \in \mathbb{F}^n \mid \forall x \in C, x_1y_1 + \dots + x_ny_n = 0\}$. We say that a generator matrix of the dual code C^\perp is a *parity check matrix* of the original code C .

We say that C is *maximum distance separable (MDS)* if a generator matrix $G \in \mathbb{F}^{k \times n}$ of C has every k columns linearly independent. This is equivalent to the more traditional definition of MDS that every nonzero codeword of C has Hamming weight at least $n - k + 1$.

Given a matrix $G \in \mathbb{F}^{k \times n}$, for $i \in [n] := \{1, 2, \dots, n\}$, we let G_i denote the i th column of G . For a subset $A \subseteq [n]$, we let $G|_A$ denote the submatrix of G with columns from A . We also let $G_A := \text{span}\{G_i \mid i \in A\}$. Note that some previous works like [BGM22] use $G|_A$ and G_A interchangeably. In this paper, we often use a code and its generator matrix interchangeable if the property of the matrix is independent of the choice of generator matrix. For example, the rank of $G|_A$ is such a property.

If we have subsets $S \subseteq [k]$ and $T \subseteq [n]$, we let $G|_{S \times T}$ denote the submatrix with entries $(i, j) \in S \times T$.

Given two matrices U and V with the same number of rows, we let $[U \mid V]$ denote the matrix with the columns of U and V concatenated. We let I_k denote the $k \times k$ identity matrix. If k is implicit from context, we write just I (e.g., $[I \mid V]$).

Given a field \mathbb{F} and parameters (n, k) , we define a *generic (n, k) -matrix* as follows. Let \mathbb{F}_{ex} be an extension of \mathbb{F} with kn transcendental⁴ variables $\{w_{i,j} : (i, j) \in [k] \times [n]\}$. Our generic matrix is $W \in \mathbb{F}_{\text{ex}}^{k \times n}$ where $W_{i,j} = w_{i,j}$ for all $(i, j) \in [k] \times [n]$. Given a polynomial code such as in (4), we say that it is a *generic polynomial code* if $\alpha_1, \dots, \alpha_n$ are algebraically independent, transcendental variables in a suitable extension of \mathbb{F} .

2.2 Properties of Higher Order MDS Codes

We now state a few properties of higher order MDS codes that did not appear in the introduction.

Definition 2.1 ([BGM23]). *We say that $A_1, \dots, A_\ell \subseteq [n]$ have the dimension k null intersection property if for all partitions $P_1 \cup \dots \cup P_s = [\ell]$, we have that*

$$\sum_{i=1}^s \left| \bigcap_{j \in P_i} A_j \right| \leq (s-1)k.$$

Theorem 2.1 ([BGM23]). *Let V be a $k \times n$ matrix. Then, V is (n, k) -MDS(ℓ) if and only if for all $A_1, \dots, A_\ell \subseteq [n]$ with $|A_i| \leq k$ and $|A_1| + \dots + |A_\ell| = (\ell-1)k$ with the dimension k null intersection property, we have that $V_{A_1} \cap \dots \cap V_{A_\ell} = 0$.*

Theorem 2.2 (Lemma 2.8 [BGM23], adapted). *Let $A_1, \dots, A_\ell \subseteq [n]$ be subsets of size at most k . Let $d \in \{0, 1, \dots, k\}$, the following are equivalent*

- (a) *In any MDS(ℓ) code V , we have that $\dim(V_{A_1} \cap \dots \cap V_{A_\ell}) \leq d$.*
- (b) *For all partitions $P_1 \sqcup \dots \sqcup P_s = [\ell]$, we have that*

$$\sum_{i=1}^s \left| \bigcap_{j \in P_i} A_j \right| \leq (s-1)k + d.$$

- (c) *There exist $\delta_1, \dots, \delta_\ell \geq 0$ with $\delta_1 + \dots + \delta_\ell = k - d$ such that for all $J \subseteq [n]$ nonempty, we have that*

$$\left| \bigcap_{j \in J} A_j \right| \leq k - \sum_{j \in J} \delta_j.$$

⁴That is, there is no nonzero polynomial equation in the $w_{i,j}$'s with coefficients in \mathbb{F} which evaluates to 0. In other words, the $w_{i,j}$'s can be thought of as "symbolic" variables.

We shall also use the following matrix identity for checking MDS(ℓ) conditions.

Lemma 2.3 ([Tia19, BGM22]). *Let V be a $k \times n$ matrix. Consider $A_1, \dots, A_\ell \subseteq [n]$ with $|A_i| \leq k$. We have that*

$$\dim(V_{A_1} \cap \dots \cap V_{A_\ell}) = k + \sum_{i=1}^{\ell} \dim(V_{A_i}) - \text{rank} \begin{pmatrix} I_k & V|_{A_1} & & & \\ I_k & & V|_{A_2} & & \\ \vdots & & & \ddots & \\ I_k & & & & V|_{A_\ell} \end{pmatrix}. \quad (7)$$

Furthermore, if $|A_1| + \dots + |A_\ell| = (\ell - 1)k$ and $\dim(V_{A_i}) = |A_i|$ for all $i \in [\ell]$, then we have that $V_{A_1} \cap \dots \cap V_{A_\ell} = 0$ if and only if the block matrix in (7) has nonzero determinant.

3 Higher Order MDS Codes with a Basis

In this section, we generalize the definition of higher order MDS codes to also allow some vectors to be chosen in a fixed order from a basis. As mentioned in the technical overview, this will allow us to substantially generalize the “repeated roots” trick in the proof of the GM-MDS theorem.

3.1 Definition

Consider $n, k, b \geq 1$ with $b \leq k$. Let $U \in \mathbb{F}^{k \times b}$ and $V \in \mathbb{F}^{k \times n}$ be matrices. We define an *order- ℓ configuration* to be ℓ pairs $(\sigma_1, A_1), \dots, (\sigma_\ell, A_\ell)$, where for each $i \in [\ell]$ we have that $0 \leq \sigma_i \leq b$ and $A_i \subseteq [n]$. For all $J \subseteq [\ell]$ nonempty, let

$$\sigma_J := \min_{i \in J} \sigma_i$$

$$A_J := \bigcap_{i \in J} A_i.$$

Let $U_{\leq \sigma} = \text{span}\{u_i : i \in [\sigma]\}$, and $U|_{\leq \sigma}$ denote the submatrix of the first σ columns of U .

Definition 3.1. *Consider $n, k, b \geq 1$ with $b \leq k$. Let $U \in \mathbb{F}^{k \times b}$ and $V \in \mathbb{F}^{k \times n}$ be matrices. We say that (U, V) is a (b, n, k) -order- ℓ higher order MDS code with a basis, or (b, n, k) -MDSb(ℓ), if for all configurations $(\sigma_1, A_1), \dots, (\sigma_\ell, A_\ell)$, we have that*

$$\dim \left(\bigcap_{i=1}^{\ell} (U_{\leq \sigma_i} + V_{A_i}) \right) = \dim \left(\bigcap_{i=1}^{\ell} (W_{\leq \sigma_i} + W'_{A_i}) \right), \quad (8)$$

where $W \in \mathbb{F}^{k \times b}$ and $W' \in \mathbb{F}^{k \times n}$ are generic matrices.

3.2 Comparison to MDS(ℓ) Codes

To help the reader get intuition for Definition 1.4, we compare and contrast with Definition 1.2.

Like in [BGM22], this definition is monotone in ℓ . That is, if (U, V) is MDSb(ℓ), then it is MDSb(ℓ') for all $\ell' \leq \ell$. This can be observed by setting some of the pairs (σ_i, A_i) in (8) equal to each other (c.f. [BGM22]).

Another key observation is the following:

Proposition 3.1. *If (U, V) is MDSb(ℓ), then V is MDS(ℓ).*

Proof. In (8), set $\sigma_i = 0$ for all $i \in [\ell]$, we then derive the condition in Definition 1.2. □

In particular, technique for proving that a code V is $\text{MDS}(\ell)$ is to find a suitable basis U such that (U, V) is $\text{MDSb}(\ell)$. This stronger statement can lend itself to an inductive proof, see Section 3.4

However, there are codes for which (U, V) is $\text{MDSb}(1)$ but $[U \mid V]$ is not MDS . For instance, let the ambient field be \mathbb{F}_2 , have $U = I_3$, and V be the matrix

$$V = \begin{pmatrix} 1 & 1 \\ 1 & 1 \\ 0 & 1 \end{pmatrix},$$

then one can verify that (U, V) is $\text{MDSb}(1)$, but $[U \mid V]$ is not MDS , as the columns of V and the third column vector of U are linearly dependent.

As more nontrivial example, have $U = I_4$ and $V \in \mathbb{F}^{2 \times 4}$ be a generic Reed-Solomon code:

$$V = \begin{pmatrix} 1 & 1 \\ \alpha & \beta \\ \alpha^2 & \beta^2 \\ \alpha^3 & \beta^3 \end{pmatrix}.$$

The original proofs of the GM-MDS theorem [Lov21, YH19] show that (U, V) is $\text{MDSb}(4)$. However, $[U \mid V]$ is not $\text{MDS}(4)$. To see why, let $A_1 = \{1, 2, 5\}$, $A_2 = \{3, 4, 5\}$, $A_3 = \{1, 3, 6\}$, and $A_4 = \{2, 4, 6\}$, then one can show

$$[U \mid V]_{A_1} \cap [U \mid V]_{A_2} \cap [U \mid V]_{A_3} \cap [U \mid V]_{A_4} \neq 0,$$

as can be checked via Theorem 2.1, despite this identity being true for a generic code.

Although an $\text{MDSb}(1)$ code (U, V) is not necessarily MDS , every subset of the columns of $[U \mid V]$ which shows up in computing (8) *does* behave like an MDS code. This is formalized in the following proposition.

Proposition 3.2. *Let (U, V) be a (b, n, k) - $\text{MDSb}(1)$ code. For any configuration (σ, A) , we have that*

$$\dim(U_{\leq \sigma} + V_A) = \min(\sigma + |A|, k).$$

Proof. For generic codes, W and W' , we must have that $\dim(U_{\leq \sigma} + V_A) = \dim(W_{\leq \sigma} + W'_A) = \min(\sigma + |A|, k)$. \square

As a corollary, we have that the \geq direction of (8) follows whenever (U, V) is $\text{MDSb}(1)$ (c.f., [BGM22] or Lemma 3.1 of [BGM23]).

Corollary 3.3. *Let (U, V) is $\text{MDSb}(1)$, then for all configurations $(\sigma_1, A_1), \dots, (\sigma_\ell, A_\ell)$, we have that*

$$\dim \left(\bigcap_{i=1}^{\ell} (U_{\leq \sigma_i} + V_{A_i}) \right) \geq \dim \left(\bigcap_{i=1}^{\ell} (W_{\leq \sigma_i} + W'_{A_i}) \right), \quad (9)$$

where W and W' are generic matrices of the same shape as U and V , respectively.

Proof. By Lemma 2.3 and the fact that $\dim(U_{\leq \sigma_i} + V_{A_i}) = \dim(W_{\leq \sigma_i} + W'_{A_i})$ for all $i \in [\ell]$, we have that

$$\begin{aligned} & \dim \left(\bigcap_{i=1}^{\ell} (U_{\leq \sigma_i} + V_{A_i}) \right) - \dim \left(\bigcap_{i=1}^{\ell} (W_{\leq \sigma_i} + W'_{A_i}) \right) \\ &= \text{rank} \begin{pmatrix} I_k & W|_{\leq \sigma_1} & W'|_{A_1} & & & & & & & & \\ I_k & & & W|_{\leq \sigma_2} & W'|_{A_2} & & & & & & \\ \vdots & & & & & \ddots & & & & & \\ I_k & & & & & & & & W|_{\leq \sigma_\ell} & W'|_{A_\ell} & \end{pmatrix} \\ & - \text{rank} \begin{pmatrix} I_k & U|_{\leq \sigma_1} & V|_{A_1} & & & & & & & & \\ I_k & & & U|_{\leq \sigma_2} & V|_{A_2} & & & & & & \\ \vdots & & & & & \ddots & & & & & \\ I_k & & & & & & & & U|_{\leq \sigma_\ell} & V|_{A_\ell} & \end{pmatrix}. \end{aligned}$$

Since (W, W') is generic, the substitution $(W, W') \rightarrow (U, V)$ can only decrease the rank⁵ of the block matrix. Thus, the final expression is nonnegative. \square

3.3 Properties of $\text{MDSb}(\ell)$ Codes

In this section, we shall prove various properties about $\text{MDSb}(\ell)$ codes, which shall lead to an inductive framework for proving that various codes are $\text{MDSb}(\ell)$, see Section 3.4.

We say that a configuration $(\sigma_1, A_1), \dots, (\sigma_\ell, A_\ell)$ is *proper* if $\sigma_i + |A_i| \leq k$ for all $i \in [\ell]$. We observe that it suffices to check that a code is $\text{MDSb}(\ell)$ by considering only proper configurations.

Proposition 3.4. *If (8) for all proper configurations, then (8) holds for all configurations.*

Proof. We prove this by induction on ℓ . The case $\ell = 1$ follows from Proposition 3.2 and that $\dim(U_{\leq \sigma} + V_A)$ is monotone in (σ, A) . For $\ell \geq 2$, note that if we have a configuration with some $\sigma_i + |A_i| > k$, then by the induction hypothesis, we have that $\dim(U_{\leq \sigma_i} + V_{A_i}) = k$, so $U_{\leq \sigma_i} + V_{A_i} = \mathbb{F}^k$. In particular, we can replace (σ_i, A_i) with any (σ'_i, A'_i) with $\sigma'_i + |A'_i| = k$ and whether (8) holds will not change. We can repeat this procedure until we are left with a proper configuration. \square

By adapting the generic intersection dimension formula of [BGM23], we can get an explicit combinatorial formula for the RHS of (8) for proper configurations.

Proposition 3.5. *Let $W \in \mathbb{F}^{k \times b}$ and $W' \in \mathbb{F}^{k \times n}$ be generic matrices. Let $(\sigma_1, A_1), \dots, (\sigma_\ell, A_\ell)$ be a proper configuration. Then,*

$$\dim \left(\bigcap_{i=1}^{\ell} (W_{\leq \sigma_i} + W'_{A_i}) \right) = \max_{P_1 \sqcup \dots \sqcup P_s = [\ell]} \left[-n(s-1) + \sum_{i=1}^s (\sigma_{P_i} + |A_{P_i}|) \right], \quad (10)$$

where $P_1 \sqcup \dots \sqcup P_s = [\ell]$ denotes a partition of $[\ell]$ into s nonempty pieces.

Proof. Let $B := \{i_1, \dots, i_b\}$ be some set of b indices disjoint from $[n]$. For all $i \in [\ell]$, let

$$C_i := A_i \cup \{i_1, \dots, i_{\sigma_i}\}.$$

It is easy to see that for all $J \subseteq [\ell]$ nonempty that $|C_J| = |\sigma_J| + |A_J|$. Thus, (10) is equivalent to

$$\dim \left(\bigcap_{i=1}^{\ell} [W|W']_{C_i} \right) = \max_{P_1 \sqcup \dots \sqcup P_s = [\ell]} \left[-n(s-1) + \sum_{i=1}^s |C_{P_i}| \right],$$

and this follows from the generic intersection dimension formula of [BGM23] (i.e., (a) \iff (b) of Theorem 2.2). \square

For convenience, we let $\text{gid}_k((\sigma_1, A_1), \dots, (\sigma_\ell, A_\ell))$ denote the RHS of (10).

Definition 3.2. *We call a proper configuration $(\sigma_1, A_1), \dots, (\sigma_\ell, A_\ell)$ to be maximal if we additionally have the following properties*

- (a) $\sum_{i=1}^{\ell} \sigma_i + |A_i| = (\ell - 1)k$.
- (b) For all partitions $P_1 \sqcup \dots \sqcup P_s = [\ell]$, we have that

$$\sum_{i=1}^s \sigma_{P_i} + |A_{P_i}| \leq (s - 1)k.$$

⁵More formally, the condition that a matrix has rank at least r is equivalent to whether there exists a $r \times r$ submatrix with nonzero determinant. Since the entries of (W, W') are transcendental, any such determinant in the block matrix is necessarily nonzero for (W, W') if it is nonzero for (U, V) .

(c) There exists $\delta_1, \dots, \delta_\ell \geq 0$ with $\sum_{i=1}^\ell \delta_i = k$ and for all $J \subseteq [n]$ nonempty, we have that

$$\sigma_J + |A_J| \leq k - \sum_{i \in J} \delta_i.$$

By adapting Theorem 2.2, we can prove that conditions (b) and (c) are equivalent.

Proposition 3.6. *Let $(\sigma_1, A_1), \dots, (\sigma_\ell, A_\ell)$ be tuples such that for all $i \in [\ell]$, we have that $|\sigma_i| + |A_i| \leq k$. Then, conditions (b) and (c) of Definition 3.2 are equivalent.*

Proof. Let $b = \max_{i \in [\ell]} \sigma_i$. Let $B = \{i_1, \dots, i_b\}$ be a set of size σ disjoint from each of the A_i 's. As in the proof of Proposition 3.5, for all $i \in [\ell]$, define

$$C_i = A_i \cup \{i_1, \dots, i_{\sigma_i}\}.$$

It is easy to see that for all $J \subseteq [\ell]$ nonempty that $|C_J| = |\sigma_J| + |A_J|$. Thus it suffices to show that the following two statements are equivalent:

(b) For all partitions $P_1 \sqcup \dots \sqcup P_s = [\ell]$, we have that

$$\sum_{i=1}^s |C_{P_i}| \leq (s-1)k.$$

(c) There exists $\delta_1, \dots, \delta_\ell \geq 0$ with $\sum_{i=1}^\ell \delta_i = k$ and for all $J \subseteq [n]$ nonempty, we have that

$$|C_J| \leq k - \sum_{i \in J} \delta_i.$$

This equivalence follows immediately from the (b) \iff (c) equivalence of Theorem 2.2 with $d = 0$. \square

By adapting a result of [BGM22] (c.f., their Lemma B.5), we can show that any proper configuration can be “mutated” into a maximal configuration. Given $(\sigma_1, A_1), \dots, (\sigma_\ell, A_\ell)$, we say that $(\sigma'_1, A'_1), \dots, (\sigma'_\ell, A'_\ell)$ is a *contraction* if $\sigma_i \leq \sigma'_i$ and $A'_i \subseteq A_i$ for all $i \in [\ell]$. We say this contraction is *minimal* if $\sum_{i=1}^\ell (\sigma_i - \sigma'_i) + (|A_i| - |A'_i|) = 1$. Likewise, we say that $(\sigma'_1, A'_1), \dots, (\sigma'_\ell, A'_\ell)$ is an *expansion* if $\sigma_i \geq \sigma'_i$ and $A'_i \supseteq A_i$ for all $i \in [\ell]$. We say this expansion is *minimal* if $\sum_{i=1}^\ell (\sigma'_i - \sigma_i) + (|A'_i| - |A_i|) = 1$.

Lemma 3.7. *Let $(\sigma_1, A_1), \dots, (\sigma_\ell, A_\ell)$ be a proper configuration with $d = \text{gid}_k((\sigma_1, A_1), \dots, (\sigma_\ell, A_\ell))$. Then,*

(a) *If $d \geq 1$, then there is a minimal contraction $(\sigma'_1, A'_1), \dots, (\sigma'_\ell, A'_\ell)$ with*

$$d - 1 = \text{gid}_k((\sigma'_1, A'_1), \dots, (\sigma'_\ell, A'_\ell)).$$

(b) *If $d = 0$ and $\sum_{i=1}^\ell \sigma_i + |A_i| < (\ell - 1)k$, then there is a minimal expansion $(\sigma'_1, A'_1), \dots, (\sigma'_\ell, A'_\ell)$ with*

$$0 = \text{gid}_k((\sigma'_1, A'_1), \dots, (\sigma'_\ell, A'_\ell)).$$

As the details of this proof do not deviate substantially from those of [BGM22], we leave the proof in the Appendix A. We thus, have the following generalization of Proposition 3.4.

Lemma 3.8. *Let (U, V) be a (n, k, b) -MDSb(1) code. The following are equivalent.*

(a) (U, V) is MDSb(ℓ).

(b) For all maximal configurations $(\sigma_1, A_1), \dots, (\sigma_\ell, A_\ell)$, we have that

$$\dim \left(\bigcap_{i=1}^{\ell} (U_{\leq \sigma_i} + V_{A_i}) \right) = 0. \quad (11)$$

Proof. Note that (a) implies (b) follows immediately from Proposition 3.5 and condition (b) of Definition 3.2. We now focus on showing that (b) implies (a).

Assume for sake of contradiction there exists a proper configuration $(\sigma_1, A_1), \dots, (\sigma_\ell, A_\ell)$ such that (8) is violated. Since (U, V) is MDSb(1), by Corollary 3.3, there are two possibilities

1. $\text{gid}_k((\sigma_1, A_1), \dots, (\sigma_\ell, A_\ell)) = d \geq 1$ but $\dim \left(\bigcap_{i=1}^{\ell} (U_{\leq \sigma_i} + V_{A_i}) \right) \geq d + 1$.
2. $\text{gid}_k((\sigma_1, A_1), \dots, (\sigma_\ell, A_\ell)) = 0$, but $\dim \left(\bigcap_{i=1}^{\ell} (U_{\leq \sigma_i} + V_{A_i}) \right) \geq 1$.

In Case 1, we apply part (a) of Lemma 3.7 to find a minimal contraction $(\sigma'_1, A'_1), \dots, (\sigma'_\ell, A'_\ell)$ with $\text{gid}_k((\sigma'_1, A'_1), \dots, (\sigma'_\ell, A'_\ell)) = d - 1$. Since the contraction is minimal, by Proposition A.2, we have that

$$\dim \left(\bigcap_{i=1}^{\ell} (U_{\leq \sigma'_i} + V_{A'_i}) \right) \geq \dim \left(\bigcap_{i=1}^{\ell} (U_{\leq \sigma_i} + V_{A_i}) \right) - 1 \geq d,$$

where $Y = (U_{\leq \sigma_i} + V_{A_i})$ and $Y' = U_{\leq \sigma'_i} + V_{A'_i}$ for the pair which changed and X is the intersection of the remaining spaces. By repeating this argument d times, we eventually reach a proper configuration $(\sigma''_1, A''_1), \dots, (\sigma''_\ell, A''_\ell)$ for which $\text{gid}_k((\sigma''_1, A''_1), \dots, (\sigma''_\ell, A''_\ell)) = 0$ but $\dim \left(\bigcap_{i=1}^{\ell} (U_{\leq \sigma''_i} + V_{A''_i}) \right) \geq 1$. Thus, we may now assume we are in Case 2.

For Case 2, by Proposition 3.5, we must have that $\sum_{i=1}^{\ell} \sigma''_i + |A''_i| \leq (\ell - 1)k$. However, if we have equality, then we violate our assumption (b). Thus, we may assume the inequality is strict. We then apply part (b) of Lemma 3.7 to find a minimal expansion $(\sigma'_1, A'_1), \dots, (\sigma'_\ell, A'_\ell)$ with $\text{gid}_k((\sigma'_1, A'_1), \dots, (\sigma'_\ell, A'_\ell)) = 0$. Since we performed an expansion,

$$\dim \left(\bigcap_{i=1}^{\ell} (U_{\leq \sigma'_i} + V_{A'_i}) \right) \geq \dim \left(\bigcap_{i=1}^{\ell} (U_{\leq \sigma''_i} + V_{A''_i}) \right) \geq 1.$$

Thus, we still lie in Case 2. We repeat until we reach a maximal configuration $(\sigma''_1, A''_1), \dots, (\sigma''_\ell, A''_\ell)$ for which $\dim \left(\bigcap_{i=1}^{\ell} (U_{\leq \sigma''_i} + V_{A''_i}) \right) \geq 1$, a contradiction. \square

3.4 Inductive Framework

Assume we are seeking to prove that (U, V) is MDSb(ℓ) for $\ell \geq 2$, and assume we have already established that (U, V) is MDSb(1). By Lemma 3.8 and Proposition 3.6, it suffices to consider tuples $(\sigma_1, A_1, \delta_1), \dots, (\sigma_\ell, A_\ell, \delta_\ell)$ which satisfy the following conditions:

- (a*) For all $i \in [\ell]$, $\sigma_i, \delta_i \geq 0$
- (b*) $\sum_{i=1}^{\ell} \sigma_i + |A_i| = (\ell - 1)k$
- (c*) $\sum_{i=1}^{\ell} \delta_i = k$
- (d*) For all $J \subseteq [\ell]$ nonempty,

$$\sigma_J + |A_J| \leq k - \sum_{i \in J} \delta_i \quad (12)$$

Note that conditions (b*), (c*), and (d*) together imply a fifth condition:

(e*) For all $i \in [\ell]$, that $\sigma_i + |A_i| + \delta_i = k$.

Note by (12) with $J = [\ell]$, we have that $\sigma_i = 0$ for at least one i , and for all $j \in [n]$ there is at least one $i \in [\ell]$ such that $j \notin A_i$.

For each set of tuples satisfying the previous constraints, we seek to show that

$$\dim \left(\bigcap_{i=1}^{\ell} (U_{\leq \sigma_i} + V_{A_i}) \right) = 0. \quad (13)$$

By Lemma 2.3, we have that (13) is equivalent to

$$\det \begin{pmatrix} I_k & U|_{\leq \sigma_1} & V|_{A_1} & & & \\ I_k & & & U|_{\leq \sigma_2} & V|_{A_2} & \\ \vdots & & & & & \ddots \\ I_k & & & & & & U|_{\leq \sigma_\ell} & V|_{A_\ell} \end{pmatrix} \neq 0 \quad (14)$$

As appears in the original proofs [Lov21, YH19] of the GM-MDS Theorem, an important special case is when there exists $J \subseteq [\ell]$ with $2 \leq |J| \leq \ell - 1$ for which (12) is tight. It turns out these cases follow only from the fact that (U, V) is MDSb($\ell - 1$).

Lemma 3.9. *Let (U, V) be MDSb($\ell - 1$) for $\ell \geq 2$. Let $(\sigma_1, A_1, \delta_1), \dots, (\sigma_\ell, A_\ell, \delta_\ell)$ be a configuration satisfying conditions (a*)-(d*). Further assume there exists $J \subseteq [\ell]$ with $2 \leq |J| \leq \ell - 1$ for which (12) is tight. Then, (13) holds.*

Proof. Let J^* be a specific set of size between 2 and $n - 1$ for which (12) is tight. Like in Case 1 of the proof of the GM-MDS theorem, we can verify that for generic matrices $W \in \mathbb{F}^{k \times n}$ and $W' \in \mathbb{F}^{n \times n}$, we have that

$$\bigcap_{i \in J^*} (W_{\leq \sigma_i} + W'_{A_i}) = W_{\leq \sigma_{J^*}} + W'_{A_{J^*}} \quad (15)$$

$$(W_{\leq \sigma_{J^*}} + W'_{A_{J^*}}) \cap \bigcap_{i \notin J^*} (W_{\leq \sigma_i} + W'_{A_i}) = 0. \quad (16)$$

In particular, to check (15), the \supseteq inclusion is straightforward, so we just need to check both sides have the same dimension. For any partition $P_1 \sqcup \dots \sqcup P_s = J^*$ we have that by (12) that

$$\begin{aligned} -n(s-1) + \sum_{i=1}^s (\sigma_{P_i} + |A_{P_i}|) &\leq -k(s-1) + \sum_{i=1}^s \left[n - \sum_{j \in P_i} \delta_j \right] \\ &= k - \sum_{j \in J^*} \delta_j \\ &= |\sigma_{J^*}| + |A_{J^*}|, \end{aligned} \quad (\text{assumption on } J^*)$$

where equality is achieved if $s = 1$.

To check (16), it suffices to check (12) (and invoke Proposition 3.6) for the tuple of sets $\{(\sigma_i, A_i, \delta_i) : i \notin J^*\} \cup \{(\sigma_{J^*}, A_{J^*}, \delta_{J^*})\}$, where $\delta_{J^*} := \sum_{i \in J^*} \delta_i$. The intersections that do not include $(\sigma_{J^*}, A_{J^*}, \delta_{J^*})$ follow by assumption on the structure of the original configuration. Otherwise, for any $J' \subseteq J^*$ (possibly empty), we need to check that

$$\sigma_{J' \cup J^*} + |A_{J' \cup J^*}| \leq k - \sum_{i \in J'} \delta_i + \delta_{J^*} = k - \sum_{i \in J' \cup J^*} \delta_i,$$

which again follows from assumption.

To conclude this case, by the induction hypothesis, we have that (U, V) is MDSb(ℓ') for all $\ell' < \ell$. Thus, we have that both (15) and (16) hold with (W, W') replaced by (U, V) . Note that for (15) we are using the fact that the \supseteq direction holds unconditionally. Therefore, (13) holds. \square

We use this inductive template in both Section 4 and Section 5 to prove various generalizations of GM-MDS.

4 Algebraic GM-MDS for Monomial Codes

The goal of this section is to prove the following result. In Section 6, we shall show how to extend this result to irreducible varieties: Theorem 6.4.

Theorem 4.1. *Let $e_1 < e_2 < e_3 < \dots < e_k$ be integers. Let $\alpha_1, \dots, \alpha_n$ be generic points in an algebraically closed field \mathbb{F} . Then, for all $\ell \geq 2$, the matrix*

$$V := \begin{pmatrix} \alpha_1^{e_1} & \alpha_2^{e_1} & \dots & \alpha_n^{e_1} \\ \alpha_1^{e_2} & \alpha_2^{e_2} & \dots & \alpha_n^{e_2} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_1^{e_k} & \alpha_2^{e_k} & \dots & \alpha_n^{e_k} \end{pmatrix} \quad (17)$$

is MDS(ℓ)

By Proposition 3.1, it suffices to show that (I, V) is MDSb(ℓ) for all $\ell \geq 1$. We proceed by the inductive approach of Section 3.4.

Consider a configuration $(\sigma_1, A_1, \delta_1), \dots, (\sigma_\ell, A_\ell, \delta_\ell)$ satisfying conditions (a*)-(e*) in Section 3.4. We seek to show (13) that

$$\dim \left(\bigcap_{i=1}^{\ell} (V_{\leq \sigma_i}^T + I_{A_i}) \right) = 0.$$

We shall prove this by induction on ℓ , k and $\sum_{i=1}^{\ell} |A_i|$. We first consider a few base cases.

Base cases. For the base case of $\ell = 1$, it suffices to prove that for all $\sigma \in \{0, 1, \dots, k\}$, we have that

$$\det \begin{pmatrix} 1 & 0 & \dots & \alpha_1^{e_1} & \alpha_2^{e_1} & \dots & \alpha_{k-\sigma}^{e_1} \\ 0 & 1 & \dots & \alpha_1^{e_2} & \alpha_2^{e_2} & \dots & \alpha_{k-\sigma}^{e_2} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \alpha_1^{e_k} & \alpha_2^{e_k} & \dots & \alpha_{k-\sigma}^{e_k} \end{pmatrix} \neq 0,$$

where the first σ columns are standard basis vectors. This follows from the fact that the coefficient of $\alpha_1^{e_{\sigma+1}} \dots \alpha_{k-\sigma}^{e_k}$ in the determinant expansion is ± 1 .

The base case of $k = 1$ is obvious, as the generator matrix of any $(n, 1)$ -code with every entry nonzero is MDS(1) and thus MDS(ℓ) for all ℓ (c.f., [BGM22]).

The final base case is $\sum_{i=1}^{\ell} |A_i| = 0$, or each $A_i = \emptyset$. In this case,

$$\dim \left(\bigcap_{i=1}^{\ell} V_{\leq \sigma_i}^T \right) = \dim V_{\leq \sigma_{[n]}}^T = \sigma_{[n]},$$

as desired.

Inductive step. We now divide into three cases.

1. There is $J \subseteq [\ell]$ with $2 \leq |J| \leq \ell - 1$ for which (12) is tight.
2. $\sigma_i = 0$ for exactly one $i \in [\ell]$.
3. None of the above.

Case 1. This case follows immediately from Lemma 3.9.

Case 2. Without loss of generality we may assume that $\sigma_1, \dots, \sigma_{\ell-1} \geq 1$ and $\sigma_\ell = 0$. Recall we seek to verify that

$$\det \begin{pmatrix} I_k & I_{k|\leq\sigma_1} & V|_{A_1} & & & & \\ I_k & & & I_{k|\leq\sigma_2} & V|_{A_2} & & \\ \vdots & & & & & \ddots & \\ I_k & & & & & & I_{k|\leq\sigma_{\ell-1}} & V|_{A_{\ell-1}} \\ I_k & & & & & & & V|_{A_\ell} \end{pmatrix} \neq 0$$

Let V' be V with the first row removed. We can expand the determinant along the $\ell-1$ columns corresponding to the first column of I_k . We can then also expand along the leftmost column. It is thus equivalent to check whether

$$\det \begin{pmatrix} I_{k-1} & I_{k-1|\leq\sigma_1-1} & V'|_{A_1} & & & & \\ I_{k-1} & & & I_{k-1|\leq\sigma_2-1} & V'|_{A_2} & & \\ \vdots & & & & & \ddots & \\ I_{k-1} & & & & & & I_{k-1|\leq\sigma_{\ell-1}-1} & V'|_{A_{\ell-1}} \\ I_{k-1} & & & & & & & V'|_{A_\ell} \end{pmatrix} \neq 0. \quad (18)$$

Consider $(\sigma'_1, A'_1, \delta'_1), \dots, (\sigma'_\ell, A'_\ell, \delta'_\ell)$, where for $i \in [\ell]$, we have that $(\sigma'_i, A'_i, \delta'_i) = (\sigma_i - 1, A_i, \delta_i)$ and $(\sigma'_\ell, A'_\ell, \delta'_\ell) = (0, A_\ell, \delta_\ell - 1)$. It is straightforward to check that: (a*), (b*), (c*) hold in $k-1$ dimensional space. For (d*), note that for any $J \subseteq [\ell-1]$ nonempty, we have that

$$\sigma'_J + |A'_J| = \sigma_J - 1 + |A_J| \leq k-1 - \sum_{i \in J} \delta_i = k-1 - \sum_{i \in J} \delta'_i.$$

Thus, by the induction hypothesis, we can deduce that (14) holds for (I_{k-1}, V') , which is precisely (18).

Case 3. We may assume that $\sigma_{\ell-1} = \sigma_\ell = 0$. By (e*) of the inductive framework, we know then that $|A_{\ell-1}| = k - \delta_{\ell-1}$ and $|A_\ell| = k - \delta_\ell$. Further by (d*), we have that $|A_{\ell-1} \cap A_\ell| \leq k - \delta_\ell - \delta_{\ell-1}$. In particular, we must have that $\delta_{\ell-1} = \delta_\ell = 0$ or $A_{\ell-1} \neq A_\ell$. In the former case, it is straightforward to check that $(\sigma_1, A_1, \delta'_1), \dots, (\sigma_{\ell-1}, A_{\ell-1}, \delta_{\ell-1})$ satisfy conditions (a*)-(d*), and so by the induction hypothesis

$$\bigcap_{i=1}^{\ell} (I_{\leq\sigma_i} + V_{A_i}) = \bigcap_{i=1}^{\ell-1} (I_{\leq\sigma_i} + V_{A_i}) = 0.$$

In the latter case, we may without loss of generality assume there is $j^* \in A_{\ell-1} \setminus A_\ell$. Now define $(\sigma'_1, A'_1, \delta'_1), \dots, (\sigma'_\ell, A'_\ell, \delta'_\ell)$ as follows. For all $i \in [\ell]$, let

$$(\sigma'_i, A'_i, \delta'_i) = \begin{cases} (\sigma_i, A_i, \delta_i) & j^* \notin A_i \\ (\sigma_i + 1, A_i \setminus \{j^*\}, \delta_i) & \text{otherwise.} \end{cases}$$

It is clear this new system satisfies conditions (a*)-(c*) of the inductive framework. Further, (d*) holds if $|J| = 1$ or $|J| = \ell$. Now assume $2 \leq |J| \leq \ell - 1$, observe that

$$\sigma'_J + |A'_J| \leq \sigma_J + |A_J| + 1 \leq k - \sum_{i \in J} \delta_i = k - \sum_{i \in J} \delta'_i,$$

where the middle inequality follows from us not being in Case 1. Note that $\sum_{i=1}^{\ell} |A'_i| < \sum_{i=1}^{\ell} |A_i|$. Therefore, by the induction hypothesis, we have that

$$\bigcap_{i=1}^{\ell} (I_{k|\leq\sigma'_i} + V_{A'_i}) = 0 \quad (19)$$

To finish, it suffices to prove the following proposition

5.1 Reduction

Our plan for proving Theorem 5.1 is to (non-obviously) reduce the statement to the following fact.

Theorem 5.2. *Let V be as in (17), then (V^T, I_n) is MDSb(ℓ) for all $\ell \geq 1$.*

Rather unintuitively, the basis is the rows of V^T and the “normal” part is I_n . We shall explain how this structure arises shortly, but first we shall prove the following fact.

Proposition 5.3. *Let V be an (n, k) -MDS matrix. Then, V^\perp is MDS(ℓ) if and only if for any $A_1, \dots, A_\ell \subseteq [n]$ with the dimension $n - k$ null intersection property, $\sum_{i=1}^\ell |A_i| = (\ell - 1)(n - k)$, and each set has size at most $n - k$, we have that*

$$\begin{pmatrix} I|_{\bar{A}_1} & I|_{\bar{A}_2} & \cdots & I|_{\bar{A}_\ell} \\ V|_{\bar{A}_1} & & & \\ & V|_{\bar{A}_2} & & \\ & & \ddots & \\ & & & V|_{\bar{A}_\ell} \end{pmatrix} \quad (22)$$

has rank $n + (\ell - 1)k$, where $\bar{A}_i = [n] \setminus A_i$.

Note that (22) has $n + (\ell - 1)k$ rows but $n + \ell k$ columns.

Proof. Let $H \in \mathbb{F}^{(\ell-1) \times \ell}$ be a parity check matrix. By Theorem 1.3, we have that V^\perp is MDS(ℓ) if and only if $H \otimes V^\perp$ is an $(\ell, n, 1, k)$ -MR tensor code. More precisely, we have that $V_{A_1} \cap \cdots \cap V_{A_\ell} = 0$ if and only if the erasure pattern $E = \bigcup_{i=1}^\ell \{i\} \times \bar{A}_i$ is recoverable for $H \otimes V^\perp$. Note that since we assume that $\sum_{i=1}^\ell |A_i| = (\ell - 1)(n - k)$, we have that

$$|E| = \sum_{i=1}^\ell |\bar{A}_i| = n\ell - (\ell - 1)(n - k) = n + (\ell - 1)k.$$

Observe that the all-ones row vector 1^T is the parity check matrix of H and V is the parity-check matrix of V^\perp . Consider the matrix

$$\begin{pmatrix} 1^T \otimes I_n \\ I_\ell \otimes V \end{pmatrix} = \begin{pmatrix} I_n & I_n & \cdots & I_n \\ V & & & \\ & V & & \\ & & \ddots & \\ & & & V \end{pmatrix} \quad (23)$$

Note that every row of (23) is a parity-check condition for the code $H \otimes V^\perp$. Further the rows of (23) span the parity check conditions of $H \otimes V^\perp$. Thus, the pattern E is recoverable if and only if the columns of (23) corresponding to E are linearly independent (c.f., [BGM22]). Observe that (23) restrict to the columns indexed by E is precisely (22). Thus, (22) has rank $|E| = n + (\ell - 1)k$ if and only if $V_{A_1} \cap \cdots \cap V_{A_\ell} = 0$, as desired. \square

We now show how Theorem 5.2 and Proposition 5.3 complete the proof of Theorem 5.1. This follows from the following theorem, and that fact that V is MDS, as proved in the base case of the proof of Theorem 4.1.

Theorem 5.4. *Let V be an (n, k) -MDS matrix such that (V^T, I_n) is MDSb(ℓ). Then, V^\perp is MDS(ℓ). That is, V is LD-MDS($\leq \ell - 1$).*

Proof of Theorem 5.4. Consider $A_1, \dots, A_\ell \subseteq [n]$ with the dimension $n - k$ null intersection property, $\sum_{i=1}^{\ell} |A_i| = (\ell - 1)(n - k)$, and each set has size at most $n - k$. We seek to show that (22) has rank $n + (\ell - 1)k$. Consider the proper order- ℓ configuration $(\sigma_1, A_1), \dots, (\sigma_\ell, A_\ell)$ where each $\sigma_i = k$. Observe that for any partition $P_1 \sqcup \dots \sqcup P_s = [\ell]$, we have that

$$\begin{aligned} -n(s-1) + \sum_{i=1}^s (\sigma_{P_i} + |A_{P_i}|) &= -n(s-1) + ks + \sum_{i=1}^s |A_{P_i}| \\ &= k + \left[-(n-k)(s-1) + \sum_{i=1}^s |A_{P_i}| \right] \\ &\geq k, \end{aligned}$$

where equality for some partition as A_1, \dots, A_ℓ have the dimension $n - k$ null intersection property. Therefore, by Proposition 3.5 and Theorem 5.2, we have that

$$\dim \left(\bigcap_{i=1}^{\ell} (V_{\leq k}^T + I_{A_i}) \right) = k.$$

Equivalently, by Lemma 2.3, we have that the matrix

$$\begin{pmatrix} I_n & V^T & I|_{A_1} & & & & \\ I_n & & & V^T & I|_{A_2} & & \\ \vdots & & & & & \ddots & \\ I_n & & & & & & V^T & I|_{A_\ell} \end{pmatrix} \quad (24)$$

has rank

$$\begin{aligned} -k + n + \sum_{i=1}^{\ell} \dim(V^T + I_{A_i}) &= (n - k) + (\ell - 1)(n - k) + \ell k \\ &= n\ell, \end{aligned}$$

where for the first line we use Proposition 3.2 and the fact that (V^T, I) is MDSb(1).

Each $I|_{A_i}$ lets us eliminate $|A_i|$ rows. Doing this and permutating some rows/columns, we get that (24) has the same rank as

$$\left[\begin{array}{cccc|cccc} I|_{\bar{A}_1 \times [n]} & V^T|_{\bar{A}_1 \times [k]} & & & & & & \\ I|_{\bar{A}_2 \times [n]} & & V^T|_{\bar{A}_2 \times [k]} & & & & & \\ \vdots & & & \ddots & & & & \\ I|_{\bar{A}_\ell \times [n]} & & & & V^T|_{\bar{A}_\ell \times [k]} & & & \\ \hline & & & & & I|_{A_1 \times A_1} & & \\ & & & & & & \ddots & \\ & & & & & & & I|_{A_\ell \times A_\ell} \end{array} \right] \quad (25)$$

Since the bottom-right submatrix of (25) is an identity matrix, we have that the top-left submatrix of (25) has rank $n\ell - \sum_{i=1}^{\ell} |A_i| = n\ell - (n - k)(\ell - 1) = n + k(\ell - 1)$. Since the top-left submatrix of (25) is the transpose of (22), we have that V^\perp is MDS(ℓ) by Proposition 5.3, where we use that V is MDS. \square

5.2 Proof of Theorem 5.2

In this section, we prove that (V^T, I_n) is $\text{MDSb}(\ell)$. We proceed by the inductive approach of Section 3.4. However there is a key difference between this proof and the others present so far. Consider a configuration $(\sigma_i, A_i, \delta_i)$ which occurs in the induction. The GM-MDS theorem (and Theorem 6.4) seeks to “shrink” the A_i ’s and grow the corresponding σ_i ’s. In our case, we will shrink the σ_i ’s and grow the A_i ’s. In the end, when $\sigma_i = 0$ for all $i \in [\ell]$, we can directly compute the intersection dimension, as I_n is just the standard basis and thus is $\text{MDS}(\ell)$.

Consider a configuration $(\sigma_1, A_1, \delta_1), \dots, (\sigma_\ell, A_\ell, \delta_\ell)$ satisfying conditions (a*)-(e*) in Section 3.4. We seek to show (13) that

$$\dim \left(\bigcap_{i=1}^{\ell} (V_{\leq \sigma_i}^T + I_{A_i}) \right) = 0.$$

We shall prove this by induction on ℓ and $\sum_{i=1}^{\ell} \sigma_i$. We first consider a few base cases.

Base cases. For the base of $\ell = 1$, we prove the stronger statement that $[V^T \mid I_n]$ is MDS. Pick a subset of n columns of this matrix, and expand the determinant along the columns corresponding to I_n . Up to sign, we are left with an expression of the general form

$$\det \begin{pmatrix} \alpha_1^{e_1} & \alpha_1^{e_2} & \cdots & \alpha_1^{e_\sigma} \\ \alpha_2^{e_1} & \alpha_2^{e_2} & \cdots & \alpha_2^{e_\sigma} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_\sigma^{e_1} & \alpha_\sigma^{e_2} & \cdots & \alpha_\sigma^{e_\sigma} \end{pmatrix} \neq 0,$$

which follows from the proof of Theorem 4.1.

For the other base case of $\sum_{i=1}^{\ell} \sigma_i = 0$, we have that $\sigma_i = 0$ for all i . In that case, we have that

$$\begin{aligned} \dim \left(\bigcap_{i=1}^{\ell} (V_{\leq \sigma_i}^T + I_{A_i}) \right) &= \dim \left(\bigcap_{i=1}^{\ell} I_{A_i} \right) \\ &= \dim I_{A_{[\ell]}} = |A_{[\ell]}| \\ &= \max_{P_1 \sqcup \cdots \sqcup P_s = [\ell]} \left[-n(s-1) + \sum_{i=1}^s (\sigma_{P_i} + |A_{P_i}|) \right], \end{aligned}$$

where the last step uses the partition $P_1 = [\ell]$, as desired.

Inductive step. We now divide into three cases.

1. There is J with $2 \leq |J| \leq \ell - 1$ for which (12) is tight.
2. $\sigma_i = \sigma_{i'} = 0$ for some $i \neq i'$
3. None of the above.

Case 1. This case follows immediately from Lemma 3.9.

Case 2. Without loss of generality assume that $\sigma_{\ell-1} = \sigma_\ell = 0$. Note that

$$\bigcap_{i=1}^{\ell} (V_{\leq \sigma_i}^T + I_{A_i}) = \bigcap_{i=1}^{\ell-2} (V_{\leq \sigma_i}^T + I_{A_i}) \cap I_{A_{\ell-1}} \cap I_{A_\ell}. \quad (26)$$

The RHS corresponds to the configuration $(\sigma_1, A_1, \delta_1), \dots, (\sigma_{\ell-2}, A_2, \delta_{\ell-2}), (0, A_{\ell-1} \cap A_\ell, \delta_{\ell-1} + \delta_i)$. This configuration might not satisfy condition (b*), but it satisfies conditions (a*), (c*), (d*). Thus, by Proposition 3.6 and the induction hypothesis that (V^T, I_n) is MDSb($\ell - 1$), we have that the RHS of (26) equals 0, as desired.

Case 3. Without loss of generality, we may assume that $\sigma_1, \dots, \sigma_{\ell-1} \geq 1$ but $\sigma_\ell = 0$. From (12) with $J = [\ell - 1]$, we have that

$$\begin{aligned} 1 + |A_{[\ell-1]}| &\leq \sigma_{[\ell-1]} + |A_{[\ell-1]}| \\ &\leq n - \sum_{i=1}^{\ell-1} \delta_i \\ &= \delta_\ell \\ &= n - |A_\ell|. \end{aligned}$$

Thus, there exists $j^* \in [n] \setminus (A_\ell \cup A_{[\ell-1]})$. For all $i \in [\ell]$, define $(\sigma'_i, A'_i, \delta'_i)$ as follows

$$(\sigma'_i, A'_i, \delta'_i) = \begin{cases} (\sigma_i, A_i, \delta_i) & j^* \in A_i \text{ or } i = \ell \\ (\sigma_i - 1, A_i \cup \{j^*\}, \delta_i) & \text{otherwise.} \end{cases}$$

We seek to show that $\{(\sigma'_i, A'_i, \delta'_i) : i \in [\ell]\}$ satisfies conditions (a*)–(d*). Clearly conditions (a*)–(c*) are satisfied. To check (d*), we split into cases based on the size of $J \subseteq [\ell]$. If $|J| = 1$, then (12) is still satisfied as $|\sigma'_i| + |A'_i| = |\sigma_i| + |A_i|$ for all $i \in [\ell]$. If $|J| = \ell$, then the LHS of (12) is still 0 as $\sigma_\ell = 0$ and $j^* \notin A_\ell$. Otherwise, we have that

$$\begin{aligned} \sigma'_J + |A'_J| &\leq \sigma_J + |A_J| + 1 && \text{(only } j^* \text{ possibly added)} \\ &\leq n - \sum_{i \in J} \delta_i && \text{(not in Case 1)} \\ &= n - \sum_{i \in J} \delta'_i. \end{aligned}$$

Since $j^* \notin A_{[\ell-1]}$, we must have that $\sum_{i=1}^{\ell} \sigma'_i < \sum_{i=1}^{\ell} \sigma_i$. Therefore, by the induction hypothesis we have that

$$\bigcap_{i=1}^{\ell} (V_{\leq \sigma'_i}^T + I_{A'_i}) = 0.$$

To complete the case, we need to show algebraically this implies (13).

Proposition 5.5. *We have that*

$$\bigcap_{i=1}^{\ell} (V_{\leq \sigma'_i}^T + I_{A'_i}) = 0 \implies \bigcap_{i=1}^{\ell} (V_{\leq \sigma_i}^T + I_{A_i}) = 0. \quad (27)$$

Proof. By Lemma 2.3, we have that the RHS of (27) is equivalent to

$$\det \begin{pmatrix} I_n & V^T|_{\leq \sigma_1} & I|_{A_1} & & & & \\ I_n & & & V^T|_{\leq \sigma_2} & I|_{A_2} & & \\ \vdots & & & & & \ddots & \\ I_n & & & & & & V^T|_{\leq \sigma_\ell} & I|_{A_\ell} \end{pmatrix} \neq 0. \quad (28)$$

A polynomial f is said to be generically non-vanishing on a variety X if the open set $U_f = \{x | f(x) \neq 0, x \in X\}$ is dense for the Zariski topology on V . For an irreducible V that simply means $f \notin I$. Equivalently, there is at least one point in V on which f does not vanish. For a general X , it means that f does not vanish on each of the irreducible components of X .

We need a simple fact about the product of two varieties.

Lemma 6.1 (Product of varieties (See 10.4.H in [Vak24])). *If $X_1 \subseteq \mathbb{F}^{k_1}$ and $X_2 \subseteq \mathbb{F}^{k_2}$ are two varieties then $X_1 \times X_2 \subseteq \mathbb{F}^{k_1+k_2}$ is also a variety. Furthermore, if X_1 and X_2 are irreducible then so is $X_1 \times X_2$.*

In general if X_1 and X_2 have irreducible components $X_{1,1}, \dots, X_{1,m_1}$ and $X_{2,1}, \dots, X_{2,m_2}$ respectively then $X_{1,i} \times X_{2,j}, i \in [m_1], j \in [m_2]$ are the irreducible components of $X_1 \times X_2$.

Lemma 6.2. *For an algebraically closed \mathbb{F} and an irreducible variety $X \subseteq \mathbb{F}^k$, there exists a d (d will be the dimension of X) such that we can find functions f_1, \dots, f_k (depending on X) in the formal power series ring $\mathbb{F}[[z_1, \dots, z_d]]$ such that for any polynomial $g \in \mathbb{F}[x_1, \dots, x_k]$ over \mathbb{F}^k it is generically non-vanishing over X if and only if $g(f_1, \dots, f_k) \in \mathbb{F}[[z_1, \dots, z_d]]$ is non-zero.*

We also have that for $X^n \subseteq \mathbb{F}^{kn}$ any polynomial $g \in \mathbb{F}[x_{1,1}, \dots, x_{1,k}, \dots, x_{n,1}, \dots, x_{n,k}]$ is generically non-vanishing over X^n if and only if $g(f_1(z_1), \dots, f_k(z_1), \dots, f_1(z_n), \dots, f_k(z_n))$ is non-zero.

The above follows from standard facts in algebraic geometry. We give a proof and references in the appendix.

6.2 Algebraic GM-MDS

Definition 6.1 (MDS(ℓ) property for varieties). *For an algebraic closed \mathbb{F} , we say a variety $X \subseteq \mathbb{F}^k$ satisfies the MDS(ℓ) property if and only if for every $k \times n$ matrix V^X with columns of V^X initialized from X , V^X is generically a $[n, k]$ -MDS(ℓ) code.*

Concretely, this means for any $A_1, \dots, A_\ell \subseteq [n]$ with $|A_i| \leq k$ and $|A_1| + \dots + |A_\ell| = (\ell - 1)k$, we have that $V_{A_1} \cap \dots \cap V_{A_\ell} = 0$ if and only if

$$\det \begin{pmatrix} I_k & V^X|_{A_1} & & & \\ I_k & & V^X|_{A_2} & & \\ \vdots & & & \ddots & \\ I_k & & & & V^X|_{A_\ell} \end{pmatrix},$$

is generically non-vanishing on $X^{(\ell-1)k}$, where V is a $k \times n$ generic matrix (that is a matrix whose every entry is an independent formal variable).

A MDS(2) variety we just call MDS.

We also note that being MDS equivalently means that the determinant of a $k \times k$ matrix where each column is a point in \mathbb{F}^k is generically non-vanishing over X^k . This allows us to give another simple characterization of MDS varieties.

Theorem 6.3 (MDS varieties are not contained in hyper-planes passing through the origin). *A variety $X \subseteq \mathbb{F}^k$ is MDS if and only if each of its irreducible components is not contained in any hyper-plane passing through the origin. In particular, an irreducible variety is MDS if and only if it is not contained in any hyperplane passing through the origin.*

Proof. Let f be the determinant of a $k \times k$ matrix M where the i th column is a point $(x_{1,i}, \dots, x_{k,i})$ in \mathbb{F}^k and X have irreducible components X_1, \dots, X_m . The irreducible components of X^k are simply $X_{j_1} \times \dots \times X_{j_k}$ with $j_1, \dots, j_k \in [m]$.

Say some irreducible component X_i of X lies in a hyperplane passing through the origin. Then f vanishes on $(X_i)^k$ which means X is not MDS (because M has a kernel corresponding to the hyperplane).

Say X is not MDS which means f vanishes on some $Y = X_{j_1} \times \dots \times X_{j_k}$. We now consider the determinants of the sub-matrices of M . Our goal will be to show some X_{j_i} lies in a hyperplane passing through the origin which would complete the proof.

If any of the co-ordinate functions (say $x_{1,1}$) (corresponding to size 1 sub-matrices) vanishes on Y then X_{j_1} lies in the co-ordinate hyperplane $x_{1,1} = 0$. Say the largest-size sub-determinant which does not vanish on Y has size $t > 0$. This means the determinant of some $t \times t$ sub-matrix M_1 of M does not vanish on Y and all the determinants of $(t+1) \times (t+1)$ sub-matrices of M vanish on Y . Pick a $(t+1) \times (t+1)$ sub-matrix M_2 of M containing M_1 . Without loss of generality we can assume the columns of M_1 come from X_{j_1}, \dots, X_{j_t} and the extra column in M_2 comes from $X_{j_{t+1}}$ (the choice of rows is not important). We know that $\det(M_2)$ vanishes on Y . Let R be the index of rows in M_2 . Expanding the determinant of M_2 along the $t+1$ th column will give us

$$\det(M_2) = \sum_{i \in R} x_{i,t+1} \det(M'_2(i)),$$

where $M'_2(i)$ is the sub-matrix of M_2 obtained by removing the i th row and $t+1$ th column. Note, $M'_2(t+1) = M_1$ which does not vanish on Y . This means it is possible to find points in $y_1 \in X_{j_1}, \dots, y_t \in X_{j_t}$ such that $g = \det(M_2)(x_1 = y_1, \dots, x_t = y_t)$ is a non-zero polynomial. But as $\det(M_2)$ vanishes on Y so does g . But that implies $X_{j_{t+1}}$ is contained in $g = 0$ which is a hyperplane passing through the origin. \square

We also generalize the definition of LD-MDS($\leq L$) to the setting of varieties.

Definition 6.2 (LD-MDS($\leq \ell$) property for varieties). *For an algebraic closed \mathbb{F} , we say a variety $X \subseteq \mathbb{F}^k$ satisfies the LD-MDS($\leq \ell$) property if and only if for ever $k \times n$ matrix V^X with columns of V^X initialized from X , the dual of V^X is generically a $[n, n-k]$ -MDS($\ell+1$) code.*

Concretely this means that for any $A_1, \dots, A_{\ell+1} \subseteq [n]$ with $|A_i| \leq n-k$ and $|A_1| + \dots + |A_{\ell+1}| = \ell(n-k)$, we have that $V_{A_1} \cap \dots \cap V_{A_{\ell+1}} = 0$ if and only if

$$\det \begin{pmatrix} I_{n-k} & (V^X)^\perp|_{A_1} & & & \\ I_{n-k} & & (V^X)^\perp|_{A_2} & & \\ \vdots & & & \ddots & \\ I_{n-k} & & & & (V^X)^\perp|_{A_{\ell+1}} \end{pmatrix},$$

is generically non-vanishing on $X^{\ell(n-k)}$ where V is a $n-k \times n$ generic matrix.

Our main theorems for varieties are the following.

Theorem 6.4. *For an algebraic closed \mathbb{F} if an irreducible X is MDS then it is MDS(ℓ) for all $\ell > 2$.*

Theorem 6.5. *For an algebraic closed \mathbb{F} if an irreducible X is MDS then it is LD-MDS($\leq \ell$) for all $\ell > 2$.*

6.3 Reduction to Power Series

We want to reduce the problem to a problem about power series.

Definition 6.3 (MDS(ℓ)-power series). *Let V be $k \times (\ell-1)k$ generic matrix (that is a matrix whose every entry is an independent formal variable). Given a tuple $F = (f_1, \dots, f_k) \in (\mathbb{F}[[z_1, \dots, z_d]])^k$, let $V^F = [F(\mathbf{z}_1), \dots, F(\mathbf{z}_{(\ell-1)k})]$ be a $k \times (\ell-1)k$ matrix where $\mathbf{z}_1, \dots, \mathbf{z}_{(\ell-1)k} = (z_{i,1}, \dots, z_{i,d}), \dots, \mathbf{z}_{(\ell-1)k}$ are distinct d tuple of variables.*

F is said to MDS(ℓ) if and only if for any $A_1, \dots, A_\ell \subseteq [(\ell-1)k]$ with $|A_i| \leq k$ and $|A_1| + \dots + |A_\ell| = (\ell-1)k$, we have that $V_{A_1} \cap \dots \cap V_{A_\ell} = 0$ if and only if

$$\det \begin{pmatrix} I_k & V^F|_{A_1} & & & \\ I_k & & V^F|_{A_2} & & \\ \vdots & & & \ddots & \\ I_k & & & & V^F|_{A_\ell} \end{pmatrix} \neq 0,$$

A MDS(2) power series we just call MDS.

It is easy to show that $F = (f_1, \dots, f_k) \in (\mathbb{F}[[z_1, \dots, z_d]])^k$ is MDS if and only if

$$\text{Det} \begin{pmatrix} f_1(\mathbf{z}_1) & f_1(\mathbf{z}_2) & \cdots & f_1(\mathbf{z}_n) \\ f_2(\mathbf{z}_1) & f_2(\mathbf{z}_2) & \cdots & f_2(\mathbf{z}_n) \\ f_3(\mathbf{z}_1) & f_3(\mathbf{z}_2) & \cdots & f_3(\mathbf{z}_n) \\ \vdots & \vdots & \ddots & \vdots \\ f_k(\mathbf{z}_1) & f_k(\mathbf{z}_2) & \cdots & f_k(\mathbf{z}_n) \end{pmatrix} \neq 0,$$

where $\mathbf{z}_1, \dots, \mathbf{z}_i = (z_{i,1}, \dots, z_{i,d}), \dots, \mathbf{z}_k$ are distinct d tuples of variables.

Definition 6.4 (LD-MDS($\leq \ell$)-power series). *Let V be $(n-k) \times \ell(n-k)$ generic matrix (that is a matrix whose every entry is an independent formal variable). Given a tuple $F = (f_1, \dots, f_k) \in (\mathbb{F}[[z_1, \dots, z_d]])^k$, let $V^F = [F(\mathbf{z}_1), \dots, F(\mathbf{z}_{\ell-1,k})]$ be a $k \times (\ell-1)k$ matrix where $\mathbf{z}_1, \dots, \mathbf{z}_i = (z_{i,1}, \dots, z_{i,d}), \dots, \mathbf{z}_{\ell-1,k}$ are distinct d tuple of variables.*

F is said to LD-MDS($\leq \ell$) if and only if for any $A_1, \dots, A_{\ell+1} \subseteq [\ell(n-k)]$ with $|A_i| \leq n-k$ and $|A_1| + \dots + |A_{\ell+1}| = \ell(n-k)$, we have that $V_{A_1} \cap \dots \cap V_{A_{\ell+1}} = 0$ if and only if

$$\det \begin{pmatrix} I_{n-k} & (V^F)^\perp|_{A_1} & & & \\ I_{n-k} & & (V^F)^\perp|_{A_2} & & \\ \vdots & & & \ddots & \\ I_{n-k} & & & & (V^F)^\perp|_{A_{\ell+1}} \end{pmatrix} \neq 0,$$

The main theorems about power series is the following.

Theorem 6.6. *For a tuple $F = (f_1, \dots, f_k) \in (\mathbb{F}[[z_1, \dots, z_d]])^k$, F is MDS if and only if F is MDS(ℓ) for all $\ell > 2$.*

Theorem 6.7. *For an algebraically closed \mathbb{F} if an irreducible X is MDS then it is LD-MDS($\leq \ell$) for all $\ell > 2$.*

Proof of Theorem 6.4 assuming Theorem 6.6. Recall, \mathbb{F} is algebraically closed. By Lemma 6.2 given an irreducible variety $X \subseteq \mathbb{F}^k$ of dimension d we have elements f_1, \dots, f_k in the power series ring $\mathbb{F}[[z_1, \dots, z_d]]$ such that a polynomial g over \mathbb{F}^{kn} is generically non-vanishing over X^n if and only if

$$g(f_1(\mathbf{z}_1), \dots, f_k(\mathbf{z}_1), \dots, f_1(\mathbf{z}_n), \dots, f_k(\mathbf{z}_n)) \neq 0.$$

□

A similar argument holds for the LD-MDS theorem.

6.4 Reduction of Power Series to Univariate Monomials

The goal will be to reduce Theorems 6.4 and 6.5 to equivalent theorems about curves of the form $x \rightarrow (x^{d_1}, \dots, x^{d_k}), d_1 < \dots < d_k$.

For $\mathbf{i} \in \mathbb{Z}_{\geq 0}^d$ we let $\mathbf{z}^{\mathbf{j}} = \prod_{k=1}^d z_k^{j_k}$. If we treat $F = (f_1, \dots, f_k) \in (\mathbb{F}[[z_1, \dots, z_d]])^k$ as a column vector of power series it is clear that,

$$F(\mathbf{z}) = \sum_{\mathbf{j} \in \mathbb{Z}_{\geq 0}^d} a_{\mathbf{j}} \mathbf{z}^{\mathbf{j}},$$

where $a_{\mathbf{j}}$ is a column vector in \mathbb{F}^k . We also define a lexicographic order on elements in $\mathbb{Z}_{\geq 0}^r$ and similarly a lexicographic order on monomials $\mathbf{z}^{\mathbf{i}}, \mathbf{i} \in \mathbb{Z}_{\geq 0}^r$.

We first give a simple characterization of MDS power series.

Lemma 6.8. A k -tuple $F = (f_1, \dots, f_k) \in \mathbb{F}[[z_1, \dots, z_d]]^k$ is k -MDS if and only if f_1, \dots, f_k are linearly independent as power series over \mathbb{F} .

Proof. If f_1, \dots, f_k are linearly dependent then it is clear that f_1, \dots, f_k are not k -MDS.

Let us now consider the case where f_1, \dots, f_k are linearly independent.

By Lemma 6.9 there exist $\mathbf{j}_1^* < \dots < \mathbf{j}_k^*$ such that $a_{\mathbf{j}_1^*}, \dots, a_{\mathbf{j}_k^*}$ are linearly independent. Also, we have $a_{\mathbf{j}} \in \text{span}(a_{\mathbf{j}_1^*}, \dots, a_{\mathbf{j}_{\beta-1}^*})$ when $\mathbf{j} < \mathbf{j}_\beta^*$.

To prove $F = (f_1, \dots, f_k)$ is MDS we want to show that,

$$\text{Det}(F(\mathbf{z}_1), \dots, F(\mathbf{z}_k)) = \text{Det} \begin{pmatrix} f_1(\mathbf{z}_1) & f_1(\mathbf{z}_2) & \cdots & f_1(\mathbf{z}_n) \\ f_2(\mathbf{z}_1) & f_2(\mathbf{z}_2) & \cdots & f_2(\mathbf{z}_n) \\ f_3(\mathbf{z}_1) & f_3(\mathbf{z}_2) & \cdots & f_3(\mathbf{z}_n) \\ \vdots & \vdots & \ddots & \vdots \\ f_k(\mathbf{z}_1) & f_k(\mathbf{z}_2) & \cdots & f_k(\mathbf{z}_n) \end{pmatrix} \neq 0,$$

where $\mathbf{z}_1, \dots, \mathbf{z}_i = (z_{i,1}, \dots, z_{i,d}), \dots, \mathbf{z}_k$ are distinct d tuples of variables.

By the multilinearity of the determinant we have,

$$\text{Det}(F(\mathbf{z}_1), \dots, F(\mathbf{z}_k)) = \sum_{\mathbf{j}_1, \dots, \mathbf{j}_k \in \mathbb{Z}_{\geq 0}^d} \text{Det}(a_{\mathbf{j}_1}, \dots, a_{\mathbf{j}_k}) \prod_{w=1}^k z_w^{\mathbf{i}_w}. \quad (30)$$

Consider the lexicographic order over $(\mathbf{j}_1, \dots, \mathbf{j}_k) \in (\mathbb{Z}_{\geq 0}^d)^k$. We note that the coefficient of $\prod_{w=1}^k z_w^{\mathbf{j}_w^*}$ is $\text{Det}(a_{\mathbf{j}_1^*}, \dots, a_{\mathbf{j}_k^*})$ which is non-zero. We also have that this is the smallest non-zero term as well but that is not important for this proof. \square

Before proving the above lemma we prove another helpful property of linearly independent power series.

Lemma 6.9. Given a k -tuple of power series $F = (f_1, \dots, f_k) \in \mathbb{F}[[z_1, \dots, z_d]]^k$ if f_1, \dots, f_k are linearly independent then for

$$F(\mathbf{z}) = \sum_{\mathbf{j} \in \mathbb{Z}_{\geq 0}^d} a_{\mathbf{j}} \mathbf{z}^{\mathbf{j}}$$

satisfies two properties,

1. There exist $\mathbf{j}_1^* < \dots < \mathbf{j}_k^*$ such that $a_{\mathbf{j}_1^*}, \dots, a_{\mathbf{j}_k^*}$ are linearly independent.
2. We also have, that $a_{\mathbf{j}}$ is in the $\text{span}(a_{\mathbf{j}_1^*}, \dots, a_{\mathbf{j}_{\beta-1}^*})$ for all $\mathbf{j}_{\beta-1}^* < \mathbf{j} < \mathbf{j}_\beta^*$.

Proof. We have

$$F(\mathbf{z}) = \sum_{\mathbf{j} \in \mathbb{Z}_{\geq 0}^d} a_{\mathbf{j}} \mathbf{z}^{\mathbf{j}},$$

with $a_{\mathbf{j}} \in \mathbb{F}^k, \mathbf{j} \in \mathbb{Z}_{\geq 0}^d$.

Let \mathbf{j}_1^* be the smallest index such that $a_{\mathbf{j}_1^*}$ is non-zero. Now inductively we define \mathbf{j}_i^* as the smallest index such that $a_{\mathbf{j}_i^*}$ does not lie in $\text{span}(a_{\mathbf{j}_1^*}, \dots, a_{\mathbf{j}_{i-1}^*})$.

We claim that it is possible to define $\mathbf{j}_1^*, \dots, \mathbf{j}_k^*$ as f_1, \dots, f_k are linearly independent. If the procedure fails then for some $i < k$, $a_{\mathbf{j}} \in \text{span}(a_{\mathbf{j}_1^*}, \dots, a_{\mathbf{j}_i^*})$ for all $\mathbf{j} \notin \{\mathbf{j}_1^*, \dots, \mathbf{j}_{i-1}^*\}$ where $a_{\mathbf{j}_1^*}, \dots, a_{\mathbf{j}_i^*}$ are linearly independent. If $a_{\mathbf{j}_1^*}, \dots, a_{\mathbf{j}_i^*}$ are column vectors then wlog let us assume the first i rows are linearly independent. After an invertible linear transformation over f_1, \dots, f_k we can assume $a_{\mathbf{j}_1^*}, \dots, a_{\mathbf{j}_i^*}$ are elementary basis vectors but that would now imply $f_{i+1} = \dots = f_k = 0$ which gives us a contradiction. \square

Proof of Theorems 6.6 and 6.7 assuming Theorems 4.1 and 5.1. We will now proceed with the reduction in two steps. First we show that it suffices to prove Theorems 6.6 and 6.7 for $F = (f_1, \dots, f_k)$ such that the f_i are monomials. The second step shows that the case of multivariate monomial can be reduced to univariate ones.

Claim 6.10. *Theorems 6.6 and 6.7 are respectively implied by Theorems 6.6 and 6.7 for $F = (f_1, \dots, f_k)$ where f_i are monomials.*

Proof. Given a k -tuple of power series $F = (f_1, \dots, f_k) \in \mathbb{F}[[z_1, \dots, z_d]]^k$ which is MDS, we apply Lemma 6.9 to find $\mathbf{j}_1^* < \dots < \mathbf{j}_k^*, \mathbf{j}_i^* \in \mathbb{Z}_{\geq 0}^d$ which satisfy the two properties in the statement of the lemma. For $\mathbf{i} = (i_1, \dots, i_t)$ in $\mathbb{Z}_{\geq 0}^d$ we let $|\mathbf{i}| = \sum_{t=1}^d i_t$.

Without loss of generality we can perform a change of basis over \mathbb{F}^k and assume $a_{\mathbf{j}_1^*}, \dots, a_{\mathbf{j}_k^*}$ are the elementary basis vectors e_1, \dots, e_k . The second property in Lemma 6.9 then implies that the leading coefficient in $f_i(\mathbf{z})$ is $\mathbf{z}^{\mathbf{j}_i^*}$ for $i \in [k]$.

We see that $F' = (\mathbf{z}^{\mathbf{j}_1^*}, \dots, \mathbf{z}^{\mathbf{j}_k^*})$ is MDS and by our assumption is also MDS(ℓ) and LD-MDS($\leq \ell$) for all $\ell > 0$.

Given any $A_1, \dots, A_\ell \subseteq [(\ell - 1)k]$ with the dimension k null intersection property. As F' is MDS(ℓ) we know that

$$\det \begin{pmatrix} I_k & V^{F'}|_{A_1} & & & \\ & I_k & V^{F'}|_{A_2} & & \\ & \vdots & & \ddots & \\ & I_k & & & V^{F'}|_{A_\ell} \end{pmatrix} \quad (31)$$

is non-zero.

We want to show that the determinant of

$$\begin{pmatrix} I_k & V^F|_{A_1} & & & \\ & I_k & V^F|_{A_2} & & \\ & \vdots & & \ddots & \\ & I_k & & & V^F|_{A_\ell} \end{pmatrix} \quad (32)$$

is non-zero. The matrix in (32) has ℓ blocks of rows with k rows in each block. Intuitively speaking we will show that the ‘lowest degree’ part of the determinant of (32) is (31) which will complete the proof.

In (32) we multiply the t th column with $\beta^{|\mathbf{j}_t^*|}$ for $t \in [k]$ to get the matrix.

$$\begin{pmatrix} \text{diag}(\beta^{|\mathbf{j}_1^*|}, \dots, \beta^{|\mathbf{j}_k^*|}) & V^F|_{A_1} & & & \\ \text{diag}(\beta^{|\mathbf{j}_1^*|}, \dots, \beta^{|\mathbf{j}_k^*|}) & & V^F|_{A_2} & & \\ & \vdots & & \ddots & \\ \text{diag}(\beta^{|\mathbf{j}_1^*|}, \dots, \beta^{|\mathbf{j}_k^*|}) & & & & V^F|_{A_\ell} \end{pmatrix} \quad (33)$$

In (33) we replace the variables $\mathbf{z}_i = (z_{i,1}, \dots, z_{i,d}), i \in [(\ell - 1)k]$ by $(\beta z_{i,1}, \dots, \beta z_{i,d})$ where β is a extra variable. Next we divide the t th row in each block by $\beta^{|\mathbf{j}_t^*|}$. The new matrix we get is

$$\begin{pmatrix} I_k & V^{F''}|_{A_1} & & & \\ & I_k & V^{F''}|_{A_2} & & \\ & \vdots & & \ddots & \\ & I_k & & & V^{F''}|_{A_\ell} \end{pmatrix} \quad (34)$$

where $F''(\mathbf{z}) = (f_1(\beta\mathbf{z})\beta^{-|\mathbf{j}_1^*|}, \dots, f_k(\beta\mathbf{z})\beta^{-|\mathbf{j}_k^*|})$. We note $f_i(\beta\mathbf{z})\beta^{-|\mathbf{j}_i^*|}$ is a polynomial in β and \mathbf{z} for all i . As we only performed row and column scaling operations. The determinant of (34) is non-zero if and

only if the determinant of (32) is non-zero. If we set $\beta = 0$ in (34) we get the matrix in (31) which completes the proof.

The reduction for the LD-MDS theorems follows similarly. \square

We note that the above argument would not work if the code was generated with different power series in each column.

Claim 6.11. *Theorems 6.6 and 6.7 for $F = (f_1, \dots, f_k)$ where f_i are multivariate monomials are respectively implied by Theorems 6.6 and 6.7 for $F = (f_1, \dots, f_k)$ where f_i are univariate monomials.*

Proof. Let $F = (\mathbf{z}^{\mathbf{i}_1}, \dots, \mathbf{z}^{\mathbf{i}_k})$, $\mathbf{z} = (z_1, \dots, z_k)$ be MDS. Without loss of generality we can assume $\mathbf{i}_1 < \dots < \mathbf{i}_k$.

The proof uses a standard reduction which encodes the degrees of multiple variables in the N -ary expansion of a number where N is large enough in comparison to the individual degrees of monomials appearing in F .

Take $N > |\mathbf{i}_k|$. Let $|\mathbf{i}|_N = \sum_{t=1}^d i_t N^{t-1}$ for $\mathbf{i} \in \mathbb{Z}^d$. We now substitute z_t with $\alpha^{N^{t-1}}$ for $t \in [d]$ in F , to get $F' = (\alpha^{|\mathbf{i}_1|_N}, \dots, \alpha^{|\mathbf{i}_k|_N})$. By our choice of N we have $|\mathbf{i}_1|_N < \dots < |\mathbf{i}_k|_N$ so F' is MDS. By our assumption it is also MDS(ℓ) and LD-MDS($\leq \ell$).

We are done as the determinants which need to be non-vanishing for F to be MDS(ℓ) and LD-MDS(ℓ) on substituting $\mathbf{z}_i = (z_{i,1}, \dots, z_{i,d})$ with $(\alpha_i^{N^0}, \dots, \alpha_i^{N^{d-1}})$ gives us the corresponding determinants for F' which are non-zero by our assumption. \square

\square

References

- [AGL24] Omar Alrabiah, Venkatesan Guruswami, and Ray Li. Randomly Punctured Reed–Solomon Codes Achieve List-Decoding Capacity over Linear-Sized Fields. In *Proceedings of the 56th Annual ACM Symposium on Theory of Computing*, pages 1458–1469, Vancouver BC Canada, June 2024. ACM.
- [BDG23] Joshua Brakensiek, Manik Dhar, and Sivakanth Gopi. Improved Field Size Bounds for Higher Order MDS Codes. In *2023 IEEE International Symposium on Information Theory (ISIT)*, pages 1243–1248, June 2023.
- [BDGZ24] Joshua Brakensiek, Manik Dhar, Sivakanth Gopi, and Zihan Zhang. AG Codes Achieve List Decoding Capacity over Constant-Sized Fields. In *Proceedings of the 56th Annual ACM Symposium on Theory of Computing*, pages 740–751, Vancouver BC Canada, June 2024. ACM.
- [BGM22] Joshua Brakensiek, Sivakanth Gopi, and Visu Makam. Lower Bounds for Maximally Recoverable Tensor Codes and Higher Order MDS Codes. *IEEE Transactions on Information Theory*, 68(11):7125–7140, November 2022.
- [BGM23] Joshua Brakensiek, Sivakanth Gopi, and Visu Makam. Generic Reed-Solomon Codes Achieve List-Decoding Capacity. In *Proceedings of the 55th Annual ACM Symposium on Theory of Computing*, STOC 2023, pages 1488–1501, New York, NY, USA, June 2023. Association for Computing Machinery.
- [DSDY13] Son Hoang Dau, Wentu Song, Zheng Dong, and Chau Yuen. Balanced Sparsest Generator Matrices for MDS Codes. In *2013 IEEE International Symposium on Information Theory*, pages 1889–1893, July 2013.
- [DSY14] Son Hoang Dau, Wentu Song, and Chau Yuen. On the existence of MDS codes over small fields with constrained generator matrices. In *2014 IEEE International Symposium on Information Theory*, pages 1787–1791, June 2014.

- [Eli57] P Elias. List Decoding for Noisy Channels. In *IRE WESCON Convention Record, 1957*, volume 2, pages 94–104, 1957.
- [Gab21] Ernst M. Gabidulin. *Rank Codes*. Technical University of Munich, March 2021.
- [GHK⁺17] Parikshit Gopalan, Guangda Hu, Swastik Kopparty, Shubhangi Saraf, Carol Wang, and Sergey Yekhanin. Maximally Recoverable Codes for Grid-like Topologies. In *Proceedings of the 2017 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, Proceedings, pages 2092–2108. Society for Industrial and Applied Mathematics, January 2017.
- [GS19] Gary Greaves and Jeven Syatriadi. Reed–Solomon Codes Over Small Fields With Constrained Generator Matrices. *IEEE Transactions on Information Theory*, 65(8):4764–4770, August 2019.
- [GST24] Eitan Goldberg, Chong Shangguan, and Itzhak Tamo. Singleton-type bounds for list-decoding and list-recovery, and related results. *Journal of Combinatorial Theory, Series A*, 203:105835, April 2024.
- [GZ23] Zeyu Guo and Zihan Zhang. Randomly Punctured Reed-Solomon Codes Achieve the List Decoding Capacity over Polynomial-Size Alphabets. In *2023 IEEE 64th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 164–176, November 2023.
- [Hal] JI Hall. Notes on Coding Theory (Chapter 5—Generalized Reed-Solomon Codes). *Department of Mathematics, Michigan State University available online at <http://users.math.msu.edu/users/jhall/classes/codenotes/coding-notes.html> (Jan. 7, 2015 revision)*.
- [HHYD14] Wael Halbawi, Tracey Ho, Hongyi Yao, and Iwan Duursma. Distributed Reed-Solomon Codes for Simple Multiple Access Networks. In *2014 IEEE International Symposium on Information Theory*, pages 651–655, June 2014.
- [Lov21] Shachar Lovett. Sparse MDS Matrices over Small Fields: A Proof of the GM-MDS Conjecture. *SIAM Journal on Computing*, 50(4):1248–1262, January 2021.
- [LWWS23] Hedongliang Liu, Hengjia Wei, Antonia Wachter-Zeh, and Moshe Schwartz. Linearized Reed-Solomon Codes with Support-Constrained Generator Matrix. In *2023 IEEE Information Theory Workshop (ITW)*, pages 7–12, April 2023.
- [MLR⁺14] Subramanian Muralidhar, Wyatt Lloyd, Sabyasachi Roy, Cory Hill, Ernest Lin, Weiwen Liu, Satadru Pan, Shiva Shankar, Viswanath Sivakumar, Linpeng Tang, and Sanjeev Kumar. f4: Facebook’s Warm BLOB Storage System. In *11th USENIX Symposium on Operating Systems Design and Implementation (OSDI 14)*, pages 383–398, 2014.
- [MS77] Florence Jessie MacWilliams and N. J. A. Sloane. *The Theory of Error Correcting Codes*. North-Holland Mathematical Library ; v. 16. North-Holland Pub. Co. ; sole distributors for the U.S.A. and Canada, Elsevier/North-Holland, Amsterdam ; New York : New York, 1977.
- [Rot22] Ron M. Roth. Higher-Order MDS Codes. *IEEE Transactions on Information Theory*, 68(12):7798–7816, December 2022.
- [RS60] I. S. Reed and G. Solomon. Polynomial Codes Over Certain Finite Fields. *Journal of the Society for Industrial and Applied Mathematics*, 8(2):300–304, June 1960.
- [Sin64] R. Singleton. Maximum Distance Q-Nary Codes. *IEEE Transactions on Information Theory*, 10(2):116–118, April 1964.
- [ST23] Chong Shangguan and Itzhak Tamo. Generalized Singleton Bound and List-Decoding Reed–Solomon Codes Beyond the Johnson Radius. *SIAM Journal on Computing*, 52(3):684–717, June 2023.

- [Tia19] Yongge Tian. Formulas for calculating the dimensions of the sums and the intersections of a family of linear subspaces with applications. *Beiträge zur Algebra und Geometrie / Contributions to Algebra and Geometry*, 60(3):471–485, September 2019.
- [Vak24] Ravi Vakil. *The Rising Sea: Foundations of Algebraic Geometry*. July 2024.
- [Woz58] John M Wozencraft. List Decoding. *Quarterly Progress Report*, 48:90–95, 1958.
- [YH19] Hikmet Yildiz and Babak Hassibi. Optimum Linear Codes With Support-Constrained Generator Matrices Over Small Fields. *IEEE Transactions on Information Theory*, 65(12):7868–7875, December 2019.
- [YH20] Hikmet Yildiz and Babak Hassibi. Gabidulin Codes With Support Constrained Generator Matrices. *IEEE Transactions on Information Theory*, 66(6):3638–3649, June 2020.
- [YS13] Muxi Yan and Alex Sprintson. Algorithms for Weakly Secure Data Exchange. In *2013 International Symposium on Network Coding (NetCod)*, pages 1–6, June 2013.

A Omitted Proofs

A.1 Proof of Lemma 3.7

We shall need the following facts implicit in [BGM22].

Proposition A.1 ([BGM22]). *Let V be a $k \times n$ matrix. Let $A_1, \dots, A_\ell \subseteq [n]$ be subsets such that $\dim(V_{A_i}) = |A_i|$. Let H be a $(\ell, \ell - 1)$ parity-check code. Then, the following are equivalent*

- (a) $\dim(V_{A_1} \cap \dots \cap V_{A_\ell}) = d$.
- (b) $\dim(\sum_{i=1}^{\ell} H_i \otimes V_{A_i}) = -d + \sum_{i=1}^{\ell} |A_i|$.

Proposition A.2. *Let X, Y, Y', Y'' be linear spaces such that $Y' \subseteq Y \subseteq Y''$ and each inclusion has codimension 1. Then,*

$$\begin{aligned} \dim(X \cap Y) &\geq \dim(X \cap Y') \geq \dim(X \cap Y) - 1 \\ \dim(X \cap Y) &\leq \dim(X \cap Y'') \leq \dim(X \cap Y) + 1 \end{aligned}$$

Proof. For the first line of inequalities. Note that the first one is trivial. For the second, observe that

$$\begin{aligned} \dim(X \cap Y') &= \dim((X \cap Y) \cap Y') \\ &= \dim(X \cap Y) + \dim Y' - \dim(Y' + (X \cap Y)) \\ &\geq \dim(X \cap Y) + \dim Y' - \dim Y \\ &= \dim(X \cap Y) - 1. \end{aligned}$$

The second inequality follows by similar logic. Or, we can use the first inequality by substituting (Y, Y'') for (Y', Y) . \square

Proof of Lemma 3.7. We begin with part (a). Assume without loss of generality that $\sigma_1 \geq \dots \geq \sigma_\ell$. If $\sigma_\ell \geq 1$, consider the minimal contraction $(\sigma'_1, A'_1), \dots, (\sigma'_\ell, A'_\ell)$ with $\sigma'_i = \sigma_i$ and $A'_i = A_i$ for all $i \in [\ell]$ except $\sigma'_\ell = \sigma_\ell - 1$. Now for any partition $P_1 \sqcup \dots \sqcup P_s = [\ell]$ with $\ell \in P_s$, then $\sigma'_{P_s} = \sigma'_\ell = \sigma_\ell - 1 = \sigma_{P_s} - 1$. Thus, we have that

$$\left[-n(s-1) + \sum_{i=1}^s (\sigma_{P_i} + |A_{P_i}|) \right] = \left[-n(s-1) + \sum_{i=1}^s (\sigma'_{P_i} + |A'_{P_i}|) \right] - 1.$$

Therefore, by taking the maximum over all such partitions and invoking Proposition 3.5, we have that $d - 1 = \text{gid}_k((\sigma'_1, A'_1), \dots, (\sigma'_\ell, A'_\ell))$.

Now assume that $\sigma_\ell = 0$ and consider $W \in \mathbb{F}^{k \times b}$ and $W' \in \mathbb{F}^{k \times n}$ generic. Let $X := \bigcap_{i=1}^{\ell} (W_{\leq \sigma_i} + W'_{A_i})$. Since $d \geq 1$, we have that $Y \neq 0$. Since $X \subseteq W'_{A_\ell}$ and $|A_\ell| \leq k$, each nonzero vector $x \in X$ is a unique nonzero linear combination of the columns of W'_{A_ℓ} . This implies there is at least one index $i^* \in A_\ell$ such that some vector $x \in X$ requires i^* in its linear combination. If we let $A'_\ell = A_\ell \setminus \{i^*\}$ and otherwise let $\sigma'_i = \sigma_i$ and $A'_i = A_i$, then $\dim(X \cap W'_{A'_\ell}) \leq \dim(X) - 1 = d - 1$. Likewise by Proposition 3.5 we must have that

$$\text{gid}_k((\sigma'_1, A'_1), \dots, (\sigma'_\ell, A'_\ell)) = \dim(X \cap W'_{A'_\ell}) = d - 1.$$

We now move on to part (b). Let $h_1, \dots, h_\ell \in \mathbb{F}^{\ell-1}$ be nonzero vectors forming an $(\ell, \ell - 1)$ -MDS code. By Proposition A.1 (and adapting it like in Proposition 3.5), we have that $\text{gid}_k((\sigma_1, A_1), \dots, (\sigma_\ell, A_\ell)) = 0$ if and only if

$$\dim \sum_{i=1}^{\ell} h_i \otimes (W_{\leq \sigma_i} + W'_{A_i}) = \sum_{i=1}^{\ell} \sigma_i + |A_i|.$$

Since $n \geq k$, we also have that

$$\dim \sum_{i=1}^{\ell} h_i \otimes (W_{\leq \sigma_i} + W'_{[n]}) = \dim(\mathbb{F}^{\ell-1} \otimes \mathbb{F}^k) = (\ell - 1)k.$$

Thus, since $\sum_{i=1}^{\ell} \sigma_i + |A_i| < (\ell - 1)k$, we argue there exists $i \in [\ell]$ and $j \in [n]$ with $j \notin A_i$ such that

$$\dim \left[h_i \otimes W'_j + \sum_{i=1}^{\ell} h_i \otimes (W_{\leq \sigma_i} + W'_{A_i}) \right] = 1 + \dim \sum_{i=1}^{\ell} h_i \otimes (W_{\leq \sigma_i} + W'_{A_i}).$$

If no such $h_i \otimes W'_j$ exists, then every $h_i \otimes W'_j$ must be a linear combination of vectors in $h_i \otimes (W_{\leq \sigma_i} + W'_{A_i})$, implying that $\dim \sum_{i=1}^{\ell} h_i \otimes (W_{\leq \sigma_i} + W'_{A_i}) = \dim \sum_{i=1}^{\ell} h_i \otimes (W_{\leq \sigma_i} + W'_{[n]})$, a contradiction.

Let $(\sigma'_1, A'_1), \dots, (\sigma'_\ell, A'_\ell)$ be the minimal expansion for which j is added to set A_i . We then have that $0 = \text{gid}_k((\sigma'_1, A'_1), \dots, (\sigma'_\ell, A'_\ell))$, as desired. \square

A.2 Proof of Lemma 6.2

\mathbb{F} throughout is algebraically closed.

At a high level the idea is that we can take a local view at a single point on the irreducible variety and that will give us a power series expansion.

Definition A.1 (Stalk of a variety at a point). *Given a variety $X \subseteq \mathbb{F}^k$ cut out by a radical ideal $I \subseteq \mathbb{F}[x_1, \dots, x_k]$ and a point $x \in X$ the stalk of X at x is constructed by taking the ring $\mathbb{F}[x_1, \dots, x_k]/I$ and inverting every polynomial f which does not vanish on x .*

The stalk corresponds to a local view of the variety X around x .

By construction for a variety X cut out by I there is a map from $\mathbb{F}[x_1, \dots, x_k]/I$ to the stalk at a point $x \in X$. It is easy to see that this map is injective if X is irreducible (equivalently I is prime). This means, for an irreducible variety looking at the stalk is enough to check if a polynomial is generically non-vanishing.

We next need the notion of smoothness.

Definition A.2 (Smooth point). *A point x on a variety $X \subseteq \mathbb{F}^k$ of dimension d cut out by a radical ideal $I = \langle f_1, \dots, f_m \rangle$ is said to be smooth when the Jacobian of the polynomials f_1, \dots, f_m is of rank $k - d$ at x .*

The stalks of smooth points have very nice properties. We will need the following one.

Theorem A.3 (Stalk at a smooth point (See 13.2.J and Section 28.2 in [Vak24])). *Given a smooth point x on an irreducible variety $X \subseteq \mathbb{F}^k$ of dimension d , the stalk at x injects into the power series ring $\mathbb{F}[[z_1, \dots, z_d]]$.*

A priori it is not clear if every variety will have a smooth points. Almost all points on a variety are smooth.

Theorem A.4 (Generic Smoothness of varieties (see 21.6.1 in [Vak24])). *If X is an irreducible variety then there is a dense open subset of X which only contains smooth points.*

We are now ready to prove Lemma 6.2.

Proof. Any irreducible variety will have at least one smooth point on it as the set of smooth points is dense (Lemma A.4). By Lemma A.3 the stalk at a smooth point on a dimension d irreducible variety maps injectively into $\mathbb{F}[[z_1, \dots, z_d]]$. This means the coordinate functions x_1, \dots, x_k on the stalk map to power series f_1, \dots, f_k .

For an irreducible variety checking a function is generically non-vanishing is equivalent to checking if the function is non-zero on a stalk of any point. As the stalk maps injectively into the power series ring we are done with the first part of the theorem.

The second part simply follows from the fact that the product of irreducible varieties is irreducible and we can take the product of the stalk of a point on X to get a stalk on X^n . \square