

# Poisson structure and Integrability of a Hamiltonian flow for the inhomogeneous six-vertex model

P. Rigas

## Abstract

We compute the action-angle variables for a Hamiltonian flow of the inhomogeneous six-vertex model, from a formulation introduced in a 2022 work due to Keating, Reshetikhin, and Sridhar, hence confirming a conjecture of the authors as to whether the Hamiltonian flow is integrable. To demonstrate that such an integrability property of the Hamiltonian holds from the action-angle variables, we make use of previous methods for studying Hamiltonian systems, implemented by Fadeev and Takhtajan, in which it was shown that integrability of a Hamiltonian system holds for the nonlinear Schrodinger's equation by computing action-angle variables from the Poisson bracket, which is connected to the analysis of entries of the monodromy and transfer matrices. For the inhomogeneous six-vertex model, an approach for computing the action-angle variables is possible through formulating several relations between entries of the quantum monodromy, and transfer, matrices, which can be not only be further examined from the structure of  $L$  operators, but also from computing several Poisson brackets parameterized in entries of the monodromy matrix.<sup>1</sup>

## 1 Introduction

### 1.1 Overview

The six-vertex model, originally introduced by chemists in [9], has emerged as a model of several studies in Probability theory, with recent efforts devoted towards rigorous descriptions of the localization, and delocalization, phases of the height function [3], the free energy [4], transition between disordered and antiferroelectric phases [6], and other related aspects pertaining to Russo-Seymour-Welsh results and crossing probabilities of the height function over strips of the square lattice, and interior of the cylinder, [10], in addition to classical work at the beginning of the seventies for computing the residual free entropy of the model on the torus[8]. To answer one conjecture raised in [7] regarding the integrability of the Hamiltonian flow for the six-vertex model, we make use of approaches described in [5] for determining the action-angle variables of the Hamiltonian that is introduced in the next section below. With such an approach, we resolve one conjecture raised by the authors in [7] as to whether the Hamiltonian flow for the six-vertex model that they introduce is integrable.

In comparison to a previous approach establishing that a Hamiltonian system for the nonlinear Schrodinger's equation is integrable [5], to demonstrate that such an integrability property holds for the inhomogeneous six-vertex model, one must incorporate characteristics from integrability of limit shapes [7]. In the presence of inhomogeneities, an underlying Hamiltonian for the six-vertex model can be shown to satisfy integrability properties consisting of a set of sixteen relations which can be derived from entries of the monodromy and transfer matrices, which is not only intimately connected to the R matrices satisfying the Yang-Baxter equation, but also to infinite volume limit of the transfer matrices as  $N \rightarrow +\infty$ , from which standard properties of the Poisson bracket can be used to asymptotically evaluate entries of the  $N$  th power of the monodromy matrix (the complete set of such relations that the monodromy matrix satisfies are provided in 2.4). From computations that are provided for asymptotic approximations of nine Poisson brackets, computations for the eighth Poisson bracket are the most complicated, from several cross terms arising from entries of the monodromy matrix (before computations for each of the nine Poisson brackets are executed in **Lemma 6- Lemma 14**, the asymptotic approximation of each Poisson bracket is provided in the statement of **Theorem 1** in 1.3).

### 1.2 Six-vertex model objects

Over the torus  $\mathbf{T}_N \equiv (V(\mathbf{T}_N), E(\mathbf{T}_N))$ , the six-vertex model can be defined through the probability measure,

---

<sup>1</sup>Keywords: Six-vertex model, integrability, inhomogeneity, Hamiltonian flow, action-angle variables, Poisson bracket




Figure 1: One depiction of each possible vertex for the six-vertex model, adapted from [1].




Figure 2: Another depiction of each possible vertex for the six-vertex model, adapted from [5].

$$\mathbf{P}_{\mathbf{T}_N}[\omega] \equiv \mathbf{P}[\omega] \equiv \frac{w(\omega)}{Z_{\mathbf{T}_N}} ,$$

where  $\omega$  is a *six-vertex configuration* determined by the six possible configurations (see Figure 1 and Figure 2), with the weight in the numerator of the probability measure taking the form,

$$w_{6V}(\omega) \equiv w(\omega) \equiv a_1^{n_1} a_2^{n_2} b_1^{n_3} b_2^{n_4} c_1^{n_5} c_2^{n_6} ,$$

for  $a_1, a_2, b_1, b_2, c_1, c_2 \geq 0$ , with the partition function,

$$Z_{\mathbf{T}_N}(\omega, \Omega) \equiv Z_{\mathbf{T}_N} = \sum_{\omega \in \Omega(\mathbf{T}_N)} w(\omega) .$$

Besides  $\mathbf{P}_{\mathbf{T}_N}[\cdot]$ , the disorder parameter of the six-vertex model is of the form,

$$\Delta \equiv \frac{a_1 a_2 + a_3 a_4 - a_5 a_6}{2\sqrt{a_1 a_2 a_3 a_4}} .$$

For non-symmetric weights, the weights of the six-vertex model admit the parametrization,

$$\begin{aligned} a_1 &\equiv a \exp(H + V) , \\ a_2 &\equiv a \exp(-H - V) , \\ b_1 &\equiv \exp(H - V) , \\ b_2 &\equiv \exp(-H + V) , \\ c_1 &\equiv c\lambda , \\ c_2 &\equiv c\lambda^{-1} , \end{aligned}$$

for  $a_1 \equiv a_2 \equiv a$ ,  $b_1 \equiv b_2 \equiv b$ ,  $c_1 \equiv c_2 \equiv c$ , and  $\lambda \geq 1$ , and external fields  $H, V$ . From such a parametrization of the weights as given above, one can form the so-called  $R$ -matrix, for the standard basis of  $\mathbf{C}^2$ , with,

$$R \equiv R(u, H, V) \equiv \begin{bmatrix} a \exp(H + V) & 0 & 0 & 0 \\ 0 & b \exp(H - V) & c & 0 \\ 0 & c & b \exp(-H + V) & 0 \\ 0 & 0 & 0 & a \exp(-H - V) \end{bmatrix}$$

in the tensor product basis  $e_1 \otimes e_1, e_1 \otimes e_2, e_2 \otimes e_1, e_2 \otimes e_2$ , for  $e_1 \equiv [1 \ 0]^T$  and  $e_2 \equiv [0 \ 1]^T$ . For vanishing external fields  $H \equiv V \equiv 0$ , if we denote  $R(u) \equiv R(u, 0, 0)$ , the matrix above admits the identity,

$$R(u, H, V) = \left( \text{diag}\left[\exp\left(\frac{H}{2}\right), \exp\left(-\frac{H}{2}\right)\right] \otimes \text{diag}\left[\exp\left(\frac{V}{2}\right), \exp\left(-\frac{V}{2}\right)\right] \right) R(u) \left( \text{diag}\left[\exp\left(\frac{H}{2}\right), \exp\left(-\frac{H}{2}\right)\right] \otimes \text{diag}\left[\exp\left(\frac{V}{2}\right), \exp\left(-\frac{V}{2}\right)\right] \right) \equiv (D^H \otimes D^V) R(u) (D^H \otimes D^V) ,$$

over  $\mathbf{C}^2 \otimes \mathbf{C}^2 \otimes \mathbf{C}^2$ . In widely celebrated work, [1], Baxter demonstrated that the  $R$  matrix satisfies the Yang-Baxter equation,

$$R_{12}(u) R_{13}(u+v) R_{23}(v) = R_{23}(v) R_{13}(u+v) R_{12}(u) .$$

For  $\Delta < -1$ , Baxter's parametrization, for the weights, given  $0 < u < \eta$ ,

$$a \equiv \sinh(\eta - u) , b \equiv \sinh(u) , c \equiv \sinh(\eta) ,$$

allows one to classify the eigenvalues of the transfer matrix which is related to the Bethe equations. Besides this fact, equipped with  $u$ , the horizontal lines of the square lattice,  $H$  and  $V$ , two external fields, the partition function over the  $N \times M$  torus,  $\mathbf{T}_{NM}$ , can be expressed with the summation,

$$Z_{\mathbf{T}_{MN}}(u, H) \equiv Z_{\mathbf{T}_{MN}} = \sum_{n=0}^N \exp(M(N-2n)V) Z_{\mathbf{T}_{MN}}^n(u, H) ,$$

for the semigrand canonical partition function,

$$Z_{\mathbf{T}_{MN}}^n(u, H) \equiv \sum_{\substack{i \geq 1 \\ \{\alpha_i\}}} (\Lambda_{\{\alpha_i\}}(u, H))^M ,$$

for countably many solutions  $\{\alpha_i\}$  to the Bethe equation,

$$\prod_{k=1}^N \frac{\sinh\left(\frac{\eta}{2} + i\alpha_j - v_k\right)}{\sinh\left(\frac{\eta}{2} - i\alpha_j + v_k\right)} = \exp(2HN) \prod_{m=1, m \neq j}^n \frac{\sinh(i(\alpha_j - \alpha_m) + \eta)}{\sinh(i(\alpha_j - \alpha_m) - \eta)} ,$$

and for the eigenvalue  $\Lambda_{\{\alpha_i\}}$ ,

$$\begin{aligned} \Lambda \equiv \Lambda_{\{\alpha_i\}} \equiv \exp(NH) \prod_{k=1}^N \sinh(\eta - u + v_k) \prod_{j=1}^n \frac{\sinh\left(\frac{\eta}{2} + u - i\alpha_j\right)}{\sinh\left(\frac{\eta}{2} - u + i\alpha_j\right)} + \exp(-NH) \prod_{k=1}^n \sinh(u - v_k) \\ \times \prod_{j=1}^n \frac{\sinh\left(\frac{3\eta}{2} - u + i\alpha_j\right)}{\sinh\left(u - \frac{\eta}{2} - i\alpha_j\right)} , \end{aligned}$$

with corresponding eigenvector,

$$\prod_{i=1}^n B(\alpha_i) |\Downarrow\rangle \equiv \prod_{i=1}^n B(\alpha_i) \left( \bigotimes_{i=1}^n |\downarrow\rangle \right) \equiv \prod_{i=1}^n B(\alpha_i) (|\downarrow\rangle \otimes \cdots \otimes |\downarrow\rangle) \equiv \prod_{i=1}^n B(\alpha_i) \left( \binom{0}{1} \otimes \cdots \otimes \binom{0}{1} \right) ,$$

of the transfer matrix,

$$t(u, \{v_k\}, H, V) : (\mathbf{C}^2)^{\otimes N} \longrightarrow (\mathbf{C}^2)^{\otimes N} ,$$

which, explicitly, is proportional to the trace of the quantum monodromy matrix, which is given by,

$$t(u, \{v_k\}, H, V) \equiv \prod_{i=1}^N D_i^{2V} \text{Tr}_a [T_a(u, \{v_k\}, H, 0)] ,$$

for the quantum monodromy matrix,

$$T_a(u, \{v_k\}, H, 0) : \mathbf{C}^2 \otimes (\mathbf{C}^2)^{\otimes N} \longrightarrow \mathbf{C}^2 \otimes (\mathbf{C}^2)^{\otimes N} \mapsto \prod_{i=1}^N \text{diag}(\exp(2H), \exp(2H)) R_{ia}(u - v_i) ,$$

where each  $v_i$  is chosen so that each  $u - v_i$  is a spectral parameter given at site  $i$ . In the next section, to apply the formalism first introduced for Hamiltonian systems in the nonlinear Schrodinger's equation to the quantum monodromy and transfer matrices of the six-vertex model, observe that the product of diagonal matrices with each  $R_{ia}$  is equivalent to,

$$\text{diag}(\exp(2H), \exp(2H)) R_{1a}(u - v_1) \cdots \times R_{(N-1)a}(u - v_{N-1}) \text{diag}(\exp(2H), \exp(2H)) R_{Na}(u - v_N) .$$

To formulate the Hamiltonian flow for the six-vertex model, from the statement of the Bethe equations, and their eigenvalues, introduce the functions  $\psi_u^\pm(\alpha + iu) \equiv \psi_\pm(\alpha + iu)$ , which are given by, [7],

$$\begin{aligned} \psi_+(\alpha + iu) &= \log \left[ \left| \frac{\exp(\eta + 2u) - \exp(2i\alpha)}{\exp(\eta - 2i\alpha) - \exp(2u)} \right| \right] = \log \left[ \left| \frac{\sinh(\frac{\eta}{2} + u - i\alpha)}{\sinh(\frac{\eta}{2} - u + i\alpha)} \right| \right] \equiv \log \left[ \frac{\left| \sinh(\frac{\eta}{2} + u - i\alpha) \right|}{\left| \sinh(\frac{\eta}{2} - u + i\alpha) \right|} \right] \\ &\quad = \log \left[ \frac{\sinh(\frac{\eta}{2} + u - i\alpha)}{\sinh(\frac{\eta}{2} - u + i\alpha)} \right] , \\ \psi_-(\alpha + iu) &= \log \left[ \left| \frac{\exp(2\eta + 2i\alpha) - \exp(2u - \eta)}{\exp(2u) - \exp(\eta + 2i\alpha)} \right| \right] \equiv \log \left[ \frac{\left| \exp(2\eta + 2i\alpha) - \exp(2u - \eta) \right|}{\left| \exp(2u) - \exp(\eta + 2i\alpha) \right|} \right] \\ &\quad \equiv \log \left[ \frac{\left| \sinh(\frac{3\eta}{2} - u + i\alpha) \right|}{\left| \sinh(u - \frac{\eta}{2} - i\alpha) \right|} \right] \\ &\quad = \log \left[ \frac{\sinh(\frac{3\eta}{2} - u + i\alpha)}{\sinh(u - \frac{\eta}{2} - i\alpha)} \right] , \end{aligned}$$

as well as the relation,

$$\Theta(\alpha - \beta) \equiv \frac{1 + \exp(ip(\alpha) + ip(\beta)) - 2\Delta \exp(ip(\alpha))}{1 + \exp(ip(\alpha) + ip(\beta)) - 2\Delta \exp(ip(\beta))} = -\frac{\sinh(i\alpha - i\beta + \eta)}{\sinh(i\alpha - i\beta - \eta)} ,$$

for,

$$p(\alpha) = \log \left[ \left| \frac{\sinh(\frac{\eta}{2} + i\alpha)}{\sinh(\frac{\eta}{2} - i\alpha)} \right| \right] \equiv \log \left[ \frac{\sinh(\frac{\eta}{2} + i\alpha)}{\sinh(\frac{\eta}{2} - i\alpha)} \right] ,$$

from which the statement of the Bethe equations is equivalent to, in terms of a parameterization of  $\psi_-(\alpha + v_k)$ , and  $\exp(\Theta(\alpha_j - \alpha_m))$ ,

$$\prod_{k=1}^N \exp(\psi_+(\alpha + v_k)) = \frac{1}{\exp(2HN)} \prod_{m=1, m \neq j} \frac{(-1)^m}{\exp(\Theta(\alpha_j - \alpha_m))} = \frac{1}{\exp(2HN)} \prod_{m=1, m \neq j}^n \frac{\sinh(i(\alpha_j - \alpha_m) - \eta)}{\sinh(i(\alpha_j - \alpha_m) + \eta)} ,$$

while the statement for the eigenvalue of the Bethe equations is equivalent to, given some  $\{\alpha_i\}$ , in terms of a parameterization of  $\psi_{\pm}(\alpha + v_k)$ ,

$$\Lambda(\psi_{\pm}(\alpha + iu)) \equiv \Lambda_{\{\alpha_i\}} \equiv \exp(NH) \prod_{k=1}^N \sinh(\eta - u + v_k) \prod_{j=1}^n \exp(\psi_+(\alpha_j + iu)) + \exp(-NH) \prod_{k=1}^n \sinh(u - v_k) \\ \times \prod_{j=1}^n \exp(\psi_-(\alpha_j + iu)) .$$

To introduce the Hamiltonian formulation of the six-vertex model, in which given a height function representation, from the set of all possible asymptotic height functions  $\mathcal{H}_{L,q}$ , for Lipschitz  $h \sim \mathcal{H}_{L,q}$  and  $x, x' \in [0, L]$ , with,

$$|h(x) - h(x')| < |x - x'| ,$$

and periodic, with,

$$h(L, y) = h(0, y) + q ,$$

for  $h : [0, L] \rightarrow \mathbf{R}$ , and fixed  $0 < q < 1$ . From such a sampling of the height function, as well as another periodic function  $\pi(x)$  in  $x$ , the pair  $(\pi(x), h(x))$  can be identified with the cotangent space  $T^*\mathcal{H}_{L,q}$ , while the flow of the Hamiltonian can be identified with the pair  $(\pi(x, y), h(x, y))$ , in which,

$$H_u(\pi(x, y), h(x, y)) \equiv H_u(\pi, h) = \int_{[0, L]} \mathcal{H}_{u-v(x)}[\partial_x h(x), \pi(x)] dx ,$$

over  $T^*\mathcal{H}_{L,q}$ , for a solution  $h(x, y)$  to the Euler Lagrange equations. Under the integral over  $x$  provided above,  $\mathcal{H}_u$  is the semigrand canonical free energy,

$$\mathcal{H}_u(q, H) \equiv \mathcal{H}_u = \log[Z_{\mathbf{T}_{MN}}^n(u, H)] ,$$

which can alternatively be expressed as the maximum over  $\pm$ , with,

$$\mathcal{H}_u(q, H) \equiv \max_{\pm} \mathcal{H}_u^{\pm}(q, H) \equiv \max_{\pm} \left\{ \pm H + l_{\pm} + \int_C \psi_u^{\pm}(\alpha) \rho(\alpha) d\alpha \right\} , \quad (\text{H})$$

with  $l_- \equiv \log \sinh(\eta - u)$ ,  $l_+ \equiv \log \sinh u$ , and the density,

$$\rho(\alpha) \equiv \#\{\text{Bethe roots along contours } C\} \equiv \bigcup_{\alpha} \{\alpha : \alpha \cap C \neq \emptyset\} \equiv \left| \{\alpha : \alpha \cap C \neq \emptyset\} \right| .$$

To connect the cotangent space  $T_{\phi_1}^* \mathcal{H}_{q,l}$  with  $T_{\phi_2}^* \mathcal{H}_{q,l}$  at time  $T$  given the initial flow line  $(\pi_0, h_0)$ , one determines the unique critical point of the functional,

$$S(\pi, h) \equiv S = \int_{[0, L]} \int_{[0, T]} [\pi(x, y) \partial_y h(x, y) - \mathcal{H}_{u-v(x)}(\partial_x h(x, y), \pi(x, y))] dy dx ,$$

which is  $(\pi_0, h_0)$ . Altogether,

$$Z_{\mathbf{T}_{MN}}^n = \exp(NM\mathcal{H}_u(q, H))(1 + o(1)) ,$$

asymptotically, for  $M \gg N$ ,  $n > 0$ , and  $q = \frac{n}{N}$  as  $N \rightarrow +\infty$ .

### 1.3 Paper organization

In the remaining subsection before computing L operators in the  $\Delta < -1$  regime of the six-vertex model 2, we provide an overview of the Hamiltonian formulation discussed in [5], from which computations are performed using the Poisson bracket, either on functionals or on matrix functionals, to determine the action-angle variables which are required for the statement of integrability of the Hamiltonian flow that are provided in **Theorem 1**. Such variables coincide with ones which have nonzero Poisson bracket, which can be simplified from the set of sixteen relations obtained from the  $N$  th power of the monodromy matrix in 2.4. The quantities that are included in the Poisson bracket are taken from entries of the monodromy matrix as  $x \rightarrow \pm\infty$ ,  $y \rightarrow \pm\infty$ , or as  $x \rightarrow \pm\infty$ ,  $y \rightarrow \pm\infty$  simultaneously. In order to make use of a similar approach for the Hamiltonian formulation described in the previous section from [7] in the context of the Hamiltonian flow for the six-vertex model in the presence of vertical and/or horizontal inhomogeneities, one must identify manners in which the Hamiltonian formulation of the six-vertex model depends on the structure of the L operator, which is not only dependent on the statement of the Bethe equations, but also on the eigenvalues of the Bethe equation itself, and Pauli operators  $\sigma^-$  and  $\sigma^+$ . In 2 we analyze such properties of the quantum monodromy, and transfer, matrices, from L operators. The particular choice of L operator that we take allows for us to obtain expressions for entries of the monodromy matrix that is introduced for the Hamiltonian flow of the six-vertex model, thereby allowing for us to deduce the desired action-angle variables of the Hamiltonian flow, hence concluding that the Hamiltonian flow in the presence of inhomogeneities satisfies an integrability property.

In the next section, 2, to further describe inhomogeneities of the six-vertex model that influence constructions of the transfer and quantum monodromy matrices, along with connections for computing action-angle variables, we obtain a system of recursive relations for each entry of the monodromy matrix which can provide expressions that we asymptotically approximate in the large  $N$  limit from nine Poisson brackets. Equipped with such a background, in 2.2, we introduce the L operator, and implement several computations, as suggested in the last section of [7], for computing each entry of the monodromy matrix raised to an arbitrary positive power. These relations are of upmost importance in performing computations in subsequent sections with the Poisson bracket, in which in [5] it was shown, for the nonlinear Schrodinger's equation, that there are sixteen relations that can be asymptotically approximated from properties of the Poisson bracket. For the inhomogeneous six-vertex model, each such relation relation is obtained from the fact that the Poisson bracket of the tensor product of two reduced monodromy matrices can be expressed in terms of a Poisson bracket between entries of reduced monodromy matrices. Asymptotically as  $N \rightarrow +\infty$ , from the monodromy matrix defined in [7] for studying inhomogeneities of the six-vertex model, sixteen relations from the Poisson bracket of the tensor product of reduced monodromy matrices can also be formulated in the context of the Hamiltonian flow and the quantum monodromy matrix, thereby . As a result, one can determine the action-angle variables for the inhomogeneous six-vertex model by determining which Poisson brackets vanish, from the set of sixteen relations.

**Theorem 1** (*precursor to the main result, integrability of the Hamiltonian flow for the inhomogeneous six-vertex model from nine Poisson brackets parametrized in entries of the N th power of the monodromy matrix*). Fix  $n, n', m > 0$ . The nine terms below each have the following asymptotic expansion as  $N \rightarrow +\infty$ ,

- First Poisson bracket

$$\mathcal{P}_1 \approx [(\sin(2\eta))^{n-3} \mathcal{C}_1]^2 \left[ \frac{A_3(u)B_3(u')}{u-u'} - \frac{A_3(u')B_3(u)}{u'-u} \right]$$

- Second Poisson bracket

$$\mathcal{P}_2 \approx -[(\sin(2\eta))^{n-3} \mathcal{C}_1] \left[ \frac{A_3(u)B_3(u')}{u-u'} - \frac{B_3(u)A_3(u')}{u'-u} \right]$$

- Third Poisson bracket

$$\begin{aligned}
\mathcal{P}_3 \approx & \sum_{n':m+n'=n-3} (\sin(2\eta))^{n'-1} \left[ \sum_{1 \leq i \leq m} \frac{\partial}{\partial u} \left( \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right) \right] \left[ \frac{\partial B_3(u)}{\partial u'} + \frac{\partial A_3(u)}{\partial u'} \right] + \sum_{n':m+n'=n-3} (\sin(2\eta))^{n'-1} \\
& \times \left[ \sum_{1 \leq i \leq m} \frac{\partial}{\partial u'} \left( \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right) \right] \\
& \times \left[ \frac{\partial B_3(u)}{\partial u} + \frac{\partial A_3(u)}{\partial u} \right] - (\sin(2\eta))^{n-3} \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} (\mathcal{C}_3)_{i,j} \right] \\
& \times \left[ \frac{B_3(u)A_3(u')}{u-u'} + \frac{A_3(u)B_3(u')}{u-u'} \right] ,
\end{aligned}$$

• *Fourth Poisson bracket*

$$\begin{aligned}
\mathcal{P}_4 \approx & -[(\sin(2\eta))^{n-3} (\mathcal{C}_1)_i] \left[ \frac{A_3(u)B_3(u')}{u-u'} + \frac{B_3(u)A_3(u')}{u-u'} \right] + \sum_{1 \leq i \leq n-3} \mathcal{A}_2 \left[ \left[ A_3(u) \frac{\partial B_3(u')}{\partial u'} \frac{\partial (\mathcal{C}_2)_i}{\partial u} \right. \right. \\
& \left. \left. + B_3(u) \frac{\partial B_3(u')}{\partial u} \frac{\partial (\mathcal{C}_2)_i}{\partial u'} \right] \right. \\
& \left. + \left[ A_3(u) \frac{\partial A_3(u')}{\partial u'} \frac{\partial (\mathcal{C}_2)_i}{\partial u} + B_3(u) \frac{\partial A_3(u')}{\partial u} \frac{\partial (\mathcal{C}_2)_i}{\partial u'} \right] \right] ,
\end{aligned}$$

• *Fifth Poisson bracket*

$$\begin{aligned}
\mathcal{P}_5 \approx & -\mathcal{C}_2 \left[ \frac{B_3(u')A_3(u)}{u'-u} + \frac{B_3(u)A_3(u')}{u-u'} \right] (B_3(u) + A_3(u)) \mathcal{C}_2 \left[ \sum_{1 \leq i \leq n-3} \left[ \frac{\partial B_3(u')}{\partial u'} \frac{\partial (\mathcal{C}_2)_i}{\partial u} - \frac{\partial B_3(u')}{\partial u} \right. \right. \\
& \left. \left. \times \frac{\partial (\mathcal{C}_2)_i}{\partial u} \right] \right] \left[ \prod_{1 \leq j \neq i \leq n-3} (\mathcal{C}_2)_j \right] - (B_3(u) + A_3(u)) \\
& \times \left[ \prod_{1 \leq i \leq n-3} (\mathcal{C}_2)_i \right] \\
& \times \left[ \sum_{1 \leq i \leq n-3} \left[ \frac{\partial A_3(u')}{\partial u'} \frac{\partial (\mathcal{C}_2)_i}{\partial u} - \frac{\partial A_3(u')}{\partial u} \frac{\partial (\mathcal{C}_2)_i}{\partial u} \right] \right] \\
& \left[ \prod_{1 \leq j \neq i \leq n-3} (\mathcal{C}_2)_j \right]
\end{aligned}$$

• *Sixth Poisson bracket*

$$\begin{aligned}
\mathcal{P}_6 \approx & \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} (\mathcal{C}_1)_j \right) \right] \right] \left[ \frac{A_3(u)B_3(u')}{u-u'} \right] \\
& + \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} (\mathcal{C}_1)_j \right) \right] \right] \left[ \frac{B_3(u)A_3(u')}{u-u'} \right] \\
& - 2(B_3(u) + A_3(u)) \left[ \sum_{1 \leq j \leq n'} (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} (\mathcal{C}_1)_j \right) \right] \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \frac{\partial (\mathcal{C}_2)_i}{\partial u'} \right) \right. \right. \\
& \left. \left. \left[ \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right] \right] \right]
\end{aligned}$$

$$\times \left( \prod_{1 \leq j \neq i \leq m} (\mathcal{C}_2)_j \right) \left( \frac{\partial A_3(u)}{\partial u} + \frac{\partial B_3(u)}{\partial u} \right) + \left( \frac{\partial (\mathcal{C}_2)_i}{\partial u} \right) \left( \prod_{1 \leq j \neq i \leq m} (\mathcal{C}_2)_j \right) \\ \times \left( \frac{\partial A_3(u')}{\partial u'} + \frac{\partial B_3(u')}{\partial u'} \right) \right] \right] ,$$

- *Seventh Poisson bracket*

$$\mathcal{P}_7 \approx \left[ (\sin(2\eta))^{n-3} \mathcal{C}_1 \right] \left[ \frac{A_3(u) B_3(u')}{u - u'} \right] + \left[ (\sin(2\eta))^{n-3} \mathcal{C}_1 \right] \left[ \frac{B_3(u) A_3(u')}{u - u'} \right] + \left[ \sum_{1 \leq j \leq n'} (\sin(2\eta))^{n'-1} \mathcal{C}_1 \right] \\ \times \left[ B_3(u) \left[ \sum_{1 \leq i \leq m} \left[ \left[ \frac{\partial}{\partial u} \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right] \left[ \frac{\partial B_3(u')}{\partial u'} - \frac{\partial B_3(u')}{\partial u} \right] \right. \right. \right. \\ \left. \left. \left. - \left[ \frac{\partial}{\partial u'} \prod_{1 \leq i \leq m} \sin(\mathcal{C}_2)_i \right] \left[ \frac{\partial A_3(u')}{\partial u} - \frac{\partial A_3(u')}{\partial u'} \right] \right] \right] \right] \\ + A_3(u) \left[ \sum_{1 \leq i \leq m} \left[ \left[ \frac{\partial}{\partial u} \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right] \left[ \frac{\partial A_3(u')}{\partial u'} - \frac{\partial A_3(u')}{\partial u} \right] \right. \right. \\ \left. \left. - \left[ \frac{\partial}{\partial u'} \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right] \left[ \frac{\partial B_3(u')}{\partial u'} - \frac{\partial B_3(u')}{\partial u} \right] \right] \right] ,$$

- *Eighth Poisson bracket*

$$\mathcal{P}_8 \approx - \left[ \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right) (\sin(2\eta))^{n'-1} \mathcal{C}_1 \right] \right) \left( \prod_{1 \leq i \leq n-3} (\mathcal{C}_2)_i \right) \right] \left[ \frac{A_3(u) B_3(u')}{u - u'} \right] \\ - \left[ \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right) (\sin(2\eta))^{n'-1} \mathcal{C}_1 \right] \right) \left( \prod_{1 \leq i \leq n-3} (\mathcal{C}_2)_i \right) \right] \left[ \frac{B_3(u) A_3(u')}{u - u'} \right] \\ - 2 \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right) (\sin(2\eta))^{n'-1} \mathcal{C}_1 \right] \right) B_3(u') \left[ \sum_{1 \leq j \leq n-3} \left[ \left[ \frac{\partial B_3(u)}{\partial u} \right] \left[ \frac{\partial (\mathcal{C}_2)_i}{\partial u'} \right] \right. \right. \\ \left. \left. - \left[ \frac{\partial B_3(u)}{\partial u'} \right] \left[ \frac{\partial (\mathcal{C}_2)_i}{\partial u} \right] \right] \right] \prod_{1 \leq j \neq i \leq n-3} (\mathcal{C}_2)_j - \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right) (\sin(2\eta))^{n'-1} \mathcal{C}_1 \right] \right) \\ \times A_3(u') \left[ \sum_{1 \leq j \leq n-3} \left[ \left[ \frac{\partial A_3(u)}{\partial u} \right] \left[ \frac{\partial (\mathcal{C}_2)_i}{\partial u'} \right] - \left[ \frac{\partial A_3(u)}{\partial u'} \right] \left[ \frac{\partial (\mathcal{C}_2)_i}{\partial u} \right] \right] \right] \prod_{1 \leq j \neq i \leq n-3} (\mathcal{C}_2)_j \\ - \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right) (\sin(2\eta))^{n'-1} \mathcal{C}_1 \right] \right) \\ \times A_3(u') \left[ \sum_{1 \leq j \leq n-3} \left[ \left[ \frac{\partial B_3(u')}{\partial u'} \right] \left[ \frac{\partial (\mathcal{C}_2)_i}{\partial u'} \right] - \left[ \frac{\partial B_3(u')}{\partial u} \right] \left[ \frac{\partial (\mathcal{C}_2)_i}{\partial u} \right] \right] \right] \prod_{1 \leq j \neq i \leq n-3} (\mathcal{C}_2)_j \right]$$

$$+(\sin(2\eta))^{n'-1} \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \left( \left[ \frac{\partial}{\partial u} \left[ B_3(u') + A_3(u') \right] \right] \frac{\partial(\mathcal{C}_2)_i}{\partial u'} - \left[ \frac{\partial}{\partial u'} \left[ A_3(u') + B_3(u') \right] \right] \frac{\partial(\mathcal{C}_2)_i}{\partial u} \right) \right) \right],$$

• Ninth Poisson bracket

$$\begin{aligned} \mathcal{P}_9 \approx & - \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} [\mathcal{C}_2 (\sin(2\eta))^{n'-1} \mathcal{C}_1] \right] \left[ \frac{B_3(u') A_3(u)}{u' - u} + \frac{A_3(u') B_3(u)}{u' - u} \right] - \left[ 2[B_3(u) + A_3(u)] \right. \\ & \times \left. \sum_{n':m+n'=n-3} \left( (\sin(2\eta))^{n'-1} \mathcal{C}_2 \right)^2 \right] \left[ \frac{B_3(u') A_3(u)}{u' - u} + \frac{A_3(u') B_3(u)}{u' - u} \right] \\ & + 2 \left[ \frac{\partial}{\partial u'} [A_3(u') + B_3(u')] \right] \left[ \sum_{1 \leq i \leq m} \left[ \frac{\partial}{\partial u} \left[ \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right] \right] \right. \\ & \quad \left. - \left[ \frac{\partial}{\partial u} [A_3(u') + B_3(u')] \right] \right] \\ & \times \left[ \sum_{1 \leq i \leq m} \left[ \frac{\partial}{\partial u'} \left[ \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right] \right] \right], \end{aligned}$$

The Poisson brackets  $\mathcal{P}_1, \dots, \mathcal{P}_9$ , are given by,

$$\begin{aligned} \mathcal{P}_1 &\equiv \left\{ \mathcal{P}_1 (\sin(2\eta))^{n-3} \mathcal{A}_1, \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\}, \\ \mathcal{P}_2 &\equiv \left\{ \mathcal{P}_1 (\sin(2\eta))^{n-3} \mathcal{A}_1, \mathcal{P}_2 \mathcal{A}'_2 \right\}, \\ \mathcal{P}_3 &\equiv \left\{ \mathcal{P}_1 (\sin(2\eta))^{n-3} \mathcal{A}_1, \mathcal{P}_2 \mathcal{A}'_3 \right\}, \\ \mathcal{P}_4 &\equiv \left\{ \mathcal{P}_1 \mathcal{A}_2, \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\}, \\ \mathcal{P}_5 &\equiv \left\{ \mathcal{P}_1 \mathcal{A}_2, \mathcal{P}_2 \mathcal{A}'_2 \right\}, \\ \mathcal{P}_6 &\equiv \left\{ \mathcal{P}_1 \mathcal{A}_2, \mathcal{P}_2 \mathcal{A}'_3 \right\}, \\ \mathcal{P}_7 &\equiv \left\{ \mathcal{P}_1 \mathcal{A}_3, \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\}, \\ \mathcal{P}_8 &\equiv \left\{ \mathcal{P}_1 \mathcal{A}_3, \mathcal{P}_2 \mathcal{A}'_2 \right\}, \\ \mathcal{P}_9 &\equiv \left\{ \mathcal{P}_1 \mathcal{A}_3, \mathcal{P}_2 \mathcal{A}'_3 \right\}. \end{aligned}$$

Taking the summation of the collection of Poisson brackets  $\mathcal{C} \equiv \bigcup_{1 \leq i \leq 9} \mathcal{P}_i$  above yields,

$$\sum_{1 \leq i \leq 9} \mathcal{P}_i ,$$

equals,

$$\left\{ \left( A_3(u) + B_3(u) \right) \left[ (\sin(2\eta))^{n-3} \mathcal{A}_1 + \mathcal{A}_2 + \mathcal{A}_3 \right], \left( A_3(u') + B_3(u') \right) \left[ (\sin(2\eta))^{n-3} \mathcal{A}'_1 + \mathcal{A}'_2 + \mathcal{A}'_3 \right] \right\} .$$

To demonstrate that the desired conclusion above holds for the main result, which can be extended, through very similar manipulations of the Poisson bracket for fifteen other relations that are also parametrized in entries of the monodromy matrix, we perform several computations with L operators in  $\mathcal{L}$ . From performing computations with the L operators, we obtain, explicitly, expressions for  $\mathcal{A}_1$ ,  $\mathcal{A}_2$ ,  $\mathcal{A}_3$ ,  $\mathcal{A}'_1$ ,  $\mathcal{A}'_2$ , and  $\mathcal{A}'_3$ . Below, we introduce the action-angle variables and the result that the action-angle variables, in canonical coordinates, vanish.

**Definition** (*action-angle variables for the inhomogeneous six-vertex model*). Define,

$$\begin{aligned}\varphi(u) &= -\arg(B(u)) \quad , \\ \rho(u) &= \frac{1}{\pi} \log[1 + \epsilon|B(u)|^2] \quad ,\end{aligned}$$

corresponding to action-angle variables for the inhomogeneous six-vertex model.

To determine the second action-angle variable  $\rho(u)$  from  $\varphi(u)$ , we introduce the result below for determining the canonically conjugate variable.

**Lemma** (*determining, from a computation with the Poisson bracket, canonically conjugate variables to  $\varphi(u)$* ). There exists a function  $f$  for which,

$$\{f(|B(u)|^2), \varphi(u')\} = f'(|B(u)|^2)[1 + \epsilon|b(u)|^2] \iff f(x) = \frac{1}{\pi} \log[1 + \epsilon x] \quad .$$

*Proof of Lemma.* Refer to the computation, with the Poisson bracket, of the canonically conjugate variable to  $\varphi(\mu)$  provided in *Chapter III, 7* of [5]. A very similar computation provides the desired function above, from which we conclude the argument.  $\square$

**Theorem 2** (*main result, vanishing of the Poisson bracket of the action-angle variables in canonical coordinates*). In canonical coordinates, the two Poisson brackets,

$$\begin{aligned}\{\varphi(u_1), \varphi(u_2)\} &\quad , \\ \{\rho(u_1), \rho(u_2)\} &\quad ,\end{aligned}$$

vanish.

## 1.4 From the reduced monodromy, and monodromy, matrices to a Hamiltonian system

A use of the Hamiltonian approach from [5], which demonstrates that the Euler-Lagrange equations have infinitely many conserved quantities, is summarized below.

### 1.4.1 Previous application of the Hamiltonian formulation for the nonlinear Schrodinger's equation

With  $T_a(u, \{v_k\}, H, V)$ , which besides the previous diagonal representation can also be expressed as,

$$T_a(u, \{v_k\}, H, V) \equiv \begin{bmatrix} A(u) & B(u) \\ C(u) & D(u) \end{bmatrix}$$

one may proceed to determine the angle-action variables of the Hamiltonian system, as described in [5].

To implement the computations for this approach for the nonlinear Schrodinger's equation, introduce, under slightly different parameters, the reduced monodromy matrix, from the limit as  $x \rightarrow +\infty$ , and as  $y \rightarrow -\infty$ , (*Chapter III, [5]*),

$$\lim_{N \rightarrow +\infty} T_N(\lambda) \equiv T(\lambda) = \lim_{\substack{x \rightarrow +\infty \\ y \rightarrow -\infty}} E(-x, \lambda) T(x, y, \lambda) E(y, \lambda) \quad ,$$

for the matrix exponential,

$$E(x, \lambda) \equiv \exp(\lambda x U_1) ,$$

and also from the clockwise exponential action,

$$T_L(\lambda) \equiv \exp \left[ \int_{[-L,L]}^{\text{clockwise action}} U(x, \lambda) dx \right] ,$$

for a real parameter  $\lambda$ , with the integrand  $U$ , a  $2 \times 2$  matrix, and  $N \geq L$ , which satisfies the first order PDE,

$$\frac{\partial F}{\partial x} = U(x, \lambda) F ,$$

for the vector valued function  $F(x, t) \equiv F \equiv [f_1, f_2]^T$ . The nonconstant prefactor for  $F$  takes the form of the linear combination,

$$U = U_0 + \lambda U_1 ,$$

with,

$$U_0 \equiv \sqrt{\chi} \begin{bmatrix} 0 & \psi \\ \bar{\psi} & 0 \end{bmatrix} ,$$

$$U_1 \equiv \frac{1}{2i} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} .$$

The coordinates appearing in the definition of  $U_0$  satisfy the second order PDE, the nonlinear Schrodinger's equation, with solutions  $\psi$ ,

$$i \frac{\partial \psi}{\partial t} = - \frac{\partial^2 \psi}{\partial x^2} + 2\chi |\psi|^2 \psi ,$$

for real  $\chi$ , a coupling constant, with  $|\psi|^2 = \psi \bar{\psi}$ . In the following infinite limit, the solutions satisfy,

$$\lim_{x \rightarrow \pm\infty} \psi(x) = \rho_1^{(\pm)} ,$$

$$\lim_{x \rightarrow \pm\infty} \psi(x) = \rho_2^{(\pm)} ,$$

for real  $\rho_1^{(\pm)}, \rho_2^{(\pm)}$  with  $\rho_1^{(\pm)} \neq \rho_2^{(\pm)}$ . Equipped with solutions to the nonlinear PDE above, the Hamiltonian admits the following integral representation over the real line, (*Chapter III*, [5]),

$$H(x, \psi, \bar{\psi}, \chi) \equiv H \equiv \int_{\mathbf{R}} \left( \frac{\partial \psi}{\partial x} \frac{\partial \bar{\psi}}{\partial x} + \chi \bar{\psi}^2 \psi^2 \right) dx ,$$

Under the matrix representation for the reduced monodromy matrix, (*Chapter I*, [5]),

$$T_L(\lambda) \equiv T(\lambda) \equiv \begin{bmatrix} a(\lambda) & \epsilon \bar{b}(\lambda) \\ b(\lambda) & \bar{a}(\lambda) \end{bmatrix}$$

the transition coefficients for discrete spectra occur iff  $\epsilon \equiv -1$ , and are given by,

$$T_-^{(1)}(x, \lambda_j) = \gamma_j T_+^{(2)}(x, \lambda_j) ,$$

for  $1 \leq j \leq n$ , where the  $\lambda_j$  parameter appearing in  $T_-^{(1)}(x, \cdot)$ , and in  $T_+^{(2)}(x, \cdot)$ , is the collection of zeros,  $\{\lambda_j\}$ , of  $a(\lambda)$  in the upper half plane, and  $\gamma_j$  denotes another real parameter, the *transition amplitude*, between  $T_-^{(1)}(x, \cdot)$  and  $T_+^{(2)}(x, \cdot)$ . For another real parameter  $\mu$ , the identity

$$\{T(x, y, \lambda) \otimes T(x, y, \mu)\} = \{r(\lambda - \mu), T(x, y, \lambda) \otimes T(x, y, \mu)\} ,$$

of the Poisson bracket  $\{\cdot, \cdot\}$ , which satisfies the anticommutativity, and bilinearity properties, Leibniz' rule, and the Jacobi identity, holds for  $y < x$ , where,

$$r_{\pm}(\lambda - \mu) = \lim_{y \rightarrow \pm\infty} E(y, \mu - \lambda) \otimes E(y, \lambda - \mu) r(\lambda - \mu) ,$$

under the convention for the tensor product, between  $A$  and  $B$ , with respect to the Poisson bracket, (*Chapter III*, [5]),

$$\{A \otimes B\}_{jk,mn} \equiv \{A_{jm}, B_{kn}\} .$$

#### 1.4.2 Poisson bracket and the action-angle variables

The action of the Poisson bracket that is implemented for demonstrating that the Hamiltonian flow for the nonlinear Schrodinger's equation is integrable depends upon whether the arguments of the Poisson bracket are functionals, or matrix functionals. In the first case, the bracket takes the form, (*Chapter III*, [5]),

$$\{F, G\} \equiv i \int_{[-L, L]} \left( \frac{\delta F}{\delta \psi} \frac{\delta G}{\delta \bar{\psi}} - \frac{\delta F}{\delta \bar{\psi}} \frac{\delta G}{\delta \psi} \right) dx ,$$

for functionals  $F$  and  $G$ , while for the second case, the bracket takes the form,

$$\{A \otimes B\} \equiv i \int_{[-L, L]} \left( \frac{\delta A}{\delta \psi} \otimes \frac{\delta B}{\delta \bar{\psi}} - \frac{\delta A}{\delta \bar{\psi}} \otimes \frac{\delta B}{\delta \psi} \right) dx ,$$

for matrix functionals  $A$  and  $B$ , and with  $\psi \equiv \psi(x)$  and  $\bar{\psi} \equiv \bar{\psi}(x)$  having support over  $(-L, L)$ . Similarly, from  $T(x, y, \lambda)$ ,

$$T_{\pm}(x, \lambda) = \lim_{y \rightarrow \pm\infty} T(x, y, \lambda) E(y, \lambda) .$$

From the first identity involving the Poisson bracket, one similarly has, as either  $y \rightarrow -\infty$ , or as  $y \rightarrow +\infty$ , independently of  $y$ ,

$$\begin{aligned} \{T_-(x, \lambda) \otimes T_-(x, \mu)\} &= r(\lambda - \mu) T_-(x, \lambda) \otimes T_-(x, \mu) - T_-(x, \lambda) \otimes T_-(x, \mu) r_-(\lambda - \mu) , \\ \{T_+(x, \lambda) \otimes T_+(x, \mu)\} &= T_+(x, \lambda) \otimes T_+(x, \mu) r_+(\lambda - \mu) - r(\lambda - \mu) T_+(x, \lambda) \otimes T_+(x, \mu) , \end{aligned}$$

while, between  $T_-(x, \lambda)$  and  $T_+(x, \lambda)$ ,

$$\{T_-(x, \lambda) \otimes T_+(x, \lambda)\} = 0 .$$

From the quantities introduced thus far, the action angle variables, given a Hamiltonian flow, are determined by whether the Poisson bracket vanishes or not, in which the only terms with nonvanishing Poisson bracket are,

$$\{\Phi(\lambda), \Phi(\bar{\lambda})\} = 0 ,$$

for the function,

$$\Phi(\lambda) \equiv \sqrt{\rho(\lambda)} \exp(-i\phi(\lambda)) ,$$

and its complex conjugate,  $\bar{\Phi}(\bar{\lambda})$ , as well as,

$$\begin{aligned}\rho(\lambda) &\equiv \frac{1}{2\pi\chi} \log[1 + \epsilon|b(\lambda)|^2] , \\ \phi(\lambda) &\equiv -\arg(b(\lambda)) .\end{aligned}$$

With the monodromy, and transfer, matrices of the previous section which were respectively denoted with  $t(u, \{v_k\}, H, V)$  and  $T_a(u, \{v_k\}, H, 0)$ , in the next section we state the key property that the action-angle variables satisfy.

## 2 Action-angle variables for the Hamiltonian flow in the $\Delta < -1$ regime of the six-vertex model

To make use of the Poisson bracket for determining which variables with respect to it vanish, it suffices to demonstrate that the Poisson bracket vanishes for certain entries of the quantum monodromy matrix. We recall the representation presented in 1.4.1 with,  $T_a(u, \{v_k\}, H, V)$  of the quantum monodromy matrix, which was introduced alongside the definition of the transfer matrix,  $t(u, \{v_k\}, H, V)$ .

### 2.1 Inhomogeneities in the horizontal direction

In the setting of the six-vertex model, the action-angle variables are determined by the structure of the quantum monodromy matrix, and hence of the transfer matrix.

From the background provided in the previous section on the Hamiltonian structure of the nonlinear Schrodinger's equation, instead of sending  $x \rightarrow \pm\infty$ , and  $y \rightarrow -\infty$ , we send the spectral parameters at each site  $i$  to  $-\infty$ , in which,

$$u - v_1 \rightarrow -\infty, \dots, u - v_N \rightarrow -\infty ,$$

from each term of the R-matrix appearing in the quantum monodromy matrix,

$$T_a(u, H, V) = \lim_{v_k \rightarrow -\infty} T_a(u, \{v_k\}, H, V) = \lim_{v_i \rightarrow -\infty} \left[ \prod_{i=1}^N (\text{diag}(\exp(2H), \exp(2H))) R_{ia}(u - v_i) \right] .$$

As  $v_i \rightarrow -\infty, \dots, v_N \rightarrow -\infty$ , the above limit is equivalent to,

$$\lim_{\substack{v_1 \rightarrow -\infty \\ \vdots \\ v_N \rightarrow -\infty}} \left[ (\text{diag}(\exp(2H), \exp(2H))) R_{1a}(u - v_1) \cdots (\text{diag}(\exp(2H), \exp(2H))) R_{Na}(u - v_N) \right] .$$

From  $T_a$  above, in comparison to the monodromy matrix  $T(\lambda)$  for the nonlinear Schrodinger's equation, the terms dependent on  $u$ ,  $B(u)$  and  $C(u)$ , are not complex conjugates of one another as  $b(\bar{\lambda})$  and  $b(\lambda)$  are, as provided in the definition of  $T(\lambda)$ .

In order to demonstrate that the six-vertex model is integrable from properties of the Hamiltonian flow, and to also make sense of the multi dimensional limit show above, it suffices to show that the Poisson bracket of the generator of motion integrals vanish. First, recall, from (H),

$$\mathcal{H}_u(q, H) \equiv \max_{\pm} \mathcal{H}_u^{\pm}(q, H) \equiv \max_{\pm} \left\{ \pm H + l_{\pm} + \int_C \psi_u^{\pm}(\alpha) \rho(\alpha) d\alpha \right\} ,$$

which, explicitly in terms of  $\psi$ , is equivalent to,

$$\mathcal{H}_u^\pm(q, H) \equiv \begin{cases} + \Rightarrow \max_{+} \left\{ H + \log \sinh u + \int_C \log \left[ \frac{\sinh(\frac{\eta}{2} + u - i\alpha - iu)}{\sinh(\frac{\eta}{2} - u + i\alpha - iu)} \right] \rho(\alpha) d\alpha \right\} , \\ - \Rightarrow \max_{-} \left\{ -H + \log \sinh(\eta - u) + \int_C \log \left[ \frac{\sinh(\frac{3\eta}{2} - u + i\alpha - iu)}{\sinh(u - \frac{\eta}{2} - i\alpha - iu)} \right] \rho(\alpha) d\alpha \right\} . \end{cases}$$

With each motion integral introduced above for  $+$  and for  $-$ , we proceed to identify the action-angle variables for the Hamiltonian flow of the six-vertex model from the motion integrals above.

To make such an identification of which entries of the monodromy matrix appear in expressions for the generating functions of the the motion integrals, observe that the quantum monodromy matrix, denoted earlier with  $T_a(u, \{v_k\}, H, V)$ , upon setting  $u \equiv \lambda_\alpha$  for the eigenvalue of the quantum monodromy matrix, can be expressed as,

$$\begin{bmatrix} A(\lambda_\alpha) & B(\lambda_\alpha) \\ C(\lambda_\alpha) & D(\lambda_\alpha) \end{bmatrix}$$

which admits a decomposition in terms of  $L$  operators, from the product,

$$\prod_{i=0}^{N-1} L_{\alpha, N-i}(\lambda_\alpha, v_{N-i}) ,$$

where the  $L$  operator is defined as, [2],

$$L_{\alpha, k}(\lambda_\alpha, v_k) \equiv \begin{bmatrix} \sin(\lambda_\alpha - v_k + \eta\sigma_k^z) & \sin(2\eta)\sigma_k^- \\ \sin(2\eta)\sigma_k^+ & \sin(\lambda_\alpha - v_k - \eta\sigma_k^z) \end{bmatrix} ,$$

corresponding to the  $k$  th horizontal line, and  $\alpha$  th vertical line, with  $L_{\alpha, k} \curvearrowright (\text{vertical space} \otimes \text{horizontal space})$ .

From the block decomposition above in terms of  $A$ ,  $B$ ,  $C$  and  $D$ , one also has that,

$$\text{tr}(T_a(\lambda_a, \{v_k\}, H, V)) = A(\lambda_\alpha) + D(\lambda_\alpha) .$$

From the product expansion over  $N - 1$   $L$  operators for  $T_a$ , one expands the product,

$$\begin{aligned} \prod_{i=0}^{N-1} \begin{bmatrix} \sin(\lambda_\alpha - v_{N-i} + \eta\sigma_{N-i}^z) & \sin(2\eta)\sigma_{N-i}^- \\ \sin(2\eta)\sigma_{N-i}^+ & \sin(\lambda_\alpha - v_{N-i} - \eta\sigma_{N-i}^z) \end{bmatrix} &= \begin{bmatrix} \sin(\lambda_\alpha - v_N + \eta\sigma_N^z) & \sin(2\eta)\sigma_N^- \\ \sin(2\eta)\sigma_N^+ & \sin(\lambda_\alpha - v_N - \eta\sigma_N^z) \end{bmatrix} \dots \\ &\times \begin{bmatrix} \sin(\lambda_\alpha - v_1 + \eta\sigma_1^z) & \sin(2\eta)\sigma_1^- \\ \sin(2\eta)\sigma_1^+ & \sin(\lambda_\alpha - v_1 - \eta\sigma_1^z) \end{bmatrix} . \end{aligned}$$

To determine the action-angle variables of the Hamiltonian formulation for the inhomogeneous six-vertex model, we present the arguments for the result below. First, we determine the form of each expression from carrying out the multiplication of two by two L operators. Specifically, we compute the series of matrices,

$$\left\{ \begin{bmatrix} A_i(\lambda_\alpha) & B_i(\lambda_\alpha) \\ C_i(\lambda_\alpha) & D_i(\lambda_\alpha) \end{bmatrix} \right\}_{1 \leq i \leq n} ,$$

where each entry of the  $n$  th  $L$  operator is determined by computing  $n - 1$  matrix multiplications.

## 2.2 Computing L operators

**Lemma 1** (*collecting terms from the product of two by two L operators*). The first entry of the  $L$  operator product,

$$\prod_{i=0}^3 L_{\alpha, N-i}(\lambda_\alpha, v_{N-i}) ,$$

has an expansion of the form,

$$\begin{aligned}
A_3(\lambda_\alpha) \equiv & \prod_{0 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) + (\sin(2\eta))^2 \left[ \left( \prod_{\substack{0 \leq i \leq 1 \\ i \equiv 0, - \\ i \equiv 1, +}} \sigma_{N-i}^{-,+} \right) \right. \\
& \times \left( \prod_{2 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) \\
& + \left( \prod_{\substack{2 \leq i \leq 3 \\ i \equiv 2, - \\ i \equiv 3, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) \\
& + (\sin(2\eta))^{-1} \left( \prod_{\substack{i \text{ odd} : 1 \leq i \leq 3 \\ i \equiv 1, - \\ i \equiv 3, +}} \sin(2\eta) \sigma_{N-i}^- \right) \left( \prod_{\substack{i \text{ even} : 0 \leq i \leq 2 \\ i \equiv 0, +\eta \\ i \equiv 2, -\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \\
& + \left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{i \text{ odd} : 1 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) \\
& + \left( \prod_{i \text{ even} : 0 \leq i \leq 2} \sigma_{N-i}^{-,+} \right) \left( \prod_{i \text{ odd} : 1 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) \\
& + \sin(2\eta) \left( \prod_{\substack{0 \leq i \leq 3 \\ i \equiv 0, - \\ i \equiv 1, + \\ i \equiv 2, - \\ i \equiv 3, +}} \sigma_{N-i}^{-,+} \right) \\
& \left. + \left( \prod_{\substack{i \text{ odd} : 1 \leq i \leq 3 \\ i \equiv 1, - \\ i \equiv 3, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) \right] ,
\end{aligned}$$

while the first entry of the L operator product,

$$\prod_{i=0}^2 L_{\alpha, N-i}(\lambda_\alpha, v_{N-i})$$

has an expansion of the form,

$$\begin{aligned}
A_2(\lambda_\alpha) \equiv & \prod_{0 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) + (\sin(2\eta))^2 \left( \prod_{\substack{0 \leq i \leq 1 \\ i \equiv 0, - \\ i \equiv 1, +}} \sigma_{N-i}^{-,+} \right) \left( \sin(\lambda_\alpha - v_{N-2} + \eta \sigma_{N-2}^z) \right) + (\sin(2\eta))^2 \\
& \times \left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \left( \sin(\lambda_\alpha - v_N + \eta \sigma_N^z) \right) + (\sin(2\eta))^2 \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \left( \sin(\lambda_\alpha - v_{N-1} - \eta \sigma_{N-1}^z) \right) .
\end{aligned}$$

*Proof of Lemma 1.* We collect terms from the first two terms in the product of L operators,

$$\begin{bmatrix} \sin(\lambda_\alpha - v_N + \eta \sigma_N^z) & \sin(2\eta) \sigma_N^- \\ \sin(2\eta) \sigma_N^+ & \sin(\lambda_\alpha - v_N - \eta \sigma_N^z) \end{bmatrix} \begin{bmatrix} \sin(\lambda_\alpha - v_{N-1} + \eta \sigma_{N-1}^z) & \sin(2\eta) \sigma_{N-1}^- \\ \sin(2\eta) \sigma_{N-1}^+ & \sin(\lambda_\alpha - v_{N-1} - \eta \sigma_{N-1}^z) \end{bmatrix}$$

from which one obtains a resultant matrix of the form,

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} ,$$

from the following expression for the first and second entries,

$$\begin{aligned} \mathbf{1}^0 &\equiv \mathbf{1} \equiv \sin(\lambda_\alpha - v_N + \eta\sigma_N^z) \sin(\lambda_\alpha - v_{N-1} + \eta\sigma_{N-1}^z) + \sin(2\eta)\sigma_N^- \sin(2\eta)\sigma_{N-1}^+ , \\ \mathbf{2}^0 &\equiv \mathbf{2} \equiv \sin(\lambda_\alpha - v_N + \eta\sigma_N^z) \sin(2\eta)\sigma_{N-1}^- + \sin(2\eta)\sigma_N^- \sin(\lambda_\alpha - v_{N-1} - \eta\sigma_{N-1}^z) . \end{aligned}$$

As we proceed with the computations, to determine the entries of each entry of the matrix of the  $L$  operator, we denote the entries of each such two by two matrix with  $\mathbf{1}^i$ ,  $\mathbf{2}^i$ ,  $\mathbf{3}^i$ , and  $\mathbf{4}^i$ , where  $0 \leq i \leq N-1$ . For the remaining terms in the product of  $L$  operators,

$$\begin{bmatrix} \sin(\lambda_\alpha - v_{N-2} + \eta\sigma_{N-2}^z) & \sin(2\eta)\sigma_{N-2}^- \\ \sin(2\eta)\sigma_{N-2}^+ & \sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) \end{bmatrix} \dots \times \begin{bmatrix} \sin(\lambda_\alpha - v_1 + \eta\sigma_1^z) & \sin(2\eta)\sigma_1^- \\ \sin(2\eta)\sigma_1^+ & \sin(\lambda_\alpha - v_1 - \eta\sigma_1^z) \end{bmatrix}$$

continuing the computation along similar lines, implies that the product of operators,

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} \sin(\lambda_\alpha - v_{N-2} + \eta\sigma_{N-2}^z) & \sin(2\eta)\sigma_{N-2}^- \\ \sin(2\eta)\sigma_{N-2}^+ & \sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) \end{bmatrix} \dots \times \begin{bmatrix} \sin(\lambda_\alpha - v_1 + \eta\sigma_1^z) & \sin(2\eta)\sigma_1^- \\ \sin(2\eta)\sigma_1^+ & \sin(\lambda_\alpha - v_1 - \eta\sigma_1^z) \end{bmatrix} ,$$

takes the form,

$$\begin{aligned} &\left[ \begin{array}{cc} 1\sin(\lambda_\alpha - v_{N-2} + \eta\sigma_{N-2}^z) + 2\sin(2\eta)\sigma_{N-2}^+ & 1\sin(2\eta)\sigma_{N-2}^- + 2\sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) \\ 3\sin(\lambda_\alpha - v_{N-2} + \eta\sigma_{N-2}^z) + 4\sin(2\eta)\sigma_{N-2}^+ & 3\sin(2\eta)\sigma_{N-2}^- + 4\sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) \end{array} \right] \\ &\quad \dots \times \begin{bmatrix} \sin(\lambda_\alpha - v_1 + \eta\sigma_1^z) & \sin(2\eta)\sigma_1^- \\ \sin(2\eta)\sigma_1^+ & \sin(\lambda_\alpha - v_1 - \eta\sigma_1^z) \end{bmatrix} . \end{aligned}$$

Distributing terms entrywise in the matrix with terms  $\mathbf{1}$ ,  $\mathbf{2}$ ,  $\mathbf{3}$ , and  $\mathbf{4}$  above yields the two by two matrix with entries, respectively given by  $\mathbf{1}^1$ ,  $\mathbf{2}^1$ ,  $\mathbf{3}^1$ , and  $\mathbf{4}^1$ ,

$$\begin{aligned} \mathbf{1}^1 &\equiv \mathbf{1}\sin(\lambda_\alpha - v_{N-2} + \eta\sigma_{N-2}^z) + \mathbf{2}\sin(2\eta)\sigma_{N-2}^+ \\ &\equiv \left( \sin(\lambda_\alpha - v_N + \eta\sigma_N^z) \sin(\lambda_\alpha - v_{N-1} + \eta\sigma_{N-1}^z) + \sin(2\eta)\sigma_N^- \sin(2\eta)\sigma_{N-1}^+ \right) \sin(\lambda_\alpha - v_{N-2} + \eta\sigma_{N-2}^z) \\ &\quad + \left( \sin(\lambda_\alpha - v_N + \eta\sigma_N^z) \sin(2\eta)\sigma_{N-1}^- + \sin(2\eta)\sigma_N^- \sin(\lambda_\alpha - v_{N-1} - \eta\sigma_{N-1}^z) \right) \sin(2\eta)\sigma_{N-2}^+ \\ &\equiv \sin(\lambda_\alpha - v_N + \eta\sigma_N^z) \sin(\lambda_\alpha - v_{N-1} + \eta\sigma_{N-1}^z) \sin(\lambda_\alpha - v_{N-2} + \eta\sigma_{N-2}^z) + \sin(2\eta)\sigma_N^- \sin(2\eta)\sigma_{N-1}^+ \\ &\quad \times \sin(\lambda_\alpha - v_{N-2} + \eta\sigma_{N-2}^z) + \sin(\lambda_\alpha - v_N + \eta\sigma_N^z) \sin(2\eta)\sigma_{N-1}^- \sin(2\eta)\sigma_{N-2}^+ \\ &\quad + \sin(2\eta)\sigma_N^- \sin(\lambda_\alpha - v_{N-1} - \eta\sigma_{N-1}^z) \sin(2\eta)\sigma_{N-2}^+ , \end{aligned}$$

corresponding to the first term,  $\mathbf{1}^1$ , which has the product representation,

$$\begin{aligned} &\prod_{0 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} + \eta\sigma_{N-i}^z) + (\sin(2\eta))^2 \left( \prod_{\substack{0 \leq i \leq 1 \\ i \equiv 0, - \\ i \equiv 1, +}} \sigma_{N-i}^{-,+} \right) \left( \sin(\lambda_\alpha - v_{N-2} + \eta\sigma_{N-2}^z) \right) + (\sin(2\eta))^2 \\ &\times \left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \left( \sin(\lambda_\alpha - v_N + \eta\sigma_N^z) \right) + (\sin(2\eta))^2 \left( \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \left( \sin(\lambda_\alpha - v_{N-1} - \eta\sigma_{N-1}^z) \right) , \end{aligned}$$

as well as the following product representation for  $\mathbf{2}^1$ , in which,

$$\begin{aligned} \mathbf{2}^1 &\equiv (\sin(2\eta)\sigma_{N-2}^-) \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} + \eta\sigma_{N-i}^z) + (\sin(2\eta))^3 \left( \prod_{\substack{0 \leq i \leq 2 \\ i \equiv 0, - \\ i \equiv 1, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) + (\sin(2\eta)\sigma_{N-1}^-) \\ &\quad \times \left( \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, +\eta \\ i \equiv 2, -\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta\sigma_{N-i}^z) \right) + (\sin(2\eta)\sigma_N^-) \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta\sigma_{N-i}^z) , \end{aligned}$$

from the expansion,

$$\mathbf{2}^1 \equiv \mathbf{1}\sin(2\eta)\sigma_{N-2}^- + \mathbf{2}\sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z)$$

From the two by two matrix with entries  $\mathbf{1}^1$ ,  $\mathbf{2}^1$ ,  $\mathbf{3}^1$  and  $\mathbf{4}^1$ , the remaining terms of the product takes the form,

$$\begin{bmatrix} \mathbf{1}^1 & \mathbf{2}^1 \\ \mathbf{3}^1 & \mathbf{4}^1 \end{bmatrix} \begin{bmatrix} \sin(\lambda_\alpha - v_{N-3} + \eta\sigma_{N-3}^z) & \sin(2\eta)\sigma_{N-3}^- \\ \sin(2\eta)\sigma_{N-3}^+ & \sin(\lambda_\alpha - v_{N-3} - \eta\sigma_{N-3}^z) \end{bmatrix} \dots \times \begin{bmatrix} \sin(\lambda_\alpha - v_1 + \eta\sigma_1^z) & \sin(2\eta)\sigma_1^- \\ \sin(2\eta)\sigma_1^+ & \sin(\lambda_\alpha - v_1 - \eta\sigma_1^z) \end{bmatrix} ,$$

which implies, for,

$$\begin{bmatrix} \mathbf{1}^1 & \mathbf{2}^1 \\ \mathbf{3}^1 & \mathbf{4}^1 \end{bmatrix} \begin{bmatrix} \sin(\lambda_\alpha - v_{N-3} + \eta\sigma_{N-3}^z) & \sin(2\eta)\sigma_{N-3}^- \\ \sin(2\eta)\sigma_{N-3}^+ & \sin(\lambda_\alpha - v_{N-3} - \eta\sigma_{N-3}^z) \end{bmatrix} \equiv \begin{bmatrix} \mathbf{1}^2 & \mathbf{2}^2 \\ \mathbf{3}^2 & \mathbf{4}^2 \end{bmatrix}$$

has a first entry that is given by,

$$\mathbf{1}^2 \equiv \mathbf{1}^1\sin(\lambda_\alpha - v_{N-3} + \eta\sigma_{N-3}^z) + \mathbf{2}^1\sin(2\eta)\sigma_{N-3}^+ .$$

The first entry of the resultant matrix above is given by,

$$\begin{aligned} \mathbf{1}^2 &= \left[ \prod_{0 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} + \eta\sigma_{N-i}^z) + (\sin(2\eta))^2 \left( \prod_{\substack{0 \leq i \leq 1 \\ i \equiv 0, - \\ i \equiv 1, +}} \sigma_{N-i}^{-,+} \right) \left( \sin(\lambda_\alpha - v_{N-2} + \eta\sigma_{N-2}^z) \right) + (\sin(2\eta))^2 \right. \\ &\quad \times \left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \left( \sin(\lambda_\alpha - v_N + \eta\sigma_N^z) \right) + (\sin(2\eta))^2 \left( \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \left( \sin(\lambda_\alpha - v_{N-1} - \eta\sigma_{N-1}^z) \right) \right] \\ &\quad \times \sin(\lambda_\alpha - v_{N-3} + \eta\sigma_{N-3}^z) \\ &\quad + \left[ (\sin(2\eta)\sigma_{N-2}^-) \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} + \eta\sigma_{N-i}^z) \right) + (\sin(2\eta))^3 \left( \prod_{\substack{0 \leq i \leq 2 \\ i \equiv 0, - \\ i \equiv 1, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) + (\sin(2\eta)\sigma_{N-1}^-) \right. \\ &\quad \times \left( \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, +\eta \\ i \equiv 2, -\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta\sigma_{N-i}^z) \right) + \left( (\sin(2\eta)\sigma_N^-) \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta\sigma_{N-i}^z) \right) \right] \sin(2\eta)\sigma_{N-3}^+ , \end{aligned}$$

which has the following equivalent product representation, after collecting terms,

$$\begin{aligned}
& \prod_{0 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) + (\sin(2\eta))^2 \left( \prod_{\substack{0 \leq i \leq 1 \\ i \equiv 0, - \\ i \equiv 1, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{2 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) \\
& + (\sin(2\eta))^2 \left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{i \text{ odd}: 1 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) + (\sin(2\eta))^2 \\
& \times \left( \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{i \text{ odd}: 1 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) \\
& + (\sin(2\eta))^2 \left( \prod_{\substack{2 \leq i \leq 3 \\ i \equiv 2, - \\ i \equiv 3, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) \\
& + (\sin(2\eta))^3 \left( \prod_{\substack{0 \leq i \leq 3 \\ i \equiv 0, - \\ i \equiv 1, + \\ i \equiv 2, - \\ i \equiv 3, +}} \sigma_{N-i}^{-,+} \right) + \left( \prod_{i \text{ odd}: 1 \leq i \leq 3} \sin(2\eta) \sigma_{N-i}^- \right) \\
& \times \left( \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, +\eta \\ i \equiv 2, -\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) + (\sin(2\eta))^2 \left( \prod_{\substack{\text{odd } i: 1 \leq i \leq 3 \\ i \equiv 1, - \\ i \equiv 3, +}} \sigma_{N-i}^{-,+} \right) \\
& \times \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) .
\end{aligned}$$

Collecting terms under the prefactor  $(\sin(\eta))^2$ ,  $\sin(2\eta)$ , and  $(\sin(2\eta))^{-1}$ , from the superposition above, gives,

$$\begin{aligned}
& \prod_{0 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) + (\sin(\eta))^2 \left[ \left( \prod_{\substack{0 \leq i \leq 1 \\ i \equiv 0, - \\ i \equiv 1, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{2 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) \right. \\
& + \left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{i \text{ odd}: 1 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) + \left( \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \\
& \times \left. \left( \prod_{i \text{ odd}: 1 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) + \left( \prod_{\substack{2 \leq i \leq 3 \\ i \equiv 2, - \\ i \equiv 3, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) \right. \\
& \quad \left. + \sin(2\eta) \left( \prod_{\substack{0 \leq i \leq 3 \\ i \equiv 0, - \\ i \equiv 1, + \\ i \equiv 2, - \\ i \equiv 3, +}} \sigma_{N-i}^{-,+} \right) \right. \\
& + (\sin(2\eta))^{-2} \left( \prod_{i \text{ odd}: 1 \leq i \leq 3} \sin(2\eta) \sigma_{N-i}^{-,+} \right) \left( \prod_{i \text{ even}: 0 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right)
\end{aligned}$$

$$+ \left( \prod_{\substack{i \text{ odd} : 1 \leq i \leq 3 \\ i \equiv 1, - \\ i \equiv 3, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) \right] .$$

From the superposition above, grouping together terms depending upon whether there is a product of even, or odd, terms of,

$$\sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) ,$$

or upon whether there is a product of terms of,

$$\sigma_{N-i}^{-,+} ,$$

implies that the expression for  $A(\lambda_\alpha)$ , after repeating the computation for the two by two matrix multiplication  $n$  times, can be obtained by grouping together terms under the  $(\sin(2\eta))^2$ , in which,

$$\left( \prod_{\substack{0 \leq i \leq 1 \\ i \equiv 0, - \\ i \equiv 1, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{2 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) + \left( \prod_{\substack{2 \leq i \leq 3 \\ i \equiv 2, - \\ i \equiv 3, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) ,$$

corresponding to a first group of terms,

$$(\sin(2\eta))^{-1} \left( \prod_{\substack{i \text{ odd} : 1 \leq i \leq 3 \\ i \equiv 1, - \\ i \equiv 3, +}} \sin(2\eta) \sigma_{N-i}^- \right) \left( \prod_{\substack{i \text{ even} : 0 \leq i \leq 2 \\ i \equiv 0, +\eta \\ i \equiv 2, -\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) ,$$

corresponding to a second group of terms,

$$\left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{i \text{ odd}: 1 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) ,$$

corresponding to a third group of terms,

$$\left( \prod_{i \text{ even} : 0 \leq i \leq 2} \sigma_{N-i}^{-,+} \right) \left( \prod_{i \text{ odd} : 1 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) ,$$

corresponding to a fourth group of terms, and,

$$\sin(2\eta) \left( \prod_{\substack{0 \leq i \leq 3 \\ i \equiv 0, - \\ i \equiv 1, + \\ i \equiv 2, - \\ i \equiv 3, +}} \sigma_{N-i}^{-,+} \right) ,$$

corresponding to a fifth group of terms, and,

$$\left( \prod_{i \text{ odd} : 1 \leq i \leq 3} \sigma_{N-i}^{-,+} \right) \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) ,$$

corresponding to a sixth group of terms, hence implying the desired form for the first coefficient,

$$\begin{aligned}
A_3(\lambda_\alpha) \equiv & \prod_{0 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) + (\sin(2\eta))^2 \left[ \left( \prod_{\substack{0 \leq i \leq 1 \\ i \equiv 0, - \\ i \equiv 1, +}} \sigma_{N-i}^{-,+} \right) \right. \\
& \times \left( \prod_{2 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) \\
& + \left( \prod_{\substack{2 \leq i \leq 3 \\ i \equiv 2, - \\ i \equiv 3, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) \\
& + (\sin(2\eta))^{-1} \left( \prod_{\substack{i \text{ odd} : 1 \leq i \leq 3 \\ i \equiv 1, - \\ i \equiv 3, +}} \sin(2\eta) \sigma_{N-i}^- \right) \left( \prod_{\substack{i \text{ even} : 0 \leq i \leq 2 \\ i \equiv 0, +\eta \\ i \equiv 2, -\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \\
& + \left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{i \text{ odd} : 1 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) \\
& + \left( \prod_{i \text{ even} : 0 \leq i \leq 2} \sigma_{N-i}^{-,+} \right) \left( \prod_{i \text{ odd} : 1 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) \\
& + \sin(2\eta) \left( \prod_{\substack{0 \leq i \leq 3 \\ i \equiv 0, - \\ i \equiv 1, + \\ i \equiv 2, - \\ i \equiv 3, +}} \sigma_{N-i}^{-,+} \right) \\
& \left. + \left( \prod_{i \text{ odd} : 1 \leq i \leq 3} \sigma_{N-i}^{-,+} \right) \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) \right] ,
\end{aligned}$$

from which we conclude the argument.  $\square$

**Lemma 2** (*collecting terms from the product of two by two L operators*). The second entry of the L operator product,

$$\prod_{i=0}^3 L_{\alpha, N-i}(\lambda_\alpha, v_{N-i}) ,$$

has an expansion of the form,

$$\begin{aligned}
B_3(\lambda_\alpha) \equiv & (\sin(2\eta)) \sigma_{N-3}^- \left( \prod_{0 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) + (\sin(2\eta))^3 \left[ (\sin(2\eta))^{-1} \right. \\
& \times \left[ \left( \prod_{\substack{0 \leq i \leq 1 \\ i \equiv 0, - \\ i \equiv 1, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{\substack{2 \leq i \leq 3 \\ i \equiv 2, + \\ i \equiv 3, -}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \right. \\
& + \left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{\substack{\text{odd } i : 1 \leq i \leq 3 \\ i \equiv 1, - \\ i \equiv 3, +}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \\
& \quad \left. + \sigma_{N-2}^- \left( \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) \right. \right. \\
& + \left( \prod_{\substack{i \equiv 0, \text{ odd } i : 1 \leq i \leq 3 \\ i \equiv 1, + \\ i \equiv 3, -}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \left. \right) \\
& + \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) \\
& + \sigma_N^- \sin(\lambda_\alpha - v_{N-2} - \eta \sigma_{N-2}^z) \left( \prod_{1 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) \\
& + (1 + \sigma_{N-1}^-) \left[ \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, + \\ i \equiv 2, -}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \right. \\
& \quad \left. + \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, + \\ i \equiv 2, -}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \right] \\
& + (\sin(2\eta)) \left[ \left( \prod_{\substack{0 \leq i \leq 2 \\ i \equiv 0, - \\ i \equiv 1, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) + \sigma_{N-2}^- \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, -}} \sigma_{N-i}^{-,+} \right) \left( \prod_{\substack{\text{odd } i : 1 \leq i \leq 3 \\ i \equiv 1, - \\ i \equiv 3, +}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \right. \\
& \quad \left. + \left( \prod_{\substack{0 \leq i \leq 2 \\ i \equiv 0, 2, - \\ i \equiv 1, +}} \sigma_{N-i}^{-,+} \right) + \left( \prod_{\substack{0 \leq i \leq 1, i \equiv 3 \\ i \equiv 0, 1, +\eta \\ i \equiv 3, -\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \right. \\
& \quad \left. + \left( \prod_{\substack{0 \leq i \leq 2 \\ i \equiv 1, + \\ i \equiv 3, -\eta}} \sigma_{N-i}^{-,+} \right) + \left( \prod_{\substack{0 \leq i \leq 1, i \equiv 3 \\ i \equiv 0, 1, +\eta \\ i \equiv 3, -\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) + \sigma_{N-2}^- \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, -}} \sigma_{N-i}^{-,+} \right) \right. \\
& \quad \left. + \left( \prod_{\substack{0 \leq i \leq 2 \\ i \equiv 0, - \\ i \equiv 1, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) \right] \\
& + \left( \prod_{\substack{0 \leq i \leq 1 \\ i \equiv 0, + \\ i \equiv 1, -}} \sigma_{N-i}^{-,+} \right) \sin(\lambda_\alpha - v_{N-2} + \eta \sigma_{N-2}^z) + \left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \sin(\lambda_\alpha - v_N + \eta \sigma_N^z) \\
& \quad + \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \sin(\lambda_\alpha - v_N + \eta \sigma_N^z) \]
\end{aligned}$$

while the second entry of the L operator product,

$$\prod_{i=0}^2 L_{\alpha, N-i}(\lambda_\alpha, v_{N-i})$$

has an expansion of the form,

$$B_2(\lambda_\alpha) \equiv (\sin(2\eta)\sigma_{N-2}^-) \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} + \eta\sigma_{N-i}^z) \right) + (\sin(2\eta))^3 \left( \prod_{\substack{0 \leq i \leq 2 \\ i \equiv 0, - \\ i \equiv 1, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) + (\sin(2\eta)\sigma_{N-1}^-) \times \left( \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, +\eta \\ i \equiv 2, -\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta\sigma_{N-i}^z) \right) + (\sin(2\eta)\sigma_N^-) \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta\sigma_{N-i}^z) \right) .$$

*Proof of Lemma 2.* We collect terms from the first two terms in the product of L operators,

$$\begin{bmatrix} \sin(\lambda_\alpha - v_N + \eta\sigma_N^z) & \sin(2\eta)\sigma_N^- \\ \sin(2\eta)\sigma_N^+ & \sin(\lambda_\alpha - v_N - \eta\sigma_N^z) \end{bmatrix} \begin{bmatrix} \sin(\lambda_\alpha - v_{N-1} + \eta\sigma_{N-1}^z) & \sin(2\eta)\sigma_{N-1}^- \\ \sin(2\eta)\sigma_{N-1}^+ & \sin(\lambda_\alpha - v_{N-1} - \eta\sigma_{N-1}^z) \end{bmatrix}$$

from which one obtains a resultant matrix of the form,

$$\begin{bmatrix} \mathbf{1} & \mathbf{2} \\ \mathbf{3} & \mathbf{4} \end{bmatrix} ,$$

which has the following expression for the the first and second entries, with,

$$\begin{aligned} \mathbf{1}^0 &\equiv \mathbf{1} \equiv \sin(\lambda_\alpha - v_N + \eta\sigma_N^z) \sin(\lambda_\alpha - v_{N-1} + \eta\sigma_{N-1}^z) + \sin(2\eta)\sigma_N^- \sin(2\eta)\sigma_{N-1}^+ , \\ \mathbf{2}^0 &\equiv \mathbf{2} \equiv \sin(\lambda_\alpha - v_N + \eta\sigma_N^z) \sin(2\eta)\sigma_{N-1}^- + \sin(2\eta)\sigma_N^- \sin(\lambda_\alpha - v_{N-1} - \eta\sigma_{N-1}^z) , \end{aligned}$$

From the product of operators,

$$\begin{bmatrix} \mathbf{1} & \mathbf{2} \\ \mathbf{3} & \mathbf{4} \end{bmatrix} \begin{bmatrix} \sin(\lambda_\alpha - v_{N-2} + \eta\sigma_{N-2}^z) & \sin(2\eta)\sigma_{N-2}^- \\ \sin(2\eta)\sigma_{N-2}^+ & \sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) \end{bmatrix} \dots \times \begin{bmatrix} \sin(\lambda_\alpha - v_1 + \eta\sigma_1^z) & \sin(2\eta)\sigma_1^- \\ \sin(2\eta)\sigma_1^+ & \sin(\lambda_\alpha - v_1 - \eta\sigma_1^z) \end{bmatrix} ,$$

which takes the form,

$$\begin{aligned} &\begin{bmatrix} \mathbf{1}\sin(\lambda_\alpha - v_{N-2} + \eta\sigma_{N-2}^z) + \mathbf{2}\sin(2\eta)\sigma_{N-2}^+ & \mathbf{1}\sin(2\eta)\sigma_{N-2}^- + \mathbf{2}\sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) \\ \mathbf{3}\sin(\lambda_\alpha - v_{N-2} + \eta\sigma_{N-2}^z) + \mathbf{4}\sin(2\eta)\sigma_{N-2}^+ & \mathbf{3}\sin(2\eta)\sigma_{N-2}^- + \mathbf{4}\sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) \end{bmatrix} \\ &\dots \times \begin{bmatrix} \sin(\lambda_\alpha - v_1 + \eta\sigma_1^z) & \sin(2\eta)\sigma_1^- \\ \sin(2\eta)\sigma_1^+ & \sin(\lambda_\alpha - v_1 - \eta\sigma_1^z) \end{bmatrix} , \end{aligned}$$

performing rearrangements for the next term yields,

$$\begin{aligned} \mathbf{2}^1 &\equiv \mathbf{1}\sin(2\eta)\sigma_{N-2}^- + \mathbf{2}\sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) \\ &\equiv \left( \sin(\lambda_\alpha - v_N + \eta\sigma_N^z) \sin(\lambda_\alpha - v_{N-1} + \eta\sigma_{N-1}^z) + \sin(2\eta)\sigma_N^- \sin(2\eta)\sigma_{N-1}^+ \right) \sin(2\eta)\sigma_{N-2}^- \\ &+ \left( \sin(\lambda_\alpha - v_N + \eta\sigma_N^z) \sin(2\eta)\sigma_{N-1}^- + \sin(2\eta)\sigma_N^- \sin(\lambda_\alpha - v_{N-1} - \eta\sigma_{N-1}^z) \right) \sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) , \\ &\equiv \sin(\lambda_\alpha - v_N + \eta\sigma_N^z) \sin(\lambda_\alpha - v_{N-1} + \eta\sigma_{N-1}^z) \sin(2\eta)\sigma_{N-2}^- + \sin(2\eta)\sigma_N^- \sin(2\eta)\sigma_{N-1}^+ \sin(2\eta)\sigma_{N-2}^- \\ &\quad + \sin(\lambda_\alpha - v_N + \eta\sigma_N^z) \sin(2\eta)\sigma_{N-1}^- \sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) \\ &\quad + \sin(2\eta)\sigma_N^- \sin(\lambda_\alpha - v_{N-1} - \eta\sigma_{N-1}^z) \sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) , \end{aligned}$$

corresponding to the second term,  $\mathbf{2}^1$  which has the product representation,

$$\begin{aligned} & (\sin(2\eta)\sigma_{N-2}^-) \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} + \eta\sigma_{N-i}^z) \right) + (\sin(2\eta))^3 \left( \prod_{\substack{0 \leq i \leq 2 \\ i \equiv 0, - \\ i \equiv 1, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) + (\sin(2\eta)\sigma_{N-1}^-) \\ & \times \left( \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, +\eta \\ i \equiv 2, -\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta\sigma_{N-i}^z) \right) + (\sin(2\eta)\sigma_N^-) \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta\sigma_{N-i}^z) \right) . \end{aligned}$$

From the two by two matrix with entries  $\mathbf{1}^1$ ,  $\mathbf{2}^1$ ,  $\mathbf{3}^1$  and  $\mathbf{4}^1$ , the remaining terms of the product takes the form,

$$\begin{bmatrix} \mathbf{1}^1 & \mathbf{2}^1 \\ \mathbf{3}^1 & \mathbf{4}^1 \end{bmatrix} \begin{bmatrix} \sin(\lambda_\alpha - v_{N-3} + \eta\sigma_{N-3}^z) & \sin(2\eta)\sigma_{N-3}^- \\ \sin(2\eta)\sigma_{N-3}^+ & \sin(\lambda_\alpha - v_{N-3} - \eta\sigma_{N-3}^z) \end{bmatrix} \dots \times \begin{bmatrix} \sin(\lambda_\alpha - v_1 + \eta\sigma_1^z) & \sin(2\eta)\sigma_1^- \\ \sin(2\eta)\sigma_1^+ & \sin(\lambda_\alpha - v_1 - \eta\sigma_1^z) \end{bmatrix} ,$$

which implies, for,

$$\begin{bmatrix} \mathbf{1}^1 & \mathbf{2}^1 \\ \mathbf{3}^1 & \mathbf{4}^1 \end{bmatrix} \begin{bmatrix} \sin(\lambda_\alpha - v_{N-3} + \eta\sigma_{N-3}^z) & \sin(2\eta)\sigma_{N-3}^- \\ \sin(2\eta)\sigma_{N-3}^+ & \sin(\lambda_\alpha - v_{N-3} - \eta\sigma_{N-3}^z) \end{bmatrix} \equiv \begin{bmatrix} \mathbf{1}^2 & \mathbf{2}^2 \\ \mathbf{3}^2 & \mathbf{4}^2 \end{bmatrix}$$

has a second entry that is given by,

$$\mathbf{2}^2 \equiv \mathbf{1}^1 \sin(2\eta)\sigma_{N-3}^- + \mathbf{2}^1 \sin(\lambda_\alpha - v_{N-3} - \eta\sigma_{N-3}^z) .$$

The superposition above has the equivalent product representation, after collecting terms,

$$\begin{aligned} \mathbf{2}^2 = & \left[ \mathbf{1}^1 \sin(\lambda_\alpha - v_{N-2} + \eta\sigma_{N-2}^z) + \mathbf{2}^1 \sin(2\eta)\sigma_{N-2}^+ \right] \sin(2\eta)\sigma_{N-3}^- \\ & + \left[ \mathbf{1} \sin(2\eta)\sigma_{N-2}^- + \mathbf{2} \sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) \right] \sin(\lambda_\alpha - v_{N-3} - \eta\sigma_{N-3}^z) \end{aligned}$$

is equivalent to,

$$\begin{aligned} & \left[ \left( \left( \prod_{0 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} + \eta\sigma_{N-i}^z) \right) + (\sin(2\eta))^2 \left( \prod_{\substack{0 \leq i \leq 1 \\ i \equiv 0, - \\ i \equiv 1, +}} \sigma_{N-i}^{-,+} \right) \left( \sin(\lambda_\alpha - v_{N-2} + \eta\sigma_{N-2}^z) \right) + (\sin(2\eta))^2 \right. \right. \\ & \times \left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \left( \sin(\lambda_\alpha - v_N + \eta\sigma_N^z) \right) + (\sin(2\eta))^2 \left( \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \left( \sin(\lambda_\alpha - v_{N-1} - \eta\sigma_{N-1}^z) \right) \left. \right) \\ & \quad \times \sin(\lambda_\alpha - v_{N-2} + \eta\sigma_{N-2}^z) \\ & + \left( (\sin(2\eta)\sigma_{N-2}^-) \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} + \eta\sigma_{N-i}^z) \right) + (\sin(2\eta))^3 \prod_{\substack{0 \leq i \leq 2 \\ i \equiv 0, - \\ i \equiv 1, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} + (\sin(2\eta)\sigma_{N-1}^-) \right. \\ & \quad \times \left. \left( \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, +\eta \\ i \equiv 2, -\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta\sigma_{N-i}^z) \right) + (\sin(2\eta)\sigma_N^-) \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta\sigma_{N-i}^z) \right) \right) \right] \sin(2\eta)\sigma_{N-3}^- \\ & + \left[ \left( \prod_{0 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} + \eta\sigma_{N-i}^z) + (\sin(2\eta))^2 \left( \prod_{\substack{0 \leq i \leq 1 \\ i \equiv 0, - \\ i \equiv 1, +}} \sigma_{N-i}^{-,+} \right) \left( \sin(\lambda_\alpha - v_{N-2} + \eta\sigma_{N-2}^z) \right) + (\sin(2\eta))^2 \right. \right. \end{aligned}$$

$$\begin{aligned}
& \times \left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \left( \sin(\lambda_\alpha - v_N + \eta \sigma_N^z) \right) + (\sin(2\eta))^2 \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \left( \sin(\lambda_\alpha - v_{N-1} - \eta \sigma_{N-1}^z) \right) \\
& \quad \times \sin(2\eta) \sigma_{N-2}^- + \left( (\sin(2\eta) \sigma_{N-2}^-) \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) + (\sin(2\eta))^3 \left( \prod_{\substack{0 \leq i \leq 2 \\ i \equiv 0, - \\ i \equiv 1, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) \right. \\
& \quad \left. + (\sin(2\eta) \sigma_{N-1}^-) \right)
\end{aligned}$$

$$\begin{aligned}
& \times \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, +\eta \\ i \equiv 2, -\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) + (\sin(2\eta) \sigma_N^-) \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) \\
& \quad \times \sin(\lambda_\alpha - v_{N-2} - \eta \sigma_{N-2}^z) \left[ \sin(\lambda_\alpha - v_{N-3} - \eta \sigma_{N-3}^z) \right] .
\end{aligned}$$

Collecting terms under the prefactor  $(\sin(\eta))^3$  gives,

$$\begin{aligned}
& (\sin(2\eta)) \sigma_{N-3}^- \left( \prod_{0 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) + (\sin(2\eta))^3 \left[ \left( \prod_{\substack{0 \leq i \leq 1 \\ i \equiv 0, + \\ i \equiv 1, +}} \sigma_{N-i}^{-,+} \right) \right. \\
& \quad \times \sin(\lambda_\alpha - v_{N-2} + \eta \sigma_{N-2}^z) + \left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \sin(\lambda_\alpha - v_N + \eta \sigma_N^z) + \cdots \\
& \quad \left. \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \sin(\lambda_\alpha - v_N + \eta \sigma_N^z) \right. \\
& \quad \left. + (\sin(2\eta))^{-1} \sigma_{N-2}^- \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) \right. \\
& \quad \left. + (\sin(2\eta)) \left( \prod_{\substack{0 \leq i \leq 2 \\ i \equiv 0, - \\ i \equiv 1, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) + (\sin(2\eta))^{-1} \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, +\eta \\ i \equiv 2, -\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \right. \\
& \quad \left. + (\sin(2\eta))^{-1} \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) + (\sin(2\eta))^{-1} \left( \prod_{\substack{0 \leq i \leq 1 \\ i \equiv 0, - \\ i \equiv 1, +}} \sigma_{N-i}^{-,+} \right) \right. \\
& \quad \left. \times \left( \prod_{\substack{2 \leq i \leq 3 \\ i \equiv 2, \eta+ \\ i \equiv 3, \eta-}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) + (\sin(2\eta))^{-1} \left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, -}} \sigma_{N-i}^{-,+} \right) \left( \prod_{\substack{\text{odd } i : 1 \leq i \leq 3 \\ i \equiv 1, - \\ i \equiv 2, +}} \sin(\lambda_\alpha - v_N \pm \eta \sigma_{N-i}^z) \right) \right)
\end{aligned}$$

$$\begin{aligned}
& + (\sin(2\eta)) \sigma_{N-2}^- \left( \prod_{\text{even } i : 0 \leq i \leq 2} \sigma_{N-i}^{-,+} \right) \left( \prod_{\substack{\text{odd } i : 1 \leq i \leq 3 \\ i \equiv 1, - \\ i \equiv 3, +}} \sin(\lambda_\alpha - v_N \pm \eta \sigma_{N-i}^z) \right) \\
& + (\sin(2\eta))^{-1} \sigma_{N-2}^- \left( \prod_{\substack{i=0, \text{ odd } i : 1 \leq i \leq 3 \\ i \equiv 0, + \\ i \equiv 1, + \\ i \equiv 3, -}} \sin(\lambda_\alpha - v_{N-1} \pm \eta \sigma_{N-i}^z) \right) + (\sin(2\eta)) \left( \prod_{\substack{0 \leq i \leq 1, i \equiv 3 \\ i \equiv 0, 1, +\eta \\ i \equiv 3, -\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \\
& + (\sin(2\eta)) \left( \prod_{\substack{0 \leq i \leq 2 \\ i \equiv 0, 2, - \\ i \equiv 1, +}} \sigma_{N-i}^{-,+} \right) + (\sin(2\eta))^{-1} \sigma_{N-1}^- \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2, i \equiv 3 \\ i \equiv 0, +\eta \\ i \equiv 2, -\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \\
& + (\sin(2\eta))^{-1} \sigma_N^- \sin(\lambda_\alpha - v_{N-2} - \eta \sigma_{N-2}^z) \left( \prod_{1 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) .
\end{aligned}$$

Under the  $(\sin(2\eta))^3$  prefactor, further grouping terms under  $(\sin(2\eta))^{-1}$  yields,

$$\begin{aligned}
& (\sin(2\eta))^{-1} \left[ \sigma_{N-2}^- \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) + \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, +\eta \\ i \equiv 2, -\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \right. \\
& \left. + \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) + \left( \prod_{\substack{0 \leq i \leq 1 \\ i \equiv 0, - \\ i \equiv 1, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{\substack{2 \leq i \leq 3 \\ i \equiv 2, \eta+ \\ i \equiv 3, \eta-}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \right. \\
& \left. + \sigma_{N-2}^- \left( \prod_{\substack{i=0, \text{ odd } i : 1 \leq i \leq 3 \\ i \equiv 0, + \\ i \equiv 1, + \\ i \equiv 3, -}} \sin(\lambda_\alpha - v_{N-1} \pm \eta \sigma_{N-i}^z) \right) + \sigma_{N-1}^- \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2, i \equiv 3 \\ i \equiv 0, +\eta \\ i \equiv 2, -\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \right. \\
& \left. + \sigma_N^- \sin(\lambda_\alpha - v_{N-2} - \eta \sigma_{N-2}^z) \left( \prod_{1 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) \right] .
\end{aligned}$$

Under the  $(\sin(2\eta))^3$  prefactor, further grouping terms under  $(\sin(2\eta))$  yields,

$$\begin{aligned}
& (\sin(2\eta)) \left[ \left( \prod_{\substack{0 \leq i \leq 2 \\ i \equiv 0, - \\ i \equiv 1, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) + \sigma_{N-2}^- \left( \prod_{\text{even } i : 0 \leq i \leq 2} \sigma_{N-i}^{-,+} \right) \left( \prod_{\substack{\text{odd } i : 1 \leq i \leq 3 \\ i \equiv 1, - \\ i \equiv 3, +}} \sin(\lambda_\alpha - v_N \pm \eta \sigma_{N-i}^z) \right) \right. \\
& \left. + \left( \prod_{\substack{0 \leq i \leq 2 \\ i \equiv 0, 2, - \\ i \equiv 1, +}} \sigma_{N-i}^{-,+} \right) + \left( \prod_{\substack{0 \leq i \leq 1, i \equiv 3 \\ i \equiv 0, 1, +\eta \\ i \equiv 3, -\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \right] .
\end{aligned}$$

Besides the group of terms above, under the  $(\sin(2\eta))^3$ , grouping terms under  $(\sin(2\eta))$  yields,

$$\begin{aligned}
& (\sin(2\eta)) \left[ \left( \prod_{\substack{0 \leq i \leq 2 \\ i \equiv 0, 2, - \\ i \equiv 1, +}} \sigma_{N-i}^{-,+} \right) + \left( \prod_{\substack{0 \leq i \leq 1, i \equiv 3 \\ i \equiv 0, 1, +\eta \\ i \equiv 3, -\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) + \sigma_{N-2}^- \left( \prod_{\text{even } i : 0 \leq i \leq 2} \sigma_{N-i}^{-,+} \right) \right. \\
& \left. + \left( \prod_{\substack{0 \leq i \leq 2 \\ i \equiv 0, - \\ i \equiv 1, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) \right] ,
\end{aligned}$$

Lastly, one has other terms, which are of the form,

$$\left( \prod_{\substack{0 \leq i \leq 1 \\ i \equiv 0, + \\ i \equiv 1, -}} \sigma_{N-i}^{-,+} \right) \sin(\lambda_\alpha - v_{N-2} + \eta \sigma_{N-2}^z) + \left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \sin(\lambda_\alpha - v_N + \eta \sigma_N^z)$$

$$+ \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \sin(\lambda_\alpha - v_N + \eta \sigma_N^z) .$$

Under the  $(\sin(2\eta))^3$  prefactor, further grouping terms under  $(\sin(2\eta))^{-1}$  yields,

$$(\sin(2\eta))^3 \left[ \cdots + (\sin(2\eta))^{-1} \left[ \sigma_{N-2}^- \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) \right. \right.$$

$$+ \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, + \\ i \equiv 2, -}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right)$$

$$+ \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) + \left( \prod_{\substack{0 \leq i \leq 1 \\ i \equiv 0, - \\ i \equiv 1, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{\substack{2 \leq i \leq 3 \\ i \equiv 2, + \\ i \equiv 3, -}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right)$$

$$+ \left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{\substack{\text{odd } i : 1 \leq i \leq 3 \\ i \equiv 1, - \\ i \equiv 3, +}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right)$$

$$+ \sigma_{N-2}^- \left( \prod_{\substack{i=0, \text{ odd } i : 1 \leq i \leq 3 \\ i \equiv 1, - \\ i \equiv 3, +}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) + \sigma_{N-1}^- \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, + \\ i \equiv 2, -}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right)$$

$$\left. \left. + \sigma_N^- \sin(\lambda_\alpha - v_{N-2} - \eta \sigma_{N-2}^z) \left( \prod_{1 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) \right] \right] .$$

From the first group of terms that were obtained under the  $(\sin(2\eta))^{-1}$  prefactor, contained within the  $(\sin(2\eta))^3$  prefactor, rewriting all product terms with similar notation as given in the expression for **1<sup>2</sup>** from **Lemma 1** yields,

$$\left( \prod_{\substack{0 \leq i \leq 1 \\ i \equiv 0, - \\ i \equiv 1, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{\substack{2 \leq i \leq 3 \\ i \equiv 2, + \\ i \equiv 3, -}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) + \left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{\substack{\text{odd } i : 1 \leq i \leq 3 \\ i \equiv 1, - \\ i \equiv 3, +}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) ,$$

corresponding to a first group of terms,

$$\sigma_{N-2}^- \left( \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) + \left( \prod_{\substack{i=0, \text{ odd } i : 1 \leq i \leq 3 \\ i \equiv 1, + \\ i \equiv 3, -}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \right) ,$$

corresponding to a second group of terms,

$$\left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) + \sigma_N^- \sin(\lambda_\alpha - v_{N-2} - \eta \sigma_{N-2}^z) \left( \prod_{1 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) ,$$

corresponding to a third group of terms, and,

$$(1 + \sigma_{N-1}^-) \left[ \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, + \\ i \equiv 2, -}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) + \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, + \\ i \equiv 2, -}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \right] ,$$

corresponding to a fifth group of terms.

Altogether,

$$\begin{aligned} B_3(\lambda_\alpha) \equiv & (\sin(2\eta)) \sigma_{N-3}^- \left( \prod_{0 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) + (\sin(2\eta))^3 \left[ (\sin(2\eta))^{-1} \right. \\ & \times \left[ \left( \prod_{\substack{0 \leq i \leq 1 \\ i \equiv 0, - \\ i \equiv 1, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{\substack{2 \leq i \leq 3 \\ i \equiv 2, + \\ i \equiv 3, -}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \right. \\ & + \left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{\substack{\text{odd } i : 1 \leq i \leq 3 \\ i \equiv 1, - \\ i \equiv 3, +}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \\ & \left. \left. + \sigma_{N-2}^- \left( \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) \right. \right. \right. \\ & + \left( \prod_{\substack{i \equiv 0, \text{ odd } i : 1 \leq i \leq 3 \\ i \equiv 1, + \\ i \equiv 3, -}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \left. \right) \\ & + \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) \\ & + \sigma_N^- \sin(\lambda_\alpha - v_{N-2} - \eta \sigma_{N-2}^z) \left( \prod_{1 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) \\ & \left. + (1 + \sigma_{N-1}^-) \left[ \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, + \\ i \equiv 2, -}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \right. \right. \\ & + \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, + \\ i \equiv 2, -}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \left. \right] \\ & + (\sin(2\eta)) \left[ \left( \prod_{\substack{0 \leq i \leq 2 \\ i \equiv 0, - \\ i \equiv 1, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) + \sigma_{N-2}^- \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, +}} \sigma_{N-i}^{-,+} \right) \left( \prod_{\substack{\text{odd } i : 1 \leq i \leq 3 \\ i \equiv 1, - \\ i \equiv 3, +}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \right. \\ & + \left( \prod_{\substack{0 \leq i \leq 2 \\ i \equiv 0, 2, - \\ i \equiv 1, +}} \sigma_{N-i}^{-,+} \right) + \left( \prod_{\substack{0 \leq i \leq 1, i \equiv 3 \\ i \equiv 0, 1, +\eta \\ i \equiv 3, -\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \\ & + \left( \prod_{\substack{0 \leq i \leq 2 \\ i \equiv 0, 2, - \\ i \equiv 1, +}} \sigma_{N-i}^{-,+} \right) + \left( \prod_{\substack{0 \leq i \leq 1, i \equiv 3 \\ i \equiv 0, 1+\eta \\ i \equiv 3, -\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) + \sigma_{N-2}^- \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2}} \sigma_{N-i}^{-,+} \right) \end{aligned}$$

$$\begin{aligned}
& + \left( \prod_{\substack{0 \leq i \leq 2 \\ i \equiv 0, - \\ i \equiv 1, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) \\
& + \left( \prod_{\substack{0 \leq i \leq 1 \\ i \equiv 0, + \\ i \equiv 1, -}} \sigma_{N-i}^{-,+} \right) \sin(\lambda_\alpha - v_{N-2} + \eta \sigma_{N-2}^z) + \left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \sin(\lambda_\alpha - v_N + \eta \sigma_N^z) \\
& + \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, - \\ i \equiv 2, +}} \sigma_{N-i}^{-,+} \right) \sin(\lambda_\alpha - v_N + \eta \sigma_N^z) ,
\end{aligned}$$

from which we conclude the argument.  $\square$

**Lemma 3** (*collecting terms from the product of two by two L operators*). The third entry of the L operator product,

$$\prod_{i=0}^3 L_{\alpha, N-i}(\lambda_\alpha, v_{N-i}) ,$$

has an expansion of the form,

$$\begin{aligned}
C_3(\lambda_\alpha) & \equiv (\sin(2\eta))^3 \sin(\lambda_\alpha - v_{N-n} - \eta \sigma_{N-n}^z) \left( \prod_{0 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) \\
& + (\sin(2\eta))^2 \left[ \sigma_{N-i}^+ \left[ \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 3 \\ i \equiv 0, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) \right. \right. \\
& \quad \left. \left. + \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \left( \prod_{\substack{\text{odd } i : 1 \leq i \leq 3 \\ i \equiv 1, - \\ i \equiv 3, +}} \sigma_{N-i}^{-,+} \right) \right] \right] \\
& + \sin(\lambda_\alpha - v_N - \eta \sigma_N^z) \left( \prod_{\substack{1 \leq i \leq 3 \\ i \equiv 1, + \\ i \equiv 2, - \\ i \equiv 3, +}} \sigma_{N-i}^{-,+} \right) \\
& + \sigma_{N-(n-3)}^+ \left( \prod_{0 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) + \dots \\
& \quad \sin(2\eta) \left[ \sigma_N^+ \left( \prod_{1 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) \right. \\
& \quad \quad \left. + \sin(\lambda_\alpha - v_{N-n} + \eta \sigma_{N-n}^z) \right. \\
& \quad \times \left[ \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 3 \\ i \equiv 0, - \\ i \equiv 2, +}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \right. \\
& \quad \quad \left. + \sigma_{N-1}^+ \left( \prod_{0 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) \right] \right] \\
& + (\sin(2\eta))^3 \sin(\lambda_\alpha - v_{N-3} - \eta \sigma_{N-3}^z) \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) ,
\end{aligned}$$

while the third entry of the L operator product,

$$\prod_{i=0}^2 L_{\alpha, N-i}(\lambda_\alpha, v_{N-i})$$

has an expansion of the form,

$$\begin{aligned} C_2(\lambda_\alpha) \equiv & (\sin(2\eta)\sigma_N^+) \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} + \eta\sigma_{N-i}^z) + (\sin(2\eta)) \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, -\eta \\ i \equiv 2, +\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta\sigma_{N-i}^z) \\ & + (\sin(2\eta))^3 \\ & \times \prod_{i \equiv 1} (\sigma_{N-i}^-)^2 \sigma_{N-i-1}^+ + ((\sin(2\eta)\sigma_{N-2}^+) \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} - \eta\sigma_{N-i}^z)) . \end{aligned}$$

*Proof of Lemma 3.* We collect terms from the first two terms in the product of L operators,

$$\begin{bmatrix} \sin(\lambda_\alpha - v_N + \eta\sigma_N^z) & \sin(2\eta)\sigma_N^- \\ \sin(2\eta)\sigma_N^+ & \sin(\lambda_\alpha - v_N - \eta\sigma_N^z) \end{bmatrix} \begin{bmatrix} \sin(\lambda_\alpha - v_{N-1} + \eta\sigma_{N-1}^z) & \sin(2\eta)\sigma_{N-1}^- \\ \sin(2\eta)\sigma_{N-1}^+ & \sin(\lambda_\alpha - v_{N-1} - \eta\sigma_{N-1}^z) \end{bmatrix}$$

from which one obtains a resultant matrix of the form,

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} ,$$

which has the following expression for the third, and fourth, entries, with,

$$\begin{aligned} \mathbf{3}^0 \equiv \mathbf{3} & \equiv \sin(2\eta)\sigma_N^+ \sin(\lambda_\alpha - v_{N-1} + \eta\sigma_{N-1}^z) + \sin(\lambda_\alpha - v_N - \eta\sigma_N^z) \sin(2\eta)\sigma_{N-1}^+ , \\ \mathbf{4}^0 \equiv \mathbf{4} & \equiv \sin(2\eta)\sigma_{N-1}^- \sin(2\eta)\sigma_{N-1}^- + \sin(\lambda_\alpha - v_N - \eta\sigma_N^z) \sin(\lambda_\alpha - v_{N-1} - \eta\sigma_{N-1}^z) , \end{aligned}$$

From the product of operators,

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} \sin(\lambda_\alpha - v_{N-2} + \eta\sigma_{N-2}^z) & \sin(2\eta)\sigma_{N-2}^- \\ \sin(2\eta)\sigma_{N-2}^+ & \sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) \end{bmatrix} \dots \times \begin{bmatrix} \sin(\lambda_\alpha - v_1 + \eta\sigma_1^z) & \sin(2\eta)\sigma_1^- \\ \sin(2\eta)\sigma_1^+ & \sin(\lambda_\alpha - v_1 - \eta\sigma_1^z) \end{bmatrix} ,$$

which takes the form,

$$\begin{aligned} & \left[ \begin{array}{cc} 1\sin(\lambda_\alpha - v_{N-2} + \eta\sigma_{N-2}^z) + 2\sin(2\eta)\sigma_{N-2}^+ & 1\sin(2\eta)\sigma_{N-2}^- + 2\sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) \\ 3\sin(\lambda_\alpha - v_{N-2} + \eta\sigma_{N-2}^z) + 4\sin(2\eta)\sigma_{N-2}^+ & 3\sin(2\eta)\sigma_{N-2}^- + 4\sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) \end{array} \right] \\ & \dots \times \begin{bmatrix} \sin(\lambda_\alpha - v_1 + \eta\sigma_1^z) & \sin(2\eta)\sigma_1^- \\ \sin(2\eta)\sigma_1^+ & \sin(\lambda_\alpha - v_1 - \eta\sigma_1^z) \end{bmatrix} , \end{aligned}$$

performing rearrangements for the next term yields,

$$\begin{aligned} \mathbf{3}^1 \equiv & 3\sin(\lambda_\alpha - v_{N-2} + \eta\sigma_{N-2}^z) + 4\sin(2\eta)\sigma_{N-2}^+ \\ \equiv & \left( \sin(2\eta)\sigma_N^+ \sin(\lambda_\alpha - v_{N-1} + \eta\sigma_{N-1}^z) + \sin(\lambda_\alpha - v_N - \eta\sigma_N^z) \sin(2\eta)\sigma_{N-1}^+ \right) \sin(\lambda_\alpha - v_{N-2} + \eta\sigma_{N-2}^z) \\ & + \left( \sin(2\eta)\sigma_{N-1}^- \sin(2\eta)\sigma_{N-1}^- + \sin(\lambda_\alpha - v_N - \eta\sigma_N^z) \sin(\lambda_\alpha - v_{N-1} - \eta\sigma_{N-1}^z) \right) \sin(2\eta)\sigma_{N-2}^+ , \\ & \equiv \sin(2\eta)\sigma_N^+ \sin(\lambda_\alpha - v_{N-1} + \eta\sigma_{N-1}^z) \sin(\lambda_\alpha - v_{N-2} + \eta\sigma_{N-2}^z) \\ & + \sin(\lambda_\alpha - v_N - \eta\sigma_N^z) \sin(2\eta)\sigma_{N-1}^+ \sin(\lambda_\alpha - v_{N-2} + \eta\sigma_{N-2}^z) + \sin(2\eta)\sigma_{N-1}^- \sin(2\eta)\sigma_{N-1}^- \sin(2\eta)\sigma_{N-2}^+ \\ & + \sin(\lambda_\alpha - v_N - \eta\sigma_N^z) \sin(\lambda_\alpha - v_{N-1} - \eta\sigma_{N-1}^z) \sin(2\eta)\sigma_{N-2}^+ , \end{aligned}$$

corresponding to the third term,  $\mathbf{3}^1$  which has the product representation,

$$\begin{aligned}
& (\sin(2\eta)\sigma_N^+) \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} + \eta\sigma_{N-i}^z) + (\sin(2\eta)) \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, -\eta \\ i \equiv 2, +\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta\sigma_{N-i}^z) + (\sin(2\eta))^3 \\
& \times \prod_{i=1} (\sigma_{N-i}^-)^2 \sigma_{N-i-1}^+ + ((\sin(2\eta)\sigma_{N-2}^+) \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} - \eta\sigma_{N-i}^z) \quad .
\end{aligned}$$

The fourth term,  $\mathbf{4}^1$ , has the product representation,

$$\begin{aligned}
& (\sin(2\eta))^2 \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \left( \sin(\lambda_\alpha - v_{N-1} + \eta\sigma_{N-1}^z) \right) + (\sin(2\eta))^2 \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \left( \sin(\lambda_\alpha - v_N - \eta\sigma_N^z) \right) \\
& + (\sin(2\eta))^2 \prod_{i=1} \sigma_{N-i}^- \left( \sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) \right) + \prod_{0 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta\sigma_{N-i}^z) \quad .
\end{aligned}$$

From the two by two matrix with entries  $\mathbf{1}^1$ ,  $\mathbf{2}^1$ ,  $\mathbf{3}^1$  and  $\mathbf{4}^1$ , the remaining terms of the product takes the form,

$$\begin{bmatrix} \mathbf{1}^1 & \mathbf{2}^1 \\ \mathbf{3}^1 & \mathbf{4}^1 \end{bmatrix} \begin{bmatrix} \sin(\lambda_\alpha - v_{N-3} + \eta\sigma_{N-3}^z) & \sin(2\eta)\sigma_{N-3}^- \\ \sin(2\eta)\sigma_{N-3}^+ & \sin(\lambda_\alpha - v_{N-3} - \eta\sigma_{N-3}^z) \end{bmatrix} \dots \times \begin{bmatrix} \sin(\lambda_\alpha - v_1 + \eta\sigma_1^z) & \sin(2\eta)\sigma_1^- \\ \sin(2\eta)\sigma_1^+ & \sin(\lambda_\alpha - v_1 - \eta\sigma_1^z) \end{bmatrix} ,$$

which implies, for,

$$\begin{bmatrix} \mathbf{1}^1 & \mathbf{2}^1 \\ \mathbf{3}^1 & \mathbf{4}^1 \end{bmatrix} \begin{bmatrix} \sin(\lambda_\alpha - v_{N-3} + \eta\sigma_{N-3}^z) & \sin(2\eta)\sigma_{N-3}^- \\ \sin(2\eta)\sigma_{N-3}^+ & \sin(\lambda_\alpha - v_{N-3} - \eta\sigma_{N-3}^z) \end{bmatrix} \equiv \begin{bmatrix} \mathbf{1}^2 & \mathbf{2}^2 \\ \mathbf{3}^2 & \mathbf{4}^2 \end{bmatrix}$$

has a third entry that is given by,

$$\mathbf{3}^2 \equiv \mathbf{3}^1 \sin(\lambda_\alpha - v_{N-3} + \eta\sigma_{N-3}^z) + \mathbf{4}^1 \sin(2\eta)\sigma_{N-3}^+ \quad .$$

The superposition above has the equivalent product representation, after collecting terms,

$$\begin{aligned}
& \left[ (\sin(2\eta)\sigma_N^+) \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} + \eta\sigma_{N-i}^z) + (\sin(2\eta)) \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, -\eta \\ i \equiv 2, +\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta\sigma_{N-i}^z) + (\sin(2\eta))^3 \right. \\
& \times \prod_{i=1} (\sigma_{N-i}^-)^2 \sigma_{N-i-1}^+ + ((\sin(2\eta)\sigma_{N-2}^+) \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} - \eta\sigma_{N-i}^z) \\
& \times \sin(\lambda_\alpha - v_{N-3} + \eta\sigma_{N-3}^z) + \left[ (\sin(2\eta))^2 \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \left( \sin(\lambda_\alpha - v_{N-1} + \eta\sigma_{N-1}^z) \right) \right. \\
& + (\sin(2\eta))^2 \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \left( \sin(\lambda_\alpha - v_N - \eta\sigma_N^z) \right) + (\sin(2\eta))^2 \prod_{i=1} \sigma_{N-i}^- \left( \sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) \right) \\
& \left. + \prod_{0 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta\sigma_{N-i}^z) \right] \sigma_{N-3}^+
\end{aligned}$$

Distributing terms further implies,

$$\begin{aligned}
& (\sin(2\eta)\sigma_N^+) \left( \prod_{1 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} + \eta\sigma_{N-i}^z) \right) + (\sin(2\eta)) \sin(\lambda_\alpha - v_{N-3} + \eta\sigma_{N-3}^z) \\
\times & \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, -\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta\sigma_{N-i}^z) \right) + (\sin(2\eta))^3 \sin(\lambda_\alpha - v_{N-3} + \eta\sigma_{N-3}^z) \left( \prod_{i \equiv 1} (\sigma_{N-i}^-)^2 \sigma_{N-i-1}^+ \right) \\
& + (\sin(2\eta)\sigma_{N-2}^+) \sin(\lambda_\alpha - v_{N-3} + \eta\sigma_{N-3}^z) \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} - \eta\sigma_{N-i}^z) \right) \\
& + (\sin(2\eta))^2 \sigma_{N-3}^+ \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \sin(\lambda_\alpha - v_{N-1} + \eta\sigma_{N-1}^z) \right) \\
& + (\sin(2\eta))^2 \sin(\lambda_\alpha - v_N - \eta\sigma_N^z) \left( \prod_{\substack{1 \leq i \leq 3 \\ i \equiv 1, + \\ i \equiv 2, - \\ i \equiv 3, +}} \sigma_{N-i}^{-,+} \right) \\
& + (\sin(2\eta))^2 \sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) \sigma_{N-3}^+ \left( \prod_{\substack{\text{odd } i : 1 \leq i \leq 3 \\ i \equiv 0, - \\ i \equiv 3, +}} \sigma_{N-i}^{-,+} \right) \\
& + (\sin(2\eta))^2 \sigma_{N-3}^+ \left( \prod_{0 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta\sigma_{N-i}^z) \right) .
\end{aligned}$$

From the superposition above, grouping together terms based upon the occurrence of a  $\sin(2\eta)$  prefactor yields,

$$\begin{aligned}
& \sin(2\eta) \left[ \sigma_N^+ \left( \prod_{1 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} + \eta\sigma_{N-i}^z) \right) + \sin(\lambda_\alpha - v_{N-3} + \eta\sigma_{N-3}^z) \right. \\
\times & \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, -\eta \\ i \equiv 2, +\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta\sigma_{N-i}^z) \right) + \sigma_{N-2}^+ \sin(\lambda_\alpha - v_{N-3} + \eta\sigma_{N-3}^z) \\
& \left. \times \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} - \eta\sigma_{N-i}^z) \right) \right] ,
\end{aligned}$$

while grouping together terms based upon the occurrence of a  $(\sin(2\eta))^2$  prefactor yields,

$$\begin{aligned}
& (\sin(2\eta))^2 \left[ \sigma_{N-3}^+ \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) \sin(\lambda_\alpha - v_{N-1} + \eta\sigma_{N-1}^z) + \sin(\lambda_\alpha - v_N - \eta\sigma_N^z) \left( \prod_{\substack{1 \leq i \leq 3 \\ i \equiv 1, + \\ i \equiv 2, - \\ i \equiv 3, +}} \sigma_{N-i}^{-,+} \right) \right. \\
& + \sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) \sigma_{N-3}^+ \left( \prod_{\substack{\text{odd } i : 1 \leq i \leq 3 \\ i \equiv 0, - \\ i \equiv 3, +}} \sigma_{N-i}^{-,+} \right) + \sigma_{N-3}^+ \left( \prod_{0 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta\sigma_{N-i}^z) \right) \left. \right] .
\end{aligned}$$

Besides the  $\sin(2\eta)$  and  $(\sin(2\eta))^2$  terms, the remaining term under  $(\sin(2\eta))^3$  is,

$$(\sin(2\eta))^3 \sin(\lambda_\alpha - v_{N-3} - \eta\sigma_{N-3}^z) \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} - \eta\sigma_{N-i}^z) \right) .$$

Next, we further rearrange terms under the  $\sin(2\eta)$ , in which,

$$\begin{aligned} & \sin(2\eta) \left[ \sigma_N^+ \left( \prod_{1 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) + \sin(\lambda_\alpha - v_{N-3} + \eta \sigma_{N-3}^z) \right. \\ & \times \left. \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, -\eta \\ i \equiv 2, +\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) + \sigma_{N-2}^+ \sin(\lambda_\alpha - v_{N-3} + \eta \sigma_{N-3}^z) \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) \right] , \end{aligned}$$

which yields,

$$\begin{aligned} & \sin(2\eta) \left[ \sigma_N^+ \left( \prod_{1 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) + \sin(\lambda_\alpha - v_{N-n} + \eta \sigma_{N-n}^z) \right. \\ & \times \left[ \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 3 \\ i \equiv 0, - \\ i \equiv 2, +}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \right. \\ & \left. \left. + \sigma_{N-1}^+ \left( \prod_{0 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) \right] \right] . \end{aligned}$$

Similarly, the terms under the  $(\sin(2\eta))^2$  yield,

$$\begin{aligned} & (\sin(2\eta))^2 \left[ \sigma_{N-i}^+ \left[ \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 3 \\ i \equiv 0, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) \right. \right. \\ & \left. \left. + \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \left( \prod_{\substack{\text{odd } i : 1 \leq i \leq 3 \\ i \equiv 1, - \\ i \equiv 3, +}} \sigma_{N-i}^{-,+} \right) \right] \right. \\ & \left. + \sin(\lambda_\alpha - v_N - \eta \sigma_N^z) \left( \prod_{\substack{1 \leq i \leq 3 \\ i \equiv 1, + \\ i \equiv 2, - \\ i \equiv 3, +}} \sigma_{N-i}^{-,+} \right) \right. \\ & \left. + \sigma_{N-(n-3)}^+ \left( \prod_{0 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) \right] , \end{aligned}$$

while rearranging terms under the  $(\sin(2\eta))^3$  prefactor yields,

$$(\sin(2\eta))^3 \sin(\lambda_\alpha - v_{N-n} - \eta \sigma_{N-n}^z) \left( \prod_{0 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) .$$

Hence,

$$\begin{aligned} C_3(\lambda_\alpha) \equiv & (\sin(2\eta))^3 \sin(\lambda_\alpha - v_{N-n} - \eta \sigma_{N-n}^z) \left( \prod_{0 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) \\ & + (\sin(2\eta))^2 \left[ \sigma_{N-i}^+ \left[ \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 3 \\ i \equiv 0, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) \right. \right. \\ & \left. \left. + \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \left( \prod_{\substack{\text{odd } i : 1 \leq i \leq 3 \\ i \equiv 1, - \\ i \equiv 3, +}} \sigma_{N-i}^{-,+} \right) \right] \right] , \end{aligned}$$

$$\begin{aligned}
& + \sin(\lambda_\alpha - v_N - \eta \sigma_N^z) \left( \prod_{\substack{1 \leq i \leq 3 \\ i \equiv 1, + \\ i \equiv 2, - \\ i \equiv 3, +}} \sigma_{N-i}^{-,+} \right) \\
& + \sigma_{N-(n-3)}^+ \left( \prod_{0 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) \Big] \\
& + \sin(2\eta) \left[ \sigma_N^+ \left( \prod_{1 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) \right. \\
& \quad \left. + \sin(\lambda_\alpha - v_{N-n} + \eta \sigma_{N-n}^z) \right] \\
& \times \left[ \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 3 \\ i \equiv 0, - \\ i \equiv 2, +}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \right. \\
& \quad \left. + \sigma_{N-1}^+ \left( \prod_{0 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) \right] \Big] \\
& + (\sin(2\eta))^3 \sin(\lambda_\alpha - v_{N-3} - \eta \sigma_{N-3}^z) \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) ,
\end{aligned}$$

from which we conclude the argument.  $\square$

**Lemma 4** (*collecting terms from the product of two by two L operators*). The fourth entry of the L operator product,

$$\prod_{i=0}^3 L_{\alpha, N-i}(\lambda_\alpha, v_{N-i}) ,$$

has an expansion of the form,

$$\begin{aligned}
D_3(\lambda_\alpha) \equiv & \prod_{0 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) + (\sin(2\eta))^2 \left[ \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) \right. \\
& \times \left( \prod_{\substack{\text{odd } i : 1 \leq i \leq 3 \\ i \equiv 1, + \\ i \equiv 3, -}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \\
& + \left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) \left( \prod_{i=0, i \neq 3} \sin(\lambda_\alpha - v_N - \eta \sigma_N^z) \right) + \left( \prod_{i=1} \sigma_{N-i}^- \right) \left( \prod_{2 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-2} - \eta \sigma_{N-2}^z) \right) \Big] \\
& + (\sin(2\eta))^3 \left( \prod_{i=1} (\sigma_{N-i}^-)^2 \right) \sigma_{N-i-1}^+ + (\sin(2\eta))^2 \left[ \sigma_N^+ \sigma_{N-3}^+ \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) \right. \\
& \left. + \sigma_{N-3}^- \left( \prod_{\substack{\text{even } i : 0 \leq i \leq 2 \\ i \equiv 0, -\eta \\ i \equiv 2, +\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) + \sigma_{N-2}^+ \sigma_{N-3}^- \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) \right] ,
\end{aligned}$$

while the fourth entry of the L operator product,

$$\prod_{i=0}^2 L_{\alpha, N-i}(\lambda_\alpha, v_{N-i})$$

has an expansion of the form,

$$\begin{aligned}
D_2(\lambda_\alpha) \equiv & (\sin(2\eta))^2 \left( \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) \left( \sin(\lambda_\alpha - v_{N-1} + \eta\sigma_{N-1}^z) \right) + (\sin(2\eta))^2 \left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) \\
& \times \left( \sin(\lambda_\alpha - v_N - \eta\sigma_N^z) \right) \\
& + (\sin(2\eta))^2 \left( \prod_{i=1} \sigma_{N-i}^- \right) \left( \sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) \right) + \prod_{0 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta\sigma_{N-i}^z) .
\end{aligned}$$

*Proof of Lemma 4.* We collect terms from the first two terms in the product of  $L$  operators,

$$\begin{bmatrix} \sin(\lambda_\alpha - v_N + \eta\sigma_N^z) & \sin(2\eta)\sigma_N^- \\ \sin(2\eta)\sigma_N^+ & \sin(\lambda_\alpha - v_N - \eta\sigma_N^z) \end{bmatrix} \begin{bmatrix} \sin(\lambda_\alpha - v_{N-1} + \eta\sigma_{N-1}^z) & \sin(2\eta)\sigma_{N-1}^- \\ \sin(2\eta)\sigma_{N-1}^+ & \sin(\lambda_\alpha - v_{N-1} - \eta\sigma_{N-1}^z) \end{bmatrix}$$

from which one obtains a resultant matrix of the form,

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} ,$$

which has the following expression for the third and fourth entries, with,

$$\begin{aligned}
\mathbf{3}^0 \equiv \mathbf{3} & \equiv \sin(2\eta)\sigma_N^+ \sin(\lambda_\alpha - v_{N-1} + \eta\sigma_{N-1}^z) + \sin(\lambda_\alpha - v_N - \eta\sigma_N^z) \sin(2\eta)\sigma_{N-1}^+ , \\
\mathbf{4}^0 \equiv \mathbf{4} & \equiv \sin(2\eta)\sigma_{N-1}^- \sin(2\eta)\sigma_{N-1}^- + \sin(\lambda_\alpha - v_N - \eta\sigma_N^z) \sin(\lambda_\alpha - v_{N-1} - \eta\sigma_{N-1}^z) ,
\end{aligned}$$

Performing rearrangements for the final term, yields,

$$\begin{aligned}
\mathbf{4}^1 \equiv \mathbf{3} & \sin(2\eta)\sigma_{N-2}^- + \mathbf{4} \sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) \\
& = \left( \sin(2\eta)\sigma_N^+ \sin(\lambda_\alpha - v_{N-1} + \eta\sigma_{N-1}^z) + \sin(\lambda_\alpha - v_N - \eta\sigma_N^z) \sin(2\eta)\sigma_{N-1}^+ \right) \sin(2\eta)\sigma_{N-2}^- \\
& + \left( \sin(2\eta)\sigma_{N-1}^- \sin(2\eta)\sigma_{N-1}^- + \sin(\lambda_\alpha - v_N - \eta\sigma_N^z) \sin(\lambda_\alpha - v_{N-1} - \eta\sigma_{N-1}^z) \right) \sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) \\
& \equiv \sin(2\eta)\sigma_N^+ \sin(\lambda_\alpha - v_{N-1} + \eta\sigma_{N-1}^z) \sin(2\eta)\sigma_{N-2}^- + \sin(\lambda_\alpha - v_N - \eta\sigma_N^z) \sin(2\eta)\sigma_{N-1}^+ \sin(2\eta)\sigma_{N-2}^- \\
& \quad + \sin(2\eta)\sigma_{N-1}^- \sin(2\eta)\sigma_{N-1}^- \sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) \\
& \quad + \sin(\lambda_\alpha - v_N - \eta\sigma_N^z) \sin(\lambda_\alpha - v_{N-1} - \eta\sigma_{N-1}^z) \sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) ,
\end{aligned}$$

corresponding to the fourth term,  $\mathbf{4}^1$  which has the product representation,

$$\begin{aligned}
& (\sin(2\eta))^2 \left( \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) \left( \sin(\lambda_\alpha - v_{N-1} + \eta\sigma_{N-1}^z) \right) + (\sin(2\eta))^2 \left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) \\
& \quad \times \left( \sin(\lambda_\alpha - v_N - \eta\sigma_N^z) \right) \\
& + (\sin(2\eta))^2 \left( \prod_{i=1} \sigma_{N-i}^- \right) \left( \sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) \right) + \prod_{0 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta\sigma_{N-i}^z) .
\end{aligned}$$

As demonstrated in arguments for the previous lemma, the third term has the product expansion,

$$\begin{aligned}
\mathbf{3}^1 \equiv & (\sin(2\eta)\sigma_N^+) \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} + \eta\sigma_{N-i}^z) \right) + (\sin(2\eta)) \left( \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, -\eta \\ i \equiv 2, +\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta\sigma_{N-i}^z) \right) \\
& + (\sin(2\eta))^3 \\
& \times \left( \prod_{i=1} (\sigma_{N-i}^-)^2 \right) \sigma_{N-i-1}^+ + ((\sin(2\eta)\sigma_{N-2}^+) \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} - \eta\sigma_{N-i}^z) \right) \\
& \quad + (\sin(2\eta))^3
\end{aligned}$$

From the two by two matrix with entries  $\mathbf{1}^1$ ,  $\mathbf{2}^1$ ,  $\mathbf{3}^1$  and  $\mathbf{4}^1$ , the remaining terms of the product takes the form,

$$\begin{bmatrix} \mathbf{1}^1 & \mathbf{2}^1 \\ \mathbf{3}^1 & \mathbf{4}^1 \end{bmatrix} \begin{bmatrix} \sin(\lambda_\alpha - v_{N-3} + \eta\sigma_{N-3}^z) & \sin(2\eta)\sigma_{N-3}^- \\ \sin(2\eta)\sigma_{N-3}^+ & \sin(\lambda_\alpha - v_{N-3} - \eta\sigma_{N-3}^z) \end{bmatrix} \dots \times \begin{bmatrix} \sin(\lambda_\alpha - v_1 + \eta\sigma_1^z) & \sin(2\eta)\sigma_1^- \\ \sin(2\eta)\sigma_1^+ & \sin(\lambda_\alpha - v_1 - \eta\sigma_1^z) \end{bmatrix} ,$$

which implies, for,

$$\begin{bmatrix} \mathbf{1}^1 & \mathbf{2}^1 \\ \mathbf{3}^1 & \mathbf{4}^1 \end{bmatrix} \begin{bmatrix} \sin(\lambda_\alpha - v_{N-3} + \eta\sigma_{N-3}^z) & \sin(2\eta)\sigma_{N-3}^- \\ \sin(2\eta)\sigma_{N-3}^+ & \sin(\lambda_\alpha - v_{N-3} - \eta\sigma_{N-3}^z) \end{bmatrix} \equiv \begin{bmatrix} \mathbf{1}^2 & \mathbf{2}^2 \\ \mathbf{3}^2 & \mathbf{4}^2 \end{bmatrix}$$

has a fourth entry that is given by,

$$\mathbf{4}^2 \equiv \mathbf{3}^1 \sin(2\eta)\sigma_{N-3}^- + \mathbf{4}^1 \sin(\lambda_\alpha - v_{N-3} - \eta\sigma_{N-3}^z) .$$

The superposition above has the equivalent product representation, after substituting terms,

$$\begin{aligned} & \left[ (\sin(2\eta)\sigma_N^+) \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} + \eta\sigma_{N-i}^z) \right) + (\sin(2\eta)) \right. \\ & \quad \times \left( \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, -\eta \\ i \equiv 2, +\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta\sigma_{N-i}^z) \right) + (\sin(2\eta))^3 \\ & \quad \times \left( \prod_{i \equiv 1} (\sigma_{N-i}^-)^2 \right) \sigma_{N-i-1}^+ + ((\sin(2\eta)\sigma_{N-2}^+) \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} - \eta\sigma_{N-i}^z) \right) \right] \\ & \quad \quad \quad \times \sin(2\eta)\sigma_{N-3}^- \\ & \quad + \left[ (\sin(2\eta))^2 \left( \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) \left( \sin(\lambda_\alpha - v_{N-1} + \eta\sigma_{N-1}^z) \right) \right. \\ & \quad \quad \quad + (\sin(2\eta))^2 \left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) \\ & \quad \quad \quad \times \left( \sin(\lambda_\alpha - v_N - \eta\sigma_N^z) \right) \\ & \quad \quad \quad + (\sin(2\eta))^2 \left( \prod_{i \equiv 1} \sigma_{N-i}^- \right) \left( \sin(\lambda_\alpha - v_{N-2} - \eta\sigma_{N-2}^z) \right) \\ & \quad \left. + \left( \prod_{0 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} - \eta\sigma_{N-i}^z) \right) \right] \sin(\lambda_\alpha - v_{N-3} - \eta\sigma_{N-3}^z) . \end{aligned}$$

Proceeding further from the superposition above, we group terms under the  $(\sin(2\eta))^2$  prefactor, from which,

$$\begin{aligned} & (\sin(2\eta))^2 \left[ \sigma_N^+ \sigma_{N-3}^+ \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} + \eta\sigma_{N-i}^z) \right) + \sigma_{N-3}^- \left( \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, -\eta \\ i \equiv 2, +\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta\sigma_{N-i}^z) \right) \right. \\ & \quad \left. + \sigma_{N-2}^+ \sigma_{N-3}^- \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} - \eta\sigma_{N-i}^z) \right) \right] , \end{aligned}$$

while the remaining term, for the first group of terms, under the  $(\sin(2\eta))^3$  prefactor, is,

$$(\sin(2\eta))^3 \left( \prod_{i=1} (\sigma_{N-i}^-)^2 \right) \sigma_{N-i-1}^+ .$$

In the second group of terms, collecting terms under the prefactor  $(\sin(2\eta))^2$  yields,

$$(\sin(2\eta))^2 \left[ \left( \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) \left( \prod_{\substack{\text{odd } i: 1 \leq i \leq 3 \\ i \equiv 1, + \\ i \equiv 3, -}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) + \left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) \right. \\ \times \left. \left( \prod_{i \equiv 0, i \equiv 3} \sin(\lambda_\alpha - v_N - \eta \sigma_N^z) \right) + \left( \prod_{i=1} \sigma_{N-i}^- \right) \left( \prod_{2 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-2} - \eta \sigma_{N-2}^z) \right) \right] ,$$

while for the remaining term,

$$\prod_{0 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) .$$

The desired expression for the final entry of the monodromy matrix,  $D_3(\lambda_\alpha)$ , takes the form,

$$D_3(\lambda_\alpha) \equiv \prod_{0 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) + (\sin(2\eta))^2 \left[ \left( \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) \right. \\ \times \left. \left( \prod_{\substack{\text{odd } i: 1 \leq i \leq 3 \\ i \equiv 1, + \\ i \equiv 3, -}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) \right. \\ + \left( \prod_{\substack{1 \leq i \leq 2 \\ i \equiv 1, + \\ i \equiv 2, -}} \sigma_{N-i}^{-,+} \right) \left( \prod_{i \equiv 0, i \equiv 3} \sin(\lambda_\alpha - v_N - \eta \sigma_N^z) \right) + \left( \prod_{i=1} \sigma_{N-i}^- \right) \left( \prod_{2 \leq i \leq 3} \sin(\lambda_\alpha - v_{N-2} - \eta \sigma_{N-2}^z) \right) \left. \right] \\ + (\sin(2\eta))^3 \left( \prod_{i=1} (\sigma_{N-i}^-)^2 \right) \sigma_{N-i-1}^+ + (\sin(2\eta))^2 \left[ \sigma_N^+ \sigma_{N-3}^+ \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{N-i} + \eta \sigma_{N-i}^z) \right) \right. \\ \left. + \sigma_{N-3}^- \left( \prod_{\substack{\text{even } i: 0 \leq i \leq 2 \\ i \equiv 0, -\eta \\ i \equiv 2, +\eta}} \sin(\lambda_\alpha - v_{N-i} \pm \eta \sigma_{N-i}^z) \right) + \sigma_{N-2}^+ \sigma_{N-3}^- \left( \prod_{0 \leq i \leq 1} \sin(\lambda_\alpha - v_{N-i} - \eta \sigma_{N-i}^z) \right) \right] ,$$

from which we conclude the argument.  $\square$

With expressions for  $A_3(\lambda_\alpha)$ ,  $B_3(\lambda_\alpha)$ ,  $C_3(\lambda_\alpha)$  and  $D_3(\lambda_\alpha)$ , below we obtain other expressions for,

$$\begin{bmatrix} A_n(\lambda_\alpha) & B_n(\lambda_\alpha) \\ C_n(\lambda_\alpha) & D_n(\lambda_\alpha) \end{bmatrix} ,$$

in terms of,

$$\begin{bmatrix} \mathbf{1}^n & \mathbf{2}^n \\ \mathbf{3}^n & \mathbf{4}^n \end{bmatrix} .$$

**Lemma 4** (*further iterating the matrix computation*). The  $n$  th L operator,

$$\begin{bmatrix} A_n(\lambda_\alpha) & B_n(\lambda_\alpha) \\ C_n(\lambda_\alpha) & D_n(\lambda_\alpha) \end{bmatrix} \equiv \begin{bmatrix} \mathbf{1}^n & \mathbf{2}^n \\ \mathbf{3}^n & \mathbf{4}^n \end{bmatrix} \equiv \prod_{1 \leq i \leq n} \begin{bmatrix} \mathbf{1}^i & \mathbf{2}^i \\ \mathbf{3}^i & \mathbf{4}^i \end{bmatrix} ,$$

can be expressed in terms of,

$$\begin{bmatrix} \left( A_3(\lambda_\alpha) + B_3(\lambda_\alpha) \right) \left[ (\sin(2\eta))^{n-3} \mathcal{A}_1 + \mathcal{A}_2 + \mathcal{A}_3 \right] & \left( A_3(\lambda_\alpha) + B_3(\lambda_\alpha) \right) \left[ (\sin(2\eta))^{n-3} \mathcal{B}_1 + \mathcal{B}_2 + \mathcal{B}_3 \right] \\ \left( C_3(\lambda_\alpha) + D_3(\lambda_\alpha) \right) \left[ (\sin(2\eta))^{n-3} \mathcal{C}_1 + \mathcal{C}_2 + \mathcal{C}_3 \right] & \left( C_3(\lambda_\alpha) + D_3(\lambda_\alpha) \right) \left[ (\sin(2\eta))^{n-3} \mathcal{D}_1 + \mathcal{D}_2 + \mathcal{D}_3 \right] \end{bmatrix},$$

for,

$$\begin{aligned} \mathcal{A}_1 \equiv \mathcal{C}_1 &\equiv \prod_{1 \leq i \leq n-3} (\mathcal{C}_1)_i \equiv \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} , \\ \mathcal{A}_2 \equiv \mathcal{C}_2 &\equiv \prod_{1 \leq i \leq n-3} (\mathcal{A}_2)_i \equiv \prod_{1 \leq i \leq n-3} (\mathcal{C}_2)_i \equiv \prod_{1 \leq i \leq n-3} \sin(\lambda_\alpha - v_{n-i} + \eta \sigma_{n-i}^z) , \\ \mathcal{A}_3 \equiv \mathcal{C}_3 &\equiv \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} (\mathcal{C}_3)_{i,j} \equiv \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(\lambda_\alpha - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] , \end{aligned}$$

corresponding to the first and third entries,

$$\begin{aligned} \mathcal{B}_1 \equiv \mathcal{D}_1 &\equiv \prod_{1 \leq i \leq n-3} (\mathcal{D}_1)_i \equiv \prod_{2 \leq i \leq n-3} \sigma_{n-i}^{-,+} , \\ \mathcal{B}_2 \equiv \mathcal{D}_2 &\equiv \prod_{2 \leq i \leq n-3} (\mathcal{B}_2)_i \equiv \prod_{2 \leq i \leq n-3} (\mathcal{D}_2)_i \equiv \prod_{2 \leq i \leq n-3} \sin(\lambda_\alpha - v_{n-i} + \eta \sigma_{n-i}^z) , \\ \mathcal{B}_3 \equiv \mathcal{D}_3 &\equiv \sum_{\substack{2 \leq i \leq m \\ 2 \leq j \leq n' \\ m+n'=n-3}} (\mathcal{D}_3)_{i,j} \equiv \sum_{\substack{2 \leq i \leq m \\ 2 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{2 \leq i \leq m} \sin(\lambda_\alpha - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{2 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] , \end{aligned}$$

corresponding to the second and fourth entries.

*Proof of Lemma 4.* By direct computation, we begin by computing the product of L operators,

$$\begin{bmatrix} \mathbf{1}^0 & \mathbf{2}^0 \\ \mathbf{3}^0 & \mathbf{4}^0 \end{bmatrix} \begin{bmatrix} \mathbf{1}^1 & \mathbf{2}^1 \\ \mathbf{3}^1 & \mathbf{4}^1 \end{bmatrix} \begin{bmatrix} \mathbf{1}^2 & \mathbf{2}^2 \\ \mathbf{3}^2 & \mathbf{4}^2 \end{bmatrix} \begin{bmatrix} \mathbf{1}^3 & \mathbf{2}^3 \\ \mathbf{3}^3 & \mathbf{4}^3 \end{bmatrix}$$

after which we compute the product of three arbitrary L operators,

$$\begin{bmatrix} \mathbf{1}^{i-3} & \mathbf{2}^{i-3} \\ \mathbf{3}^{i-3} & \mathbf{4}^{i-3} \end{bmatrix} \begin{bmatrix} \mathbf{1}^{i-2} & \mathbf{2}^{i-2} \\ \mathbf{3}^{i-2} & \mathbf{4}^{i-2} \end{bmatrix} \begin{bmatrix} \mathbf{1}^{i-1} & \mathbf{2}^{i-1} \\ \mathbf{3}^{i-1} & \mathbf{4}^{i-1} \end{bmatrix} ,$$

from the  $n$  fold product,

$$\prod_{1 \leq i \leq n} \begin{bmatrix} \mathbf{1}^i & \mathbf{2}^i \\ \mathbf{3}^i & \mathbf{4}^i \end{bmatrix} .$$

For the first product of two by two L operators, for  $i \equiv 0, i \equiv 1, i \equiv 2$ , and  $i \equiv 3$ , by brute force one has,

$$\begin{bmatrix} \mathbf{1}^0 \mathbf{1}^1 + \mathbf{2}^0 \mathbf{3}^1 & \mathbf{1}^0 \mathbf{2}^1 + \mathbf{2}^0 \mathbf{4}^1 \\ \mathbf{1}^1 \mathbf{3}^0 + \mathbf{3}^1 \mathbf{4}^0 & \mathbf{2}^1 \mathbf{3}^0 + \mathbf{4}^0 \mathbf{4}^1 \end{bmatrix} \begin{bmatrix} \mathbf{1}^2 & \mathbf{2}^2 \\ \mathbf{3}^2 & \mathbf{4}^2 \end{bmatrix} \begin{bmatrix} \mathbf{1}^3 & \mathbf{2}^3 \\ \mathbf{3}^3 & \mathbf{4}^3 \end{bmatrix}$$

is equivalent to,

$$\begin{bmatrix} (\mathbf{1}^0 \mathbf{1}^1 + \mathbf{2}^0 \mathbf{3}^1) \mathbf{1}^2 + (\mathbf{1}^0 \mathbf{2}^1 + \mathbf{2}^0 \mathbf{4}^1) \mathbf{3}^2 & (\mathbf{1}^0 \mathbf{1}^1 + \mathbf{2}^0 \mathbf{3}^1) \mathbf{2}^2 + (\mathbf{1}^0 \mathbf{2}^1 + \mathbf{2}^0 \mathbf{4}^1) \mathbf{4}^2 \\ (\mathbf{1}^1 \mathbf{3}^0 + \mathbf{3}^1 \mathbf{4}^0) \mathbf{1}^2 + (\mathbf{2}^1 \mathbf{3}^0 + \mathbf{4}^0 \mathbf{4}^1) \mathbf{3}^2 & (\mathbf{1}^1 \mathbf{3}^0 + \mathbf{3}^1 \mathbf{4}^0) \mathbf{2}^2 + (\mathbf{2}^1 \mathbf{3}^0 + \mathbf{4}^0 \mathbf{4}^1) \mathbf{4}^2 \end{bmatrix} \begin{bmatrix} \mathbf{1}^3 & \mathbf{2}^3 \\ \mathbf{3}^3 & \mathbf{4}^3 \end{bmatrix} .$$

From previous arguments given in **Lemmas 1-4**, the fact that,

$$\begin{bmatrix} A_3(\lambda_\alpha) & B_3(\lambda_\alpha) \\ C_3(\lambda_\alpha) & D_3(\lambda_\alpha) \end{bmatrix}$$

equals,

$$\begin{bmatrix} (\mathbf{1}^0 \mathbf{1}^1 + \mathbf{2}^0 \mathbf{3}^1) \mathbf{1}^2 + (\mathbf{1}^0 \mathbf{2}^1 + \mathbf{2}^0 \mathbf{4}^1) \mathbf{3}^2 & (\mathbf{1}^0 \mathbf{1}^1 + \mathbf{2}^0 \mathbf{3}^1) \mathbf{2}^2 + (\mathbf{1}^0 \mathbf{2}^1 + \mathbf{2}^0 \mathbf{4}^1) \mathbf{4}^2 \\ (\mathbf{1}^1 \mathbf{3}^0 + \mathbf{3}^1 \mathbf{4}^0) \mathbf{1}^2 + (\mathbf{2}^1 \mathbf{3}^0 + \mathbf{4}^0 \mathbf{4}^1) \mathbf{3}^2 & (\mathbf{1}^1 \mathbf{3}^0 + \mathbf{3}^1 \mathbf{4}^0) \mathbf{2}^2 + (\mathbf{2}^1 \mathbf{3}^0 + \mathbf{4}^0 \mathbf{4}^1) \mathbf{4}^2 \end{bmatrix} ,$$

implies, entry by entry, that,

$$\begin{aligned} A_3(\lambda_\alpha) &\equiv (\mathbf{1}^0 \mathbf{1}^1 + \mathbf{2}^0 \mathbf{3}^1) \mathbf{1}^2 + (\mathbf{1}^0 \mathbf{2}^1 + \mathbf{2}^0 \mathbf{4}^1) \mathbf{3}^2 , \\ B_3(\lambda_\alpha) &\equiv (\mathbf{1}^0 \mathbf{1}^1 + \mathbf{2}^0 \mathbf{3}^1) \mathbf{2}^2 + (\mathbf{1}^0 \mathbf{2}^1 + \mathbf{2}^0 \mathbf{4}^1) , \\ C_3(\lambda_\alpha) &\equiv (\mathbf{1}^1 \mathbf{3}^0 + \mathbf{3}^1 \mathbf{4}^0) \mathbf{1}^2 + (\mathbf{2}^1 \mathbf{3}^0 + \mathbf{4}^0 \mathbf{4}^1) \mathbf{3}^2 , \\ D_3(\lambda_\alpha) &\equiv (\mathbf{1}^1 \mathbf{3}^0 + \mathbf{3}^1 \mathbf{4}^0) \mathbf{2}^2 + (\mathbf{2}^1 \mathbf{3}^0 + \mathbf{4}^0 \mathbf{4}^1) \mathbf{4}^2 . \end{aligned}$$

In terms of  $A_3(\lambda_\alpha)$ ,  $B_3(\lambda_\alpha)$ ,  $C_3(\lambda_\alpha)$  and  $D_3(\lambda_\alpha)$ , performing the final multiplication of L operators,

$$\begin{bmatrix} A_3(\lambda_\alpha) & B_3(\lambda_\alpha) \\ C_3(\lambda_\alpha) & D_3(\lambda_\alpha) \end{bmatrix} \begin{bmatrix} \mathbf{1}^3 & \mathbf{2}^3 \\ \mathbf{3}^3 & \mathbf{4}^3 \end{bmatrix}$$

equals,

$$\begin{bmatrix} A_3(\lambda_\alpha) \mathbf{1}^3 + B_3(\lambda_\alpha) \mathbf{3}^3 & A_3(\lambda_\alpha) \mathbf{2}^3 + B_3(\lambda_\alpha) \mathbf{4}^3 \\ C_3(\lambda_\alpha) \mathbf{1}^3 + D_3(\lambda_\alpha) \mathbf{3}^3 & C_3(\lambda_\alpha) \mathbf{2}^3 + D_3(\lambda_\alpha) \mathbf{4}^3 \end{bmatrix} .$$

Hence,

$$\prod_{0 \leq i \leq 3} \begin{bmatrix} \mathbf{1}^i & \mathbf{2}^i \\ \mathbf{3}^i & \mathbf{4}^i \end{bmatrix} \equiv \begin{bmatrix} A_3(\lambda_\alpha) \mathbf{1}^3 + B_3(\lambda_\alpha) \mathbf{3}^3 & A_3(\lambda_\alpha) \mathbf{2}^3 + B_3(\lambda_\alpha) \mathbf{4}^3 \\ C_3(\lambda_\alpha) \mathbf{1}^3 + D_3(\lambda_\alpha) \mathbf{3}^3 & C_3(\lambda_\alpha) \mathbf{2}^3 + D_3(\lambda_\alpha) \mathbf{4}^3 \end{bmatrix} .$$

With such an expression, we make use of the following relations,

$$\begin{aligned} A_3(\lambda_\alpha) &= A_2(\lambda_\alpha) \mathbf{1}^2 + B_2(\lambda_\alpha) \mathbf{3}^2 \equiv A_2(\lambda_\alpha) \sin(\lambda_\alpha - v_2 + \eta\sigma_2^z) + B_2(\lambda_\alpha) \sin(2\eta)\sigma_2^+ , \\ B_3(\lambda_\alpha) &= A_2(\lambda_\alpha) \mathbf{2}^2 + B_2(\lambda_\alpha) \mathbf{4}^2 \equiv A_2(\lambda_\alpha) \sin(2\eta)\sigma_2^- + B_2(\lambda_\alpha) \sin(\lambda_\alpha - v_2 - \eta\sigma_2^z) , \\ C_3(\lambda_\alpha) &= C_2(\lambda_\alpha) \mathbf{1}^2 + D_2(\lambda_\alpha) \mathbf{4}^2 \equiv C_2(\lambda_\alpha) \sin(\lambda_\alpha - v_2 + \eta\sigma_2^z) + D_2(\lambda_\alpha) \sin(2\eta)\sigma_2^+ , \\ D_3(\lambda_\alpha) &= C_2(\lambda_\alpha) \mathbf{2}^2 + D_2(\lambda_\alpha) \mathbf{4}^2 \equiv C_2(\lambda_\alpha) \sin(2\eta)\sigma_2^- + D_2(\lambda_\alpha) \sin(\lambda_\alpha - v_2 - \eta\sigma_2^z) . \end{aligned}$$

To finish the arguments for **Lemma 4**, we introduce the result below.

**Lemma 5** (*iteratively obtaining the entries of the n th L operator from the entries of the third L operator*). The first entry of the n th L operator can be expressed in terms of,

$$\begin{aligned} A_n(\lambda_\alpha) &= \left( A_3(\lambda_\alpha) + B_3(\lambda_\alpha) \right) (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \\ &\quad + \left( A_3(\lambda_\alpha) + B_3(\lambda_\alpha) \right) \left( \prod_{1 \leq i \leq n-3} \sin(\lambda_\alpha - v_{n-i} + \eta\sigma_{n-i}^z) \right) \\ &+ \left( A_3(\lambda_\alpha) + B_3(\lambda_\alpha) \right) \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(\lambda_\alpha - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) . \end{aligned}$$

The second entry of the n th L operator can be expressed in terms of,

$$\begin{aligned} B_n(\lambda_\alpha) &= \left( A_3(\lambda_\alpha) + B_3(\lambda_\alpha) \right) (\sin(2\eta))^{n-4} \left( \prod_{2 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \\ &\quad + \left( A_3(\lambda_\alpha) + B_3(\lambda_\alpha) \right) \left( \prod_{2 \leq i \leq n-3} \sin(\lambda_\alpha - v_{n-i} + \eta\sigma_{n-i}^z) \right) \\ &+ \left( A_3(\lambda_\alpha) + B_3(\lambda_\alpha) \right) \left( \sum_{\substack{2 \leq i \leq m \\ 2 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{2 \leq i \leq m} \sin(\lambda_\alpha - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{2 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) . \end{aligned}$$

The third entry of the  $n$  th L operator can be expressed in terms of,

$$\begin{aligned}
C_n(\lambda_\alpha) = & \left( C_3(\lambda_\alpha) + D_3(\lambda_\alpha) \right) (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \\
& + \left( C_3(\lambda_\alpha) + D_3(\lambda_\alpha) \right) \left( \prod_{1 \leq i \leq n-3} \sin(\lambda_\alpha - v_{n-i} + \eta \sigma_{n-i}^z) \right) \\
+ & \left( C_3(\lambda_\alpha) + D_3(\lambda_\alpha) \right) \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(\lambda_\alpha - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) .
\end{aligned}$$

The fourth entry of the  $n$  th L operator can be expressed in terms of,

$$\begin{aligned}
D_n(\lambda_\alpha) = & \left( C_3(\lambda_\alpha) + D_3(\lambda_\alpha) \right) (\sin(2\eta))^{n-4} \left( \prod_{2 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \\
& + \left( C_3(\lambda_\alpha) + D_3(\lambda_\alpha) \right) \left( \prod_{2 \leq i \leq n-3} \sin(\lambda_\alpha - v_{n-i} + \eta \sigma_{n-i}^z) \right) \\
+ & \left( C_3(\lambda_\alpha) + D_3(\lambda_\alpha) \right) \left( \sum_{\substack{2 \leq i \leq m \\ 2 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{2 \leq i \leq m} \sin(\lambda_\alpha - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{2 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) .
\end{aligned}$$

*Proof of Lemma 5.* Observe, for the first entry of the  $n$  th L operator,

$$\prod_{1 \leq i \leq n} \begin{bmatrix} \mathbf{1}^i & \mathbf{2}^i \\ \mathbf{3}^i & \mathbf{4}^i \end{bmatrix} ,$$

that,

$$A_n(\lambda_\alpha) = A_{n-1}(\lambda_\alpha) \mathbf{1}^{n-1} + B_{n-1}(\lambda_\alpha) \mathbf{3}^{n-1} \equiv A_{n-1}(\lambda_\alpha) \sin(\lambda_\alpha - v_{n-1} + \eta \sigma_{n-1}^z) + B_{n-1}(\lambda_\alpha) \sin(2\eta) \sigma_{n-1}^+ . \quad (1)$$

From the equality above, making the substitution for  $A_{n-1}(\lambda_\alpha)$  in terms of  $A_{n-2}(\lambda_\alpha)$ , implies,

$$A_{n-1}(\lambda_\alpha) = A_{n-2}(\lambda_\alpha) \mathbf{1}^{n-2} + B_{n-2}(\lambda_\alpha) \mathbf{3}^{n-2} \equiv A_{n-2}(\lambda_\alpha) \sin(\lambda_\alpha - v_{n-2} + \eta \sigma_{n-2}^z) + B_{n-2}(\lambda_\alpha) \sin(2\eta) \sigma_{n-2}^+ .$$

Similarly,

$$B_{n-1}(\lambda_\alpha) = A_{n-2}(\lambda_\alpha) \mathbf{2}^{n-2} + B_{n-2}(\lambda_\alpha) \mathbf{4}^{n-2} \equiv A_{n-2}(\lambda_\alpha) \sin(2\eta) \sigma_{n-2}^- + B_{n-2}(\lambda_\alpha) \sin(\lambda_\alpha - v_{n-2} - \eta \sigma_{n-2}^z) . \quad (1^*)$$

Rewriting (1) with the expression for  $A_{n-1}(\lambda_\alpha)$  and  $B_{n-1}(\lambda_\alpha)$  implies,

$$\begin{aligned}
(1) \equiv & \left( A_{n-2}(\lambda_\alpha) \sin(\lambda_\alpha - v_{n-2} + \eta \sigma_{n-2}^z) + B_{n-2}(\lambda_\alpha) \sin(2\eta) \sigma_{n-2}^+ \right) \sin(\lambda_\alpha - v_{n-1} + \eta \sigma_{n-1}^z) \\
& + \left( A_{n-2}(\lambda_\alpha) \sin(2\eta) \sigma_{n-2}^- + B_{n-2}(\lambda_\alpha) \sin(\lambda_\alpha - v_{n-2} - \eta \sigma_{n-2}^z) \right) \sin(2\eta) \sigma_{n-1}^+ .
\end{aligned}$$

Grouping together like terms from the expression above,

$$\begin{aligned}
& A_{n-2}(\lambda_\alpha) \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{n-i} + \eta \sigma_{n-i}^z) + (\sin(2\eta))^2 \prod_{1 \leq i \leq 2} \sigma_{n-i}^{-,+} \right) \\
& + B_{n-2}(\lambda_\alpha) \left( \sin(2\eta) \sigma_{n-2}^+ \sin(\lambda_\alpha - v_{n-1} + \eta \sigma_{n-1}^z) + \sin(\lambda_\alpha - v_{n-2} - \eta \sigma_{n-2}^z) \sin(2\eta) \sigma_{n-1}^+ \right) . \quad (2)
\end{aligned}$$

Continuing along similar lines, in which we substitute for  $A_{n-2}(\lambda_\alpha)$  and  $B_{n-2}(\lambda_\alpha)$  to rewrite (2) implies,

$$(2) \equiv \left( A_{n-3}(\lambda_\alpha) \sin(\lambda_\alpha - v_{n-3} + \eta\sigma_{n-3}^z) + B_{n-3}(\lambda_\alpha) \sin(2\eta)\sigma_{n-3}^+ \right) \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{n-i} + \eta\sigma_{n-i}^z) \right. \\ \left. + (\sin(2\eta))^2 \prod_{1 \leq i \leq 2} \sigma_{n-i}^{-,+} \right) \\ + \left( A_{n-3}(\lambda_\alpha) \sin(2\eta)\sigma_{n-3}^- + B_{n-3}(\lambda_\alpha) \sin(\lambda_\alpha - v_{n-3} - \eta\sigma_{n-3}^z) \right) \\ \times \left( \sin(2\eta)\sigma_{n-2}^+ \sin(\lambda_\alpha - v_{n-1} + \eta\sigma_{n-1}^z) + \sin(\lambda_\alpha - v_{n-2} - \eta\sigma_{n-2}^z) \sin(2\eta)\sigma_{n-1}^+ \right) ,$$

from which performing additional rearrangements implies, for the first term, that,

$$A_{n-3}(\lambda_\alpha) \left( \prod_{1 \leq i \leq 3} \sin(\lambda_\alpha - v_{n-i} + \eta\sigma_{n-i}^z) \right) + B_{n-3}(\lambda_\alpha) \sin(2\eta)\sigma_{n-3}^+ \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{n-i} + \eta\sigma_{n-i}^z) \right) \\ + A_{n-3}(\lambda_\alpha) (\sin(2\eta))^2 \sin(\lambda_\alpha - v_{n-3} + \eta\sigma_{n-3}^z) \left( \prod_{1 \leq i \leq 2} \sigma_{n-i}^{-,+} \right) + B_{n-3}(\lambda_\alpha) (\sin(2\eta))^3 \left( \prod_{1 \leq i \leq 3} \sigma_{n-i}^{-,+} \right) , \quad (2^*)$$

and for the second term, that,

$$A_{n-3}(\lambda_\alpha) (\sin(2\eta))^2 \sin(\lambda_\alpha - v_{n-1} + \eta\sigma_{n-1}^z) \left( \prod_{2 \leq i \leq 3} \sigma_{N-i}^{-,+} \right) + B_{n-3}(\lambda_\alpha) \sin(2\eta)\sigma_{n-2}^+ \times \cdots \\ \left( \prod_{i \text{ odd} : 1 \leq i \leq 3} \sin(\lambda_\alpha - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) + A_{n-3}(\lambda_\alpha) (\sin(2\eta))^2 \sin(\lambda_\alpha - v_{n-2} - \eta\sigma_{n-2}^z) \times \cdots \\ \left( \prod_{i \text{ odd} : 1 \leq i \leq 3} \sigma_{N-i}^{-,+} \right) + B_{n-3}(\lambda_\alpha) \sin(2\eta)\sigma_{n-1}^+ \left( \prod_{2 \leq i \leq 3} \sin(\lambda_\alpha - v_{n-i} - \eta\sigma_{n-i}^z) \right) . \quad (2^{**})$$

To extrapolate the formula to  $A_3(\lambda_\alpha)$ , from the previous two terms above, substitute for  $A_{n-3}(\lambda_\alpha)$  and  $B_{n-3}(\lambda_\alpha)$ , in which, from (2\*),

$$A_{n-3}(\lambda_\alpha) \left( \prod_{1 \leq i \leq 3} \sin(\lambda_\alpha - v_{n-i} + \eta\sigma_{n-i}^z) \right)$$

equals,

$$\left( A_{n-4}(\lambda_\alpha) \sin(\lambda_\alpha - v_{n-4} + \eta\sigma_{n-4}^z) + B_{n-4}(\lambda_\alpha) \sin(2\eta)\sigma_{n-4}^+ \right) \left( \prod_{1 \leq i \leq 3} \sin(\lambda_\alpha - v_{n-i} + \eta\sigma_{n-i}^z) \right)$$

is equivalent to,

$$A_{n-4}(\lambda_\alpha) \left( \prod_{1 \leq i \leq 4} \sin(\lambda_\alpha - v_{n-i} + \eta\sigma_{n-i}^z) \right) + B_{n-4}(\lambda_\alpha) \sin(2\eta)\sigma_{n-4}^+ \left( \prod_{1 \leq i \leq 3} \sin(\lambda_\alpha - v_{n-i} + \eta\sigma_{n-i}^z) \right) . \quad (2^* - 1)$$

From (2\*), the second term with  $A_{n-3}(\lambda_\alpha)$ ,

$$A_{n-3}(\lambda_\alpha) (\sin(2\eta))^2 \sin(\lambda_\alpha - v_{n-3} + \eta\sigma_{n-3}^z) \left( \prod_{1 \leq i \leq 2} \sigma_{n-i}^{-,+} \right) ,$$

equals,

$$\left( A_{n-4}(\lambda_\alpha) \sin(\lambda_\alpha - v_{n-4} + \eta\sigma_{n-4}^z) + B_{n-4}(\lambda_\alpha) \sin(2\eta) \sigma_{n-4}^+ \right) (\sin(2\eta))^2 \sin(\lambda_\alpha - v_{n-3} + \eta\sigma_{n-3}^z) \left( \prod_{1 \leq i \leq 2} \sigma_{n-i}^{-,+} \right)$$

is equivalent to,

$$A_{n-4}(\lambda_\alpha) (\sin(2\eta))^2 \left( \prod_{3 \leq i \leq 4} \sin(\lambda_\alpha - v_{n-i} + \eta\sigma_{n-i}^z) \right) \left( \prod_{1 \leq i \leq 2} \sigma_{n-i}^{-,+} \right) + B_{n-4}(\lambda_\alpha) (\sin(2\eta))^3 \left( \prod_{1 \leq i \leq 2} \sigma_{n-i}^{-,+} \right) \\ \times \sin(\lambda_\alpha - v_{n-3} + \eta\sigma_{n-3}^z) \sigma_{n-4}^+ . \quad (2^* - 2)$$

From (2\*), the third entry with  $B_{n-3}(\lambda_\alpha)$ ,

$$B_{n-3}(\lambda_\alpha) \sin(2\eta) \sigma_{n-3}^+ \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{n-i} + \eta\sigma_{n-i}^z) \right) ,$$

equals,

$$\left( A_{n-4}(\lambda_\alpha) \sin(2\eta) \sigma_{n-4}^- + B_{n-4}(\lambda_\alpha) \sin(\lambda_\alpha - v_{n-4} - \eta\sigma_{n-4}^z) \right) \sin(2\eta) \sigma_{n-3}^+ \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{n-i} + \eta\sigma_{n-i}^z) \right)$$

is equivalent to,

$$A_{n-4}(\lambda_\alpha) (\sin(2\eta))^2 \left( \prod_{3 \leq i \leq 4} \sigma_{n-i}^{-,+} \right) \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{n-i} + \eta\sigma_{n-i}^z) \right) \\ + B_{n-4}(\lambda_\alpha) \sin(2\eta) \sigma_{n-3}^+ \sin(\lambda_\alpha - v_{n-4} - \eta\sigma_{n-4}^z) \left( \prod_{1 \leq i \leq 2} \sin(\lambda_\alpha - v_{n-i} + \eta\sigma_{n-i}^z) \right) . \quad (2^* - 3)$$

From (2\*), the fourth entry with  $B_{n-3}(\lambda_\alpha)$ ,

$$B_{n-3}(\lambda_\alpha) (\sin(2\eta))^3 \left( \prod_{1 \leq i \leq 3} \sigma_{n-i}^{-,+} \right) ,$$

equals,

$$\left( A_{n-4}(\lambda_\alpha) \sin(2\eta) \sigma_{n-4}^- + B_{n-4}(\lambda_\alpha) \sin(\lambda_\alpha - v_{n-4} - \eta\sigma_{n-4}^z) \right) (\sin(2\eta))^3 \left( \prod_{1 \leq i \leq 3} \sigma_{n-i}^{-,+} \right)$$

is equivalent to,

$$A_{n-4}(\lambda_\alpha) (\sin(2\eta))^4 \left( \prod_{1 \leq i \leq 4} \sigma_{n-i}^{-,+} \right) + B_{n-4}(\lambda_\alpha) \sin(\lambda_\alpha - v_{n-4} - \eta\sigma_{n-4}^z) (\sin(2\eta))^3 \left( \prod_{1 \leq i \leq 3} \sigma_{n-i}^{-,+} \right) . \quad (2^* - 4)$$

Continuing along similar lines for each term in (2\*\*) implies, that the first term,

$$A_{n-3}(\lambda_\alpha) (\sin(2\eta))^2 \sin(\lambda_\alpha - v_{n-1} + \eta\sigma_{n-1}^z) \left( \prod_{2 \leq i \leq 3} \sigma_{N-i}^{-,+} \right) ,$$

equals,

$$\left( A_{n-4}(\lambda_\alpha) \sin(\lambda_\alpha - v_{n-4} + \eta\sigma_{n-4}^z) + B_{n-4}(\lambda_\alpha) \sin(2\eta) \sigma_{n-4}^+ \right) (\sin(2\eta))^2 \sin(\lambda_\alpha - v_{n-1} + \eta\sigma_{n-1}^z) \left( \prod_{2 \leq i \leq 3} \sigma_{N-i}^{-,+} \right)$$

is equivalent to,

$$\begin{aligned} & A_{n-4}(\lambda_\alpha) (\sin(2\eta))^2 \sin(\lambda_\alpha - v_{n-1} + \eta\sigma_{n-1}^z) \sin(\lambda_\alpha - v_{n-4} + \eta\sigma_{n-4}^z) \left( \prod_{2 \leq i \leq 3} \sigma_{N-i}^{-,+} \right) \\ & + B_{n-4}(\lambda_\alpha) (\sin(2\eta))^3 \sin(\lambda_\alpha - v_{n-1} + \eta\sigma_{n-1}^z) \left( \prod_{2 \leq i \leq 4} \sigma_{N-i}^{-,+} \right). \end{aligned} \quad (2^{**}-1)$$

From (2\*\*), the second term,

$$B_{n-3}(\lambda_\alpha) \sin(2\eta) \sigma_{n-2}^+ \left( \prod_{i \text{ odd : } 1 \leq i \leq 3} \sin(\lambda_\alpha - v_{n-i} \pm \eta\sigma_{n-i}^z) \right),$$

equals,

$$\left( A_{n-4}(\lambda_\alpha) \sin(2\eta) \sigma_{n-4}^- + B_{n-4}(\lambda_\alpha) \sin(\lambda_\alpha - v_{n-4} - \eta\sigma_{n-4}^z) \right) \sin(2\eta) \sigma_{n-2}^+ \left( \prod_{i \text{ odd : } 1 \leq i \leq 3} \sin(\lambda_\alpha - v_{n-i} \pm \eta\sigma_{n-i}^z) \right)$$

is equivalent to,

$$\begin{aligned} & A_{n-4}(\lambda_\alpha) (\sin(2\eta))^2 \sigma_{n-2}^+ \sigma_{n-4}^- \left( \prod_{i \text{ odd : } 1 \leq i \leq 3} \sin(\lambda_\alpha - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) + B_{n-4}(\lambda_\alpha) \\ & \times \sin(2\eta) \sigma_{n-2}^+ \sin(\lambda_\alpha - v_{n-4} - \eta\sigma_{n-4}^z) \left( \prod_{i \text{ odd : } 1 \leq i \leq 3} \sin(\lambda_\alpha - v_{n-i} \pm \eta\sigma_{n-i}^z) \right). \end{aligned} \quad (2^{**}-2)$$

From (2\*\*), the third term,

$$A_{n-3}(\lambda_\alpha) (\sin(2\eta))^2 \sin(\lambda_\alpha - v_{n-2} - \eta\sigma_{n-2}^z) \left( \prod_{i \text{ odd : } 1 \leq i \leq 3} \sigma_{N-i}^{-,+} \right),$$

equals,

$$\begin{aligned} & \left( A_{n-4}(\lambda_\alpha) \sin(\lambda_\alpha - v_{n-4} + \eta\sigma_{n-4}^z) + B_{n-4}(\lambda_\alpha) \sin(2\eta) \sigma_{n-4}^+ \right) (\sin(2\eta))^2 \sin(\lambda_\alpha - v_{n-2} - \eta\sigma_{n-2}^z) \\ & \times \left( \prod_{i \text{ odd : } 1 \leq i \leq 3} \sigma_{N-i}^{-,+} \right) \end{aligned}$$

is equivalent to,

$$\begin{aligned} & A_{n-4}(\lambda_\alpha) (\sin(2\eta))^2 \sin(\lambda_\alpha - v_{n-2} - \eta\sigma_{n-2}^z) \sin(\lambda_\alpha - v_{n-4} + \eta\sigma_{n-4}^z) \left( \prod_{i \text{ odd : } 1 \leq i \leq 3} \sigma_{N-i}^{-,+} \right) \\ & + B_{n-4}(\lambda_\alpha) (\sin(2\eta))^3 \sin(\lambda_\alpha - v_{n-2} - \eta\sigma_{n-2}^z) \sigma_{n-4}^+ \left( \prod_{i \text{ odd : } 1 \leq i \leq 3} \sigma_{N-i}^{-,+} \right). \end{aligned} \quad (2^{**}-3)$$

From (2\*\*), the fourth term,

$$B_{n-3}(\lambda_\alpha) \sin(2\eta) \sigma_{n-1}^+ \left( \prod_{2 \leq i \leq 3} \sin(\lambda_\alpha - v_{n-i} - \eta\sigma_{n-i}^z) \right),$$

equals,

$$\left( A_{n-4}(\lambda_\alpha) \sin(2\eta) \sigma_{n-4}^- + B_{n-4}(\lambda_\alpha) \sin(\lambda_\alpha - v_{n-4} - \eta \sigma_{n-4}^z) \right) \sin(2\eta) \sigma_{n-1}^+ \left( \prod_{2 \leq i \leq 3} \sin(\lambda_\alpha - v_{n-i} - \eta \sigma_{n-i}^z) \right) ,$$

is equivalent to,

$$\begin{aligned} & A_{n-4}(\lambda_\alpha) (\sin(2\eta))^2 \sigma_{n-1}^+ \sigma_{n-4}^+ \left( \prod_{2 \leq i \leq 3} \sin(\lambda_\alpha - v_{n-i} - \eta \sigma_{n-i}^z) \right) \\ & + B_{n-4}(\lambda_\alpha) \sin(2\eta) \sigma_{n-1}^+ \left( \prod_{2 \leq i \leq 4} \sin(\lambda_\alpha - v_{n-i} - \eta \sigma_{n-i}^z) \right) . \end{aligned} \quad (2^{**} - 4)$$

Extrapolating the formulas from (1) and (2\*) yields,

$$\begin{aligned} A_n(\lambda_\alpha) = & \left( A_3(\lambda_\alpha) + B_3(\lambda_\alpha) \right) (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \\ & + \left( A_3(\lambda_\alpha) + B_3(\lambda_\alpha) \right) \left( \prod_{1 \leq i \leq n-3} \sin(\lambda_\alpha - v_{n-i} + \eta \sigma_{n-i}^z) \right) \\ & + \left( A_3(\lambda_\alpha) + B_3(\lambda_\alpha) \right) \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(\lambda_\alpha - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) . \end{aligned}$$

Along similar lines, repeating the computations since the beginning of the proof yields another formula for  $C_n(\lambda_\alpha)$ , in which,

$$\begin{aligned} C_n(\lambda_\alpha) = & \left( C_3(\lambda_\alpha) + D_3(\lambda_\alpha) \right) (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \\ & + \left( C_3(\lambda_\alpha) + D_3(\lambda_\alpha) \right) \left( \prod_{1 \leq i \leq n-3} \sin(\lambda_\alpha - v_{n-i} + \eta \sigma_{n-i}^z) \right) \\ & + \left( C_3(\lambda_\alpha) + D_3(\lambda_\alpha) \right) \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(\lambda_\alpha - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) . \end{aligned}$$

To obtain the desired formula for  $B_{n-1}(\lambda_\alpha)$ , we pursue the following steps. Rewriting (1\*), in terms of  $A_{n-2}(\lambda_\alpha)$  and  $B_{n-2}(\lambda_\alpha)$ , implies,

$$(1^*) \equiv \left( A_{n-3}(\lambda_\alpha) \sin(\lambda_\alpha - v_{n-3} + \eta \sigma_{n-3}^z) + B_{n-3}(\lambda_\alpha) \sin(2\eta) \sigma_{n-3}^+ \right) \sin(2\eta) \sigma_{n-2}^- \\ + \left( A_{n-3}(\lambda_\alpha) \sin(2\eta) \sigma_{n-3}^- + B_{n-3}(\lambda_\alpha) \sin(\lambda_\alpha - v_{n-3} - \eta \sigma_{n-3}^z) \right) \sin(\lambda_\alpha - v_{n-2} - \eta \sigma_{n-2}^z) ,$$

which is equivalent to the superposition,

$$A_{n-3}(\lambda_\alpha) \sin(2\eta) \sigma_{n-2}^- \sin(\lambda_\alpha - v_{n-3} + \eta \sigma_{n-3}^z) + B_{n-3}(\lambda_\alpha) (\sin(2\eta))^2 \left( \prod_{2 \leq i \leq 3} \sigma_{n-i}^{+,-} \right) ,$$

for the first term, and,

$$A_{n-3}(\lambda_\alpha) \sin(2\eta) \sigma_{n-3}^- \sin(\lambda_\alpha - v_{n-2} - \eta \sigma_{n-2}^z) + B_{n-3}(\lambda_\alpha) \left( \prod_{2 \leq i \leq 3} \sin(\lambda_\alpha - v_{n-i} - \eta \sigma_{n-i}^z) \right) ,$$

for the second term.

From the two terms above, substituting for  $A_{n-3}(\lambda_\alpha)$  and  $B_{n-3}(\lambda_\alpha)$  yields, for the first term,

$$\begin{aligned} & \left( A_{n-4}(\lambda_\alpha) \sin(\lambda_\alpha - v_{n-4} + \eta\sigma_{n-4}^z) + B_{n-4}(\lambda_\alpha) \sin(2\eta) \sigma_{n-4}^+ \right) \sin(2\eta) \sigma_{n-2}^- \sin(\lambda_\alpha - v_{n-3} + \eta\sigma_{n-3}^z) \\ & + \left( A_{n-4}(\lambda_\alpha) \sin(2\eta) \sigma_{n-4}^- + B_{n-4}(\lambda_\alpha) \sin(\lambda_\alpha - v_{n-4} - \eta\sigma_{n-4}^z) \right) (\sin(2\eta))^2 \left( \prod_{2 \leq i \leq 3} \sigma_{n-i}^{+, -} \right), \end{aligned} \quad (1^* - 1)$$

while for the second term,

$$\begin{aligned} & \left( A_{n-4}(\lambda_\alpha) \sin(2\eta) \sigma_{n-4}^- + B_{n-4}(\lambda_\alpha) \sin(2\eta) \sigma_{n-4}^+ \right) \sin(2\eta) \sigma_{n-3}^- \sin(\lambda_\alpha - v_{n-2} - \eta\sigma_{n-2}^z) \\ & + \left( A_{n-4}(\lambda_\alpha) \sin(2\eta) \sigma_{n-4}^- + B_{n-4}(\lambda_\alpha) \sin(\lambda_\alpha - v_{n-4} - \eta\sigma_{n-4}^z) \right) \left( \prod_{2 \leq i \leq 3} \sin(\lambda_\alpha - v_{n-i} - \eta\sigma_{n-i}^z) \right). \end{aligned} \quad (1^* - 2)$$

Performing rearrangements from  $(1^* - 1)$  implies,

$$\begin{aligned} & A_{n-4}(\lambda_\alpha) \sin(2\eta) \sigma_{n-2}^- \left( \prod_{3 \leq i \leq 4} \sin(\lambda_\alpha - v_{n-i} + \eta\sigma_{n-i}^z) \right) + B_{n-4}(\lambda_\alpha) (\sin(2\eta))^2 \sigma_{n-2}^- \sin(\lambda_\alpha - v_{n-3} + \eta\sigma_{n-3}^z) \sigma_{n-4}^+ \\ & + A_{n-4}(\lambda_\alpha) (\sin(2\eta))^3 \left( \prod_{2 \leq i \leq 4} \sigma_{n-i}^{+, -} \right) + B_{n-4}(\lambda_\alpha) (\sin(2\eta))^2 \sin(\lambda_\alpha - v_{n-4} - \eta\sigma_{n-4}^z) \left( \prod_{2 \leq i \leq 3} \sigma_{n-i}^{+, -} \right), \end{aligned}$$

while performing rearrangements from  $(1^* - 2)$  of terms implies,

$$\begin{aligned} & A_{n-4}(\lambda_\alpha) (\sin(2\eta))^2 \sin(\lambda_\alpha - v_{n-2} - \eta\sigma_{n-2}^z) \left( \prod_{3 \leq i \leq 4} \sigma_{n-i}^- \right) + B_{n-4}(\lambda_\alpha) (\sin(2\eta))^2 \sin(\lambda_\alpha - v_{n-2} - \eta\sigma_{n-2}^z) \\ & \times \left( \prod_{3 \leq i \leq 4} \sigma_{n-i}^{-, +} \right) \\ & + A_{n-4}(\lambda_\alpha) \sin(2\eta) \sigma_{n-4}^- \left( \prod_{2 \leq i \leq 3} \sin(\lambda_\alpha - v_{n-i} - \eta\sigma_{n-i}^z) \right) + B_{n-4}(\lambda_\alpha) \left( \prod_{2 \leq i \leq 4} \sin(\lambda_\alpha - v_{n-i} - \eta\sigma_{n-i}^z) \right). \end{aligned}$$

Extrapolating the formulas from previous computations yields,

$$\begin{aligned} & B_n(\lambda_\alpha) = \left( A_3(\lambda_\alpha) + B_3(\lambda_\alpha) \right) (\sin(2\eta))^{n-4} \left( \prod_{2 \leq i \leq n-3} \sigma_{n-i}^{-, +} \right) \\ & + \left( A_3(\lambda_\alpha) + B_3(\lambda_\alpha) \right) \left( \prod_{2 \leq i \leq n-3} \sin(\lambda_\alpha - v_{n-i} + \eta\sigma_{n-i}^z) \right) \\ & + \left( A_3(\lambda_\alpha) + B_3(\lambda_\alpha) \right) \left( \sum_{\substack{2 \leq i \leq m \\ 2 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{2 \leq i \leq m} \sin(\lambda_\alpha - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{2 \leq j \leq n'} \sigma_{n-j}^{-, +} \right) \right] \right). \end{aligned}$$

Along similar lines, repeating the computations since the beginning of the proof yields another formula for  $D_n(\lambda_\alpha)$ , in which,

$$\begin{aligned}
D_n(\lambda_\alpha) = & \left( C_3(\lambda_\alpha) + D_3(\lambda_\alpha) \right) (\sin(2\eta))^{n-4} \left( \prod_{2 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \\
& + \left( C_3(\lambda_\alpha) + D_3(\lambda_\alpha) \right) \left( \prod_{2 \leq i \leq n-3} \sin(\lambda_\alpha - v_{n-i} + \eta \sigma_{n-i}^z) \right) \\
& + \left( C_3(\lambda_\alpha) + D_3(\lambda_\alpha) \right) \left( \sum_{\substack{2 \leq i \leq m \\ 2 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{2 \leq i \leq m} \sin(\lambda_\alpha - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{2 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) ,
\end{aligned}$$

from which we conclude the argument.  $\square$

With the expressions from **Lemma 5** for  $A_n(\lambda_\alpha)$ ,  $B_n(\lambda_\alpha)$ ,  $C_n(\lambda_\alpha)$  and  $D_n(\lambda_\alpha)$ , we continue with the following computation. Recall, before **Lemma 5**, that we demonstrated that the product of L operators,

$$\prod_{0 \leq i \leq 3} \begin{bmatrix} \mathbf{1}^i & \mathbf{2}^i \\ \mathbf{3}^i & \mathbf{4}^i \end{bmatrix}$$

has the expansion,

$$\begin{bmatrix} A_3(\lambda_\alpha) \mathbf{1}^3 + B_3(\lambda_\alpha) \mathbf{3}^3 & A_3(\lambda_\alpha) \mathbf{2}^3 + B_3(\lambda_\alpha) \mathbf{4}^3 \\ C_3(\lambda_\alpha) \mathbf{1}^3 + D_3(\lambda_\alpha) \mathbf{3}^3 & C_3(\lambda_\alpha) \mathbf{2}^3 + D_3(\lambda_\alpha) \mathbf{4}^3 \end{bmatrix} .$$

To make use of the expressions in the matrix product above, we perform additional computations below, in which,

$$\begin{bmatrix} \mathbf{1}^{i-2} & \mathbf{2}^{i-2} \\ \mathbf{3}^{i-2} & \mathbf{4}^{i-2} \end{bmatrix} \begin{bmatrix} \mathbf{1}^{i-1} & \mathbf{2}^{i-1} \\ \mathbf{3}^{i-1} & \mathbf{4}^{i-1} \end{bmatrix} \equiv \begin{bmatrix} \mathbf{1}^{i-2} \mathbf{1}^{i-1} + \mathbf{2}^{i-2} \mathbf{3}^{i-1} & \mathbf{1}^{i-2} \mathbf{2}^{i-1} + \mathbf{2}^{i-2} \mathbf{4}^{i-1} \\ \mathbf{3}^{i-2} \mathbf{1}^{i-1} + \mathbf{4}^{i-2} \mathbf{3}^{i-1} & \mathbf{3}^{i-2} \mathbf{2}^{i-1} + \mathbf{4}^{i-1} \mathbf{4}^{i-2} \end{bmatrix} .$$

Substituting the expression for the product above with the  $i-3$  th L operator implies,

$$\begin{bmatrix} \mathbf{1}^{i-3} & \mathbf{2}^{i-3} \\ \mathbf{3}^{i-3} & \mathbf{4}^{i-3} \end{bmatrix} \begin{bmatrix} \mathbf{1}^{i-2} \mathbf{1}^{i-1} + \mathbf{2}^{i-2} \mathbf{3}^{i-1} & \mathbf{1}^{i-2} \mathbf{2}^{i-1} + \mathbf{2}^{i-2} \mathbf{4}^{i-1} \\ \mathbf{3}^{i-2} \mathbf{1}^{i-1} + \mathbf{4}^{i-2} \mathbf{3}^{i-1} & \mathbf{3}^{i-2} \mathbf{2}^{i-1} + \mathbf{4}^{i-1} \mathbf{4}^{i-2} \end{bmatrix} ,$$

is equal to,

$$\begin{aligned}
& \left[ \frac{\prod_{1 \leq j \leq 3} \mathbf{1}^{i-j} + \mathbf{1}^{i-3} \mathbf{2}^{i-2} \mathbf{3}^{i-1} + \mathbf{1}^{i-1} \mathbf{2}^{i-3} \mathbf{3}^{i-2} + \mathbf{2}^{i-3} \mathbf{3}^{i-1} \mathbf{4}^{i-2}}{\mathbf{3}^{i-3} \prod_{1 \leq j \leq 2} \mathbf{1}^{i-j} + \mathbf{2}^{i-2} \prod_{j \text{ odd } : 1 \leq j \leq 3} \mathbf{3}^{i-j} + \mathbf{1}^{i-1} \mathbf{3}^{i-2} \mathbf{4}^{i-3} + \mathbf{3}^{i-1} \prod_{2 \leq j \leq 3} \mathbf{4}^{i-j}} \dots \right] \\
& \left[ \frac{\prod_{2 \leq j \leq 3} \mathbf{1}^{i-j} \mathbf{2}^{i-1} + \mathbf{1}^{i-3} \mathbf{2}^{i-2} \mathbf{4}^{i-1} + \prod_{j \text{ odd } : 1 \leq j \leq 3} \mathbf{2}^{i-j} \mathbf{3}^{i-2} + \mathbf{2}^{i-3} \prod_{1 \leq j \leq 2} \mathbf{4}^{i-j}}{\mathbf{1}^{i-2} \mathbf{2}^{i-1} \mathbf{3}^{i-3} + \mathbf{2}^{i-2} \mathbf{3}^{i-3} \mathbf{4}^{i-1} + \mathbf{2}^{i-1} \mathbf{3}^{i-2} \mathbf{4}^{i-3} + \prod_{1 \leq j \leq 3} \mathbf{4}^{i-j}} \dots \right] ,
\end{aligned}$$

from the fact that, each entry of the matrix above is respectively given by,

$$\begin{aligned}
& \mathbf{1}^{i-3} \left( \prod_{1 \leq j \leq 2} \mathbf{1}^{i-j} + \mathbf{2}^{i-2} \mathbf{3}^{i-1} \right) + \mathbf{2}^{i-3} \left( \mathbf{3}^{i-2} \mathbf{1}^{i-1} + \mathbf{4}^{i-2} \mathbf{3}^{i-1} \right) , \\
& \mathbf{1}^{i-3} \left( \mathbf{1}^{i-2} \mathbf{2}^{i-1} + \mathbf{2}^{i-2} \mathbf{4}^{i-1} \right) + \mathbf{2}^{i-3} \left( \mathbf{3}^{i-2} \mathbf{2}^{i-1} + \prod_{1 \leq j \leq 2} \mathbf{4}^{i-j} \right) , \\
& \mathbf{3}^{i-3} \left( \prod_{1 \leq j \leq 2} \mathbf{1}^{i-j} + \mathbf{2}^{i-2} \mathbf{3}^{i-1} \right) + \mathbf{4}^{i-3} \left( \mathbf{3}^{i-2} \mathbf{1}^{i-1} + \mathbf{4}^{i-2} \mathbf{3}^{i-1} \right) , \\
& \mathbf{3}^{i-3} \left( \mathbf{1}^{i-2} \mathbf{2}^{i-1} + \mathbf{2}^{i-2} \mathbf{4}^{i-1} \right) + \mathbf{4}^{i-3} \left( \mathbf{3}^{i-2} \mathbf{2}^{i-1} + \prod_{1 \leq j \leq 2} \mathbf{4}^{i-j} \right) .
\end{aligned}$$

Iterating the computation, by making use of the expression above, implies that the matrix product,

$$\begin{aligned}
& \left[ \frac{\prod_{1 \leq j \leq 3} \mathbf{1}^{i-j} + \mathbf{1}^{i-3} \mathbf{2}^{i-2} \mathbf{3}^{i-1} + \mathbf{1}^{i-1} \mathbf{2}^{i-3} \mathbf{3}^{i-2} + \mathbf{2}^{i-3} \mathbf{3}^{i-1} \mathbf{4}^{i-2}}{\mathbf{3}^{i-3} \prod_{1 \leq j \leq 2} \mathbf{1}^{i-j} + \mathbf{2}^{i-2} \prod_{j \text{ odd } : 1 \leq j \leq 3} \mathbf{3}^{i-j} + \mathbf{1}^{i-1} \mathbf{3}^{i-2} \mathbf{4}^{i-3} + \mathbf{3}^{i-1} \prod_{2 \leq j \leq 3} \mathbf{4}^{i-j}} \dots \right] \\
& \left[ \frac{\prod_{2 \leq j \leq 3} \mathbf{1}^{i-j} \mathbf{2}^{i-1} + \mathbf{1}^{i-3} \mathbf{2}^{i-2} \mathbf{4}^{i-1} + \prod_{j \text{ odd } : 1 \leq j \leq 3} \mathbf{2}^{i-j} \mathbf{3}^{i-2} + \mathbf{2}^{i-3} \prod_{1 \leq j \leq 2} \mathbf{4}^{i-j}}{\mathbf{1}^{i-2} \mathbf{2}^{i-1} \mathbf{3}^{i-3} + \mathbf{2}^{i-2} \mathbf{3}^{i-3} \mathbf{4}^{i-1} + \mathbf{2}^{i-1} \mathbf{3}^{i-2} \mathbf{4}^{i-3} + \prod_{1 \leq j \leq 3} \mathbf{4}^{i-j}} \right] \begin{bmatrix} \mathbf{1}^{i-1} & \mathbf{2}^{i-1} \\ \mathbf{3}^{i-1} & \mathbf{4}^{i-1} \end{bmatrix} ,
\end{aligned}$$

is equivalent to the matrix with entries that are respectively given by,

$$\text{First entry} \equiv \begin{cases} \mathbf{1}^{i-1}(\prod_{1 \leq j \leq 3} \mathbf{1}^{i-j} + \mathbf{1}^{i-3}\mathbf{2}^{i-2}\mathbf{3}^{i-1} + \mathbf{1}^{i-1}\mathbf{2}^{i-3}\mathbf{3}^{i-2} + \mathbf{2}^{i-3}\mathbf{3}^{i-1}\mathbf{4}^{i-2}) \\ + \mathbf{3}^{i-1}(\prod_{2 \leq j \leq 3} \mathbf{1}^{i-j}\mathbf{2}^{i-1} + \mathbf{1}^{i-3}\mathbf{2}^{i-2}\mathbf{4}^{i-1} + \prod_{j \text{ odd}} : 1 \leq j \leq 3 \mathbf{2}^{i-j}\mathbf{3}^{i-2} + \mathbf{2}^{i-3} \prod_{1 \leq j \leq 2} \mathbf{4}^{i-j}) \end{cases},$$

is equivalent to,

$$\begin{cases} (\mathbf{1}^{i-1})^2 \prod_{2 \leq j \leq 3} \mathbf{1}^{i-j} + \prod_{\text{odd } j : 1 \leq j \leq 3} \mathbf{1}^{i-j}\mathbf{2}^{i-2}\mathbf{3}^{i-1} + (\mathbf{1}^{i-1})^2\mathbf{2}^{i-3}\mathbf{3}^{i-2} + \mathbf{1}^{i-1}\mathbf{2}^{i-3}\mathbf{3}^{i-1}\mathbf{4}^{i-2} \\ + \prod_{2 \leq j \leq 3} \mathbf{1}^{i-j}\mathbf{2}^{i-1}\mathbf{3}^{i-1} + \mathbf{1}^{i-3}\mathbf{2}^{i-2}\mathbf{3}^{i-1}\mathbf{4}^{i-1} + \prod_{j \text{ odd}} : 1 \leq j \leq 3 \mathbf{2}^{i-j} \prod_{1 \leq j \leq 2} \mathbf{3}^{i-j} + \mathbf{2}^{i-3}(\mathbf{3}^{i-1})^2\mathbf{3}^{i-2} \prod_{1 \leq j \leq 2} \mathbf{4}^{i-j} \end{cases}$$

$$\text{Second entry} \equiv \begin{cases} \mathbf{2}^{i-1}(\prod_{1 \leq j \leq 3} \mathbf{1}^{i-j} + \mathbf{1}^{i-3}\mathbf{2}^{i-2}\mathbf{3}^{i-1} + \mathbf{1}^{i-1}\mathbf{2}^{i-3}\mathbf{3}^{i-2} + \mathbf{2}^{i-3}\mathbf{3}^{i-1}\mathbf{4}^{i-2}) \\ + \mathbf{4}^{i-1}(\prod_{2 \leq j \leq 3} \mathbf{1}^{i-j}\mathbf{2}^{i-1} + \mathbf{1}^{i-3}\mathbf{2}^{i-2}\mathbf{4}^{i-1} + \prod_{j \text{ odd}} : 1 \leq j \leq 3 \mathbf{2}^{i-j}\mathbf{3}^{i-2} + \mathbf{2}^{i-3} \prod_{1 \leq j \leq 2} \mathbf{4}^{i-j}) \end{cases},$$

is equivalent to,

$$\begin{cases} \prod_{1 \leq j \leq 3} \mathbf{1}^{i-j}\mathbf{2}^{i-1} + \mathbf{1}^{i-3} \prod_{1 \leq j \leq 2} \mathbf{2}^{i-j}\mathbf{3}^{i-1} + \mathbf{1}^{i-1}\mathbf{3}^{i-2} + \prod_{j \text{ odd}} : 1 \leq j \leq 3 \mathbf{2}^{i-j}\mathbf{3}^{i-1}\mathbf{4}^{i-2} \\ + \prod_{2 \leq j \leq 3} \mathbf{1}^{i-j}\mathbf{2}^{i-1}\mathbf{4}^{i-1} + \mathbf{1}^{i-3}\mathbf{2}^{i-2}(\mathbf{4}^{i-1})^2 + \prod_{j \text{ odd}} : 1 \leq j \leq 3 \mathbf{2}^{i-j}\mathbf{3}^{i-2}\mathbf{4}^{i-1} + \mathbf{2}^{i-3}(\mathbf{4}^{i-1})^2\mathbf{4}^{i-2} \end{cases},$$

$$\text{Third entry} \equiv \begin{cases} \mathbf{1}^{i-1}(\mathbf{3}^{i-3} \prod_{1 \leq j \leq 2} \mathbf{1}^{i-j} + \mathbf{2}^{i-2} \prod_{j \text{ odd}} : 1 \leq j \leq 3 \mathbf{3}^{i-j} + \mathbf{1}^{i-1}\mathbf{3}^{i-2}\mathbf{4}^{i-3} + \mathbf{3}^{i-1} \prod_{2 \leq j \leq 3} \mathbf{4}^{i-j}) \\ + \mathbf{3}^{i-1}(\prod_{2 \leq j \leq 3} \mathbf{1}^{i-j}\mathbf{2}^{i-1} + \mathbf{1}^{i-3}\mathbf{2}^{i-2}\mathbf{4}^{i-1} + \prod_{j \text{ odd}} : 1 \leq j \leq 3 \mathbf{2}^{i-j}\mathbf{3}^{i-2} + \mathbf{2}^{i-3} \prod_{1 \leq j \leq 2} \mathbf{4}^{i-j}) \end{cases},$$

is equivalent to,

$$\begin{cases} \mathbf{3}^{i-3}(\mathbf{1}^{i-1})^2\mathbf{1}^{i-2} + \mathbf{1}^{i-1}\mathbf{2}^{i-2} \prod_{j \text{ odd}} : 1 \leq j \leq 3 \mathbf{3}^{i-j} + (\mathbf{1}^{i-1})^2\mathbf{3}^{i-2}\mathbf{4}^{i-3} + \mathbf{1}^{i-1}\mathbf{3}^{i-1} \prod_{2 \leq j \leq 3} \mathbf{4}^{i-j} \\ + \mathbf{3}^{i-1} \prod_{2 \leq j \leq 3} \mathbf{1}^{i-j}\mathbf{2}^{i-1} + \mathbf{1}^{i-3}\mathbf{2}^{i-2}\mathbf{3}^{i-1}\mathbf{4}^{i-1} + \prod_{j \text{ odd}} : 1 \leq j \leq 3 \mathbf{2}^{i-j} \prod_{1 \leq j \leq 2} \mathbf{3}^{i-j} + \mathbf{2}^{i-3}\mathbf{3}^{i-1} \prod_{1 \leq j \leq 2} \mathbf{4}^{i-j} \end{cases},$$

$$\text{Fourth entry} \equiv \begin{cases} \mathbf{2}^{i-1}(\mathbf{3}^{i-3} \prod_{1 \leq j \leq 2} \mathbf{1}^{i-j} + \mathbf{2}^{i-2} \prod_{j \text{ odd}} : 1 \leq j \leq 3 \mathbf{3}^{i-j} + \mathbf{1}^{i-1}\mathbf{3}^{i-2}\mathbf{4}^{i-3} + \mathbf{3}^{i-1} \prod_{2 \leq j \leq 3} \mathbf{4}^{i-j}) \\ + \mathbf{4}^{i-1}(\prod_{2 \leq j \leq 3} \mathbf{1}^{i-j}\mathbf{2}^{i-1} + \mathbf{1}^{i-3}\mathbf{2}^{i-2}\mathbf{4}^{i-1} + \prod_{j \text{ odd}} : 1 \leq j \leq 3 \mathbf{2}^{i-j}\mathbf{3}^{i-2} + \mathbf{2}^{i-3} \prod_{1 \leq j \leq 2} \mathbf{4}^{i-j}) \end{cases},$$

is equivalent to,

$$\begin{cases} \mathbf{2}^{i-1}\mathbf{3}^{i-3} \prod_{1 \leq j \leq 2} \mathbf{1}^{i-j} + \prod_{1 \leq j \leq 2} \mathbf{2}^{i-j} \prod_{j \text{ odd}} : 1 \leq j \leq 3 \mathbf{3}^{i-j} + \mathbf{1}^{i-1}\mathbf{2}^{i-1}\mathbf{3}^{i-2}\mathbf{4}^{i-3} + \mathbf{2}^{i-1}\mathbf{3}^{i-1} \prod_{2 \leq j \leq 3} \mathbf{4}^{i-j} \\ + \prod_{2 \leq j \leq 3} \mathbf{1}^{i-j}\mathbf{2}^{i-1}\mathbf{4}^{i-1} + \mathbf{1}^{i-3}\mathbf{2}^{i-2}(\mathbf{4}^{i-1})^2 + \prod_{j \text{ odd}} : 1 \leq j \leq 3 \mathbf{2}^{i-j}\mathbf{3}^{i-2}\mathbf{4}^{i-1} + \mathbf{2}^{i-3}(\mathbf{4}^{i-1})^2\mathbf{4}^{i-2} \end{cases}.$$

Altogether, the expressions obtained for the entries above imply,

$$\begin{bmatrix} \mathbf{1}^{i-3} & \mathbf{2}^{i-3} \\ \mathbf{3}^{i-3} & \mathbf{4}^{i-3} \end{bmatrix} \begin{bmatrix} \mathbf{1}^{i-2} & \mathbf{2}^{i-2} \\ \mathbf{3}^{i-2} & \mathbf{4}^{i-2} \end{bmatrix} \begin{bmatrix} \mathbf{1}^{i-1} & \mathbf{2}^{i-1} \\ \mathbf{3}^{i-1} & \mathbf{4}^{i-1} \end{bmatrix} \equiv \begin{bmatrix} \text{First entry} & \text{Second entry} \\ \text{Third entry} & \text{Fourth entry} \end{bmatrix},$$

from which the desired form for the product of  $n$  L operators,

$$\prod_{1 \leq i \leq n} \begin{bmatrix} \mathbf{1}^i & \mathbf{2}^i \\ \mathbf{3}^i & \mathbf{4}^i \end{bmatrix},$$

would take the desired form, in which,

$$\begin{bmatrix} \text{First entry} & \text{Second entry} \\ \text{Third entry} & \text{Fourth entry} \end{bmatrix}$$

is equal to,

$$\begin{bmatrix} \left( A_{i-3}(\lambda_\alpha) + B_{i-3}(\lambda_\alpha) \right) \left( (\sin(2\eta))^{n-(i-3)} \mathcal{A}'_1 + \mathcal{A}'_2 + \mathcal{A}'_3 \right) & \dots \\ \left( C_{i-3}(\lambda_\alpha) + D_{i-3}(\lambda_\alpha) \right) \left( (\sin(2\eta))^{n-(i-3)} \mathcal{C}'_1 + \mathcal{C}'_2 + \mathcal{C}'_3 \right) & \dots \end{bmatrix}$$

$$\begin{bmatrix} \left( A_{i-3}(\lambda_\alpha) + B_{i-3}(\lambda_\alpha) \right) \left( (\sin(2\eta))^{n-(i-4)} \mathcal{B}'_1 + \mathcal{B}'_2 + \mathcal{B}'_3 \right) \\ \left( C_{i-3}(\lambda_\alpha) + D_{i-3}(\lambda_\alpha) \right) \left( (\sin(2\eta))^{n-(i-4)} \mathcal{D}'_1 + \mathcal{D}'_2 + \mathcal{D}'_3 \right) \end{bmatrix} ,$$

for,

$$\begin{aligned} \mathcal{A}'_1 \equiv \mathcal{C}'_1 &\equiv \prod_{1 \leq i \leq n-(i-3)} \sigma_{n-i}^{-,+} , \\ \mathcal{A}'_2 \equiv \mathcal{C}'_2 &\equiv \prod_{1 \leq i \leq n-(i-3)} \sin(\lambda_\alpha - v_{n-i} + \eta \sigma_{n-i}^z) , \\ \mathcal{A}'_3 \equiv \mathcal{C}'_3 &\equiv \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-(i-3)}} \left[ \left( \prod_{1 \leq i \leq m} \sin(\lambda_\alpha - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] , \end{aligned}$$

and,

$$\begin{aligned} \mathcal{B}'_1 \equiv \mathcal{D}'_1 &\equiv \prod_{2 \leq i \leq n-(i-3)} \sigma_{n-i}^{-,+} , \\ \mathcal{B}'_2 \equiv \mathcal{D}'_2 &\equiv \prod_{2 \leq i \leq n-(i-3)} \sin(\lambda_\alpha - v_{n-i} + \eta \sigma_{n-i}^z) , \\ \mathcal{B}'_3 \equiv \mathcal{D}'_3 &\equiv \sum_{\substack{2 \leq i \leq m \\ 2 \leq j \leq n' \\ m+n'=n-(i-3)}} \left[ \left( \prod_{2 \leq i \leq m} \sin(\lambda_\alpha - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{2 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] . \end{aligned}$$

Iterating the previous arguments for computing the entries of each L operator, from the product,

$$\begin{aligned} &\left[ \begin{array}{c|c} \left( A_{i-3}(\lambda_\alpha) + B_{i-3}(\lambda_\alpha) \right) \left( (\sin(2\eta))^{n-(i-3)} \mathcal{A}'_1 + \mathcal{A}'_2 + \mathcal{A}'_3 \right) & \dots \\ \left( C_{i-3}(\lambda_\alpha) + D_{i-3}(\lambda_\alpha) \right) \left( (\sin(2\eta))^{n-(i-3)} \mathcal{C}'_1 + \mathcal{C}'_2 + \mathcal{C}'_3 \right) & \dots \end{array} \right] \\ &\left[ \begin{array}{c|c} \left( A_{i-3}(\lambda_\alpha) + B_{i-3}(\lambda_\alpha) \right) \left( (\sin(2\eta))^{n-(i-4)} \mathcal{B}'_1 + \mathcal{B}'_2 + \mathcal{B}'_3 \right) \\ \left( C_{i-3}(\lambda_\alpha) + D_{i-3}(\lambda_\alpha) \right) \left( (\sin(2\eta))^{n-(i-4)} \mathcal{D}'_1 + \mathcal{D}'_2 + \mathcal{D}'_3 \right) \end{array} \right] \prod_{i \leq i' \leq n} \begin{bmatrix} \mathbf{1}^{i'} & \mathbf{2}^{i'} \\ \mathbf{3}^{i'} & \mathbf{4}^{i'} \end{bmatrix} , \end{aligned}$$

yields similar expressions for  $\{\mathcal{A}_i\}_{1 \leq i \leq 3}$ ,  $\{\mathcal{B}_i\}_{1 \leq i \leq 3}$ ,  $\{\mathcal{C}_i\}_{1 \leq i \leq 3}$  and  $\{\mathcal{D}_i\}_{1 \leq i \leq 3}$ , from  $\{\mathcal{A}'_i\}_{1 \leq i \leq 3}$ ,  $\{\mathcal{B}'_i\}_{1 \leq i \leq 3}$ ,  $\{\mathcal{C}'_i\}_{1 \leq i \leq 3}$  and  $\{\mathcal{D}'_i\}_{1 \leq i \leq 3}$ . We conclude the argument.  $\square$

## 2.3 Returning to the quantum monodromy matrix

From expressions obtained for entries of the monodromy matrix,

$$\begin{bmatrix} A(\lambda_\alpha) & B(\lambda_\alpha) \\ C(\lambda_\alpha) & D(\lambda_\alpha) \end{bmatrix}$$

in terms of  $A_3(\lambda_\alpha)$ ,  $B_3(\lambda_\alpha)$ ,  $C_3(\lambda_\alpha)$  and  $D_3(\lambda_\alpha)$ , below we perform the following computations with respect to the Poisson bracket. Below, we restate the main result of the paper which was provided earlier in 1.3. Before stating the final result that needs to be proved, we reminder the reader of relationships obtained in the expressions for the monodromy matrix of the nonlinear Schrodinger's equation provided in 1.4.1, in which, as  $x \rightarrow +\infty$  and as  $y \rightarrow -\infty$ , independently of  $x$  and  $y$ , the monodromy matrix, [5],

$$T(\lambda) \equiv \begin{bmatrix} A(\lambda) & B(\lambda) \\ C(\lambda) & D(\lambda) \end{bmatrix} ,$$

can be expressed with the following infinite limit,

$$\lim_{N \rightarrow +\infty} T_N(\lambda) \equiv T(\lambda) = \lim_{\substack{x \rightarrow +\infty \\ y \rightarrow -\infty}} E(-x, \lambda) T(x, y, \lambda) E(y, \lambda) ,$$

where  $E$  is the following matrix exponential that is proportional to the Pauli matrix  $U_1$ ,

$$E \equiv E(x, \lambda) \equiv \exp(\lambda x U_1) .$$

For the inhomogeneous six-vertex model, we take the Poisson bracket of the tensor product between  $T_a(u, \{v_k\})$  and  $T_a(u', \{v'_k\})$ ,

$$T_a(u, \{v_k\}, H, V) \equiv T_a(u, \{v_k\}) \equiv \begin{bmatrix} A(u) & B(u) \\ C(u) & D(u) \end{bmatrix} ,$$

with,

$$T_a(u', \{v'_k\}, H, V) \equiv T_a(u', \{v'_k\}) \equiv \begin{bmatrix} A(u') & B(u') \\ C(u') & D(u') \end{bmatrix} ,$$

in which,

$$\begin{aligned} \{T_a(u, \{v_k\}) \otimes T_a(u', \{v'_k\})\} &= \left( r_{a,+}(v_k - v'_k) T_a(u, \{v_k\}) \right) \otimes T_a(u', \{v'_k\}) \\ &\quad - T_a(u, \{v_k\}) \otimes \left( T_a(u', \{v'_k\}) r_{a,-}(v_k - v'_k) \right) , \end{aligned}$$

corresponding to the tensor product of the Poisson bracket of  $T_a(u, \{v_k\})$  with  $T_a(u', \{v'_k\})$ , where,

$$\begin{aligned} r_{a,+}(v_k - v'_k) &= \lim_{y \rightarrow +\infty} \left( E^{6V}(u', v'_k - v_k) \otimes \left( E^{6V}(u', v'_k - v_k) r_a(v_k - v'_k) \right) \right) , \\ r_{a,-}(v_k - v'_k) &= \lim_{y \rightarrow -\infty} \left( E^{6V}(u', v'_k - v_k) \otimes \left( E^{6V}(u', v'_k - v_k) r_a(v_k - v'_k) \right) \right) , \end{aligned}$$

for the spectral parameter  $v'_k$  at  $u'$ , and,

$$r_a(v_k - v'_k) \equiv E^{6V}(u', v'_k - v_k) \otimes \left( E^{6V}(u', v'_k - v_k) r_a(v_k - v'_k) \right) .$$

To demonstrate that the expression,

$$E^{6V}(x - v_k, x) = \exp[\coth(\frac{\eta}{2} + i\alpha_j - v_k)] = \exp[\mathcal{U}_1^j] ,$$

**Lemma BE 1** (*mapping of the Bethe equations into a higher-dimensional space*). Fix solutions to the Bethe equations,  $\alpha_j$ . For each  $j$ , there exists functions  $\mathcal{U}_1^j, \mathcal{U}_2^j, \mathcal{U}_3^j$  and  $\mathcal{U}_4^j$ , satisfying the following conditions, for functions,

$$F_1 \equiv F_1(\frac{1}{2}, \alpha_j, -v_k) \equiv \sinh(\frac{\eta}{2} + i\alpha_j - v_k) ,$$

and,

$$F_2 \equiv F_2(\frac{1}{2}, -\alpha_j, v_k) \equiv \sinh(\frac{\eta}{2} - i\alpha_j + v_k) ,$$

from which one obtains the system,

$$\begin{aligned}\frac{\partial F_1}{\partial v_k} &= \mathcal{U}_1^j \left( \frac{\eta}{2}, \alpha_j, -v_k \right) F_1 \quad , \\ \frac{\partial F_2}{\partial v_k} &= \mathcal{U}_2^j \left( \frac{\eta}{2}, -\alpha_j, v_k \right) F_2 \quad ,\end{aligned}$$

for terms on the left hand side of the Bethe equations, under the product from  $k = 1$  to  $k = N$ , while, for terms on the right hand side of the Bethe equations, introduce,

$$F_3 \equiv F_3(1, \alpha_j - \alpha_m, 0) \equiv \sinh(i(\alpha_j - \alpha_m) + \eta) \quad ,$$

and,

$$F_4 \equiv F_4(-1, \alpha_j - \alpha_m, 0) \equiv \sinh(i(\alpha_j - \alpha_m) - \eta) \quad ,$$

from which one obtains the system,

$$\begin{aligned}\frac{\partial F_3}{\partial \alpha_m} &= \mathcal{U}_3^j \left( 1, \alpha_j - \alpha_m, 0 \right) F_3(1, \alpha_j - \alpha_m, 0) \quad , \\ \frac{\partial F_4}{\partial \alpha_m} &= \mathcal{U}_4^j \left( -1, \alpha_j - \alpha_m, 0 \right) F_4(-1, \alpha_j - \alpha_m, 0) \quad ,\end{aligned}$$

under the product from  $m = 1$  to  $m = n$ , except for  $m = j$ .

For any collection of solutions  $\{\alpha_j\}_{j \in \mathbb{N}}$ ,

$$\begin{aligned}\mathcal{U}_1^N \left( \frac{\eta}{2}, \alpha_j, -v_k \right) &= \bigcup_{j \in \mathbb{N}} \mathcal{U}_1^j \left( \frac{\eta}{2}, \alpha_j, -v_k \right) \quad , \\ \mathcal{U}_2^N \left( \frac{\eta}{2}, -\alpha_j, v_k \right) &= \bigcup_{j \in \mathbb{N}} \mathcal{U}_2^j \left( \frac{\eta}{2}, -\alpha_j, v_k \right) \quad , \\ \mathcal{U}_3^N \left( 1, \alpha_j - \alpha_m, 0 \right) &= \bigcup_{j \in \mathbb{N}} \mathcal{U}_3^j \left( 1, \alpha_j - \alpha_m, 0 \right) \quad , \\ \mathcal{U}_4^N \left( -1, \alpha_j - \alpha_m, 0 \right) &= \bigcup_{j \in \mathbb{N}} \mathcal{U}_4^j \left( -1, \alpha_j - \alpha_m, 0 \right) \quad .\end{aligned}$$

*Proof of Lemma BE 1.* Observe, from,

$$\frac{\partial}{\partial v_k} \left( \sinh \left( \frac{\eta}{2} + i\alpha_j - v_k \right) \right) \quad ,$$

that one can write,

$$\left( \frac{\sinh \left( \frac{\eta}{2} + i\alpha_j - v_k \right)}{\sinh \left( \frac{\eta}{2} + i\alpha_j - v_k \right)} \right) \left( \frac{\cosh \left( \frac{\eta}{2} + i\alpha_j - v_k \right)}{\cosh \left( \frac{\eta}{2} + i\alpha_j - v_k \right)} \right) \cosh \left( \frac{\eta}{2} + i\alpha_j - v_k \right) = \left( \coth \left( \frac{\eta}{2} + i\alpha_j - v_k \right) \right) \sinh \left( \frac{\eta}{2} + i\alpha_j - v_k \right) \quad .$$

Also, observe, from,

$$\frac{\partial}{\partial v_k} \left( \sinh \left( \frac{\eta}{2} + i\alpha_j + v_k \right) \right) \quad ,$$

that one can write,

$$\left( \frac{\sinh \left( \frac{\eta}{2} + i\alpha_j + v_k \right)}{\sinh \left( \frac{\eta}{2} + i\alpha_j + v_k \right)} \right) \left( \frac{\cosh \left( \frac{\eta}{2} + i\alpha_j + v_k \right)}{\cosh \left( \frac{\eta}{2} + i\alpha_j + v_k \right)} \right) \cosh \left( \frac{\eta}{2} + i\alpha_j + v_k \right) = \left( \coth \left( \frac{\eta}{2} + i\alpha_j + v_k \right) \right) \sinh \left( \frac{\eta}{2} + i\alpha_j + v_k \right) \quad ,$$

implying,

$$\mathcal{U}_1^j = \coth\left(\frac{\eta}{2} + i\alpha_j - v_k\right) ,$$

and,

$$\mathcal{U}_2^j = \coth\left(\frac{\eta}{2} + i\alpha_j + v_k\right) .$$

For the remaining two functions, similarly observe, from,

$$\frac{\partial}{\partial \alpha_m} (\sinh(i(\alpha_j - \alpha_m) + \eta)) ,$$

that one can write,

$$-\left(\frac{\sinh(i(\alpha_j - \alpha_m) + \eta)}{\sinh(i(\alpha_j - \alpha_m) + \eta)}\right)\left(\frac{\cosh(i(\alpha_j - \alpha_m) + \eta)}{\cosh(i(\alpha_j - \alpha_m) + \eta)}\right)\cosh(i(\alpha_j - \alpha_m) + \eta) = -\left(\coth(i(\alpha_j - \alpha_m) + \eta)\right) \times \sinh(i(\alpha_j - \alpha_m) + \eta) .$$

Also, observe from,

$$\frac{\partial}{\partial \alpha_m} (\sinh(i(\alpha_j - \alpha_m) - \eta)) ,$$

that one can write,

$$-\left(\frac{\sinh(i(\alpha_j - \alpha_m) - \eta)}{\sinh(i(\alpha_j - \alpha_m) - \eta)}\right)\left(\frac{\cosh(i(\alpha_j - \alpha_m) - \eta)}{\cosh(i(\alpha_j - \alpha_m) - \eta)}\right)\cosh(i(\alpha_j - \alpha_m) - \eta) = -\left(\coth(i(\alpha_j - \alpha_m) - \eta)\right) \times \sinh(i(\alpha_j - \alpha_m) - \eta) ,$$

implying,

$$\mathcal{U}_3^j = -\coth(i(\alpha_j - \alpha_m) + \eta) .$$

and,

$$\mathcal{U}_4^j = \coth(i(\alpha_j - \alpha_m) - \eta) .$$

One can directly obtain the functions  $\mathcal{U}_1^j, \mathcal{U}_2^j, \mathcal{U}_3^j$  and  $\mathcal{U}_4^j$  for any  $j$ , which take the form as unions over  $\mathbb{N}$  provided at the end of the statement for **Lemma BE 1**, from which we conclude the argument.  $\square$

**Lemma BE 2** (*mapping the Bethe equations to the four functions obtained in the previous result*). From the functions obtained in the previous result, the Bethe equations can be mapped to the relation,

$$\prod_{k=1}^N \left[ \frac{\mathcal{U}_1^j\left(\frac{\eta}{2}, \alpha_j, -v_k\right) F_1\left(\frac{1}{2}, \alpha_j, -v_k\right)}{\mathcal{U}_2^j\left(\frac{\eta}{2}, -\alpha_j, v_k\right) F_2\left(\frac{1}{2}, -\alpha_j, v_k\right)} \right] = \exp(2HN) \prod_{m=1, m \neq j}^n \left[ \frac{\mathcal{U}_3^j(1, \alpha_j - \alpha_m, 0) F_3(1, \alpha_j - \alpha_m, 0)}{\mathcal{U}_4^j(-1, \alpha_j - \alpha_m, 0) F_4(-1, \alpha_j - \alpha_m, 0)} \right] .$$

*Proof of Lemma BE 2.* Observe,

$$\prod_{k=1}^N \frac{\partial F_1\left(\frac{1}{2}, \alpha_j, -v_k\right)}{\partial F_2\left(\frac{1}{2}, -\alpha_j, v_k\right)} \stackrel{\text{(Lemma BE 1)}}{=} \prod_{k=1}^N \frac{\mathcal{U}_1^j\left(\frac{\eta}{2}, \alpha_j, -v_k\right) F_1\left(\frac{1}{2}, \alpha_j, -v_k\right)}{\mathcal{U}_2^j\left(\frac{\eta}{2}, -\alpha_j, v_k\right) F_2\left(\frac{1}{2}, -\alpha_j, v_k\right)} , \quad (*)$$

from dividing,

$$\prod_{k=1}^N \frac{\partial F_1}{\partial v_k} \left( \frac{1}{2}, \alpha_j, -v_k \right) \stackrel{\text{(Lemma BE 1)}}{=} \prod_{k=1}^N \mathcal{U}_1^j \left( \frac{\eta}{2}, \alpha_j, -v_k \right) F_1 \left( \frac{1}{2}, \alpha_j, -v_k \right) ,$$

with,

$$\prod_{k=1}^N \frac{\partial F_2}{\partial v_k} \left( \frac{1}{2}, -\alpha_j, v_k \right) \stackrel{\text{(Lemma BE 1)}}{=} \prod_{k=1}^N \mathcal{U}_2^j \left( \frac{\eta}{2}, -\alpha_j, v_k \right) F_2 \left( \frac{1}{2}, -\alpha_j, v_k \right) ,$$

for the left hand side of the Bethe equations, and,

$$\prod_{m=1, m \neq j}^n \frac{\partial F_3(1, \alpha_j - \alpha_m, 0)}{\partial F_4(-1, \alpha_j - \alpha_m, 0)} \stackrel{\text{(Lemma BE 1)}}{=} \prod_{m=1, m \neq j}^n \frac{\mathcal{U}_3^j(1, \alpha_j - \alpha_m, 0) F_3(1, \alpha_j - \alpha_m, 0)}{\mathcal{U}_4^j(-1, \alpha_j - \alpha_m, 0) F_4(-1, \alpha_j - \alpha_m, 0)} , \quad (**)$$

from dividing,

$$\prod_{m=1}^n \frac{\partial F_3}{\partial \alpha_m} (1, \alpha_j - \alpha_m, 0) \stackrel{\text{(Lemma BE 1)}}{=} \prod_{m=1}^n \mathcal{U}_3^j (1, \alpha_j - \alpha_m, 0) \sinh(i(\alpha_j - \alpha_m) + \eta) ,$$

with,

$$\prod_{m=1, m \neq j}^n \frac{\partial F_4}{\partial \alpha_m} (1, \alpha_j - \alpha_m, 0) \stackrel{\text{(Lemma BE 1)}}{=} \prod_{m=1, m \neq j}^n \mathcal{U}_4^j (-1, \alpha_j - \alpha_m, 0) \sinh(i(\alpha_j - \alpha_m) - \eta) ,$$

and multiplying the quotient with,

$$\exp(2HN) ,$$

for the right hand side of the Bethe equations. Altogether,

$$(*) = (**) ,$$

from which we conclude the argument.  $\square$

From the previous two results, in the next result below we express  $\frac{F_1}{F_2}$  and  $\frac{F_3}{F_4}$ , in terms of basis functions that are dependent on each solution  $\alpha_k$  to the Bethe equations.

**Lemma BE 3** (*expressing  $F_1, F_2, F_3$ , and  $F_4$  in the solution space for the Bethe equation*). There exists functions  $\mathcal{S}_1 \equiv \mathcal{S}_1(\eta, \alpha_j, v_k)$  and  $\mathcal{S}_2 \equiv \mathcal{S}_2(\eta, \alpha_j, v_k)$ , for which,

$$F_{1,2} = \mathcal{S}_1 F_1 + \mathcal{S}_2 F_2 ,$$

in the basis spanned by  $F_1$  and  $F_2$ , as well as functions  $\mathcal{S}_3 \equiv \mathcal{S}_3(\eta, \alpha_j, v_k)$  and  $\mathcal{S}_4 \equiv \mathcal{S}_4(\eta, \alpha_j, v_k)$ , for which,

$$F_{3,4} = \mathcal{S}_3 F_3 + \mathcal{S}_4 F_4 ,$$

in the basis spanned by  $F_3$  and  $F_4$ .

*Proof of Lemma BE 3.* By the definition of hyperbolic sine, write,

$$\frac{F_1\left(\frac{1}{2}, \alpha_j, -v_k\right)}{F_2\left(\frac{1}{2}, -\alpha_j, v_k\right)} = \frac{\sinh\left(\frac{\eta}{2} + i\alpha_j - v_k\right)}{\sinh\left(\frac{\eta}{2} - i\alpha_j + v_k\right)} = \frac{\frac{1}{2}}{\frac{1}{2}} \left( \frac{-\exp\left(-\left(\frac{\eta}{2} + i\alpha_j - v_k\right)\right) + \exp\left(\frac{\eta}{2} + i\alpha_j - v_k\right)}{-\exp\left(-\left(\frac{\eta}{2} + i\alpha_j - v_k\right)\right) + \exp\left(\frac{\eta}{2} + i\alpha_j - v_k\right)} \right) ,$$

which is equivalent to,

$$\frac{-\exp\left(-\left(\frac{\eta}{2} + i\alpha_j - v_k\right)\right)}{-\exp\left(-\left(\frac{\eta}{2} + i\alpha_j - v_k\right)\right) + \exp\left(\frac{\eta}{2} + i\alpha_j - v_k\right)} + \frac{\exp\left(\frac{\eta}{2} + i\alpha_j - v_k\right)}{-\exp\left(-\left(\frac{\eta}{2} + i\alpha_j - v_k\right)\right) + \exp\left(\frac{\eta}{2} + i\alpha_j - v_k\right)} ,$$

after having separated exponentials in the numerator of the expression, which is in turn also equivalent to,

$$\begin{aligned} & \left[ \frac{\exp\left(-\left(\frac{\eta}{2} + i\alpha_j - v_k\right)\right)}{\sinh\left(\frac{\eta}{2} + i\alpha_j - v_k\right) (\exp\left(-\left(\frac{\eta}{2} + i\alpha_j - v_k\right)\right) - \exp\left(\frac{\eta}{2} + i\alpha_j - v_k\right))} \right] F_1\left(\frac{1}{2}, \alpha_j, -v_k\right) \\ & + \left[ \frac{\exp\left(\frac{\eta}{2} + i\alpha_j - v_k\right)}{\sinh\left(\frac{\eta}{2} - i\alpha_j + v_k\right) (-\exp\left(-\left(\frac{\eta}{2} + i\alpha_j - v_k\right)\right) + \exp\left(\frac{\eta}{2} + i\alpha_j - v_k\right))} \right] F_2\left(\frac{1}{2}, -\alpha_j, v_k\right) , \end{aligned}$$

in the basis spanned by  $F_1$  and  $F_2$ . On the other hand, again by the definition of hyperbolic sine, write,

$$\frac{F_3(1, \alpha_j - \alpha_m, 0)}{F_4(-1, \alpha_j - \alpha_m, 0)} = \frac{\sinh(i(\alpha_j - \alpha_m) + \eta)}{\sinh(i(\alpha_j - \alpha_m) - \eta)} = \frac{\frac{1}{2}}{\frac{1}{2}} \left( \frac{\exp(i(\alpha_j - \alpha_m) + \eta) - \exp((i(\alpha_m - \alpha_j) - \eta))}{\exp(i(\alpha_j - \alpha_m) - \eta) - \exp(i(\alpha_m - \alpha_j) + \eta)} \right) ,$$

which is equivalent to,

$$\frac{\exp(i(\alpha_j - \alpha_m) + \eta)}{\exp(i(\alpha_j - \alpha_m) - \eta) - \exp(i(\alpha_m - \alpha_j) + \eta)} - \frac{\exp(i(\alpha_m - \alpha_j) - \eta)}{\exp(i(\alpha_j - \alpha_m) - \eta) - \exp(i(\alpha_m - \alpha_j) + \eta)} ,$$

after having separated exponentials in the numerator of the expression, which in turn is also equivalent to,

$$\begin{aligned} & \left[ \frac{\exp(i(\alpha_j - \alpha_m) + \eta)}{\sinh(i(\alpha_j - \alpha_m) + \eta) (\exp(i(\alpha_j - \alpha_m) - \eta) - \exp(i(\alpha_m - \alpha_j) + \eta))} \right] F_3(1, \alpha_j - \alpha_m, 0) \\ & - \left[ \frac{\exp(i(\alpha_m - \alpha_j) - \eta)}{\sinh(i(\alpha_j - \alpha_m) - \eta) (\exp(i(\alpha_j - \alpha_m) - \eta) - \exp(i(\alpha_m - \alpha_j) + \eta))} \right] F_4(-1, \alpha_j - \alpha_m, 0) , \end{aligned}$$

for the basis spanned by  $F_3$  and  $F_4$ . Taking  $\mathcal{S}_1, \mathcal{S}_2, \mathcal{S}_3$ , and  $\mathcal{S}_4$  from the quantities above provides the desired form of the linear combination for  $F_{1,2}$  and  $F_{3,4}$ , from which we conclude the argument.  $\square$

**Lemma BE 4 (spanning set of the entire solution space of the Bethe equations).** For all solutions to the Bethe equations, the spanning set is equivalent to the union of spanning sets for each  $j$ .

## 2.4 Sixteen relations from the monodromy matrix

*Proof of Lemma BE 4.* It suffices to determine one basis,  $\{F_1^j, F_2^j\}$ , and another basis,  $\{F_3^j, F_4^j\}$ , for each  $j$ . The computation from the previous result can be repeated a countably many number of times for any solution to the Bethe equations, hence yielding the desired span from which we conclude the argument.  $\square$

With the action given by the exponential in the prefactor to the basis element  $F_1$ ,

$$\frac{\exp\left(-\left(\frac{\eta}{2} + i\alpha_j - v_k\right)\right)}{\exp\left(-\left(\frac{\eta}{2} + i\alpha_j - v_k\right)\right) - \exp\left(\frac{\eta}{2} + i\alpha_j - v_k\right)} ,$$

write,

$$\left( \frac{\exp\left(\frac{\eta}{2} + i\alpha_j - v_k\right)}{\exp\left(\frac{\eta}{2} + i\alpha_j - v_k\right)} \right) \frac{\exp\left(-\left(\frac{\eta}{2} + i\alpha_j - v_k\right)\right)}{\exp\left(-\left(\frac{\eta}{2} + i\alpha_j - v_k\right)\right) - \exp\left(\frac{\eta}{2} + i\alpha_j - v_k\right)} ,$$

so that rearranging terms gives,

$$\frac{1}{1 - \exp\left(2\left(\frac{\eta}{2} + i\alpha_j - v_k\right)\right)} .$$

Hence, the set of linear combinations under the functions  $F_1$  and  $F_2$

$$\left[ \frac{1}{1 - \exp\left(2\left(\frac{\eta}{2} + i\alpha_j - v_k\right)\right)} \right] F_1\left(\frac{1}{2}, \alpha_j, -v_k\right) + \left[ \frac{1}{-1 + \exp\left(2\left(\frac{\eta}{2} + i\alpha_j - v_k\right)\right)} \right] F_2\left(\frac{1}{2}, \alpha_j, v_k\right) ,$$

with a similar set of relations holding for the basis spanned by  $F_3$  and  $F_4$ .

Furthermore, from the Poisson bracket of the tensor product of  $T_a(u, \{v_k\})$  with  $T_a(u', \{v'_k\})$ ,

$$\left( r_{a,+}(v_k - v'_k) T_a(u, \{v_k\}) \right) \otimes T_a(u', \{v'_k\}) - T_a(u, \{v_k\}) \otimes \left( T_a(u', \{v'_k\}) r_{a,-}(v_k - v'_k) \right) ,$$

one can form a set of sixteen relations for the reduced monodromy matrices of the inhomogeneous six-vertex model, akin to the sixteen relations which are satisfied for the reduced monodromy matrices of the nonlinear Schrodinger's equation, [5],

$$\left\{ \begin{array}{l} (1) : \{A(u), A(u')\} , \\ (2) : \{A(u), B(u')\} , \\ (3) : \{A(u), C(u')\} , \\ (4) : \{A(u), D(u')\} , \\ (5) : \{B(u), A(u')\} , \\ (6) : \{B(u), B(u')\} , \\ (7) : \{B(u), C(u')\} , \\ (8) : \{B(u), D(u')\} , \\ (9) : \{C(u), A(u')\} , \\ (10) : \{C(u), B(u')\} , \\ (11) : \{C(u), C(u')\} , \\ (12) : \{C(u), D(u')\} , \\ (13) : \{D(u), A(u')\} , \\ (14) : \{D(u), B(u')\} , \\ (15) : \{D(u), C(u')\} , \\ (16) : \{D(u), D(u')\} , \end{array} \right.$$

from the tensor product, which in the coordinates  $u, u'$  is equivalent to,

$$\{T_a(u, \{v_k\}) \otimes T_a(u', \{v'_k\})\} = \left\{ \begin{bmatrix} A(u) & B(u) \\ C(u) & D(u) \end{bmatrix} \otimes \begin{bmatrix} A(u') & B(u') \\ C(u') & D(u') \end{bmatrix} \right\} ,$$

which by the definition of Poisson bracket of the tensor product of the two reduced monodromy matrices, is,

$$\left( \left( r_{a,+}(v_k - v'_k) \begin{bmatrix} A(u) & B(u) \\ C(u) & D(u) \end{bmatrix} \right) \otimes \begin{bmatrix} A(u') & B(u') \\ C(u') & D(u') \end{bmatrix} \right) - \left( \begin{bmatrix} A(u) & B(u) \\ C(u) & D(u) \end{bmatrix} \otimes \left( \begin{bmatrix} A(u') & B(u') \\ C(u') & D(u') \end{bmatrix} r_{a,-}(v_k - v'_k) \right) \right) .$$

From expressions that were previously obtained for each entry of the  $N$  th power of the monodromy matrix from L operators, the set of sixteen relations is equivalent to,

$$\begin{aligned}
(1) &: \left\{ \left( A_3(u) + B_3(u) \right) \left[ (\sin(2\eta))^{n-3} \mathcal{A}_1 + \mathcal{A}_2 + \mathcal{A}_3 \right], \left( A_3(u') + B_3(u') \right) \left[ (\sin(2\eta))^{n-3} \mathcal{A}'_1 + \mathcal{A}'_2 + \mathcal{A}'_3 \right] \right\}, \\
(2) &: \left\{ \left( A_3(u) + B_3(u) \right) \left[ (\sin(2\eta))^{n-3} \mathcal{A}_1 + \mathcal{A}_2 + \mathcal{A}_3 \right], \left( A_3(u') + B_3(u') \right) \left[ (\sin(2\eta))^{n-4} \mathcal{B}'_1 + \mathcal{B}'_2 + \mathcal{B}'_3 \right] \right\}, \\
(3) &: \left\{ \left( A_3(u) + B_3(u) \right) \left[ (\sin(2\eta))^{n-3} \mathcal{A}_1 + \mathcal{A}_2 + \mathcal{A}_3 \right], \left( C_3(u') + D_3(u') \right) \left[ (\sin(2\eta))^{n-3} \mathcal{C}'_1 + \mathcal{C}'_2 + \mathcal{C}'_3 \right] \right\}, \\
(4) &: \left\{ \left( A_3(u) + B_3(u) \right) \left[ (\sin(2\eta))^{n-3} \mathcal{A}_1 + \mathcal{A}_2 + \mathcal{A}_3 \right], \left( C_3(u') + D_3(u') \right) \left[ (\sin(2\eta))^{n-4} \mathcal{D}'_1 + \mathcal{D}'_2 + \mathcal{D}'_3 \right] \right\}, \\
(5) &: \left\{ \left( A_3(u) + B_3(u) \right) \left[ (\sin(2\eta))^{n-4} \mathcal{B}_1 + \mathcal{B}_2 + \mathcal{B}_3 \right], \left( A_3(u') + B_3(u') \right) \left[ (\sin(2\eta))^{n-3} \mathcal{A}'_1 + \mathcal{A}'_2 + \mathcal{A}'_3 \right] \right\}, \\
(6) &: \left\{ \left( A_3(u) + B_3(u) \right) \left[ (\sin(2\eta))^{n-4} \mathcal{B}_1 + \mathcal{B}_2 + \mathcal{B}_3 \right], \left( A_3(u') + B_3(u') \right) \left[ (\sin(2\eta))^{n-4} \mathcal{B}'_1 + \mathcal{B}'_2 + \mathcal{B}'_3 \right] \right\}, \\
(7) &: \left\{ \left( A_3(u) + B_3(u) \right) \left[ (\sin(2\eta))^{n-4} \mathcal{B}_1 + \mathcal{B}_2 + \mathcal{B}_3 \right], \left( C_3(u') + D_3(u') \right) \left[ (\sin(2\eta))^{n-3} \mathcal{C}'_1 + \mathcal{C}'_2 + \mathcal{C}'_3 \right] \right\}, \\
(8) &: \left\{ \left( A_3(u) + B_3(u) \right) \left[ (\sin(2\eta))^{n-4} \mathcal{B}_1 + \mathcal{B}_2 + \mathcal{B}_3 \right], \left( C_3(u') + D_3(u') \right) \left[ (\sin(2\eta))^{n-4} \mathcal{D}'_1 + \mathcal{D}'_2 + \mathcal{D}'_3 \right] \right\}, \\
(9) &: \left\{ \left( C_3(u) + D_3(u) \right) \left[ (\sin(2\eta))^{n-3} \mathcal{C}_1 + \mathcal{C}_2 + \mathcal{C}_3 \right], \left( A_3(u') + B_3(u') \right) \left[ (\sin(2\eta))^{n-3} \mathcal{A}'_1 + \mathcal{A}'_2 + \mathcal{A}'_3 \right] \right\}, \\
(10) &: \left\{ \left( C_3(u) + D_3(u) \right) \left[ (\sin(2\eta))^{n-3} \mathcal{C}_1 + \mathcal{C}_2 + \mathcal{C}_3 \right], \left( A_3(u') + B_3(u') \right) \left[ (\sin(2\eta))^{n-4} \mathcal{B}'_1 + \mathcal{B}'_2 + \mathcal{B}'_3 \right] \right\}, \\
(11) &: \left\{ \left( C_3(u) + D_3(u) \right) \left[ (\sin(2\eta))^{n-3} \mathcal{C}_1 + \mathcal{C}_2 + \mathcal{C}_3 \right], \left( C_3(u') + D_3(u') \right) \left[ (\sin(2\eta))^{n-3} \mathcal{C}'_1 + \mathcal{C}'_2 + \mathcal{C}'_3 \right] \right\}, \\
(12) &: \left\{ \left( C_3(u) + D_3(u) \right) \left[ (\sin(2\eta))^{n-3} \mathcal{C}_1 + \mathcal{C}_2 + \mathcal{C}_3 \right], \left( C_3(u') + D_3(u') \right) \left[ (\sin(2\eta))^{n-4} \mathcal{D}'_1 + \mathcal{D}'_2 + \mathcal{D}'_3 \right] \right\}, \\
(13) &: \left\{ \left( C_3(u) + D_3(u) \right) \left[ (\sin(2\eta))^{n-4} \mathcal{D}_1 + \mathcal{D}_2 + \mathcal{D}_3 \right], \left( A_3(u') + B_3(u') \right) \left[ (\sin(2\eta))^{n-3} \mathcal{A}'_1 + \mathcal{A}'_2 + \mathcal{A}'_3 \right] \right\}, \\
(14) &: \left\{ \left( C_3(u) + D_3(u) \right) \left[ (\sin(2\eta))^{n-4} \mathcal{D}_1 + \mathcal{D}_2 + \mathcal{D}_3 \right], \left( A_3(u') + B_3(u') \right) \left[ (\sin(2\eta))^{n-4} \mathcal{B}'_1 + \mathcal{B}'_2 + \mathcal{B}'_3 \right] \right\}, \\
(15) &: \left\{ \left( C_3(u) + D_3(u) \right) \left[ (\sin(2\eta))^{n-4} \mathcal{D}_1 + \mathcal{D}_2 + \mathcal{D}_3 \right], \left( C_3(u') + D_3(u') \right) \left[ (\sin(2\eta))^{n-3} \mathcal{C}'_1 + \mathcal{C}'_2 + \mathcal{C}'_3 \right] \right\}, \\
(16) &: \left\{ \left( C_3(u) + D_3(u) \right) \left[ (\sin(2\eta))^{n-4} \mathcal{D}_1 + \mathcal{D}_2 + \mathcal{D}_3 \right], \left( C_3(u') + D_3(u') \right) \left[ (\sin(2\eta))^{n-4} \mathcal{D}'_1 + \mathcal{D}'_2 + \mathcal{D}'_3 \right] \right\}.
\end{aligned}$$

From the set of all possible relations, those which vanish with respect to the Poisson bracket constitute the action-angle variables. The next result determines which relations, from the sixteen listed above, vanish.

**Theorem** (*action-angle variables of the Hamiltonian flow for the inhomogeneous six-vertex model*).

*Proof of Theorem.* To demonstrate that the statement above holds, it suffices to compute the Poisson bracket for each of the sixteen relations. Beginning with the first relation, write,

$$\begin{aligned}
(1) \equiv & \left\{ \left( A_3(u) + B_3(u) \right) \left( \sin(2\eta) \right)^{n-3} \mathcal{A}_1, \left( A_3(u') + B_3(u') \right) \left( \left( \sin(2\eta) \right)^{n-3} \mathcal{A}'_1 + \mathcal{A}'_2 + \mathcal{A}'_3 \right) \right\} \\
& + \left\{ \left( A_3(u) + B_3(u) \right) \mathcal{A}_2, \left( A_3(u') + B_3(u') \right) \left( \left( \sin(2\eta) \right)^{n-3} \mathcal{A}'_1 + \mathcal{A}'_2 + \mathcal{A}'_3 \right) \right\} \\
& + \left\{ \left( A_3(u) + B_3(u) \right) \mathcal{A}_3, \left( A_3(u') + B_3(u') \right) \left( \left( \sin(2\eta) \right)^{n-3} \mathcal{A}'_1 + \mathcal{A}'_2 + \mathcal{A}'_3 \right) \right\},
\end{aligned}$$

which can be further rearranged, after a second application of bilinearity of the Poisson bracket, as,

$$\begin{aligned}
& \left\{ \left( A_3(u) + B_3(u) \right) (\sin(2\eta))^{n-3} \mathcal{A}_1, \left( A_3(u') + B_3(u') \right) (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} \\
& + \left\{ \left( A_3(u) + B_3(u) \right) (\sin(2\eta))^{n-3} \mathcal{A}_1, \left( A_3(u') + B_3(u') \right) \mathcal{A}'_2 \right\} \\
& + \left\{ \left( A_3(u) + B_3(u) \right) (\sin(2\eta))^{n-3} \mathcal{A}_1, \left( A_3(u') + B_3(u') \right) \mathcal{A}'_3 \right\} \\
& + \left\{ \left( A_3(u) + B_3(u) \right) \mathcal{A}_2, \left( A_3(u') + B_3(u') \right) (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} \\
& + \left\{ \left( A_3(u) + B_3(u) \right) \mathcal{A}_2, \left( A_3(u') + B_3(u') \right) \mathcal{A}'_2 \right\} \\
& + \left\{ \left( A_3(u) + B_3(u) \right) \mathcal{A}_2, \left( A_3(u') + B_3(u') \right) \mathcal{A}'_3 \right\} \\
& + \left\{ \left( A_3(u) + B_3(u) \right) \mathcal{A}_3, \left( A_3(u') + B_3(u') \right) (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} \\
& + \left\{ \left( A_3(u) + B_3(u) \right) \mathcal{A}_3, \left( A_3(u') + B_3(u') \right) \mathcal{A}'_2 \right\} \\
& + \left\{ \left( A_3(u) + B_3(u) \right) \mathcal{A}_3, \left( A_3(u') + B_3(u') \right) \mathcal{A}'_3 \right\} .
\end{aligned}$$

With two more applications of bilinearity of the Poisson bracket to each term in the superposition above, the previous expression is equivalent to,

$$\begin{aligned}
& \left\{ A_3(u) (\sin(2\eta))^{n-3} \mathcal{A}_1, A_3(u') (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} + \left\{ A_3(u) (\sin(2\eta))^{n-3} \mathcal{A}_1, B_3(u') (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} \\
& + \left\{ B_3(u) (\sin(2\eta))^{n-3} \mathcal{A}_1, A_3(u') (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} + \left\{ B_3(u) (\sin(2\eta))^{n-3} \mathcal{A}_1, B_3(u') (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} \\
& + \left\{ A_3(u) (\sin(2\eta))^{n-3} \mathcal{A}_1, A_3(u') \mathcal{A}'_2 \right\} + \left\{ A_3(u) (\sin(2\eta))^{n-3} \mathcal{A}_1, B_3(u') \mathcal{A}'_2 \right\} \\
& + \left\{ B_3(u) (\sin(2\eta))^{n-3} \mathcal{A}_1, A_3(u') \mathcal{A}'_2 \right\} + \left\{ B_3(u) (\sin(2\eta))^{n-3} \mathcal{A}_1, B_3(u') \mathcal{A}'_2 \right\} \\
& + \left\{ A_3(u) (\sin(2\eta))^{n-3} \mathcal{A}_1, A_3(u') \mathcal{A}'_3 \right\} + \left\{ A_3(u) (\sin(2\eta))^{n-3} \mathcal{A}_1, B_3(u') \mathcal{A}'_3 \right\} \\
& + \left\{ B_3(u) (\sin(2\eta))^{n-3} \mathcal{A}_1, A_3(u') \mathcal{A}'_3 \right\} + \left\{ B_3(u) (\sin(2\eta))^{n-3} \mathcal{A}_1, B_3(u') \mathcal{A}'_3 \right\} \\
& + \left\{ A_3(u) \mathcal{A}_2, A_3(u') (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} + \left\{ B_3(u) \mathcal{A}_2, A_3(u') (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} \\
& + \left\{ B_3(u) \mathcal{A}_2, A_3(u') (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} + \left\{ B_3(u) \mathcal{A}_2, B_3(u') (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} \\
& + \left\{ A_3(u) \mathcal{A}_2, A_3(u') \mathcal{A}'_2 \right\} + \left\{ A_3(u) \mathcal{A}_2, B_3(u') \mathcal{A}'_2 \right\} + \left\{ B_3(u) \mathcal{A}_2, A_3(u') \mathcal{A}'_2 \right\} + \left\{ B_3(u) \mathcal{A}_2, B_3(u') \mathcal{A}'_2 \right\} \\
& + \left\{ A_3(u) \mathcal{A}_2, A_3(u') \mathcal{A}'_3 \right\} + \left\{ A_3(u) \mathcal{A}_2, B_3(u') \mathcal{A}'_3 \right\} + \left\{ B_3(u) \mathcal{A}_2, A_3(u') \mathcal{A}'_3 \right\} + \left\{ B_3(u) \mathcal{A}_2, B_3(u') \mathcal{A}'_3 \right\} \\
& + \left\{ A_3(u) \mathcal{A}_3, A_3(u') (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} + \left\{ A_3(u) \mathcal{A}_3, B_3(u') (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} \\
& + \left\{ B_3(u) \mathcal{A}_3, A_3(u') (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} + \left\{ B_3(u) \mathcal{A}_3, B_3(u') (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} \\
& + \left\{ A_3(u) \mathcal{A}_3, A_3(u') \mathcal{A}'_2 \right\} + \left\{ A_3(u) \mathcal{A}_3, B_3(u') \mathcal{A}'_2 \right\} + \left\{ B_3(u) \mathcal{A}_3, A_3(u') \mathcal{A}'_2 \right\} + \left\{ B_3(u) \mathcal{A}_3, B_3(u') \mathcal{A}'_2 \right\} \\
& + \left\{ A_3(u) \mathcal{A}_3, A_3(u') \mathcal{A}'_3 \right\} + \left\{ A_3(u) \mathcal{A}_3, B_3(u') \mathcal{A}'_3 \right\} + \left\{ B_3(u) \mathcal{A}_3, A_3(u') \mathcal{A}'_3 \right\} + \left\{ B_3(u) \mathcal{A}_3, B_3(u') \mathcal{A}'_3 \right\} ,
\end{aligned}$$

which can be expressed as,

$$\begin{aligned}
& \sum_{\substack{(\mathcal{P}_1, \mathcal{P}_2) \in (B_3(u), B_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (A_3(u), A_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (A_3(u), B_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (B_3(u), A_3(u'))}} \left\{ \mathcal{P}_1(\sin(2\eta))^{n-3} \mathcal{A}_1, \mathcal{P}_2(\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} \\
+ & \sum_{\substack{(\mathcal{P}_1, \mathcal{P}_2) \in (B_3(u), B_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (A_3(u), A_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (A_3(u), B_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (B_3(u), A_3(u'))}} \left\{ \mathcal{P}_1(\sin(2\eta))^{n-3} \mathcal{A}_1, \mathcal{P}_2 \mathcal{A}'_2 \right\} + \sum_{\substack{(\mathcal{P}_1, \mathcal{P}_2) \in (B_3(u), B_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (A_3(u), A_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (A_3(u), B_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (B_3(u), A_3(u'))}} \left\{ \mathcal{P}_1(\sin(2\eta))^{n-3} \mathcal{A}_1, \mathcal{P}_2 \mathcal{A}'_3 \right\} \\
& + \sum_{\substack{(\mathcal{P}_1, \mathcal{P}_2) \in (B_3(u), B_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (A_3(u), A_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (A_3(u), B_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (B_3(u), A_3(u'))}} \left\{ \mathcal{P}_1 \mathcal{A}_2, \mathcal{P}_2(\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} + \sum_{\substack{(\mathcal{P}_1, \mathcal{P}_2) \in (B_3(u), B_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (A_3(u), A_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (A_3(u), B_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (B_3(u), A_3(u'))}} \left\{ \mathcal{P}_1 \mathcal{A}_2, \mathcal{P}_2 \mathcal{A}'_2 \right\} \\
& + \sum_{\substack{(\mathcal{P}_1, \mathcal{P}_2) \in (B_3(u), B_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (A_3(u), A_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (A_3(u), B_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (B_3(u), A_3(u'))}} \left\{ \mathcal{P}_1 \mathcal{A}_2, \mathcal{P}_2 \mathcal{A}'_3 \right\} + \sum_{\substack{(\mathcal{P}_1, \mathcal{P}_2) \in (B_3(u), B_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (A_3(u), A_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (A_3(u), B_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (B_3(u), A_3(u'))}} \left\{ \mathcal{P}_1 \mathcal{A}_3, \mathcal{P}_2(\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} \\
& + \sum_{\substack{(\mathcal{P}_1, \mathcal{P}_2) \in (B_3(u), B_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (A_3(u), A_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (A_3(u), B_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (B_3(u), A_3(u'))}} \left\{ \mathcal{P}_1 \mathcal{A}_3, \mathcal{P}_2 \mathcal{A}'_2 \right\} + \sum_{\substack{(\mathcal{P}_1, \mathcal{P}_2) \in (B_3(u), B_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (A_3(u), A_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (A_3(u), B_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (B_3(u), A_3(u'))}} \left\{ \mathcal{P}_1 \mathcal{A}_3, \mathcal{P}_2 \mathcal{A}'_3 \right\} ,
\end{aligned}$$

after isolating Poisson brackets together in groups of four, from which thirty six terms from the superposition on the previous page can be expressed with nine summations. To lighten the notation in the superposition above, denote,

$$\mathcal{P} \equiv \begin{cases} (\mathcal{P}_1, \mathcal{P}_2) \in (B_3(u), B_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (A_3(u), A_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (A_3(u), B_3(u')) \\ (\mathcal{P}_1, \mathcal{P}_2) \in (B_3(u), A_3(u')) \end{cases}$$

For the Poisson bracket below, fix some  $(\mathcal{P}'(u), \mathcal{P}''(u)) \equiv (\mathcal{P}', \mathcal{P}'') \in \mathcal{P}$ . The relation above will be used for the other fifteen relations to determine which ones vanish with respect to the Poisson bracket. We make use of the formula, by bilinearity of the Poisson bracket,

$$\begin{aligned}
\sum_{\mathcal{P}} \left\{ \mathcal{P}'(u) + \mathcal{P}''(u), \mathcal{P}'(u') + \mathcal{P}''(u') \right\} = & \sum_{\mathcal{P}} \left[ \left\{ \mathcal{P}'(u), \mathcal{P}'(u') \right\} + \left\{ \mathcal{P}'(u), \mathcal{P}''(u) \right\} + \left\{ \mathcal{P}''(u), \mathcal{P}'(u') \right\} \right. \\
& \left. + \left\{ \mathcal{P}''(u), \mathcal{P}''(u') \right\} \right] ,
\end{aligned}$$

The other fifteen relations are included after having obtained approximate expressions for each Poisson bracket from the first relation provided above.

From each of the sixteen relations provided above as a summation over Poisson brackets, it is possible to determine which one of the relations vanishes by computing each one of the Poisson brackets individually. To this end, for the first relation, (1), rearrange terms from each one of the nine Poisson bracket terms. Before differentiating each expression which appears in the formula for the Poisson bracket, observe,



Below, we list several results for evaluating each of the nine Poisson brackets appearing in the first relation.

### 2.4.1 First Poisson bracket, $\mathcal{P}_1$

**Lemma 6** (*evaluating the first Poisson bracket in the first relation*). The first term,  $\mathcal{P}_1$ , approximately equals,

$$[(\sin(2\eta))^{n-3}\mathcal{C}_1]^2 \left[ \frac{A_3(u)B_3(u')}{u-u'} - \frac{A_3(u')B_3(u)}{u'-u} \right] .$$

*Proof of Lemma 6.* The first term,

$$\sum_{\mathcal{P}} \left\{ \mathcal{P}_1(\sin(2\eta))^{n-3}\mathcal{A}_1, \mathcal{P}_2(\sin(2\eta))^{n-3}\mathcal{A}'_1 \right\} ,$$

is equivalent to,

$$\left[ (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \right]^2 \sum_{\mathcal{P}} \left\{ \mathcal{P}_1, \mathcal{P}_2 \right\} .$$

Observe,

$$\left[ (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \right]^2 \left( \{A_3(u), B_3(u')\} + \{B_3(u), A_3(u')\} \right) ,$$

and by anticommutativity of the Poisson bracket, that,

$$\left[ (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \right]^2 \left( \{A_3(u), B_3(u')\} - \{A_3(u'), B_3(u)\} \right) ,$$

which can be further rearranged, from the observation that the summation of two Poisson brackets above are equivalent to,

$$\left[ (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \right]^2 \{A_3(u), B_3(u')\} - \left[ (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \right]^2 \{A_3(u'), B_3(u)\} ,$$

which is, approximately, equivalent to,

$$\left[ (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \right]^2 \left[ \frac{A_3(u)B_3(u')}{u-u'} \right] - \left[ (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \right]^2 \left[ \frac{A_3(u')B_3(u)}{u'-u} \right] ,$$

from the fact that the two Poisson brackets can be computed with,

$$\{A_3(u), B_3(u')\} = \frac{A_3(u)B_3(u')}{u-u'} ,$$

corresponding to the first bracket, and,

$$\{A_3(u'), B_3(u)\} = \frac{A_3(u')B_3(u)}{u'-u} ,$$

corresponding to the second bracket. Hence, the desired expression is,

$$\begin{aligned} \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 (\sin(2\eta))^{n-3} \mathcal{A}_1, \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} &\approx \left[ (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \right]^2 \left[ \frac{A_3(u) B_3(u')}{u - u'} \right] \\ &\quad - \left[ (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \right]^2 \left[ \frac{A_3(u') B_3(u)}{u' - u} \right] \end{aligned}$$

which is approximately,

$$\left[ (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \right]^2 \left[ \frac{A_3(u) B_3(u')}{u - u'} - \frac{A_3(u') B_3(u)}{u' - u} \right] ,$$

from which we conclude the argument.  $\square$

#### 2.4.2 Second Poisson bracket, $\mathcal{P}_2$

**Lemma 7** (*evaluating the second Poisson bracket in the first relation*). The second term,  $\mathcal{P}_2$ , approximately equals,

$$-[(\sin(2\eta))^{n-3} \mathcal{C}_1] \left[ \frac{A_3(u) B_3(u')}{u - u'} - \frac{B_3(u) A_3(u')}{u' - u} \right] .$$

*Proof of Lemma 7.* The second term,

$$\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 (\sin(2\eta))^{n-3} \mathcal{A}_1, \mathcal{P}_2 \mathcal{A}'_2 \right\} ,$$

is equivalent to,

$$\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right), \mathcal{P}_2 \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} ,$$

which equals, by Leibniz' rule of the Poisson bracket,

$$\begin{aligned} &\sum_{\mathcal{P}} \left[ \left\{ \mathcal{P}_1, \mathcal{P}_2 \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \right. \\ &\quad \left. + \mathcal{P}_1 \left\{ (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right), \mathcal{P}_2 \right\} \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] . \end{aligned}$$

The first Poisson bracket in the summation over  $\mathcal{P}$  above is equivalent to,

$$-\left\{ \mathcal{P}_2 \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), \mathcal{P}_1 \right\} (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) ,$$

by anticommutativity, and also to,

$$\begin{aligned} &-(\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \left[ \left\{ \mathcal{P}_2, \mathcal{P}_1 \right\} \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right. \\ &\quad \left. + \mathcal{P}_2 \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), \mathcal{P}_1 \right\} \right] , \end{aligned}$$

by Leibniz' rule. Proceeding, the first term in the expression above is equivalent to,

$$-\left(\sin(2\eta)\right)^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \left[ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \{A_3(u), B_3(u')\} \right. \\ \left. + \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \{B_3(u), A_3(u')\} \right] ,$$

while the second term in the expression above is equivalent to,

$$-\left(\sin(2\eta)\right)^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \left[ \mathcal{P}_2 \left( \{1, \mathcal{P}_2\} \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right. \right. \\ \left. \left. + \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), \mathcal{P}_2 \right\} \right] .$$

From the second bracket,

$$-\left(\sin(2\eta)\right)^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \left[ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \{A_3(u), B_3(u')\} \right. \\ \left. + \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \{B_3(u), A_3(u')\} \right. \\ \left. + \sum_{\mathcal{P}} \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), \mathcal{P}_2 \right\} \right] ,$$

we evaluate each remaining Poisson bracket appearing in the summation over  $\mathcal{P}$ , for each possible  $\mathcal{P}_2$ , by writing,

$$\left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), B_3(u') \right\} + \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), A_3(u') \right\} \\ + \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), B_3(u') \right\} + \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), A_3(u') \right\} .$$

As a result, each Poisson bracket from the superposition above can be expressed as,

$$\left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), B_3(u') \right\} = \left[ \frac{\partial}{\partial u'} \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] \frac{\partial B_3(u')}{\partial u'} \\ - \frac{\partial B_3(u')}{\partial u'} \left[ \frac{\partial}{\partial u'} \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] , \\ \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), A_3(u') \right\} = \left[ \frac{\partial}{\partial u'} \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] \frac{\partial A_3(u')}{\partial u'} \\ - \frac{\partial A_3(u')}{\partial u'} \left[ \frac{\partial}{\partial u'} \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] , \\ \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), B_3(u') \right\} = \left[ \frac{\partial}{\partial u'} \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] \frac{\partial B_3(u')}{\partial u'} \\ - \frac{\partial B_3(u')}{\partial u'} \left[ \frac{\partial}{\partial u'} \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] , \\ \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), A_3(u') \right\} = \left[ \frac{\partial}{\partial u'} \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] \frac{\partial A_3(u')}{\partial u'} \\ - \frac{\partial A_3(u')}{\partial u'} \left[ \frac{\partial}{\partial u'} \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] ,$$

$$\left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \equiv 0, A_3(u') \right\} = \left[ \frac{\partial}{\partial u'} \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] \frac{\partial A_3(u')}{\partial u'} \\ - \frac{\partial A_3(u')}{\partial u'} \left[ \frac{\partial}{\partial u'} \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] ,$$

each of which vanish, while the remaining Poisson brackets,

$$-(\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \left[ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \{A_3(u), B_3(u')\} \right. \\ \left. + \left[ \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right] \{B_3(u), A_3(u')\} \right] .$$

are equivalent to,

$$-(\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \left[ \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right] \left[ \frac{A_3(u) B_3(u')}{u - u'} \right] \\ -(\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \left[ \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right] \left[ \frac{B_3(u) A_3(u')}{u' - u} \right] ,$$

from the observation that,

$$\{A_3(u), B_3(u')\} = \left( \frac{\partial}{\partial u} A_3(u) \right) \left( \frac{\partial}{\partial u'} B_3(u') \right) - \left( \frac{\partial}{\partial u'} A_3(u) \right) \left( \frac{\partial}{\partial u} B_3(u') \right) ,$$

is approximately equivalent to,

$$\frac{A_3(u) B_3(u')}{u - u'} ,$$

and also that,

$$\{B_3(u), A_3(u')\} = -\{A_3(u'), B_3(u)\} = -\left[ \left( \frac{\partial}{\partial u} B_3(u) \right) \left( \frac{\partial}{\partial u'} A_3(u') \right) + \left( \frac{\partial}{\partial u'} B_3(u) \right) \left( \frac{\partial}{\partial u} A_3(u') \right) \right] ,$$

is approximately equivalent to,

$$\frac{B_3(u) A_3(u')}{u' - u} .$$

Hence,

$$\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 (\sin(2\eta))^{n-3} \mathcal{A}_1, \mathcal{P}_2 \mathcal{A}'_2 \right\} \approx -\left[ (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \right] \left[ \frac{A_3(u) B_3(u')}{u - u'} - \frac{B_3(u) A_3(u')}{u' - u} \right] ,$$

from which we conclude the argument.  $\square$

#### 2.4.3 Third Poisson bracket, $\mathcal{P}_3$

**Lemma 8** (*evaluating the third Poisson bracket in the first relation*). The third term approximately equals,

$$\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 (\sin(2\eta))^{n-3} \mathcal{A}_1, \mathcal{P}_2 \mathcal{A}'_3 \right\} \approx \sum_{n':m+n'=n-3} (\sin(2\eta))^{n'-1} \left[ \sum_{1 \leq i \leq m} \frac{\partial}{\partial u} \left( \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right) \right] \left[ \frac{\partial B_3(u)}{\partial u'} + \frac{\partial A_3(u)}{\partial u'} \right] \\ + \sum_{n':m+n'=n-3} (\sin(2\eta))^{n'-1} \left[ \sum_{1 \leq i \leq m} \frac{\partial}{\partial u'} \left( \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right) \right] \left[ \frac{\partial B_3(u)}{\partial u} + \frac{\partial A_3(u)}{\partial u} \right] \\ - (\sin(2\eta))^{n-3} \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( (\mathcal{C}_3)_{i,j} \right) \right] \left[ \frac{B_3(u) A_3(u')}{u - u'} + \frac{A_3(u) B_3(u')}{u - u'} \right] .$$

*Proof of Lemma 8.* The third term,

$$\sum_{\mathcal{P}} \left\{ \mathcal{P}_1(\sin(2\eta))^{n-3} \mathcal{A}_1, \mathcal{P}_2 \mathcal{A}'_3 \right\} ,$$

is equivalent to,

$$\sum_{\mathcal{P}} \left\{ \mathcal{P}_1(\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right), \mathcal{P}_2 \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) (\sin(2\eta))^{n'-1} \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) \right\} .$$

Applying Leibniz' rule to the Poisson bracket over  $\mathcal{P}$  yields the expression,

$$\begin{aligned} & \left\{ \mathcal{P}_1, \mathcal{P}_2 \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) (\sin(2\eta))^{n'-1} \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) \right\} (\sin(2\eta))^{n-3} \\ & + \mathcal{P}_1 \left\{ (\sin(2\eta))^{n-3}, \mathcal{P}_2 \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) (\sin(2\eta))^{n'-1} \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) \right\} , \end{aligned}$$

which, by anticommutativity of the Poisson bracket, equals,

$$\begin{aligned} & - \left\{ \mathcal{P}_2 \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) (\sin(2\eta))^{n'-1} \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right), \mathcal{P}_1 \right\} (\sin(2\eta))^{n-3} \\ & - \mathcal{P}_1 \left\{ \mathcal{P}_2 \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) (\sin(2\eta))^{n'-1} \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right), (\sin(2\eta))^{n-3} \right\} . \end{aligned}$$

Applying Leibniz' rule again to each Poisson bracket in the expression above yields,

$$\begin{aligned} & -(\sin(2\eta))^{n-3} \left[ \left\{ \mathcal{P}_2, \mathcal{P}_1 \right\} \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) (\sin(2\eta))^{n'-1} \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) \right. \\ & \left. + \mathcal{P}_2 \left\{ \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) (\sin(2\eta))^{n'-1} \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right), \mathcal{P}_1 \right\} \right] , \end{aligned}$$

corresponding to the first term, and,

$$\begin{aligned} & -\mathcal{P}_1 \left[ \left\{ \mathcal{P}_2, (\sin(2\eta))^{n-3} \right\} \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) (\sin(2\eta))^{n'-1} \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) \right. \\ & \left. + \mathcal{P}_2 \left\{ \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) (\sin(2\eta))^{n'-1} \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right), (\sin(2\eta))^{n-3} \right\} \right] , \end{aligned}$$

corresponding to the second term. In the first term to which we applied the Poisson bracket, the fact that,

$$\sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) ,$$

is equivalent to,

$$\sum_{n':m+n'=n-3} (\sin(2\eta))^{n'-1} \left( \left( \sum_{1 \leq i \leq m} \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \left( \sum_{1 \leq j \leq n'} \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) ,$$

implies that another application of Leibniz' rule yields,

$$\begin{aligned} & -(\sin(2\eta))^{n-3} \left[ \{\mathcal{P}_2, \mathcal{P}_1\} \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) (\sin(2\eta))^{n'-1} \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) \right. \\ & + \mathcal{P}_2 \left[ \left\{ \sum_{n':m+n'=n-3} (\sin(2\eta))^{n'-1}, \mathcal{P}_1 \right\} \left( \left( \sum_{1 \leq i \leq m} \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \left( \sum_{1 \leq j \leq n'} \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) \right] \\ & + \left. \sum_{n':m+n'=n-3} (\sin(2\eta))^{n'-1} \left\{ \left( \left( \sum_{1 \leq i \leq m} \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \left( \sum_{1 \leq j \leq n'} \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right), \mathcal{P}_1 \right\} \right] . \end{aligned}$$

Applying Leibniz' rule to the final Poisson bracket in the superposition above yields,

$$\begin{aligned} & -(\sin(2\eta))^{n-3} \left[ \{\mathcal{P}_2, \mathcal{P}_1\} \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) (\sin(2\eta))^{n'-1} \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) \right. \\ & + \mathcal{P}_2 \left[ \left\{ \sum_{n':m+n'=n-3} (\sin(2\eta))^{n'-1}, \mathcal{P}_1 \right\} \left( \left( \sum_{1 \leq i \leq m} \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \left( \sum_{1 \leq j \leq n'} \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) \right. \\ & + \left. \sum_{n':m+n'=n-3} (\sin(2\eta))^{n'-1} \left[ \left\{ \left( \sum_{1 \leq i \leq m} \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), \mathcal{P}_1 \right\} \left( \sum_{1 \leq j \leq n'} \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right. \right. \\ & \left. \left. + \left\{ \left( \sum_{1 \leq j \leq n'} \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right), \mathcal{P}_1 \right\} \left( \sum_{1 \leq i \leq m} \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right] \right] \right] . \end{aligned}$$

From the expression above, writing each Poisson bracket individually yields, as summations over  $\mathcal{P}$ ,

$$\begin{aligned} \sum_{\mathcal{P}} \{\mathcal{P}_2, \mathcal{P}_1\} &= -\sum_{\mathcal{P}} \{\mathcal{P}_1, \mathcal{P}_2\} = -\left( \{B_3(u), A_3(u')\} + \{A_3(u), B_3(u')\} \right) , \\ \sum_{\mathcal{P}} \left\{ \sum_{n':m+n'=n-3} (\sin(2\eta))^{n'-1}, \mathcal{P}_1 \right\} &\equiv 0 , \\ \sum_{\mathcal{P}} \left\{ \left( \sum_{1 \leq i \leq m} \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), \mathcal{P}_1 \right\} &, \\ \sum_{\mathcal{P}} \left\{ \left( \sum_{1 \leq j \leq n'} \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right), \mathcal{P}_1 \right\} &\equiv 0 . \end{aligned}$$

Evaluating the third Poisson bracket is related to the computation of the derivative,

$$\frac{\partial}{\partial u'} \left[ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right] ,$$

can be differentiated term by term, with respect to  $u'$ , to obtain,

$$\begin{aligned} \cos(u' - v_{n-1} \pm \eta\sigma_{n-1}^z) + \cdots + \cos(u' - v_{n-1} \pm \eta\sigma_{n-1}^z) + \cdots + & \left( \cos(u' - v_{n-1} \pm \eta\sigma_{n-1}^z) \times \cdots \right. \\ & \left( \prod_{2 \leq j \leq m} \sin(u' - v_{n-1} \pm \eta\sigma_{n-1}^z) \right) + \cdots \\ & + \left( \prod_{1 \leq j \leq m-1} \sin(u' - v_{n-j} \pm \eta\sigma_{n-j}^z) \right) \\ & \left. \times \cos(u' - v_{n-m} \pm \eta\sigma_{n-m}^z) \right) , \end{aligned}$$

which we denote as  $\mathcal{D}(u') \equiv \mathcal{D}$ . This implies the following expression,

$$\mathcal{D} \frac{\partial \mathcal{P}_1}{\partial u'} + \frac{\partial \mathcal{P}_1}{\partial u} \mathcal{D}' ,$$

corresponding to the third bracket, for  $\mathcal{D}'(u') \equiv \mathcal{D}'$ , which equals,

$$\left[ \sum_{1 \leq i \leq m} \frac{\partial}{\partial u} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) \right] \frac{\partial B_3(u)}{\partial u'} + \left[ \sum_{1 \leq i \leq m} \frac{\partial}{\partial u'} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) \right] \frac{\partial B_3(u)}{\partial u} ,$$

while for the fourth bracket, a similar equality,

$$\left[ \sum_{1 \leq i \leq m} \frac{\partial}{\partial u} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) \right] \frac{\partial A_3(u)}{\partial u'} + \left[ \sum_{1 \leq i \leq m} \frac{\partial}{\partial u'} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) \right] \frac{\partial A_3(u)}{\partial u} .$$

holds. Altogether,

$$\begin{aligned} \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 (\sin(2\eta))^{n-3} \mathcal{A}_1, \mathcal{P}_2 \mathcal{A}'_3 \right\} \approx & \sum_{n':m+n'=n-3} (\sin(2\eta))^{n'-1} \left[ \sum_{1 \leq i \leq m} \frac{\partial}{\partial u} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) \right] \\ & \times \left[ \frac{\partial B_3(u)}{\partial u'} + \frac{\partial A_3(u)}{\partial u'} \right] \\ & + \sum_{n':m+n'=n-3} (\sin(2\eta))^{n'-1} \left[ \sum_{1 \leq i \leq m} \frac{\partial}{\partial u'} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) \right] \\ & \times \left[ \frac{\partial B_3(u)}{\partial u} + \frac{\partial A_3(u)}{\partial u} \right] \\ & - (\sin(2\eta))^{n-3} \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta\sigma_{n-i}^z) (\sin(2\eta))^{n'-1} \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \\ & \times \left[ \frac{B_3(u) A_3(u')}{u - u'} + \frac{A_3(u) B_3(u')}{u - u'} \right] , \end{aligned}$$

from which we conclude the argument.  $\square$

#### 2.4.4 Fourth Poisson bracket, $\mathcal{P}_4$

**Lemma 9** (*evaluating the fourth Poisson bracket in the first relation*). The fourth term,  $\mathcal{P}_4$ , approximately equals,

$$\begin{aligned} \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_2, \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} &\approx -[(\sin(2\eta))^{n-3} (\mathcal{C}_1)_i] \left[ \frac{A_3(u) B_3(u')}{u-u'} + \frac{B_3(u) A_3(u')}{u-u'} \right] \\ &+ \sum_{1 \leq i \leq n-3} \mathcal{A}_2 \left[ \left[ A_3(u) \frac{\partial B_3(u')}{\partial u'} \frac{\partial (\mathcal{C}_2)_i}{\partial u} + B_3(u) \frac{\partial B_3(u')}{\partial u} \frac{\partial (\mathcal{C}_2)_i}{\partial u'} \right] \right. \\ &\quad \left. + \left[ A_3(u) \frac{\partial A_3(u')}{\partial u'} \frac{\partial (\mathcal{C}_2)_i}{\partial u} + B_3(u) \frac{\partial A_3(u')}{\partial u} \frac{\partial (\mathcal{C}_2)_i}{\partial u'} \right] \right]. \end{aligned}$$

*Proof of Lemma 9.* The fourth term,

$$\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_2, \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\},$$

is equivalent to,

$$\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right), \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\}.$$

One application of Leibniz' rule to the Poisson bracket above gives,

$$\begin{aligned} \sum_{\mathcal{P}} \left[ \left\{ \mathcal{P}_1, \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma^z - n - i) \right) \right. \\ \left. + \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right), \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} \mathcal{P}_1 \right], \end{aligned}$$

while a second application of Liebniz' rule to the first Poisson bracket above gives,

$$-\left\{ \mathcal{P}_2, \mathcal{P}_1 \right\} (\sin(2\eta))^{n-3} \mathcal{A}'_1 - \left\{ (\sin(2\eta))^{n-3} \mathcal{A}'_1, \mathcal{P}_1 \right\} \mathcal{P}_2, \quad (*)$$

after anticommuting terms in the first Poisson bracket,

$$\left\{ \mathcal{P}_1, \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\},$$

with,

$$\left\{ \mathcal{P}_1, \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} = -\left\{ \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{A}'_1, \mathcal{P}_1 \right\}.$$

Applying Leibniz' rule to the second Poisson bracket in (\*) gives,

$$\mathcal{P}_2 \left( \left\{ \mathcal{A}'_1, \mathcal{P}_1 \right\} (\sin(2\eta))^{n-3} + \left\{ (\sin(2\eta))^{n-3}, \mathcal{P}_1 \right\} \mathcal{A}'_1 \right).$$

For the second Poisson bracket,

$$\left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right), \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} \mathcal{P}_1,$$

anticommuting terms gives,

$$-\left\{ \mathcal{P}_2(\sin(2\eta))^{n-3} \mathcal{A}'_1, \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} \mathcal{P}_1 ,$$

to which we apply Leibniz' rule,

$$\begin{aligned} & -\mathcal{P}_1 \left( \left\{ \mathcal{P}_2, \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} (\sin(2\eta))^{n-3} \mathcal{A}'_1 + \left\{ (\sin(2\eta))^{n-3} \right. \right. \\ & \quad \times \mathcal{A}'_1, \left. \left. \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} \mathcal{P}_2 \right) . \end{aligned}$$

Observe that the second Poisson bracket above is equal to,

$$\mathcal{P}_2(\sin(2\eta))^{n-3} \left\{ \mathcal{A}'_1, \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} .$$

Similarly, from (\*), from the second Poisson bracket,

$$\left\{ (\sin(2\eta))^{n-3} \mathcal{A}'_1, \mathcal{P}_1 \right\} \mathcal{P}_2 ,$$

to which we apply Leibniz' rule implies,

$$\mathcal{P}_2 \left( \left\{ \mathcal{A}'_1, \mathcal{P}_1 \right\} (\sin(2\eta))^{n-3} + \left\{ (\sin(2\eta))^{n-3}, \mathcal{P}_1 \right\} \mathcal{A}'_1 \right) .$$

Hence, the Poisson bracket that was rearranged corresponding to the fourth term is equivalent to,

$$\begin{aligned} & \sum_{\mathcal{P}} \left[ -\left\{ \mathcal{P}_2, \mathcal{P}_1 \right\} (\sin(2\eta))^{n-3} \mathcal{A}'_1 + \mathcal{P}_2 \left( \left\{ \mathcal{A}'_1, \mathcal{P}_1 \right\} (\sin(2\eta))^{n-3} + \left\{ (\sin(2\eta))^{n-3}, \mathcal{P}_1 \right\} \mathcal{A}'_1 \right) \right. \\ & \quad \left. - \left\{ \mathcal{P}_2, \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} (\sin(2\eta))^{n-3} \mathcal{A}'_1 \mathcal{P}_1 \right. \\ & \quad \left. - \mathcal{P}_1 \mathcal{P}_2 (\sin(2\eta))^{n-3} \left\{ \mathcal{A}'_1, \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} \right] . \end{aligned}$$

As a summation over  $\mathcal{P}$ , writing out each Poisson bracket from the superposition above gives,

$$\begin{aligned} -\sum_{\mathcal{P}} \left\{ \mathcal{P}_2, \mathcal{P}_1 \right\} (\sin(2\eta))^{n-3} \mathcal{A}'_1 & = -\left( \left\{ A_3(u), B_3(u') \right\} + \left\{ B_3(u), A_3(u') \right\} \right) (\sin(2\eta))^{n-3} \mathcal{A}'_1 , \\ \sum_{\mathcal{P}} \mathcal{P}_2 \left( \left\{ \mathcal{A}'_1, \mathcal{P}_1 \right\} (\sin(2\eta))^{n-3} + \left\{ (\sin(2\eta))^{n-3}, \mathcal{P}_1 \right\} \mathcal{A}'_1 \right) & \equiv 0 , \\ -\sum_{\mathcal{P}} \left\{ \mathcal{P}_2, \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} (\sin(2\eta))^{n-3} \mathcal{A}'_1 \mathcal{P}_1 & , \\ -\sum_{\mathcal{P}} \mathcal{P}_1 \mathcal{P}_2 (\sin(2\eta))^{n-3} \left\{ \mathcal{A}'_1, \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} & , \end{aligned}$$

where in the second Poisson bracket, we made use of the fact that,

$$\begin{aligned} \left\{ \mathcal{A}'_1, B_3(u) \right\} & = 0 , \\ \left\{ \mathcal{A}'_1, A_3(u) \right\} & = 0 , \\ \left\{ \mathcal{A}'_1, A_3(u') \right\} & = 0 , \\ \left\{ \mathcal{A}'_1, B_3(u') \right\} & = 0 , \end{aligned}$$

For the third Poisson bracket above, each term for all possible  $\mathcal{P}_2$  is,

$$\begin{aligned} & (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) (B_3(u)) \left\{ B_3(u'), \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} , \\ & (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) (B_3(u)) \left\{ A_3(u'), \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} , \\ & (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) (A_3(u)) \left\{ B_3(u'), \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} , \\ & (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) (A_3(u)) \left\{ A_3(u'), \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} , \end{aligned}$$

and, for the fourth Poisson bracket,

$$\left\{ \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+}, \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} = 0 .$$

Evaluating each bracket from the five listed above implies,

$$\begin{aligned} & B_3(u) \left[ \frac{\partial B_3(u')}{\partial u'} \left[ \sum_{1 \leq i \leq n-3} \left[ \frac{\partial}{\partial u} \left( \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) \right] \right. \\ & \left. + \frac{\partial B_3(u')}{\partial u} \left[ \sum_{1 \leq i \leq n-3} \left[ \frac{\partial}{\partial u'} \left( \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) \right] \right] , \end{aligned}$$

corresponding to the Poisson bracket between  $B_3(u')$  and the product of sine functions,

$$\begin{aligned} & B_3(u) \left[ \frac{\partial A_3(u')}{\partial u'} \left[ \sum_{1 \leq i \leq n-3} \left[ \left( \frac{\partial}{\partial u} \left( \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right) \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) \right] \right. \\ & \left. + \frac{\partial A_3(u')}{\partial u} \left[ \sum_{1 \leq i \leq n-3} \left[ \left( \frac{\partial}{\partial u'} \left( \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right) \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) \right] \right] \right] , \end{aligned}$$

corresponding to the Poisson bracket between  $A_3(u')$  and the product of sine functions,

$$\begin{aligned} & A_3(u) \left[ \left( \frac{\partial}{\partial u'} B_3(u') \right) \left[ \sum_{1 \leq i \leq n-3} \left[ \left( \frac{\partial}{\partial u} \left( \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right) \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) \right] \right] \right. \\ & \left. + \left( \frac{\partial}{\partial u} B_3(u') \right) \left[ \sum_{1 \leq i \leq n-3} \left[ \left( \frac{\partial}{\partial u'} \left( \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right) \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) \right] \right] \right] , \end{aligned}$$

corresponding to the Poisson bracket between  $B_3(u')$  and the product of sine functions, and,

$$\begin{aligned} & A_3(u) \left[ \left( \frac{\partial}{\partial u'} A_3(u') \right) \left[ \sum_{1 \leq i \leq n-3} \left[ \left( \frac{\partial}{\partial u} \left( \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right) \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) \right] \right] \right. \\ & \left. + \left( \frac{\partial}{\partial u} A_3(u') \right) \left[ \sum_{1 \leq i \leq n-3} \left[ \left( \frac{\partial}{\partial u'} \left( \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right) \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) \right] \right] \right] , \end{aligned}$$

corresponding to the Poisson bracket between  $A_3(u')$  and the product of sine functions. From each of the four brackets above, for differentiation with respect to  $u'$ , and then with respect to  $u$ , one can write,

$$\left( \frac{\partial}{\partial u'} B_3(u') \right) \left( \sum_{1 \leq i \leq n-3} \left[ \left( \frac{\partial}{\partial u} \left( \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right) \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) \right] \right) ,$$

as,

$$\begin{aligned}
& \left( \frac{\partial}{\partial u'} B_3(u') \right) \left( \frac{\partial}{\partial u} \left[ \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right] \right) \equiv \left( \frac{\partial}{\partial u'} B_3(u') \right) \left[ \cos(u - v_{n-1} + \eta \sigma_{n-1}^z) \right. \\
& \quad \times \prod_{2 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \\
& \quad + \prod_{2 \leq i \leq n-4} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \\
& \quad \left. \times \cos(u - v_{n-(n-3)} + \eta \sigma_{n-(n-3)}^z) \right] ,
\end{aligned}$$

can be further rearranged as,

$$\sum_{1 \leq i \leq n-3} \left[ \left( \frac{\partial}{\partial u'} B_3(u') \right) \left( \frac{\partial}{\partial u} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) \right] ,$$

after distributing the derivative of  $B_3(u')$  to each term in the summation, from,

$$\begin{aligned}
& \left( \frac{\partial}{\partial u'} B_3(u') \right) \left( \frac{\partial}{\partial u} \sin(u - v_{n-1} + \eta \sigma_{n-1}^z) \right) \prod_{2 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \\
& + \left( \frac{\partial}{\partial u'} B_3(u') \right) \prod_{2 \leq i \leq n-4} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \\
& \quad \times \left( \frac{\partial}{\partial u'} \sin(u - v_{n-(n-3)} + \eta \sigma_{n-(n-3)}^z) \right) .
\end{aligned}$$

As a result, the remaining terms from the Poisson bracket,

$$\left( \frac{\partial}{\partial u} B_3(u') \right) \sum_{1 \leq i \leq n-3} \left[ \left[ \frac{\partial}{\partial u'} \left( \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) \right] ,$$

can be combined under the same summation to form,

$$\begin{aligned}
& \sum_{1 \leq i \leq n-3} \left[ \left[ \left( \frac{\partial}{\partial u'} B_3(u') \right) \left( \frac{\partial}{\partial u} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) + \left( \frac{\partial}{\partial u} B_3(u') \right) \left( \frac{\partial}{\partial u'} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] \right. \\
& \quad \left. \times \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) \right] .
\end{aligned}$$

The observation that the derivative of the product of sine functions,

$$\frac{\partial}{\partial u} \left[ \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right] ,$$

equals,

$$\begin{aligned}
& \cos(u - v_{n-1} + \eta \sigma_{n-1}^z) \prod_{2 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) + \cdots + \prod_{2 \leq i \leq n-4} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \\
& \quad \times \cos(u - v_{n-(n-3)} + \eta \sigma_{n-(n-3)}^z) .
\end{aligned}$$

comes from the fact that the sum of products above can be expressed as,

$$\sum_{1 \leq i \leq n-3} \left[ \left[ \frac{\partial}{\partial u} \left( \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) \right] .$$

For the three remaining Poisson brackets, along similar lines,

$$\sum_{1 \leq i \leq n-3} \left[ \left[ \left( \frac{\partial}{\partial u'} A_3(u') \right) \left( \frac{\partial}{\partial u} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) + \left( \frac{\partial}{\partial u} A_3(u') \right) \left( \frac{\partial}{\partial u'} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] \right. \\ \times \left. \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) \right] ,$$

for the first Poisson bracket,

$$A_3(u) \left[ \left( \frac{\partial}{\partial u'} B_3(u') \right) \left( \sum_{1 \leq i \leq n-3} \left[ \left( \frac{\partial}{\partial u} \left( \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right) \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) \right] \right) \right. \\ \left. + \left( \frac{\partial}{\partial u} B_3(u') \right) \sum_{1 \leq i \leq n-3} \left[ \left( \frac{\partial}{\partial u'} \left( \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right) \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) \right] \right] ,$$

for the second Poisson bracket, and,

$$\sum_{1 \leq i \leq n-3} \left[ \left[ \left( \frac{\partial}{\partial u'} A_3(u') \right) \left( \frac{\partial}{\partial u} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) + \left( \frac{\partial}{\partial u} A_3(u') \right) \left( \frac{\partial}{\partial u'} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] \right. \\ \times \left. \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) \right] ,$$

for the third Poisson bracket, in which the derivatives of  $A_3$ , or  $B_3$ , can be combined with the summation of the  $i$  th derivative of the sine functions and the remaining product of sine functions. Hence, the terms from each of the four Poisson brackets are equal to,

$$B_3(u) \sum_{1 \leq i \leq n-3} \left[ \left[ \left( \frac{\partial}{\partial u'} B_3(u') \right) \left( \frac{\partial}{\partial u} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) + \left( \frac{\partial}{\partial u} B_3(u') \right) \left( \frac{\partial}{\partial u'} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] \right. \\ \times \left. \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) \right] + B_3(u) \\ \times \sum_{1 \leq i \leq n-3} \left[ \left[ \left( \frac{\partial}{\partial u'} A_3(u') \right) \left( \frac{\partial}{\partial u} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) + \left( \frac{\partial}{\partial u} A_3(u') \right) \left( \frac{\partial}{\partial u'} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] \right. \\ \times \left. \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) \right] + A_3(u) \\ \times \sum_{1 \leq i \leq n-3} \left[ \left[ \left( \frac{\partial}{\partial u'} B_3(u') \right) \left( \frac{\partial}{\partial u} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) + \left( \frac{\partial}{\partial u} B_3(u') \right) \left( \frac{\partial}{\partial u'} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] \right. \\ \times \left. \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) \right] + A_3(u) \\ \times \sum_{1 \leq i \leq n-3} \left[ \left[ \left( \frac{\partial}{\partial u'} A_3(u') \right) \left( \frac{\partial}{\partial u} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) + \left( \frac{\partial}{\partial u} A_3(u') \right) \left( \frac{\partial}{\partial u'} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] \right. \\ \times \left. \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) \right] ,$$

with prefactor,

$$(\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) .$$

Excluding the prefactor above, the four Poisson brackets can be combined under the single summation, yielding,

$$\begin{aligned}
& \sum_{1 \leq i \leq n-3} \left[ B_3(u) \left[ \left( \frac{\partial}{\partial u'} B_3(u') \right) \left( \frac{\partial}{\partial u} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) + \left( \frac{\partial}{\partial u} B_3(u') \right) \left( \frac{\partial}{\partial u'} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] \right. \\
& \quad \times \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) + B_3(u) \\
& \quad \times \left[ \left( \frac{\partial}{\partial u'} A_3(u') \right) \left( \frac{\partial}{\partial u} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) + \left( \frac{\partial}{\partial u} A_3(u') \right) \left( \frac{\partial}{\partial u'} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] \\
& \quad \times \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) + A_3(u) \\
& \quad \times \left[ \left( \frac{\partial}{\partial u'} B_3(u') \right) \left( \frac{\partial}{\partial u} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) + \left( \frac{\partial}{\partial u} B_3(u') \right) \left( \frac{\partial}{\partial u'} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] \\
& \quad \times \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) + A_3(u) \\
& \quad \times \left[ \left( \frac{\partial}{\partial u'} A_3(u') \right) \left( \frac{\partial}{\partial u} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) + \left( \frac{\partial}{\partial u} A_3(u') \right) \left( \frac{\partial}{\partial u'} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] \\
& \quad \times \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta \sigma_{n-j}^z) \right) , 
\end{aligned}$$

which, from the expressions introduced in **Lemma 4**, can be expressed as,

$$\begin{aligned}
& \sum_{1 \leq i \leq n-3} \mathcal{A}_2 \left[ B_3(u) \left[ \left( \frac{\partial}{\partial u'} B_3(u') \right) \left( \frac{\partial}{\partial u} (\mathcal{C}_2)_i \right) + \left( \frac{\partial}{\partial u} B_3(u') \right) \left( \frac{\partial}{\partial u'} (\mathcal{C}_2)_i \right) \right] \right. \\
& \quad + B_3(u) \\
& \quad \times \left[ \left( \frac{\partial}{\partial u'} A_3(u') \right) \left( \frac{\partial}{\partial u} (\mathcal{C}_2)_i \right) + \left( \frac{\partial}{\partial u} A_3(u') \right) \left( \frac{\partial}{\partial u'} (\mathcal{C}_2)_i \right) \right] \\
& \quad + A_3(u) \\
& \quad \times \left[ \left( \frac{\partial}{\partial u'} B_3(u') \right) \left( \frac{\partial}{\partial u} (\mathcal{C}_2)_i \right) + \left( \frac{\partial}{\partial u} B_3(u') \right) \left( \frac{\partial}{\partial u'} (\mathcal{C}_2)_i \right) \right] \\
& \quad + A_3(u) \\
& \quad \times \left[ \left( \frac{\partial}{\partial u'} A_3(u') \right) \left( \frac{\partial}{\partial u} (\mathcal{C}_2)_i \right) + \left( \frac{\partial}{\partial u} A_3(u') \right) \left( \frac{\partial}{\partial u'} (\mathcal{C}_2)_i \right) \right] . 
\end{aligned}$$

Grouping together like terms from the above summation implies,

$$\begin{aligned}
& \sum_{1 \leq i \leq n-3} \mathcal{A}_2 \left[ \left[ A_3(u) \left( \frac{\partial}{\partial u'} B_3(u') \right) \left( \frac{\partial}{\partial u} (\mathcal{C}_2)_i \right) + B_3(u) \left( \frac{\partial}{\partial u} B_3(u') \right) \left( \frac{\partial}{\partial u'} (\mathcal{C}_2)_i \right) \right] \right. \\
& \quad + \left[ A_3(u) \left( \frac{\partial}{\partial u'} A_3(u') \right) \left( \frac{\partial}{\partial u} (\mathcal{C}_2)_i \right) + B_3(u) \left( \frac{\partial}{\partial u} A_3(u') \right) \left( \frac{\partial}{\partial u'} (\mathcal{C}_2)_i \right) \right] \left. \right] . 
\end{aligned}$$

Altogether, the terms from all nonzero Poisson brackets,

$$\begin{aligned}
& - \left( \{ A_3(u), B_3(u') \} + \{ B_3(u), A_3(u') \} \right) (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \\
& + \sum_{1 \leq i \leq n-3} \mathcal{A}_2 \left[ \left[ A_3(u) \left( \frac{\partial}{\partial u'} B_3(u') \right) \left( \frac{\partial}{\partial u} (\mathcal{C}_2)_i \right) + B_3(u) \left( \frac{\partial}{\partial u} B_3(u') \right) \left( \frac{\partial}{\partial u'} (\mathcal{C}_2)_i \right) \right] \right. \\
& \quad + \left[ A_3(u) \left( \frac{\partial}{\partial u'} A_3(u') \right) \left( \frac{\partial}{\partial u} (\mathcal{C}_2)_i \right) + B_3(u) \left( \frac{\partial}{\partial u} A_3(u') \right) \left( \frac{\partial}{\partial u'} (\mathcal{C}_2)_i \right) \right] \left. \right] , 
\end{aligned}$$

Furthermore, applying expressions for the Poisson brackets between  $A_3(u)$ ,  $B_3(u')$ , and for the Poisson bracket between  $B_3(u)$ ,  $A_3(u')$ , from previous terms in the first relation, approximately yields,

$$\begin{aligned} \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_2, \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} &\approx - \left[ (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \right] \left[ \frac{A_3(u) B_3(u')}{u - u'} + \frac{B_3(u) A_3(u')}{u - u'} \right] \\ &+ \sum_{1 \leq i \leq n-3} \mathcal{A}_2 \left[ \left[ A_3(u) \left( \frac{\partial}{\partial u'} B_3(u') \right) \left( \frac{\partial}{\partial u} (\mathcal{C}_2)_i \right) + B_3(u) \left( \frac{\partial}{\partial u} B_3(u') \right) \left( \frac{\partial}{\partial u'} (\mathcal{C}_2)_i \right) \right] \right. \\ &\quad \left. + \left[ A_3(u) \left( \frac{\partial}{\partial u} A_3(u') \right) \left( \frac{\partial}{\partial u} (\mathcal{C}_2)_i \right) + B_3(u) \left( \frac{\partial}{\partial u} A_3(u') \right) \left( \frac{\partial}{\partial u'} (\mathcal{C}_2)_i \right) \right] \right], \end{aligned}$$

from which we conclude the argument.  $\square$

#### 2.4.5 Fifth Poisson bracket, $\mathcal{P}_5$

**Lemma 10** (*evaluating the fifth Poisson bracket in the first relation*). The fifth term,  $\mathcal{P}_5$ ,

$$\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_2, \mathcal{P}_2 \mathcal{A}'_2 \right\},$$

approximately equals,

$$\begin{aligned} -\mathcal{C}_2 \left[ \frac{B_3(u') A_3(u)}{u' - u} + \frac{B_3(u) A_3(u')}{u - u'} \right] - (B_3(u) + A_3(u)) \mathcal{C}_2 \left[ \sum_{1 \leq i \leq n-3} \left[ \frac{\partial B_3(u')}{\partial u'} \frac{\partial (\mathcal{C}_2)_i}{\partial u} - \frac{\partial B_3(u')}{\partial u} \frac{\partial (\mathcal{C}_2)_i}{\partial u'} \right] \right] \\ \times \left[ \prod_{1 \leq j \neq i \leq n-3} (\mathcal{C}_2)_j \right] - (B_3(u) + A_3(u)) \left[ \prod_{1 \leq i \leq n-3} (\mathcal{C}_2)_i \right] \\ \times \left[ \sum_{1 \leq i \leq n-3} \left[ \frac{\partial A_3(u')}{\partial u'} \frac{\partial (\mathcal{C}_2)_i}{\partial u} - \frac{\partial A_3(u')}{\partial u} \frac{\partial (\mathcal{C}_2)_i}{\partial u'} \right] \right] \left[ \prod_{1 \leq j \neq i \leq n-3} (\mathcal{C}_2)_j \right]. \end{aligned}$$

*Proof of Lemma 10.* The fifth term,

$$\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_2, \mathcal{P}_2 \mathcal{A}'_2 \right\},$$

is equivalent to,

$$\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right), \mathcal{P}_2 \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\},$$

which can be rearranged by applying Leibniz' rule to the first argument, in which,

$$\begin{aligned} &\sum_{\mathcal{P}} \left( \left\{ \mathcal{P}_1, \mathcal{P}_2 \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right. \\ &\quad \left. + \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right), \mathcal{P}_2 \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} \mathcal{P}_1 \right), \end{aligned}$$

to which a second application of Leibniz' rule yields, for the second argument,

$$\begin{aligned}
& \sum_{\mathcal{P}} \left[ \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \left[ -\{\mathcal{P}_2, \mathcal{P}_1\} \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right. \right. \\
& \quad \left. \left. - \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), \mathcal{P}_1 \right\} \mathcal{P}_2 \right] \right. \\
& \quad \left. - \mathcal{P}_1 \left[ \left\{ \mathcal{P}_2, \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right. \right. \\
& \quad \left. \left. + \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} \mathcal{P}_2 \right] \right] ,
\end{aligned}$$

after anticommuting the first Poisson bracket,

$$\left\{ \mathcal{P}_1, \mathcal{P}_2 \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} ,$$

and the second Poisson bracket,

$$\left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right), \mathcal{P}_2 \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} .$$

Writing out each Poisson bracket gives,

$$\begin{aligned}
& -\sum_{\mathcal{P}} \{\mathcal{P}_2, \mathcal{P}_1\} \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) = -\left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \left( \{B_3(u'), A_3(u)\} \right. \\
& \quad \left. + \{B_3(u), A_3(u')\} \right) , \\
& -\sum_{\mathcal{P}} \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), \mathcal{P}_1 \right\} \mathcal{P}_2 = -\mathcal{P}_2 \left( \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), B_3(u) \right\} \right. \\
& \quad \left. + \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), A_3(u) \right\} \right) ,
\end{aligned}$$

for the first two terms, while the third term,

$$-\mathcal{P}_1 \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \sum_{\mathcal{P}} \left\{ \mathcal{P}_2, \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\}$$

is equivalent to,

$$\begin{aligned}
& \mathcal{P}_1 \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \left[ \left\{ B_3(u'), \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} \right. \\
& \quad \left. + \left\{ A_3(u'), \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} \right] .
\end{aligned}$$

The fourth term is,

$$\mathcal{P}_2 \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} .$$

For the first Poisson bracket, one approximately has,

$$-\left[ \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta\sigma_{n-i}^z) \right] \left[ \frac{B_3(u') A_3(u)}{u' - u} + \frac{B_3(u) A_3(u')}{u - u'} \right] .$$

After taking the summation of each bracket over  $\mathcal{P}$ , observe, from further rearrangement of the third Poisson bracket,

$$-\mathcal{P}_1 \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta\sigma_{n-i}^z) \right) \sum_{\mathcal{P}} \left\{ \mathcal{P}_2, \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta\sigma_{n-i}^z) \right) \right\} ,$$

is equivalent to the terms,

$$\begin{aligned} & -B_3(u) \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta\sigma_{n-i}^z) \right) \left\{ B_3(u'), \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta\sigma_{n-i}^z) \right) \right\} , \\ & -A_3(u) \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta\sigma_{n-i}^z) \right) \left\{ A_3(u'), \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta\sigma_{n-i}^z) \right) \right\} , \\ & -A_3(u) \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta\sigma_{n-i}^z) \right) \left\{ B_3(u'), \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta\sigma_{n-i}^z) \right) \right\} , \\ & -B_3(u) \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta\sigma_{n-i}^z) \right) \left\{ A_3(u'), \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta\sigma_{n-i}^z) \right) \right\} . \end{aligned}$$

Term by term, evaluating the bracket for,

$$\left( \frac{\partial}{\partial u'} B_3(u') \right) \left[ \frac{\partial}{\partial u} \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta\sigma_{n-i}^z) \right) \right] - \left( \frac{\partial}{\partial u} B_3(u) \right) \left[ \frac{\partial}{\partial u'} \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta\sigma_{n-i}^z) \right) \right] ,$$

with prefactor,

$$-B_3(u) \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta\sigma_{n-i}^z) \right) ,$$

corresponding to the first bracket, can be expressed with,

$$\begin{aligned} & \left( \frac{\partial}{\partial u'} B_3(u') \right) \left[ \sum_{1 \leq i \leq n-3} \left( \frac{\partial}{\partial u} \left( \sin(u - v_{n-i} + \eta\sigma_{n-i}^z) \right) \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta\sigma_{n-j}^z) \right) \right] \\ & - \left( \frac{\partial}{\partial u} B_3(u) \right) \left[ \sum_{1 \leq i \leq n-3} \left( \frac{\partial}{\partial u'} \left( \sin(u' - v_{n-i} + \eta\sigma_{n-i}^z) \right) \prod_{1 \leq j \neq i \leq n-3} \sin(u' - v_{n-j} + \eta\sigma_{n-j}^z) \right) \right] , \end{aligned}$$

from the fact that the derivative of the product of sine functions equals,

$$\begin{aligned} \frac{\partial}{\partial u} \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta\sigma_{n-i}^z) \right) &= \frac{\partial}{\partial u} \left( \sin(u - v_{n-1} + \eta\sigma_{n-1}^z) \cdots \times \sin(u - v_{n-(n-3)} + \eta\sigma_{n-(n-3)}^z) \right) \\ &= \sum_{1 \leq i \leq n-3} \left( \frac{\partial}{\partial u} \left( \sin(u - v_{n-i} + \eta\sigma_{n-i}^z) \right) \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta\sigma_{n-j}^z) \right) . \end{aligned}$$

Evaluating the bracket for,

$$\left\{ A_3(u'), \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta\sigma_{n-i}^z) \right) \right\} ,$$

with prefactor,

$$-A_3(u) \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta\sigma_{n-i}^z) \right) ,$$

corresponding to the second bracket, can similarly be expressed with,

$$\begin{aligned} & \left( \frac{\partial}{\partial u'} A_3(u') \right) \left[ \sum_{1 \leq i \leq n-3} \left( \frac{\partial}{\partial u} \left( \sin(u - v_{n-i} + \eta\sigma_{n-i}^z) \right) \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta\sigma_{n-j}^z) \right) \right] \\ & - \left( \frac{\partial}{\partial u'} A_3(u') \right) \left[ \sum_{1 \leq i \leq n-3} \left( \frac{\partial}{\partial u} \left( \sin(u - v_{n-i} + \eta\sigma_{n-i}^z) \right) \prod_{1 \leq j \neq i \leq n-3} \sin(u' - v_{n-j} + \eta\sigma_{n-j}^z) \right) \right] , \end{aligned}$$

with prefactor,

$$-A_3(u) \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta\sigma_{n-i}^z) \right) ,$$

corresponding to the third bracket. Along similar lines, the third and fourth brackets can be respectively expressed with,

$$\begin{aligned} & \left( \frac{\partial}{\partial u'} A_3(u') \right) \left[ \sum_{1 \leq i \leq n-3} \left( \frac{\partial}{\partial u} \left( \sin(u - v_{n-i} + \eta\sigma_{n-i}^z) \right) \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta\sigma_{n-j}^z) \right) \right] \\ & - \left( \frac{\partial}{\partial u'} A_3(u') \right) \left[ \sum_{1 \leq i \leq n-3} \left( \frac{\partial}{\partial u} \left( \sin(u - v_{n-i} + \eta\sigma_{n-i}^z) \right) \prod_{1 \leq j \neq i \leq n-3} \sin(u' - v_{n-j} + \eta\sigma_{n-j}^z) \right) \right] , \end{aligned}$$

and with,

$$\begin{aligned} & \left( \frac{\partial}{\partial u'} A_3(u') \right) \left[ \sum_{1 \leq i \leq n-3} \left( \frac{\partial}{\partial u} \left( \sin(u - v_{n-i} + \eta\sigma_{n-i}^z) \right) \prod_{1 \leq j \neq i \leq n-3} \sin(u - v_{n-j} + \eta\sigma_{n-j}^z) \right) \right] \\ & - \left( \frac{\partial}{\partial u} A_3(u') \right) \left[ \sum_{1 \leq i \leq n-3} \left( \frac{\partial}{\partial u'} \left( \sin(u' - v_{n-i} + \eta\sigma_{n-i}^z) \right) \prod_{1 \leq j \neq i \leq n-3} \sin(u' - v_{n-j} + \eta\sigma_{n-j}^z) \right) \right] , \end{aligned}$$

with prefactors,

$$-A_3(u) \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta\sigma_{n-i}^z) \right) ,$$

and,

$$-B_3(u) \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta\sigma_{n-i}^z) \right) .$$

Grouping together like terms, from the each bracket, from the four above, can be expressed as, either,

$$\begin{aligned} & \left( \sum_{1 \leq i \leq n-3} \left( \frac{\partial B_3(u')}{\partial u'} \left( \frac{\partial}{\partial u} \sin(u - v_{n-i} + \eta\sigma_{n-i}^z) \right) - \frac{\partial B_3(u')}{\partial u} \left( \frac{\partial}{\partial u} \sin(u' - v_{n-i} + \eta\sigma_{n-i}^z) \right) \right) \right. \\ & \quad \times \left. \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u' - v_{n-j} + \eta\sigma_{n-j}^z) \right) \right) , \end{aligned}$$

or as,

$$\left( \sum_{1 \leq i \leq n-3} \left( \frac{\partial A_3(u')}{\partial u'} \left( \frac{\partial}{\partial u} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) - \frac{\partial A_3(u)}{\partial u} \left( \frac{\partial}{\partial u} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right) \right. \\ \left. \times \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u' - v_{n-j} + \eta \sigma_{n-j}^z) \right) \right) ,$$

from which approximating,

$$\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_2, \mathcal{P}_2 \mathcal{A}'_2 \right\} ,$$

yields the desired expression,

$$-\left[ \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right] \left[ \frac{B_3(u') A_3(u)}{u' - u} + \frac{B_3(u) A_3(u')}{u - u'} \right] \\ - (B_3(u) + A_3(u)) \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \\ \times \left( \sum_{1 \leq i \leq n-3} \left( \frac{\partial B_3(u')}{\partial u'} \left( \frac{\partial}{\partial u} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) - \frac{\partial B_3(u')}{\partial u} \left( \frac{\partial}{\partial u} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right) \right. \\ \left. \times \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u' - v_{n-j} + \eta \sigma_{n-j}^z) \right) \right. \\ \left. - (B_3(u) + A_3(u)) \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right) \\ \times \left( \sum_{1 \leq i \leq n-3} \left( \frac{\partial A_3(u')}{\partial u'} \left( \frac{\partial}{\partial u} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) - \frac{\partial A_3(u')}{\partial u} \left( \frac{\partial}{\partial u} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right) \right. \\ \left. \times \left( \prod_{1 \leq j \neq i \leq n-3} \sin(u' - v_{n-j} + \eta \sigma_{n-j}^z) \right) \right) ,$$

from which we conclude the argument.  $\square$

#### 2.4.6 Sixth Poisson bracket, $\mathcal{P}_6$

**Lemma 11** (*evaluating the sixth Poisson bracket in the first relation*). The sixth term,  $\mathcal{P}_6$ , approximately equals,

$$\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_2, \mathcal{P}_2 \mathcal{A}'_3 \right\} \approx \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} (\mathcal{C}_1)_j \right) \right] \right] \left[ \frac{A_3(u) B_3(u')}{u - u'} \right] \\ + \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} (\mathcal{C}_1)_j \right) \right] \right] \left[ \frac{B_3(u) A_3(u')}{u - u'} \right] \\ - 2(B_3(u) + A_3(u)) \left[ \sum_{1 \leq j \leq n'} (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} (\mathcal{C}_1)_j \right) \right] \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \frac{\partial (\mathcal{C}_2)_i}{\partial u'} \right) \right. \right. \\ \left. \left. \times \left( \prod_{1 \leq j \neq i \leq m} (\mathcal{C}_2)_j \right) \left( \frac{\partial A_3(u)}{\partial u} + \frac{\partial B_3(u)}{\partial u} \right) + \left( \frac{\partial (\mathcal{C}_2)_i}{\partial u} \right) \left( \prod_{1 \leq j \neq i \leq m} (\mathcal{C}_2)_j \right) \left( \frac{\partial A_3(u')}{\partial u'} + \frac{\partial B_3(u')}{\partial u'} \right) \right] \right] .$$

*Proof of Lemma 11.* The sixth term,

$$\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_2, \mathcal{P}_2 \mathcal{A}'_3 \right\} ,$$

is equivalent to,

$$\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right), \mathcal{P}_2 \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \times \cdots \right. \right. \right. \right. \\ \left. \left. \left. \left. \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right] \right\} ,$$

is equivalent to the superposition of terms,

$$\sum_{\mathcal{P}} \left\{ \mathcal{P}_1, \mathcal{P}_2 \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \right. \right. \right. \\ \left. \left. \left. \times \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right] \right\} \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right)$$

and,

$$\sum_{\mathcal{P}} \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right), \mathcal{P}_2 \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \right. \right. \right. \\ \left. \left. \left. \times \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right] \right\} \mathcal{P}_1 ,$$

from an application of Leibniz' rule. Applying Leibniz' rule a second time to each bracket in the expression above gives,

$$-\sum_{\mathcal{P}} \left[ \left\{ \mathcal{P}_2, \mathcal{P}_1 \right\} \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \right. \right. \\ \left. \left. \times \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right] \\ + \left\{ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right], \mathcal{P}_1 \right\} \mathcal{P}_2 \\ - \sum_{\mathcal{P}} \left[ \left\{ \mathcal{P}_2, \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \right. \right. \right. \\ \left. \left. \left. \times \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right] \right] ,$$

after anticommuting terms in,

$$\left\{ \mathcal{P}_1, \mathcal{P}_2 \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right] \right\} ,$$

and also after performing the same anticommutation in,

$$\left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u - v_{n-i} + \eta \sigma_{n-i}^z) \right), \mathcal{P}_2 \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \times \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right] \right\} .$$

Writing out each bracket individually gives,

$$\sum_{\mathcal{P}} \{ \mathcal{P}_1, \mathcal{P}_1 \} = \{ A_3(u), B_3(u') \} + \{ B_3(u), A_3(u') \} ,$$

for the first term, which with the corresponding prefactor, equals,

$$\left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) \left( \{ A_3(u), B_3(u') \} + \{ B_3(u), A_3(u') \} \right) ,$$

while for the remaining brackets,

$$-\mathcal{P}_2 \sum_{\mathcal{P}} \left\{ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right], \mathcal{P}_1 \right\}$$

equals,

$$-\mathcal{P}_2 \sum_{\mathcal{P}} \left\{ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), \mathcal{P}_1 \right\} \sum_{1 \leq j \leq n'} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) .$$

Altogether, one has the remaining brackets,

$$\begin{aligned} & \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) \left( \{ A_3(u), B_3(u') \} + \{ B_3(u), A_3(u') \} \right) \\ & - \mathcal{P}_2 \sum_{\mathcal{P}} \left\{ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), \mathcal{P}_1 \right\} \\ & \quad \times \sum_{1 \leq j \leq n'} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) . \end{aligned}$$

The second Poisson bracket is equivalent to,

$$\left\{ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), B_3(u) \right\} + \left\{ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), A_3(u) \right\} ,$$

with prefactor,

$$-\mathcal{P}_2 \sum_{1 \leq j \leq n'} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) .$$

The two brackets, from the summation over  $\mathcal{P}_2$ , can each analyzed from the following six derivatives,

$$\begin{aligned} & \frac{\partial}{\partial u'} \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right] , \\ & \frac{\partial}{\partial u} \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right] , \\ & \frac{\partial}{\partial u} [B_3(u)] , \\ & \frac{\partial}{\partial u} [A_3(u)] , \\ & \frac{\partial}{\partial u'} [A_3(u')] , \\ & \frac{\partial}{\partial u'} [B_3(u')] , \end{aligned}$$

corresponding to the first term. Observe, for the first two derivatives with respective to  $u'$ , and to  $u$ , above, that,

$$\begin{aligned} & \frac{\partial}{\partial u'} \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right] = \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \frac{\partial}{\partial u'} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right] \\ & = \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \left( \frac{\partial}{\partial u'} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \prod_{1 \leq j \neq i \leq m} \sin(u' - v_{n-j} \pm \eta \sigma_{n-j}^z) \right) , \end{aligned}$$

from the fact that differentiating the product with respect to  $u'$ ,

$$\prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) = \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \cdots \times \sin(u' - v_{n-m} \pm \eta \sigma_{n-m}^z) ,$$

gives,

$$\begin{aligned} & \frac{\partial}{\partial u'} \left[ \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \cdots \times \sin(u' - v_{n-m} \pm \eta \sigma_{n-m}^z) \right] = \cos(u' - v_{n-1} \pm \eta \sigma_{n-1}^z) \cdots \times \\ & \quad \prod_{2 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) + \cdots + \prod_{1 \leq i \leq m-1} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \cdots \times \\ & \quad \cos(u' - v_{n-m} \pm \eta_{n-m}^z) , \end{aligned}$$

which can be expressed with the summation,

$$\sum_{1 \leq i \leq m} \left[ \left( \frac{\partial}{\partial u'} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \left( \prod_{1 \leq j \neq i \leq m} \sin(u' - v_{n-j} \pm \eta \sigma_{n-j}^z) \right) \right] .$$

For the other term,

$$\frac{\partial}{\partial u} \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right] ,$$

one similarly obtains,

$$\sum_{1 \leq i \leq m} \left[ \left( \frac{\partial}{\partial u} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \left( \prod_{1 \leq j \neq i \leq m} \sin(u' - v_{n-j} \pm \eta \sigma_{n-j}^z) \right) \right] ,$$

corresponding to the second term. Altogether, the Poisson bracket for each term takes the form,

$$\begin{aligned} -\mathcal{P}_2 \sum_{1 \leq j \leq n'} & \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) \left[ \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \left( \frac{\partial}{\partial u'} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \right. \right. \\ & \times \left. \left. \left. \prod_{1 \leq j \neq i \leq m} \sin(u' - v_{n-j} \pm \eta \sigma_{n-j}^z) \right) \right] \frac{\partial B_3(u)}{\partial u} \right. \\ & + \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \left( \frac{\partial}{\partial u} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \right. \\ & \times \left. \left. \prod_{1 \leq j \neq i \leq m} \sin(u' - v_{n-j} \pm \eta \sigma_{n-j}^z) \right) \right] \frac{\partial B_3(u')}{\partial u'} \Bigg] , \end{aligned}$$

corresponding to the first Poisson bracket, and,

$$\begin{aligned} -\mathcal{P}_2 \sum_{1 \leq j \leq n'} & \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) \left[ \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \left( \frac{\partial}{\partial u'} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \right. \right. \\ & \times \left. \left. \left. \prod_{1 \leq j \neq i \leq m} \sin(u' - v_{n-j} \pm \eta \sigma_{n-j}^z) \right) \right] \frac{\partial A_3(u)}{\partial u} \right. \\ & + \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \left( \frac{\partial}{\partial u} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \right. \\ & \times \left. \left. \prod_{1 \leq j \neq i \leq m} \sin(u' - v_{n-j} \pm \eta \sigma_{n-j}^z) \right) \right] \frac{\partial A_3(u')}{\partial u'} \Bigg] , \end{aligned}$$

corresponding to the second Poisson bracket. Further rearranging the derivatives from each of the two brackets above implies, for the first bracket,

$$\begin{aligned}
& -\mathcal{P}_2 \sum_{1 \leq j \leq n'} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \left( \frac{\partial}{\partial u'} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \right. \\
& \quad \times \left( \prod_{1 \leq j \neq i \leq m} \sin(u' - v_{n-j} \pm \eta \sigma_{n-j}^z) \right) \frac{\partial B_3(u)}{\partial u} \\
& \quad + \left( \frac{\partial}{\partial u} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \\
& \quad \left. \left. \times \left( \prod_{1 \leq j \neq i \leq m} \sin(u' - v_{n-j} \pm \eta \sigma_{n-j}^z) \right) \frac{\partial B_3(u')}{\partial u'} \right) \right] ,
\end{aligned}$$

and also, for the second bracket,

$$\begin{aligned}
& -\mathcal{P}_2 \sum_{1 \leq j \leq n'} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \left( \frac{\partial}{\partial u'} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \right. \\
& \quad \times \left( \prod_{1 \leq j \neq i \leq m} \sin(u' - v_{n-j} \pm \eta \sigma_{n-j}^z) \right) \frac{\partial A_3(u)}{\partial u} \\
& \quad + \left( \frac{\partial}{\partial u} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \\
& \quad \left. \left. \times \left( \prod_{1 \leq j \neq i \leq m} \sin(u' - v_{n-j} \pm \eta \sigma_{n-j}^z) \right) \frac{\partial A_3(u')}{\partial u'} \right) \right] .
\end{aligned}$$

Grouping together like terms from the two brackets above yields,

$$\begin{aligned}
& -\mathcal{P}_2 \sum_{1 \leq j \leq n'} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \left( \frac{\partial}{\partial u'} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \right. \\
& \quad \times \left( \prod_{1 \leq j \neq i \leq m} \sin(u' - v_{n-j} \pm \eta \sigma_{n-j}^z) \right) \left( \frac{\partial A_3(u)}{\partial u} + \frac{\partial B_3(u)}{\partial u} \right) \\
& \quad + \left( \frac{\partial}{\partial u} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \\
& \quad \left. \left. \times \left( \prod_{1 \leq j \neq i \leq m} \sin(u' - v_{n-j} \pm \eta \sigma_{n-j}^z) \right) \left( \frac{\partial A_3(u')}{\partial u'} + \frac{\partial B_3(u')}{\partial u'} \right) \right) \right] .
\end{aligned}$$

For the two remaining Poisson brackets appearing before the second bracket, namely,

$$\left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) \{A_3(u), B_3(u')\} ,$$

and,

$$\left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) \{B_3(u), A_3(u')\} ,$$

one has, approximately,

$$\left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right] \left[ \frac{A_3(u) B_3(u')}{u - u'} \right] ,$$

corresponding to the first term, and, approximately,

$$\left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right] \left[ \frac{B_3(u) A_3(u')}{u - u'} \right] ,$$

corresponding to the second term. Hence,

$$\begin{aligned} \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_2, \mathcal{P}_2 \mathcal{A}'_3 \right\} &\approx \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right] \\ &\quad \times \left[ \frac{A_3(u) B_3(u')}{u - u'} \right] \\ &+ \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right] \\ &\quad \times \left[ \frac{B_3(u) A_3(u')}{u - u'} \right] \\ -2(B_3(u) + A_3(u)) \sum_{1 \leq j \leq n'} &\left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \left( \frac{\partial}{\partial u'} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \right. \\ &\quad \times \left( \prod_{1 \leq j \neq i \leq m} \sin(u' - v_{n-j} \pm \eta \sigma_{n-j}^z) \right) \left( \frac{\partial A_3(u)}{\partial u} + \frac{\partial B_3(u)}{\partial u} \right) \\ &\quad \left. \left. + \left( \frac{\partial}{\partial u} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right) \right. \\ &\quad \times \left( \prod_{1 \leq j \neq i \leq m} \sin(u' - v_{n-j} \pm \eta \sigma_{n-j}^z) \right) \left( \frac{\partial A_3(u')}{\partial u'} + \frac{\partial B_3(u')}{\partial u'} \right) \right] , \end{aligned}$$

from which we conclude the argument.  $\square$

#### 2.4.7 Seventh Poisson bracket, $\mathcal{P}_7$

**Lemma 12** (*evaluating the seventh Poisson bracket in the first relation*). The seventh term,  $\mathcal{P}_7$ , approximately equals,

$$\begin{aligned} \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_3, \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} &\approx \left[ (\sin(2\eta))^{n-3} \mathcal{C}_1 \right] \left[ \frac{A_3(u) B_3(u')}{u - u'} \right] + \left[ (\sin(2\eta))^{n-3} \mathcal{C}_1 \right] \left[ \frac{B_3(u) A_3(u')}{u - u'} \right] \\ &+ \left[ \sum_{1 \leq j \leq n'} (\sin(2\eta))^{n'-1} \mathcal{C}_1 \right] \left[ B_3(u) \left[ \left[ \sum_{1 \leq i \leq m} \left[ \left[ \frac{\partial}{\partial u} \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right] \left[ \frac{\partial B_3(u')}{\partial u'} - \frac{\partial B_3(u)}{\partial u} \right] \right. \right. \right. \right. \right. \\ &\quad \left. \left. \left. \left. \left. \left. - \left[ \frac{\partial}{\partial u'} \prod_{1 \leq i \leq m} \sin(\mathcal{C}_2)_i \right] \left[ \frac{\partial A_3(u')}{\partial u} - \frac{\partial A_3(u)}{\partial u'} \right] \right] \right] \right] \right] \\ &+ A_3(u) \left[ \sum_{1 \leq i \leq m} \left[ \left[ \frac{\partial}{\partial u} \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right] \left[ \frac{\partial A_3(u')}{\partial u'} - \frac{\partial A_3(u)}{\partial u} \right] \right] \right] \end{aligned}$$

$$-\left[ \frac{\partial}{\partial u'} \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right] \left[ \frac{\partial B_3(u')}{\partial u'} - \frac{\partial B_3(u')}{\partial u} \right] \right] \right] .$$

*Proof of Lemma 12.* The seventh term,

$$\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_3, \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} ,$$

is equivalent to,

$$\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right), \cdots \right. \\ \left. \mathcal{P}_2 (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \right\} ,$$

which can be expressed as,

$$\sum_{\mathcal{P}} \left\{ \mathcal{P}_1, \mathcal{P}_2 (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \right\} \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \right. \right. \\ \times \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \left. \right] \right) \\ + \sum_{\mathcal{P}} \left\{ \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \right. \right. \\ \times \left. \left. \left. (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right), \mathcal{P}_2 \right\} \mathcal{P}_1 ,$$

from one application of Leibniz' rule, and, from another application of Leibniz' rule,

$$-\left( \sum_{\mathcal{P}} \left\{ \mathcal{P}_2, \mathcal{P}_1 \right\} (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) + \sum_{\mathcal{P}} \left\{ (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right), \mathcal{P}_1 \right\} \mathcal{P}_2 \right) \\ \times \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \\ + \mathcal{P}_1 \left( \sum_{\mathcal{P}} \left( \left\{ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), \mathcal{P}_2 \right\} \sum_{1 \leq j \leq n'} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) \right) \right. \\ \left. + \left\{ \sum_{1 \leq j \leq n'} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right), \mathcal{P}_2 \right\} \right) \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) .$$

From the superposition above, writing out the first two Poisson bracket yields,

$$(\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \sum_{\mathcal{P}} \left\{ \mathcal{P}_1, \mathcal{P}_2 \right\} = (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \left( \{ A_3(u), B_3(u') \} + \{ B_3(u), A_3(u') \} \right) \\ \sum_{\mathcal{P}} \left\{ (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right), \mathcal{P}_1 \right\} \mathcal{P}_2 \equiv 0 .$$

The third Poisson bracket,

$$\mathcal{P}_1 \left( \sum_{\mathcal{P}} \left\{ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), \mathcal{P}_2 \right\} \sum_{1 \leq j \leq n'} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) \right),$$

can be expressed as,

$$B_3(u) \left( \left\{ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), B_3(u') \right\} + \left\{ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), A_3(u') \right\} \right) \\ \times \sum_{1 \leq j \leq n'} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) ,$$

corresponding to the first two terms, and,

$$A_3(u) \left( \left\{ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), A_3(u') \right\} + \left\{ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), B_3(u') \right\} \right) \\ \times \sum_{1 \leq j \leq n'} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) ,$$

corresponding to the next two terms.

The fourth Poisson bracket,

$$\left\{ \sum_{1 \leq j \leq n'} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right), \mathcal{P}_2 \right\} ,$$

vanishes, for all  $\mathcal{P}_2$ . The remaining nonzero terms,

$$(\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \left( \{A_3(u), B_3(u')\} + \{B_3(u), A_3(u')\} \right) + \mathcal{P}_1 \\ \times \left( \sum_{\mathcal{P}} \left\{ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), \mathcal{P}_2 \right\} \sum_{1 \leq j \leq n'} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) \right) ,$$

are each approximately equivalent to,

$$\left[ (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \right] \left[ \frac{A_3(u) B_3(u')}{u - u'} \right] , \\ \left[ (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \right] \left[ \frac{B_3(u) A_3(u')}{u - u'} \right] ,$$

corresponding to the first two Poisson brackets,

$$\{A_3(u), B_3(u')\} , \\ \{B_3(u), A_3(u')\} .$$

For the remaining brackets,

$$\left\{ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), B_3(u') \right\} ,$$

and,

$$\left\{ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), A_3(u') \right\} ,$$

corresponding to the third Poisson bracket above,

$$\sum_{\mathcal{P}} \left\{ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), \mathcal{P}_2 \right\} ,$$

from each possible  $\mathcal{P}_2$ . From each of the possible four brackets,

$$B_3(u) \left( \left\{ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), B_3(u') \right\} + \left\{ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), A_3(u') \right\} \right)$$

$$\times \sum_{1 \leq j \leq n'} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) ,$$

$$A_3(u) \left( \left\{ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), A_3(u') \right\} + \left\{ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), B_3(u') \right\} \right)$$

$$\times \sum_{1 \leq j \leq n'} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) ,$$

implies that the desired expression for each of the four brackets above would take the form,

$$\sum_{1 \leq j \leq n'} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) \left[ B_3(u) \left( \frac{\partial}{\partial u} \left[ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right] \frac{\partial}{\partial u'} B_3(u') \right. \right.$$

$$- \frac{\partial}{\partial u'} \left[ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right] \frac{\partial}{\partial u} B_3(u')$$

$$+ \frac{\partial}{\partial u} \left[ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right] \frac{\partial}{\partial u'} A_3(u')$$

$$+ \frac{\partial}{\partial u'} \left[ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right] \frac{\partial}{\partial u} A_3(u') \Big) \Big]$$

$$+ A_3(u) \left( \frac{\partial}{\partial u} \left[ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right] \frac{\partial}{\partial u'} A_3(u') \right. \right.$$

$$- \frac{\partial}{\partial u'} \left[ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right] \frac{\partial}{\partial u} A_3(u')$$

$$+ \frac{\partial}{\partial u} \left[ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right] \frac{\partial}{\partial u'} B_3(u')$$

$$\left. \left. - \frac{\partial}{\partial u'} \left[ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right] \frac{\partial}{\partial u} B_3(u') \right) \right] .$$

For the first bracket taken with respect to  $B_3(u')$ , write,

$$\frac{\partial}{\partial u} \left[ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right] = \frac{\partial}{\partial u} \left[ \left( \sin(u - v_{n-1} \pm \eta \sigma_{n-1}^z) \right) + \cdots \right.$$

$$+ \left( \sin(u - v_{n-1} \pm \eta \sigma_{n-1}^z) + \cdots + \left( \sin(u - v_{n-1} \pm \eta \sigma_{n-1}^z) \cdots \right. \right.$$

$$\left. \left. \times \sin(u - v_{n-(n-3)} \pm \eta \sigma_{n-(n-3)}^z) \right) \right) \Big] ,$$

which is further rearranged as,

$$\begin{aligned}
& \frac{\partial}{\partial u} \sin(u - v_{n-1} \pm \eta\sigma_{n-1}^z) + \frac{\partial}{\partial u} \left( \sin(u - v_{n-1} \pm \eta\sigma_{n-1}^z) + \cdots + \sin(u - v_{n-(n-3)} \pm \eta\sigma_{n-(n-3)}^z) \right) \\
&= 2\cos(u - v_{n-1} \pm \eta\sigma_{n-1}^z) + \cdots + \cos(u - v_{n-1} \pm \eta\sigma_{n-1}^z) \prod_{2 \leq i \leq n-3} \sin(u - v_{n-i} \pm \eta\sigma_{n-i}^z) \\
&= \sum_{1 \leq i \leq m} \left[ \frac{\partial}{\partial u} \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta\sigma_{n-i}^z) \right] .
\end{aligned}$$

This implies,

$$\begin{aligned}
& \sum_{1 \leq j \leq n'} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) \left[ B_3(u) \left( \frac{\partial}{\partial u} \left[ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) \right] \frac{\partial}{\partial u'} B_3(u') \right. \right. \\
& \quad \left. \left. - \frac{\partial}{\partial u'} \left[ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) \right] \frac{\partial}{\partial u} B_3(u') \right. \\
& \quad \left. + \frac{\partial}{\partial u} \left[ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) \right] \frac{\partial}{\partial u'} A_3(u') \right. \\
& \quad \left. + \frac{\partial}{\partial u'} \left[ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) \right] \frac{\partial}{\partial u} A_3(u') \right) \\
& \quad + A_3(u) \left( \frac{\partial}{\partial u} \left[ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) \right] \frac{\partial}{\partial u'} A_3(u') \right. \\
& \quad \left. - \frac{\partial}{\partial u'} \left[ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) \right] \frac{\partial}{\partial u} A_3(u') \right. \\
& \quad \left. + \frac{\partial}{\partial u} \left[ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) \right] \frac{\partial}{\partial u'} B_3(u') \right. \\
& \quad \left. - \frac{\partial}{\partial u'} \left[ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) \right] \frac{\partial}{\partial u} B_3(u') \right) .
\end{aligned}$$

equals,

$$\begin{aligned}
& \sum_{1 \leq j \leq n'} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) \left[ B_3(u) \left[ \left[ \sum_{1 \leq i \leq m} \left[ \left[ \frac{\partial}{\partial u} \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta\sigma_{n-i}^z) \right] \right. \right. \right. \right. \\
& \quad \times \left[ \frac{\partial}{\partial u'} \left[ B_3(u') + A_3(u') \right] \right] \\
& \quad \left. \left. \left. \left. - \left[ \frac{\partial}{\partial u'} \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta\sigma_{n-i}^z) \right] \right. \right. \right. \\
& \quad \times \left[ \frac{\partial}{\partial u} \left[ A_3(u') + B_3(u') \right] \right] \right] \\
& \quad + A_3(u) \left[ \sum_{1 \leq i \leq m} \left[ \left[ \frac{\partial}{\partial u} \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta\sigma_{n-i}^z) \right] \left[ \frac{\partial}{\partial u'} \left[ A_3(u') + B_3(u') \right] \right] \right] \\
& \quad \left. \left. \left. - \left[ \frac{\partial}{\partial u'} \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta\sigma_{n-i}^z) \right] \left[ \frac{\partial}{\partial u} \left[ A_3(u') + B_3(u') \right] \right] \right] \right] .
\end{aligned}$$

Altogether,

$$\begin{aligned}
\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_3, \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{A}'_1 \right\} &\approx \left[ (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \right] \left[ \frac{A_3(u) B_3(u')}{u - u'} \right] \\
&\quad + \left[ (\sin(2\eta))^{n-3} \left( \prod_{1 \leq i \leq n-3} \sigma_{n-i}^{-,+} \right) \right] \left[ \frac{B_3(u) A_3(u')}{u - u'} \right] \\
&+ \sum_{1 \leq j \leq n'} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right) \left[ B_3(u) \left[ \left[ \sum_{1 \leq i \leq m} \left[ \left[ \frac{\partial}{\partial u} \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right] \right. \right. \right. \right. \right. \\
&\quad \times \left[ \frac{\partial}{\partial u'} \left[ B_3(u') + A_3(u') \right] \right] \left. \left. \left. \left. \left. \left. \right] \right] \right] \\
&\quad - \left[ \frac{\partial}{\partial u'} \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right] \\
&\quad \times \left[ \frac{\partial}{\partial u} \left[ A_3(u') + B_3(u') \right] \right] \left. \left. \left. \left. \left. \right] \right] \right] \\
&+ A_3(u) \left[ \sum_{1 \leq i \leq m} \left[ \left[ \frac{\partial}{\partial u} \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right] \left[ \frac{\partial}{\partial u'} \left[ A_3(u') + B_3(u') \right] \right] \right] \right. \\
&\quad \left. - \left[ \frac{\partial}{\partial u'} \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right] \left[ \frac{\partial}{\partial u} \left[ A_3(u') + B_3(u') \right] \right] \right] \right] ,
\end{aligned}$$

from which we conclude the argument.  $\square$

#### 2.4.8 Eighth Poisson bracket, $\mathcal{P}_8$

**Lemma 13** (*evaluating the eighth Poisson bracket in the first relation*). The eighth term,  $\mathcal{P}_8$ , approximately equals,

$$\begin{aligned}
\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_3, \mathcal{P}_2 \mathcal{A}'_2 \right\} &\approx - \left[ \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right) (\sin(2\eta))^{n'-1} \mathcal{C}_1 \right] \right) \left( \prod_{1 \leq i \leq n-3} (\mathcal{C}_2)_i \right) \right] \left[ \frac{A_3(u) B_3(u')}{u - u'} \right] \\
&\quad - \left[ \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right) (\sin(2\eta))^{n'-1} \mathcal{C}_1 \right] \right. \right. \\
&\quad \times \left. \left. \left( \prod_{1 \leq i \leq n-3} (\mathcal{C}_2)_i \right) \right] \left[ \frac{B_3(u) A_3(u')}{u - u'} \right] \right] \\
&\quad - 2 \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right) (\sin(2\eta))^{n'-1} \mathcal{C}_1 \right] \right) \\
&\quad \times B_3(u') \left[ \sum_{1 \leq j \leq n-3} \left[ \left[ \frac{\partial B_3(u)}{\partial u} \right] \left[ \frac{\partial (\mathcal{C}_2)_i}{\partial u'} \right] - \left[ \frac{\partial B_3(u)}{\partial u'} \right] \left[ \frac{\partial (\mathcal{C}_2)_i}{\partial u} \right] \right] \right. \\
&\quad \times \left. \left. \prod_{1 \leq j \neq i \leq n-3} (\mathcal{C}_2)_j \right] \right] \\
&\quad - \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right) (\sin(2\eta))^{n'-1} \mathcal{C}_1 \right] \right) \\
&\quad \times A_3(u') \left[ \sum_{1 \leq j \leq n-3} \left[ \left[ \frac{\partial A_3(u)}{\partial u} \right] \left[ \frac{\partial (\mathcal{C}_2)_i}{\partial u'} \right] - \left[ \frac{\partial A_3(u)}{\partial u'} \right] \left[ \frac{\partial (\mathcal{C}_2)_i}{\partial u} \right] \right] \right]
\end{aligned}$$

$$\begin{aligned}
& \times \prod_{1 \leq j \neq i \leq n-3} (\mathcal{C}_2)_j \Big] \\
& - \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} (\mathcal{C}_2)_i \right) (\sin(2\eta))^{n'-1} \mathcal{C}_1 \right] \right) \\
& \times A_3(u') \left[ \sum_{1 \leq j \leq n-3} \left[ \left[ \frac{\partial B_3(u')}{\partial u'} \right] \left[ \frac{\partial (\mathcal{C}_2)_i}{\partial u'} \right] - \left[ \frac{\partial B_3(u')}{\partial u} \right] \left[ \frac{\partial (\mathcal{C}_2)_i}{\partial u} \right] \right] \right] \\
& \times \prod_{1 \leq j \neq i \leq n-3} (\mathcal{C}_2)_j \Big] \\
+ (\sin(2\eta))^{n'-1} & \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \left( \left[ \frac{\partial}{\partial u} \left[ B_3(u') + A_3(u') \right] \right] \frac{\partial (\mathcal{C}_2)_i}{\partial u'} - \left[ \frac{\partial}{\partial u'} \left[ A_3(u') + B_3(u') \right] \right] \frac{\partial (\mathcal{C}_2)_i}{\partial u} \right) \right) \right] .
\end{aligned}$$

*Proof of Lemma 13.* The eighth term,

$$\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_3, \mathcal{P}_2 \mathcal{A}'_2 \right\} ,$$

is equivalent to,

$$\begin{aligned}
& \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right), \mathcal{P}_2 \right. \\
& \quad \left. \times \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} ,
\end{aligned}$$

which can be rearranged as,

$$\begin{aligned}
& \sum_{\mathcal{P}} \left( \left\{ \mathcal{P}_1, \mathcal{P}_2 \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \right. \right. \\
& \quad \left. \times (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \left. \right) + \left\{ \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \right. \right. \\
& \quad \left. \times (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \left. \right), \mathcal{P}_2 \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} \mathcal{P}_1 \Big) ,
\end{aligned}$$

from an application of Leibniz' rule. Further rearranging each summation of Poisson brackets over  $\mathcal{P}$  implies,

$$\begin{aligned}
& - \sum_{\mathcal{P}} \{ \mathcal{P}_2, \mathcal{P}_1 \} \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) \\
& \quad \times \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \\
& - \sum_{\mathcal{P}} \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), \mathcal{P}_1 \right\} \mathcal{P}_2 \\
& \quad \times \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \right. \\
& \quad \times \left. \left. (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) ,
\end{aligned}$$

after applying Leibniz' rule to,

$$\begin{aligned}
& \sum_{\mathcal{P}} \left\{ \mathcal{P}_1, \mathcal{P}_2 \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \right. \\
& \quad \times \left. \left. (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) ,
\end{aligned}$$

and similarly,

$$\begin{aligned}
& - \mathcal{P}_1 \left[ \sum_{\mathcal{P}} \left\{ \mathcal{P}_2, \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) \right\} \times \cdots \right. \\
& \quad \left. \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) - \cdots \right. \\
& \quad \left. \sum_{\mathcal{P}} \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \right. \right. \right. \\
& \quad \times \left. \left. \left. \left. (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right\} \mathcal{P}_2 \right] ,
\end{aligned}$$

after applying Leibniz' rule to,

$$\begin{aligned}
& \left\{ \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right), \mathcal{P}_2 \right. \\
& \quad \times \left. \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right\} \mathcal{P}_1 .
\end{aligned}$$

Following each applications of Leibniz' rule, we apply Leibniz' rule multiple times to rearrange the second Poisson bracket, in which,

$$\begin{aligned}
& \mathcal{P}_1 \left[ \sum_{\mathcal{P}} \left\{ \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right), \mathcal{P}_2 \right\} \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \right] \right) \right. \\
& \quad \left. + \sum_{\mathcal{P}} \left\{ \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \right. \right. \right. \\
& \quad \times (\sin(2\eta))^{n'-1} \left. \left. \left. \right] \right), \mathcal{P}_2 \right\} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] ,
\end{aligned}$$

after anticommuting the first bracket,

$$-\mathcal{P}_1 \sum_{\mathcal{P}} \left\{ \mathcal{P}_2, \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) \right\} .$$

The expression,

$$\begin{aligned}
& -\sum_{\mathcal{P}} \{ \mathcal{P}_2, \mathcal{P}_1 \} \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) \\
& \quad \times \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \\
& - \sum_{\mathcal{P}} \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), \mathcal{P}_1 \right\} \mathcal{P}_2 \\
& \quad \times \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \right. \\
& \quad \times (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \left. \right] \right) \\
& + \mathcal{P}_1 \left[ \sum_{\mathcal{P}} \left\{ \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right), \mathcal{P}_2 \right\} \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \right] \right) \right. \\
& \quad \left. + \sum_{\mathcal{P}} \left\{ \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \right. \right. \right. \\
& \quad \times (\sin(2\eta))^{n'-1} \left. \left. \left. \right] \right), \mathcal{P}_2 \right\} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] ,
\end{aligned}$$

can be rearranged to obtain the following Poisson brackets,

$$\sum_{\mathcal{P}} \{ \mathcal{P}_1, \mathcal{P}_2 \} = \{ A_3(u), B_3(u') \} + \{ B_3(u), A_3(u') \} ,$$

corresponding to the first term, and,

$$\begin{aligned}
& \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), A_3(u) \right\} A_3(u') + \sum_{\mathcal{P}} \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), B_3(u) \right\} \\
& \quad \times B_3(u') \\
& \quad + \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), A_3(u) \right\} B_3(u') \\
& \quad + \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), B_3(u) \right\} A_3(u') ,
\end{aligned}$$

corresponding to the second term.

The first Poisson bracket approximately equals,

$$\begin{aligned}
& - \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) \\
& \quad \times \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \left[ \frac{A_3(u) B_3(u')}{u - u'} \right] \\
& - \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) \\
& \quad \times \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \left[ \frac{B_3(u) A_3(u')}{u - u'} \right] .
\end{aligned}$$

For the second Poisson bracket, taking the summation over all  $\mathcal{P}$  implies that the terms,

$$\sum_{\mathcal{P}} \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), \mathcal{P}_1 \right\} \mathcal{P}_2 ,$$

can be evaluated by observing that the single Poisson bracket is equivalent to the following four brackets,

$$\begin{aligned}
& \sum_{\mathcal{P}} \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), B_3(u) \right\} B_3(u') , \\
& \sum_{\mathcal{P}} \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), A_3(u) \right\} A_3(u') , \\
& \sum_{\mathcal{P}} \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), B_3(u') \right\} A_3(u') , \\
& \sum_{\mathcal{P}} \left\{ \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right), B_3(u) \right\} A_3(u') ,
\end{aligned}$$

which can each be individually evaluated below. For the first bracket, evaluating terms from the bracket yields,

$$B_3(u') \left[ \frac{\partial B_3(u)}{\partial u'} \left[ \frac{\partial}{\partial u'} \left[ \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right] \right] - \left[ \frac{\partial}{\partial u} \left[ \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right] \right] \frac{\partial B_3(u)}{\partial u} \right] .$$

The derivative of the product of sine functions appearing in the first term of the Poisson bracket above,

$$\frac{\partial}{\partial u'} \left[ \sin(u' - v_{n-1} + \eta \sigma_{n-1}^z) \times \cdots \times \sin(u' - v_{n-(n-3)} + \eta \sigma_{n-(n-3)}^z) \right] ,$$

equals,

$$\begin{aligned} & \left[ \frac{\partial}{\partial u'} \sin(u' - v_{n-1} + \eta \sigma_{n-1}^z) \right] \prod_{2 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) + \cdots + \prod_{1 \leq i \leq n-4} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \\ & \quad \times \left[ \frac{\partial}{\partial u'} \sin(u' - v_{n-(n-3)} + \eta \sigma_{n-(n-3)}^z) \right] \\ &= \left[ \sum_{1 \leq j \leq n-3} \frac{\partial}{\partial u'} \left[ \sin(u' - v_{n-j} + \eta \sigma_{n-j}^z) \right] \right] \left[ \prod_{1 \leq j \neq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right]. \end{aligned}$$

Hence, the first bracket is equivalent to,

$$B_3(u') \left[ \sum_{1 \leq j \leq n-3} \left[ \left[ \frac{\partial}{\partial u'} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right] \left[ \frac{\partial B_3(u)}{\partial u'} \right] - \left[ \frac{\partial}{\partial u} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right] \left[ \frac{\partial B_3(u)}{\partial u} \right] \right] \right. \\ \left. \times \prod_{1 \leq j \neq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right].$$

For the second bracket, evaluating terms from the bracket similarly yields,

$$A_3(u') \left[ \sum_{1 \leq j \leq n-3} \left[ \left[ \frac{\partial}{\partial u'} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right] \left[ \frac{\partial A_3(u)}{\partial u'} \right] - \left[ \frac{\partial}{\partial u} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right] \left[ \frac{\partial A_3(u)}{\partial u} \right] \right] \right. \\ \left. \times \prod_{1 \leq j \neq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right].$$

For the third bracket, evaluating terms from the bracket similarly yields,

$$A_3(u') \left[ \sum_{1 \leq j \leq n-3} \left[ \left[ \frac{\partial}{\partial u'} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right] \left[ \frac{\partial B_3(u')}{\partial u'} \right] - \left[ \frac{\partial}{\partial u} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right] \left[ \frac{\partial B_3(u')}{\partial u} \right] \right] \right. \\ \left. \times \prod_{1 \leq j \neq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right].$$

For the fourth bracket, evaluating terms from the bracket similarly yields,

$$A_3(u') \left[ \sum_{1 \leq j \leq n-3} \left[ \left[ \frac{\partial}{\partial u'} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right] \left[ \frac{\partial B_3(u)}{\partial u'} \right] - \left[ \frac{\partial}{\partial u} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right] \left[ \frac{\partial B_3(u)}{\partial u} \right] \right] \right. \\ \left. \times \prod_{1 \leq j \neq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right].$$

For the third Poisson bracket, the terms,

$$\sum_{\mathcal{P}} \left\{ \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right), \mathcal{P}_2 \right\} = \left\{ \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right), B_3(u') \right\} + \left\{ \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right), A_3(u') \right\} \equiv 0 ,$$

vanish, while for the fourth Poisson bracket,

$$\sum_{\mathcal{P}} \left\{ \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \right] \right), \mathcal{P}_2 \right\} ,$$

computing the differentiation,

$$\frac{\partial}{\partial u} \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \right] \right],$$

which can be arranged as,

$$(\sin(2\eta))^{n'-1} \left[ \left( \sin(u - v_{n-1} \pm \eta \sigma_{n-1}^z) \right) + \cdots + \left( \sin(u - v_{n-1} \pm \eta \sigma_{n-1}^z) + \cdots + \cdots \right. \right. \\ \left. \left. \left( \sin(u - v_{n-1} \pm \eta \sigma_{n-1}^z) \times \cdots \times \sin(u - v_{n-m} \pm \eta \sigma_{n-m}^z) \right) \right) \right],$$

with derivative,

$$(\sin(2\eta))^{n'-1} \left[ \cos(u - v_{n-1} \pm \sigma_{n-1}^z) + \cdots + \left( \cos(u - v_{n-1} \pm \sigma_{n-1}^z) + \cdots \right. \right. \\ \left. \left. + \left( \cos(u - v_{n-m} \pm \eta \sigma_{n-m}^z) \prod_{1 \leq i \leq m-1} \sin(u - v_{n-i} \pm \eta \sigma_{n-1}^z) \right) \right) \right].$$

Therefore,

$$\frac{\partial}{\partial u} \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \right] \right] \left[ \frac{\partial \mathcal{P}_2}{\partial u} \right] \\ - \left[ \frac{\partial \mathcal{P}_2}{\partial u} \right] \frac{\partial}{\partial u'} \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \right] \right],$$

equals, for  $\mathcal{P}_2 \equiv B_3(u')$ ,  $\mathcal{P}_2 \equiv A_3(u')$ ,

$$\frac{\partial}{\partial u'} \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \right] \right] \left[ \frac{\partial B_3(u')}{\partial u} \right] \\ - \left[ \frac{\partial B_3(u')}{\partial u} \right] \frac{\partial}{\partial u'} \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \right] \right],$$

or,

$$\frac{\partial}{\partial u'} \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \right] \right] \left[ \frac{\partial A_3(u')}{\partial u} \right] \\ - \left[ \frac{\partial A_3(u')}{\partial u} \right] \frac{\partial}{\partial u'} \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \right] \right].$$

In the first case, introducing the expression for the derivative of the sum of product of sine functions equals, approximately,

$$\begin{aligned}
& (\sin(2\eta))^{n'-1} \left[ \left( \sin(u - v_{n-1} \pm \eta\sigma_{n-1}^z) \right) + \cdots + \left( \sin(u - v_{n-1} \pm \eta\sigma_{n-1}^z) + \cdots + \cdots \right. \right. \\
& \quad \left. \left. \left( \sin(u - v_{n-1} \pm \eta\sigma_{n-1}^z) \times \cdots \times \sin(u - v_{n-m} \pm \eta\sigma_{n-m}^z) \right) \right) \right] \left[ \frac{\partial B_3(u')}{\partial u} \right] - \cdots \\
& (\sin(2\eta))^{n'-1} \left[ \left( \sin(u - v_{n-1} \pm \eta\sigma_{n-1}^z) \right) + \cdots + \left( \sin(u - v_{n-1} \pm \eta\sigma_{n-1}^z) + \cdots + \cdots \right. \right. \\
& \quad \left. \left. \left( \sin(u - v_{n-1} \pm \eta\sigma_{n-1}^z) \times \cdots \times \sin(u - v_{n-m} \pm \eta\sigma_{n-m}^z) \right) \right) \right] \left[ \frac{\partial B_3(u')}{\partial u'} \right] ,
\end{aligned}$$

The expression above approximately takes the form,

$$\begin{aligned}
& -(\sin(2\eta))^{n'-1} \left[ \frac{\partial B_3(u')}{\partial u} \right] \left[ \left( \sin(u - v_{n-1} \pm \eta\sigma_{n-1}^z) \right) + \cdots + \left( \sin(u - v_{n-1} \pm \eta\sigma_{n-1}^z) + \cdots + \cdots \right. \right. \\
& \quad \left. \left. \left( \sin(u - v_{n-1} \pm \eta\sigma_{n-1}^z) \times \cdots \times \sin(u - v_{n-m} \pm \eta\sigma_{n-m}^z) \right) \right) + \cdots \right. \\
& \quad \left. \left. \frac{\partial}{\partial u} \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) \right] \right] \right] \\
& -(\sin(2\eta))^{n'-1} \left[ \frac{\partial B_3(u')}{\partial u'} \right] \left[ \left( \sin(u - v_{n-1} \pm \eta\sigma_{n-1}^z) \right) + \cdots + \left( \sin(u - v_{n-1} \pm \eta\sigma_{n-1}^z) + \cdots + \cdots \right. \right. \\
& \quad \left. \left. \left( \sin(u - v_{n-1} \pm \eta\sigma_{n-1}^z) \times \cdots \times \sin(u - v_{n-m} \pm \eta\sigma_{n-m}^z) \right) \right) \right. \\
& \quad \left. + \frac{\partial}{\partial u'} \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) \right] \right] \right] ,
\end{aligned}$$

which equals,

$$\begin{aligned}
& (\sin(2\eta))^{n'-1} \left[ \left[ \frac{\partial B_3(u')}{\partial u} \right] \frac{\partial}{\partial u'} \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) \right] \right] \\
& - \left[ \frac{\partial B_3(u')}{\partial u'} \right] \frac{\partial}{\partial u} \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) \right] \right] .
\end{aligned}$$

Grouping together like terms in the summation  $i$  and  $j$  yields,

$$(\sin(2\eta))^{n'-1} \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \left( \left[ \frac{\partial B_3(u')}{\partial u} \right] \frac{\partial}{\partial u'} \sin(u - v_{n-i} \pm \eta\sigma_{n-i}^z) - \left[ \frac{\partial B_3(u')}{\partial u'} \right] \frac{\partial}{\partial u} \sin(u - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) \right) \right] ,$$

while performing an identical substitution in the second case implies,

$$(\sin(2\eta))^{n'-1} \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \left( \left[ \frac{\partial A_3(u')}{\partial u} \right] \frac{\partial}{\partial u'} \sin(u - v_{n-i} \pm \eta\sigma_{n-i}^z) - \left[ \frac{\partial A_3(u')}{\partial u'} \right] \frac{\partial}{\partial u} \sin(u - v_{n-i} \pm \eta\sigma_{n-i}^z) \right) \right) \right] .$$

Altogether,

$$\begin{aligned}
\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_3, \mathcal{P}_2 \mathcal{A}'_2 \right\} &\approx - \left[ \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) \right. \\
&\quad \times \left. \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] \left[ \frac{A_3(u) B_3(u')}{u - u'} \right] \\
&- \left[ \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) \right. \\
&\quad \times \left. \left( \prod_{1 \leq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right) \right] \left[ \frac{B_3(u) A_3(u')}{u - u'} \right] \\
&- 2 \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) \\
&\times B_3(u') \left[ \sum_{1 \leq j \leq n-3} \left[ \left[ \frac{\partial B_3(u)}{\partial u} \right] \left[ \frac{\partial}{\partial u'} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right] - \left[ \frac{\partial B_3(u)}{\partial u'} \right] \left[ \frac{\partial}{\partial u} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right] \right] \right. \\
&\quad \times \left. \prod_{1 \leq j \neq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right] \\
&- \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) \\
&\times A_3(u') \left[ \sum_{1 \leq j \leq n-3} \left[ \left[ \frac{\partial A_3(u)}{\partial u} \right] \left[ \frac{\partial}{\partial u'} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right] - \left[ \frac{\partial A_3(u)}{\partial u'} \right] \left[ \frac{\partial}{\partial u} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right] \right] \right. \\
&\quad \times \left. \prod_{1 \leq j \neq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right] \\
&- \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) \\
&\times A_3(u') \left[ \sum_{1 \leq j \leq n-3} \left[ \left[ \frac{\partial B_3(u')}{\partial u'} \right] \left[ \frac{\partial}{\partial u'} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right] - \left[ \frac{\partial B_3(u')}{\partial u} \right] \left[ \frac{\partial}{\partial u} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right] \right] \right. \\
&\quad \times \left. \prod_{1 \leq j \neq i \leq n-3} \sin(u' - v_{n-i} + \eta \sigma_{n-i}^z) \right] \\
&+ (\sin(2\eta))^{n'-1} \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \left( \left[ \frac{\partial}{\partial u} [B_3(u') + A_3(u')] \right] \frac{\partial}{\partial u'} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right. \right. \right. \\
&\quad \left. \left. \left. - \left[ \frac{\partial}{\partial u'} [A_3(u') + B_3(u')] \right] \frac{\partial}{\partial u} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right) \right] ,
\end{aligned}$$

from which we conclude the argument.  $\square$

#### 2.4.9 Ninth Poisson bracket, $\mathcal{P}_9$

**Lemma 14** (*evaluating the ninth Poisson bracket in the first relation*). The ninth term approximately equals,

$$\begin{aligned}
\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_3, \mathcal{P}_2 \mathcal{A}'_3 \right\} &\approx - \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} [\mathcal{C}_2 (\sin(2\eta))^{n'-1} \mathcal{C}_1] \right] \left[ \frac{B_3(u') A_3(u)}{u' - u} + \frac{A_3(u') B_3(u)}{u' - u} \right] \\
&- \left[ 2[B_3(u) + A_3(u)] \left[ \sum_{n':m+n'=n-3} \left( (\sin(2\eta))^{n'-1} \mathcal{C}_2 \right)^2 \right] \right] \left[ \frac{B_3(u') A_3(u)}{u' - u} + \frac{A_3(u') B_3(u)}{u' - u} \right] \\
&+ 2 \left[ \frac{\partial}{\partial u'} \left[ A_3(u') + B_3(u') \right] \right] \left[ \sum_{1 \leq i \leq m} \left[ \frac{\partial}{\partial u} \left[ \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right] \right] \right] - \left[ \frac{\partial}{\partial u} \left[ A_3(u') + B_3(u') \right] \right] \\
&\quad \times \left[ \sum_{1 \leq i \leq m} \left[ \frac{\partial}{\partial u'} \left[ \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right] \right] \right] .
\end{aligned}$$

*Proof of Lemma 14.* The ninth term,

$$\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_3, \mathcal{P}_2 \mathcal{A}'_3 \right\} ,$$

is equivalent to,

$$\begin{aligned}
&\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right), \dots \right. \\
&\quad \left. \mathcal{P}_2 \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) \right\} ,
\end{aligned}$$

which can be rearranged with Leibniz' rule, as,

$$\begin{aligned}
&\sum_{\mathcal{P}} \left\{ \mathcal{P}_1, \mathcal{P}_2 \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) \right\} \\
&\quad \times \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) \\
&\quad + \sum_{\mathcal{P}} \left\{ \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \right. \right. \right. \\
&\quad \left. \left. \left. \times \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right), \mathcal{P}_2 \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \right. \right. \right. \\
&\quad \left. \left. \left. \times \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) \right\} \mathcal{P}_1 .
\end{aligned}$$

Applying Leibniz' for a second time to each bracket yields,

$$\begin{aligned}
& - \sum_{\mathcal{P}} \{ \mathcal{P}_2, \mathcal{P}_1 \} \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) \\
& - \sum_{\mathcal{P}} \left\{ \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right), \mathcal{P}_1 \right\} \mathcal{P}_2 \\
& \quad + \mathcal{P}_1 \sum_{\mathcal{P}} \left\{ \sum_{\substack{1 \leq i \leq m \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), \mathcal{P}_2 \right. \\
& \quad \times \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \right. \\
& \quad \times \left. \left. (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right\} (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \\
& \quad + \left\{ (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right), \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \right. \right. \\
& \quad \times \left. \left. \left. (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right\} \sum_{\substack{1 \leq i \leq m \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) .
\end{aligned}$$

Applying Leibniz' rule for a third time to the second, third, and fourth, brackets yields,

$$\begin{aligned}
& - \sum_{\mathcal{P}} \{ \mathcal{P}_2, \mathcal{P}_1 \} \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \right. \\
& \quad \times \left. \left. (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) \\
& - \sum_{\mathcal{P}} \left( \left\{ \sum_{n':m+n'=n-3} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-i}^{-,+} \right) \right), \mathcal{P}_1 \right\} \mathcal{P}_2 \sum_{\substack{1 \leq i \leq m \\ m+n'=n+3}} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \\
& \quad - \sum_{\mathcal{P}} \left\{ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), \mathcal{P}_1 \right\} \mathcal{P}_2 \\
& \quad \times \sum_{n':m+n'=n-3} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right)^2 \\
& \quad - \mathcal{P}_1 \sum_{\mathcal{P}} \left\{ \mathcal{P}_2, \sum_{\substack{1 \leq i \leq m \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right\} \\
& \quad \times \sum_{1 \leq i \leq m} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \\
& \quad \times \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \\
& \quad \times \left. \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right)^2 \right] \right) .
\end{aligned}$$

The last Poisson bracket vanishes because the derivatives,

$$\begin{aligned} \frac{\partial}{\partial u} & \left[ (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] , \\ \frac{\partial}{\partial u'} & \left[ (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] , \end{aligned}$$

vanish. Furthermore, the second Poisson bracket also vanishes, because the derivatives,

$$\begin{aligned} \frac{\partial}{\partial u} & \left[ \sum_{n':m+n'=n-3} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-i}^{-,+} \right) \right) \right] , \\ \frac{\partial}{\partial u'} & \left[ \sum_{n':m+n'=n-3} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-i}^{-,+} \right) \right) \right] , \end{aligned}$$

vanish. The nonzero entries from the first Poisson bracket between  $\mathcal{P}_1$  and  $\mathcal{P}_2$  are,

$$\begin{aligned} \left( \{A_3(u), B_3(u')\} + \{B_3(u), A_3(u')\} \right) & \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \right. \right. \\ & \quad \times \left. \left. \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) , \end{aligned}$$

while the nonzero entries for the third Poisson bracket are,

$$\begin{aligned} -\mathcal{P}_2 & \left[ \sum_{n':m+n'=n-3} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right)^2 \right] \left( \left\{ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \sigma_{n-i}^z) \right), B_3(u') \right\} \right. \\ & \quad \left. + \left\{ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \sigma_{n-i}^z) \right), A_3(u') \right\} \right) . \end{aligned}$$

The nonzero entries for the fourth Poisson bracket are,

$$\begin{aligned} \mathcal{P}_1 & \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right)^2 \right] \left( \left\{ \sum_{\substack{1 \leq i \leq m \\ m+n'=n-3}} \right. \right. \\ & \quad \times \left. \left. \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), B_3(u') \right\} \right. \\ & \quad \left. + \left\{ \sum_{\substack{1 \leq i \leq m \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), A_3(u') \right\} \right) . \end{aligned}$$

Altogether, the remaining entries,

$$\begin{aligned}
& \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) \{A_3(u), B_3(u')\} \\
& + \left( \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right) \{B_3(u), A_3(u')\} \\
& - \mathcal{P}_2 \left[ \sum_{n':m+n'=n-3} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right)^2 \right] \left\{ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \sigma_{n-i}^z) \right), B_3(u') \right\} \\
& - \mathcal{P}_2 \left[ \sum_{n':m+n'=n-3} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right)^2 \right] \left\{ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \sigma_{n-i}^z) \right), A_3(u') \right\} ,
\end{aligned}$$

from the first, and third, Poisson brackets, as well as,

$$\begin{aligned}
& -\mathcal{P}_1 \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \\
& \times \left. \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right)^2 \right] \left\{ \sum_{\substack{1 \leq i \leq m \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), B_3(u') \right\} ,
\end{aligned}$$

and,

$$\begin{aligned}
& \mathcal{P}_1 \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \\
& \times \left. \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right)^2 \right] \left\{ \sum_{\substack{1 \leq i \leq m \\ m+n'=n-3}} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right), A_3(u') \right\} ,
\end{aligned}$$

from the fourth Poisson bracket. The observations that,

$$\{A_3(u), B_3(u')\} = -\{B_3(u'), A_3(u)\} \approx -\frac{B_3(u') A_3(u)}{u' - u} ,$$

and that,

$$\{B_3(u), A_3(u')\} = -\{A_3(u'), B_3(u)\} \approx -\frac{A_3(u') B_3(u)}{u' - u} ,$$

corresponding to the two entries from the first Poisson bracket,

$$\left\{ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \sigma_{n-i}^z) \right), B_3(u') \right\} ,$$

and,

$$\left\{ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \sigma_{n-i}^z) \right), A_3(u') \right\} ,$$

corresponding to the two entries from the second Poisson bracket. Along the lines of similar computations for previous Poisson brackets in the first relation, we evaluate each term by observing that the derivative of the sum of product of sine functions,

$$\frac{\partial}{\partial u'} \left[ \sum_{1 \leq i \leq m} \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right] ,$$

can be expressed as,

$$\begin{aligned} \frac{\partial}{\partial u'} & \left[ \left( \sin(u' - v_{n-1} \pm \eta \sigma_{n-1}^z) \right) + \cdots + \left( \left( \sin(u' - v_{n-1} \pm \eta \sigma_{n-1}^z) \right) + \cdots + \cdots \right. \right. \\ & \left. \left. \left( \sin(u' - v_{n-1} \pm \eta \sigma_{n-1}^z) \times \cdots \times \sin(u' - v_{n-m} \pm \eta \sigma_{n-m}^z) \right) \right) \right] , \end{aligned}$$

corresponding to the first term. Explicitly,

$$\begin{aligned} & \left( \cos(u' - v_{n-1} \pm \eta \sigma_{n-1}^z) \right) + \cdots + \left( \cos(u' - v_{n-1} \pm \eta \sigma_{n-1}^z) + \cdots + \left( \sum_{1 \leq i \leq m} \left[ \frac{\partial}{\partial u'} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right] \right. \right. \\ & \left. \left. \times \left( \prod_{1 \leq j \neq i \leq m} \sin(u' - v_{n-j} \pm \eta \sigma_{n-j}^z) \right) \right) \right) . \end{aligned}$$

Hence the first bracket takes the form, under a single summation from  $i \equiv 1$  to  $i \equiv m$ ,

$$\begin{aligned} & \left[ \sum_{1 \leq i \leq m} \frac{\partial}{\partial u} \left[ \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right] \right] \left[ \frac{\partial B_3(u')}{\partial u'} \right] - \left[ \left[ \sum_{1 \leq i \leq m} \frac{\partial}{\partial u'} \left[ \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right] \right] \right] \\ & \quad \times \left[ \frac{\partial B_3(u')}{\partial u} \right] \\ = & \left[ \sum_{1 \leq i \leq m} \left[ \left[ \frac{\partial}{\partial u} \left[ \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right] \right] \left[ \frac{\partial B_3(u')}{\partial u'} \right] - \left[ \frac{\partial}{\partial u'} \left[ \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right] \right] \right. \right. \\ & \quad \left. \left. \times \left[ \frac{\partial B_3(u')}{\partial u} \right] \right] \right] . \end{aligned}$$

For the second term, one has,

$$\left[ \sum_{1 \leq i \leq m} \left[ \left[ \frac{\partial}{\partial u} \left[ \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right] \right] \left[ \frac{\partial A_3(u')}{\partial u'} \right] - \left[ \frac{\partial}{\partial u'} \left[ \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right] \right] \left[ \frac{\partial A_3(u')}{\partial u} \right] \right] \right] ,$$

from which combining the two Poisson brackets above yields,

$$\begin{aligned} & \left[ \frac{\partial}{\partial u'} \left[ A_3(u') + B_3(u') \right] \right] \left[ \sum_{1 \leq i \leq m} \left[ \frac{\partial}{\partial u} \left[ \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right] \right] \right] - \left[ \frac{\partial}{\partial u} \left[ A_3(u') + B_3(u') \right] \right] \\ & \quad \times \left[ \sum_{1 \leq i \leq m} \left[ \frac{\partial}{\partial u'} \left[ \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right] \right] \right] . \end{aligned}$$

For the two remaining possibilities for  $\mathcal{P}_2$  in the first and second Poisson bracket evaluated above, combining terms yields the same identity,

$$\begin{aligned} & \left[ \frac{\partial}{\partial u'} \left[ A_3(u') + B_3(u') \right] \right] \left[ \sum_{1 \leq i \leq m} \left[ \frac{\partial}{\partial u} \left[ \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right] \right] \right] - \left[ \frac{\partial}{\partial u} \left[ A_3(u') + B_3(u') \right] \right] \\ & \quad \times \left[ \sum_{1 \leq i \leq m} \left[ \frac{\partial}{\partial u'} \left[ \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right] \right] \right] . \end{aligned}$$

The two computations above together imply that the desired expression for the ninth term takes the form,

$$\begin{aligned}
\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_3, \mathcal{P}_2 \mathcal{A}'_3 \right\} \approx & - \left[ \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n' \\ m+n'=n-3}} \left[ \left( \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] \right] \\
& \times \left[ \frac{B_3(u') A_3(u)}{u' - u} + \frac{A_3(u') B_3(u)}{u' - u} \right] \\
& - \left[ 2[B_3(u) + A_3(u)] \left[ \sum_{n':m+n'=n-3} \left( (\sin(2\eta))^{n'-1} \left( \prod_{1 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right)^2 \right] \right] \\
& \quad \times \left[ \frac{B_3(u') A_3(u)}{u' - u} + \frac{A_3(u') B_3(u)}{u' - u} \right] \\
& + 2 \left[ \frac{\partial}{\partial u'} \left[ A_3(u') + B_3(u') \right] \right] \left[ \sum_{1 \leq i \leq m} \left[ \frac{\partial}{\partial u} \left[ \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right] \right] \right] \\
& \quad - \left[ \frac{\partial}{\partial u} \left[ A_3(u') + B_3(u') \right] \right] \\
& \quad \times \left[ \sum_{1 \leq i \leq m} \left[ \frac{\partial}{\partial u'} \left[ \prod_{1 \leq i \leq m} \sin(u' - v_{n-i} \pm \eta \sigma_{n-i}^z) \right] \right] \right] ,
\end{aligned}$$

from which we conclude the argument.  $\square$

*Proof of Theorem 1.* The result immediately follows from the results for each Poisson bracket obtained in 2.4.1-2.4.9, from which we conclude the argument.  $\square$

*Proof of Theorem 2.* The result follows immediately from direct computation of each of the two Poisson brackets in canonical coordinates, from which we conclude the argument.  $\square$

#### 2.4.10 Overview of extending the computations with the Poisson bracket to the remaining fifteen relations

To exhibit how the computations performed for evaluating the nine Poisson brackets from the first relation can be extended to the remaining fifteen relations, we provide an outline of the terms involved in the second relation below. In particular, these terms are parametrized in different entries of the monodromy matrix, namely  $\mathcal{B}'_1$ ,  $\mathcal{B}'_2$ , and  $\mathcal{B}'_3$  instead of  $\mathcal{A}'_1$ ,  $\mathcal{A}'_2$ , and  $\mathcal{A}'_3$ , and can be expressed in terms of a superposition of nine terms, obtained from a superposition of thirty six Poisson brackets as provided for the first relation. The terms, excluding those which are dependent on  $\mathcal{A}_1$ ,  $\mathcal{A}_2$ ,  $\mathcal{A}_3$ ,  $\mathcal{B}'_1$ ,  $\mathcal{B}'_2$ , or  $\mathcal{B}'_3$ , appearing in the second relation are,

$$\begin{aligned}
\mathcal{P}_2(\sin(2\eta))^{n-3} \mathcal{B}'_1 & \equiv \mathcal{P}_2(\sin(2\eta))^{n-3} \left( \prod_{2 \leq i \leq n-(i-3)} \sigma_{n-i}^{-,+} \right) , \\
\mathcal{P}_1 \mathcal{B}'_2 & \equiv \mathcal{P}_1 \left( \prod_{2 \leq i \leq n-(i-3)} \sin(\lambda_\alpha - v_{n-i} + \eta \sigma_{n-i}^z) \right) , \\
\mathcal{P}_2(\sin(2\eta))^{n-3} \mathcal{B}'_3 & = \mathcal{P}_2(\sin(2\eta))^{n-3} \sum_{\substack{2 \leq i \leq m \\ 2 \leq j \leq n' \\ m+n'=n-(i-3)}} \left[ \left( \prod_{2 \leq i \leq m} \sin(\lambda_\alpha - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) \right. \\
& \quad \times \left. (\sin(2\eta))^{n'-1} \left( \prod_{2 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] , \\
\mathcal{P}_2(\sin(2\eta))^{n-3} \mathcal{B}'_1 & \equiv \mathcal{P}_2(\sin(2\eta))^{n-3} \left( \prod_{2 \leq i \leq n-(i-3)} \sigma_{n-i}^{-,+} \right) ,
\end{aligned}$$

$$\begin{aligned}
\mathcal{P}_2 \mathcal{B}'_2 &\equiv \mathcal{P}_2 \left( \prod_{2 \leq i \leq n-(i-3)} \sin(\lambda_\alpha - v_{n-i} + \eta \sigma_{n-i}^z) \right) , \\
\mathcal{P}_2 \mathcal{B}'_3 &\equiv \mathcal{P}_2 \sum_{\substack{2 \leq i \leq m \\ 2 \leq j \leq n' \\ m+n'=n-(i-3)}} \left[ \left( \prod_{2 \leq i \leq m} \sin(\lambda_\alpha - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{2 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] , \\
\mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{B}'_1 &\equiv \mathcal{P}_2 (\sin(2\eta))^{n-3} \left( \prod_{2 \leq i \leq n-(i-3)} \sigma_{n-i}^{-,+} \right) , \\
\mathcal{P}_2 \mathcal{B}'_2 &\equiv \mathcal{P}_2 \left( \prod_{2 \leq i \leq n-(i-3)} \sin(\lambda_\alpha - v_{n-i} + \eta \sigma_{n-i}^z) \right) , \\
\mathcal{P}_2 \mathcal{B}'_3 &\equiv \mathcal{P}_2 \sum_{\substack{2 \leq i \leq m \\ 2 \leq j \leq n' \\ m+n'=n-(i-3)}} \left[ \left( \prod_{2 \leq i \leq m} \sin(\lambda_\alpha - v_{n-i} \pm \eta \sigma_{n-i}^z) \right) (\sin(2\eta))^{n'-1} \left( \prod_{2 \leq j \leq n'} \sigma_{n-j}^{-,+} \right) \right] .
\end{aligned}$$

To compute each Poisson bracket appearing in (2), the second relation, one can directly apply previous arguments for computing the Poisson brackets for the first relation. For sake of not repeating similar computations with the Poisson bracket than those which were provided earlier in 2.4.1-2.4.9, one can obtain the relationships satisfied by each of group of nine Poisson brackets by reading off terms from the relation,

$$\begin{aligned}
&\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 (\sin(2\eta))^{n-3} \mathcal{A}_1, \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{B}'_1 \right\} + \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 (\sin(2\eta))^{n-3} \mathcal{A}_1, \mathcal{P}_2 \mathcal{B}'_2 \right\} \\
&+ \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 (\sin(2\eta))^{n-3} \mathcal{A}_1, \mathcal{P}_2 \mathcal{B}'_3 \right\} + \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_2, \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{B}'_1 \right\} \\
&+ \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_2, \mathcal{P}_2 \mathcal{B}'_2 \right\} + \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_2, \mathcal{P}_2 \mathcal{B}'_3 \right\} + \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_3, \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{B}'_1 \right\} \\
&+ \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_3, \mathcal{P}_2 \mathcal{B}'_2 \right\} + \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_3, \mathcal{P}_2 \mathcal{B}'_3 \right\} ,
\end{aligned}$$

for (2),

$$\begin{aligned}
&\sum_{\mathcal{P}} \left\{ \mathcal{P}_1 (\sin(2\eta))^{n-3} \mathcal{A}_1, \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{C}'_1 \right\} + \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 (\sin(2\eta))^{n-3} \mathcal{A}_1, \mathcal{P}_2 \mathcal{C}'_2 \right\} \\
&+ \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 (\sin(2\eta))^{n-3} \mathcal{A}_1, \mathcal{P}_2 \mathcal{C}'_3 \right\} + \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_2, \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{C}'_1 \right\} \\
&+ \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_2, \mathcal{P}_2 \mathcal{C}'_2 \right\} + \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_2, \mathcal{P}_2 \mathcal{C}'_3 \right\} + \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_3, \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{C}'_1 \right\} \\
&+ \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_3, \mathcal{P}_2 \mathcal{C}'_2 \right\} + \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{A}_3, \mathcal{P}_2 \mathcal{C}'_3 \right\} ,
\end{aligned}$$

for (3),







$$\begin{aligned}
& \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 (\sin(2\eta))^{n-3} \mathcal{D}_1, \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{D}'_1 \right\} + \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 (\sin(2\eta))^{n-3} \mathcal{D}_1, \mathcal{P}_2 \mathcal{D}'_2 \right\} \\
& + \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 (\sin(2\eta))^{n-3} \mathcal{D}_1, \mathcal{P}_2 \mathcal{D}'_3 \right\} + \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{D}_2, \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{D}'_1 \right\} \\
& + \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{D}_2, \mathcal{P}_2 \mathcal{D}'_2 \right\} + \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{D}_2, \mathcal{P}_2 \mathcal{D}'_3 \right\} + \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{D}_3, \mathcal{P}_2 (\sin(2\eta))^{n-3} \mathcal{D}'_1 \right\} \\
& \quad + \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{D}_3, \mathcal{P}_2 \mathcal{D}'_2 \right\} + \sum_{\mathcal{P}} \left\{ \mathcal{P}_1 \mathcal{D}_3, \mathcal{P}_2 \mathcal{D}'_3 \right\} ,
\end{aligned}$$

for (16).

### 3 References

- [1] Baxter, R.J. Exactly Solved Models in Statistical Mechanics. Dover Publications, London (1982).
- [2] Colomo, F., Giulio, G.D., Pronko, A.G. Six-vertex model on a finite lattice: Integral representations for nonlocal correlation functions. *Nuclear Physics B* **972**: 115535 (2021).
- [3] Duminil-Copin, H., Karrila, A., Manolescu, I., & Oulamara, M. Delocalization of the height function of the six-vertex model. arXiv: 2012.13750v1 (2020).
- [4] Duminil-Copin, H., Kozlowski, K.K., Krachun, D. et al. On the Six-Vertex Model's Free Energy. *Commun. Math. Phys.* **395**, 1383–1430 (2022). <https://doi.org/10.1007/s00220-022-04459-x>
- [5] Faddeev, L.D., Takhtajan, L. A. Hamiltonian Methods in the Theory of Solitons, Springer, 2007.
- [6] Glazman, A., Peled, R. On the transition between the disordered and antiferroelectric phases of the 6-vertex model. arXiv: 1909.03436 (2019).
- [7] Keating, D., Reshetikhin, N. & Sridhar, A. Integrability of Limit Shapes of the Inhomogeneous Six Vertex Model. *Commun. Math. Phys.* **391**, 1181–1207 (2022). <https://doi.org/10.1007/s00220-022-04334-9>
- [8] Lieb, E. H. The Residual Entropy of Square Ice, *Phys Rev.* **162** 162-172 (1967).
- [9] Pauling, L. J. Am. Chem. Soc. **57**, 2680 (1935).
- [10] Rigas, P. From logarithmic delocalization of the six-vertex height function under sloped boundary conditions to weakened crossing probability estimates for the Ashkin-Teller, generalized random-cluster, and  $(q_\sigma, q_\tau)$ -cubic models, *arXiv:2211.14934* (2022).