# Quest for the golden ratio universality class

Vladislav Popkov<sup>1,2</sup> and G.M. Schütz<sup>3</sup>

<sup>1</sup>Department of Physics, University of Wuppertal, Gaussstraße 20, 42119 Wuppertal, Germany <sup>2</sup>Faculty of Mathematics and Physics, University of Ljubljana, Jadranska 19, SI-1000 Ljubljana, Slovenia <sup>3</sup>Departamento de Matemática, Instituto Superior Técnico, Av. Rovisco Pais 1, 1049-001 Lisboa, Portugal

Using mode coupling theory the conditions for all allowed dynamical universality classes for the conserved modes in one-dimensional driven systems are presented in closed form as a function of the stationary currents and their derivatives. With a view on the search for the golden ratio universality class the existence of some families of microscopic models is ruled out a priori by using an Onsager-type macroscopic current symmetry. In particular, if the currents are symmetric or antisymmetric under the interchange of the conserved densities, then at equal mean densities the golden modes can only appear in the antisymmetric case and if the conserved quantities are correlated, but not in the symmetric case where at equal densities one mode is always diffusive and the second may be either Kardar-Parisi-Zhang (KPZ), modified KPZ, 3/2-Lévy, or also diffusive. We also show that the predictions of mode coupling theory for a noisy chain of harmonic oscillators are exact.

## I. INTRODUCTION

The golden ratio  $\varphi = (1 + \sqrt{5})/2 \approx 1.618$  is undoubtedly one of the most fascinating irrational numbers. It shows up in completely different circumstances, from in the arrangement of seeds in sunflowers or petals in flowers [1, 2], in the shape of snowflakes [3], the statistics of signals in the human brain [4] to art and architectural masterpieces [5]. From the mathematical viewpoint, the golden ratio is the "most" irrational number, in the sense that the number for which rational approximations like the continued fraction [6] perform the worst.

In the realm of physics the golden ratio is somewhat less prominent. In particular, in the theory of phase transitions – where *rational* numbers are ubiquitous in the characteristics of universality classes – specific irrational numbers have not been encountered until recently. Indeed, all critical indices coming from Landau theory of phase transitions [7] are rational numbers. Also the critical exponents produced in conformal field theories are all rational (unless continuously varying). Generically, since various scaling relations relating different critical exponents have the form of a ratio, the rationality of critical exponents seems to be a robust feature, leaving little room for irrationality.

It has therefore come as a surprise that the golden ratio can characterize universal fluctuations of long-lived relaxation modes of interacting many-body systems with conservation laws, governed by the dynamical exponent z, i.e., with stationary space-time correlations of the form

$$\langle \sigma(0,0)\sigma(x,t)\rangle \sim f\left(\frac{|x-vt|^z}{t}\right),$$
(1)

at large space and time scales. Here  $\sigma(x, t)$  is a conserved fluctuation field and v is its characteristic mode velocity. The theory of the Fibonacci universality classes [8], based on mode-coupling theory for nonlinear flucuating hydrodynamics, predicts the possibility of coexistence of modes with different dynamical exponents from the discrete infinite set of Kepler ratios  $z_k = C_{k+1}/C_k$  where  $C_k$  are the Fibonacci numbers  $1, 1, 2, 3, 5, \ldots$ . The ubiquitous diffusive Edwards-Wilkinson (EW) mode k = 2 [9] and the celebrated superdiffusive Kardar-Parisi-Zhang (KPZ) mode k = 3 [10] appear already in systems with one conservation law, while to reach the more exotic modes with  $k \ge 4$  at least k - 2 conservation laws (and hence relaxation modes) are necessary. The golden ratio  $\varphi = \lim_{k\to\infty} z_k = (1 + \sqrt{5})/2$  turns out to be a remarkable exception: This truly irrational dynamical exponent can be generated in (1) already in a system with just two long-lived relaxation modes.

With this knowledge in hand, and using fine tuning of interaction parameters we were able to demonstrate the existence of the irrational critical exponent  $\varphi$  in driven lattice gases with two and three conservation laws [8, 11]. Also in anharmonic chains with two conservation laws the golden universality class appears for fine-tuned parameter values [12]. However, ideally, one would like to have the means to produce a system from the remarkable golden universality class in a robust way, based on symmetries, instead of fine-tuning. Our present work presents a systematic study of properties of the interactions needed to generate one or another universality class, including the golden ratio, for a system with two conservation laws. In order to emphasize that under the given assumptions some of the results are mathematically rigorous, we use a mathematical presentation in terms of propositions, theorems and corollaries when appropriate.

## **II. FLUCTUATIONS IN SYSTEMS WITH LOCAL CONSERVATION LAWS**

To set the stage we describe the setting that we have in mind. In this introductory section we keep the number of conserved quantities arbitrary (but finite) and only point to the case of two conservation laws in some instances to facilitate the detailed discussion of two conservation laws in the subsequent sections.

#### A. Nonequilibrium steady state (NESS)

Consider a one-dimensional many-body dynamics with translation invariant short-range interactions with n locally conserved densities  $\rho_{\alpha}$ ,  $\alpha \in \{1, \ldots, n\}$  where a driving force produces macroscopic stationary currents denoted by  $j^{\alpha}$ . The resulting non-equilibrium steady state (NESS) is assumed to be unique for given values of the conserved densities, to have non-vanishing global fluctuations of the conserved densities, and exhibit a decay of local density correlations that is sufficiently fast to guarantee that the total stationary density fluctuations are finite. These properties are generic since in one space dimension stationary correlations between local observables usually decay rapidly with distance [13]. Indeed, for a single conservation law and translation invariant short-range interactions no counter examples to these properties are known except for some very special cases, viz., in systems with long-range interactions [14–16] or with facilitated dynamics [17–21]. For two conservation laws long-range correlations accompanied by phase separation have been reported and analysed numerically and by mean field theories [22–24]. All these atypical types of behaviour as well as frozen systems without density fluctuations are ruled out in the present investigation.

The total stationary density fluctuations are encoded in the compressibility matrix K in terms of the variances  $K_{\alpha\alpha}$  of the conserved quantities and their covariances  $K_{\alpha\beta}$ . Hence K is symmetric and all matrix elements are in general functions of all the conserved quantities  $\rho_{\alpha}$ . The cross correlations given by the covariances may be positive or negative or even vanish for all pairs of densities while the variances  $K_{\alpha\alpha}$  strictly positive real numbers (excluding the atypical situations mentioned above). Specifically, for two conservation laws we denote the matrix elements of the compressibility matrix K by

$$K = \begin{pmatrix} \kappa_1 & \bar{\kappa} \\ \bar{\kappa} & \kappa_2 \end{pmatrix}. \tag{2}$$

rather than by  $K_{\alpha\beta}$  and we usually omit the dependence of these quantities on  $\rho_1$  and  $\rho_2$ .

Also the stationary currents  $j^{\alpha}$  are in general functions of all densities  $\rho_{\alpha}$ . Their derivatives w.r.t. the densities are denoted by subscripts  $j^{\alpha}_{\beta}$ ,  $j^{\alpha}_{\beta\gamma}$ , and so on. The current Jacobian J has matrix elements  $J_{\alpha\beta} = j^{\alpha}_{\beta}$  and the Hessians  $H^{\alpha}$  associated with the current-density relation have matrix elements  $H^{\alpha}_{\beta\gamma} = j^{\alpha}_{\beta\gamma}$ . Thus for two conservation laws the current Jacobian and the Hessians are the matrices

$$J = \begin{pmatrix} j_1^1 & j_2^1 \\ j_1^2 & j_2^2 \end{pmatrix}, \quad H^{\alpha} = \begin{pmatrix} j_{11}^{\alpha} & j_{12}^{\alpha} \\ j_{21}^{\alpha} & j_{22}^{\alpha} \end{pmatrix}$$
(3)

with the dependence on  $\rho_1$  and  $\rho_2$  also usually omitted.

### B. Large scale dynamics

The long-time relaxation of the system is assumed to be determined by the long-wave length Fourier modes of the conserved quantities in the sense that at sufficiently large times the system is locally (on a microscopic scale) in a stationary state with densities  $\rho_{\alpha}$ , which vary only slowly with space and time.

The generic decay of correlations ensures bounded density fluctuations in the NESS and thus in driven nonequilibrium systems with non-vanishing stationary currents the validity of a coarse-grained hydrodynamic description of the conserved densities  $\rho_{\alpha}(x,t)$  in terms of a continuity equation  $\partial_t \rho_{\alpha}(x,t) + \partial_x J_{\alpha}(x,t) = 0$  where the currents  $J_{\alpha}(x,t)$ depend on macroscopic space and time only via the densities  $\rho_1(x,t), \ldots, \rho_n(x,t)$  [13, 25]. Thus the hydrodynamic limit is given by a system of hyperbolic conservation laws

$$\partial_t \rho_\alpha(x,t) + \sum_{\beta=1}^n J_{\alpha\beta} \partial_x \rho_\beta(x,t) = 0 \tag{4}$$

where the current Jacobian  $J_{\alpha\beta} = \partial_{\rho\beta} j^{\alpha}(\rho_1, \dots, \rho_n)$  is given by the stationary current-density relation  $j^{\alpha}(\rho_1, \dots, \rho_n)$  evaluated at the space-time point (x, t).

Fluctuations are not captured in this deterministic large-scale description. They are expected to be described by the theory of nonlinear fluctuating hydrodynamics [26, 27] for fluctuation fields which are centered around a fixed stationary mean  $\rho_{\alpha}$  of the conserved quantities. To analyze the large-scale behaviour of these fluctuations it is most convenient to express them as the eigenmodes  $\sigma_{\alpha}(x,t)$  that appear in generic form in (1) without mode index  $\alpha$ , i.e., as those linear combinations of the conserved fields for which the current Jacobian J is diagonal for the chosen stationary densities  $\rho_{\alpha}$ . According to nonlinear fluctuating hydrodynamics these eigenmodes then satisfy the system of coupled stochastic Burgers equations

$$\partial_t \sigma_\alpha + \partial_x \left( v_\alpha \sigma_\alpha + \sum_{\beta,\gamma=1}^2 G^\alpha_{\beta\gamma} \sigma_\beta \sigma_\gamma - \sum_{\beta=1}^2 D_{\alpha\beta} \partial_x \sigma_\beta + \xi_\alpha \right) = 0 \tag{5}$$

with the mode velocities  $v_{\alpha}$  which are the eigenvalues of the current Jacobian, with phenomenological diffusion matrix D and with Gaussian white noises  $\xi_{\alpha}$ . The strength of the nonlinearity is given by the mode coupling coefficients  $G^{\alpha}_{\beta\gamma}$  which depend primarily on the current Jacobian and also – in an insubstantial way – on the static compressibilities  $K_{\alpha\beta}$  (see below).

These nonlinear stochoastic partial differential equations (spde's) can be treated by mode coupling theory as described in [26]. With this approach we found in [8, 28] together with our collaborators the exact analytical scaling solution for the dynamical structure function (1) for all modes  $\alpha$ . This yields the Fibonacci dynamical universality classes as follows:

- In the absence of the diagonal mode coupling term, i.e.,  $G^{\alpha}_{\beta\beta} = 0$  for all modes  $\beta$  (including the mode  $\beta = \alpha$ ) the mode  $\alpha$  is diffusive and belongs to the Edwards-Wilkinson universality class with dynamical exponent  $z_{\alpha} = 2$ .
- For non-vanishing self-coupling term, i.e.,  $G^{\alpha}_{\alpha\alpha} \neq 0$ , the mode  $\alpha$  is superdiffusive and belongs to the Kardar-Parisi-Zhang universality class or a modification thereof, both with dynamical exponent  $z_{\alpha} = 3/2$ .
- If  $G^{\alpha}_{\alpha\alpha} = 0$  but  $G^{\alpha}_{\beta\beta} \neq 0$  for some mode  $\beta \neq \alpha$  then mode  $\alpha$  is in a subdiffusive Lévy universality class with the dynamical exponent  $z_{\alpha}$  being a Kepler ratio of two consecutive Fibonacci numbers or the golden ratio.
- The non-diagonal mode coupling terms  $G^{\alpha}_{\beta\gamma}$  with  $\beta \neq \gamma$  are irrelevant for these scaling properties.

The results of [8, 28] for the scaling functions are valid for strict hyperbolicity of the underlying deterministic pde (4), i.e, when the eigenvalues of the current Jacobian are non-degenerate.

It is worth stressing that the mode coupling coefficients  $G^{\alpha}_{\beta\beta}$  appearing in the macroscopic phenomenological spde (4) are fully given by the stationary distribution of the underlying dynamics, viz., by the current Jacobian J and by the static compressibility matrix K. In fact, whether  $G^{\alpha}_{\beta\beta}$  vanishes or not is determined by the current Jacobian J alone, the compressibilities encoded in K only renormalize the amplitudes. This implies the very remarkable fact that all the *dynamical* universality classes of a system can be read off from the *stationary* currents alone. An explicit example is provided in the Appendix B.

Specifically for two conservation laws, treated independently in [11] and [12] in the same issue of J. Stat. Phys., the elusive golden ratio occurs if and only if  $G_{11}^1 = G_{22}^2 = 0$  and  $G_{22}^1 \neq 0$  and  $G_{11}^2 \neq 0$ . In this case both modes belong to the golden universality class with  $z_1 = z_2 = \varphi$ . However, the question that has been left open and which is addressed here are the circumstances under which these conditions may be satisfied or not, i.e., which current-density relations that arise from some microscopic dynamics admit or forbid the occurrence of two golden modes.

## C. Onsager-type current symmetry and microscopic dynamics

Before attacking this problem directly we point out a further general consequence of the generic assumption of sufficiently rapidly decaying stationary correlations. This is the Onsager-type current symmetry

$$JK = KJ^T \tag{6}$$

proved rigorously first for a family of Markovian particle systems with fully factorized stationary product measure in [29] and later generally under very mild assumptions concerning the decay of stationary correlations and range of microscopic interactions for classical dynamics in [30] and subsequently for quantum systems in [31, 32]. This current symmetry appears in many contexts in hydrodynamic theory, see e.g. [26, 27] for a review and [33–35] for recent applications in generalized hydrodynamics for integrable quantum systems. As pointed out in [29] the symmetry (6) can be seen as a non-equilibrium version of the Onsager reciprocity relations in so far as its validity relies only on general time-reversal properties of the time evolution and on an extremely mild assumption on the decay of correlations as discussed in the rigorous proofs of [30, 32]. Indeed, the proof of (6) follows arguments analogous to those for the Onsager relations. We shall call one-dimensional physical systems with these generic properties *regular one-dimensional hydrodynamic systems*.

Remarkably and perhaps also surprisingly, this *macroscopic* current symmetry directly imposes constraints on the existence of specific *microscopic* dynamics: For a given family of invariant measures (parametrized by the conserved densities  $\rho_{\alpha}$  which fix K) the only physically permissible microscopic dynamics are such that the transition rates yield currents (which fix J) compatible with (6).

To appreciate the significance of this constraint in the search for physical systems that exhibit the golden universality class (or any other specific Fibonacci universality class) one must recall that for predicting universality classes for some microscopic dynamics one needs to know which diagonal mode coupling matrix elements  $G^{\alpha}_{\beta\beta}$  vanish. This requires knowledge of the *exact* stationary current-density relation, which in turn requires knowledge of the exact invariant measure of the dynamics. However, away from thermal equilibrium detailed balance does not hold, and there is no simple recipe for obtaining the exact invariant measure for a given dynamics. Therefore, computing the invariant invariant for some "nice" physical dynamics is usually a hopeless endeavor.

Instead, in the search for specific universality classes one usually works in opposite direction:

(i) One starts by defining a measure with a simple structure (such as product or matrix product measures [36, 37]) that allows for computing expectation values like the stationary currents in explicit form as functions of the densities,
(ii) proposes a physically motivated dynamics (such as short-range interactions) that respects the conservation laws,
(iii) proves invariance of the measure w.r.t. this dynamics and/or adjusts parameters accordingly,

(iv) computes stationary currents and compressibilities from (i) and (ii).

However, there is no obvious starting point for guessing such a hypothetical dynamics (which might not even exist) and then checking invariance w.r.t. the measure may be cumbersome due to the absence of detailed balance. Therefore, being able to rule out *a priori* certain microscopic dynamics as unphysical for a given invariant measure can be useful in the search for dynamical universality classes.

Here the current symmetry helps. It does not provide a sufficient condition for a microscopic dynamics to have the specified measure as invariant measure, but a necessary condition that can be used to rule out a proposed dynamics. This property can be utilized by taking step (iv) *before* step (iii): If step (iv) yields a current-density relation that is incompatible with (6) then there is no need to go through the potentially daunting task of step (iii). Unlike step (iii) (which may actually be an unsoluble problem), step (iv) is easy since the measure has been proposed in step (i) so as to facilitate the computation of the stationary currents.

To illustrate the point we focus on two conservation laws and note that the current symmetry (6) then becomes

$$J_{21}\kappa_1 - J_{12}\kappa_2 = (J_{11} - J_{22})\bar{\kappa}.$$
(7)

Evidently, this relation constitutes a necessary condition for the existence of microscopic dynamics that would be compatible with a predefined invariant measure – without the need to perform an explicit check of invariance of the measure under the proposed dynamics. Only the static compressibilities and the stationary currents of the model need to be computed using the invariant measure.

This is highlighted in the following theorem for partially factorized invariant measures of the form  $\pi_{\rho_1,\rho_2} = \pi_{\rho_1}^1 \pi_{\rho_2}^2$ where  $\pi_{\rho_{\alpha}}^{\alpha}$  are measures that do not necessarily factorize further such as matrix product measures. Frequently studied models of this type are two-lane systems [38–40] where particles of species  $\alpha$  jump on along a one-dimensional lattice called lane  $\alpha$  like in a system with only one conservation law, but with a rate that depends on the particle configuration on the other lane.

**Theorem II.1** Any regular hydrodynamic two-component system with a partially factorized invariant measure

$$\pi_{\rho_1,\rho_2} = \pi_{\rho_1}^1 \pi_{\rho_2}^2 \tag{8}$$

has stationary currents of the form

$$j^{1}(\rho_{1},\rho_{2}) = f_{1}(\rho_{1}) + \kappa_{1}(\rho_{1}) \int^{\rho_{2}} \mathrm{d}y \, g(\rho_{1},y), \tag{9}$$

$$j^{2}(\rho_{1},\rho_{2}) = f_{2}(\rho_{2}) + \kappa_{2}(\rho_{2}) \int^{\rho_{1}} \mathrm{d}x \, g(x,\rho_{2}), \tag{10}$$

with the compressibilities  $\kappa_{\alpha}(x) \in \mathbb{R}^+$  for  $x \in \mathbb{R}$ , functions  $f_{\alpha} : \mathbb{R} \to \mathbb{R}$  and interaction kernel  $g(x, y) \in \mathbb{R}$  for  $(x, y) \in \mathbb{R}^2$  that is the same for both currents. Moreover, for any pair of densities  $(\rho_1, \rho_2)$  the off-diagonal elements of the current Jacobian either both vanish or have equal sign.

5

*Proof:* The lower integration limits in (9) and (10) do not need to be specified as they can be absorbed into the functions  $f_{\alpha}(\cdot)$ . By writing the currents in a generic fashion as

$$j^{1}(\rho_{1},\rho_{2}) = f_{1}(\rho_{1}) + \kappa_{1}(\rho_{1}) \int^{\rho_{2}} dy \, g_{1}(\rho_{1},y)$$
$$j^{2}(\rho_{1},\rho_{2}) = f_{2}(\rho_{2}) + \kappa_{2}(\rho_{2}) \int^{\rho_{1}} dx \, g_{2}(x,\rho_{2})$$

in terms of functions  $f_{\alpha}(\cdot), g_{\alpha}(\cdot, \cdot)$ , yields

$$J_{12}(\rho_1, \rho_2) = \kappa_1(\rho_1)g_1(\rho_1, \rho_2), \quad J_{21}(\rho_1, \rho_2) = \kappa_2(\rho_2)g_2(\rho_1, \rho_2).$$

For a factorized invariant measure one has  $\bar{\kappa} = \partial_1 \kappa_2 = \partial_2 \kappa_1 = 0$  so that the current symmetry reduces to

$$J_{21}(\rho_1, \rho_2)\kappa_1(\rho_1) = J_{12}(\rho_1, \rho_2)\kappa_2(\rho_2).$$
(11)

Since  $\kappa_{\alpha}(\cdot) > 0$  by assumption, this implies  $g_2(\rho_1, \rho_2) = g_1(\rho_1, \rho_2)$  for all  $(\rho_1, \rho_2) \in \mathbb{R}^2$  and proves also the assertion regarding the off-diagonal elements of the current Jacobian. 

For a monotone increase or decrease of the current  $j^{\alpha}$  as a function of the other density  $\rho_{\beta}$  with  $\beta \neq \alpha$  as in the models of [38–40] and also assuming the existence of symmetries between the densities the current symmetry II.1 provides the following rather general no-go corollary.

Corollary II.2 A regular one-dimensional hydrodynamic two-component system with the three properties (i) partially factorized invariant measure  $\pi_{\rho_1,\rho_2} = \pi_{\rho_1}^1 \pi_{\rho_2}^2$ (ii) antisymmetric currents  $j^2(\rho_1,\rho_2) = -j^1(\rho_2,\rho_1)$ (iii) monotonicity  $j_2^1 > 0$  and  $j_1^2 > 0$  (or  $j_2^1 < 0$  and  $j_1^2 < 0$ ) for all densities  $\rho_1,\rho_2$ 

does not exist.

This holds since on the one hand monotonicity for all  $(\rho_1, \rho_2)$  requires that  $g(\cdot, \cdot)$  does not change sign while on the other hand the antisymmetry requires  $g(\rho_1, \rho_2)\kappa_2(\rho_2) = -g(\rho_2, \rho_1)\kappa_1(\rho_2)$  which is contradictory since  $\kappa_1$  and  $\kappa_2$  are both strictly positive. How the current symmetry rules out specific microscopic dynamics for a product measure is demonstrated for a concrete example in the Appendix A.

#### DYNAMICAL UNIVERSALITY CLASSES FOR TWO CONSERVATION LAWS III.

In the remainder of this paper we restrict ourselves to two conservation laws and discuss criteria for the occurrence of the golden and other universality classes. To this end, the mode coupling coefficients  $G^{\alpha}_{\beta\beta}$  are computed in terms of the matrix elements of K and J. Related results of [11, 12] do not allow for immediate predictions from the stationary currents for a given particle system since in these works the mode coupling matrices are not directly expressed in term of the stationary currents. This is achieved below.

#### Eigenmodes Α.

The system of stochastic Burgers equations (5) is expressed in terms of eigenmodes for which the current Jacobian J is diagonal. For the diagonalization of J we use as parameters the reduced trace  $\theta$ , the asymmetry  $\omega$ , and the signed square root  $\delta$  of the discriminant which are defined by

$$\theta := J_{11} - J_{22}, \quad \omega := \frac{J_{12}}{J_{21}}, \quad \delta := \begin{cases} \theta \sqrt{1 + \frac{4J_{12}J_{21}}{\theta^2}} & \theta \neq 0\\ \sqrt{4J_{12}J_{21}} & \theta = 0. \end{cases}$$
(12)

Only current Jacobians with non-degenerate eigenvalues

$$\lambda_{\pm} = \frac{1}{2} \left( J_{11} + J_{22} \pm \delta \right), \tag{13}$$

i.e., with non-zero discriminant, are considered below which ensures a strictly hyperbolic conservation law (4).

The following proposition concerning the diagonalization of a  $2 \times 2$  matrix is high school math. It serves to introduce notation for the eigenmodes and some of their basic properties.

**Proposition III.1** Let  $j^{\alpha}$  be the stationary currents of a two-component system satisfying the current symmetry (6) with compressibility matrix K (2) with strictly positive diagonal elements and current Jacobian J with nonvanishing discriminant. With the functions

$$u_{+} := \frac{\delta - \theta}{2J_{21}}, \quad u_{-} := -\frac{u_{+}}{\omega}, \quad v := 1 - u_{+}u_{-},$$
(14)

the normalization factors

$$z_{+}^{2} := \kappa_{1} + 2u_{+}\bar{\kappa} + u_{+}^{2}\kappa_{2}, \quad z_{-}^{2} := u_{-}^{2}\kappa_{1} + 2u_{-}\bar{\kappa} + \kappa_{2}, \tag{15}$$

and the matrix elements

$$u_1^+ := \frac{1}{z_+}, \quad u_2^+ := \frac{1}{z_+}u_+, \quad u_1^- := \frac{1}{z_-}u_-, \quad u_2^- := \frac{1}{z_-}$$
(16)

$$v_1^+ := \frac{z_+}{v}, \quad v_2^+ := -\frac{z_+}{v}u_-, \quad v_1^- := -\frac{z_-}{v}u_+, \quad v_2^- := \frac{z_-}{v}$$
 (17)

the matrices

$$R := \begin{pmatrix} u_1^+ & u_2^+ \\ u_1^- & u_2^- \end{pmatrix}, \quad R^{-1} := \begin{pmatrix} v_1^+ & v_1^- \\ v_2^+ & v_2^- \end{pmatrix}$$
(18)

satisfy

$$1 = RR^{-1} = RKR^{T}, \quad RJR^{-1} = \begin{pmatrix} \lambda_{+} & 0\\ 0 & \lambda_{-} \end{pmatrix}$$
(19)

where 1 is the two-dimensional unit matrix.

*Proof:* The matrices  $R, R^{-1}$  read in terms of the quantities  $z_{\pm}, u_{\pm}$ , and v

$$R = \begin{pmatrix} z_{+}^{-1} & z_{+}^{-1}u_{+} \\ z_{-}^{-1}u_{-} & z_{-}^{-1} \end{pmatrix}, \quad R^{-1} = v^{-1} \begin{pmatrix} z_{+} & -z_{-}u_{+} \\ -z_{+}u_{-} & z_{-} \end{pmatrix}.$$

The definition (14) of v yields immediately the first equality in (19). By the definitions (15) of the normalization factors the diagonal elements of  $RKR^T$  are equal to 1. The function  $u_+$  satisfies the quadratic equation

$$J_{21}u_+ - J_{12}u_+^{-1} + \theta = 0$$

which implies the two equalities

$$1 + u_+ u_- = \frac{\theta u_+}{J_{12}}, \quad 1 - u_+ u_- = \frac{J_{12}}{\delta u_+}.$$

The first equality together with the current symmetry (37) yields  $(RKR^T)_{12} = (RKR^T)_{21} = 0$ . The second equality proves the diagonalization of J.

Corollary III.2 The column and row vectors

$$\vec{v}_{\pm} := \begin{pmatrix} v_1^{\pm} \\ v_2^{\pm} \end{pmatrix}, \quad \vec{u}_{\pm} := \begin{pmatrix} u_1^{\pm}, u_2^{\pm} \end{pmatrix}$$

are right and left eigenvectors of J with eigenvalues  $\lambda_{\pm}$  and they satisfy the biorthogonality relation

$$\vec{u}_s \cdot \vec{v}_{s'} := u_1^s v_1^{s'} + u_2^s v_2^{s'} = \delta_{s,s'}.$$

for  $s, s' \in \{+, -\}$ .

Since  $\delta > 0$  by assumption, the current Jacobian is diagonalizable for all permissible parameter values. In particular, the tridiagonal case  $J_{12}J_{21} = 0$  is covered by the proposition by taking appropriate limits: For  $J_{12} = 0$  one gets  $\delta = \theta$ ,  $u_+ = 0$ ,  $u_- = -J_{21}/\theta$ , while for  $J_{21} = 0$  one has  $\delta = -\theta$ ,  $u_+ = -J_{12}/\theta$ ,  $u_- = 0$ .

## B. Mode-coupling matrices

To detail the conditions that lead to the various dynamical universality classes predicted by mode coupling theory we express in explicit form the diagonal elements of the mode coupling matrices in terms of the stationary currents and compressibilities.

**Definition III.3** The mode-coupling matrices  $G^{\gamma}$  are the linear combinations

$$G^{\gamma} := \frac{1}{2} \sum_{\lambda} R_{\gamma\lambda} (R^{-1})^T H^{\lambda} R^{-1}$$
(20)

of transformed Hessians  $H^{\lambda}$ .

They satisfy  $G_{\alpha\beta}^{\gamma} = G_{\beta\alpha}^{\gamma}$  since the Hessians  $H^{\lambda}$  are symmetric by construction. It is also possible to express the Hessians in terms of the mode-coupling matrices, see Appendix C.

According to mode coupling theory only the diagonal elements of the mode coupling matrices determine the largescale behaviour of the fluctuations. Hence we ignore the off-diagonal elements and we also recall that the normalization factors  $z_{\pm}$  of the eigenvectors are non-zero for any permissible choice of parameters. Therefore the ratios

$$g := \frac{\sqrt{z_+ z_-}}{v^2}, \quad y := \sqrt{\frac{z_+}{z_-}}$$
 (21)

are non-zero and well-defined. The theorem below expresses the diagonal elements  $G^{\alpha}_{\beta\beta}$  of the mode coupling matrices in terms of the stationary currents and compressibilities. We stress that the compressibilities enter only the non-zero overall amplitude given in terms of g and y. Whether a matrix element vanishes or not is solely determined by the current-density relation via the Hessians and the unnormalized eigenvector components of the current Jacobian.

**Theorem III.4** The diagonal elements of the mode coupling matrices are given by

$$G_{11}^{1} = \frac{g}{2} y \left[ H_{11}^{1} + u_{+} H_{11}^{2} - 2u_{-} (H_{12}^{1} + u_{+} H_{12}^{2}) + u_{-}^{2} (H_{22}^{1} + u_{+} H_{22}^{2}) \right]$$
(22)

$$G_{22}^{1} = \frac{g}{2}y^{-3} \left[ u_{+}^{2} (H_{11}^{1} + u_{+}H_{11}^{2}) - 2u_{+} (H_{12}^{1} + u_{+}H_{12}^{2}) + H_{22}^{1} + u_{+}H_{22}^{2} \right]$$
(23)

$$G_{11}^{2} = \frac{g}{2}y^{3} \left[ u_{-}H_{11}^{1} + H_{11}^{2} - 2u_{-}(u_{-}H_{12}^{1} + H_{12}^{2}) + u_{-}^{2}(u_{-}H_{22}^{1} + H_{22}^{2}) \right]$$
(24)

$$G_{22}^{2} = \frac{g}{2}y^{-1} \left[ u_{+}^{2} (u_{-}H_{11}^{1} + H_{11}^{2}) - 2u_{+} (u_{-}H_{12}^{1} + H_{12}^{2}) + u_{-}H_{22}^{1} + H_{22}^{2} \right]$$
(25)

*Proof:* To compute  $G^{\gamma}$  we note that with the diagonal normalization matrix

$$Z = \left(\begin{array}{cc} z_+ & 0\\ 0 & z_- \end{array}\right)$$

and the rescaled diagonalization matrices

$$\tilde{R} = \begin{pmatrix} 1 & u_+ \\ u_- & 1 \end{pmatrix}, \quad \tilde{R}^{-1} = v^{-1} \begin{pmatrix} 1 & -u_+ \\ -u_- & 1 \end{pmatrix}$$

one has  $R = Z^{-1}\tilde{R}$ . Defining furthermore

$$S := v\tilde{R}^{-1} = \begin{pmatrix} 1 & -u_+ \\ -u_- & 1 \end{pmatrix}$$

yields the modified mode coupling matrix

$$\tilde{G}^{\gamma} := 2v^2 Z^{-1} G^{\gamma} Z^{-1} = \sum_{\lambda} R_{\gamma\lambda} S^T H^{\lambda} S$$

with matrix elements

$$\tilde{G}_{11}^{\gamma} = \frac{2v^2}{z_+^2} G_{11}^{\gamma}, \quad \tilde{G}_{22}^{\gamma} = \frac{2v^2}{z_-^2} G_{22}^{\gamma}$$

The matrix multiplication can now be performed in a straightforward fashion and yields the matrix elements

$$(S^T H^{\lambda} S)_{11} = H_{11}^{\lambda} - 2u_- H_{12}^{\lambda} + u_-^2 H_{22}^{\lambda} (S^T H^{\lambda} S)_{12} = (1 + u_+ u_-) H_{12}^{\lambda} - u_+ H_{11}^{\lambda} - u_- H_{22}^{\lambda} (S^T H^{\lambda} S)_{22} = H_{22}^{\lambda} - 2u_+ H_{12}^{\lambda} + u_+^2 H_{11}^{\lambda}$$

and  $(S^T H^{\lambda} S)_{21} = (S^T H^{\lambda} S)_{12}$  by symmetry. Using the definition (20) therefore gives

$$\begin{split} G^1_{11} &= \frac{z_+}{2v^2} \left[ H^1_{11} - 2u_- H^1_{12} + u^2_- H^1_{22} + u_+ \left( H^2_{11} - 2u_- H^2_{12} + u^2_- H^2_{22} \right) \right] \\ G^1_{22} &= \frac{z_-^2}{2v^2 z_+} \left[ H^1_{22} - 2u_+ H^1_{12} + u^2_+ H^1_{11} + u_+ \left( H^2_{22} - 2u_+ H^2_{12} + u^2_+ H^2_{11} \right) \right] \\ G^2_{11} &= \frac{z_+^2}{2v^2 z_-} \left[ u_- \left( H^1_{11} - 2u_- H^1_{12} + u^2_- H^2_{12} \right) + H^2_{11} - 2u_- H^2_{12} + u^2_- H^2_{22} \right] \\ G^2_{22} &= \frac{z_-}{2v^2} \left[ u_- \left( H^1_{22} - 2u_+ H^1_{12} + u^2_+ H^1_{11} \right) + H^2_{22} - 2u_+ H^2_{12} + u^2_+ H^2_{11} \right]. \end{split}$$

Regrouping terms yield (22) - (25).

## C. Scenarios for dynamical universality classes

Generically, i.e., for two vanishing self-coupling coefficients  $G_{11}^1 \neq 0$  and  $G_{22}^2 \neq 0$  mode coupling theory predicts two KPZ modes. To search for the golden universality class we consider the mode coupling scenarios with less than two KPZ modes in more detail. We note that  $u_+u_- \neq 1$  since  $v = 1 - u_+u_- \neq 0$  by construction. It follows that  $u_-^{-1} \neq u_+$ .

## 1. A) One vanishing self-coupling coefficient

The following cases are permitted by mode coupling theory.

(i) Mode 1 is KPZ and mode 2 is 5/3-Lévy: This requires  $G_{11}^1 \neq 0$ ,  $G_{22}^2 = 0$ ,  $G_{11}^2 \neq 0$ , with arbitrary  $G_{22}^1$ . Thus

$$2(H_{12}^{1} + u_{+}H_{12}^{2}) \neq u_{-}^{-1}(H_{11}^{1} + u_{+}H_{11}^{2}) + u_{-}(H_{22}^{1} + u_{+}H_{22}^{2}),$$
  

$$0 \neq \omega(u_{-}H_{11}^{1} + H_{11}^{2}) + (u_{-}H_{12}^{1} + H_{22}^{2}),$$
  

$$2(u_{-}H_{12}^{1} + H_{12}^{2}) = u_{+}(u_{-}H_{11}^{1} + H_{11}^{2}) + u_{+}^{-1}(u_{-}H_{22}^{1} + H_{22}^{2}).$$

(ii) Mode 1 is KPZ and mode 2 is diffusive: This requires  $G_{11}^1 \neq 0$ ,  $G_{22}^1 = G_{22}^2 = G_{11}^2 = 0$ . Thus

$$\begin{split} &H_{22}^{1}+u_{+}H_{22}^{2} \neq -\omega(H_{11}^{1}+u_{+}H_{11}^{2}), \\ &2(H_{12}^{1}+u_{+}H_{12}^{2}) = u_{+}(H_{11}^{1}+u_{+}H_{11}^{2})+u_{+}^{-1}(H_{22}^{1}+u_{+}H_{22}^{2}), \\ &u_{-}H_{22}^{1}+H_{22}^{2} = -\omega(u_{-}H_{11}^{1}+H_{11}^{2}), \\ &2(u_{-}H_{12}^{1}+H_{12}^{2}) = (u_{+}+u_{-}^{-1})(u_{-}H_{11}^{1}+H_{11}^{2}). \end{split}$$

(iii) Mode 1 is modified KPZ and mode 2 is diffusive: This requires  $G_{11}^1 \neq 0$ ,  $G_{22}^1 \neq 0$ , and  $G_{22}^2 = G_{11}^2 = 0$ . Thus

$$\begin{aligned} & 2(H_{12}^1 + u_+ H_{12}^2) \neq u_-^{-1}(H_{11}^1 + u_+ H_{11}^2) + u_-(H_{22}^1 + u_+ H_{22}^2), \\ & 2(H_{12}^1 + u_+ H_{12}^2) \neq u_+(H_{11}^1 + u_+ H_{11}^2) + u_+^{-1}(H_{22}^1 + u_+ H_{22}^2), \\ & u_- H_{22}^1 + H_{22}^2 = -\omega(u_- H_{11}^1 + H_{11}^2), \\ & 2(u_- H_{12}^1 + H_{12}^2) = (u_+ + u_-^{-1})(u_- H_{11}^1 + H_{11}^2). \end{aligned}$$

This requires  $G_{22}^2 = G_{11}^1 = 0$  and therefore

$$2(H_{12}^{1} + u_{+}H_{12}^{2}) = u_{-}^{-1}(H_{11}^{1} + u_{+}H_{11}^{2}) + u_{-}(H_{22}^{1} + u_{+}H_{22}^{2})$$
(26)

$$2(u_{-}H_{12}^{1} + H_{12}^{2}) = u_{+}(u_{-}H_{11}^{1} + H_{11}^{2}) + u_{+}^{-1}(u_{-}H_{22}^{1} + H_{22}^{2}).$$

$$(27)$$

The remaining diagonal mode coupling coefficients take the form

$$G_{22}^{1} = \frac{z_{-}^{2}}{2vz_{+}} \left[ \omega (H_{11}^{1} + u_{+}H_{11}^{2}) + (H_{22}^{1} + u_{+}H_{22}^{2}) \right]$$
(28)

$$G_{11}^2 = \frac{z_+^2}{2vz_-\omega} \left[ \omega(u_-H_{11}^1 + H_{11}^2) + (u_-H_{22}^1 + H_{22}^2) \right].$$
(29)

(i) Two modes are golden Lévy:

$$0 \neq \omega(H_{11}^1 + u_+ H_{11}^2) + (H_{22}^1 + u_+ H_{22}^2)$$
(30)

$$0 \neq \omega(u_{-}H_{11}^{1} + H_{11}^{2}) + (u_{-}H_{22}^{1} + H_{22}^{2}).$$
(31)

(ii) Mode 1 is 3/2-Lévy and mode 2 is EW:

$$0 \neq \omega(H_{11}^1 + u_+ H_{11}^2) + (H_{22}^1 + u_+ H_{22}^2)$$
(32)

$$0 = \omega(u_{-}H_{11}^{1} + H_{11}^{2}) + (u_{-}H_{22}^{1} + H_{22}^{2}).$$
(33)

(iii) Two modes are EW:

$$0 = \omega(H_{11}^1 + u_+ H_{11}^2) + (H_{22}^1 + u_+ H_{22}^2)$$
(34)

$$0 = \omega (u_{-}H_{11}^{1} + H_{11}^{2}) + (u_{-}H_{22}^{1} + H_{22}^{2}).$$
(35)

We conclude that the conditions (26), (27), (30), and (31) must be met for the occurrence of the golden ratio. In this case both modes are golden Lévy.

**Remark III.5** The current symmetry imposes the constraints

$$H_{12}^{1} \det (K) = H_{11}^{2} \kappa_{1}^{2} - H_{11}^{1} \kappa_{1} \bar{\kappa} + H_{22}^{1} \bar{\kappa} \kappa_{2} - H_{22}^{2} \bar{\kappa}^{2} + J_{21} (\kappa_{1} \partial_{1} - \bar{\kappa} \partial_{2}) \kappa_{1} - J_{12} (\kappa_{1} \partial_{1} - \bar{\kappa} \partial_{2}) \kappa_{2} - (J_{11} - J_{22}) (\kappa_{1} \partial_{1} - \bar{\kappa} \partial_{2}) \bar{\kappa}$$
(36)

and

$$H_{12}^{2} \det (K) = H_{11}^{2} \bar{\kappa} \kappa_{1} - H_{11}^{1} \bar{\kappa}^{2} + H_{22}^{2} \bar{\kappa}^{2} - H_{22}^{2} \bar{\kappa} \kappa_{2} + J_{12} (\kappa_{2} \partial_{2} - \bar{\kappa} \partial_{1}) \kappa_{2} - J_{21} (\kappa_{2} \partial_{2} - \bar{\kappa} \partial_{1}) \kappa_{1} + (J_{11} - J_{22}) (\kappa_{2} \partial_{2} - \bar{\kappa} \partial_{1}) \bar{\kappa}$$
(37)

on the matrix elements of the Hessians. Hence the diagonal elements (22) - (25) of the mode coupling matrix and therefore the mode scenarios discussed above can be expressed entirely in terms of J, K, and the diagonal elements of the Hessians rather than in terms of J, K, and the full Hessians.

## IV. SYMMETRIC OR ANTISYMMETRIC CURRENTS AT EQUAL DENSITIES

The golden universality class has been observed in [11, 12] on a special parameter manifold that requires fine-tuning of densities and/or interaction parameters. We investigate here whether it can occur in more generic circumstances, viz., for the symmetry property of the current-density relation  $j^2(\rho_1, \rho_2) = \pm j^1(\rho_2, \rho_1)$  at the point of equal densities.  $\rho_1 = \rho_2$  where

$$J_{11} = \pm J_{22} \tag{38}$$

$$J_{12} = \pm J_{21}. \tag{39}$$

These properties of J are sufficient to show which mode coupling coefficients vanish.

## A. Antisymmetric case

In the antisymmetric case one has for equal densities  $\theta \neq 0$  and  $\omega = -1$  which gives  $u_{-} = u_{+}$  and  $\lambda_{-} = -\lambda_{+}$ . The current symmetry implies  $\bar{\kappa} \neq 0$  which means that a microscopic physical dynamics with a partially factorized invariant measure of the form (8) and antisymmetric current-density-relation at equal densities does not exist.

However, non-vanishing cross correlations between the conserved densities are not ruled out. The Hessians then have the symmetry relations

$$H_{11}^2 = -H_{22}^1, \quad H_{12}^2 = -H_{12}^1, \quad H_{22}^2 = -H_{11}^1.$$

This yields the mode coupling coefficients

$$\begin{split} G_{11}^{1} &= \frac{u_{+}z_{+}}{2v^{2}} \left[ u_{+}^{-1}(1-u_{+}^{3})H_{11}^{1} - 2(1-u_{+})H_{12}^{1} - (1-u_{+})H_{22}^{1} \right] \\ G_{22}^{1} &= \frac{u_{+}z_{-}^{2}}{2v^{2}z_{+}} \left[ -(1-u_{+})H_{11}^{1} - 2(1-u_{+})H_{12}^{1} + u_{+}^{-1}(1-u_{+}^{3})H_{22}^{1} \right] \\ G_{11}^{2} &= -\left(\frac{z_{+}}{z_{-}}\right)^{3}G_{22}^{1} \\ G_{22}^{2} &= -\frac{z_{-}}{z_{+}}G_{11}^{1} \end{split}$$

Hence the following scenarios are possible:

(i) 2 KPZ modes:  $u_+ \neq 1$  and

$$u_{+}(H_{22}^{1} + 2H_{12}^{1}) \neq (1 + u_{+} + u_{+}^{2})H_{11}^{1}$$

$$\tag{40}$$

(ii) 2 golden modes:  $u_+ \neq 1$  and

$$u_{+}(H_{22}^{1} + 2H_{12}^{1}) = (1 + u_{+} + u_{+}^{2})H_{11}^{1}$$

$$(41)$$

$$H_{11}^1 \neq H_{22}^1 \tag{42}$$

(iii) 2 EW modes: All diagonal mode coupling coefficients vanish if either  $u_{+} = 1$  or

$$2u_{+}H_{12}^{1} = (1+u_{+}^{2})H_{11}^{1}$$
(43)

$$H_{11}^1 = H_{22}^1 \tag{44}$$

In all allowed cases both modes are the same. These four cases correspond to the diagonal of the mode coupling table of [11].

## B. Symmetric case

In the symmetric case the properties of the Jacobian correspond to  $\theta = 0$  and  $\omega = 1$  and therefore  $u_{\pm} = \pm 1$ . The non-degeneracy of the eigenvalues  $\lambda_{\pm}$  implies  $J_{12}J_{21} \neq 0$ . There is no conflict with the current symmetry. The Hessians have the symmetry relations

$$H_{11}^2 = H_{22}^1, \quad H_{12}^2 = H_{12}^1, \quad H_{22}^2 = H_{11}^1$$

From (22) - (25) one gets

$$\begin{aligned} G_{11}^{1} &= \frac{1}{4(z_{+}^{-1})} \left[ H_{11}^{1} + H_{22}^{1} + 2H_{12}^{1} \right], \\ G_{22}^{1} &= \frac{(z_{+}^{-1})}{4(z_{-}^{-1})^{2}} \left[ H_{11}^{1} + H_{22}^{1} - 2H_{12}^{1} \right], \\ G_{11}^{2} &= G_{22}^{2} = 0. \end{aligned}$$

Hence in all three cases mode 2 is diffusive, thus ruling out the golden mode. Mode 1 can be KPZ, mKPZ, 3/2L, or EW. These four cases correspond to the bottom row (rightmost column) of the mode coupling table of [11].

## V. CONCLUSIONS

So far the golden universality class for fluctuations in driven systems with two conservation laws has remained somewhat elusive. It was only found by fine-tuning of the conserved densities or interaction parameters which determine the form of the stationary current-density relations [11, 12]. In the quest for a natural occurrence that does not require fine-tuning we have shown that systems with currents that are symmetric at equal densities cannot exhibit golden modes. In fact, in this case at least one of the two mode is diffusive with dynamical exponent z = 2 and belongs to the EW universality whereas the other mode is KPZ, modified KPZ or 3/2-Lévy with dynamical exponent z = 3/2 in each case.

However, the golden modes can appear with systems with two conservation laws if the currents are antisymmetric at equal densities. In this case, both modes belong to the same universality class which may be KPZ, EW or golden. However, such systems cannot have invariant measures that factorize over the two conserved quantities. This property may explain the rarity of analytical and simulation results for the golden modes, as one often considers such systems to make them amenable to explicit computations. On the other hand, from a physical perspective such a factorization is highly exceptional which gives hope that the golden universality class might be less unusual as it appears at the present state of the art.

The nonexistence of physical systems with antisymmetric currents at equal densities and an invariant measure that factorizes over the conserved quantities derives from the current symmetry that arises like Onsager's reciprocity relations from time-reversal. Thus the current symmetry is not only of fundamental significance but also helps ruling out microscopic dynamics in the search for other modes without having to check whether an invariant measure that is amenable to exact computation of the current is invariant for a candidate microscopic dynamics. Potential applications include two-lane versions of zero-range processes [38] or KLS-models [41–44] which are exclusion processes with next-nearest-neighbour interactions with the invariant measure of a one-dimensional Ising model for each lane.

## ACKNOWLEDGMENTS

It is a pleasure to acknowledge stimulating discussions with P. Gonçalves. This work is financially supported by FCT/Portugal through CAMGSD, IST-ID, Projects UIDB/04459/2020 and UIDB/04459/2020 and by the FCT Grant 2022.09232.PTDC. V. P. acknowledges financial support from the Deutsche Forschungsgemeinschaft through DFG project KL 645/20-2, and thanks IST Lisbon for hospitality and support.

- [1] Livio, M. (2008) The Golden Ratio: The Story of Phi, the World's Most Astonishing Number, Broadway Books, New York.
- [2] Hyun-Woo Lee and Leonid S. Levitov, Universality in Phyllotaxis: a Mechanical Theory, published in: Symmetry in Plants, Series in Mathematical Biology and Medicine, eds. R. V. Jean, D. Barabé, World Scientific Pub Co Inc, 1998, also in https://arxiv.org/pdf/2101.02652.pdf
- [3] J. Kepler, De Nive sexangula (Godfrey Tampach, Frankfurt, 1611); English translation: J. Kepler, On the Six-Cornered Snowflake (Clarendon Press, Oxford, 1966)
- Belinda Pletzer, Hubert Kerschbaum, Wolfgang Klimesch, When frequencies never synchronize: The golden mean and the resting EEG, Brain Research, vol. 1335, 2010, Pages 91-102, https://doi.org/10.1016/j.brainres.2010.03.074
- [5] Gary B. Meisner and Rafael Araujo, The Golden Ratio: The Divine Beauty of Mathematics, Race Point Publishing, 2018
- [6] A.Y. Khinchin, Continued Fractions, 1949; (reprint: Dover Books on Mathematics, 1964)
- [7] L. D. Landau, On the theory of phase transitions, Zh. Eksp. Teor. Fiz. 7, pp. 19–32 (1937)
- [8] V. Popkov, A. Schadschneider, J. Schmidt, and G.M. Schütz, Fibonacci family of dynamical universality classes, Proc. Natl. Acad. Science (USA) 112 12645–12650 (2015).
- [9] J. Krug, Origins of scale invariance in growth processes. Adv. Phys. 46, 139–282 (1997).
- [10] T. Halpin-Healy, K.A. Takeuchi, A KPZ Cocktail-Shaken, not Stirred... J. Stat. Phys. 160, 794–814 (2015).
- [11] V. Popkov, J. Schmidt, and G.M. Schütz, Universality classes in two-component driven diffusive systems. J. Stat. Phys. 160, 835–860 (2015).
- [12] H. Spohn and G. Stoltz, Nonlinear fluctuating hydrodynamics in one dimension: The case of two conserved fields, J. Stat. Phys. 160, 861–884 (2015).
- [13] H. Spohn, Large Scale Dynamics of Interacting Particles. (Springer, Berlin, 1991)
- [14] Spohn, H.: Bosonization, vicinal surfaces, and hydrodynamic fluctuation theory. Phys. Rev. E 60, 6411–6420 (1999).
- [15] Popkov, V. and Schütz, G.M.: Transition Probabilities and Dynamic Structure Function in the ASEP Conditioned on Strong Flux, J. Stat. Phys. 142, 627–639 (2011).

- [16] Karevski, D., Schütz, G.M.: Conformal invariance in driven diffusive systems at high currents. Phys. Rev. Lett. 118, 030601 (2017)
- [17] D. Helbing, D. Mukamel, and G. M. Schütz, Phys. Rev. Lett. 82, 10–13 (1999).
- [18] M. Rossi, R. Pastor-Satorras, and A. Vespignani. Universality class of absorbing phase transitions with a conserved field. Phys. Rev. Lett. 85 1803–1806 (2000).
- [19] A. Gabel, P. L. Krapivsky, and S. Redner. Facilitated asymmetric exclusion. Phys. Rev. Lett. 105 210603 (2010).
- [20] O. Blondel, C. Erignoux, M. Sasada, and M. Simon. Hydrodynamic limit for a facilitated exclusion process. Annales de l'Institut Henri Poincaré, Probabilités et Statistiques 56, 667–714 (2020).
- [21] C. Erignoux and L. Zhao, Stationary fluctuations for the facilitated exclusion process, arXiv:2305.13853 [math.PR] (2023)
- [22] Ramaswamy, S., Barma, M., Das, D., Basu, A.: Phase Diagram of a Two-Species Lattice Model with a Linear Instability. Phase Transit. 75, 363–375 (2002)
- [23] Kafri, Y., Levine, E., Mukamel, D., Schütz, G.M., Willmann R.D.: Phase-separation transition in one-dimensional driven models. Phys. Rev. E 68, 035101(R) (2003)
- [24] Chakraborty, S., Pal, S., Chatterjee, S., Barma, M.: Large compact clusters and fast dynamics in coupled nonequilibrium systems. Phys. Rev. E 93, 050102(R) (2016)
- [25] Kipnis, C., Landim, C.: Scaling limits of interacting particle systems. Springer, Berlin (1999).
- [26] H. Spohn, Nonlinear Fluctuating hydrodynamics for anharmonic chains. J. Stat. Phys. 154, 1191–1227 (2014).
- [27] Spohn, H.: Fluctuating Hydrodynamics Approach to Equilibrium Time Correlations for Anharmonic Chains. In: S. Lepri (ed.), Thermal Transport in Low Dimensions. From Statistical Physics to Nanoscale Heat Transfer, Lecture Notes in Physics 921, 107–158 (Springer, Switzerland, 2016)
- [28] V. Popkov, A. Schadschneider, J. Schmidt, G.M. Schütz, Exact scaling solution of the mode coupling equations for nonlinear fluctuating hydrodynamics in one dimension, J. Stat. Mech. 093211 (2016).
- [29] B. Tóth and B. Valkó, J. Stat. Phys. **112**, 497–521 (2003).
- [30] R. Grisi and G.M. Schütz, Current symmetries for particle systems with several conservation laws, J. Stat. Phys. 145, 1499–1512 (2011).
- [31] Castro-Alvaredo, O. A., Doyon, B., Yoshimura, T.: Emergent Hydrodynamics in Integrable Quantum Systems Out of Equilibrium. Phys. Rev. X 6, 041065 (2016)
- [32] D. Karevski and G. M. Schütz, Charge-current correlation equalities for quantum systems far from equilibrium. SciPost Phys. 6, 068 (2019)
- [33] B. Doyon and H. Spohn, Drude Weight for the Lieb-Liniger Bose Gas, SciPost Phys. 3, 039 (2017).
- [34] V. Alba, B. Bertini, M., L. Piroli, P. Ruggiero, Generalized-hydrodynamic approach to inhomogeneous quenches: correlations, entanglement and quantum effects, J. Stat. Mech. (2021) 114004
- [35] J. De Nardis, B. Doyon, M. Medenjak, M. Panfil, Correlation functions and transport coefficients in generalised hydrodynamics, J. Stat. Mech. (2022) 014002
- [36] Blythe, R.A., Evans, M.R.: Nonequilibrium steady states of matrix-product form: a solver's guide. J. Phys. A: Math. Theor. 40 R333–R441 (2007)
- [37] L. Cantini and A. Zahra, Hydrodynamic behavior of the two-TASEP, J. Phys. A: Math. Theor. 55 305201 (2022)
- [38] Grosskinsky S, Schütz GM, Spohn H, Condensation in the zero range process: Stationary and dynamical properties J. Stat. Phys. 113 (3-4): 389-410 (2003).
- [39] V. Popkov and M. Salerno, Hydrodynamic limit of multichain driven diffusive models, Phys. Rev. E 69, 046103 (2004).
- [40] Schmidt, J., Schütz, G.M., and van Beijeren, H.: A lattice Gas Model for Generic One-Dimensional Hamiltonian Systems. J. Stat. Phys. 183, 8 (2021).
- [41] S. Katz, J. L. Lebowitz, and H. Spohn, Nonequilibrium steady states of stochastic lattice gas models of fast ionic conductors J. Stat. Phys. 34, 497–537 (1984)
- [42] G.M. Schütz, Diffusion-annihilation in the presence of a driving field. J. Phys. A: Math. Gen. 28, 3405–3415 (1995).
- [43] J.S. Hager, J. Krug, V. Popkov and G.M. Schütz, Phys. Rev. E 63, 056110 (2001).
- [44] M. Dierl, M. Einax, P. Maass, One-dimensional transport of interacting particles: Currents, density profiles, phase diagrams and symmetries Phys. Rev. E 87, 062126 (2013).
- [45] B. Derrida, An exactly soluble nonequilibrium system: The asymmetric simple exclusion process. Phys. Rep. 301, 65–83 (1998).
- [46] Liggett, T.M.: Stochastic Interacting Systems: Contact, Voter and Exclusion Processes. Springer, Berlin (1999)
- [47] Schütz, G.M.: Exactly solvable models for many-body systems far from equilibrium, in: Phase Transitions and Critical Phenomena Vol. 19, C. Domb and J. Lebowitz (eds.), 1–251, Academic Press, London (2001).
- [48] H. Guiol, Some Properties of k-Step Exclusion Processes, J. Stat. Phys. 94, 495–511 (1999).
- [49] M.V. Freire, H. Guiol, K. Ravishankar, and E. Saada, Microscopic structure of the k-step exclusion process, Bull. Braz. Math. Soc. 33, 319–339 (2002).
- [50] C. Bernardin and S. Olla, Non-equilibrium macroscopic dynamics of chains of anharmonic oscillators, http://cermics.enpc.fr/~stoltz/Olla/lecture\_1.pdf (2009)
- [51] C. Bernardin and G. Stoltz, Anomalous diffusion for a class of systems with two conserved quantities, Nonlinearity 25, 1099–1133 (2012).
- [52] C. Bernardin, P. Gonçalves, and M. Jara, 3/4-fractional superdiffusion in a system of harmonic oscillators perturbed by a conservative noise, Arch. Rational Mech. Anal. 220, 505–542 (2016).
- [53] After completing this computation, P. Gonçalves mentioned to us that the correctness of the scale factor  $C_0$  was communicated to her by H. Spohn some years ago. Since we are not aware of a published version of this observation we decided

to keep it in this manuscript, without claiming originality.

## Appendix A: Current symmetry and constraints on the microscopic dynamics

We illustrate how the current symmetry rules out microscopic dynamics for some two-lane exclusion processes. These driven lattice gases are defined on two one-dimensional lattices (that we call lanes), that are indexed by  $\alpha \in \{1,2\}$  and have L sites each. We view the two lanes as being arranged in parallel, with a pair of sites on the two lanes labeled by an integer k.

Particles jump along each lane (but not to the other) obeying the exclusion rule, i.e., a jump attempt fails if the target site is already occupied. Thus the state of each lattice site k is described by a pair of occupation numbers  $n_k^k \in \{0, 1\}$ , indicating whether lane  $\alpha$  is occupied by a particle or not.

To be definite, we look for jump dynamics such that the NESS has a particularly simple form given by an invariant measure that is a product measure factorizing over all sites of the two lattices, i.e.,

$$\pi_{\rho_1,\rho_2}(n_1^1,\dots,n_L^1,n_1^2,\dots,n_L^2) = \prod_{\alpha=1}^2 \prod_{k=1}^L \left[ (1-\rho_\alpha)(1-n_k^\alpha) + \rho_\alpha n_k^\alpha \right]$$
(A1)

with average density  $\rho_{\alpha}$  on each lane, compressibilities  $\kappa_{\alpha} = \rho_{\alpha}(1 - \rho_{\alpha})$  and no correlations between different sites which implies  $\bar{\kappa} = 0$ . Well-studied examples for single-lane dynamics with these properties dynamics include the paradigmatic asymmetric simple exclusion process [45–47], or the k-step exclusion processes with long-range jumps [48, 49], to mention just a few.

We denote the stationary currents arising from these dynamics in the absence of any coupling between the lanes as  $j_0^{\alpha}(\rho_{\alpha})$ . It is well-known that fluctuations in such systems generically belong to the KPZ universality class [10]. To study non-trivial coupled systems that may exhibit other fluctuation patterns we allow the transition rates in each lane to depend on the occupation numbers of the other lane and ask whether such models with the invariant product measure (A1) can exist.

To be specific, we consider three cases:

(i) Totally asymmetric dynamics with equal dynamics on both lanes such that the rate for a forward jump from site k to a site k + r on lane 1 is enhanced by the number of particles on sites k and k + r on lane 2 and the rate of a backward jump from site k to a site k - r on lane 2 is enhanced by the number of particles on sites k and k - r.

(ii) Totally asymmetric dynamics with equal dynamics on both lanes but opposite directionality such that the rate for a forward jump from site k to a site k + r on lane 1 is enhanced by the number of particles on sites k and k + r on lane 2 and the rate of a backward jump from site k to a site k - r on lane 2 is enhanced by the number of particles on sites k and k - r.

(iii) Facilitated totally asymmetric dynamics, with parallel directionality on both lanes and rates such that a forward jump from site k to a site k + r on lane  $\alpha$  is enhanced by a factor  $b_{\alpha}$  when a site k' is occupied on both lanes (and k' different from k and the target site k + r).

In case (i) the factorization of the invariant measure yields the exact currents

$$j^{1}(\rho_{1},\rho_{2}) = j_{0}(\rho_{1})(1+2b_{1}\rho_{2}), \quad j^{2}(\rho_{1},\rho_{2}) = j_{0}(\rho_{2})(1+2b_{2}\rho_{1})$$
(A2)

and therefore  $J_{12}(\rho_1, \rho_2) = 2b_1 j_0^1(\rho_1)$  and  $J_{21}(\rho_1, \rho_2) = 2b_2 j_0^2(\rho_2)$ . For such dynamics the current symmetry (11) requires

$$b_2 j_0(\rho_2) \rho_1(1-\rho_1) = b_1 j_0(\rho_1) \rho_2(1-\rho_2).$$
(A3)

Therefore such dynamics exist if and only if  $b_1 = b_2$  and  $j_0(x) = cx(1-x)$  as in the conventional TASEP. For r = 1 this is the two-lane model of [39]. Once still needs to prove that for  $b_1 = b_2$  the product measure (A1) is invariant (as is done in [39]) but there is no reason to check invariance in the more general case  $b_1 \neq b_2$ : The macroscopic current symmetry asserts that such a microscopic model does not exist.

In case (ii) the factorization of the invariant measure yields the exact currents

$$j^{1}(\rho_{1},\rho_{2}) = j_{0}(\rho_{1})(1+2b_{1}\rho_{2}), \quad j^{2}(\rho_{1},\rho_{2}) = -j_{0}(\rho_{2})(1+2b_{2}\rho_{1})$$
(A4)

and therefore  $J_{12}(\rho_1, \rho_2) = 2b_1 j_0^1(\rho_1)$  and  $J_{21}(\rho_1, \rho_2) = -2b_2 j_0^2(\rho_2)$ . For such dynamics to exist the current symmetry (11) requires

$$b_2 j_0(\rho_2) \rho_1(1-\rho_1) = -b_1 j_0(\rho_1) \rho_2(1-\rho_2).$$
(A5)

Therefore such dynamics exist if and only if  $b_1 = -b_2$  and  $j_0(x) = cx(1-x)$  as in the conventional TASEP. Hence a model with current enhancement on both lanes due the presence of particles on the other lane need not be considered. The macroscopic current symmetry asserts that such a microscopic model does not exist.

In case (iii) the factorization of the invariant measure yields the exact currents

$$j^{1}(\rho_{1},\rho_{2}) = j^{1}_{0}(\rho_{1})(1+b_{1}\rho_{1}\rho_{2}), \quad j^{2}(\rho_{1},\rho_{2}) = j^{2}_{0}(\rho_{2})(1+b_{2}\rho_{1}\rho_{2})$$
(A6)

Thus  $J_{12}(\rho_1, \rho_2) = b_1 \rho_1 j_0^1(\rho_1)$  and  $J_{21}(\rho_1, \rho_2) = b_2 \rho_2 j_0^2(\rho_2)$ . For such dynamics to exist the current symmetry (11) requires

$$b_2 j_0^2(\rho_2)(1-\rho_1) = b_1 j_0^1(\rho_1)(1-\rho_2).$$
(A7)

This is valid for all  $(\rho_1, \rho_2)$  if and only if  $b_1 = b_2 = 0$ , i.e., no such coupled facilitated dynamics for which the fully factorized measure is invariant exists.

#### Appendix B: On the reliability of mode coupling theory for nonlinear fluctuating hydrodynamics

Consider an infinite chain of particles of unit mass at positions  $q_j$  interacting through a pair potential  $V(r_j)$  where  $r_j = q_j - q_{j-1}$  is interparticle distance. With the momentum  $p_j$  of the particles the total energy is given by

$$E = \sum_{j \in \mathbb{Z}} \left[ \frac{1}{2} p_j^2 + V(r_j) \right]$$
(B1)

and the deterministic Hamiltonian dynamics for such a system are given by

$$\frac{\mathrm{d}}{\mathrm{d}t}r_j(t) = p_j(t) - p_{j-1}(t) \tag{B2}$$

$$\frac{\mathrm{d}}{\mathrm{d}t}p_j(t) = V'(r_{j+1}(t)) - V'(r_j(t)).$$
(B3)

Evidently, besides the energy E also the total momentum P and "volume" V, i.e., the quantities

$$P = \sum_{j \in \mathbb{Z}} p_j, \quad V = \sum_{j \in \mathbb{Z}} r_j \tag{B4}$$

are conserves under the dynamics.

For the existence of such dynamics for general potentials see [50]. In the special case of harmonic oscillators with  $V(x) = x^2/2$  the system is completely integrable. This is most easily seen by introducing the variables

$$\eta_{2j-1}(t) := r_j(t), \quad \eta_{2j}(t) := p_j(t).$$
 (B5)

Then the two coupled equations (B2) and (B3) can be expressed as the single linear equation

$$\frac{\mathrm{d}}{\mathrm{d}t}\eta_j(t) = \eta_{j+1}(t) - \eta_{j-1}(t) \tag{B6}$$

which is explicitly solvable by Fourier transformation.

To introduce noise an exchange of the variables  $\eta_j(t)$ ,  $\eta_{j+1}(t)$  at exponential random times was introduced in [51] for more general potentials and studied in detail for the harmonic case in [52]. This random exchange violates momentum conservation but leaves the total energy E and the pseudo volume W, expressed in terms of the variables  $\eta_j$  as

$$E = \sum_{j \in \mathbb{Z}} \eta_j^2, \quad W = \sum_{j \in \mathbb{Z}} \eta_j, \tag{B7}$$

are conserved.

To write the generator of the stochastic dynamics for this randomized chain of harmonic oscillators we denote by  $\boldsymbol{\eta} := (\dots, \eta_{j-1}, \eta_j, \eta_{j+1}, \dots)$  the infinite set of variables  $\eta_j \in \mathbb{R}$  and introduce the swapped state variable  $\boldsymbol{\eta}^{kk+1}$  by the local states variables

$$\eta_l^{kk+1} = \eta_l + (\eta_{l+1} - \eta_l)\delta_{l,k} + (\eta_{l-1} - \eta_l)\delta_{l,k+1}.$$
(B8)

which the state reached from  $\eta$  after an exchange of  $\eta_k$  and  $\eta_{k+1}$ . The deterministic part of the generator that yields the Hamiltonian dynamics (B6) is denoted by  $\mathcal{A}$  and the generator for the random exchange is denoted by  $\mathcal{S}$ . The full generator  $\mathcal{L}$  acting on measurable functions  $f(\cdot)$  is then given by  $\mathcal{L} = \mathcal{S} + \mathcal{A}$  with

$$\mathcal{S}f(\eta) = \sum_{k \in \mathbb{Z}} [f(\eta^{kk+1}) - f(\eta)], \quad \mathcal{A}f(\eta) = \sum_{k \in \mathbb{Z}} (\eta_{k+1} - \eta_{k-1}) \frac{\partial}{\partial \eta_k} f(\eta).$$
(B9)

The invariant measure of the process is a product measure

$$\mu(\boldsymbol{\eta}) = \prod_{k \in \mathbb{Z}} I(\eta_k) \tag{B10}$$

with marginals

$$I(x) = \sqrt{\frac{\beta}{2\pi}} e^{-\frac{\beta}{2}(x-\rho)^2}.$$
(B11)

The quantities  $\beta$  and  $\rho$  are free parameters that can be expressed in terms of the densities of the conserved quantities as

$$\rho_1 := \langle \eta_k \rangle = \rho, \quad \rho_2 := \langle \eta_k^2 \rangle = \rho^2 + \frac{1}{\beta}.$$
(B12)

The static compressibilities are given by

$$\kappa_{11} := \langle (\eta_k - \rho_1)W \rangle = \langle \eta_k^2 \rangle - \langle \eta_k \rangle^2 = \frac{1}{\beta} = \rho_2 - \rho_1^2$$
(B13)

$$\kappa_{12} := \langle (\eta_k - \rho_1)E \rangle = \langle \eta_k^3 \rangle - \rho_1 \rho_2 = 2\rho_1(\rho_2 - \rho_1^2) \tag{B14}$$

$$\kappa_{22} := \langle \left( \eta_k^2 - \rho_2 \right) E \rangle = \langle \eta_k^4 \rangle - \rho_2^2 = 2(\rho_2 - \rho_1^2) \left( \rho_1^2 + \rho_2 \right).$$
(B15)

Hence the compressibility matrix reads

$$K = (\rho_2 - \rho_1^2) \begin{pmatrix} 1 & 2\rho_1 \\ 2\rho_1 & 2(\rho_1^2 + \rho_2) \end{pmatrix}$$
(B16)

From the action of the generator on the locally conserved quantities one obtains the discrete continuity equations

$$\mathcal{L}\eta_k = 2\eta_{k+1} - 2\eta_k = j_k^1 - j_{k+1}^1 \tag{B17}$$

$$\mathcal{L}\eta_k^2 = \eta_{k+1}^2 + \eta_{k-1}^2 - 2\eta_k^2 + 2(\eta_{k+1} - \eta_{k-1})\eta_k = j_k^2 - j_{k+1}^2.$$
(B18)

with the instantaneous currents

$$j_k^1 = -2\eta_k, \quad j_k^2 = \eta_{k-1}^2 - \eta_k^2 - 2\eta_{k-1}\eta_k.$$
 (B19)

The stationary currents are given by

$$j^{1} := \langle j_{k}^{v} \rangle = -2\rho_{1}, \quad j^{2} := \langle j_{k}^{e} \rangle = -2\rho_{1}^{2}$$
(B20)

which yields the current Jacobian

$$J = -2 \begin{pmatrix} 1 & 0\\ 2\rho_1 & 0 \end{pmatrix}.$$
 (B21)

Notice that the action of  $\mathcal{L}$  on  $\eta_k$  is linear and therefore the time evolution of  $\rho_k(t) := \langle \eta_k(t) \rangle$  can be integrated by Fourier transformation.

Using the results derived above these stationary currents yield the mode velocities

$$v_1 = -2, \quad v_2 = 0$$
 (B22)

and the mode coupling matrices

$$G^{(1)} = 0$$
 (B23)

$$G^{(2)} = -\sqrt{2} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}.$$
 (B24)

Mode coupling theory then predicts that mode 1 is diffusive and mode 2 belongs to the 3/2-Lévy universality class.

To compute the scaling form of the dynamical structure function [8, 11, 12] one needs the diffusion coefficient of mode 1. This can be computed directly and rigorously from the linear evolution of the conserved density  $\eta_k(t)$  by Fourier transformation and yields the large-scale behaviour in the moving reference frame with velocity  $v_1$  the scaling form

$$S_1(p,t) = \frac{1}{\sqrt{2\pi}} e^{-D_1 p^2 t}$$
(B25)

with  $D_1 = 1$ . Mode coupling theory then predicts the Fourier transform of the scaling function for mode 2 to be [8]

$$S_2(p,t) = \frac{1}{\sqrt{2\pi}} e^{-C_0 p^{3/2} [1 - i \operatorname{sign}(p(v_1 - v_2))]t}$$
(B26)

which is a maximally asymmetric 3/2-Lévy distribution with the scale factor

$$C_0 = \frac{(G_{11}^2)^2}{2\sqrt{D_1|v_1 - v_2|}} = \frac{1}{\sqrt{2}}$$
(B27)

On the other hand, in [52] the dynamical structure function was proved rigorously to be a fundamental solution of the fractional pde

$$\partial_t u(x,t) = -\frac{1}{\sqrt{2}} \left[ (-\Delta)^{3/4} - \nabla (-\Delta)^{1/4} \right] u(x,t).$$
(B28)

A Fourier transformation reproduces the mode coupling prediction (B26), thus proving that the mode coupling approximation is exact for this model and yields even the correct scale factor  $C_0$  [53].

## Appendix C: Inverse relationship between the mode coupling matrices and the Hessians

The inverse relationship between the mode coupling matrices and the Hessians is

$$H^{\gamma} = 2 \sum_{\lambda} (R^{-1})_{\gamma\lambda} R^T G^{\lambda} R.$$

Decomposing the mode coupling matrices into their two diagonal components  $G^{\lambda}_{\alpha\alpha}$  and the symmetric off-diagonal part  $G^{\lambda}_{12}$  yields after straightforward computation

$$\begin{aligned} H^{\gamma} &= f_{1}^{\gamma} \begin{pmatrix} 1 & u_{+} \\ u_{+} & u_{+}^{2} \end{pmatrix} + \bar{f}^{\gamma} \begin{pmatrix} 2u_{-} & 1 + u_{+}u_{-} \\ 1 + u_{+}u_{-} & 2u_{+} \end{pmatrix} + f_{2}^{\gamma} \begin{pmatrix} u_{-}^{2} & u_{-} \\ u_{-} & 1 \end{pmatrix} \\ &= u_{+} f_{1}^{\gamma} \begin{pmatrix} \frac{\delta + \theta}{2J_{12}} & 1 \\ 1 & \frac{\delta - \theta}{2J_{21}} \end{pmatrix} + \frac{2u_{+}}{J_{12}} \bar{f}^{\gamma} \begin{pmatrix} -J_{21} & \theta \\ \theta & J_{12} \end{pmatrix} + \frac{u_{+}J_{21}}{J_{12}} f_{2}^{\gamma} \begin{pmatrix} \frac{\delta - \theta}{2J_{12}} & -1 \\ -1 & \frac{\delta + \theta}{2J_{21}} \end{pmatrix} \end{aligned}$$

where the amplitudes

$$f_{1}^{\gamma} = \frac{2}{z_{+}^{2}} \sum_{\lambda} (R^{-1})_{\gamma \lambda} G_{11}^{\lambda}$$
  
$$\bar{f}^{\gamma} = \frac{2}{z_{+}z_{-}} \sum_{\lambda} (R^{-1})_{\gamma \lambda} G_{12}^{\lambda}$$
  
$$f_{2}^{\gamma} = \frac{2}{z_{-}^{2}} \sum_{\lambda} (R^{-1})_{\gamma \lambda} G_{22}^{\lambda}.$$

depend on the mode coupling coefficients.