

CHARACTERIZATION OF ENTIRE SOLUTIONS OF SYSTEMS OF QUADRATIC TRINOMIAL DIFFERENCE AND PARTIAL DIFFERENTIAL DIFFERENCE EQUATIONS IN \mathbb{C}^n

GOUTAM HALDAR

ABSTRACT. In this paper we establish some results about the existence and precise forms of finite order entire solutions of some systems of quadratic trinomial functional equations one of which in \mathbb{C}^n , $n \in \mathbb{N}$ and other two in \mathbb{C}^2 . Our results are the generalizations and improvements of the previous theorems given by Xu-Cao [42, 43] and Xu-Jiang [45]. Moreover, we exhibit some examples in support of our claims.

1. INTRODUCTION

It has a long history on the study of entire and meromorphic solutions of Fermat type equations by employing Nevanlinna theory as a tool. For detail study, we insist the readers to go through [5–7, 11, 12, 19, 32, 32, 40, 41, 48, 50] and the references therein. We assume the readers are familiar with the notations and basic definitions of Nevanlinna theory (see [17, 21]). In 1939, Iyer [19] proved that the entire solutions of Fermat type equation

$$f(z)^2 + g(z)^2 = 1 \quad (1.1)$$

are $f = \cos a(z)$, $g = \sin a(z)$, where $a(z)$ is an entire function.

Utilizing difference Nevanlinna theory of meromorphic functions developed by Halburd-Korhonen [13, 14], and Chiang-Feng [4], independently, Liu *et al.* [29], in 2012, proved that any finite order transcendental entire solutions of

$$f(z)^2 + f(z+c)^2 = 1, \quad (1.2)$$

has the form $f(z) = \sin(az + b)$, where b is a constant and $a = (4k + 1)\pi/2c$, with k is an integer. In the same paper, they also proved that any finite order transcendental entire solution of

$$f'(z)^2 + f(z+c)^2 = 1 \quad (1.3)$$

has the form $f(z) = \sin(z \pm Bi)$, where B is a constant and $c = 2p\pi$ or $c = (2p + 1)\pi$, $p \in \mathbb{Z}$, whereas any transcendental entire solutions with finite order of

$$f'(z)^2 + [f(z+c) - f(z)]^2 = 1, \quad (1.4)$$

Date: Received: xxxxxx; Revised: yyyyyy; Accepted: zzzzzz.

* Corresponding author.

2020 *Mathematics Subject Classification.* 39A45, 30D35, 32H30, 39A14, 35A20.

Key words and phrases. entire solutions, Fermat-type, differential difference equations, Nevanlinna theory.

must take the form $f(z) = 12 \sin(2z + bi)$, where $c = (2p+1)\pi$, p being an integer, and b is a constant.

Although there are many important and interesting results on the existence of entire and meromorphic solutions of complex difference as well as differential-difference equations (see [25–31,39]), only a few number of results are there on the existence and forms of entire and meromorphic solutions of quadratic trinomial functional equation. To the best of our knowledge, Saleeby [37] first investigated the existence and forms of entire and meromorphic solutions of quadratic trinomial functional equation

$$f^2 + 2wfg + g^2 = 1, \quad (1.5)$$

where w is a constant. Observe that (1.1) can be seen as the case of $w = 0$. For $w \neq 0$, Saleeby [37] proved the following classical results about the Fermat type functional equations (1.5).

Theorem A. [37] *Let $\alpha \in \mathbb{C}$ with $\alpha^2 \neq 0, 1$. Then any transcendental entire solution of (1.5) has the form $f(z) = \frac{1}{\sqrt{2}} \left(\frac{\cos(h(z))}{\sqrt{1+w}} + \frac{\sin(h(z))}{\sqrt{1-w}} \right)$ and $\frac{1}{\sqrt{2}} \left(\frac{\cos(h(z))}{\sqrt{1+w}} - \frac{\sin(h(z))}{\sqrt{1-w}} \right)$, where h is entire \mathbb{C}^n . The meromorphic solutions of (1.5) is of the form $f(z) = \frac{\alpha_1 - \alpha_2 \beta(z)^2}{(\alpha_1 - \alpha_2)\beta(z)}$ and $g(z) = \frac{1 - \beta(z)^2}{(\alpha_1 - \alpha_2)\beta(z)}$, where $\beta(z)$ is meromorphic in \mathbb{C}^n and $\alpha_1 = -w + \sqrt{w^2 - 1}$, $\alpha_2 = -w - \sqrt{w^2 - 1}$.*

By utilizing Theorem A, in 2016, Liu-Yang [31] obtained more precise properties of entire and meromorphic solutions of

$$f(z)^2 + 2wf(z)f'(z) + f'(z)^2 = 1 \quad (1.6)$$

and

$$f(z)^2 + 2wf(z)f(z+c) + f(z+c)^2 = 1, \quad w \neq 0, \pm 1. \quad (1.7)$$

To the best of our knowledge, there are only a few number of results on the system of complex difference and differential-difference equations in the literature (see [9]). It was Gao [9], who, in 2016, first discussed the existence and form of transcendental entire solutions of system of differential difference equations and obtained the result as follows: If (f, g) be a pair of finite order transcendental entire solution of the system of differential-difference equations

$$\begin{cases} f'(z)^2 + g(z+c)^2 = 1, \\ g'(z)^2 + f(z+c)^2 = 1. \end{cases} \quad (1.8)$$

Then (f, g) must be of the form

$$(f, g) = (\sin(z - ib), \sin(z - ib_1)), \quad \text{or} \quad (\sin(z + ib), \sin(z + ib_1)),$$

where b, b_1 are constants and $c = k\pi$, k is a integer.

2. SOLUTIONS OF FERMAT TYPE EQUATIONS IN SEVERAL COMPLEX VARIABLES

Recently, the study of entire and meromorphic solutions of Fermat type partial differential equations in several complex variables has received considerable attention in the literature (see [22–24, 35, 36]). The study partial differential equations which is a generalizations of the well-known eikonal equation in real variable case has a long history. We insist the readers to go through [8, 10, 33] and the references therein. In 1995, Khavinson pointed out that in \mathbb{C}^2 , the entire solution of the Fermat type partial differential equations $f_{z_1}^2 + f_{z_2}^2 = 1$ must necessarily be linear. After the development of the difference Nevanlinna theory in several complex variables, specially the difference version of logarithmic derivative lemma (see [1–3, 20]), many Researchers started studying the existence and the precise form of entire and meromorphic solutions of different variants of Fermat type difference and partial differential difference equations, and obtained very remarkable and interesting results (see [15, 16, 34, 38, 42–44, 46]).

Hereinafter, we denote by $z + w = (z_1 + w_1, z_2 + w_2, \dots, z_n + w_n)$, where $z = (z_1, z_2, \dots, z_n), w = (w_1, w_2, \dots, w_n) \in \mathbb{C}^n$, n being a positive integer.

In 2018, Xu-Cao [42] established the existence of entire solutions for some Fermat type complex difference and partial differential difference equations with several variables. We list some of them below.

Theorem B. [42] *Let $c = (c_1, c_2, \dots, c_n) \in \mathbb{C}^n \setminus \{(0, 0, \dots, 0)\}$. If $f : \mathbb{C}^n \rightarrow \mathbb{P}^1(\mathbb{C})$ be an entire solution with finite order of the Fermat type difference equation*

$$f(z)^2 + f(z + c)^2 = 1, \quad (2.1)$$

the $f(z)$ must assume the form $f(z) = \cos(L(z) + B)$, where L is a linear function of the form $L(z) = a_1 z_1 + \dots + a_n z_n$ on \mathbb{C}^n such that $L(c) = -\pi/2 - 2k\pi$ ($k \in \mathbb{Z}$), and B is a constant on \mathbb{C} .

Theorem C. [42] *Let $c = (c_1, c_2) \in \mathbb{C}^2$. If $f(z)$ be transcendental entire solution with finite order of the Fermat type partial differential-difference equation*

$$\left(\frac{\partial f(z_1, z_2)}{\partial z_1} \right)^2 + f(z_1 + c_1, z_2 + c_2)^2 = 1, \quad (2.2)$$

f must be of the form $f(z_1, z_2) = \sin(Az_1 + Bz_2 + \phi(z_2))$, Where A, B are constant on \mathbb{C} satisfying $A^2 = 1$ and $Ae^{i(Ac_1 + Bc_2)} = 1$, and $\phi(z_2)$ is a polynomial in one variable z_2 such that $\phi(z_2) \equiv \phi(z_2 + c_2)$. In the special case whenever $c_2 \neq 0$, we have $f(z_1, z_2) = \sin(Az_1 + Bz_2 + \text{Constant})$.

Very recently, corresponding to Equations (2.5) and (1.6), Xu-Jiang [45] considered the following system of trinomial difference and differential-difference equations in \mathbb{C}^2

$$\begin{cases} f(z)^2 + 2wf(z)g(z + c) + g(z + c)^2 = 1 \\ g(z)^2 + 2wg(z)f(z + c) + f(z + c)^2 = 1 \end{cases} \quad (2.3)$$

and

$$\begin{cases} \left(\frac{\partial f}{\partial z_1}\right)^2 + 2w\frac{\partial f}{\partial z_1}g(z+c) + g(z+c)^2 = 1 \\ \left(\frac{\partial g}{\partial z_1}\right)^2 + 2w\frac{\partial g}{\partial z_1}f(z+c) + f(z+c)^2 = 1, \end{cases} \quad (2.4)$$

and obtained the results as follows.

Theorem D. [45] Let $c = (c_1, c_2) \in \mathbb{C}^2 \setminus \{(0, 0)\}$. Then any pair of finite order transcendental entire solutions of (2.3) must assume one of the following forms:

(i)

$$f(z) = \frac{1}{\sqrt{2}} \left(\frac{\cos(\gamma(z) + b_1)}{\sqrt{1+w}} + \frac{\sin(\gamma(z) + b_1)}{\sqrt{1-w}} \right),$$

$$g(z) = \frac{1}{\sqrt{2}} \left(\frac{\cos(\gamma(z) + b_2)}{\sqrt{1+w}} + \frac{\sin(\gamma(z) + b_2)}{\sqrt{1-w}} \right),$$

where $\gamma(z) = L(z) + H(s)$, $L(z) = a_1z_1 + a_2z_2$, $H(s)$ is a polynomial in $s := c_2z_1 - c_1z_2$, $a_i, b_i, c_i, i = 1, 2$ satisfy the relations

$$e^{2iL(c)} = \frac{-w + \sqrt{w^2 - 1}}{-w - \sqrt{w^2 - 1}}, \quad e^{2i(b_1 - b_2)} = 1.$$

(ii)

$$f(z) = \frac{1}{\sqrt{2}} \left(\frac{\cos(\gamma(z) + b_1)}{\sqrt{1+w}} + \frac{\sin(\gamma(z) + b_1)}{\sqrt{1-w}} \right),$$

$$g(z) = \frac{1}{\sqrt{2}} \left(\frac{\cos(\gamma(z) + b_2)}{\sqrt{1+w}} - \frac{\sin(\gamma(z) + b_2)}{\sqrt{1-w}} \right),$$

where $\gamma(z)$ is the same as in (i), and $a_i, b_i, c_i, i = 1, 2$ satisfy the relations

$$e^{2iL(c)} = 1, \quad e^{2i(b_1 - b_2)} = 1.$$

Theorem E. [45] Let $c = (c_1, c_2) \in \mathbb{C}^2 \setminus \{(0, 0)\}$ with $c_2 \neq 0$. Then any pair of finite order transcendental entire solutions of (2.4) must assume one of the following forms:

(i)

$$f(z) = \frac{1}{\sqrt{2}} \left(\frac{\cos(L(z) - L(c) + b_1)}{\sqrt{1+w}} + \frac{\sin(L(z) - L(c) + b_1)}{\sqrt{1-w}} \right),$$

$$g(z) = \frac{1}{\sqrt{2}} \left(\frac{\cos(L(z) - L(c) + b_2)}{\sqrt{1+w}} + \frac{\sin(L(z) - L(c) + b_2)}{\sqrt{1-w}} \right),$$

where $L(z) = a_1z_1 + a_2z_2$, $a_i, b_i, c_i, i = 1, 2$ satisfy the relations

$$a_1^2 = 1, \quad e^{2iL(c)} = \frac{-w + \sqrt{w^2 - 1}}{-w - \sqrt{w^2 - 1}}, \quad e^{2i(b_1 - b_2)} = 1.$$

(ii)

$$f(z) = \frac{1}{\sqrt{2}} \left(\frac{\cos(L(z) - L(c) + b_1)}{\sqrt{1+w}} + \frac{\sin(L(z) - L(c) + b_1)}{\sqrt{1-w}} \right),$$

$$g(z) = \frac{1}{\sqrt{2}} \left(\frac{\cos(L(z) - L(c) + b_2)}{\sqrt{1+w}} - \frac{\sin(L(z) - L(c) + b_2)}{\sqrt{1-w}} \right),$$

where $L(z)$ is the same as in (i), and $a_i, b_i, c_i, i = 1, 2$ satisfy the relations

$$a_1^2 = 1, \quad e^{2iL(c)} = 1, \quad e^{2i(b_1 - b_2)} = 1.$$

Observing Theorems B and E, it is inevitable to ask the following open questions.

Question 2.1. *What can be said about the entire solutions of (2.3) when one replace the right hand sides of each equation by $e^{g_1(z)}$ and $e^{g_2(z)}$, respectively, where $g_1(z), g_2(z)$ are any two polynomials in \mathbb{C}^n , n being an integer?*

Question 2.2. *What can be said about the existence and precise form of transcendental entire solutions of (2.4) when one replace (a) first order partial derivative by k -th order partial derivative for any $k \in \mathbb{N}$ and (b) right hand sides of each of the equations by $e^{g_1(z_1, z_2)}$ and $e^{g_2(z_1, z_2)}$, respectively, where $g_1(z_1, z_2), g_2(z_1, z_2)$ are any two polynomials in \mathbb{C}^2 ?*

Question 2.3. *What can be said about the existence and precise form of transcendental entire solutions of*

$$\begin{cases} \left[\frac{\partial^k f}{\partial z_1^k} \right]^2 + 2w \frac{\partial^k f}{\partial z_1^k} [g(z+c) - g(z)] + [g(z+c) - g(z)]^2 = 1, \\ \left[\frac{\partial^k g}{\partial z_1^k} \right]^2 + 2w \frac{\partial^k g}{\partial z_1^k} [f(z+c) - f(z)] + [f(z+c) - f(z)]^2 = 1? \end{cases} \quad (2.5)$$

3. RESULTS AND EXAMPLES

Inspired by the above results and questions, we investigate the existence and form of transcendental entire solutions of the system of equations

$$\begin{cases} f(z)^2 + 2wf(z)g(z+c) + g(z+c)^2 = e^{g_1(z)} \\ g(z)^2 + 2wg(z)f(z+c) + f(z+c)^2 = e^{g_2(z)}, \end{cases} \quad (3.1)$$

$$\begin{cases} \left[\frac{\partial^k f}{\partial z_1^k} \right]^2 + 2w \frac{\partial^k f}{\partial z_1^k} g(z+c) + g(z+c)^2 = e^{g_1(z_1, z_2)} \\ \left[\frac{\partial^k g}{\partial z_1^k} \right]^2 + 2w \frac{\partial^k g}{\partial z_1^k} f(z+c) + f(z+c)^2 = e^{g_2(z_1, z_2)}, \end{cases} \quad (3.2)$$

where $g_1(z_1, z_2), g_2(z_1, z_2)$ are any two polynomials and $k \in \mathbb{N}$ and Eq. (2.5). Our results are the generalizations of Theorems B–C to a large extent.

Before we state our main results, let us denote two non-zero complex numbers A_1, A_2 as follows.

$$A_1 = \frac{1}{2\sqrt{1+w}} + \frac{1}{2i\sqrt{1-w}} \quad \text{and} \quad A_2 = \frac{1}{2\sqrt{1+w}} - \frac{1}{2i\sqrt{1-w}}.$$

Before we state our main results, let us set the following.

$$\begin{aligned} \Phi(z) &= \sum_{i_1=1}^{nC_2} H_{i_1}^2(s_{i_1}^2) + \sum_{i_2=1}^{nC_3} H_{i_2}^3(s_{i_2}^3) + \sum_{i_3=1}^{nC_4} H_{i_3}^4(s_{i_3}^4) + \cdots \\ &+ \sum_{i_{n-2}=1}^{nC_{n-1}} H_{i_{n-2}}^{n-1}(s_{i_{n-2}}^{n-1}) + H_{n-1}^n(s_{n-1}^n), \end{aligned} \quad (3.3)$$

$$\begin{aligned} \Psi(z) &= \sum_{i_1=1}^{nC_2} L_{i_1}^2(t_{i_1}^2) + \sum_{i_2=1}^{nC_3} L_{i_2}^3(t_{i_2}^3) + \sum_{i_3=1}^{nC_4} L_{i_3}^4(t_{i_3}^4) + \cdots \\ &+ \sum_{i_{n-2}=1}^{nC_{n-1}} L_{i_{n-2}}^{n-1}(t_{i_{n-2}}^{n-1}) + L_{n-1}^n(t_{n-1}^n), \end{aligned} \quad (3.4)$$

where $H_{i_1}^2$ is a polynomial in $s_{i_1}^2 := d_{i_1 j_1}^2 z_{j_1} + d_{i_1 j_2}^2 z_{j_2}$ with $d_{i_1 j_1}^2 c_{j_1} + d_{i_1 j_2}^2 c_{j_2} = 0$, $1 \leq i_1 \leq {}^nC_2$, $1 \leq j_1 < j_2 \leq n$, $H_{i_2}^3$ is a polynomial in $s_{i_2}^3 := d_{i_2 j_1}^3 z_{j_1} + d_{i_2 j_2}^3 z_{j_2} + d_{i_2 j_3}^3 z_{j_3}$ with $d_{i_2 j_1}^3 c_{j_1} + d_{i_2 j_2}^3 c_{j_2} + d_{i_2 j_3}^3 c_{j_3} = 0$, $1 \leq i_2 \leq {}^nC_3$, $1 \leq j_1 < j_2 < j_3 \leq n \dots$, $H_{i_{n-2}}^{n-1}$ is a polynomial in $s_{i_{n-2}}^{n-1} := d_{i_{n-2} j_1} z_{j_1} + d_{i_{n-2} j_2} z_{j_2} + \cdots + d_{i_{n-2} j_{n-1}} z_{j_{n-1}}$ with $d_{i_{n-2} j_1} c_{j_1} + d_{i_{n-2} j_2} c_{j_2} + \cdots + d_{i_{n-2} j_{n-1}} c_{j_{n-1}} = 0$, $1 \leq i_{n-2} \leq {}^nC_{n-1}$, $1 \leq j_1 < j_2 < \cdots < j_{n-1} \leq n$ and H_{n-1}^n is a polynomial in $s_{n-1}^n := d_{n-1 1} z_1 + d_{n-1 2} z_2 + \cdots + d_{n-1 n} z_n$ with $d_{n-1 1} c_1 + d_{n-1 2} c_2 + \cdots + d_{n-1 n} c_n = 0$, $L_{i_1}^2$ is a polynomial in $t_{i_1}^2 := e_{i_1 j_1}^2 z_{j_1} + e_{i_1 j_2}^2 z_{j_2}$ with $e_{i_1 j_1}^2 c_{j_1} + e_{i_1 j_2}^2 c_{j_2} = 0$, $1 \leq i_1 \leq {}^nC_2$, $1 \leq j_1 < j_2 \leq n$, $H_{i_2}^3$ is a polynomial in $t_{i_2}^3 := e_{i_2 j_1}^3 z_{j_1} + e_{i_2 j_2}^3 z_{j_2} + e_{i_2 j_3}^3 z_{j_3}$ with $e_{i_2 j_1}^3 c_{j_1} + e_{i_2 j_2}^3 c_{j_2} + e_{i_2 j_3}^3 c_{j_3} = 0$, $1 \leq i_2 \leq {}^nC_3$, $1 \leq j_1 < j_2 < j_3 \leq n \dots$, $L_{i_{n-2}}^{n-1}$ is a polynomial in $t_{i_{n-2}}^{n-1} := e_{i_{n-2} j_1} z_{j_1} + e_{i_{n-2} j_2} z_{j_2} + \cdots + e_{i_{n-2} j_{n-1}} z_{j_{n-1}}$ with $e_{i_{n-2} j_1} c_{j_1} + e_{i_{n-2} j_2} c_{j_2} + \cdots + e_{i_{n-2} j_{n-1}} c_{j_{n-1}} = 0$, $1 \leq i_{n-2} \leq {}^nC_{n-1}$, $1 \leq j_1 < j_2 < \cdots < j_{n-1} \leq n$ and L_{n-1}^n is a polynomial in $t_{n-1}^n := e_{n-1 1} z_1 + e_{n-1 2} z_2 + \cdots + e_{n-1 n} z_n$ with $e_{n-1 1} c_1 + e_{n-1 2} c_2 + \cdots + e_{n-1 n} c_n = 0$, where for each k , the representation of s_i^k and t_i^k in terms of the conditions of j_1, j_2, \dots, j_k are unique.

Now, we list our main results as follows.

Theorem 3.1. *Let $c = (c_1, c_2, \dots, c_n) \in \mathbb{C}^n$, $w \in \mathbb{C}$ such that $w^2 \neq 0, 1$ and $g_1(z)$, $g_2(z)$ be any two polynomials in \mathbb{C}^n . If $(f(z), g(z))$ be a pair of transcendental entire solution with finite order of the system Fermat type difference equation (3.1), then one of the following must occur.*

$$(i) \quad g_1(z) = L(z) + \Phi(z) + d_1, \quad g_2(z) = L(z) + \Phi(z) + d_2,$$

$$f(z) = \frac{A_1 \xi_1 + A_2 \xi_1^{-1}}{\sqrt{2}} e^{\frac{1}{2}[L(z) + \Phi(z) + d_1]}, \quad g(z) = \frac{A_1 \xi_2 + A_2 \xi_2^{-1}}{\sqrt{2}} e^{\frac{1}{2}[L(z) + \Phi(z) + d_2]},$$

where $L(z) = \sum_{j=1}^n a_j z_j$, $\Phi(z)$ is defined as in (3.3) and satisfy the relations

$$e^{L(c)} = \frac{(A_2 \xi_2 + A_1 \xi_2^{-1})(A_2 \xi_1 + A_1 \xi_1^{-1})}{(A_1 \xi_1 + A_2 \xi_1^{-1})(A_1 \xi_2 + A_2 \xi_2^{-1})},$$

$$e^{d_1 - d_2} = \frac{(A_2 \xi_2 + A_1 \xi_2^{-1})(A_1 \xi_2 + A_2 \xi_2^{-1})}{(A_1 \xi_1 + A_2 \xi_1^{-1})(A_2 \xi_1 + A_1 \xi_1^{-1})},$$

$a_j, d_1, d_2, \xi_1 (\neq 0), \xi_2 (\neq 0) \in \mathbb{C}$ for $j = 1, 2, \dots, n$.

(ii)

$$g_1(z) = L_1(z) + L_2(z) + \Phi(z) + \Psi(z) + d_1 + d_3,$$

$$g_2(z) = L_1(z) + L_2(z) + \Phi(z) + \Psi(z) + d_2 + d_4,$$

$$f(z) = \frac{1}{\sqrt{2}} [A_1 e^{L_1(z) + \Phi(z) + d_1} + A_2 e^{L_2(z) + \Psi(z) + d_3}],$$

$$g(z) = \frac{1}{\sqrt{2}} [A_1 e^{L_2(z) + \Psi(z) + d_4} + A_2 e^{L_1(z) + \Phi(z) + d_2}],$$

where $L_1(z) = \sum_{j=1}^n a_j z_j$, $L_2(z) = \sum_{j=1}^n b_j z_j$, $\Phi(z)$, $\Psi(z)$ are defined in (3.3) and (3.4), respectively, $a_j, b_j, d_i \in \mathbb{C}$ for $j = 1, 2, \dots, n$, $i = 1, \dots, 4$ such that $L_1(z) + \Phi(z) \neq L_2(z) + \Psi(z)$ and satisfy one of the following relations:

(a) $e^{L_1(c)} = 1$, $e^{L_2(c)} = 1$, $e^{d_1 - d_2} = 1$ and $e^{d_3 - d_4} = 1$,

(b) $e^{L_1(c)} = 1$, $e^{L_2(c)} = -1$, $e^{d_1 - d_2} = 1$ and $e^{d_3 - d_4} = -1$,

(c) $e^{L_1(c)} = -1$, $e^{L_2(c)} = 1$, $e^{d_1 - d_2} = -1$ and $e^{d_3 - d_4} = 1$,

(d) $e^{L_1(c)} = -1$, $e^{L_2(c)} = -1$, $e^{d_1 - d_2} = -1$ and $e^{d_3 - d_4} = -1$.

(iii) $g_1(z)$ and $g_2(z)$ have the same form as in (ii),

$$f(z) = \frac{1}{\sqrt{2}} [A_1 e^{L_1(z) + \Phi(z) + d_1} + A_2 e^{L_2(z) + \Psi(z) + d_3}],$$

$$g(z) = \frac{1}{\sqrt{2}} [A_1 e^{L_1(z) + \Phi(z) + d_2} + A_2 e^{L_2(z) + \Psi(z) + d_4}],$$

where L_1, L_2, Φ, Ψ are same as in (ii) with $L_1(z) + \Phi(z) \neq L_2(z) + \Psi(z)$, and satisfy one of the following relations:

(a) $e^{L_1(c)} = A_2/A_1$, $e^{L_2(c)} = A_1/A_2$, $e^{d_1 - d_2} = 1$ and $e^{d_3 - d_4} = 1$,

(b) $e^{L_1(c)} = A_2/A_1$, $e^{L_2(c)} = -A_1/A_2$, $e^{d_1 - d_2} = 1$ and $e^{d_3 - d_4} = -1$,

(c) $e^{L_1(c)} = -A_2/A_1$, $e^{L_2(c)} = A_1/A_2$, $e^{d_1 - d_2} = -1$ and $e^{d_3 - d_4} = 1$,

(d) $e^{L_1(c)} = -A_2/A_1$, $e^{L_2(c)} = -A_1/A_2$, $e^{d_1 - d_2} = -1$ and $e^{d_3 - d_4} = -1$.

The following examples show the existence of transcendental entire solutions with finite order of the system (3.1).

Example 3.1. Let $n = 2$, $w = 3$, $\xi_1 = \xi_2 = 1$, $c_1 = c_2 = \pi i$, $d_1 = \pi i$, $d_2 = -\pi i$, $g_1(z) = z_1 + z_2 + (z_1 - z_2)^3 + d_1$, $g_2(z) = z_1 + z_2 + (z_1 - z_2)^3 + d_2$. Then in view of the conclusion (i) of Theorem 3.1, we can see that

$$(f(z), g(z)) = \left(\frac{1}{2\sqrt{2}} e^{\frac{1}{2}[z_1+z_2+(z_1-z_2)^3+d_1]}, \frac{1}{2\sqrt{2}} e^{\frac{1}{2}[z_1+z_2+(z_1-z_2)^3+d_2]} \right)$$

is a solution of (3.1).

Example 3.2. Let $w = 2$, $n = 3$, $L(z) = z_1 + z_2 + z_3$, $L_2(z) = z_1 + z_2 - z_3$, $c_1 = \pi i$, $c_2 = 2\pi i$, $c_3 = -\pi i$, $\Phi(z) = -\pi^2(z_1 - z_2)^2 + i\pi^3(z_1 + z_2)^3 + \pi^4(z_2 - z_3)^4$, $\Psi(z) = \pi^4(z_1 + 2z_2 - z_3)^2$. Choose $d_j \in \mathbb{C}$ such that $d_1 - d_2 = 2\pi i$ and $d_3 - d_4 = 4\pi i$. Then, in view of the conclusion (ii) of Theorem 3.1, $(f(z), g(z))$, where

$$f(z) = \frac{1}{\sqrt{2}} \left[-\frac{3 - \sqrt{3}}{6} e^{L_1(z) + \Phi(z) + d_1} + \frac{3 + \sqrt{3}}{6} e^{L_2(z) + \Psi(z) + d_3} \right] \text{ and}$$

$$g(z) = \frac{1}{\sqrt{2}} \left[-\frac{3 - \sqrt{3}}{6} e^{L_2(z) + \Psi(z) + d_3} + \frac{3 + \sqrt{3}}{6} e^{L_1(z) + \Phi(z) + d_1} \right]$$

is a solution of (3.1).

Theorem 3.2. Let $k \in \mathbb{N}$, $c = (c_1, c_2) \in \mathbb{C}^2$, $w \in \mathbb{C}$ such that $w^2 \neq 0, 1$, $g_1(z_1, z_2), g_2(z_1, z_2)$ be any two polynomials in \mathbb{C}^2 . If (f, g) is a pair of finite order transcendental entire solution of (3.2), then one of the following cases must occur.

(i) $g_1(z_1, z_2) = \mathcal{L}(z) + \chi(z_2) + d_1$, $g_2(z_1, z_2) = \mathcal{L}(z) + \chi(z_2) + d_2$,

$$f(z) = \frac{A_2 \xi_2 + A_1 \xi_2^{-1}}{\sqrt{2}} e^{\frac{1}{2}[\mathcal{L}(z) + \chi(z_2) - \mathcal{L}(c) + d_2]}, \quad g(z) = \frac{A_2 \xi_1 + A_1 \xi_1^{-1}}{\sqrt{2}} e^{\frac{1}{2}[\mathcal{L}(z) + \chi(z_2) - \mathcal{L}(c) + d_1]},$$

where $\mathcal{L}(z) = a_1 z_1 + a_2 z_2$, $\chi(z_2)$ is a polynomial in z_2 , only such that $\chi(z_2 + c_2) = \chi(z_2)$, $\xi_j \neq 0$, $a_j, d_j \in \mathbb{C}$, $j = 1, 2$, $a_1 \neq 0$, and satisfy the following:

$$\begin{cases} e^{\mathcal{L}(c)} = \left(\frac{a_1}{2}\right)^{2k} \frac{(A_2 \xi_2 + A_1 \xi_2^{-1})(A_2 \xi_1 + A_1 \xi_1^{-1})}{(A_1 \xi_1 + A_2 \xi_1^{-1})(A_1 \xi_2 + A_2 \xi_2^{-1})}, \\ e^{\frac{1}{2}[\mathcal{L}(c) + d_1 - d_2]} = \left(\frac{a_1}{2}\right)^k \frac{A_2 \xi_2 + A_1 \xi_2^{-1}}{A_1 \xi_1 + A_2 \xi_1^{-1}}. \end{cases}$$

In particular, if $c_2 \neq 0$, then $\chi(z_2)$ becomes linear in z_2 .

(ii)

$$g_1(z_1, z_2) = L_1(z) + L_2(z) + \Phi_1(z_2) + \Psi_1(z_2) + d_1 + d_3,$$

$$g_2(z_1, z_2) = L_1(z) + L_2(z) + \Phi_1(z_2) + \Psi_1(z_2) + d_2 + d_4,$$

$$f(z_1, z_2) = \frac{A_2 e^{L_2(z) + \Psi_1(z_2) - L_2(c) + d_4} + A_1 e^{L_1(z) + \Phi_1(z_2) - L_1(c) + d_2}}{\sqrt{2}},$$

$$g(z_1, z_2) = \frac{A_2 e^{L_1(z) + \Phi_1(z_2) - L_1(c) + d_1} + A_1 e^{L_2(z) + \Psi_1(z_2) - L_2(c) + d_3}}{\sqrt{2}},$$

where $L_1(z) = a_1z_1 + a_2z_2$, $L_2(z) = b_1z_1 + b_2z_2$, Φ_1, Ψ_1 are two polynomials in z_2 , only with $\Phi_1(z_2 + c_2) = \Phi_1(z_2)$, $\Psi_1(z_2 + c_2) = \Psi_1(z_2)$ such that $L_1(z) + \Phi_1(z_2) \neq L_2(z) + \Psi_1(z_2)$, $a_i, b_i, d_j \in \mathbb{C}$ for $i = 1, 2, j = 1, 2, 3, 4$, and satisfy one of the following relations:

- (a) $e^{L_1(c)} = a_1^k$, $e^{L_2(c)} = b_1^k$, $e^{d_1-d_2} = 1$ and $e^{d_3-d_4} = 1$,
- (b) $e^{L_1(c)} = a_1^k$, $e^{L_2(c)} = -b_1^k$, $e^{d_1-d_2} = 1$ and $e^{d_3-d_4} = -1$,
- (c) $e^{L_1(c)} = -a_1^k$, $e^{L_2(c)} = b_1^k$, $e^{d_1-d_2} = -1$ and $e^{d_3-d_4} = 1$,
- (d) $e^{L_1(c)} = -a_1^k$, $e^{L_2(c)} = -b_1^k$, $e^{d_1-d_2} = -1$ and $e^{d_3-d_4} = -1$.

(iii) The form of $g_1(z_1, z_2)$ and $g_2(z_1, z_2)$ are the same as in (ii),

$$f(z_1, z_2) = \frac{A_2 e^{L_1(z) + \Phi_1(z_2) - L_1(c) + d_2} + A_1 e^{L_2(z) + \Psi_1(z_2) - L_2(c) + d_4}}{\sqrt{2}},$$

$$g(z_1, z_2) = \frac{A_2 e^{L_1(z) + \Phi_1(z_2) - L_1(c) + d_1} + A_1 e^{L_2(z) + \Psi_1(z_2) - L_2(c) + d_3}}{\sqrt{2}},$$

where L_1, L_2, Φ_1, Ψ_1 are the same as in (ii), and satisfy one of the following conditions:

- (a) $e^{L_1(c)} = A_2 a_1^k / A_1$, $e^{L_2(c)} = A_1 b_1^k / A_2$, $e^{d_1-d_2} = 1$ and $e^{d_3-d_4} = 1$,
- (b) $e^{L_1(c)} = A_2 a_1^k / A_1$, $e^{L_2(c)} = -A_1 b_1^k / A_2$, $e^{d_1-d_2} = 1$ and $e^{d_3-d_4} = -1$,
- (c) $e^{L_1(c)} = -A_2 a_1^k / A_1$, $e^{L_2(c)} = A_1 b_1^k / A_2$, $e^{d_1-d_2} = -1$ and $e^{d_3-d_4} = 1$,
- (d) $e^{L_1(c)} = -A_2 a_1^k / A_1$, $e^{L_2(c)} = -A_1 b_1^k / A_2$, $e^{d_1-d_2} = -1$ and $e^{d_3-d_4} = -1$.

The following examples show the existence of transcendental entire solutions with finite order of the system (3.2).

Example 3.3. Let $k = 1$, $L(z) = z_1 + z_2$, $\xi_1 = 1$, $\xi_2 = -1$ and w be a complex number such that $w \neq 0, \pm 1$. Choose $c = (c_1, c_2) \in \mathbb{C}$ such that $e^{L(c)/2} = 1/2$ and $d_1, d_2 \in \mathbb{C}$ such that $e^{(d_1-d_2)/2} = -1$. Then, in view of the conclusion (i) of Theorem 3.2, we easily see that (f, g) , where

$$f(z_1, z_2) = -\frac{2}{\sqrt{2(1+w)}} e^{\frac{1}{2}[z_1+z_2+d_2]} \quad \text{and} \quad g(z_1, z_2) = \frac{2}{\sqrt{2(1+w)}} e^{\frac{1}{2}[z_1+z_2+d_1]}$$

is a solution of (3.2) with $g_1(z_1, z_2) = z_1 + z_2 + d_1$ and $g_2(z_1, z_2) = z_1 + z_2 + d_2$.

Example 3.4. Let $k = 1$, $w = 2$, $L_1(z) = z_1 + z_2$, $L_2(z) = z_1 - 2z_2$, $c_1 = 8\pi i/3$, $c_2 = -2\pi i/3$. Choose $d_j \in \mathbb{C}$ for $j = 1, \dots, 4$ such that $d_1 - d_2 = 2m\pi i$ and $d_3 - d_4 = 2p\pi i$, $m, p \in \mathbb{Z}$. Then, in view of the conclusion (ii) of Theorem 3.2, we easily see that $(f(z), g(z))$, where

$$f(z_1, z_2) = \frac{1}{6\sqrt{2}} \left[-(3 - \sqrt{3})e^{z_1+z_2+d_2} + (3 + \sqrt{3})e^{z_1-2z_2+d_4} \right] \quad \text{and}$$

$$g(z_1, z_2) = \frac{1}{6\sqrt{2}} \left[(3 + \sqrt{3})e^{z_1+z_2+d_1} - (3 - \sqrt{3})e^{z_1-2z_2+d_3} \right]$$

is a solution of (3.2) with $g_1(z_1, z_2) = 2z_1 - z_2 + d_1 + d_3$ and $g_2(z_1, z_2) = 2z_1 - z_2 + d_2 + d_4$.

Theorem 3.3. *Let k be a positive integer, $c = (c_1, c_2) \in \mathbb{C}^2$, $w \in \mathbb{C}$ such that $w^2 \neq 0, 1$ and $f(z), g(z)$ are transcendental entire functions of finite order not of c -periodic. If $(f(z), g(z))$ be a solution of (2.5), then one of the following conclusions must hold.*

(i)

$$f(z) = \frac{A_1 e^{L(z)+\Phi_1(z_2)+\gamma} + A_2 e^{-(L(z)+\Phi_1(z_2)+\gamma)}}{\sqrt{2}\alpha^k} \quad \text{and}$$

$$g(z) = \frac{A_1 e^{L(z)+\Phi_1(z_2)+\gamma+\eta} + A_2 e^{-(L(z)+\Phi_1(z_2)+\gamma+\eta)}}{\sqrt{2}\alpha^k}.$$

where $L(z) = \alpha z_1 + \beta z_2 + \gamma$, $\Phi_1(z_2)$ is a polynomial in z_2 with $\Phi_1(z_2 + c_2) = \Phi_1(z_2)$, $\alpha, \beta, \gamma, \eta \in \mathbb{C}$, and satisfy the following relations:

$$e^{L(c)} = \frac{A_2 e^\eta}{(-1)^k A_1 \alpha^k + A_2 e^\eta}, \quad e^{2\eta} = 1 \quad \text{and} \quad \alpha^k = -\frac{(-1)^k A_1^2 + A_2^2}{(-1)^k A_1 A_2 e^\eta}.$$

In particular, if $c_2 \neq 0$, then the polynomial $\Phi_1(z_2)$ becomes linear in z_2 .

(ii)

$$f(z) = \frac{A_1 e^{L(z)+\Phi_1(z_2)+\gamma} + A_2 e^{-(L(z)+\Phi_1(z_2)+\gamma)}}{\sqrt{2}\alpha^k},$$

$$g(z) = \frac{A_1 e^{-(L(z)+\Phi_1(z_2)+\gamma)+\eta} + A_2 e^{L(z)+\Phi_1(z_2)+\gamma-\eta}}{\sqrt{2}\alpha^k},$$

where k is an even positive integer, $L(z), \Phi_1(z_2)$ are the same as in (i), and satisfy the following relations:

$$e^{L(c)} = 1 + e^{-\eta}\alpha^k, \quad e^{2\eta} = 1 \quad \text{and} \quad \alpha^k = -2e^\eta.$$

The following example shows that the existence of transcendental entire solutions of (2.5) is precise.

Example 3.5. *Let $k = 2, w = 2$ and choose $\alpha, \eta, c_1, c_2 \in \mathbb{C}$ such that $\alpha^2 = -2, \eta = 2p\pi i$ and $c_1 + c_2 = (2m + 1)\pi i$, $p, m \in \mathbb{Z}$. Then in view of the conclusion (ii) of Theorem 3.3, we see that (f, g) , where*

$$f(z_1, z_2) = \frac{1}{12\sqrt{2}} \left[(3 - \sqrt{3})e^{\alpha z_1 + z_2 + 1} - (3 + \sqrt{3})e^{-(\alpha z_1 + z_2 + 1)} \right] \quad \text{and}$$

$$g(z_1, z_2) = \frac{1}{12\sqrt{2}} \left[(3 - \sqrt{3})e^{-(\alpha z_1 + z_2 + 1)} - (3 + \sqrt{3})e^{\alpha z_1 + z_2 + 1} \right] \quad \text{and}$$

is a solution of (2.5).

4. PROOF OF THE MAIN RESULTS

Before we starting the proof of the main results, we present here some necessary lemmas which will play key role to prove the main results of this paper.

Lemma 4.1. [49] *Let $f_j \not\equiv 0$ ($j = 1, 2, \dots, m$; $m \geq 3$) be meromorphic functions such that f_1, \dots, f_{m-1} are not constants, $f_1 + f_2 + \dots + f_m = 1$ and such that*

$$\sum_{j=1}^m \left\{ N_{n-1} \left(r, \frac{1}{f_j} \right) + (m-1) \overline{N}(r, f_j) \right\} < \lambda T(r, f_j) + O(\log^+ T(r, f_j))$$

holds for $j = 1, \dots, m-1$ and all r outside possibly a set with finite logarithmic measure, where $\lambda < 1$ is a positive number. Then $f_m = 1$.

Lemma 4.2. [18] *Let $f_j \not\equiv 0$ ($j = 1, 2, 3$) be meromorphic functions on such that f_1 are not constant, $f_1 + f_2 + f_3 = 1$, and such that*

$$\sum_{j=1}^3 \left\{ N_2 \left(r, \frac{1}{f_j} \right) + 2 \overline{N}(r, f_j) \right\} < \lambda T(r, f_j) + O(\log^+ T(r, f_j))$$

holds for all r outside possibly a set with finite logarithmic measure, where $\lambda < 1$ is a positive number. Then, either $f_2 = 1$ or $f_3 = 1$.

Lemma 4.3. [49] *Let $a_0(z), a_1(z), \dots, a_n(z)$ ($n \geq 1$) be meromorphic functions on and $g_0(z), g_1(z), \dots, g_n(z)$ are entire functions such that $g_j(z) - g_k(z)$ are not constants for $0 \leq j < k \leq n$. If $\sum_{j=0}^n a_j(z) e^{g_j(z)} \equiv 0$, and $T(r, a_j) = o(T(r))$, where $T(r) = \min_{0 \leq j < k \leq n} T(r, e^{g_j - g_k})$ for $j = 0, 1, \dots, n$, then $a_j(z) \equiv 0$ for each $j = 0, 1, \dots, n$.*

Proof of Theorem 3.1. Let $(f(z), g(z))$ be a pair of finite order transcendental entire solution of (3.1). Let us first make a transformation

$$f(z) = \frac{1}{\sqrt{2}}(u + v) \quad \text{and} \quad g(z + c) = \frac{1}{\sqrt{2}}(u - v). \tag{4.1}$$

Then first equation of (3.1) yields

$$(1 + w)u^2 + (1 - w)v^2 = e^{g_1(z)},$$

which can be rewritten as

$$\left(\frac{\sqrt{1+w}u}{e^{g_1(z)/2}} + i \frac{\sqrt{1-w}v}{e^{g_1(z)/2}} \right) \left(\frac{\sqrt{1+w}u}{e^{g_1(z)/2}} - i \frac{\sqrt{1-w}v}{e^{g_1(z)/2}} \right) = 1.$$

In view of the above equation and assumption, it follows that there exists a polynomial $p_1(z)$ in \mathbb{C}^n such that

$$\frac{u\sqrt{1+w}}{e^{g_1(z)/2}} + i \frac{v\sqrt{1-w}}{e^{g_1(z)/2}} = e^{p_1(z)} \quad \text{and} \quad \frac{u\sqrt{1+w}}{e^{g_1(z)/2}} - i \frac{v\sqrt{1-w}}{e^{g_1(z)/2}} = e^{-p_1(z)}. \tag{4.2}$$

Let

$$\gamma_1(z) = p_1(z) + \frac{1}{2}g_1(z), \quad \gamma_2(z) = -p_1(z) + \frac{1}{2}g_1(z). \tag{4.3}$$

Then from (4.2) and (4.3), we obtain

$$u = \frac{1}{2\sqrt{1+w}} [e^{\gamma_1} + e^{\gamma_2}], \quad v = \frac{1}{2i\sqrt{1-w}} [e^{\gamma_1} - e^{\gamma_2}]. \quad (4.4)$$

Therefore, in view of (4.1) and (4.4), we obtain

$$f(z) = \frac{1}{\sqrt{2}} [A_1 e^{\gamma_1} + A_2 e^{\gamma_2}], \quad g(z+c) = \frac{1}{\sqrt{2}} [A_2 e^{\gamma_1} + A_1 e^{\gamma_2}]. \quad (4.5)$$

Similarly, in view of the second equation of (3.1), there exists a polynomial $p_2(z)$ such that

$$g(z) = \frac{1}{\sqrt{2}} [A_1 e^{\gamma_3} + A_2 e^{\gamma_4}], \quad f(z+c) = \frac{1}{\sqrt{2}} [A_2 e^{\gamma_3} + A_1 e^{\gamma_4}], \quad (4.6)$$

where

$$\gamma_3(z) = p_2(z) + \frac{1}{2}g_2(z), \quad \gamma_4(z) = -p_2(z) + \frac{1}{2}g_2(z). \quad (4.7)$$

From (4.5) and (4.6), we easily get

$$\begin{cases} A_1 e^{\gamma_1(z+c)-\gamma_4(z)} + A_2 e^{\gamma_2(z+c)-\gamma_4(z)} - A_2 e^{\gamma_3(z)-\gamma_4(z)} = A_1, \\ A_1 e^{\gamma_3(z+c)-\gamma_2(z)} + A_2 e^{\gamma_4(z+c)-\gamma_2(z)} - A_2 e^{\gamma_1(z)-\gamma_2(z)} = A_1. \end{cases} \quad (4.8)$$

Now we discuss the following four possible cases.

Case 1: Let $\gamma_1(z) - \gamma_2(z) = \zeta_1$ and $\gamma_3(z) - \gamma_4(z) = \zeta_2$, where ζ_1, ζ_2 are two complex constants.

Then, it clearly follows from (4.3) and (4.7) that $p_1(z)$ and $p_2(z)$ both are constants in \mathbb{C} . Let $e^{p_1} = \xi_1$ and $e^{p_2} = \xi_2$.

Then, from (4.5) and (4.6), it follows that

$$\begin{cases} f(z) = \frac{A_1 \xi_1 + A_2 \xi_1^{-1}}{\sqrt{2}} e^{\frac{1}{2}g_1(z)}, \\ g(z+c) = \frac{A_2 \xi_1 + A_1 \xi_1^{-1}}{\sqrt{2}} e^{\frac{1}{2}g_1(z)}, \\ g(z) = \frac{A_1 \xi_2 + A_2 \xi_2^{-1}}{\sqrt{2}} e^{\frac{1}{2}g_2(z)}, \\ f(z+c) = \frac{A_2 \xi_2 + A_1 \xi_2^{-1}}{\sqrt{2}} e^{\frac{1}{2}g_2(z)}. \end{cases} \quad (4.9)$$

Since f, g are transcendental entire, it follows from (4.9) that $A_1 \xi_1 + A_2 \xi_1^{-1}$, $A_2 \xi_1 + A_1 \xi_1^{-1}$, $A_1 \xi_2 + A_2 \xi_2^{-1}$ and $A_2 \xi_2 + A_1 \xi_2^{-1}$ are all non zero constants. Since p_1 and p_2 are constants and f, g are transcendental entire, it follows from (4.9) that g_1 and g_2 are non-constants.

After simple calculations, we have from (4.9) that

$$\begin{cases} (A_1 \xi_1 + A_2 \xi_1^{-1}) e^{\frac{1}{2}[g_1(z+c)-g_2(z)]} = A_2 \xi_2 + A_1 \xi_2^{-1}, \\ (A_1 \xi_2 + A_2 \xi_2^{-1}) e^{\frac{1}{2}[g_2(z+c)-g_1(z)]} = A_2 \xi_1 + A_1 \xi_1^{-1}. \end{cases} \quad (4.10)$$

Since g_1, g_2 are non-constant polynomials in \mathbb{C}^n , from (4.10), we see that $g_1(z+c) - g_2(z) = \eta_1$ and $g_2(z+c) - g_1(z) = \eta_2$, where $\eta_1, \eta_2 \in \mathbb{C}$. Otherwise, left hand sides of both the equations are transcendental, whereas right hand sides will be

polynomial, a contradiction. Therefore, $g_1(z+2c) - g_2(z) = g_2(z+2c) - g_1(z) = \zeta_1 + \zeta_2$. Thus,

$$g_1(z) = L(z) + \Phi(z) + d_1 \quad \text{and} \quad g_2(z) = L(z) + \Phi(z) + d_2,$$

where $L(z) = \sum_{j=1}^n a_j z_j$ and $\Phi(z)$ is a polynomial in \mathbb{C}^n defined as in (3.3), $a_j, d_1, d_2 \in \mathbb{C}$, $j = 1, 2, \dots, n$.

Thus, from (4.10), we get

$$e^{\frac{1}{2}[L(c)+d_1-d_2]} = \frac{A_2\xi_2 + A_1\xi_2^{-1}}{A_1\xi_1 + A_2\xi_1^{-1}} \quad \text{and} \quad e^{\frac{1}{2}[L(c)+d_2-d_1]} = \frac{A_2\xi_1 + A_1\xi_1^{-1}}{A_1\xi_2 + A_2\xi_2^{-1}}, \quad (4.11)$$

which imply

$$e^{L(c)} = \frac{(A_2\xi_2 + A_1\xi_2^{-1})(A_2\xi_1 + A_1\xi_1^{-1})}{(A_1\xi_1 + A_2\xi_1^{-1})(A_1\xi_2 + A_2\xi_2^{-1})} \quad \text{and} \quad e^{d_1-d_2} = \frac{(A_2\xi_2 + A_1\xi_2^{-1})(A_1\xi_2 + A_2\xi_2^{-1})}{(A_1\xi_1 + A_2\xi_1^{-1})(A_2\xi_1 + A_1\xi_1^{-1})}.$$

Therefore, from (4.9), we obtain

$$f(z) = \frac{A_1\xi_1 + A_2\xi_1^{-1}}{\sqrt{2}} e^{\frac{1}{2}[L(z)+\Phi(z)+d_1]} \quad \text{and} \quad g(z) = \frac{A_1\xi_2 + A_2\xi_2^{-1}}{\sqrt{2}} e^{\frac{1}{2}[L(z)+\Phi(z)+d_2]}.$$

Case 2. Let $\gamma_1(z) - \gamma_2(z)$ and $\gamma_3(z) - \gamma_4(z)$ both are non-constants. Then, by Lemma 4.2, it follows from (4.8) that

$$A_1 e^{\gamma_1(z+c)-\gamma_4(z)} = A_1 \quad \text{or} \quad A_2 e^{\gamma_2(z+c)-\gamma_4(z)} = A_1$$

and

$$A_1 e^{\gamma_3(z+c)-\gamma_2(z)} = A_1 \quad \text{or} \quad A_2 e^{\gamma_4(z+c)-\gamma_2(z)} = A_1.$$

Now, we consider the following four possible subcases.

Subcase 2.1. Let

$$A_1 e^{\gamma_1(z+c)-\gamma_4(z)} = A_1 \quad \text{and} \quad A_1 e^{\gamma_3(z+c)-\gamma_2(z)} = A_1. \quad (4.12)$$

Then, from (4.8) and (4.12), we obtain

$$A_2 e^{\gamma_2(z+c)-\gamma_3(z)} = A_2 \quad \text{and} \quad A_2 e^{\gamma_4(z+c)-\gamma_1(z)} = A_2. \quad (4.13)$$

Since γ_j 's, $j = 1, 2, 3, 4$ are all polynomials, it follows from (4.9) and (4.10) that $\gamma_1(z+c) - \gamma_4(z) = \zeta_1$, $\gamma_3(z+c) - \gamma_2(z) = \zeta_2$, $\gamma_2(z+c) - \gamma_3(z) = \zeta_3$ and $\gamma_4(z+c) - \gamma_1(z) = \zeta_4$, $\zeta_j \in \mathbb{C}$ for $j = 1, 2, 3, 4$. This implies that

$$\gamma_1(z+2c) - \gamma_1(z) = \gamma_4(z+2c) - \gamma_4(z) = \zeta_1 + \zeta_2 \quad \text{and}$$

$$\gamma_2(z+2c) - \gamma_2(z) = \gamma_3(z+2c) - \gamma_3(z) = \zeta_3 + \zeta_4.$$

Therefore, we must get

$$\gamma_1(z) = L_1(z) + \Phi(z) + d_1, \quad \gamma_4(z) = L_1(z) + \Phi(z) + d_2,$$

$$\gamma_2(z) = L_2(z) + \Psi(z) + d_3, \quad \text{and} \quad \gamma_3(z) = L_2(z) + \Psi(z) + d_4,$$

where $L_1(z) = \sum_{j=1}^n a_j z_j$, $L_2(z) = \sum_{j=1}^n b_j z_j$, $\Phi(z)$ and $\Psi(z)$ are defined in (3.3) and (3.4), respectively, $a_j, b_j, d_i \in \mathbb{C}$ for $i = 1, 2$ and $j = 1, \dots, n$. As $\gamma_4(z) - \gamma_3(z)$ and $\gamma_2(z) - \gamma_1(z)$ both are non-constants, we must have

$$L_1(z) + \Phi(z) \neq L_2(z) + \Psi(z).$$

Therefore, in view of (4.3) and (4.7), we have

$$g_1(z) = L_1(z) + L_2(z) + \Phi(z) + \Psi(z) + d_1 + d_3 \quad \text{and}$$

$$g_2(z) = L_1(z) + L_2(z) + \Phi(z) + \Psi(z) + d_2 + d_4.$$

Therefore, from (4.12) and (4.13), we obtain

$$\begin{cases} e^{L_1(c)+d_1-d_2} = 1, & e^{L_1(c)+d_2-d_1} = 1, \\ e^{L_2(c)+d_3-d_4} = 1, & e^{L_2(c)+d_4-d_3} = 1. \end{cases} \quad (4.14)$$

After simple computation, we obtain from (4.14) that

$$e^{2L_1(c)} = 1, \quad e^{2L_2(c)} = 1, \quad e^{2(d_1-d_2)} = 1 \quad \text{and} \quad e^{2(d_3-d_4)} = 1,$$

which yield

$$e^{L_1(c)} = \pm 1, \quad e^{d_1-d_2} = \pm 1, \quad e^{L_2(c)} = \pm 1 \quad \text{and} \quad e^{d_3-d_4} = \pm 1.$$

If $e^{L_1(c)} = 1$, then from (4.14), we obtain $e^{d_1-d_2} = 1$. If $e^{L_1(c)} = -1$, then from (4.14), we obtain $e^{d_1-d_2} = -1$. Similarly, If $e^{L_2(c)} = 1$, then from (4.14), it follows that $e^{d_3-d_4} = 1$. If $e^{L_2(c)} = -1$, then it follows from (4.14) that $e^{d_3-d_4} = -1$.

Hence, from (4.5) and (4.6), we get

$$f(z) = \frac{1}{\sqrt{2}} [A_1 e^{L_1(z)+\Phi(z)+d_1} + A_2 e^{L_2(z)+\Psi(z)+d_3}] \quad \text{and}$$

$$g(z) = \frac{1}{\sqrt{2}} [A_1 e^{L_2(z)+\Psi(z)+d_4} + A_2 e^{L_1(z)+\Phi(z)+d_2}].$$

Subcase 2.2. Let

$$A_1 e^{\gamma_1(z+c)-\gamma_4(z)} = A_1 \quad \text{and} \quad A_2 e^{\gamma_4(z+c)-\gamma_2(z)} = A_1. \quad (4.15)$$

Therefore, from (4.3) and (4.15), we obtain

$$A_2 e^{\gamma_2(z+c)-\gamma_3(z)} = A_2 \quad \text{and} \quad A_1 e^{\gamma_3(z+c)-\gamma_1(z)} = A_2. \quad (4.16)$$

Since γ_j 's are polynomials, $j = 1, 2, 3, 4$, it follows from (4.15) and (4.16) that $\gamma_1(z+c) - \gamma_4(z) = \zeta_1$, $\gamma_4(z+c) - \gamma_2(z) = \zeta_2$, $\gamma_2(z+c) - \gamma_3(z) = \zeta_3$ and $\gamma_3(z+c) - \gamma_1(z) = \zeta_4$, $\zeta_j \in \mathbb{C}$ for $j = 1, 2, 3, 4$. This implies that $\gamma_1(z+4c) - \gamma_1(z) = \gamma_2(z+4c) - \gamma_2(z) = \sum_{j=1}^4 \zeta_j$. Hence, we can have $\gamma_1(z) = L(z) + \Phi(z) + k_1$ and $\gamma_2(z) = L(z) + \Phi(z) + k_2$, where $L(z)$ and $\Phi(z)$ are the same as defined in Case 1. But, then we must have $\gamma_2(z) - \gamma_1(z) = k_2 - k_1 \in \mathbb{C}$, a contradiction.

Subcase 2.3. Let

$$A_2 e^{\gamma_2(z+c)-\gamma_4(z)} = A_1 \quad \text{and} \quad A_1 e^{\gamma_3(z+c)-\gamma_2(z)} = A_1.$$

By similar argument as used in **Subcase 2.2**, we can easily get a contradiction.

Subcase 2.4. Let

$$A_2 e^{\gamma_2(z+c)-\gamma_4(z)} = A_1 \quad \text{and} \quad A_2 e^{\gamma_4(z+c)-\gamma_2(z)} = A_1. \quad (4.17)$$

In view of (4.3) and (4.17), it follows that

$$A_1 e^{\gamma_1(z+c)-\gamma_3(z)} = A_2 \quad \text{and} \quad A_1 e^{\gamma_3(z+c)-\gamma_1(z)} = A_2. \quad (4.18)$$

Since $\gamma_1, \gamma_2, \gamma_3, \gamma_4$ are all polynomials, from (4.17) and (4.18), we get $\gamma_2(z+c) - \gamma_4(z) = l_1$, $\gamma_4(z+c) - \gamma_2(z) = l_2$, $\gamma_1(z+c) - \gamma_3(z) = l_3$ and $\gamma_3(z+c) - \gamma_1(z) = l_4$, $l_j \in \mathbb{C}$ for $j = 1, 2, 3, 4$. This implies that $\gamma_1(z+2c) - \gamma_1(z) = \gamma_3(z+2c) - \gamma_3(z) = l_3 + l_4$ and $\gamma_2(z+2c) - \gamma_2(z) = \gamma_4(z+2c) - \gamma_4(z) = l_1 + l_2$.

Therefore, we must have

$$\gamma_1(z) = L_1(z) + \Phi(z) + d_1, \quad \gamma_3(z) = L_1(z) + \Phi(z) + d_2,$$

$$\gamma_2(z) = L_2(z) + \Psi(z) + d_3, \quad \text{and} \quad \gamma_4(z) = L_2(z) + \Psi(z) + d_4,$$

where $L_1(z), L_2(z), \Phi(z), \Psi(z)$ are defined in Subcase 2.1. As $\gamma_2 - \gamma_1$ and $\gamma_4 - \gamma_3$ are non-constants, we have $L_1(z) + \Phi(z) \neq L_2(z) + \Psi(z)$.

Therefore, in view of (4.3) and (4.7), we obtain

$$g_1(z) = L_1(z) + L_2(z) + \Phi(z) + \Psi(z) + d_1 + d_3 \quad \text{and}$$

$$g_2(z) = L_1(z) + L_2(z) + \Phi(z) + \Psi(z) + d_2 + d_4.$$

Hence, from (4.17) and (4.18), we have

$$\begin{cases} A_2 e^{L_2(c)+d_3-d_4} = A_1, & A_2 e^{L_2(c)+d_4-d_3} = A_1 \\ A_1 e^{L_1(c)+d_1-d_2} = A_2, & A_1 e^{L_1(c)+d_2-d_1} = A_2. \end{cases} \quad (4.19)$$

From (4.19), it follows that

$$e^{2L_1(c)} = A_2^2/A_1^2, \quad e^{2L_2(c)} = A_1^2/A_2^2, \quad e^{2(d_1-d_4)} = 1 \quad \text{and} \quad e^{2(d_3-d_4)} = 1,$$

which yield

$$e^{L_1(c)} = \pm A_2/A_1, \quad e^{L_2(c)} = \pm A_1/A_2, \quad e^{d_1-d_4} = \pm 1 \quad \text{and} \quad e^{2(d_3-d_4)} = \pm 1.$$

If $e^{L_1(c)} = A_2/A_1$, then from (4.19), we get $e^{d_1-d_2} = 1$. If $e^{L_1(c)} = -A_2/A_1$, then it follows from (4.19) that $e^{d_1-d_2} = -1$. Similarly, If $e^{L_2(c)} = A_1/A_2$, then from (4.19), we get $e^{d_3-d_4} = 1$. If $e^{L_2(c)} = -A_1/A_2$, then from (4.19), we get $e^{d_3-d_4} = -1$.

Hence, from (4.5) and (4.6), we get

$$f(z) = \frac{1}{\sqrt{2}} [A_1 e^{L_1(z)+\Phi(z)+d_1} + A_2 e^{L_2(z)+\Psi(z)+d_3}] \quad \text{and}$$

$$g(z) = \frac{1}{\sqrt{2}} [A_1 e^{L_1(z)+\Phi(z)+d_2} + A_2 e^{L_2(z)+\Psi(z)+d_4}].$$

Case 3. Let $\gamma_3(z) - \gamma_4(z) = \eta \in \mathbb{C}$ and $\gamma_1(z) - \gamma_2(z)$ be non-constant. Then, in view of (4.7), we see that $p_2(z)$ is a constant.

Since $\gamma_1 - \gamma_2$ is non-constant, in view of Lemma 4.2, it follows from the second equation of (4.8) that either $A_1 e^{\gamma_3(z+c)-\gamma_2(z)} = A_1$ or $A_2 e^{\gamma_4(z+c)-\gamma_2(z)} = A_1$.

First suppose that $A_1 e^{\gamma_3(z+c)-\gamma_2(z)} = A_1$. Then, from the second equation of (4.8), we have $A_2 e^{\gamma_4(z+c)-\gamma_1(z)} = A_2$. As all γ_j , $j = 1, 2, 3, 4$ are polynomials, it follows that $\gamma_3(z+c) - \gamma_2(z) = m_1$ and $\gamma_4(z+c) - \gamma_1(z) = m_2$, where $m_1, m_2 \in \mathbb{C}$. Therefore, $\gamma_1(z) - \gamma_2(z) = m_1 - m_2 - \eta$, a constant in \mathbb{C} , which contradicts to our assumption.

Similarly, we can get a contradiction for the case $A_2 e^{\gamma_4(z+c)-\gamma_2(z)} = A_1$.

Case 4. Let $\gamma_2 - \gamma_1 = \eta_1 \in \mathbb{C}$ and $\gamma_4 - \gamma_3$ be non-constant. Then, by similar argument as used in **Subcase 3**, we easily get a contradiction. \square

Proof of Theorem 3.2. Let (f, g) is a pair of finite transcendental entire solutions of the system (3.2).

Then, in a similar manner as in Theorem 3.1, we obtain that

$$\begin{cases} \frac{\partial^k f}{\partial z_1^k} = \frac{1}{\sqrt{2}} [A_1 e^{\gamma_1(z)} + A_2 e^{\gamma_2(z)}] \\ g(z+c) = \frac{1}{\sqrt{2}} [A_2 e^{\gamma_1(z)} + A_1 e^{\gamma_2(z)}] \\ \frac{\partial^k g}{\partial z_1^k} = \frac{1}{\sqrt{2}} [A_1 e^{\gamma_3(z)} + A_2 e^{\gamma_4(z)}] \\ f(z+c) = \frac{1}{\sqrt{2}} [A_2 e^{\gamma_3(z)} + A_1 e^{\gamma_4(z)}], \end{cases} \quad (4.20)$$

where $\gamma_1, \gamma_2, \gamma_3, \gamma_4$ are defined in (4.3) and (4.7).

Differentiating partially k times with respect to z_1 , second and fourth equations of (4.20), we obtain

$$\begin{cases} \frac{\partial^k f(z+c)}{\partial z_1^k} = \frac{1}{\sqrt{2}} [A_2 h_1(z) e^{\gamma_3(z)} + A_1 h_2(z) e^{\gamma_4(z)}], \\ \frac{\partial^k g(z+c)}{\partial z_1^k} = \frac{1}{\sqrt{2}} [A_2 h_3(z) e^{\gamma_1(z)} + A_1 h_4(z) e^{\gamma_2(z)}], \end{cases} \quad (4.21)$$

where h_1, h_2, h_3, h_4 are given by

$$\begin{cases} h_1(z) = \left(\frac{\partial \gamma_3}{\partial z_1} \right)^k + M_k \left(\frac{\partial^k \gamma_3}{\partial z_1^k}, \dots, \frac{\partial \gamma_3}{\partial z_1} \right), \\ h_2(z) = \left(\frac{\partial \gamma_4}{\partial z_1} \right)^k + N_k \left(\frac{\partial^k \gamma_4}{\partial z_1^k}, \dots, \frac{\partial \gamma_4}{\partial z_1} \right), \\ h_3(z) = \left(\frac{\partial \gamma_1}{\partial z_1} \right)^k + O_k \left(\frac{\partial^k \gamma_1}{\partial z_1^k}, \dots, \frac{\partial \gamma_1}{\partial z_1} \right), \\ h_4(z) = \left(\frac{\partial \gamma_2}{\partial z_1} \right)^k + R_k \left(\frac{\partial^k \gamma_2}{\partial z_1^k}, \dots, \frac{\partial \gamma_2}{\partial z_1} \right), \end{cases}$$

where M_k is the partial differential polynomial in $\frac{\partial \gamma_3}{\partial z_1}$ of order less than k in which $\frac{\partial \gamma_3}{\partial z_1}$ appears in product with at least one more partial derivative of $\gamma_3(z)$ of order greater than or equal to 2. Similar definitions for N_k, O_k and R_k .

Now, from the first and third equations of (4.20) and (4.21), we obtain

$$\begin{cases} A_1 e^{\gamma_1(z+c)-\gamma_4(z)} + A_2 e^{\gamma_2(z+c)-\gamma_4(z)} - A_2 h_1(z) e^{\gamma_3(z)-\gamma_4(z)} = A_1 h_2(z), \\ A_1 e^{\gamma_3(z+c)-\gamma_2(z)} + A_2 e^{\gamma_4(z+c)-\gamma_2(z)} - A_2 h_3(z) e^{\gamma_1(z)-\gamma_2(z)} = A_1 h_4(z). \end{cases} \quad (4.22)$$

Now, we consider the following four possible cases.

Case 1. Let $\gamma_2(z) - \gamma_1(z) = \eta_1$ and $\gamma_4(z) - \gamma_3(z) = \eta_2$, where $\eta_1, \eta_2 \in \mathbb{C}$. Then, in view of (4.3) and (4.7), it follows that $h_1(z)$ and $h_2(z)$ are both constants. Set

$e^{h_1} = \xi_1$ and $e^{h_2} = \xi_2$, ξ_1, ξ_2 are non-zero constants in \mathbb{C} . Therefore, in view of (4.20), we obtain

$$\begin{cases} \frac{\partial^k f}{\partial z_1^k} = \frac{A_1 \xi_1 + A_2 \xi_1^{-1}}{\sqrt{2}} e^{\frac{1}{2} g_1(z)}, \\ g(z+c) = \frac{A_2 \xi_1 + A_1 \xi_1^{-1}}{\sqrt{2}} e^{\frac{1}{2} g_1(z)}, \\ \frac{\partial^k g}{\partial z_1^k} = \frac{A_1 \xi_2 + A_2 \xi_2^{-1}}{\sqrt{2}} e^{\frac{1}{2} g_2(z)}, \\ f(z+c) = \frac{A_2 \xi_2 + A_1 \xi_2^{-1}}{\sqrt{2}} e^{\frac{1}{2} g_2(z)}. \end{cases} \quad (4.23)$$

As f, g are transcendental entire, in view of (4.23), we see that $g_1(z), g_2(z)$ both are non-constant polynomials and $A_1 \xi_1 + A_2 \xi_1^{-1}$, $A_2 \xi_1 + A_1 \xi_1^{-1}$, $A_1 \xi_2 + A_2 \xi_2^{-1}$, $A_2 \xi_2 + A_1 \xi_2^{-1}$ are all non-zero constants.

Also, after simple computations, we obtain from (4.23) that

$$\begin{cases} e^{\frac{1}{2}[g_1(z+c)-g_2(z)]} = \frac{A_2 \xi_2 + A_1 \xi_2^{-1}}{A_1 \xi_1 + A_2 \xi_1^{-1}} r_1(z), \\ e^{\frac{1}{2}[g_2(z+c)-g_1(z)]} = \frac{A_2 \xi_1 + A_1 \xi_1^{-1}}{A_1 \xi_2 + A_2 \xi_2^{-1}} r_2(z) \end{cases} \quad (4.24)$$

with

$$r_1(z) = \left(\frac{1}{2} \frac{\partial g_2(z)}{\partial z_1} \right)^k + G_1 \left(\frac{\partial^k g_2(z)}{\partial z_1^k}, \dots, \frac{\partial g_2(z)}{\partial z_1} \right) \quad \text{and}$$

$$r_2(z) = \left(\frac{1}{2} \frac{\partial g_1(z)}{\partial z_1} \right)^k + G_2 \left(\frac{\partial^k g_1(z)}{\partial z_1^k}, \dots, \frac{\partial g_1(z)}{\partial z_1} \right),$$

where G_1 is a differential polynomial in $g_2(z)$ of order less than k in which $g_2'(z)$ appears in the product with at least one more partial derivative of order greater than or equal to 2, and similar definition for G_2 .

Since $g_1(z), g_2(z)$ are non-constant polynomials, from (4.24), conclude that $r_1, r_2, g_1(z+c)-g_2(z)$ and $g_2(z+c)-g_1(z)$ are all constants. Let $g_1(z+c)-g_2(z) = \eta_3$ and $g_2(z+c)-g_1(z) = \eta_4$, $\eta_3, \eta_4 \in \mathbb{C}$. This implies that $g_1(z+2c)-g_1(z) = g_2(z+2c)-g_2(z) = \eta_3 + \eta_4$. Hence, we can get

$$g_1(z_1, z_2) = \mathcal{L}(z) + \chi(z_2) + d_1, \quad g_2(z_1, z_2) = \mathcal{L}(z) + \chi(z_2) + d_2,$$

where $\mathcal{L}(z) = a_1 z_1 + a_2 z_2$, $\chi(z_2)$ is a polynomial in z_2 such that $\chi(z_2 + c_2) = \chi(z_2)$, $a_1, a_2, d_1, d_2 \in \mathbb{C}$. Since, r_1, r_2 both are constants, it follows that

$$r_1 = r_2 = \left(\frac{a_1}{2} \right)^k.$$

Therefore, from (4.24), we see that

$$\begin{cases} e^{\frac{1}{2}[\mathcal{L}(c)+d_1-d_2]} = \left(\frac{a_1}{2}\right)^k \frac{A_2\xi_2 + A_1\xi_2^{-1}}{A_1\xi_1 + A_2\xi_1^{-1}}, \\ e^{\frac{1}{2}[\mathcal{L}(c)+d_2-d_1]} = \left(\frac{a_1}{2}\right)^k \frac{A_2\xi_1 + A_1\xi_1^{-1}}{A_1\xi_2 + A_2\xi_2^{-1}}. \end{cases} \quad (4.25)$$

Hence, from (4.25), it follows that

$$e^{\mathcal{L}(c)} = \left(\frac{a_1}{2}\right)^{2k} \frac{(A_2\xi_2 + A_1\xi_2^{-1})(A_2\xi_1 + A_1\xi_1^{-1})}{(A_1\xi_1 + A_2\xi_1^{-1})(A_1\xi_2 + A_2\xi_2^{-1})}.$$

Thus, from (4.23), we get

$$f(z_1, z_2) = \frac{A_2\xi_2 + A_1\xi_2^{-1}}{\sqrt{2}} e^{\frac{1}{2}[\mathcal{L}(z)+\chi(z_2)-\mathcal{L}(c)+d_2]} \quad \text{and}$$

$$g(z_1, z_2) = \frac{A_2\xi_1 + A_1\xi_1^{-1}}{\sqrt{2}} e^{\frac{1}{2}[\mathcal{L}(z)+\chi(z_2)-\mathcal{L}(c)+d_1]}.$$

Case 2. Let $\gamma_1(z) - \gamma_2(z)$ and $\gamma_3(z) - \gamma_4(z)$ both are non-constant.

Observe that $h_1(z)$ and $h_2(z)$ both can not be simultaneously zero. Otherwise, from the first equation of (4.22), we obtain $e^{\gamma_1(z+c)-\gamma_2(z+c)} = -1$, which is a contradiction as $\gamma_2(z) - \gamma_1(z)$ is non-constant.

Now, let $h_1(z) \not\equiv 0$ and $h_2(z) \equiv 0$. Then, first equation of (4.22) reduces to

$$A_1 e^{\gamma_1(z+c)-\gamma_3(z)} + A_2 e^{\gamma_2(z+c)-\gamma_3(z)} = A_2 h_1(z). \quad (4.26)$$

As $\gamma_2(z) - \gamma_1(z)$ is a non-constant polynomial, in view of (4.26), it follows that $\gamma_1(z+c) - \gamma_3(z)$ and $\gamma_2(z+c) - \gamma_3(z)$ both are non-constants.

Rewrite (4.26) as

$$A_1 e^{\gamma_1(z+c)} + A_2 e^{\gamma_2(z+c)} - A_2 h_1 e^{\gamma_3(z)} \equiv 0. \quad (4.27)$$

Therefore, in view of Lemma 4.3, we can easily get a contradiction from (4.27).

In a similar manner, we can get a contradiction when $h_1(z) \equiv 0$ and $h_2(z) \not\equiv 0$. Hence, $h_1(z) \not\equiv 0$ and $h_2(z) \not\equiv 0$. By similar arguments, we also get $h_3(z) \not\equiv 0$ and $h_4(z) \not\equiv 0$.

Therefore, by (4.22) and Lemma 4.2, we obtain

$$\text{either } A_1 e^{\gamma_1(z+c)-\gamma_4(z)} = A_1 h_2(z), \quad \text{or } A_2 e^{\gamma_2(z+c)-\gamma_4(z)} = A_1 h_2(z)$$

and

$$\text{either } A_1 e^{\gamma_3(z+c)-\gamma_2(z)} = A_1 h_4(z), \quad \text{or } A_2 e^{\gamma_4(z+c)-\gamma_2(z)} = A_1 h_4(z).$$

Now, we discuss the four possible subcases.

Subcase 2.1. Let

$$A_1 e^{\gamma_1(z+c)-\gamma_4(z)} = A_1 h_2(z) \quad \text{and} \quad A_1 e^{\gamma_3(z+c)-\gamma_2(z)} = A_1 h_4(z). \quad (4.28)$$

Then, by (4.22) and (4.28), we get

$$A_2 e^{\gamma_2(z+c)-\gamma_3(z)} = A_2 h_1(z) \quad \text{and} \quad A_2 e^{\gamma_4(z+c)-\gamma_1(z)} = A_2 h_3(z). \quad (4.29)$$

Since γ_j 's are polynomials, $j = 1, \dots, 4$, (4.28) and (4.29) yield that $h_1, h_2, h_3, h_4, \gamma_1(z+c) - \gamma_4(z), \gamma_3(z+c) - \gamma_2(z), \gamma_2(z+c) - \gamma_3(z), \gamma_4(z+c) - \gamma_1(z)$ are all constants. Let $\gamma_1(z+c) - \gamma_4(z) = k_1, \gamma_3(z+c) - \gamma_2(z) = k_2, \gamma_2(z+c) - \gamma_3(z) = k_3$ and $\gamma_4(z+c) - \gamma_1(z) = k_4$, where $k_j \in \mathbb{C}, j = 1, \dots, 4$. These imply $\gamma_1(z+2c) - \gamma_1(z) = \gamma_4(z+2c) - \gamma_4(z) = k_1 + k_4$ and $\gamma_2(z+2c) - \gamma_2(z) = \gamma_3(z+2c) - \gamma_3(z) = k_2 + k_3$. Thus, we have

$$\gamma_1(z) = L_1(z) + \Phi_1(z_2) + d_1, \quad \gamma_4(z) = L_1(z) + \Phi_1(z_2) + d_2,$$

$$\gamma_2(z) = L_2(z) + \Psi_1(z_2) + d_3, \quad \text{and} \quad \gamma_3(z) = L_2(z) + \Psi_1(z_2) + d_4,$$

where $L_1(z) = a_1z_1 + a_2z_2, L_2(z) = b_1z_1 + b_2z_2, \Phi_1(z_2)$ and $\Psi_1(z_2)$ are polynomials in z_2 such that $\Phi_1(z_2 + c_2) = \Phi_1(z_2)$ and $\Psi_1(z_2 + c_2) = \Psi_1(z_2), a_j, b_j, e_j, t_j \in \mathbb{C}$ for $j = 1, 2$.

Since $\gamma_1(z) - \gamma_2(z)$ and $\gamma_3(z) - \gamma_4(z)$ are non-constants, we conclude that

$$L_1(z) + \Phi_1(z_2) \neq L_2(z) + \Psi_1(z_2).$$

Therefore, in view of (4.3) and (4.7), we have

$$\begin{cases} g_1(z_1, z_2) = L_1(z) + L_2(z) + \Phi_1(z_2) + \Psi_1(z_2) + d_1 + d_3, \\ g_2(z_1, z_2) = L_1(z) + L_2(z) + \Phi_1(z_2) + \Psi_1(z_2) + d_2 + d_4. \end{cases} \quad (4.30)$$

Hence, from (4.28) and (4.29), we obtain

$$\begin{cases} e^{L_1(c)+d_1-d_2} = a_1^k, & e^{L_1(c)+d_2-d_1} = a_1^k, \\ e^{L_2(c)+d_3-d_4} = b_1^k, & e^{L_2(c)+d_4-d_3} = b_1^k. \end{cases} \quad (4.31)$$

From (4.31), we obtain

$$e^{2L_1(c)} = a_1^{2k}, \quad e^{2(d_1-d_2)} = 1, \quad e^{2L_2(c)} = b_1^{2k} \quad \text{and} \quad e^{2(d_3-d_4)} = 1.$$

Therefore, we have $e^{L_1(c)} = \pm a_1^k, e^{L_2(c)} = \pm b_1^k, e^{d_1-d_2} = \pm 1$ and $e^{d_3-d_4} = \pm 1$.

If $e^{L_1(c)} = a_1^k$, then from (4.31), we get $e^{d_1-d_2} = 1$. If $e^{L_1(c)} = -a_1^k$, then from (4.31), we get $e^{d_1-d_2} = -1$. If $e^{L_2(c)} = b_1^k$, then from (4.31), we get $e^{d_3-d_4} = 1$ and if $e^{L_2(c)} = -b_1^k$, then we get $e^{d_3-d_4} = -1$.

Thus, from fourth and second equations of (4.20), we obtain

$$\begin{cases} f(z) = \frac{1}{\sqrt{2}} [A_2 e^{L_2(z)+\Psi_1(z_2)-L_2(c)+d_4} + A_1 e^{L_1(z)+\Phi_1(z_2)-L_1(c)+d_2}] \\ g(z) = \frac{1}{\sqrt{2}} [A_2 e^{L_1(z)+\Phi_1(z_2)-L_1(c)+d_1} + A_1 e^{L_2(z)+\Psi_1(z_2)-L_2(c)+d_3}]. \end{cases}$$

Subcase 2.2. Let

$$A_1 e^{\gamma_1(z+c)-\gamma_4(z)} = A_1 h_2(z) \quad \text{and} \quad A_2 e^{\gamma_4(z+c)-\gamma_2(z)} = A_1 h_4(z). \quad (4.32)$$

Then, in view of (4.22) and (4.32), we obtain

$$A_2 e^{\gamma_2(z+c)-\gamma_3(z)} = A_2 h_1(z) \quad \text{and} \quad A_1 e^{\gamma_3(z+c)-\gamma_1(z)} = A_2 h_3(z). \quad (4.33)$$

Since $\gamma_1(z), \gamma_2(z), \gamma_3(z), \gamma_4(z)$ are polynomials, we conclude from (4.32) and (4.33) that $\gamma_1(z+c) - \gamma_3(z) = m_1, \gamma_4(z+c) - \gamma_1(z) = m_2, \gamma_2(z+c) - \gamma_4(z) = m_3$ and $\gamma_3(z+c) - \gamma_2(z) = m_4$, where $m_j \in \mathbb{C}, j = 1, \dots, 4$. These imply $\gamma_1(z+2c) - \gamma_1(z) = \gamma_2(z+4c) - \gamma_2(z) = \sum_{j=1}^4 m_j$. Therefore, $\gamma_1(z) = L_1(z) + \Phi_1(z_2) + d_1$

and $\gamma_2(z) = L_1(z) + \Phi_1(s_1) + d_2$, where $L_1(z)$ and $\Phi_1(s_1)$ are defined in Subcase 2.1, $d_1, d_2 \in \mathbb{C}$. Hence, $\gamma_1(z) - \gamma_2(z) = d_1 - d_2 \in \mathbb{C}$, a contradiction.

Subcase 2.3. Let

$$A_2 e^{\gamma_2(z+c)-\gamma_4(z)} = A_1 h_2(z) \quad \text{and} \quad A_1 e^{\gamma_3(z+c)-\gamma_2(z)} = A_1 h_4(z).$$

Then, by similar arguments as in Subcase 2.2, we easily get a contradiction.

Subcase 2.4. Let

$$A_2 e^{\gamma_2(z+c)-\gamma_4(z)} = A_1 h_2(z) \quad \text{and} \quad A_2 e^{\gamma_4(z+c)-\gamma_2(z)} = A_1 h_4(z). \quad (4.34)$$

Then, by (4.22) and (4.34), we get

$$A_1 e^{\gamma_1(z+c)-\gamma_3(z)} = A_2 h_1(z) \quad \text{and} \quad A_1 e^{\gamma_3(z+c)-\gamma_1(z)} = A_2 h_3(z). \quad (4.35)$$

As $\gamma_1(z), \gamma_2(z), \gamma_3(z), \gamma_4(z)$ are polynomials, from (4.34) and (4.35), we conclude that $h_1, h_2, h_3, h_4, \gamma_2(z+c) - \gamma_4(z), \gamma_4(z+c) - \gamma_2(z), \gamma_1(z+c) - \gamma_3(z), \gamma_3(z+c) - \gamma_1(z)$ are all constants. Let $\gamma_2(z+c) - \gamma_4(z) = n_1, \gamma_4(z+c) - \gamma_2(z) = n_2, \gamma_1(z+c) - \gamma_3(z) = n_3$ and $\gamma_3(z+c) - \gamma_1(z) = n_4$, where $n_j \in \mathbb{C}, j = 1, \dots, 4$. Thus, we get $\gamma_1(z+2c) - \gamma_1(z) = \gamma_3(z+2c) - \gamma_3(z) = n_1 + n_2$ and $\gamma_2(z+2c) - \gamma_2(z) = \gamma_4(z+2c) - \gamma_4(z) = n_3 + n_4$. Hence, by similar arguments as in Subcase 2.1, we obtain

$$\gamma_1(z) = L_1(z) + \Phi_1(z_2) + d_1, \quad \gamma_3(z) = L_1(z) + \Phi_1(z_2) + d_2,$$

$$\gamma_2(z) = L_2(z) + \Psi_1(z_2) + d_3, \quad \text{and} \quad \gamma_4(z) = L_2(z) + \Psi_1(z_2) + d_4,$$

where $L(z), L_2(z), \Phi_1(z_2), \Psi_1(z_2)$ are same as in Subcase 2.1 with $L_1(z) + \Phi_1(z_2) \neq L_2(z) + \Psi_1(z_2)$. Thus, in view of (4.3) and (4.7), we obtain the same form of $g_1(z_1, z_2)$ and $g_2(z_1, z_2)$ as in (4.30).

Therefore, in view of (4.34) and (4.35), we have

$$\begin{cases} A_2 e^{L_2(c)+d_3-d_4} = A_1 b_1^k, & A_2 e^{L_2(c)+d_4-d_3} = A_1 b_1^k, \\ A_1 e^{L_1(c)+d_1-d_2} = A_2 a_1^k, & A_1 e^{L_1(c)+d_2-d_1} = A_2 a_1^k. \end{cases} \quad (4.36)$$

From (4.36), we obtain

$$e^{2L_1(c)} = \frac{A_2^2}{A_1^2} a_1^{2k} \quad e^{2(d_1-d_2)} = 1, \quad e^{2L_2(c)} = \frac{A_1^2}{A_2^2} b_1^{2k} \quad \text{and} \quad e^{2(d_3-d_4)} = 1,$$

which imply

$$e^{L_1(c)} = \pm \frac{A_2}{A_1} a_1^k \quad e^{d_1-d_2} = \pm 1, \quad e^{L_2(c)} = \pm \frac{A_1}{A_2} b_1^k \quad \text{and} \quad e^{d_3-d_4} = \pm 1.$$

If $e^{L_1(c)} = A_2 a_1^k / A_1$, then from (4.36), we get $e^{d_1-d_2} = 1$. If $e^{L_1(c)} = -A_2 a_1^k / A_1$, then we have $e^{d_1-d_2} = -1$. If $e^{L_2(c)} = A_1 b_1^k / A_2$, then from (4.36), we get $e^{d_3-d_4} = 1$. If $e^{L_2(c)} = -A_1 b_1^k / A_2$, then we have $e^{d_3-d_4} = -1$.

Hence, from fourth and second equations of (4.20), we get

$$\begin{cases} f(z) = \frac{1}{\sqrt{2}} [A_2 e^{L_1(z)+\Phi_1(z_2)-L_1(c)+d_2} + A_1 e^{L_2(z)+\Psi_1(z_2)-L_2(c)+d_4}], \\ g(z) = \frac{1}{\sqrt{2}} [A_2 e^{L_1(z)+\Phi_1(z_2)-L_1(c)+d_1} + A_1 e^{L_2(z)+\Psi_1(z_2)-L_2(c)+d_3}]. \end{cases}$$

Case 3. Let $\gamma_3(z) - \gamma_4(z) = \eta \in \mathbb{C}$ and $\gamma_1(z) - \gamma_2(z)$ be non-constant. Then the first equation of (4.22) reduces to

$$A_1 e^{\gamma_1(z+c)-\gamma_4(z)} + A_2 e^{\gamma_2(z+c)-\gamma_4(z)} = A_1 h_2(z) + A_2 h_1(z) e^\eta. \quad (4.37)$$

Note that $A_1 h_2(z) + A_2 h_1(z) e^\eta \not\equiv 0$. Otherwise from (4.37), we have

$$A_1 e^{\gamma_1(z+c)-\gamma_2(z+c)} = -A_2,$$

which implies that $\gamma_1(z+c) - \gamma_2(z+c)$ and hence $\gamma_1(z) - \gamma_2(z)$ becomes constant. This is a contradiction.

Now, in view of (4.37), it follows that

$$T(r, e^{\gamma_1(z+c)-\gamma_4(z)}) = T(r, e^{\gamma_2(z+c)-\gamma_4(z)}) + O(1). \quad (4.38)$$

We note that

$$N(r, e^{\gamma_1(z+c)-\gamma_4(z)}) = N\left(r, \frac{1}{e^{\gamma_1(z+c)-\gamma_4(z)}}\right) = S(r, e^{\gamma_1(z+c)-\gamma_4(z)})$$

and

$$N\left(r, \frac{1}{e^{\gamma_1(z+c)-\gamma_4(z)} - w}\right) = N\left(r, \frac{1}{e^{\gamma_2(z+c)-\gamma_4(z)}}\right) = S(r, e^{\gamma_2(z+c)-\gamma_4(z)}),$$

where $w = h_2 + A_2 h_1 e^\eta / A_1$.

Now by the second fundamental theorem of Nevanlinna for several complex variables, we get

$$\begin{aligned} & T(r, e^{\gamma_1(z+c)-\gamma_4(z)}) \\ & \leq N(r, e^{\gamma_1(z+c)-\gamma_4(z)}) + N\left(r, \frac{1}{e^{\gamma_1(z+c)-\gamma_4(z)}}\right) + N\left(r, \frac{1}{e^{\gamma_1(z+c)-\gamma_4(z)} - w}\right) \\ & \quad + S(r, e^{\gamma_1(z+c)-\gamma_4(z)}) \\ & \leq S(r, e^{\gamma_1(z+c)-\gamma_4(z)}) + S(r, e^{\gamma_2(z+c)-\gamma_4(z)}). \end{aligned}$$

This implies that $\gamma_1(z+c) - \gamma_4(z)$ is constant. In view of (4.38), we see that $\gamma_2(z+c) - \gamma_4(z)$ is also constant. These imply $\gamma_1(z+c) - \gamma_2(z+c)$ and hence $\gamma_1(z) - \gamma_2(z)$ is a constant. This is a contradiction.

Case 4. Let $\gamma_1 - \gamma_2 = \eta_1 \in \mathbb{C}$ and $\gamma_3 - \gamma_4$ be non-constant. Then, by similar arguments as used in Subcase 3, we can get a contradiction. So, we omit the details. \square

Proof of Theorem 3.3. Let $(f(z), g(z))$ be a pair of finite order transcendental entire solution of (2.5).

Then, by similar argument as in Theorem 3.1, we obtain

$$\begin{cases} \frac{\partial^k f}{\partial z_1^k} = \frac{1}{\sqrt{2}} [A_1 e^{p_1(z)} + A_2 e^{-p_1(z)}], \\ g(z+c) - g(z) = \frac{1}{\sqrt{2}} [A_2 e^{p_1(z)} + A_1 e^{-p_1(z)}], \\ \frac{\partial^k g}{\partial z_1^k} = \frac{1}{\sqrt{2}} [A_1 e^{p_2(z)} + A_2 e^{-p_2(z)}], \\ f(z+c) - f(z) = \frac{1}{\sqrt{2}} [A_2 e^{p_2(z)} + A_1 e^{-p_2(z)}], \end{cases} \quad (4.39)$$

where $p_1(z)$ and $p_2(z)$ are two non-constant polynomials.

Differentiating fourth equation of (4.39) k times, we get

$$\frac{\partial^k f(z+c)}{\partial z_1^k} - \frac{\partial^k f}{\partial z_1^k} = \frac{1}{\sqrt{2}} [A_2 H_1 e^{p_2(z)} + A_1 H_2 e^{-p_2(z)}], \quad (4.40)$$

where $H_1 = \left(\frac{\partial p_2}{\partial z_1}\right)^k + M_k \left(\frac{\partial^k p_2}{\partial z_1^k}, \dots, \frac{\partial p_2}{\partial z_1}\right)$ and $H_2 = \left(-\frac{\partial p_2}{\partial z_1}\right)^k + N_k \left(\frac{\partial^k p_2}{\partial z_1^k}, \dots, \frac{\partial p_2}{\partial z_1}\right)$, where M_k is a partial differential polynomial in $p_2(z)$ of degree less than k in which $\frac{\partial p_2}{\partial z_1}$ appears in the product with at least one more partial derivative of p_2 of order greater than 1. Similar definition is for N_k .

Now in view of the first equation of (4.39) and (4.40), we obtain

$$\begin{aligned} A_1 e^{p_1(z+c)+p_2(z)} + A_2 e^{-p_1(z+c)+p_2(z)} - A_1 e^{p_1(z)+p_2(z)} - A_2 e^{-p_1(z)+p_2(z)} \\ - A_2 H_1 e^{2p_2(z)} = A_1 H_2. \end{aligned} \quad (4.41)$$

Similarly in view of the second and third equations of (4.39), we obtain

$$\begin{aligned} A_1 e^{p_2(z+c)+p_1(z)} + A_2 e^{-p_2(z+c)+p_1(z)} - A_1 e^{p_2(z)+p_1(z)} - A_2 e^{-p_2(z)+p_1(z)} \\ - A_2 H_3 e^{2p_1(z)} = A_1 H_4, \end{aligned} \quad (4.42)$$

where $H_3 = \left(\frac{\partial p_1}{\partial z_1}\right)^k + O_k \left(\frac{\partial^k p_1}{\partial z_1^k}, \dots, \frac{\partial p_1}{\partial z_1}\right)$ and $H_4 = \left(-\frac{\partial p_1}{\partial z_1}\right)^k + R_k \left(\frac{\partial^k p_1}{\partial z_1^k}, \dots, \frac{\partial p_1}{\partial z_1}\right)$, where O_k is a differential polynomial in $p_1(z)$ of degree less than k in which $\frac{\partial p_1}{\partial z_1}$ appears in the product with at least one higher order derivative of p_1 . Similar definition is for R_k .

Now, we consider the following two possible cases.

Case 1. Let $p_2(z) - p_1(z) = \eta$, $\eta \in \mathbb{C}$. Then (4.41) and (4.42) yield

$$\begin{cases} A_1 e^\eta e^{p_1(z+c)+p_1(z)} + A_2 e^\eta e^{p_1(z)-p_1(z+c)} - e^\eta (A_1 + A_2 H_1 e^\eta) e^{2p_1(z)} \\ = A_1 H_2 + A_2 e^\eta, \\ A_1 e^\eta e^{p_1(z+c)+p_1(z)} + A_2 e^{-\eta} e^{p_1(z)-p_1(z+c)} - (A_1 e^\eta + A_2 H_3) e^{2p_1(z)} \\ = A_1 H_4 + A_2 e^{-\eta}. \end{cases} \quad (4.43)$$

Note that $A_1 + A_2 H_1 e^\eta$ and $A_1 H_2 + A_2 e^\eta$ can not be zero simultaneously. Otherwise, from the first equation of (4.43), we get $p_1(z+c)$, and hence $p_1(z)$ is constant, a contradiction.

Next, suppose that $A_1 + A_2H_1e^\eta \neq 0$ and $A_1H_2 + A_2e^\eta \equiv 0$. The the first equation of (4.43) reduces to

$$A_1e^{p_1(z+c)-p_1(z)} + A_2e^{-(p_1(z)+p_1(z+c))} = A_1 + A_2H_1e^\eta. \quad (4.44)$$

Now, observe that

$$N\left(r, e^{-(p_1(z+c)+p_1(z))}\right) = N\left(r, \frac{1}{e^{-(p_1(z+c)+p_1(z))}}\right) = S\left(r, e^{-(p_1(z+c)+p_1(z))}\right)$$

and in view of (4.44), we have

$$N\left(r, \frac{1}{e^{-(p_1(z+c)+p_1(z))} - w}\right) = N\left(r, \frac{1}{e^{p_1(z+c)-p_1(z)}}\right) = S\left(r, e^{-(p_1(z+c)+p_1(z))}\right),$$

where $w = (A_1 + A_2H_1e^\eta)/A_2$.

By the second fundamental theorem of Nevanlinna, we obtain

$$\begin{aligned} & T\left(r, e^{-(p_1(z+c)+p_1(z))}\right) \\ & \leq \overline{N}\left(r, e^{-(p_1(z+c)+p_1(z))}\right) + \overline{N}\left(r, \frac{1}{e^{-(p_1(z+c)+p_1(z))}}\right) \\ & \quad + \overline{N}\left(r, \frac{1}{e^{-(p_1(z+c)+p_1(z))} - w}\right) + S\left(r, e^{-(p_1(z+c)+p_1(z))}\right) \\ & \leq S\left(r, e^{-(p_1(z+c)+p_1(z))}\right) + S\left(r, e^{p_1(z+c)-p_1(z)}\right). \end{aligned}$$

This implies that $p_1(z+c) + p_1(z)$ is constant, and hence $p_1(z)$ is constant, which is a contradiction.

Similarly, we can get a contradiction for the case $A_1 + A_2H_1e^\eta \equiv 0$ and $A_1H_2 + A_2e^\eta \neq 0$. Hence, $A_1 + A_2H_1e^\eta \neq 0$ and $A_1H_2 + A_2e^\eta \neq 0$.

In a similar manner, we can show that $A_1e^\eta + A_2H_3 \neq 0$ and $A_1H_4 + A_2e^{-\eta} \neq 0$.

Therefore, in view of Lemma 4.1 and (4.43), we obtain

$$\begin{cases} A_2e^\eta e^{p_1(z)-p_1(z+c)} = A_1H_2 + A_2e^\eta, \\ A_2e^{-\eta} e^{p_1(z)-p_1(z+c)} = A_1H_4 + A_2e^{-\eta}. \end{cases} \quad (4.45)$$

By (4.43) and (4.45), we get

$$\begin{cases} A_1e^{p_1(z+c)-p_1(z)} = A_1 + A_2H_1e^\eta, \\ A_1e^\eta e^{p_1(z+c)-p_1(z)} = A_2H_3 + A_1e^\eta. \end{cases} \quad (4.46)$$

Since $p_1(z)$ is a non-constant polynomial, it follows from (4.45) and (4.46) that $p_1(z+c) - p_1(z)$, H_1, H_2, H_3, H_4 are all constants. Otherwise, L.H.S. of both the equations of (4.45) and (4.46) are transcendental entire, whereas R.H.S.'s are polynomials. Therefore, we must have $p_1(z) = L(z) + \Phi_1(z_2) + \gamma$, where $L(z) = \alpha z_1 + \beta z_2$, $\Phi_1(z_2)$ is a polynomial in z_2 with $\Phi_1(z_2 + c_2) = \Phi_1(z_2)$, $\alpha, \beta, \gamma \in \mathbb{C}$. Thus, $p_2(z) = L(z) + \Phi_1(z_2) + \gamma + \eta$.

Therefore, in view of (4.45), (4.46) and the form of H_1 , H_2 , H_3 and H_4 , we obtain

$$\begin{cases} A_2 e^\eta e^{-\alpha c} = A_1 (-\alpha)^k + A_2 e^\eta, & A_2 e^{-\eta} e^{-\alpha c} = A_1 (-\alpha)^k + A_2 e^{-\eta}, \\ A_1 e^{\alpha c} = A_1 + A_2 \alpha^k e^\eta, & A_1 e^\eta e^{\alpha c} = A_2 \alpha^k + A_1 e^\eta. \end{cases} \quad (4.47)$$

From (4.47), we obtain

$$e^{L(c)} = \frac{A_2 e^\eta}{(-1)^k A_1 \alpha^k + A_2 e^\eta}, \quad e^{2\eta} = 1 \quad \text{and} \quad \alpha^k = -\frac{(-1)^k A_1^2 + A_2^2}{(-1)^k A_1 A_2 e^\eta}$$

Thus, from (4.39), we obtain

$$f(z) = \frac{A_1 e^{L(z)+\Phi_1(z_2)+\gamma} + A_2 e^{-(L(z)+\Phi_1(z_2)+\gamma)}}{\sqrt{2}\alpha^k} \quad \text{and}$$

$$g(z) = \frac{A_1 e^{L(z)+\Phi_1(z_2)+\gamma+\eta} + A_2 e^{-(L(z)+\Phi_1(z_2)+\gamma+\eta)}}{\sqrt{2}\alpha^k}.$$

Case 2. Let $p_2(z) - p_1(z)$ be non-constant.

Now we consider two possible subcases.

Subcase 2.1. Let $p_2(z) + p_1(z) = \eta$, where η is a constant in \mathbb{C} .

Then, (4.41) and (4.42) yield

$$\begin{cases} A_1 e^\eta e^{p_1(z+c)-p_1(z)} + A_2 e^\eta e^{-(p_1(z+c)+p_1(z))} - A_2 e^\eta (1 + H_1 e^\eta) e^{-2p_1(z)} \\ = A_1 (H_2 + e^\eta), \\ A_1 e^\eta e^{p_1(z)-p_1(z+c)} + A_2 e^{-\eta} e^{p_1(z+c)+p_1(z)} - A_2 (H_3 + e^{-\eta}) e^{2p_1(z)} \\ = A_1 (H_4 + e^\eta). \end{cases} \quad (4.48)$$

Now, by similar argument as in Case 1, we can prove that $1 + H_1 e^\eta$, $H_2 + e^\eta$, $H_3 + e^{-\eta}$ and $H_4 + e^\eta$ are all non-zero.

Therefore, by Lemma 4.1 and (4.48), we have

$$e^{p_1(z+c)-p_1(z)} = H_2 e^{-\eta} + 1 \quad \text{and} \quad e^{p_1(z)-p_1(z+c)} = H_4 e^{-\eta} + 1. \quad (4.49)$$

From (4.48) and (4.49), we obtain

$$e^{-p_1(z+c)+p_1(z)} = 1 + H_1 e^\eta \quad \text{and} \quad e^{p_1(z+c)-p_1(z)} = 1 + e^\eta H_3. \quad (4.50)$$

Then by similar arguments as in the Case 1, we obtain that

$$p_1(z) = L(z) + \Phi_1(z_2) + \gamma \quad \text{and} \quad p_2(z) = -(L(z) + \Phi_1(z_2) + \gamma) + \eta.$$

Therefore, from (4.49) and (4.50), we obtain

$$\begin{cases} e^{L(c)} = 1 + e^{-\eta} \alpha^k, & e^{-L(c)} = e^{-\eta} (-\alpha)^k + 1, \\ e^{-L(c)} = 1 + e^\eta (-\alpha)^k, & e^{L(c)} = 1 + e^\eta \alpha^k. \end{cases} \quad (4.51)$$

From first and fourth equations of (4.51), we get

$$e^{2\eta} = 1.$$

From the above two equations of (4.51), we obtain

$$(-1)^k e^{-\eta} \alpha^k + 1 + (-1)^k = 0. \quad (4.52)$$

If k is odd, then in view of (4.52), we can get a contradiction. If k is even, then from (4.52), we get

$$\alpha^k = -2e^\eta.$$

Therefore, from (4.39), we have

$$f(z) = \frac{A_1 e^{L(z)+\Phi_1(z_2)+\gamma} + A_2 e^{-(L(z)+\Phi_1(z_2)+\gamma)}}{\sqrt{2}\alpha^k},$$

$$g(z) = \frac{A_1 e^{-(L(z)+\Phi_1(z_2)+\gamma)+\eta} + A_2 e^{L(z)+\Phi_1(z_2)+\gamma-\eta}}{\sqrt{2}\alpha^k}.$$

Subcase 2.2. Let $p_2(z) + p_1(z)$ be a non-constant polynomial.

Now we consider four possible cases below.

Subcase 2.2.1. Let $p_1(z+c) + p_2(z) = \zeta_1$ and $p_2(z+c) + p_1(z) = \zeta_2$, where $\zeta_1, \zeta_2 \in \mathbb{C}$. Then, we easily see that $p_1(z+2c) - p_1(z) = \zeta_1 - \zeta_2$ and $p_2(z+2c) - p_2(z) = \zeta_2 - \zeta_1$. This implies that $p_1(z) = \alpha z + \beta_1$ and $p_2(z) = -\alpha z + \beta_2$, where $\alpha, \beta_1, \beta_2 \in \mathbb{C}$. But, then $p_1(z) + p_2(z) = \beta_1 + \beta_2 = \text{constant}$, a contradiction.

Subcase 2.2.2. Let $p_1(z+c) + p_2(z) = \zeta_1 \in \mathbb{C}$ and $p_2(z+c) + p_1(z)$ is non-constant.

Then, by Lemma 4.1 and (4.42), we get $A_2 e^{-p_2(z+c)+p_1(z)} = A_1 H_4$. This implies that $-p_2(z+c) + p_1(z) = \zeta_2 \in \mathbb{C}$. Therefore, we must have $p_1(z+2c) + p_1(z) = \zeta_1 + \zeta_2$, and hence $p_1(z)$ must be a constant, a contradiction.

Subcase 2.2.3. Let $p_2(z+c) + p_1(z) = \zeta_1 \in \mathbb{C}$ and $p_1(z+c) + p_2(z)$ is non-constant.

Then, by similar argument as in subcase 2.2.2, we obtain a contradiction.

Subcase 2.2.4. Let $p_2(z+c)+p_1(z)$ and $p_1(z+c)+p_2(z)$ both are non-constant.

Then, using Lemma 4.1, we obtain from (4.41) and (4.42) that

$$A_2 e^{-p_1(z+c)+p_2(z)} = A_1 H_2 \quad \text{and} \quad A_2 e^{-p_2(z+c)+p_1(z)} = A_1 H_4.$$

As $p_1(z)$ and $p_2(z)$ are non-constant polynomials, we must have $-p_1(z+c) + p_2(z) = \xi_1$ and $-p_2(z+c) + p_1(z) = \xi_2$, where $\xi_1, \xi_2 \in \mathbb{C}$. This implies that $-p_1(z+c) + p_1(z) = -p_2(z+c) + p_2(z) = \xi_1 + \xi_2$. Therefore, we must get $p_1(z) = \alpha z + \beta_1$ and $p_2(z) = \alpha z + \beta_2$, where $\alpha, \beta_1, \beta_2 \in \mathbb{C}$. But, then we get $p_2(z) - p_1(z) = \beta_2 - \beta_1 = \text{constant}$, which is a contradiction. \square

5. STATEMENTS AND DECLARATIONS

Conflict of interest The authors declare that there are no conflicts of interest regarding the publication of this paper.

Funding There is no funding received from any organizations for this research work.

Data availability statement Data sharing is not applicable to this article as no database were generated or analyzed during the current study.

Acknowledgment: The authors would like to thank the referee(s) for the helpful suggestions and comments to improve the exposition of the paper.

REFERENCES

- [1] Biancofiore, A, Stoll, W.: *Another proof of the lemma of the logarithmic derivative in several complex variables*, *Recent developments in several complex variables*, Princeton University Press, Princeton, 29–45 (1981)
- [2] Cao, T.B., Korhonen, R.J.: *A new version of the second main theorem for meromorphic mappings intersecting hyperplanes in several complex variables*, *J. Math. Anal. Appl.* **444**(2), 1114–1132 (2016)
- [3] T. B. Cao and L. Xu, *Logarithmic difference lemma in several complex variables and partial difference equations*, *Ann. Math. Pure Appl.* **199**, 767–794 (2020)
- [4] Chiang, Y.M., Feng, S.J.: *On the Nevanlinna characteristic of $f(z + \eta)$ and difference equations in the complex plane*, *Ramanujan J.* **16**(1), 105–129 (2008).
- [5] Chiang, Y.M., Feng, S.J.: *On the growth of logarithmic differences, difference quotients and logarithmic derivatives of meromorphic functions*, *Trans. Am. math. Soc.* **361**, 3767–3791 (2009).
- [6] Chen, Z.X.: *Growth and zeros of meromorphic solution of some linear difference equations*, *J. Math. Anal. Appl.* **373**, 235–241 (2011).
- [7] Chen, Z.X.: *Zeros of entire solutions to complex linear difference equations*. *Acta Math. Sci. Ser. B Engl. Ed.* **32**, 1141–1148 (2012).
- [8] Courant, R., Hilbert, D.: *Methods of Mathematical Physics, vol. II: Partial Differential Equations*, Interscience (New York, 1962)
- [9] L. Y. Gao: *Entire solutions of two types of systems of complex differential-difference equations*, *Acta Math. Sinica (Chin. Ser.)* **59** (2016), 677–685.
- [10] Garabedian, P.R.: *Partial Differential Equations*, Wiley (New York, 1964).
- [11] Gross, F.: *On the equation $f^n(z) + g^n(z) = 1$* , *Bull. Amer. Math. Soc.* **72**, 86–88 (1966).
- [12] Gross, F.: *On the equation $f^n(z) + g^n(z) = h^n(z)$* , *Amer. Math. Monthly.* **73**, 1093–1096 (1966).
- [13] Halburd, R.G., Korhonen, R.J.: *Difference analogue of the lemma on the logarithmic derivative with applications to difference equations*, *J. Math. Anal. Appl.* **314**, 477–487 (2006).
- [14] Halburd, R.G., Korhonen, R.J.: *Nevanlinna theory for the difference operator*, *Ann. Acad. Sci. Fenn. Math.* **31**, 463–478 (2006).
- [15] Haldar, G.: *Solutions of Fermat-type partial differential difference equations in \mathbb{C}^2* , *Mediterr. J. Math.*, **20** (2023), 50, DOI: <https://doi.org/10.1007/s00009-022-02180-6>
- [16] Haldar, G., Ahamed, M.B.: *Entire solutions of several quadratic binomial and trinomial partial differential-difference equations in \mathbb{C}^2* , *Anal. Math. Phys.* **12** (2022), Article number: 113, DOI: <https://doi.org/10.1007/s13324-022-00722-5>
- [17] Hayman, W.K.: *Meromorphic Functions*, The Clarendon Press, Oxford, 1964.
- [18] Hu, P.C., Li, P., Yang, C. C.: *Unicity of Meromorphic Mappings, Advances in Complex Analysis and Its Applications*, vol. 1, Kluwer Academic Publishers, Dordrecht, Boston, London (2003).
- [19] Iyer, G.: *On certain functional equations*. *J. Indian. Math. Soc.* **3**, 312–315 (1939).
- [20] Korhonen, R.J.: *A difference Picard theorem for meromorphic functions of several variables*, *Comput. Methods Funct. Theory* **12**(1), 343–361 (2012)
- [21] Laine, I.: *Nevanlinna Theory and Complex Differential Equations*, Walter de Gruyter, Berlin, (1993).
- [22] Li, B.Q.: *On reduction of functional-differential equations*, *Complex Var.* **31**, 311–324 (1996)

- [23] Li, B.Q.: *On entire solutions of Fermat type partial differential equations*, Int. J. Math. **15**, 473–485 (2004)
- [24] B. Q. Li, *On certain non-linear differential equations in complex domains*, Arch. Math. **91** (2008), 344–353.
- [25] Liu, M.L., Gao, L.Y.: *Transcendental solutions of systems of complex differential-difference equations*, Sci. Sin. Math. **49**, 1–22 (2019).
- [26] Liu, K.: *Meromorphic functions sharing a set with applications to difference equations*. J. Math. Anal. Appl. **359**, 384–393 (2009).
- [27] Liu, K., Yang, L.Z.: *On Entire Solutions of Some Differential-Difference Equations*. Comput. Methods Funct. Theory **13**, 433–447 (2012).
- [28] Liu, K., Cao, T.B.: *Entire solutions of Fermat type q -difference-differential equations*. Electron. J. Diff. Equ. **59**, 1–10 (2013).
- [29] Liu, K., Cao, T.B., Cao, H.Z.: *Entire solutions of Fermat-type differential-difference equations*. Arch. Math. **99**, 147–155 (2012).
- [30] Liu, K., Dong, X.: *Fermat-type differential and difference equations*, Electron. J. Differ. Equ. **2015**, 1–10 (2015).
- [31] Liu, K., Yang, L.Z.: *A note on meromorphic solutions of Fermat types equations*, An. Stiint. Univ. Al. I. Cuza Iasi Mat. (N. S.). **1**, 317–325 (2016).
- [32] Montel, P.: *Lecons sur les familles de normales fonctions analytiques et leurs applications*. Gauthier-Viuars Paris, 135–136 (1927).
- [33] Rauch, J.: *Partial Differential Equations*, Springer-Verlag (New York, 1991)
- [34] Ronkin, L.I.: *Introduction to the Theory of Entire Functions of Several Variables*, Moscow: Nauka 1971 (Russian), American Mathematical Society, Providence (1974)
- [35] Saleeby, E.G.: *Entire and meromorphic solutions of Fermat-type partial differential equations*, Analysis (Munich) **19**, 369–376 (1999)
- [36] Saleeby, E.G.: *On entire and meromorphic solutions of $\lambda u^k + \sum_{i=1}^n u_{z_i}^m = 1$* , Complex Var. Theory Appl. **49**, 101–107 (2004)
- [37] E. G. Saleeby, *On complex analytic solutions of certain trinomial functional and partial differential equations*, Aequationes Math. **85** (2013), 553–562.
- [38] Stoll, W.: *Holomorphic Functions of Finite Order in Several Complex Variables*, American Mathematical Society, Providence, 1974
- [39] Tang, J.F., Liao, L.W.: *The transcendental meromorphic solutions of a certain type of non-linear differential equations*, J. Math. Anal. Appl. **334**, 517–527 (2007).
- [40] Taylor, R., Wiles, A.: *Ring-theoretic properties of certain Hecke algebra*, Ann. Math. **141**, 553–572 (1995)
- [41] Wiles, A.: *Modular elliptic curves and Fermat last theorem*, Ann. Math. **141**, 443–551 (1995)
- [42] Xu, L., Cao, T.B.: *Solutions of complex Fermat-type partial difference and differential-difference equations*, Mediterr. J. Math. **15**, 1–14 (2018)
- [43] Xu, L., Cao, T.B.: *Correction to: Solutions of Complex Fermat-Type Partial Difference and Differential-Difference Equations*, Mediterr. J. Math. **17**, 1–4 (2020)
- [44] Xu, H.Y., Haldar, G.: *Solutions of complex nonlinear functional equations including second order partial differential and difference equations in \mathbb{C}^2* , Electron. J. Differ. Equ., **43**(2023), 1–18
- [45] Xu, H.Y., Jiang, Y.Y.: *Results on entire and meromorphic solutions for several systems of quadratic trinomial functional equations with two complex variables*, RACSAM (2022). <https://doi.org/10.1007/s13398-021-01154-9>
- [46] Xu, X.Y., Liu, S.Y., Li, Q.P.: *Entire solutions for several systems of nonlinear difference and partial differential-difference equations of Fermat-type*, J. Math. Anal. Appl. **483**, 123–641 (2020)
- [47] Yang, C.C.: *A generalization of a theorem of P. Montel on entire functions*, Proc. Amer. Math. Soc. **26**, 332–334 (1970)

- [48] Yang, C.C., Li, P.: *On the transcendental solutions of a certain type of non-linear differential equations*, Arch. Math. **82**, 442–448 (2004)
- [49] Yang, C.C., Yi, H.X.: *Uniqueness Theory of Meromorphic Functions*, Kluwer Acad. Publ. (2003)
- [50] Zheng, X.M., Tu, J.: *Growth of meromorphic solutions of linear difference equations*, J. Math. Anal. Appl. **384**, 349–356 (2011)

DEPARTMENT OF MATHEMATICS, MALDA COLLEGE, MALDA, WEST BENGAL 732101,
INDIA.

Email address: goutamiit1986@gmail.com