Large deviations and conditioning for chaotic non-invertible deterministic maps: analysis via the forward deterministic dynamics and the backward stochastic dynamics

Cécile Monthus

Université Paris-Saclay, CNRS, CEA, Institut de Physique Théorique, 91191 Gif-sur-Yvette, France

The large deviations properties of trajectory observables for chaotic non-invertible deterministic maps as studied recently by N. R. Smith, Phys. Rev. E 106, L042202 (2022) and by R. Gutierrez, A. Canella-Ortiz, C. Perez-Espigares, arXiv:2304.13754 are revisited in order to analyze in detail the similarities and the differences with the case of stochastic Markov chains. To be concrete, we focus on the simplest example displaying the two essential properties of local-stretching and global-folding, namely the doubling map $x_{t+1} = 2x_t \mod 1$ on the real-space interval $x \in [0, 1]$ that can be also analyzed via the decomposition $x = \sum_{l=1}^{+\infty} \frac{\sigma_l}{2^l}$ into binary coefficients $\sigma_l = 0, 1$. The large deviations properties of trajectory observables can be studied either via deformations of the forward deterministic dynamics or via deformations of the backward stochastic dynamics. Our main conclusions concerning the construction of the corresponding Doob canonical conditioned processes are: (i) nontrivial conditioned dynamics can be constructed only in the backward stochastic perspective where the reweighting of existing transitions is possible, and not in the forward deterministic perspective; (ii) the corresponding conditioned steady state is not smooth on the real-space interval $x \in [0, 1]$ and can be better characterized in the binary space $\sigma_{l=1,2,\dots+\infty}$. As a consequence, the backward stochastic dynamics in the binary space is also the most appropriate framework to analyze higher levels of large deviations, and we obtain the explicit large deviations at level 2 for the probability of the empirical density of long backward trajectories.

I. INTRODUCTION

A. Large deviations for dynamical trajectories : deterministic chaotic systems versus Markov processes

The study of large deviations properties of deterministic dynamical systems has a long history (see the review [1] and the PhD thesis [2] with references therein), in parallel with the general theory of large deviations in statistical physics [1, 3, 4] and with the recent developments for stochastic processes (see the reviews with different scopes [5–13], the PhD Theses [14–19] and the Habilitation Thesis [20]). For Markov processes, the large deviations properties of time-local trajectory observables over a large time-window [0, T] have been much studied recently via the appropriate deformations of the Markov generators [5, 10–13, 15, 20–63, 65], and the construction of the corresponding canonical conditioned processes [38, 39].

The same idea of deformed generators has been applied recently to analyze the statistics of trajectory observables in various deterministic dynamical systems, either in continuous time [2, 66, 67] with the development of population cloning algorithms, or in discrete time for chaotic non-invertible maps [68, 69], that will be the focus of the present paper. Many analytical and numerical results with figures can be found in [68, 69] for various trajectory observables of the most famous one-dimensional maps, namely the doubling map, the tent map, and the logistic map (see also [68] for results on the two-dimensional cat map), with the following slightly different perspectives : [68] focuses mainly on the dominant right-eigenvector eigenvalue equation of the deformed Frobenius-Perron operators and on the Monte-Carlo algorithm to construct the biased backward stochastic trajectories, while [69] considers also the dominant left-eigenvectors of the deformed Frobenius-Perron operators and the construction of the Doob conditioned dynamics making rare events typical.

B. Goals of the present work

The main goal of the present work is to revisit these two recent papers [68, 69] in order to stress the similarities and the differences with the theory of large deviations at various levels for Markov processes, as summarized in Appendix A. In particular, we will discuss in detail the three following issues :

(1) What are the differences between the forward and the backward perspectives for chaotic non-invertible maps?

Chaotic non-invertible maps can be analyzed either from the point of view of the forward deterministic dynamics or from the point of view of the backward stochastic dynamics. As a consequence, the appropriate deformations of these dynamics needed to analyze the statistics of trajectory observables will have completely different qualitative properties, and it is thus important to understand the advantages and the drawbacks of the forward and the backward perspectives In particular, we will stress that the construction of the Doob canonical conditioned processes, where the various existing trajectories are reweighted, is possible only within the backward stochastic perspective, and not within the forward deterministic perspective.

(2) What properties are singular in real-space for chaotic non-invertible maps?

While for Markov processes, one is used to eigenvectors that are regular in real space, one needs to be aware that for chaotic non-invertible maps, various properties are singular in real-space, in particular some dominant eigenvectors of the deformed dynamics : for the deformed forward dynamics, the right-eigenvectors are smooth in real space, but the left-eigenvectors are singular in real space, as already found numerically on the example in Fig S1 of [69]; for the deformed backward dynamics, the left-eigenvectors are smooth but the right-eigenvectors are singular, so that their products corresponding to the Doob conditioned steady states are always singular in real space. We will stress that all the observables that are singular in real-space can be better analyzed via the symbolic dynamics, and explain how explicit expressions can be obtained in simple cases.

(3) What higher levels of large deviations can be written explicitly for chaotic non-invertible maps?

As the large deviations of trajectory observables are not explicit in general, since one is not able to solve eigenvalues equations for arbitrary deformations of the generators, it is interesting to consider higher levels of large deviations in order to see what is the smallest level that can be written explicitly. Indeed for Markov processes (see the reminder in Appendix A), a major progress has been the formulation of the so-called level 2.5, where one can write explicit rates functions for the joint distribution of the empirical density and of the empirical flows, in particular for discrete-time Markov chains [4, 14, 70–75], for continuous-time Markov jump processes with discrete configuration space [14, 17, 20, 45, 72–88], for diffusion processes in continuous space [17, 20, 54, 73, 75, 76, 80, 88–90], and for jump-diffusion or jump-drift processes [72, 91–93]. All the lower levels can be then obtained via the optimization of the explicit rate function 2.5 in the presence of the appropriate constraints, but the solutions of these constrained optimizations cannot be written explicitly in general. As a consequence, the large deviation at level 2 for the distribution of the empirical density alone, as well as the large deviations of trajectory observables that can be rewritten in terms of the empirical density and of the empirical flows, are not explicit in general.

For chaotic non-invertible maps, the backward dynamics is stochastic, but remains nevertheless very special since the corresponding forward dynamics is deterministic. We will stress that this property induces level reductions in the large deviations of backward trajectories with respect to the case of Markov chains : the application of the explicit level 2.5 for Markov chains actually reduces to an explicit expression for the level 2 concerning the distribution of the empirical density alone; the general trajectory observables of Markov chains that involve both the empirical density and the empirical flows can be rewritten in terms of the empirical density alone, and thus reduce to the trajectory observables of the so-called Level 1, that can be obtained from the Level 2 via contraction.

C. Organization of the present paper and main results

In order to discuss more concretely the three issues summarized above, we will focus on the simplest chaotic noninvertible displaying the two essential properties of local-stretching and global-folding, namely the doubling map on the real-space interval $x \in [0, 1[$. Let us now describe the purpose of each section and mention the equation numbers where the main results can be found.

• In section II, the essential properties of the forward deterministic dynamics associated to the doubling map are recalled, both in real space and in binary space. The spectral properties governing the convergence towards the uniform steady state actually involve a lot of subtleties (see [94–96] and references therein). For our present purposes, the most important property is that the non-dominant left-eigenvectors of the Frobenius-Perron are singular in real space (Eqs 14 and 15), as a consequence of the sensitivity with respect to initial conditions.

• In section III, the appropriate deformations of this forward deterministic dynamics are analyzed to study large deviations of trajectory observables, that actually reduce to the observables of the so-called Level 1 (Eq. 27) as a consequence of the deterministic character of the forward dynamics. We stress that the dominant left-eigenvector of the deformed dynamics is singular in real space, already at the first order of perturbation theory that involves the non-dominant left eigenvectors of the unperturbed dynamics (Eq. 40). As a consequence, the Doob conditioned steady state is also singular in real space, while the Doob conditional kernel coincides with the initial kernel (Eq. 34), since the Doob canonical conditioning can only change the weights of the possible transitions but cannot produce new transitions.

• In section IV, the backward stochastic dynamics associated to the doubling map is described, both in real space and in binary space, in order to stress the differences with respect to the forward deterministic dynamics discussed in section III.

• In section IV, the large deviations of trajectory observables are analyzed via the deformations of this backward stochastic dynamics. We stress that for this deformed backward dynamics, the dominant left-eigenvector is regular in real space, but the dominant right-eigenvector is singular in real space, so that their product corresponding to the Doob

conditioned steady state is again always singular in real space. However here the Doob conditional backward kernel is different from the initial backward kernel and corresponds to the appropriate reweighting of the two pre-images (Eqs 62 and 63 in real space, or Eqs 72 and 73 in binary space).

• In section VI, all these previous different perspectives are illustrated on the simplest trajectory observable of the doubling map already considered in the two recent papers [68, 69]. We first recall the explicit solution [68] for the dominant eigenvalue (Eq 75) of the deformed forward dynamics with its corresponding right eigenvector that is smooth in real space (Eq. 76) and factorized in binary space (Eq. 85). We write the corresponding Doob conditioned backward kernel in real-space (Eq. 89) as well as in binary space (Eq. 92). The conditioned steady state is given by a simple explicit expression in binary space (Eq. 93) while its translation in real space is singular. We also give the explicit expression in binary space of the dominant left eigenvector of the deformed forward dynamics (Eq. 100), whose translation in real space is singular as already found numerically on Fig S1 of [69].

• In section VII, we turn to the third issue (3) described in the previous subsection concerning higher levels of large deviations beyond the level of trajectory observables. We stress that the application of the explicit large deviations at level 2.5 for arbitrary Markov chains (Eqs 101 and 102) to the specific case of the stochastic backward dynamics associated to the doubling map produces explicit large deviations at level 2 for the empirical density alone (Eqs 109 and 110). This level reduction from the level 2.5 towards the level 2 can be understood from the very special type of stochasticity of the backward dynamics that is associated to a forward deterministic dynamics.

• In VIII, we summarize our conclusions concerning the doubling map and discuss the generalizations to other chaotic non-invertible maps or other chaotic deterministic dynamics.

• Three appendices contain complementary material with respect to the main text: Appendix A summarizes the large deviations properties at various levels for Markov chains, while Appendix B recalls the properties of the backward Markov chain associated to a given forward Markov chain; Appendix C contains the properties of the doubling map for the Fourier coefficients of the density on the real space interval $x \in [0, 1[$.

II. DOUBLING-MAP DETERMINISTIC DYNAMICS IN REAL-SPACE AND IN BINARY-SPACE

In this section, some important properties of the Doubling-map deterministic dynamics are summarized in order to stress the similarities and differences with Markov chains described in section A 1 of Appendix A.

A. Deterministic trajectories on the real-space interval [0,1]

The doubling map on the interval $x \in [0, 1]$

$$x_{t+1} = f(x_t) \equiv 2x_t [\text{mod1}] \equiv \begin{cases} 2x_t & \text{for } 0 \le x_t < \frac{1}{2} \\ (2x_t - 1) & \text{for } \frac{1}{2} \le x_t < 1 \end{cases}$$
(1)

is the simplest example of chaotic dynamics displaying the two essential properties of local-stretching and globalfolding:

(i) the uniform local-stretching of amplitude f'(x) = 2 ensures the exponential divergence in time of the separation between two nearby trajectories, and thus the sensitivity to initial conditions.

(ii) the global-folding corresponds to the non-invertibility of the doubling map, where each value $x_{t+1} \in [0, 1]$ has two pre-images

$$\begin{aligned}
x_t^- &= \frac{x_{t+1}}{2} \in [0, \frac{1}{2}] \\
x_t^+ &= \left(\frac{1}{2} + \frac{x_{t+1}}{2}\right) \in [\frac{1}{2}, 1[
\end{aligned}$$
(2)

so that some information about the past is lost at each time-step. As a consequence, when one tries to reconstruct the past trajectories backward in time, one obtains a stochastic dynamics as will be described in section IV, while in the present section, we focus on the forward deterministic point of view.

B. Frobenius-Perron dynamics for the probability density $\rho_t(x)$ on the real space interval $x \in [0, 1]$

The Heaviside function $\theta(z)$ defined as

$$\theta(z) = \begin{cases} 1 \text{ for } z \ge 0\\ 0 \text{ for } z < 0 \end{cases}$$
(3)

is useful to write the forward deterministic kernel $w(\tilde{x}|x)$ associated to the doubling map of Eq. 1

$$w(\tilde{x}|x) \equiv \delta(\tilde{x} - f(x)) = \delta\left(\tilde{x} - 2x\right) \left[\theta\left(x\right) - \theta\left(x - \frac{1}{2}\right)\right] + \delta\left(\tilde{x} - (2x - 1)\right) \left[\theta\left(x - \frac{1}{2}\right) - \theta\left(x - 1\right)\right] = \frac{\delta\left(x - \frac{\tilde{x}}{2}\right) + \delta\left(x - \frac{\tilde{x} + 1}{2}\right)}{2}$$

$$(4)$$

that governs the Frobenius-Perron dynamics for the probability density $\rho_t(.)$

$$\rho_t(\tilde{x}) = \int_0^1 dx w(\tilde{x}|x) \rho_{t-1}(x) = \frac{\rho_{t-1}(\frac{\tilde{x}}{2}) + \rho_{t-1}(\frac{\tilde{x}+1}{2})}{2}$$
(5)

which is the analog of the Markov chain evolution of Eq. A1. The physical interpretation is that the new histogram $\rho_t(.)$ represents the average of the two rescaled half-histograms of $\rho_{t-1}(.)$ on the two intervals $[0, \frac{1}{2}]$ and $[\frac{1}{2}, 1]$. The iteration of Eq. 5 up to the initial density $\rho_{t=0}(.)$ at t = 0

$$\rho_t(x) = \frac{\rho_{t-2}(\frac{x}{4}) + \rho_{t-2}(\frac{x+1}{4}) + \rho_{t-2}(\frac{x+2}{4}) + \rho_{t-2}(\frac{x+3}{4})}{4} = \dots = \frac{1}{2^t} \sum_{j=0}^{2^t-1} \rho_0\left(\frac{x}{2^t} + \frac{j}{2^t}\right) \tag{6}$$

yields that the histogram $\rho_t(.)$ at time t represents the average of the 2^t rescaled histograms $\rho_0(.)$ on the 2^t intervals $\left[\frac{j}{2^t}, \frac{j+1}{2^t}\right]$ labelled by $j = 0, 1, ..., 2^t - 1$. If the initial density $\rho_{t=0}$ is smooth, one thus expect the convergence towards the uniform invariant density

$$\rho_t(x) \underset{t \to +\infty}{\simeq} \rho_*(x) = 1 \tag{7}$$

that corresponds to the steady solution for the dynamics of Eq. 5. As for Markov chains, one would like to characterize the convergence convergence towards this steady state via the spectral decomposition of the kernel analogous to Eq. A4.

C. Spectral decomposition when the initial density $\rho_{t=0}(x)$ is infinitely differentiable on the interval $x \in [0, 1]$

Let us summarize the spectral properties of the kernel $w(\tilde{x}|x)$ in the space of infinitely differentiable densities on the interval $x \in [0, 1]$ (see [94–96] and references therein). If the density $\rho_t(x)$ on the whole interval $x \in [0, 1]$ can be reproduced by its Taylor expansion around the origin

$$\rho_t(x) = \sum_{p=0}^{+\infty} \frac{\rho_t^{(p)}(0)}{p!} x^p \tag{8}$$

the dynamics can be analyzed via the action of the kernel $w(\tilde{x}|x)$ on the powers x^p for $p=1,2,...+\infty$

$$\int_{0}^{1} dx w(\tilde{x}|x) x^{p} = \frac{\left(\frac{\tilde{x}}{2}\right)^{p} + \left(\frac{\tilde{x}+1}{2}\right)^{p}}{2} = \frac{\tilde{x}^{p}}{2^{p}} + \sum_{q=0}^{p-1} \tilde{x}^{q} \frac{\binom{p}{q}}{2^{p+1}}$$
(9)

Since it produces the same power \tilde{x}^p and the smaller powers \tilde{x}^q with q = 0, ..., p - 1, the diagonal coefficients directly give the eigenvalues $\lambda_{p\neq 0}$ of the kernel w(.|.)

$$\lambda_p = 2^{-p} \quad p = 1, 2, .., +\infty$$
 (10)

The corresponding right eigenvectors satisfying Eq. A5

$$\lambda_p \ r_p(x) = \frac{r_p(\frac{x}{2}) + r_p(\frac{x+1}{2})}{2} \tag{11}$$

are the Bernoulli polynomials $r_p(x) = B_p(x)$ of degrees p [94, 95]: beyond $r_0(x) = B_0(x) = 1 = \rho_*(x)$ corresponding to the uniform steady state discussed in Eq. 7, the first members read

$$r_{1}(x) = x - \frac{1}{2} \equiv B_{1}(x)$$

$$r_{2}(x) = x^{2} - x + \frac{1}{6} \equiv B_{2}(x)$$

$$r_{3}(x) = x^{3} - \frac{3}{2}x^{2} + \frac{1}{2}x \equiv B_{3}(x)$$
(12)

while their Fourier definition is given in Eq. C11 for arbitrary p. Beyond $l_0(x) = 1$, the left eigenvectors satisfying Eq. A5 with the eigenvalues $\lambda_p = 2^{-p}$ of Eq. 10 with p = 1, 2, ...

$$0 \le x < \frac{1}{2}: \quad \lambda_p l_p(x) = l_p(2x)$$

$$\frac{1}{2} \le x < 1: \quad \lambda_p l_p(x) = l_p(2x-1)$$
 (13)

are however singular distributions $D_p(x)$ that read in terms of the delta functions and their derivatives of arbitrary orders at the two boundaries x = 0 and x = 1 [94, 95]

$$l_p(x) = \frac{(-1)^{p-1}}{p!} \left[\delta^{(p-1)}(x-1) - \delta^{(p-1)}(x) \right] \equiv D_p(x)$$
(14)

with the first members

$$l_{1}(x) = \delta(x-1) - \delta(x) \equiv D_{1}(x)$$

$$l_{2}(x) = -\frac{1}{2} \left[\delta'(x-1) - \delta'(x) \right] \equiv D_{2}(x)$$

$$l_{3}(x) = \frac{1}{6} \left[\delta''(x-1) - \delta''(x) \right] \equiv D_{3}(x)$$
(15)

The scalar product of the distribution $D_p(x)$ for p > 0 with another function g(x) can be evaluated using (p-1) integrations by parts

$$\begin{split} \langle D_p | g \rangle &= \int_0^1 dx D_p(x) g(x) \quad = \int_0^1 dx g(x) \frac{(-1)^{p-1}}{p!} \frac{d^{(p-1)}}{dx^{p-1}} \left[\delta(x-1) - \delta(x) \right] = \int_0^1 dx \left[\delta(x-1) - \delta(x) \right] \frac{1}{p!} \frac{d^{(p-1)}g(x)}{dx^{p-1}} \\ &= \frac{g^{(p-1)}(1) - g^{(p-1)}(0)}{p!} \end{split}$$
(16)

In particular, the spectral decomposition of Eq. A8

$$\rho_t(\tilde{x}) = 1 + \sum_{p=1}^{+\infty} 2^{-pt} B_p(\tilde{x}) \int_0^1 dx_0 D_p(x_0) \rho_0(x_0)$$

= $1 + \sum_{p=1}^{+\infty} 2^{-pt} B_p(\tilde{x}) \left[\frac{\rho_0^{(p-1)}(1) - \rho_0^{(p-1)}(0)}{p!} \right]$ (17)

that corresponds to the Euler-Maclaurin formula for t = 0 [94], can be used only for if the initial density $\rho_0(x)$ is infinitely differentiable. Note that when the initial density $\rho_0(x)$ is not infinitely differentiable, the discussion of the spectral properties is much more involved and depends on the precise space of densities that one wishes to consider (see the very detailed discussion in [95]). Besides the analysis of the dynamics on the real space interval [0, 1[discussed above, it is useful to consider other formulations (see the very detailed discussion in [95]), in particular in Fourier space as recalled in Appendix C and in the binary-space as recalled in the next subsection.

D. Properties of the dynamics in terms of the binary coefficients $\sigma_l = 0, 1$ parametrizing $x = \sum_{l=1}^{+\infty} \frac{\sigma_l}{2^l}$

When $x \in [0, 1]$ is represented by its binary coefficients $\sigma_l = 0, 1$ with $l = 1, 2, ... + \infty$

$$x = \sum_{l=1}^{+\infty} \frac{\sigma_l}{2^l} = \frac{\sigma_1}{2} + \frac{\sigma_2}{4} + \frac{\sigma_3}{8} + \dots$$
(18)

the dynamics of the doubling map of Eq. 1

$$\tilde{x} \equiv \sum_{l=1}^{+\infty} \frac{\tilde{\sigma}_l}{2^l} = 2x [\text{mod}1] = \sum_{l=1}^{+\infty} \frac{\sigma_{l+1}}{2^l}$$
(19)

translates into the shift dynamics for the binary coefficients

$$\tilde{\sigma}_l = \sigma_{l+1} \quad \text{for } l = 1, 2, \dots \tag{20}$$

while the first coefficient σ_1 disappears.

The real-space kernel w(.|.) of Eq. 4 translates for the binary coefficients into the kernel

$$W(\tilde{\sigma}_{.}|\sigma_{.}) \equiv \prod_{l=1}^{+\infty} \delta_{\tilde{\sigma}_{l},\sigma_{l+1}}$$
(21)

The normalization of this kernel over $\tilde{\sigma}$ corresponds to the left eigenvector unity for any configuration $\tilde{\sigma}$.

$$L_0(\tilde{\sigma}_{.}) = \prod_{l=1}^{+\infty} \left(\delta_{\tilde{\sigma}_l,0} + \delta_{\tilde{\sigma}_l,1}\right) \tag{22}$$

The correspondence between the probability density $\rho_t(x)$ on the interval [0,1] and the probability $P_t(\sigma_{\cdot})$ of the binary variables reads

$$\rho_t(x) = \sum_{\sigma_{\perp}} P_t(\sigma_{\perp}) \delta\left(x - \sum_{l=1}^{+\infty} \frac{\sigma_l}{2^l}\right)$$
(23)

The steady uniform density $\rho_*(x) = 1$ of Eq. 7 translates into the steady uniform probability for the binary variables $\sigma_l = 0, 1$ drwn with probabilities $(\frac{1}{2}, \frac{1}{2})$

$$P_*(\sigma_{\cdot}) = \prod_{l=1}^{+\infty} \left(\frac{\delta_{\sigma_l,0} + \delta_{\sigma_l,1}}{2} \right) = R_0(\sigma_{\cdot}) \tag{24}$$

that is the right eigenvector $R_0(\sigma)$ of the kernel of Eq. 21 associated to the highest eigenvalue $\lambda_0 = 1$.

The Frobenius-Perron dynamics for the probability $P_t(.)$ of the binary variables

$$P_{t}(\tilde{\sigma}_{1}, \tilde{\sigma}_{2}, ...) = \sum_{\sigma_{.}} w(\tilde{\sigma}_{.} | \sigma_{.}) P_{t-1}(\sigma_{.}) = \sum_{\sigma_{1} = \pm 1} P_{t-1}(\sigma_{1}, \tilde{\sigma}_{1}, \tilde{\sigma}_{2}, ...)$$
(25)

means that one just integrates the previous probability $P_{t-1}(.)$ over its first variable σ_1 . The iteration up to the initial probability $P_{t=0}(.)$ at time t = 0

$$P_t(\tilde{\sigma}_1, \tilde{\sigma}_2, ...) = \sum_{\sigma_1 = \pm 1; \sigma_2 = \pm 1; ...; \sigma_t = \pm 1} P_0(\sigma_1, \sigma_2, ..., \sigma_t, \tilde{\sigma}_1, \tilde{\sigma}_2, ...)$$
(26)

means that one just integrates the initial probability $P_{t=0}(.)$ over its first t variables $(\sigma_1; \sigma_2; ...; \sigma_t)$. This formulation shows even more clearly the issues of the convergence towards the uniform distribution of Eq. 24 for large t depending on the initial distribution $P_{t=0}(.)$: the convergence found in Eq. 7 in real space for initial densities that were sufficiently smooth means for the binary variables that the coefficients σ_l for large enough l should be independent and take the values (0, 1) with the equal probabilities $(\frac{1}{2}, \frac{1}{2})$.

E. Discussion : similarities and differences with Markov chains

In summary, if the initial density $\rho_{t=0}(.)$ is infinitely differentiable on the real-space interval $x \in [0, 1]$, the convergence towards the uniform steady state via the spectral decomposition of Eq. 17 is similar to the analog property of Eq. A8 for Markov chains. However, the fact that the left eigenvectors of Eq. 14 are singular distributions that reflects the sensitivity with respect to initial conditions will have very important consequences in the study of large deviations for trajectory observables via appropriate deformations of the kernel, as discussed in the next section.

III. LARGE DEVIATIONS OF TRAJECTORY OBSERVABLES VIA THE DEFORMED DYNAMICS

In this section, the analysis of large deviations of trajectory observables for chaotic non-invertible maps described in the two recent papers [68, 69] is revisited on the concrete example of the doubling map in order to analyze in detail the similarities and differences with the case of stochastic Markov chains summarized in subsection A 2 of Appendix A.

The first important difference for any deterministic dynamics $x_{t+1} = f(x_t)$ is that the general trajectory observable of Eq. A9 which is constructed from the elementary transitions $x_t \to x_{t+1}$ during the long trajectory

$$\mathcal{O}^{Traj}[x_0, x_1, ..., x_T] = \frac{1}{T} \sum_{t=0}^{T-1} O(x_{t+1}, x_t) = \frac{1}{T} \sum_{t=0}^{T-1} O(f(x_t), x_t)$$
$$\equiv \frac{1}{T} \sum_{t=0}^{T-1} \omega(x_t) \quad \text{with} \ \omega(x_t) = O(f(x_t), x_t)$$
(27)

reduces to an observable that involves only the function $\omega(x_t) = \Omega(f(x_t), x_t)$ of the single position x_t .

A. Dynamics of the generating function $Z_T^{[k]}(x_T|x_0)$ governed by the k-deformed kernel $w^{[k]}(\tilde{x}|x)$

As recalled in details in subsection A 2 of Appendix A, the standard method to analyze the large deviations of Eq. A22 for trajectory observables is based on the generating function $Z_T^{[k]}(x_T|x_0)$ of Eq. A10 for the trajectory observable of Eq. 27

$$Z_{T}^{[k]}(x_{T}|x_{0}) \equiv \langle e^{kT\mathcal{O}^{Traj}} \rangle = \int dx_{1} \int dx_{2} \dots \int dx_{T-1} \prod_{t=0}^{T-1} \left[w(x_{t+1}, x_{t})e^{k\omega(x_{t})} \right]$$
$$\equiv \int dx_{1} \int dx_{2} \dots \int dx_{T-1} \prod_{t=0}^{T-1} w^{[k]}(x_{t+1}, x_{t}) \equiv \langle x_{T} | \left(w^{[k]} \right)^{T} | x_{0} \rangle$$
(28)

whose dynamics can be analyzed via the k-deformation of Eq. A11 that reads for the doubling-map kernel of Eq. 4

$$w^{[k]}(\tilde{x}|x) \equiv \delta(\tilde{x} - f(x))e^{k\omega(x)} = \delta\left(\tilde{x} - 2x\right)e^{k\omega(x)}\left[\theta\left(x\right) - \theta\left(x - \frac{1}{2}\right)\right] + \delta\left(\tilde{x} - (2x - 1)\right)e^{k\omega(x)}\left[\theta\left(x - \frac{1}{2}\right) - \theta\left(x - 1\right)\right]$$
$$= \frac{\delta(x - \frac{\tilde{x}}{2}) + \delta(x - \frac{\tilde{x} + 1}{2})}{2}e^{k\omega(x)}\left[\theta\left(\tilde{x}\right) - \theta\left(\tilde{x} - 1\right)\right]$$
(29)

The dynamics for the generating function

$$Z_T^{[k]}(\tilde{x}) = \int_0^1 dx w^{[k]}(\tilde{x}|x) Z_{T-1}^{[k]}(x) = \frac{Z_{T-1}^{[k]}(\frac{\tilde{x}}{2}) e^{k\omega(\frac{\tilde{x}}{2})} + Z_{T-1}^{[k]}(\frac{\tilde{x}}{2}) e^{k\omega(\frac{\tilde{x}+1}{2})}}{2}$$
(30)

corresponds to the deformation of the Frobenius-Perron dynamics of Eq. 5 for the probability density $\rho_t(x)$.

B. Eigenvalue problem governing the generating function $Z_T^{[k]}(x_T|x_0)$ for large time $T \to +\infty$

As recalled around Eq. A12 for Markov chains, the generating function for large time T is governed by the highest eigenvalue $\lambda_0^{[k]}$ associated to the positive right eigenvector $r_0^{[k]}(\tilde{x}) \ge 0$ and to the positive left eigenvector $l_0^{[k]}(\tilde{x}) \ge 0$

satisfying Eq. A13 that reads for the deformed doubling-map kernel of Eq. 29

$$\begin{aligned} \lambda_0^{[k]} r_0^{[k]}(\tilde{x}) &= \int_0^1 dx \; w^{[k]}(\tilde{x}|x) r_0^{[k]}(x) = \frac{e^{k\omega(\frac{\tilde{x}}{2})} r_0^{[k]}(\frac{\tilde{x}}{2}) + e^{k\omega(\frac{\tilde{x}+1}{2})} r_0^{[k]}(\frac{\tilde{x}+1}{2})}{2} \\ \lambda_0^{[k]} l_0^{[k]}(x) &= \int_0^1 d\tilde{x} l_0^{[k]}(\tilde{x}) w^{[k]}(\tilde{x}|x) \\ &= l_0^{[k]}(2x) e^{k\omega(x)} \left[\theta\left(x\right) - \theta\left(x - \frac{1}{2}\right) \right] + l_0^{[k]}(2x-1) e^{k\omega(x)} \left[\theta\left(x - \frac{1}{2}\right) - \theta\left(x - 1\right) \right] \end{aligned}$$
(31)

However, as already discussed for the undeformed case k = 0 around Eq. 7, the convergence of $\rho_t(x)$ towards the uniform distribution $\rho_*(x) = 1$ requires the smoothness of the initial condition $\rho_{t=0}(x_0)$. Similarly, the convergence of the dynamics Eq. 30 towards

$$Z_T^{[k]}(\tilde{x}) \underset{T \to +\infty}{\simeq} [\lambda_0^{[k]}]^T r_0^{[k]}(x_T) \int dx_0 l_0^{[k]}(x_0) Z_0^{[k]}(x_0)$$
(32)

requires the smoothness of the initial condition $Z_{T=0}^{[k]}(x_0) = \rho_{t=0}(x_0)$. Then if one constructs the probability of the end-point analog to Eq. A15, it will converge for large time $T \to +\infty$

$$P_T^{[k]End}(x) \equiv \frac{Z_T^{[k]}(x)}{\int dy Z_T^{[k]}(y)} \underset{T \to +\infty}{\simeq} \frac{r_0^{[k]}(x)}{\int dy r_0^{[k]}(y)} \equiv P_*^{[k]End}(x)$$
(33)

towards the distribution $P_*^{[k]End}(x)$ determined by the smooth right eigenvector $r_0^{[k]}(.)$.

However, if one tries to construct the conditioned kernel of Eq. A18

$$w^{C[k]}(\tilde{x}|x) \equiv \frac{1}{\lambda_0^{[k]}} l_0^{[k]}(\tilde{x}) w^{[k]}(\tilde{x}|x) \frac{1}{l_0^{[k]}(x)} = \frac{l_0^{[k]}(\tilde{x}) w(\tilde{x}|x) e^{k\Omega(\tilde{x},x)}}{\int_0^1 dz l_0^{[k]}(z) w(z|x) e^{k\Omega(z,x)}}$$
$$= \frac{l_0^{[k]}(\tilde{x}) \delta(\tilde{x} - f(x)) e^{k\omega(x)}}{\int_0^1 dz l_0^{[k]}(z) \delta(z - f(x)) e^{k\omega(x)}} = \frac{l_0^{[k]}(f(x)) \delta(\tilde{x} - f(x)) e^{k\omega(x)}}{l_0^{[k]}(f(x)) e^{k\omega(x)}} = \delta(\tilde{x} - f(x)) = w(\tilde{x}|x)$$
(34)

one obtains that it coincides with the initial deterministic kernel $w(\tilde{x}|x) = \delta(\tilde{x} - f(x))$, as expected since the Doob canonical conditioning can only change the weights of the possible transitions as recalled after Eq. A18, but cannot produce new transitions.

On the other hand, the conditioned density given by Eq. A19 in terms of the deformed eigenvectors of Eq. 31

$$\rho_*^{C[k]}(x) \equiv l_0^{[k]}(x) r_0^{[k]}(x) \tag{35}$$

should be different from the undeformed uniform steady density $\rho_*(x) = 1$. In order to clarify what is really going on, it is thus useful to consider the perturbation theory in the deformation parameter k.

С. Perturbation theory in k for any trajectory observable

The perturbation theory in the deformation parameter k is recalled for Markov chains in subsection A 2 f of Appendix A. For the doubling map where the invariant density is uniform $\rho_*(x) = 1$, the first-order correction of Eq. A29 reads for $\omega(x) = \Omega(f(x), x)$ of Eq. 27 using the normalization over \tilde{x} of the kernel $w(\tilde{x}|x)$

$$\lambda^{(1)} = \int_0^1 dx \left[\int_0^1 d\tilde{x} w(\tilde{x}|x) \right] \omega(x) \rho_*(x) = \int_0^1 dx \omega(x)$$
(36)

The second-order correction $\lambda^{(2)}$ of Eq. A36 reads using the other eigenvalues $\lambda_p = 2^{-p}$ of Eq. 10 for $p = 1, 2, .., +\infty$, with their right eigenvectors given by the Bernoulli polynomials $r_p(.) = B_p(.)$ of Eq. 12 and their left eigenvectors given by the singular distributions $l_p(.) = D_p(.)$ of Eq. 14

$$\lambda^{(2)} = \int_0^1 dx \frac{\omega^2(x)}{2} \rho_*(x) + \sum_{p=1}^{+\infty} \frac{\lambda_p}{1 - \lambda_p} \left(\int_0^1 dx_2 \omega(x_2) r_p(x_2) \right) \left(\int_0^1 dx_1 l_p(x_1) \omega(x_1) \rho_*(x_1) \right)$$
$$= \int_0^1 dx \frac{\omega^2(x)}{2} + \sum_{p=1}^{+\infty} \frac{1}{2^p - 1} \left(\int_0^1 dx_2 \omega(x_2) B_p(x_2) \right) \left(\int_0^1 dx_1 D_p(x_1) \omega(x_1) \right)$$
(37)

where the last scalar product reads using Eq. 16

$$\int_0^1 dx_1 D_p(x_1) \omega(x_1) = \frac{\omega^{(p-1)}(1) - \omega^{(p-1)}(0)}{p!}$$
(38)

The first-order perturbation theory of Eq. A33 for the right eigenvector yields that the correction $r_0^{(1)}(\tilde{x})$ with respect to the uniform invariant density $\rho_*(x) = 1$ can be decomposed onto the Bernoulli polynomials $B_p(.)$ of Eq. 12

$$r_{0}^{(1)}(\tilde{x}) = \sum_{p=1}^{+\infty} B_{p}(\tilde{x}) \left[\frac{1}{2^{p} - 1} \int_{0}^{1} dx D_{p}(x) \omega(x) \right]$$
$$= \sum_{p=1}^{+\infty} B_{p}(\tilde{x}) \left[\frac{\omega^{(p-1)}(1) - \omega^{(p-1)}(0)}{(2^{p} - 1)p!} \right]$$
(39)

while the coefficients using Eq. 38 involve derivatives of arbitrary order of $\omega(.)$ at x = 0 and x = 1.

The first-order perturbation theory of Eq. A33 for the left eigenvector yields that the correction $l_0^{(1)}(\tilde{x})$ with respect to the trivial eigenvector $l_0(x) = 1$ can be decomposed onto the singular distributions $D_p(.)$ of Eq. 14

$$l_0^{(1)}(\tilde{x}) = \sum_{p=1}^{+\infty} D_p(\tilde{x}) \left[\frac{1}{1 - 2^{-p}} \int_0^1 dx \omega(x) B_p(x) \right]$$
(40)

where the coefficients can be computed as integrals of $\omega(x)$ with the Bernoulli polynomials $B_p(x)$ of Eq. 12.

The first-order correction of Eq. A35 for the conditioned density of Eq. A20

$$\rho_*^{C(1)}(\tilde{x}) = r_0^{(1)}(\tilde{x}) + l_0^{(1)}(\tilde{x}) = \sum_{p=1}^{+\infty} B_p(\tilde{x}) \left[\frac{\omega^{(p-1)}(1) - \omega^{(p-1)}(0)}{(2^p - 1)p!} \right] + \sum_{p=1}^{+\infty} D_p(\tilde{x}) \left[\frac{1}{1 - 2^{-p}} \int_0^1 dx \omega(x) B_p(x) \right]$$
(41)

then contains both the regular contributions given by the Bernoulli polynomials $B_p(\tilde{x})$ of Eq. 12 and the singular contributions given by the singular distributions $D_p(.)$ of Eq. 14.

D. Discussion : similarities and differences with Markov chains

The method to analyze large deviations of trajectory observables via the appropriate deformations of the dynamics is similar to the same approach for Markov chains, but there are two essential differences:

(i) the conditioned kernel $w^{C[k]}(\tilde{x}|x)$ constructed via Eq. 34 coincides with the initial deterministic kernel $w(\tilde{x}|x) = \delta(\tilde{x} - f(x))$, since the Doob canonical conditioning can only change the weights of the possible transitions but not create new transitions.

(ii) the first-order theory in k indicates that both the deformed left eigenvector (Eq. 40) and the conditioned density (Eq. 41) become very singular in real space as soon as $k \neq 0$.

In order to overcome the first difficulty (i), the natural idea is to replace the forward deterministic dynamics by the backward stochastic dynamics described in the next section. The singular character (ii) of various observables in real-space for $k \neq 0$ suggests that it will be useful to analyze this backward stochastic dynamics not only in real-space but also in the binary space where its deformations will be easier to characterize.

IV. STOCHASTIC BACKWARD DYNAMICS IN REAL-SPACE AND IN BINARY-SPACE

In this section, we describe the backward stochastic dynamics associated to the doubling map that is obviously closer to the Markov chains described in Appendix A than the deterministic forward dynamics considered in the previous sections.

A. Stochastic backward dynamics governed by the kernel $w_B(x|\tilde{x})$

As recalled in Appendix B for Markov chains, the trajectories can be alternatively constructed backward in time via the appropriate backward kernel. For chaotic non-invertible maps, the advantages of this backward perspective have been already stressed in [68]. For the doubling map, the backward kernel $w_B(x|\tilde{x})$ of Eq. B3 associated to the forward deterministic kernel of Eq. 4 and to its uniform steady state $\rho_*(x) = 1$ of Eq. 7 reads

$$w_B(x|\tilde{x}) \equiv w(\tilde{x}|x) = \delta\left(\tilde{x} - 2x\right) \left[\theta\left(x\right) - \theta\left(x - \frac{1}{2}\right)\right] + \delta\left(\tilde{x} - (2x - 1)\right) \left[\theta\left(x - \frac{1}{2}\right) - \theta\left(x - 1\right)\right]$$
$$= \frac{\delta\left(x - \frac{\tilde{x}}{2}\right) + \delta\left(x - \frac{\tilde{x} + 1}{2}\right)}{2} \left[\theta\left(\tilde{x}\right) - \theta\left(\tilde{x} - 1\right)\right]$$
(42)

The last expression means that the backward dynamics is a stochastic Markov chain, where for each $\tilde{x} \in [0, 1[$, one chooses one of the two pre-images $\left(x_{-} = \frac{\tilde{x}}{2}; x_{+} = \frac{\tilde{x}+1}{2}\right)$ with the equal probabilities $\left(\frac{1}{2}, \frac{1}{2}\right)$.

So, even if the probability of a trajectory $\mathcal{P}_T^{Traj}(x_0, x_1, ..., x_T)$ in the steady state can be written either with the forward kernel or with the backward kernel via Eq. B8, one obtains a completely different perspective :

(i) once the initial point x_0 is drawn with $\rho_*(x_0)$, the forward trajectory is then completely determined by the deterministic forward rules $x_{t+1} = f(x_t)$

$$\mathcal{P}_T^{Forward(x_0)}(x_1, ..., x_T) = \prod_{t=0}^{T-1} w(x_{t+1}|x_t) = \prod_{t=0}^{T-1} \delta(x_{t+1} - f(x_t))$$
(43)

(ii) once the final point x_T is drawn with $\rho_*(x_T)$, there are 2^T possible backward trajectories that all have the same probability 2^{-T}

$$\mathcal{P}_T^{Backward(x_T)}(x_{T-1}, x_{T-2}, ..., x_0) = \prod_{t=0}^{T-1} W_B(x_t | x_{t+1}) = \prod_{t=0}^{T-1} \frac{\delta(x_t - \frac{x_{t+1}}{2}) + \delta(x_t - \frac{x_{t+1}+1}{2})}{2}$$
(44)

B. Backward stochastic dynamics for probability density $\rho_{\tau}^{B}(x)$ on the real space interval $x \in [0, 1]$

When considering the backward trajectory, it is useful to replace the forward-time t = 0, 1, .., T by the backward-time $\tau = T - t = 0, 1, .., T$ (see around Eq. B9 for more details), so that the evolution for the probability density $\rho_{\tau}^{B}(x)$ in the growing backward time $\tau - 1 \rightarrow \tau$ is governed by the backward kernel of Eq. 42

$$\rho_{\tau}^{B}(x) = \int_{0}^{1} d\tilde{x} W_{B}(x|\tilde{x}) \rho_{\tau-1}^{B}(\tilde{x}) = \rho_{\tau-1}^{B}(2x) \left[\theta\left(x\right) - \theta\left(x - \frac{1}{2}\right)\right] + \rho_{\tau-1}^{B}(2x-1) \left[\theta\left(x - \frac{1}{2}\right) - \theta\left(x - 1\right)\right]$$
(45)

The physical interpretation is that the new histogram $\rho_{\tau}^B(.)$ corresponds to the gluing of two rescaled histograms $\rho_{\tau-1}^B(.)$ on the two intervals $[0, \frac{1}{2}]$ and $[\frac{1}{2}, 1]$. The iteration of Eq. 45 up to the initial density $\rho_{\tau=0}(.)$ at $\tau = 0$

$$\rho_{\tau}^{B}(x) = \rho_{\tau-2}^{B}(4x) \left[\theta\left(x\right) - \theta\left(x - \frac{1}{4}\right) \right] + \rho_{\tau-2}^{B}(4x-1) \left[\theta\left(x - \frac{1}{4}\right) - \theta\left(x - \frac{1}{2}\right) \right] \\
+ \rho_{\tau-2}^{B}(4x-2) \left[\theta\left(x - \frac{1}{2}\right) - \theta\left(x - \frac{3}{4}\right) \right] + \rho_{\tau-2}^{B}(4x-3) \left[\theta\left(x - \frac{3}{4}\right) - \theta\left(x - 1\right) \right] = \dots \\
= \sum_{j=0}^{2^{\tau}-1} \rho_{0}^{B} \left(2^{\tau}x - j\right) \left[\theta\left(x - \frac{j}{2^{\tau}}\right) - \theta\left(x - \frac{j+1}{2^{\tau}}\right) \right]$$
(46)

means that the histogram $\rho_{\tau}^{B}(.)$ at time τ corresponds to the gluing of 2^{τ} rescaled histograms $\rho_{\tau=0}(.)$ on the 2^{τ} intervals $\left[\frac{j}{2\tau}, \frac{j+1}{2\tau}\right]$ labelled by $j = 0, 1, ..., 2^{\tau} - 1$. In particular, the weight of each of these 2^{τ} intervals $\left[\frac{j}{2\tau}, \frac{j+1}{2\tau}\right]$ is $\frac{1}{2\tau}$

$$\int_{\frac{j}{2^{\tau}}}^{\frac{j+1}{2^{\tau}}} dx \rho_{\tau}^{B}(x) = \int_{\frac{j}{2^{\tau}}}^{\frac{j+1}{2^{\tau}}} dx \rho_{0}^{B} \left(2^{\tau} x - j\right) = \int_{0}^{1} \frac{du}{2^{\tau}} \rho_{0}^{B}(u) = \frac{1}{2^{\tau}}$$
(47)

As a consequence, the backward dynamics will convergence towards the backward uniform steady state

$$\rho_{\tau}^{B}(\tilde{x}) \underset{\tau \to +\infty}{\simeq} \rho_{*}^{B}(\tilde{x}) = 1 = \rho_{*}(\tilde{x})$$

$$\tag{48}$$

that coincides with the forward uniform steady state $\rho_*(\tilde{x}) = 1$ as expected from Eq. B7, but this convergence will be rather weird, since the gluing of 2^{τ} rescaled initial histograms in Eq. 46 will introduce discontinuities of the density or of its derivatives at all positions $\frac{j}{2\tau}$.

C. Spectral decomposition of the backward dynamics

As explained around Eqs B4, the spectral decomposition of the backward kernel

$$w_B(x|\tilde{x}) = \sum_{p=0}^{+\infty} \lambda_p r_p^B(x) l_p^B(\tilde{x})$$
(49)

involves the same eigenvalues $\lambda_p = 2^{-p}$ of Eq. 10 as the forward kernel $w(\tilde{x}|x)$, while the right and left eigenvectors for the backward kernel given by Eq. B5 read in terms of the Bernouilli polynomials of Eq. 12 and in terms of the singular distributions of Eq. 14

$$r_{p}^{B}(x) = l_{p}(x)\rho_{*}(x) = \frac{(-1)^{p-1}}{p!} \left[\delta^{(p-1)}(x-1) - \delta^{(p-1)}(x)\right] \equiv D_{p}(x)$$

$$l_{p}^{B}(\tilde{x}) = \frac{r_{p}(\tilde{x})}{\rho_{*}(\tilde{x})} = B_{p}(\tilde{x})$$
(50)

So the right and left eigenvectors have been exchanged with respect to the forward kernel. The spectral decomposition of Eq. A8 for the backward dynamics

$$\rho_{\tau}^{B}(x) = 1 + \sum_{p=1}^{+\infty} 2^{-pt} D_{p}(x) \int_{0}^{1} d\tilde{x} B_{p}(\tilde{x}) \rho_{0}^{B}(\tilde{x})$$
(51)

is thus very singular, since the distributions $D_p(x)$ of Eq. 14 appear directly and not as scalar products as in the forward spectral decomposition of Eq. 17. These singularities in real space suggest that it is useful to analyze the backward dynamics from the point of view of the binary variables as described in the next subsection.

D. Backward dynamics for the binary variables

The real-space backward kernel of Eq. 42 translates for the binary variables σ_l of Eq. 18 into

$$W_B(\sigma_{\cdot}|\tilde{\sigma}_{\cdot}) = \frac{(\delta_{\sigma_1,0} + \delta_{\sigma_1,1})}{2} \prod_{l=2}^{+\infty} \delta_{\sigma_l,\tilde{\sigma}_{l-1}}$$
(52)

The normalization over σ is obvious, while the uniform steady density of Eq. 24 of the forward dynamics is also the steady density of the backward dynamics

$$P_*^B(\sigma_{\cdot}) = P_*(\sigma_{\cdot}) = \prod_{l=1}^{+\infty} \left(\frac{\delta_{\sigma_l,0} + \delta_{\sigma_l,1}}{2} \right)$$
(53)

by construction (Eq. B7).

The dynamics for the probability $P_{\tau}^{B}(\sigma)$ governed by the backward kernel of Eq. 52

$$P_{\tau}^{B}(\sigma_{1},\sigma_{2},\sigma_{3},...) = \sum_{\tilde{\sigma}_{.}} W_{B}(\sigma_{.}|\tilde{\sigma}_{.}) P_{\tau-1}^{B}(\tilde{\sigma}_{.}) = \frac{(\delta_{\sigma_{1},0} + \delta_{\sigma_{1},1})}{2} \prod_{l=2}^{+\infty} \delta_{\sigma_{l},\tilde{\sigma}_{l-1}} P_{\tau-1}^{B}(\tilde{\sigma}_{.}) \\ = \frac{(\delta_{\sigma_{1},0} + \delta_{\sigma_{1},1})}{2} P_{\tau-1}^{B}(\sigma_{2},\sigma_{3},...)$$
(54)

means that one adds the first coefficient $\sigma_1 = 0, 1$ drawn with the probabilities $(\frac{1}{2}, \frac{1}{2})$, while all the other binary coefficients $(\sigma_2, \sigma_3, ...)$ are distributed with the shifted probability $P^B_{\tau-1}(.)$ at the previous time $(\tau - 1)$. The iteration up to $\tau = 0$

$$P_{\tau}^{B}(\sigma_{1},\sigma_{2},\sigma_{3},...) = \left[\prod_{l=1}^{\tau} \frac{\delta_{\sigma_{l},0} + \delta_{\sigma_{l},1}}{2}\right] P_{0}^{B}(\sigma_{\tau+1},\sigma_{\tau+2},...)$$
(55)

means that the first τ binary coefficients $\sigma_1 = 0, 1$ are independently drawn with the probabilities $(\frac{1}{2}, \frac{1}{2})$, while the higher coefficients $l > \tau$ appear with their initial probability $P_0^B(\sigma_{\tau+1}, \sigma_{\tau+2}, ...)$ at $\tau = 0$. For large τ , Eq. 55 converges towards the uniform distribution of Eq. 53, but the information on the initial distribution $P_0^B(.)$ is still intact and simply shifted towards the higher coefficients $l > \tau$.

V. TRAJECTORY OBSERVABLES VIA THE DEFORMED STOCHASTIC BACKWARD DYNAMICS

In this section, the large deviations of trajectory observables are analyzed via deformations of the stochastic backward dynamics described in the previous section IV. We stress the similarities and the differences with the deformations of the forward deterministic dynamics of section III.

A. Backward dynamics for the generating function via the k-deformed backward kernel $w_B^{[k]}(x|\tilde{x})$

As discussed in Appendix B, the large deviations of trajectories observables can be alternatively studied via the k-deformed backward kernel $w_B(x|\tilde{x})$ of Eq. B10 that reads for the forward doubling kernel of Eq. 4 and its uniform steady density $\rho_*(x) = 1$

$$w_{B}^{[k]}(x|\tilde{x}) = \delta(\tilde{x} - f(x))e^{k\Omega(f(x),x)} = \delta(\tilde{x} - f(x))e^{k\omega(x)}$$

$$= \delta(\tilde{x} - 2x)e^{k\omega(x)} \left[\theta(x) - \theta\left(x - \frac{1}{2}\right)\right] + \delta(\tilde{x} - (2x - 1))e^{k\omega(x)} \left[\theta\left(x - \frac{1}{2}\right) - \theta(x - 1)\right]$$

$$= \frac{\delta(x - \frac{\tilde{x}}{2})e^{k\omega(\frac{\tilde{x}}{2})} + \delta(x - \frac{\tilde{x}+1}{2})e^{k\omega(\frac{\tilde{x}+1}{2})}}{2} \equiv \delta\left(x - \frac{\tilde{x}}{2}\right)w_{B-}^{[k]}(\tilde{x}) + \delta\left(x - \frac{\tilde{x}+1}{2}\right)w_{B+}^{[k]}(\tilde{x})$$
(56)

The last line shows that with respect to the undeformed backward kernel $w_B(x|\tilde{x})$ of Eq. B3, the two pre-images $\left(x_- = \frac{\tilde{x}}{2}; x_+ = \frac{\tilde{x}+1}{2}\right)$ of \tilde{x} that were chosen with the equal probabilities $\left(\frac{1}{2}, \frac{1}{2}\right)$ are now chosen with the k-deformed weights

$$w_{B-}^{[k]}(\tilde{x}) = \frac{e^{k\omega\left(\frac{\tilde{x}}{2}\right)}}{2}$$
$$w_{B+}^{[k]}(\tilde{x}) = \frac{e^{k\omega\left(\frac{\tilde{x}+1}{2}\right)}}{2}$$
(57)

The backward dynamics for the generating function

$$Z_{\tau}^{B[k]}(x) = \int_{0}^{1} d\tilde{x} w_{B}^{[k]}(x|\tilde{x}) Z_{\tau-1}^{B[k]}(\tilde{x})$$

$$= e^{k\omega(x)} Z_{\tau-1}^{B[k]}(2x) \left[\theta\left(x\right) - \theta\left(x - \frac{1}{2}\right) \right] + e^{k\omega(x)} Z_{\tau-1}^{B[k]}(2x-1) \left[\theta\left(x - \frac{1}{2}\right) - \theta\left(x - 1\right) \right]$$
(58)

corresponds to the k-deformation of the backward dynamics of Eq. 45 for the probability density $\rho_{\tau}^{B}(x)$.

B. Eigenvalue problem governing the generating function for large time $\tau \to +\infty$

As explained around Eq. B12, the highest deformed eigenvalue $\lambda_0^{[k]}$ is the same for the backward and the forward kernels, while the positive right and left eigenvectors of the backward kernel satisfying

$$\lambda_{0}^{[k]} r_{0}^{B[k]}(x) = \int d\tilde{x} \, w_{B}^{[k]}(x|\tilde{x}) r_{0}^{B[k]}(\tilde{x}) = r_{0}^{B[k]}(2x) e^{k\omega(x)} \left[\theta\left(x - \frac{1}{2}\right) \right] + r_{0}^{B[k]}(2x - 1) e^{k\omega(x)} \left[\theta\left(x - \frac{1}{2}\right) - \theta\left(x - 1\right) \right]$$

$$\lambda_{0}^{[k]} l_{0}^{B[k]}(\tilde{x}) = \int dx l_{0}^{B[k]}(x) w_{B}^{[k]}(x|\tilde{x}) = l_{0}^{B[k]} \left(\frac{\tilde{x}}{2}\right) w_{B-}^{[k]}(\tilde{x}) + l_{0}^{B[k]} \left(\frac{\tilde{x} + 1}{2}\right) w_{B+}^{[k]}(\tilde{x})$$

$$= l_{0}^{B[k]} \left(\frac{\tilde{x}}{2}\right) \frac{e^{k\omega\left(\frac{\tilde{x}}{2}\right)}}{2} + l_{0}^{B[k]} \left(\frac{\tilde{x} + 1}{2}\right) \frac{e^{k\omega\left(\frac{\tilde{x} + 1}{2}\right)}}{2}$$
(59)

are related to the forward eigenvectors via Eq. B12 that reads for the doubling map with uniform density $\rho_*(x) = 1$

$$r_0^{B[k]}(\tilde{x}) = l_0^{[k]}(\tilde{x}) l_0^{B[k]}(x) = r_0^{[k]}(x)$$
(60)

This exchange between the deformed right and left eigenvectors have very important consequences : (i) the backward right eigenvector $r_0^{B[k]}(\tilde{x}) = l_0^{[k]}(\tilde{x})$ corresponding to the forward left eigenvector is not a smooth function of x. As a consequence, if one construct the probability of the end-point analog to Eq. A15

$$P_{\tau}^{B[k]End}(x) \equiv \frac{Z_{\tau}^{B[k]}(x)}{\int dy Z_{\tau}^{B[k]}(y)} \underset{\tau \to +\infty}{\simeq} \frac{r_0^{B[k]}(x)}{\int dy r_0^{B[k]}(y)} = \frac{l_0^{[k]}(x)}{\int dy l_0^{[k]}(y)} \equiv P_*^{B[k]End}(x)$$
(61)

it will converge towards a singular distribution $P_*^{[k]End}(x)$. (ii) the backward left eigenvector $l_0^{B[k]}(x) = r_0^{[k]}(x)$ corresponding to the forward right eigenvector is a smooth function of x and can be used to construct the canonical conditioned backward kernel as described in the next subsection.

Canonical conditioned backward kernel $w_B^{C[k]}(x|\tilde{x})$ with k-deformed probabilities for the two pre-images С.

The conditioned backward kernel $w_B^{C[k]}(x|\tilde{x})$ of Eq. B14 reads using the backward kernel of Eq. 56

$$w_{B}^{C[k]}(x|\tilde{x}) \equiv \frac{1}{\lambda_{0}^{[k]}} l_{0}^{B[k]}(x) w^{B[k]}(x|\tilde{x}) \frac{1}{l_{0}^{B[k]}(\tilde{x})} = \frac{l_{0}^{B[k]}(x) w^{B[k]}(x|\tilde{x})}{\int dz l_{0}^{B[k]}(z) w^{B[k]}(z|\tilde{x})}$$
$$= \frac{\delta \left(x - \frac{\tilde{x}}{2}\right) l_{0}^{B[k]}\left(\frac{\tilde{x}}{2}\right) w_{B-}^{[k]}(\tilde{x}) + \delta \left(x - \frac{\tilde{x}+1}{2}\right) l_{0}^{B[k]}\left(\frac{\tilde{x}+1}{2}\right) w_{B+}^{[k]}(\tilde{x})}{l_{0}^{B[k]}\left(\frac{\tilde{x}}{2}\right) w_{B-}^{[k]}(\tilde{x}) + l_{0}^{B[k]}\left(\frac{\tilde{x}+1}{2}\right) w_{B+}^{[k]}(\tilde{x})}$$
$$\equiv \delta \left(x - \frac{\tilde{x}}{2}\right) p_{B-}^{C[k]}(\tilde{x}) + \delta \left(x - \frac{\tilde{x}+1}{2}\right) p_{B+}^{C[k]}(\tilde{x}) \tag{62}$$

The last line shows that with respect to the undeformed backward kernel $w_B(x|\tilde{x})$ of Eq. B3, the two pre-images ($x_{-} = \frac{\tilde{x}}{2}; x_{+} = \frac{\tilde{x}+1}{2}$) of \tilde{x} that were chosen with the equal probabilities $(\frac{1}{2}, \frac{1}{2})$ are now chosen with the k-deformed complementary probabilities $p_{B-}^{C[k]}(\tilde{x}) + p_{B+}^{C[k]}(\tilde{x}) = 1$ that involve the deformed weights $w_{B\pm}^{[k]}(.)$ of Eq. 57 and the deformed backward left eigenvector $l_0^{B[k]}(.)$ of Eq. 59

$$p_{B_{+}}^{C[k]}(\tilde{x}) = \frac{l_{0}^{B[k]}\left(\frac{\tilde{x}}{2}\right)w_{B_{-}}^{[k]}(\tilde{x})}{l_{0}^{B[k]}\left(\frac{\tilde{x}}{2}\right)w_{B_{-}}^{[k]}(\tilde{x}) + l_{0}^{B[k]}\left(\frac{\tilde{x}+1}{2}\right)w_{B_{+}}^{[k]}(\tilde{x})} = \frac{l_{0}^{B[k]}\left(\frac{\tilde{x}}{2}\right)e^{k\omega\left(\frac{\tilde{x}}{2}\right)}}{l_{0}^{B[k]}\left(\frac{\tilde{x}}{2}\right)e^{k\omega\left(\frac{\tilde{x}+1}{2}\right)}e^{k\omega\left(\frac{\tilde{x}+1}{2}\right)}}$$
$$p_{B_{+}}^{C[k]}(\tilde{x}) = \frac{l_{0}^{B[k]}\left(\frac{\tilde{x}+1}{2}\right)w_{B_{+}}^{[k]}(\tilde{x})}{l_{0}^{B[k]}\left(\frac{\tilde{x}}{2}\right)e^{k\omega\left(\frac{\tilde{x}}{2}\right)}e^{k\omega\left(\frac{\tilde{x}+1}{2}\right)}}$$
$$= \frac{l_{0}^{B[k]}\left(\frac{\tilde{x}+1}{2}\right)e^{k\omega\left(\frac{\tilde{x}+1}{2}\right)}}{l_{0}^{B[k]}\left(\frac{\tilde{x}}{2}\right)e^{k\omega\left(\frac{\tilde{x}+1}{2}\right)}e^{k\omega\left(\frac{\tilde{x}+1}{2}\right)}}$$
(63)

The delta-function identity yields that Eq. 62 can be rewritten as

$$w_{B}^{C[k]}(x|\tilde{x}) \equiv \delta\left(x - \frac{\tilde{x}}{2}\right) p_{B-}^{C[k]}(\tilde{x}) + \delta\left(x - \frac{\tilde{x}+1}{2}\right) p_{B+}^{C[k]}(\tilde{x}) \\ = \delta\left(\tilde{x} - 2x\right) p_{B-}^{C[k]}(\tilde{x}) \left[\theta\left(x\right) - \theta\left(x - \frac{1}{2}\right)\right] + \delta\left(\tilde{x} - (2x - 1)\right) p_{B+}^{C[k]}(\tilde{x}) \left[\theta\left(x - \frac{1}{2}\right) - \theta\left(x - 1\right)\right]$$
(64)

The conditioned backward steady state $\rho_*^{BC[k]}(x)$ corresponding its right eigenvector associated to the eigenvalue unity

$$\rho_*^{BC[k]}(x) = \int_0^1 d\tilde{x} w_B^{C[k]}(x|\tilde{x}) \rho_*^{BC[k]}(\tilde{x})$$

= $p_{B-}^{C[k]}(2x) \rho_*^{BC[k]}(2x) \left[\theta\left(x\right) - \theta\left(x - \frac{1}{2}\right) \right] + p_{B+}^{C[k]}(2x-1) \rho_*^{BC[k]}(2x-1) \left[\theta\left(x - \frac{1}{2}\right) - \theta\left(x - 1\right) \right] (65)$

is given by Eq. B13

$$\rho_*^{BC[k]}(x) \equiv l_0^{B[k]}(x) r_0^{B[k]}(x) = l_0^{[k]}(x) r_0^{[k]}(x)$$
(66)

and is thus expected to be singular in x. It is thus useful to consider the formulation of the backward deformed and conditioned dynamics in terms of the binary variables.

D. Analysis via the backward deformed and conditioned dynamics for the binary variables σ_l

1. Properties of the backward deformed kernel in the binary variables σ_l

With the notation

$$\Omega(\sigma_{.}) \equiv \omega \left(x = \sum_{l=1}^{+\infty} \frac{\sigma_l}{2^l} \right)$$
(67)

for the parametrization of the trajectory observable of Eq. 27 in terms of the binary observables σ_l , the k-deformation of the backward kernel of Eq. 52

$$W_B^{[k]}(\sigma_{.}|\tilde{\sigma}_{.}) = e^{k\Omega(\sigma_{.})} W_B(\sigma_{.}|\tilde{\sigma}_{.}) = e^{k\Omega(\sigma_{.})} \frac{(\delta_{\sigma_1,0} + \delta_{\sigma_1,1})}{2} \prod_{l=2}^{+\infty} \delta_{\sigma_l,\tilde{\sigma}_{l-1}}$$
$$\equiv \left[\delta_{\sigma_1,0} W_{B-}^{[k]}(\tilde{\sigma}) + \delta_{\sigma_1,1} W_{B+}^{[k]}(\tilde{\sigma}) \right] \prod_{l=2}^{+\infty} \delta_{\sigma_l,\tilde{\sigma}_{l-1}}$$
(68)

involve the two following weights are the analog of the real-space weights of Eq. 57

$$W_{B-}^{[k]}(\tilde{\sigma}) = \frac{e^{k\Omega(0,\tilde{\sigma}_{1},\tilde{\sigma}_{2},..)}}{2} \\ W_{B+}^{[k]}(\tilde{\sigma}) = \frac{e^{k\Omega(1,\tilde{\sigma}_{1},\tilde{\sigma}_{2},..)}}{2}$$
(69)

The eigenvalues equations for the backward deformed eigenvectors read

$$\lambda_{0}^{[k]} L_{0}^{B[k]}(\tilde{\sigma}_{1}, \tilde{\sigma}_{2}, ...) = \sum_{\sigma_{.}} L_{0}^{B[k]}(\sigma_{.}) W_{B}^{[k]}(\sigma_{.}|\tilde{\sigma}_{.})$$
$$= \frac{L_{0}^{B[k]}(0, \tilde{\sigma}_{1}, \tilde{\sigma}_{2}, ...) e^{k\Omega(0, \tilde{\sigma}_{1}, \tilde{\sigma}_{2}, ...)} + L_{0}^{B[k]}(1, \tilde{\sigma}_{1}, \tilde{\sigma}_{2}, ...) e^{k\Omega(1, \tilde{\sigma}_{1}, \tilde{\sigma}_{2}, ...)}}{2}$$
(70)

and

$$\lambda_{0}^{[k]} R_{0}^{B[k]}(\sigma_{1}, \sigma_{2}, ...) = \sum_{\tilde{\sigma}} W_{B}^{[k]}(\sigma_{.} | \tilde{\sigma}_{.}) R_{0}^{B[k]}(\tilde{\sigma}_{.})$$
$$= \frac{\delta_{\sigma_{1},0} e^{k\Omega(0, \sigma_{2}, \sigma_{3}, ...)} + \delta_{\sigma_{1},1} e^{k\Omega(1, \sigma_{2}, \sigma_{3}, ...)}}{2} R_{0}^{B[k]}(\sigma_{2}, \sigma_{3}, ...)$$
(71)

2. Construction of the backward conditioned kernel in the binary variables σ_l

The deformed left eigenvector $L_0^{B[k]}(\sigma_{\cdot})$ of Eq. 70 is useful to construct the backward conditioned kernel via Eq. B14

$$W_{B}^{C[k]}(\sigma_{.}|\tilde{\sigma}_{.}) \equiv \frac{1}{\lambda_{0}^{[k]}} L_{0}^{B[k]}(\sigma_{.}) W_{B}^{[k]}(\sigma_{.}|\tilde{\sigma}_{.}) \frac{1}{L_{0}^{B[k]}(\tilde{\sigma}_{.})} = \frac{L_{0}^{B[k]}(\sigma_{.}) W_{B}^{[k]}(\sigma_{.}|\tilde{\sigma}_{.})}{\sum_{\sigma'_{.}} L_{0}^{B[k]}(\sigma'_{.}) W_{B}^{[k]}(\sigma'_{.}|\tilde{\sigma}_{.})} = \left[\delta_{\sigma_{1},0} p_{B^{-}}^{C[k]}(\tilde{\sigma}_{.}) + \delta_{\sigma_{1},1} p_{B^{+}}^{C[k]}(\tilde{\sigma}_{.})\right] \prod_{l=2}^{+\infty} \delta_{\sigma_{l},\tilde{\sigma}_{l-1}}$$
(72)

where the two complementary probabilities $p_{B-}^{C[k]}(\tilde{\sigma}_{.}) + p_{B+}^{C[k]}(\tilde{\sigma}_{.}) = 1$ read

$$p_{B_{-}}^{C[k]}(\tilde{\sigma}_{.}) \equiv \frac{L_{0}^{B[k]}(0,\tilde{\sigma}_{1},\tilde{\sigma}_{2},..)e^{k\Omega(0,\tilde{\sigma}_{1},\tilde{\sigma}_{2},..)}}{L_{0}^{B[k]}(0,\tilde{\sigma}_{1},\tilde{\sigma}_{2},..)e^{k\Omega(0,\tilde{\sigma}_{1},\tilde{\sigma}_{2},..)} + L_{0}^{B[k]}(1,\tilde{\sigma}_{1},\tilde{\sigma}_{2},..)e^{k\Omega(1,\tilde{\sigma}_{1},\tilde{\sigma}_{2},..)}}$$
$$p_{B_{+}}^{C[k]}(\tilde{\sigma}_{.}) \equiv \frac{L_{0}^{B[k]}(1,\tilde{\sigma}_{1},\tilde{\sigma}_{2},..)e^{k\Omega(1,\tilde{\sigma}_{1},\tilde{\sigma}_{2},..)}}{L_{0}^{B[k]}(0,\tilde{\sigma}_{1},\tilde{\sigma}_{2},..)e^{k\Omega(0,\tilde{\sigma}_{1},\tilde{\sigma}_{2},..)} + L_{0}^{B[k]}(1,\tilde{\sigma}_{1},\tilde{\sigma}_{2},..)e^{k\Omega(1,\tilde{\sigma}_{1},\tilde{\sigma}_{2},..)}}$$
(73)

This formulation of the backward conditioned dynamics in the binary variables σ_l is indeed helpful to clarify the properties that are singular in the real-space variable $x \in [0, 1[$, as shown explicitly on an example of trajectory observable in the next section.

VI. EXAMPLE OF THE TRAJECTORY OBSERVABLE $\omega(x) = x$ FROM VARIOUS PERSPECTIVES

In this section, the different perspectives described in sections III and V are illustrated on the simplest trajectory observable $\omega(x) = x$ for the doubling map [68].

A. Analysis via the k-deformed forward dynamics on the real-space interval $x \in [0, 1]$

1. Exact solution for the deformed eigenvalue $\lambda_0^{[k]}$ and the deformed right eigenvector $r_0^{[k]}(.)$ [68]

The example of the trajectory observable $\omega(x) = x$ where the right eigenvalue Eq. 31 reads

$$\lambda_0^{[k]} r_0^{[k]}(\tilde{x}) = \frac{e^{k\frac{\tilde{x}}{2}} r_0^{[k]}(\frac{\tilde{x}}{2}) + e^{k\frac{\tilde{x}+1}{2}} r_0^{[k]}(\frac{\tilde{x}+1}{2})}{2}$$
(74)

was solved in [68] with the k-deformed eigenvalue

$$\lambda_0^{[k]} = \frac{1 + e^k}{2} \tag{75}$$

and the k-deformed right eigenvector

$$r_0^{[k]}(\tilde{x}) = e^{k\left(\tilde{x} - \frac{1}{2}\right)} \tag{76}$$

The corresponding left eigenvalue Eq. 31

$$0 \le x < \frac{1}{2}: \quad \lambda_0^{[k]} l_0^{[k]}(x) = l_0^{[k]}(2x) e^{kx}$$

$$\frac{1}{2} \le x < 1: \quad \lambda_0^{[k]} l_0^{[k]}(x) = l_0^{[k]}(2x-1) e^{kx}$$
(77)

is expected to be singular in real space as soon as $k \neq 0$, as indicated by the perturbation theory in the next subsection.

2. Comparison with the general perturbation theory in k

Let us compare with the perturbation theory of subsection IIIC. The series expansion of the exact eigenvalue of Eq. 75

$$\lambda_0^{[k]} = \frac{1+e^k}{2} = 1 + \frac{k}{2} + \frac{k^2}{4} + O(k^3) \tag{78}$$

is in agreement with the first-order correction of Eq 36 and the second-order correction Eq. 37

$$\lambda^{(1)} = \int_0^1 dx x = \frac{1}{2}$$

$$\lambda^{(2)} = \int_0^1 dx \frac{x^2}{2} + \int_0^1 dx x \left(x - \frac{1}{2}\right) = \frac{1}{4}$$
(79)

The series expansion of the exact right eigenvector of Eq. 76

$$r_0^{[k]}(\tilde{x}) = e^{k\left(\tilde{x} - \frac{1}{2}\right)} = 1 + k\left(\tilde{x} - \frac{1}{2}\right) + O(k^2)$$
(80)

is in agreement with the first-order correction $r_0^{(1)}(\tilde{x})$ Eq. 39 for $\omega(x) = x$ that reduces to the first Bernoulli polynomial $B_1(\tilde{x})$ of Eq. 12

$$r_0^{(1)}(\tilde{x}) = B_1(\tilde{x}) = \left(\tilde{x} - \frac{1}{2}\right)$$
 (81)

The first-order perturbation theory of Eq. 40 for the left eigenvector

$$l_0^{(1)}(\tilde{x}) = \sum_{p=1}^{+\infty} D_p(\tilde{x}) \left[\frac{1}{1-2^{-p}} \int_0^1 dx x B_p(x) \right] = \sum_{p=1}^{+\infty} D_p(\tilde{x}) \left[\frac{1}{1-2^{-p}} \int_0^1 dx B_1(x) B_p(x) \right]$$
(82)

involves the singular distributions $D_p(.)$ of Eq. 14 for all $p = 1, 2, .., +\infty$.

B. Analysis via the k-deformed forward kernel for the binary variables σ_l

For $\omega(x) = x$, the k-deformed kernel

$$W^{[k]}(\tilde{\sigma}_{.}|\sigma_{.}) \equiv e^{k\Omega(\sigma_{.})} \prod_{l=1}^{+\infty} \delta_{\tilde{\sigma}_{l},\sigma_{l+1}}$$
(83)

is factorized in the binary variables

$$W^{[k]}(\tilde{\sigma}_{.}|\sigma_{.}) = e^{k \sum_{l=1}^{+\infty} \frac{\sigma_{l}}{2^{l}}} \prod_{l=1}^{+\infty} \delta_{\tilde{\sigma}_{l},\sigma_{l+1}} = e^{k \frac{\sigma_{1}}{2}} \prod_{l=1}^{+\infty} \left[e^{k \frac{\sigma_{l+1}}{2^{l+1}}} \delta_{\tilde{\sigma}_{l},\sigma_{l+1}} \right]$$
(84)

The k-deformed right eigenvector of Eq. 76 in real space translates for the binary variables into the factorized form

$$R_0^{[k]}(\sigma_{\cdot}) = e^{-\frac{k}{2}} \prod_{l=1}^{+\infty} \frac{e^{k\frac{\sigma_l}{2^l}}}{2}$$
(85)

The solution for the corresponding left eigenvector

$$\frac{1+e^{k}}{2} L_{0}^{[k]}(\sigma_{1},\sigma_{2},...) = \sum_{\tilde{\sigma}_{.}} L_{0}^{[k]}(\tilde{\sigma}) W^{[k]}(\tilde{\sigma}_{.}|\sigma_{.}) = \sum_{\tilde{\sigma}_{.}} L_{0}^{[k]}(\tilde{\sigma}) e^{k\frac{\sigma_{1}}{2}} \prod_{l=1}^{+\infty} \left[e^{k\frac{\sigma_{l+1}}{2^{l+1}}} \delta_{\tilde{\sigma}_{l},\sigma_{l+1}} \right] \\
= L_{0}^{[k]}(\sigma_{2},\sigma_{3},...) \prod_{l=1}^{+\infty} e^{k\frac{\sigma_{l}}{2^{l}}}$$
(86)

will be easier to derive later with its appropriate normalization via the backward perspective (see Eq. 100).

C. Analysis via the k-conditioned backward kernel $w_B^{C[k]}(x|\tilde{x})$ on the real-space interval $x \in [0, 1[$

The eigenvalue $\lambda_0^{[k]} = \frac{1+e^k}{2}$ of Eq. 75 that is associated to the k-deformed forward right eigenvector of Eq. 76 is now associated to the k-deformed backward left eigenvector of Eq. 60

$$l_0^{B[k]}(\tilde{x}) = r_0^{[k]}(\tilde{x}) = e^{k\left(\tilde{x} - \frac{1}{2}\right)}$$
(87)

As a consequence, the two k-deformed probabilities of Eq. 63

$$p_{B-}^{C[k]}(\tilde{x}) = \frac{1}{1+e^k}$$

$$p_{B+}^{C[k]}(\tilde{x}) = \frac{e^k}{1+e^k}$$
(88)

are independent of \tilde{x} , and the conditioned backward kernel of Eq. 64 reduces to

$$w_{B}^{C[k]}(x|\tilde{x}) = \frac{\delta\left(x - \frac{\tilde{x}}{2}\right) + \delta\left(x - \frac{\tilde{x}+1}{2}\right)e^{k}}{1 + e^{k}} \\ = \frac{\delta\left(\tilde{x} - 2x\right)}{1 + e^{k}} \left[\theta\left(x\right) - \theta\left(x - \frac{1}{2}\right)\right] + \frac{\delta\left(\tilde{x} - (2x - 1)\right)e^{k}}{1 + e^{k}} \left[\theta\left(x - \frac{1}{2}\right) - \theta\left(x - 1\right)\right]$$
(89)

However the conditioned backward steady state $\rho_*^{BC[k]}(x)$ of Eq. 66 is expected to be singular in x, and it is thus useful to turn to the formulation of the backward conditioned dynamics in terms of the binary variables.

D. Analysis via the backward deformed and conditioned kernel for the binary variables σ_l

1. Backward deformed kernel for the binary variables σ_l

For $\omega(x) = x$, the backward deformed kernel of Eq. 68 is factorized in the binary variables

$$W_{B}^{[k]}(\sigma_{.}|\tilde{\sigma}_{.}) = e^{k \sum_{l=1}^{+\infty} \frac{\sigma_{l}}{2^{l}}} \left(\frac{\delta_{\sigma_{1},0} + \delta_{\sigma_{1},1}}{2} \right) \prod_{l=2}^{+\infty} \delta_{\sigma_{l},\tilde{\sigma}_{l-1}} = \left(\frac{\delta_{\sigma_{1},0} + e^{\frac{k}{2}} \delta_{\sigma_{1},1}}{2} \right) \prod_{l=2}^{+\infty} \left[e^{k \frac{\sigma_{l}}{2^{l}}} \delta_{\sigma_{l},\tilde{\sigma}_{l-1}} \right]$$
(90)

The backward left eigenvector of Eq. 87 in real space translates for the binary variables into

$$L_0^{B[k]}(\sigma_{.}) = e^{-\frac{k}{2}} \prod_{l=1}^{+\infty} e^{k\frac{\sigma_l}{2l}}$$
(91)

2. Backward conditioned kernel for the binary variables σ_l with its explicit conditioned steady density $P_B^{C[k]}(\sigma_l)$

The backward conditioned kernel of Eq. 89 in real space translates for the binary variables into

$$W_B^{C[k]}(\sigma_{\cdot}|\tilde{\sigma}_{\cdot}) = \left(\frac{\delta_{\sigma_1,0} + e^k \delta_{\sigma_1,1}}{1 + e^k}\right) \prod_{l=2}^{+\infty} \delta_{\sigma_l,\tilde{\sigma}_{l-1}}$$
(92)

The corresponding conditioned steady probability is now obvious and factorized in these binary variables

$$P_B^{C[k]}(\sigma_{.}) = \prod_{l=1}^{+\infty} \left(\frac{\delta_{\sigma_l,0} + e^k \delta_{\sigma_l,1}}{1 + e^k} \right) = \prod_{l=1}^{+\infty} \left(\frac{e^{k\sigma_l}}{1 + e^k} \right)$$
(93)

Since the translation in real-space $x \in [0, 1]$

$$\rho_B^{C[k]}(x) = \sum_{\sigma_{\cdot}} P_B^{C[k]}(\sigma_{\cdot}) \delta\left(x - \sum_{l=1}^{+\infty} \frac{\sigma_l}{2^l}\right)$$
(94)

is not smooth, it is useful to characterize this real-space density $\rho_B^{C[k]}(x)$ via its Fourier coefficients of Eq. C2 for $n \in \mathbb{Z}$

$$\hat{\rho}_{B}^{C[k]}(n) = \int_{0}^{1} dx \rho_{B}^{C[k]}(x) e^{-i2\pi nx} = \sum_{\sigma} P_{B}^{C[k]}(\sigma) e^{-i2\pi nx} \sum_{l=1}^{+\infty} \frac{\sigma_{l}}{2^{l}} = \prod_{l=1}^{+\infty} \left(\sum_{\sigma_{l}=0,1} \frac{e^{\left(k-i2\pi \frac{n}{2^{l}}\right)\sigma_{l}}}{1+e^{k}} \right)$$
$$= \prod_{l=1}^{+\infty} \left(\frac{1+e^{\left(k-i2\pi \frac{n}{2^{l}}\right)}}{1+e^{k}} \right)$$
(95)

3. Explicit expression of the backward deformed right eigenvector $R_0^{B[k]}(\sigma_1)$

Since the backward conditioned steady state $P_B^{C[k]}(\sigma)$ given by the explicit expression of Eq. 93 coincides with the product $L_0^{B[k]}(\sigma)R_0^{B[k]}(\sigma)$ of the backward deformed eigenvectors, we can use the explicit expression of Eq. 91 for the left eigenvector $L_0^{B[k]}(\sigma)$ to obtain that the right eigenvector $R_0^{B[k]}(\sigma)$ reads

$$R_0^{B[k]}(\sigma_{.}) = \frac{P_B^{C[k]}(\sigma_{.})}{L_0^{B[k]}(\sigma_{.})} = \prod_{l=1}^{+\infty} \frac{\left(\frac{\delta_{\sigma_l,0} + e^k \delta_{\sigma_l,1}}{1 + e^k}\right)}{e^{k\frac{\sigma_l}{2^l}}} = \prod_{l=1}^{+\infty} \left[\frac{\left(\delta_{\sigma_l,0} + e^{k(1 - \frac{1}{2^l})} \delta_{\sigma_l,1}\right)}{1 + e^k}\right]$$
(96)

and satisfies indeed the eigenvalue equation involving the backward deformed kernel of Eq. 90

$$\sum_{\tilde{\sigma}_{.}} W_{B}^{[k]}(\sigma_{.}|\tilde{\sigma}_{.}) R_{0}^{B[k]}(\tilde{\sigma}_{.}) = \frac{1+e^{k}}{2} R_{0}^{B[k]}(\sigma_{.})$$
(97)

Since the translation in real-space $x \in [0, 1]$

$$r_0^{B[k]}(x) = \sum_{\sigma_{\cdot}} R_0^{B[k]}(\sigma_{\cdot}) \delta\left(x - \sum_{l=1}^{+\infty} \frac{\sigma_l}{2^l}\right)$$
(98)

is not smooth, it is useful to to characterize this real-space eigenvector $r_0^{B[k]}(x)$ via its Fourier coefficients of Eq. C2 for $n \in \mathbb{Z}$

$$\hat{r}_{0}^{B[k]}(n) = \int_{0}^{1} dx r_{0}^{B[k]}(x) e^{-i2\pi nx} = \sum_{\sigma} r_{0}^{B[k]}(\sigma) e^{-i2\pi n \sum_{l=1}^{+\infty} \frac{\sigma_{l}}{2^{l}}}$$
$$= \prod_{l=1}^{+\infty} \left(\frac{1 + e^{\left[k(1 - \frac{1}{2^{l}}) - i2\pi \frac{n}{2^{l}}\right]}}{1 + e^{k}} \right)$$
(99)

4. Explicit expression of the forward deformed left eigenvector $L_0^{[k]}(\sigma_{\cdot})$

Similarly, the forward deformed left eigenvector $L_0^{[k]}(\sigma)$ can be computed from the ratio between the conditioned density $\rho^{C[k]}(\sigma) = \rho_B^{C[k]}(\sigma)$ of Eq. 93 and the forward deformed right eigenvector $R_0^{[k]}(\sigma)$ of Eq. 85,

$$L_{0}^{[k]}(\sigma_{\cdot}) = \frac{P^{C[k]}(\sigma_{\cdot})}{R_{0}^{[k]}(\sigma_{\cdot})} = \prod_{l=1}^{+\infty} \frac{\left(\frac{\delta_{\sigma_{l},0} + e^{k}\delta_{\sigma_{l},1}}{1+e^{k}}\right)}{\left(\frac{e^{k\frac{\sigma_{l}}{2}}}{2}\right)} = \prod_{l=1}^{+\infty} \left[\frac{2}{1+e^{k}}\left(\delta_{\sigma_{l},0} + e^{k(1-\frac{1}{2^{l}})}\delta_{\sigma_{l},1}\right)\right] = \prod_{l=1}^{+\infty} \left[\frac{2e^{k\sigma_{l}(1-\frac{1}{2^{l}})}}{1+e^{k}}\right]$$
(100)

that satisfies indeed the eigenvalue Eq. 86.

In real space, the forward deformed left eigenvector $l_0^{[k]}(x) = r_0^{B[k]}(x)$ coincides with the backward deformed right eigenvector $r_0^{B[k]}(x)$ of Eq. 98 with its Fourier coefficients of Eq. 99 : the singular character of $l_0^{[k]}(x)$ in real space was already found numerically on Fig S1 of [69].

VII. LARGE DEVIATIONS FOR THE BACKWARD EMPIRICAL DENSITY IN THE BINARY SPACE

In this section, the explicit large deviations at level 2.5 for Markov chains (see the reminder in subsection A 3 of Appendix A) are applied to the stochastic backward dynamics associated to the doubling map in order to obtain the explicit large deviations at level 2 for the empirical density alone.

A. Reminder on the explicit large deviations at level 2.5 for an arbitrary Markov chain in binary space

For an arbitrary Markov chain in binary space governed by the kernel $W(\sigma_{.}|\tilde{\sigma}_{.})$, the joint distribution $P_{T}^{[2.5]}\left[\mathring{P}(.);\mathring{W}(.|.)\right]$ of the empirical probability $\mathring{P}(\sigma_{.})$ and of the empirical kernel $\mathring{W}(\sigma_{.}|\tilde{\sigma}_{.})$ seen during a trajectory over the time-window T follows the large deviation form at the so-called level 2.5 (see the reminder in Appendix A around Eq. A42 in real-space variables)

$$P_{T}^{[2.5]} \begin{bmatrix} \mathring{P}(.); \mathring{W}(.|.) \end{bmatrix} \xrightarrow{\simeq}_{T \to +\infty} \delta \left(\sum_{\sigma_{.}} \mathring{P}(\sigma_{.}) - 1 \right) \left[\prod_{\tilde{\sigma}_{.}} \delta \left(\sum_{\sigma_{.}} \mathring{W}(\sigma_{.}|\tilde{\sigma}_{.}) - 1 \right) \right] \left[\prod_{\sigma_{.}} \delta \left(\sum_{\tilde{\sigma}_{.}} \mathring{W}(\sigma_{.}|\tilde{\sigma}_{.}) \mathring{P}(\tilde{\sigma}_{.}) - \mathring{P}(\sigma_{.}) \right) \right] e^{-TI_{2.5}} \begin{bmatrix} \mathring{P}(.); \mathring{W}(.|.) \end{bmatrix}$$

$$(101)$$

where the rate function $I_{2.5}\left[\mathring{P}(.); \mathring{W}(.|.)\right]$ at level 2.5 is given by the explicit relative entropy

$$I_{2.5}\left[\mathring{P}(.);\mathring{W}(.|.)\right] = \sum_{\sigma_{.}}\sum_{\tilde{\sigma}_{.}}\mathring{W}(\sigma_{.}|\tilde{\sigma}_{.})\mathring{P}(\tilde{\sigma}_{.})\ln\left(\frac{\mathring{W}(\sigma_{.}|\tilde{\sigma}_{.})}{W(\sigma_{.}|\tilde{\sigma}_{.})}\right)$$
(102)

while the prefactors in front of the exponential in Eq. 101 impose the three following constitutive constraints for the empirical observables (see the reminder of Appendix A around Eqs A38 A40 A41 in real-space variables)

(i) the empirical probability $P(\sigma_{\cdot})$ has to be normalized over σ_{\cdot}

(ii) the empirical kernel $W(\sigma_{\cdot}|\tilde{\sigma}_{\cdot})$ has to be normalized over σ_{\cdot} for any $\tilde{\sigma}_{\cdot}$

(iii) the empirical probability $\check{P}(\sigma_{\cdot})$ should be steady with respect to the dynamics governed by the empirical kernel $\mathring{W}(\sigma_{\cdot}|\tilde{\sigma}_{\cdot})$.

As recalled in Appendix A, all the lower levels can be obtained via optimization of the level 2.5 in the presence of constraints, in particular the level 2 concerning the distribution $P_T^{[2]}\left[\mathring{P}(.)\right]$ of the empirical probability $\mathring{P}(\sigma_.)$ alone, that can be obtained from the integration of the joint distribution $P_T^{[2.5]}\left[\mathring{P}(.); \mathring{W}(.|.)\right]$ of Eq. 101 over the empirical kernel $\mathring{W}(\sigma_.|\tilde{\sigma}_.)$ as follows

$$P_{T}^{[2]}\left[\mathring{P}(.)\right] \equiv \int d\mathring{W}(.|.)P_{T}^{[2.5]}\left[\mathring{P}(.);\mathring{W}(.|.)\right]$$

$$\sum_{T \to +\infty} \delta\left(\sum_{\sigma_{.}} \mathring{P}(\sigma_{.}) - 1\right) \int d\mathring{W}(.|.)\left[\prod_{\tilde{\sigma}_{.}} \delta\left(\sum_{\sigma_{.}} \mathring{W}(\sigma_{.}|\tilde{\sigma}_{.}) - 1\right)\right] \left[\prod_{\sigma_{.}} \delta\left(\sum_{\tilde{\sigma}_{.}} \mathring{W}(\sigma_{.}|\tilde{\sigma}_{.})\mathring{P}(\tilde{\sigma}_{.}) - \mathring{P}(\sigma_{.})\right)\right] e^{-TI_{2.5}\left[\mathring{P}(.);\mathring{W}(.|.)\right]}$$

$$\sum_{T \to +\infty} \delta\left(\sum_{\sigma_{.}} \mathring{P}(\sigma_{.}) - 1\right) e^{-TI_{2}\left[\mathring{P}(.)\right]}$$

$$(103)$$

So the rate function $I_2\left[\mathring{P}(.)\right]$ at level 2 corresponds to the optimization of the rate function $I_{2.5}\left[\mathring{P}(.);\mathring{W}(.|.)\right]$ over the empirical kernel $\mathring{W}(\sigma_.|\tilde{\sigma}_.)$ satisfying the constitutive constraints, but for an arbitrary Markov chain, the solution of this constrained optimization cannot be written as an explicit function of the empirical probability $\mathring{P}(\sigma_.)$.

In the next section, we return to the special case of the backward stochastic dynamics of the doubling map in the binary variables in order to obtain what levels of large deviations can be written explicitly.

B. Application to the backward stochastic dynamics of the doubling map in the binary variables

Let us apply Eq. 101 to the special case of the backward kernel $W_B(\sigma_.|\tilde{\sigma}_.)$ of Eq. 52 in the binary variables associated to the the doubling map. In order for the rate function $I_{2.5}\left[\mathring{P}(.); \mathring{W}(.|.)\right]$ of Eq. 102 to remain finite, the empirical backward kernel $\mathring{W}_B(\sigma_.|\tilde{\sigma}_.)$ has to be of the form

$$\mathring{W}_B(\sigma_{\cdot}|\tilde{\sigma}_{\cdot}) = \left[\delta_{\sigma_1,0}\mathring{p}_{B-}(\tilde{\sigma}_{\cdot}) + \delta_{\sigma_1,1}\mathring{p}_{B+}(\tilde{\sigma}_{\cdot})\right] \prod_{l=2}^{+\infty} \delta_{\sigma_l,\tilde{\sigma}_{l-1}}$$
(104)

where the normalization condition of the empirical kernel $W(\sigma_{\perp}|\tilde{\sigma}_{\perp})$ over σ_{\perp} for any $\tilde{\sigma}_{\perp}$ reads

$$1 = \sum_{\sigma_{.}} \mathring{W}_{B}(\sigma_{.}|\tilde{\sigma}_{.}) = \mathring{p}_{B-}(\tilde{\sigma}_{.}) + \mathring{p}_{B+}(\tilde{\sigma}_{.})$$
(105)

The condition (iii) imposing that the empirical density $\check{P}_B(\sigma_{\cdot})$ is steady with respect to the dynamics governed by the empirical kernel $\mathring{W}_B(\sigma_{\cdot}|\tilde{\sigma}_{\cdot})$

$$\mathring{P}_{B}(\sigma_{1},\sigma_{2},...) = \sum_{\tilde{\sigma}_{.}} \mathring{W}_{B}(\sigma_{.}|\tilde{\sigma}_{.}) \mathring{P}_{B}(\tilde{\sigma}_{.}) = [\delta_{\sigma_{1},0} \mathring{p}_{B-}(\sigma_{2},\sigma_{3},...) + \delta_{\sigma_{1},1} \mathring{p}_{B+}(\sigma_{2},\sigma_{3},...)] \mathring{P}_{B}(\sigma_{2},\sigma_{3},...)$$
(106)

can be used to compute the two complementary empirical probabilities $\mathring{p}_{B\pm}(\tilde{\sigma}_{.})$ that parametrize the empirical kernel of Eq. 104 in terms of the empirical probability $\mathring{P}(.)$

$$\dot{p}_{B-}(\sigma_2, \sigma_3, ...) = \frac{\dot{P}_B(0, \sigma_2, \sigma_3, ...)}{\dot{P}_B(\sigma_2, \sigma_3, ...)}$$

$$\dot{p}_{B+}(\sigma_2, \sigma_3, ...) = \frac{\dot{P}_B(1, \sigma_2, \sigma_3, ...)}{\dot{P}_B(\sigma_2, \sigma_3, ...)}$$
(107)

so that the constraint of Eq. 105 becomes

$$1 = \mathring{p}_{B-}(\tilde{\sigma}_{.}) + \mathring{p}_{B+}(\tilde{\sigma}_{.}) = \frac{\mathring{P}_{B}(0, \sigma_{2}, \sigma_{3}, ...) + \mathring{P}_{B}(1, \sigma_{2}, \sigma_{3}, ...)}{\mathring{P}_{B}(\sigma_{2}, \sigma_{3}, ...)}$$
(108)

in terms of the empirical probability.

As a consequence, in our present very specific case, the empirical backward kernel $\check{W}_B(\sigma_{-}|\check{\sigma}_{-})$ of Eq. 104 parametrized by the two complementary empirical probabilities $\mathring{p}_{B\pm}(\check{\sigma}_{-})$ can be rewritten in terms of the empirical empirical probability $\mathring{P}(.)$ using Eq. 107. So the application of the level 2.5 of Eq. 101 for arbitrary Markov chains actually gives, for our present very specific case, the following explicit large deviation form at level 2 for the probability of the empirical probability $\mathring{P}_B(.)$ alone

$$P_{T}^{[2]} \begin{bmatrix} \mathring{P}_{B}(.) \end{bmatrix} \xrightarrow[T \to +\infty]{} \delta \left(\sum_{\sigma_{.}} \mathring{P}_{B}(\sigma_{.}) - 1 \right) \left[\prod_{\sigma_{2}, \sigma_{3}, \ldots} \delta \left(\mathring{P}_{B}(0, \sigma_{2}, \sigma_{3}, \ldots) + \mathring{P}_{B}(1, \sigma_{2}, \sigma_{3}, \ldots) - \mathring{P}_{B}(\sigma_{2}, \sigma_{3},) \right) \right]$$

$$e^{-TI_{2}} \begin{bmatrix} \mathring{P}_{B}(.) \end{bmatrix}$$

$$(109)$$

where the constraint of Eq. 108 has to be added to the usual normalization constraint of the empirical probability, while the rate function $I_2 \begin{bmatrix} \dot{P}_B(.) \end{bmatrix}$ at level 2 obtained from Eq. 102 reads using Eqs 104 and 107

$$I_{2}\left[\mathring{P}_{B}(.)\right] = \sum_{\sigma_{-}} \left[\delta_{\sigma_{1},0} \frac{\mathring{P}_{B}(0,\sigma_{2},\sigma_{3},...)}{\mathring{P}_{B}(\sigma_{2},\sigma_{3},...)} + \delta_{\sigma_{1},1} \frac{\mathring{P}_{B}(1,\sigma_{2},\sigma_{3},...)}{\mathring{P}_{B}(\sigma_{2},\sigma_{3},...)} \right] \mathring{P}_{B}(\sigma_{2},\sigma_{3},...) \ln \left[\delta_{\sigma_{1},0} 2 \frac{\mathring{P}_{B}(0,\sigma_{2},\sigma_{3},...)}{\mathring{P}_{B}(\sigma_{2},\sigma_{3},...)} + \delta_{\sigma_{1},1} 2 \frac{\mathring{P}_{B}(1,\sigma_{2},\sigma_{3},...)}{\mathring{P}_{B}(\sigma_{2},\sigma_{3},...)} \right] \\ = \sum_{\sigma_{2},\sigma_{3},...} \left(\mathring{P}_{B}(0,\sigma_{2},\sigma_{3},...) \ln \left[2 \frac{\mathring{P}_{B}(0,\sigma_{2},\sigma_{3},...)}{\mathring{P}(\sigma_{2},\sigma_{3},...)} \right] + \left[\mathring{P}_{B}(1,\sigma_{2},\sigma_{3},...) \right] \ln \left[2 \frac{\mathring{P}_{B}(1,\sigma_{2},\sigma_{3},...)}{\mathring{P}_{B}(\sigma_{2},\sigma_{3},...)} \right] \right) \\ = \sum_{\sigma_{1},\sigma_{2},\sigma_{3},...} \mathring{P}_{B}(\sigma_{1},\sigma_{2},\sigma_{3},...) \ln \left[2 \frac{\mathring{P}_{B}(\sigma_{1},\sigma_{2},\sigma_{3},...)}{\mathring{P}_{B}(\sigma_{2},\sigma_{3},...)} \right] \right]$$
(110)

In summary, while for arbitrary Markov chains only the level 2.5 for the joint probability $P_T^{[2.5]}\left[\mathring{P}(.);\mathring{W}(.|.)\right]$ of Eq. 101 is always explicit, we have obtained that for the specific case of the backward stochastic Markov chains associated to the forward deterministic doubling map, the empirical kernel $\mathring{W}(.|.)$ is actually determined by the empirical probability $\mathring{P}(.)$ so that the explicit level 2.5 produces an explicit level 2 concerning the distribution of $P_T^{[2]}\left[\mathring{P}_B(.)\right]$ of the empirical probability $\mathring{P}_B(.)$ alone. This level reduction from 2.5 towards 2 is expected to be general for chaotic non-invertible maps, and can be considered as the counterpart of the reduction of Eq. 27 concerning trajectory observables.

VIII. CONCLUSIONS

In this paper, we have revisited the recent studies [68, 69] concerning the large deviations properties of trajectory observables for chaotic non-invertible deterministic maps in order to analyze in detail the similarities and the differences with the case of stochastic Markov chains recalled in Appendix A. To be concrete, we have focused on the simplest example of the doubling map on the real-space interval $x \in [0, 1]$ that can be also analyzed via the decomposition $x = \sum_{l=1}^{+\infty} \frac{\sigma_l}{2^l}$ into binary coefficients $\sigma_l = 0, 1$. Since the large deviations properties of trajectory observables can be studied either via deformations of the forward deterministic dynamics or via deformations of the backward stochastic dynamics, either in real space or in binary space, we have discussed the advantages and the drawbacks of these various perspectives, and we have illustrated them with the explicit example of the simplest trajectory observable $\omega(x) = x$ already considered in [68, 69]. Let us now summarize the main conclusions and discuss what happens in other deterministic chaotic systems:

(1) The k-deformation of the forward deterministic dynamics can be used to obtain the deformed dominant eigenvalue $\lambda_0^{[k]}$ governing the scaled cumulants of trajectory observables, with its corresponding left and right eigenvectors:

(1a) We have stressed that the dominant deformed left eigenvector $l_0^{[k]}(.)$ is singular in real space, since it involves at first order in perturbation theory in k the non-dominant left eigenvector $l_p^{[0]}(.)$ of the unperturbed forward dynamics, that are expected to be always singular in real space as a consequence of the sensitivity with respect to initial conditions of any deterministic chaotic dynamics.

(1b) For the doubling map, we have stressed that the corresponding dominant deformed right eigenvector $r_0^{[k]}(.)$ is regular in real space, as are the non-dominant right eigenvectors $r_p^{[0]}(.)$ of the unperturbed forward dynamics. This property should hold for other non-invertible maps, whose backward dynamics is stochastic at each time step, since this stochastic noise will erase the information on the initial condition and will separate trajectories independently of the details of their initial conditions. However for invertible maps whose backward dynamics is deterministic chaotic, both the dominant deformed right eigenvector $r_0^{[k]}(.)$ and the non-dominant undeformed right eigenvectors $r_p^{[0]}(.)$ will be be singular in real space. Note that for deterministic chaotic systems with strange attractors, even the dominant undeformed right eigenvector $r_0^{[0]}(.) = \rho_*(.)$ is already singular in real space.

(2) For chaotic non-invertible maps, we have stressed two important advantages of the backward stochastic dynamics in the symbolic space:

(2a) For each trajectory observable, an appropriate Doob canonical conditioned dynamics different from the initial dynamics can be constructed only in the backward stochastic perspective where the reweighting of existing transitions is possible, and not in the forward deterministic perspective. The corresponding conditioned steady state $\rho_*^{C[k]}(.) = l_0^{[k]}(.)r_0^{[k]}(.)$ given by the product of the dominant deformed left and right eigenvectors is always singular in real space, as a consequence of the singular character of the left eigenvector $l_0^{[k]}(.)$ any deterministic chaotic dynamics as discussed in 1a) above. As a consequence, it is more appropriate to analyze this conditioned dynamics in symbolic space, as shown on an explicit example for the doubling map where the symbolic space corresponds to infinite binary strings.

(2b) The backward stochastic dynamics in symbolic space is also the most appropriate framework to analyze large deviations at higher levels beyond the level of trajectory observables. We have explained that the application of the explicit large deviations at level 2.5 for arbitrary Markov chains to the specific case of the stochastic backward dynamics associated to forward chaotic deterministic non-invertible maps produces explicit large deviations at level 2 for the empirical density alone. In accordance with this level reduction from the level 2.5 towards the level 2, the general trajectory observables of Markov chains that involve both the empirical density and the empirical flows can be rewritten in terms of the empirical density alone, and thus reduce to the trajectory observables of the so-called Level 1, that can be obtained from the Level 2 via contraction.

Note that for chaotic invertible maps, whose backward dynamics is not stochastic but deterministic, one looses these advantages summarized in (2a) and (2b), but the symbolic dynamics is even more necessary to analyze the large deviations of trajectory observables via deformed generators, since both the dominant deformed right and left eigenvectors are singular in real space as discussed in points (1a) and (1b) above.

Appendix A: Reminder on large deviations at various levels for stochastic Markov chains

In this Appendix, we recall the large deviations properties at various levels for stochastic Markov chains, in order to compare with the main text concerning chaotic maps.

1. Analysis of the Markov chain dynamics via the spectral decomposition of its kernel $w(\tilde{x}|x)$

The evolution of the probability density $\rho_t(x)$ to be at position x at time t

$$\rho_{t+1}(\tilde{x}) = \int dx \ w(\tilde{x}|x)\rho_t(x) \tag{A1}$$

is governed by the Markov-Chain kernel $w(x, \tilde{x})$ satisfying the normalization

$$\int d\tilde{x} \ w(\tilde{x}|x) = 1 \quad \text{for any } x \tag{A2}$$

We focus on cases where the dynamics of Eq. A1 converges towards some normalizable steady-state $\rho_*(.)$ satisfying the time-independent version of Eq. A1

$$\rho_*(\tilde{x}) = \int dx \ w(\tilde{x}|x)\rho_*(x) \tag{A3}$$

The convergence towards this steady state can be analyzed via the spectral decomposition of the kernel $w(\tilde{x}|x)$

$$w(\tilde{x}|x) = \langle \tilde{x}| \left(\sum_{p=0}^{+\infty} \lambda_p |r_p\rangle \langle l_p|\right) |x\rangle = \sum_{p=0}^{+\infty} \lambda_p r_p(\tilde{x}) l_p(x)$$
(A4)

in terms of its eigenvalues λ_p with their right eigenvectors $r_p(.)$ and left eigenvectors $l_p(.)$ satisfying

$$\lambda_p r_p(\tilde{x}) = \int dx \ w(\tilde{x}|x) r_p(x)$$

$$\lambda_p l_p(x) = \int d\tilde{x} l_p(\tilde{x}) w(\tilde{x}|x)$$
(A5)

with the orthonormalization

$$\delta_{p,p'} = \langle l_p | r_{p'} \rangle = \int dx l_p(x) r_{p'}(x) \tag{A6}$$

Eqs A1 and A2 mean that the highest eigenvalue is $\lambda_{p=0} = 1$ with the corresponding positive right and left eigenvectors

$$r_0(\tilde{x}) = \rho_*(\tilde{x})$$

$$l_0(x) = 1$$
(A7)

Then the density $\rho_t(.)$ at time t can be written in terms of the initial density $\rho_{t=0}(.)$ using the spectral decomposition of Eq. A4

$$\rho_t(\tilde{x}) = \sum_{p=0}^{+\infty} \lambda_p^t r_p(\tilde{x}) \int dx_0 l_p(x_0) \rho_0(x_0)$$

= $\rho_*(\tilde{x}) + \sum_{p=1}^{+\infty} \lambda_p^t r_p(\tilde{x}) \int dx_0 l_p(x_0) \rho_0(x_0)$ (A8)

2. Large deviations of trajectory observables via appropriate deformations of Markov kernel

Let us now consider an observable $\mathcal{O}^{Traj}[x_0, x_1, ..., x_T]$ of the trajectory $(x_0, x_1, ..., x_T)$ that can be parametrized by a function $O(\tilde{x}, x)$

$$\mathcal{O}^{Traj}\left[x_0, x_1, \dots, x_T\right] = \frac{1}{T} \sum_{t=0}^{T-1} O(x_{t+1}, x_t)$$
(A9)

i.e. it is a time-average of the function $O(x_{t+1}, x_t)$ involving two consecutive positions.

a. Generating function $Z_T^{[k]}(x_T|x_0)$ of the trajectory observable via via the appropriate deformed kernel $w^{[k]}(\tilde{x}|x)$

The standard method to analyze the statistics of the trajectory observable of Eq. A9 is based on the generating function of \mathcal{O}^{Traj} over the trajectories starting at $x(t=0) = x_0$ and ending at $x(t=T) = x_T$

$$Z_{T}^{[k]}(x_{T}|x_{0}) \equiv \langle e^{kT\mathcal{O}^{Traj}} \rangle = \int dx_{1} \int dx_{2} \dots \int dx_{T-1} \prod_{t=0}^{T-1} \left[w(x_{t+1}, x_{t}) e^{kO(x_{t+1}, x_{t})} \right]$$
$$\equiv \int dx_{1} \int dx_{2} \dots \int dx_{T-1} \prod_{t=0}^{T-1} w^{[k]}(x_{t+1}, x_{t}) \equiv \langle x_{T} | \left(w^{[k]} \right)^{T} | x_{0} \rangle$$
(A10)

that is governed by the k-deformed kernel

$$w^{[k]}(\tilde{x}|x) \equiv w(\tilde{x}|x)e^{kO(\tilde{x},x)}$$
(A11)

b. Eigenvalue problem for $w^{[k]}(\tilde{x}|x)$ to obtain the scaled-cumulant generating function of the trajectory observable

For large time $T \to +\infty$, the generating function of Eq. A10 is governed

$$Z_T^{[k]}(x_T|x_0) = \langle x_T | \left(w^{[k]} \right)^T | x_0 \rangle \underset{T \to +\infty}{\simeq} [\lambda_0^{[k]}]^T r_0^{[k]}(x_T) l_0^{[k]}(x_0)$$
(A12)

by the highest eigenvalue $\lambda_0^{[k]}$ associated to the positive right eigenvector $r_0^{[k]}(\tilde{x}) \ge 0$ and to the positive left eigenvector $l_0^{[k]}(\tilde{x}) \ge 0$ satisfying

$$\lambda_{0}^{[k]} r_{0}^{[k]}(\tilde{x}) = \int dx \ w^{[k]}(\tilde{x}|x) r_{0}^{[k]}(x) = \int dx \ w(\tilde{x}|x) e^{kO(\tilde{x},x)} r_{0}^{[k]}(x)$$

$$\lambda_{0}^{[k]} l_{0}^{[k]}(x) = \int d\tilde{x} l_{0}^{[k]}(\tilde{x}) w^{[k]}(\tilde{x}|x) = \int d\tilde{x} l_{0}^{[k]}(\tilde{x}) w(\tilde{x}|x) e^{kO(\tilde{x},x)}$$
(A13)

with the normalization

$$\int dx l_0^{[k]}(x) r_0^{[k]}(x) = 1 \tag{A14}$$

while for k = 0 one recovers the eigenvectors of Eq A7 associated the highest eigenvalue $\lambda_0^{[k=0]} = 1$ with its eigenvectors of Eq. A7.

From the generating function $Z_T^{[k]}(x_T|x_0)$ that will grow or decay in time as $[\lambda_0^{[k]}]^T$, there are two simple ways to construct processes that conserve the probability as we now recall.

c. Construction of the normalized density
$$\rho_T^{[k]End}(x|x_0) = \frac{Z_T^{[k]}(x|x_0)}{\int dy Z_T^{[k]}(y|x_0)}$$
 for the end-point x

The simplest way to construct a normalized probability out of the generation function $Z_T^{[k]}(x|x_0)$ is to consider the probability of the end-point

$$\rho_T^{[k]End}(x|x_0) \equiv \frac{Z_T^{[k]}(x|x_0)}{\int dy Z_T^{[k]}(y|x_0)} \tag{A15}$$

The asymptotic behavior of Eq. A12 for the generating function yields that the probability of the end point x

$$\rho_T^{[k]End}(x|x_0) \underset{T \to +\infty}{\simeq} \frac{r_0^{[k]}(x)}{\int dy r_0^{[k]}(y)} \equiv \rho_*^{[k]End}(x)$$
(A16)

converges, independently of the initial point x_0 towards the distribution $\rho_*^{[k]End}(x)$ determined by the appropriate normalization right eigenvector $r_0^{[k]}(.)$.

d. Construction of the conditioned kernel $w^{C[k]}(\tilde{x}|x)$ via the Doob's transform involving the left eigenvector $l_0^{[k]}(.)$

Another interesting way to construct a normalized density out of the generation function $Z_T^{[k]}(x|x_0)$ is via the Doob's conditioning similarity transform involving the left eigenvector $l_0^{[k]}(.)$ of Eq. Eq. A13

$$\rho_T^{C[k]}(x|x_0) \equiv \frac{1}{[\lambda_0^{[k]}]^T} l_0^{[k]}(x) Z_T^{[k]}(x|x_0) \frac{1}{l_0^{[k]}(x_0)} = \frac{1}{[\lambda_0^{[k]}]^T} l_0^{[k]}(x) \langle x| \left(w^{[k]}\right)^T |x_0\rangle \frac{1}{l_0^{[k]}(x_0)} \\ \equiv \langle x| \left(w^{C[k]}\right)^T |x_0\rangle \tag{A17}$$

where the conditioned kernel

$$w^{C[k]}(\tilde{x}|x) \equiv \frac{1}{\lambda_0^{[k]}} l_0^{[k]}(\tilde{x}) w^{[k]}(\tilde{x}|x) \frac{1}{l_0^{[k]}(x)}$$
$$= \frac{l_0^{[k]}(\tilde{x}) w^{[k]}(\tilde{x}|x)}{\int dz l_0^{[k]}(z) w^{[k]}(z|x)} = \frac{l_0^{[k]}(\tilde{x}) w(\tilde{x}|x) e^{kO(\tilde{x},x)}}{\int dz l_0^{[k]}(z) w(z|x) e^{kO(z,x)}}$$
(A18)

has been rewritten using the eigenvalue Eq. A13 for the left eigenvector $l_0^{[k]}(x)$ in the denominator to obtain the last expression that makes obvious the normalization over \tilde{x} for any x. The physical meaning of Eq. A18 is that the transitions $x \to \tilde{x}$ that were possible with the undeformed kernel $w(\tilde{x}|x) \neq 0$ are reweighted, while the transitions $x \to \tilde{x}$ that were not possible with the undeformed kernel $w(\tilde{x}|x) = 0$ are still impossible.

Plugging the asymptotic behavior of Eq. A12 for the generating function into Eq. A17 yields that for large T, the density $\rho_T^{C[k]}(x|x_0)$

$$\rho_T^{C[k]}(x|x_0) \equiv \frac{1}{[\lambda_0^{[k]}]^T} l_0^{[k]}(x) Z_T^{[k]}(x|x_0) \frac{1}{l_0^{[k]}(x_0)} \underset{T \to +\infty}{\simeq} l_0^{[k]}(x) r_0^{[k]}(x) \equiv \rho_*^{C[k]}(x)$$
(A19)

converges, independently of the initial point x_0 towards the conditioned steady distribution $\rho_*^{C[k]}(x)$ given by the product of the left eigenvector $l_0^{[k]}(.)$ and of the right eigenvector $r_0^{[k]}(.)$, whose normalization over x is ensured by Eq. A14.

In order to better understand the different physical meaning with respect to the end-point distribution of Eqs A15 and A16, it is useful to consider the density of position x at some interior time $0 \ll t \ll T$ using the asymptotic expression of Eq. A12 for the three generating functions on the time-intervals [0, t], [t, T] and [0, T]

$$\rho_t^{[k]Interior}(x) \equiv \frac{Z_{T-t}^{[k]}(x_T|x)Z_t^{[k]}(x|x_0)}{Z_T^{[k]}(x_T|x_0)} \\
\sum_{T \to +\infty} \frac{[\lambda_0^{[k]}]^{T-t}r_0^{[k]}(x_T)l_0^{[k]}(x)[\lambda_0^{[k]}]^t r_0^{[k]}(x)l_0^{[k]}(x_0)}{[\lambda_0^{[k]}]^T r_0^{[k]}(x_T)l_0^{[k]}(x_0)} = l_0^{[k]}(x)r_0^{[k]}(x) \equiv \rho_*^{C[k]}(x)$$
(A20)

So the conditioned steady state $\rho_*^{C[k]}(x)$ corresponds to the distribution of the position x seen at interior times $0 \ll t \ll T$ independently of the initial position x_0 and of the final position x_T .

e. Physical meaning of the largest deformed eigenvalue $\lambda_0^{[k]}$

The dominant behavior of Eq. A12 that is independent of the two positions $(x_0; x_T)$

$$Z_T^{[k]} \underset{T \to +\infty}{\simeq} [\lambda_0^{[k]}]^T \equiv e^{Tc(k)} \quad \text{with} \ c(k) \equiv \ln \lambda_0^{[k]}$$
(A21)

means that $c(k) \equiv \ln \lambda_0^{[k]}$ represents the generating function of the scaled-cumulants of the trajectory observable $\mathcal{O}^{Traj}[x_0, x_1, ..., x_T]$ of Eq. A9 (see the review [4] for more details).

The link with the rate function $I(\mathcal{O})$ governing the large deviations for large T of the probability $P_T(\mathcal{O})$ to see the value \mathcal{O} for the trajectory observable $\mathcal{O}^{Traj}[x_0, x_1, ..., x_T]$ of Eq. A9

$$P_T(\mathcal{O}) \underset{T \to +\infty}{\simeq} e^{-TI(\mathcal{O})}$$
(A22)

can be understood via the computation of the generating function of Eqs A10 and A21 via the saddle-point evaluation for large T

$$Z_T^{[k]} \equiv \langle e^{kT\mathcal{O}} \rangle = \int d\mathcal{O}P_T(\mathcal{O})e^{kT\mathcal{O}}$$
$$\underset{T \to +\infty}{\simeq} \int d\mathcal{O}e^{T} \left[k\mathcal{O} - I(\mathcal{O})\right] \underset{T \to +\infty}{\simeq} e^{Tc(k)}$$
(A23)

that leads to the Legendre transformation

$$k\mathcal{O} - I(\mathcal{O}) = c(k)$$

$$k - I'(\mathcal{O}) = 0$$
(A24)

with its inverse

$$I(\mathcal{O}) = k\mathcal{O} - c(k)$$

$$0 = \mathcal{O} - c'(k)$$
(A25)

In summary, the largest deformed eigenvalue $\lambda_0^{[k]}$ gives directly the scaled cumulant generating function c(k) = $\left[\ln \lambda_0^{[k]}\right]$, whose Legendre transform is the rate function $I(\mathcal{O})$ governing the large deviations of Eq. A22 for the trajectory observable.

Perturbation theory in k of the eigenvalue $\lambda_0^{[k]}$ and for the eigenvectors f.

When it is not possible to solve exactly the eigenvalue equations of Eq. A13 for the deformed kernel $w^{[k]}(\tilde{x}|x)$, it is useful to consider the perturbation theory with respect to the undeformed kernel $w^{[k=0]}(\tilde{x}|x) = w(\tilde{x}|x)$ where the eigenvalue $\lambda_0^{[k=0]} = 1$ is known with its eigenvectors of Eq. A7. The series expansion in k for the deformed kernel of Eq. A11

$$w^{[k]}(\tilde{x}|x) = w(\tilde{x}|x)e^{kO(\tilde{x},x)} = w(\tilde{x}|x) + kw^{(1)}(\tilde{x}|x) + k^2w^{(2)}(\tilde{x}|x) + o(k)$$
(A26)

involves the first-order and second-order corrections

$$w^{(1)}(\tilde{x}|x) = w(\tilde{x}|x)O(\tilde{x},x) w^{(2)}(\tilde{x}|x) = w(\tilde{x}|x)\frac{O^{2}(\tilde{x},x)}{2}$$
(A27)

The perturbation theory for the highest eigenvalue

$$\lambda_0^{[k]} = 1 + k\lambda^{(1)} + k^2\lambda^{(2)} + o(k) \tag{A28}$$

is very similar to the standard theory of quantum mechanics with the following conclusions.

The fist-order correction $\lambda^{(1)}$ only involves the first-order perturbation $w^{(1)}(\tilde{x}|x)$ and the unperturbed eigenvectors of Eq. A7

$$\lambda^{(1)} = \langle l_0 | w^{(1)} | r_0 \rangle = \int d\tilde{x} \int dx l_0(\tilde{x}) w^{(1)}(\tilde{x} | x) r_0(x) = \int dx \int d\tilde{x} w(\tilde{x} | x) O(\tilde{x}, x) \rho_*(x)$$
(A29)

and coincides with the steady value of the trajectory observable that can be computed in terms of the steady state $\rho_*(x).$

The first-order corrections for the right and left eigenvectors with respect to their unperturbed values of Eqs A7

$$r_0^{[k]}(x) = r_0(x) + kr_0^{(1)}(x) + o(k) = \rho_*(x) + kr_0^{(1)}(x) + o(k)$$

$$l_0^{[k]}(x) = l_0(x) + kl_0^{(1)}(x) + o(k) = 1 + kl_0^{(1)}(x) + o(k)$$
(A30)

can be written

in terms of the Green function

$$G \equiv (\mathbb{1} - |r_0\rangle\langle l_0|) \frac{1}{\mathbb{1} - w} \left(\mathbb{1} - |r_0\rangle\langle l_0|\right) = \sum_{p=1}^{+\infty} \frac{|r_p\rangle\langle l_p|}{1 - \lambda_p}$$
(A32)

The first expression means that the Green function G is the inverse of the operator $(\mathbb{1}-w)$ in the subspace $(\mathbb{1}-|r_0\rangle\langle l_0|)$ orthogonal to the subspace $|r_0\rangle\langle l_0|$ associated to the unity eigenvalue $\lambda_0 = 1$ The second expression involves all the unperturbed other eigenvalues $\lambda_{p\geq 1}$ with their right and left eigenvectors that were discussed for the spectral decomposition of Eq. A8.

So the first-order corrections of Eqs A31

$$r_{0}^{(1)}(\tilde{x}) = \int dz \int dx G(\tilde{x}, z) w^{(1)}(z|x) r_{0}(x) = \sum_{p=1}^{+\infty} r_{p}(\tilde{x}) \left[\frac{1}{1-\lambda_{p}} \int dx \int dz l_{p}(z) w(z|x) O(z, x) \rho_{*}(x) \right]$$

$$l_{0}^{(1)}(z) = \int dx \int d\tilde{x} l_{0}(x) w^{(1)}(x|\tilde{x}) G(\tilde{x}, z) = \sum_{p=1}^{+\infty} l_{p}(z) \left[\frac{1}{1-\lambda_{p}} \int d\tilde{x} \int dx l_{0}(x) w(x|\tilde{x}) O(x, \tilde{x}) r_{p}(\tilde{x}) \right]$$
(A33)

are decomposed onto all the other unperturbed right eigenvectors $r_p(\tilde{x})$ and on all the other unperturbed left eigenvectors $l_p(z)$ respectively.

The first-order correction for the conditioned density of Eq. A20

$$\rho_*^{C[k]}(\tilde{x}) = l_0^{[k]}(\tilde{x}) r_0^{[k]}(\tilde{x}) = \rho_*(\tilde{x}) + k \rho_*^{C(1)}(\tilde{x}) + o(k)$$
(A34)

is given by

$$\rho_*^{C(1)}(\tilde{x}) = l_0(\tilde{x})r_0^{(1)}(\tilde{x}) + l_0^{(1)}(\tilde{x})r_0(\tilde{x}) = r_0^{(1)}(\tilde{x}) + l_0^{(1)}(\tilde{x})\rho_*(\tilde{x})$$
(A35)

so that it involves both corrections $r_0^{(1)}(\tilde{x})$ and $l_0^{(1)}(\tilde{x})$ of Eq. A33. The second-order correction $\lambda^{(2)}$ also involves the Green function of Eq. A32

$$\begin{aligned} \lambda^{(2)} &= \langle l_0 | w^{(2)} | r_0 \rangle + \langle l_0 | w^{(1)} G w^{(1)} | r_0 \rangle \\ &= \int d\tilde{x} \int dx l_0(\tilde{x}) w^{(2)}(\tilde{x} | x) r_0(x) + \int d\tilde{x}_2 \int dx_2 \int d\tilde{x}_1 \int dx_1 l_0(\tilde{x}_2) w^{(1)}(\tilde{x}_2, x_2) w^{(1)}(\tilde{x}_1 | x_1) r_0(x_1) \\ &= \int d\tilde{x} \int dx w(\tilde{x} | x) \frac{O^2(\tilde{x}, x)}{2} \rho_*(x) + \int d\tilde{x}_2 \int dx_2 \int d\tilde{x}_1 \int dx_1 w(\tilde{x}_2, x_2) O(\tilde{x}_2, x_2) W(\tilde{x}_1 | x_1) O(\tilde{x}_1, x_1) \rho_*(x_1) \\ \end{aligned}$$

3. Large deviations at higher levels

Explicit large deviations at level 2.5 for the empirical density $\mathring{\rho}(.)$ and the empirical kernel $\mathring{w}(.,.)$ a.

If one observes a trajectory $(x_0, x_1, ..., x_T)$ over a long-time T, it is interesting to consider (i) the empirical 1-point density

$$\mathring{\rho}(x) \equiv \frac{1}{T} \sum_{t=0}^{T-1} \delta(x_t - x)$$
(A37)

satisfying the normalization

$$\int dx \dot{\rho}(x) = 1 \tag{A38}$$

(ii) the empirical kernel

$$\dot{w}(\tilde{x},x) = \frac{\frac{1}{T} \sum_{t=0}^{T-1} \delta(x_{t+1} - \tilde{x}) \delta(x_t - x)}{\frac{1}{T} \sum_{t=0}^{T-1} \delta(x_t - x)} = \frac{\frac{1}{T} \sum_{t=0}^{T-1} \delta(x_{t+1} - \tilde{x}) \delta(x_t - x)}{\dot{\rho}(x)}$$
(A39)

satisfying the normalization

$$\int d\tilde{x} \dot{w}(\tilde{x}, x) \equiv \frac{\frac{1}{T} \sum_{t=0}^{T-1} \delta(x_t - x)}{\dot{\rho}(x)} = 1 \quad \text{for any } x$$
(A40)

and having the empirical density $\dot{\rho}(.)$ of Eq. A37 as steady distribution

$$\int dx \mathring{w}(\tilde{x}, x) \mathring{\rho}(x) = \frac{1}{T} \sum_{t=0}^{T-1} \delta(x_{t+1} - \tilde{x}) = \mathring{\rho}(\tilde{x}) + \frac{\delta(x_T - \tilde{x}) - \delta(x_0 - \tilde{x})}{T} = \mathring{\rho}(\tilde{x}) + O\left(\frac{1}{T}\right)$$
(A41)

up to boundary terms of order 1/T that are negligible for large T.

The joint distribution $P_T^{[2.5]}[\dot{\rho}(.); \dot{w}(.,.)]$ of the empirical density $\dot{\rho}(.)$ of Eq. A37 and of the empirical kernel $\dot{w}(.,.)$ of Eq. A39 displays the large deviation form for large T [4, 14, 70–75]

$$P_T^{[2.5]}\left[\mathring{\rho}(.); \mathring{w}(.,.)\right] \xrightarrow[T \to +\infty]{} \delta\left(\int dx \mathring{\rho}(x) - 1\right) \left[\prod_x \delta\left(\int d\tilde{x} \mathring{w}(\tilde{x}, x) - 1\right)\right] \left[\prod_{\tilde{x}} \delta\left(\int dx \mathring{w}(\tilde{x}, x) \mathring{\rho}(x) - \mathring{\rho}(\tilde{x})\right)\right] e^{-TI_{2.5}\left[\mathring{\rho}(.); \mathring{w}(.,.)\right]}$$
(A42)

where the factors in front of the exponential correspond to the constitutive constraints for the empirical observables discussed in Eqs A38 A40 A41, while the rate function $I_{2.5} [\dot{\rho}(.); \dot{w}(.,.)]$ at level 2.5 appearing in the exponential has for explicit expression

$$I_{2.5}[\mathring{\rho}(.); \mathring{w}(.,.)] = \int dx \mathring{\rho}(x) \int d\tilde{x} \mathring{w}(\tilde{x}, x) \ln\left(\frac{\mathring{w}(\tilde{x}, x)}{w(\tilde{x}, x)}\right)$$
(A43)

This level 2.5 for large deviations plays an essential role since it is the lowest level that is explicit for an arbitrary Markov chain. Indeed, all the lower levels can be obtained via optimization of this level 2.5 in the presence of constraints, but the solution of these constrained optimizations cannot be written explicitly in general, as recalled below.

b. Large deviations at level 2 for the empirical density $\mathring{\rho}(.)$ alone

The level 2 concerning the distribution of the empirical density $\mathring{\rho}(.)$ alone can be obtained from the integration over Eq. A42 over the empirical kernel $\mathring{w}(.,.)$

$$P_{T}^{[2]}[\mathring{\rho}(.)] \equiv \int \mathcal{D}\mathring{w}(.,.)(.,.)P_{T}^{[2.5]}[\mathring{\rho}(.);\mathring{w}(.,.)]$$

$$\underset{T \to +\infty}{\simeq} \delta\left(\int dx\mathring{\rho}(x) - 1\right) \int \mathcal{D}\mathring{w}(.,.)(.,.) \left[\prod_{x} \delta\left(\int d\tilde{x}\mathring{w}(\tilde{x},x) - 1\right)\right] \left[\prod_{y} \delta\left(\int dx\mathring{w}(\tilde{x},x)\mathring{\rho}(x) - \mathring{\rho}(\tilde{x})\right)\right] e^{-TI_{2.5}}[\mathring{\rho}(.);\mathring{w}(.,.)]$$

$$\underset{T \to +\infty}{\simeq} \delta\left(\int dx\mathring{\rho}(x) - 1\right) e^{-TI_{2}}[\mathring{\rho}(.)]$$
(A44)

So the rate function $I_2[\dot{\rho}(.)]$ at level 2 for the empirical density alone corresponds to the optimization of the rate function $I_{2.5}[\dot{\rho}(.); \dot{w}(.,.)]$ at level 2.5 over the empirical kernel $\dot{w}(.,.)$ satisfying the appropriate constraints, but cannot be written as an explicit function of $\dot{\rho}(.)$ in general, since the solution of this constrained optimization is not explicit for an arbitrary Markov chain.

c. Link with the large deviations of trajectory observables

The trajectory observable $\mathcal{O}^{Traj}[x_0, x_1, ..., x_T]$ of Eq. A9 can be rewritten as a function $\mathcal{O}[\dot{\rho}(.); \dot{w}(.,.)]$ of the empirical density $\dot{\rho}(.)$ of Eq. A37 and of the empirical kernel $\dot{w}(.,.)$ of Eq. A39

$$\mathcal{O}^{Traj}\left[x_0, x_1, ..., x_T\right] = \int d\tilde{x} \int dx \Omega(\tilde{x}, x) \mathring{w}(\tilde{x}, x) \mathring{\rho}(x) \equiv \mathcal{O}\left[\mathring{\rho}(.); \mathring{w}(., .)\right]$$
(A45)

As a consequence, the probability distribution $P_T(\mathcal{O})$ of the trajectory observable already discussed in Eq. A22 can be alternatively evaluated from the probability $P_T^{[2.5]}[\dot{\rho}(.); \dot{w}(.,.)]$ at the Level 2.5 of Eq. A42 via the integral

$$P_{T}(\mathcal{O}) = \int \mathcal{D}\mathring{\rho}(.,.)(.) \int \mathcal{D}\mathring{w}(.,.)(.,.)P_{T}^{[2.5]}[\mathring{\rho}(.);\mathring{w}(.,.)] \,\delta\left(\mathcal{O} - \mathcal{O}\left[\mathring{\rho}(.);\mathring{w}(.,.)\right]\right) \\ \xrightarrow{\simeq}_{T \to +\infty} \int \mathcal{D}\mathring{\rho}(.,.)(.) \int \mathcal{D}\mathring{w}(.,.)(.,.)\delta\left(\int dx\mathring{\rho}(x) - 1\right) \left[\prod_{x} \delta\left(\int d\tilde{x}\mathring{w}(\tilde{x},x) - 1\right)\right] \left[\prod_{y} \delta\left(\int dx\mathring{w}(\tilde{x},x)\mathring{\rho}(x) - \mathring{\rho}(\tilde{x})\right)\right] \\ \delta\left(\mathcal{O} - \mathcal{O}\left[\mathring{\rho}(.);\mathring{w}(.,.)\right]\right) e^{-TI_{2.5}\left[\mathring{\rho}(.);\mathring{w}(.,.)\right]} \xrightarrow{\simeq}_{T \to +\infty} e^{-TI(\mathcal{O})}$$
(A46)

The daddle-point evaluation of the integral for large T means that the rate function $I(\mathcal{O})$ for the trajectory observable \mathcal{O} already discussed around Eq. A24 also corresponds to the optimization of the explicit rate function $I^{[2.5]}[\dot{\rho}(.); \dot{w}(.,.)]$ over the empirical density $\dot{\rho}(.)$ and the empirical kernel $\dot{w}(.,.)$ satisfying their constitutive constraints, as well as the supplementary constraint $\delta(\mathcal{O} - \mathcal{O}[\dot{\rho}(.); \dot{w}(.,.)])$ reproducing the appropriate value \mathcal{O} of the trajectory observable. Again, the solution of this constrained optimization is not explicit for an arbitrary observable of an arbitrary Markov chain, so the rate function $I(\mathcal{O})$ can be computed explicitly only in special cases.

Appendix B: Alternative analysis of the large deviations and conditioning via the backward Markov chain

In this Appendix, we describe how the large deviations and conditioning analyzed via the forward Markov chain in the previous Appendix A can be alternatively studied from the point of view of the backward Markov chain, since this change of perspective is very useful in the main text concerning chaotic non-invertible maps.

1. Backward Markov chain associated to the forward Markov chain

When the initial condition x_0 is drawn with the steady state distribution $\rho_*(x_0)$, the probability $\mathcal{P}^{Forward}_*(x_{0\leq t\leq T})$ of the forward trajectory $(x_0, x_1, ..., x_T)$ over the time-window [0, T] reads in terms of the forward-kernel w(.|.)

$$\mathcal{P}_T^{Traj}(x_0, x_1, ..., x_T) = \rho_*(x_0) \prod_{t=0}^{T-1} w(x_{t+1}|x_t)$$
(B1)

For T = 1, the joint probability of the two consecutive positions (x_0, x_1) can be also rewritten as

$$\mathcal{P}_{T=1}^{Traj}(x_0, x_1) = w(x_1|x_0)\rho_*(x_0) \equiv w_B(x_0|x_1)\rho_*(x_1)$$
(B2)

where the backward-kernel $w^B(x|\tilde{x})$ represents the probability to see the previous point x when the next point is \tilde{x} and can be obtained from the forward kernel $w(\tilde{x}|x)$ via the similarity transformation involving the steady state ρ_*

$$w_B(x|\tilde{x}) = \frac{1}{\rho_*(\tilde{x})} w(\tilde{x}|x)\rho_*(x)$$
(B3)

The spectral decomposition of Eq. A4 of the forward kernel $w(\tilde{x}|x)$ yields that the spectral decomposition of the backward kernel

$$w_B(x|\tilde{x}) = \sum_{p=0}^{+\infty} \lambda_p \left[\rho_*(x)l_p(x)\right] \left[\frac{r_p(\tilde{x})}{\rho_*(\tilde{x})}\right] \equiv \sum_{p=0}^{+\infty} \lambda_p r_p^B(x)l_p^B(\tilde{x})$$
(B4)

involves the same eigenvalues λ_p as the forward kernel $w(\tilde{x}|x)$, while the right and left eigenvectors for the backward kernel read

$$r_p^B(x) = l_p(x)\rho_*(x)$$

$$l_p^B(\tilde{x}) = \frac{r_p(\tilde{x})}{\rho_*(\tilde{x})}$$
(B5)

In particular, for the highest eigenvalue $\lambda_{p=0} = 1$, plugging Eq. A7 into Eq. B5 yields

$$r_0^B(\tilde{x}) = \rho_*(\tilde{x})$$

$$l_0^B(x) = 1$$
(B6)

This means that the backward-kernel of Eq. B3 is normalized over x for any \tilde{x} and that the backward kernel of Eq. B3 has the same steady state $\rho_*(.)$ as the forward kernel w(.|.)

$$\rho_*(x) = \int d\tilde{x} w_B(x|\tilde{x}) \rho_*(\tilde{x}) \tag{B7}$$

The backward-kernel $w^B(x|\tilde{x})$ can be used to construct the trajectory going backward since the trajectory probability of Eq. B1 can be rewritten as

$$\mathcal{P}_{T}^{Traj}(x_{0}, x_{1}, ..., x_{T}) = \rho_{*}(x_{0}) \prod_{t=0}^{T-1} w(x_{t+1}|x_{t}) = \rho_{*}(x_{0}) \prod_{t=0}^{T-1} \left[w_{B}(x_{t}|x_{t+1}) \frac{\rho_{*}(x_{t+1})}{\rho_{*}(x_{t})} \right]$$
$$= \rho_{*}(x_{T}) \prod_{t=0}^{T-1} w_{B}(x_{t}|x_{t+1})$$
(B8)

When considering the backward trajectory, it is thus useful to replace the forward-time t = 0, 1, ..., T by the backward-time

$$\tau = T - t = 0, 1, .., T \tag{B9}$$

and to relabel the positions $x_t = y_{(\tau=T-t)}$: then the backward time $\tau = 0, 1, ..., T$ grows from the minimal value $\tau = 0$ associated the initial position $x_T = y_0$ of the backward trajectory towards the maximal value $\tau = T$ associated the final position $x_0 = y_T$ of the backward trajectory.

2. Large deviations of trajectory observables via the deformed backward dynamics

a. Generating function via the k-deformed backward kernel $w_B^{[k]}(x|\tilde{x})$

The k-deformed kernel of Eq. A11 associated to the backward kernel $w_B(x|\tilde{x})$ of Eq. B3

$$w_B^{[k]}(x|\tilde{x}) \equiv w_B(x|\tilde{x})e^{kO(\tilde{x},x)} = w(\tilde{x}|x)\frac{\rho_*(x)}{\rho_*(\tilde{x})}e^{kO(\tilde{x},x)} = \frac{1}{\rho_*(\tilde{x})}w^{[k]}(\tilde{x}|x)\rho_*(x)$$
(B10)

This similarity transformation between the deformed forward kernel $w_B^{[k]}(\tilde{x}|x)$ and the deformed backward kernel $w_B^{[k]}(x|\tilde{x})$ yields that their highest eigenvalue $\lambda_0^{[k]}$ is the same, while the positive right and left eigenvectors of the backward kernel satisfying

$$\lambda_{0}^{[k]} r_{0}^{B[k]}(x) = \int d\tilde{x} w_{B}^{[k]}(x|\tilde{x}) r_{0}^{B[k]}(\tilde{x})$$

$$\lambda_{0}^{[k]} l_{0}^{B[k]}(\tilde{x}) = \int dx l_{0}^{B[k]}(x) w_{B}^{[k]}(x|\tilde{x})$$
(B11)

are related to the forward eigenvectors via

$$r_0^{B[k]}(\tilde{x}) = l_0^{[k]}(\tilde{x})\rho_*(\tilde{x})$$

$$l_0^{B[k]}(x) = \frac{r_0^{[k]}(x)}{\rho_*(x)}$$
(B12)

b. Construction of the backward conditioned steady state $\rho_*^{BC[k]}(x)$ and of the backward conditioned kernel $w^{BC[k]}(\tilde{x}|x)$

Using Eq. B12, one obtains that the backward conditioned steady state $\rho_*^{BC[k]}(x)$ coincides with the forward conditioned steady state $\rho_*^{C[k]}(x)$ of Eq. A20

$$\rho_*^{BC[k]}(x) \equiv l_0^{B[k]}(x) r_0^{B[k]}(x) = l_0^{[k]}(x) r_0^{[k]}(x) \equiv \rho_*^{C[k]}(x)$$
(B13)

The backward conditioned kernel $w_B^{C[k]}(x|\tilde{x})$ constructed via the analog to Eq. A18

$$w_B^{C[k]}(x|\tilde{x}) \equiv \frac{1}{\lambda_0^{[k]}} l_0^{B[k]}(x) w^{B[k]}(x|\tilde{x}) \frac{1}{l_0^{B[k]}(\tilde{x})} = \frac{l_0^{B[k]}(x) w^{B[k]}(x|\tilde{x})}{\int dz l_0^{B[k]}(z) w^{B[k]}(z|\tilde{x})} = \frac{l_0^{B[k]}(x) w_B(x|\tilde{x}) e^{kO(\tilde{x},x)}}{\int dz l_0^{B[k]}(z) w(z|\tilde{x}) e^{kO(\tilde{x},z)}}$$
(B14)

is related to the conditioned forward kernel $w^{C[k]}(\tilde{x}|x)$ of Eq. A18 via the similarity transformation involving common conditioned steady density of Eq. B13

$$w_B^{C[k]}(x|\tilde{x})\rho_*^{C[k]}(\tilde{x}) = w^{C[k]}(\tilde{x}|x)\rho_*^{C[k]}(x)$$
(B15)

This equation represents the conditioned probability of two consecutive positions as computed from the backward or forward perspectives, and is the analog of Eq. B2 for the initial dynamics.

Appendix C: Doubling map : Properties of the forward and backward dynamics for the Fourier coefficients

In order to better understand the forward and backward dynamics that were discussed for the density $\rho_{\cdot}(.)$ on the real-space interval $x \in [0, 1[$ in sections II and IV of the main text, it is useful in the present Appendix to describe these dynamics from the point of view of the Fourier coefficients of the density $\rho_{\cdot}(.)$ (see [95] for more details).

1. Forward dynamics for the Fourier coefficients $\hat{\rho}_t(n) = \int_0^1 d\tilde{x} \rho_t(x) e^{-i2\pi nx}$ of the density $\rho_t(x)$

a. General properties of the Fourier coefficients $\hat{\rho}_t(n) = \int_0^1 d\tilde{x} \rho_t(x) e^{-i2\pi nx}$ with $n \in \mathbf{Z}$

The Fourier series representation of the density $\rho_t(x)$

$$\rho_t(x) = \sum_{n=-\infty}^{+\infty} \hat{\rho}_t(n) e^{i2\pi nx}$$
(C1)

involves the Fourier coefficients

$$\hat{\rho}_t(n) = \int_0^1 dx \rho_t(x) e^{-i2\pi nx}$$
(C2)

that satisfy the complex-conjugate relations (since the density is real $\rho_t(\tilde{x}) = \overline{\rho_t(\tilde{x})}$)

$$\hat{\rho}_t(-n) = \int_0^1 dx \rho_t(x) e^{i2\pi nx} = \overline{\hat{\rho}_t(n)}$$
(C3)

while the coefficient for n = 0 is fixed by the normalization

$$\hat{\rho}_t(n=0) = \int_0^1 dx \rho_t(x) = 1$$
(C4)

b. Frobenius-Perron dynamics for the Fourier coefficients $\hat{\rho}_t(n)$ with $n \in \mathbb{Z}$

The Frobenius-Perron evolution of Eq. 5 translates for the Fourier coefficients into

$$\hat{\rho}_t(\tilde{n}) = \sum_{n=-\infty}^{+\infty} \hat{W}(\tilde{n}|n)\hat{\rho}_{t-1}(n)$$
(C5)

where the matrix $\hat{W}(\tilde{n}|n)$ can be computed from the real-space kernel $w(\tilde{x}|x)$ of Eq. 4

$$\hat{W}(\tilde{n}|n) \equiv \int_{0}^{1} dx e^{i2\pi nx} \int_{0}^{1} d\tilde{x} e^{-i2\pi \tilde{n}\tilde{x}} w(\tilde{x}|x) = \delta_{2\tilde{n},n}$$
(C6)

So the dynamics of Eq. C5 reduces to

$$\hat{\rho}_t(\tilde{n}) = \sum_{n = -\infty}^{+\infty} \delta_{2\tilde{n}, n} \hat{\rho}_{t-1}(n) = \hat{\rho}_{t-1}(2\tilde{n})$$
(C7)

and corresponds to the Fourier translation of the real-space evolution of Eq. 5 : all odd Fourier coefficients $\hat{\rho}_{t-1}(n = 2m + 1)$ of the density at time (t - 1) disappear, while the even Fourier coefficients $\hat{\rho}_{t-1}(2\tilde{n})$ become $\hat{\rho}_t(\tilde{n})$. The iteration of Eq. C7 up to the initial condition at t = 0

$$\hat{\rho}_t(n) = \hat{\rho}_{t-1}(2n) = \hat{\rho}_{t-2}(4n) = \dots = \hat{\rho}_0(2^t n) \tag{C8}$$

shows that only the Fourier coefficients $\hat{\rho}_0(n_0)$ of the initial density with indices $n_0 = 2^t n$ have not yet disappeared at time t. If the Fourier coefficients $\hat{\rho}_0(n_0)$ of the initial condition decay to zero for large n_0 , then only the Fourier coefficient n = 0 survives for large time $t \to +\infty$

$$\hat{\rho}_t(n) \underset{t \to +\infty}{\simeq} \hat{\rho}_*(n) = \delta_{n=0} \tag{C9}$$

in correspondence with the uniform real-space steady state of Eq. 7.

In Fourier space, the right eigenvectors $\hat{r}_p(n)$ associated to the eigenvalues $\lambda_p = 2^{-p}$ of Eq. 10 satisfy

$$\lambda_p \ \hat{r}_p(n) = \hat{r}_p(2n) \tag{C10}$$

and are given by the Fourier coefficients of the Bernoulli polynomials $r_p(x) = B_p(x)$ of Eq. 12

$$\hat{r}_p(n) = \hat{B}_p(n) = -\frac{p!}{(i2\pi n)^p} \quad \text{for} \ n \neq 0$$
 (C11)

with the first members

$$\hat{r}_1(n) = \hat{B}_1(n) = -\frac{1}{i2\pi n} \quad \text{for} \quad n \neq 0$$
$$\hat{r}_2(n) = \hat{B}_2(n) = \frac{1}{2\pi^2 n^2} \quad \text{for} \quad n \neq 0$$
(C12)

However, if one tries to translate the real-space left eigenvectors $l_p(x) = D_p(x)$ of Eq. 14 for p = 1, 2, ... in the Fourier basis, one obtains that all coefficients vanish. And if one writes the left eigenvalue equation directly in terms of Fourier coefficients

$$\lambda_p \ \hat{l}_p(m) = \sum_{n=-\infty}^{+\infty} \hat{l}_p(n) \delta_{2n,m} = \begin{cases} 0 \text{ for odd } m \\ \hat{l}^{[p]}(\frac{m}{2}) \text{ for even } m \end{cases}$$
(C13)

one cannot obtain any solution for $p \neq 0$.

2. Backward dynamics for the Fourier coefficients

In Fourier space with Eqs C1 and C2, the backward dynamics of Eq. 45 translates for the Fourier coefficients into

$$\hat{\rho}_{\tau}^{B}(n) = \sum_{\tilde{n}=-\infty}^{+\infty} \hat{W}_{B}(n|\tilde{n})\hat{\rho}_{\tau-1}^{B}(\tilde{n})$$
(C14)

with the matrix

$$\hat{W}^{B}(n|\tilde{n}) \equiv \int_{0}^{1} dx e^{i2\pi nx} \int_{0}^{1} d\tilde{x} e^{-i2\pi \tilde{n}\tilde{x}} w(\tilde{x}|x) = \delta_{n,2\tilde{n}}$$
(C15)

so that Eq. C14 reduces to

$$\hat{\rho}_{\tau}^{B}(n) = \sum_{\tilde{n}=-\infty}^{+\infty} \delta_{n,2\tilde{n}} \hat{\rho}_{\tau-1}^{B}(\tilde{n}) = \begin{cases} \hat{\rho}_{\tau-1}^{B}(\frac{n}{2}) \text{ for even } n\\ 0 \text{ for odd } n \end{cases}$$
(C16)

The iteration up to the initial condition at $\tau = 0$ yields

$$\hat{\rho}_{\tau}^{B}(n) = \begin{cases} \hat{\rho}_{0}^{B}(\frac{n}{2\tau}) \text{ if } \frac{n}{2\tau} = n_{0} \text{ is an integer} \\ 0 \text{ if } \frac{n}{2\tau} \text{ is not an integer} \end{cases}$$
(C17)

The physical meaning is that the Fourier coefficients $\hat{\rho}_0^B(n_0)$ of the initial condition at $\tau = 0$ are transferred to the Fourier coefficients $\hat{\rho}_{\tau}^B(n = 2^{\tau} n_0)$ while all the other coefficients where $\frac{n}{2^{\tau}}$ is not an integer are zero.

For large time $\tau \to +\infty$, only the Fourier coefficient n = 0 survives

$$\hat{\rho}^B_{\tau}(n) \underset{\tau \to +\infty}{\simeq} \hat{\rho}^B_*(n) = \delta_{n,0} \tag{C18}$$

in correspondence with the uniform real-space steady state of Eq. 48. However this convergence is very weird, since for any finite large τ , there are finite isolated Fourier coefficients for $n = 2^{\tau} n_0$ in Eq. C16.

Via the exchange of Eq. B5 between the right and left eigenvectors between the forward and backward kernels, one obtains that the left eigenvectors of the backward Fourier kernel $\hat{W}^B(n|\tilde{n})$ of Eq. C15 are given by Eq. C11

$$\hat{l}_{p}^{B}(n) = \hat{r}_{p}(n) = \hat{B}_{p}(n) = -\frac{p!}{(i2\pi n)^{p}} \quad \text{for} \ n \neq 0$$
 (C19)

and correspond to the Fourier coefficients of the Bernoulli polynomials $B_p(x)$ of Eq. 12, while there are no right eigenvectors for $p \neq 0$, since there are no left eigenvectors for the Fourier forward kernel as discussed around Eq. C13.

- [1] Y. Oono, Progress of Theoretical Physics Supplement 99, 165 (1989).
- [2] J. Tailleur, "Grandes déviations, physique statistique et systèmes dynamiques PhD in Paris (2007).
- [3] R.S. Ellis, Physica D 133, 106 (1999).
- [4] H. Touchette, Phys. Rep. 478, 1 (2009);
 H. Touchette, Modern Computational Science 11: Lecture Notes from the 3rd International Oldenburg Summer School, BIS-Verlag der Carl von Ossietzky Universitat Oldenburg, (2011).
- [5] B. Derrida, J. Stat. Mech. P07023 (2007).
- [6] R J Harris and G M Schütz, J. Stat. Mech. P07020 (2007).
- [7] E.M. Sevick, R. Prabhakar, S. R. Williams and D. J. Searles, Ann. Rev. of Phys. Chem. Vol 59, 603 (2008).
- [8] H. Touchette and R.J. Harris, chapter "Large deviation approach to nonequilibrium systems" of the book "Nonequilibrium Statistical Physics of Small Systems: Fluctuation Relations and Beyond", Wiley (2013).
- [9] L. Bertini, A. De Sole, D. Gabrielli, G. Jona-Lasinio and C. Landim Rev. Mod. Phys. 87, 593 (2015).
- [10] R. L. Jack and P. Sollich, The European Physical Journal Special Topics 224, 2351 (2015).
- [11] A. Lazarescu, J. Phys. A: Math. Theor. 48 503001 (2015).
- [12] A. Lazarescu, J. Phys. A: Math. Theor. 50 254004 (2017).
- [13] R. L. Jack, Eur. Phy. J. B 93, 74 (2020).
- [14] A. de La Fortelle, PhD Thesis (2000) "Contributions to the theory of large deviations and applications" INRIA Rocquencourt.

- [15] V. Lecomte, PhD Thesis (2007) "Thermodynamique des histoires et fluctuations hors d'équilibre" Université Paris 7.
- [16] R. Chétrite, PhD Thesis (2008) "Grandes déviations et relations de fluctuation dans certains modèles de systèmes hors d'équilibre" ENS Lyon.
- [17] B. Wynants, arXiv:1011.4210, PhD Thesis (2010), "Structures of Nonequilibrium Fluctuations", Catholic University of Leuven.
- [18] L. Chabane, PhD Thesis (2021) "From rarity to typicality : the improbable journey of a large deviation", Université Paris-Saclay.
- [19] J. du Buisson, PhD Thesis (2022) "Dynamical large deviations of diffusions" Stellenbosch University, South Africa arXiv:2210.09040.
- [20] R. Chétrite, HDR Thesis (2018) "Pérégrinations sur les phénomènes aléatoires dans la nature", Laboratoire J.A. Dieudonné, Université de Nice.
- [21] C. Giardina, J. Kurchan and L. Peliti, Phys. Rev. Lett. 96, 120603 (2006).
- [22] V. Lecomte, C. Appert-Rolland and F. van Wijland, Phys. Rev. Lett. 95 010601 (2005).
- [23] V. Lecomte, C. Appert-Rolland and F. van Wijland, J. Stat. Phys. 127 51-106 (2007).
- [24] V. Lecomte, C. Appert-Rolland and F. van Wijland, Comptes Rendus Physique 8, 609 (2007).
- [25] J.P. Garrahan, R.L. Jack, V. Lecomte, E. Pitard, K. van Duijvendijk and F. van Wijland, Phys. Rev. Lett. 98, 195702 (2007).
- [26] J.P. Garrahan, R.L. Jack, V. Lecomte, E. Pitard, K. van Duijvendijk and F. van Wijland, J. Phys. A 42, 075007 (2009).
- [27] K. van Duijvendijk, R.L. Jack and F. van Wijland, Phys. Rev. E 81, 011110 (2010).
- [28] R. L. Jack and P. Sollich, Prog. Theor. Phys. Supp. 184, 304 (2010).
- [29] D. Simon, J. Stat. Mech. (2009) P07017.
- [30] V. Popkov, G. M. Schuetz and D. Simon, J. Stat. Mech. P10007 (2010).
- [31] C. Giardina, J. Kurchan, V. Lecomte and J. Tailleur, J. Stat. Phys. 145, 787 (2011).
- [32] D. Simon, J. Stat. Phys. 142, 931 (2011).
- [33] V. Popkov and G. M. Schuetz, J. Stat. Phys 142, 627 (2011).
- [34] V. Belitsky and G. M. Schuetz, J. Stat. Phys. 152, 93 (2013).
- [35] O. Hirschberg, D. Mukamel and G. M. Schuetz, J. Stat. Mech. P11023 (2015).
- [36] G. M. Schuetz, From Particle Systems to Partial Differential Equations II, Springer Proceedings in Mathematics and Statistics Volume 129, pp 371-393, P. Gonçalves and A.J. Soares (Eds.), (Springer, Cham, 2015).
- [37] R. Chétrite and H. Touchette, Phys. Rev. Lett. 111, 120601 (2013).
- [38] R. Chétrite and H. Touchette Ann. Henri Poincare 16, 2005 (2015).
- [39] R. Chétrite and H. Touchette, J. Stat. Mech. P12001 (2015).
- [40] P. T. Nyawo and H. Touchette, Phys. Rev. E 94, 032101 (2016).
- [41] H. Touchette, Physica A 504, 5 (2018).
- [42] F. Angeletti and H. Touchette, Journal of Mathematical Physics 57, 023303 (2016).
- [43] P. T. Nyawo and H. Touchette, Europhys. Lett. 116, 50009 (2016);
 P. T. Nyawo and H. Touchette, Phys. Rev. E 98, 052103 (2018).
- [44] J.P. Garrahan, Physica A 504, 130 (2018).
- [45] C. Monthus, J. Stat. Mech. (2019) 023206.
- [46] C. Monthus, J. Phys. A: Math. Theor. 52, 485001 (2019).
- [47] E. Roldan and P. Vivo, Phys. Rev. E 100, 042108 (2019).
- [48] A. Lazarescu, T. Cossetto, G. Falasco and M. Esposito, J. Chem. Phys. 151, 064117 (2019).
- [49] B. Derrida and T. Sadhu, Journal of Statistical Physics 176, 773 (2019);
 B. Derrida and T. Sadhu, Journal of Statistical Physics 177, 151 (2019).
- [50] K. Proesmans and B. Derrida, J. Stat. Mech. (2019) 023201.
- [51] N. Tizon-Escamilla, V. Lecomte and E. Bertin, J. Stat. Mech. (2019) 013201.
- [52] J. du Buisson and H. Touchette, Phys. Rev. E 102, 012148 (2020).
- [53] E. Mallmin, J. du Buisson and H. Touchette, J. Phys. A: Math. Theor. 54 295001 (2021).
- [54] C. Monthus, J. Stat. Mech. (2021) 033303.
- [55] F. Carollo, J. P. Garrahan, I. Lesanovsky and C. Perez-Espigares, Phys. Rev. A 98, 010103 (2018).
- [56] F. Carollo, R.L. Jack and J.P. Garrahan, Phys. Rev. Lett. 122, 130605 (2019).
- [57] F. Carollo, J.P. Garrahan and R.L. Jack, J. Stat. Phys. 184, 13 (2021).
- [58] C. Monthus, J. Stat. Mech. (2021) 063301.
- [59] A. Lapolla, D. Hartich and A. Godec, Phys. Rev. Research 2, 043084 (2020).
- [60] C. Monthus, Eur. Phys. J. B 95, 32 (2022).
- [61] L. Chabane, A. Lazarescu and G. Verley, Journal of Statistical Physics 187, 6 (2022).
- [62] A. Mazzolo and C. Monthus, Phys. Rev. E 107, 014101 (2023).
- [63] J. du Buisson and H. Touchette, Phys. Rev. E 107, 054111, 2023.
- [64] J. du Buisson, T. D. P. Mnyulwa, H. Touchette, Phys. Rev. E 108, 044136, 2023.
- [65] C. Monthus, J. Stat. Mech. (2023) 083204.
- [66] J. Tailleur and J. Kurchan, Nature Physics 3, 203 (2007)
- [67] T. Laffargue, K.-D. N. T. Lam, J. Kurchan, J. Tailleur, J. Phys. A: Math. Theor. 46 254002 (2013).
- [68] N. R. Smith, Phys. Rev. E 106, L042202 (2022)
- [69] R. Gutierrez, A. Canella-Ortiz, C. Perez-Espigares, arXiv:2304.13754

- [70] G. Fayolle and A. de La Fortelle, Problems of Information Transmission 38, 354 (2002).
- [71] C. Monthus, Eur. Phys. J. B 92, 149 (2019) in the topical issue "Recent Advances in the Theory of Disordered Systems" edited by F. Igloi and H. Rieger.
- [72] C. Monthus, J. Stat. Mech. (2021) 033201.
- [73] C. Monthus, J. Stat. Mech. (2021) 063211
- [74] C. Monthus, Eur. Phys. J. B 95, 139 (2022).
- [75] C. Monthus,arXiv:2306.10932.
- [76] C. Monthus, arxiv 2308.12638
- [77] A. de La Fortelle, Problems of Information Transmission 37, 120 (2001).
- [78] C. Maes and K. Netocny, Europhys. Lett. 82, 30003 (2008).
- [79] C. Maes, K. Netocny and B. Wynants, Markov Proc. Rel. Fields. 14, 445 (2008).
- [80] A. C. Barato and R. Chétrite, J. Stat. Phys. 160, 1154 (2015).
- [81] L. Bertini, A. Faggionato and D. Gabrielli, Ann. Inst. Henri Poincare Prob. and Stat. 51, 867 (2015).
- [82] L. Bertini, A. Faggionato and D. Gabrielli, Stoch. Process. Appli. 125, 2786 (2015).
- [83] C. Monthus, J. Phys. A: Math. Theor. 52, 135003 (2019).
- [84] C. Monthus, J. Phys. A: Math. Theor. 52, 025001 (2019).
- [85] A. C. Barato, R. Chétrite, J. Stat. Mech. (2018) 053207.
- [86] L. Chabane, R. Chétrite, G. Verley, J. Stat. Mech. (2020) 033208.
- [87] C. Monthus, J. Stat. Mech. (2022) 013206.
- [88] C. Monthus, J. Stat. Mech. (2023) 063206.
- [89] C. Maes, K. Netocny and B. Wynants, Physica A 387, 2675 (2008).
- [90] J. Hoppenau, D. Nickelsen and A. Engel, New J. Phys. 18 083010 (2016).
- [91] C. Monthus, J. Stat. Mech. (2021) 083212.
- [92] C. Monthus, J. Stat. Mech. (2021) 083205.
- [93] C. Monthus, J. Stat. Mech. (2021) 103202.
- [94] P. Gaspard, Phys. A Math. Gen.25 (1992) L483-L485
- [95] L. Vepstas, https://www.linas.org/math/bernoulli.pdf
- [96] X.-Y. Hu and V. Rosenhaus, arxiv 2204.13655