

Evaluating Binary Outcome Classifiers Estimated from Survey Data

Adway S. Wadekar ^{*1} and Jerome P. Reiter¹

¹Department of Statistical Science, Duke University

May 2024

Abstract

Surveys are commonly used to facilitate research in epidemiology, health, and the social and behavioral sciences. Often, these surveys are not simple random samples, and respondents are given weights reflecting their probability of selection into the survey. We show that using survey weights can be beneficial for evaluating the quality of predictive models when splitting data into training and test sets. In particular, we characterize model assessment statistics, such as sensitivity and specificity, as finite population quantities and compute survey-weighted estimates of these quantities with test data comprising a random subset of the original data. Using simulations with data from the National Survey on Drug Use and Health and the National Comorbidity Survey, we show that unweighted metrics estimated with sample test data can misrepresent population performance, but weighted metrics appropriately adjust for the complex sampling design. We also show that this conclusion holds for models trained using upsampling for mitigating class imbalance. The results suggest that weighted metrics should be used when evaluating performance on test data derived from complex surveys.

Key Words: logistic, random forest, selection, tree, weight

1 Introduction

Empirical research in diverse fields, particularly within epidemiology and the social and behavioral sciences, relies on data collected via sample surveys (Safdar, Abbo, Knobloch, & Seo, 2016; Bennett et al., 1967; Wright & Marsden, 2010; Fabric, Choi, & Bird, 2012). Frequently, and especially for large-scale surveys run by government agencies, the surveys

*adway.wadekar@duke.edu

use complex design features like stratification, clustering, and unequal probability sampling. Each individual has a survey weight that, loosely, indicates how many individuals it represents in the population (Pfeffermann, 1996). Generally, the weight for any individual equals the inverse of that individual’s probability of selection into the survey, possibly with some multiplicative adjustment to account for survey nonresponse or to ensure weighted estimates of certain quantities are close to known population totals (Mansournia & Altman, 2016).

It is well-known that analysts should account for the survey design when estimating finite population quantities; for example, analysts can estimate population means or totals using the unbiased estimator of Horvitz and Thompson (1952). Analysts also can account for survey designs in explanatory models like linear and logistic regressions, for example, by using survey-weighted versions of those models (Dumouchel & Duncan, 1983; Reiter, Zanutto, & Hunter, 2005; Faiella, 2010). Often, however, analysts using data from complex survey designs do not use survey weights when fitting explanatory models (Bell et al., 2012). Indeed, there is debate as to whether or not survey weights are needed in model-based analyses at all (Bollen, Biemer, Karr, Tueller, & Berzofsky, 2016).

Independent of this debate, epidemiological, public health, and other social science researchers increasingly are using predictive modeling as an alternative to explanatory analyses (Naimi, Platt, & Larkin, 2018; Jiang, Gradus, & Rosellini, 2020; Bzdok, Varoquaux, & Steyerberg, 2021; Allen et al., 2024). As examples, Vázquez et al. (2020) use classification techniques to predict substance users in a sample of Mexican children; Wadekar (2020a, 2020b) uses machine learning methods to identify adults at risk for opioid or other substance use disorder; and, Borowski, Rosellini, Street, Gradus, and Vogt (2022) use predictive modeling to identify veterans at risk for suicidal ideation after a year of service. With the predictive analysis paradigm, the primary goal is to develop models that can be used to identify individuals in the population who might be at risk for a disease or other outcome, as opposed to identifying factors that might be associated with the outcome. To do so, researchers have expanded beyond parametric models to using classification and regression trees (CART) and random forests as well as black-box models like neural networks, sometimes in ensemble (Hastie, Tibshirani, & Friedman, 2009).

As with parametric models, the use of survey weights in the development of predictive models is idiosyncratic. Some analysts use them and some do not; see Section 2.2 for examples. The incorporation of survey weights can be complicated by some predictive modeling strategies. For instance, in data with imbalanced distributions of outcomes, it is common practice to go against the grain of representative survey weighting and up-weight, down-weight, or even synthetically generate observations to balance the outcome distribution in the training data to assist with model development (Jiang et al., 2020). Similarly, and of primary relevance here, some analysts use survey weights when assessing the performance of predictive models and others do not (Wieczorek, Guerin, & McMahon, 2022; Iparragirre, Barrio, Arostegui, & Arostegui, 2022; Iparragirre, Lumley, Barrio, & Arostegui, 2023).

In this article, we investigate the role of survey weights in the testing stage of predictive analysis for binary outcome classifiers. We do so for a common data science pipeline: construct a model on a random subsample of the survey data using, when necessary, procedures

like downsampling or SMOTE (Chawla, Bowyer, Hall, & Kegelmeyer, 2002) to adjust for class imbalance, use the model to predict outcomes for a holdout sample from the survey, and compute estimates of sensitivity and specificity using the holdout sample. We propose finite population quantities corresponding to the sensitivity and specificity that one would obtain by applying the predictive model to all individuals in the population, and we define survey-weighted estimators of those finite population quantities. Using simulation studies, we show that these survey-weighted estimators can approximate the corresponding finite population quantities more accurately than unweighted estimators. These findings apply for any predictive analysis on survey data, including those that use pre-processing steps like upsampling in model development. R code for computing the survey-weighted metrics is available at <https://github.com/.../....>

The remainder of the article is organized as follows. Section 2 briefly reviews survey weights and summarizes related work from the machine learning literature. Section 3 presents the survey-weighted estimators for predictive model evaluation. Section 4 describes simulation studies used to evaluate these estimators. These simulations use data from the National Survey on Drug Use and Health (U.S. Department of Health and Human Services, 2021) as well as data from the National Comorbidity Survey 2004 replication (Kessler & Merikangas, 2004). Section 5 summarizes the findings. An online supplement provides supporting material.

2 Background

In Section 2.1, we review key concepts from survey sampling. In Section 2.2, we review literature related to weighting in machine learning methods.

2.1 Survey Weights

Let P be a finite population comprising N individuals. Each individual $i = 1, \dots, N$ is measured on h survey variables, which we write as $\mathbf{z}_i = (z_{i1}, \dots, z_{ih})$. Thus, $P = \{\mathbf{z}_i : i = 1, \dots, N\}$. Let D comprise n randomly sampled individuals from P ; these are the data used for predictive modeling. For $i = 1, \dots, N$, we define the survey inclusion indicator $d_i = 1$ when individual i is sampled for the survey and $d_i = 0$ when not. Each individual has some nonzero probability π_i of being included in D , that is, $Pr(d_i = 1) = \pi_i > 0$. The values of π_i are determined by the randomization scheme used to select the sample.

Many survey analyses make use of the inverse probability of selection, $w_i = 1/\pi_i$, i.e., the survey weight. Each w_i can be interpreted as the number of individuals in P represented by individual i . Some national surveys adjust survey weights to $w_i = c_i(1/\pi_i)$, where c_i is some multiplicative factor, for example, to account for nonresponse, reduce standard errors of survey-weighted estimates, or to calibrate to auxiliary totals using raking (Little, 1986; Deville & Särndal, 1992; Brick, 2013; Da Silva & Opsomer, 2004; Phipps & Toth, 2012; Kott, 2016).

For the case of $w_i = 1/\pi_i$ for all i , Horvitz and Thompson (1952) present an estimator of population totals that can be applied for any sampling design. For any variable z_k , let $t_{z_k} = \sum_{i=1}^N z_{ik}$ be the total of z_k in P . We seek to estimate t_{z_k} from D . The estimator is

$$\hat{t}_{z_k} = \sum_{i \in D} w_i z_{ik}. \quad (1)$$

Horvitz and Thompson (1952) prove that $\mathbb{E}[\hat{t}_{z_k}] = t_{z_k}$, that is, the estimator in (1) is unbiased for any survey design. In these expectations, all individuals' w_i and z_{ik} are considered fixed features of that individual. The random variables are the indicators d_i of who is randomly selected to be in the sample. The estimator in (1) also can be unbiased when the w_i are adjusted weights under certain conditions, for example, on the reasons why individuals do not respond to the survey (Little & Rubin, 2020). We assume that the analyst uses whatever survey weights w_i are available in the data files.

2.2 Weights in Predictive Modeling

An array of literature discusses weighting and non-representative data in machine learning frameworks for model training. This literature generally does not address survey weights for computing evaluation metrics like sensitivity and specificity. Rather, much of it focuses on using analyst-specified weights to train models accurately, with an ultimate goal of mitigating the class imbalance problem (Huang, Li, Loy, & Tang, 2016; Byrd & Lipton, 2019; Xu, Ye, & Ruan, 2021). Steingrimsson, Gatsonis, Li, and Dahabreh (2023) study how models should be altered when training on data from one source and testing on another. Keilwagen, Grosse, and Grau (2014) discuss the use of weights to train and evaluate models when outcome labels are assigned not with certainty but with a degree of confidence.

Zadrozny (2004) shows that machine learning classifiers fit to nonrepresentative data can result in biased estimates of classification. Additionally, Zadrozny (2004) presents a theoretical approach to adjust machine learning model fits for nonrepresentative data. The basic idea is to introduce weights to the loss function used for model fitting. Each individual's weight is the ratio of an estimated marginal probability that the individual is in the training data over an estimated conditional probability that the individual is in the training data, given some analyst-specified set of variables. Although similar in objectives, our contributions differ from Zadrozny (2004) in that we use weights that account for the probability of selection into the survey and subsequently the subsample of cases in the evaluation set. These weights differ from ratios of estimated marginal and conditional probabilities that do not necessarily account for the two-stage sampling process. We also present metrics specifically to estimate sensitivity and specificity in complex samples rather than arbitrary loss functions with analyst-specified weights. Finally, we perform repeated sampling simulations using training and testing data with genuine complex survey data, comparing both weighted and unweighted performance metrics against defined, population-level results.

Turning to the roles of complex survey weights in predictive modeling, Yao, Li, and Graubard (2015) present survey-weighted estimators for area-under-the-curve (AUC) metrics. Their approach is targeted at estimating the AUC of a fitted model estimated with

all the training data; they do not discuss splitting into training and evaluation data. They also do not consider procedures like downsampling to mitigate class imbalance. Tsujimoto (2022) also studies survey-weighted estimates of AUCs of this form.

MacNell et al. (2023) examine the use of weights when estimating and evaluating gradient boosting classifiers. They treat a survey-weighted version of the classifier coupled with survey-weighted estimates of performance as a gold standard, and compare the performance of unweighted classifiers against this standard. Wieczorek et al. (2022) examine the use of weights when evaluating classifiers using k -fold cross validation and ultimately recommend using survey-weighted estimates for evaluations. While the topics and findings of these two papers are similar to ours, our work contributes to the literature in ways not addressed by these papers. Principally, we formally define finite population versions of sensitivity and specificity for general classifiers, whereas these two papers leave such quantities unspecified. Formal definitions of finite population quantities enable theoretical arguments for why estimators of them are (or are not) consistent, regardless of the classifier used to make predictions. In addition, these quantities facilitate considerations of variances of the estimators and for computation of estimators of the variances. They also facilitate benchmarking the weighted and unweighted performance metrics against a truth, as opposed to comparing against one another. Thus, we can assess which, if any, of the metrics is a reliable estimator of population-level accuracy.

3 Methodology

We begin by defining some standard performance metrics that are commonly used in assessing the predictive performance of classifiers. Let TP refer to the number of true positives predicted by a model, and let TN refer to the number of true negatives predicted by a model. Similarly, let FP and FN be the number of false positives and false negatives predicted by the model, respectively.

Definition 3.1. *Sensitivity is the percentage of observations with a positive outcome that the predictive model identifies correctly.*

$$\text{Sensitivity} = \frac{TP}{TP + FN}. \quad (2)$$

Definition 3.2. *Specificity is the percentage of observations with a negative outcome that the predictive model identifies correctly.*

$$\text{Specificity} = \frac{TN}{TN + FP}. \quad (3)$$

To compute sensitivity and specificity, often analysts partition the collected data into a training set D_t and test set D_e , where $D = (D_t, D_e)$. We assume the analyst determines D_e (and thus D_t) by taking a simple random sample of D , although the analyst may choose some other design (Wieczorek et al., 2022). For $i = 1, \dots, N$, let $e_i = 1$ when individual

i is selected into D_e and $e_i = 0$ otherwise. Note that $e_i = 1$ only if $d_i = 1$, and $e_i = 0$ automatically for individuals not in D .

We conceive of values of sensitivity and specificity computed from D_e as estimates of quantities defined over P . Our goal, then, is to obtain approximately unbiased estimates of these population quantities, as we now describe. To begin, we slightly modify the notation for the survey variables. For $i = 1, \dots, N$, we write $\mathbf{z}_i = (\mathbf{x}_i, y_i)$, where y_i represents an outcome of interest and \mathbf{x}_i represents a set of predictor variables of interest. Here, we assume that each $y_i \in \{0, 1\}$. The analyst fits a predictive model using D_t that can take any \mathbf{x}_i and produce a predicted value \hat{y}_i . We refer to this model as M so that $\hat{y}_i = M(\mathbf{x}_i)$.

Suppose that we could apply M to every individual in P . Then, we would have a collection of predictions \hat{y}_i for all N individuals in P . As a result, for $i = 1, \dots, N$, we could define each individual in P as a true positive, true negative, false positive, or false negative under M using the indicator variables,

$$TP_i = \mathbb{1}(\hat{y}_i = y_i \text{ and } y_i = 1) \quad (4)$$

$$TN_i = \mathbb{1}(\hat{y}_i = y_i \text{ and } y_i = 0) \quad (5)$$

$$FP_i = \mathbb{1}(\hat{y}_i \neq y_i \text{ and } y_i = 0) \quad (6)$$

$$FN_i = \mathbb{1}(\hat{y}_i \neq y_i \text{ and } y_i = 1). \quad (7)$$

Here, $\mathbb{1}(\cdot) = 1$ when the event inside the parentheses is true, and $\mathbb{1}(\cdot) = 0$ otherwise. For a fixed M , each individual in P can have only one of these indicators equal to 1.

Using these indicators, we define the finite population quantities,

$$N_{TP} = \sum_{i=1}^N TP_i \quad (8)$$

$$N_{TN} = \sum_{i=1}^N TN_i \quad (9)$$

$$N_{FP} = \sum_{i=1}^N FP_i \quad (10)$$

$$N_{FN} = \sum_{i=1}^N FN_i. \quad (11)$$

Thus, we define a finite population sensitivity under M as $N_{TP}/(N_{TP} + N_{FN})$ and a finite population specificity as $N_{TN}/(N_{TN} + N_{FP})$.

We next consider how to estimate these finite population quantities. In doing so, we conceive of each \hat{y}_i as a fixed characteristic of individual i , where $i = 1, \dots, N$. Put another way, we pretend that the analyst would like to apply M and hence compute \hat{y}_i for all individuals in P , and that the analyst would not use some other M with different samples of D or D_t . Obviously, this may not be true in practice. If the analyst had obtained a different M from another D or D_t , they would want to evaluate the quality of those values of $M(\mathbf{x}_i)$

for the individuals in P . However, conceiving of M as fixed simplifies understanding of the finite population quantities in (8)–(11).

Because we use random sampling to create D_e , we can use Horvitz and Thompson (1952) estimators for (8)–(11). We need the probability π_i^* that any individual i is in D_e , i.e., $\pi_i^* = P(e_i = 1)$. For $i = 1, \dots, N$, we have

$$\pi_i^* = \mathbb{P}(d_i = 1)\mathbb{P}(e_i = 1 \mid d_i = 1) = \pi_i\pi_{ei}. \quad (12)$$

Typically, we obtain D_e as a simple random sample of n_e individuals from the n in D , so that $\mathbb{P}(e_i = 1 \mid d_i = 1) = \pi_{ei} = n_e/n$. Thus, for each individual $i = 1, \dots, n$ in D , we define

$$w_i^* = w_i(n/n_e), \quad (13)$$

where w_i is the original survey weight.

With these weights, we can use Horvitz and Thompson (1952) estimators for (8)–(11). We have

$$\widehat{N}_{TP} = \sum_{i \in D_e} w_i^* TP_i \quad (14)$$

$$\widehat{N}_{TN} = \sum_{i \in D_e} w_i^* TN_i \quad (15)$$

$$\widehat{N}_{FP} = \sum_{i \in D_e} w_i^* FP_i \quad (16)$$

$$\widehat{N}_{FN} = \sum_{i \in D_e} w_i^* FN_i. \quad (17)$$

The expectations of the estimators in (14)–(17) are equal to the corresponding quantities in (8)–(11), assuming that M is fixed. We show that this is the case for \widehat{N}_{TP} ; proofs for the other estimators are similar. Assuming for simplicity that each $w_i = 1/\pi_i$, we have

$$\mathbb{E} \left[\widehat{N}_{TP} \right] = \mathbb{E} \left[\mathbb{E} \left[\widehat{N}_{TP} \mid D \right] \right] = \mathbb{E} \left[\mathbb{E} \left[\sum_{i \in D_e} w_i^* TP_i \mid D \right] \right] = \mathbb{E} \left[\mathbb{E} \left[\sum_{i \in D_e} (1/(\pi_i\pi_{ei})) TP_i \mid D \right] \right] \quad (18)$$

$$= \mathbb{E} \left[\sum_{i \in D} \mathbb{E} \left[(1/(\pi_i\pi_{ei})) e_i TP_i \mid D \right] \right] = \mathbb{E} \left[\sum_{i \in D} (1/\pi_i) TP_i \right] \quad (19)$$

$$= \mathbb{E} \left[\sum_{i \in P} (1/\pi_i) d_i TP_i \right] = \sum_{i \in P} TP_i \quad (20)$$

since $\mathbb{E}(e_i \mid D) = \pi_{ei}$ and $\mathbb{E}(d_i) = \pi_i$ for all $i = 1, \dots, N$. The random variables in these expectations are the indicators d_i and e_i ; all TP_i are considered fixed attributes of the individual i , for fixed M .

We plug in the estimates from (14)–(17) to obtain the estimates of the sensitivity (SN) and specificity (SP). We have

$$\widehat{SN} = \frac{\widehat{N}_{TP}}{\widehat{N}_{TP} + \widehat{N}_{FN}} \quad (21)$$

$$\widehat{SP} = \frac{\widehat{N}_{TN}}{\widehat{N}_{TN} + \widehat{N}_{FP}}. \quad (22)$$

Ratios of Horvitz and Thompson (1952) estimators are consistent for the corresponding population ratios (Lohr, 2021). Estimates of standard errors of ratio estimators can be obtained using Taylor series approximations. These are available in survey sampling software like the “survey” package in R (Lumley, 2004). We note that survey-weighted estimates of sensitivity and specificity like (21) and (22) have been proposed by Iparragirre et al. (2022) and Iparragirre, Barrio, and Arostegui (2023).

4 Simulation Studies

We present three simulation studies to examine the repeated sampling properties of \widehat{SN} and \widehat{SP} . In the first, we construct a population using data from the National Survey on Drug Use and Health (NSDUH), from which we take stratified samples. In the second and third, we use data from the NSDUH and the 2004 replication of the National Comorbidity Survey (NCS-R). We use the adjusted survey weights available on the data files. We do not know the values of \mathbf{z}_i for records not in the surveys, so that we only can note differences in the values of weighted and unweighted estimates.

4.1 Simulation using constructed population from NSDUH

We treat data from the NSDUH as a population, and take samples from it. Here, we are motivated by analyses in Wadekar (2020b), who uses data from the NSDUH to predict opioid use disorder. He uses logistic regression, random forests, and decision trees as classifiers, and reports unweighted estimates of performance metrics.

Specifically, we combine data from the 2017, 2018, and 2019 editions of the NSDUH to construct a population of $N = 116,761$ individuals. We ensure that all N individuals have values for all variables of interest. This necessitates discarding 11,558 individuals with partially complete data from the original data files. As the outcome variable, we set $y_i = 1$ if individual i has any illicit drug use in the past year, and $y_i = 0$ otherwise. Marginally, 31,807 (27.2%) people have illicit drug use and 84,954 (72.8%) people do not. As the predictors, we use sex, race, age category, income, an indicator of any mental illness within the last year, education level, self-reported health rating, employment status, age of first use of alcohol, age of first use of marijuana, parole status, probation status, approached by someone using drugs, obesity, whether heroin is fairly or very easy to obtain, perception of great risk trying heroin once or twice, and perception of great risk trying heroin weekly.

One-stage Stratified Design Schematic




Figure 1: Sampling design schematic for the stratified sampling simulation in Section 4.1.

Figure 1 displays a schematic for any one run of the simulation. First, we sample $n = 10,000$ records from the constructed P according to a stratified design. The strata are age categories defined in the NSDUH data files, namely 18–25 years, 26–34 years, 35–49 years, 50–64 years, and over 65 years. Labeling the strata in increasing age order, the strata sizes in P are $N_1 = 38,475$, $N_2 = 23,953$, $N_3 = 30,787$, $N_4 = 13,371$, and $N_5 = 10,175$. As sample sizes for D , we use $n_1 = 2,300$, $n_2 = 1,500$, $n_3 = 1,950$, $n_4 = 1,750$ and $n_5 = 2,500$. Thus, the sampling rates are not proportional to the population sizes. We oversample the oldest and youngest groups to demonstrate a design where ignoring weights could be problematic for evaluation.

We split the stratified sample D into training and test sets. We train three classifiers on an 80% sample of D : a baseline logistic regression classifier, a random forest classifier that does not adjust for class imbalance in the outcome variable, and a random forest classifier that does adjust for class imbalance using a standard upsampling approach (Japkowicz & Stephen, 2002). Specifically, we randomly sample and duplicate observations in the minority class in the 80% training set until the number of observations associated with each outcome is balanced. The resulting data are used to fit the classifier. The test set, D_e , remains unbalanced. For the random forests, we use the “randomForest” package in R (Liaw & Wiener, 2002). We use the default splitting Gini impurity splitting criteria and set the number of trees in the forest to 1,000.

We repeat the process of sampling D and fitting the classifiers 500 times. In each run, we apply the classifiers trained on D_t to all N records in P , including those in D . In this way, in each run we can compute $(N_{TP}, N_{TN}, N_{FP}, F_{FN})$, since we know y_i for all units in

Metric	Population	Unweighted	Weighted
Logistic regression			
Sensitivity	0.477 (± 0.0004)	0.426 (± 0.001)	0.478 (± 0.001)
Specificity	0.914 (± 0.0001)	0.940 (± 0.0002)	0.914 (± 0.0004)
AUROC	0.814 (± 0.00002)	0.825 (± 0.0005)	0.815 (± 0.0004)
Random forest			
Sensitivity	0.460 (± 0.0005)	0.399 (± 0.001)	0.444 (± 0.001)
Specificity	0.925 (± 0.0002)	0.944 (± 0.0003)	0.922 (± 0.0004)
AUROC	0.812 (± 0.00006)	0.807 (± 0.005)	0.799 (± 0.0005)
Balanced random forest			
Sensitivity	0.670 (± 0.0004)	0.602 (± 0.001)	0.650 (± 0.001)
Specificity	0.800 (± 0.0003)	0.842 (± 0.0005)	0.791 (± 0.0006)
AUROC	0.808 (± 0.00007)	0.804 (± 0.0004)	0.792 (± 0.0005)

Table 1: Results from simulation in Section 4.1. Entries are average values over 500 independent runs. Monte Carlo standard deviations are in parentheses.

P . These population quantities allow us to compute SN and SP for each fitted classifier. Although the 10,000 values in D are included in each population quantity in each run, the population quantities are determined mainly by the estimated classifier’s performance on the 106,761 out-of-sample cases. By averaging the population quantities over the 500 runs, we also approximate their sampling distributions over repeated draws of D .

In each of the 500 runs, we also compute \widehat{SN} using (21) and \widehat{SP} using (22) for the D_e in that run. For comparisons, we compute unweighted estimates of sensitivity and specificity using the usual expressions applied to D_e as in Definitions 3.1 and 3.2. Finally, in each run we compute the area under the Receiver Operating Characteristic curves (AUROC). We compute weighted version of the AUROC by collecting the values of $(\widehat{SN}, \widehat{SP})$ over a grid of decision thresholds ranging from 0 to 1.

Table 1 summarizes the results over the 500 runs. For sensitivity and specificity at a 50% decision threshold, the averages of the weighted estimates are closer to the averages of SN and SP than the unweighted estimates. In particular, an analyst interpreting the unweighted estimates would be overly optimistic on specificity but unduly pessimistic on sensitivity; this is not the case for \widehat{SN} and \widehat{SP} . For AUROC, the averages for both the weighted and unweighted estimates are not substantially different from the average population AUROC. We also see this in the simulations of Section 4.2, where we discuss it further. Overall, the results suggest that using survey-weighted metrics, especially at commonly-used decision thresholds, can provide better indications of how a model would perform in the population than unweighted metrics.

	Unweighted	Weighted
NSDUH analysis		
Sensitivity	0.71 (± 0.003)	0.63 (± 0.004)
Specificity	0.76 (± 0.001)	0.81 (± 0.001)
AUROC	0.80 (± 0.001)	0.80 (± 0.001)
NCS-R analysis		
Sensitivity	0.66 (± 0.02)	0.67 (± 0.02)
Specificity	0.72 (± 0.01)	0.73 (± 0.01)
AUROC	0.73 (± 0.003)	0.74 (± 0.004)

Table 2: Weighted and unweighted performance metrics for the NSDUH and NCS-R analyses. Monte Carlo standard deviations are in parentheses.

4.2 Applications to surveys

We next turn to analyzing two genuine surveys without constructing populations. They comprise 38,813 individuals from the 2017 edition of the NSDUH and 9,205 individuals from the NCS-R. While we do not have population truths as benchmarks, we can compare the weighted and unweighted performance metrics. We also estimate the standard errors of the ratio estimators in (21) and (22) using the “survey” package in R (Lumley, 2004).

Unlike our simulation in Section 4.1, these surveys are not stratified samples; rather, they use multi-stage sampling designs. With the NSDUH data, we use the outcome and predictors described in Section 4.1. For the NCS-R data, we use the outcome and predictors from analyses done by Jiang, Gradus, Lash, and Fox (2021). They use random forests to predict whether or not someone has attempted suicide from an array of mental illness and substance use variables, including panic disorder, agoraphobia, specific phobia, social phobia, post-traumatic stress disorder, major depressive disorder, alcohol abuse, alcohol dependence, drug abuse, and drug dependence. Among the 9,205 individuals in the sample, 8,794 have not attempted suicide, while 411 have, making this an outcome variable with significant imbalance in the classes.

For both analyses, we proceed with 50 train-test cycles using balanced random forests. That is, in each of 50 runs, we use simple random samples to split the survey data into training and test sets comprising 80% of records and 20% of records, respectively. We use the upsampling approach of Japkowicz and Stephen (2002) to make each D_t , which we use to train the random forest using 1,000 trees and the Gini impurity splitting criteria. We compute weighted and unweighted estimates of the sensitivities, specificities, and AUROCs on each D_e . The 50 runs serve a different purpose than those in the previous section. Specifically, we use the 50 train/test splits to assess the stability of the estimated performance metrics over different splits. In practice, the analyst may well train the predictive model using a single train/test split.

Table 2 displays results for the NSDUH and NCS-R analyses. The differences in the unweighted and weighted metrics vary by survey. With the NSDUH data, the differences are large, with \widehat{SN} nearly 0.08 lower than the unweighted estimate of sensitivity and \widehat{SP} nearly

0.05 higher than the unweighted estimate of specificity. In contrast, in the NCS-R analysis, the weighted and unweighted metrics have nearly identical values. For both surveys, the Monte Carlo standard deviations indicate that the variations across the 50 runs are small, suggesting the results are stable.


To understand the contrast in results for the NSDUH and NCS-R analyses, we compare the distributions of the survey weights in the two surveys; these are displayed in Figure 2a and Figure 2b. The coefficient of variation for the weight distribution in the NSDUH data is 1.09, whereas for the NCS-R data it is 0.52. Evidently, the deviations in individuals' weights from their average are much larger in the NSDUH than in the NCS-R. When weights are dispersed, as in the NSDUH, we expect weighted estimates to differ more noticeably from their corresponding unweighted estimates. When weights are close together, as in the NCS-R, weighted estimates are not much different than multiplying each observation's indicator variable in (14)–(17) by a constant, which cancels from the numerator and denominator of the ratio estimators used to make \widehat{SN} and \widehat{SP} .

For both surveys, the weighted and unweighted averages of AUROC are nearly identical. We conjecture that this arises because the weighted and unweighted values of sensitivity and specificity are not likely to differ at threshold values far from the predicted probabilities generated by the classifier. For example, suppose a classifier mostly gives predicted probabilities near 0.5. Suppose we consider the threshold for classifying someone as having a disease as a 95% predicted probability or higher. For most individuals, TP_i in (4) likely equals zero, since few individuals if any will have a large enough predicted probability to be classified by the model as having the disease. Whether or not one multiplies values of TP_i by w_i does not strongly affect the estimated sensitivity value, which we expect to be low. We expect the opposite to hold for very low decision thresholds as well. Thus, in data where many threshold values are outside the range of most predicted probabilities, we do not expect the use of weights to have noticeable impact on the AUROC.


For any run, we also can compute standard errors for the survey-weighted, estimated population quantities. To give a sense of the magnitudes of the estimated standard errors, for each analysis we compute typical estimated standard errors as the square root of the average of the 50 estimated variances obtained from the “survey” package. For the NSDUH analysis, these averages are 0.02 and 0.007 for the sensitivity and specificity, respectively. For the NCS-R analysis, these averages are 0.06 and 0.01 for the sensitivity and specificity, respectively. Thus, we expect the evaluation metrics to be estimated reasonably accurately, with the highest uncertainty about the sensitivity in the NCS-R. Of note, these estimated standard deviations also reflect uncertainty from sampling D , which is not present in the Monte Carlo standard deviations of Table 2.

5 Discussion

By defining finite population quantities for sensitivity and specificity, analysts can use techniques from survey sampling to estimate how a classifier would perform if applied to the full population. The survey-weighted estimators have appealing theoretical properties and per-



(a) NSDUH weight distribution



(b) NCS-R weight distribution

Figure 2: Weight distributions in the two surveys used in Section 4.2.

form well in the simulation studies presented here. They can be used in typical data science pipelines that split data into training and test sets. They also can be used with any classifier; for example, an analyst can employ methods to mitigate class imbalance, and use the survey weights in the model assessment phase to gauge population-level accuracy. More broadly, the analyst can use whatever steps needed to improve in-sample accuracy of the classifier, and so long as survey weights are used when assessing predictive performance in the test set, the accuracy measures reflect how the model predictions would perform if applied to all units in the population. Given the findings presented here, we recommend that analysts use weighted metrics when reporting results from machine learning studies that make use of survey data.

References

- Allen, B., Schell, R. C., Jent, V. A., Krieger, M., Pratty, C., Hallowell, B. D., . . . Neill, D. B. (2024). PROVIDENT: Development and validation of a machine learning model to predict neighborhood-level overdose risk in Rhode Island. *Epidemiology*, to appear.
- Bell, B. A., Onwuegbuzie, A. J., Ferron, J. M., Jiao, Q. G., Hibbard, S. T., & Kromrey, J. D. (2012). Use of design effects and sample weights in complex health survey data: A review of published articles using data from 3 commonly used adolescent health surveys. *American Journal of Public Health*, *102*, 1399–1405.
- Bennett, J. W., Katz, D., Massarik, F., McClosky, H., Morgan, J. N., Suchman, E. A., . . . Trow, M. (1967). *Survey research in the social sciences* (C. Y. Glock, Ed.). Russell Sage Foundation.
- Bollen, K. A., Biemer, P. P., Karr, A. F., Tueller, S., & Berzofsky, M. E. (2016). Are survey weights needed? a review of diagnostic tests in regression analysis. *Annual Review of Statistics and Its Application*, *3*, 375–392.
- Borowski, S., Rosellini, A. J., Street, A. E., Gradus, J. L., & Vogt, D. (2022). The first year after military service: Predictors of U.S. veterans’ suicidal ideation. *American Journal of Preventive Medicine*, *63*, 233–241.
- Brick, J. M. (2013). Unit nonresponse and weighting adjustments: A critical review. *Journal of Official Statistics*, *29*, 329–353.
- Byrd, J., & Lipton, Z. (2019). What is the effect of importance weighting in deep learning? In *Proceedings of the 36th International Conference on Machine Learning* (pp. 872–881). PMLR.
- Bzdok, D., Varoquaux, G., & Steyerberg, E. W. (2021). Prediction, not association, paves the road to precision medicine. *JAMA Psychiatry*, *78*, 127–128.
- Chawla, N. V., Bowyer, K. W., Hall, L. O., & Kegelmeyer, W. P. (2002, June). SMOTE: Synthetic Minority Over-sampling Technique. *Journal of Artificial Intelligence Research*, *16*, 321–357. doi: 10.1613/jair.953
- Da Silva, D. N., & Opsomer, J. D. (2004). Properties of the weighting cell estimator under a nonparametric response mechanism. *Survey Methodology*, *30*, 45–55.
- Deville, J.-C., & Särndal, C.-E. (1992). Calibration estimators in survey sampling. *Journal of the American Statistical Association*, *87*(418), 376–382.
- Dumouchel, W. H., & Duncan, G. J. (1983). Using sample survey weights in multiple regression analyses of stratified samples. *Journal of the American Statistical Association*, *78*, 535–543.
- Fabic, M. S., Choi, Y., & Bird, S. (2012). A systematic review of demographic and health surveys: data availability and utilization for research. *Bulletin of the World Health Organization*, *90*, 604–612.
- Faiella, I. (2010). *The Use of Survey Weights in Regression Analysis* (SSRN Scholarly Paper No. 1601936). Rochester, NY: Social Science Research Network.
- Hastie, T., Tibshirani, R., & Friedman, J. (2009). *The Elements of Statistical Learning: Data mining, Inference, and Prediction* (Vol. 2). New York: Springer.

- Horvitz, D. G., & Thompson, D. J. (1952). A generalization of sampling without replacement from a finite universe. *Journal of the American Statistical Association*, *47*, 663–685.
- Huang, C., Li, Y., Loy, C. C., & Tang, X. (2016). Learning deep representation for imbalanced classification. In *2016 IEEE Conference on Computer Vision and Pattern Recognition (CVPR)* (pp. 5375–5384).
- Iparragirre, A., Barrio, I., & Arostegui, I. (2023). Estimation of the roc curve and the area under it with complex survey data. *Stat*, *12*(1), e635.
- Iparragirre, A., Barrio, I., Arostegui, J., & Arostegui, I. (2022). Estimation of cut-off points under complex-sampling design data. *SORT-Statistics and Operations Research Transactions*, *46*(1), 137–158.
- Iparragirre, A., Lumley, T., Barrio, I., & Arostegui, I. (2023). Variable selection with LASSO regression for complex survey data. *Stat*, *12*, e578.
- Japkowicz, N., & Stephen, S. (2002). The class imbalance problem: A systematic study. *Intelligent Data Analysis*, *6*, 429–449.
- Jiang, T., Gradus, J. L., Lash, T. L., & Fox, M. P. (2021). Addressing measurement error in random forests using quantitative bias analysis. *American Journal of Epidemiology*, *190*, 1830–1840.
- Jiang, T., Gradus, J. L., & Rosellini, A. J. (2020). Supervised machine learning: A brief primer. *Behavior Therapy*, *51*, 675–687.
- Keilwagen, J., Grosse, I., & Grau, J. (2014). Area under precision-recall curves for weighted and unweighted data. *PLoS ONE*, *9*, e92209.
- Kessler, R. C., & Merikangas, K. R. (2004). The National Comorbidity Survey Replication (NCS-R): Background and aims. *International Journal of Methods in Psychiatric Research*, *13*, 60–68.
- Kott, P. S. (2016). Calibration weighting in survey sampling. *Wiley Interdisciplinary Reviews: Computational Statistics*, *8*(1), 39–53.
- Liaw, A., & Wiener, M. (2002). Classification and regression by randomforest. *R News*, *2*(3), 18–22. Retrieved from <https://CRAN.R-project.org/doc/Rnews/>
- Little, R. J. A. (1986). Survey nonresponse adjustments for estimates of means. *International Statistical Review*, *54*, 139–157.
- Little, R. J. A., & Rubin, D. B. (2020). *Statistical Analysis with Missing Data*, third edition. Hoboken, NJ: John Wiley & Sons, Inc.
- Lohr, S. L. (2021). *Sampling: Design and Analysis*. New York: CRC Press.
- Lumley, T. (2004). Analysis of complex survey samples. *Journal of Statistical Software*, *9*, 1–19.
- MacNell, N., Feinstein, L., Wilkerson, J., Salo, P. M., Molsberry, S. A., Fessler, M. B., . . . Zeldin, D. C. (2023, January). Implementing machine learning methods with complex survey data: Lessons learned on the impacts of accounting sampling weights in gradient boosting. *PLOS ONE*, *18*, e0280387.
- Mansournia, M. A., & Altman, D. G. (2016). Inverse probability weighting. *BMJ*, *352*, i189.
- Naimi, A. I., Platt, R. W., & Larkin, J. C. (2018). Machine learning for fetal growth

- prediction. *Epidemiology*, 29, 290–298.
- Pfeffermann, D. (1996, September). The use of sampling weights for survey data analysis. *Statistical Methods in Medical Research*, 5, 239–261.
- Phipps, P., & Toth, D. (2012). Analyzing establishment nonresponse using an interpretable regression tree model with linked administrative data. *The Annals of Applied Statistics*, 6, 772–794.
- Reiter, J. P., Zanutto, E. L., & Hunter, L. W. (2005). Analytical modeling in complex surveys of work practices. *ILR Review*, 59, 82–100.
- Safdar, N., Abbo, L. M., Knobloch, M. J., & Seo, S. K. (2016). Research methods in health-care epidemiology: survey and qualitative research. *Infection Control and Hospital Epidemiology*, 37, 1272–1277.
- Steingrimsson, J. A., Gatsonis, C., Li, B., & Dahabreh, I. J. (2023). Transporting a prediction model for use in a new target population. *American Journal of Epidemiology*, 192(2), 296–304. Retrieved from <https://doi.org/10.1093/aje/kwac128> doi: 10.1093/aje/kwac128
- Tsujimoto, T. H. M. (2022). *Statistical methods for receiver operating characteristics (ROC) curves in complex sampling surveys* (Unpublished doctoral dissertation). University of North Carolina at Chapel Hill.
- U.S. Department of Health and Human Services. (2021). *National Survey on Drug Use and Health 2019*. Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality.
- Vázquez, A. L., Domenech Rodríguez, M. M., Barrett, T. S., Schwartz, S., Amador Buenabad, N. G., Bustos Gamiño, M. N., . . . Villatoro Velázquez, J. A. (2020). Innovative identification of substance use predictors: Machine learning in a national sample of Mexican children. *Prevention Science*, 21, 171–181.
- Wadekar, A. S. (2020a). A psychosocial approach to predicting substance use disorder (SUD) among adolescents. In *2020 IEEE/ACM International Conference on Advances in Social Networks Analysis and Mining (ASONAM)* (pp. 819–826). IEEE.
- Wadekar, A. S. (2020b). Understanding opioid use disorder (oud) using tree-based classifiers. *Drug and Alcohol Dependence*, 208, 107839.
- Wieczorek, J., Guerin, C., & McMahon, T. (2022). K-fold cross-validation for complex sample surveys. *Stat*, 11.
- Wright, J. D., & Marsden, P. V. (2010). Survey research and social science: History, current practice, and future prospects. In *Handbook of survey research* (pp. 3–26). Emerald Group Publishing Limited.
- Xu, D., Ye, Y., & Ruan, C. (2021, March). *Understanding the role of importance weighting for deep learning* (No. arXiv:2103.15209). arXiv.
- Yao, W., Li, Z., & Graubard, B. I. (2015). Estimation of ROC curve with complex survey data. *Statistics in Medicine*, 34, 1293–1303.
- Zadrozny, B. (2004). Learning and evaluating classifiers under sample selection bias. In *In International Conference on Machine Learning ICML'04* (pp. 903–910).