

TRANSITIVITY AND RELATED NOTIONS FOR GRAPH INDUCED SYMBOLIC SYSTEMS

PRASHANT KUMAR AND PUNEET SHARMA

ABSTRACT. In this paper, we investigate the dynamical behavior of a two dimensional shift X_G (generated by a two dimensional graph $G = (\mathcal{H}, \mathcal{V})$) using the adjacency matrices of the generating graph G . In particular, we investigate properties such as transitivity, directional transitivity, weak mixing, directional weak mixing and mixing for the shift space X_G . We prove that if $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ (for all i, j), while doubly transitivity (weak mixing) of X_H (or X_V) ensures the same for two dimensional shift generated by the graph G , directional transitivity (in the direction (r, s)) can be characterized through the block representation of $H^r V^s$. We provide necessary and sufficient criteria to establish horizontal (vertical) transitivity for the shift space X_G . We also provide examples to establish the necessity of the conditions imposed. Finally, we investigate the decomposability of a given graph into product of graphs with reduced complexity.

1. INTRODUCTION

Symbolic dynamics has widely been used to investigate the dynamics of a various physical and natural processes around us. As any general dynamical system can be embedded in a symbolic system with appropriate number of symbols [27], it is sufficient to study the shift spaces and its subsystems to investigate the dynamics of a general discrete dynamical system. As the study of a general symbolic system can be used to model the dynamics of various systems arising in different fields of sciences and engineering, the area has found applications in various areas of science and engineering such as automata theory, data transmission, data storage and communication systems [2, 3, 5].

2020 *Mathematics Subject Classification.* 37B10, 37B20, 37B51.

Key words and phrases. multidimensional shift spaces, shifts of finite type, transitivity, weak mixing, strong mixing.

In recent years, multidimensional symbolic dynamics has emerged as a topic of interest and has captivated the attention of several researchers across the globe. The convenience of symbol-based representation allows a system with higher complexity to be studied through multidimensional symbolic systems. The easier computability and wide applicability of such systems has enabled the topic to find applications in different fields of sciences and engineering including mathematics, computer science, physics and many more. In particular, area has found applications areas such as computational biology, tiling spaces and study of quasicrystals [7, 22, 24, 26]. In recent times, several researchers have extensively investigated various dynamical and structural aspects of multidimensional shifts and interesting results have been obtained. In [21], the author established the undecidability of the non-emptiness problem for multidimensional shift spaces. In [19], the authors characterized any multidimensional shift of finite type using an infinite matrix. In [20], the authors provide an algorithm for generating the elements of the shift space using a sequence of finite matrices (of increasing order). The authors establish that the sequence generated yields precisely the elements of the shift space under consideration and hence characterizes the elements of the shift space. In [23], the authors established that any strongly irreducible two dimensional shift space has a dense set of periodic points. In [1], the authors proved that any multidimensional shifts of finite type exhibiting positive topological entropy cannot be minimal. In particular, they established that any shift of finite type with positive topological entropy contains a shift which is not of finite type and hence contains infinitely many shifts of finite type. In [14], the authors investigate mixing Z^d shifts of finite type and sofic shifts with large entropy. They give examples to prove that there exists Z^d -mixing systems such that no non-trivial full shift is a factor for such systems. They also establish existence of sofic systems such that the only minimal subsystem is a singleton. In [13], authors prove that the set of entropies for Z^d shifts of finite type ($d \geq 2$) coincides with the set of the infimum of all possible recursive sequences of rational numbers. In [12], the author established that any $h \geq 0$ is the entropy of a Z^d effective dynamical system if and only if it is the \liminf of a recursive sequence of rational numbers. In [18], authors discuss the non-emptiness problem and the existence of periodic points for a multidimensional shift space. They derive conditions under which the shift space is non-empty and possesses periodic points. The authors investigate the structure of the shift space

using the generating matrices. They prove that any d -dimensional shift of finite type is finite if and only if it is conjugate to a shift generated through permutation matrices. In [10], authors investigated the structure and mixing properties of two-dimensional shift spaces generated by consistent matrices. In [11], the authors discuss the entropy and ergodicity of a two-dimensional shift space using consistent matrices. In [16], authors gave generalized definitions of transitivity for two-dimensional shift spaces and studied various mixing notions for Dot Systems. Before we move further, we provide some of the basic definitions and concepts required.

Let $A = \{a_i : i \in I\}$ be a finite set, d be a positive integer and let A be equipped with the discrete metric. Let $A^{\mathbb{Z}^d}$ be equipped with the product topology. Then, the metric $\mathcal{D} : A^{\mathbb{Z}^d} \times A^{\mathbb{Z}^d} \rightarrow \mathbb{R}^+$ defined as $\mathcal{D}(x, y) = \frac{1}{n+1}$ (where n is the least non-negative integer such that $x \neq y$ in $R_n = [-n, n]^d$) generates the product topology. For any $a \in \mathbb{Z}^d$, the map $\sigma^a : A^{\mathbb{Z}^d} \rightarrow A^{\mathbb{Z}^d}$ defined as $(\sigma^a(x))(k) = x(k+a)$ is a d -dimensional shift and is a homeomorphism. Any $X \subset A^{\mathbb{Z}^d}$ is called shift invariant if $\sigma^a(X) \subset X$ for all $a \in \mathbb{Z}^d$. Any non-empty, closed, shift invariant subset of $A^{\mathbb{Z}^d}$ is called a **subshift**. Let \mathcal{F} be a given set of finite patterns and let $X = \{x \in A^{\mathbb{Z}^d} : \text{any pattern from } \mathcal{F} \text{ does not appear in } x\}$. Then X defines a subshift of $A^{\mathbb{Z}^d}$ generated by set of forbidden patterns \mathcal{F} . We say that the shift space X is a **shift of finite type** if it can be generated by a finite forbidden set of finite patterns. We say that a pattern is **allowed** if it is not an extension of any forbidden pattern. See [2, 4, 5, 8, 9, 15, 19] for details.

A subshift X is called **transitive** if for any pair of non-empty open sets U, V in X , there exists $n \in \mathbb{Z}^d$ such that $\sigma^n(U) \cap V \neq \emptyset$. A one-dimensional subshift X is called **doubly transitive** if both the left and right shifts are transitive for positive time. For any $r \in \mathbb{Z}^d$, a subshift X is called **r-transitive** if for any non-empty open sets U, V in X , there exists $k \in \mathbb{Z}$ such that $\sigma^{kr}(U) \cap V \neq \emptyset$. It may be noted that a subshift X is r -transitive if the system (X, σ^r) is transitive (as a one dimensional system). A shift space X is called **totally transitive** if (X, σ^r) is transitive for all $r \in \mathbb{Z}^d \setminus \{0\}$. A subshift X is called **weakly mixing** if for any pairs of non-empty open sets U_1, U_2 and V_1, V_2 in X , there exists $n \in \mathbb{Z}^d$ such that $\sigma^n(U_i) \cap V_i \neq \emptyset$ for $i = 1, 2$. For any $r \in \mathbb{Z}^d$, a subshift X is called **r-weakly mixing** if for any pairs of non-empty open sets U_1, U_2 and V_1, V_2 in X , there exists $k \in \mathbb{Z}$ such that $\sigma^{kr}(U_i) \cap V_i \neq \emptyset$

for $i = 1, 2$. Once again, it may be noted that a subshift X is r -weakly mixing if the system (X, σ^r) is weakly mixing (as a one dimensional system). A subshift X is called **mixing** if for any non-empty open sets U and V in X , the set $\{n \in \mathbb{Z}^d : \sigma^n(U) \cap V \neq \emptyset\}$ is cofinite. Refer [2, 9, 16, 25] for details.

Let $G = (V, E)$ be a graph (where V and E are the set of vertices and edges respectively). It may be noted that the set of bi-infinite walks over the graph G is a one dimensional (one step) shift of finite type. Also, as any given shift of finite type X can be coded as a higher block shift (conjugate to X and can be generated by a finite graph G), every one dimensional shift of finite type can be visualized as a shift generated from some graph. Also, if $\{G_1, G_2, \dots, G_d\}$ is a set of graphs with a common set of vertices V , the collection naturally induces a d -dimensional shift of finite type (where i -th graph determines the compatibility of the vertices in the i -th direction). In such a case, we refer (G_1, G_2, \dots, G_d) as a **d -dimensional graph**. We say that a two dimensional graph $G = (H, V)$ is **connected** if for every pair of vertices $i, j \in \mathcal{V}(G)$, $\exists (r, s) \in (\mathbb{Z}^+ \times \mathbb{Z}^+) \setminus \{0, 0\}$ such that $(H^r V^s)_{ij} > 0$. It may be noted that if $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ for all i, j , then any $1 \times r$ strip can extended to a $2 \times r$ (and hence to $n \times m$ block for any $n, m \in \mathbb{N}$) block allowed for the shift space. Consequently, the condition ensures extension of a given $r \times s$ block to a valid configuration for the shift space under discussion. See [2, 5, 10, 11, 18] for details.

An $n \times n$ real, non-negative, square matrix A is called **irreducible** if for any pair of indices $i, j \in \{1, 2, \dots, n\}$, there exists $k \in \mathbb{N}$ such that $A_{ij}^k \neq 0$. Such a matrix A is called **primitive** if there exists $k \in \mathbb{N}$ such that $A_{ij}^k \neq 0 \forall i, j \in \{1, 2, \dots, n\}$. It can be seen that a one dimensional shift generated by the matrix A is doubly transitive (strongly mixing) if and only if A is irreducible (primitive). Further, it is known that any one dimensional shift of finite type X is totally transitive $\Leftrightarrow X$ is weakly mixing $\Leftrightarrow X$ is strongly mixing. Refer [2, 25] for details.

Let G_1 and G_2 be two directed graphs. Then, the Cartesian (Tensor) product of G_1 and G_2 is defined as the graph with $\mathcal{V}(G_1) \times \mathcal{V}(G_2)$ as the set of vertices. The vertices (a, b) and (c, d) are connected in **Cartesian Graph Product** (denoted by $G_1 \square G_2$), if $a = c$ ($ac \in \mathcal{E}(G_1)$) and $bd \in \mathcal{E}(G_2)$ ($b = d$). In other words, two tuples are connected in Cartesian graph product if one of the indices (of the tuple) coincide and the other is connected (in their respective graphs). The vertices

(a, b) and (c, d) are connected in **Tensor Graph Product** (denoted by $G_1 \times G_2$) if $ac \in \mathcal{E}(G_1)$ and $bd \in \mathcal{E}(G_2)$. We define the graph product (Cartesian and Tensor) of two-dimensional graphs G_1 and G_2 to be **component-wise graph product**. See [6, 17] for details.

In this paper, we investigate the dynamical behavior of a two dimensional shift generated by the graph $G = (\mathcal{H}, \mathcal{V})$. For the sake of simplicity, we use H and V for the adjacency matrices of \mathcal{H} and \mathcal{V} respectively. In particular, we investigate transitivity, r -transitivity, total transitivity, weakly mixing, r -weakly mixing and mixing for shift space under consideration. We prove that if $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0, \forall i, j$, while doubly transitivity (weak mixing) of X_H (or X_V) ensures the same for two dimensional shift generated by the graph G , irreducibility of $H^r V^s$ provides a necessary and sufficient criteria to establish directional doubly transitivity (in (r, s) direction ($rs > 0$)) for the multidimensional shift space under discussion. We provide examples to show that the results derived do not hold if any of the conditions imposed fails to hold. We also provide necessary and sufficient criteria to establish horizontal (vertical) transitivity for the shift space X_G . We also investigate the decomposability of a given graph into product of graphs with reduced complexity.

2. MAIN RESULTS

Proposition 1. *Let X_G be a shift space generated by the two dimensional graph $G = (\mathcal{H}, \mathcal{V})$. If H and V are irreducible permutation matrices such that $HV = VH$, then X_G comprises of a single periodic point (with finite orbit).*

Proof. Let X_G be shift space generated by the two dimensional graph $G = (\mathcal{H}, \mathcal{V})$. Firstly, note that if H and V are permutation matrices, then the horizontal (vertical) neighbor of any symbol is uniquely determined. Further, if $HV = VH$, fixing a symbol at $(0, 0)$ determines the symbols at all other positions in the plane and hence yields a unique configuration in the shift space X_G . Finally, as H and V are irreducible, any two symbols are connected by a horizontal (vertical) path. Consequently, the shift space comprises of a single periodic orbit (with finite orbit) and the proof is complete. \square

Remark 1. The above proof provides a sufficient condition for the shift space to be finite (in fact a single periodic orbit) when the generating matrices commute with each other. It may be noted that as the above arguments hold even when only one of the generating matrix

is irreducible, the above result holds under a weaker condition (irreducibility of one of the generating matrices). In [10], the authors establish transitivity of the shift space under irreducibility of one of the generating matrices when the generating matrices are “consistent”. However, as irreducibility of H (or V) ensures connectivity of any two vertices through a horizontal (vertical) path, $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ for all i, j ensures that any two allowed rectangular blocks can be connected (horizontally or vertically) and hence can be extended to a configuration in the shift space X_G . Consequently, the shift space X_G is transitive for a bigger class of subshifts (of the full shift) and we get the following result.

Proposition 2. *Let X_G be a shift space generated by the two dimensional graph $G = (\mathcal{H}, \mathcal{V})$ such that $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ holds for all i, j . If H (or V) is irreducible then X_G is transitive.*

Proof. Let H be an irreducible matrix and let $x, y \in X_G$. For any

central blocks $x_n = \begin{matrix} x_{-n,n} & \cdots & x_{n,n} \\ \vdots & & \vdots \end{matrix}$ and $y_n = \begin{matrix} y_{-n,n} & \cdots & y_{n,n} \\ \vdots & & \vdots \end{matrix}$ of x and y respectively, irreducibility of H ensures that $x_{n,n}$ and $y_{-n,-n}$ can be connected through a horizontal path. Consequently, there exists symbols a_1, a_2, \dots, a_k such that

$$\begin{array}{ccccccc} & & & & y_{-n,n} & \cdots & y_{n,n} \\ & & & & \vdots & & \vdots \\ x_{-n,n} & \cdots & x_{n,n} & a_1 \cdots a_k & y_{-n,-n} & \cdots & y_{n,-n} \\ \vdots & & \vdots & & & & \\ x_{-n,-n} & \cdots & x_{n,-n} & & & & \end{array}$$

is an allowed pattern for the shift space X_G . Further, as $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ holds for all i, j , such an allowed pattern can be extended to a $(4n+k+2) \times (4n+k+2)$ square and hence to a configuration in X_G . As the argument holds for any central blocks of any size n , for any neighborhoods U and V of x and y respectively, there exists $r, s \in \mathbb{N}$ such that $\sigma^{(r,s)}(U) \cap V \neq \emptyset$. As the arguments hold for any $x, y \in X_G$, X_G is transitive and the proof is complete. \square

Remark 2. The above results establish non-emptiness and transitivity under suitable conditions for a two dimensional shift space. While Proposition 1 establishes the shift space to comprise of a single periodic orbit when generating matrices are irreducible permutation

matrices commuting with each other, Proposition 2 provides sufficient conditions to establish transitivity for the two dimensional shift space respectively. However, the conditions derived are sufficient in nature and the conclusions derived may hold even in the absence of the conditions imposed. We now provide examples in support of our arguments.

Example 1. Let X_G be a shift space generated by the adjacency matrices given below:

$$H = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix} \end{matrix} \quad V = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 1 & 0 & 1 \\ 0 & 1 & 0 & 0 \end{pmatrix} \end{matrix}$$

It may be noted that H and V are neither irreducible nor permutation matrices. Further, as $HV \neq VH$, none of the conditions (imposed in Proposition 1) holds. However, as the shift space comprises of a single periodic orbit of the point

$$\begin{matrix} \dots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \dots \\ \dots & 1 & 2 & 1 & 2 & 1 & 2 & 1 & 2 & 1 & 2 & 1 & 2 & 1 & 2 & 1 & 2 & \dots \\ \dots & 3 & 4 & 3 & 4 & 3 & 4 & 3 & 4 & 3 & 4 & 3 & 4 & 3 & 4 & 3 & 4 & \dots \\ \dots & 1 & 2 & 1 & 2 & 1 & 2 & 1 & 2 & 1 & 2 & 1 & 2 & 1 & 2 & 1 & 2 & \dots \\ \dots & 3 & 4 & 3 & 4 & 3 & 4 & 3 & 4 & 3 & 4 & 3 & 4 & 3 & 4 & 3 & 4 & \dots \\ \dots & 1 & 2 & 1 & 2 & 1 & 2 & 1 & 2 & 1 & 2 & 1 & 2 & 1 & 2 & 1 & 2 & \dots \\ \dots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \dots \end{matrix}$$

X_G comprises of a single periodic point (even when none of the conditions imposed in Proposition 1 hold).

Example 2. Let X_G be the shift space corresponding to graph $G = (\mathcal{H}, \mathcal{V})$, where the adjacency matrices for the generating graph are given by

$$H = \begin{matrix} & \begin{matrix} 0 & 1 & 2 & e & f & g \end{matrix} \\ \begin{matrix} 0 \\ 1 \\ 2 \\ e \\ f \\ g \end{matrix} & \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 \end{pmatrix} \end{matrix} \quad V = \begin{matrix} & \begin{matrix} 0 & 1 & 2 & e & f & g \end{matrix} \\ \begin{matrix} 0 \\ 1 \\ 2 \\ e \\ f \\ g \end{matrix} & \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 \end{pmatrix} \end{matrix}$$

It may be noted that H and V are irreducible but $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ for all i, j does not hold. Further, as the pattern $2e$ cannot be extended vertically to form a valid pattern for the shift space, any valid configuration for X_G comprises either of $0, 1, 2$ or e, f, g and hence the two dimensional shift space X_G is not transitive. Thus, the shift space may fail to be transitive even when each of the generating matrices are irreducible (It is worth mentioning that if $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ holds for all i, j , irreducibility of any of the generating matrices is sufficient to ensure transitivity of the two dimensional shift X_G).

Corollary 1. *Let X_G be a shift space generated by the two dimensional graph $G = (\mathcal{H}, \mathcal{V})$ such that $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ holds for all i, j . If X_H (or X_V) is weak mixing then X_G exhibits weak mixing.*

Proof. As proof of Proposition 2 establishes that if H (or V) is irreducible then $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ for all i, j ensures transitivity of the shift space X_G . As a similar set of arguments can be repeated for any pair of open sets in the shift space, if X_H is weakly mixing then X_G is weak mixing (under the imposed condition). \square

Remark 3. The above result proves that if the imposed conditions hold, then the two dimensional shift is weak mixing when the one dimensional shift generated by any of the generating matrices is weak mixing. However, the result holds strictly under the imposed conditions and may not be true otherwise even when the one dimensional shifts generated by each of the generating matrices are weak mixing. We now give an example in support of our claim.

Example 3. *Let X_G be the two dimensional shift space generated by graph $G = (\mathcal{H}, \mathcal{V})$, where corresponding adjacency matrices are*

$$H = \begin{matrix} & \begin{matrix} 0 & 1 & 2 \end{matrix} \\ \begin{matrix} 0 \\ 1 \\ 2 \end{matrix} & \begin{pmatrix} 1 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} \end{matrix} \quad V = \begin{matrix} & \begin{matrix} 0 & 1 & 2 \end{matrix} \\ \begin{matrix} 0 \\ 1 \\ 2 \end{matrix} & \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \\ 1 & 0 & 1 \end{pmatrix} \end{matrix}$$

Firstly, it may be noted that H and V are primitive matrices (hence generate one dimensional weak mixing shifts) and the condition $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ for all i, j fails to hold. However, the two dimensional shift generated by G comprises of a single periodic orbit generated by the point

$$\begin{array}{cccccccccccccccccccc}
 \dots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \dots \\
 \dots & 1 & 2 & 0 & 0 & 1 & 2 & 0 & 0 & 1 & 2 & 0 & 0 & 1 & 2 & 0 & 0 & \dots & \dots \\
 \dots & 0 & 0 & 1 & 2 & 0 & 0 & 1 & 2 & 0 & 0 & 1 & 2 & 0 & 0 & 1 & 2 & \dots & \dots \\
 \dots & 1 & 2 & 0 & 0 & 1 & 2 & 0 & 0 & 1 & 2 & 0 & 0 & 1 & 2 & 0 & 0 & \dots & \dots \\
 \dots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \dots
 \end{array}$$

Consequently, the shift space X_G does not exhibit weak mixing. Thus, the shift space X_G generated by a graph G need not exhibit weak mixing even when the shift spaces generated by each of the generating matrices are weak mixing.

Proposition 3. *Let $G = (H, V)$ be a two dimensional connected graph satisfying $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ for all i, j , then X_G is transitive.*

Proof. Let U, V be $\frac{1}{n}$ -neighborhoods of configurations $x, y \in X_G$. If x_n and y_n are central blocks of length $(2n + 1)$ of the elements x and y respectively, as G is connected, there exists a path from $x_{n,n}$ (top right corner of x_n) to $y_{-n,-n}$ (bottom left corner of y_n). As $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ (for all i, j), the extended pattern can further be extended to an allowed rectangular pattern in $\mathcal{B}(X_G)$ (and finally to a valid configuration $z \in X_G$). Thus, there exists $(r, s) \in (\mathbb{Z}^+ \times \mathbb{Z}^+) \setminus \{0, 0\}$ such that $\sigma^{(r,s)}(U) \cap V \neq \emptyset$. Hence X_G is transitive and the proof is complete. \square

Proposition 4. *Let X_G be a shift space and let $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ holds for all i, j . Then, X_G is doubly (r, s) -transitive ($rs > 0$) if and only if $H^r V^s$ is irreducible.*

Proof. Without loss of generality, assume that $r, s > 0$ (as $rs > 0$). Let X_G be doubly (r, s) -transitive ($rs > 0$), $x, y \in X_G$ and U, V be $\frac{1}{n}$ -neighborhoods of x and y respectively. Let x_n, y_n be central blocks (of size $2n + 1$) of the elements x and y respectively. As X_G is transitive in the direction (r, s) , there exists $u \in U, v \in V$ and $k \in \mathbb{N}$ such that $\sigma^{(kr, ks)}(u) = v$. As U, V are $\frac{1}{n}$ -neighborhoods of x, y respectively, x_n, y_n are central blocks of u, v respectively and hence $(H^r V^s)_{x_n, y_{-n, -n}}^k \neq 0$. As the proof holds with central blocks of arbitrary sizes (and hence with arbitrary corner points), $H^r V^s$ is irreducible and the proof of forward part is complete.

Conversely, let $H^r V^s$ be irreducible ($r, s > 0$), $x, y \in X_G$, U and V be $\frac{1}{n}$ -neighborhoods of x and y respectively and x_n, y_n be central blocks (of size $2n + 1$) of the elements x, y respectively. For any central blocks

Example 4. Let X_G be a shift space arising from the following adjacency matrices:

$$H = \begin{array}{c} \begin{array}{cccc} & 1 & 2 & 3 & 4 \\ 1 & (0 & 1 & 0 & 0) \\ 2 & (1 & 0 & 1 & 0) \\ 3 & (0 & 0 & 0 & 1) \\ 4 & (0 & 0 & 1 & 0) \end{array} \end{array} \quad V = \begin{array}{c} \begin{array}{cccc} & 1 & 2 & 3 & 4 \\ 1 & (1 & 0 & 0 & 0) \\ 2 & (0 & 1 & 0 & 0) \\ 3 & (0 & 0 & 1 & 0) \\ 4 & (0 & 0 & 0 & 1) \end{array} \end{array}$$

Then, it can be seen that X_G is (1,1)-transitive (is transitive along the line $y = x$). However, as

$$HV = VH = \begin{array}{c} \begin{array}{cccc} & 1 & 2 & 3 & 4 \\ 1 & (0 & 1 & 0 & 0) \\ 2 & (1 & 0 & 1 & 0) \\ 3 & (0 & 0 & 0 & 1) \\ 4 & (0 & 0 & 1 & 0) \end{array} \end{array}$$

The matrix HV fails to be irreducible and hence (r, s) -transitivity of the shift space X_G need not imply irreducibility of the matrix $H^r V^s$.

Corollary 2. Let X_G be a shift space where condition $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ holds for all i, j . Then, $H^r V^s$ is primitive $\Leftrightarrow X_G$ is weak mixing in direction (r, s) ($rs > 0$).

Proof. The proof follows from the discussions in the Remark 4. \square

Proposition 5. Let X_G be a one dimensional shift space generated by a one dimensional graph G (with adjacency matrix A). Then X_G is transitive $\Leftrightarrow A$ is irreducible or A has exactly two irreducible, permutation sub-matrices A_1, A_2 such that for any $i \in A_1, j \in A_2$, there exists a unique path connecting the two vertices (which traverses through all the vertices of $A \setminus (A_1 \cup A_2)$).

Proof. Let X_G be a one dimensional transitive shift space generated by a one dimensional graph G (with adjacency matrix A). Note that if A is not irreducible then there exists $i, j \in \{1, 2, \dots, n\}$ such that $A_{ij}^k = 0 \forall k \in \mathbb{N}$. However, as X_G is transitive, there exists a block of the form $ja_1 a_2 \dots a_r i$ allowed for the shift space. Further, as existence of two blocks of the form $ja_1 a_2 \dots a_r i$ ensures $A_{ij}^k \neq 0$ for some $k \in \mathbb{N}$ (as X_G is transitive), there exists a unique block of the form $ja_1 a_2 \dots a_r i$. Thus, fixing position of j fixes every position before j in a unique manner. Also, as existence of two blocks of the form $ja_1 a_2 \dots a_r i b_1 b_2 \dots b_m$ ensures $A_{ij}^k \neq 0$ for some $k \in \mathbb{N}$ (again by transitivity of X_G), fixing position of j fixes the entire sequence and the proof of forward part is complete.

Conversely, if hypothesis for the converse holds then any non-periodic point in the shift space is of the form $\overline{x_1 x_2 \dots x_k j c_1 c_2 \dots c_l y_1 y_2 \dots y_m}$. Also, as $\overline{x_1 x_2 \dots x_k}$ and $\overline{y_1 y_2 \dots y_m}$ are the only other elements in the shift space, any two blocks allowed for the shift space interact over time and the shift space is transitive. \square

Remark 5. The above result characterizes the transitivity for a one dimensional shift space generated by a graph G . However, as directional shifts for a multidimensional shift spaces can be visualized as a one dimensional shift, a similar result can be derived to characterize the transitivity (in a particular direction) for a multidimensional shift space. In particular, for a two dimensional shift space generated by $G = (\mathcal{H}, \mathcal{V})$, if $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ holds (for all i, j) then a similar argument establishes (r, s) -transitivity for the shift space if $H^r V^s$ can be represented using the similar block form (as derived in Proposition 5). Thus we get the following corollary.

Corollary 3. *Let X_G be a two dimensional shift space generated by the graph $G = (\mathcal{H}, \mathcal{V})$ such that $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ holds for all i, j . Then X_G is (r, s) -transitive $\Leftrightarrow H^r V^s$ is irreducible or $H^r V^s$ has exactly two irreducible, permutation sub-matrices A_1, A_2 such that for any $i \in A_1, j \in A_2$, there exists a unique path connecting the two vertices (which traverses through all the vertices of $H^r V^s \setminus (A_1 \cup A_2)$).*

Proof. The proof follows from discussions in Remark 5. \square

Proposition 6. *Let X_G be a two dimensional shift space and let $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ holds for all i, j . If H or V are irreducible matrices, then X_G is doubly (r, s) -transitive (for some direction $(r, s) \in \mathbb{N} \times \mathbb{N}$).*

Proof. Let X_G be the shift generated by the two dimensional graph G such that $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ holds for all i, j and let H be an irreducible matrix. As V generates a non-empty one dimensional shift, there exists a vertex $i \in \{1, 2, \dots, n\}$ and $m \in \mathbb{N}$ such that $V_{ii}^m \neq 0$. Also, as H is irreducible, there exists a finite path $W = a_0 a_1 \dots a_r$ such that $a_0 = a_r = i$ and each symbol j appears atleast once in the path W . Consequently, for the pattern P of the form

$$\begin{array}{ccccccc} & & i & a_1 & a_2 & \dots & a_{r-1} & i \\ & & & & & & & \vdots \\ & & & & & & & i \\ i & a_1 & a_2 & \dots & a_{r-1} & i & & \end{array}$$

is allowed for the shift space X_G . Further, note that diagonal repetition (R times, where R is more than the number of symbols) is allowed for the shift space and can be extended to a configuration in

X_G . Also, as $\sigma^{(s(r+1),sm)}(i) = a_s$, $H^{r+1}V^m$ is irreducible. Consequently, X_G is doubly $(r+1, m)$ -transitive and the proof is complete. \square

Remark 6. The above proof establishes that if $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ holds for all i, j , then irreducibility of any of the generating matrices ensures directional transitivity (in some direction (r, s)) and thus transitivity cannot happen in the shift space independently (without directional transitivity). However, in general, the transitivity of the shift space X_G need not imply directional transitivity and the notions are indeed distinct. Also, as weakly mixing ensures interaction between any two pairs of symbols, if any of the shift spaces generated by H or V is weakly mixing, $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ for all i, j ensures weak mixing in some direction (r, s) for the two dimensional shift space. It may be noted that as a shift is (r, s) -weak mixing if and only if H^rV^s is primitive (for $rs > 0$), (r, s) -weak mixing of a system is equivalent to $(1, 1)$ -weak mixing. It is worth mentioning that directional mixing of a system indeed depends on the direction and existence of mixing notions in one direction for a shift space X_G need not guarantee existence of mixing notion in the other. Consequently, total transitivity and weak mixing are indeed distinct notions for higher dimensional shifts of finite type (recall that the two notions coincide for shifts of finite type in one dimensional case). We now establish our claims below.

Corollary 4. *Let X_G be a shift space such that $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ for all i, j . If H or V is primitive, then X_G is (r, s) -weak mixing (for some direction $(r, s) \in \mathbb{N} \times \mathbb{N}$). Further, X_G is (r, s) -weak mixing (for $rs > 0$) if and only if X_G is $(1, 1)$ -weak mixing.*

Proof. The proof follows from the discussions in Remark 6. \square

Example 5. *Let X_G be a shift space arising from following adjacency matrices.*

$$H = V = \begin{matrix} & \begin{matrix} 0 & 1 & 2 & 3 \end{matrix} \\ \begin{matrix} 0 \\ 1 \\ 2 \\ 3 \end{matrix} & \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \end{pmatrix} \end{matrix}$$

It may be noted that as H is primitive ($H_{ij}^4 > 0 \forall i, j$), X_G is $(1, 1)$ -weak mixing. However, $HV^T = V^TH$ fails to hold. Further, as

$$HV^T = \begin{matrix} & 0 & 1 & 2 & 3 \\ \begin{matrix} 0 \\ 1 \\ 2 \\ 3 \end{matrix} & \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & 2 & 0 & 0 \\ 0 & 0 & 2 & 2 \\ 0 & 0 & 2 & 2 \end{pmatrix} \end{matrix} \quad V^T H = \begin{matrix} & 0 & 1 & 2 & 3 \\ \begin{matrix} 0 \\ 1 \\ 2 \\ 3 \end{matrix} & \begin{pmatrix} 2 & 0 & 0 & 2 \\ 0 & 2 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 2 & 0 & 0 & 2 \end{pmatrix} \end{matrix}$$

fixing any symbol at $(0,0)$ enforces the same symbol along the line $y = -x$ and hence X_G does not exhibit $(1,-1)$ -weak mixing. Consequently, any of the mixing notions in one direction need not ensure the mixing notions in the other. Also, as the example exhibits weak mixing in $(1,1)$ -direction, the case provides an example of weak mixing shift which fails to be totally transitive. Consequently, the two notions are distinct for higher dimensional shifts of finite type and an analogous extension of the result valid for one dimensional case does not hold.

Let P_k and Q_k denote the collection of valid $1 \times k$ and $k \times 1$ patterns respectively and let $V_k(H_k)$ denote the matrix characterizing vertical (horizontal) compatibility of elements of P_k (Q_k). It may be noted that $H_1 = H$ and $V_1 = V$. In general, H_k and V_k generate valid patterns of size $k \times r$ and $r \times k$ respectively (for any $r \in \mathbb{N}$). Such matrices have been defined in [11] and have been used to investigate the dynamical properties of “shifts of finite type” (as defined in [10, 11]). We now investigate notions of horizontal (vertical) mixing for the shift space X_G in terms of the matrices H_k and V_k .

Proposition 7. *Let X_G be a two dimensional shift space generated by a graph G and let $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ holds for all i, j . Then, X_G is horizontally doubly transitive $\Leftrightarrow H_k$ is irreducible $\forall k \in \mathbb{N}$.*

Proof. Let X_G be a two dimensional horizontally doubly transitive

$$\text{shift space. Let } x, y \in X_G, x_k = \begin{matrix} x_{-k,k} & \cdots & x_{k,k} \\ \vdots & & \vdots \\ x_{-k,-k} & \cdots & x_{k,-k} \end{matrix} \text{ and } y_k = \begin{matrix} y_{-k,k} & \cdots & y_{k,k} \\ \vdots & & \vdots \\ y_{-k,-k} & \cdots & y_{k,-k} \end{matrix},$$

be central blocks and let U, V be $\frac{1}{k}$ -neighborhoods of x, y respectively. As X_G is horizontally doubly transitive, there exists $z \in U, w \in V$ and $r \in \mathbb{N}$ such that $\sigma^{(r,0)}(z) = w$. It may be noted that as $z \in U$ and $w \in V$, the central blocks of z and w are x_k and y_k respectively. Further, as

$$\sigma^{(r,0)}(z) = w, a = \begin{matrix} x_{k,k} \\ \vdots \\ x_{k,-k} \end{matrix} \text{ and } b = \begin{matrix} y_{-k,k} \\ \vdots \\ y_{-k,-k} \end{matrix} \text{ can be connected horizontally}$$

(through a finite path). As the arguments hold for any $a, b \in H_k$, H_k is irreducible. Further, as the argument holds for any $k \in \mathbb{N}$, H_k is

irreducible for all $k \in \mathbb{N}$ and the proof of forward part is complete.

Conversely, let H_k be irreducible for all $k \in \mathbb{N}$. Let $x, y \in X_G$ and let U and V be $\frac{1}{k}$ -neighborhoods of x and y respectively. For any

central blocks $x_k = \begin{matrix} x_{-k,k} & \cdots & x_{k,k} \\ \vdots & & \vdots \\ x_{-k,-k} & \cdots & x_{k,-k} \end{matrix}$ and $y_k = \begin{matrix} y_{-k,k} & \cdots & y_{k,k} \\ \vdots & & \vdots \\ y_{-k,-k} & \cdots & y_{k,-k} \end{matrix}$ of x and y respectively, irreducibility of H_k ensures that $a = \begin{matrix} x_{k,k} \\ \vdots \\ x_{k,-k} \end{matrix}$ and

$b = \begin{matrix} y_{-k,k} \\ \vdots \\ y_{-k,-k} \end{matrix}$ can be connected horizontally through a finite path (say of length l) and hence there exists a valid pattern of the form

$$\begin{matrix} x_{-k,k} & \cdots & x_{k,k} & \cdots & y_{-k,k} & \cdots & y_{k,k} \\ \vdots & & \vdots & & \vdots & & \vdots \\ x_{-k,-k} & \cdots & x_{k,-k} & \cdots & y_{-k,-k} & \cdots & y_{k,-k} \end{matrix}$$

As $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ holds for all i, j , the above pattern can be extended to a valid configuration in X_G (say z). Then, $z \in U$, $\sigma^{(2k+1+l,0)}(z) \in V$. As the proof holds for any $x, y \in X_G$, the shift space is horizontally doubly transitive and the proof is complete. \square

Remark 7. The above result establishes that for any shift space X_G , X_G is horizontally doubly transitive if and only if each H_k is irreducible. However, transitivity of X_G need not ensure irreducibility for all H_k . Once again, as horizontal transitivity can be visualized as a one dimensional shift, horizontal transitivity of the shift space can be characterized via the block representations of the matrices H_k . Finally, as similar arguments establish the equivalence of vertical transitivity with irreducibility of V_k (for all k), we get the following corollary.

Corollary 5. *Let X_G be two dimensional shift space generated by the graph $G = (\mathcal{H}, \mathcal{V})$ such that $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ for all i, j . Then, X_G is horizontally transitive \Leftrightarrow either all H_k are irreducible or all H_k have exactly two irreducible, permutation sub-matrices A_{1k}, A_{2k} such that for any $i \in A_{1k}, j \in A_{2k}$, there exists a unique path connecting the two vertices (which traverses through all the vertices of $H_k \setminus (A_{1k} \cup A_{2k})$).*

Proof. The proof follows from discussions in Remark 7. \square

Proposition 8. *Let X_G be a two dimensional shift generated by the graph G and let $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ holds for all i, j . If X_H is weakly mixing then X_G is vertically doubly transitive $\Leftrightarrow X_G$ is vertically weakly mixing.*

Proof. Let X_G be vertically doubly transitive shift space and let $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ holds for all i, j . Let $x, y, z, w \in X_G$ and

$$x_k = \begin{array}{ccc} x_{-k,k} & \dots & x_{k,k} \\ \vdots & & \vdots \\ x_{-k,-k} & \dots & x_{k,-k} \end{array}, y_k = \begin{array}{ccc} y_{-k,k} & \dots & y_{k,k} \\ \vdots & & \vdots \\ y_{-k,-k} & \dots & y_{k,-k} \end{array}, z_k = \begin{array}{ccc} z_{-k,k} & \dots & z_{k,k} \\ \vdots & & \vdots \\ z_{-k,-k} & \dots & z_{k,-k} \end{array}$$

and $w_k = \begin{array}{ccc} w_{-k,k} & \dots & w_{k,k} \\ \vdots & & \vdots \\ w_{-k,-k} & \dots & w_{k,-k} \end{array}$ be central blocks of size k (squares of size

$$w_{-k,-k} \dots w_{k,-k}$$

$2k+1$) of x, y, z and w respectively. Let U, V, Z, W be $\frac{1}{k}$ -neighborhoods of x, y, z and w respectively. As X_H is weakly mixing, the symbols $x_{k,k}$ and $z_{k,-k}$ can be connected to $y_{-k,k}$ and $w_{-k,-k}$ respectively through horizontal paths of same length (say l). Also, as V_k 's are irreducible for all k , the strips $x_{-k,k} \dots x_{k,k} \dots y_{-k,k} \dots y_{k,k}$ and $z_{-k,-k} \dots z_{k,-k} \dots w_{-k,-k} \dots w_{k,-k}$ can be connected vertically (say through a path of length p). Consequently, there exists patterns of the form

$$\begin{array}{ccc} z_{-k,k} & \dots & z_{k,k} \\ \vdots & & \vdots \\ z_{-k,-k} & \dots & z_{k,-k} \\ \\ \vdots & & \vdots \\ \vdots & & \vdots \\ \\ x_{-k,k} & \dots & x_{k,k} \\ \vdots & & \vdots \\ x_{-k,-k} & \dots & x_{k,-k} \end{array} \quad \begin{array}{ccc} w_{-k,k} & \dots & w_{k,k} \\ \vdots & & \vdots \\ w_{-k,-k} & \dots & w_{k,-k} \\ \\ \vdots & & \vdots \\ \vdots & & \vdots \\ \\ y_{-k,k} & \dots & y_{k,k} \\ \vdots & & \vdots \\ y_{-k,-k} & \dots & y_{k,-k} \end{array}$$

which are valid for the shift space X_G and hence can be extended to configurations in X_G (say r and s respectively). Finally, note that while $r \in U, s \in V$ (as central blocks of r and s are x_k and y_k respectively), we also have $\sigma^{(0,2k+1+p)}(r) \in Z$ and $\sigma^{(0,2k+1+p)}(s) \in W$ (as their central blocks are z_k and w_k respectively). Consequently, X_G exhibits weak mixing and the proof of forward part is complete.

Conversely, if X_G exhibits vertical weak mixing then it exhibits vertical doubly transitivity and the proof is complete. \square

Remark 8. The above results establish the equivalence of vertical doubly transitivity and vertical weak mixing under the condition $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ for all i, j when H exhibits weak mixing. For stronger notions of mixing in any direction (r, s) , it may be noted that as shift in a particular direction is a one dimensional map and stronger notions of mixing (such as total transitivity, weak mixing and strong mixing) are equivalent for one dimensional shifts, similar arguments establish analogous results for various notions of directional mixing. It may be noted that while any of the mixing notions for one of the generating matrices ensures the same for X_G under $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0$ for all i, j , any of the mixing notions for X_G need not ensure the same for any of the generating matrices under the conditions imposed. We now give an example in support of our claim.

Example 6. Let X_G be a shift space arising from following adjacency matrices:

$$H = \begin{array}{c} \\ 0 \\ 1 \\ 2 \\ 3 \end{array} \begin{array}{cccc} & 0 & 1 & 2 & 3 \\ \left(\begin{array}{cccc} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 \end{array} \right) \end{array} \quad V = \begin{array}{c} \\ 0 \\ 1 \\ 2 \\ 3 \end{array} \begin{array}{cccc} & 0 & 1 & 2 & 3 \\ \left(\begin{array}{cccc} 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{array} \right) \end{array}$$

It may be noted that H and V are not irreducible. However, as $(HV)_{ij} \neq 0 \Leftrightarrow (VH)_{ij} \neq 0 \forall i, j$ holds and HV is primitive, the shift space X_G exhibits $(1, 1)$ -weak mixing. Consequently, weak mixing (directional weak mixing) for X_G need not ensure weak mixing (any form of mixing) for any of the generating matrices.

We now turn our attention to the graph shifts arising from graph products between two-dimensional graphs $G_i = (H_i, V_i)$. It may be noted that if a two-dimensional graph G can be expressed as a graph product of some two-dimensional graphs, then dynamics of shift space X_G can be investigated through graphs of reduced complexity and hence provides a more efficient way of investigating the dynamics of the shift space under discussion. Before we move further, we first provide an example in support of our claim.

Example 7. Let G be a two-dimensional graph generated by matrices

$$H = \begin{matrix} & a & b & c & d & e & f \\ \begin{matrix} a \\ b \\ c \\ d \\ e \\ f \end{matrix} & \begin{pmatrix} 1 & 1 & 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \end{matrix} \quad V = \begin{matrix} & a & b & c & d & e & f \\ \begin{matrix} a \\ b \\ c \\ d \\ e \\ f \end{matrix} & \begin{pmatrix} 0 & 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \end{matrix}$$

Next, consider matrices

$$H_1 = \begin{matrix} & 0 & 1 & 2 \\ \begin{matrix} 0 \\ 1 \\ 2 \end{matrix} & \begin{pmatrix} 1 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \quad V_1 = \begin{matrix} & 0 & 1 & 2 \\ \begin{matrix} 0 \\ 1 \\ 2 \end{matrix} & \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}$$

and

$$H_2 = \begin{matrix} & 3 & 4 \\ \begin{matrix} 3 \\ 4 \end{matrix} & \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}, \quad V_2 = \begin{matrix} & 3 & 4 \\ \begin{matrix} 3 \\ 4 \end{matrix} & \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}$$

It can be observed that \exists indices i, j such that $(HV)_{ij} = 0$ but $(VH)_{ij} \neq 0$ and \exists indices k, l such that $(VH)_{kl} = 0$ but $(HV)_{kl} \neq 0$. However, as $H = H_1 \times H_2$ and $V = V_1 \times V_2$, the two-dimensional graph $G = (H, V)$ can be expressed as the Tensor product $G_1 \times G_2$, where $G_i = (H_i, V_i)$ for $i \in \{1, 2\}$. Consequently, X_G is conjugate to $X_{G_1 \times G_2}$ and thus the qualitative nature of the shift space can be investigated through dynamics of "much simpler spaces" X_{G_1} and X_{G_2} . In particular, it can be seen that X_{G_1} and X_{G_2} are non-empty, have horizontal (vertical) periodic points and are $(1, 1)$ -transitive. Consequently, the shift space X_G is indeed non-empty, has horizontal (vertical) periodic points and is also $(1, 1)$ -transitive. Consequently, the dynamics of the shift space X_G can be investigated through dynamics of "much simpler spaces" X_{G_1} and X_{G_2} .

Proposition 9. For any given graph $G = (V, E)$ with n vertices, G can be expressed as a Tensor product of two graphs if and only if there exists $r_1, r_2 \in \mathbb{N}$ such that $n = r_1 r_2$, a $r_2 \times r_2$ matrix B and a permutation $\{v_{\sigma(1)}, v_{\sigma(2)}, \dots, v_{\sigma(n)}\}$ of the vertex set V such that adjacency matrix A_G (with respect to $\{v_{\sigma(1)}, v_{\sigma(2)}, \dots, v_{\sigma(n)}\}$) is of the form

$$\begin{pmatrix} S_{11} & S_{12} & \dots & S_{1r_2} \\ S_{21} & S_{22} & \dots & S_{2r_2} \\ \vdots & \vdots & \ddots & \vdots \\ S_{r_1 1} & S_{r_1 2} & \dots & S_{r_1 r_2} \end{pmatrix} \quad \text{where } S_{ij} \text{ are either } B \text{ or } 0 \text{ (the zero matrix)}$$

Proof. First, consider that graph G can be expressed as a Tensor product of two graphs G_1, G_2 , where $\mathcal{V}(G_1) = \{a_1, a_2 \dots a_{r_1}\}$ and $\mathcal{V}(G_2) = \{b_1, b_2 \dots b_{r_2}\}$. Then, if the vertex set for Tensor product is ordered as $(a_1, b_1), (a_1, b_2) \dots (a_1, b_{r_2}), (a_2, b_1), (a_2, b_2) \dots (a_2, b_{r_2}) \dots (a_{r_1}, b_1), (a_{r_1}, b_2) \dots (a_{r_1}, b_{r_2})$, as $(a_i, b_j)(a_k, b_l) \in \mathcal{E}(G_1 \times G_2)$ if and only if $a_i a_k \in \mathcal{E}(G_1)$ and $b_j b_l \in \mathcal{E}(G_2)$, the adjacency matrix for the Tensor product is given by

$$\begin{pmatrix} S_{11} & S_{12} & \dots & S_{1r_2} \\ S_{21} & S_{22} & \dots & S_{2r_2} \\ \vdots & \vdots & \vdots & \vdots \\ S_{r_1 1} & S_{r_1 2} & \dots & S_{r_1 r_2} \end{pmatrix} \quad \text{where } S_{ij} = A_{G_2} \text{ if } a_i \text{ and } a_j \text{ are adjacent}$$

in G_1 (and 0 otherwise). As G is the Tensor product of G_1 and G_2 , $A_G = A_{G_1 \times G_2}$ and hence the proof of forward part is complete.

Conversely, let the adjacency matrix of G be of the form

$$\begin{pmatrix} S_{11} & S_{12} & \dots & S_{1r_2} \\ S_{21} & S_{22} & \dots & S_{2r_2} \\ \vdots & \vdots & \vdots & \vdots \\ S_{r_1 1} & S_{r_1 2} & \dots & S_{r_1 r_2} \end{pmatrix} \quad \text{where } S_{ij} \text{ are either } B \text{ or } 0 \text{ (the zero}$$

matrix). Define $G_1 = (V_1, E_1)$, where $V_1 = \{a_1, a_2, \dots a_{r_1}\}$ and $(a_i a_j) \in E_1$ if $B_{ij} = B$ and $G_2 = (V_2, E_2)$ where $V_2 = \{b_1, b_2, \dots, b_{r_2}\}$ and E_2 be defined such that $A_{G_2} = B$. Then, as $n = r_1 r_2$ and $A_G = A_{G_1 \times G_2}$, the converse holds and the proof is complete. \square

Remark 9. It may be noted that for any graph G , $G = G_1 \times G_2$ where $G_1 = G$ and $G_2 = (V_2, E_2)$ with $V_2 = \{v_2\}$ is a singleton and E_2 consists of a self loop at v_2 . Consequently, any graph G can be written as a Tensor product of two graphs. However, such a factorization does not reduce the computational complexity of the shift space induced by G and the reduction of the computational complexity indeed depends on the structure of the adjacency matrix of the underlying graph. Further, for given graphs $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$, note that $(a_i, b_j)(a_k, b_l) \in \mathcal{E}(G_1 \square G_2)$ if and only if $a_i a_k \in \mathcal{E}(G_1)$ ($a_i = a_k$) and $b_j = b_l$ ($b_j b_l \in \mathcal{E}(G_2)$). Further, if $\mathcal{V}(G_1) = \{a_1, a_2 \dots a_n\}$, $\mathcal{V}(G_2) = \{b_1, b_2 \dots b_m\}$ and the adjacency matrix of $(G_1 \square G_2)$ is indexed using natural order then for $a_i \neq a_j$, while $a_i a_j \notin \mathcal{E}(G_1)$ forces the corresponding $m \times m$ block to be zero matrix, $a_i a_j \in \mathcal{E}(G_1)$ forces the corresponding $m \times m$ block to be an $m \times m$ identity matrix (for $a_i \neq a_j$). Finally, while $a_i = a_j$ then, $a_i a_i \notin \mathcal{E}(G_1)$ forces the corresponding $m \times m$ block to be B , and $a_i a_i \in \mathcal{E}(G_1)$ forces the corresponding block to be $B_{ii} = B * I$ (where

B is a binary matrix and $*$ is a matrix operation such that $(P * Q)_{ij} = \max(P_{ij}, Q_{ij})$. Consequently, the adjacency matrix of the Cartesian product of graphs can be characterized using the corresponding block representation and we get the following corollary.

Corollary 6. *For any given graph $G = (V, E)$ with n vertices, G can be expressed as a Cartesian product of two graphs if and only if there exists $r_1, r_2 \in \mathbb{N}$ such that $n = r_1 r_2$, a $r_2 \times r_2$ matrix B and a permutation $\{v_{\sigma(1)}, v_{\sigma(2)}, \dots, v_{\sigma(n)}\}$ of the vertex set V such that adjacency matrix A_G (with respect to $\{v_{\sigma(1)}, v_{\sigma(2)}, \dots, v_{\sigma(n)}\}$) is of the form*

$$\begin{pmatrix} B_{11} & B_{12} & \dots & B_{1r_2} \\ B_{21} & B_{22} & \dots & B_{2r_2} \\ \vdots & \vdots & \vdots & \vdots \\ B_{r_1 1} & B_{r_1 2} & \dots & B_{r_1 r_2} \end{pmatrix} \quad \text{where } B_{ij} = \begin{cases} 0 \text{ or } I, & i \neq j; \\ B \text{ or } (B * I) & i = j; \end{cases}$$

Proof. The proof follows from the discussion in Remark 9. □

REFERENCES

- [1] **A. Quas, P. Trow**, Subshifts of Multidimensional shifts of finite type, *Ergodic Theory and Dynamical Systems*, 20 (03), 2000, 859 - 874.
- [2] **Bruce P. Kitchens**, Symbolic Dynamics: One-Sided, Two-Sided and Countable State Markov Shifts, *Universitext. Springer-Verlag*, Berlin Heidelberg, 1998.
- [3] **C.E. Shannon**, A mathematical theory of communication, *The Bell System Technical Journal*, vol. 27, no. 3, 1948, pp. 379-423.
- [4] **Coven, E., Johnson, A., Jonoska, N., and Madden, K.**, The Symbolic Dynamics Of Multidimensional Tiling Systems, *Ergodic Theory And Dynamical Systems*, 23(2), 2003, pp. 447-460.
- [5] **D. Lind, B. Marcus**, An Introduction to Symbolic Dynamics and Coding, *Cambridge University Press*, Cambridge, 1995.
- [6] **Gert Sabidussi**, Graph Multiplication, *Mathematische Zeitschrift*, vol. 72, 1959/60, pp. 446-457.
- [7] **Izhikevich Eugene M.**, Dynamical Systems in Neuroscience : The Geometry of Excitability and Bursting, *Cambridge Mass: MIT Press.*, 2007.
- [8] **Jan De Vries**, Topological Dynamical system: An Introduction to the Dynamics of Continuous Mappings. *De Gruyter*, Berlin, Boston, 2014.
- [9] **M. Hochman**, On the automorphism groups of multidimensional shifts of finite type, *Ergodic Theory and Dynamical Systems*, 30(3), 2010, pp. 809-840, doi:10.1017/S0143385709000248.
- [10] **Markley, Nelson G. and Paul, Michael E.**, Matrix Subshifts for \mathbb{Z}^v Symbolic Dynamics, *Proceedings of the London Mathematical Society*, Volume s3-43, Issue 2, September 1981, pp. 251-272.
- [11] **Markley, Nelson G. and Paul, Michael E.**, Maximal Measures and Entropy for \mathbb{Z}^v Subshifts of Finite Type, *Classical Mechanics and Dynamical Systems*, 1981, pp. 135-157.

- [12] **Michael Hochman**, On Dynamics and Recursive Properties of Multidimensional Symbolic Dynamics, *Inventiones Mathematica*, Volume 176, Number 1, April, 2009.
- [13] **Michael Hochman, Tom Meyerovitch**, A Characterization of the Entropies of Multidimensional Shifts of Finite type, *Annals of Mathematics*, Vol. 171 (2010), No. 3, 2011-2038.
- [14] **Mike Boyle, Ronnie Pavlov, and Michael Schraudner**, Multidimensional Sofic Shifts without Separation and their Factors, *Transactions of the American Mathematical Society*, Volume 362, Number 9, September 2010, Pages 4617-4653.
- [15] **Morse, Marston, and Gustav A. Hedlund**, Symbolic Dynamics II. Sturmian Trajectories, *American Journal of Mathematics*, vol. 62, no. 1, 1940, pp. 1–42.
- [16] **Natasa Jonoska, Joni Burnette Pirnot**, Transitivity in two-dimensional Local Languages Defined by Dot Systems, *International Journal of Foundations of Computer Science*, vol. 17, no. 2, 2006, pp. 435-463, <https://doi.org/10.1142/S0129054106003917>.
- [17] **Paul M. Weichsel**, The Kronecker product of graphs, *Proceedings of the American Mathematical Society*, vol 13 (1962), pp. 47-52.
- [18] **Prashant Kumar and Puneet Sharma**, On Graph Induced Symbolic Systems, *Appl. Gen. Topol.*, vol. 24, no. 2, pp. 359–378, Oct. 2023.
- [19] **Puneet Sharma and Dileep Kumar** Matrix characterization of multidimensional subshifts of finite type, *Appl. Gen. Topol.*, vol. 20, no. 2, pp. 407–418, Oct. 2019.
- [20] **Puneet Sharma and Dileep Kumar** Multidimensional Shifts and Finite Matrices, *Topology Proceedings*, vol. 57, pp. 241–257, 2021.
- [21] **R. Berger** The undecidability of the Domino Problem, *Mem. Amer. Math. Soc.*, 66, 1966.
- [22] **Robinson, R.M**, Undecidability and nonperiodicity for tilings of the plane, *Inventiones mathematicae*, 12, 1971, pp. 177-209.
- [23] **S.Lightwood**, Morphisms from non-periodic Z^2 -subshifts I: Constructing embeddings from homomorphisms, *Ergodic Theory Dynamical Systems*, 23 (2), 2003, pp. 587-609.
- [24] **Song M, Ouyang Z, Liu ZL**, Discrete dynamical system modelling for gene regulatory networks of 5-hydroxymethylfurfural tolerance for ethanologenic yeast, *IET Syst Biol.*, 2009 May;3(3):203-18. doi: 10.1049/iet-syb.2008.0089.
- [25] **T. K. Subrahmonian Moothathu**, Stronger forms of sensitivity for dynamical systems, *Nonlinearity*, vol. 20, no. 9, 2007, pp. 2115 -2126.
- [26] **Ted Janssen and Marc de Boissieu**, Dynamics of quasicrystals, *Comptes Rendus Physique*, vol. 15, no. 1, 2014, pp. 58-69.
- [27] **Xin-Chu Fu, WeiPing Lu, Peter Ashwin, Jinqiao Duan**, Symbolic Representations of Iterated Maps, *Topological Methods in Nonlinear Analysis*, volume 18, no. 1, 2001, pp. 119-147.

DEPARTMENT OF MATHEMATICS, I.I.T. JODHPUR, NH 65, NAGAU ROAD, KARWAR, JODHPUR-342030, INDIA

Email address: puneet@iitj.ac.in, kumar.48@iitj.ac.in