

Detection of X-ray polarization in the high synchrotron peaked blazar 1ES 1959+650

Athira M. Bharathan,^{1,*} C. S. Stalin,^{2,†} Rwitika Chatterjee,³ S. Sahayanathan,⁴ Indrani Pal,² Blesson Mathew,¹ and Vivek K. Agrawal⁵

¹Department of Physics, CHRIST (Deemed to be University), Bangalore, India

²Indian Institute of Astrophysics, Block II, Koramangala, Bangalore 560 034 India

³Space Astronomy Group, ISITE Campus, U. R. Rao Satellite Centre, Bangalore 560 037, India

⁴Astrophysical Sciences Division, Bhabha Atomic Research Centre, Mumbai-400085, India

Homi Bhabha National Institute, Mumbai-400094, India

⁵Space Astronomy Group, ISITE Campus, U. R. Rao Satellite Centre, Bangalore 560 037, India

(Dated: November 6, 2023)

We report the first time measurement of X-ray polarization in the 2–8 keV band from the high synchrotron peaked (HSP) blazar 1ES 1959+650. The low energy hump in the broadband spectral energy distribution (SED) of blazars in the $\log\nu$ versus νF_ν plane is believed to be due to the synchrotron emission process from relativistic particles in their jets. In HSP blazars, the observed X-ray emission lies in the high energy tail of the synchrotron part of the SED, and X-ray polarization measurements can constrain the processes by which particles are accelerated in their jets. We present here our results on the X-ray polarization characteristics of 1ES 1959+650 using observations carried out by the *Imaging X-ray Polarimetry Explorer (IXPE)*. *IXPE* observed this source for four epochs between May 2022 and August 2023, with observation time ranging from about 50 to 300 ksec. Of the four epochs of observations, we detected polarization on two epochs, significantly larger than the minimum detectable polarization values. From model-independent analysis during the observations on 28 October 2022, in the 2–8 keV band, we found the degree of polarization of $\Pi_X = 9.0 \pm 1.6$ % and an electric vector position angle of $\psi_X = 53 \pm 5$ deg. Similarly, from the observations on 14 August 2023, we found Π_X and ψ_X values as 12.5 ± 0.7 % and 20 ± 2 deg respectively in the 2–8 keV band. These values obtained from model-independent analysis are also in agreement with the values obtained from spectro-polarimetric analysis of the I,Q, and U spectra in the 2–8 keV range. The measured X-ray polarization is larger than that known in the optical that ranges between 2.5–9% during the period 2008 to 2018. This result can be inferred as the shock acceleration of the particles in the jet of 1ES 1959+650.

I. INTRODUCTION

Blazars are a peculiar category of active galactic nuclei (AGN) that are powered by the accretion of matter onto super massive black holes (SMBHs $10^6 - 10^{10} M_\odot$) residing at the centers of galaxies [1–5]. These blazars with their relativistic jets oriented close to the line of sight to the observer emit over the entire electromagnetic spectrum from the low energy radio to the high energy γ -rays [6]. The emission from them is dominated by the boosted non-thermal emission from particles in their relativistic jets, however, have sub-dominant thermal emission from the accretion disk and emission lines from their broad line region (BLR) [7].

The broadband spectral energy distribution (SED) of these sources shows a double hump structure. The low energy hump that peaks between the UV and X-ray frequencies is attributed to synchrotron emission process from relativistic electrons that gyrate around the magnetic field in their relativistic jets. The high energy hump peaks in the GeV to TeV energy range and the physical process that gives rise to the high energy hump is debated. Both leptonic and hadronic models are proposed

to explain the high energy emission process in blazars, however, we still do not understand the origin of high energy emission in them.

In the leptonic model of emission from blazar jets, the high energy emission is attributed to inverse Compton scattering of low energy photons by relativistic electrons. The seed photons can be the same synchrotron photons produced by the electrons via the synchrotron mechanism a process called synchrotron self Compton [8, 9]. Alternatively, the seed photons can originate from external to the jet such as the accretion disk, the BLR and the torus a process called the external Compton [10, 11]. Alternatively, in the hadronic scenario, the high energy emission can be due to synchrotron radiation by relativistic protons and photon-pion production [12–14]. Based on the peak frequency (ν_p) of the synchrotron component of the broadband SED, blazars are further classified in low synchrotron peaked (LSP; $\nu_p < 10^{15}$ Hz), intermediate synchrotron peaked (ISP; $10^{14} < \nu_p < 10^{15}$ Hz) and high synchrotron peaked (HSP; $\nu_p > 10^{15}$ Hz) blazars.

Leptonic, hadronic, and/or lepto-hadronic models are able to provide a good fit to broadband SED of blazars [14, 15]. A key observable to distinguish among these two models to explain the broadband SED is via flux variability observations across wavelengths. Ideally, correlated optical and γ -ray flux variations will favor a leptonic scenario, while a γ -ray flare without a counterpart at lower energies in the optical/IR cannot be produced in

* athira.bharathan@res.christuniversity.in

† stalin@iiap.res.in

the one zone leptonic scenario. However, observationally the correlations between optical and γ -ray flux variations are found to be complex [16–18]. Another observable that can constrain these two models is X-ray polarization [19, 20]. Also, X-ray polarization can provide strong diagnostics of the particle acceleration processes [21].

Optical polarization is a characteristic of blazars and monitoring observations have found polarization variations, and these polarization changes are often correlated with flares at other wavelengths [22–27]. Polarization observations can thus be the key to the emission processes in the central regions of blazars as well as provide the diagnosis of the orientation of the magnetic field in the jet. Though blazars have been studied extensively for polarization in the optical and radio regimes, we have no knowledge of their X-ray polarization characteristics till the launch of the *Imaging X-ray Polarimetry Explorer (IXPE)* [28] in December 2021. Since its launch, IXPE has observed both Seyfert type AGN as well as blazars. As of today, X-ray polarization characteristic is known for a dozen blazars, namely, Mrk 421 [29, 30], 1ES 0229+200 [31], PG 1553+113 [32], BL Lacertae [33, 34], Mrk 501 [35], 3C 273, 3C 279, 3C 454.3 and S5 0716+714 [36].

1ES 1959+650, the BL Lac object at a redshift $z = 0.048$ [37] is a HSP blazar wherein the synchrotron peak in its broadband SED falls in the UV to X-ray region [38, 39]. It was first detected in the radio band with the Green Bank Telescope [40]. In the X-rays it was first detected in the Einstein Slew Survey [41] and later in BeppoSAX, *Swift* and XMM-Newton [42, 43] It is also detected in GeV-TeV energies [44, 45]. This source was found to show uncorrelated flux variations between X-rays and TeV γ -rays [38, 46], which is inconsistent with the simple one zone synchrotron self Compton (SSC) models. Alternatively, such uncorrelated flux variations can be explained by multi-zone SSC mechanism [47], external (EC) mechanism [38] or hadronic processes [48]. 1ES 1959+650 is observed by IXPE for four epochs between May 2022 and August 2023 and in this work we report the first time detection of X-ray polarization in 1ES 1959+650. The paper is organized as follows, In Section 2, we described the observations and reduction of data from IXPE, the X-ray polarization properties are discussed in Section 3 and the results are discussed in Section 4, followed by the summary in the final Section.

II. OBSERVATIONS AND DATA REDUCTION

IXPE observed 1ES 1959+650 on four epochs between May 2022 and August 2023 with its three detector units (DU) for a net exposure ranging between ~ 50 and ~ 300 ksec. The log of the IXPE observation is given in Table I. We used the cleaned and calibrated level 2 data for the scientific analysis.

We analyzed the publicly available level 2 data using IXPEOBSSIM software v30.0.0 [49]. We generated a count

TABLE I. Log of IXPE observation. The details are the observational ID (OBSID), Date of observation and the exposure time.

OBSID	Date	Exposure Time (secs)
01006201	03 May 2022	53519
01006001	09 June 2022	200432
02004801	28 October 2022	194844
02240801	14 August 2023	312477

map in sky coordinates using the CMAP algorithm within the `xpbinview` task. We adopted a circular region with a radius of $70''$ for the source extraction from the three DUs, and a source-free region with a radius of $100''$ for the background extraction for each DU. We then used the `xpselect` task to generate the filtered source and background regions for the polarimetric analysis. For spectropolarimetric analysis, we generated the I, Q and U source and background spectra using the PHA1, PHA1Q and PHA1U algorithm using `xpbin` task within IXPEOBSSIM for the three DUs.

III. ANALYSIS

A. Polarimetry

We analyzed the polarimetric signal from 1ES 1959+650 for all the four OBSIDs using PCUBE algorithm in the `xpbinview` task. We generated the three polarization cubes for the three DUs to extract information such as the Stokes parameters (I, Q, U); the minimum detectable polarization (MDP); the polarization degree (Π_X); the polarization angle (Ψ_X) and their associated errors. We first generated the three polarization cubes corresponding to three DUs in the entire 2–8 keV energy band. The combined polarization parameters from the three DUs in the 2–3 keV, 3–5 keV, 5–8 keV and 2–8 keV bands are given in Table III. Of the four OBSIDs we detected significant polarization on two epochs, namely, 28 October 2022 and 14 August 2023. In these two epochs, the measured polarization is beyond the MDP values. The normalized U/I and Q/I Stokes parameters obtained from the combined cube for the observations of all the epochs are shown in Fig. 1.

To check for the energy dependence of the polarization parameters, we also derived the polarization parameters in three energy bins of 2–3 keV, 3–5 keV and 5–8 keV using the PCUBE algorithm. The derived parameters are given in Table II and Table III. The polarization values for the total energy range of 2–8 keV as well as for the different energy ranges are given in Fig. 2 for both the epochs where significant polarization is detected. From Table II and Table III it is evident polarization is not detected in 1ES 1959+650 during observations carried out on May 2022 and June 2022. However, polarization was significantly detected in the later two epochs of observa-




FIG. 1. *IXPE* I, Q, U Stokes best-fit spectra with residuals in the 2–8 keV band for the observation of 14 August 2023 (left) and normalized U/I and Q/I Stokes parameter in the total 2–8 keV band of *IXPE* for 03 May 2022 (orange triangle), 09 June 2022 (green asterisk), 28 October 2022 (orange circle), and 14 August 2023 (green square)(right).

tions were carried out on 28 October 2022 and 14 August 2023. On the observations of 28 October 2022, Π_X and ψ_X are found to be similar within errors in the energy range of 2–3, 3–5 and the 2–8 keV. However, from the observations of 14 August 2023, there is a tendency for an increase in Π_X from the lower energy range to the higher energy range, with ψ_X remaining constant at ~ 20 deg between different energy bands.

B. Spectro-polarimetry

Of the four epochs of observations, X-ray polarization was detected on two epochs from model independent analysis. For those two epochs, we carried out the spectro-polarimetric analysis of the *IXPE* I, Q, U spectra in the 2–8 keV energy band. For the spectral fitting, we used an absorbed *powerlaw*, modified by a multiplicative constant polarization model *polconstant* in XSPEC V [50]. This model assumes a constant polarization degree and a constant polarization angle over the specific energy band. In XSPEC the model takes the following form,

$$\text{constant} \times TBabs \times (\text{polconst} \times \text{po}) \quad (1)$$

Here, the *const* represents the inter-calibration constant for each detector. *TBabs* was used to model the Milky Way Galactic hydrogen column density, which was taken from [51]. We fixed the column density (N_H) during the fit. This model fits the I, Q, and U spectra (from the three detectors) well. The best fit I, Q and U spectra along the residuals are given in Fig. 1. All the model parameters for each of the detectors were tied during the

fit. The errors were calculated at the 90% confidence ($\chi^2 = 2.71$ criterion). The best-fit parameters are given in Table IV.

IV. DISCUSSION

Observations with *IXPE* have enabled the detection of X-ray polarization in the HSP blazar 1ES 1959+650. Of the four epochs of observations, that span a duration of about 15 months, significant polarization was detected on two epochs, namely 28 October 2022 and 14 August 2023 separated by about 10 months. On October 2022, in the 2–8 keV band, we measured the polarization of $\Pi_X = 9.0 \pm 1.6\%$ and $\psi_X = 53 \pm 5$ deg, while on 14 August 2023, we measured values of $\Pi_X = 12.5 \pm 0.7\%$ and $\psi_X = 20 \pm 2$ deg. Thus, there is significant variation both in polarization degree and the polarization angle between the two epochs. The high Π_X measured on 14 August 2023, is similar to such high values measured for the blazars 1ES 0229+200 [31] with $\Pi_X = 17.9\%$ along $\psi_X \sim 25$ deg and the first *IXPE* observation of another HSP blazar Mrk 421 [52] with $\Pi_X \sim 15\%$. From optical polarization observations carried out by the Steward Observatory during the period 2008 and 2018, [53] found the degree of optical polarization (Π_o) to range between 2.5 – 9% with a mean value of $\Pi_o = 4.17 \pm 0.12$ and a stable polarization angle. The degree of X-ray polarization is thus higher than the average degree of polarization in the optical band which is similar to that seen in other HSP blazars such as Mrk 501 [35], Mrk 421 [52] and 1ES 0229+2000 [31] based on simultaneous X-ray and optical polarization observations. Such energy dependent polarization with larger polarization degree in




FIG. 2. The position of the measured polarization between Π_X and Ψ_X in the 2–8 keV band (left) and in three different energy bands (right). The top panels are for the observation done on 28 October 2022, while the bottom panels pertain to the observation carried out on 14 August 2023.

TABLE II. The measured polarization parameters in different energy bands.

OBSID	Date	Π_X (%)			MDP (%)				
		2-3 keV	3-5 keV	5-8 keV	2-8 keV	2-3 keV	3-5 keV	5-8 keV	2-8 keV
01006201	03 May 2022	3.7 ± 3.1	9.5 ± 2.8	12.3 ± 6.8	4.6 ± 2.4	9.3	8.6	20.5	7.2
01006001	09 June 2022	2.1 ± 1.5	2.9 ± 1.5	3.5 ± 3.8	2.1 ± 1.2	4.7	4.5	11.4	3.7
20048001	28 October 2022	7.8 ± 1.9	11.7 ± 1.9	8.4 ± 5.2	9.0 ± 1.6	6.0	6.1	15.7	2.1
22508001	14 August 2023	11.8 ± 0.1	12.6 ± 0.9	15.9 ± 2.6	12.5 ± 0.7	2.7	2.8	7.9	2.1

X-rays compared to the mean value of polarization degree in the optical band, suggests that the X-ray emission must be produced by energetic electrons accelerated at shock fronts [54].

The narrow band spectral study of the source in X-rays suggests the X-ray spectral index in the energy range 2–10 keV is ~ 2.197 [55]. Such a steep X-ray spectral index suggests this energy regime falls on the high energy end of the synchrotron component of the SED. Hence, this emission demands high electron energies as compared to the ones responsible for low energy emission at the optical waveband. If we assume these electrons are accelerated at the shock front under the Fermi process, then the high energy electrons populate in close vicinity to the shock front owing to their short cooling timescales. On the other hand, the longer cooling timescale of the low energy electrons lets them diffuse deep into the jet medium from the shock front. The shearing of the plasma at the shock front aligns the magnetic field [56] [35] which will result in strong polarised emission. Consistently, the emissions arising from the close vicinity to the shock front will be more polarised than the ones emerging from farther regions. Our detailed polarization analysis of 1ES 1959+650 suggests, that the X-ray emission during both the epochs of high polarization are significantly larger than its optical counterpart. This result supports that the efficient acceleration of electrons at the shock

front is responsible for the broadband emission during these epochs.

The two epochs of high X-ray polarization observed from 1ES 1959+650 are separated by ~ 10 months with significant variation in Π_X and Ψ_X . Interestingly, the Π_X increased by a factor of ~ 1.4 while the Ψ_X is reduced by ~ 2.5 . The change in polarization angle could possibly be due to the changes in the position angle of the jet with time and such variation in the jet position angle is already known in blazars [57–59]. The high degree of polarization and modified polarization angle may also be associated with strong oblique shocks which are not normal to the jet flow (e.g. collimation shock)[60].

V. CONCLUSIONS

We carried out the analysis of the X-ray polarimetric observations carried out by *IXPE* on 1ES 1959+650. The observations were carried out on four epochs that span from May 2022 to August 2023. The findings of the polarimetric study are as follows,

1. Of the four epochs of observations, we detected significant polarization on two epochs, namely 28 October 2022 and 14 August 2023
2. From a model-independent analysis, we found a

TABLE III. The measured polarization parameters in different energy bands.

OBSID	Date	ψ_X (deg)			
		2-3 keV	3-5 keV	5-8 keV	2-8 keV
01006201	03 May 2022	-69±23	-55±9	62±16	-73±15
01006001	09 June 2022	-53±21	-67±14	-17±31	-53±16
20048001	28 October 2022	56±7	46±5	67±18	53±5
22508001	14 August 2023	20±2	21±2	18±5	20±2

value of $\Pi_X = 9.0 \pm 1.6\%$ and $\Psi_X = 53 \pm 5$ deg in the 2–8 keV band for the observations on 28 October 2022. Similarly, from the observations on 14 August 2023, using model independent analysis, we found $\Pi_X = 12.5 \pm 0.7\%$ and $\psi_X = 20 \pm 2$ deg in the 2–8 keV band. We thus found polarization to vary between the two epochs of observations.

- From spectro-polarimetric analysis of model fits to the 2–8 keV spectra of 28 October 2022, we found values of Π_X and Ψ_X of $9.6^{+0.03}_{-0.1}\%$ and 49^{+39}_{-32} respectively. Similarly, for the observations done on 14 August 2023, we found $\Pi_X = 12.6 \pm 1.1$ and $\psi_X = 20 \pm 2$ deg. Thus, the polarimetric measurements obtained from both model-independent and spectro-polarimetric analyses agree with each other.
- To check for energy dependent polarization, we derived the polarization parameters in different energy bands. For the epoch of 28 October 2022, polarization is significantly detected in the 2–3 keV, 3–5 keV and 2–8 keV bands. Both Π_X and ψ_X are found to be similar within error bars. We thus did not find evidence of changes in polarization between energy bands. However, for the epoch of 14 August 2023, polarization is significantly detected in all energy bands, and while ψ_X is found to be similar among energy bands, Π_X is larger in the 5–8 keV band than that at 2–3 keV band. There is thus energy dependent polarization.
- Though optical polarization observations are not available during the epoch of IXPE observations, the measured X-ray polarization during the epochs of 28 October 2022 and 14 August 2023 is larger than the average optical polarization degree over a period of 10 years between 2008 and 2018. These observations tend to favor a scenario of shock acceleration of electrons that produce the synchrotron optical and X-rays in 1ES 1959+650.

- We found larger Π_X on 14 August 2023, relative to 28 October 2022, however, ψ_X is lower on 14 August 2023 compared to 28 October 2022. This could be associated with the variation in electric vector direction caused by the change in the position angle of the jet with time.

With the results reported in this work, the number of blazars with X-ray polarization measurements from IXPE observations have increased to 10. X-ray polarimetric observations of more blazars simultaneous with observations at other wavelengths such as optical, and radio are needed to better constrain the physical processes close to their central regions.

ACKNOWLEDGMENTS

The Imaging X-ray Polarimetry Explorer (IXPE) is a joint US and Italian mission. The US contribution is supported by the National Aeronautics and Space Administration (NASA) and led and managed by its Marshall Space Flight Center (MSFC), with industry partner Ball Aerospace (contract NNM15AA18C). The Italian contribution is supported by the Italian Space Agency (Agenzia Spaziale Italiana, ASI) through contract ASI-OHBI-2017-12-I.0, agreements ASI-INAF-2017-12-H0 and ASI-INFN-2017.13-H0, and its Space Science Data Center (SSDC) with agreements ASI-INAF-2022-14-HH.0 and ASI-INFN 2021-43-HH.0, and by the Istituto Nazionale di Astrofisica (INAF) and the Istituto Nazionale di Fisica Nucleare (INFN) in Italy. This research used data products provided by the IXPE Team (MSFC, SSDC, INAF, and INFN) and distributed with additional software tools by the High-Energy Astrophysics Science Archive Research Center (HEASARC), at NASA Goddard Space Flight Center (GSFC). Athira M Bharathan acknowledges the Department of Science and Technology (DST) for the INSPIRE FELLOWSHIP (IF200255). Special appreciation is extended to Soumya Gupta for her dedicated assistance.

[1] D. Lynden-Bell, Galactic Nuclei as Collapsed Old Quasars, *Nature (London)* **223**, 690 (1969).
[2] N. I. Shakura and R. A. Sunyaev, Black holes in binary systems. Observational appearance., *Astronomy and As-*

trophysics **24**, 337 (1973).
[3] M. J. Rees, Black Hole Models for Active Galactic Nuclei, *Annual Review of Astron and Astrophys* **22**, 471 (1984).
[4] R. Antonucci, Unified models for active galactic nuclei

TABLE IV. Comparison between the derived parameters via model independent and spectropolarimetric analysis carried out in the 2-8 keV band for the two epochs where polarization is detected. Here, π_x is in percentage, ψ_x is in degrees, Γ is the photon index and N_H is the column density in units of 10^{21}cm^{-2}

OBSID	Date	Model Independent		Spectropolarimetry				
		Π_X (2-8 keV)	ψ_X (2-8 keV)	Π_X (2-8 keV)	ψ_X (2-8 keV)	Γ	N_H	χ^2/dof
20048001	28-10-2022	9.0 \pm 1.6	53 \pm 5	9.6 $^{+0.03}_{-0.1}$	49 $^{+39}_{-32}$	2.45 \pm 0.02	1.01	1.1
22508001	14-08-2023	12.5 \pm 0.7	20 \pm 2	12.6 \pm 1.1	20 \pm 2	2.69 \pm 0.01	1.01	1.3

- and quasars., *Annual Review of Astron and Astrophys* **31**, 473 (1993).
- [5] C. M. Urry and P. Padovani, Unified Schemes for Radio-Loud Active Galactic Nuclei, *Publications of the ASP* **107**, 803 (1995), [arXiv:astro-ph/9506063 \[astro-ph\]](#).
- [6] T. Hovatta and E. Lindfors, Relativistic Jets of Blazars, *New Astronomy Review* **87**, 101541 (2019), [arXiv:2003.06322 \[astro-ph.HE\]](#).
- [7] M. Cerruti, Leptonic and Hadronic Radiative Processes in Supermassive-Black-Hole Jets, *Galaxies* **8**, 72 (2020), [arXiv:2012.13302 \[astro-ph.HE\]](#).
- [8] L. Maraschi, G. Ghisellini, and A. Celotti, A Jet Model for the Gamma-Ray-emitting Blazar 3C 279, *Astrophysical Journal, Letters* **397**, L5 (1992).
- [9] A. Mastichiadis and J. G. Kirk, Variability in the synchrotron self-Compton model of blazar emission., *Astronomy and Astrophysics* **320**, 19 (1997), [arXiv:astro-ph/9610058 \[astro-ph\]](#).
- [10] C. D. Dermer, R. Schlickeiser, and A. Mastichiadis, High-energy gamma radiation from extragalactic radio sources., *Astronomy and Astrophysics* **256**, L27 (1992).
- [11] M. Sikora, M. C. Begelman, and M. J. Rees, Comptonization of Diffuse Ambient Radiation by a Relativistic Jet: The Source of Gamma Rays from Blazars?, *Astrophys. J.* **421**, 153 (1994).
- [12] K. Mannheim and P. L. Biermann, Gamma-ray flaring of 3C 279 : a proton-initiated cascade in the jet ?, *Astronomy and Astrophysics* **253**, L21 (1992).
- [13] K. Mannheim, The proton blazar., *Astronomy and Astrophysics* **269**, 67 (1993), [arXiv:astro-ph/9302006 \[astro-ph\]](#).
- [14] M. Böttcher, A. Reimer, K. Sweeney, and A. Prakash, Leptonic and Hadronic Modeling of Fermi-detected Blazars, *Astrophys. J.* **768**, 54 (2013), [arXiv:1304.0605 \[astro-ph.HE\]](#).
- [15] V. S. Paliya, H. Zhang, M. Böttcher, M. Ajello, A. Domínguez, M. Joshi, D. Hartmann, and C. S. Stalin, Leptonic and Hadronic Modeling of Fermi-LAT Hard Spectrum Quasars and Predictions for High-energy Polarization, *Astrophys. J.* **863**, 98 (2018), [arXiv:1807.02085 \[astro-ph.HE\]](#).
- [16] B. Rajput, C. S. Stalin, S. Sahayanathan, S. Rakshit, and A. K. Mandal, Temporal correlation between the optical and γ -ray flux variations in the blazar 3C 454.3, *Mon. Not. Roy. Astr. Soc.* **486**, 1781 (2019), [arXiv:1903.12637 \[astro-ph.HE\]](#).
- [17] B. Rajput, C. S. Stalin, and S. Sahayanathan, Correlation between optical and γ -ray flux variations in bright flat spectrum radio quasars, *Mon. Not. Roy. Astr. Soc.* **498**, 5128 (2020), [arXiv:2009.02055 \[astro-ph.HE\]](#).
- [18] B. Rajput, Z. Shah, C. S. Stalin, S. Sahayanathan, and S. Rakshit, Correlation between optical and γ -ray flux variations in BL Lacs, *Mon. Not. Roy. Astr. Soc.* **504**, 1772 (2021), [arXiv:2104.00402 \[astro-ph.HE\]](#).
- [19] H. Zhang and M. Böttcher, X-Ray and Gamma-Ray Polarization in Leptonic and Hadronic Jet Models of Blazars, *Astrophys. J.* **774**, 18 (2013), [arXiv:1307.4187 \[astro-ph.HE\]](#).
- [20] A. L. Peirson, I. Liodakis, and R. W. Romani, Testing High-energy Emission Models for Blazars with X-Ray Polarimetry, *Astrophys. J.* **931**, 59 (2022), [arXiv:2204.11803 \[astro-ph.HE\]](#).
- [21] H. Sol and A. Zech, Blazars at Very High Energies: Emission Modelling, *Galaxies* **10**, 105 (2022), [arXiv:2211.03580 \[astro-ph.HE\]](#).
- [22] T. D. Kinman, E. Lamla, and C. A. Wirtanen, The Optical Brightness Variations and Polarization of the Quasi-Stellar Radio Source 3c 446, *Astrophys. J.* **146**, 964 (1966).
- [23] J. R. P. Angel and H. S. Stockman, Optical and infrared polarization of active extragalactic objects, *Annual Review of Astron and Astrophys* **18**, 321 (1980).
- [24] R. Itoh, K. Nalewajko, Y. Fukazawa, M. Uemura, Y. T. Tanaka, K. S. Kawabata, G. M. Madejski, F. K. Schinzel, Y. Kanda, K. Shiki, H. Akitaya, M. Kawabata, Y. Moritani, T. Nakaoka, T. Ohsugi, M. Sasada, K. Takaki, K. Takata, T. Ui, M. Yamanaka, and M. Yoshida, Systematic Study of Gamma-ray-bright Blazars with Optical Polarization and Gamma-Ray Variability, *Astrophys. J.* **833**, 77 (2016), [arXiv:1610.04313 \[astro-ph.HE\]](#).
- [25] S. Rakshit, C. S. Stalin, S. Muneer, S. Neha, and V. S. Paliya, Flux and Polarization Variability of OJ 287 during the Early 2016 Outburst, *Astrophys. J.* **835**, 275 (2017), [arXiv:1612.07464 \[astro-ph.GA\]](#).
- [26] B. Rajput, A. Pandey, C. S. Stalin, and B. Mathew, Study of correlation between optical flux and polarization variations in BL Lac objects, *Mon. Not. Roy. Astr. Soc.* **517**, 3236 (2022), [arXiv:2209.02444 \[astro-ph.HE\]](#).
- [27] A. Pandey, B. Rajput, and C. S. Stalin, Correlation between optical flux and polarization variations in flat-spectrum radio quasars on diverse time-scales, *Mon. Not. Roy. Astr. Soc.* **510**, 1809 (2022), [arXiv:2111.08247 \[astro-ph.HE\]](#).
- [28] M. C. Weisskopf, P. Soffitta, L. Baldini, B. D. Ramsey, S. L. O'Dell, R. W. Romani, G. Matt, W. D. Deinger, W. H. Baumgartner, R. Bellazzini, E. Costa, J. J. Kolodziejczak, L. Latronico, H. L. Marshall, F. Muleri, S. D. Bongiorno, A. Tennant, N. Bucciantini, M. Dovciak, F. Marin, A. Marscher, J. Poutanen, P. Slane, R. Turolla, W. Kalinowski, A. Di Marco, S. Fabiani, M. Minuti, F. La Monaca, M. Pinchera, J. Rankin, C. Sgro', A. Trois, F. Xie, C. Alexander, D. Z. Allen, F. Amici, J. Andersen, A. Antonelli, S. Antoniak, P. Attinà, M. Barbanera, M. Bachetti, R. M. Baggett, J. Bladt, A. Brez, R. Bonino, C. Boree, F. Borotto, S. Breeding, D. Brienza, H. K. Bygott,

- C. Caporale, C. Cardelli, R. Carpentiero, S. Castellano, M. Castronuovo, L. Cavalli, E. Cavazzuti, M. Ceccanti, M. Centrone, S. Citraro, F. D'Amico, E. D'Alba, L. Di Gesu, E. Del Monte, K. L. Dietz, N. Di Lalla, G. D. Persio, D. Dolan, I. Donnarumma, Y. Evangelista, K. Ferrant, R. Ferrazzoli, M. Ferrie, J. Footdale, B. Forsyth, M. Foster, B. Garelick, S. Gunji, E. Gurnee, M. Head, G. Hibbard, S. Johnson, E. Kelly, K. Kilaru, C. Lefevre, S. L. Roy, P. Loffredo, P. Lorenzi, L. Lucchesi, T. Maddox, G. Magazzu, S. Maldera, A. Manfreda, E. Mangraviti, M. Marengo, A. Marrocchesi, F. Massaro, D. Mauger, J. McCracken, M. McEachen, R. Mize, P. Mereu, S. Mitchell, I. Mitsuishi, A. Morbidini, F. Mosti, H. Nasimi, B. Negri, M. Negro, T. Nguyen, I. Nitschke, A. Nuti, M. Onizuka, C. Oppedisano, L. Orsini, D. Osborne, R. Pacheco, A. Paggi, W. Painter, S. D. Pavelitz, C. Pentz, R. Piazzolla, M. Perri, M. Pesce-Rollins, C. Peterson, M. Pilia, A. Profeti, S. Puccetti, J. Ranganathan, A. Ratheesh, L. Reedy, N. Root, A. Rubini, S. Ruswick, J. Sanchez, P. Sarra, F. Santoli, E. Scalise, A. Sciortino, C. Schroeder, T. Seek, K. Sodian, G. Spandre, C. O. Speegle, T. Tamagawa, M. Tardiola, A. Tobia, N. E. Thomas, R. Valerie, M. Vimercati, A. L. Walden, B. Weddendorf, J. Wedmore, D. Welch, D. Zanetti, and F. Zanetti, The Imaging X-Ray Polarimetry Explorer (IXPE): Pre-Launch, *Journal of Astronomical Telescopes, Instruments, and Systems* **8**, 026002 (2022), [arXiv:2112.01269 \[astro-ph.IM\]](#).
- [29] D. E. Kim, L. Di Gesu, I. Liodakis, A. P. Marscher, S. G. Jorstad, R. Midde, H. L. Marshall, L. Pacciani, I. Agudo, F. Tavecchio, N. Cibrario, S. Tugliani, R. Bonino, M. Negro, S. Puccetti, F. Tombesi, E. Costa, I. Donnarumma, P. Soffitta, T. Mizuno, Y. Fukazawa, K. S. Kawabata, T. Nakaoka, M. Uemura, R. Imazawa, M. Sasada, H. Akitaya, F. Josè Aceituno, G. Bonnoli, V. Casanova, I. Myserlis, A. Sievers, E. Angelakis, A. Kraus, W. Y. Cheong, H.-W. Jeong, S. Kang, S.-H. Kim, S.-S. Lee, B. Agís-González, A. Sota, J. Escudero, M. Gurwell, G. K. Keating, R. Rao, P. M. Kouch, E. Lindfors, I. G. Bourbah, S. Kiehlmann, E. Kontopodis, N. Mandarakas, S. Romanopoulos, R. Skalidis, A. Vervelaki, S. S. Savchenko, L. A. Antonelli, M. Bachetti, L. Baldini, W. H. Baumgartner, R. Bellazzini, S. Bianchi, S. D. Bongiorno, A. Brez, N. Bucciantini, F. Capitanio, S. Castellano, E. Cavazzuti, C.-T. Chen, S. Ciprini, A. De Rosa, E. Del Monte, N. Di Lalla, A. Di Marco, V. Doroshenko, M. Dovčiak, S. R. Ehlert, T. Enoto, Y. Evangelista, S. Fabiani, R. Ferrazzoli, J. A. Garcia, S. Gunji, K. Hayashida, J. Heyl, W. Iwakiri, P. Kaaret, V. Karas, F. Kislat, T. Kitaguchi, J. J. Kolodziejczak, H. Krawczynski, F. La Monaca, L. Latronico, S. Maldera, A. Manfreda, F. Marin, A. Marinucci, F. Massaro, G. Matt, I. Mitsuishi, F. Muleri, C. Y. Ng, S. L. O'Dell, N. Omodei, C. Oppedisano, A. Papitto, G. G. Pavlov, A. L. Peirson, M. Perri, M. Pesce-Rollins, P.-O. Petrucci, M. Pilia, A. Possenti, J. Poutanen, B. D. Ramsey, J. Rankin, A. Ratheesh, O. Roberts, R. W. Romani, C. Sgrò, P. Slane, G. Spandre, D. Swartz, T. Tamagawa, R. Taverna, Y. Tawara, A. F. Tennant, N. E. Thomas, A. Trois, S. S. Tsygankov, R. Turolla, J. Vink, M. C. Weisskopf, K. Wu, F. Xie, and S. Zane, Magnetic Field Properties inside the Jet of Mrk 421: Multiwavelength Polarimetry Including the Imaging X-ray Polarimetry Explorer, [arXiv e-prints](#), [arXiv:2310.06097 \(2023\)](#), [arXiv:2310.06097 \[astro-ph.HE\]](#).
- [30] L. Di Gesu, H. L. Marshall, S. R. Ehlert, D. E. Kim, I. Donnarumma, F. Tavecchio, I. Liodakis, S. Kiehlmann, I. Agudo, S. G. Jorstad, F. Muleri, A. P. Marscher, S. Puccetti, R. Middei, M. Perri, L. Pacciani, M. Negro, R. W. Romani, A. Di Marco, D. Blinov, I. G. Bourbah, E. Kontopodis, N. Mandarakas, S. Romanopoulos, R. Skalidis, A. Vervelaki, C. Casadio, J. Escudero, I. Myserlis, M. A. Gurwell, R. Rao, G. K. Keating, P. M. Kouch, E. Lindfors, F. J. Aceituno, M. I. Bernardos, G. Bonnoli, V. Casanova, M. García-Comas, B. Agís-González, C. Husillos, A. Marchini, A. Sota, R. Imazawa, M. Sasada, Y. Fukazawa, K. S. Kawabata, M. Uemura, T. Mizuno, T. Nakaoka, H. Akitaya, S. S. Savchenko, A. A. Vasilyev, J. L. Gómez, L. A. Antonelli, T. Barnouin, R. Bonino, E. Cavazzuti, L. Costamante, C.-T. Chen, N. Cibrario, A. De Rosa, F. Di Pierro, M. Errando, P. Kaaret, V. Karas, H. Krawczynski, L. Lisalda, G. Madejski, C. Malacaria, F. Marin, A. Marinucci, F. Massaro, G. Matt, I. Mitsuishi, S. L. O'Dell, A. Paggi, A. L. Peirson, P.-O. Petrucci, B. D. Ramsey, A. F. Tennant, K. Wu, M. Bachetti, L. Baldini, W. H. Baumgartner, R. Bellazzini, S. Bianchi, S. D. Bongiorno, A. Brez, N. Bucciantini, F. Capitanio, S. Castellano, S. Ciprini, E. Costa, E. Del Monte, N. Di Lalla, V. Doroshenko, M. Dovčiak, T. Enoto, Y. Evangelista, S. Fabiani, R. Ferrazzoli, J. A. Garcia, S. Gunji, K. Hayashida, J. Heyl, W. Iwakiri, F. Kislat, T. Kitaguchi, J. J. Kolodziejczak, F. La Monaca, L. Latronico, S. Maldera, A. Manfreda, C. Y. Ng, N. Omodei, C. Oppedisano, A. Papitto, G. G. Pavlov, M. Pesce-Rollins, M. Pilia, A. Possenti, J. Poutanen, J. Rankin, A. Ratheesh, O. J. Roberts, C. Sgrò, P. Slane, P. Soffitta, G. Spandre, D. A. Swartz, T. Tamagawa, R. Taverna, Y. Tawara, N. E. Thomas, F. Tombesi, A. Trois, S. S. Tsygankov, R. Turolla, J. Vink, M. C. Weisskopf, F. Xie, and S. Zane, Discovery of X-ray polarization angle rotation in the jet from blazar Mrk 421, *Nature Astronomy* **7**, 1245 (2023), [arXiv:2305.13497 \[astro-ph.HE\]](#).
- [31] S. R. Ehlert, I. Liodakis, R. Middei, A. P. Marscher, F. Tavecchio, I. Agudo, P. M. Kouch, E. Lindfors, K. Nilsson, I. Myserlis, M. Gurwell, R. Rao, F. J. Aceituno, G. Bonnoli, V. Casanova, B. Agíz-González, J. Escudero, J. O. Santos, A. Sota, E. Angelakis, A. Kraus, G. K. Keating, L. A. Antonelli, M. Bachetti, L. Baldini, W. H. Baumgartner, R. Bellazzini, S. Bianchi, S. D. Bongiorno, R. Bonino, A. Brez, N. Bucciantini, F. Capitanio, S. Castellano, E. Cavazzuti, C.-T. Chen, S. Ciprini, E. Costa, A. De Rosa, E. Del Monte, L. Di Gesu, N. Di Lalla, A. Di Marco, I. Donnarumma, V. Doroshenko, M. Dovčiak, T. Enoto, Y. Evangelista, S. Fabiani, R. Ferrazzoli, J. A. Garcia, S. Gunji, K. Hayashida, J. Heyl, W. Iwakiri, S. G. Jorstad, P. Kaaret, V. Karas, F. Kislat, T. Kitaguchi, J. J. Kolodziejczak, H. Krawczynski, F. La Monaca, L. Latronico, S. Maldera, A. Manfreda, F. Marin, A. Marinucci, H. L. Marshall, F. Massaro, G. Matt, I. Mitsuishi, T. Mizuno, F. Muleri, M. Negro, C. Y. Ng, S. L. O'Dell, N. Omodei, C. Oppedisano, A. Papitto, G. G. Pavlov, A. L. Peirson, M. Perri, M. Pesce-Rollins, P.-O. Petrucci, M. Pilia, A. Possenti, J. Poutanen, S. Puccetti, B. D. Ramsey, J. Rankin, A. Ratheesh, O. J. Roberts, R. W. Romani, C. Sgrò, P. Slane, P. Soffitta, G. Spandre, D. A. Swartz, T. Tamagawa, R. Taverna,

- Y. Tawara, A. F. Tennant, N. E. Thomas, F. Tombesi, A. Trois, S. S. Tsygankov, R. Turolla, J. Vink, M. C. Weisskopf, K. Wu, F. Xie, and S. Zane, X-ray Polarization of the BL Lac Type Blazar 1ES 0229+200, [arXiv e-prints](#), [arXiv:2310.01635](#) (2023), [arXiv:2310.01635 \[astro-ph.HE\]](#).
- [32] R. Middei, M. Perri, S. Puccetti, I. Liodakis, L. Di Gesu, A. P. Marscher, N. Rodriguez Cavero, F. Tavecchio, I. Donnarumma, M. Laurenti, S. G. Jorstad, I. Agudo, H. L. Marshall, L. Pacciani, D. E. Kim, F. J. Aceituno, G. Bonnoli, V. Casanova, B. Agís-González, A. Sota, C. Casadio, J. Escudero, I. Myserlis, A. Sievers, P. M. Kouch, E. Lindfors, M. Gurwell, G. K. Keating, R. Rao, S. Kang, S.-S. Lee, S.-H. Kim, W. Y. Cheong, H.-W. Jeong, E. Angelakis, A. Kraus, L. A. Antonelli, M. Bachetti, L. Baldini, W. H. Baumgartner, R. Bellazzini, S. Bianchi, S. D. Bongiorno, R. Bonino, A. Brez, N. Bucciantini, F. Capitanio, S. Castellano, E. Cavazzuti, C.-T. Chen, S. Ciprini, E. Costa, A. De Rosa, E. Del Monte, N. Di Lalla, A. Di Marco, V. Doroshenko, M. Dovčiak, S. R. Ehlert, T. Enoto, Y. Evangelista, S. Fabiani, R. Ferrazzoli, J. A. García, S. Gunji, K. Hayashida, J. Heyl, W. Iwakiri, P. Kaaret, V. Karas, F. Kislak, T. Kitaguchi, J. J. Kolodziejczak, H. Krawczynski, F. La Monaca, L. Latronico, S. Maldera, A. Manfreda, F. Marin, A. Marinucci, F. Massaro, G. Matt, I. Mitsuishi, T. Mizuno, F. Muleri, M. Negro, C.-Y. Ng, S. L. O'Dell, N. Omodei, C. Oppedisano, A. Papitto, G. G. Pavlov, A. L. Peirson, M. Pesce-Rollins, P.-O. Petrucci, M. Pilia, A. Possenti, J. Poutanen, B. D. Ramsey, J. Rankin, A. Ratheesh, O. J. Roberts, R. W. Romani, C. Sgrò, P. Slane, P. Soffitta, G. Spandre, D. A. Swartz, T. Tamagawa, R. Taverna, Y. Tawara, A. F. Tennant, N. E. Thomas, F. Tombesi, A. Trois, S. S. Tsygankov, R. Turolla, J. Vink, M. C. Weisskopf, K. Wu, F. Xie, and S. Zane, IXPE and Multiwavelength Observations of Blazar PG 1553+113 Reveal an Orphan Optical Polarization Swing, *Astrophysical Journal, Letters* **953**, L28 (2023), [arXiv:2308.00039 \[astro-ph.HE\]](#).
- [33] A. L. Peirson, M. Negro, I. Liodakis, R. Middei, D. E. Kim, A. P. Marscher, H. L. Marshall, L. Pacciani, R. W. Romani, K. Wu, A. Di Marco, N. Di Lalla, N. Omodei, S. G. Jorstad, I. Agudo, P. M. Kouch, E. Lindfors, F. J. Aceituno, M. I. Bernardos, G. Bonnoli, V. Casanova, M. García-Comas, B. Agís-González, C. Husillos, A. Marchini, A. Sota, C. Casadio, J. Escudero, I. Myserlis, A. Sievers, M. Gurwell, R. Rao, R. Imazawa, M. Sasada, Y. Fukazawa, K. S. Kawabata, M. Uemura, T. Mizuno, T. Nakaoka, H. Akitaya, Y. Cheong, H.-W. Jeong, S. Kang, S.-H. Kim, S.-S. Lee, E. Angelakis, A. Kraus, N. Cibrario, I. Donnarumma, J. Poutanen, F. Tavecchio, L. A. Antonelli, M. Bachetti, L. Baldini, W. H. Baumgartner, R. Bellazzini, S. Bianchi, S. D. Bongiorno, R. Bonino, A. Brez, N. Bucciantini, F. Capitanio, S. Castellano, E. Cavazzuti, C.-T. Chen, S. Ciprini, E. Costa, A. De Rosa, E. Del Monte, L. Di Gesu, V. Doroshenko, M. Dovčiak, S. R. Ehlert, T. Enoto, Y. Evangelista, S. Fabiani, R. Ferrazzoli, J. A. Garcia, S. Gunji, K. Hayashida, J. Heyl, W. Iwakiri, P. Kaaret, V. Karas, T. Kitaguchi, J. J. Kolodziejczak, H. Krawczynski, F. La Monaca, L. Latronico, G. Madejski, S. Maldera, A. Manfreda, F. Marin, A. Marinucci, F. Massaro, G. Matt, I. Mitsuishi, F. Muleri, C. Y. Ng, S. L. O'Dell, C. Oppedisano, A. Papitto, G. G. Pavlov, M. Perri, M. Pesce-Rollins, P.-O. Petrucci, M. Pilia, A. Possenti, B. D. Ramsey, J. Rankin, A. Ratheesh, R. W. Romani, C. Sgrò, P. Slane, P. Soffitta, G. Spandre, T. Tamagawa, R. Taverna, Y. Tawara, A. F. Tennant, N. E. Thomas, F. Tombesi, A. Trois, S. Tsygankov, R. Turolla, J. Vink, M. C. Weisskopf, K. Wu, F. Xie, and S. Zane, X-Ray Polarization Observations of BL Lacertae, *Astrophysical Journal, Letters* **942**, L10 (2023), [arXiv:2211.13764 \[astro-ph.HE\]](#).
- [34] R. Middei, I. Liodakis, M. Perri, S. Puccetti, E. Cavazzuti, L. Di Gesu, S. R. Ehlert, G. Madejski, A. P. Marscher, H. L. Marshall, F. Muleri, M. Negro, S. G. Jorstad, B. Agís-González, I. Agudo, G. Bonnoli, M. I. Bernardos, V. Casanova, M. García-Comas, C. Husillos, A. Marchini, A. Sota, P. M. Kouch, E. Lindfors, G. A. Borman, E. N. Kopatskaya, E. G. Larionova, D. A. Morozova, S. S. Savchenko, A. A. Vasilyev, A. V. Zhovtan, C. Casadio, J. Escudero, I. Myserlis, A. Hales, S. Kamenno, R. Kneissl, H. Messiah, H. Nagai, D. Blinov, I. G. Bourbah, S. Kiehlmann, E. Kontopodis, N. Mandarakas, S. Romanopoulos, R. Skalidis, A. Vervelaki, J. R. Masiero, D. Mawet, M. A. Millar-Blanchaer, G. V. Panopoulou, S. Tinyanont, A. V. Berdyugin, M. Kagitani, V. Kravtsov, T. Sakanoi, R. Imazawa, M. Sasada, Y. Fukazawa, K. S. Kawabata, M. Uemura, T. Mizuno, T. Nakaoka, H. Akitaya, M. Gurwell, R. Rao, N. Di Lalla, N. Cibrario, I. Donnarumma, D. E. Kim, N. Omodei, L. Pacciani, J. Poutanen, F. Tavecchio, L. A. Antonelli, M. Bachetti, L. Baldini, W. H. Baumgartner, R. Bellazzini, S. Bianchi, S. D. Bongiorno, R. Bonino, A. Brez, N. Bucciantini, F. Capitanio, S. Castellano, S. Ciprini, E. Costa, A. De Rosa, E. Del Monte, A. Di Marco, V. Doroshenko, M. Dovčiak, T. Enoto, Y. Evangelista, S. Fabiani, R. Ferrazzoli, J. A. Garcia, S. Gunji, K. Hayashida, J. Heyl, W. Iwakiri, V. Karas, T. Kitaguchi, J. J. Kolodziejczak, H. Krawczynski, F. La Monaca, L. Latronico, S. Maldera, A. Manfreda, F. Marin, A. Marinucci, F. Massaro, G. Matt, I. Mitsuishi, C. Y. Ng, S. L. O'Dell, C. Oppedisano, A. Papitto, G. G. Pavlov, A. L. Peirson, M. Pesce-Rollins, P.-O. Petrucci, M. Pilia, A. Possenti, B. D. Ramsey, J. Rankin, A. Ratheesh, R. W. Romani, C. Sgrò, P. Slane, P. Soffitta, G. Spandre, T. Tamagawa, R. Taverna, Y. Tawara, A. F. Tennant, N. E. Thomas, F. Tombesi, A. Trois, S. Tsygankov, R. Turolla, J. Vink, M. C. Weisskopf, K. Wu, F. Xie, and S. Zane, X-Ray Polarization Observations of BL Lacertae, *Astrophysical Journal, Letters* **942**, L10 (2023), [arXiv:2211.13764 \[astro-ph.HE\]](#).
- [35] I. Liodakis, A. P. Marscher, I. Agudo, A. V. Berdyugin, M. I. Bernardos, G. Bonnoli, G. A. Borman, C. Casadio, V. Casanova, E. Cavazzuti, N. Rodriguez Cavero, L. Di Gesu, N. Di Lalla, I. Donnarumma, S. R. Ehlert, M. Errando, J. Escudero, M. García-Comas, B. Agís-González, C. Husillos, J. Jormanainen, S. G. Jorstad, M. Kagitani, E. N. Kopatskaya, V. Kravtsov, H. Krawczynski, E. Lindfors, E. G. Larionova, G. M. Madejski, F. Marin, A. Marchini, H. L. Marshall, D. A. Morozova, F. Massaro, J. R. Masiero, D. Mawet, R. Middei, M. A. Millar-Blanchaer, I. Myserlis, M. Negro, K. Nilsson, S. L. O'Dell, N. Omodei, L. Pacciani, A. Paggi, G. V. Panopoulou, A. L. Peirson, M. Perri, P.-O. Petrucci, J. Poutanen, S. Puccetti, R. W. Romani, T. Sakanoi, S. S. Savchenko, A. Sota, F. Tavecchio, G. G. Pavlov, M. Perri, M. Pesce-Rollins, P.-O. Petrucci, M. Pilia, A. Possenti, B. D. Ramsey, J. Rankin, A. Ratheesh, O. J. Roberts, C. Sgrò, P. Slane, P. Soffitta, G. Spandre, D. A. Swartz, T. Tamagawa, R. Taverna, Y. Tawara, A. F. Tennant, N. E. Thomas, F. Tombesi, A. Trois, S. Tsygankov, R. Turolla, J. Vink, M. C. Weisskopf, F. Xie, and S. Zane, X-Ray Polarization of BL Lacertae in Outburst, *Astrophysical Journal, Letters* **948**, L25 (2023), [arXiv:2305.13898 \[astro-ph.HE\]](#).

- chio, S. Tinyanont, A. A. Vasilyev, Z. R. Weaver, A. V. Zhovtan, L. A. Antonelli, M. Bachetti, L. Baldini, W. H. Baumgartner, R. Bellazzini, S. Bianchi, S. D. Bongiorno, R. Bonino, A. Brez, N. Bucciantini, F. Capitanio, S. Castellano, S. Ciprini, E. Costa, A. De Rosa, E. Del Monte, A. Di Marco, V. Doroshenko, M. Dovciak, T. Enoto, Y. Evangelista, S. Fabiani, R. Ferrazzoli, J. A. Garcia, S. Gunji, K. Hayashida, J. Heyl, W. Iwakiri, V. Karas, T. Kitaguchi, J. J. Kolodziejczak, F. La Monaca, L. Latronico, S. Maldera, A. Manfreda, A. Marinucci, G. Matt, I. Mitsuishi, T. Mizuno, F. Muleri, S. C. Y. Ng, C. Oppedisano, A. Papitto, G. G. Pavlov, M. Pesce-Rollins, M. Pilia, A. Possenti, B. D. Ramsey, J. Rankin, A. Ratheesh, C. Sgró, P. Slane, P. Soffitta, G. Spandre, T. Tamagawa, R. Taverna, Y. Tawara, A. F. Tennant, N. E. Thomas, F. Tombesi, A. Trois, S. Tsygankov, R. Turolla, J. Vink, M. C. Weisskopf, K. Wu, F. Xie, and S. Zane, Polarized blazar X-rays imply particle acceleration in shocks, *Nature (London)* **611**, 677 (2022), [arXiv:2209.06227 \[astro-ph.HE\]](https://arxiv.org/abs/2209.06227).
- [36] H. L. Marshall, I. Liodakis, A. P. Marscher, N. Di Lalla, S. G. Jorstad, D. E. Kim, R. Middei, M. Negro, N. Omodei, A. L. Peirson, M. Perri, S. Puccetti, I. Agudo, G. Bonnoli, A. V. Berdyugin, E. Cavazzuti, N. Rodriguez Caverio, I. Donnarumma, L. Di Gesu, J. Jormanainen, H. Krawczynski, E. Lindfors, F. Marin, F. Massaro, L. Pacciani, J. Poutanen, F. Tavecchio, P. M. Kouch, F. J. Aceituno, M. I. Bernardos, G. Bonnoli, V. Casanova, M. Garcia-Comas, B. Agis-Gonzalez, C. Husillos, A. Marchini, A. Sota, D. Blinov, I. G. Bourbah, S. Kielhmann, E. Kontopodis, N. Mandarakas, S. Romanopoulos, R. Skalidis, A. Vervelaki, G. A. Borman, E. N. Kopatskaya, E. G. Larionova, D. A. Morozova, S. S. Savchenko, A. A. Vasilyev, A. V. Zhovtan, C. Casadio, J. Escudero, J. Kramer, I. Myserlis, E. Trainou, R. Imazawa, M. Sasada, Y. Fukazawa, K. S. Kawabata, M. Uemura, T. Mizuno, T. Nakaoka, H. Akitaya, J. R. Masiero, D. Mawet, M. A. Millar-Blanchaer, G. V. Panopoulou, S. Tinyanont, A. V. Berdyugin, M. Kagitani, V. Kravtsov, T. Sakanoi, L. A. Antonelli, M. Bachetti, L. Baldini, W. H. Baumgartner, R. Bellazzini, S. Bianchi, S. D. Bongiorno, R. Bonino, A. Brez, N. Bucciantini, F. Capitanio, S. Castellano, E. Cavazzuti, C.-T. Chen, S. Ciprini, E. Costa, A. De Rosa, E. Del Monte, L. Di Gesu, A. Di Marco, I. Donnarumma, V. Doroshenko, M. Dovciak, S. R. Ehler, T. Enoto, Y. Evangelista, S. Fabiani, R. Ferrazzoli, J. A. Garcia, S. Gunji, K. Hayashida, J. Heyl, W. Iwakiri, P. Kaaret, V. Karas, F. Kislak, T. Kitaguchi, J. J. Kolodziejczak, H. Krawczynski, F. La Monaca, L. Latronico, S. Maldera, A. Manfreda, F. Marin, A. Marinucci, G. Matt, I. Mitsuishi, T. Mizuno, F. Muleri, C. Y. Ng, S. L. O'Dell, C. Oppedisano, A. Papitto, G. G. Pavlov, M. Pesce-Rollins, P.-O. Petrucci, M. Pilia, A. Possenti, J. Poutanen, S. Puccetti, B. D. Ramsey, J. Rankin, A. Ratheesh, O. J. Roberts, R. W. Romani, C. Sgro, P. Slane, P. Soffitta, G. Spandre, D. A. Swartz, T. Tamagawa, R. Taverna, Y. Tawara, A. F. Tennant, N. E. Thomas, F. Tombesi, A. Trois, S. S. Tsygankov, R. Turolla, J. Vink, M. C. Weisskopf, K. Wu, F. Xie, and S. Zane, Observations of Low and Intermediate Spectral Peak Blazars with the Imaging X-ray Polarimetry Explorer, [arXiv e-prints](https://arxiv.org/abs/2310.11510), [arXiv:2310.11510 \(2023\)](https://arxiv.org/abs/2310.11510), [arXiv:2310.11510 \[astro-ph.HE\]](https://arxiv.org/abs/2310.11510).
- [37] E. S. Perlman, J. T. Stocke, J. F. Schachter, M. Elvis, E. Ellingson, C. M. Urry, M. Potter, C. D. Impey, and P. Kolchinsky, The Einstein Slew Survey Sample of BL Lacertae Objects, *Astrophysical Journal, Supplement* **104**, 251 (1996).
- [38] H. Krawczynski, S. B. Hughes, D. Horan, F. Aharonian, M. F. Aller, H. Aller, P. Boltwood, J. Buckley, P. Coppi, G. Fossati, N. Göting, J. Holder, D. Horns, O. M. Kurtanidze, A. P. Marscher, M. Nikolashvili, R. A. Remillard, A. Sadun, and M. Schröder, Multiwavelength Observations of Strong Flares from the TeV Blazar 1ES 1959+650, *Astrophys. J.* **601**, 151 (2004), [arXiv:astro-ph/0310158 \[astro-ph\]](https://arxiv.org/abs/astro-ph/0310158).
- [39] B. Kapanadze, D. Dorner, S. Vercellone, P. Romano, S. Kapanadze, and T. Mdzinarishvili, A recent strong X-ray flaring activity of 1ES 1959+650 with possibly less efficient stochastic acceleration, *Mon. Not. Roy. Astr. Soc.* **461**, L26 (2016).
- [40] P. C. Gregory and J. J. Condon, The 87GB Catalog of Radio Sources Covering 0 degrees \leq δ \leq +75 degrees at 4.85 GHz, *Astrophysical Journal, Supplement* **75**, 1011 (1991).
- [41] M. Elvis, D. Plummer, J. Schachter, and G. Fabbiano, The Einstein Slew Survey, *Astrophysical Journal, Supplement* **80**, 257 (1992).
- [42] V. Beckmann, A. Wolter, A. Celotti, L. Costamante, G. Ghisellini, T. Maccacaro, and G. Tagliaferri, BeppoSAX spectral survey of BL Lacs - New spectra and results, *Astronomy and Astrophysics* **383**, 410 (2002), [arXiv:astro-ph/0112311 \[astro-ph\]](https://arxiv.org/abs/astro-ph/0112311).
- [43] F. Massaro, A. Tramacere, A. Cavaliere, M. Perri, and P. Giommi, X-ray spectral evolution of TeV BL Lacertae objects: eleven years of observations with BeppoSAX, XMM-Newton and Swift satellites, *Astronomy and Astrophysics* **478**, 395 (2008), [arXiv:0712.2116 \[astro-ph\]](https://arxiv.org/abs/0712.2116).
- [44] T. Nishiyama, Detection of a new TeV gamma-ray source of BL Lac object 1ES 1959+650, in *26th International Cosmic Ray Conference (ICRC26), Volume 3*, International Cosmic Ray Conference, Vol. 3 (1999) p. 370.
- [45] J. Holder, I. H. Bond, P. J. Boyle, S. M. Bradbury, J. H. Buckley, D. A. Carter-Lewis, W. Cui, C. Dowdall, C. Duke, I. de la Calle Perez, A. Falcone, D. J. Fegan, S. J. Fegan, J. P. Finley, L. Fortson, J. A. Gaidos, K. Gibbs, S. Gammell, J. Hall, T. A. Hall, A. M. Hillas, D. Horan, M. Jordan, M. Kertzman, D. Kieda, J. Kildea, J. Knapp, K. Kosack, H. Krawczynski, F. Krennrich, S. LeBohec, E. T. Linton, J. Lloyd-Evans, P. Moriarty, D. Müller, T. N. Nagai, R. Ong, M. Page, R. Palladini, D. Petry, B. Power-Mooney, J. Quinn, P. Rebillot, P. T. Reynolds, H. J. Rose, M. Schroedter, G. H. Sembroski, S. P. Swordy, V. V. Vassiliev, S. P. Wakely, G. Walker, and T. C. Weekes, Detection of TeV Gamma Rays from the BL Lacertae Object 1ES 1959+650 with the Whipple 10 Meter Telescope, *Astrophysical Journal, Letters* **583**, L9 (2003), [arXiv:astro-ph/0212170 \[astro-ph\]](https://arxiv.org/abs/astro-ph/0212170).
- [46] E. Aliu, S. Archambault, T. Arlen, T. Aune, A. Barnacka, M. Beilicke, W. Benbow, K. Berger, R. Bird, A. Bouvier, J. H. Buckley, V. Bugaev, M. Cerruti, X. Chen, L. Ciupik, E. Collins-Hughes, M. P. Connolly, W. Cui, J. Dumm, J. D. Eisch, A. Falcone, S. Federici, Q. Feng, J. P. Finley, H. Fleischhack, P. Fortin, L. Fortson, A. Furniss, N. Galante, G. H. Gillanders, S. Griffin, S. T. Griffiths, J. Grube, G. Gyuk, N. Håkansson, D. Hanna, J. Holder, G. Hughes, Z. Hughes, T. B. Hu-

- mensky, C. A. Johnson, P. Kaaret, P. Kar, M. Kertzman, Y. Khassen, D. Kieda, H. Krawczynski, F. Krennrich, M. J. Lang, A. S. Madhavan, P. Majumdar, S. McArthur, A. McCann, K. Meagher, J. Millis, P. Moriarty, R. Mukherjee, T. Nelson, D. Nieto, A. O’Faoláin de Bhróithe, R. A. Ong, A. N. Otte, N. Park, J. S. Perkins, M. Pohl, A. Popkow, H. Prokoph, J. Quinn, K. Ragan, J. Rajotte, L. C. Reyes, P. T. Reynolds, G. T. Richards, E. Roache, A. Sadun, M. Santander, G. H. Sembroski, K. Shahinyan, F. Sheidaei, A. W. Smith, D. Staszak, I. Tezhinsky, M. Theiling, J. Tyler, A. Varlotta, V. V. Vassiliev, S. Vincent, S. P. Wakely, T. C. Weekes, A. Weinstein, R. Welsing, A. Wilhelm, D. A. Williams, B. Zitzer, VERITAS Collaboration, M. Böttcher, and M. Fumagalli, Investigating Broadband Variability of the TeV Blazar 1ES 1959+650, *Astrophys. J.* **797**, 89 (2014), [arXiv:1412.1031 \[astro-ph.HE\]](#).
- [47] P. B. Graff, M. Georganopoulos, E. S. Perlman, and D. Kazanas, A Multizone Model for Simulating the High-Energy Variability of TeV Blazars, *Astrophys. J.* **689**, 68 (2008), [arXiv:0808.2135 \[astro-ph\]](#).
- [48] M. Böttcher, A Hadronic Synchrotron Mirror Model for the “Orphan” TeV Flare in 1ES 1959+650, *Astrophys. J.* **621**, 176 (2005), [arXiv:astro-ph/0411248 \[astro-ph\]](#).
- [49] L. Baldini, N. Bucciantini, N. D. Lalla, S. Ehlert, A. Manfreda, M. Negro, N. Omodei, M. Pesce-Rollins, C. Sgrò, and S. Silvestri, ixpeobssim: A simulation and analysis framework for the imaging X-ray polarimetry explorer, *SoftwareX* **19**, 101194 (2022), [arXiv:2203.06384 \[astro-ph.IM\]](#).
- [50] K. A. Arnaud, XSPEC: The First Ten Years, in *Astronomical Data Analysis Software and Systems V*, Astronomical Society of the Pacific Conference Series, Vol. 101, edited by G. H. Jacoby and J. Barnes (1996) p. 17.
- [51] R. Willingale, R. L. C. Starling, A. P. Beardmore, N. R. Tanvir, and P. T. O’Brien, Calibration of X-ray absorption in our Galaxy, *Mon. Not. Roy. Astr. Soc.* **431**, 394 (2013), [arXiv:1303.0843 \[astro-ph.HE\]](#).
- [52] L. Di Gesu, I. Donnarumma, F. Tavecchio, I. Agudo, T. Barnounin, N. Cibrario, N. Di Lalla, A. Di Marco, J. Escudero, M. Errando, S. G. Jorstad, D. E. Kim, P. M. Kouch, I. Liodakis, E. Lindfors, G. Madejski, H. L. Marshall, A. P. Marscher, R. Middei, F. Muleri, I. Myserlis, M. Negro, N. Omodei, L. Pacciani, A. Paggi, M. Perri, S. Puccetti, L. A. Antonelli, M. Bachetti, L. Baldini, W. H. Baumgartner, R. Bellazzini, S. Bianchi, S. D. Bongiorno, R. Bonino, A. Brez, N. Bucciantini, F. Capitanio, S. Castellano, E. Cavazzuti, S. Ciprini, E. Costa, A. De Rosa, E. Del Monte, V. Doroshenko, M. Dovčiak, S. R. Ehlert, T. Enoto, Y. Evangelista, S. Fabiani, R. Ferrazzoli, J. A. Garcia, S. Gunji, K. Hayashida, J. Heyl, W. Iwakiri, V. Karas, T. Kitaguchi, J. J. Kolodziejczak, H. Krawczynski, F. La Monaca, L. Latronico, S. Maldera, A. Manfreda, F. Marin, A. Marinucci, F. Massaro, G. Matt, I. Mitsuishi, T. Mizuno, C. Y. Ng, S. L. O’Dell, C. Oppedisano, A. Papitto, G. G. Pavlov, A. L. Peirson, M. Pesce-Rollins, P.-O. Petrucci, M. Pilia, A. Possenti, J. Poutanen, B. D. Ramsey, J. Rankin, A. Ratheesh, R. W. Romani, C. Sgrò, P. Slane, P. Soffitta, G. Spandre, T. Tamagawa, R. Taverna, Y. Tawara, A. F. Tennant, N. E. Thomas, F. Tombesi, A. Trois, S. Tsygankov, R. Turolla, J. Vink, M. C. Weisskopf, K. Wu, F. Xie, and S. Zane, The X-Ray Polarization View of Mrk 421 in an Average Flux State as Observed by the Imaging X-Ray Polarimetry Explorer, *Astrophysical Journal, Letters* **938**, L7 (2022), [arXiv:2209.07184 \[astro-ph.HE\]](#).
- [53] J. Otero-Santos, J. A. Acosta-Pulido, J. Becerra González, C. M. Raiteri, M. I. Carnerero, N. Castro Segura, O. González-Martín, and A. Luashvili, Variability and evolution of the optical polarization of a sample of gamma-ray blazars, *Mon. Not. Roy. Astr. Soc.* **523**, 4504 (2023), [arXiv:2306.03919 \[astro-ph.HE\]](#).
- [54] F. Tavecchio, M. Landoni, L. Sironi, and P. Coppi, Probing shock acceleration in BL Lac jets through X-ray polarimetry: the time-dependent view, *Mon. Not. Roy. Astr. Soc.* **498**, 599 (2020), [arXiv:2007.08353 \[astro-ph.HE\]](#).
- [55] G. Tagliaferri, L. Foschini, G. Ghisellini, L. Maraschi, pre=”and” post=””, a. and” affil=”4”¿G. Tosti, J. Albert, E. Aliu, H. Anderhub, P. Antoranz, C. Baixeras, J. A. Barrio, H. Bartko, D. Bastieri, J. K. Becker, W. Bednarek, A. Bedyugin, K. Berger, C. Bigongiari, A. Biland, R. K. Bock, P. Bordas, V. Bosch-Ramon, T. Bretz, I. Britvitch, M. Camara, E. Carmona, A. Chilingarian, J. A. Coarasa, S. Commichau, J. L. Contreras, J. Cortina, M. T. Costado, V. Curtef, V. Danielyan, F. Dazzi, A. De Angelis, C. Delgado, R. de los Reyes, B. De Lotto, D. Dorner, M. Doro, M. Errando, M. Fagiolini, D. Ferenc, E. Fernández, R. Firpo, M. V. Fonseca, L. Font, M. Fuchs, N. Galante, R. J. García-López, M. Garczarczyk, M. Gaug, M. Giller, F. Goebel, D. Hakobyan, M. Hayashida, T. Hengstebeck, A. Herrero, D. Höhne, J. Hose, S. Huber, C. C. Hsu, P. Jacon, T. Jogler, R. Kosyra, D. Kranich, R. Kritzer, A. Laille, E. Lindfors, S. Lombardi, F. Longo, M. López, E. Lorenz, P. Majumdar, G. Maneva, K. Mannheim, M. Maricotti, M. Martínez, D. Mazin, C. Merck, M. Meucci, M. Meyer, J. M. Miranda, R. Mirzoyan, S. Mizobuchi, A. Moralejo, D. Nieto, K. Nilsson, J. Ninkovic, E. Oña-Wilhelmi, N. Otte, I. Oya, M. Panniello, R. Paoletti, J. M. Paredes, M. Pasanen, D. Pascoli, F. Pauss, R. Pegna, M. Persic, L. Peruzzo, A. Piccioli, E. Prandini, N. Puchades, A. Raymers, W. Rhode, M. Ribó, J. Rico, M. Rissi, A. Robert, S. Rügemer, A. Saggion, T. Y. Saito, A. Sánchez, P. Sartori, V. Scalzotto, V. Scapin, R. Schmitt, T. Schweizer, M. Shayduk, K. Shinozaki, S. N. Shore, N. Sidro, A. Sillanpää, D. Sobczynska, F. Spanier, A. Stamerra, L. S. Stark, L. Takalo, F. Tavecchio, P. Temnikov, D. Tescaro, M. Teshima, D. F. Torres, N. Turini, H. Vankov, A. Venturini, V. Vitale, R. M. Wagner, T. Wibig, W. Wittek, F. Zandanel, R. Zanin, J. Zapatero, and MAGIC Collaboration”, Simultaneous Multiwavelength Observations of the Blazar 1ES 1959+650 at a Low TeV Flux, *Astrophys. J.* **679**, 1029 (2008), [arXiv:0801.4029 \[astro-ph\]](#).
- [56] F. D. Kahn, A turbulent radio jet, *Mon. Not. Roy. Astr. Soc.* **202**, 553 (1983).
- [57] S. Britzen, C. Fendt, G. Witzel, S. J. Qian, I. N. Pashchenko, O. Kurtanidze, M. Zajacek, G. Martinez, V. Karas, M. Aller, H. Aller, A. Eckart, K. Nilsson, P. Arévalo, J. Cuadra, M. Subroweit, and A. Witzel, OJ287: deciphering the ‘Rosetta stone of blazars, *Mon. Not. Roy. Astr. Soc.* **478**, 3199 (2018).
- [58] R. Lico, J. Liu, M. Giroletti, M. Orienti, J. L. Gómez, B. G. Piner, N. R. MacDonald, F. D’Ammando, and A. Fuentes, A parsec-scale wobbling jet in the high-synchrotron peaked blazar PG 1553+113, *Astronomy and Astrophysics* **634**, A87 (2020), [arXiv:2001.01753](#)

- [59] [\[astro-ph.HE\]](#).
Z. R. Weaver, S. G. Jorstad, A. P. Marscher, D. A. Morozova, I. S. Troitsky, I. Agudo, J. L. Gómez, A. Lähteenmäki, J. Tammi, and M. Tornikoski, Kinematics of Parsec-scale Jets of Gamma-Ray Blazars at 43 GHz during 10 yr of the VLBA-BU-BLAZAR Program, *Astrophysical Journal, Supplement* **260**, 12 (2022), [arXiv:2202.12290 \[astro-ph.HE\]](#).
- [60] A. P. Marscher, S. G. Jorstad, F. D. D’Arcangelo, P. S. Smith, G. G. Williams, V. M. Larionov, H. Oh, A. R. Olmstead, M. F. Aller, H. D. Aller, I. M. McHardy, A. Lähteenmäki, M. Tornikoski, E. Valtaoja, V. A. Hagen-Thorn, E. N. Kopatskaya, W. K. Gear, G. Tosti, O. Kurtanidze, M. Nikolashvili, L. Sigua, H. R. Miller, and W. T. Ryle, The inner jet of an active galactic nucleus as revealed by a radio-to- γ -ray outburst, *Nature (London)* **452**, 966 (2008).