

UNBIASED KINETIC LANGEVIN MONTE CARLO WITH INEXACT GRADIENTS

BY NEIL K. CHADA^{1,a}, BENEDICT LEIMKUHNER^{2,b}, DANIEL PAULIN^{2,c}
AND PETER A. WHALLEY^{2,d}

¹*Department of Actuarial Mathematics and Statistics, Heriot Watt University, neilchada123@gmail.com*

²*School of Mathematics, University of Edinburgh, b.leimkuhler@ed.ac.uk; dpaulin@ed.ac.uk; p.a.whalley@sms.ed.ac.uk*

We present an unbiased method for Bayesian posterior means based on kinetic Langevin dynamics that combines advanced splitting methods with enhanced gradient approximations. Our approach avoids Metropolis correction by coupling Markov chains at different discretization levels in a multilevel Monte Carlo approach. Theoretical analysis demonstrates that our proposed estimator is unbiased, attains finite variance, and satisfies a central limit theorem. It can achieve accuracy $\epsilon > 0$ for estimating expectations of Lipschitz functions in d dimensions with $\mathcal{O}(d^{1/4}\epsilon^{-2})$ expected gradient evaluations, without assuming warm start. We exhibit similar bounds using both approximate and stochastic gradients, and our method's computational cost is shown to scale independently of the size of the dataset. The proposed method is tested using a multinomial regression problem on the MNIST dataset and a Poisson regression model for soccer scores. Experiments indicate that the number of gradient evaluations per effective sample is independent of dimension, even when using inexact gradients. For product distributions, we give dimension-independent variance bounds. Our results demonstrate that the unbiased algorithm we present can be much more efficient than the "gold-standard" randomized Hamiltonian Monte Carlo.

CONTENTS

1	Introduction	2
1.1	Unbiased estimation without accept/reject steps	2
1.2	Proposed methodology	3
1.3	Organization	5
1.4	Notation	6
2	Background & preliminary material	6
2.1	Splitting methods	7
2.2	Extension to stochastic gradients	8
3	Unbiased multilevel Monte Carlo methods	8
3.1	UBUBU with exact gradients	11
3.2	UBUBU with stochastic gradients	16
3.3	UBUBU with approximate gradients	22
4	Numerical results	25
4.1	Gaussian target	25
4.2	Bayesian multinomial regression	27
4.3	Poisson regression model	30
5	Conclusion	31
	Funding	32

MSC2020 subject classifications: 65C05, 65C30, 65C40, 62F15.

Keywords and phrases: unbiased estimation, kinetic Langevin dynamics, multilevel Monte Carlo, stochastic gradient.

A	Discussion and outline of results	32
B	Unbiased multilevel estimators	33
C	Convergence results	36
D	Variance bounds for UBUBU estimator with exact gradients	47
	D.1 Variance bound of $D_{l,l+1}$	47
	D.2 Variance bound of D_0	54
	D.3 Variance of $S(c_R)$	59
E	Initialization and Gaussian approximation	62
	E.1 OHO scheme	62
	E.2 Initialization and bounds	66
F	Variance bounds for UBU with SVRG	66
	F.1 Variance bound of $D_{l,l+1}$	67
	F.2 Variance of $S(c_R)$	77
G	Variance bounds for UBUBU estimator with approximate gradients	78
	G.1 Non-asymptotic guarantees	79
	G.2 Variance bound of $D_{l,l+1}$	85
	G.3 Variance bound of $S(c_R)$	87
H	Auxiliary results & RHMC algorithm	87
	References	93

1. Introduction. Markov chain Monte Carlo (MCMC) methods are standard computational tools for high-dimensional Bayesian inference [82]. They enable the computation of posterior means and variances and other observable averages by replacing ensemble calculations with Monte Carlo sums over discrete Markov processes. A limitation to the broader uptake of Bayesian inference is the scaling of the computational cost of MCMC algorithms with model dimension and dataset size. Typical MCMC methods (Metropolis Adjusted Langevin Algorithm [6, 83], Hamiltonian Monte Carlo [34, 74]) employ Metropolis-Hastings correction steps to ensure convergence to the desired invariant distribution. The cost of implementing such corrections scales linearly with dataset size. Even worse, in order to maintain a high acceptance rate, stepsizes must decrease as a function of the model dimension, which implies that convergence rates are also dependent on dimension [7, 24, 84].

By contrast, optimization methods typically have convergence rates that are independent of the dimension and can make use of stochastic gradients based on a subset of the data instead of the entire dataset [50]. For these reasons, optimization algorithms are much more scalable than sampling methods, so practitioners often prefer machine-learning approaches. The relative inefficiency of sampling compared to optimization also limits the uptake of uncertainty quantification techniques (typically built on a Bayesian foundation) in high-dimensional machine learning applications.

1.1. *Unbiased estimation without accept/reject steps.* This paper describes a technique for performing Bayesian inference based on unbiased unadjusted Markov chain Monte Carlo that does not rely on Metropolis-Hastings accept/reject steps. Our algorithm is based on a multilevel scheme [42] that combines several different unadjusted MCMC chains to eliminate bias efficiently. Our approach is related to a recent paper [85] that introduced an unbiased unadjusted MCMC method, however we employ state-of-the-art integrators, and we extend the method with modifications for handling incomplete (or approximate) gradients, thus obtaining a procedure with improved scalability and competitiveness compared to state-of-the-art algorithms such as randomized Hamiltonian Monte Carlo (RHMC) [16, 24].

Unbiased Monte Carlo methods have been widely studied in the recent literature; see Section 2.1 of [49] for an overview. The goal of the methods of [26, 44, 46, 49, 80] is to remove

burn-in bias via couplings. [52] proposed an alternative method for eliminating burn-in bias by considering a burn-in period of random length. The cited papers above all require that the stationary distribution of the Markov chain has no bias (hence, these methods typically involve Metropolization) and are not able to remove discretization bias in SDEs such as (1.1) treated using numerical methods. Middleton et al [68] extended unbiased methods to intractable likelihoods, and [33] created unbiased estimators of MCMC asymptotic variances.

There have been several proposals for creating computationally efficient estimators for functions of SDE paths based on numerical discretization using multilevel Monte Carlo variance reduction techniques. Our scheme relates to the method of Muller et al [73] for approximating functions of whole paths of kinetic Langevin dynamics using integrators based on splitting. Unlike our approach, that work did not address the stationary distribution; moreover, the burn-in bias was not eliminated, and they did not consider the incorporation of approximate or stochastic gradients. More recently, Giles et al [43] introduced a general framework for multilevel approximation of expectations with respect to the stationary distribution of overdamped Langevin dynamics and also considered stochastic gradients. However, their approach does not produce unbiased samples, and overdamped Langevin dynamics generally appear less efficient at exploring distributions with high condition numbers than well-tuned kinetic Langevin dynamics [78], as considered here. Until this work, multilevel approaches have not been shown to be competitive with Hamiltonian Monte Carlo methods for high-dimensional sampling.

We also mention that, in the area of molecular simulation, unadjusted numerical discretizations of kinetic Langevin dynamics have been employed for sampling from complex distributions for many years [19, 48, 58, 59]. Even though such discretizations introduce bias, this is often dominated by the Monte Carlo error—even at substantially larger stepsizes than would typically be used in Metropolized calculations [58]. On the other hand, the magnitude of the sampling bias due to finite stepsize is problem-dependent and can be difficult to quantify; thus, there are situations where the ability to ameliorate the discretization bias is crucial. Some authors have proposed reducing the discretization bias by decreasing stepsize asymptotically [36, 99]. However, such a procedure can slow convergence or introduce heuristic schedules into the sampling apparatus.

1.2. *Proposed methodology.* We consider kinetic Langevin dynamics (also referred to as underdamped Langevin dynamics [25, 30]):

$$(1.1) \quad \begin{aligned} dX_t &= V_t dt, \\ dV_t &= -\nabla U(X_t) dt - \gamma V_t dt + \sqrt{2\gamma} dW_t, \end{aligned}$$

where $U : \mathbb{R}^d \rightarrow \mathbb{R}$ is a potential energy function, $\{W_t\}_{t \geq 0}$ is a standard d -dimensional Brownian motion, and $\gamma > 0$ is a friction coefficient. Under fairly weak assumptions, the unique invariant measure of the process $\{X_t, V_t\}_{t \geq 0}$ is of the form

$$(1.2) \quad \pi(dx dv) \propto \exp\left(-U(x) - \frac{\|v\|^2}{2}\right) dx dv.$$

This dynamics forms the basis of many sampling methods [19, 63], and it has a dimension-independent convergence rate for a large class of distributions [20]. In this paper, we expand on the work of [85] and develop a comprehensive and practical framework for unbiased estimation. Specifically, we consider using a splitting integrator called UBU [86], which is strongly second-order accurate, where the unbiased estimator we introduce is referred to as UBUBU (Unbiased-UBU). Figure 1 illustrates the synchronously coupled paths of UBU discretizations of kinetic Langevin dynamics.




Fig 1: Coupled sample paths based on synchronous coupling from UBU (Section 2) discretization scheme of kinetic Langevin diffusion for a Gaussian target at stepsizes $h = 1.5, 0.75$ and $h = 0.75, 0.375$. UBU is strong order 2, so the typical distance between coupled paths is $\mathcal{O}(h^2)$.

In Figure 1 we see that UBU discretization can be pathwise accurate even at large step-size. Nevertheless, there is always some residual bias, and the stationary distribution of the discretization with stepsize h , π_h , differs from the target distribution π . The idea of unbiased estimation as proposed in [85] was to consider a sequence of discretization levels $h_l = 2^{-l}h_0$ for $l = 0, 1, 2, \dots$ to create an estimator of the form

$$(1.3) \quad \hat{\pi}(f) = \hat{\pi}_{h_0}(f) + \sum_{l=0}^{\infty} \hat{\pi}_{h_{l+1}, h_l}(f),$$

where f is some arbitrary quantity of interest, $\hat{\pi}_{h_0}(f)$ is an unbiased estimator of $\pi_{h_0}(f)$, and $\hat{\pi}_{h_{l+1}, h_l}(f)$ is an unbiased estimator of $\pi_{h_{l+1}}(f) - \pi_{h_l}(f)$. A sophisticated coupling construction was used for defining $\hat{\pi}_{h_{l+1}, h_l}(f)$ based on four Markov chains using Euler–Maruyama discretization of (1.1). Under certain weak assumptions, the estimator (1.3) was shown to have no bias, finite variance and finite expected computational cost.

In our algorithm:

- (i) The burn-in bias is eliminated differently, resulting in simpler couplings. Our estimator is still of the form (1.3). However, instead of estimating $\pi_{h_0}(f)$ and $\pi_{h_{l+1}}(f) - \pi_{h_l}(f)$, which requires eliminating the burn-in bias for both discretization levels, we let $\hat{\pi}_{h_0}(f)$ be an unbiased estimator of $\tilde{\pi}_{h_0}(f)$, and $\hat{\pi}_{h_{l+1}, h_l}(f)$ be an unbiased estimator $\tilde{\pi}_{h_{l+1}}(f) - \tilde{\pi}_{h_l}(f)$. Here $\tilde{\pi}_{h_l}(f)$ denotes the expected value of f according to the empirical distribution of a Markov chain using discretization stepsize h_l , thinning 2^l , and burn-in period of length $(B_0 + l \cdot B)/h_l$, for some constants $B_0, B > 0$. See Figure 2 for an illustration. Due to the increasing burn-in periods at smaller stepsizes, the bias of $\tilde{\pi}_{h_l}(f)$ shrinks to zero as $l \rightarrow \infty$. With this approach, we only need to couple two chains for creating unbiased estimators of $\tilde{\pi}_{h_{l+1}}(f) - \tilde{\pi}_{h_l}(f)$, and simple synchronous couplings can be used.
- (ii) We use UBU discretization instead of Euler-Maruyama. The higher accuracy of UBU means that the differences between consecutive discretization levels h_l and h_{l+1} are smaller, and as a result, our estimator has a lower variance. We show that under certain assumptions, it is unbiased, has finite variance and finite expected computational cost.
- (iii) In our method, the number of samples per level is deterministic (except at very small stepsize), and we can use Richardson extrapolation [81] to further lower the variance.
- (iv) We show unbiasedness and finite variance even when using approximate or stochastic gradients. This dramatically improves the scalability of our method to large datasets.




Fig 2: Elimination of bias by increasing burn-in lengths at higher discretization levels.

(v) The usual unbiased estimator of Rhee and Glynn takes the form

$$(1.4) \quad \hat{\pi}(f) = \frac{\xi_L}{\mathbb{P}_L(L)},$$

such that

$$\begin{aligned} \mathbb{E}[\xi_{l_*}] &= \pi_{l_*}(f), \\ \mathbb{E}[\xi_l] &= \pi_l(f) - \pi_{l-1}(f) \quad l \in \{l_* + 1, l_* + 2, \dots\}, \end{aligned}$$

where L is a random variable with probability mass function \mathbb{P}_L on $\mathbb{N}_{l_*} := \{l_*, l_* + 1, \dots\}$ that is independent of the sequence $\{\xi_l\}_{l \in \mathbb{N}_{l_*}}$. This approach was also used in [85]. Various alternative schemes with lower variance were proposed in [93].

1.3. Organization. This article is organized as follows. In Section 2, we provide the necessary background material related to this work, including a discussion of splitting methods for kinetic Langevin dynamics, in particular the UBU discretization, as well as others such as BAOAB. We then discuss variants of our algorithm based on UBU, which includes an extension to stochastic gradients.

Section 3 is devoted to introducing our unbiased algorithms. We first provide some simple conditions for creating unbiased estimators with finite variance based on telescopic sums, together with a central limit theorem for such estimators. We then present our method using exact gradients and discuss necessary assumptions for unbiasedness and finite variance including showing that the variance of the estimator is finite. In addition to exact gradients, we also state versions of our method using stochastic and approximate gradients, with theoretical analysis.

Numerical experiments are provided in Section 4 on a range of model problems, including a simple Gaussian problem, an MNIST multinomial regression problem and a Poisson regression model applied to soccer game outcome prediction. Our unbiased methods are compared to RHMC, and demonstrate gains in terms of accuracy, and computational efficiency for high-dimensional problems, while eliminating bias.

We conclude our work and findings in Section 5, while suggesting future directions for research. Finally, we provide detailed proofs of all theorems in the appendices, as summarized in A.

Table 1 compares various Metropolized methods of the literature with our approach. [24] states that the warm start assumption cannot be removed as [57] has shown a lower bound of $\mathcal{O}(d^{1/2})$ without it. [1] proposes an algorithmic warm start using unadjusted kinetic Langevin dynamics at $\mathcal{O}(d^{1/2})$ gradient evaluations. For Gaussian targets, [2] has shown that it is possible to achieve a warm start using $\mathcal{O}(d^{1/4})$ gradient evaluations.

Algorithm	Gradient Evaluations	Conditions	Reference
MALA	$\mathcal{O}(d^{3/7})$	$h = \mathcal{O}(d^{-3/7})$, warm start, strongly Hessian Lipschitz	[24]
HMC	$\mathcal{O}(d^{1/4})$	$h = \mathcal{O}(d^{-1/4})$, warm start, strongly Hessian Lipschitz	[24]
RHMC	$\mathcal{O}(d^{1/4})$	$h = \mathcal{O}(d^{-1/4})$, warm start, Gaussian target	[2]
UBUBU	$\mathcal{O}(d^{1/4})$	$h_0 = \mathcal{O}(d^{-1/4})$, strongly Hessian Lipschitz	this work

TABLE 1

Dimension dependency of gradient evaluations per effective sample for different algorithms for m -strongly convex and M - ∇ Lipschitz potentials, in comparison to UBUBU.

1.4. Notation.

- We let $z_{0:k} = (z_0, z_1, \dots, z_k)$ denote a sequence of variables.
- Let 0_d denote the d -dimensional vector of zeros.
- Let I_d denote the d -dimensional identity matrix.
- Let C denote an absolute constant (whose value may differ in each proposition or theorem).
- Let $C(\text{var}_1, \dots, \text{var}_n)$ denote a constant that is a function of variables $\text{var}_1, \dots, \text{var}_n$ (this function may differ in each proposition or theorem).
- Let G , SG and A denote an abbreviation for gradient, stochastic gradient and ‘‘approximate gradient’’.
- We let $l \in \mathbb{R}^+$ denote the level of discretization with respect to our discretized ULD, with stepsize h_l defined at each level.
- Let D_0 denote the empirical average of samples at level 0.
- Let $D_{l,l+1}$ denote the difference of empirical averages of samples at levels $l+1$ and l , which are generated jointly via a synchronous coupling.
- N denotes the number of samples taken at level 0.
- $N_{l,l+1}$ denotes the number of samples taken from the coupling of levels l and $l+1$.
- N_D is the size of the dataset (number of terms in potential $U(x) = U_0(x) + \sum_{i=1}^{N_D} U_i(x)$).
- Let $z_k = (x_k, v_k)$ denotes step k in a numerical discretization of kinetic Langevin dynamics with time step h (specified each time this notation is used). Similarly Z_t is the solution of (1.1) initialized at the invariant measure with synchronously coupled Brownian motion. $Z^k = Z_{kh}$ denotes the value of the continuous time process at the same time as z_k .
- \hat{x}_k denotes the point where the last gradient is evaluated for SVRG
- $\|\cdot\|_{L^2} := (\mathbb{E}\|\cdot\|^2)^{1/2}$ and $\|\cdot\|_{L^2,a,b} := \left(\mathbb{E}\|\cdot\|_{a,b}^2\right)^{1/2}$.

2. Background & preliminary material. In this section, we provide the essential background material on kinetic (underdamped) Langevin dynamics and a splitting-type scheme called UBU. We then discuss the extension to stochastic gradients and state assumptions required in the remainder of the article.

For this work, we consider Langevin dynamics as defined by Equation (1.1) under temporal discretization. The simplest discretization is the Euler-Maruyama scheme. For a given stepsize $h > 0$, this proceeds, after initialization of x_0, v_0 , with the following recursion:

$$(2.1) \quad \begin{aligned} x_{k+1} &= x_k + hv_k, \\ v_{k+1} &= v_k - h\nabla U(x_k) - h\gamma v_k + \sqrt{2\gamma h}\xi_{k+1}, \end{aligned}$$

where $(\xi_k)_{k \in \mathbb{N}}$ are i.i.d. $\mathcal{N}(0_d, I_d)$ random variables. Under suitable assumptions on the potential U , for h small enough, the discrete-time Markov chain expressed as $\{x_k, v_k\}_{k \in \mathbb{N}}$ admits a unique invariant measure π_h and moreover converges geometrically, meaning that for

suitable classes of functions $f : \mathbb{R}^{2d} \rightarrow \mathbb{R}$,

$$\mathbb{E} \left[\left(K^{-1} \sum_{k=1}^K f(x_k, v_k) - \int f(x, v) \pi_h(dx dv) \right)^2 \right] = \mathcal{O}(K^{-1}),$$

see [35]. In addition to this, π_h converges to π in distribution, as $h \rightarrow 0$. These types of results regarding convergence and accuracy can be extended to other numerical discretizations for the underdamped system, which we next discuss.

2.1. Splitting methods. Improved discretization methods with a high order of accuracy in both the weak and strong senses can be constructed by *splitting* [9, 59, 89], in which the SDE is broken into parts that can be either be solved analytically or which are in some way easier to handle numerically.

An accurate splitting method was introduced in [102] and was also studied in [86]. This splitting method only requires one gradient evaluation per iteration but has strong order two. The method is based on splitting the SDE (1.1) as follows

$$\begin{pmatrix} dx \\ dv \end{pmatrix} = \underbrace{\begin{pmatrix} 0 \\ -\nabla U(x) dt \end{pmatrix}}_{\mathcal{B}} + \underbrace{\begin{pmatrix} v dt \\ -\gamma v dt + \sqrt{2\gamma} dW_t \end{pmatrix}}_{\mathcal{U}},$$

which can be integrated exactly over a step of size h . Given $\gamma > 0$, let $\eta = \exp(-\gamma h/2)$, and for ease of notation, we define the following operators

$$(2.2) \quad \mathcal{B}(x, v, h) = (x, v - h\nabla U(x)),$$

and

$$(2.3) \quad \begin{aligned} \mathcal{U}(x, v, h/2, \xi^{(1)}, \xi^{(2)}) &= \left(x + \frac{1-\eta}{\gamma} v + \sqrt{\frac{2}{\gamma}} \left(\mathcal{Z}^{(1)}(h/2, \xi^{(1)}) - \mathcal{Z}^{(2)}(h/2, \xi^{(1)}, \xi^{(2)}) \right), \right. \\ &\quad \left. \eta v + \sqrt{2\gamma} \mathcal{Z}^{(2)}(h/2, \xi^{(1)}, \xi^{(2)}) \right), \end{aligned}$$

where

$$(2.4) \quad \begin{aligned} \mathcal{Z}^{(1)}(h/2, \xi^{(1)}) &= \sqrt{\frac{h}{2}} \xi^{(1)}, \\ \mathcal{Z}^{(2)}(h/2, \xi^{(1)}, \xi^{(2)}) &= \sqrt{\frac{1-\eta^2}{2\gamma}} \left(\sqrt{\frac{1-\eta}{1+\eta}} \cdot \frac{4}{\gamma h} \xi^{(1)} + \sqrt{1 - \frac{1-\eta}{1+\eta}} \cdot \frac{4}{\gamma h} \xi^{(2)} \right). \end{aligned}$$

The \mathcal{B} operator indicated here is as given previously, whereas \mathcal{U} as defined above is the exact solution in the weak sense of the remainder of the dynamics when $\xi^{(1)}, \xi^{(2)} \sim \mathcal{N}(0, I_d)$ are independent random vectors. Different orders of composition of \mathcal{B} and \mathcal{U} can be taken to define different numerical integrators of kinetic Langevin dynamics, two such methods considered in [102] are BUB, a half step in \mathcal{B} , followed by a full step in \mathcal{U} and a further half step in \mathcal{B} and UBU, a half step in \mathcal{U} followed by a full \mathcal{B} step, followed by a half \mathcal{U} step.

The Markov kernel for an UBU step with stepsize h will be denoted by P_h , which can be described by (2.5) as follows.

$$(2.5) \quad \begin{aligned} &\left(\xi_{k+1}^{(i)} \right)_{i=1}^4, \quad \xi_{k+1}^{(i)} \sim \mathcal{N}(0_d, I_d) \text{ for all } i = 1, \dots, 4. \\ &(x_{k+1}, v_{k+1}) = \mathcal{UBU} \left(x_k, v_k, h, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}, \xi_{k+1}^{(3)}, \xi_{k+1}^{(4)} \right) \\ &= \mathcal{U} \left(\mathcal{B} \left(\mathcal{U} \left(x_k, v_k, h/2, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)} \right), h \right), h/2, \xi_{k+1}^{(3)}, \xi_{k+1}^{(4)} \right). \end{aligned}$$

We have found that the strong second-order property and generally high accuracy of UBU makes it suitable for unbiased estimation, as described in Section 3.

The BAOAB method is an alternative splitting scheme that is known to be second-order weakly accurate and has small bias (see [15, 59–61]). BAOAB is exact for Gaussian targets and has a robustness property for large values of the friction parameter γ (see [63]), but its strong order is one. Theorem 3.3 of [92] claims that the stochastic velocity Verlet (SVV) method is, like UBU, also strongly second-order accurate. Despite their strengths as raw sampling schemes, both BAOAB and SVV exhibited worse performance than UBU in our preliminary numerical experiments in the setting of unbiased estimation. For this reason, we focus on UBU in this paper. Nevertheless, it is important to note that the unbiased estimation approach of this paper is by no means limited to the UBU integrator, and its performance could be further improved by more accurate integrators developed in the future.

2.2. Extension to stochastic gradients. In this subsection, we consider extending splitting methods with the use of stochastic gradients. We use the following definition from [62].

DEFINITION 2.1. *A stochastic gradient approximation of a potential U is defined by a function $\mathcal{G} : \mathbb{R}^d \times \Omega \rightarrow \mathbb{R}^d$ and a probability distribution ρ on a Polish space Ω , such that for every $x \in \mathbb{R}^d$, $\mathcal{G}(x, \cdot)$ is measurable on (Ω, \mathcal{F}) , and for $\omega \sim \rho$,*

$$\mathbb{E}(\mathcal{G}(x, \omega)) = \nabla U(x).$$

The function \mathcal{G} and the distribution ρ together define the stochastic gradient, which we denote as (\mathcal{G}, ρ) .

Replacing the exact gradients with such stochastic gradients in the \mathcal{B} step yields

$$(2.6) \quad \mathcal{B}_{\mathcal{G}}(x, v, h, \omega) = (x, v - h\mathcal{G}(x, \omega)),$$

and we can use this inside BAOAB and UBU to obtain stochastic gradient variants.

[62] has proven convergence bounds for BAOAB with stochastic gradients in Wasserstein distance and also shown that some widely used stochastic gradient schemes (random sampling with replacement, control variate gradient estimator).

3. Unbiased multilevel Monte Carlo methods. In this section, we introduce and motivate our proposed algorithm, which we refer to as Unbiased UBU (UBUBU). We first describe the basic unbiased Monte Carlo scheme and introduce some essential assumptions. We then give relevant results which help to motivate our estimator, including a central limit theorem, a non-asymptotic bound on the variance with exact gradients, and other related results. Finally, we state our algorithm.

Suppose that for each $h \in (0, h_{\max}]$ (stepsize parameter), Q_h is a Markov kernel on some Polish state space Λ with stationary distribution μ_h such that μ_h converges to μ in distribution as $h \rightarrow 0$ (for example, these might be discretizations of a diffusion with different time stepsizes). Assume that we are interested in computing the expectation $\mu(f)$ of a function f satisfying $\pi_h(f^2) < \infty$ for every $h \in (0, h_{\max}]$ and $\mu(f^2) < \infty$. [85] suggested a multilevel estimation method based on stepsizes

$$(3.1) \quad h_0 \in (0, h_{\max}] \text{ and } h_l = h_0 \cdot 2^{-l} \text{ for } l = 1, 2, \dots,$$

using a telescopic sum of the form

$$\mu(f) = \mu_{h_0}(f) + \sum_{j=1}^{\infty} (\mu_{h_j}(f) - \mu_{h_{j-1}}(f)).$$

Unbiased estimators of each term in the sum can be constructed via coupling. A challenge with this approach is that obtaining an unbiased estimator for $\mu_{h_0}(f)$ already requires two chains to be coupled based on the approach proposed in the papers [21, 44, 46, 49]. Estimating the expectations $\mu_{h_j}(f) - \mu_{h_{j-1}}(f)$ is even more challenging, requiring the coupling of four chains. The nature of the couplings means that it is not straightforward to use splitting methods such as UBU or BAOAB (as Markov kernels from different starting points need to be coupled closely in total variation distance, and this is difficult unless the distributions are Gaussian).

To overcome such issues, we propose a different telescoping sum for estimating $\mu(f)$,

$$(3.2) \quad \mu(f) = \tilde{\mu}_{h_0}(f) + \sum_{l=0}^{\infty} (\tilde{\mu}_{h_{l+1}}(f) - \tilde{\mu}_{h_l}(f)).$$

Here $\tilde{\mu}_{h_l}$ are created using some empirical averages, which will be defined in the rest of this section for exact, stochastic, and approximate gradients.

Suppose that D_0 is a random variable satisfying that $\mathbb{E}(D_0) = \tilde{\mu}_0(f)$. Let $\{D_0^{(r)}\}_{r=1}^N$ be N i.i.d. copies of D_0 , and we define

$$(3.3) \quad S_0 = \frac{1}{N} \sum_{r=1}^N D_0^{(r)}.$$

Then it is clear that $\mathbb{E}(S_0) = \mathbb{E}(D_0) = \tilde{\mu}_{h_0}(f)$.

Let $D_{l,l+1}$ be a random variable such that

$$\mathbb{E}D_{l,l+1} = \tilde{\mu}_{h_{l+1}}(f) - \tilde{\mu}_{h_l}(f).$$

Let $c_{0,1}, c_{1,2}, \dots$ be positive constants such that $c_{l,l+1} \rightarrow 0$ as $l \rightarrow \infty$, and let

$$(3.4) \quad \begin{aligned} L(N) &= \max \{l \in \mathbb{N} : c_{l,l+1}N \geq 0.5\}, \\ N_{l,l+1} &= \lceil c_{l,l+1}N \rceil \text{ for } l \leq L(N), \\ N_{l,l+1} &\sim \text{Bernoulli}(c_{l,l+1}N) \text{ for } l > L(N). \end{aligned}$$

For each $l \geq 1$, let $\{D_{l,l+1}^{(r)}\}_{r=1}^{N_{l,l+1}}$ be $N_{l,l+1}$ i.i.d. copies of $D_{l,l+1}$, and

$$(3.5) \quad S_{l,l+1} = \frac{1}{\mathbb{E}(N_{l,l+1})} \sum_{r=1}^{N_{l,l+1}} D_{l,l+1}^{(r)}.$$

It is clear from the definitions and Wald's equation that

$$\mathbb{E}S_{l,l+1} = \mathbb{E}D_{l,l+1} = \tilde{\mu}_{h_{l+1}}(f) - \tilde{\mu}_{h_l}(f).$$

Our first estimator is defined as

$$(3.6) \quad S = S_0 + \sum_{l=0}^{\infty} S_{l,l+1},$$

where the terms $S_0, S_{0,1}, S_{1,2}, \dots$ are independent.

The random $D_{l,l+1}$ variable will play a key role in our approach, as it is going to link two different discretization levels with stepsizes h_l and h_{l+1} . $\text{Var}(S)$ depends on $\text{Var}(D_{l,l+1})$, which is determined by how closely we couple the two discretizations. This is closely related to the strong order of the discretizations, determining how close they are to the underlying diffusion.

It is possible to improve estimator (3.6) slightly by the use of Richardson extrapolation [81]. The idea is that when h is sufficiently small, for Q_h defined in terms of an SDE discretization, the differences $\mu_h(f) - \mu(f)$ tend to follow a certain asymptotic behaviour in h , which can be characterized by an asymptotic expansion [59, 61]. For symmetric splittings like BAOAB it is known that $\mu_h(f) - \mu(f) = c_{f,\mu} h^2 (1 + \mathcal{O}(h))$ for some constant $c_{f,\mu}$ depending on f and μ . The same property can be established for UBU, using similar arguments. Based on this observation, and taking into account that such behaviour may only be valid at small stepsizes, our refined estimator is defined as

$$(3.7) \quad S(c_R) = S_0 + \sum_{l=0}^{L(N)-1} S_{l,l+1} + \frac{S_{L(N),L(N)+1}}{1-c_R} + \sum_{l=L(N)+1}^{\infty} \bar{S}_{l,l+1},$$

$$\bar{S}_{l,l+1} = \frac{1}{\mathbb{E}(N_{l,l+1})} \sum_{r=1}^{N_{l,l+1}} \left[D_{l,l+1}^{(r)} - S_{L(N),L(N)+1} \cdot c_R^{l-L(N)} \right],$$

where $c_R \in [0, 1)$ can be any number (we state the recommended choice of this in our algorithms). Our first estimator S is a special case since $S(0) = S$.

The key assumptions we make on the variances are as follows:

ASSUMPTION 3.1. $f : \Lambda \rightarrow \mathbb{R}$ is a measurable function. $(\tilde{\mu}_{h_l})_{l \geq 0}$ is a sequence of distributions satisfying that $\tilde{\mu}_{h_l}(f) \rightarrow \mu(f)$ as $l \rightarrow \infty$. The random variable D_0 satisfies that $\mathbb{E}(D_0) = \mu_{h_0}(f)$, $\text{Var}(D_0) < \infty$, for every $l \geq 0$, the random variable $D_{l,l+1}$ satisfies that $\mathbb{E}(D_{l,l+1}) = \tilde{\mu}_{l+1}(f) - \tilde{\mu}_{l+1}(f)$ and $|\mathbb{E}(D_{l,l+1}^2)| \leq V_D \phi_D^{-l}$ for some finite constants $V_D > 0$, $\phi_D > 2$.

ASSUMPTION 3.2. The constants $c_{l,l+1}$ controlling $N_{l,l+1}$ satisfy

$$\underline{c}_N \phi_N^{-l} \leq c_{l,l+1} \leq \bar{c}_N \phi_N^{-l},$$

for some finite constants $0 < \underline{c}_N \leq \bar{c}_N$, $\phi_N > 2$.

ASSUMPTION 3.3. The computational cost of generating a sample from $D_{l,l+1}$ is $\mathcal{O}(2^l(K + lB + B_0))$ for some finite constants B , B_0 , and generating a sample from D_0 has a finite computational cost.

PROPOSITION 3.4. Suppose that Assumptions 3.1, 3.2 and 3.3 hold, and that $2 < \phi_N < \phi_D$. Then S as defined in (3.6) is an unbiased estimator of $\mu(f)$ that has finite variance

$$\text{Var}(S) \leq \frac{\text{Var}(D_0)}{N} + \frac{V_D}{N \underline{c}_N \left(1 - \frac{\phi_D}{\phi_N}\right)},$$

and finite expected computational cost.

Similarly, for any $c_R \in [0, 1)$, $S(c_R)$ as defined in (3.7) is also an unbiased estimator of $\mu(f)$ with finite variance

$$\text{Var}(S(c_R)) \leq \frac{\text{Var}(D_0)}{N} + \frac{V_D}{N \underline{c}_N \left(1 - \frac{\phi_N}{\phi_D}\right)} + \frac{V_D}{N \underline{c}_N \left(1 - \frac{\phi_N}{\phi_D}\right)} \frac{2}{(1-c_R)^2} \left(\frac{\phi_N}{\phi_D}\right)^{\log(2\underline{c}_N N / \phi_N) / \log(\phi_N)},$$

and finite expected computational cost.

PROOF. See Section B of the Appendix. \square

We show below that a Central Limit Theorem (CLT) holds for these estimators.

THEOREM 3.5. *Under the assumptions of Proposition 3.4, we have that, as $N \rightarrow \infty$,*

$$\sqrt{N}(S - \mu(f)) \Rightarrow \mathcal{N}(0, \sigma_S^2) \quad \text{and} \quad \sqrt{N}(S(c_R) - \mu(f)) \Rightarrow \mathcal{N}(0, \sigma_S^2),$$

where

$$(3.8) \quad \sigma_S^2 := \text{Var}(D_0) + \sum_{l=0}^{\infty} \frac{\text{Var}(D_{l,l+1})}{c_{l,l+1}}.$$

PROOF. See Appendix B. \square

3.1. *UBUBU with exact gradients.* Now, we will specify the way D_0 and $D_{l,l+1}$ are defined based on UBU discretization of (1.1) with exact gradients, as defined in (2.5). Let μ_0 be an initial distribution on Λ that we can readily sample from, for example, a Dirac- δ at the maximum-a-posteriori (MAP) estimator. Let

$$(3.9) \quad R_0 = P_{h_0} \text{ and } R_l = P_{h_l}^{2^l} \text{ for } l = 1, 2, \dots$$

These Markov kernels correspond to the same amount of time h_0 in the timescale of the limiting diffusion (and clearly, R_l still has μ_{h_l} as its stationary distribution). Consider B_0 burn-in steps with kernel R_0 at level 0, and $B_l = B_0 + lB$ steps with kernel R_l at level l . Define the approximate versions of μ_{h_l} as

$$(3.10) \quad \tilde{\mu}_{h_l} = \frac{1}{K} \sum_{i=1}^K \mu_0 R_l^{B_l+i}.$$

Estimates with respect to this can be computed by taking B_l burn-in steps according to R_l (equivalently $2^l B_l$ burn-in steps according to P_{h_l}), and then K additional steps that are used for computing an empirical average. In this way, we can compute expectations with respect to $\tilde{\mu}_{h_l}$ without the use of couplings. Moreover, given that at the diffusion time scale, the burn-in time tends to infinity as l grows, it is reasonable to expect that under suitable assumptions, $\tilde{\mu}_{h_l}$ converges to μ as $l \rightarrow \infty$.

Let D_0 be the empirical average of a function f based on K samples from Markov chain with kernel R_0 with burn-in B_0 initiated from μ_0 , i.e. for the Markov chain $z_{-B_0}^{(0)} \sim \mu_0$, $z_{-B_0+1}^{(0)} \sim R_0(z_{-B_0}^{(0)}, \cdot)$, \dots , $z_K^{(0)} \sim R_0(z_{K-1}^{(0)}, \cdot)$. Let ν_0 denote the joint distribution of $z_{-B_0}^{(0)}, \dots, z_K^{(0)}$, and define

$$(3.11) \quad D_0 = \frac{1}{K} \sum_{i=1}^K f(z_i^{(0)}).$$

Let $\{D_0^{(r)}\}_{r=1}^N$ be N i.i.d. copies of D_0 , and define

$$(3.12) \quad S_0 = \frac{1}{N} \sum_{r=1}^N D_0^{(r)}.$$

Then it is clear that $\mathbb{E}(S_0) = \mathbb{E}(D_0) = \tilde{\mu}_{h_0}(f)$. For $l \geq 0$, let $z_{-B_l}^{(l,l+1)}, \dots, z_K^{(l,l+1)}, z_{-B_{l+1}}^{(l,l+1)}, \dots, z_K^{(l,l+1)}$ be Λ valued random variables defined on the same probability space (i.e. coupled) such that

- $z_{-B_l}^{(l,l+1)}, \dots, z_K^{(l,l+1)}$ is a Markov chain with kernel R_l initiated as $z_{-B_l}^{(l,l+1)} \sim \mu_0$, and
- $z'_{-B_{l+1}}^{(l,l+1)}, \dots, z'_K{}^{(l,l+1)}$ is a Markov chain with kernel R_{l+1} initiated $z'_{-B_{l+1}}{}^{(l,l+1)} \sim \mu_0$.

Let

$$(3.13) \quad D_{l,l+1} = \frac{1}{K} \sum_{i=1}^K [f(z'_i{}^{(l,l+1)}) - f(z_i^{(l,l+1)})].$$

From the definitions, it follows that

$$\mathbb{E}D_{l,l+1} = \tilde{\mu}_{h_{l+1}}(f) - \tilde{\mu}_{h_l}(f),$$

hence $D_{l,l+1}$ is an unbiased estimator of the difference $\tilde{\mu}_{h_{l+1}}(f) - \tilde{\mu}_{h_l}(f)$.

When these Markov chains are discretizations of the same diffusion, it is natural to create synchronous couplings by using the same Brownian noise to generate the Gaussian random variables used during the periods z_{-B_l}, \dots, z_K and z'_{-B_l}, \dots, z'_K . Such couplings can significantly reduce the variance of $D_{l,l+1}$. Let \mathcal{B} and \mathcal{U} be as in (2.2-2.3). Further we define \mathcal{U}^2 to be

$$(3.14) \quad \mathcal{U}^2(x, v, h, \xi^{(1)}, \xi^{(2)}, \xi^{(3)}, \xi^{(4)}) = \mathcal{U}\left(\mathcal{U}\left(x, v, h/2, \xi^{(1)}, \xi^{(2)}\right), h/2, \xi^{(3)}, \xi^{(4)}\right).$$

As \mathcal{U} is an exact solution in the weak sense to its respective component in the splitting, this is an exact solution in the weak sense which uses Brownian increments $(\xi^{(1)}, \xi^{(2)})$ in the first half step $h/2$ and $(\xi^{(3)}, \xi^{(4)})$ in the second half step $h/2$. The \mathcal{U}^2 operator is an exact solution over stepsize h .

A coupling can be constructed between discretization levels so that the two discretization levels share Brownian motion in the exact integration of the \mathcal{U} steps. This is done by using the Brownian increments from two respective \mathcal{U} solutions at the higher level and concatenating them using the \mathcal{U}^2 operator at the lower level. Next, the stochastic integrals in the two levels are coupled by sharing the same Brownian noise. The Markov kernel $P_{h,h/2}$ for the two discretization levels $h, h/2$ is defined as follows.

(3.15)

$$\begin{aligned} & \left(\xi_{k+1}^{(i)}\right)_{i=1}^8 \sim \mathcal{N}(0_d, I_d) \text{ for all } i = 1, \dots, 8. \\ & \left(x'_{k+1/2}, v'_{k+1/2}\right) = \mathcal{U}\left(\mathcal{B}\left(\mathcal{U}\left(x'_k, v'_k, h/4, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}\right), h/2\right), h/4, \xi_{k+1}^{(3)}, \xi_{k+1}^{(4)}\right) \\ & \left(x'_{k+1}, v'_{k+1}\right) = \mathcal{U}\left(\mathcal{B}\left(\mathcal{U}\left(x'_{k+1/2}, v'_{k+1/2}, h/4, \xi_{k+1}^{(5)}, \xi_{k+1}^{(6)}\right), h/2\right), h/4, \xi_{k+1}^{(7)}, \xi_{k+1}^{(8)}\right) \\ & \left(x_{k+1}, v_{k+1}\right) = \\ & \mathcal{U}^2\left(\mathcal{B}\left(\mathcal{U}^2\left(x_k, v_k, h/2, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}, \xi_{k+1}^{(3)}, \xi_{k+1}^{(4)}\right), h\right), h/2, \xi_{k+1}^{(5)}, \xi_{k+1}^{(6)}, \xi_{k+1}^{(7)}, \xi_{k+1}^{(8)}\right). \end{aligned}$$

This Markov chain acts on the state space $\mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^d$, moving from (x_k, v_k, x'_k, v'_k) to $(x_{k+1}, v_{k+1}, x'_{k+1}, v'_{k+1})$ via the steps in (3.15). When looking at the individual components, $(x_k, v_k) \rightarrow (x_{k+1}, v_{k+1})$ corresponds to one UBU step at stepsize h , while $(x'_k, v'_k) \rightarrow (x'_{k+1}, v'_{k+1})$ corresponds to two UBU steps at stepsize $h/2$. A key property here is that the stochastic integrals between two steps are synchronously coupled, which ensures that these two chains approximate the same underlying diffusion (in the strong sense). Hence, they are expected to remain close, which was observed in our numerical simulations.

We now create a coupling between levels l and $l+1$, denoted by $\nu_{l,l+1}$.

 $\nu_{l,l+1}$ coupling

- 1: For given initial distribution μ_0 on Λ , we define $z_{-B_l}^{(l,l+1)} \sim \mu_0$ and $z_{-B_{l+1}}^{(l,l+1)} \sim \mu_0$ as independent random variables.
 - 2: We let $z_{-B_{l+1}}^{(l,l+1)}, \dots, z_{-B_l}^{(l,l+1)}$ be a Markov chain evolving according to $R_{l+1} = (P_{h_{l+1}})^{2^{l+1}}$.
 - 3: Let $(z_{-B_l}^{(l,l+1)}, z_{-B_l}^{(l,l+1)}), (z_{-B_{l+1}}^{(l,l+1)}, z_{-B_{l+1}}^{(l,l+1)}), \dots, (z_K^{(l,l+1)}, z_K^{(l,l+1)})$ be a Markov chain evolving according to $R_{l,l+1} = (P_{h_l, h_{l+1}})^{2^l}$.
 - 4: Let $\nu_{l,l+1}$ denote the joint distribution of $z_{-B_l}^{(l,l+1)}, \dots, z_K^{(l,l+1)}, z_{-B_{l+1}}^{(l,l+1)}, \dots, z_K^{(l,l+1)}$.
-

The motivation for this $\nu_{l,l+1}$ coupling is that if two coupled chains are driven by the same noise and approximate the same diffusion, they are expected to be close most of the time. Given a sufficiently long burn-in, they will likely stay close during the iterations $1, 2, \dots, K$ used for computing the differences in their empirical averages, reducing the variance of $D_{l,l+1}$. Let $c_N > 0$ and $\phi_N > 2$ be constants, and let

$$(3.16) \quad c_{l,l+1} = c_N \phi_N^{-l} \text{ for } l \in \mathbb{N}.$$

We call the overall estimator $S(c_R)$ based on formula (3.7) with $D_{l,l+1}$ defined based on coupling construction $\nu_{l,l+1}$ as *Unbiased UBU* (or *UBUBU*, for short). The steps for constructing this estimator are summarized in Algorithm 1.

Now, we will state our theoretical results for this algorithm. To prove unbiasedness and finite variance for our estimator $S(c_R)$, we require several assumptions, which we state below. These include assumptions on the smoothness and strong convexity of our potential, as well as restrictions on various parameters of the algorithm.

ASSUMPTION 3.6 (*M- ∇ Lipschitz*). $U : \mathbb{R}^d \rightarrow \mathbb{R}$ is twice continuously differentiable and there exists $M > 0$ such that for all $x, y \in \mathbb{R}^d$

$$\|\nabla U(x) - \nabla U(y)\| \leq M\|x - y\|.$$

ASSUMPTION 3.7 (*m-strong convexity*). $U : \mathbb{R}^d \rightarrow \mathbb{R}$ is continuously differentiable and there exists $m > 0$ such that for all $x, y \in \mathbb{R}^d$

$$\langle \nabla U(x) - \nabla U(y), x - y \rangle \geq m|x - y|^2.$$

The strongly Hessian Lipschitz property relies on the following tensor norm from [24], which we require in our setup.

DEFINITION 3.8. For $A \in \mathbb{R}^{d \times d \times d}$, let

$$\|A\|_{\{1,2\}\{3\}} = \sup_{x \in \mathbb{R}^{d \times d}, y \in \mathbb{R}^d} \left\{ \sum_{i,j,k=1}^d A_{ijk} x_{ij} y_k \left| \sum_{i,j=1}^d x_{ij}^2 \leq 1, \sum_{k=1}^d y_k^2 \leq 1 \right. \right\}.$$

REMARK 3.9. The $\|A\|_{\{1,2\}\{3\}}$ norm in Definition 3.8 can be equivalently written as

$$(3.17) \quad \|A\|_{\{12\}\{3\}} = \left\| \sum_{i_1} A_{i_1, \cdot, \cdot}^T \cdot A_{i_1, \cdot, \cdot} \right\|^{1/2},$$

where $A_{i_1, \cdot, \cdot} = (A_{i_1, i_2, i_3})_{1 \leq i_2 \leq d, 1 \leq i_3 \leq d}$ is a $d \times d$ matrix, see the proof of Lemma 7 of [77].

Algorithm 1 Unbiased-UBU (UBUBU)

1: **Input:**

- Maximum stepsize h_0 .
- Friction parameter $\gamma > 0$.
- Initial distribution μ_0 on $\mathbb{R}^d \times \mathbb{R}^d$ for $l \geq 0$.
- Potential function $U : \mathbb{R}^d \rightarrow \mathbb{R}$ of target distribution.
- Burn-in length parameters B_0 and B .
- Number of samples parameter K .
- Number of parallel chains parameters N , c_N and ϕ_N .
- Richardson extrapolation parameter $c_R \in [0, 1)$ (default value $c_R = \frac{1}{4}$).
- Test function f .

2: **Averages from level 0:**

3: **for** $r = 1, \dots, N$ **do**

4: Sample $z_{-B_0}^{(0,r)}, \dots, z_K^{(0,r)}$ from ν_0 .

5: Compute $D_0^{(r)}$ based on (3.11) using the samples $z_1^{(0,r)}, \dots, z_K^{(0,r)}$.

6: **end for**

7: Compute S_0 using (3.12).

8: **Generate number of chains:**

9: Sample $N_{l,l+1}$ according to (3.4), let $l_{\max} = \max\{l : N_{l,l+1} > 0\}$.

10: **Averages of differences** $D_{l,l+1}$ **from** $l = 0, \dots, l_{\max}$:

11: **for** $l = 0, \dots, l_{\max}$ **do**

12: **for** $r = 1, \dots, N_{l,l+1}$ **do**

13: Sample $z_{-B_l}^{(l,l+1,r)}, \dots, z_K^{(l,l+1,r)}, z_{-B_{l+1}}^{l(l,l+1,r)}, \dots, z_K^{l(l,l+1,r)}$ according to $\nu_{l,l+1}$.

14: Compute $D_{l,l+1}^{(r)}$ based on (3.13) using $z_1^{(r,l,l+1)}, \dots, z_K^{(r,l,l+1)}, z_1^{l(r,l,l+1)}, \dots, z_K^{l(r,l,l+1)}$.

15: **end for**

16: Compute $S_{l,l+1}$ using (3.5).

17: **end for**

18: Compute $S(c_R)$ using (3.7).

19: **Output:**

20: Unbiased estimator $S(c_R)$,

21: Samples $z_1^{(0,r)}, \dots, z_K^{(0,r)}$ for parallel chains $1 \leq r \leq N$,

22: Samples $z_1^{(l,l+1,r)}, \dots, z_K^{(l,l+1,r)}, z_1^{l(r,l,l+1)}, \dots, z_K^{l(r,l,l+1)}$ for $0 \leq l \leq l_{\max}$, chains $1 \leq r \leq N_{l,l+1}$.

ASSUMPTION 3.10 (M_1^s -strongly Hessian Lipschitz). $U : \mathbb{R}^d \rightarrow \mathbb{R}$ is three times continuously differentiable and M_1^s -strongly Hessian Lipschitz if there exists $M_1^s > 0$ such that

$$\|\nabla^3 U(x)\|_{\{1,2\}\{3\}} \leq M_1^s$$

for all $x \in \mathbb{R}^d$.

REMARK 3.11. In Lemma H.6 in the Appendix, we show that Bayesian multinomial regression satisfies this assumption.

ASSUMPTION 3.12 (1-Lipschitzness of f). f is a 1-Lipschitz function with respect to the Euclidean distance on \mathbb{R}^{2d} , that only depends on x , not v (i.e. $f(x, v) = f(x, v')$ for any $x, v, v' \in \mathbb{R}^d$).

ASSUMPTION 3.13 (Distance of initial distribution from target). The initial distribution on $\Lambda = \mathbb{R}^{2d}$ satisfy that $\mathcal{W}_2(\pi, \mu_0) \leq c_{\mu_0} \sqrt{\frac{d}{m}}$, for some $c_{\mu_0} > 0$.

REMARK 3.14. *It is easy to show that under Assumption (3.7), for $\mu_0 = \delta_{x^*} \times \mathcal{N}(0_d, I_d)$, and for $\mu_0 = \mathcal{N}(x^*, (\nabla^2 U(x^*))^{-1}) \times \mathcal{N}(0_d, I_d)$ (Gaussian approximation), this condition holds with $c_{\mu_0} = 2$.*

The heart of the work is related to demonstrating unbiasedness and finite variance as a result of the multilevel scheme presented in Figure 2. We now state our first main result, which is a non-asymptotic bound on the variance of our estimator (3.7).

THEOREM 3.15. *Suppose that Assumptions 3.6, 3.7, 3.10, 3.12, 3.13 hold, and in addition,*

$$\gamma \geq \sqrt{8M}, \quad h_0 \leq \frac{1}{\gamma} \cdot \frac{m}{264M}, \quad B \geq \frac{16 \log(4)\gamma}{mh_0}, \quad B_0 \geq \frac{16\gamma}{mh_0} \log \left(\frac{c_{\mu_0} + 1}{\sqrt{M}\gamma h_0^2} \right).$$

Suppose that $c_R \in [0, 1)$, and $2 < \phi_N < 16$. Then for any $N \geq 1$, the UBUBU estimator $S(c_R)$ has finite expected computational cost, $\mathbb{E}S(c_R) = \pi(f)$, and it has finite variance. Moreover, it satisfies a CLT as $N \rightarrow \infty$, and the asymptotic variance σ_S^2 defined in (3.8) can be bounded as

$$\sigma_S^2 \leq \frac{C(m, M, M_1^s, \gamma, c_N, \phi_N)}{Kh_0} \left(1 + \frac{1}{h_0 K} + dh_0^4 \right).$$

PROOF. See Section D.3 in the Appendix. \square

REMARK 3.16. *In particular, when setting $h_0 = \mathcal{O}(d^{-1/4})$, and $K > 1/h_0$, the bound simplifies to $\text{Var}(S(c_R)) \leq \frac{C(\gamma, m, M, M_1)}{NKh_0}$. This indicates that the overall number of gradient evaluations per effective sample in this setting is $\mathcal{O}(d^{1/4})$, which matches the best available bounds for HMC in [24], without the warm start assumption required in that paper.*

The following proposition shows dimension-free bounds for product distributions. We are going to use an assumption on the initial distribution μ_0 .

ASSUMPTION 3.17. *Suppose that μ_0 and the target distribution π are of product form*

$$\mu_0(dx, dv) = \prod_{i=1}^d \mu_{0,i}(dx_i, dv_i) \quad \text{for all } l \geq 0, \quad \pi(dx, dv) = \prod_{i=1}^d \tilde{\pi}_i(dx_i) \frac{e^{-v_i^2/2} dv_i}{\sqrt{2\pi}},$$

for $x = (x_1, \dots, x_d) \in \mathbb{R}^d$, $v = (v_1, \dots, v_d) \in \mathbb{R}^d$, and that

$$\max_{1 \leq i \leq d} \mathcal{W}_2(\pi_i, \mu_{0,i}) \leq c_{\mu_0} \sqrt{\frac{1}{m}},$$

for some finite constant c_{μ_0} , where $\pi_i(dx_i, dv_i) = \tilde{\pi}_i(dx_i) \frac{e^{-v_i^2/2}}{\sqrt{2\pi}} dv_i$ is the joint distribution of (x_i, v_i) according to the target π .

PROPOSITION 3.18. *Suppose that Assumption 3.17 holds, and denote the potential U as $U(x) = \sum_{i=1}^d U_i(x_i)$. Suppose that Assumptions 3.6, 3.7, and 3.10 hold for each component $(U_i)_{1 \leq i \leq d}$, and that*

$$\gamma \geq \sqrt{8M}, \quad h_0 \leq \frac{1}{\gamma} \cdot \frac{m}{264M}, \quad B \geq \frac{16 \log(4)\gamma}{mh_0}, \quad B_0 \geq \frac{16\gamma}{mh_0} \log \left(\frac{c_{\mu_0} + 1}{\sqrt{M}\gamma h_0^2} \right).$$

Suppose that f is of the form

$$(3.18) \quad f(x, v) = g(\langle w^{(1)}, x \rangle, \dots, \langle w^{(r)}, x \rangle),$$

where $g : \mathbb{R}^r \rightarrow \mathbb{R}$ is 1-Lipschitz, and $w^{(1)}, \dots, w^{(r)} \in \mathbb{R}^d$. Suppose that $c_R \in [0, 1)$ and $2 < \phi_N < 16$. Then for any $N \geq 1$, the UBUBU estimator $S(c_R)$ has finite expected computational cost, $\mathbb{E}S(c_R) = \pi(f)$, and it has finite variance. Moreover, it satisfies a CLT as $N \rightarrow \infty$, and the asymptotic variance can be bounded as

$$\sigma_S^2 \leq \frac{C(m, M, M_1, \gamma, r, c_N, \phi_N)}{Kh_0} \sum_{1 \leq i \leq r} \|w^{(i)}\|^2.$$

PROOF. See Section D.3 in the Appendix. \square

REMARK 3.19. *These bounds are independent of the dimension d . This is not surprising as the different components evolve independently according to the kinetic Langevin diffusion (1.1), and we do not introduce any dependencies in the UBUBU algorithm. This is in contrast with Metropolized methods, where the accept/reject steps introduce dependencies in the evolution of the components. The results could be generalized to potentials which are separable into independent groups of coordinates, i.e. $U(x) = \sum_{i=1}^s U_i(x_{G_i})$, where G_1, \dots, G_s is a partition of $[d]$, and the size of each group $|G_i|$ is small.*

3.2. *UBUBU with stochastic gradients.* In this section, we extend the unbiased estimation methods of the previous section to the setting where we have instead stochastic or approximate gradient evaluations, combined with a control variate approach that occasionally computes full gradients for variance reduction, as in [50, 101].

In many applications, particularly in data science and machine learning, gradient computations are computationally expensive due to large datasets and the need to iterate through the entire dataset at each gradient evaluation. A common approach for reducing the cost of the gradient-based methods is to use stochastic gradient approximations based on subsampling the dataset to compute unbiased estimates (see [5, 18, 50, 79, 91, 95]).

In these applications the potential $U : \mathbb{R}^d \rightarrow \mathbb{R}$ is typically of the form

$$(3.19) \quad U(x) = U_0(x) + \sum_{i=1}^{N_D} U_i(x),$$

where $x \in \mathbb{R}^d$, the dataset is of size $N_D \in \mathbb{N}$. U_0 can be chosen as the negative log density of the prior distribution or some other term that does not require accessing the data. In our examples, U_0 can be taken to be a quadratic function, for example, a quadratic matching the Hessian at the minimizer (which can be computed before sampling).

We remark that one of the most efficient samplers in the big data regime is the Zig-Zag sampler [8] whose complexity is independent of the data size according to a limiting argument (although as stated in [8], some logarithmic factors were ignored). [27] is another recent paper that proposes a Metropolis-Hastings-type MCMC algorithm based on subsampling that only accesses $\mathcal{O}(1)$ or even $\mathcal{O}(1/\sqrt{N_D})$ data points per step. Although this method was shown to have state-of-the-art performance on a 10-dimensional logistic regression example, its efficiency on high-dimensional models has not yet been demonstrated.

In this section, we will develop a version of UBUBU using stochastic gradients. We are going to use random variables of the form $\omega \in [N_D]^{N_b}$, which is a random selection of N_b indices to be selected uniformly on $[N_D] = \{1, \dots, N_D\}$, i.i.d. with replacement [5]. We denote the distribution of ω here as $\mathcal{SWR}(N_D, N_b)$.

DEFINITION 3.20. *The sub-sampled stochastic gradient of U at x with respect to \hat{x} is*

$$(3.20) \quad \mathcal{G}(x, \omega | \hat{x}) = \nabla U_0(x) + \sum_{i=1}^{N_D} \nabla U_i(\hat{x}) + \frac{N_D}{N_b} \sum_{i \in \omega} [\nabla U_i(x) - \nabla U_i(\hat{x})],$$

where $\omega \sim \mathcal{SWR}(N_D, N_b)$.

$\mathcal{G}(\cdot | \hat{x}) : \mathbb{R}^d \times \Omega \rightarrow \mathbb{R}^d$ is an unbiased estimator of $\nabla U(x)$ in the sense of Definition 2.1. We can use this estimator in UBU by replacing the \mathcal{B} step with

$$(3.21) \quad \mathcal{B}_{\mathcal{G}}(x, v, h, \omega | \hat{x}) = (x, v - h\mathcal{G}(x, \omega | \hat{x})).$$

Let $x^* \in \mathbb{R}^d$ be the minimizer of the potential U , then the selection $\hat{x} = x^*$ at each step corresponds to the control variate gradient estimator, see [5]. When approximating the step \mathcal{B} in UBU using this control variate approach, we can only achieve strong order 1/2.

Another possibility is to update \hat{x} every $\tau = \lceil N_D/N_b \rceil$ iterations with the latest position where the gradient was evaluated (this is not x_k for UBU as the gradients are evaluated after moving forward by a \mathcal{U} step with stepsize $h/2$). We refer to this as the stochastic variance reduced gradient (SVRG) approach (see [50, 101]). The overall computational cost of this approach is approximately twice that of the control variate approach (due to the need for a full gradient evaluation). Since the gradient is reevaluated every τ iterations, when h is small, the position \hat{x} becomes closer to the positions x that are considered, and the approximate dynamics provide a better approximation of the underlying diffusion (1.1). We will show that the SVRG discretization has strong order 3/2, which is better than the control variate estimator, hence we will use it within our unbiasing scheme. The evolution of SVRG steps can be written as follows,

$$(3.22) \quad \begin{aligned} & \left(\xi_{k+1}^{(i)} \right)_{i=1}^4 \sim \mathcal{N}(0_d, I_d) \text{ for all } i = 1, \dots, 4. \\ & \omega_{k+1} \sim \mathcal{SWR}(N_D, N_b) \\ & (\bar{x}_k, \bar{v}_k) = \mathcal{U} \left(x_k, v_k, h/2, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)} \right) \\ & \hat{x}_k = \bar{x}_{\lfloor k/\tau \rfloor \tau} \\ & (x_{k+1}, v_{k+1}) = \mathcal{U} \left(\mathcal{B}_{\mathcal{G}}(\bar{x}_k, \bar{v}_k, h, \omega_{k+1} | \hat{x}_k), h/2, \xi_{k+1}^{(3)}, \xi_{k+1}^{(4)} \right). \end{aligned}$$

Let P_h^{SVRG} denote the time inhomogenous Markov kernel describing the evolution of (x_k, \hat{x}_k, v_k) according to the SVRG steps (3.22).

We use a Gaussian approximation of the target. Let

$$(3.23) \quad \mu_G = \mathcal{N}(x^*, (H^*)^{-1}) \times \mathcal{N}(0_d, I_d) \quad \text{with} \quad H^* = \nabla^2 U(x^*),$$

and define

$$(3.24) \quad \mathcal{H}_*(x, v, h) = \begin{pmatrix} x^* \\ 0_d \end{pmatrix} + \exp \left(h \begin{pmatrix} 0_d & I_d \\ -H^* & 0_d \end{pmatrix} \right) \begin{pmatrix} x - x^* \\ v \end{pmatrix},$$

$$(3.25) \quad \mathcal{O}(x, v, h/2, \xi^{(1)}, \xi^{(2)}) = \left(x, \eta v + \sqrt{2\gamma} \mathcal{Z}^{(2)} \left(h/2, \xi^{(1)}, \xi^{(2)} \right) \right),$$

$$(3.26) \quad \mathcal{O}^2(x, v, h, \xi^{(1)}, \xi^{(2)}, \xi^{(3)}, \xi^{(4)}) = \mathcal{O} \left(\mathcal{O} \left(x, v, h/2, \xi^{(1)}, \xi^{(2)} \right), h/2, \xi^{(3)}, \xi^{(4)} \right).$$

with $\mathcal{H}_*(x, v, h)$ corresponding the solution of the Hamiltonian dynamics on target $\mu_G \times \mathcal{N}(0_d, I_d)$ initiated in (x, v) at time h . It follows from (2.4) that $\sqrt{2\gamma} \mathcal{Z}^{(2)}(h/2, \xi^{(1)}, \xi^{(2)}) \sim$

$\mathcal{N}(0_d, (1 - \eta^2)I_d)$, so this \mathcal{O} steps keeps the target invariant. We are going to use the OHO scheme as part of our algorithm.

$$(3.27) \quad \begin{aligned} (\xi_{k+1}^{(i)})_{i=1}^4 &\sim \mathcal{N}(0_d, I_d) \text{ for all } i = 1, \dots, 4. \\ (\bar{x}_k, \bar{v}_k) &= \mathcal{O} \left(x_k, v_k, h/2, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)} \right) \\ (x_{k+1}, v_{k+1}) &= \mathcal{O} \left(\mathcal{H}_* (\bar{x}_k, \bar{v}_k, h), h/2, \xi_{k+1}^{(3)}, \xi_{k+1}^{(4)} \right). \end{aligned}$$

Let P_h^{OHO} denote the time homogeneous Markov kernel describing the evolution of (x_k, v_k) according to the OHO steps (3.27).

Two chains evolving according to SVRG with step size h and SVRG with step size $h/2$ can be coupled as follows.

$$(3.28) \quad \begin{aligned} (\xi_{k+1}^{(i)})_{i=1}^8 &\sim \mathcal{N}(0_d, I_d) \text{ for all } i = 1, \dots, 8. \\ \omega'_{k+1/2}, \omega'_{k+1} &\sim \mathcal{SWR}(N_D, N_b), \\ (\bar{x}_k, \bar{v}_k) &= \mathcal{U}^2 \left(x_k, v_k, h/2, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}, \xi_{k+1}^{(3)}, \xi_{k+1}^{(4)} \right) \\ \hat{x}_k &= \bar{x}_{\lfloor k/\tau \rfloor \tau} \\ (x_{k+1}, v_{k+1}) &= \mathcal{U}^2 \left(\mathcal{B}_G (\bar{x}_k, \bar{v}_k, h, \omega_{k+1} | \hat{x}_k), h/2, \xi_{k+1}^{(5)}, \xi_{k+1}^{(6)}, \xi_{k+1}^{(7)}, \xi_{k+1}^{(8)} \right), \\ (\bar{x}'_k, \bar{v}'_k) &= \mathcal{U} \left(x'_k, v'_k, h/4, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)} \right) \\ \hat{x}'_k &= \bar{x}'_{\lfloor 2k/\tau \rfloor \tau/2} \\ (x'_{k+1/2}, v'_{k+1/2}) &= \mathcal{U} \left(\mathcal{B}_G (\bar{x}'_k, \bar{v}'_k, h/2, \omega'_{k+1/2}, v | \hat{x}'_k), h/4, \xi_{k+1}^{(3)}, \xi_{k+1}^{(4)} \right) \\ (\bar{x}'_{k+1/2}, \bar{v}'_{k+1/2}) &= \mathcal{U} \left(x'_{k+1/2}, v'_{k+1/2}, h/4, \xi_{k+1}^{(5)}, \xi_{k+1}^{(6)} \right) \\ \hat{x}'_{k+1/2} &= \bar{x}'_{\lfloor (2k+1)/\tau \rfloor \tau/2} \\ (x'_{k+1}, v'_{k+1}) &= \mathcal{U} \left(\mathcal{B}_G (\bar{x}'_{k+1/2}, \bar{v}'_{k+1/2}, h/2, \omega'_{k+1} | \hat{x}'_{k+1/2}), h/4, \xi_{k+1}^{(7)}, \xi_{k+1}^{(8)} \right) \end{aligned}$$

Let $P_{h,h/2}^{SVRG}$ denote the time inhomogenous Markov kernel describing the evolution of $(x_k, \hat{x}_k, v_k, x'_k, \hat{x}'_k, v'_k)$ according to the SVRG steps (3.28).

Finally, we will also need to couple one chain with step size h running OHO on the Gaussian approximation μ_G , and another chain running SVRG on the target with step size $h/2$.

$$\begin{aligned}
& \left(\xi_{k+1}^{(i)} \right)_{i=1}^8 \sim \mathcal{N}(0_d, I_d) \text{ for all } i = 1, \dots, 8, \\
& \omega'_{k+1/2}, \omega'_{k+1} \sim \mathcal{SWR}(N_D, N_b), \\
& (\bar{x}_k, \bar{v}_k) = \mathcal{O}^2 \left(x_k, v_k, h/2, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}, \xi_{k+1}^{(3)}, \xi_{k+1}^{(4)} \right) \\
& (x_{k+1}, v_{k+1}) = \mathcal{O}^2 \left(\mathcal{H}_* (\bar{x}_k, \bar{v}_k, h), h/2, \xi_{k+1}^{(5)}, \xi_{k+1}^{(6)}, \xi_{k+1}^{(7)}, \xi_{k+1}^{(8)} \right), \\
(3.29) \quad & (\bar{x}'_k, \bar{v}'_k) = \mathcal{U} \left(x'_k, v'_k, h/4, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)} \right) \\
& \hat{x}'_k = \bar{x}'_{\lfloor 2k/\tau \rfloor \tau/2} \\
& \left(x'_{k+1/2}, v'_{k+1/2} \right) = \mathcal{U} \left(\mathcal{B}_{\mathcal{G}} \left(\bar{x}'_k, \bar{v}'_k, h/2, \omega'_{k+1/2}, v \mid \hat{x}'_k \right), h/4, \xi_{k+1}^{(3)}, \xi_{k+1}^{(4)} \right) \\
& \left(\bar{x}'_{k+1/2}, \bar{v}'_{k+1/2} \right) = \mathcal{U} \left(x'_{k+1/2}, v'_{k+1/2}, h/4, \xi_{k+1}^{(5)}, \xi_{k+1}^{(6)} \right) \\
& \hat{x}'_{k+1/2} = \bar{x}'_{\lfloor (2k+1)/\tau \rfloor \tau/2} \\
& \left(x'_{k+1}, v'_{k+1} \right) = \mathcal{U} \left(\mathcal{B}_{\mathcal{G}} \left(\bar{x}'_{k+1/2}, \bar{v}'_{k+1/2}, h/2, \omega'_{k+1} \mid \hat{x}'_{k+1/2} \right), h/4, \xi_{k+1}^{(7)}, \xi_{k+1}^{(8)} \right)
\end{aligned}$$

Let $P_{h,h/2}^{OH O/SV R G}$ denote the time inhomogenous Markov kernel describing the evolution of $(x_k, v_k, x'_k, \hat{x}'_k, v'_k)$ according to the steps (3.29).

We now create a coupling between levels 0 and 1, denoted by $\nu_{0,1}^{SG}$.

$\nu_{0,1}^{SG}$ coupling

- 1: Define $z_{-B_1}^{(0,1)} \sim \mu_G$ and let $z'_{-B_1}{}^{(0,1)} = z_{-B_1}^{(0,1)}$.
 - 2: Let $(z_{-B_1}^{(0,1)}, z'_{-B_1}{}^{(0,1)}, \hat{x}'_{-B_1}{}^{(0,1)}), (z_{-B_1+1}^{(0,1)}, z'_{-B_1+1}{}^{(0,1)}, \hat{x}'_{-B_1+1}{}^{(0,1)}), \dots, (z_K^{(0,1)}, z'_K{}^{(0,1)}, \hat{x}'_K{}^{(0,1)})$ be a Markov chain with kernel $R_{0,1}^{OH O/SV R G} = P_{h_0, h_1}^{OH O/SV R G}$ (satisfying that $z_k^{(0,1)} \sim \mu_G$ for all k).
 - 3: Let $\nu_{0,1}$ denote the joint distribution of $z_{-B_0}^{(0,1)}, \dots, z_K^{(0,1)}, z'_{-B_1}{}^{(0,1)}, \dots, z'_K{}^{(0,1)}$.
-

We now create a coupling between levels l and $l+1$ for $l \geq 1$, denoted by $\nu_{l,l+1}^{SG}$.

$\nu_{l,l+1}^{SG}$ coupling

- 1: Define $z_{-B_{l+1}}^{(l,l+1)} \sim \mu_G$ and let $z'_{-B_{l+1}}{}^{(l,l+1)} = z_{-B_{l+1}}^{(l,l+1)}$.
 - 2: Let $(z_{-B_{l+1}}^{(l,l+1)}, z'_{-B_{l+1}}{}^{(l,l+1)}, \hat{x}'_{-B_{l+1}}{}^{(l,l+1)}), \dots, (z_{-B_l}^{(l,l+1)}, z'_{-B_l}{}^{(l,l+1)}, \hat{x}'_{-B_l}{}^{(l,l+1)})$ be a Markov chain with kernel $R_{l,l+1}^{OH O/SV R G} = (P_{h_l, h_{l+1}}^{OH O/SV R G})^{2^l}$ (satisfying that $z_{-B_l}^{(l,l+1)} \sim \mu_G$).
 - 3: Let $(z_{-B_l}^{(l,l+1)}, z'_{-B_l}{}^{(l,l+1)}, \hat{x}'_{-B_l}{}^{(l,l+1)}), (z_{-B_{l+1}}^{(l,l+1)}, z'_{-B_{l+1}}{}^{(l,l+1)}, \hat{x}'_{-B_{l+1}}{}^{(l,l+1)}), \dots, (z_K^{(l,l+1)}, z'_K{}^{(l,l+1)}, \hat{x}'_K{}^{(l,l+1)})$ be a Markov chain evolving according to $R_{l,l+1}^{SV R G} = (P_{h_l, h_{l+1}}^{SV R G})^{2^l}$.
 - 4: Let $\nu_{l,l+1}^{SG}$ denote the joint distribution of $z_{-B_l}^{(l,l+1)}, \dots, z_K^{(l,l+1)}, z'_{-B_{l+1}}{}^{(l,l+1)}, \dots, z'_K{}^{(l,l+1)}$.
-




Fig 3: Coupling scheme for UBUBU-SG.

Figure 3 illustrates our couplings between different levels using OHO/UBU discretizations. Given some constants $c_N > 0$, $\phi_N > 2$, we let

$$(3.30) \quad c_{l,l+1} = c_N \phi_N^{-l} \text{ for } l \in \mathbb{N}.$$

Our stochastic gradient-based method (UBUBU-SG) proceeds as stated in Algorithm 2. We recommend setting the Richardson extrapolation parameter $c_R = \frac{1}{2\sqrt{2}}$ in this case (as SVRG has strong order 3/2).

In order to show variance bounds for this algorithm, we make the following assumptions.

ASSUMPTION 3.21 (∇ Lipschitz property). For every $1 \leq i \leq N_D$, $U_i : \mathbb{R}^d \rightarrow \mathbb{R}$ is twice differentiable and there exists a $\tilde{M} > 0$ such that for all $x, y \in \mathbb{R}^d$,

$$\|\nabla U_i(x) - \nabla U_i(y)\| \leq \tilde{M} \|x - y\|,$$

for every $1 \leq i \leq N_D$ and moreover,

$$\|\nabla U(x) - \nabla U(y)\| \leq M \|x - y\| \quad \text{for } M = N_D \tilde{M}.$$

ASSUMPTION 3.22 ($N_D \tilde{m}$ -strong convexity). There exists a $\tilde{m} > 0$ such that for all $x, y \in \mathbb{R}^d$

$$\langle \nabla U(x) - \nabla U(y), x - y \rangle \geq m \|x - y\|^2 \quad \text{for } m = N_D \tilde{m}.$$

ASSUMPTION 3.23 (strongly Hessian Lipschitz property). $U : \mathbb{R}^d \rightarrow \mathbb{R}$ is three times continuously differentiable and M_1^s -strongly Hessian Lipschitz if there exists $M_1^s > 0$ such that

$$\|\nabla^3 U(x)\|_{\{1,2\}\{3\}} \leq M_1^s \quad \text{for } M_1^s = N_D \tilde{M}_1^s,$$

for all $x \in \mathbb{R}^d$.

The next theorem states our bounds on the asymptotic variance for this algorithm.

Algorithm 2 Unbiased-UBU with stochastic gradients (UBUBU-SG)

-
- 1: **Input:**
- Maximum stepsize h_0 .
 - Friction parameter $\gamma > 0$.
 - Individual potential terms $(U_i)_{0 \leq i \leq N_D}$.
 - Minimizer x^* of Potential $U(x)$ and its Hessian $H^* = \nabla^2 U(x^*)$.
 - Batch size parameter N_b (related to $\tau = \lceil N_D/N_b \rceil$).
 - Burn-in length parameters B_0 and B .
 - Number of samples parameter K .
 - Number of parallel chains parameters N , c_N and ϕ_N .
 - Richardson extrapolation parameter $c_R \in [0, 1)$ (default value $c_R = \frac{1}{2\sqrt{2}}$).
 - Test function f .
- 2: **Samples from Gaussian approximation at level 0:**
- 3: Sample NK i.i.d. samples $z_1^{(0)}, \dots, z_{NK}^{(0)}$ from μ_G .
- 4: Compute $S_0 := \frac{1}{NK} \sum_{i=1}^{NK} f(z_i^{(0)})$.
- 5: **Generate number of chains:**
- 6: Sample $(N_{l,l+1})_{l \geq 0}$ according to (3.4) and (3.30), let $l_{\max} = \max\{l : N_{l,l+1} > 0\}$.
- 7: **Averages of differences $D_{l,l+1}$ from $l = 0, \dots, l_{\max}$:**
- 8: **for** $l = 0, \dots, l_{\max}$ **do**
- 9: **for** $r = 1, \dots, N_{l,l+1}$ **do**
- 10: Sample $z_{-B_l}^{(l,l+1,r)}, \dots, z_K^{(l,l+1,r)}, z_{-B_{l+1}}^{(l,l+1,r)}, \dots, z_K^{(l,l+1,r)}$ from $\nu_{l,l+1}^{SG}$.
- 11: Compute $D_{l,l+1}^{(r)}$ based on (3.13) using $z_1^{(r,l,l+1)}, \dots, z_K^{(r,l,l+1)}, z_1^{(r,l,l+1)}, \dots, z_K^{(r,l,l+1)}$.
- 12: **end for**
- 13: Compute $S_{l,l+1}$ using (3.5).
- 14: **end for**
- 15: Compute $S(c_R)$ using (3.7).
- 16: **Output:**
- 17: Unbiased estimator $S(c_R)$,
- 18: Samples $z_1^{(0)}, \dots, z_{NK}^{(0)}$,
- 19: Samples $z_1^{(l,l+1,r)}, \dots, z_K^{(l,l+1,r)}, z_1^{(r,l,l+1)}, \dots, z_K^{(r,l,l+1)}$ for levels $0 \leq l \leq l_{\max}$, parallel chains $1 \leq r \leq N_{l,l+1}$.
-

THEOREM 3.24. *Considering UBUBU with stochastic gradients, suppose that Assumptions 3.12, 3.21, 3.22, 3.23 hold, and in addition $\gamma \geq \sqrt{8M}$,*

$$h_0 \leq \frac{C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tau/N_D, N_b)}{N_D^{3/2}}, \quad B \geq \frac{16 \log(2^{3/2})\tilde{\gamma}}{\tilde{m}h_0N_D^{1/2}}, \quad B_0 \geq \frac{16\tilde{\gamma}}{\tilde{m}h_0N_D^{1/2}} \log\left(\frac{1}{N_D^{9/4}h_0^{3/2}}\right).$$

Suppose that $c_R \in [0, 1)$ and $2 < \phi_N < 8$. Then for any $N \geq 1$, the UBUBU estimator $S(c_R)$ has finite expected computational cost, $\mathbb{E}S(c_R) = \pi(f)$, and it has finite variance. Moreover, it satisfies a CLT as $N \rightarrow \infty$, and the asymptotic variance σ_S^2 defined in (3.8) can be bounded as

$$\sigma_S^2 \leq \frac{1}{\tilde{m}N_DK} + C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s, \tau/N_D, N_b, \phi_N) \frac{d^2}{c_N N_D^2}.$$

PROOF. See Section F.2 in the Appendix. □

REMARK 3.25. *With the choice $c_N = \mathcal{O}\left(\frac{1}{N_D}\right)$ and $K = \mathcal{O}(1)$, we get a bound $\sigma_S^2 \leq \mathcal{O}\left(\frac{d^2}{\tilde{m}N_D}\right)$, which, except for the dimension dependence, is similar to the variance of a 1-Lipschitz function according to the target. Hence obtaining an effective sample only requires*

evaluating a full gradient only once per $\mathcal{O}(N_D)$ iteration, so there is no increase in computational cost as the dataset size N_D increases. The dimension dependency $\mathcal{O}(d^2)$ in our bound is likely not sharp as we have not observed any dimension dependency in our simulations.

REMARK 3.26. Although we have used Gaussian approximation at level 0 in Theorem 3.24 as this allows us to obtain better computational complexity in terms of N_D , one could also consider using UBU discretizations with SVRG gradients starting from level 0. This might be advantageous when the Gaussian approximation is not yet accurate. One could also consider different initial distributions. It is straightforward to adapt the proofs of Theorem 3.24 to show that even in such situations, under appropriate assumptions on the burn-in times, the UBUBU-SG method produces unbiased estimators with finite variance.

3.3. *UBUBU with approximate gradients.* Stochastic gradients are not the only possible approach for computing accurate approximations of the gradient. In case the potential is close to a Gaussian (which is typical in the big data regime due to the Bernstein-von-Mises theorem), the following approximation can be quite accurate.

DEFINITION 3.27. The quadratic approximate gradient of U at x with respect to \hat{x} is defined by

$$(3.31) \quad \mathcal{Q}(x|\hat{x}) = \nabla U(\hat{x}) + \nabla^2 U(x^*)(x - \hat{x}),$$

where x^* is the minimizer of U .

When using this approximation for the gradient, the \mathcal{B} step becomes

$$(3.32) \quad \mathcal{B}_{\mathcal{Q}}(x, v, h|\hat{x}) = (x, v - h\mathcal{Q}(x|\hat{x})).$$

The UBU iterations in this case become

$$(3.33) \quad \begin{aligned} & \left(\xi_{k+1}^{(i)} \right)_{i=1}^4 \sim \mathcal{N}(0_d, I_d) \text{ for all } i = 1, \dots, 4. \\ & (\bar{x}_k, \bar{v}_k) = \mathcal{U} \left(x_k, v_k, h/2, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)} \right), \\ & \hat{x}_k = \bar{x}_{\lfloor k/\tau \rfloor \tau} \\ & (x_{k+1}, v_{k+1}) = \mathcal{U} \left(\mathcal{B}_{\mathcal{Q}}(\bar{x}_k, \bar{v}_k, h|\hat{x}_k), h/2, \xi_{k+1}^{(3)}, \xi_{k+1}^{(4)} \right). \end{aligned}$$

Let P_h^A denote the time inhomogenous Markov kernel describing the evolution of (x_k, \hat{x}_k, v_k) according to the approximate gradient steps (3.33).

The reference point \hat{x} is updated after every τ iterations for some $\tau \geq 1$. We only need to evaluate the full gradient once per τ iterations, and use an approximation based on the Hessian at the minimizer otherwise. Since the Hessian $H^* = \nabla^2 U(x^*)$ only has to be computed once, this does not affect overall efficiency when the number of samples N is sufficiently high. For many potentials of interest, the approximation steps in (3.31) can be computed at a much smaller cost than the gradient of U . Moreover, when thinning is used (such at levels $l = 1$ and higher), multiple steps according to (3.33) can be combined into one using the fact that this is a linear system, further reducing the number of matrix-vector products required.

We follow a similar strategy as in the UBUBU-SG case (see Figure 3). We use Gaussian samples at level 0, and couplings involving both OHO and UBU discretizations.

Two chains evolving according to approximate gradients with step sizes h and $h/2$ can be coupled as follows.

$$\begin{aligned}
& \left(\xi_{k+1}^{(i)} \right)_{i=1}^8 \sim \mathcal{N}(0_d, I_d) \text{ for all } i = 1, \dots, 8. \\
& (\bar{x}_k, \bar{v}_k) = \mathcal{U}^2 \left(x_k, v_k, h/2, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}, \xi_{k+1}^{(3)}, \xi_{k+1}^{(4)} \right), \\
& \hat{x}_k = \bar{x}_{\lfloor k/\tau \rfloor \tau} \\
& (x_{k+1}, v_{k+1}) = \mathcal{U}^2 \left(\mathcal{B}_{\mathcal{Q}}(\bar{x}_k, \bar{v}_k, h | \hat{x}_k), h/2, \xi_{k+1}^{(5)}, \xi_{k+1}^{(6)}, \xi_{k+1}^{(7)}, \xi_{k+1}^{(8)} \right). \\
(3.34) \quad & (\bar{x}'_k, \bar{v}'_k) = \mathcal{U} \left(x'_k, v'_k, h/2, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)} \right), \\
& \hat{x}'_k = \bar{x}'_{\lfloor 2k/\tau \rfloor \tau/2} \\
& \left(x'_{k+1/2}, v'_{k+1/2} \right) = \mathcal{U} \left(\mathcal{B}_{\mathcal{Q}}(\bar{x}'_k, \bar{v}'_k, h/2, v | \hat{x}'_k), h/4, \xi_{k+1}^{(3)}, \xi_{k+1}^{(4)} \right), \\
& \left(\bar{x}'_{k+1/2}, \bar{v}'_{k+1/2} \right) = \mathcal{U} \left(x'_{k+1/2}, v'_{k+1/2}, h/2, \xi_{k+1}^{(5)}, \xi_{k+1}^{(6)} \right), \\
& \hat{x}'_{k+1/2} = \bar{x}'_{\lfloor (2k+1)/\tau \rfloor \tau/2} \\
& \left(x'_{k+1}, v'_{k+1} \right) = \mathcal{U} \left(\mathcal{B}_{\mathcal{Q}} \left(\bar{x}'_{k+1/2}, \bar{v}'_{k+1/2}, h/2 | \hat{x}'_{k+1/2} \right), h/4, \xi_{k+1}^{(7)}, \xi_{k+1}^{(8)} \right),
\end{aligned}$$

Let $P_{h,h/2}^A$ denote the time inhomogenous Markov kernel describing the evolution of $(x_k, \hat{x}_k, v_k, x'_k, \hat{x}'_k, v'_k)$ according to the coupled approximate gradient steps (3.34).

As with stochastic gradients, will also need to couple one chain with step size h running OHO on the Gaussian approximation μ_G , and another chain based on UBU with approximate gradients on the target with step size $h/2$.

$$\begin{aligned}
& \left(\xi_{k+1}^{(i)} \right)_{i=1}^8 \sim \mathcal{N}(0_d, I_d) \text{ for all } i = 1, \dots, 8. \\
& (\bar{x}_k, \bar{v}_k) = \mathcal{O}^2 \left(x_k, v_k, h/2, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}, \xi_{k+1}^{(3)}, \xi_{k+1}^{(4)} \right) \\
& (x_{k+1}, v_{k+1}) = \mathcal{O}^2 \left(\mathcal{H}_*(\bar{x}_k, \bar{v}_k, h), h/2, \xi_{k+1}^{(5)}, \xi_{k+1}^{(6)}, \xi_{k+1}^{(7)}, \xi_{k+1}^{(8)} \right), \\
& (\bar{x}'_k, \bar{v}'_k) = \mathcal{U} \left(x'_k, v'_k, h/4, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)} \right) \\
(3.35) \quad & \hat{x}'_k = \bar{x}'_{\lfloor 2k/\tau \rfloor \tau/2} \\
& \left(x'_{k+1/2}, v'_{k+1/2} \right) = \mathcal{U} \left(\mathcal{B}_{\mathcal{Q}}(\bar{x}'_k, \bar{v}'_k, h/2 | \hat{x}'_k), h/4, \xi_{k+1}^{(3)}, \xi_{k+1}^{(4)} \right) \\
& \left(\bar{x}'_{k+1/2}, \bar{v}'_{k+1/2} \right) = \mathcal{U} \left(x'_{k+1/2}, v'_{k+1/2}, h/4, \xi_{k+1}^{(5)}, \xi_{k+1}^{(6)} \right) \\
& \hat{x}'_{k+1/2} = \bar{x}'_{\lfloor (2k+1)/\tau \rfloor \tau/2} \\
& \left(x'_{k+1}, v'_{k+1} \right) = \mathcal{U} \left(\mathcal{B}_{\mathcal{Q}}(\bar{x}'_{k+1/2}, \bar{v}'_{k+1/2}, h/2 | \hat{x}'_{k+1/2}), h/4, \xi_{k+1}^{(7)}, \xi_{k+1}^{(8)} \right)
\end{aligned}$$

Let $P_{h,h/2}^{OHO/A}$ denote the time inhomogenous Markov kernel describing the evolution of $(x_k, v_k, x'_k, \hat{x}'_k, v'_k)$ according to the steps (3.35).

We define $\nu_{0,1}^A$ as joint distribution of $z_{-B_0}^{(0,1)}, \dots, z_K^{(0,1)}, z_{-B_1}^{(0,1)}, \dots, z_K^{(0,1)}$ that is similar to the $\nu_{0,1}^{SG}$ coupling for UBUBU-SG, but using inhomogenous Markov kernels $P_{h,h/2}^{OHO/A}$ instead of $P_{h,h/2}^{OHO/SVRG}$. Similarly, we let $\nu_{l,l+1}^A$ denote the joint distribution of $z_{-B_l}^{(l,l+1)}, \dots, z_K^{(l,l+1)}, z_{-B_{l+1}}^{(l,l+1)}, \dots, z_K^{(l,l+1)}$, defined analogously to $\nu_{l,l+1}^{SG}$ for UBUBU-SG, but using $P_{h,h/2}^{OHO/A}$ and $P_{h,h/2}^A$ in place of $P_{h,h/2}^{OHO/SVRG}$ and $P_{h,h/2}^{SVRG}$. We choose $c_{l,l+1}$ as

$$(3.36) \quad c_{l,l+1} = c_N \phi_N^{-l} \text{ for } l \in \mathbb{N}.$$

The UBUBU-Approx method follows similar steps as in Algorithm 2, but it uses the couplings ν_0^A and $\nu_{l,l+1}^A$ instead of ν_0^{SG} , and $\nu_{l,l+1}^{SG}$. In terms of input, unlike in Algorithm 2, we do not use individual potential terms $U_i(x)$ and batch size N_b , but require gradient calculation frequency τ . We recommend setting the Richardson extrapolation parameter $c_R = \frac{1}{2}$ in this case (as this approximate gradient scheme has strong order 1). A slightly weaker form of Assumption 3.21 will suffice here.

ASSUMPTION 3.28 (∇ Lipschitz property). *There is a $\tilde{M} > 0$ such that for all $x, y \in \mathbb{R}^d$,*

$$\|\nabla U(x) - \nabla U(y)\| \leq M \|x - y\| \quad \text{for } M = N_D \tilde{M}.$$

Our results for this scheme are stated in Theorem 3.29.

THEOREM 3.29. *Considering UBUBU-Approx method, suppose that Assumptions 3.12, 3.22, 3.23, 3.28 hold, and in addition $\gamma \geq \sqrt{8\tilde{M}}$,*

$$h_0 \leq \frac{C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tau/N_D)}{N_D^{3/2}}, \quad B \geq \frac{16 \log(2) \tilde{\gamma}}{\tilde{m} h_0 N_D^{1/2}}, \quad B_0 \geq \frac{16 \tilde{\gamma}}{\tilde{m} N_D^{1/2} h_0} \log \left(\frac{1}{N_D^3 h_0^2} \right).$$

Suppose that $c_R \in [0, 1)$ and $2 < \phi_N < 4$. Then for any $N \geq 1$, $S(c_R)$ has finite expected computational cost, $\mathbb{E}S(c_R) = \pi(f)$, and it has finite variance. Moreover, it satisfies a CLT as $N \rightarrow \infty$, and the asymptotic variance σ_S^2 defined in (3.8) can be bounded as

$$\sigma_S^2 \leq \frac{1}{\tilde{m} N_D K} + \frac{C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s, \tau/N_D, \phi_N) d^2}{c_N N_D^2}.$$

PROOF. See Section G.3 in the Appendix. □

REMARK 3.30. *To control the asymptotic variance of Theorem 3.24 and Theorem 3.29 for large d we would need to set $h_0 < \mathcal{O}(d^{-2})$; the dimension dependency in this bound might not be sharp, and we did not observe such limitations in our simulations. UBU iterations with AG and SVRG gradient approximations no longer form a time homogeneous Markov chain (unless the state space is extended), so it is challenging to establish $\mathcal{O}(1/K)$ scaling in the bound on σ_S^2 , like in Theorem 3.15. If we select $h_0 \sim \mathcal{O}(1/N_D^{3/2})$, then for large N_D , the total computational cost of the approximate and stochastic gradient methods scales like $\mathcal{O}(N)$ due to Proposition 3.4. This is a significant improvement over UBUBU with exact gradients, which has a computational cost of $\mathcal{O}(N_D N)$.*

REMARK 3.31. *Although we have used Gaussian approximation at level 0 in Theorem 3.29 to understand the complexity in terms of N_D , one could also consider using UBU discretizations with approximate gradients starting from level 0. We could also use different initial distributions. It is not difficult to adapt the proof to show that even in such situations, under appropriate assumptions on the burn-in times, the UBUBU-Approx method produces unbiased estimators with finite variance.*

Algorithm	Computational Cost
UBUBU (Exact gradients)	$\mathcal{O}(N_D N)$
UBUBU (stochastic gradients)	$\mathcal{O}(N)$
UBUBU (approximate gradients)	$\mathcal{O}(N)$

TABLE 2

Comparison of the computational cost of the various UBUBU methods in terms of N and N_D .

4. Numerical results. In this section, we provide numerical examples to demonstrate the effectiveness of our unbiased estimator UBUBU with exact, approximate and stochastic gradients. We test this on a range of problems, including a Gaussian example, a multinomial regression problem on the MNIST dataset, and a Poisson regression model for soccer scores; these computations serve to highlight the comparisons of our method with RHMC, which we view as the gold standard. We briefly describe the latter in Algorithm 3, stated in Section H of the Appendix.

For RHMC, we have used a partial refreshment parameter of $\alpha = 0.7$, which typically performed 50% – 70% better than doing full velocity refreshment ($\alpha = 0$). We choose parameters E_L (expected number of leapfrog steps) and h (stepsize) such that the acceptance rate is approximately 0.65 (as recommended in [7]), and that $E_L h \approx \frac{1}{\sqrt{m}}$ (m is the minimal eigenvalue of the Hessian at the mode), in line with the theoretical results for optimal convergence of the continuous time RHMC process [64]. We found that the effective sample sizes obtained in all of our experiments are in line with the continuous convergence rates of [64] scaled by the stepsize h , so we do not think that other parameter choices can significantly improve the performance of RHMC.

Our numerical experiments with unbiased estimators are specific to the UBU splitting method, as was the analysis. We also ran some preliminary numerical experiments with an unbiased version of BAOAB, but found that UBUBU was more efficient in all cases.

We estimated the ESS values based on at least 16 parallel runs in each simulation. For UBUBU, the number of parallel chains N was chosen in the range $N \in [64, 256]$, and we set $c_N = 1/16$, $\phi_N = 2\sqrt{2}$ in each case.

We will post the Matlab code of our simulations at <https://github.com/paulindani>.

4.1. *Gaussian target.* Here we consider a Gaussian target in d dimensions whose precision matrix has eigenvalues

$$1, 1 + \frac{\kappa - 1}{d - 1}, 1 + \frac{2(\kappa - 1)}{d - 1}, \dots, \kappa.$$

Theorem 4 of [57] has shown that for some Gaussian targets with condition number κ , the inverse spectral gap of HMC taking K leapfrog steps per iteration was shown to be at least $\mathcal{O}(K\kappa\sqrt{d}/\log(d))$. More recently, it has been shown that randomizing the integration time can substantially improve the performance of HMC [16]. In continuous time, sharp convergence results have been obtained for RHMC in [64]. Moreover, for Gaussians with condition number κ , RHMC can approximate the target distribution with $\mathcal{O}(\sqrt{\kappa}d^{1/4})$ queries under a warm-start assumption [2]. In our preliminary experiments, RHMC significantly outperformed HMC on high-dimensional problems, so we only consider RHMC here.

We provided RHMC with the advantage of being initialized from the Gaussian target distribution, while UBUBU was initialized in $\mu_0(x, v) = \delta_0(x)\mathcal{N}(v, I_d)$. Our numerical simulations are presented in Figures 4-6.

Figure 4 shows the maximum number gradient evaluations per effective sample (ESS) among all components $f(x) = x_i$ for $1 \leq i \leq d$ as a function of the dimension $d = 10, 10^2, \dots, 10^5$, for condition number $\kappa \in \{4, 100\}$. Figure 5 shows the number of gradient

evaluations per ESS for the norm test function $f(x) = \|x\|$ as a function of the dimension d . As we can see, in both scenarios, UBUBU does not show any dimension dependence, while the number of gradient evaluations per ESS scales as $\mathcal{O}(d^{1/4})$ for RHMC. In our experiments, UBUBU is 20-40 times more efficient than RHMC for $d = 100000$.

Figure 6 presents the histograms of the number of gradient evaluations per effective sample size (ESS) amongst test functions $f(x) = x_1, \dots, f(x) = x_d$, when comparing UBUBU with RHMC. This experiment is for a specific dimensions size of $d = 10^5$ and condition numbers $\kappa \in \{4, 100\}$. As we can observe, UBUBU outperforms RHMC in terms of gradient evaluations per ESS.




Fig 4: Dimensional dependence of gradients/ESS over all components for Gaussian targets.




Fig 5: Dimension dependence of gradients/ESS for test function $\|x\|$ for Gaussian targets.

An important question related to this example is the dimension dependence of the original unbiased kinetic Langevin scheme based on Euler–Maruyama discretization presented in [85]. Due to the different estimator proposed there, the number of samples $N_{l,l+1}$ is random for every l , and the variance of the term equivalent to $S_{l,l+1} = \frac{1}{N_{l,l+1}} \sum_{i=1}^{N_{l,l+1}} D_{l,l+1}^{(r)}$ will be proportional to $\mathbb{E}(D_{l,l+1}^2)$, not $\text{Var}(D_{l,l+1})$ like in our case. For functions like the norm $f(x, v) = \|x\|$, in general, using the strong order one property of the Euler–Maruyama scheme ([86]), $\mathbb{E}(D_{l,l+1}) = \mathcal{O}(\sqrt{dh_l})$ and $\mathbb{E}(D_{l,l+1}^2) = \mathcal{O}(dh_l^2)$. So the asymptotic variance of the final estimator is $\mathcal{O}(1 + dh_0^2)$, and by choosing $h_0 = \mathcal{O}(d^{-1/2})$, we expect that this will require $\mathcal{O}(d^{1/2})$ gradient evaluations per effective sample.




Fig 6: Gradient/ESS over all components for the Gaussian target example.

4.2. *Bayesian multinomial regression.* Our second numerical example is to consider a Bayesian multinomial regression (BMR) problem. BMR is a generalized linear regression model which estimates probabilities for r different categories of dependent variable y using a set of explanatory variables x . Here, provided m classes, we let $q = (q^1, \dots, q^m) \in \mathbb{R}^d$ with $d = md_o$ and $q^i \in \mathbb{R}^{d_o}$. The likelihood associated with the problem is given as

$$(4.1) \quad p(y^j | q) = \frac{\exp(\langle x^j, q^{y^j} \rangle)}{\sum_{1 \leq k \leq m} \exp(\langle x^j, q^k \rangle)}.$$

Our focus is on estimating a posterior distribution, where the posterior potential is given as

$$(4.2) \quad U(q) = -\log(p_0(q)) - \sum_{k=1}^{N_D} \log(p(y^j | q)).$$

Here we chose p_0 as a Gaussian prior $p_0(q) = \frac{\exp(-\|q\|^2 / (2\sigma_0^2))}{(\pi\sigma_0^2)^{d/2}}$. In Lemma H.6 in Appendix H, we show that the gradient-Lipschitz and strongly Hessian Lipschitz conditions (Assumptions 3.6 and 3.10) hold for this example.

We are interested in applying our BMR model to the MNIST dataset [56] about classifying handwritten digits from 0 to 9, which are shown as examples in Figure 7. The dataset contains 60,000 training data points and 10,000 test data points where the images are of size 28 by 28 pixels. The covariate vectors x^j are obtained by flattening the images into vectors taking values on the interval $[0, 1]$, and adding a 1 in the end for the intercept term. Hence $d_0 = 28^2 + 1 = 785$, $m = 10$, and $d = d_0 m = 7850$. We set the prior variance $\sigma_0^2 = 0.1$ (this was tested to provide good prediction performance).

For our numerical simulations, we will present two different scenarios: one without preconditioning (Figure 9) and one with preconditioning (Figure 10). In both figures, we evaluated the efficiency of the methods in terms of gradient evaluations per ESS for the coordinate test functions $f(x) = x_1, \dots, f(x) = x_d$.




Fig 7: MNIST datasets containing images of handwritten digits from 0 to 9.

To compare the posterior distribution with a Gaussian approximation, we have selected a component with a relatively large third derivative. Figure 8 illustrates the potential function and the Gaussian approximation with precision $\nabla^2 U(x^*)$ along the line $x^* + te_i$. Here $e_i = (0, \dots, 0, 1, 0, \dots, 0)$ is the unit vector of the chosen component ($i = 7491$ in our implementation), and t is chosen to cover up to 3 times the standard deviation difference from x_i^* . As we can see, the distribution of this component has a significant skewness, and the density values can differ by up to 40% even in the bulk of the distribution.




Fig 8: MNIST example. Left: Comparison between potential and quadratic approximation. Right: Difference between the potential and quadratic approximation.

In the first scenario (no preconditioning), the condition number of the Hessian at the mode $\nabla^2 U(x^*)$ is $\kappa \approx 7.2 \times 10^3$. We included simulation results with RHMC, UBUBU, and UBUBU-SG. For UBUBU-SG, we used a 10% batch ($N_b = 6000$, $N_D = 60000$), and set the maximum level with control variate stochastic gradient approximation as $s_{\max} = 2$. As we can see, UBUBU improves upon RHMC, and this is further improved by UBUBU-SG.

By preconditioning, we mean that we obtain samples from a transformed potential $U(Ax)$ for some matrix A , which may have a better condition number than the original potential. It is easy to see that if X follows a distribution with density proportional to $\exp(-U(x))$, then $X' = A^{-1}X$ has a density proportional to $\exp(-U(Ax))$.

In the case of RHMC, the best performance was obtained by preconditioning using the matrix square root of the Hessian at the mode, $A = (\nabla^2 U(x^*))^{-1/2}$. For UBUBU, this same approach worked reasonably well, but the best performance was obtained by only preconditioning in the eigenvectors corresponding to the largest 1000 eigenvalues of $\nabla^2 U(x^*)$ (i.e. shrinking them to the same size as the 1000th largest eigenvalue), and keeping the other directions unchanged. This resulted in a condition number of $\kappa \approx 4.5$ for the Hessian of the transformed potential at its mode.

We also included the implementation of the approximate gradient version UBUBU-Approx with the same preconditioning as for UBUBU and set the frequency of full gradient evaluations as $\tau = 15$. This has drastically reduced the number of gradient evaluations without hurting performance, and it shows approximately 100 times improvement over RHMC.




Fig 9: Gradient/ESS over all components for MNIST dataset without preconditioning.




Fig 10: Gradient/ESS over all components for MNIST dataset with preconditioning.

In addition to the coordinate test functions, we have also evaluated the efficiency of these methods for the posterior predictive probability of digits 0, 1 . . . , 9 on the test dataset (10000 images, 100000 test functions in total). Figure 11 presents experiments comparing RHMC and UBUBU-Approx on these test functions. The experiments show approximately 60 times improvement in efficiency for UBUBU-Approx compared to RHMC, which is in line with our theory proving that UBUBU does not exhibit dimension dependency (Proposition 3.18).




Fig 11: Gradient/ESS for probabilities of all 10 digits over 10000 test images for MNIST dataset with preconditioning.

4.3. *Poisson regression model.* Our final example is a Poisson regression model for predicting soccer scores taken from [55].

Let $g = 1, \dots, G$ be the index of games. Let S_g^H denote the number of goals scored by the home team at game g , and let S_g^A denote the number of goals scored by the away team. The independent Poisson model [65] assumes that these scores are distributed as

$$S_g^H \sim \text{Poisson}(\lambda_g^H), \quad S_g^A \sim \text{Poisson}(\lambda_g^A),$$

conditionally independently given the rates λ_g^H and λ_g^A .

In our implementation, the rates are connected to the linear predictors η_g^H and η_g^A using the function $\text{softplus}(x) = \log(1 + \exp(x))$ (see Figure 12), i.e.

$$(4.3) \quad \lambda_g^A = \text{softplus}(\eta_g^A), \quad \lambda_g^H = \text{softplus}(\eta_g^H).$$

This function is Lipschitz and also gradient Lipschitz, which is desirable given our theory. Although this is less frequently used in the literature than the log link function, it was shown to be more robust and less sensitive to outliers [98, 100]. The linear predictors are modelled based on a random effect model with time-dependent attacking and defending strengths for each team. Let $w(g)$ denote the week of game g , then we set

$$(4.4) \quad \eta_g^H = a_{\text{home.team}(g),w(g)} + d_{\text{away.team}(g),w(g)}, \quad \eta_g^A = a_{\text{away.team}(g),w(g)} + d_{\text{home.team}(g),w(g)}$$

Let \mathbf{a} be all attacking strengths of all teams over the whole period, and \mathbf{d} denote all defending strengths. Then the log-likelihood is of the form

$$\log(p(\mathbf{a}, \mathbf{d})) = C(S_1^H, \dots, S_G^H, S_1^A, \dots, S_G^A) + \sum_{g=1}^G (-\lambda_g^H + S_g^H \log(\lambda_g^H) - \lambda_g^A + S_g^A \log(\lambda_g^A)),$$

which can be written as a function of \mathbf{a} and \mathbf{d} using (4.3) and (4.4).

We used a Gaussian random walk prior for the attacking/defending strengths $a_{\text{team},w}$ and $d_{\text{team},w}$, together with a weak Gaussian prior on every attacking and defending strength. Let




Fig 12: Softplus and ReLU activation functions.

\mathcal{T} denote the set of teams during the whole period considered (teams change from season to season due to relegation/promotion), then the overall log prior is of the form

$$\begin{aligned} \log p_0(\mathbf{a}, \mathbf{d}) = & C(\sigma, \sigma_0) - \sum_{\text{team} \in \mathcal{T}} \left(\sum_{w=w(1)}^{w(G)} \frac{a_{\text{team},w}^2}{2\sigma_0^2} + \sum_{w=w(1)}^{w(G)-1} \frac{(a_{\text{team},w+1} - a_{\text{team},w})^2}{2\sigma^2} \right) \\ & - \sum_{\text{team} \in \mathcal{T}} \left(\sum_{w=w(1)}^{w(G)} \frac{d_{\text{team},w}^2}{2\sigma_0^2} + \sum_{w=w(1)}^{w(G)-1} \frac{(d_{\text{team},w+1} - d_{\text{team},w})^2}{2\sigma^2} \right), \end{aligned}$$

We set $\sigma^2 = 0.01$ (this means a strong correlation for about two years), and $\sigma_0^2 = 10$ (weakly informative prior).

We considered 20 years of Premier League data (7600 games) from 19/08/2000 until 26/07/2020. Our model has $d = 89526$ parameters, and the condition number of the Hessian at the mode is $\kappa \approx 4 \cdot 10^3$.

We have implemented RHMC, UBUBU and UBUBU-Approx with $\tau = 20$ for this model. In the UBUBU-Approx algorithm, the target at level 0 was chosen as the Gaussian approximation (with mean x^* , and precision matrix $\nabla^2 U(x^*)$), meaning that gradient evaluations were only used from level 1 onwards. The test functions were chosen as $f(x) = x_1, \dots, f(x) = x_d$. Our numerical simulations are presented in Figure 13. As we can see, UBUBU uses approximately 30 times fewer gradient evaluations per effective sample than RHMC, and UBUBU-Approx uses 2000 times fewer gradient evaluations than RHMC.

5. Conclusion. In this article, we presented a new unbiased estimator which can exploit high strong-order numerical integrators for underdamped Langevin dynamics. We refer to our estimator as UBUBU which does not rely on the Metropolis acceptance/reject step. Our estimator is influenced by the work of [85], and instead is constructed using a telescoping sum for different discretization levels [44, 80]. We were able to show various theoretical insights, which include showing unbiasedness and finite variance, a central limit theorem, and asymptotic and non-asymptotic bounds on the variance for three algorithms, based on exact, stochastic, and approximate gradients. We have studied the behaviour of our algorithm for product target distributions and shown that for a large class of test functions, it has dimension-independent computational complexity. For stochastic gradients, we also considered the dependency on the size of the data in the big data limit and shown that our method is very efficient in such situations. The proof of these results relies on Wasserstein contraction results for the UBU dynamics. We provided numerical experiments verifying our theory and demonstrating the performance gains over other well-known methods such as randomized HMC. We have considered a range of model problems including an MNIST multinomial




Fig 13: Gradient/ESS over all components of a Poisson regression model for soccer scores.

regression problem, and a Poisson regression model tested on a real-world dataset. Our comparisons are based on gradient evaluations per effective sample size.

In terms of future work, there are various directions which could be taken up. One of them is related to exploiting higher-order schemes, which were provided in [40, 41]. Numerically, strong orders of up to 4 have been observed. [40] have proven strong order $3/2$, $5/2$ and 3 under gradient Lipschitz, Hessian Lipschitz and third-order Lipschitz assumptions, respectively. However, the dimensional dependence obtained under each of these assumptions has not been shown to improve on the UBU scheme in [86]. Furthermore, such splitting schemes typically require more than one gradient evaluation per step, unlike our strategy. In a different direction, one could consider integrators adapted to potentials that do not have the gradient-Lipschitz property (such as in the case of sparsity-inducing priors [76] or log link functions). Other potential directions are nested expectations [97] and static parameter estimation [3, 31]. Finally, one could consider the setting where one does not assume convexity [22, 38, 66, 88].

Funding. The authors acknowledge the support of the Engineering and Physical Sciences Research Council Grant EP/S023291/1 (MAC-MIGS Centre for Doctoral Training). NKC is supported by an EPSRC-UKRI AI for Net Zero Grant: “Enabling CO2 Capture And Storage Projects Using AI”, (Grant EP/Y006143/1).

APPENDIX A: DISCUSSION AND OUTLINE OF RESULTS

The beginning of this appendix is devoted to providing a road-map for our results. In Appendix B, which follows, we provide variance estimates of the full gradient multilevel UBUBU method. The approach we use is to bound $\text{Var}(D_0)$ using Theorem 2 of [51] and to use the strong error estimates of [86] for UBU to estimate $\text{Var}(D_{l,l+1})$. [51] requires Ricci curvature of the UBU Markov chain and extending [86] to global strong error estimates in Appendix D.1 requires Wasserstein convergence. We provide this in Appendix C in the full gradient setting using the methods of [63]. We provide L^4 Lyapunov drift inequalities

in the full gradient setting. We can then bound the average distance to the minimizer non-asymptotically, the key result needed to get complexity bounds in the big data setting. We also provide the proof of the central limit theorem of the estimator in Appendix B.

In Appendix E we describe the initialization and the OHO scheme for the approximate and stochastic gradient methods and some estimates of the distance between the initial measure and the target measure. We then use the techniques of [101] to provide global strong error estimates of the SVRG method. We combine and extend the techniques of [86] and [101] to prove new non-asymptotic stochastic gradient error bounds for the UBU integrator. From this we obtain in Appendix D variance bounds and estimates on our estimator UBUBU with exact gradients. This is then extended to providing estimates of the variance of our multilevel estimator in the SVRG stochastic gradient setting in Appendix F.

We further develop bounds for our new approximate gradient UBU method in Appendix G using the same techniques, in the approximate gradient setting. In general, Appendices F, G follow similarly where one requires bounds on the variance of the quantity D_0 and $D_{l,l+1}$. However, we use an interpolation argument to improve the results in Appendix G as opposed to the methods used in Appendix F. We also use some classical results from the theory of ODEs to establish bounds between continuous diffusions to establish the variance of $D_{0,1}$ in Appendix F and G. Finally, we provide some auxiliary results in Appendix H.

APPENDIX B: UNBIASED MULTILEVEL ESTIMATORS

PROPOSITION 3.4. *Suppose that Assumptions 3.1, 3.2 and 3.3 hold, and that $2 < \phi_N < \phi_D$. Then S as defined in (3.6) is an unbiased estimator of $\mu(f)$ that has finite variance*

$$\text{Var}(S) \leq \frac{\text{Var}(D_0)}{N} + \frac{V_D}{N \underline{c}_N \left(1 - \frac{\phi_D}{\phi_N}\right)},$$

and finite expected computational cost.

Similarly, for any $c_R \in [0, 1)$, $S(c_R)$ as defined in (3.7) is also an unbiased estimator of $\mu(f)$ with finite variance

$$\text{Var}(S(c_R)) \leq \frac{\text{Var}(D_0)}{N} + \frac{V_D}{N \underline{c}_N \left(1 - \frac{\phi_N}{\phi_D}\right)} + \frac{V_D}{N \underline{c}_N \left(1 - \frac{\phi_N}{\phi_D}\right)} \frac{2}{(1 - c_R)^2} \left(\frac{\phi_N}{\phi_D}\right)^{\log(2 \underline{c}_N N / \phi_N) / \log(\phi_N)},$$

and finite expected computational cost.

PROOF OF PROPOSITION 3.4. From Assumption 3.3, and the definition of S , it follows that the expected computational cost of S is upper bounded as follows:

$$\begin{aligned} & \mathcal{O} \left(N + \sum_{l=0}^{\infty} \mathbb{E}(N_{l,l+1}) 2^l (K + lB + B_0) \right) \\ & \leq \mathcal{O} \left(N \left(1 + \bar{c}_N \sum_{l=0}^{\infty} \left(\frac{2}{\phi_N}\right)^l (K + lB + B_0) \right) \right) < \infty. \end{aligned}$$

From Assumption 3.1, and the independence of the terms, we have that

$$\begin{aligned} \text{Var}(S) & \leq \frac{\text{Var}(D_0)}{N} + \sum_{l=0}^{\infty} \frac{\mathbb{E}(D_{l,l+1}^2)}{\mathbb{E}N_{l,l+1}} \leq \frac{\text{Var}(D_0)}{N} + \frac{V_D}{\underline{c}_N N} \sum_{l=0}^{\infty} \left(\frac{\phi_D}{\phi_N}\right)^{-l} \\ & = \frac{\text{Var}(D_0)}{N} + \frac{V_D}{N \underline{c}_N \left(1 - \frac{\phi_D}{\phi_N}\right)} < \infty. \end{aligned}$$

By Jensen's inequality, and Assumption 3.1, $\mathbb{E}(|S_0| + \sum_{l=0}^{\infty} |S_{l,l+1}|) < \infty$, hence by the dominated convergence theorem,

$$\mathbb{E}(S) = \mathbb{E}(S_0) + \sum_{l=0}^{\infty} \mathbb{E}(S_{l,l+1}) = \tilde{\mu}_{h_0}(f) + \sum_{l=0}^{\infty} \tilde{\mu}_{h_{l+1}}(f) - \tilde{\mu}_{h_l}(f) = \mu(f),$$

which concludes the proof for S .

For $S(c_R)$, the computational cost is the same as for S , so it has finite expectation. For the variance, we have

$$\text{Var}(S(c_R)) \leq \frac{\text{Var}(D_0)}{N} + \sum_{l=0}^{L(N)-1} \frac{\mathbb{E}(D_{l,l+1}^2)}{\mathbb{E}N_{l,l+1}} + \text{Var} \left(\frac{S_{L(N),L(N)+1}}{1-c_R} + \sum_{l=L(N)+1}^{\infty} \bar{S}_{l,l+1} \right).$$

The last term can be bounded as

$$\begin{aligned} & \text{Var} \left(\frac{S_{L(N),L(N)+1}}{1-c_R} + \sum_{l=L(N)+1}^{\infty} \bar{S}_{l,l+1} \right) \\ &= \text{Var} \left(S_{L(N),L(N)+1} \left(\frac{1}{1-c_R} - \sum_{l=L(N)+1}^{\infty} \frac{\mathbb{1}[N_{l,l+1}=1]}{\mathbb{E}(N_{l,l+1})} c_R^{l-L(N)} \right) + \sum_{l=L(N)+1}^{\infty} S_{l,l+1} \right) \\ &\leq 2 \cdot \text{Var} \left(S_{L(N),L(N)+1} \left(\frac{1}{1-c_R} - \sum_{l=L(N)+1}^{\infty} \frac{\mathbb{1}[N_{l,l+1}=1]}{\mathbb{E}(N_{l,l+1})} c_R^{l-L(N)} \right) \right) \\ &+ 2 \cdot \text{Var} \left(\sum_{l=L(N)+1}^{\infty} S_{l,l+1} \right) \leq \frac{2}{(1-c_R)^2} \mathbb{E}(S_{L(N),L(N)+1}^2) + 2 \sum_{l=L(N)+1}^{\infty} \mathbb{E}(S_{l,l+1}^2). \end{aligned}$$

As before, we have $\mathbb{E}(S_{l,l+1}^2) \leq \frac{\mathbb{E}(D_{l,l+1}^2)}{\mathbb{E}N_{l,l+1}} \leq \frac{V_D}{c_N N} \cdot \left(\frac{\phi_D}{\phi_N} \right)^{-l}$ for any $l \geq 0$. Using the fact that $\phi_N^{-L(N)-1} \leq \frac{1}{2c_N N}$, we have $\phi_N^{L(N)} \geq \frac{2c_N N}{\phi_N}$, hence $L(N) \geq \frac{\log(2c_N N/\phi_N)}{\log(\phi_N)}$. After some rearrangement, we obtain that

$$\begin{aligned} & \text{Var}(S(c_R)) \leq \\ & \frac{\text{Var}(D_0)}{N} + \frac{V_D}{N c_N \left(1 - \frac{\phi_N}{\phi_D}\right)} + \frac{V_D}{N c_N \left(1 - \frac{\phi_N}{\phi_D}\right)} \frac{2}{(1-c_R)^2} \left(\frac{\phi_N}{\phi_D} \right)^{\log(2c_N N/\phi_N)/\log(\phi_N)}, \end{aligned}$$

where the factor $\left(\frac{\phi_N}{\phi_D} \right)^{\log(2c_N N/\phi_N)/\log(\phi_N)}$ tends to 0 as $N \rightarrow \infty$. Finally, unbiasedness can be shown as before using the dominated convergence theorem. \square

We show below that a central limit theorem holds for these estimators.

THEOREM 3.5. *Under the assumptions of Proposition 3.4, we have that, as $N \rightarrow \infty$,*

$$\sqrt{N}(S - \mu(f)) \Rightarrow \mathcal{N}(0, \sigma_S^2) \quad \text{and} \quad \sqrt{N}(S(c_R) - \mu(f)) \Rightarrow \mathcal{N}(0, \sigma_S^2),$$

where

$$(3.8) \quad \sigma_S^2 := \text{Var}(D_0) + \sum_{l=0}^{\infty} \frac{\text{Var}(D_{l,l+1})}{c_{l,l+1}}.$$

PROOF OF THEOREM 3.5. First, we prove the result for $H := \sqrt{N}(S - \mu(f))$. For $l_{\max} \geq 0$, let

$$\begin{aligned} H^{l_{\max}} &:= \sqrt{N}(S_0 - \mathbb{E}(S_0)) + \sum_{l=0}^{l_{\max}} (S_{l,l+1} - \mathbb{E}(S_{l,l+1})) \\ &= \frac{1}{\sqrt{N}} \sum_{r=1}^N (D_0^{(r)} - \mathbb{E}(D_0^{(r)})) + \sum_{l=0}^{l_{\max}} \frac{\sqrt{N}}{\mathbb{E}(N_{l,l+1})} \sum_{r=1}^{N_{l,l+1}} D_{l,l+1}^{(r)} - \mathbb{E}(D_{l,l+1}) \\ &:= H_0 + \sum_{l=0}^{l_{\max}} H_{l,l+1}. \end{aligned}$$

Then by using independence, and the fact that $\left(\frac{\sqrt{N}}{\mathbb{E}(N_{l,l+1})}\right) / \left(\frac{1}{\sqrt{N}}\right) \rightarrow \frac{1}{c_{l,l+1}}$, by the proof of the central limit theorem (see Sections 3.3-3.4 of [37]), for every $t \in \mathbb{R}$, H_0 and $H_{l,l+1}$ satisfies

$$\begin{aligned} \mathbb{E}(e^{itH_0}) &\rightarrow e^{-t^2\mathcal{V}_0/2} \text{ as } N \rightarrow \infty \text{ for } \mathcal{V}_0 = \text{Var}(D_0), \\ \mathbb{E}(e^{itH_l}) &\rightarrow e^{-t^2\mathcal{V}_{l,l+1}/2} \text{ as } N \rightarrow \infty \text{ for } \mathcal{V}_{l,l+1} = \frac{\text{Var}(D_{l,l+1})}{c_{l,l+1}}. \end{aligned}$$

Using independence, we can multiply these together to obtain that for any $t \in \mathbb{R}$,

$$\mathbb{E}\left(e^{itH^{l_{\max}}}\right) \rightarrow e^{-t^2(\mathcal{V}_0 + \sum_{l=0}^{l_{\max}} \mathcal{V}_{l,l+1})/2} \text{ as } N \rightarrow \infty.$$

By Lemma 3.3.19 of [37], it follows that for a random variable X with $\mathbb{E}(X) = 0$ and $\mathbb{E}(X^2) < \infty$, we have

$$|\mathbb{E}(e^{itX}) - 1| \leq \frac{t^2\mathbb{E}(X^2)}{2}.$$

For $X = \sqrt{N}(S - \mu(f)) - H^{l_{\max}}$, we have

$$\mathbb{E}(X^2) = \text{Var}(X) \leq N \sum_{l=l_{\max}+1}^{\infty} \frac{\mathbb{E}(D_{l,l+1}^2)}{\mathbb{E}N_{l,l+1}} \leq \frac{V_D}{c_N} \sum_{l=l_{\max}+1}^{\infty} \left(\frac{\phi_D}{\phi_N}\right)^{-l} \leq \frac{V_D}{c_N} \frac{\left(\frac{\phi_N}{\phi_D}\right)^{l_{\max}}}{1 - \frac{\phi_N}{\phi_D}}.$$

Using independence of $H^{l_{\max}}$ and $H - H^{l_{\max}}$, $\mathbb{E}(e^{itH}) = \mathbb{E}(e^{itH^{l_{\max}}}) \cdot \mathbb{E}(e^{it(H - H^{l_{\max}})})$, so

$$\begin{aligned} &\limsup_{N \rightarrow \infty} \left| \mathbb{E}(e^{itH}) - e^{-t^2(\mathcal{V}_0 + \sum_{l=0}^{l_{\max}} \mathcal{V}_{l,l+1})/2} \right| \\ &\leq e^{-t^2(\mathcal{V}_0 + \sum_{l=0}^{l_{\max}} \mathcal{V}_{l,l+1})/2} \cdot \frac{V_D}{c_N} \sum_{l=l_{\max}+1}^{\infty} \left(\frac{\phi_D}{\phi_N}\right)^{-l} \leq \frac{V_D}{c_N} \frac{\left(\frac{\phi_N}{\phi_D}\right)^{l_{\max}}}{1 - \frac{\phi_N}{\phi_D}}. \end{aligned}$$

By letting $l_{\max} \rightarrow \infty$, it follows that $\limsup_{N \rightarrow \infty} \mathbb{E}(e^{itH}) = e^{-t^2\sigma_S^2}$, hence the convergence follows by the Lévy-Cramér continuity theorem (see Theorem 3.3.17 of [37]).

The proof for $S(c_R)$ follows the same lines, except that the variances of the terms for $l \geq L(N)$ need to be controlled separately using the same bounds as in the proof of Proposition 3.4, we omit the details. \square

APPENDIX C: CONVERGENCE RESULTS

The first set of results we prove are provided below for the convergence of the UBU scheme. Proving contraction of a coupling has been a popular method for establishing convergence rates both in the continuous time setting and for the discretization for Langevin dynamics (underdamped/kinetic) and Hamiltonian Monte Carlo (see for example [10–12, 29, 30, 32, 36, 39, 45, 70, 71, 86, 87] and many more).

Our approach to obtain convergence rates is based on proving contraction for a synchronous coupling. We need an appropriate metric to attain convergence, and contraction of the UBU scheme. We introduce the Wasserstein distance in this metric.

DEFINITION C.1 (Weighted Euclidean norm). For $z = (x, v) \in \mathbb{R}^{2d}$ we introduce the weighted Euclidean norm

$$\|z\|_{a,b}^2 = \|x\|^2 + 2b \langle x, v \rangle + a \|v\|^2,$$

for $a, b > 0$ with $b^2 < a$.

REMARK C.2. Using the assumption $b^2 < a$, we can show that this is equivalent to the Euclidean norm on \mathbb{R}^{2d} . Under the condition $b^2 \leq a/2$, we have

$$(C.1) \quad \frac{1}{2} \min(a, 1) \|z\|^2 \leq \frac{1}{2} \|z\|_{a,0}^2 \leq \|z\|_{a,b}^2 \leq \frac{3}{2} \|z\|_{a,0}^2 \leq \frac{3}{2} \max(a, 1) \|z\|^2.$$

DEFINITION C.3 (p -Wasserstein distance). Let us define $\mathcal{P}_p(\mathbb{R}^{2d})$ to be the set of probability measures which have p -th moment for $p \in [1, \infty)$ (i.e. $\mathbb{E}(\|Z\|^p) < \infty$). Then the p -Wasserstein distance in norm $\|\cdot\|_{a,b}$ between two measures $\mu, \nu \in \mathcal{P}_p(\mathbb{R}^{2d})$ is defined as

$$(C.2) \quad \mathcal{W}_{p,a,b}(\nu, \mu) = \left(\inf_{\xi \in \Gamma(\nu, \mu)} \int_{\mathbb{R}^{2d}} \|z_1 - z_2\|_{a,b}^p d\xi(z_1, z_2) \right)^{1/p},$$

where $\|\cdot\|_{a,b}$ is the norm introduced before and that $\Gamma(\nu, \mu)$ is the set of measures with respective marginals of ν and μ .

Before we proceed, we need to introduce the concept of Wasserstein convergence, which most of the results rely upon.

LEMMA C.4 (Wasserstein convergence). Let $1 \leq p \leq \infty$, $\mu, \nu \in \mathcal{P}_p(\mathbb{R}^{2d})$, and $a, b > 0$ with $b^2 < a$. Let us assume that $(z_k)_{k \geq 0} = (x_k, v_k)_{k \geq 0}$ and $(\tilde{z}_k)_{k \geq 0} = (\tilde{x}_k, \tilde{v}_k)_{k \geq 0}$ are two Markov chains with state space Λ and kernel P_h defined on the same probability space (a coupling) such that $z_0 \sim \nu$, $\tilde{z}_0 \sim \mu$, and $\mathbb{E}(\|z_0 - \tilde{z}_0\|^p) = [\mathcal{W}_{p,a,b}(\nu, \mu)]^p$. If the following contractive property holds,

$$(C.3) \quad \left[\mathbb{E}(\|\tilde{z}_{k+1} - z_{k+1}\|_{a,b}^p | z_{0:k}, \tilde{z}_{0:k}) \right]^{1/p} \leq (1 - c(h)) \|\tilde{z}_k - z_k\|_{a,b} \quad \text{for every } k \geq 0,$$

then we have

$$\mathcal{W}_{p,a,b}(\nu P_h^n, \mu P_h^n) \leq (1 - c(h))^n \mathcal{W}_{p,a,b}(\nu, \mu) \quad \text{for every } n \geq 0.$$

REMARK C.5. The existence of an optimal coupling satisfying that $\mathbb{E}(\|z_0 - \tilde{z}_0\|_{a,b}^p) = [\mathcal{W}_{p,a,b}(\nu, \mu)]^p$ follows by Theorem 4.1 of [94].

PROOF. By induction, we have $\mathbb{E}(\|\tilde{z}_n - z_n\|_{a,b}^p | z_0, \tilde{z}_0) \leq (1 - c(h))^n \|z_0 - \tilde{z}_0\|_{a,b}^p$, and the result follows by taking expectations and using Definition C.2. \square

Now, we present our first proposition, a convergence result of the UBU scheme with full gradients.

PROPOSITION C.6. *Suppose that U is m -strongly convex and M - ∇ Lipschitz. Let*

$$(C.4) \quad a = \frac{1}{M}, \quad b = \frac{1}{\gamma}, \quad c_2(h) = \frac{mh}{4\gamma}, \quad c(h) = \frac{mh}{8\gamma}.$$

Let P_h denote the transition kernel for a step of UBU with stepsize h . For all $\gamma \geq \sqrt{8M}$, $h < \frac{1}{2\gamma}$, $1 \leq p \leq \infty$, $\mu, \nu \in \mathcal{P}_p(\mathbb{R}^{2d})$, (C.2) holds. Hence for all $n \in \mathbb{N}$,

$$\mathcal{W}_{p,a,b}(\nu P_h^n, \mu P_h^n) \leq (1 - c_2(h))^{n/2} \mathcal{W}_{p,a,b}(\nu, \mu) \leq (1 - c(h))^n \mathcal{W}_{p,a,b}(\nu, \mu).$$

Further to this, P_h has a unique invariant measure π_h satisfying that $\pi_h \in \mathcal{P}_p(\mathbb{R}^{2d})$ for all $1 \leq p \leq \infty$.

REMARK C.7. *We are going to use the same choices of a and b as stated in (C.4) everywhere in the paper.*

COROLLARY C.8. *Suppose that U is an m -strongly convex M - ∇ Lipschitz potential, $\gamma \geq \sqrt{8M}$, $1 \leq p \leq 2$, $\mu, \nu \in \mathcal{P}_p(\mathbb{R}^{2d})$. Suppose that $(X_0, V_0) \sim \mu$, then the solution of (1.1) exists in the strong sense for any $t \geq 0$, and the corresponding Markov kernel P_t^{cont} satisfies*

$$(C.5) \quad \mathcal{W}_{p,a,b}(\nu P_t^{\text{cont}}, \mu P_t^{\text{cont}}) \leq \exp\left(-\frac{mt}{8\gamma}\right) \mathcal{W}_{p,a,b}(\nu, \mu) \quad \text{for} \quad a = \frac{1}{M}, b = \frac{1}{\gamma}.$$

REMARK C.9. *One can improve the restriction on γ slightly by writing the potential as a perturbation of a quadratic as in [87]. Due to the restrictions on the stepsize h and the friction parameter γ in Proposition C.6, $c(h) = \mathcal{O}\left(\frac{m}{M}\right)$ for all allowed parameter choices. In general, for ∇ Lipschitz, strongly-convex potentials, it may be impossible to prove contraction using such a quadratic form argument and synchronous coupling for $\gamma \leq \mathcal{O}(\sqrt{M})$ as explained in [71]. In the continuous time dynamics, $\gamma = \mathcal{O}(\sqrt{m})$ seems to yield the fastest convergence rate, as explained in [20]. In Example C.10 in the Appendix, we show that for Gaussian targets, UBU has an accelerated convergence rate $c(h) = \mathcal{O}(\sqrt{\frac{m}{M}})$ with the choice $\gamma = \mathcal{O}(\sqrt{m})$ and $h = \mathcal{O}(1/\sqrt{M})$.*

PROOF OF PROPOSITION C.6. We follow the approach of [70][Corollary 20]. It is sufficient to prove contraction of a synchronous coupling of Markov chains in an appropriate norm, we will use the $\|\cdot\|_{a,b}$ norm of Definition C.1 with $a = \frac{1}{M}$, $b = \frac{1}{\gamma}$. Based on the assumptions, we have $b^2 < a/2$. Hence, (C.1) holds.

We aim to show that contraction occurs in this norm for two Markov chains simulated by the same discretization $z_n = (x_n, v_n) \in \mathbb{R}^{2d}$ and $\tilde{z}_n = (\tilde{x}_n, \tilde{v}_n) \in \mathbb{R}^{2d}$ that are synchronously coupled (i.e. share the same Gaussian random variables $\xi^{(1)}, \dots, \xi^{(4)}$ in (2.5)), that is,

$$(C.6) \quad \|\tilde{z}_{k+1} - z_{k+1}\|_{a,b}^2 < (1 - c(h))^2 \|\tilde{z}_k - z_k\|_{a,b}^2.$$

Let $c_2(h) = 1 - (1 - c(h))^2$, $z_j^\Delta = \tilde{z}_j - z_j$ for $j \in \mathbb{N}$, then (C.6) is equivalent to showing that

$$(C.7) \quad \left(z_k^\Delta\right)^T \left((1 - c_2(h)) \mathcal{M} - \mathcal{P}^T \mathcal{M} \mathcal{P}\right) z_k^\Delta > 0, \quad \text{where} \quad \mathcal{M} = \begin{pmatrix} I_d & bI_d \\ bI_d & aI_d \end{pmatrix},$$

and $z_{k+1}^\Delta = \mathcal{P} z_k^\Delta$ (\mathcal{P} depends on z_k and \tilde{z}_k , but we omit this in the notation).

Proving contraction for a general scheme is equivalent to showing that the matrix $\mathcal{H} := (1 - c_2(h))\mathcal{M} - \mathcal{P}^T\mathcal{M}\mathcal{P} \succ 0$ is positive definite. The matrix \mathcal{H} is symmetric and hence of the block form

$$(C.8) \quad \mathcal{H} = \begin{pmatrix} A & B \\ B^T & C \end{pmatrix},$$

where A, B, C are $d \times d$ matrices, then

$$(C.9) \quad \mathcal{H} \succ 0 \Leftrightarrow A \succ 0 \quad \text{and} \quad C - BA^{-1}B \succ 0,$$

as shown in Theorem 7.7.7 of [47]. Further it is straightforward to show that if A, B and C commute then

$$(C.10) \quad \mathcal{H} \succ 0 \Leftrightarrow A \succ 0 \quad \text{and} \quad AC - B^2 \succ 0.$$

Considering two synchronously coupled trajectories of the UBU scheme, such that they have common noise and consider the difference process $x^\Delta := (\tilde{x}_j - x_j)$, $v^\Delta = (\tilde{v}_j - v_j)$ and $z^\Delta = (x^\Delta, v^\Delta)$, where $z_j^\Delta = (x_j^\Delta, v_j^\Delta)$ for $j = k, k+1$ for $k \in \mathbb{N}$. Let $\eta = \exp\{-\gamma h/2\}$, and

$$Q = \int_0^1 \nabla^2 U(\tilde{x}_k + t(x_k - \tilde{x}_k)) dt.$$

By convexity, we have $mI_d \preceq Q \preceq MI_d$. Using the definition of the UBU scheme in (2.5), we can show that $z_{k+1}^\Delta = \mathcal{P}z_k^\Delta$ and $\mathcal{H} := (1 - c_2(h))\mathcal{M} - \mathcal{P}^T\mathcal{M}\mathcal{P} = \begin{pmatrix} A & B \\ B^T & C \end{pmatrix}$ has elements of the form

$$\begin{aligned} A &= -c_2(h)I_d + Q \left(2bh\eta + \frac{2h(1-\eta)}{\gamma} \right) + Q^2 \left(-ah^2\eta^2 - \frac{h^2(1-\eta)^2}{\gamma^2} - \frac{2bh^2\eta(1-\eta)}{\gamma} \right) \\ B &= \left((1-\eta^2) \left(b - \frac{1}{\gamma} \right) - bc_2(h) \right) I_d + Q^2 \left(-\frac{ah^2\eta^2(1-\eta)}{\gamma} - \frac{2bh^2\eta(1-\eta)^2}{\gamma^2} - \frac{h^2(1-\eta)^3}{\gamma^3} \right) \\ &\quad + Q \left(ah\eta^3 + \frac{h(\eta+1)(1-\eta)^2}{\gamma^2} + \frac{h(1-\eta)^2}{\gamma^2} + \frac{bh\eta^2(1-\eta)}{\gamma} + \frac{bh\eta(1-\eta)}{\gamma} + \frac{bh\eta(1-\eta^2)}{\gamma} \right) \\ C &= \left(a(1-\eta^4) - \frac{2b\eta^2(1-\eta^2)}{\gamma} - \frac{(1-\eta^2)^2}{\gamma^2} - ac_2(h) \right) I_d \\ &\quad + Q^2 \left(-\frac{ah^2\eta^2(1-\eta)^2}{\gamma^2} - \frac{2bh^2\eta(1-\eta)^3}{\gamma^3} - \frac{h^2(1-\eta)^4}{\gamma^4} \right) \\ &\quad + Q \left(\frac{2ah\eta^3(1-\eta)}{\gamma} + \frac{2bh\eta^2(1-\eta)^2}{\gamma^2} + \frac{2bh\eta(\eta+1)(1-\eta)^2}{\gamma^2} + \frac{2h(\eta+1)(1-\eta)^3}{\gamma^3} \right). \end{aligned}$$

We will now check that $\mathcal{H} \succ 0$ using (C.10). By firstly considering A we wish to show that all its eigenvalues are positive which can be precisely stated as

$$\begin{aligned} P_A(\lambda) &\geq -c_2(h) + \frac{2h\lambda}{\gamma} + \left(-\frac{1}{M} - \frac{2h}{\gamma} \right) h^2\lambda^2 \\ &\geq \frac{7h\lambda}{4\gamma} + \left(-\frac{1}{M} - \frac{1}{\gamma^2} \right) h^2\lambda^2 > 0, \end{aligned}$$

where λ is an eigenvalue of Q ($m \leq \lambda \leq M$), $P_A(\lambda)$ denotes the eigenvalue of A according to the same eigenvector (Q, A, B, C are all symmetric and have the same eigenvectors here). We used our assumptions that $\gamma^2 \geq M$, $1 - \eta \leq h\gamma/2$, and $h < \frac{1}{2\gamma}$. Hence, we have $A \succ 0$.

Now it remains to prove that $AC - B^2 \succ 0$, now we have that $AC - B^2$ is a polynomial of Q , which we denote $P_{AC-B^2}(Q)$ and hence has eigenvalues dictated by the eigenvalues λ of Q . Because the terms are more complicated than the previous discretizations, we choose a convenient way of expanding the expression, which can obtain positive definiteness. That is to expand the expression in terms of a . Therefore one can show that $P_{AC-B^2}(\lambda) = c_0 + c_1a + c_2a^2$, where

$$\begin{aligned} c_1 + c_2a &= \frac{h^2c_2(h)\lambda^2\eta^4}{\gamma^2} - \frac{2h^2c_2(h)\lambda^2\eta^2}{\gamma^2} - \frac{h^2\lambda^2\eta^4}{\gamma^2} + \frac{2h^2\lambda^2\eta^2}{\gamma^2} + \frac{2hc_2(h)\lambda\eta^4}{\gamma} - \frac{2h\lambda\eta^4}{\gamma} \\ &+ c_2(h)\eta^4 + \frac{h^2c_2(h)\lambda^2}{\gamma^2} - \frac{h^2\lambda^2}{\gamma^2} - \frac{2hc_2(h)\lambda}{\gamma} + \frac{2h\lambda}{\gamma} + c_2(h)^2 - c_2(h) \\ &+ a(-\eta^2h^2\lambda^2 + \eta^2h^2c_2(h)\lambda^2) \\ &\geq \frac{h\lambda}{\gamma}(1 - c_2(h)) \left(\frac{7}{4}(1 - \eta^4) - \frac{4h\lambda}{\gamma} - h\gamma \right). \end{aligned}$$

Furthermore, we have that

$$\begin{aligned} c_0 &= \frac{h^2(1 - c_2(h))\lambda^2\eta^4}{\gamma^4} - \frac{2h^2(1 - c_2(h))\lambda^2\eta^2}{\gamma^4} + \frac{2h(1 - c_2(h))\lambda\eta^4}{\gamma^3} + \frac{c_2(h)(1 - \eta^4)}{\gamma^2} \\ &- \frac{c_2(h)^2}{\gamma^2} + \frac{h^2\lambda^2(1 - c_2(h))}{\gamma^4} - \frac{2h\lambda(1 - c_2(h))}{\gamma^3} \\ &> \frac{h\lambda}{\gamma^3}(1 - c_2(h)) \left(\frac{h\lambda}{\gamma^3}(1 - \eta^2)^2 - 2(1 - \eta^4) \right), \end{aligned}$$

where now we combine this with the previous estimate

$$P_{AC-B^2}(\lambda) > \frac{h(1 - c_2(h))}{\gamma} \left(\frac{7}{4}(1 - \eta^4) - \frac{4h\lambda}{\gamma} - h\gamma - \frac{2\lambda(1 - \eta^4)}{\gamma^2} \right) > 0,$$

which is true when $\gamma \geq \sqrt{8M}$ and we have used the fact that $1 - \eta^4 \geq h\gamma$. Hence $AC - B^2 \succ 0$ and our contraction results hold. All computations can be checked using *Mathematica*. The first claim follows by Lemma C.4 using (C.6). The existence of a unique invariant distribution $\pi_h \in \mathcal{P}_p(\mathbb{R}^{2d})$ follows by the same argument as in [70][Corollary 20]. \square

PROOF OF COROLLARY C.8. By the triangle inequality, we have that for $a = \frac{1}{M}$, $b = \frac{1}{\gamma}$, any $n \in \mathbb{N}$ such that $n > t \cdot 2\gamma$,

$$\begin{aligned} &\mathcal{W}_{p,a,b}(\nu P_t^{\text{cont}}, \mu P_t^{\text{cont}}) \\ &\leq \mathcal{W}_{p,a,b}(\nu P_{t/n}^n, \mu P_{t/n}^n) + \mathcal{W}_{p,a,b}(\nu P_t^{\text{cont}}, \nu P_{t/n}^n) + \mathcal{W}_{p,a,b}(\mu P_t^{\text{cont}}, \mu P_{t/n}^n). \end{aligned}$$

The first term can be bounded using Proposition C.6, and the upper bound can be shown to converge to $\exp\left(-\frac{mt}{8\gamma}\right)$ as $n \rightarrow \infty$. The second and third terms can be shown to converge to 0 as $n \rightarrow \infty$ using the strong convergence of the UBU discretization towards the diffusion (strong order 1 under these assumptions), which was established in Section 7.7 of [86], and the claim of the corollary now follows. \square

EXAMPLE C.10. Considering the anisotropic Gaussian distribution on \mathbb{R}^2 with a m -strongly convex and M - ∇ Lipschitz potential $U : \mathbb{R}^2 \mapsto \mathbb{R}$ given by

$$U(x, y) = \frac{1}{2}mx^2 + \frac{1}{2}My^2.$$

For the BU scheme the transition matrix for the difference chain of synchronously coupled chains is given by the matrix

$$P = \begin{pmatrix} I - h \left(\frac{1-\eta^2}{\gamma} \right) Q & \frac{1-\eta^2}{\gamma} I \\ -h\eta^2 Q & \eta^2 I \end{pmatrix}, \text{ where } Q = \begin{pmatrix} m & 0 \\ 0 & M \end{pmatrix},$$

with eigenvalues

$$\frac{1 + \eta^2 - h \frac{1-\eta^2}{\gamma} \lambda \pm \sqrt{-4\eta^2 + \left(1 + \eta^2 - h \frac{1-\eta^2}{\gamma} \lambda\right)^2}}{2},$$

for $\lambda = m, M$. For stability and contraction, we require that

$$(C.11) \quad \lambda_{\max} := \max_{\lambda \in \{m, M\}} \left| \frac{1 + \eta^2 - h \frac{1-\eta^2}{\gamma} \lambda \pm \sqrt{-4\eta^2 + \left(1 + \eta^2 - h \frac{1-\eta^2}{\gamma} \lambda\right)^2}}{2} \right| < 1.$$

From this, we can compute the stepsize restrictions and the best convergence rate as, by Gelfand's formula, the asymptotic contraction rate exactly equals $1 - \lambda_{\max}$. Due to the convexity of the absolute value function it is necessary that $\frac{1}{2}|1 + \eta^2 - h \frac{1-\eta^2}{\gamma} M| < 1$, therefore $h < \sqrt{\frac{8}{M}}$, when $h < \frac{1}{2\gamma}$. In the moderate to high friction regime, the contraction rate can be written as

$$c = \frac{1 - \eta^2 + h \frac{1-\eta^2}{\gamma} m - \sqrt{-4h \left(\frac{1-\eta^2}{\gamma} \right) m + \left(1 - \eta^2 + h \frac{1-\eta^2}{\gamma} m\right)^2}}{2}$$

which can be shown to be $\mathcal{O}(mh/\gamma)$ for $\gamma \geq \mathcal{O}(\sqrt{M})$ and $h < \mathcal{O}(\frac{1}{\gamma})$ for appropriate constants. In the low friction regime, we set γ such that $-4\eta^2 + \left(1 + \eta^2 - h \frac{1-\eta^2}{\gamma} m\right)^2 = 0$, noting that the solution to this yields γ to be $\mathcal{O}(\sqrt{m})$. In this case, the eigenvalues of P are

$$\eta, \quad \frac{1}{2} \left(1 + \eta^2 - h \frac{1-\eta^2}{\gamma} M \pm \sqrt{-4\eta^2 + \left(1 + \eta^2 - h \frac{1-\eta^2}{\gamma} M\right)^2} \right),$$

with modulus η when $\left(1 + \eta^2 - h \frac{1-\eta^2}{\gamma} M\right)^2 < 4\eta^2$. This restriction implies that h is $\mathcal{O}(1/\sqrt{M})$. The contraction rate is therefore given by

$$c = 1 - \eta \geq \frac{h\gamma}{4} = \mathcal{O} \left(\sqrt{\frac{m}{M}} \right),$$

where h is $\mathcal{O}(1/\sqrt{M})$. We have the corresponding contraction rate results for UBU as well due to the fact that $(UBU)^n = U(BU)^{n-1}U$ and U is Lipschitz.

A key ingredient to establishing some variance bounds for the inexact gradient methods is to establish non-asymptotic bounds on the fourth moment of the distance to the minimizer. To do this we use a Lyapunov function similar to the one used for kinetic Langevin dynamics in [39] and inspired by [67]. Related Lyapunov functions have also been used in [35] for discretized kinetic Langevin dynamics and [53] for optimizers based on Langevin dynamic methods. These bounds provide novel drift conditions in L^4 for UBU scheme and can be extended to the case of stochastic gradients.

The following lemma will be useful for the argument.

LEMMA C.11 (Convexity bound). *For all $x \in \mathbb{R}^d$ and for a m -strongly convex, M - ∇ Lipschitz potential $U : \mathbb{R}^d \rightarrow \mathbb{R}$ with minimizer $x^* \in \mathbb{R}^d$ such that $\nabla U(x^*) = 0$, we have*

$$(x - x^*) \cdot (\nabla U(x) - \nabla U(x^*)) / 2 \geq \lambda (U(x) - U(x^*) + \gamma^2 \|x - x^*\|^2 / 4)$$

for

$$(C.12) \quad \lambda = \min \left(\frac{1}{4}, \frac{m}{\gamma^2} \right).$$

PROOF. By convexity, it follows that $(x - x^*) \cdot (\nabla U(x) - \nabla U(x^*)) / 4 \geq (U(x) - U(x^*)) / 4$, and by m -strong convexity, we have $(x - x^*) \cdot (\nabla U(x) - \nabla U(x^*)) / 4 \geq m \|x - x^*\|^2 / 4$. We obtain the result by adding up these two inequalities. \square

PROPOSITION C.12. *Consider the UBU scheme with the underlying potential $U : \mathbb{R}^d \rightarrow \mathbb{R}$ is M - ∇ Lipschitz and m -strongly convex. Denote $x^* \in \mathbb{R}^d$ to be the minimizer of U such that $\nabla U(x^*) = 0$ and $(x_k, v_k, \bar{x}_k)_{k \in \mathbb{N}}$ to be defined by (F.4)-(F.6) the iterates of the full gradient UBU scheme and the points of gradient evaluation within each iteration. Further assume that $h < \min \left(1, \frac{1}{2\gamma}, \frac{\lambda}{8\gamma(4+\lambda)} \right)$ and $\gamma^2 \geq M$, then we have*

$$\begin{aligned} \mathbb{E} [\|\bar{x}_k - x^*\|^4 \mid x_0, v_0] &\leq \frac{4}{m^2} \left[4 \left(1 - \frac{c_4(h)}{2} \right)^k (\gamma^4 \|x_0 - x^*\|^4 + \|v_0\|^4 + 122\gamma^2 h^2 d^2) \right. \\ &\quad \left. + 2 \frac{(6h\gamma d + 160h\gamma(1+\lambda^2))^2}{4c_4(h)} + 24h^2\gamma^2 d^2 \right], \end{aligned}$$

where

$$(C.13) \quad c_4(h) := h\lambda\gamma - 8h^2\gamma^2(4 + \lambda)$$

PROOF. Using the fact that $(UBU)^n = \mathcal{U}(\mathcal{BU})^{n-1}\mathcal{BU}$ we can consider convergence of \mathcal{BU} . We have that the \mathcal{BU} function can be written as the update rule

$$(C.14) \quad \bar{x}_{k+1} = \bar{x}_k + \frac{1 - \eta^2}{\gamma} (\bar{v}_k - h\nabla U(\bar{x}_k)) + \sqrt{\frac{2}{\gamma}} \left(\mathcal{Z}^{(1)} \left(h, \xi_{k+1}^{(1)} \right) - \mathcal{Z}^{(2)} \left(h, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)} \right) \right),$$

$$(C.15) \quad \bar{v}_{k+1} = \eta^2 (\bar{v}_k - h\nabla U(\bar{x}_k)) + \sqrt{2\gamma} \mathcal{Z}^{(2)} \left(h, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)} \right),$$

where we used the notation $(\bar{x}_k)_{k \in \mathbb{N}}$ because this is the point of the gradient evaluation at each step of UBU and is the same as the $(\bar{x}_k)_{k \in \mathbb{N}}$ in (F.5). As a reminder,

$$\mathcal{Z}^{(1)} \left(h, \xi_{k+1}^{(1)} \right) = \sqrt{h} \xi_{k+1}^{(1)}$$

$$\mathcal{Z}^{(2)}\left(h, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}\right) = \sqrt{\frac{1-\eta^4}{2\gamma}} \left(\sqrt{\frac{1-\eta^2}{1+\eta^2}} \cdot \frac{2}{\gamma h} \xi_{k+1}^{(1)} + \sqrt{1 - \frac{1-\eta^2}{1+\eta^2}} \cdot \frac{2}{\gamma h} \xi_{k+1}^{(2)} \right).$$

We choose our Lyapunov function $\mathcal{V} : \mathbb{R}^{2d} \rightarrow \mathbb{R}$, defined for $(x, v) \in \mathbb{R}^{2d}$ by

$$(C.16) \quad \mathcal{V}(x, v) := U(x) - U(x^*) + \frac{1}{4}\gamma^2 (\|x - x^* + \gamma^{-1}v\|^2 + \|\gamma^{-1}v\|^2 - \lambda\|x - x^*\|^2).$$

It is easy to check that for all $(x, v) \in \mathbb{R}^{2d}$, $\|x - x^* + \gamma^{-1}v\|^2 + \|\gamma^{-1}v\|^2 \geq \frac{1}{2}\|x - x^*\|^2$ and hence using (C.12),

$$(C.17) \quad \mathcal{V}(x, v) \geq \left(\frac{m}{2} + \frac{1}{16}\gamma^2\right) \|x - x^*\|^2.$$

In order to have control over fourth moments $\mathbb{E}[\|\bar{x}_k - x^*\|^4]$, we start with

$$\begin{aligned} & \mathbb{E} [\mathcal{V}(\bar{x}_{k+1}, \bar{v}_{k+1})^2 \mid \bar{x}_k, \bar{v}_k] = \\ & \mathbb{E} \left[\left(U(\bar{x}_{k+1}) - U(x^*) + \frac{1}{4}\gamma^2 (\|\bar{x}_{k+1} - x^* + \gamma^{-1}\bar{v}_{k+1}\|^2 + \|\gamma^{-1}\bar{v}_{k+1}\|^2 - \lambda\|\bar{x}_{k+1} - x^*\|^2) \right)^2 \mid \bar{x}_k, \bar{v}_k \right], \end{aligned}$$

and using [75][Lemma 1.2.3] we have

$$U(\bar{x}_{k+1}) - U(x^*) \leq U(\bar{x}_k) - U(x^*) + [\nabla U(\bar{x}_k) \cdot (\bar{x}_{k+1} - \bar{x}_k)] + \frac{M}{2} \|\bar{x}_{k+1} - \bar{x}_k\|^2$$

and

$$\begin{aligned} \mathbb{E} [\mathcal{V}(\bar{x}_{k+1}, \bar{v}_{k+1})^2 \mid \bar{x}_k, \bar{v}_k] & \leq \mathbb{E} \left[\left(U(\bar{x}_k) - U(x^*) + [\nabla U(\bar{x}_k) \cdot (\bar{x}_{k+1} - \bar{x}_k)] + \frac{M}{2} \|\bar{x}_{k+1} - \bar{x}_k\|^2 \right. \right. \\ & \left. \left. + \frac{1}{4}\gamma^2 (\|\bar{x}_{k+1} - x^* + \gamma^{-1}\bar{v}_{k+1}\|^2 + \|\gamma^{-1}\bar{v}_{k+1}\|^2 - \lambda\|\bar{x}_{k+1} - x^*\|^2) \right)^2 \mid \bar{x}_k, \bar{v}_k \right]. \end{aligned}$$

Now, we can decompose the right-hand side in the form

$$\mathbb{E} \left(\left(r(\bar{x}_k, \bar{v}_k) + \mathbf{s}(\bar{x}_k, \bar{v}_k) \cdot (\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}) + (\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)})^T \mathcal{T}(\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}) \right)^2 \mid \bar{x}_k, \bar{v}_k \right),$$

for $r : \mathbb{R}^{2d} \rightarrow \mathbb{R}$, $\mathbf{s} : \mathbb{R}^{2d} \rightarrow \mathbb{R}^{2d}$ and $\mathcal{T} \in \mathbb{R}^{2d \times 2d}$. We then have

$$\begin{aligned} \mathbb{E} [\mathcal{V}(\bar{x}_{k+1}, \bar{v}_{k+1})^2 \mid \bar{x}_k, \bar{v}_k] & \leq r^2(\bar{x}_k, \bar{v}_k) + \mathbb{E}_{\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}} \left(\left(\mathbf{s}(\bar{x}_k, \bar{v}_k) \cdot (\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}) \right)^2 \right. \\ & \left. + \left((\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)})^T \mathcal{T}(\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}) \right)^2 + 2r(\bar{x}_k, \bar{v}_k) (\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)})^T \mathcal{T}(\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}) \right), \end{aligned}$$

using the fact that $\xi_{k+1}^{(1)}$ and $\xi_{k+1}^{(2)}$ are independently distributed and have zero first and third moments. The terms r , \mathbf{s} and \mathcal{T} are given by

$$\begin{aligned} r(\bar{x}_k, \bar{v}_k) & = \mathcal{V}(\bar{x}_k, \bar{v}_k) - \frac{h\gamma}{2} \nabla U(\bar{x}_k) \cdot (\bar{x}_k - x^* + \gamma^{-1}\bar{v}_k) + \frac{1-\eta^2}{\gamma} \bar{v}_k \cdot \nabla U(\bar{x}_k) - \frac{\lambda(1-\eta^2)\gamma}{2} \langle \bar{x}_k - x^*, \bar{v}_k \rangle \\ & - \frac{1-\eta^4}{4} \|\bar{v}_k\|^2 - \frac{h\eta^4}{2} \bar{v}_k \cdot \nabla U(\bar{x}_k) - h \frac{1-\eta^2}{\gamma} \nabla U(\bar{x}_k) \cdot \nabla U(\bar{x}_k) + h^2 \frac{(1+\eta^4)}{4} \|\nabla U(\bar{x}_k)\|^2 \\ & + \left(\frac{M}{2} - \frac{\gamma^2\lambda}{4} \right) \left(\frac{1-\eta^2}{\gamma} \right)^2 \|\bar{v}_k - h\nabla U(\bar{x}_k)\|^2 + h \frac{\lambda(1-\eta^2)\gamma}{2} \langle \bar{x}_k - x^*, \nabla U(\bar{x}_k) \rangle, \end{aligned}$$

$$\begin{aligned}
\mathbf{s}(\bar{x}_k, \bar{v}_k) \cdot (\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}) &= ((\sqrt{h} - a_1)\xi_{k+1}^{(1)} - a_2\xi_{k+1}^{(2)}) \cdot \left(\frac{M\sqrt{2\gamma}}{2\gamma} \frac{1-\eta^2}{\gamma} (\bar{v}_k - h\nabla U(\bar{x}_k)) + \sqrt{\frac{2}{\gamma}} \nabla U(\bar{x}_k) \right) \\
&+ \frac{\sqrt{2\gamma h}}{4} \gamma \left(\bar{x}_k - x^* + \gamma^{-1}\bar{v}_k - \frac{h}{\gamma} \nabla U(\bar{x}_k) \right) \cdot \xi_{k+1}^{(1)} + \frac{\eta^2\sqrt{2\gamma}}{4} (\bar{v}_k - h\nabla U(\bar{x}_k)) \cdot (a_1\xi_{k+1}^{(1)} + a_2\xi_{k+1}^{(2)}) \\
&- \frac{\lambda\gamma\sqrt{2\gamma}}{4} \left(\bar{x}_k - x^* + \frac{1-\eta^2}{\gamma} (\bar{v}_k - h\nabla U(\bar{x}_k)) \cdot ((\sqrt{h} - a_1)\xi_{k+1}^{(1)} - a_2\xi_{k+1}^{(2)}) \right), \\
(\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)})^T \mathcal{T}(\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}) &= \left(\frac{M}{\gamma} - \frac{\lambda\gamma}{2} \right) \left\| (\sqrt{h} - a_1)\xi_{k+1}^{(1)} - a_2\xi_{k+1}^{(2)} \right\|^2 \\
&+ \frac{h\gamma}{2} \left\| \xi_{k+1}^{(1)} \right\|^2 + \frac{\gamma}{2} \left\| a_1\xi_{k+1}^{(1)} + a_2\xi_{k+1}^{(2)} \right\|^2,
\end{aligned}$$

where we have defined $\mathcal{Z}^{(2)}(h, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}) := a_1\xi_{k+1}^{(1)} + a_2\xi_{k+1}^{(2)}$ and $\mathcal{Z}^{(1)}(h, \xi_{k+1}^{(1)}) := \sqrt{h}\xi_{k+1}^{(1)}$ and $\mathcal{Z}^{(1)}(h, \xi_{k+1}^{(1)}) - \mathcal{Z}^{(2)}(h, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}) = (\sqrt{h} - a_1)\xi_{k+1}^{(1)} - a_2\xi_{k+1}^{(2)}$ with $|\sqrt{h} - a_1| \leq 2\sqrt{h}$, $|a_2| \leq \sqrt{h}$ and $|a_1| \leq \sqrt{h}$.

We start by bounding the deterministic component r :

$$\begin{aligned}
r(\bar{x}_k, \bar{v}_k) &= \mathcal{V}(\bar{x}_k, \bar{v}_k) - \frac{h\gamma}{2} \nabla U(\bar{x}_k) \cdot (\bar{x}_k - x^* + \gamma^{-1}\bar{v}_k) + \frac{1-\eta^2}{\gamma} \bar{v}_k \cdot \nabla U(\bar{x}_k) - \frac{\lambda(1-\eta^2)\gamma}{2} \langle \bar{x}_k - x^*, \bar{v}_k \rangle \\
&- \frac{1-\eta^4}{4} \|\bar{v}_k\|^2 - \frac{h\eta^4}{2} \bar{v}_k \cdot \nabla U(\bar{x}_k) + \mathcal{O}(h^2)
\end{aligned}$$

where the higher-order terms are given by

$$\begin{aligned}
&- h \frac{1-\eta^2}{\gamma} \nabla U(\bar{x}_k) \cdot \nabla U(\bar{x}_k) + \left(\frac{M}{2} - \frac{\gamma^2\lambda}{4} \right) \left(\frac{1-\eta^2}{\gamma} \right)^2 \|\bar{v}_k - h\nabla U(\bar{x}_k)\|^2 \\
&+ h \frac{\lambda(1-\eta^2)\gamma}{2} \langle \bar{x}_k - x^*, \nabla U(\bar{x}_k) \rangle + h^2 \frac{(1+\eta^4)}{4} \|\nabla U(\bar{x}_k)\|^2.
\end{aligned}$$

Using Lemma C.11 we have

$$\begin{aligned}
r(\bar{x}_k, \bar{v}_k) &\leq \mathcal{V}(\bar{x}_k, \bar{v}_k) - h\gamma\lambda \left(U(\bar{x}_k) - U(x^*) + \frac{\gamma^2}{4} \|\bar{x}_k - x^*\|^2 + \frac{1-\eta^2}{2h} \langle \bar{x}_k - x^*, \bar{v}_k \rangle + \frac{1-\eta^4}{4h\gamma\lambda} \|\bar{v}_k\|^2 \right) \\
&+ \mathcal{O}(h^2) \\
&\leq (1 - h\lambda\gamma) \mathcal{V}(\bar{x}_k, \bar{v}_k) + h\gamma\lambda \left(\frac{1-\eta^4}{4h\gamma\lambda} - \frac{1}{2\lambda} \right) \|\bar{v}_k\|^2 \\
&+ h\gamma\lambda \left(\frac{1-\eta^2}{2h} - \frac{\gamma}{2} \right) \langle \bar{x}_k - x^*, \bar{v}_k \rangle + \mathcal{O}(h^2) \\
&\leq (1 - h\lambda\gamma) \mathcal{V}(\bar{x}_k, \bar{v}_k) + h\gamma\lambda \left(\frac{1-\eta^2}{2h} - \frac{\gamma}{2} \right) \langle \bar{x}_k - x^*, \bar{v}_k \rangle + \mathcal{O}(h^2),
\end{aligned}$$

where we have used

$$\left(-\frac{h}{2} + \frac{1-\eta^2}{\gamma} - \frac{h\eta^4}{2} \right) \bar{v}_k \cdot \nabla U(\bar{x}_k) \leq \frac{h^2\gamma}{2} |\bar{v}_k \cdot \nabla U(\bar{x}_k)|,$$

due to the fact that for all $0 < x < 1$, $0 \leq -x + 2(1 - e^{-x}) - xe^{-2x} \leq x^2$ and $0 < h\gamma < 1$.

We group this term into higher-order terms and use the fact that $1 - \eta^2 \geq h\gamma - \frac{(h\gamma)^2}{2}$ to arrive

at

$$h\gamma\lambda\left(\frac{1-\eta^2}{2h}-\frac{\gamma}{2}\right)\langle\bar{x}_k-x^*,\bar{v}_k\rangle\leq h\gamma\lambda\left(\frac{\gamma}{2}-\frac{1-\eta^2}{2h}\right)|\langle\bar{x}_k-x^*,\bar{v}_k\rangle|\leq\lambda\frac{h^2\gamma^3}{4}|\langle\bar{x}_k-x^*,\bar{v}_k\rangle|.$$

We again group this into the higher-order terms. Assuming $h < 1$, we find that the second-order terms are bounded by

$$\begin{aligned} & Mh^2(\|\bar{v}_k\|^2+h^2M^2\|\bar{x}_k-x^*\|^2)+h^2\frac{\gamma^2\lambda}{2}M\|\bar{x}_k-x^*\|^2 \\ & +\frac{h^2\gamma}{2}\left(\sqrt{M}\|\bar{v}_k\|^2+M^{3/2}\|\bar{x}_k-x^*\|^2\right)+\lambda\frac{h^2\gamma^3}{4}\left(\gamma\|\bar{x}_k-x^*\|^2+\frac{1}{\gamma}\|\bar{v}_k\|^2\right)+\frac{h^2M^2}{2}\|\bar{x}_k-x^*\|^2. \end{aligned}$$

Assuming that $\lambda \leq \frac{1}{4}$ we have, for all $x, v \in \mathbb{R}^d$,

$$8\mathcal{V}(x, v) \geq \|v\|^2 \quad 16\mathcal{V}(x, v) \geq \gamma^2\|x-x^*\|^2$$

and using $h < \frac{1}{2\sqrt{M}}$, the $\mathcal{O}(h^2)$ terms are bounded by

$$\begin{aligned} & h^2\left(\gamma^2+\frac{\gamma^2\lambda}{4}+\frac{\gamma^2}{2}\right)\|\bar{v}_k\|^2+\gamma^2h^2\left(\frac{M}{4}+\frac{M\lambda}{2}+\frac{M}{2}+\frac{\gamma^2\lambda}{4}+\frac{M^2}{2\gamma^2}\right)\|\bar{x}_k-x^*\|^2 \\ & \leq 8h^2\gamma^2(4+\lambda)\mathcal{V}(\bar{x}_k, \bar{v}_k). \end{aligned}$$

Therefore

$$r(\bar{x}_k, \bar{v}_k) \leq (1-h\lambda\gamma+8h^2\gamma^2(4+\lambda))\mathcal{V}(\bar{x}_k, \bar{v}_k).$$

Now let us define $c_4(h) := h\lambda\gamma - 8h^2\gamma^2(4+\lambda)$, then we have that

$$r^2(\bar{x}_k, \bar{v}_k) \leq (1-c_4(h))^2\mathcal{V}^2(\bar{x}_k, \bar{v}_k)$$

and

$$\begin{aligned} & 2r(\bar{x}_k, \bar{v}_k)\mathbb{E}_{\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}}\left[(\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)})^T\mathcal{T}(\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)})\right] \leq \\ & 2(1-c_4(h))\mathcal{V}(\bar{x}_k, \bar{v}_k)\mathbb{E}_{\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}}\left[(\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)})^T\mathcal{T}(\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)})\right]. \end{aligned}$$

From the fact that $\lambda\gamma/2 \leq M/\gamma$ (due to Lemma C.11) and $\gamma^2 \geq 8M$ we have the estimates

$$\begin{aligned} & \mathbb{E}_{\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}}\left[(\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)})^T\mathcal{T}(\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)})\right] = \mathbb{E}_{\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}}\left[\left(\frac{M}{\gamma}-\frac{\lambda\gamma}{2}\right)\|(\sqrt{h}-a_1)\xi_{k+1}^{(1)}-a_2\xi_{k+1}^{(2)}\|^2\right] \\ & +\frac{h\gamma}{2}\|\xi_{k+1}^{(1)}\|^2+\frac{\gamma}{2}\|a_1\xi_{k+1}^{(1)}+a_2\xi_{k+1}^{(2)}\|^2 \leq 3h\gamma d \\ & \mathbb{E}_{\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}}\left[\left((\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)})^T\mathcal{T}(\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)})\right)^2\right] \\ & \leq \mathbb{E}_{\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}}\left[\left(2h\gamma\|\xi_{k+1}^{(1)}\|^2+2h\gamma\|\xi_{k+1}^{(2)}\|^2\right)^2\right] \leq 24h^2\gamma^2d^2. \end{aligned}$$

Therefore the remaining term we need to bound is $\mathbb{E}_{\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}}\left(\mathbf{s}(\bar{x}_k, \bar{v}_k) \cdot (\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)})\right)^2 = \|\mathbf{s}(\bar{x}_k, \bar{v}_k)\|^2$, where

$$\mathbf{s}(\bar{x}_k, \bar{v}_k) \cdot (\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)})$$

$$\begin{aligned}
&= ((\sqrt{h} - a_1)\xi_{k+1}^{(1)} - a_2\xi_{k+1}^{(2)}) \cdot \left(\frac{M\sqrt{2\gamma}}{2\gamma} \frac{1-\eta^2}{\gamma} (\bar{v}_k - h\nabla U(\bar{x}_k)) + \sqrt{\frac{2}{\gamma}} \nabla U(\bar{x}_k) \right) \\
&+ \frac{\sqrt{2\gamma h}}{4} \gamma \left(\bar{x}_k - x^* + \gamma^{-1}\bar{v}_k - \frac{h}{\gamma} \nabla U(\bar{x}_k) \right) \cdot \xi_{k+1}^{(1)} + \frac{\eta^2\sqrt{2\gamma}}{4} (\bar{v}_k - h\nabla U(\bar{x}_k)) \cdot (a_1\xi_{k+1}^{(1)} + a_2\xi_{k+1}^{(2)}) \\
&- \frac{\lambda\gamma\sqrt{2\gamma}}{4} \left(\bar{x}_k - x^* + \frac{1-\eta^2}{\gamma} (\bar{v}_k - h\nabla U(\bar{x}_k)) \right) \cdot ((\sqrt{h} - a_1)\xi_{k+1}^{(1)} - a_2\xi_{k+1}^{(2)}),
\end{aligned}$$

using that $\gamma^2\|x - x^*\|^2 \leq 16\mathcal{V}(x, v)$ and $\|v\|^2 \leq 8\mathcal{V}(x, v)$ for all $x, v \in \mathbb{R}^d$ we have

$$\|\mathbf{s}(\bar{x}_k, \bar{v}_k)\|^2 = \|s_1(\bar{x}_k, \bar{v}_k)\|^2 + \|s_2(\bar{x}_k, \bar{v}_k)\|^2,$$

where

$$\begin{aligned}
\|s_1(\bar{x}_k, \bar{v}_k)\|^2 &\leq h \left(\left(2\frac{M(1-\eta^2)}{\gamma\sqrt{2\gamma}} + \frac{\eta^2\sqrt{2\gamma}}{4} + 2\frac{\lambda\sqrt{2\gamma}(1-\eta^2)}{4} \right) \|\bar{v}_k\| \right. \\
&+ \left(2\frac{hM^2(1-\eta^2)}{\gamma\sqrt{2\gamma}} + 2\sqrt{\frac{2}{\gamma}}M + \frac{\sqrt{2\gamma}hM}{4} + \frac{\eta^2\sqrt{2\gamma}hM}{4} + 2\frac{\lambda\sqrt{2\gamma}(1-\eta^2)hM}{4} \right) \|\bar{x}_k - x^*\| \\
&+ \left. \frac{\sqrt{2\gamma}^{3/2}}{4} \|\bar{x}_k - x^* + \gamma^{-1}\bar{v}_k\| \right)^2 \\
&\leq h \left((2\sqrt{\gamma} + \lambda\sqrt{\gamma})\sqrt{\mathcal{V}(\bar{x}_k, \bar{v}_k)} + \left(\frac{2\gamma^{3/2}}{\sqrt{8}} + \frac{\lambda\gamma^{3/2}}{32} \right) \frac{4}{\gamma} \sqrt{\mathcal{V}(\bar{x}_k, \bar{v}_k)} + \frac{5}{2}\sqrt{\gamma}\sqrt{\mathcal{V}(\bar{x}_k, \bar{v}_k)} \right)^2 \\
&\leq 110h\gamma(1 + \lambda^2)\mathcal{V}(\bar{x}_k, \bar{v}_k)
\end{aligned}$$

for $\gamma^2 \geq \sqrt{8M}$ and $h < \frac{1}{2\gamma}$ and

$$\begin{aligned}
\|s_2(\bar{x}_k, \bar{v}_k)\|^2 &\leq h \left(\left(2\frac{M(1-\eta^2)}{\gamma\sqrt{2\gamma}} + \frac{\eta^2\sqrt{2\gamma}}{4} + 2\frac{\lambda\sqrt{2\gamma}(1-\eta^2)}{4} \right) \|\bar{v}_k\| \right. \\
&+ \left. \left(2\frac{hM^2(1-\eta^2)}{\gamma\sqrt{2\gamma}} + 2\sqrt{\frac{2}{\gamma}}M + \frac{\eta^2\sqrt{2\gamma}hM}{4} + 2\frac{\lambda\sqrt{2\gamma}(1-\eta^2)hM}{4} \right) \|\bar{x}_k - x^*\| \right)^2 \\
&\leq h \left((2\sqrt{\gamma} + \lambda\sqrt{\gamma})\sqrt{\mathcal{V}(\bar{x}_k, \bar{v}_k)} + \left(\frac{2\gamma^{3/2}}{\sqrt{8}} + \frac{\lambda\gamma^{3/2}}{32} \right) \frac{4}{\gamma} \sqrt{\mathcal{V}(\bar{x}_k, \bar{v}_k)} \right)^2 \\
&\leq 50h\gamma(1 + \lambda^2)\mathcal{V}(\bar{x}_k, \bar{v}_k).
\end{aligned}$$

Therefore $\mathbb{E}_{\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}} \left[\left(\mathbf{s}(\bar{x}_k, \bar{v}_k) \cdot (\xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}) \right)^2 \right] \leq 160h\gamma(1 + \lambda^2)\mathcal{V}(\bar{x}_k, \bar{v}_k)$. Combining estimates, we have the drift inequality

$$\begin{aligned}
\mathbb{E} \left[\mathcal{V}(\bar{x}_{k+1}, \bar{v}_{k+1})^2 \mid \bar{x}_k, \bar{v}_k \right] &\leq (1 - c_4(h))^2 \mathcal{V}^2(\bar{x}_k, \bar{v}_k) + 6h\gamma d(1 - c_4(h)) \mathcal{V}(\bar{x}_k, \bar{v}_k) \\
&+ 160h\gamma(1 + \lambda^2)\mathcal{V}(\bar{x}_k, \bar{v}_k) + 24h^2\gamma^2d^2.
\end{aligned}$$

We will now use the quadratic property that states, for $b_1, b_2 > 0$,

$$b_2x^2 + \frac{b_1}{4b_2} \geq b_1x,$$

for all $x \in \mathbb{R}$ and therefore

$$c_4(h)\mathcal{V}^2(\bar{x}_k, \bar{v}_k) + \frac{(6h\gamma d + 160h\gamma(1 + \lambda^2))^2}{4c_4(h)} \geq 6h\gamma d(1 - c_4(h))\mathcal{V}(\bar{x}_k, \bar{v}_k) + 160h\gamma(1 + \lambda^2)\mathcal{V}(\bar{x}_k, \bar{v}_k)$$

and therefore for $c_4(h) < \frac{1}{2}$ (which is satisfied when $h < \frac{1}{2\gamma}$ and $\lambda < 1$, which is satisfied as $\lambda \leq M/2\gamma^2 \leq 1$ for $\gamma^2 \geq \frac{M}{2}$) we have

(C.18)

$$\mathbb{E} [\mathcal{V}(\bar{x}_{k+1}, \bar{v}_{k+1})^2 | \bar{x}_k, \bar{v}_k] \leq \left(1 - \frac{c_4(h)}{2}\right) \mathcal{V}^2(\bar{x}_k, \bar{v}_k) + \frac{(6h\gamma d + 160h\gamma(1 + \lambda^2))^2}{4c_4(h)} + 24h^2\gamma^2 d^2,$$

then globally, we have

$$\begin{aligned} \frac{m^2}{4} \mathbb{E} [\|\bar{x}_k - x^*\|^4 | y_0, v_0] &\leq \mathbb{E} [\mathcal{V}(\bar{x}_k, v_k)^2 | y_0, v_0] \\ &\leq \left(1 - \frac{c_4(h)}{2}\right)^k \mathcal{V}^2(\bar{x}_0, \bar{v}_0) + 2 \frac{(6h\gamma d + 160h\gamma(1 + \lambda^2))^2}{4c_4(h)} + 24h^2\gamma^2 d^2. \end{aligned}$$

Now, we have proved this for the iterates of \mathcal{BU} , where we wish to use the relation $(\mathcal{UBU})^k = \mathcal{U}(\mathcal{BU})^{k-1}\mathcal{BU}$. In this case, we have that \bar{x}_k , the $(k+1)$ -th point of approximate gradient/full gradient evaluation, is precisely the position after $\mathcal{U}(\mathcal{BU})^k$. It follows that

$$\frac{m^2}{4} \mathbb{E} [\|\bar{x}_k - x^*\|^4 | \bar{x}_0, \bar{v}_0] \leq \left(1 - \frac{c_4(h)}{2}\right)^k \mathcal{V}^2(\bar{x}_0, \bar{v}_0) + 2 \frac{(6h\gamma d + 160h\gamma(1 + \lambda^2))^2}{4c_4(h)} + 24h^2\gamma^2 d^2,$$

where $(\bar{x}_0, \bar{v}_0) = \mathcal{U}(x_0, v_0, h/2, \xi_0^{(1)}, \xi_0^{(2)})$. It is easy to show that $\mathcal{V}(x, v) \leq \gamma^2\|x - x^*\|^2 + \|v\|^2$ for all $(x, v) \in \mathbb{R}^{2d}$ using [75][Lemma 1.2.3] and that $\gamma^2 \geq M$. Therefore

$$\begin{aligned} \mathbb{E} [\mathcal{V}^2(\mathcal{U}(x_0, v_0, h/2, \xi_0^{(1)}, \xi_0^{(2)})) | x_0, v_0] &\leq \mathbb{E} [2\gamma^4\|\bar{x}_0 - x^*\|^4 + 2\|\bar{v}_0\|^4 | x_0, v_0] \\ &\leq 2\gamma^4 \mathbb{E} \left\| x_0 - x^* + \frac{1 - \eta^2}{\gamma} v_0 + \sqrt{\frac{2}{\gamma}} \left(\mathcal{Z}^{(1)}(h/2, \xi_0^{(1)}) - \mathcal{Z}^{(2)}(h/2, \xi_0^{(1)}, \xi_0^{(2)}) \right) \right\|^4 \\ &\quad + 2\mathbb{E} \left\| \eta v_0 + \sqrt{2\gamma} \mathcal{Z}^{(2)}(h/2, \xi_0^{(1)}, \xi_0^{(2)}) \right\|^4 \\ &\leq 4\gamma^4\|x_0 - x^*\|^4 + 4\|v_0\|^4 + \mathbb{E} \left[32\gamma^2\|\mathcal{Z}^{(1)}(h/2, \xi_0^{(1)})\|^4 + 40\gamma^2\|\mathcal{Z}^{(2)}(h/2, \xi_0^{(1)}, \xi_0^{(2)})\|^4 \right] \\ &\leq 4\gamma^4\|x_0 - x^*\|^4 + 4\|v_0\|^4 + 8\gamma^2 h^2 d^2 + 480\gamma^2 h^2 d^2, \end{aligned}$$

where we have used that $U(\bar{x}_0) - U(x^*) \leq \frac{M}{2}\|\bar{x}_0 - x^*\|^2$ in the first inequality and naive bounds on the fourth moments of the Gaussian increments. Hence, we arrive at the estimate

$$\begin{aligned} \mathbb{E} [\|\bar{x}_k - x^*\|^4 | x_0, v_0] &\leq \frac{4}{m^2} \left[4 \left(1 - \frac{c_4(h)}{2}\right)^k (\gamma^4\|x_0 - x^*\|^4 + \|v_0\|^4 + 122\gamma^2 h^2 d^2) \right. \\ &\quad \left. + 2 \frac{(6h\gamma d + 160h\gamma(1 + \lambda^2))^2}{4c_4(h)} + 24h^2\gamma^2 d^2 \right], \end{aligned}$$

for the UBU scheme with full gradients. \square

APPENDIX D: VARIANCE BOUNDS FOR UBUBU ESTIMATOR WITH EXACT GRADIENTS

D.1. Variance bound of $D_{l,l+1}$. To bound the variance of $D_{l,l+1}$ we use strong error estimates for the UBU integrator using the results of [86].

In this analysis we define for random vectors $z_1, z_2 \in \mathbb{R}^{2d}$ the L^2 norm $\|z_1\|_{L^2, a, b} = \mathbb{E} \left(\|z_1\|_{a, b}^2 \right)^{1/2}$ and respective inner product $\langle z_1, z_2 \rangle_{L^2, a, b} = \mathbb{E} (z_1^T \mathcal{M} z_2)$, where

$$\mathcal{M} = \begin{pmatrix} I_d & bI_d \\ bI_d & aI_d \end{pmatrix}.$$

ASSUMPTION D.1 (Local Strong Error [86]). *Let $\phi(z, t, (W_s)_{s=0}^t)$ be the solution of the continuous dynamics (1.1) with initial condition $z \in \mathbb{R}^{2d}$ up to time t , with Brownian motion $(W_s)_{s=0}^t$. Let $\psi_h(z, t, (W_s)_{s=0}^t)$ be the solution of a numerical discretization with initial condition $z \in \mathbb{R}^{2d}$ up to time t , with Brownian motion $(W_s)_{s=0}^t$ and stepsize h . Let $z' \sim \pi$, then we assume that*

$$\psi_h(z', h, (W_s)_{s=0}^h) - \phi(z', h, (W_s)_{s=0}^h) = \alpha_h(z', (W_s)_{s=0}^h) + \beta_h(z', (W_s)_{s=0}^h),$$

where

$$\|\alpha_h(z', (W_s)_{s=0}^h)\|_{L^2, a, b} \leq C_1 h^{q+1/2},$$

$$\|\beta_h(z', (W_s)_{s=0}^h)\|_{L^2, a, b} \leq C_2 h^{q+1},$$

and

$$\begin{aligned} & \left| \left\langle \psi_h(z', h, (W_s)_{s=0}^h) - \psi_h(z, h, (W_s)_{s=0}^h), \alpha_h(z', (W_s)_{s=0}^h) \right\rangle_{L^2, a, b} \right| \\ & \leq C_0 h \|z' - z\|_{L^2, a, b} \|\alpha_h(z', (W_s)_{s=0}^h)\|_{L^2, a, b}. \end{aligned}$$

for some $C_0, C_1, C_2 > 0$.

We restate Assumptions 3.6-3.13 here for easier readability.

ASSUMPTION 3.6 (M - ∇ Lipschitz). $U : \mathbb{R}^d \rightarrow \mathbb{R}$ is twice continuously differentiable and there exists $M > 0$ such that for all $x, y \in \mathbb{R}^d$

$$\|\nabla U(x) - \nabla U(y)\| \leq M \|x - y\|.$$

ASSUMPTION 3.7 (m -strong convexity). $U : \mathbb{R}^d \rightarrow \mathbb{R}$ is continuously differentiable and there exists $m > 0$ such that for all $x, y \in \mathbb{R}^d$

$$\langle \nabla U(x) - \nabla U(y), x - y \rangle \geq m |x - y|^2.$$

ASSUMPTION 3.10 (M_1^s -strongly Hessian Lipschitz). $U : \mathbb{R}^d \rightarrow \mathbb{R}$ is three times continuously differentiable and M_1^s -strongly Hessian Lipschitz if there exists $M_1^s > 0$ such that

$$\|\nabla^3 U(x)\|_{\{1,2\}\{3\}} \leq M_1^s$$

for all $x \in \mathbb{R}^d$.

ASSUMPTION 3.12 (1-Lipschitzness of f). f is a 1-Lipschitz function with respect to the Euclidean distance on \mathbb{R}^{2d} , that only depends on x , not v (i.e. $f(x, v) = f(x, v')$ for any $x, v, v' \in \mathbb{R}^d$).

ASSUMPTION 3.13 (Distance of initial distribution from target). The initial distribution on $\Lambda = \mathbb{R}^{2d}$ satisfy that $\mathcal{W}_2(\pi, \mu_0) \leq c_{\mu_0} \sqrt{\frac{d}{m}}$, for some $c_{\mu_0} > 0$.

We make use of the following proposition, essentially due to [86].

PROPOSITION D.2. Suppose a numerical scheme approximating (1.1) satisfies Assumption D.1, with a potential which satisfies Assumptions 3.6-3.10, and $\psi_h(z, h, (W_s)_0^h) \sim P_h(z, \cdot)$ satisfies the Wasserstein contractivity condition (C.3) for $p = 2$, and some $a, b > 0$, $a < b^2$.

Let $\phi(z, t, (W_s)_{s=0}^t)$ be the solution of the continuous dynamics (1.1) with initial condition $z \in \mathbb{R}^{2d}$ up to time t , with Brownian motion $(W_s)_{s=0}^t$. Let $\psi_h(z, t, (W_s)_{s=0}^t)$ be the solution of a numerical discretization with initial condition $z \in \mathbb{R}^{2d}$ up to time t , with Brownian motion $(W_s)_{s=0}^t$ and stepsize $h > 0$ satisfying that

$$(D.1) \quad (1 - c(h))^2 + C_0^2 h^2 < 1.$$

Then for any $k \geq 0$, any z_0 such that $\|z_0\|_{L^2, a, b} < \infty$, and $Z^0 \sim \pi$, we have

$$\begin{aligned} & \left\| \psi_h \left(z_0, kh, (W_s)_{s=0}^{kh} \right) - \phi \left(Z^0, kh, (W_s)_{s=0}^{kh} \right) \right\|_{L^2, a, b} \\ & \leq (1 - R(h))^k \|z_0 - Z^0\|_{L^2, a, b} + \sqrt{2} C_1 \frac{h^{q+1/2}}{\sqrt{R(h)}} + \frac{2C_2 h^{q+1}}{R(h)}, \end{aligned}$$

where $R(h) = 1 - \sqrt{(1 - c(h))^2 + C_0^2 h^2}$.

In particular, the discretization scheme admits a stationary distribution π_h , and its bias can be bounded as

$$(D.2) \quad \mathcal{W}_{2, a, b}(\pi_h, \pi) \leq \sqrt{2} C_1 \frac{h^{q+1/2}}{\sqrt{R(h)}} + \frac{2C_2 h^{q+1}}{R(h)}.$$

PROOF. Introduce the notation

$$Z^n := \phi \left(Z^0, nh, (W_s)_{s=0}^{nh} \right), \quad z_n := \psi_h \left(z_0, nh, (W_s)_{s=0}^{nh} \right)$$

for all $n \in \mathbb{N}$. Using the assumption $Z^0 \sim \pi$, we also have $Z^n \sim \pi$, since the kinetic Langevin dynamics keeps π invariant. By Assumption D.1, we then have

(D.3)

$$\begin{aligned}
& \left\| z_k - Z^k \right\|_{L^2, a, b} = \left\| \psi_h \left(z_{k-1}, h, (W_s)_{(k-1)h}^{kh} \right) - \phi \left(Z^{k-1}, h, (W_s)_{(k-1)h}^{kh} \right) \right\|_{L^2, a, b} \\
& = \left\| \psi_h \left(z_{k-1}, h, (W_s)_{(k-1)h}^{kh} \right) - \psi_h \left(Z^{k-1}, h, (W_s)_{(k-1)h}^{kh} \right) \right. \\
& \quad \left. + \psi_h \left(Z^{k-1}, h, (W_s)_{(k-1)h}^{kh} \right) - \phi \left(Z^{k-1}, h, (W_s)_{(k-1)h}^{kh} \right) \right\|_{L^2, a, b} \\
& = \left\| \psi_h \left(z_{k-1}, h, (W_s)_{(k-1)h}^{kh} \right) - \psi_h \left(Z^{k-1}, h, (W_s)_{(k-1)h}^{kh} \right) \right. \\
& \quad \left. + \alpha_h \left(Z^{k-1}, (W_s)_{(k-1)h}^{kh} \right) + \beta_h \left(Z^{k-1}, (W_s)_{(k-1)h}^{kh} \right) \right\|_{L^2, a, b} \\
& \leq \left\| \beta^{k-1} \right\|_{L^2, a, b} + \left\| \psi_h \left(z_{k-1}, h, (W_s)_{(k-1)h}^{kh} \right) - \psi_h \left(Z^{k-1}, h, (W_s)_{(k-1)h}^{kh} \right) + \alpha^{k-1} \right\|_{L^2, a, b},
\end{aligned}$$

where α^{k-1} and β^{k-1} are defined as

$$\begin{aligned}
& \psi_h \left(Z^{k-1}, h, (W_s)_{(k-1)h}^{kh} \right) - \phi \left(Z^{k-1}, h, (W_s)_{(k-1)h}^{kh} \right) \\
& = \alpha_h \left(Z^{k-1}, (W_s)_{(k-1)h}^{kh} \right) + \beta_h \left(Z^{k-1}, (W_s)_{(k-1)h}^{kh} \right) \\
& := \alpha^{k-1} + \beta^{k-1}.
\end{aligned}$$

Assumption D.1, and the Wasserstein contractivity condition (C.3) then together imply

$$\begin{aligned}
& \left\| \psi_h \left(z_{k-1}, h, (W_s)_{(k-1)h}^{kh} \right) - \psi_h \left(Z^{k-1}, h, (W_s)_{(k-1)h}^{kh} \right) + \alpha^{k-1} \right\|_{L^2, a, b} \\
& = \left(\left\| \alpha^{k-1} \right\|_{L^2, a, b}^2 + \left\| \psi_h \left(z_{k-1}, h, (W_s)_{(k-1)h}^{kh} \right) - \psi_h \left(Z^{k-1}, h, (W_s)_{(k-1)h}^{kh} \right) \right\|_{L^2, a, b}^2 \right. \\
& \quad \left. + 2 \left\langle \alpha^{k-1}, \psi_h \left(z_{k-1}, h, (W_s)_{(k-1)h}^{kh} \right) - \psi_h \left(Z^{k-1}, h, (W_s)_{(k-1)h}^{kh} \right) \right\rangle_{L^2, a, b} \right)^{1/2} \\
& \leq \left(\left\| \alpha^{k-1} \right\|_{L^2, a, b}^2 + (1 - c(h))^2 \left\| z_{k-1} - Z^{k-1} \right\|_{L^2, a, b}^2 \right. \\
& \quad \left. + 2C_0 h \left\| \alpha^{k-1} \right\|_{L^2, a, b} \left\| z_{k-1} - Z^{k-1} \right\|_{L^2, a, b} \right)^{1/2} \\
& \leq \left(2 \left\| \alpha^{k-1} \right\|_{L^2, a, b}^2 + ((1 - c(h))^2 + C_0^2 h^2) \left\| z_{k-1} - Z^{k-1} \right\|_{L^2, a, b}^2 \right)^{1/2} \\
& \leq \left(2C_1^2 h^{2q+1} + ((1 - c(h))^2 + C_0^2 h^2) \left\| z_{k-1} - Z^{k-1} \right\|_{L^2, a, b}^2 \right)^{1/2}.
\end{aligned}$$

Lemma 28 of [86] states that if a sequence of nonnegative real numbers $(a_n)_{n \geq 0}$ satisfies that $a_{n+1} \leq \sqrt{(1-A)^2 a_n^2 + B} + C$ with $A \in (0, 1)$, $B \geq 0$, $C \geq 0$, then for every $n \geq 0$,

$$a_n \leq (1-A)^n a_0 + \sqrt{\frac{B}{A}} + \frac{C}{A}.$$

Using this for $a_n = \|z_n - Z^n\|_{L^2, a, b}$, we have that

$$\|z_k - Z^k\|_{L^2, a, b} \leq (1 - R(h))^k \|z_0 - Z^0\|_{L^2, a, b} + \sqrt{2} \frac{C_1 h^{q+1/2}}{\sqrt{R(h)}} + \frac{2C_2 h^{q+1}}{R(h)},$$

where $R(h) = 1 - \sqrt{(1 - c(h))^2 + C_0^2 h^2}$, which is our first claim.

The existence of a stationary distribution π_h follows from Lemma C.4. The bound on the bias follows by letting $k \rightarrow \infty$. \square

We now are in a position to present our first result related to the variance of our unbiased scheme, which is a bound on the variance related to the global strong error or convergence. This is given below.

PROPOSITION D.3. *Suppose a numerical scheme approximating (1.1) satisfies the same assumptions as in Proposition D.2, and f satisfies Assumption 3.12. If we have two chains at coarser and finer discretization levels l and $l+1$ using stepsizes h_l and $h_{l+1} = \frac{h_l}{2}$ satisfying (D.1) with synchronously coupled Brownian motions $(z_k)_{k \in \mathbb{N}}$ and $(z'_k)_{k \in \mathbb{N}}$, such that $z_0 \sim \pi_0$ and $z'_0 \sim \pi'_0$, then we have*

$$\begin{aligned} \text{Var}(f(z'_k) - f(z_k)) &\leq \mathbb{E} \left[(f(z'_k) - f(z_k))^2 \right] \leq \mathbb{E} \|z'_k - z_k\|_{a, b}^2 \\ &\leq \left(\exp \left(-\frac{mkh_l}{8\gamma} \right) (\|z'_0 - z_0\|_{L^2, a, b} + \mathcal{W}_{2, a, b}(\pi_0, \pi) + \mathcal{W}_{2, a, b}(\pi'_0, \pi)) \right. \\ &\quad + (1 - R(h_l))^k \mathcal{W}_{2, a, b}(\pi_0, \pi) + (1 - R(h_{l+1}))^{2k} \mathcal{W}_{2, a, b}(\pi'_0, \pi) \\ &\quad \left. + \sqrt{2} C_1 \left(\frac{h_{l+1}^{q+1/2}}{\sqrt{R(h_{l+1})}} + \frac{h_l^{q+1/2}}{\sqrt{R(h_l)}} \right) + 2C_2 \left(\frac{h_{l+1}^{q+1}}{R(h_{l+1})} + \frac{h_l^{q+1}}{R(h_l)} \right) \right)^2, \end{aligned}$$

where $R(h_i) = 1 - \sqrt{(1 - c(h_i))^2 + C_0^2 h_i^2}$ for $i = l, l+1$.

PROOF OF PROPOSITION D.3. Consider the following variance bound:

$$\text{Var}(f(z'_k) - f(z_k)) \leq \mathbb{E} \left[(f(z'_k) - f(z_k))^2 \right] \leq \mathbb{E} \|z'_k - z_k\|_{a, b}^2.$$

Let $\tilde{Z}_0 \sim \pi$ be such that $\|\tilde{Z}_0 - z_0\|_{L^2, a, b} = \mathcal{W}_{2, a, b}(\pi_0, \pi)$, and $\tilde{Z}'_0 \sim \pi$ be such that $\|\tilde{Z}'_0 - z'_0\|_{L^2, a, b} = \mathcal{W}_{2, a, b}(\pi'_0, \pi)$ (the existence of optimal couplings was shown in Theorem 4.1 of [94]). We use the estimate

$$\begin{aligned} \sqrt{\mathbb{E} \|z'_k - z_k\|_{a, b}^2} &= \|z'_k - z_k\|_{L^2, a, b} \\ &\leq \left\| z_k - \phi \left(\tilde{Z}_0, kh_l, (W_s)_{s=0}^{kh_l} \right) \right\|_{L^2, a, b} \\ &\quad + \left\| \phi \left(\tilde{Z}_0, kh_l, (W_s)_{s=0}^{kh_l} \right) - \phi \left(\tilde{Z}'_0, kh_l, (W_s)_{s=0}^{kh_l} \right) \right\|_{L^2, a, b} \\ &\quad + \left\| z'_k - \phi \left(\tilde{Z}'_0, kh_l, (W_s)_{s=0}^{kh_l} \right) \right\|_{L^2, a, b} \\ &=: \text{(I)} + \text{(II)} + \text{(III)}. \end{aligned}$$

We split this into two global error terms (I) and (III) and a contraction term (II). We estimate the second term by Corollary C.8 as

$$\begin{aligned} \text{(II)} &\leq \exp\left(-\frac{mkh_l}{8\gamma}\right) \|\tilde{Z}'_0 - \tilde{Z}_0\|_{L^2,a,b} \\ &\leq \exp\left(-\frac{mkh_l}{8\gamma}\right) (\|z'_0 - z_0\|_{L^2,a,b} + \mathcal{W}_{2,a,b}(\pi_0, \pi) + \mathcal{W}_{2,a,b}(\pi'_0, \pi)). \end{aligned}$$

By Proposition D.2, we have

$$\text{(I)} \leq (1 - R(h_l))^k \mathcal{W}_{2,a,b}(\pi_0, \pi) + \sqrt{2}C_1 \frac{2h_l^{q+1/2}}{\sqrt{R(h_l)}} + \frac{2C_2 h_l^{q+1}}{R(h_l)}.$$

The same argument can be applied to (III) to obtain

$$\text{(III)} \leq (1 - R(h_{l+1}))^{2k} \mathcal{W}_{2,a,b}(\pi'_0, \pi) + \sqrt{2}C_1 \frac{2h_{l+1}^{q+1/2}}{\sqrt{R(h_{l+1})}} + \frac{2C_2 h_{l+1}^{q+1}}{R(h_{l+1})}.$$

Combining these we get the required result. \square

Below are a number of useful remarks to highlight from the above theorem.

REMARK D.4. *The local error, which arises from [86] is demonstrated through the bound on $\alpha_h + \beta_h$ from Assumption D.1. This indicates there is an order of local strong order $q + 1/2$. However, when we go to the global strong order, the order is only reduced by $1/2$ as it is order q . As stated in [86], this is similar to the Euler–Maruyama scheme with local strong order $3/2$, but global strong order 1 [69][Theorem 1.1].*

REMARK D.5. *Proposition D.3 holds for $q = 2$ for the UBU scheme; [86] showed that the assumptions are true. For the UBU scheme we have for $\gamma^2 \geq M$ and $h < \frac{1}{2\gamma}$ that $C_2 \leq \sqrt{d} \left(\frac{7}{10}\gamma^2 + \frac{M_1^s}{10\sqrt{M}} \right)$, $C_1 = \frac{\sqrt{6dM\gamma}}{24}$ and $C_0 \leq 4\sqrt{2M}$. These constants can be computed by following [86][Section 7.6] where all computations are done with arbitrary γ , the constant c we consider to be set to 1 in their estimates. Constants C_1 and C_2 are estimated in the second and third step, whilst C_0 is estimated in the fourth step and fifth step. We remark that there is a missing term and a stronger assumption is needed in [86][Section 7.6, fifth step] which has been corrected in [77]. The additional term can be treated by the same argument as in the fourth step to arrive at the C_0 bound.*

COROLLARY D.6. *Suppose that Assumptions 3.6, 3.7, 3.10, and 3.13 hold, $\gamma \geq \sqrt{8M}$ and*

$$\text{(D.4)} \quad h_0 \leq \frac{1}{\gamma} \cdot \frac{m}{264M}.$$

Assume that the burn-in periods $B \geq \frac{16\log(4)\gamma}{mh_0}$, $B_0 \geq \frac{16\gamma}{mh_0} \log\left(\frac{c_{\mu_0}+1}{\sqrt{M\gamma}h_0^2}\right)$. Then for every $l \geq 0$, $1 \leq k \leq K$, the UBUBU samples satisfy

$$\begin{aligned} \text{Var}\left(f(z_k^{(l,l+1)}) - f(z_k^{(l,l+1)})\right) &\leq \mathbb{E}\left[\left(f(z_k^{(l,l+1)}) - f(z_k^{(l,l+1)})\right)^2\right] \leq \mathbb{E}\|z_k^{(l,l+1)} - z_k^{(l,l+1)}\|_{a,b}^2 \\ &\leq Cd \left(\left(\gamma^4 + \frac{(M_1^s)^2}{M} \right) \left(\frac{\gamma}{m} \right)^2 + \frac{M\gamma^2}{m} \right) h_l^4. \end{aligned}$$

PROOF OF COROLLARY D.6. We have $(B_0 + Bl)2^l$ burn-in steps at level l , and $(B_0 + B(l+1))2^{l+1}$ burn-in steps at level $l+1$. Let $\delta_* = \delta_{x^*} \times \delta_{0_d}$ be a distribution on Λ that fixes $x = x^*$ and $v = 0_d$. Using the assumptions, we have

$$\begin{aligned} R(h_i) &= 1 - \sqrt{(1 - c(h_i))^2 + C_0^2 h_i^2} = 1 - \sqrt{\left(1 - \frac{mh_i}{8\gamma}\right)^2 + C_0^2 h_i^2} \\ &= 1 - \sqrt{1 - \frac{mh_i}{4\gamma} + \left(\left(\frac{m}{8\gamma}\right)^2 + C_0^2\right) h_i^2} \geq 1 - \sqrt{1 - \frac{mh_i}{8\gamma}} \geq \frac{mh_i}{16\gamma}, \\ \mathcal{W}_{2,a,b}(\pi_0, \pi) &= \mathcal{W}_{2,a,b}(\mu_0, \pi) \leq c_{\mu_0} \sqrt{\frac{d}{m}}, \\ \mathcal{W}_{2,a,b}(\pi'_0, \pi) &\leq \mathcal{W}_{2,a,b}(\mu_0 R_{l+1}^B, \pi_{h_{l+1}}) + \mathcal{W}_{2,a,b}(\pi_{h_{l+1}}, \pi), \\ &\leq \mathcal{W}_{2,a,b}(\mu_0, \pi) + 2\mathcal{W}_{2,a,b}(\pi_{h_{l+1}}, \pi) \leq c_{\mu_0} \sqrt{\frac{d}{m}} + 2\sqrt{2}C_1 \frac{h_{l+1}^{q+1/2}}{\sqrt{R(h_{l+1})}} + \frac{4C_2 h_{l+1}^{q+1}}{R(h_{l+1})}, \end{aligned}$$

and

$$\begin{aligned} \|z'_0 - z_0\|_{L^2,a,b} &\leq \mathcal{W}_{2,a,b}(\pi'_0, \delta_*) + \mathcal{W}_{2,a,b}(\pi_0, \delta_*) \\ &\leq \mathcal{W}_{2,a,b}(\pi'_0, \pi) + \mathcal{W}_{2,a,b}(\mu_0, \pi) + 2\mathcal{W}_{2,a,b}(\pi, \delta_*) \\ &\leq (3c_{\mu_0} + 3) \sqrt{\frac{d}{m}} + 2\sqrt{2}C_1 \frac{h_{l+1}^{q+1/2}}{\sqrt{R(h_{l+1})}} + \frac{4C_2 h_{l+1}^{q+1}}{R(h_{l+1})}. \end{aligned}$$

It is easy to check that (D.4) together with $C_0 \leq 4\sqrt{2M}$ implies that the condition (D.1) of Proposition D.3 is satisfied, and we have

$$\begin{aligned} \text{Var} \left(f(z'_k{}^{(l,l+1)}) - f(z_k{}^{(l,l+1)}) \right) &\leq \mathbb{E} \left[\left(f(z'_k{}^{(l,l+1)}) - f(z_k{}^{(l,l+1)}) \right)^2 \right] \\ &\leq \left(\exp \left(-\frac{m(B_0 + lB)h_0}{8\gamma} \right) (\|z'_0 - z_0\|_{L^2,a,b} + \mathcal{W}_{2,a,b}(\pi_0, \pi) + \mathcal{W}_{2,a,b}(\pi'_0, \pi)) \right. \\ &\quad \left. + \exp \left(-\frac{m(B_0 + lB)h_0}{16\gamma} \right) (\mathcal{W}_{2,a,b}(\pi_0, \pi) + \mathcal{W}_{2,a,b}(\pi'_0, \pi)) \right. \\ &\quad \left. + \sqrt{2}C_1 \left(\frac{h_{l+1}^{q+1/2}}{\sqrt{R(h_{l+1})}} + \frac{h_l^{q+1/2}}{\sqrt{R(h_l)}} \right) + 2C_2 \left(\frac{h_{l+1}^{q+1}}{R(h_{l+1})} + \frac{h_l^{q+1}}{R(h_l)} \right) \right)^2 \\ &\leq \left(\exp \left(-\frac{m(B_0 + lB)h_0}{16\gamma} \right) (7c_{\mu_0} + 3) \sqrt{\frac{d}{m}} + 10\sqrt{2}C_1 \left(\frac{h_l^{5/2}}{\sqrt{\frac{mh_l}{16\gamma}}} \right) + 20C_2 \left(\frac{h_l^3}{\frac{mh_l}{16\gamma}} \right) \right)^2 \end{aligned}$$

using the assumptions on B_0 and B

$$\leq C \left(C_1^2 \frac{\gamma}{m} + C_2^2 \left(\frac{\gamma}{m} \right)^2 \right) h_l^4 \leq Cd \left(\left(\gamma^4 + \frac{(M_1^s)^2}{M} \right) \left(\frac{\gamma}{m} \right)^2 + \frac{M\gamma^2}{m} \right) h_l^4.$$

□

PROPOSITION D.7. Suppose that the assumptions of Proposition C.6 hold for $h = h_l$. Let $R_{l,l+1} = (P_{h_l, h_{l+1}})^{2^l}$ be the Markov kernel defined in Section 3.1 for two synchronously

coupled UBU chains at discretization levels $l, l+1$. This chain is moving on state space Λ^2 . Let $\bar{z}_1, \dots, \bar{z}_K$ be a Markov chain with kernel $R_{l, l+1}$. Let $F : \Lambda^2 \rightarrow \mathbb{R}$ be 1-Lipschitz in norm $\|\cdot\|_{a,b}$ on Λ^2 , defined as $\|z_1, z_2\|_{a,b}^2 = \|z_1\|_{a,b}^2 + \|z_2\|_{a,b}^2$. Then we have

$$\begin{aligned} \text{Var} \left(\frac{\sum_{i=1}^K F(\bar{z}_i)}{K} \right) &\leq \frac{2}{K^2} \sum_{i=1}^K \sum_{k=0}^{K-i} \min \left(\frac{\text{Var}(F(\bar{z}_i)) + \text{Var}(F(\bar{z}_{i+k}))}{2}, \right. \\ &\quad \left. \sqrt{\text{Var}(F(\bar{z}_i)) \mathbb{E} \left[\|\bar{z}_i - \mathbb{E}\bar{z}_i\|_{a,b}^2 \right]} \cdot \exp \left(-\frac{mh_0}{8\gamma} \cdot k \right) \right), \end{aligned}$$

PROOF. We need to bound

$$\text{Cov}(F(\bar{z}_i), F(\bar{z}_{i+k})) = \mathbb{E}[(F(\bar{z}_i) - \mathbb{E}(F(\bar{z}_i)))(F(\bar{z}_{i+k}) - \mathbb{E}(F(\bar{z}_{i+k})))].$$

Let \tilde{z}_i be an independent identically distributed copy of \bar{z}_i . For $0 \leq l \leq K - i - 1$, and assume that conditioned on $\tilde{z}_{i:l+i}$ and $\bar{z}_{1:i+l}$, $\tilde{z}_{i+l+1} \sim P(\tilde{z}_{i+l}, \cdot)$, and $(\bar{z}_{i+l+1}, \tilde{z}_{i+l+1})$ are synchronously coupled, i.e. \tilde{z}_{i+l+1} is defined based on (3.15) using the same Gaussian variables that were used to move from \bar{z}_{i+l} to \bar{z}_{i+l+1} . Since we have also used synchronous couplings in the proof of Proposition C.6, it follows from Proposition C.6 that

$$\begin{aligned} &\mathbb{E} \left(\|\tilde{z}_{i+l+1} - \bar{z}_{i+l+1}\|_{a,b}^2 \mid \bar{z}_{i:i+l}, \tilde{z}_{i:i+l} \right) \\ &\leq \max \left(\left(1 - \frac{mh_l}{8\gamma} \right)^{2 \cdot 2^l}, \left(1 - \frac{mh_{l+1}}{8\gamma} \right)^{2 \cdot 2^{l+1}} \right) \|\tilde{z}_{i+l} - \bar{z}_{i+l}\|_{a,b}^2 \end{aligned}$$

using that $1 - x \leq \exp(-x)$ for $x \geq 0$,

$$\leq \exp \left(-\frac{mh_0}{4\gamma} \right) \|\tilde{z}_{i+l} - \bar{z}_{i+l}\|_{a,b}^2.$$

By using this bound recursively, we have

$$\mathbb{E} \left(\|\tilde{z}_{i+k} - \bar{z}_{i+k}\|_{a,b}^2 \mid \bar{z}_i, \tilde{z}_i \right) \leq \exp \left(-\frac{mh_0}{4\gamma} \cdot k \right) \|\bar{z}_i - \tilde{z}_i\|_{a,b}^2.$$

Since \tilde{z}_i is independent of \bar{z}_i , and $\tilde{z}_{i+1}, \dots, \tilde{z}_{i+k}$ was constructed using \tilde{z}_i and Gaussians that are independent of \bar{z}_i (synchronous coupling with $\bar{z}_{i+1}, \dots, \bar{z}_{i+k}$), it follows that \tilde{z}_{i+k} is still independent of \bar{z}_i . Using this and the 1-Lipschitz property of F , we have

$$\begin{aligned} \text{Cov}(F(\bar{z}_i), F(\bar{z}_{i+k})) &= \mathbb{E}[(F(\bar{z}_i) - \mathbb{E}(F(\bar{z}_i)))(F(\bar{z}_{i+k}) - \mathbb{E}(F(\bar{z}_{i+k})))] \\ &= \mathbb{E}[(F(\bar{z}_i) - \mathbb{E}(F(\bar{z}_i)))(F(\bar{z}_{i+k}) - F(\tilde{z}_{i+k}))] \\ &= \mathbb{E}[(F(\bar{z}_i) - \mathbb{E}(F(\bar{z}_i)))\mathbb{E}(F(\bar{z}_{i+k}) - F(\tilde{z}_{i+k}) \mid \bar{z}_i, \tilde{z}_i)] \\ &\leq \sqrt{\text{Var}(F(\bar{z}_i)) \mathbb{E} \left[\|\bar{z}_{i+k} - \tilde{z}_{i+k}\|_{a,b}^2 \right]} \\ &\leq \exp \left(-\frac{mh_0}{8\gamma} \cdot k \right) \sqrt{\text{Var}(F(\bar{z}_i)) \cdot \mathbb{E} \left[\|\bar{z}_i - \tilde{z}_i\|_{a,b}^2 \right]} \\ &= \exp \left(-\frac{mh_0}{8\gamma} \cdot k \right) \sqrt{2\text{Var}(F(\bar{z}_i)) \cdot \mathbb{E} \left[\|\bar{z}_i - \mathbb{E}\bar{z}_i\|_{a,b}^2 \right]}, \end{aligned}$$

and the claim follows by summation. \square

PROPOSITION D.8. *Under the same assumptions as in Corollary D.6, the UBUBU samples satisfy that*

$$\text{Var}(D_{l,l+1}) \leq \frac{1}{K} C(\gamma, m, M, M_1^s) dh_l^4 \left(C(\gamma, m, M, M_1^s) - 2\log(h_0) + \log(4)l + \frac{4\gamma}{mh_0} \right).$$

PROOF. Note that the function $F(z_1, z_2) = f(z_1) - f(z_2)$ is 1-Lipschitz with respect to $\|(z_1, z_2)\|_{a,b} = \|z_1\|_{a,b} + \|z_2\|_{a,b}$. Let $\bar{z}_i = (z_i^{(l,l+1)}, z_i'^{(l,l+1)})$, then by Proposition D.7, we have that

$$\begin{aligned} \text{Var}(D_{l,l+1}) &\leq \frac{1}{K^2} \sum_{i=1}^K \sum_{k=0}^{K-i} \min \left(\frac{\text{Var}(F(\bar{z}_i)) + \text{Var}(F(\bar{z}_{i+k}))}{2}, \right. \\ &\quad \left. \sqrt{\text{Var}(F(\bar{z}_i)) \mathbb{E} \left[\|\bar{z}_i - \mathbb{E}\bar{z}_i\|_{a,b}^2 \right]} \cdot \exp \left(-\frac{mh_0}{8\gamma} \cdot k \right) \right). \end{aligned}$$

By a similar argument as in the proof of Corollary D.6, using our assumptions on B and B_0 , we can show that

$$\left(\mathbb{E} \left[\|\bar{z}_i - \mathbb{E}\bar{z}_i\|_{a,b}^2 \right] \right)^{1/2} \leq \left(\mathbb{E} \left[\|\bar{z}_i - (x^*, 0_d, x^*, 0_d)\|_{a,b}^2 \right] \right)^{1/2} \leq C \sqrt{\frac{d}{m}},$$

and by Proposition D.3, we have

$$\text{Var}(F(\bar{z}_i)) \leq C(\gamma, m, M, M_1^s) dh_l^4.$$

Let

$$k^*(l) := \max \left(\log \left(C \sqrt{\frac{1}{m}} \right) - \frac{1}{2} \log(C(\gamma, m, M, M_1) h_l^4), 0 \right),$$

then for $k \geq \lceil k^*(l) \rceil$, we have

$$\begin{aligned} &\sqrt{\text{Var}(F(\bar{z}_i)) \mathbb{E} \left[\|\bar{z}_i - \mathbb{E}\bar{z}_i\|_{a,b}^2 \right]} \cdot \exp \left(-\frac{mh_0}{8\gamma} \cdot k \right) \\ &\leq C(\gamma, m, M, M_1^s) dh_l^4 \exp \left(-\frac{mh_0}{8\gamma} \cdot (k - \lceil k^*(l) \rceil) \right). \end{aligned}$$

It is clear that $\lceil k^*(l) \rceil \leq C(\gamma, m, M, M_1^s) - 2\log(h_0) + \log(4)l$, and after some rearrangement, we have

$$\text{Var}(D_{l,l+1}) \leq \frac{1}{K} C(\gamma, m, M, M_1^s) dh_l^4 \left(C(\gamma, m, M, M_1^s) - 2\log(h_0) + \log(4)l + \frac{4\gamma}{mh_0} \right). \quad \square$$

D.2. Variance bound of D_0 .

PROPOSITION D.9. *Consider an m -strongly convex M - ∇ Lipschitz potential U and let P_h be the transition kernel of UBU with stepsize h . Suppose that $f : \Omega \rightarrow \mathbb{R}$ only depends on x and is a 1-Lipschitz function. Suppose that $\gamma \geq \sqrt{8M}$, and $h < \frac{1}{2\gamma}$. Let μ_0 be a distribution on Λ , and the Markov chain $z_{-B_0}^{(0)} \sim \mu_0$, $z_{-B_0+1}^{(0)} \sim P_h(z_{-B_0}^{(0)}, \cdot)$, \dots , $z_K \sim P_h(z_{K-1}^{(0)}, \cdot)$. Then D_0 as defined in (3.11) satisfies that*

$$\text{Var}(D_0) \leq \frac{C}{c(h)K} \left(1 + \frac{1}{c(h)K} \right) \left(\frac{1}{\gamma} + \frac{\gamma}{M} \right) \frac{h}{c(h)} + \frac{(1-c(h))^{2(B_0+1)}}{2c(h)^2 K^2} \sigma_{\mu_0}^2,$$

where

$$c(h) = \frac{mh}{8\gamma}, \quad \sigma_{\mu_0}^2 = \int \int \|w - \tilde{w}\|_{a,b}^2 d\mu_0(w) d\mu_0(\tilde{w}),$$

for some absolute constant C .

PROOF. The bound is based on Theorem 2 of [51]. We need to control the following quantities for every $z \in \Lambda$:

$$(D.5) \quad \sigma(z)^2 := \frac{1}{2} \int \int \|w - \tilde{w}\|_{a,b}^2 P_h(z, dw) P_h(z, d\tilde{w}),$$

$$(D.6) \quad n(z) := \inf_{g: \Lambda \rightarrow \mathbb{R}, \|g\|_{a,b, Lip} \leq 1} \frac{\int \int \|w - \tilde{w}\|_{a,b}^2 P_h(z, dw) P_h(z, d\tilde{w})}{\int \int (g(w) - g(\tilde{w}))^2 P_h(z, dw) P_h(z, d\tilde{w})}.$$

Here we choose $a = \frac{1}{M}$, and $b = \frac{1}{\gamma}$ as in Proposition C.6. To control $\sigma^2(z)$, let us define two independent identically distributed random variables $w(z) \sim P_h(z, \cdot)$ and $\tilde{w}(z) \sim P_h(z, \cdot)$. Using the definition of UBU in (2.5), we have

$$\begin{aligned} \sigma(z)^2 &= \frac{1}{2} \mathbb{E}(\|w(z) - \tilde{w}(z)\|_{a,b}^2) \\ &= \frac{1}{2} \mathbb{E} \left(\left\| \mathcal{UBU} \left(z, h, \xi^{(1)}, \xi^{(2)}, \xi^{(3)}, \xi^{(4)} \right) - \mathcal{UBU} \left(z, h, \tilde{\xi}^{(1)}, \tilde{\xi}^{(2)}, \tilde{\xi}^{(3)}, \tilde{\xi}^{(4)} \right) \right\|_{a,b}^2 \right) \\ &\leq \mathbb{E} \left(\left\| \mathcal{U} \left(\mathcal{B} \left(\mathcal{U} \left(z, h/2, \xi^{(1)}, \xi^{(2)} \right), h \right), h/2, \xi^{(3)}, \xi^{(4)} \right) \right. \right. \\ &\quad \left. \left. - \mathcal{U} \left(\mathcal{B} \left(\mathcal{U} \left(z, h/2, \xi^{(1)}, \xi^{(2)} \right), h \right), h/2, \tilde{\xi}^{(3)}, \tilde{\xi}^{(4)} \right) \right\|_{a,b}^2 \right) \\ &\quad + \mathbb{E} \left(\left\| \mathcal{U} \left(\mathcal{B} \left(\mathcal{U} \left(z, h/2, \xi^{(1)}, \xi^{(2)} \right), h \right), h/2, \tilde{\xi}^{(3)}, \tilde{\xi}^{(4)} \right) \right. \right. \\ &\quad \left. \left. - \mathcal{U} \left(\mathcal{B} \left(\mathcal{U} \left(z, h/2, \tilde{\xi}^{(1)}, \tilde{\xi}^{(2)} \right), h \right), h/2, \tilde{\xi}^{(3)}, \tilde{\xi}^{(4)} \right) \right\|_{a,b}^2 \right). \end{aligned}$$

Recalling the definitions of \mathcal{U} and \mathcal{B} from equations (2.2-2.3), we have

$$\begin{aligned} \mathcal{B}(x, v, h) &= (x, v - h\nabla U(x)), \\ \mathcal{U}(x, v, h/2, \xi^{(1)}, \xi^{(2)}) &= \left(x + \frac{1-\eta}{\gamma} v + \sqrt{\frac{2}{\gamma}} \left(\mathcal{Z}^{(1)}(h/2, \xi^{(1)}) - \mathcal{Z}^{(2)}(h/2, \xi^{(1)}, \xi^{(2)}) \right), \right. \\ &\quad \left. \eta v + \sqrt{2\gamma} \mathcal{Z}^{(2)}(h/2, \xi^{(1)}, \xi^{(2)}) \right), \\ \mathcal{Z}^{(1)}(h/2, \xi^{(1)}) &= \sqrt{\frac{h}{2}} \xi^{(1)}, \\ \mathcal{Z}^{(2)}(h/2, \xi^{(1)}, \xi^{(2)}) &= \sqrt{\frac{1-\eta^2}{2\gamma}} \left(\sqrt{\frac{1-\eta}{1+\eta} \cdot \frac{4}{\gamma h}} \xi^{(1)} + \sqrt{1 - \frac{1-\eta}{1+\eta} \cdot \frac{4}{\gamma h}} \xi^{(2)} \right). \end{aligned}$$

First,

$$\begin{aligned} & \mathbb{E} \left(\left\| \mathcal{U} \left(\mathcal{B} \left(\mathcal{U} \left(z, h/2, \xi^{(1)}, \xi^{(2)} \right), h \right), h/2, \xi^{(3)}, \xi^{(4)} \right) \right. \right. \\ & \quad \left. \left. - \mathcal{U} \left(\mathcal{B} \left(\mathcal{U} \left(z, h/2, \xi^{(1)}, \xi^{(2)} \right), h \right), h/2, \tilde{\xi}^{(3)}, \tilde{\xi}^{(4)} \right) \right\|_{a,b}^2 \right) \\ &= \mathbb{E} \left(\left\| \left(\sqrt{\frac{2}{\gamma}} \left(\mathcal{Z}^{(1)} \left(\frac{h}{2}, \xi^{(1)} \right) - \mathcal{Z}^{(2)} \left(\frac{h}{2}, \xi^{(1)}, \xi^{(2)} \right) \right) - \sqrt{\frac{2}{\gamma}} \left(\mathcal{Z}^{(1)} \left(\tilde{\xi}^{(1)} \right) - \mathcal{Z}^{(2)} \left(\frac{h}{2}, \xi^{(1)}, \tilde{\xi}^{(2)} \right) \right) \right. \right. \\ & \quad \left. \left. \sqrt{2\gamma} \mathcal{Z}^{(2)} \left(\frac{h}{2}, \xi^{(1)}, \xi^{(2)} \right) - \sqrt{2\gamma} \mathcal{Z}^{(2)} \left(\frac{h}{2}, \tilde{\xi}^{(1)}, \tilde{\xi}^{(2)} \right) \right\|_{a,b}^2 \right) \end{aligned}$$

using (C.1), and the fact that $a = \frac{1}{M}$

$$\begin{aligned} & \leq \frac{6}{\gamma} \mathbb{E} \left(\left\| \mathcal{Z}^{(1)} \left(\frac{h}{2}, \xi^{(1)} \right) - \mathcal{Z}^{(2)} \left(\frac{h}{2}, \xi^{(1)}, \tilde{\xi}^{(2)} \right) \right\|^2 \right) + \frac{6\gamma}{M} \mathbb{E} \left(\left\| \mathcal{Z}^{(2)} \left(\frac{h}{2}, \xi^{(1)}, \tilde{\xi}^{(2)} \right) \right\|^2 \right) \\ & \leq \left(\frac{3}{\gamma} + \frac{3\gamma}{M} \right) dh. \end{aligned}$$

Second, using the assumptions $\gamma \geq \sqrt{8M}$ and $h \leq \frac{1}{\sqrt{M}}$, for any x, v, x', v' ,

$$\begin{aligned} & \left\| \mathcal{U} \left(x, v, h/2, \tilde{\xi}^{(3)}, \tilde{\xi}^{(4)} \right) - \mathcal{U} \left(x', v', h/2, \tilde{\xi}^{(3)}, \tilde{\xi}^{(4)} \right) \right\|_{a,b}^2 \\ & \leq \frac{3}{2} \left\| \mathcal{U} \left(x, v, h/2, \tilde{\xi}^{(3)}, \tilde{\xi}^{(4)} \right) - \mathcal{U} \left(x', v', h/2, \tilde{\xi}^{(3)}, \tilde{\xi}^{(4)} \right) \right\|_{a,0}^2 \\ & \leq \frac{3}{2} \left(\frac{1}{M} \exp(-\gamma h) \|v - v'\|^2 + 2\|x - x'\|^2 + \frac{2(1 - \exp(-\gamma h/2))^2}{\gamma^2} \|v - v'\|^2 \right) \\ \text{(D.7)} \quad & \leq 3\|x - x'\|^2 + \frac{3}{2} \frac{1}{M} \|v - v'\|^2 \leq 6\|(x - x', v - v')\|_{a,b}^2, \end{aligned}$$

$$\begin{aligned} & \|\mathcal{B}(x, v, h) - \mathcal{B}(x', v', h)\|_{a,b}^2 \leq \frac{3}{2} \|(x - x', v - v' + h\nabla U(x') - h\nabla U(x))\|_{0,a}^2 \\ & \leq \frac{3}{2} \|x - x'\|^2 + \frac{3}{M} \|v - v'\|^2 + \frac{3h^2}{M} \|\nabla U(x') - \nabla U(x)\|^2 \end{aligned}$$

$$\text{(D.8)} \quad \leq \left(\frac{3}{2} + 3h^2M \right) \|x - x'\|^2 + \frac{3}{M} \|v - v'\|^2 \leq 6\|(x - x', v - v')\|_{a,b}^2$$

hence

$$\begin{aligned} & \mathbb{E} \left(\left\| \mathcal{U} \left(\mathcal{B} \left(\mathcal{U} \left(z, h/2, \xi^{(1)}, \xi^{(2)} \right), h \right), h/2, \tilde{\xi}^{(3)}, \tilde{\xi}^{(4)} \right) \right. \right. \\ & \quad \left. \left. - \mathcal{U} \left(\mathcal{B} \left(\mathcal{U} \left(z, h/2, \tilde{\xi}^{(1)}, \tilde{\xi}^{(2)} \right), h \right), h/2, \tilde{\xi}^{(3)}, \tilde{\xi}^{(4)} \right) \right\|_{a,b}^2 \right) \\ & \leq 36 \mathbb{E} \left(\left\| \mathcal{U} \left(z, h/2, \xi^{(1)}, \xi^{(2)} \right) - \mathcal{U} \left(z, h/2, \tilde{\xi}^{(1)}, \tilde{\xi}^{(2)} \right) \right\|_{a,b}^2 \right) \end{aligned}$$

$$\begin{aligned}
&\leq 36\mathbb{E}\left(\left\|\left(\sqrt{\frac{2}{\gamma}}\left(\mathcal{Z}^{(1)}\left(\frac{h}{2}, \xi^{(1)}\right) - \mathcal{Z}^{(2)}\left(\frac{h}{2}, \xi^{(1)}, \xi^{(2)}\right)\right) - \sqrt{\frac{2}{\gamma}}\left(\mathcal{Z}^{(1)}\left(\tilde{\xi}^{(1)}\right) - \mathcal{Z}^{(2)}\left(\frac{h}{2}, \xi^{(1)}, \tilde{\xi}^{(2)}\right)\right)\right\|_{a,b}\right)^2 \\
&\quad \left\|\sqrt{2\gamma}\mathcal{Z}^{(2)}\left(\frac{h}{2}, \xi^{(1)}, \xi^{(2)}\right) - \sqrt{2\gamma}\mathcal{Z}^{(2)}\left(\frac{h}{2}, \tilde{\xi}^{(1)}, \tilde{\xi}^{(2)}\right)\right\|_{a,b}^2\right) \leq 36\left(\frac{3}{\gamma} + \frac{3\gamma}{M}\right)dh,
\end{aligned}$$

using the same argument as for the previous term. Hence by summing up the above bounds, we have

$$(D.9) \quad \sigma(z)^2 \leq 37\left(\frac{3}{\gamma} + \frac{3\gamma}{M}\right)dh.$$

Now, we will lower bound $n(z)$ as defined in (D.6). By (C.1), we have

$$\mathbb{E}(\|w(z) - \tilde{w}(z)\|_{a,b}^2) \geq \frac{1}{2}\mathbb{E}(\|w(z) - \tilde{w}(z)\|_{a,0}^2)$$

(D.10)

$$= \mathbb{E}\left(\left\|\mathcal{U}\left(\mathcal{B}\left(\mathcal{U}\left(z, h/2, \xi^{(1)}, \xi^{(2)}\right), h\right), h/2, \xi^{(3)}, \xi^{(4)}\right)\right\|_{a,0}^2\right)$$

by Jensen's inequality

$$\geq \mathbb{E}\left(\left\|\left(\sqrt{\frac{2}{\gamma}}\left(\mathcal{Z}^{(1)}\left(\frac{h}{2}, \xi^{(1)}\right) - \mathcal{Z}^{(2)}\left(\frac{h}{2}, \xi^{(1)}, \xi^{(2)}\right)\right), \sqrt{2\gamma}\mathcal{Z}^{(2)}\left(\frac{h}{2}, \xi^{(1)}, \xi^{(2)}\right)\right)\right\|_{a,0}^2\right)$$

(D.11)

$$\geq \frac{\gamma}{M}dh.$$

For the denominator, we have

$$\begin{aligned}
&\int \int (g(w) - g(\tilde{w}))^2 P_h(z, dw) P_h(z, d\tilde{w}) = 2 \cdot \text{Var}_{w \sim P_h(z, \cdot)}(g(w)) \\
&= 2 \cdot \text{Var}\left(g\left(\mathcal{U}\left(\mathcal{B}\left(\mathcal{U}\left(z, h/2, \xi^{(1)}, \xi^{(2)}\right), h\right), h/2, \xi^{(3)}, \xi^{(4)}\right)\right)\right)
\end{aligned}$$

by the Efron-Stein inequality [17, 90]

$$\begin{aligned}
&\leq 2\mathbb{E}(\text{Var}_{\xi^{(1)}, \xi^{(2)}}\left(g\left(\mathcal{U}\left(\mathcal{B}\left(\mathcal{U}\left(z, h/2, \xi^{(1)}, \xi^{(2)}\right), h\right), h/2, \xi^{(3)}, \xi^{(4)}\right)\right)\right) \\
&\quad + 2\mathbb{E}(\text{Var}_{\xi^{(3)}, \xi^{(4)}}\left(g\left(\mathcal{U}\left(\mathcal{B}\left(\mathcal{U}\left(z, h/2, \xi^{(1)}, \xi^{(2)}\right), h\right), h/2, \xi^{(3)}, \xi^{(4)}\right)\right)\right),
\end{aligned}$$

where $\text{Var}_{\xi^{(1)}, \xi^{(2)}}(\cdot)$ means that we compute the conditional variance with respect to $\xi^{(3)}, \xi^{(4)}$ (so the $\xi^{(3)}, \xi^{(4)}$ are kept constant, and only the variance with respect to $\xi^{(1)}, \xi^{(2)}$ is considered). Let

$$\begin{aligned}
J_{\mathcal{U}}(h) &:= \frac{\partial \mathcal{U}(x, v, h, \xi^{(1)}, \xi^{(2)})}{\partial (\xi^{(1)}, \xi^{(2)})} \\
&= \left(\begin{array}{c} \left(\sqrt{\frac{2h}{\gamma}} - \frac{\sqrt{2(1-e^{-\gamma h})}}{\gamma^{3/2}\sqrt{h}}\right) I_d, \quad -\frac{(1-e^{-\gamma h})\sqrt{2}}{\sqrt{\gamma h}} I_d \\ -\frac{1}{\gamma}\sqrt{1-e^{-2\gamma h}} - \frac{2(1-e^{-\gamma h})^2}{\gamma h} I_d, \quad \sqrt{1-e^{-2\gamma h}} - \frac{2(1-e^{-\gamma h})^2}{\gamma h} I_d \end{array} \right), \\
\tilde{g}_h(z) &:= g\left(\mathcal{U}\left(\mathcal{B}(z, h), h/2, \xi^{(3)}, \xi^{(4)}\right)\right).
\end{aligned}$$

Using the assumption that g is 1-Lipschitz in (D.6), and the bounds (D.7-D.8), it follows that \tilde{g}_h is a 6-Lipschitz function in $\|\cdot\|_{a,b}$, and (C.1) implies that it is a 12-Lipschitz function in $\|\cdot\|_{a,0}$. Since the continuously differentiable Lipschitz functions are dense amongst Lipschitz functions (see [4]), we can assume without loss of generality that g and thus \tilde{g}_h are continuously differentiable. Note that

$$\begin{aligned}\tilde{g}_h(z) - \tilde{g}_h(z') &= \langle \nabla \tilde{g}_h(z), z - z' \rangle + o(\|z - z'\|_{a,0}) \\ &= \left\langle \begin{pmatrix} I_d & 0 \\ 0 & a^{-1/2} I_d \end{pmatrix} \nabla \tilde{g}_h(z), \begin{pmatrix} I_d & 0 \\ 0 & a^{1/2} I_d \end{pmatrix} (z - z') \right\rangle + o(\|z - z'\|_{a,0}).\end{aligned}$$

Using this, it is easy to show that the 12-Lipschitz property of \tilde{g}_h in $\|\cdot\|_{a,0}$ implies that $\|\nabla \tilde{g}_h(z)\|_{1/a,0} \leq 12$ for every $z \in \Lambda$. Hence, we obtain

$$\begin{aligned}& \left\| \frac{\partial}{\partial(\xi^{(1)}, \xi^{(2)})} \left(g \left(\mathcal{U} \left(\mathcal{B} \left(\mathcal{U} \left(z, h/2, \xi^{(1)}, \xi^{(2)} \right), h \right), h/2, \xi^{(3)}, \xi^{(4)} \right) \right) \right) \right\| \\ &= \left\| \frac{\partial}{\partial(\xi^{(1)}, \xi^{(2)})} \tilde{g}_h \left(\mathcal{U} \left(z, h/2, \xi^{(1)}, \xi^{(2)} \right) \right) \right\| = \left\| J_U(h/2) \nabla \tilde{g}_h \left(\mathcal{U} \left(z, h/2, \xi^{(1)}, \xi^{(2)} \right) \right) \right\| \\ &\leq 12 \sup_{w \in \Lambda: \|w\|_{1/a,0} \leq 1} \|J_U(h/2)w\| = 12 \sup_{w \in \Lambda: \|w\| \leq 1} \left\| J_U(h/2) \begin{pmatrix} I_d & 0_d \\ 0_d & \frac{1}{\sqrt{M}} I_d \end{pmatrix} w \right\| \\ &= 12 \left\| J_U(h/2) \begin{pmatrix} I_d & 0_d \\ 0_d & \frac{1}{\sqrt{M}} I_d \end{pmatrix} \right\| \\ &= 12 \left\| \left(\begin{array}{c} \left(\frac{\sqrt{h}}{\gamma} - \frac{2(1-e^{-\gamma h/2})}{\gamma^{3/2}\sqrt{h}} \right), \quad \frac{-2(1-e^{-\gamma h/2})}{\sqrt{M}\gamma h} \\ -\frac{1}{\gamma} \sqrt{1 - e^{-\gamma h} - \frac{4(1-e^{-\gamma h/2})^2}{\gamma h}}, \quad \frac{1}{\sqrt{M}} \sqrt{1 - e^{-\gamma h} - \frac{4(1-e^{-\gamma h/2})^2}{\gamma h}} \end{array} \right) \right\|,\end{aligned}$$

using the fact that $-(1 - e^{-x})^2 \leq -x^2 + x^3$ for $x \geq 0$, and that $\gamma h \leq 1$

$$\leq 12 \left(\sqrt{\frac{h}{\sqrt{M}}} + \frac{2\gamma h}{\sqrt{M}} \right).$$

From the Gaussian Poincaré inequality (see e.g. Theorem 3.20 of [17]), and the fact that $\xi^{(1)}, \xi^{(2)}$ are standard normal, it follows that

$$\begin{aligned}& 2\mathbb{E}(\text{Var}_{\xi^{(1)}, \xi^{(2)}} \left(g \left(\mathcal{U} \left(\mathcal{B} \left(\mathcal{U} \left(z, h/2, \xi^{(1)}, \xi^{(2)} \right), h \right), h/2, \xi^{(3)}, \xi^{(4)} \right) \right) \right)) \\ &\leq 2 \cdot 12^2 \left(\sqrt{\frac{h}{\sqrt{M}}} + \frac{2\gamma h}{\sqrt{M}} \right)^2 \leq 576 \left(\frac{h}{\sqrt{M}} + 4 \frac{\gamma^2 h^2}{M} \right).\end{aligned}$$

We can bound the second term similarly, since

$$\begin{aligned}& \left\| \frac{\partial}{\partial(\xi^{(3)}, \xi^{(4)})} \left(g \left(\mathcal{U} \left(\mathcal{B} \left(\mathcal{U} \left(z, h/2, \xi^{(1)}, \xi^{(2)} \right), h \right), h/2, \xi^{(3)}, \xi^{(4)} \right) \right) \right) \right\| \\ &= \left\| J_U(h/2) \nabla g \left(\mathcal{U} \left(\mathcal{B} \left(\mathcal{U} \left(z, h/2, \xi^{(1)}, \xi^{(2)} \right), h \right), h/2, \xi^{(3)}, \xi^{(4)} \right) \right) \right\| \\ &\leq \left\| J_U(h/2) \begin{pmatrix} I_d & 0_d \\ 0_d & \frac{1}{\sqrt{M}} I_d \end{pmatrix} \right\| \leq \sqrt{\frac{h}{\sqrt{M}}} + \frac{2\gamma h}{\sqrt{M}},\end{aligned}$$

and thus by the Gaussian Poincaré inequality,

$$\begin{aligned} & 2\mathbb{E}(\text{Var}_{\xi^{(3)}, \xi^{(4)}} \left(g \left(\mathcal{U} \left(\mathcal{B} \left(\mathcal{U} \left(z, h/2, \xi^{(1)}, \xi^{(2)} \right), h \right), h/2, \xi^{(3)}, \xi^{(4)} \right) \right) \right) \\ & \leq 2 \left(\sqrt{\frac{h}{\sqrt{M}}} + \frac{2\gamma h}{\sqrt{M}} \right)^2 \leq 4 \left(\frac{h}{\sqrt{M}} + 4\frac{\gamma^2 h^2}{M} \right). \end{aligned}$$

By adding these up, we obtain

$$\int \int (g(w) - g(\tilde{w}))^2 P_h(z, dw) P_h(z, d\tilde{w}) \leq 580 \left(\frac{h}{\sqrt{M}} + 4\frac{\gamma^2 h^2}{M} \right),$$

and hence by (D.6) and (D.11), we have

$$(D.12) \quad n(z) \geq \frac{\frac{\gamma}{M} dh}{580 \left(\frac{h}{\sqrt{M}} + 4\frac{\gamma^2 h^2}{M} \right)} \geq \frac{\frac{\gamma}{M} dh}{580 \cdot 5 \left(\frac{\gamma h}{M} \right)} \geq \frac{d}{2900}.$$

Combining this with (D.9), we have that

$$\sup_{z \in \Lambda} \frac{\sigma(z)^2}{n(z)} \leq \left(37 \left(\frac{3}{\gamma} + \frac{3\gamma}{M} \right) dh \right) \cdot \frac{2900}{d} \leq 321900 \left(\frac{1}{\gamma} + \frac{\gamma}{M} \right) h,$$

and the claim now follows by Theorem 2 of [51] and the bound on $\text{Var}[\mathbb{E}(\hat{\pi}(f))|X_0]$ on page 2427 of [51], using the fact that $\kappa \geq 1 - \sqrt{1 - \frac{mh}{4\gamma}} \geq \frac{mh}{8\gamma}$ by Proposition C.6. \square

D.3. Variance of $S(c_R)$.

THEOREM 3.15. *Suppose that Assumptions 3.6, 3.7, 3.10, 3.12, 3.13 hold, and in addition,*

$$\gamma \geq \sqrt{8M}, \quad h_0 \leq \frac{1}{\gamma} \cdot \frac{m}{264M}, \quad B \geq \frac{16 \log(4)\gamma}{mh_0}, \quad B_0 \geq \frac{16\gamma}{mh_0} \log \left(\frac{c_{\mu_0} + 1}{\sqrt{M}\gamma h_0^2} \right).$$

Suppose that $c_R \in [0, 1)$, and $2 < \phi_N < 16$. Then for any $N \geq 1$, the UBUBU estimator $S(c_R)$ has finite expected computational cost, $\mathbb{E}S(c_R) = \pi(f)$, and it has finite variance. Moreover, it satisfies a CLT as $N \rightarrow \infty$, and the asymptotic variance σ_S^2 defined in (3.8) can be bounded as

$$\sigma_S^2 \leq \frac{C(m, M, M_1^s, \gamma, c_N, \phi_N)}{Kh_0} \left(1 + \frac{1}{h_0 K} + dh_0^4 \right).$$

PROOF OF THEOREM 3.15. By Corollary D.6, and the fact that

$$\mathbb{E}(D_{l,l+1}^2) \leq \max_{1 \leq k \leq K} \mathbb{E} \left[\left(f(z_k^{(l,l+1)}) - f(z_k^{(l+1)}) \right)^2 \right],$$

it follows that under the assumptions of Corollary D.6, we have

$$\mathbb{E}(D_{l,l+1}^2) \leq Cd \left(\left(\gamma^4 + \frac{(M_1^s)^2}{M} \right) \left(\frac{\gamma}{m} \right)^2 + \frac{M\gamma^2}{m} \right) h_l^4 \leq V_D \phi_D^{-l},$$

for $V_D = Ch_0^4 d \left(\left(\gamma^4 + \frac{(M_1^s)^2}{M} \right) \left(\frac{\gamma}{m} \right)^2 + \frac{M\gamma^2}{m} \right)$ and $\phi_D = 16$. From Proposition D.9, and using the fact that $c(h_0) = \frac{h_0 m}{8\gamma}$, and our assumptions on B_0 , we have

$$\text{Var}(D_0) \leq \frac{C}{c(h_0)K} \left(1 + \frac{1}{c(h_0)K} \right) \left(\frac{1}{\gamma} + \frac{\gamma}{M} \right) \frac{h_0}{c(h_0)} + \frac{(1 - c(h_0))^{2(B_0+1)}}{2c(h_0)^2 K^2} \sigma_{\mu_0}^2$$

$$(D.13) \leq \frac{C}{K} \cdot \frac{1}{h_0} \left(\frac{8\gamma}{m} \right)^2 \left(\frac{1}{\gamma} + \frac{\gamma}{M} \right) \left(1 + \frac{8\gamma}{h_0 m} \cdot \frac{1}{K} \right).$$

The computational cost at each level satisfies the assumptions of Proposition 3.4, so if we fix $2 < \phi_N < 16$, all assumptions of this proposition are satisfied. Hence $S(c_R)$ is an unbiased estimator with finite variance and computational cost.

The claim about the asymptotic variance follows by using the bounds in (D.13) and in Proposition D.8, and adding up all terms according to (3.8). \square

PROPOSITION D.10. *Suppose that the assumptions of Proposition C.6 hold for $h = h_l$. Let $R_{l,l+1} = (P_{h_l, h_{l+1}})^2$ be the Markov kernel defined in Section 3.1 for two synchronously coupled UBU chains at discretization levels $l, l+1$. This chain is moving on state space Λ^2 . Let $\bar{z}_1, \dots, \bar{z}_K$ be a Markov chain with kernel $R_{l,l+1}$. Let $F : \Lambda^2 \rightarrow \mathbb{R}$ be of the form $F(z, z') = f(z) - f(z')$, where f is of the form (3.18). Suppose that the target π is a product distribution, satisfying the same conditions as in Proposition 3.18. Then we have*

$$\begin{aligned} \text{Var} \left(\frac{\sum_{i=1}^K F(\bar{z}_i)}{K} \right) &\leq \frac{2}{K^2} \sum_{i=1}^K \sum_{k=0}^{K-i} \min \left(\frac{\text{Var}(F(\bar{z}_i)) + \text{Var}(F(\bar{z}_{i+k}))}{2}, \right. \\ &\quad \left. \sqrt{4r \left(\sum_{s=1}^r \|w^{(s)}\|^2 \right)} \sqrt{\text{Var}(F(\bar{z}_i)) \max_{1 \leq j \leq d} \mathbb{E} \left[\|\bar{z}_{i,j} - \mathbb{E} \bar{z}_{i,j}\|_{a,b}^2 \right]} \cdot \exp \left(-\frac{mh_0}{8\gamma} \cdot k \right)} \right). \end{aligned}$$

PROOF. We proceed similarly to the proof of Proposition D.7.

$$\text{Cov}(F(\bar{z}_i), F(\bar{z}_{i+k})) = \mathbb{E}[(F(\bar{z}_i) - \mathbb{E}(F(\bar{z}_i)))(F(\bar{z}_{i+k}) - \mathbb{E}(F(\bar{z}_{i+k})))].$$

Let \tilde{z}_i be an independent identically distributed copy of \bar{z}_i , and define $(\bar{z}_{i:i+k}, \tilde{z}_{i:i+k})$ as synchronously coupled, in the same way as in the proof of Proposition D.7. It follows from applying Proposition C.6 on each coordinate, and using independence that for every coordinate $1 \leq j \leq d$,

$$\mathbb{E} \left(\|\tilde{z}_{i+k,j} - \bar{z}_{i+k,j}\|_{a,b}^2 \mid \bar{z}_{i,j}, \tilde{z}_{i,j} \right) \leq \exp \left(-\frac{mh_0}{4\gamma} \cdot k \right) \|\bar{z}_{i,j} - \tilde{z}_{i,j}\|_{a,b}^2.$$

With a slight abuse of notation, index j here refers to both position and velocity components, hence $\tilde{z}_{i,j} = (\tilde{x}_{i,j}, \tilde{v}_{i,j}, \tilde{x}'_{i,j}, \tilde{v}'_{i,j}) \in \mathbb{R}^4$. As previously, \bar{z}_i and \tilde{z}_{i+k} are independent, and

$$\begin{aligned} \text{Cov}(F(\bar{z}_i), F(\bar{z}_{i+k})) &= \mathbb{E}[(F(\bar{z}_i) - \mathbb{E}(F(\bar{z}_i)))(F(\bar{z}_{i+k}) - \mathbb{E}(F(\bar{z}_{i+k}))) \mid \bar{z}_i, \tilde{z}_i] \\ &\leq \sqrt{\text{Var}(F(\bar{z}_i)) \text{Var}(F(\bar{z}_{i+k}) - F(\tilde{z}_{i+k}))} \end{aligned}$$

By the Efron-Stein inequality [17, 90], and some rearrangement, we have

$$\begin{aligned} \text{Var}(F(\bar{z}_{i+k}) - F(\tilde{z}_{i+k})) &\leq 2 \sum_{j=1}^d \left(\sum_{s=1}^r |w_j^{(s)}| \right)^2 \mathbb{E} \left(\|\tilde{z}_{i+k,j} - \bar{z}_{i+k,j}\|_{a,b}^2 \right) \\ &\leq 2r \left(\sum_{s=1}^r \|w^{(s)}\|^2 \right) \exp \left(-\frac{mh_0}{4\gamma} \cdot k \right) \max_{1 \leq j \leq d} \mathbb{E} \left[\|\bar{z}_{i,j} - \tilde{z}_{i,j}\|_{a,b}^2 \right] \\ &= 4r \left(\sum_{s=1}^r \|w^{(s)}\|^2 \right) \exp \left(-\frac{mh_0}{4\gamma} \cdot k \right) \max_{1 \leq j \leq d} \mathbb{E} \left[\|\bar{z}_{i,j} - \mathbb{E} \bar{z}_{i,j}\|_{a,b}^2 \right], \end{aligned}$$

and the claim follows by rearrangement and summation. \square

PROPOSITION 3.18. *Suppose that Assumption 3.17 holds, and denote the potential U as $U(x) = \sum_{i=1}^d U_i(x_i)$. Suppose that Assumptions 3.6, 3.7, and 3.10 hold for each component $(U_i)_{1 \leq i \leq d}$, and that*

$$\gamma \geq \sqrt{8M}, \quad h_0 \leq \frac{1}{\gamma} \cdot \frac{m}{264M}, \quad B \geq \frac{16 \log(4)\gamma}{mh_0}, \quad B_0 \geq \frac{16\gamma}{mh_0} \log \left(\frac{c_{\mu_0} + 1}{\sqrt{M}\gamma h_0^2} \right).$$

Suppose that f is of the form

$$(3.18) \quad f(x, v) = g(\langle w^{(1)}, x \rangle, \dots, \langle w^{(r)}, x \rangle),$$

where $g : \mathbb{R}^r \rightarrow \mathbb{R}$ is 1-Lipschitz, and $w^{(1)}, \dots, w^{(r)} \in \mathbb{R}^d$. Suppose that $c_R \in [0, 1)$ and $2 < \phi_N < 16$. Then for any $N \geq 1$, the UBUBU estimator $S(c_R)$ has finite expected computational cost, $\mathbb{E}S(c_R) = \pi(f)$, and it has finite variance. Moreover, it satisfies a CLT as $N \rightarrow \infty$, and the asymptotic variance can be bounded as

$$\sigma_S^2 \leq \frac{C(m, M, M_1, \gamma, r, c_N, \phi_N)}{Kh_0} \sum_{1 \leq i \leq r} \|w^{(i)}\|^2.$$

PROOF OF PROPOSITION 3.18. Unbiasedness, finite variance, and finite computational cost follow from Theorem 3.15. By (3.8), the asymptotic variance can be expressed as

$$\sigma_S^2 := \text{Var}(D_0) + \sum_{l=0}^{\infty} \text{Var}(D_{l,l+1}) \cdot \frac{\phi_N^l}{c}.$$

It is easy to show that f is $\sum_{s=1}^r \|w^{(s)}\|$ -Lipschitz, so the variance term $\text{Var}(D_0)$ can be bounded using Proposition D.9, relying on the burn-in assumptions.

To control $\text{Var}(D_{l,l+1})$, we first need to bound terms of the form $\text{Var}(f(z'_k) - f(z_k))$. Let $z_{k,j} = (x_{k,j}, v_{k,j}) \in \mathbb{R}^2$ denote components j in both x and v . Using the Efron-Stein inequality [17, 90], and independence of the components, we have

$$\begin{aligned} \text{Var}(f(z'_k)^{(l,l+1)} - f(z_k)^{(l,l+1)}) &\leq 2 \sum_{j=1}^d \left(\sum_{s=1}^r |w_j^{(s)}| \right)^2 \mathbb{E} \left(\left\| z_{k,j}'^{(l,l+1)} - z_{k,j}^{(l,l+1)} \right\|_{a,b}^2 \right) \\ &\leq 2r \left(\sum_{s=1}^r \|w^{(s)}\|^2 \right) \max_{1 \leq j \leq d} \mathbb{E} \left(\left\| z_{k,j}'^{(l,l+1)} - z_{k,j}^{(l,l+1)} \right\|_{a,b}^2 \right). \end{aligned}$$

By applying Corollary D.6 component-wise, it follows that under our assumptions,

$$\max_{1 \leq j \leq d} \mathbb{E} \left(\left\| z_{k,j}'^{(l,l+1)} - z_{k,j}^{(l,l+1)} \right\|_{a,b}^2 \right) \leq C(m, M, \gamma, M_1) h_l^4,$$

hence

$$\text{Var}(f(z'_k)^{(l,l+1)} - f(z_k)^{(l,l+1)}) \leq C(m, M, M_1, \gamma, r) \sum_{1 \leq i \leq r} \|w^{(i)}\|^2 h_l^4.$$

Using this, and Proposition D.10, by a similar argument as in the proof of Theorem 3.15, we can show that

$$\text{Var}(D_{l,l+1}) \leq \frac{C(m, M, M_1, \gamma, r)}{K} \left(\sum_{1 \leq i \leq r} \|w^{(i)}\|^2 \right) h_l^4 \left(1 + \frac{4\gamma}{mh_0} + \log(4)l \right),$$

and the claim follows by summation and rearrangement. \square

APPENDIX E: INITIALIZATION AND GAUSSIAN APPROXIMATION

We will use the following assumptions in this section and sections **F** and **G**, which we restate here for easier readability. We will consider potentials of the form

$$(E.1) \quad U(x) = U_0(x) + \sum_{i=1}^{N_D} U_i(x),$$

where we aim to understand the scaling of the computational complexity when inexact gradients are used within the UBUBU framework in the large N_D case. We assume that the potential has the form (E.1) in this section and sections **F** and **G**, and we impose the following assumptions on the potential.

ASSUMPTION 3.21 (∇ Lipschitz property). *For every $1 \leq i \leq N_D$, $U_i : \mathbb{R}^d \rightarrow \mathbb{R}$ is twice differentiable and there exists a $\tilde{M} > 0$ such that for all $x, y \in \mathbb{R}^d$,*

$$\|\nabla U_i(x) - \nabla U_i(y)\| \leq \tilde{M}\|x - y\|,$$

for every $1 \leq i \leq N_D$ and moreover,

$$\|\nabla U(x) - \nabla U(y)\| \leq M\|x - y\| \quad \text{for } M = N_D \tilde{M}.$$

ASSUMPTION 3.22 ($N_D \tilde{m}$ -strong convexity). *There exists a $\tilde{m} > 0$ such that for all $x, y \in \mathbb{R}^d$*

$$\langle \nabla U(x) - \nabla U(y), x - y \rangle \geq m\|x - y\|^2 \quad \text{for } m = N_D \tilde{m}.$$

ASSUMPTION 3.23 (strongly Hessian Lipschitz property). *$U : \mathbb{R}^d \rightarrow \mathbb{R}$ is three times continuously differentiable and M_1^s -strongly Hessian Lipschitz if there exists $M_1^s > 0$ such that*

$$\|\nabla^3 U(x)\|_{\{1,2\}\{3\}} \leq M_1^s \quad \text{for } M_1^s = N_D \tilde{M}_1^s,$$

for all $x \in \mathbb{R}^d$.

For a better understanding of the scaling in terms of N_D , we also introduce

$$(E.2) \quad \tilde{\gamma} = \frac{\gamma}{\sqrt{N_D}},$$

so that $\gamma = \sqrt{N_D} \tilde{\gamma}$.

E.1. OHO scheme. In this section, we detail some results for the OHO scheme we use for initialization in (3.24)-(3.26). We state results for a potential that satisfies Assumptions 3.6 and 3.7 that can be applied in the case of Gaussian approximation. In particular, we show strong error results using similar techniques to [63] and [88].

We define the solution map \mathcal{H} to have update rule

$$(E.3) \quad \mathcal{H} : (x, v) \rightarrow \phi_h(x, v),$$

where $\phi_h(x, v)$ is the solution to the ODE

$$dX_t = V_t dt, \quad dV_t = -\nabla U(X_t) dt,$$

initialized at $(X_0, V_0) := (x, v) \in \mathbb{R}^{2d}$ at time $h > 0$. We then define the OHO scheme with stepsize $h > 0$ as a half step of \mathcal{O} with stepsize $h/2$ (defined in (3.25)), followed by a full step of \mathcal{H} with stepsize h and a half step of \mathcal{O} with stepsize $h/2$, which exactly preserves the invariant measure.

REMARK E.1. *We remark that the OHO scheme is a special case of the scheme studied in [72] using a hypocoercivity approach. It has also been considered as an exact splitting for discretization analysis in [13, 15, 45, 70]. In practice this scheme is only applicable when the Hamiltonian dynamics can be solved exactly, for example for a Gaussian target.*

PROPOSITION E.2. *Let $h > 0$ and $k \in \mathbb{N}$ and $(X_t, V_t)_{t \geq 0} := (Z_t)_{t \geq 0}$ be the solution of (1.1) and $(x_t, v_t)_{t \geq 0} := (z_t)_{t \geq 0}$ be the solution to the \mathcal{OHO} scheme with stepsize $h > 0$, with synchronously coupled Brownian motion and where both are initialized at $z_0 = Z_0 \sim \pi$. We have that*

$$\|Z_{kh} - z_{kh}\|_{L^2, a, b} \leq \sqrt{\frac{3}{2}} e^{\frac{3}{2}hk\sqrt{M}} \left(\frac{3h\gamma\sqrt{k\gamma h d} + 5k(h\gamma)^2\sqrt{d}}{\sqrt{M}} \right).$$

PROOF. Considering the OHO scheme given by

$$(x_h, v_h) := \left(x + h \left(\eta v + \sqrt{1 - \eta^2} \xi_1 \right) - \int_0^h \nabla U(x(t))(h-t) dt, \right. \\ \left. \eta \left(\eta v + \sqrt{1 - \eta^2} \xi_1 - h \nabla U(x) - \int_0^h \nabla^2 U(x(t)) \bar{v}(t)(h-t) dt \right) + \sqrt{1 - \eta^2} \xi_2 \right),$$

where the Hamiltonian dynamics $(x(t), v(t))_{t=0}^h$ is initialized at $(x, \eta v + \sqrt{1 - \eta^2} \xi_1)$. The kinetic Langevin dynamics for one step can be written as

$$(E.4) \quad V_h = \mathcal{E}(h)V_0 - \int_0^h \mathcal{E}(h-s)\nabla U(X_s) ds + \sqrt{2\gamma} \int_0^h \mathcal{E}(h-s) dW_s,$$

$$(E.5) \quad X_h = X_0 + \mathcal{F}(h)V_0 - \int_0^h \mathcal{F}(h-s)\nabla U(X_s) ds + \sqrt{2\gamma} \int_0^h \mathcal{F}(h-s) dW_s,$$

where $\mathcal{E}(h) = e^{-\gamma h}$, $\mathcal{F}(h) = \frac{1-e^{-\gamma h}}{\gamma}$, and we couple the noises such that $\sqrt{1 - \eta^2} \xi_1 = \sqrt{2\gamma} \int_0^{h/2} \mathcal{E}(h/2-s) dW_s$ and $\sqrt{1 - \eta^2} \xi_2 = \sqrt{2\gamma} \int_{h/2}^h \mathcal{E}(h-s) dW_s$. Considering the velocity component we have

$$\begin{aligned} \|V_h - v_h\|_{L^2} &\leq \left\| \eta^2 (V_0 - v_0) - \int_0^h \mathcal{E}(h-s) (\nabla U(X_s) - \nabla U(x)) ds + \sqrt{2\gamma}(1-\eta) \int_0^{h/2} \mathcal{E}(h/2-s) dW_s \right\|_{L^2} \\ &\quad + \sqrt{Md} \left| \frac{1 - \eta^2 - h\gamma\eta}{\gamma} \right| + \frac{h^2 M \sqrt{d}}{2} \\ &\leq \left\| \eta^2 (V_0 - v_0) - \int_0^h \mathcal{E}(h-s) (\nabla U(X_0) - \nabla U(x)) ds + \sqrt{2\gamma}(1-\eta) \int_0^{h/2} \mathcal{E}(h/2-s) dW_s \right\|_{L^2} \\ &\quad + \left\| \int_0^h \mathcal{E}(h-s) (\nabla U(X_s) - \nabla U(X_0)) ds \right\|_{L^2} + \left(\gamma\sqrt{M} + \frac{M}{2} \right) h^2 \sqrt{d} \\ &\leq \left\| \eta^2 (V_0 - v_0) - \int_0^h \mathcal{E}(h-s) (\nabla U(X_0) - \nabla U(x)) ds + \sqrt{2\gamma}(1-\eta) \int_0^{h/2} \mathcal{E}(h/2-s) dW_s \right\|_{L^2} \\ &\quad + \left(\gamma\sqrt{M} + 3M \right) h^2 \sqrt{d}, \end{aligned}$$

where the final estimate is a rough estimate due to (E.4), $(X_s, V_s) \sim \pi$ for all $s \in [0, h]$, the fact that U is M - ∇ Lipschitz, $h < \frac{1}{2\gamma}$ and $\gamma^2 \geq 4M$. Now considering time $kh \geq 0$ for $k \in \mathbb{N}$ and iteratively applying the argument whilst keeping Brownian components in the same L^2 norm, we have

$$\begin{aligned}
\|V_{kh} - v_{kh}\|_{L^2} &\leq \sum_{i=1}^k hMa_i + \sqrt{2\gamma} \left\| \sum_{i=1}^k \eta^{2(k-i)} (1-\eta) \int_{(i-1)h}^{(i-1/2)h} \mathcal{E}((i-1/2)h-s) dW_s \right\|_{L^2} \\
&\quad + 3k\gamma\sqrt{M}h^2\sqrt{d} \\
&\leq \sum_{i=1}^k hMa_i + \frac{h\gamma\sqrt{2\gamma}}{2} \left\| \sum_{i=1}^k \int_{(i-1)h}^{(i-1/2)h} \mathcal{E}((i-1/2)h-s) dW_s \right\|_{L^2} \\
&\quad + k(h\gamma)^2\sqrt{d} + 3k\gamma\sqrt{M}h^2\sqrt{d} \\
&\leq \sum_{i=1}^k hMa_i + h\gamma\sqrt{k\gamma h d} + 3k(h\gamma)^2\sqrt{d},
\end{aligned}$$

where $a_i := \|X_{ih} - x_{ih}\|_{L^2}$ for $i \in \mathbb{N}$ and $b_i := \|V_{ih} - v_{ih}\|_{L^2}$ for $i \in \mathbb{N}$. We have also used the independence of the Brownian motion over independent time intervals.

Now considering the position components we have

$$\begin{aligned}
\|X_h - x_h\|_{L^2} &\leq \left\| X_0 - x_0 + \mathcal{F}(h)(V_0 - v_0) - \int_0^h \mathcal{F}(h-s)(\nabla U(X_s) - \nabla U(x_s)) ds \right. \\
&\quad \left. + \int_0^h \sqrt{2\gamma} \mathcal{F}(h-s) dW_s - h\sqrt{2\gamma} \int_0^{h/2} \mathcal{E}(h/2-s) dW_s \right\|_{L^2} + 2\gamma h^2\sqrt{d} \\
&\leq \left\| X_0 - x_0 + \mathcal{F}(h)(V_0 - v_0) - \int_0^h \mathcal{F}(h-s)(\nabla U(X_0) - \nabla U(x_0)) ds \right. \\
&\quad \left. + \int_0^h \sqrt{2\gamma} \mathcal{F}(h-s) dW_s - h\sqrt{2\gamma} \int_0^{h/2} \mathcal{E}(h/2-s) dW_s \right\|_{L^2} + 3\gamma h^2\sqrt{d}.
\end{aligned}$$

Then as before we consider time $kh \geq 0$ for $k \in \mathbb{N}$ and we have

$$\begin{aligned}
\|X_{kh} - x_{kh}\|_{L^2} &\leq \sum_{i=1}^k (hb_i + h^2Ma_i) + 3k\gamma h^2\sqrt{d} \\
&\quad + \sqrt{2\gamma} \left\| \sum_{i=1}^k \int_{(i-1)h}^{ih} \mathcal{F}(ih-s) dW_s - h \int_{(i-1)h}^{(i-1/2)h} \mathcal{E}((i-1/2)h-s) dW_s \right\|_{L^2} \\
&\leq \sum_{i=1}^k (hb_i + h^2Ma_i) + 3k\gamma h^2\sqrt{d} + 2\sqrt{2\gamma}h\sqrt{hkd}.
\end{aligned}$$

In $\|\cdot\|_{L^2, a, b}$ using the preceding estimates we have

$$\|Z_{kh} - z_{kh}\|_{L^2, a, 0} \leq \frac{3}{2} \sum_{i=1}^k h\sqrt{M} \|Z_{ih} - z_{ih}\|_{L^2, a, 0} + \frac{3h\gamma\sqrt{k\gamma h d} + 5k(h\gamma)^2\sqrt{d}}{\sqrt{M}}$$

$$\leq e^{\frac{3}{2}hk\sqrt{M}} \left(\frac{3h\gamma\sqrt{k\gamma hd} + 5k(h\gamma)^2\sqrt{d}}{\sqrt{M}} \right).$$

□

THEOREM E.3. *Let $h > 0$ and $l \in \mathbb{N}$ and $(X_t, V_t)_{t \geq 0} := (Z_t)_{t \geq 0}$ be the solution of kinetic Langevin dynamics and $(x_l, v_l)_{l \in \mathbb{N}} := (z_l)_{l \in \mathbb{N}}$ be the iterates of the solution to the \mathcal{OHO} scheme with stepsize $h > 0$, where both are initialized at the same point according to the invariant measure. We have that*

$$\|Z_{lh} - z_l\|_{L^2, a, b} \leq 104h\gamma^2\sqrt{d} \left(\frac{3\sqrt{2\gamma/\sqrt{M}} + 10\gamma/\sqrt{M}}{m} \right).$$

PROOF. We use an approach used in [63] to remove the exponential constant in Proposition E.2. We define a sequence of interpolating variants $Z_l^{(k)}$ for every $k = 0, \dots, l$ as follows. We first define $Z_0^{(k)} = Z_0$, and then $(Z_i^{(k)})_{i=1}^k$ are defined by \mathcal{OHO} steps followed by $(Z_i^{(k)})_{i=k+1}^l$ steps of kinetic Langevin dynamics with stepsize $h > 0$. We break the l steps into blocks of size $\tilde{l} = \lceil \frac{2}{3h\sqrt{M}} \rceil$, then we have

$$\begin{aligned} \|Z_{lh} - z_l\|_{L^2, a, b} &= \|Z_l^{(0)} - Z_l^{(l)}\|_{L^2, a, b} \\ &\leq \sum_{j=0}^{\lceil l/\tilde{l} \rceil - 1} \left\| \left(Z_l^{(j\tilde{l})} - Z_l^{((j+1)\tilde{l})} \right) \right\|_{L^2, a, b} + \left\| \left(Z_l^{(\lceil l/\tilde{l} \rceil \tilde{l})} - Z_l^{(l)} \right) \right\|_{L^2, a, b}, \end{aligned}$$

where we bound the terms using the fact that the first $j\tilde{l}$ steps according to \mathcal{OHO} keep the stationary distribution invariant and they only deviate in the following \tilde{l} steps, where we will use Proposition E.2 with l chosen as \tilde{l} . We finally use contraction of the continuous dynamics (Corollary C.8) in the remaining steps and we have

$$\left\| \left(Z_l^{(j\tilde{l})} - Z_l^{((j+1)\tilde{l})} \right) \right\|_{L^2, a, b} \leq 4e^{-(l-1-(j+1)\tilde{l})c(h)} \left(\frac{3h\gamma\sqrt{\tilde{l}\gamma hd} + 5\tilde{l}(h\gamma)^2\sqrt{d}}{\sqrt{M}} \right).$$

Then summing up the terms we have that

$$\begin{aligned} \|Z_{lh} - z_l\|_{L^2, a, b} &\leq 4 \left(\frac{3h\gamma\sqrt{\tilde{l}\gamma hd} + 5\tilde{l}(h\gamma)^2\sqrt{d}}{\sqrt{M}} \right) \left(1 + \frac{1}{1 - e^{-c(h)\tilde{l}}} \right) \\ &\leq 8 \left(\frac{3h\gamma\sqrt{\tilde{l}\gamma hd} + 5\tilde{l}(h\gamma)^2\sqrt{d}}{\sqrt{M}} \right) \left(1 + \frac{12\gamma\sqrt{M}}{m} \right) \\ &\leq 104h\gamma^2\sqrt{d} \left(\frac{3\sqrt{2\gamma/\sqrt{M}} + 10\gamma/\sqrt{M}}{m} \right), \end{aligned}$$

as required. □

E.2. Initialization and bounds. For convex potentials, we can approximate the gradient with the Hessian at the minimizer by

$$(E.6) \quad \mathcal{Q}(x | x^*) = \nabla U(x^*) + H^*(x - x^*) = H^*(x - x^*),$$

where $x^* \in \mathbb{R}^d$ is the minimizer of U and $H^* = \nabla^2 U(x^*)$.

LEMMA E.4. *Considering the gradient approximation \mathcal{Q} given by (E.6), where the potential U satisfies Assumption 3.23, and has a minimizer $x^* \in \mathbb{R}^d$ we then have the property*

$$\mathbb{E} \|\nabla U(x) - \mathcal{Q}(x | x^*)\|^p \leq (\tilde{M}_1^s)^p N_D^p \|x - x^*\|_{L^{2p}}^{2p},$$

for any $x \in \mathbb{R}^d$.

PROOF. Follows from Taylor expansion. \square

We then define the measure $\mu_G = \mathcal{N}(x^*, (H^*)^{-1}) \times \mathcal{N}(0_d, I_d)$ to be the Gaussian approximation of the target as in (3.23), which is the invariant measure of the OHO scheme and continuous kinetic Langevin dynamics with the use of the gradient approximation (E.6).

PROPOSITION E.5. *Let $p = 2$ or 4 , then we have the following Wasserstein bound between a potential U which satisfies Assumptions 3.22, 3.23 and 3.28*

$$\mathcal{W}_{p,a,b}(\pi, \mu_G) \leq \sqrt{\frac{3}{2}} \left(\frac{(2p)!}{2^p p!} \right)^{1/p} \frac{\tilde{M}_1 d}{\tilde{m}^2 N_D}.$$

PROOF. If we let π_x denote the marginal in the position of π and $(\mu_G)_x$ denote the marginal in position of μ_G . We have from the equivalence of norms that for $p = 2, 4$

$$\begin{aligned} \mathcal{W}_{p,a,b}(\pi, \mu_G) &\leq \sqrt{\frac{3}{2}} \mathcal{W}_{p,a,0}(\pi, \mu_G) \leq \sqrt{\frac{3}{2}} \mathcal{W}_p(\pi_x, (\mu_G)_x) \\ &\leq \sqrt{\frac{3}{2}} \frac{\|\nabla U - \mathcal{Q}(\cdot | x^*)\|_{L^p}}{m} \leq \sqrt{\frac{3}{2}} \frac{M_1^s \|x - x^*\|_{L^{2p}}^2}{m} \\ &\leq \sqrt{\frac{3}{2}} \left(\frac{(2p)!}{2^p p!} \right)^{1/p} \frac{M_1^s d}{m^2} \end{aligned}$$

where the third inequality is due to Proposition 22 of [96], the fourth due to Lemma E.4 and the final inequality by Lemma H.2. \square

APPENDIX F: VARIANCE BOUNDS FOR UBU WITH SVRG

For this section, we make use of the technique of the recent work of Hu et al [101], related to using stochastic variance reduced gradient (SVRG).

A stochastic gradient version of the UBU scheme is simply constructed by replacing the \mathcal{B} operator with

$$\mathcal{B}_G(x, v, h, \omega | \hat{x}) = (x, v - h\mathcal{G}(x, \omega | \hat{x})),$$

where \mathcal{G} is a stochastic gradient approximation of the potential U as defined in the approximation given by (3.20).

We start with an alternative formula for the kinetic Langevin dynamics introduced in (1.1). This is used for the analysis of the UBU scheme in the full gradient setting in [86] and alternative schemes with the SVRG approximation in (3.20). The convenient way of expressing

kinetic Langevin dynamics is to use Itô's formula on the product $e^{\gamma t} V_t$. This results in the following set of equations for (1.1) with initial condition $(X_0, V_0) \in \mathbb{R}^{2d}$:

$$(F.1) \quad V_t = \mathcal{E}(t)V_0 - \int_0^t \mathcal{E}(t-s)\nabla U(X_s)ds + \sqrt{2\gamma} \int_0^t \mathcal{E}(t-s)dW_s,$$

$$(F.2) \quad X_t = X_0 + \mathcal{F}(t)V_0 - \int_0^t \mathcal{F}(t-s)\nabla U(X_s)ds + \sqrt{2\gamma} \int_0^t \mathcal{F}(t-s)dW_s,$$

where

$$(F.3) \quad \mathcal{E}(t) = e^{-\gamma t} \quad \mathcal{F}(t) = \frac{1 - e^{-\gamma t}}{\gamma}.$$

Then the UBU scheme (as in [86]) can be expressed as

$$(F.4) \quad v_{k+1} = \mathcal{E}(h)v_k - h\mathcal{E}(h/2)\nabla U(\bar{x}_k) + \sqrt{2\gamma} \int_{kh}^{(k+1)h} \mathcal{E}((k+1)h-s)dW_s,$$

$$(F.5) \quad \bar{x}_k = x_k + \mathcal{F}(h/2)v_k + \sqrt{2\gamma} \int_{kh}^{(k+1/2)h} \mathcal{F}((k+1/2)h-s)dW_s,$$

$$(F.6) \quad x_{k+1} = x_k + \mathcal{F}(h)v_k - h\mathcal{F}(h/2)\nabla U(\bar{x}_k) + \sqrt{2\gamma} \int_{kh}^{(k+1)h} \mathcal{F}((k+1)h-s)dW_s,$$

which can be more easily compared to the true dynamics via (F.1) and (F.2). We will refer to $(\bar{x}_k)_{k \in \mathbb{N}}$ as the gradient evaluation points of the scheme. Similarly, stochastic gradient UBU can be expressed as (F.4)-(F.6) by replacing the gradients with stochastic gradient approximations,

(F.7)

$$v_{k+1} = \mathcal{E}(h)v_k - h\mathcal{E}(h/2)\mathcal{G}(\bar{x}_k, \omega_{k+1}|\hat{x}_k) + \sqrt{2\gamma} \int_{kh}^{(k+1)h} \mathcal{E}((k+1)h-s)dW_s,$$

(F.8)

$$\bar{x}_k = x_k + \mathcal{F}(h/2)v_k + \sqrt{2\gamma} \int_{kh}^{(k+1/2)h} \mathcal{F}((k+1/2)h-s)dW_s,$$

(F.9)

$$\hat{x}_k = \bar{x}_{L(k)} \quad \text{for } L(k) = \tau \lfloor k/\tau \rfloor,$$

(F.10)

$$x_{k+1} = x_k + \mathcal{F}(h)v_k - h\mathcal{F}(h/2)\mathcal{G}(\bar{x}_k, \omega_{k+1}|\hat{x}_k) + \sqrt{2\gamma} \int_{kh}^{(k+1)h} \mathcal{F}((k+1)h-s)dW_s,$$

If we are using a stochastic gradient approximation of the UBU dynamics, additional bias is introduced by the use of gradient approximations. We wish to measure the local error caused by the stochastic gradient approximation.

F.1. Variance bound of $D_{l,l+1}$. Suppose now we have two UBU schemes, a UBU scheme $(z_k)_{k \in \mathbb{N}} = (x_k, v_k)_{k \in \mathbb{N}}$ which uses a stochastic gradient approximation as defined in Definition 2.1 with $(\omega_k)_{k \in \mathbb{N}}$ such that $\omega_k \sim \mathcal{SWR}(N_D, N_b)$ for each $k \in \mathbb{N}$. Further at iteration k define $z_k^h := (x_k^h, v_k^h) := \psi_h(z_k, h, (W_{t'})_{t'=kh}^{(k+1)h})$ to be a step of the full gradient UBU scheme at iteration z_k , with synchronously coupled Brownian motion. Then the local error after one step is

$$\mathbb{E}\|(x_{k+1} - x_k^h, v_{k+1} - v_k^h)\|^2 = h^2 (\mathcal{E}(h/2) + \mathcal{F}(h/2))^2 \mathbb{E}\|\nabla U(\bar{x}_k) - \mathcal{G}(\bar{x}_k, \omega_{k+1})\|^2,$$

and

$$\mathbb{E}\|x_{k+1} - x_k^h\|^2 \leq \frac{h^4}{4} \mathbb{E}\|\nabla U(\bar{x}_k) - \mathcal{G}(\bar{x}_k, \omega_{k+1})\|^2,$$

where expectations are taken over stochastic gradient approximation and Brownian increments. The sequence $(\bar{x}_k)_{k \in \mathbb{N}}$ are the points where the stochastic gradient approximations are evaluated defined by (F.8). We now wish to bound the term $\mathbb{E}\|\nabla U(\bar{x}_k) - \mathcal{G}(\bar{x}_k, \omega_{k+1})\|^2$, uniformly in k to control the error due to the stochastic gradient. For this, we state Lemma 1 of [101] with our notations, together with its proof.

LEMMA F.1. *Considering iterates $(x_k, v_k, \bar{x}_k)_{k \in \mathbb{N}}$ of stochastic gradient UBU with the SVRG $(\mathcal{G}, \text{SWR}(N_D, N_b))$ for a potential U which has the form (3.19), with data size N_D and batch size N_b , epoch length $\tau = \lceil N_D/N_b \rceil$, and initial condition $(x_0, v_0) \in \mathbb{R}^{2d}$, then we have the property*

$$\begin{aligned} & \mathbb{E} \left(\left\| \mathcal{G}(\bar{x}_k, \omega_{k+1} \mid \bar{x}_{L(k)}) - \nabla U(\bar{x}_k) \right\|^2 \right) \\ & \leq \frac{N_D(N_D - N_b)(\tau - 1)^2}{N_b(N_D - 1)} \cdot \max_{j < k} \sum_{i=1}^{N_D} \mathbb{E} \left(\left\| \nabla U_i(\bar{x}_{j+1}) - \nabla U_i(\bar{x}_j) \right\|^2 \right) \end{aligned}$$

COROLLARY F.2. *Suppose that Assumption 3.21 holds. For UBU with SVRG updates as defined by (F.7)-(F.10), we have*

$$(F.11) \quad \mathbb{E} \left(\left\| \mathcal{G}(\bar{x}_k, \omega_{k+1} \mid \bar{x}_{L(k)}) - \nabla U(\bar{x}_k) \right\|^2 \right) \leq \Theta \max_{j < k} \mathbb{E} \|\bar{x}_{j+1} - \bar{x}_j\|^2,$$

$$(F.12) \quad \Theta = \frac{\tilde{M}^2 N_D^2 (N_D - N_b)(\tau - 1)^2}{N_b(N_D - 1)}.$$

PROOF OF LEMMA F.1. For the potential of the form $U(x) = U_0(x) + \sum_{i=1}^{N_D} U_i(x)$ and for $k \geq 1$ we define $\bar{X}_i = \nabla U_i(\bar{x}_k) - \nabla U_i(\bar{x}_{L(k)})$ and we define $Y_i = N_D \bar{X}_i - \sum_{j=1}^{N_D} \bar{X}_j$ for $i = 1, \dots, N_D$. Then we have that $\sum_{i=1}^{N_D} Y_i = 0$ and that

$$\mathcal{G}(\bar{x}_k, \omega_{k+1} \mid \bar{x}_{L(k)}) - \nabla U(\bar{x}_k) = \frac{1}{N_b} \sum_{i \in \omega_{k+1}} Y_i.$$

Therefore our aim is to establish a bound on $\frac{1}{N_b} \sum_{i \in \omega_{k+1}} Y_i$. We have that

$$\begin{aligned} \mathbb{E}_{\omega_{k+1}} \left\| \frac{1}{N_b} \sum_{i \in \omega_{k+1}} Y_i \right\|^2 &= \frac{1}{N_b^2} \mathbb{E}_{\omega_{k+1}} \left(\sum_{i \in \omega_{k+1}} \|Y_i\|^2 + \sum_{i \neq j \in \omega_{k+1}} \langle Y_i, Y_j \rangle \right) \\ &= \frac{1}{N_b N_D} \sum_{i=1}^{N_D} \|Y_i\|^2 + \frac{b-1}{N_b N_D (N_D - 1)} \sum_{i \neq j} \langle Y_i, Y_j \rangle \\ &= \frac{N_D - N_b}{N_D - 1} \frac{1}{N_b N_D} \sum_{i=1}^{N_D} \|Y_i\|^2, \end{aligned}$$

where the last line is due to the fact that $\sum_{i=1}^{N_D} Y_i = 0$. Then using the fact that $\sum_{i=1}^{N_D} \|Y_i\|^2 \leq N_D^2 \sum_{i=1}^{N_D} \|\bar{X}_i\|^2$ and the last full gradient evaluation is at $k - \tau + 1 \leq L(k) \leq k$ we have that

$$\mathbb{E} \left(\left\| \nabla U(\bar{x}_k) - \mathcal{G}(\bar{x}_k, \omega_{k+1} \mid \bar{x}_{L(k)}) \right\|^2 \right) = \frac{N_D - N_b}{N_b N_D (N_D - 1)} \sum_{i=1}^{N_D} \mathbb{E} \|Y_i\|^2$$

$$\begin{aligned}
&\leq \frac{N_D(N_D - N_b)}{N_b(N_D - 1)} \sum_{i=1}^{N_D} \mathbb{E} \|\bar{X}_i\|^2 \\
&\leq \frac{N_D(N_D - N_b)}{N_b(N_D - 1)} \sum_{i=1}^{N_D} \mathbb{E} \|\nabla U_i(\bar{x}_k) - \nabla U_i(\bar{x}_{L(k)})\|^2 \\
&\leq \frac{N_D(N_D - N_b)(k - L(k))}{N_b(N_D - 1)} \sum_{j=L(k)}^{k-1} \sum_{i=1}^{N_D} \mathbb{E} \|\nabla U_i(\bar{x}_{j+1}) - \nabla U_i(\bar{x}_j)\|^2 \\
&\leq \frac{N_D(N_D - N_b)(\tau - 1)^2}{N_b(N_D - 1)} \max_{j < k} \sum_{i=1}^{N_D} \mathbb{E} \|\nabla U_i(\bar{x}_{j+1}) - \nabla U_i(\bar{x}_j)\|^2,
\end{aligned}$$

which concludes the proof. \square

Hence it is sufficient to bound $\mathbb{E} \|\bar{x}_{k+1} - \bar{x}_k\|^2$ uniformly in $k \in \mathbb{N}$, which will be done in the following lemma.

LEMMA F.3 (Displacement Lemma). *Let a stochastic gradient UBU integrator defined by (F.7)-(F.8) with stochastic gradient (\mathcal{G}, ρ) satisfy*

$$\mathbb{E} \left(\|\mathcal{G}(\bar{x}_k, \omega_{k+1} | \hat{x}_k) - \nabla U(\bar{x}_k)\|^2 \right) \leq \Theta \max_{j < k} \mathbb{E} \|\bar{x}_{j+1} - \bar{x}_j\|^2,$$

for some $\Theta > 0$. If U satisfies Assumptions 3.21, 3.22, $h < 1/2\gamma$ and $\gamma^2 \geq 8M$, then

$$\|\bar{x}_{k+1} - \bar{x}_k\|_{L^2} \leq h^2 \sqrt{\Theta} \max_{0 \leq i < k} \|\bar{x}_{i+1} - \bar{x}_i\|_{L^2} + 7h\sqrt{M} \|z_{k-1} - Z^{k-1}\|_{L^{2,a,b}} + 6h\sqrt{d},$$

where $Z^k := Z_{kh} = \phi(Z_0, kh, (W_t)_{t'=0}^{kh}) \in \mathbb{R}^{2d}$ is the solution to (1.1) initialized at the invariant measure $Z_0 \sim \pi$ at time kh for $k \in \mathbb{N}$.

PROOF. Then we use the following estimate

$$\begin{aligned}
\|\bar{x}_{k+1} - \bar{x}_k\|_{L^2} &= \|\bar{x}_{k+1} - x_{k+1} + x_{k+1} - x_k + x_k - \bar{x}_k\|_{L^2} \\
&\leq \|\mathcal{F}(h/2)(v_k - v_{k-1})\|_{L^2} + \|x_{k+1} - x_k\|_{L^2} \\
&\quad + \sqrt{2\gamma} \left\| \int_{(k+1)h}^{(k+3/2)h} \mathcal{F}((k+3/2)h - s) dW_s \right\|_{L^2} \\
&\quad + \sqrt{2\gamma} \left\| \int_{kh}^{(k+1/2)h} \mathcal{F}((k+1/2)h - s) dW_s \right\|_{L^2} \\
&=: \text{(I)} + \text{(II)} + \text{(III)} + \text{(IV)},
\end{aligned}$$

and we bound (I), (II), (III) and (IV) separately. (III) and (IV) can be bounded above by $\sqrt{\gamma h^3 d}$. Firstly, we will bound (II), but first we denote

$$A_j = \|\bar{x}_{j+1} - \bar{x}_j\|_{L^2}^2,$$

for $j \in \mathbb{N}$, and $z_k^h = (x_k^h, v_k^h) := \psi_h(z_k, h, (W_t)_{t'=kh}^{(k+1)h})$ is an iterate with stepsize h and initial point (x_k, v_k) of the full gradient UBU scheme and synchronously coupled Brownian motion

to the stochastic gradient scheme. We then estimate

$$\begin{aligned} \|x_{k+1} - x_k\|_{L^2} &\leq \|x_{k+1} - x_k^h\|_{L^2} + \|x_k^h - x_k\|_{L^2} \\ &\leq \frac{h^2}{2} \sqrt{\Theta} \max_{j < k} \sqrt{A_j} + \|x_k^h - x_k\|_{L^2}, \end{aligned}$$

then if we define the notation $Z_k^t = (X_k^t, V_k^t) := \phi(z_k, t, (W_{t'})_{t'=kh}^{kh+t}) \in \mathbb{R}^{2d}$ for $k \in \mathbb{N}$ and $t \geq 0$ to be the continuous dynamics solution with initial condition (x_k, v_k) at time t defined by (F.1) and (F.2). Then we can estimate the second term by splitting it up into discretization error and one-step displacement and bounding each of these terms separately as

$$\|x_k^h - x_k\|_{L^2} \leq \|x_k^h - X_k^h\|_{L^2} + \|X_k^h - x_k\|_{L^2}.$$

Then using [86][Section 7.6] we have that

$$\begin{aligned} \|x_k^h - X_k^h\|_{L^2} &\leq \left\| \int_0^h \mathcal{F}(h/2) (\nabla U(X_k^s) - \nabla U(\bar{x}_k)) ds + \int_0^h (\mathcal{F}(h-s) - \mathcal{F}(h/2)) \nabla U(X_k^s) ds \right\|_{L^2} \\ &\leq \frac{h}{2} \int_0^h \|\nabla U(X_k^s) - \nabla U(\bar{x}_k)\|_{L^2} ds + h^2 \max_{0 \leq s \leq h} \|\nabla U(X_k^s)\|_{L^2} \\ &\leq \frac{hM}{2} \int_0^h \left\| X_k^s - \left(x_k + \mathcal{F}(h/2)v_k + \sqrt{2\gamma} \int_{kh}^{(k+1/2)h} \mathcal{F}((k+1/2)h-s) dW_s \right) \right\|_{L^2} ds \\ &\quad + h^2 \max_{0 \leq s \leq h} \|\nabla U(X_k^s)\|_{L^2} \\ &\leq h^2 \max_{0 \leq s \leq h} \|\nabla U(X_k^s)\|_{L^2} + \frac{hM}{2} \int_0^h \|X_k^s - x_k\|_{L^2} ds + \frac{h^3M}{4} \max_{0 \leq s \leq h} \|V_k^s\|_{L^2} + \frac{h^{7/2}M\sqrt{\gamma d}}{4}. \end{aligned}$$

Now, we bound

$$\int_0^h \|X_k^s - x_k\|_{L^2} ds \leq h^2 \|v_k\|_{L^2} + h^3 \max_{0 \leq s \leq h} \|\nabla U(X_k^s)\|_{L^2} + h^{5/2} \sqrt{2\gamma d},$$

and using the fact that $h < \min\{\frac{1}{5\sqrt{M}}, \frac{1}{2\gamma}\}$ we have

$$\|x_k^h - X_k^h\|_{L^2} \leq \frac{3h^2}{2} \max_{0 \leq s \leq h} \|\nabla U(X_k^s)\|_{L^2} + h \max_{0 \leq s \leq h} \|V_k^s\|_{L^2} + h\sqrt{d},$$

and using (1.1) we have that

$$\begin{aligned} \|X_k^h - x_k\|_{L^2} &= \left\| \mathcal{F}(h)v_k - \int_0^h \mathcal{F}(h-s) \nabla U(X_k^s) ds + \sqrt{2\gamma} \int_0^h \mathcal{F}(h-s) dW_s \right\|_{L^2} \\ &\leq h \|v_k\|_{L^2} + h^2 \max_{0 \leq s \leq h} \|\nabla U(X_k^s)\|_{L^2} + \sqrt{2\gamma h^3 d}. \end{aligned}$$

To bound the maximum terms we introduce $(Z_t)_{t \geq 0} = (X_t, V_t)_{t \geq 0}$ to be the solution to (1.1) initialized at the invariant measure with synchronously coupled Brownian motion. We also define $Z^k := Z_{kh}$ for $k \in \mathbb{N}$. Then we have, in expectation, for any $0 \leq s \leq h$,

$$\begin{aligned} \|V_k^s\|_{L^2} &\leq \|V_k^s - V_{kh+s}\|_{L^2} + \|V_{kh+s}\|_{L^2} \\ &\leq \sqrt{2M} \|z_k - Z^k\|_{L^2, a, b} + \sqrt{d}, \end{aligned}$$

and for any $0 \leq s \leq h$ we have

$$\begin{aligned} \|\nabla U(X_k^s)\|_{L^2} &\leq \|\nabla U(X_k^s) - \nabla U(X_{kh+s})\|_{L^2} + \|\nabla U(X_{kh+s})\|_{L^2} \\ &\leq M\|X_k^s - X_{kh+s}\|_{L^2} + \sqrt{Md} \\ &\leq \sqrt{2}M\|z_k - Z^k\|_{L^2,a,b} + \sqrt{Md}, \end{aligned}$$

where we have used contraction of the continuous dynamics under synchronous coupling provided in Corollary C.8 and [28][Lemma 2] to bound $\|\nabla U(X_{kh+s})\|_{L^2}$. Therefore we have the following bound on (II)

$$(II) \leq \frac{h^2}{2} \sqrt{\Theta} \max_{0 \leq i < k} \sqrt{A_i} + 4h\sqrt{M}\|z_k - Z^k\|_{L^2,a,b} + \frac{9h\sqrt{d}}{2},$$

where $h < \frac{1}{5\sqrt{M}}$ due to the fact that $\gamma^2 \geq 8M$ and $h < \frac{1}{2\gamma}$.

Next, we consider (I) and we can estimate

$$\begin{aligned} (I) &\leq \frac{h}{2} \|v_k - v_{k-1}\|_{L^2} \leq \frac{h}{2} \|v_k - v_{k-1}^h\|_{L^2} + \frac{h}{2} \|v_{k-1}^h - v_{k-1}\|_{L^2} \\ &\leq \frac{h^2}{2} \sqrt{\Theta} \max_{0 \leq i < k} \sqrt{A_i} + \frac{h}{2} \|v_{k-1}^h - v_{k-1}\|_{L^2}, \end{aligned}$$

where

$$\|v_{k-1}^h - v_{k-1}\|_{L^2} \leq \|v_{k-1}^h - V_{k-1}^h\|_{L^2} + \|V_{k-1}^h - v_{k-1}\|_{L^2}.$$

Then we can bound

$$\begin{aligned} &\|v_{k-1}^h - V_{k-1}^h\|_{L^2} \leq \\ &\left\| \int_0^h \mathcal{E}(h/2) (\nabla U(X_{k-1}^s) ds - \nabla U(\bar{x}_{k-1})) + (\mathcal{E}(h-s) - \mathcal{E}(h/2)) \nabla U(X_{k-1}^s) ds \right\|_{L^2} \\ &\leq M \int_0^h \|X_{k-1}^s - \bar{x}_{k-1}\|_{L^2} ds + h \max_{0 \leq s \leq h} \|\nabla U(X_{k-1}^s)\|_{L^2} \\ &\leq \frac{3}{50} \max_{0 \leq s \leq h} \|V_{k-1}^s\|_{L^2} + 2\frac{51}{50}h \max_{0 \leq s \leq h} \|\nabla U(X_{k-1}^s)\|_{L^2} + \frac{2}{25}\sqrt{d}, \end{aligned}$$

where we have used the estimate of $\int_0^h \|X_{k-1}^s - \bar{x}_{k-1}\|_{L^2} ds$ from the $\|x_{k-1}^h - X_{k-1}^h\|_{L^2}$ bound and the fact that $h < 1/5\sqrt{M}$. Using (F.1) we have

$$\begin{aligned} \|V_{k-1}^h - v_{k-1}\|_{L^2} &\leq \left\| (\mathcal{E}(h) - 1)v_{k-1} - \int_0^h \mathcal{E}(h-s) \nabla U(X_{k-1}^s) ds + \sqrt{2\gamma} \int_0^h \mathcal{E}(h-s) dW_s \right\|_{L^2} \\ &\leq h\gamma \|v_{k-1}\|_{L^2} + h \max_{0 \leq s \leq h} \|\nabla U(X_{k-1}^s)\|_{L^2} + \sqrt{2\gamma hd} \\ &\leq 2h\sqrt{M} (\gamma + \sqrt{M}) \|z_{k-1} - Z^{k-1}\|_{L^2,a,b} + h\sqrt{d} (\gamma + \sqrt{M}) + \sqrt{2\gamma hd}, \end{aligned}$$

and we can combine terms to get the following bound on (I)

$$(I) \leq \frac{h^2}{2} \max_{0 \leq i < k} \sqrt{\Theta A_i} + 3h\sqrt{M}\|z_{k-1} - Z^{k-1}\|_{L^2,a,b} + \frac{5}{4}h\sqrt{d},$$

and summing all terms we have that

$$\|\bar{x}_{k+1} - \bar{x}_k\|_{L^2} \leq h^2 \max_{0 \leq i < k} \sqrt{\Theta A_i} + 7h\sqrt{M}\|z_{k-1} - Z^{k-1}\|_{L^2,a,b} + 6h\sqrt{d}.$$

□

PROPOSITION F.4. *For a stochastic gradient UBU integrator with iterates $(z_k)_{k \in \mathbb{N}}$, gradient evaluation points $(\bar{x}_k)_{k \in \mathbb{N}}$, transition kernel P_h and potential U satisfying Assumptions 3.21-3.22, where we approximate the gradient using a unbiased stochastic gradient (\mathcal{G}, ρ) satisfying*

$$\mathbb{E} \left(\|\mathcal{G}(\bar{x}_k, \omega_{k+1} | \hat{x}_k) - \nabla U(\bar{x}_k)\|^2 \right) \leq \Theta \max_{j < k} \mathbb{E} \|\bar{x}_{j+1} - \bar{x}_j\|^2.$$

Consider the continuous solution to (1.1) initialized at the invariant measure, for $k \in \mathbb{N}$ define $Z^k := Z_{kh} = \phi(Z_0, kh, (W_{t'})_{t'=0}^{kh}) \in \mathbb{R}^{2d}$ with synchronously coupled Brownian motion to $(z_k)_{k \in \mathbb{N}}$, then for all

$$h < \min \left\{ \frac{1}{2\tilde{\gamma}N_D^{1/2}}, 1/2, \frac{\tilde{m}^{1/3}N_D^{1/6}}{24(\Theta\tilde{\gamma})^{1/3}}, \frac{1}{4\Theta^{1/4}}, \frac{\tilde{m}}{256\tilde{M}\tilde{\gamma}N_D^{1/2}} \right\},$$

we have

$$\begin{aligned} \|z_k - Z^k\|_{L^2, a, b} &\leq 4(1 - R_2(h)/2)^k \left(\|z_0 - Z^0\|_{L^2, a, b} + \sqrt{2}C_1h^{5/2} \right) \\ &+ C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s)h^{3/2}d^{1/2}N_D^{-3/4}\Theta^{1/2}, \end{aligned}$$

where $R_2(h) = 1 - c_2(h) + C_0^2h^2$.

Further for all $\mu \in \mathcal{P}_2(\mathbb{R}^{2d})$, and all $k \in \mathbb{N}$,

$$\begin{aligned} \mathcal{W}_{2, a, b}(\mu P_h^k, \pi) &\leq 4(1 - R_2(h)/2)^k \left(\mathcal{W}_{2, a, b}(\mu, \pi) + \sqrt{2}C_1h^{5/2} \right) \\ &+ C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s)h^{3/2}d^{1/2}N_D^{-3/4}\Theta^{1/2}. \end{aligned}$$

PROOF. Let us define the notation $Z_k^t = (X_k^t, V_k^t) := \phi(z_k, t, (W_{t'})_{t'=kh}^{kh+t}) \in \mathbb{R}^{2d}$ for $k \in \mathbb{N}$ and $t \geq 0$ to be the continuous dynamics solution with initial condition (x_k, v_k) at time t defined by (F.1) and (F.2). Further define $z_k^h = (x_k^h, v_k^h) := \psi_h(z_k, h, (W_{t'})_{t'=kh}^{(k+1)h})$ is an iterate with stepsize h and initial point (x_k, v_k) of the full gradient UBU scheme and synchronously coupled Brownian motion to the stochastic gradient scheme.

Firstly, we split up the difference in the following way

$$\begin{aligned} \|z_k - Z^k\|_{L^2, a, b}^2 &= \left\| \left(z_k - z_{k-1}^h \right) + \left(z_{k-1}^h - Z^k \right) \right\|_{L^2, a, b}^2 \\ &= \left\| z_k - z_{k-1}^h \right\|_{L^2, a, b}^2 + 2 \left\langle z_k - z_{k-1}^h, z_{k-1}^h - Z^k \right\rangle_{L^2, a, b} + \|z_{k-1}^h - Z^k\|_{L^2, a, b}^2. \end{aligned}$$

Considering the inner product we have the expectation conditional on z_{k-1} and $(W_{t'})_{t'=(k-1)h}^{kh}$ is zero as it is independent of the Brownian motion (due to synchronous coupling) and the stochastic gradient estimator is unbiased. Therefore

$$\begin{aligned} \|z_k - Z^k\|_{L^2, a, b}^2 &\leq \|z_k - z_{k-1}^h\|_{L^2, a, b}^2 + (\|\beta_{k-1}\|_{L^2, a, b} \\ &+ \|z_{k-1}^h - \psi(Z^{k-1}, h, (W_{t'})_{t'=(k-1)h}^{kh}) + \alpha_{k-1}\|_{L^2, a, b})^2 \\ &= (\text{I})' + (\text{II})'. \end{aligned}$$

We have that

$$(\text{I})' \leq \frac{2h^2\Theta}{M} \max_{j < k} \|\bar{x}_{j+1} - \bar{x}_j\|_{L^2}^2,$$

and

$$(II)' \leq \left(\sqrt{(1 - c_2(h) + C_0^2 h^2) \|z_{k-1} - Z^{k-1}\|_{L^2, a, b}^2 + 2C_1^2 h^5 + C_2 h^3} \right)^2.$$

Let $R_2(h) := c_2(h) - C_0^2 h^2$, then assuming that $mh/8\gamma < R_2(h) < 1/2$ (which holds for $h < \frac{1}{2\gamma}$ and $h < \frac{m}{256\gamma M}$), using Lemma H.1,

$$\begin{aligned} \|z_k - Z^k\|_{L^2, a, b} &\leq \sqrt{2}(1 - R_2(h)/2)^k \left(\|z_0 - Z^0\|_{L^2, a, b} + \sqrt{2C_1^2 h^5} \right) + \frac{2\sqrt{2}C_2 h^3}{R_2(h)} \\ &\quad + 2\sqrt{\frac{\frac{2h^2\Theta}{M} \max_{j < k} \|\bar{x}_{j+1} - \bar{x}_j\|_{L^2}^2 + 2C_1^2 h^5}{R_2(h)}}, \end{aligned}$$

and now we wish to bound $\max_{j < k} \|\bar{x}_{j+1} - \bar{x}_j\|_{L^2}^2$. Considering Lemma F.3 we have that

$$\begin{aligned} \|\bar{x}_{k+1} - \bar{x}_k\|_{L^2} &\leq h^2 \sqrt{\Theta \max_{0 \leq j \leq k} \|\bar{x}_{j+1} - \bar{x}_j\|_{L^2}^2} + 6h\sqrt{d} \\ &\quad + 7h\sqrt{M} \left(\sqrt{2}(1 - R_2(h)/2)^k \left(\|z_0 - Z^0\|_{L^2} + \sqrt{2C_1^2 h^5} \right) + \frac{16\sqrt{2}\gamma C_2 h^2}{m} \right) \\ &\quad + 56h\sqrt{\frac{h\gamma\Theta \max_{j < k} \|\bar{x}_{j+1} - \bar{x}_j\|_{L^2}^2 + \gamma C_1^2 h^4}{m}}. \end{aligned}$$

If we assume that

$$h < \min \left\{ \frac{m^{1/3}}{24(\Theta\gamma)^{1/3}}, \frac{1}{4\Theta^{1/4}}, 1 \right\},$$

then

$$\begin{aligned} \max_{j < k} \|\bar{x}_{j+1} - \bar{x}_j\|_{L^2} &\leq 21h\sqrt{M} \left(\sqrt{2}(1 - R_2(h)/2)^k \left(\|z_0 - Z^0\|_{L^2} + \sqrt{2C_1^2 h^5} \right) + \frac{16\sqrt{2}\gamma C_2 h^2}{m} \right) \\ &\quad + 18h\sqrt{d} + 168h^3\sqrt{M}C_1\sqrt{\frac{\gamma}{m}}, \end{aligned}$$

and

$$\begin{aligned} \|z_k - Z^k\|_{L^2, a, b} &\leq \sqrt{2}(1 - R_2(h)/2)^k \left(\|z_0 - Z^0\|_{L^2, a, b} + \sqrt{2}C_1 h^{5/2} \right) \\ &\quad + \frac{16\sqrt{2}\gamma h^2 (C_2\sqrt{\gamma} + C_1\sqrt{m})}{m} + 2\sqrt{\frac{16h\gamma\Theta \max_{j < k} \|\bar{x}_{j+1} - \bar{x}_j\|_{L^2}^2}{mM}} \\ &\leq 4(1 - R_2(h)/2)^k \left(\|z_0 - Z^0\|_{L^2, a, b} + \sqrt{2}C_1 h^{5/2} \right) + \frac{24\sqrt{2}\gamma h^2 (C_2\sqrt{\gamma} + C_1\sqrt{m})}{m} \\ &\quad + 144h^{3/2} \left(\frac{d\gamma\Theta}{mM} \right)^{1/2} + 1344h^{7/2}C_1\frac{\gamma\sqrt{\Theta}}{m}, \end{aligned}$$

and the first claim follow by rewriting this bound in terms of \tilde{m} , \tilde{M} and $\tilde{\gamma}$. For non-asymptotic Wasserstein results, we simply replace Z^{k-1} with the continuous dynamics initialized at $\tilde{Z}_{k-1} \sim \pi$ be such that $\|\tilde{Z}_{k-1} - z_{k-1}\|_{L^2, a, b} = \mathcal{W}_{2, a, b}(\mu P_h^{k-1}, \pi)$ as in [86][Theorem 23]. We can then apply Lemma H.1 to get the required result. \square

REMARK F.5. *To get the non-asymptotic result to have discretization error which is of order $\mathcal{O}(h^{3/2})$, the gradient approximation needs to be an unbiased estimator of the gradient, without this property the discretization error reduces to order $\mathcal{O}(h)$.*

LEMMA F.6. *Suppose we have two Langevin diffusions governed by (1.1) with synchronously coupled Brownian motion, $(Z_t)_{t \geq 0}$ with potential U satisfying Assumptions 3.28-3.23 and $(\tilde{Z}_t)_{t \geq 0}$ with potential defined by (G.4), a Gaussian approximation of U . We further assume that the diffusions are initialized at their invariant measures and $\gamma \geq \sqrt{8M}$. We then have that*

$$\|Z_t - \tilde{Z}_t\|_{L^2, a, b} \leq e^{-\frac{mt}{16\gamma}} \|Z_0 - \tilde{Z}_0\|_{L^2, a, b} + \frac{dC(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s)}{N_D}.$$

PROOF. We define $(X_t, V_t)_{t \geq 0} := (Z_t)_{t \geq 0}$ to be the diffusion according to (F.1)-(F.2) with potential U and define $(\tilde{X}_t, \tilde{V}_t)_{t \geq 0} := (\tilde{Z}_t)_{t \geq 0}$ to be the diffusion according to (F.1)-(F.2) with potential \tilde{U} , defined by (G.4), and synchronously coupled Brownian motion. By the same argument as Corollary C.8 with the expectations rather than Wasserstein distance we have that for $h > 0$ up to first order in h

$$\begin{aligned} \|Z_{(k+1)h} - \tilde{Z}_{(k+1)h}\|_{L^2, a, b} &\lesssim \left(1 - \frac{mh}{16\gamma}\right) \|Z_{kh} - \tilde{Z}_{kh}\|_{L^2, a, b} \\ &\quad + h \left\| \left(0, \nabla \tilde{U}(\tilde{X}_{kh}) - \nabla U(\tilde{X}_{kh})\right) \right\|_{L^2, a, b} \\ &\leq \left(1 - \frac{mh}{16\gamma}\right) \|Z_{kh} - \tilde{Z}_{kh}\|_{L^2, a, b} + \frac{hM_1^s}{\sqrt{M}} \left\| \tilde{X}_{kh} - x^* \right\|_{L^4}^2, \end{aligned}$$

where the last inequality is due to Lemma E.4 and x^* is the minimizer of U and \tilde{U} . Then due to Proposition C.12 taking the limit as $h \rightarrow 0$

$$\|\tilde{X}_{kh} - x^*\|_{L^4}^2 \leq \frac{2}{m} \left[\sqrt{4 \left(1 - \frac{h\lambda\gamma}{2}\right)^k (\gamma^4 \|\tilde{X}_0 - x^*\|^4 + \|\tilde{V}_0\|^4) + \frac{(6d + 160(1 + \lambda^2))^2}{2\lambda^2}} \right],$$

where we choose $k = t/h$ and define

$$c_u := \frac{2M_1^s}{m\sqrt{M}} \left[\sqrt{4e^{-\frac{t\lambda\gamma}{2}} (\gamma^4 \|\tilde{X}_0 - x^*\|^4 + \|\tilde{V}_0\|^4) + \frac{(6d + 160(1 + \lambda^2))^2}{2\lambda^2}} \right],$$

we have

$$\limsup_{h \rightarrow 0} \frac{\|Z_{t+h} - \tilde{Z}_{t+h}\|_{L^2, a, b} - \|Z_t - \tilde{Z}_t\|_{L^2, a, b}}{h} \leq -\frac{m}{16\gamma} \|Z_t - \tilde{Z}_t\|_{L^2, a, b} + c_u,$$

and all terms are bounded on the right-hand side due to the assumptions on the initial condition therefore due to the Denjoy–Young–Saks theorem we have the upper Dini derivative (upper right-hand derivative) is finite. Hence considering $u : \mathbb{R} \rightarrow \mathbb{R}$ to be solution to the ODE

$$\frac{d}{dt} u(t) = -\frac{m}{16\gamma} u(t) + c_u,$$

with initial condition $u(0) = \|Z_0 - \tilde{Z}_0\|_{L^2, a, b}$ which we can solve exactly. Therefore by the comparison principle for ODEs and Dini derivatives [54][Lemma 3.4] we have

$$\begin{aligned} \|Z_t - \tilde{Z}_t\|_{L^2, a, b} &\leq u(t) \leq e^{-\frac{mt}{16\gamma}} \|Z_0 - \tilde{Z}_0\|_{L^2, a, b} + \frac{16\gamma}{m} c_u \\ &\leq e^{-\frac{mt}{16\gamma}} \|Z_0 - \tilde{Z}_0\|_{L^2, a, b} + \frac{dC(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s)}{N_D}, \end{aligned}$$

as required. \square

PROPOSITION F.7. *Suppose two stochastic gradient UBU chains at coarser and finer discretization levels l and $l+1$, with synchronously coupled Brownian motions $(z_k)_{k \in \mathbb{N}}$ and $(z'_k)_{k \in \mathbb{N}}$ and stepsizes h_l and $h_{l+1} = h_l/2$, satisfying the conditions of Proposition F.4, be such that $z_0 \sim \pi_0$ and $z'_0 \sim \pi'_0$. Then for f satisfying Assumption 3.12 we have the following variance bound*

$$\begin{aligned} \text{Var} (f(z'_k) - f(z_k)) &\leq \mathbb{E} (f(z'_k) - f(z_k))^2 \leq \\ &\left(\exp \left(-\frac{mkh_l}{8\gamma} \right) (\|z'_0 - z_0\|_{L^2, a, b} + \mathcal{W}_{2, a, b}(\pi_0, \pi) + \mathcal{W}_{2, a, b}(\pi'_0, \pi)) \right. \\ &+ 4(1 - R(h_l)/2)^k \left(\mathcal{W}_{2, a, b}(\pi_0, \pi) + \sqrt{2}C_1 h_l^{5/2} \right) \\ &+ 4(1 - R(h_{l+1})/2)^{2k} \left(\mathcal{W}_{2, a, b}(\pi'_0, \pi) + \sqrt{2}C_1 h_{l+1}^{5/2} \right) \\ &\left. + C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s) d^{1/2} N_D^{-3/4} \Theta^{1/2} h_l^{3/2} \right)^2. \end{aligned}$$

PROOF. By following the same argument as Proposition D.3 using Proposition F.4 we have the desired result. \square

PROPOSITION F.8. *Suppose that the assumptions of Proposition F.4 hold for the potential U , $h_0 > 0$ and $\gamma > 0$ and the SVRG stochastic gradient approximation. Assume that*

$$h_0 \leq \frac{C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tau/N_D, N_b)}{N_D^{3/2}}, \quad B \geq \frac{16 \log(2^{3/2}) \tilde{\gamma}}{\tilde{m} h_0 N_D^{1/2}}, \quad B_0 \geq \frac{16 \tilde{\gamma}}{\tilde{m} h_0 N_D^{1/2}} \log \left(\frac{1}{N_D^{9/4} h_0^{3/2}} \right),$$

and the levels are initialized as described in Section E. Then for every $l \geq 1$, $1 \leq k \leq K$, for a test function f which satisfies Assumption 3.12 the UBUBU samples satisfy

$$\begin{aligned} \text{Var} \left(f(z_k^{(l, l+1)}) - f(z_k^{(l, l+1)}) \right) &\leq \mathbb{E} \left[\left(f(z_k^{(l, l+1)}) - f(z_k^{(l, l+1)}) \right)^2 \right] \\ &\leq \mathbb{E} \|z_k^{(l, l+1)} - z_k^{(l, l+1)}\|_{a, b}^2 \\ &\leq C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s, \tau/N_D, N_b) N_D^{5/2} h_l^3 d, \end{aligned}$$

and further

$$(F.13) \quad \text{Var}(D_{l, l+1}) \leq C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s, \tau/N_D, N_b) N_D^{5/2} h_l^3 d.$$

PROOF OF PROPOSITION F.8. Following a similar proof as Corollary D.6. However, we need to be careful with the bounding the distance between z_0 and z'_0 , which is the reason for our construction of the initial conditions in Section E using the OHO scheme. In particular, we wish to have at most $\mathcal{O}(1/N_D)$ distance in initialization. We define $(Z_t)_{t \geq 0}$ to be defined by (1.1) with the potential U and $Z_0 \sim \pi$ such that $\|Z_0 - z'_0\|_{L^2, a, b} = \mathcal{W}_{2, a, b}(\pi, \mu_G)$ are optimally coupled. We define $(Z_t^G)_{t \geq 0}$ to be defined by (1.1) with the potential being a Gaussian approximation of the potential such that $Z_0^G = z'_0 \sim \mu_G$, $(z_t^G)_{t \geq 0}$ to be the OHO scheme with the potential being a Gaussian approximation of the potential such that $z_0^G = z'_0 \sim \mu_G$. We

therefore have $z_B^{\mathcal{G}} = z_0$ and

$$\begin{aligned} \|z_0 - z'_{B/h_{l+1}}\|_{L^2, a, b} &\leq \|z_B^{\mathcal{G}} - Z_B^{\mathcal{G}}\|_{L^2, a, b} + \|Z_B^{\mathcal{G}} - Z_B\|_{L^2, a, b} + \|Z_B - z'_{B/h_{l+1}}\|_{L^2, a, b} \\ &\leq h\sqrt{d}C(\tilde{\gamma}, \tilde{m}, \tilde{M}) + \frac{dC(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s)}{N_D} \\ &\quad + 5\|Z_0 - z'_0\|_{L^2, a, b} + C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s, \tau/N_D, N_b)N_D^{5/2}h_l^3d \\ &\leq \frac{dC(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s, \tau/N_D, N_b)}{N_D}, \end{aligned}$$

where we have used Theorem E.3 for the first term, Lemma F.6 for the second and Proposition F.4 for the third.

We also have the following rough estimates for the Wasserstein distances

$$\mathcal{W}_{2, a, b}(\mu_0^{(l+1)}, \pi) = \mathcal{W}_{2, a, b}(\pi_0, \pi) = \mathcal{W}_{2, a, b}(\mu_G, \pi) \leq \frac{dC(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s)}{N_D},$$

which follow from Proposition E.5, where the estimate of $\mathcal{W}_{2, a, b}(\mu_0^{(l+1)}, \pi)$ along with Proposition F.4 implies $\mathcal{W}_{2, a, b}(\pi'_0, \pi) \leq \frac{dC(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s, \tau/N_D, N_b)}{N_D}$.

We have $(B_0 + Bl)2^l$ burn-in steps at level l , and $(B_0 + B(l+1))2^{l+1}$ burn-in steps at level $l+1$. Using the assumption that $h_0 < \frac{\tilde{m}}{256\tilde{M}\tilde{\gamma}N_D^{1/2}}$, we have for all $i \in \mathbb{N}$ $R(h_i) \geq \frac{mh_i}{8\tilde{\gamma}}$, and using Proposition F.4 we have

$$\begin{aligned} \text{Var} \left(f(z_k^{(l, l+1)}) - f(z_k^{(l, l+1)}) \right) &\leq \mathbb{E} \left[\left(f(z_k^{(l, l+1)}) - f(z_k^{(l, l+1)}) \right)^2 \right] \\ &\leq \left(\exp \left(-\frac{\tilde{m}\sqrt{N_D}(B_0 + lB)h_0}{8\tilde{\gamma}} \right) \left(\|z'_{B/h_{l+1}} - z_0\|_{L^2, a, b} + \mathcal{W}_{2, a, b}(\pi_0, \pi) + \mathcal{W}_{2, a, b}(\pi'_0, \pi) \right) \right. \\ &\quad + 4 \exp \left(-\frac{\tilde{m}\sqrt{N_D}(B_0 + lB)h_0}{16\tilde{\gamma}} \right) \left(\mathcal{W}_{2, a, b}(\pi_0, \pi) + \mathcal{W}_{2, a, b}(\pi'_0, \pi) \right) \\ &\quad \left. + C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s, \tau/N_D, N_b)h_l^{3/2}d^{1/2}N_D^{5/4} \right)^2 \\ &\leq \left(\exp \left(-\frac{m(B_0 + lB)h_0}{16\tilde{\gamma}} \right) \frac{dC(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s, \tau/N_D)}{N_D} + C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s, \tau/N_D, N_b)h_l^{3/2}d^{1/2}N_D^{5/4} \right)^2 \end{aligned}$$

using the assumptions on B_0 and B

$$\leq C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s, \tau/N_D, N_b)N_D^{5/2}h_l^3d.$$

We now use the simple bound

$$\begin{aligned} \text{Var}(D_{l, l+1}) &\leq \mathbb{E}(D_{l, l+1}^2) \leq \max_{1 \leq k \leq K} \mathbb{E} \left[\left(f(z_k^{(l, l+1)}) - f(z_k^{(l, l+1)}) \right)^2 \right] \\ &\leq C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s, \tau/N_D, N_b)N_D^{5/2}h_l^3d \end{aligned}$$

as required. \square

REMARK F.9. *As an alternative, one can consider a coupling with a randomized mid-point scheme, which was utilized in the work of [101] and [14] in the context of kinetic*

Langevin dynamics and Hamiltonian Monte Carlo. This is beyond the scope of the work, and thus we leave this as a direction to consider for future work.

PROPOSITION F.10. *Suppose a full gradient Gaussian approximation OHO chain $(z_k)_{k \in \mathbb{N}}$ at level 0 and a stochastic gradient UBU chain $(z'_k)_{k \in \mathbb{N}}$ at level 1 using the SVRG unbiased estimator, with stepsizes h_0 and $h_1 = \frac{h_0}{2}$, respectively. Further, we assume that they have synchronously coupled Brownian motions and $z_0 \sim \pi_0 = \mu_G$ and $z'_0 \sim \pi'_0$. Assuming the same assumptions as Proposition D.3 for $(z_k)_{k \in \mathbb{N}}$ and Proposition F.4 for $(z'_k)_{k \in \mathbb{N}}$. Then for f satisfying Assumption 3.12 we have the following variance bound*

$$\begin{aligned} \text{Var}(f(z'_k) - f(z_k)) &\leq \mathbb{E} \left[(f(z'_k) - f(z_k))^2 \right] \\ &\leq \left(\exp \left(-\frac{\tilde{m}\sqrt{N_D}kh_0}{8\tilde{\gamma}} \right) (\|z'_0 - z_0\|_{L^2, a, b} + \mathcal{W}_{2, a, b}(\pi'_0, \pi)) \right. \\ &\quad + 4(1 - R(h_1)/2)^{2k} \mathcal{W}_{2, a, b}(\pi'_0, \pi) + \frac{dC(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s)}{N_D} + h_0\sqrt{d}C(\tilde{\gamma}, \tilde{m}, \tilde{M}) \\ &\quad \left. + C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s)d^{1/2}N_D^{-3/4}\Theta^{1/2}h_1^{3/2} \right)^2, \end{aligned}$$

where $R(h_1) = 1 - \sqrt{(1 - c(h_1))^2 + C_0^2 h_0^2}$.

PROOF. By following the same argument as Proposition D.3, but by using Proposition F.4 and Theorem E.3 we have the desired result. However, because level zero and level one are approximating different diffusions, we can't use the contraction results for the continuous dynamics to bound (II), so we consider an alternative argument. For this component, we use Lemma F.6. To bound (I) we use Theorem E.3 and to bound (III) we use Proposition F.4 and we have the required result. \square

PROPOSITION F.11. *Suppose that the assumptions of Proposition F.10 hold for the potential U and $h_0 > 0$. Assume that*

$$h_0 \leq \frac{C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tau/N_D, N_b)}{N_D^{3/2}}, \quad B \geq \frac{16 \log(2^{3/2})\tilde{\gamma}}{\tilde{m}h_0N_D^{1/2}}, \quad B_0 \geq \frac{16\tilde{\gamma}}{\tilde{m}N_D^{1/2}h_0} \log \left(\frac{1}{N_D^{9/4}h_0^{3/2}} \right),$$

$1 < k \leq K$, the levels are initialized as described in Section E and for a function f which satisfies Assumption 3.12 for stochastic gradient UBUBU we have

$$(F.14) \quad \text{Var}(D_{0,1}) \leq \frac{C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s)d^2}{N_D^2} + C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s, \tau/N_D, N_b)N_D^{5/2}h_0^3d.$$

PROOF. Following the same proof as Proposition F.8 using the results of Proposition F.10. \square

F.2. Variance of $S(c_R)$.

THEOREM 3.24. *Considering UBUBU with stochastic gradients, suppose that Assumptions 3.12, 3.21, 3.22, 3.23 hold, and in addition $\gamma \geq \sqrt{8M}$,*

$$h_0 \leq \frac{C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tau/N_D, N_b)}{N_D^{3/2}}, \quad B \geq \frac{16 \log(2^{3/2}) \tilde{\gamma}}{\tilde{m} h_0 N_D^{1/2}}, \quad B_0 \geq \frac{16 \tilde{\gamma}}{\tilde{m} h_0 N_D^{1/2}} \log \left(\frac{1}{N_D^{9/4} h_0^{3/2}} \right).$$

Suppose that $c_R \in [0, 1)$ and $2 < \phi_N < 8$. Then for any $N \geq 1$, the UBUBU estimator $S(c_R)$ has finite expected computational cost, $\mathbb{E}S(c_R) = \pi(f)$, and it has finite variance. Moreover, it satisfies a CLT as $N \rightarrow \infty$, and the asymptotic variance σ_S^2 defined in (3.8) can be bounded as

$$\sigma_S^2 \leq \frac{1}{\tilde{m} N_D K} + C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s, \tau/N_D, N_b, \phi_N) \frac{d^2}{c_N N_D^2}.$$

PROOF. By Propositions F.8 and F.11, we have for $l \geq 1$ that

$$\mathbb{E}(D_{l,l+1}^2) \leq C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s, \tau/N_D, N_b) h_l^3 d N_D^{5/2} \leq V_{D_1} \phi_{D_1}^{-l},$$

for $V_{D_1} = C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s, \tau/N_D, N_b) h_0^3 d N_D^{5/2}$ and $\phi_{D_1} = 8$. For $l = 0$ we have

$$\mathbb{E}(D_{l,l+1}^2) \leq C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s, \tau/N_D, N_b) \frac{d^2}{N_D^2} \leq V_{D_2} \phi_{D_2}^{-l},$$

for $V_{D_2} = C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s, \tau/N_D, N_b) \frac{d^2}{N_D^2}$ and $\phi_{D_2} = 2$.

Due to the fact that for D_0 we take K i.i.d. Gaussian samples, it is easy to show using the Gaussian Poincaré inequality that

$$\text{Var}(D_0) \leq \frac{1}{\tilde{m} N_D K}$$

The computational cost at levels $l \geq 1$ satisfies the assumptions of Proposition 3.4, so if we fix $2 < \phi_N < 8$, all assumptions of this proposition are satisfied. Hence $S(c_R)$ is an unbiased estimator with finite variance and computational cost.

For the asymptotic variance using (3.8), and the above estimates we have

$$\begin{aligned} \sigma_S^2 &\leq \frac{1}{\tilde{m} N_D} + C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s, \tau/N_D, N_b) \frac{d^2}{c_N N_D^2} + \sum_{l=1}^{\infty} \frac{V_{D_1} \phi_{D_1}^{-l}}{c_N \phi_N^{-l}} \\ &\leq \frac{1}{\tilde{m} N_D} + C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s, \tau/N_D, N_b, \phi_N) \left(\frac{d^2}{c_N N_D^2} + \frac{h_0^3 d N_D^{5/2}}{c_N} \right), \end{aligned}$$

if we choose h_0 to be of the order $\mathcal{O}(1/N_D^{3/2})$ then we have

$$\leq \frac{1}{\tilde{m} N_D} + C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s, \tau/N_D, N_b, \phi_N) \frac{d^2}{c_N N_D^2},$$

as required. □

APPENDIX G: VARIANCE BOUNDS FOR UBUBU ESTIMATOR WITH APPROXIMATE GRADIENTS

One can also approximate the gradient in a cheap way, which has bias, but such that the bias tends to zero with the stepsize. The multilevel estimator will still be an unbiased estimator from the target measure.

For convex potentials, we can approximate the gradient with the Hessian at the minimizer by

$$(G.1) \quad \mathcal{Q}(x | \hat{x}) = \nabla U(\hat{x}) + \nabla^2 U(x^*)(x - \hat{x}).$$

Despite the fact that this estimator is biased, in our multilevel approach, the overall estimator will still be unbiased.

As before, the updates in $(\bar{x}_k, \bar{v}_k)_{k \geq 0}$ form a \mathcal{BU} step, so they can be expressed as

$$(G.2) \quad \bar{x}_{k+1} = \bar{x}_k + \frac{1 - \eta^2}{\gamma} (\bar{v}_k - h\mathcal{Q}(\bar{x}_k | \bar{x}_{L(k)})) + \sqrt{\frac{2}{\gamma}} \left(\mathcal{Z}^{(1)}(h, \xi_{k+1}^{(1)}) - \mathcal{Z}^{(2)}(h, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}) \right),$$

$$(G.3) \quad \bar{v}_{k+1} = \eta^2 (\bar{v}_k - h\mathcal{Q}(\bar{x}_k | \bar{x}_{L(k)})) + \sqrt{2\gamma} \mathcal{Z}^{(2)}(h, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)}),$$

where $\hat{x}_k = \bar{x}_{L(k)}$ and $L(k) = \tau \lfloor k/\tau \rfloor$.

It turns out that at level 0, it can be advantageous to simply use the gradients of the Gaussian approximation, and never compute gradients of U . This corresponds to gradient approximation of the form

$$(G.4) \quad \mathcal{Q}^*(x) = \mathcal{Q}(x | x^*) = \nabla^2 U(x^*)(x - x^*),$$

and so (G.2)-(G.3) holds with $\hat{x}_k = x^*$ for every $k \geq 0$ in this case.

G.1. Non-asymptotic guarantees.

LEMMA G.1. *Considering iterates $(x_k, v_k, \bar{x}_k)_{k \in \mathbb{N}}$ of approximate gradient UBU, with epoch length τ and gradient approximation \mathcal{Q} given by (E.6), and initial condition $(x_0, v_0) \in \mathbb{R}^{2d}$, we have the property*

$$\mathbb{E} \|\nabla U(\bar{x}_k) - \mathcal{Q}(\bar{x}_k | \bar{x}_{L(k)})\|^2 \leq \tilde{M}_1^2 N_D^2 (\tau - 1)^2 \max_{j \leq k} \|\bar{x}_j - x^*\|_{L^4}^2 \cdot \max_{j < k} \|\bar{x}_{j+1} - \bar{x}_j\|_{L^4}^2,$$

and we also have

$$\mathbb{E} \|\nabla U(\bar{x}_k) - \mathcal{Q}(\bar{x}_k | x^*)\|^2 \leq \tilde{M}_1^2 N_D^2 \|\bar{x}_k - x^*\|_{L^4}^4.$$

PROOF. Let the last full gradient evaluation be at iteration $L(k)$, then

$$\begin{aligned} \mathbb{E} \|\nabla U(\bar{x}_k) - \mathcal{Q}(\bar{x}_k | \bar{x}_{L(k)})\|^2 &= \mathbb{E} \|\nabla U(\bar{x}_k) - \nabla U(\bar{x}_{L(k)}) - \nabla^2 U(x^*)(\bar{x}_k - \bar{x}_{L(k)})\|^2 \\ &= \mathbb{E} \left\| \left(\int_{t=0}^1 \nabla^2 U(\bar{x}_k + t(\bar{x}_{L(k)} - \bar{x}_k)) dt - \nabla^2 U(x^*) \right) (\bar{x}_k - \bar{x}_{L(k)}) \right\|^2 \\ &\leq \tilde{M}_1^2 N_D^2 \mathbb{E} \left(\left(\int_{t=0}^1 \|\bar{x}_k + t(\bar{x}_{L(k)} - \bar{x}_k) - x^*\| \right)^2 \|\bar{x}_k - \bar{x}_{L(k)}\|^2 \right) \\ &\leq \frac{\tilde{M}_1^2 N_D^2}{2} (\|\bar{x}_k - x^*\|_{L^4}^2 + \|\bar{x}_{L(k)} - x^*\|_{L^4}^2) \|\bar{x}_k - \bar{x}_{L(k)}\|_{L^4}^2 \\ &\leq \tilde{M}_1^2 N_D^2 (\tau - 1)^2 \max_{j \leq k} \|\bar{x}_j - x^*\|_{L^4}^2 \cdot \max_{j < k} \|\bar{x}_{j+1} - \bar{x}_j\|_{L^4}^2. \end{aligned}$$

The second claim follows by Taylor expansion. \square

Now, we are going to bound the terms $\|\bar{x}_j - x^*\|_{L^4}$ and $\|\bar{x}_{j+1} - \bar{x}_j\|_{L^4}$.

LEMMA G.2. *When using exact gradients, we have*

$$\|\bar{x}_{k+1} - \bar{x}_k\|_{L^4} \leq 2h\sqrt{M}\|z_k - z^*\|_{L^4,a,b} + 2h\sqrt{d}.$$

With approximate gradients, we have

$$\|\bar{x}_{k+1} - \bar{x}_k\|_{L^4} \leq 2h\sqrt{M} \left(\|\bar{z}_k - z^*\|_{L^4,a,b} + \sqrt{2}(1 + M/m)\|\bar{z}_{L(k)} - z^*\|_{L^4,a,b} \right) + 2h\sqrt{d}.$$

PROOF. In the case of exact gradients, we have

$$\begin{aligned} \bar{x}_{k+1} &= \bar{x}_k + \frac{1 - \eta^2}{\gamma} (\bar{v}_k - h\nabla U(\bar{x}_k)) + \sqrt{\frac{2}{\gamma}} \left(\mathcal{Z}^{(1)} \left(h, \xi_{k+1}^{(1)} \right) - \mathcal{Z}^{(2)} \left(h, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)} \right) \right), \\ \bar{v}_{k+1} &= \eta^2 (\bar{v}_k - h\nabla U(\bar{x}_k)) + \sqrt{2\gamma} \mathcal{Z}^{(2)} \left(h, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)} \right), \end{aligned}$$

so using $\|\nabla U(x_k)\| \leq \|x_k - x^*\|$, Lemma H.4, and the fact that $\xi \sim N(0, I_d)$ satisfies that $\mathbb{E}(\|\xi\|^4) \leq 3d^2$, we have that for $h \leq 1/\sqrt{M}$, $\gamma \geq \sqrt{8M}$,

$$\|\bar{x}_{k+1} - \bar{x}_k\|_{L^4} \leq h\|\bar{v}_k\|_{L^4} + h^2M\|x_k - x^*\|_{L^4} + 3^{1/4}2^{1/2}h\sqrt{d} \leq 2h\sqrt{M}\|z_k - z^*\|_{L^4,a,b} + 2h\sqrt{d}.$$

For approximate gradients, we have

(G.5)

$$\bar{x}_{k+1} = \bar{x}_k + \frac{1 - \eta^2}{\gamma} (\bar{v}_k - h\mathcal{Q}(\bar{x}_k|\bar{x}_{L(k)})) + \sqrt{\frac{2}{\gamma}} \left(\mathcal{Z}^{(1)} \left(h, \xi_{k+1}^{(1)} \right) - \mathcal{Z}^{(2)} \left(h, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)} \right) \right),$$

(G.6)

$$\bar{v}_{k+1} = \eta^2 (\bar{v}_k - h\mathcal{Q}(\bar{x}_k|\bar{x}_{L(k)})) + \sqrt{2\gamma} \mathcal{Z}^{(2)} \left(h, \xi_{k+1}^{(1)}, \xi_{k+1}^{(2)} \right),$$

Let $\tilde{x}_k = x_{L(k)} - (\nabla^2 U(x^*))^{-1} \nabla U(\bar{x}_{L(k)})$, and $\tilde{U}_k(x) = \frac{1}{2}(x - \tilde{x}_k)^T \nabla^2 U(x^*)(x - \tilde{x}_k)$. Then the approximate gradient step is the same as an exact gradient step with respect to the potential \tilde{U}_k . So we have by the result for exact gradients that for approximate gradients,

$$\begin{aligned} \|\bar{x}_{k+1} - \bar{x}_k\|_{L^4} &\leq h\|\bar{v}_k\|_{L^4} + h^2M\|x_k - \tilde{x}_k\|_{L^4} + 3^{1/4}2^{1/2}hd \\ &\leq 2h\sqrt{M} \left\| \bar{z}_k - \begin{pmatrix} \tilde{x}_k \\ 0_d \end{pmatrix} \right\|_{L^4,a,b} + 2h\sqrt{d}. \end{aligned}$$

Here using the triangle inequality, we have

$$\begin{aligned} \left\| z_k - \begin{pmatrix} \tilde{x}_k \\ 0_d \end{pmatrix} \right\|_{L^4,a,b} &\leq \|\bar{z}_k - z^*\|_{L^4,a,b} + \left\| \begin{pmatrix} \tilde{x}_k \\ 0_d \end{pmatrix} - z^* \right\|_{L^4,a,b} \\ &\leq \|\bar{z}_k - z^*\|_{L^4,a,b} + \sqrt{2}(1 + M/m)\|\bar{z}_{L(k)} - z^*\|_{L^4,a,b}, \end{aligned}$$

hence

$$\|\bar{x}_{k+1} - \bar{x}_k\|_{L^4} \leq 2h\sqrt{M} \left(\|\bar{z}_k - z^*\|_{L^4,a,b} + \sqrt{2}(1 + M/m)\|\bar{z}_{L(k)} - z^*\|_{L^4,a,b} \right) + 2h\sqrt{d}. \quad \square$$

We still need to control the evolution of $\|\bar{z}_k - z^*\|_{L^4,a,b}$. As before in (C.16), we define the Lyapunov function \mathcal{V} as

$$\mathcal{V}(x, v) = U(x) - U(x^*) + \frac{1}{4}\gamma^2 (\|x - x^* + \gamma^{-1}v\|^2 + \|\gamma^{-1}v\|^2 - \lambda\|x - x^*\|^2).$$

The following lemma establishes some useful properties about this.

LEMMA G.3. *Suppose that $\gamma \geq \sqrt{8M}$, and that Assumptions 3.6 and 3.7 hold for U . Then for any $z = (x, v) \in \Lambda$, $\mathcal{V}(x, v) \geq 0$, and*

$$(G.7) \quad \mathcal{V}^{1/2}(x, v) \geq \frac{1}{8}(\gamma\|x - x^*\| + \|v\|) \geq \frac{\sqrt{M}}{8}\|z - z^*\|_{a,b}.$$

Moreover, $\mathcal{V}^{1/2}$ is 8γ -Lipschitz with respect to the $\|\cdot\|_{a,b}$ norm.

PROOF. Using the strong convexity of U ,

$$\begin{aligned} \mathcal{V}(x, v) &= U(x) - U(x^*) + \frac{1}{4}\gamma^2 (\|x - x^* + \gamma^{-1}v\|^2 + \|\gamma^{-1}v\|^2 - \lambda\|x - x^*\|^2) \\ &\geq \frac{m}{2}\|x - x^*\|^2 + \frac{1}{4}\gamma^2 ((1 - \lambda)\|x - x^*\|^2 + 2\gamma^{-2}\|v\|^2 + 2\langle x - x^*, \gamma^{-1}v \rangle) \end{aligned}$$

using that $|2\langle x - x^*, \gamma^{-1}v \rangle| \leq \frac{\|x - x^*\|^2}{c} + c\|\gamma^{-1}v\|^2$ with $c = 8/5$, and that $0 < \lambda \leq \frac{1}{4}$,

$$\geq \frac{1}{4}\gamma^2 \left(\frac{1}{8}\|x - x^*\|^2 + \frac{2}{5}\|\gamma^{-1}v\|^2 \right) \geq \frac{1}{64}(\gamma\|x - x^*\| + \|v\|)^2 \geq \frac{M}{64}\|z - z^*\|_{a,b}^2,$$

and our first claim follows by taking square-root.

For the second claim, note that $\nabla \mathcal{V}^{1/2}(x, v) = \frac{1}{2} \frac{\nabla \mathcal{V}}{\mathcal{V}^{1/2}(x, v)}$. Here

$$\begin{aligned} \nabla_x \mathcal{V}(x, v) &= \nabla U(x) + \frac{1}{2}\gamma^2((1 - \lambda)(x - x^*) + \gamma^{-1}v), \\ \|\nabla_x \mathcal{V}(x, v)\| &\leq \left(M + \frac{\gamma^2(1 - \lambda)}{2} \right) \|x - x^*\| + \frac{\gamma}{2}\|v\| \leq \gamma^2\|x - x^*\| + \frac{\gamma}{2}\|v\|, \\ \nabla_v \mathcal{V}(x, v) &= \frac{1}{2}\gamma^2(\gamma^{-1}((x - x^*) + \gamma^{-1}v) + \gamma^{-2}v) \\ \|\nabla_v \mathcal{V}(x, v)\| &\leq \frac{\gamma}{2}\|x - x^*\| + \|v\|, \end{aligned}$$

so we have

$$\begin{aligned} \|\nabla_x \mathcal{V}^{1/2}(x, v)\| &= \frac{\|\nabla_x \mathcal{V}(x, v)\|}{2\mathcal{V}^{1/2}(x, v)} \leq 4\gamma, \\ \|\nabla_v \mathcal{V}^{1/2}(x, v)\| &= \frac{\|\nabla_v \mathcal{V}(x, v)\|}{2\mathcal{V}^{1/2}(x, v)} \leq 4, \end{aligned}$$

and since $\gamma \geq \sqrt{8M}$, for any $(x, v), (x', v') \in \Lambda$, we have

$$\begin{aligned} |\mathcal{V}^{1/2}(x, v) - \mathcal{V}^{1/2}(x', v')| &= \left\langle \int_{t=0}^1 \nabla \mathcal{V}^{1/2}(x + t(x' - x), v + t(v' - v)) dt, (x' - x, v' - v) \right\rangle \\ &\leq 4\sqrt{2}\gamma\|(x, v) - (x', v')\|_{a,0} \leq 8\gamma\|(x, v) - (x', v')\|_{a,b}. \end{aligned}$$

□

As previously, let $\lambda = \min\left(\frac{1}{4}, \frac{m}{\gamma^2}\right)$, and $c_4(h) = h\lambda\gamma - 8h^2\gamma^2(4 + \lambda)$. By (C.18), for the exact gradient scheme, if $c_4(h) < \frac{1}{2}$, we have

$$\mathbb{E} [\mathcal{V}(\bar{x}_{k+1}, \bar{v}_{k+1})^2 | \bar{x}_k, \bar{v}_k] \leq \left(1 - \frac{c_4(h)}{2}\right) \mathcal{V}^2(\bar{x}_k, \bar{v}_k) + \frac{(6h\gamma d + 160h\gamma(1 + \lambda^2))^2}{4c_4(h)} + 24h^2\gamma^2 d^2.$$

Let $C_{\mathcal{V}}(h) := \frac{(6h\gamma d + 160h\gamma(1+\lambda^2))^2}{4c_4(h)} + 24h^2\gamma^2 d^2$, then by applying this j times, we have

$$(G.8) \quad \mathbb{E} [\mathcal{V}(\bar{x}_{k+j}, \bar{v}_{k+j})^2 \mid \bar{x}_k, \bar{v}_k] \leq \left(1 - \frac{c_4(h)}{2}\right)^j \mathcal{V}^2(\bar{x}_k, \bar{v}_k) + C_{\mathcal{V}}(h) \frac{1 - (c_4(h)/2)^j}{1 - c_4(h)/2}.$$

Now we are going to generalise this result to the approximate gradient scheme.

LEMMA G.4. *Consider iterates $(x_k, v_k, \bar{x}_k)_{k \in \mathbb{N}}$ of approximate gradient UBU, with epoch length τ and gradient approximation \mathcal{Q} given by (E.6), and initial condition $(x_0, v_0) \in \mathbb{R}^{2d}$. Suppose that $L(k) = k$ (i.e. k is divisible by τ), and $c_4(h) > 0$, then for any $1 \leq j \leq \tau$, we have*

$$\begin{aligned} \|\mathcal{V}^{1/2}(\bar{z}_{k+j})\|_{L^4} &\leq \left[\left(1 - \frac{c_4(h)}{2}\right)^j \|\mathcal{V}^{1/2}(\bar{z}_k)\|_{L^4}^4 + C_{\mathcal{V}}(h)j \right]^{1/4} \\ &+ 8\gamma \left(48h^2\sqrt{M} \cdot j^2 (\|\mathcal{V}^{1/2}(\bar{z}_k)\|_{L^4}^4 + C_{\mathcal{V}}(h)j)^{1/4} + 6h^2j^2\sqrt{dM}\right). \end{aligned}$$

PROOF. We use an interpolation argument, inspired by the interpolation to independence coupling in [23]. For $0 \leq i \leq j$, let $\bar{z}_{k+j}^{(i)} = (\bar{x}_{k+j}^{(i)}, \bar{v}_{k+j}^{(i)})$ be defined by performing $j-i$ BU steps with exact gradients starting from (\bar{x}_k, \bar{v}_k) according to (C.14)-(C.15), followed by i steps with approximate gradients according to (G.2)-(G.3). Then we have $\bar{z}_{k+j} = \bar{z}_{k+j}^{(j)}$, and $\bar{z}_{k+j}^{(0)}$ corresponds to taking j steps with exact gradients. By the triangle inequality, we have

$$\|\bar{z}_{k+j} - \bar{z}_{k+j}^{(0)}\|_{a,b} \leq \sum_{i=0}^{j-1} \|\bar{z}_{k+j}^{(i+1)} - \bar{z}_{k+j}^{(i)}\|_{a,b}.$$

Using Proposition C.6, we have a contraction according to $\|\cdot\|_{a,b}$ with synchronous coupling when using the approximate gradients (because these are exact gradients with respect to a Gaussian), so we have

$$\|\bar{z}_{k+j}^{(i+1)} - \bar{z}_{k+j}^{(i)}\|_{L^4, a,b} \leq \|\bar{z}_{k+i+1}^{(1)} - \bar{z}_{k+i+1}^{(0)}\|_{L^4, a,b},$$

which is the one-step error of the approximate gradient scheme versus the exact gradient scheme.

$$\begin{aligned} &\|\bar{z}_{k+i+1}^{(1)} - \bar{z}_{k+i+1}^{(0)}\|_{L^4, a,b} \\ &= \left\| \left(\frac{(1-\eta^2)h}{\gamma} (\mathcal{Q}(\bar{x}_{k+i}^{(0)} \mid \bar{x}_k) - \nabla U(\bar{x}_{k+i}^{(0)})), \eta^2 h (\mathcal{Q}(\bar{x}_{k+i}^{(0)} \mid \bar{x}_k) - \nabla U(\bar{x}_{k+i}^{(0)})) \right) \right\|_{L^4, a,b} \\ &\leq \sqrt{2} \|\bar{x}_{k+i}^{(0)} - \bar{x}_k\|_{L^4} \cdot M \left(h^2 + \frac{h}{\sqrt{M}} \right). \end{aligned}$$

So, for $h < \frac{1}{\sqrt{M}}$, we have

$$\begin{aligned} \|\bar{z}_{k+j} - \bar{z}_{k+j}^{(0)}\|_{L^4, a,b} &\leq 3h\sqrt{M} \sum_{i=0}^{j-1} \|\bar{x}_{k+i}^{(0)} - \bar{x}_k\|_{L^4} \\ &\leq 3h\sqrt{M} \cdot j \cdot \sum_{0 \leq i \leq j-1} \|\bar{x}_{k+i+1}^{(0)} - \bar{x}_{k+i}^{(0)}\|_{L^4} \end{aligned}$$

using Lemma G.2,

$$\leq 6h^2 M \cdot j \cdot \sum_{0 \leq i \leq j-1} \|\bar{z}_{k+i}^{(0)} - z^*\|_{L^4, a, b} + 6h^2 j^2 \sqrt{dM}$$

using Lemma G.3,

$$\leq 48h^2 \sqrt{M} \cdot j \cdot \sum_{0 \leq i \leq j-1} \|\mathcal{V}^{1/2}(\bar{z}_{k+i}^{(0)})\|_{L^4} + 6h^2 j^2 \sqrt{dM}$$

using (G.8)

$$\leq 48h^2 \sqrt{M} \cdot j^2 (\|\mathcal{V}^{1/2}(\bar{z}_k)\|_{L^4}^4 + C_{\mathcal{V}}(h)j)^{1/4} + 6h^2 j^2 \sqrt{dM}.$$

We do know that

$$\|\mathcal{V}^{1/2}(\bar{z}_{k+j}^{(0)})\|_{L^4}^4 \leq \left(1 - \frac{c_4(h)}{2}\right)^j \|\mathcal{V}^{1/2}(\bar{z}_k)\|_{L^4}^4 + C_{\mathcal{V}}(h),$$

so by the 8γ -Lipschitz property of $\mathcal{V}^{1/2}$ in $\|\cdot\|_{a, b}$ by Lemma G.3, we have

$$\begin{aligned} \|\mathcal{V}^{1/2}(\bar{z}_{k+j})\|_{L^4} &\leq \left[\left(1 - \frac{c_4(h)}{2}\right)^j \|\mathcal{V}^{1/2}(\bar{z}_k)\|_{L^4}^4 + C_{\mathcal{V}}(h)j \right]^{1/4} \\ &+ 8\gamma \left(48h^2 \sqrt{M} \cdot j^2 (\|\mathcal{V}^{1/2}(\bar{z}_k)\|_{L^4}^4 + C_{\mathcal{V}}(h)j)^{1/4} + 6h^2 j^2 \sqrt{dM} \right). \end{aligned}$$

□

COROLLARY G.5. Consider iterates $(x_k, v_k, \bar{x}_k)_{k \in \mathbb{N}}$ of approximate gradient UBU, with epoch length τ and gradient approximation \mathcal{Q} given by (E.6) approximating a potential U which satisfies Assumptions 3.28 and 3.22 with $z_0 \sim \mu_G$. Assume that

$$h < \min \left\{ 2/\tau\gamma, 1, 1/2\gamma, \frac{\lambda\tau}{64(432\sqrt{M}\tau^2 + \gamma(1+\lambda)\tau)} \right\}, \quad \gamma \geq \sqrt{M},$$

then

$$\|\bar{z}_k - z^*\|_{L^4, a, b} \leq \frac{C(\tilde{\gamma}, \tilde{m}, \tilde{M})\sqrt{d}}{\sqrt{N_D}}.$$

PROOF. If we define $b_k := \|\mathcal{V}^{1/2}(\bar{z}_{\tau k})\|_{L^4}$, then for $\gamma \geq \sqrt{8M}$ and $h < \frac{8}{\tau\gamma}$, we have $c_4(h) \leq 2/\tau$ (here $c_4(h)$ and λ are defined as in (C.13) and (C.12)), and so

$$\begin{aligned} b_{k+1} &\leq \left[\left(1 - \frac{c_4(h)}{2}\right)^\tau b_k^4 + C_{\mathcal{V}}(h)\tau \right]^{1/4} + 384h^2 \gamma \sqrt{M} \tau^2 (b_k^4 + C_{\mathcal{V}}(h)\tau)^{1/4} + 48\gamma \sqrt{M} h^2 \tau^2 \sqrt{d} \\ \text{(G.9)} \quad &\leq \left[\left(1 - \frac{c_4(h)\tau}{4}\right) b_k^4 + C_{\mathcal{V}}(h)\tau \right]^{1/4} + 384h^2 \gamma \sqrt{M} \tau^2 (b_k^4 + C_{\mathcal{V}}(h)\tau)^{1/4} + 48\gamma \sqrt{M} h^2 \tau^2 \sqrt{d}. \end{aligned}$$

Using this, for $b_k < \max \left\{ \left(\frac{8C_{\mathcal{V}}(h)}{c_4(h)}\right)^{1/4}, \sqrt{d} \right\}$ we have that

$$\text{(G.10)} \quad b_{k+1} \leq (1 + 384h^2 \gamma \sqrt{M} \tau^2) \left[\frac{8C_{\mathcal{V}}(h)}{c_4(h)} + d^2 + C_{\mathcal{V}}(h)\tau \right]^{1/4} + 48\gamma \sqrt{M} h^2 \tau^2 \sqrt{d}.$$

For $b_k \geq \max \left\{ \left(\frac{8C_V(h)}{c_4(h)} \right)^{1/4}, \sqrt{d} \right\}$, using that $(1+x)^{1/4} \leq 1 + \frac{x}{4}$ for $x \in [-1, \infty)$, we have

$$\begin{aligned} b_{k+1} &\leq b_k \left[\left[\left(1 - \frac{c_4(h)\tau}{4} \right) + \frac{C_V(h)\tau}{b_k^4} \right]^{1/4} + 384h^2\gamma\sqrt{M}\tau^2 \left(1 + \frac{C_V(h)\tau}{b_k^4} \right)^{1/4} + \frac{48\gamma\sqrt{M}h^2\tau^2\sqrt{d}}{b_k} \right] \\ &\leq \left[1 - \frac{c_4(h)\tau}{32} + 432h^2\gamma\sqrt{M}\tau^2 \right] b_k \end{aligned}$$

using the definition $c_4(h) = h\lambda\gamma - 8h^2\gamma^2(4 + \lambda)$

$$\leq \left[1 - h\frac{\lambda\gamma\tau}{32} + h^2(432\gamma\sqrt{M}\tau^2 + \gamma^2(1 + \lambda)\tau) \right] b_k$$

using the assumption $h \leq \frac{\lambda\gamma\tau}{64(432\gamma\sqrt{M}\tau^2 + \gamma^2(1 + \lambda)\tau)}$

$$\leq \left[1 - h\frac{\lambda\gamma\tau}{64} \right] b_k.$$

Therefore we have that for all $k \in \mathbb{N}$

$$b_k \leq \left[1 - h\frac{\lambda\gamma\tau}{64} \right]^k b_0 + (1 + 384h^2\gamma\sqrt{M}\tau^2) \left[\frac{8C_V(h)}{c_4(h)} + d^2 + C_V(h)\tau \right]^{1/4} + 48\gamma\sqrt{M}h^2\tau^2\sqrt{d}.$$

Now considering $b_{k,j} := \|\mathcal{V}^{1/2}(\bar{z}_{\tau k+j})\|_{L^4}$ we have that by the same argument replacing τ by j that

$$b_{k,j} \leq \left[1 - h\frac{\lambda\gamma j}{64} \right]^j b_k + (1 + 384h^2\gamma\sqrt{M}j^2) \left[\frac{8C_V(h)}{c_4(h)} + d^2 + C_V(h)j \right]^{1/4} + 48\gamma\sqrt{M}h^2j^2\sqrt{d}.$$

Therefore considering the iterates of the approximate gradient UBU scheme we have

$$\frac{\sqrt{\tilde{M}N_D}}{8} \|\bar{z}_k - z^*\|_{L^4,a,b} \leq C(\tilde{\gamma}, \tilde{m}, \tilde{M}) \left(\|\mathcal{V}^{1/2}(\bar{z}_0)\|_{L^4} + h\tau\sqrt{N_D} + \sqrt{d} \right)$$

and therefore using Lemma H.2 for the initial distribution we have

$$\|\bar{z}_k - z^*\|_{L^4,a,b} \leq \frac{C(\tilde{\gamma}, \tilde{m}, \tilde{M})\sqrt{d}}{\sqrt{N_D}}.$$

□

PROPOSITION G.6. *For an approximate gradient UBU integrator with iterates $(z_k)_{k \in \mathbb{N}}$, transition kernel P_h and a potential U satisfying Assumptions 3.6-3.10 and $z_0 \sim \mu_G$, where we approximate the gradient using the gradient approximation \mathcal{Q} given in (E.6). Consider the continuous solution to (1.1) $(Z_t)_{t \geq 0}$, and define $Z^k := Z_{kh}$ for $k \in \mathbb{N}$, where $Z^0 \sim \pi$ is initialized at the invariant measure with synchronously coupled Brownian motion to $(z_k)_{k \in \mathbb{N}}$, then for all*

$$h < \min \left\{ 2/\tau\gamma, 1, 1/2\gamma, \frac{\lambda\tau}{64(432\sqrt{M}\tau^2 + \gamma(1 + \lambda)\tau)} \right\}, \quad \gamma \geq \sqrt{8M},$$

$k, l \in \mathbb{N}$ such that $k > l$

$$\|z_k - Z^k\|_{L^2,a,b} \leq (1 - c(h))^{k-l} \|z_l - Z^l\|_{L^2,a,b} + \frac{h \left((\tau - 1)\sqrt{d} + \sqrt{N_D} \right) C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s)\sqrt{d}}{\sqrt{N_D}},$$

and further

$$\mathcal{W}_{2,a,b}(\mu_G P_h^k, \pi) \leq (1 - c(h))^{k-l} \mathcal{W}_{2,a,b}(\mu_G P_h^l, \pi) + \frac{h \left((\tau - 1)\sqrt{d} + \sqrt{N_D d} \right) C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s) \sqrt{d}}{\sqrt{N_D}}.$$

PROOF. Firstly, we introduce the notation $z_k^h := (x_k^h, v_k^h) := \psi_h \left(z_k, h, (W_{t'})_{t'=kh}^{(k+1)h} \right)$ for all $k \in \mathbb{N}$, an iteration of the full gradient scheme with stepsize $h > 0$ and initial point z_k with synchronously coupled Brownian motion to the approximate gradient scheme. We split up the difference in the following way

$$\begin{aligned} \|z_k - Z^k\|_{L^2,a,b} &\leq \left\| z_k - z_{k-1}^h \right\|_{L^2,a,b} + \|z_{k-1}^h - Z^k\|_{L^2,a,b} \\ &\leq \|z_k - z_{k-1}^h\|_{L^2,a,b} + \|\psi(Z^{k-1}, h, (W_{t'})_{t'=(k-1)h}^{kh}) - Z^k\|_{L^2,a,b} \\ &\quad + \|z_{k-1}^h - \psi(Z^{k-1}, h, (W_{t'})_{t'=(k-1)h}^{kh})\|_{L^2,a,b} \\ &= \text{(I)'} + \text{(II)'} + \text{(III)'}. \end{aligned}$$

We have by Corollary G.5, Lemma G.2 and Lemma G.1 that

$$\begin{aligned} \text{(I)'} &\leq \frac{\sqrt{2}h}{\sqrt{M}} M_1^s (\tau - 1) \max_{j \leq k-1} \|\bar{x}_j - x^*\|_{L^4} \cdot \max_{j < k-1} \|\bar{x}_{j+1} - \bar{x}_j\|_{L^4} \\ &\leq (\tau - 1) h^2 C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s) d, \end{aligned}$$

by the discretization results in Appendix H

$$\text{(II)'} \leq \tilde{C} h^2 \leq \frac{3}{7} \sqrt{d} \left(\sqrt{M} + \gamma \right) h^2$$

and

$$\text{(III)'} \leq (1 - c(h)) \|z_{k-1} - Z^{k-1}\|_{L^2,a,b},$$

where the inequality for (II)' is shown in Appendix H. Therefore going from local to global we have that

$$\begin{aligned} \|z_k - Z^k\|_{L^2,a,b} &\leq (1 - c(h))^{k-l} \|z_l - Z^l\|_{L^2,a,b} + \frac{h^2 \left((\tau - 1)d + \sqrt{N_D d} \right) C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s)}{c(h)} \\ &= (1 - c(h))^{k-l} \|z_l - Z^l\|_{L^2,a,b} + \frac{h \left((\tau - 1)d + \sqrt{N_D d} \right) C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s)}{\sqrt{N_D}}. \end{aligned}$$

For non-asymptotic Wasserstein results, we simply replace Z^{k-1} with the continuous dynamics initialized at $\tilde{Z}_{k-1} \sim \pi$ be such that $\|\tilde{Z}_{k-1} - z_{k-1}\|_{L^2,a,b} = \mathcal{W}_{2,a,b}(\mu P_h^{k-1}, \pi)$ as in [86][Theorem 23]. We can then apply Lemma H.1 to get the required result. \square

G.2. Variance bound of $D_{l,l+1}$.

PROPOSITION G.7. *Suppose two approximate gradient UBU chains at coarser and finer discretization levels l and $l+1$, with synchronously coupled Brownian motions $(z_k)_{k \in \mathbb{N}}$ and $(z'_k)_{k \in \mathbb{N}}$ and stepsizes h_l and $h_{l+1} = h_l/2$, satisfying the conditions of Proposition G.6, be such that $z_0 \sim \pi_0 = \mu_G$ and $z'_0 \sim \pi'_0 = \mu_G(P_{h_{l+1}}^A)^{B/h_{l+1}}$. Then for f satisfying Assumption*

3.12 we have the following variance bound

$$\begin{aligned} \text{Var}(f(z'_k) - f(z_k)) &\leq \mathbb{E}(f(z'_k) - f(z_k))^2 \leq \\ &\left(\exp\left(-\frac{\tilde{m}\sqrt{N_D}kh_l}{8\tilde{\gamma}}\right) (\|z'_0 - z_0\|_{L^2,a,b} + \mathcal{W}_{2,a,b}(\pi_0, \pi) + \mathcal{W}_{2,a,b}(\pi'_0, \pi)) \right. \\ &+ (1 - c(h_l))^k \mathcal{W}_{2,a,b}(\pi_0, \pi) + (1 - c(h_{l+1}))^{2k} \mathcal{W}_{2,a,b}(\pi'_0, \pi) \\ &\left. + \frac{h_l((\tau - 1)d + \sqrt{N_D d}) C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s)}{\sqrt{N_D}} \right)^2. \end{aligned}$$

PROOF. By following the same argument as Proposition D.3 using Proposition G.6 we have the desired result. \square

COROLLARY G.8. *Suppose that the assumptions of Proposition G.7 hold for the potential U and $h_0 > 0$. Assume that*

$$h_0 \leq \frac{C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tau/N_D)}{N_D^{3/2}}, \quad B \geq \frac{16 \log(2)\tilde{\gamma}}{\tilde{m}h_0 N_D^{1/2}}, \quad B_0 \geq \frac{16\tilde{\gamma}}{\tilde{m}N_D^{1/2}h_0} \log\left(\frac{1}{N_D^3 h_0^2}\right),$$

and the levels are initialized as described in Section E, Let $l \geq 1$, $1 \leq k \leq K$, and a test function f satisfy Assumption 3.12 then for approximate gradient UBUBU with $\tau = N_D$ we have

$$(G.11) \quad \text{Var}(D_{l,l+1}) \leq C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s, \tau/N_D) d^2 h_l^2 N_D.$$

PROOF. Following the proof of Proposition F.8, but using the results of Proposition G.6 you get the required result. \square

PROPOSITION G.9. *Suppose a OHO chain at level 0 using a full gradient Gaussian approximation and a approximate gradient UBU chain at level 1, with synchronously coupled Brownian motions $(z_k)_{k \in \mathbb{N}}$ and $(z'_k)_{k \in \mathbb{N}}$ and stepsizes h_0 and $h_1 = h_0/2$, satisfying the conditions of Proposition G.6, be such that $z_0 \sim \pi_0 = \mu_G$ and $z'_0 \sim \pi'_0 = \mu_G(P_{h_1}^A)^{B/h_1}$. Then for f satisfying Assumption 3.12 we have the following variance bound*

$$\begin{aligned} \text{Var}(f(z'_k) - f(z_k)) &\leq \mathbb{E}\left[(f(z'_k) - f(z_k))^2\right] \leq \mathbb{E}\|z'_k - z_k\|_{a,b}^2 \\ &\leq \left(\exp\left(-\frac{mkh_0}{16\tilde{\gamma}}\right) (\|z'_0 - z_0\|_{L^2,a,b} + \mathcal{W}_{2,a,b}(\pi'_0, \pi)) \right. \\ &+ (1 - c(h_1))^k \mathcal{W}_{2,a,b}(\pi'_0, \pi) + \frac{dC(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s)}{N_D} + C(\tilde{\gamma}, \tilde{m}, \tilde{M})h_0\sqrt{d} \\ &\left. + \frac{h_1((\tau - 1)d + \sqrt{N_D d}) C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s)}{\sqrt{N_D}} \right)^2. \end{aligned}$$

PROOF. We aim to consider the same argument as Proposition F.10 using the results from Proposition G.6, Lemma F.6 and Proposition E.3. \square

COROLLARY G.10. *Suppose that the assumptions of Proposition G.9 hold for the potential U and $h_0 > 0$. Assume that*

$$h_0 \leq \frac{C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tau/N_D)}{N_D^{3/2}}, \quad B \geq \frac{16 \log(2) \tilde{\gamma}}{\tilde{m} h_0 N_D^{1/2}}, \quad B_0 \geq \frac{16 \tilde{\gamma}}{\tilde{m} N_D^{1/2} h_0} \log \left(\frac{1}{N_D^3 h_0^2} \right),$$

and the levels are initialized as described in Section E. Let $1 \leq k \leq K$, and a test function f satisfy Assumption 3.12 then for approximate gradient UBUBU with $\tau = N_D$ we have

$$(G.12) \quad \text{Var}(D_{0,1}) \leq \frac{C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s, \tau/N_D) d^2}{N_D^2}.$$

PROOF. Following the same argument as Corollary G.8, but using Proposition G.9 and Proposition D.3. \square

G.3. Variance bound of $S(c_R)$.

THEOREM 3.29. *Considering UBUBU-Approx method, suppose that Assumptions 3.12, 3.22, 3.23, 3.28 hold, and in addition $\gamma \geq \sqrt{8M}$,*

$$h_0 \leq \frac{C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tau/N_D)}{N_D^{3/2}}, \quad B \geq \frac{16 \log(2) \tilde{\gamma}}{\tilde{m} h_0 N_D^{1/2}}, \quad B_0 \geq \frac{16 \tilde{\gamma}}{\tilde{m} N_D^{1/2} h_0} \log \left(\frac{1}{N_D^3 h_0^2} \right).$$

Suppose that $c_R \in [0, 1]$ and $2 < \phi_N < 4$. Then for any $N \geq 1$, $S(c_R)$ has finite expected computational cost, $\mathbb{E}S(c_R) = \pi(f)$, and it has finite variance. Moreover, it satisfies a CLT as $N \rightarrow \infty$, and the asymptotic variance σ_S^2 defined in (3.8) can be bounded as

$$\sigma_S^2 \leq \frac{1}{\tilde{m} N_D K} + \frac{C(\tilde{\gamma}, \tilde{m}, \tilde{M}, \tilde{M}_1^s, \tau/N_D, \phi_N) d^2}{c_N N_D^2}.$$

PROOF. Following the same argument as Theorem 3.24 using Corollaries G.8 and G.10. \square

APPENDIX H: AUXILIARY RESULTS & RHMC ALGORITHM

LEMMA H.1. *If we have a sequence of non-negative numbers $(r_k)_{k \in \mathbb{N}}$ such that for constants $A \in (0, 1/2)$, $B, C, D \in \mathbb{R}_{\geq 0}$ such that*

$$r_{k+1}^2 \leq \left(((1-A)r_k^2 + B)^{1/2} + C \right)^2 + D$$

then

$$r_k \leq \sqrt{2} \left(1 - \frac{A}{2} \right)^k (r_0 + \sqrt{B}) + \frac{2\sqrt{2}C}{A} + 2\sqrt{\frac{D+B}{A}}.$$

PROOF. If we define $\tilde{r}_k := \sqrt{(1-A)r_k^2 + B}$, then we have that

$$\begin{aligned} \tilde{r}_{k+1}^2 &\leq (1-A)(\tilde{r}_k + C)^2 + (1-A)D + B \\ &\leq ((1-A/2)\tilde{r}_k + C)^2 + D + B. \end{aligned}$$

Then using [28][Lemma 7] we have that

$$\tilde{r}_k \leq (1-A/2)^k \tilde{r}_0 + \frac{2C}{A} + \sqrt{\frac{2(D+B)}{A}},$$

then

$$r_k \sqrt{1-A} \leq \tilde{r}_k \leq (1-A/2)^k \left(r_0 + \sqrt{B} \right) + \frac{2C}{A} + \sqrt{\frac{2(D+B)}{A}},$$

and, using the fact that $A \leq 1/2$, we obtain the required result. \square

LEMMA H.2. *If a potential $U : \mathbb{R}^d \rightarrow \mathbb{R}$ is such that $\nabla^2 U \succ mI$, and $\nabla U(x^*) = 0$ then for $x \sim \pi \propto e^{-U(x)}$ we have*

$$\mathbb{E} \left[(x - x^*)^4 \right]^{1/4} \leq 3^{1/4} \sqrt{\frac{d}{m}} \quad \text{and} \quad \mathbb{E} \left[(x - x^*)^8 \right]^{1/8} \leq 105^{1/8} \sqrt{\frac{d}{m}}.$$

PROOF. By using integration by parts and the convexity of U we have that

$$\begin{aligned} \int_{x \in \mathbb{R}^d} \|x - x^*\|^4 e^{-U(x)} dx &\leq \int_{x \in \mathbb{R}^d} \sum_{i=1}^d \sum_{j=1}^d (x_i - x_i^*)^2 (x_j - x_j^*)^2 e^{-U(x)} dx \\ &\leq d \sum_{i=1}^d \int_{x \in \mathbb{R}^d} (x_i - x_i^*)^4 e^{-U(x)} dx \\ &\leq \frac{d}{m} \sum_{i=1}^d \int_{x_{-i} \in \mathbb{R}^{d-1}} \int_{x_i \in \mathbb{R}} (x_i - x_i^*)^3 \partial_i U(x) e^{-U(x)} dx_i dx_{-i} \\ &= \frac{3d}{m} \sum_{i=1}^d \int_{x_{-i} \in \mathbb{R}^{d-1}} \int_{x_i \in \mathbb{R}} (x_i - x_i^*)^2 e^{-U(x)} dx_i dx_{-i} \\ &\leq \frac{3d}{m^2} \sum_{i=1}^d \int_{x_{-i} \in \mathbb{R}^{d-1}} \int_{x_i \in \mathbb{R}} (x_i - x_i^*) \partial_i U(x) e^{-U(x)} dx_i dx_{-i} \\ &= \frac{3d}{m^2} \sum_{i=1}^d \int_{x \in \mathbb{R}^d} e^{-U(x)} dx, \end{aligned}$$

and similarly, we have

$$\begin{aligned} \int_{x \in \mathbb{R}^d} \|x - x^*\|^8 e^{-U(x)} dx &\leq \int_{x \in \mathbb{R}^d} \sum_{i,j,k,l=1}^d (x_i - x_i^*)^2 (x_j - x_j^*)^2 (x_k - x_k^*)^2 (x_l - x_l^*)^2 e^{-U(x)} dx \\ &\leq \int_{x \in \mathbb{R}^d} d^3 \sum_{i=1}^d (x_i - x_i^*)^8 e^{-U(x)} dx \\ &\leq \frac{105d^3}{m^4} \sum_{i=1}^d \int_{x \in \mathbb{R}^d} e^{-U(x)} dx, \end{aligned}$$

as required. \square

PROPOSITION H.3 (Local error bounds for UBU). *Suppose we have a potential U which satisfies 3.6 and 3.7. Let $\phi(\xi, h, (W_t)_{t'=0}^h)$ be the solution to (1.1) at time h with initial*

condition $\xi \sim \pi$, using Brownian motion $(W_{t'})_{t'=0}^t$. Let $\psi_h(\xi, h, (W_{t'})_{t'=0}^t)$ to be the solution of the numerical discretization UBU step, defined in Sec. 2.1, with stepsize h and the same initial condition and Brownian motion. Then we have the following local error bound

$$\|\phi(\xi, h, (W_{t'})_{t'=0}^h) - \psi_h(\xi, h, (W_{t'})_{t'=0}^h)\|_{L^2, a, b} \leq \frac{3}{7} \sqrt{d} (\sqrt{M} + \gamma) h^2,$$

for $h < \min \left\{ \frac{1}{5\sqrt{M}}, \frac{1}{2\gamma} \right\}$.

PROOF. Using the method of [86] we wish to bound the local error of the UBU scheme, when initialized at the target measure of the continuous dynamics. When considering (F.1) and (F.4) we have that for $\xi \sim \pi$

$$\phi(\xi, h, (W_{t'})_{t'=0}^h) - \psi_h(\xi, h, (W_{t'})_{t'=0}^h) = (\Delta_x, \Delta_v),$$

$$\Delta_x = - \int_0^h \mathcal{F}(h-s) \nabla U(x(s)) ds + h \mathcal{F}(h/2) \nabla U(y).$$

and

$$\Delta_v = - \int_0^h \mathcal{E}(h-s) \nabla U(x(s)) ds + h \mathcal{E}(h/2) \nabla U(y).$$

Next, we use the fundamental theorem of calculus

$$\begin{aligned} \mathcal{E}(h-s) \nabla U(x(s)) &= \mathcal{E}(h/2) \nabla U(x(h/2)) \\ &+ \int_{h/2}^s (\mathcal{E}(h-s') \nabla^2 U(x(s')) v(s') + \gamma \mathcal{E}(h-s') \nabla U(x(s'))) ds'. \end{aligned}$$

Then

$$\Delta_v = -h \mathcal{E}(h/2) (\nabla U(x(h/2)) - \nabla U(y)) + \tilde{I}_1 + \tilde{I}_2,$$

where

$$\tilde{I}_1 = - \int_0^h \int_{h/2}^s \mathcal{E}(h-s') \nabla^2 U(x(s')) v(s') ds' ds,$$

and

$$\tilde{I}_2 = - \int_0^h \int_{h/2}^s \gamma \mathcal{E}(h-s') \nabla U(x(s')) ds' ds.$$

Hence

$$\|h \mathcal{E}(h/2) (\nabla U(x(h/2)) - \nabla U(y))\|_{L^2} \leq \frac{h^3 M^{3/2} \sqrt{d}}{\sqrt{48}}$$

from [86][Eq. 36]. Now, we estimate \tilde{I}_1 as

$$\begin{aligned} \mathbb{E} \left(\|\tilde{I}_1\|^2 \right) &\leq \mathbb{E} \left[\left(\int_0^h \left| \int_{h/2}^s \mathcal{E}(h-s')^2 ds' \right| ds \right) \times \left(\int_0^h \left| \int_{h/2}^s \|\nabla^2 U(x(s')) v(s')\|^2 ds' \right| ds \right) \right] \\ &\leq \frac{\mathcal{F}(h)^2}{4} \times \frac{h^2 M^2 d}{4} \leq \frac{h^4 M^2 d}{16}, \end{aligned}$$

and we estimate \tilde{I}_2 as

$$\begin{aligned} \mathbb{E} \left(\|\tilde{I}_2\|^2 \right) &\leq \gamma^2 \mathbb{E} \left[\left(\int_0^h \left| \int_{h/2}^s \mathcal{E}(h-s')^2 ds' \right| ds \right) \times \left(\int_0^h \left| \int_{h/2}^s \|\nabla U(x(s'))\|^2 ds' \right| ds \right) \right] \\ &\leq \gamma^2 \frac{\mathcal{F}(h)^2}{4} \times \frac{h^2 M d}{4} \leq \frac{h^4 M \gamma^2 d}{16}, \end{aligned}$$

then

$$\|\Delta_v\|_{L^2} \leq \frac{h^3 M^{3/2} \sqrt{d}}{\sqrt{48}} + \frac{h^2 M \sqrt{d}}{4} + \frac{h^2 \gamma \sqrt{M d}}{4}.$$

Using [86][Eq 42 Estimate] we get the bound

$$\|\Delta_x\|_{L^2} \leq \frac{h^3}{24} \left(\sqrt{3} h M^{3/2} + \left(\frac{\sqrt{42}}{2} + 1 \right) M + \gamma M^{1/2} \right) \sqrt{d}.$$

In the modified Euclidean norm we have

$$\begin{aligned} \|(\Delta_x, \Delta_v)\|_{L^2, a, b} &\leq \sqrt{\frac{3}{2}} \left(\|\Delta_x\|_{L^2} + \frac{1}{\sqrt{M}} \|\Delta_v\|_{L^2} \right) \\ &\leq \sqrt{\frac{3d}{2}} h^2 \left(\frac{h}{24} \left(\sqrt{3} h M^{3/2} + \frac{9}{2} M + \gamma M^{1/2} \right) + \frac{\sqrt{M}}{4} + \frac{\gamma}{4} \right), \end{aligned}$$

and under the assumption that $h < \min\{\frac{1}{5\sqrt{M}}, \frac{1}{2\gamma}\}$ we see that

$$\|(\Delta_x, \Delta_v)\|_{L^2, a, b} \leq \frac{3}{7} \sqrt{d} (\sqrt{M} + \gamma) h^2.$$

□

The following lemma will bound the variances of $\mathcal{Z}^{(1)}$, $\mathcal{Z}^{(2)}$ and $\mathcal{Z}^{(1)} - \mathcal{Z}^{(2)}$.

LEMMA H.4. *For $\mathcal{Z}^{(1)}$ and $\mathcal{Z}^{(2)}$ as defined in (2.4), we have*

$$\begin{aligned} \text{Cov} \left(\mathcal{Z}^{(1)} \left(h, \xi^{(1)} \right) \right) &= h I_d, \\ \text{Cov} \left(\mathcal{Z}^{(2)} \left(h, \xi^{(1)}, \xi^{(2)} \right) \right) &\preceq h I_d, \\ \text{Cov} \left(\mathcal{Z}^{(1)} \left(h, \xi^{(1)} \right) - \mathcal{Z}^{(2)} \left(h, \xi^{(1)}, \xi^{(2)} \right) \right) &\preceq \frac{\gamma h^2}{4} I_d. \end{aligned}$$

PROOF. From the definitions

$$\begin{aligned} \mathcal{Z}^{(1)} \left(h, \xi^{(1)} \right) &= \sqrt{h} \xi^{(1)}, \\ \mathcal{Z}^{(2)} \left(h, \xi^{(1)}, \xi^{(2)} \right) &= \sqrt{\frac{1-\eta^4}{2\gamma}} \left(\sqrt{\frac{1-\eta^2}{1+\eta^2}} \cdot \frac{2}{\gamma h} \xi^{(1)} + \sqrt{1 - \frac{1-\eta^2}{1+\eta^2}} \cdot \frac{2}{\gamma h} \xi^{(2)} \right), \end{aligned}$$

it is clear that $\text{Cov} \left(\mathcal{Z}^{(1)} \left(h, \xi^{(1)} \right) \right) = h I_d$ and $\text{Cov} \left(\mathcal{Z}^{(2)} \left(h, \xi^{(1)}, \xi^{(2)} \right) \right) = \frac{1-\eta^4}{2\gamma} I_d \preceq h I_d$. For the last claim, we have

$$\text{Cov} \left(\mathcal{Z}^{(2)} \left(h, \xi^{(1)}, \xi^{(2)} \right) - \mathcal{Z}^{(1)} \left(h, \xi^{(1)} \right) \right)$$

$$\begin{aligned}
&= \left(\sqrt{\frac{1-\eta^4}{2\gamma}} \sqrt{\frac{1-\eta^2}{1+\eta^2} \cdot \frac{2}{\gamma h}} - \sqrt{h} \right)^2 + \frac{1-\eta^4}{2\gamma} \left(1 - \frac{1-\eta^2}{1+\eta^2} \cdot \frac{2}{\gamma h} \right) \\
&= \frac{1-\eta^4}{2\gamma} + h - 2 \frac{(1-\eta^2)}{\gamma} = \frac{1-e^{-2\gamma h} - 4(1-e^{-\gamma h}) + 2\gamma h}{2\gamma} \leq \frac{(\gamma h)^2}{2} \cdot \frac{1}{2\gamma} \leq \frac{\gamma h^2}{4}.
\end{aligned}$$

□

LEMMA H.5. *Let $A = \sum_{l=1}^n A^{(l)}$, with $A^{(l)} \in \mathbb{R}^{d \times d}$ for every $1 \leq l \leq n$. Then we have*

$$\|A\|_{\{12\}\{3\}} = \left\| \sum_{i_1, l, m} (A_{i_1, \cdot}^{(l)})^T \cdot A_{i_1, \cdot}^{(m)} \right\|^{1/2}.$$

PROOF. This follows by expanding the formula $\|A\|_{\{12\}\{3\}} = \left\| \sum_{i_1} A_{i_1, \cdot}^T \cdot A_{i_1, \cdot} \right\|^{1/2}$ shown in Lemma 7 of [77]. □

The following lemma shows some bounds for the gradient-Lipschitz constant M and strongly Hessian Lipschitz constant M_1^s for the Bayesian multinomial regression example.

LEMMA H.6. *Consider the Bayesian multinomial regression likelihood of the form,*

$$(H.1) \quad p(y^j | q) = \frac{\exp(\langle x^j, q^{y^j} \rangle)}{\sum_{1 \leq k \leq m} \exp(\langle x^j, q^k \rangle)},$$

where the posterior potential is given as

$$(H.2) \quad U(q) = -\log(p_0(q)) - \sum_{k=1}^{N_D} \log(p(y^k | q)),$$

with $p_0(q) = \frac{\exp(-\|q\|^2/(2\sigma_0^2))}{(\pi\sigma_0^2)^{d/2}}$. This satisfies the following bounds,

$$\begin{aligned}
\sup_{q \in \mathbb{R}^d} \|\nabla^2 U(q)\| &\leq \sigma_0^{-2} + \left\| \sum_{l=1}^{N_D} (x^l)(x^l)^T \right\|, \\
\sup_{q \in \mathbb{R}^d} \|\nabla^3 U(q)\|_{\{12\}\{3\}} &\leq 6 \left\| \sum_{l=1}^{N_D} \left[(x^l)(x^l)^T \left(\sum_{m=1}^{N_D} \langle x^l, x^m \rangle^2 \right) \right] \right\|^{1/2}.
\end{aligned}$$

REMARK H.7. *If $N_D \rightarrow \infty$, and $(x^l)_{1 \leq l \leq N_D}$ are i.i.d. samples from a continuous d -dimensional distribution that is non-degenerate with $\mathbb{E}(\|x^l\|^6) = \mathcal{O}(1)$, then we would expect $\|\nabla^2 U(q)\| \propto \frac{N_D}{d}$, and $\|\nabla^3 U(q)\|_{\{12\}\{3\}} \propto \frac{N_D}{d}$.*

PROOF. For $1 \leq i \leq m$, let $E^i = \begin{pmatrix} 0_{d_o} \\ \vdots \\ I_{d_o} \\ \vdots \\ 0_{d_o} \end{pmatrix}$ be an $d \times d_o$ block matrix with an identity

matrix at block i . Let

$$S(x, q) = \sum_{1 \leq l \leq m} \exp(\langle x, q^l \rangle).$$

Let $x \otimes y \otimes z \in \mathbb{R}^{d \times d \times d}$ denote the tensor product of 3 vectors, i.e. $(x \otimes y \otimes z)_{ijk} = x_i y_j z_k$. Then we can express the likelihood term $\log(p(y|q))$ and its derivatives as follows.

$$\begin{aligned}
\log(p(y|q)) &= \langle x, q^y \rangle - \log(S(x, q)) \\
\nabla_q \log(p(y|q)) &= \sum_{i=1}^m (E^i x) (\mathbb{I}[y = i] - \exp(\langle x, q^i \rangle) (S(x, q))^{-1}) \\
\nabla_q^2 \log(p(y|q)) &= \sum_{i,j=1}^m (E^i x) (E^j x)^T \exp(\langle x, q^i \rangle + \langle x, q^j \rangle) (S(x, q))^{-2} \\
&\quad - \sum_{i=1}^m (E^i x) (E^i x)^T \exp(\langle x, q^i \rangle) (S(x, q))^{-1} \\
\nabla_q^3 \log(p(y|q)) &= - \sum_{i=1}^m (S(x, q))^{-1} \exp(\langle x, q^i \rangle) (E^i x) \otimes (E^i x) \otimes (E^i x) \\
&\quad + \sum_{i,j=1}^m (S(x, q))^{-2} \exp(\langle x, q^i \rangle + \langle x, q^j \rangle) \\
&\quad \cdot ((E^i x) \otimes (E^i x) \otimes (E^j x) + (E^i x) \otimes (E^j x) \otimes (E^i x) + (E^i x) \otimes (E^j x) \otimes (E^j x)) \\
&\quad - 2(S(x, q))^{-3} \sum_{i,j,k=1}^m (E^i x) \otimes (E^j x) \otimes (E^k x) \exp(\langle x, q^i \rangle + \langle x, q^j \rangle + \langle x, q^k \rangle).
\end{aligned}$$

The first claim of the lemma bounding $\|\nabla^2 U(q)\|$ follows from the fact that

$$0_d \preceq -\nabla_q^2 \log(p(y|q)) \preceq \sum_i^m (E^i x) (E^i x)^T \exp(\langle x, q^i \rangle) (S(x, q))^{-1} \preceq \sum_i^m (E^i x) (E^i x)^T,$$

here \preceq denotes the semidefinite order.

For the second claim, note that

$$\left\| - \sum_{l=1}^{N_D} \sum_{i=1}^m (S(x^l, q))^{-1} \exp(\langle x^l, q^i \rangle) (E^i x^l) \otimes (E^i x^l) \otimes (E^i x^l) \right\|_{\{12\}\{3\}}$$

using Lemma H.5

$$\begin{aligned}
&\leq \left\| \sum_{l,m=1}^{N_D} \langle x^l, x^m \rangle \left((x^l) (x^m)^T \right)^2 \right\|^{1/2} = \left\| \sum_{l,m=1}^{N_D} \langle x^l, x^m \rangle^2 (x^l) (x^m)^T \right\|^{1/2} \\
&\leq \left\| \frac{1}{2} \sum_{l,m=1}^{N_D} \langle x^l, x^m \rangle^2 [(x^l) (x^l)^T + (x^m) (x^m)^T] \right\|^{1/2} \\
&= \left\| \sum_{l=1}^{N_D} \left[(x^l) (x^l)^T \left(\sum_{m=1}^{N_D} \langle x^l, x^m \rangle^2 \right) \right] \right\|^{1/2}.
\end{aligned}$$

The other terms in the sum can be bounded similarly as

$$\left\| \sum_{i,j=1}^m (S(x, q))^{-2} \exp(\langle x, q^i \rangle + \langle x, q^j \rangle) \right\|$$

$$\begin{aligned}
& \cdot \left((E^i x) \otimes (E^i x) \otimes (E^j x) + (E^i x) \otimes (E^j x) \otimes (E^i x) + (E^i x) \otimes (E^j x) \otimes (E^j x) \right) \Big\|_{\{12\}\{3\}} \\
& \leq 3 \left\| \sum_{l=1}^{N_D} \left[(x^l)(x^l)^T \left(\sum_{m=1}^{N_D} \langle x^l, x^m \rangle^2 \right) \right] \right\|^{1/2}, \\
& \left\| -2(S(x, q))^{-3} \sum_{i,j,k=1}^m (E^i x) \otimes (E^j x) \otimes (E^k x) \exp(\langle x, q^i \rangle + \langle x, q^j \rangle + \langle x, q^k \rangle) \right\|_{\{12\}\{3\}} \\
& \leq 2 \left\| \sum_{l=1}^{N_D} \left[(x^l)(x^l)^T \left(\sum_{m=1}^{N_D} \langle x^l, x^m \rangle^2 \right) \right] \right\|^{1/2},
\end{aligned}$$

and the claim follows by the triangle inequality. \square

Algorithm 3 Randomized Hamiltonian Monte Carlo with Partial Refreshment (RHMC)

1: **Input:**

- stepsize h .
- Initial distribution μ_0 on $\mathbb{R}^d \times \mathbb{R}^d$.
- Potential function $U : \mathbb{R}^d \rightarrow \mathbb{R}$ of target distribution.
- Number of samples parameter K .
- Expected number of leapfrog steps parameter $E_L \geq 1$.
- Partial refreshment parameter α .

2: **Initialise** $(x_0, v_0) \sim \mu_0$.

3: **for** $i = 1, \dots, K$ **do**

4: Sample $L \sim \text{Geom}(1/E_L)$.

5: **Perform L leapfrog steps.**

6: Set $(\tilde{x}_0, \tilde{v}_0) := (x_i, v_i)$.

7: **for** $j = 0, \dots, L - 1$ **do**

8: $\tilde{v}_{j+1/2} := \tilde{v}_j - \frac{h}{2} \nabla U(\tilde{x}_j)$

9: $\tilde{x}_{j+1} := \tilde{x}_j + h\tilde{v}_{j+1/2}$

10: $\tilde{v}_{j+1} := \tilde{v}_{j+1/2} - \frac{h}{2} \nabla U(\tilde{x}_{j+1})$

11: **end for**

12: Let $(x'_i, v'_i) = (\tilde{x}_L, \tilde{v}_L)$

13: **Compute Hamiltonian.**

14: $H(x_i, v_i) = U(x_i) + \frac{1}{2} \|v_i\|^2$, $H(x'_i, v'_i) = U(x'_i) + \frac{1}{2} \|v'_i\|^2$.

15: **Perform Metropolis-Hastings accept/reject step.**

16: With probability $\min \left[1, \exp(H(x_i, v_i) - H(x'_i, v'_i)) \right]$, set $(x_{i+1}, v_{i+1}) = (x'_i, v'_i)$ (accept proposal).

17: Otherwise, set $(x_{i+1}, v_{i+1}) = (x_i, v_i)$ (reject proposal).

18: **Partial velocity refreshment.**

19: Sample $Z \sim \mathcal{N}(0_d, I_d)$ and update $v_{i+1} \rightarrow \alpha v_{i+1} + (1 - \alpha^2)^{1/2} Z$.

20: **end for**

21: **Output:**

22: Samples $(x_1, v_1), \dots, (x_K, v_K)$.

REFERENCES

- [1] Jason M. Altschuler and Sinho Chewi.
Faster high-accuracy log-concave sampling via algorithmic warm starts.
In *2023 IEEE 64th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 2169–2176,
2023.

- [2] Simon Apers, Sander Gribling, and Dániel Szilágyi.
Hamiltonian Monte Carlo for efficient Gaussian sampling: long and random steps.
arXiv preprint arXiv:2209.12771, 2022.
- [3] Elsidig Awadelkarim, Ajay Jasra, and Hamza Ruzayqat.
Unbiased parameter estimation for partially observed diffusions.
arXiv preprint arXiv:2309.10589, 2023.
- [4] Daniel Azagra, Juan Ferrera, Fernando López-Mesas, and Yenny Rangel.
Smooth approximation of Lipschitz functions on Riemannian manifolds.
Journal of Mathematical Analysis and Applications, 326(2):1370–1378, 2007.
- [5] Jack Baker, Paul Fearnhead, Emily B. Fox, and Christopher Nemeth.
Control variates for stochastic gradient MCMC.
Stat. Comput., 29(3):599–615, 2019.
- [6] J.E. Besag.
Comments on ‘Representations of knowledge in complex systems’ by U. Grenander and M.I. Miller.
J. Roy. Statist. Soc. Ser. B, (56):591–592, 1994.
- [7] Alexandros Beskos, Natesh Pillai, Gareth Roberts, Jesus-Maria Sanz-Serna, and Andrew Stuart.
Optimal tuning of the hybrid Monte Carlo algorithm.
Bernoulli, 19(5A):1501–1534, 2013.
- [8] Joris Bierkens, Paul Fearnhead, and Gareth Roberts.
The zig-zag process and super-efficient sampling for Bayesian analysis of big data.
Ann. Statist., 47(3):1288–1320, 2019.
- [9] Stephen D. Bond and Benedict J. Leimkuhler.
Molecular dynamics and the accuracy of numerically computed averages.
Acta Numer., 16:1–65, 2007.
- [10] Nawaf Bou-Rabee and Andreas Eberle.
Couplings for Andersen dynamics.
Ann. Inst. Henri Poincaré Probab. Stat., 58(2):916–944, 2022.
- [11] Nawaf Bou-Rabee and Andreas Eberle.
Mixing time guarantees for unadjusted Hamiltonian Monte Carlo.
Bernoulli, 29(1):75–104, 2023.
- [12] Nawaf Bou-Rabee, Andreas Eberle, and Raphael Zimmer.
Coupling and convergence for Hamiltonian Monte Carlo.
Ann. Appl. Probab., 30(3):1209–1250, 2020.
- [13] Nawaf Bou-Rabee and Tore Selland Kleppe.
Randomized Runge-Kutta-Nyström.
arXiv preprint arXiv:2310.07399, 2023.
- [14] Nawaf Bou-Rabee and Milo Marsden.
Unadjusted Hamiltonian MCMC with stratified Monte Carlo time integration.
arXiv preprint arXiv:2211.11003, 2022.
- [15] Nawaf Bou-Rabee and Houman Owhadi.
Long-run accuracy of variational integrators in the stochastic context.
SIAM J. Numer. Anal., 48(1):278–297, 2010.
- [16] Nawaf Bou-Rabee and Jesús María Sanz-Serna.
Randomized Hamiltonian Monte Carlo.
The Annals of Applied Probability, 27(4):2159 – 2194, 2017.
- [17] Stéphane Boucheron, Gábor Lugosi, and Pascal Massart.
Concentration inequalities.
Oxford University Press, Oxford, 2013.
A nonasymptotic theory of independence, With a foreword by Michel Ledoux.
- [18] Nicolas Brosse, Alain Durmus, and Eric Moulines.
The promises and pitfalls of stochastic gradient Langevin dynamics.
In *Advances in Neural Information Processing Systems*, pages 8268–8278, 2018.
- [19] Axel Brünger, Charles L. Brooks, and Martin Karplus.
Stochastic boundary conditions for molecular dynamics simulations of ST2 water.
Chemical Physics Letters, 105(5):495–500, 1984.
- [20] Yu Cao, Jianfeng Lu, and Lihan Wang.
On explicit L^2 -convergence rate estimate for underdamped Langevin dynamics.
Arch. Ration. Mech. Anal., 247(5):Paper No. 90, 34, 2023.

- [21] Neil K. Chada, Jordan Franks, Ajay Jasra, Kody J. Law, and Matti Vihola. Unbiased inference for discretely observed hidden Markov model diffusions. *SIAM/ASA J. Uncertain. Quantif.*, 9(2):763–787, 2021.
- [22] Martin Chak and Pierre Monmarché. Reflection coupling for unadjusted generalized Hamiltonian Monte Carlo in the nonconvex stochastic gradient case. *arXiv preprint arXiv:2310.18774*, 2023.
- [23] Louis HY Chen and Adrian Röllin. Stein couplings for normal approximation. *arXiv preprint arXiv:1003.6039*, 2010.
- [24] Yuansi Chen and Khashayar Gatmiry. When does Metropolized Hamiltonian Monte Carlo provably outperform Metropolis-adjusted Langevin algorithm? *arXiv preprint arXiv:2304.04724*, 2023.
- [25] Xiang Cheng, Niladri S Chatterji, Peter L Bartlett, and Michael I Jordan. Underdamped Langevin MCMC: A non-asymptotic analysis. In *Conference on learning theory*, pages 300–323. PMLR, 2018.
- [26] Adrien Corenflos, Matthew Sutton, and Nicolas Chopin. Debiasing piecewise deterministic Markov process samplers using couplings. *arXiv preprint arXiv:2306.15422*, 2023.
- [27] Rob Cornish, Paul Vanetti, Alexandre Bouchard-Côté, George Deligiannidis, and Arnaud Doucet. Scalable Metropolis-Hastings for exact Bayesian inference with large datasets. In *International Conference on Machine Learning*, pages 1351–1360. PMLR, 2019.
- [28] Arnak Dalalyan. Further and stronger analogy between sampling and optimization: Langevin Monte Carlo and gradient descent. In *Conference on Learning Theory*, pages 678–689. PMLR, 2017.
- [29] Arnak S. Dalalyan. Theoretical guarantees for approximate sampling from smooth and log-concave densities. *J. R. Stat. Soc. Ser. B. Stat. Methodol.*, 79(3):651–676, 2017.
- [30] Arnak S. Dalalyan and Lionel Riou-Durand. On sampling from a log-concave density using kinetic Langevin diffusions. *Bernoulli*, 26(3):1956–1988, 2020.
- [31] Pierre Del Moral, Shulan Hu, Ajay Jasra, Hamza Ruzayqat, and Xinyu Wang. Bayesian Parameter Inference for Partially Observed Diffusions using Multilevel Stochastic Runge-Kutta Methods. *arXiv preprint arXiv:2309.13557*, 2023.
- [32] George Deligiannidis, Daniel Paulin, Alexandre Bouchard-Côté, and Arnaud Doucet. Randomized Hamiltonian Monte Carlo as scaling limit of the bouncy particle sampler and dimension-free convergence rates. *Ann. Appl. Probab.*, 31(6):2612–2662, 2021.
- [33] Randall Douc, Pierre E. Jacob, Anthony Lee, and Dootika Vats. Solving the Poisson equation using coupled Markov chains. *arXiv preprint arXiv:2206.05691*, 2022.
- [34] Simon Duane, Anthony D Kennedy, Brian J Pendleton, and Duncan Roweth. Hybrid Monte Carlo. *Physics letters B*, 195(2):216–222, 1987.
- [35] Alain Durmus, Aurélien Enfroy, Éric Moulines, and Gabriel Stoltz. Uniform minorization condition and convergence bounds for discretizations of kinetic Langevin dynamics. *arXiv preprint arXiv:2107.14542*, 2021.
- [36] Alain Durmus and Éric Moulines. Nonasymptotic convergence analysis for the unadjusted Langevin algorithm. *Ann. Appl. Probab.*, 27(3):1551–1587, 2017.
- [37] Rick Durrett. *Probability: theory and examples*, volume 49 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press, Cambridge, fifth edition, 2019.
- [38] Andreas Eberle, Arnaud Guillin, and Raphael Zimmer. Couplings and quantitative contraction rates for Langevin dynamics. *Annals of Applied Probability*, 47(4):1982–2010, 2019.

- [39] Andreas Eberle, Arnaud Guillin, and Raphael Zimmer.
Couplings and quantitative contraction rates for Langevin dynamics.
Ann. Probab., 47(4):1982–2010, 2019.
- [40] James Foster, Terry Lyons, and Harald Oberhauser.
The shifted ODE method for underdamped Langevin MCMC.
arXiv preprint arXiv:2101.03446, 2021.
- [41] James M. Foster, Gonçalo dos Reis, and Calum Strange.
High order splitting methods for SDEs satisfying a commutativity condition.
SIAM J. Numer. Anal., 62(1):500–532, 2024.
- [42] Michael B. Giles.
Multilevel Monte Carlo methods.
Acta Numer., 24:259–328, 2015.
- [43] Michael B. Giles, Mateusz B. Majka, Lukasz Szpruch, Sebastian J. Vollmer, and Konstantinos C. Zygalakis.
Multi-level Monte Carlo methods for the approximation of invariant measures of stochastic differential equations.
Stat. Comput., 30(3):507–524, 2020.
- [44] Peter W Glynn and Chang-han Rhee.
Exact estimation for markov chain equilibrium expectations.
Journal of Applied Probability, 51(A):377–389, 2014.
- [45] Nicolaï Gouraud, Pierre Le Bris, Adrien Majka, and Pierre Monmarché.
HMC and underdamped Langevin united in the unadjusted convex smooth case.
arXiv preprint arXiv:2202.00977, 2023.
- [46] Jeremy Heng and Pierre E Jacob.
Unbiased Hamiltonian Monte Carlo with couplings.
Biometrika, 106(2), 2019.
- [47] Roger A. Horn and Charles R. Johnson.
Matrix analysis.
Cambridge University Press, Cambridge, second edition, 2013.
- [48] Jesús A Izaguirre, Daniel P Catarello, Justin M Wozniak, and Robert D Skeel.
Langevin stabilization of molecular dynamics.
The Journal of Chemical Physics, 114(5):2090–2098, 2001.
- [49] Pierre E. Jacob, John O’Leary, and Yves F. Atchadé.
Unbiased Markov chain Monte Carlo methods with couplings.
J. R. Stat. Soc. Ser. B. Stat. Methodol., 82(3):543–600, 2020.
- [50] Rie Johnson and Tong Zhang.
Accelerating stochastic gradient descent using predictive variance reduction.
Advances in neural information processing systems, 26, 2013.
- [51] Aldéric Joulin and Yann Ollivier.
Curvature, concentration and error estimates for Markov chain Monte Carlo.
Ann. Probab., 38(6):2418–2442, 2010.
- [52] Nabil Kahlalé.
Unbiased time-average estimators for Markov chains.
Mathematics of Operations Research, 2023.
- [53] Aikaterini Karoni, Benedict Leimkuhler, and Gabriel Stoltz.
Friction-adaptive descent: a family of dynamics-based optimization methods.
J. Comput. Dyn., 10(4):450–484, 2023.
- [54] Hassan K. Khalil.
Nonlinear systems.
Prentice-Hall: Upper Saddle River, New Jersey, 2002.
- [55] Siem Jan Koopman and Rutger Lit.
A dynamic bivariate Poisson model for analysing and forecasting match results in the English Premier League.
Journal of the Royal Statistical Society. Series A, 178(1):167–186, 2015.
- [56] Yann LeCun, Burges Cortes, Corinna, and Chris et al.
MNIST handwritten digit database.
2010.
- [57] Yin Tat Lee, Ruoqi Shen, and Kevin Tian.
Lower bounds on Metropolized sampling methods for well-conditioned distributions.
Advances in Neural Information Processing Systems, 34:18812–18824, 2021.

- [58] Ben Leimkuhler and Charles Matthews.
Molecular dynamics.
Interdisciplinary applied mathematics, 39:443, 2015.
- [59] Benedict Leimkuhler and Charles Matthews.
Rational construction of stochastic numerical methods for molecular sampling.
Appl. Math. Res. Express. AMRX, (1):34–56, 2013.
- [60] Benedict Leimkuhler and Charles Matthews.
Robust and efficient configurational molecular sampling via Langevin dynamics.
Journal of Chemical Physics, 138:174102, 2013.
- [61] Benedict Leimkuhler, Charles Matthews, and Gabriel Stoltz.
The computation of averages from equilibrium and nonequilibrium Langevin molecular dynamics.
IMA J. Numer. Anal., 36(1):13–79, 2016.
- [62] Benedict Leimkuhler, Daniel Paulin, and Peter A Whalley.
Contraction Rate Estimates of Stochastic Gradient Kinetic Langevin Integrators.
arXiv preprint arXiv:2306.08592, 2023.
- [63] Benedict J. Leimkuhler, Daniel Paulin, and Peter A. Whalley.
Contraction and Convergence Rates for Discretized Kinetic Langevin Dynamics.
SIAM Journal on Numerical Analysis, 62(3):1226–1258, 2024.
- [64] Jianfeng Lu and Lihan Wang.
On explicit L^2 -convergence rate estimate for piecewise deterministic Markov processes in MCMC algorithms.
The Annals of Applied Probability, 32(2):1333 – 1361, 2022.
- [65] Michael J Maher.
Modelling association football scores.
Statistica Neerlandica, 36(3):109–118, 1982.
- [66] Mateusz B Majka, Aleksandar Mijatović, and Lukasz Szpruch.
Non-asymptotic bounds for sampling algorithms without log-concavity.
Annals of Applied Probability, 30(4):1534–1581, 2020.
- [67] J. C. Mattingly, A. M. Stuart, and D. J. Higham.
Ergodicity for SDEs and approximations: locally Lipschitz vector fields and degenerate noise.
Stochastic Process. Appl., 101(2):185–232, 2002.
- [68] Lawrence Middleton, George Deligiannidis, Arnaud Doucet, and Pierre E. Jacob.
Unbiased Markov chain Monte Carlo for intractable target distributions.
Electron. J. Stat., 14(2):2842–2891, 2020.
- [69] G. N. Milstein and M. V. Tretyakov.
Stochastic numerics for mathematical physics.
Scientific Computation. Springer-Verlag, Berlin, 2004.
- [70] Pierre Monmarché.
High-dimensional MCMC with a standard splitting scheme for the underdamped Langevin diffusion.
Electron. J. Stat., 15(2):4117–4166, 2021.
- [71] Pierre Monmarché.
Almost sure contraction for diffusions on \mathbb{R}^d . Application to generalized Langevin diffusions.
Stochastic Process. Appl., 161:316–349, 2023.
- [72] Pierre Monmarché.
An entropic approach for Hamiltonian Monte Carlo: the idealized case.
Ann. Appl. Probab., 34(2):2243–2293, 2024.
- [73] Eike H. Müller, Rob Scheichl, and Tony Shardlow.
Improving multilevel Monte Carlo for stochastic differential equations with application to the Langevin equation.
Proc. A., 471(2176):20140679, 20, 2015.
- [74] Radford M. Neal.
MCMC using Hamiltonian dynamics.
In *Handbook of Markov chain Monte Carlo*, Chapman & Hall/CRC Handb. Mod. Stat. Methods, pages 113–162. CRC Press, Boca Raton, FL, 2011.
- [75] Yurii Nesterov et al.
Lectures on convex optimization, volume 137.
Springer, 2018.
- [76] Trevor Park and George Casella.
The Bayesian lasso.
Journal of the American Statistical Association, 103(482):681–686, 2008.

- [77] Daniel Paulin and Peter A. Whalley.
Correction to “Wasserstein distance estimates for the distributions of numerical approximations to ergodic stochastic differential equations”.
arXiv preprint arXiv:2402.08711, 2024.
- [78] Grigorios A. Pavliotis.
Stochastic processes and applications, volume 60 of *Texts in Applied Mathematics*.
Springer, New York, 2014.
Diffusion processes, the Fokker-Planck and Langevin equations.
- [79] Matias Quiroz, Robert Kohn, Mattias Villani, and Minh-Ngoc Tran.
Speeding up MCMC by efficient data subsampling.
J. Amer. Statist. Assoc., 114(526):831–843, 2019.
- [80] Chang-han Rhee and Peter W Glynn.
Unbiased estimation with square root convergence for SDE models.
Operations Research, 63(5):1026–1043, 2015.
- [81] Lewis Fry Richardson.
The approximate arithmetical solution by finite differences with an application to stresses in masonry dams.
Philosophical Transactions of the Royal Society of America, 210:307–357, 1911.
- [82] Christian P. Robert and George Casella.
Monte Carlo statistical methods.
Springer Texts in Statistics. Springer-Verlag, New York, second edition, 2004.
- [83] Gareth O Roberts and Jeffrey S Rosenthal.
Optimal scaling of discrete approximations to langevin diffusions.
Journal of the Royal Statistical Society: Series B (Statistical Methodology), 60(1):255–268, 1998.
- [84] Gareth O Roberts and Jeffrey S Rosenthal.
Optimal scaling for various Metropolis-Hastings algorithms.
Statistical science, 16(4):351–367, 2001.
- [85] Hamza Ruzayqat, Neil K. Chada, and Ajay Jasra.
Unbiased estimation using underdamped Langevin dynamics.
SIAM J. Sci. Comput., 45(6):A3047–A3070, 2023.
- [86] Jesus María Sanz-Serna and Konstantinos C. Zygalakis.
Wasserstein distance estimates for the distributions of numerical approximations to ergodic stochastic differential equations.
J. Mach. Learn. Res., 22:Paper No. 242, 37, 2021.
- [87] Katharina Schuh.
Global contractivity for Langevin dynamics with distribution-dependent forces and uniform in time propagation of chaos.
Ann. Inst. Henri Poincaré Probab. Stat., 2024.
- [88] Katharina Schuh and Peter A Whalley.
Convergence of kinetic Langevin samplers for non-convex potentials.
arXiv preprint arXiv:2405.09992, 2024.
- [89] Robert D Skeel and Jesús A Izaguirre.
An impulse integrator for Langevin dynamics.
Molecular Physics, 100(24):3885–3891, 2002.
- [90] J. Michael Steele.
An Efron-Stein inequality for nonsymmetric statistics.
Ann. Statist., 14(2):753–758, 1986.
- [91] Yee Whye Teh, Alexandre H. Thiery, and Sebastian J. Vollmer.
Consistency and fluctuations for stochastic gradient Langevin dynamics.
The Journal of Machine Learning Research, 17(1):193–225, 2016.
- [92] Adam Telatovich and Xiantao Li.
The strong convergence of operator-splitting methods for the Langevin dynamics model.
arXiv preprint arXiv:1706.04237, 2017.
- [93] Matti Vihola.
Unbiased estimators and multilevel Monte Carlo.
Operations Research, 66(2):448–462, 2018.
- [94] Cédric Villani.
Optimal transport: old and new, volume 338.
Springer, 2009.

- [95] Sebastian J. Vollmer, Konstantinos C. Zygalakis, and Yee Whye Teh.
Exploration of the (non-)asymptotic bias and variance of stochastic gradient Langevin dynamics.
J. Mach. Learn. Res., 17:Paper No. 159, 45, 2016.
- [96] Maxime Vono, Daniel Paulin, and Arnaud Doucet.
Efficient MCMC sampling with dimension-free convergence rate using ADMM-type splitting.
J. Mach. Learn. Res., 23:Paper No. [25], 69, 2022.
- [97] Tianze Wang and Guanyang Wang.
Unbiased multilevel Monte Carlo methods for intractable distributions: MLMC meets MCMC.
J. Mach. Learn. Res., 24:Paper No. [249], 40, 2023.
- [98] Christian H Weiß, Fukang Zhu, and Aisouda Hoshiyar.
Softplus ingarch models.
Statistica Sinica, 32(2):1099–1120, 2022.
- [99] M. Welling and Y. W. Teh.
Bayesian learning via stochastic gradient Langevin dynamics.
In *Proceedings of the 28th International Conference on Machine Learning*, pages 681–688, 2011.
- [100] Paul FV Wiemann, Thomas Kneib, and Julien Hambuckers.
Using the softplus function to construct alternative link functions in generalized linear models and beyond.
Statistical Papers, pages 1–26, 2023.
- [101] Hu Zhengmian, Feihu Huang, and Heng Huang.
Optimal Underdamped Langevin MCMC Method.
Advances in Neural Information Processing Systems, 34:19363–19374, 2021.
- [102] Alfonso Álamo Zapatero.
Word Series for the Numerical Integration of Stochastic Differential Equations.
PhD thesis, Universidad de Valladolid, 2021.