# FROM TODA HIERARCHY TO KP HIERARCHY

DI YANG, JIAN ZHOU

ABSTRACT. Using the matrix-resolvent method and a formula of the second-named author on the *n*-point function for a KP taufunction, we show that the tau-function of an arbitrary solution to the Toda lattice hierarchy is a KP tau-function. We then generalize this result to tau-functions for the extended Toda hierarchy (ETH) by developing the matrix-resolvent method for the ETH. As an example the partition function of Gromov–Witten invariants of the complex projective line is a KP tau-function, and an application on irreducible representations of the symmetric group is obtained.

#### Contents

1.	Introduction	1
2.	The formula for the $n$ -point function for a KP tau-function	6
3.	From the Toda lattice hierarchy to the KP hierarchy	9
4.	Extended Toda flows in the KP hierarchy	10
5.	Application to Gromov–Witten invariants of $\mathbb{P}^1$	19
References		27

# 1. INTRODUCTION

The Kadomtsev–Petviashvili (KP) hierarchy and the Toda lattice hierarchy are two important integrable hierarchies. In this paper, we will show that the tau-function of an arbitrary solution to the Toda lattice hierarchy gives an infinite family of tau-functions of the KP hierarchy. This is achieved by combining two different results: a formula [55] of the second-named author on the *n*-point function for a KP tau-function in the big cell, and a work [50] of the first-named author on the *n*-point function for a Toda tau-function.

The KP hierarchy is an infinite family of equations with infinitely many unknown functions, which can be written using the Lax pair formalism as follows:

(1) 
$$\frac{\partial L_{\rm KP}}{\partial p_k} = \left[ \left( L^k \right)_+, L \right], \quad k \ge 1,$$

where

(2) 
$$L_{\rm KP} = \partial + \sum_{j \ge 1} u_j \partial^{-j}, \quad \partial = \partial/\partial X,$$

is a pseudodifferential operator, called the *Lax operator*. For details about pseudodifferential operators and their operations see for example [15]. The independent variables  $p_1, p_2, p_3, \ldots$  are called *times*. Since  $\partial L_{\text{KP}}/\partial p_1 = \partial L/\partial X$ , we identify  $p_1$  with X.

Let  $\mathbf{p} = (p_1, p_2, ...)$  denote the infinite vector of KP times. It is well known (see e.g. [15]) that an arbitrary solution  $(u_1(\mathbf{p}), u_2(\mathbf{p}), ...)$  to the KP hierarchy (1) can be compactly represented by a single function  $\tau = \tau(\mathbf{p})$  called the *tau-function* as

(3) 
$$u_1(\mathbf{p}) = \frac{\partial^2 \log \tau}{\partial p_1^2}, \quad u_2(\mathbf{p}) = \frac{1}{2} \frac{\partial^2 \log \tau}{\partial p_1 \partial p_2} - \frac{1}{2} \frac{\partial^3 \log \tau}{\partial p_1^3}, \quad \dots,$$

and the tau-function  $\tau$  satisfies the *Hirota bilinear identities* given by

(4) 
$$\operatorname{res}_{\lambda=\infty} \tau(\mathbf{p} - [\lambda^{-1}])\tau(\mathbf{p}' + [\lambda^{-1}])e^{\sum_{k\geq 1}(p_k - p'_k)\lambda^k}d\lambda = 0.$$

Here  $[\lambda^{-1}] := (\frac{1}{\lambda}, \frac{1}{2\lambda^2}, \frac{1}{3\lambda^3}, \dots)$ . Equivalence between (4) and (1) is a standard result in the theory of integrable systems, see for example [15]. Equation (4) itself gives the defining equations for a KP tau-function (namely, we do not need to start with a solution  $(u_1(\mathbf{p}), u_2(\mathbf{p}), \dots)$  to the KP hierarchy (1)).

Let  $\Lambda : f(x) \mapsto f(x+1)$  be the shift operator and

(5) 
$$L := \Lambda + v(x) + w(x)\Lambda^{-1}$$

the *Lax operator*. The *Toda lattice hierarchy* (also known as the 1d Toda chain or 1-Toda hierarchy) can be defined using the Lax pair formalism as

(6) 
$$\frac{\partial L}{\partial m_i} = \frac{1}{(i+1)!} \left[ (L^{i+1})_+, L \right], \quad i \ge 0.$$

Here, for a difference operator P written in the form  $P = \sum_{k \in \mathbb{Z}} P_k \Lambda^k$ ,  $P_+$  is defined as  $\sum_{k \ge 0} P_k \Lambda^k$ . We also denote  $\mathbf{m} = (m_0, m_1, m_2, \dots)$ .

Let  $(v(x, \mathbf{m}), w(x, \mathbf{m}))$  be an arbitrary power-series-in-**m** solution to the Toda lattice hierarchy, and  $\tau(x, \mathbf{m})$  the tau-function [21] of this solution. Our first result of this paper is given by the following theorem.

**Theorem 1.1.** The tau-function  $\tau(x, \mathbf{m})$  for the Toda lattice hierarchy is a tau-function of the KP hierarchy for any x, where  $\mathbf{m}$  and the KP times  $\mathbf{p}$  are related by

(7) 
$$m_i = i! p_{i+1}, \quad i \ge 0.$$

Let us now recall some earlier results in the literature that motivate us to obtain this result. It is well known in matrix model theory [1, 31] that using the theory of orthogonal polynomials, the GUE partition function  $\{Z_N(\mathbf{s}; \epsilon)\}_{N\geq 1}$  is a tau-function for the truncated Toda lattice hierarchy, in the sense that the variable x in (5) is restricted to positive integers N. Here N is the size of the matrices of the Hermitian matrices used to defined the partition function. A less well-known result is that for each N,  $Z_N(\mathbf{s}; \epsilon)$  is also a tau-function of the KP hierarchy [47]. Both of these results have been revisited recently and some new perspectives naturally arise.

First of all, in [21, Definition 1.2.2], a notion of matrix resolvent for the Toda lattice hierarchy was introduced. Based on this, a definition of the tau function of Dubrovin-Zhang type of the Toda lattice taufunction was given in [21, Definition 1.2.4]. Usually a normalization constant is chosen so that the partition function of Gaussian unitary ensemble (GUE) is equal to 1 when all the coupling constant is set to be equal to zero. In [21, Appendix A] it was shown that after multiplying by a suitable correction factor, the GUE partition functions give us a tau-function of Dubrovin-Zhang type. In [21] it was also shown that the correction factor can be obtained by using the theory of Toda taufunctions, so this method is applicable to other examples.

Note the major goal of [21] is to develop a method for computing *n*-point function associated with the tau-function of the Toda lattice hierarchy based on the matrix-resolvent method, by generalizing the earlier results developed by Bertola, Dubrovin and the first-named author in the cases of KdV hierarchy [7] and Drinfeld-Sokolov hierarchies associated with simple Lie algebras [8]. Earlier the second-named author proved an explicit formula [54] for the Schur expansion of the Witten-Kontsevich tau-function of the KdV hierarchy. Balogh and the first-named author interpreted this formula in terms of the affine coordinates of the Witten-Kontsevich tau-function in [5]. Inspired by [7], the second-named author proved a formula [55] of *n*-point function associated with an arbitrary tau-function of the KP hierarchy, based on the affine coordinates of the element in the Sato Grassmannian corresponding to the tau-function. The formula in [21] have a difference from the formula in [55]: In the former matrices of power series are used whereas ordinary power series are used in the latter. To remedy the difference, the first-named author proved a formula for n-point function using only power series in [50] for tau-functions of Toda lattice hierarchy (see also [23]).

#### DI YANG, JIAN ZHOU

Secondly, it was shown in [56] that the normalized GUE partition function  $\frac{Z_N(s;\epsilon)}{Z_N(0;\epsilon)}$  gives rise to a family of KP tau-functions in the big cell parameterized by the t'Hooft coupling constant  $x = N\epsilon$  (see also [35, 43]). Furthermore, an explicit formula for the affine coordinates for this family of KP tau-function was derived in [56]. As an application, the formulas for the corresponding *n*-point functions were obtained by applying the formula in [55].

Now the GUE partition functions can be studied from two different perspectives: either as a tau-function of Dubrovin-Zhang type of the Toda lattice hierarchy, or as a family of tau-functions of the KP hierarchy. The belief that this is just a special case of general phenomenon leads us to Theorem 1.1.

There are earlier results that also lead to Theorem 1.1. It is well known that by introducing the t'Hooft coupling constant  $x = N\epsilon$ ,  $\log \frac{Z_N(\mathbf{s};\epsilon)}{Z_N(\mathbf{0};\epsilon)}$  has an expansion as weighted sum of ribbon graphs [21, 34, 56]. In other words, the enumeration of ribbon graphs gives rise to a special family of tau-functions of the KP hierarchy which become a special tau-function for the Toda lattice hierarchy by multiplying by a suitable correction factor; furthermore, GUE is a matrix model for enumerating the ribbon graphs. Recall that ribbon graphs can be regarded as *clean* dessins. As a generalization, the weighted sum of Grothendieck's dessins d'enfants has been shown by Zograf [59] (see also Kazarian and Zograf [36]) to be a tau-function of the KP hierarchy, that is referred to as the dessin partition function. Based on this fact, the second-named author had found the affine coordinates of the taufunction of the KP hierarchy associated with the dessin counting [57], and so explicit formula for the n-point functions of the dessin partition function can be written down using the general formula in [55]. The dessin partition function is a family of KP tau-functions parameterized by two parameters. It reduces to some other well-known tau-function by suitably specifying these parameters [57]. A proposal to study the dualities among different models based on the theory of KP hierarchy was then proposed in [57] and was further elaborated in [58].

The study of KP tau-functions and the study of Toda tau-functions were merged in our earlier work [52] in our study of the dessin taufunction and its role in unifying various theories. Since counting dessins are similar to counting ribbon graphs and GUE provides a matrix model for enumerating the ribbon graphs, one naturally expects a matrix model for enumerating dessins. The problem was addressed by Ambjørn and Chekhov [4], where they proposed a matrix model in the class of generalized Kontsevich models and an equivalent Hermitian

one matrix model. In [52], we found that the Laguerre unitary ensemble (LUE) gives a simpler and rigorous matrix model for the dessins counting in the sense that the dessin partition function is equal to the *normalized* LUE partition function. Furthermore, by multiplying by a suitable correction factor, the dessin partition function gives rise to a Dubrovin–Zhang type tau-function of the Toda lattice hierarchy. It was proposed in [52] to study the duality between dessin partition function with partition functions of other theories from the viewpoint of Toda lattice hierarchy. In this new approach, one can apply the theory of normal forms of integrable hierarchies and the extended Toda hierarchy [11] as developed by Dubrovin and Zhang [25, 26]. In fact, the relevant Frobenius manifold is the Frobenius manifold associated with the Gromov–Witten invariants of  $\mathbb{P}^1$ . It has already been used in [19] (cf. [21, 51]) to calculate the GUE correlators.

Based on some earlier results of Dijkgraaf and Witten [16], Eguchi and Yang [27] proposed a matrix model for the Gromov-Witten invariants of  $\mathbb{P}^1$  (see also [28]). Besides the Toda lattice hierarchy, the  $\mathbb{P}^1$ -partition function satisfies an extra family of flows introduced in [27, 32, 53] (see also [10, 11, 17, 27, 42]). These flows are called *extended flows* by Getzler [32] and Zhang [53]. They can be defined using the Lax pair formalism [11] as follows:

(8) 
$$\frac{\partial L}{\partial b_i} = \frac{2}{i!} \left[ \left( L^i (\log L - c_i) \right)_+, L \right], \quad i \ge 0,$$

where  $c_i := \sum_{j=1}^{i} \frac{1}{j}$  are harmonic numbers and for the definition of log L see [11, 42]. The flows (8) commute with the traditional flows (6) of the Toda lattice hierarchy, and they also pairwise commute. All-together, (6), (8) form the *extended Toda hierarchy* (ETH) [11, 32, 53].

Let  $(v(\mathbf{b} + x\mathbf{1}, \mathbf{m}), w(\mathbf{b} + x\mathbf{1}, \mathbf{m}))$  be an arbitrary solution to the ETH, and  $\tau(\mathbf{b}+x\mathbf{1}, \mathbf{m})$  the tau-function [11] of this solution. Here  $\mathbf{b} = (b_0, b_1, b_2, ...)$  and  $\mathbf{1} = (1, 0, 0, ...)$ . The following theorem generalizes Theorem 1.1.

**Theorem 1.2.** The tau-function  $\tau(\mathbf{b} + x\mathbf{1}, \mathbf{m})$  for the ETH is a KP tau-function for any x and b, where m and p are related by (7).

The proof is in Section 4.

It was conjectured in [17, 18, 27, 32], and was proved by Dubrovin and Zhang [26] that the partition function  $Z^{\mathbb{P}^1}(\mathbf{b} + x\mathbf{1}, \mathbf{m})$  of Gromov– Witten (GW) invariants of  $\mathbb{P}^1$  (see Section 5 for the definition) is the tau-function of a particular solution to the ETH. (Independently, Okounkov and Pandharipande [44, 45] proved that  $Z^{\mathbb{P}^1}(x\mathbf{1}, \mathbf{m})$  satisfies the bilinear equations for the 1-Toda hierarchy.) Based on the result of Dubrovin and Zhang and on Theorem 1.2 we immediately obtain the following corollary.

**Corollary 1.3.** The partition function  $Z^{\mathbb{P}^1}(\mathbf{b} + x\mathbf{1}, \mathbf{m})$  of GW invariants of  $\mathbb{P}^1$  is a KP tau-function for any x and b, where **m** is related to **p** by (7). In particular,  $Z^{\mathbb{P}^1}(x\mathbf{1}, \mathbf{m})$  is a KP tau-function for any x.

The stationary sector of the  $\mathbb{P}^1$ -partition function is  $Z^{\mathbb{P}^1}(0\mathbf{1}, \mathbf{m})$ . We compute in Section 5 its affine coordinates by two different methods: One by the results of [23], the other by the results of [44]. The former gives us an explicit closed formula, the latter gives an expression in terms of characters of the symmetric groups. We then get a closed formula for some Plancherel averages in Corallary 5.8.

The rest of the paper is organized as follows. In Section 2 we recall the formula of n-point function for a KP tau-function. In Section 3 we show that an arbitrary tau-function for the Toda lattice hierarchy is a KP tau-function with the ground ring being the ring of functions of the space variable of Toda. The generalization to ETH is given in Section 4. In Section 5 we give an application of Corollary 1.3.

Acknowledgements. One of the authors D.Y. is grateful to Marco Bertola, Boris Dubrovin and Youjin Zhang for their advice. The work is supported by NSFC (No. 12371254, No. 11890662, No. 12061131014) and CAS No. YSBR-032.

# 2. The formula for the n-point function for a KP tau-function

In this section we first review the explicit formula (see Theorem A below) obtained by the second-named author for the *n*-point function for an arbitrary KP tau-function in the big cell, and then consider the converse statement that gives a criterion for a KP tau-function.

Before entering into the details let us introduce some *notations*. Denote by  $\mathcal{R}$  a suitable ground ring. For a formal power series  $F(\mathbf{p}) \in \mathcal{R}[[\mathbf{p}]]$ , define the *n*-point function associated with  $F(\mathbf{p})$  by

(9) 
$$G_n(\xi_1,\ldots,\xi_n) = \sum_{k_1,\ldots,k_n \ge 1} \prod_{i=1}^n \frac{k_i}{\xi_i^{k_i+1}} \cdot \frac{\partial^n F(\mathbf{p})}{\partial p_{k_1} \ldots \partial p_{k_n}} \bigg|_{\mathbf{p}=\mathbf{0}}, \quad n \ge 0.$$

By a partition  $\mu = (\mu_1, \mu_2, ...)$ , we mean a sequence of weakly decreasing non-negative integers with  $\mu_k = 0$  for sufficiently large k. The length  $\ell(\mu)$  is the number of the non-zero parts of  $\mu$  and the weight  $|\mu| := \mu_1 + \mu_2 + ...$  The set of all partitions will be denoted by  $\mathcal{P}_d$ . The Schur polynomial  $s_{\mu}(\mathbf{p})$  associated to  $\mu \in \mathcal{P}$  is a polynomial in the variables  $\mathbf{p} = (p_1, p_2, \ldots)$ , defined by

(10) 
$$s_{\mu}(\mathbf{p}) := \det_{1 \le i, j \le \ell(\lambda)} (h_{\lambda_i - i + j}(\mathbf{p})),$$

where  $h_j(\mathbf{p})$  are polynomials defined by the generating function

(11) 
$$\sum_{j=0}^{\infty} h_j(\mathbf{p}) z^j := e^{\sum_{k=1}^{\infty} p_k z^k}.$$

One constructive way to describe tau-functions of the KP hierarchy is to use Sato's Grassmannian. Let  $z^{1/2} \mathcal{R}[[z, z^{-1}]]$  be the space of formal series in z of half integral powers. An element V in the big cell  $Gr_0$  of Sato's Grassmannian is specified by a sequence of series

(12) 
$$\Psi_j(z) = z^{j+1/2} + \sum_{i=0}^{\infty} A_{i,j} z^{-i-1/2}, \quad j \ge 0,$$

where the coefficients  $A_{i,j}$  are called the *affine coordinates* of V. To any element  $V \in Gr_0$ , there corresponds a particular KP tau-function  $Z_V$  defined by

(13) 
$$Z_V := \sum_{\lambda \in \mathcal{P}} c_\lambda s_\lambda(\mathbf{p}),$$

where, for a partition  $\lambda$  written in terms of the Frobenius notation [38]  $\lambda = (m_1, \ldots, m_k | n_1, \ldots, n_k),$ 

(14) 
$$c_{\lambda} := (-1)^{n_1 + \dots + n_k} \det_{1 \le i, j \le k} (A_{m_i, n_j}).$$

The following theorem was obtained in [55].

**Theorem A** ([55]). The *n*-point function associated with  $\log Z_V$  is given by the following formula: For n = 1,

(15) 
$$G_1(\xi) = \sum_{i,j\ge 0} A_{i,j} \xi^{-i-j-2},$$

and for  $n \geq 2$ ,

(16) 
$$G_n(\xi_1, \dots, \xi_n) = (-1)^{n-1} \sum_{n \text{-cycles}} \prod_{i=1}^n B(\xi_{\sigma(i)}, \xi_{\sigma(i+1)}) - \frac{\delta_{n,2}}{(\xi_1 - \xi_2)^2},$$

where

(17) 
$$B(\xi_i, \xi_j) = \begin{cases} i_{\xi_i, \xi_j} \frac{1}{\xi_i - \xi_j} + A(\xi_i, \xi_j), & i < j, \\ A(\xi_i, \xi_i), & i = j, \\ i_{\xi_j, \xi_i} \frac{1}{\xi_i - \xi_j} + A(\xi_i, \xi_j), & i > j, \end{cases}$$

and

(18) 
$$A(\xi,\eta) = \sum_{i,j\ge 0} A_{i,j}\xi^{-j-1}\eta^{-i-1}$$

Here we have used the following notation:

(19) 
$$\iota_{x,y}\frac{1}{(x-y)^n} = -\sum_{k\geq 0} \binom{-n}{k} x^{-n-k} y^k.$$

Theorem A enables one to easily compute the *n*-point function once we have found the affine coordinates  $A_{i,j}$ . For example, there is only one 2-cycle, so the two-point function is given by

$$G_2(\xi_1,\xi_2) = \frac{A(\xi_1,\xi_2) - A(\xi_2,\xi_1)}{\xi_1 - \xi_2} - A(\xi_1,\xi_2) \cdot A(\xi_2,\xi_1).$$

Let us consider the converse of the above Theorem A. Recalling the simple fact that multiplying a KP tau-function  $\tau(\mathbf{p})$  by the exponential of an arbitrary affine linear function of  $\mathbf{p}$ 

(20) 
$$\tau(\mathbf{p}) \cdot e^{C_0 + \sum_{k \ge 1} C_k p_k}, \quad a_k \in \mathcal{R} \ (k \ge 0),$$

produces again a KP-tau function (which can be verified directly using (4)), we immediately obtain from Theorem A the following

**Corollary 2.1.** Let F be a formal power series in  $\mathbf{p}$  and  $G_n(\xi_1, \ldots, \xi_n)$  its n-point function. If there are  $(A_{i,j})_{i,j\geq 0}$  such that for all  $n \geq 2$   $G_n$  are given by (16), then  $Z = e^F$  is a tau-function of the KP hierarchy.

We note that Corollary 2.1 is also given in [3].

The next proposition describes how the affine coordinates change under (20).

**Proposition 2.2.** Let V, V' be two elements in the big cell having affine coordinates  $A_{i,j}, A'_{i,j}$ , respectively, and let  $Z_V, Z_{V'}$  be the KP tau-functions associated to V, V'. If there exist  $C_1, C_2, \ldots$  such that

(21) 
$$Z_{V'} = e^{\sum_{k \ge 1} C_k p_k} Z_V,$$

then we have the following identity: (22)

$$\sum_{i,j\geq 0} A'_{i,j} \xi^{-j-1} \eta^{-i-1} + \frac{1}{\xi - \eta} = \left( \sum_{i,j\geq 0} A_{i,j} \xi^{-j-1} \eta^{-i-1} + \frac{1}{\xi - \eta} \right) \frac{e^{-\sum_{\ell\geq 1} C_\ell \xi^{-\ell}}}{e^{-\sum_{\ell\geq 1} C_\ell \eta^{-\ell}}}$$

*Proof.* By direct verifications using (15).

In this section, we show that an arbitrary tau-function for the Toda lattice hierarchy is a KP tau-function.

Let  $(v(x, \mathbf{m}), w(x, \mathbf{m}))$  be an arbitrary power-series-in-**m** solution to the Toda lattice hierarchy (6), and  $\tau(x, \mathbf{m})$  the tau-function [11, 26, 21] of this solution. The solution  $(v(x, \mathbf{m}), w(x, \mathbf{m}))$  is in one-to-one correspondence with its initial value

(23) 
$$v(x, \mathbf{0}) =: f(x), \quad w(x, \mathbf{0}) =: g(x).$$

Denote by  $L_{ini}$  the initial Lax operator:

(24) 
$$L_{\rm ini} = \Lambda + f(x) + g(x)\Lambda^{-1},$$

and denote

(25) 
$$s(x) = -(1 - \Lambda^{-1})^{-1} (\log g(x)).$$

Recall from [50] that two elements

(26) 
$$\psi_1(\lambda, x) = (1 + O(\lambda^{-1}))\lambda^x, \quad \psi_2(\lambda, x) = (1 + O(\lambda^{-1}))e^{-s(x)}\lambda^{-x}$$

are called forming a pair of wave functions of  $L_{\rm ini}$  if

(27) 
$$L_{\text{ini}}(\psi_1) = \lambda \psi_1, \quad L_{\text{ini}}(\psi_2) = \lambda \psi_2,$$

(28) 
$$d(\lambda, x) := \psi_1 \Lambda^{-1}(\psi_2) - \psi_2 \Lambda^{-1}(\psi_1) = \lambda e^{-s(x-1)}.$$

The following formula is proved in [50]:

(29) 
$$\sum_{i_1,\dots,i_n\geq 0} \frac{(i_1+1)!\cdots(i_n+1)!}{\lambda_1^{i_1+2}\dots\lambda_n^{i_n+2}} \frac{\partial^n\log\tau(x,\mathbf{m})}{\partial m_{i_1}\dots\partial m_{i_n}} \bigg|_{\mathbf{m}=\mathbf{0}}$$
$$= (-1)^{n-1} \frac{e^{ns(x-1)}}{\prod_{j=1}^n \lambda_j} \sum_{n-\text{cycles}} \prod_{j=1}^n D(\lambda_{\sigma(j)},\lambda_{\sigma(j+1)};x) - \frac{\delta_{n,2}}{(\lambda-\mu)^2},$$

where  $n \geq 2$ , and

(30) 
$$D(\lambda,\mu;x) := \frac{\psi_1(\lambda,x)\psi_2(\mu,x-1) - \psi_1(\lambda,x-1)\psi_2(\mu,x)}{\lambda - \mu}.$$

Remark 3.1. The function  $D(\lambda, \mu; x)$  looks very similar to the Christoffel– Darboux kernel in matrix models [14, 41]. We hope to study their relations in a future work.

In [50] the following expansion is also observed:

(31) 
$$\frac{e^{s(x-1)}}{\mu}D(\lambda,\mu;x)\frac{\lambda^x}{\mu^x} = \frac{1}{\lambda-\mu} + \sum_{i,j\ge 0}\frac{A_{i,j}(x)}{\lambda^{j+1}\mu^{i+1}} =: B(\lambda,\mu;x).$$

We note that the expansion given in [50] is slightly weaker than (31). In the next section we will give a complete proof of a generalized version of (31).

Using (31) and (29) we find

(32) 
$$\sum_{i_1,\dots,i_n\geq 0} \frac{(i_1+1)!\cdots(i_n+1)!}{\lambda_1^{i_1+2}\cdots\lambda_n^{i_n+2}} \frac{\partial^n\log\tau(x,\mathbf{m})}{\partial m_{i_1}\cdots\partial m_{i_n}}\Big|_{\mathbf{m}=\mathbf{0}}$$
$$= (-1)^{n-1}\sum_{n\text{-cycles}} \prod_{j=1}^n B(\lambda_{\sigma(j)},\lambda_{\sigma(j+1)}) - \frac{\delta_{n,2}}{(\lambda-\mu)^2}, \quad n\geq 2.$$

(See also the Corollary 1 of [50].)

*Proof of Theorem 1.1.* By using (31), (32) and Corollary 2.1.

If we write

(33) 
$$\tau(x,\mathbf{m}) = \tau_0(x)\tau_1(x,\mathbf{m}), \quad \tau_1(x,\mathbf{m}=\mathbf{0}) \equiv 1$$

then by Theorem 1.1 we know that  $\tau_1(x, \mathbf{m})$  is a KP tau-function in the big cell. The factor  $\tau_0(x)$  can [11, 50] be determined by

(34) 
$$\frac{\tau_0(x+1)\tau_0(x-1)}{\tau_0(x)^2} = g(x) = e^{s(x-1)-s(x)}.$$

This factor can be identified with the *correction factor* in [21, 50, 52].

Let  $\tilde{\tau}_1(x, \mathbf{m})$  be the KP tau-function associated to the point in Sato's Grassmannian having the affine coordinates  $A_{i,j}(x)$ , with  $A_{i,j}(x)$  given in (31). Then there exists  $a_i(x)$ ,  $i \geq 0$ , such that

(35) 
$$\tau_1(x,\mathbf{m}) = e^{\sum_{i\geq 0} a_i(x)m_i} \tilde{\tau}_1(x,\mathbf{m}).$$

By further applying Proposition 2.2, we get the affine coordinates for  $\tau_1(x, \mathbf{m})$ .

### 4. EXTENDED TODA FLOWS IN THE KP HIERARCHY

The goal of this section is to prove Theorem 1.2.

As in Introduction, for an arbitrary solution  $(v(\mathbf{b} + x\mathbf{1}, \mathbf{m}), w(\mathbf{b} + x\mathbf{1}, \mathbf{m}))$  to the ETH, let  $\tau(\mathbf{b} + x\mathbf{1}, \mathbf{m})$  be the tau-function (again in the sense of [11, 26]) of the solution. Let us also denote

(36) 
$$\sigma(x, \mathbf{b}, \mathbf{m}) = \log \frac{\tau(\mathbf{b} + x\mathbf{1}, \mathbf{m})}{\tau(\mathbf{b} + (x+1)\mathbf{1}, \mathbf{m})}.$$

4.1. The matrix-resolvent method. The matrix-resolvent method for studying tau-functions for the Toda lattice hierarchy was developed in [21] (see also [50]). By the locality nature of this method, the same formulation as for the Toda lattice hierarchy applies to the ETH.

Denote by

(37) 
$$\mathcal{A} = \mathbb{Z}[v(x), w(x), v(x \pm 1), w(x \pm 1), v(x \pm 2), w(x \pm 2), \dots]$$

the polynomial ring. Recall that the *basic matrix resolvent*  $R(\lambda)$  is defined [21] as the unique element in Mat $(2, \mathcal{A}[[\lambda^{-1}]])$  satisfying

(38) 
$$\Lambda(R(\lambda))U(\lambda) - U(\lambda)R(\lambda) = 0$$

(39) 
$$\operatorname{tr} R(\lambda) = 1, \quad \det R(\lambda) = 0,$$

(40) 
$$R(\lambda) - \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \in \operatorname{Mat}(2, \mathcal{A}[[\lambda^{-1}]]\lambda^{-1}),$$

where

(41) 
$$U(\lambda) := \begin{pmatrix} V(x) - \lambda & W(x) \\ -1 & 0 \end{pmatrix}.$$

Define  $S_i = \frac{1}{(i+1)!} \operatorname{Coef} \left( \Lambda(R(\lambda)_{21}), \lambda^{-i-2} \right) \in \mathcal{A}, i \geq 0$ , and a sequence of elements  $\Omega_{i,j} \in \mathcal{A}$  by

(42) 
$$\sum_{i,j\geq 0} \frac{(i+1)!(j+1)!}{\lambda^{i+2}\mu^{j+2}} \Omega_{i,j} = \frac{\operatorname{tr} R(\lambda)R(\mu)}{(\lambda-\mu)^2} - \frac{1}{(\lambda-\mu)^2}.$$

According to [21], the above-defined  $(\Omega_{i,j}, S_i)$  gives rise to part of the canonical tau-structure for the ETH in [11, 26]. Indeed, it is shown in [21] that  $(\Omega_{i,j}, S_i)$  is associated to the tau-symmetric hamiltonian densities  $h_{\alpha,p}$ ,  $\alpha = 1, 2, p \ge -1$ , [11, 26] for the Toda lattice hierarchy. More precisely, the following identities hold for an arbitrary solution  $(v(\mathbf{b} + x\mathbf{1}, \mathbf{m}), w(\mathbf{b} + x\mathbf{1}, \mathbf{m}))$  to the ETH:

$$(43)$$

$$\frac{\partial^2 \log \tau(\mathbf{b} + x\mathbf{1}, \mathbf{m})}{\partial m_i \partial m_j} = \Omega_{i,j}(\mathbf{b} + x\mathbf{1}, \mathbf{m}),$$

$$(44)$$

$$\frac{\partial}{\partial m_i} \left( \log \frac{\tau(\mathbf{b} + (x+1)\mathbf{1}, \mathbf{m})}{\tau(\mathbf{b} + x\mathbf{1}, \mathbf{m})} \right) = S_i(\mathbf{b} + x\mathbf{1}, \mathbf{m}) = h_{2,i-1}(\mathbf{b} + x\mathbf{1}, \mathbf{m}),$$

$$(45)$$

$$\frac{\tau(\mathbf{b} + (x+1)\mathbf{1}, \mathbf{m})\tau(\mathbf{b} + (x-1)\mathbf{1}, \mathbf{m})}{\tau(\mathbf{b} + x\mathbf{1}, \mathbf{m})^2} = w(\mathbf{b} + x\mathbf{1}, \mathbf{m});$$

moreover, according to [21], for any  $n \ge 2$ ,

(46) 
$$\sum_{i_1,\ldots,i_n\geq 0} \prod_{j=1}^n \frac{(i_j+1)!}{\lambda_j^{i_j+2}} \frac{\partial^n \log \tau(\mathbf{b}+x\mathbf{1},\mathbf{m})}{\partial m_{i_1}\dots \partial m_{i_n}}$$
$$= -\sum_{n\text{-cycles}} \frac{\operatorname{tr} \prod_{j=1}^n R(\lambda_{\sigma(j)})}{\prod_{j=1}^n (\lambda_{\sigma(j)} - \lambda_{\sigma(j+1)})} - \frac{\delta_{n,2}}{(\lambda - \mu)^2}.$$

Note that by (36) and (44), (45) we have

(47) 
$$-\frac{\partial\sigma(x,\mathbf{b},\mathbf{m})}{\partial m_i} = S_i(\mathbf{b} + x\mathbf{1},\mathbf{m}),$$

(48) 
$$e^{\sigma(x-1,\mathbf{b},\mathbf{m})-\sigma(x,\mathbf{b},\mathbf{m})} = w(\mathbf{b}+x\mathbf{1},\mathbf{m}).$$

It is also clear from (34) that  $\sigma(x, b_0 = 0, b_1 = 0, ..., \mathbf{m} = \mathbf{0}) = s(x)$ .

4.2. Wave functions for ETH and the KP hierarchy. It is shown in [10] (see also [12, 42]) that there exist dressing operators P, Q of the form

(49) 
$$P = \sum_{k \ge 0} P_k(x, \mathbf{b}, \mathbf{m}) \Lambda^{-k}, \quad P_0(x, \mathbf{b}, \mathbf{m}) \equiv 1,$$

(50) 
$$Q = \sum_{k \ge 0} Q_k(x, \mathbf{b}, \mathbf{m}) \Lambda^k,$$

such that

(51) 
$$L = P \circ \Lambda \circ P^{-1} = Q \circ \Lambda^{-1} \circ Q^{-1},$$
$$\partial P \qquad 1$$

(52) 
$$\frac{\partial P}{\partial m_i} = -\frac{1}{(i+1)!} \left( L^{i+1} \right)_- \circ P,$$

(53) 
$$\frac{\partial P}{\partial b_i} = -\frac{2}{i!} \left( L^i (\log L - c_i) \right)_{-} \circ P,$$

(54) 
$$\frac{\partial Q}{\partial m_i} = \frac{1}{(i+1)!} \left( L^{i+1} \right)_+ \circ Q,$$

(55) 
$$\frac{\partial Q}{\partial b_i} = \frac{2}{i!} \left( L^i (\log L - c_i) \right)_+ \circ Q.$$

Define

(56) 
$$\psi_1(\lambda; x, \mathbf{b}, \mathbf{m}) := P\left(\lambda^x e^{\sum_{i\geq 0} \frac{2}{i!} b_i \lambda^i (\log \lambda - c_i) + \sum_{i\geq 0} \frac{m_i}{(i+1)!} \lambda^{i+1}}\right),$$
  
(57) 
$$\psi_2(\lambda; x, \mathbf{b}, \mathbf{m}) := Q\left(\lambda^{-x} e^{-\sum_{i\geq 0} \frac{2}{i!} b_i \lambda^i (\log \lambda - c_i) - \sum_{i\geq 0} \frac{m_i}{(i+1)!} \lambda^{i+1}}\right).$$

Then  $\psi_a = \psi_a(\lambda; x, \mathbf{b}, \mathbf{m}), a = 1, 2$ , satisfy the following equations:

(58) 
$$L\psi_a = \lambda\psi_a, \quad a = 1, 2,$$

(59) 
$$\frac{\partial \psi_1}{\partial m_i} = \frac{1}{(i+1)!} (L^{i+1})_+ \psi_1,$$

(60) 
$$\frac{\partial \psi_1}{\partial b_i} = \frac{2}{i!} L^i (\log L - c_i)_+ \psi_1,$$

(61) 
$$\frac{\partial \psi_2}{\partial m_i} = -\frac{1}{(i+1)!} (L^{i+1})_- \psi_2,$$

(62) 
$$\frac{\partial \psi_2}{\partial b_i} = -\frac{2}{i!} L^i (\log L - c_i)_- \psi_2$$

Moreover,  $\psi_1$ ,  $\psi_2$  have the form

(63)

$$\psi_1 = \phi_1(\lambda; x, \mathbf{b}, \mathbf{m}) \lambda^x e^{\sum_{i \ge 0} \frac{2}{i!} b_i \lambda^i (\log \lambda - c_i) + \sum_{i \ge 0} \frac{m_i}{(i+1)!} \lambda^{i+1}},$$

(64)

$$\psi_2 = \phi_2(\lambda; x, \mathbf{b}, \mathbf{m}) e^{-\sigma(x, \mathbf{b}, \mathbf{m})} \lambda^{-x} e^{-\sum_{i \ge 0} \frac{2!}{i!} b_i \lambda^i (\log \lambda - c_i) - \sum_{i \ge 0} \frac{m_i}{(i+1)!} \lambda^{i+1}}$$

where

(65) 
$$\phi_1(\lambda; x, \mathbf{b}, \mathbf{m}) = \sum_{k \ge 0} P_k(x, \mathbf{b}, \mathbf{m}) \lambda^{-k},$$

(66) 
$$\phi_2(\lambda; x, \mathbf{b}, \mathbf{m}) = \sum_{k \ge 0} B_k(x, \mathbf{b}, \mathbf{m}) \lambda^{-k}.$$

We recall that  $P_k(x, \mathbf{b}, \mathbf{m})$  are given in (49), and we also note using (50) that  $B_k(x, \mathbf{b}, \mathbf{m}) = Q_k(x, \mathbf{b}, \mathbf{m})e^{\sigma(x, \mathbf{b}, \mathbf{m})}, k \ge 0$ , and  $B_0(x, \mathbf{b}, \mathbf{m}) \equiv 1$ . We call  $\psi_1$  the wave function of type A and  $\psi_2$  the wave function of type B, associated to the solution  $(v(x, \mathbf{b}, \mathbf{m}), w(x, \mathbf{b}, \mathbf{m}))$ . Denote

(67)  $d(\lambda; x, \mathbf{b}, \mathbf{m})$   $:= \psi_1(\lambda; x, \mathbf{b}, \mathbf{m})\psi_2(\lambda; x - 1, \mathbf{b}, \mathbf{m}) - \psi_1(\lambda; x - 1, \mathbf{b}, \mathbf{m})\psi_2(\lambda; x, \mathbf{b}, \mathbf{m}).$ 

The following lemma is important.

**Lemma 4.1.** For any  $i \ge 0$  we have the identities:

(68) 
$$\frac{\partial \left(e^{\Lambda^{-1}(\sigma(x,\mathbf{b},\mathbf{m}))}d(\lambda;x,\mathbf{b},\mathbf{m})\right)}{\partial b_i} = 0,$$

(69) 
$$\frac{\partial \left(e^{\Lambda^{-1}(\sigma(x,\mathbf{b},\mathbf{m}))}d(\lambda;x,\mathbf{b},\mathbf{m})\right)}{\partial m_i} = 0.$$

*Proof.* We first introduce some notations:

(70) 
$$R_j := \operatorname{res} L^j, \quad N_j = \operatorname{Coef}(L^j, \Lambda^{-1}),$$

(71)  $r_j := \operatorname{res} \log L \circ L^j, \quad n_j = \operatorname{Coef}(\log L \circ L^j, \Lambda^{-1}).$ 

(Warning: avoid from confusing the notation  $R_j$  with the notation for the matrix resolvent.) For examples [10],

(72) 
$$R_0 = 1, \quad R_1 = v, \quad N_0 = 0, \quad N_1 = w$$

(73) 
$$r_0 = \frac{1}{2}\Lambda \circ \frac{\partial}{\Lambda - 1}(v), \quad n_0 = \frac{1}{2}\frac{\partial}{\Lambda - 1}(\log w).$$

Here we omitted the arguments in  $R_j, r_j, N_j, n_j, v, w$ . Below, when no ambiguity will occur, we often do this type of omission.

As it was given in [11], the tau-symmetric hamiltonian densities  $h_{\alpha,p}$ for the ETH are related to  $R_j$ ,  $r_j$  by

(74) 
$$h_{2,p} = \frac{1}{(p+2)!} R_{p+2},$$

(75) 
$$h_{1,p} = \frac{2}{(p+1)!} r_{p+1} - \frac{2c_{p+1}}{(p+1)!} R_{p+1}, \qquad p \ge -1.$$

We also recall from [11] the following formula:

(76) 
$$h_{\alpha,i-1} = (\Lambda - 1)\frac{\partial \log \tau}{\partial t^{\alpha,i}} = -\frac{\partial \sigma}{\partial t^{\alpha,i}}, \quad i \ge 0.$$

Let us now prove (69). (Although (69) was already proved in [50], our proof here will be slightly different from [50] and can be generalized to proving (68).) We have

(77) 
$$e^{-\Lambda^{-1}(\sigma)}\frac{\partial(e^{\Lambda^{-1}(\sigma)}d)}{\partial m_i} = \frac{\partial\Lambda^{-1}(\sigma)}{\partial m_i}d + \frac{\partial d}{\partial m_i}, \quad i \ge 0.$$

Denote

(78) 
$$B_i := -(L^{i+1})_-, \quad i \ge -1.$$

Then for all  $i \ge 0$  we have

$$B_{i} = -(L^{i} \circ L)_{-} = -((L^{i})_{+} + (L^{i})_{-}) \circ L)_{-}$$
  
=  $-(R_{i}L)_{-} - ((L^{i})_{-} \circ L)_{-}$   
=  $B_{i-1} \circ L + N_{i} - R_{i}w\Lambda^{-1} = \cdots =$   
=  $\sum_{j=0}^{i} (N_{j} - R_{j}w\Lambda^{-1}) \circ L^{i-j}.$ 

(The resulting equality is also valid for i = -1 as  $B_{-1} = 0$ .) Thus,

(79) 
$$(i+1)! \frac{\partial \psi_2}{\partial m_i} = B_i(\psi_2) = \sum_{j=0}^i \lambda^{i-j} (N_j - R_j w \Lambda^{-1})(\psi_2),$$

and

(80) 
$$(i+1)! \frac{\partial \psi_1}{\partial m_i} = \lambda^{i+1} \psi_1 + \sum_{j=0}^i \lambda^{i-j} (N_j - R_j w \Lambda^{-1})(\psi_1).$$

By substituting (80), (79) in (77) and by a lengthy but straightforward calculation using also (58), (47), we obtain

$$e^{-\Lambda^{-1}(\sigma)} \frac{\partial (e^{\Lambda^{-1}(\sigma)}d)}{\partial m_i} = -\Lambda^{-1}(S_i)d + \frac{\Lambda^{-1}(R_{i+1})}{(i+1)!}d + \frac{d}{(i+1)!} \sum_{j=0}^i \lambda^{i-j} \left(N_j + \Lambda^{-1}(N_j) + \Lambda^{-1}(vR_j - R_{j+1})\right) = 0.$$

Note that in the last equality we used (44), (74) and the relation  $L^{k+1} = L \circ L^k = L^k \circ L$  (which implies the vanishing of each summand in the above  $\sum_j$ ).

We proceed to prove (68). Like in [50] we note that

$$\Lambda(e^{\Lambda^{-1}(\sigma(x,\mathbf{b},\mathbf{m}))}d(\lambda;x,\mathbf{b},\mathbf{m}))$$
  
=  $e^{\sigma}(\Lambda(\psi_1)\psi_2 - \psi_1\Lambda(\psi_2))$   
=  $e^{\sigma}\{((\lambda - v)\psi_1 - w\Lambda^{-1}(\psi_1))\psi_2 - \psi_1((\lambda - v)\psi_2 - w\Lambda^{-1}(\psi_2))\}$   
=  $e^{\Lambda^{-1}(\sigma(x,\mathbf{b},\mathbf{m}))}d(\lambda;x,\mathbf{b},\mathbf{m})$ 

(the last equality used (48)). It follows that

$$\partial_x(e^{\Lambda^{-1}(\sigma(x,\mathbf{b},\mathbf{m}))}d(\lambda;x,\mathbf{b},\mathbf{m})) = \frac{\partial_x}{\Lambda - 1} \circ (\Lambda - 1)(e^{\Lambda^{-1}(\sigma(x,\mathbf{b},\mathbf{m}))}d(\lambda;x,\mathbf{b},\mathbf{m})) = 0.$$

Thus,

$$0 = e^{-\Lambda^{-1}(\sigma(\mathbf{b},\mathbf{t}))} \partial_x (e^{\Lambda^{-1}(\sigma(\mathbf{b},\mathbf{m}))} d(\lambda;\mathbf{b},\mathbf{m}))$$
  
=  $-\Lambda^{-1}(h_{1,-1}) + \partial_x (d(\lambda;\mathbf{b},\mathbf{m})) = -2\Lambda^{-1}(r_0)d + \partial_x (d(\lambda;\mathbf{b},\mathbf{m})).$ 

Denote

(81) 
$$Q_i = -2(L^i(\log L - c_i))_- = -2(L^i\log L)_0 - 2c_iB_{i-1}, \quad i \ge 0.$$
  
In the way similar to the derivation for  $B_i$ , we find

(82) 
$$Q_i = \sum_{j=0}^{i-1} (2m_j - 2r_j w \Lambda^{-1}) \circ L^{i-j} - 2(\log L)_{-} \circ L^i - 2c_i B_{i-1}.$$

Therefore,

(83) 
$$i! \frac{\partial \psi_2}{\partial b_i} = Q_i(\psi_2) = \sum_{j=0}^{i-1} \lambda^{i-j} (2m_j - 2r_j w \Lambda^{-1})(\psi_2) - 2(\log L)_-(\psi_2) - 2c_i \sum_{j=0}^{i-1} \lambda^{i-1-j} (N_j - R_j w \Lambda^{-1})(\psi_2),$$

and

(84) 
$$i! \frac{\partial \psi_1}{\partial b_i} = 2\lambda^i (\log \lambda - c_i) \psi_1 + \sum_{j=0}^{i-1} \lambda^{i-j} (2m_j - 2r_j w \Lambda^{-1}) (\psi_1) - 2(\log L)_-(\psi_1) - 2c_i \sum_{j=0}^{i-1} \lambda^{i-1-j} (N_j - R_j w \Lambda^{-1}) (\psi_1).$$

Using these relations and using (60), (62) with i = 0, and again by a lengthy calculation, we find

$$e^{-\Lambda^{-1}(\sigma)}\frac{\partial(e^{\Lambda^{-1}(\sigma)}d)}{\partial b_i} = -2\Lambda^{-1}(r_0)d + \partial_x(d) = 0.$$

The lemma is proved.

It follows from Lemma 4.1 and (63), (64) that

(85) 
$$d(\lambda; x, \mathbf{b}, \mathbf{m})) = \lambda e^{-\Lambda^{-1}(\sigma(x, \mathbf{b}, \mathbf{m}))} \sum_{k \ge 0} d_k \lambda^{-k},$$

where  $d_k, k \ge 0$ , are constants with  $d_0 = 1$ . Therefore, for any choice  $\psi_1$  of the wave functions of type A associated to  $(v(x, \mathbf{b}, \mathbf{m}), w(x, \mathbf{b}, \mathbf{m}))$ , there exists a choice,  $\psi_2$ , of the wave functions of type B associated to  $(v(x, \mathbf{b}, \mathbf{m}), w(x, \mathbf{b}, \mathbf{m}))$ , such that

(86) 
$$d(\lambda; x, \mathbf{b}, \mathbf{m}) = \lambda e^{-\Lambda^{-1}(\sigma(x, \mathbf{b}, \mathbf{m}))}$$

As a generalization of the terminology in [50], we say that  $\psi_1, \psi_2$  form a pair of wave functions associated to  $(v(\mathbf{b} + x\mathbf{1}, \mathbf{m}), w(\mathbf{b} + x\mathbf{1}, \mathbf{m}))$ if (86) holds.

(87)

$$D(\lambda, \mu; x, \mathbf{b}, \mathbf{m})$$
  
:=  $\frac{\psi_1(\lambda; x, \mathbf{b}, \mathbf{m})\psi_2(\mu; x - 1, \mathbf{b}, \mathbf{m}) - \psi_1(\lambda; x - 1, \mathbf{b}, \mathbf{m})\psi_2(\mu; x, \mathbf{b}, \mathbf{m})}{\lambda - \mu}$ ,

which coincides with the one introduced in [50] when restricted to  $b_2 = b_3 = \cdots = 0$ .

16

We arrive at the following theorem.

**Theorem 4.2.** The following identity holds:

(88) 
$$\sum_{i_1,\dots,i_n\geq 0} \prod_{j=1}^n \frac{(i_j+1)!}{\lambda_j^{i_j+2}} \frac{\partial^n \log \tau(\mathbf{b}+x\mathbf{1},\mathbf{m})}{\partial m_{i_1}\dots\partial m_{i_n}}$$
$$= (-1)^{n-1} \sum_{n-cycles} \prod_{j=1}^n B(\lambda_{\sigma(j)},\lambda_{\sigma(j+1)};x,\mathbf{b},\mathbf{m}) - \frac{\delta_{n,2}}{(\lambda-\mu)^2},$$

where  $n \geq 2$  and

(89) 
$$B(\lambda,\mu;x,\mathbf{b},\mathbf{m})$$
$$:=\frac{e^{\sigma(x-1,\mathbf{b},\mathbf{m})}}{\mu}D(\lambda,\mu;x,\mathbf{b},\mathbf{m})\frac{e^{\sum_{i\geq 0}\frac{2}{i!}b_i\lambda^i(\log\lambda-c_i)+\sum_{i\geq 0}\frac{m_i}{(i+1)!}\lambda^{i+1}}}{e^{\sum_{i\geq 0}\frac{2}{i!}b_i\mu^i(\log\mu-c_i)+\sum_{i\geq 0}\frac{m_i}{(i+1)!}\mu^{i+1}}}.$$

Proof. Let

(90) 
$$\Psi_{\text{pair}}(\lambda; x, \mathbf{b}, \mathbf{m}) := \begin{pmatrix} \psi_1(\lambda; x, \mathbf{b}, \mathbf{m}) & \psi_2(\lambda; x, \mathbf{b}, \mathbf{m}) \\ \psi_1(\lambda; x - 1, \mathbf{b}, \mathbf{m}) & \psi_2(\lambda; x - 1, \mathbf{b}, \mathbf{m}) \end{pmatrix}$$

By using the arguments the same as in [50] one can show that

(91) 
$$R(\lambda; x, \mathbf{b}, \mathbf{m}) = \Psi_{\text{pair}}(\lambda; x, \mathbf{b}, \mathbf{m}) \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \Psi_{\text{pair}}(\lambda; x, \mathbf{b}, \mathbf{m})^{-1}$$

(see the Proposition 3 and its proof of [50]). It follows from (46), (91)that for  $n \geq 2$ ,

$$\sum_{i_1,\dots,i_n\geq 0} \prod_{j=1}^n \frac{(i_j+1)!}{\lambda_j^{i_j+2}} \frac{\partial^n \log \tau(\mathbf{b}+x\mathbf{1},\mathbf{m})}{\partial m_{i_1}\dots \partial m_{i_n}}$$
$$= (-1)^{n-1} \frac{e^{n\sigma(x-1,\mathbf{b},\mathbf{m})}}{\prod_{j=1}^n \lambda_j} \sum_{n-\text{cycles}} \prod_{j=1}^n D(\lambda_{\sigma(j)},\lambda_{\sigma(j+1)};x,\mathbf{b},\mathbf{m}) - \frac{\delta_{n,2}}{(\lambda-\mu)^2}.$$
This yields identity (88).

This yields identity (88).

By taking  $\mathbf{m} = \mathbf{0}$  we immediately obtain the following corollary.

# Corollary 4.3. We have

(93) 
$$\sum_{i_1,\dots,i_n\geq 0} \prod_{j=1}^n \frac{(i_j+1)!}{\lambda_j^{i_j+2}} \frac{\partial^n \log \tau(\mathbf{b}+x\mathbf{1},\mathbf{m})}{\partial m_{i_1}\dots \partial m_{i_n}} \bigg|_{\mathbf{m}=\mathbf{0}}$$
$$= (-1)^{n-1} \sum_{n-cycles} \prod_{j=1}^n B(\lambda_{\sigma(j)},\lambda_{\sigma(j+1)};x,\mathbf{b}) - \frac{\delta_{n,2}}{(\lambda-\mu)^2},$$

where  $B(\lambda, \mu; x, \mathbf{b}) := B(\lambda, \mu; x, \mathbf{b}, \mathbf{m} = \mathbf{0}).$ 

•

In terms of the functions  $\phi_1, \phi_2$  the pair condition (86) reads (94)  $\phi_1(\lambda; x, \mathbf{b}, \mathbf{m})\phi_2(\lambda; x-1, \mathbf{b}, \mathbf{m}) - w(x, \mathbf{b}, \mathbf{m})\lambda^{-2}\phi_2(\lambda; x, \mathbf{b}, \mathbf{m})\phi_1(\lambda; x-1, \mathbf{b}, \mathbf{m}) \equiv 1,$ and the function  $B(\lambda, \mu; \mathbf{b}, \mathbf{m})$  reads

$$B(\lambda,\mu;x,\mathbf{b},\mathbf{m})$$
  
:= 
$$\frac{\phi_1(\lambda;x,\mathbf{b},\mathbf{m})\phi_2(\mu;x-1,\mathbf{b},\mathbf{m}) - w(\mathbf{b},\mathbf{m})\lambda^{-1}\mu^{-1}\phi_1(\lambda;x-1,\mathbf{b},\mathbf{m})\phi_2(\mu;\mathbf{b},\mathbf{m})}{\lambda-\mu}$$

**Lemma 4.4.** The function  $B(\lambda, \mu; x, \mathbf{b}, \mathbf{m})$  admits the following expansion:

(96) 
$$B(\lambda,\mu;x,\mathbf{b},\mathbf{m}) = \frac{1}{\lambda-\mu} + \sum_{i,j\geq 0} \frac{A_{i,j}(x,\mathbf{b},\mathbf{m})}{\lambda^{j+1}\mu^{i+1}}$$

In particular,

(97) 
$$B(\lambda,\mu;x,\mathbf{b}) = \frac{1}{\lambda-\mu} + \sum_{i,j\geq 0} \frac{A_{i,j}(x,\mathbf{b})}{\lambda^{j+1}\mu^{i+1}},$$

where  $A_{i,j}(x, \mathbf{b}) = A_{i,j}(x, \mathbf{b}, \mathbf{m} = \mathbf{0}).$ 

*Proof.* Recall the following identity:

(98) 
$$\phi_2(\mu) = \phi_2(\lambda) + \phi'_2(\lambda)(\mu - \lambda) + (\mu - \lambda)^2 \partial_\lambda \left(\frac{\phi_2(\lambda) - \phi_2(\mu)}{\lambda - \mu}\right),$$

where we omit the arguments  $\mathbf{b}, \mathbf{m}$  from  $\phi_2(\lambda; \mathbf{b}, \mathbf{m})$ . Similarly, (99)

$$\frac{\phi_2(\mu)}{\mu} = \frac{\phi_2(\lambda)}{\lambda} + \partial_\lambda \left(\frac{\phi_2(\lambda)}{\lambda}\right) (\mu - \lambda) + (\mu - \lambda)^2 \partial_\lambda \left(\frac{\phi_2(\lambda)/\lambda - \phi_2(\mu)/\mu}{\lambda - \mu}\right).$$

Substituting these identities in (95) and using (94), (65), (66), we find the validity of the expansion (96).

Proof of Theorem 1.2. By using (93), (97) and Theorem 2.1.

Like in the previous section, if we write

(100) 
$$\tau(\mathbf{b} + x\mathbf{1}, \mathbf{m}) = \tau_0(\mathbf{b} + x\mathbf{1})\tau_1(\mathbf{b} + x\mathbf{1}, \mathbf{m}), \quad \tau_1(\mathbf{b} + x\mathbf{1}, \mathbf{0}) \equiv 1,$$

then by Theorem 1.2 we know that  $\tau_1(\mathbf{b} + x\mathbf{1}, \mathbf{m})$  is a KP tau-function in the big cell. The factor  $\tau_0(\mathbf{b} + x\mathbf{1})$  can be determined from the definition of the tau-structure [11]. Let  $\tilde{\tau}_1(x, \mathbf{b}, \mathbf{m})$  be the KP taufunction associated to the point in the infinite Grassmannian with affine

coordinates  $A_{i,j}(x, \mathbf{b})$ , with  $A_{i,j}(x, \mathbf{b})$  given in (97). Then there exists  $a_i(x, \mathbf{b}), i \ge 0$ , such that

(101) 
$$\tau_1(\mathbf{b} + x\mathbf{1}, \mathbf{m}) = e^{\sum_{i \ge 0} a_i(x, \mathbf{b})m_i} \tilde{\tau}_1(x, \mathbf{b}, \mathbf{m}).$$

By further applying Proposition 2.2, we can find a formula for the affine coordinates for  $\tau_1(\mathbf{b} + x\mathbf{1}, \mathbf{m})$ , whose dependence on  $b_0, x$  is through  $b_0 + x$ .

## 5. Application to Gromov–Witten invariants of $\mathbb{P}^1$

In this section, we give an application of the above results regarding the so-called topological solution to the ETH.

Recall that the free energy  $\mathcal{F}^{\mathbb{P}^1} = \mathcal{F}^{\mathbb{P}^1}(\mathbf{b}, \mathbf{m})$  of GW invariants of  $\mathbb{P}^1$  is defined by

(102)

$$\mathcal{F}^{\mathbb{P}^1} = \sum_{k \ge 1} \frac{1}{k!} \sum_{\substack{1 \le \alpha_1, \dots, \alpha_k \le 2\\ i_1, \dots, i_k \ge 0}} \prod_{j=1}^k t^{\alpha_j, i_j} \sum_{g \ge 0} \sum_{d \ge 0} q^d \langle \tau_{i_1}(\alpha_1) \cdots \tau_{i_k}(\alpha_k) \rangle_{g, d}^{\mathbb{P}^1}$$

where  $b_i = t^{1,i}$ ,  $m_i = t^{2,i}$ ,  $i \ge 0$ , and  $\langle \tau_{i_1}(\alpha_1) \cdots \tau_{i_k}(\alpha_k) \rangle_{g,d}^{\mathbb{P}^1}$  are genus gand degree d GW invariants of  $\mathbb{P}^1$  [23, 26, 44, 45]. When  $\alpha_1 = \cdots = \alpha_k = 2$  the GW invariants  $\langle \tau_{i_1}(\alpha_1) \cdots \tau_{i_k}(\alpha_k) \rangle_{g,d}^{\mathbb{P}^1}$  are called in the *stationary sector* [44, 45]. The exponential

(103) 
$$\exp \mathcal{F}^{\mathbb{P}^1}(\mathbf{b}, \mathbf{m}) =: Z^{\mathbb{P}^1}(\mathbf{b}, \mathbf{m})$$

is called the *partition function of GW invariants of*  $\mathbb{P}^1$ . The restrictions

(104) 
$$\mathcal{F}^{\mathbb{P}^1}(\mathbf{b}=\mathbf{0},\mathbf{m}) =: \mathcal{F}^{\mathbb{P}^1}(\mathbf{m}), \quad Z^{\mathbb{P}^1}(\mathbf{b}=\mathbf{0},\mathbf{m}) =: Z^{\mathbb{P}^1}(\mathbf{m})$$

are called the free energy and respectively the partition function of stationary GW invariants of  $\mathbb{P}^1$ , which are of particular interest due to their closed formulas [23] and their relations to the representations of the symmetric group [44].

It is proved by Dubrovin and Zhang [26] that the partition function  $Z^{\mathbb{P}^1}(\mathbf{b} + x\mathbf{1}, \mathbf{m})$  of GW invariants of  $\mathbb{P}^1$  is a particular tau-function for the ETH. Then, as given in Introduction, the result of Dubrovin–Zhang together with Theorem 1.2 implies Corollary 1.3. As before, we can write the partition function  $Z^{\mathbb{P}^1}(\mathbf{b} + x\mathbf{1}, \mathbf{m})$  as follows:

(105) 
$$Z^{\mathbb{P}^1}(\mathbf{b} + x\mathbf{1}, \mathbf{m}) = Z_0^{\mathbb{P}^1}(\mathbf{b} + x\mathbf{1}) \cdot Z_1^{\mathbb{P}^1}(\mathbf{b} + x\mathbf{1}, \mathbf{m}),$$

where the factor  $Z_0^{\mathbb{P}^1}(\mathbf{b}+x\mathbf{1})$  can be determined by using the definition of tau-function for the ETH [26], and the factor  $Z_1^{\mathbb{P}^1}(\mathbf{b}+x\mathbf{1},\mathbf{m})$ , which satisfies  $Z_1^{\mathbb{P}^1}(\mathbf{b}+x\mathbf{1},\mathbf{0}) \equiv 1$ , is a particular KP tau-function in the big cell. Denote by  $A_{i,j}^{\mathbb{P}^1}(\mathbf{b} + x\mathbf{1})$  the affine coordinates of  $Z_1^{\mathbb{P}^1}(\mathbf{b} + x\mathbf{1}, \mathbf{m})$ . The restrictions  $A_{i,j}^{\mathbb{P}^1}(\mathbf{0})$  are of particular interest, which are the affine coordinates for the partition function  $Z^{\mathbb{P}^1}(\mathbf{m})$  of stationary GW invariants of  $\mathbb{P}^1$ . In the next two subsections we will derive two formulas for  $A_{i,j}^{\mathbb{P}^1}(\mathbf{0})$ .

5.1. Explicit generating series for  $A_{i,j}^{\mathbb{P}^1}(\mathbf{0})$ . It was proved in [23] (see also [22, 40, 50]) that, for each  $n \geq 2$ ,

(106) 
$$\sum_{i_1,\dots,i_n \ge 0} \prod_{j=1}^n \frac{(i_j+1)!}{\xi_j^{i_j+2}} \sum_{g\ge 0} \sum_{d\ge 0} q^d \left\langle \prod_{j=1}^n \tau_{i_j}(2) \right\rangle_{g,d}^{\mathbb{P}^1}$$
$$= (-1)^{n-1} \sum_{n-\text{cycles}} \prod_{j=1}^n B(\lambda_{\sigma(j)}, \lambda_{\sigma(j+1)}) - \frac{\delta_{n,2}}{(\lambda-\mu)^2},$$

where

(107)  
$$B(\lambda,\mu) = \frac{1}{\lambda-\mu} - \sum_{i,j\geq 0} \frac{(-1)^{j+1}}{\lambda^{j+1}\mu^{i+1}} \sum_{n\geq 1} \frac{q^k}{k!}$$
$$\cdot \sum_{1\leq i_1,j_1\leq k} (-1)^{i_1+j_1} \frac{(i_1+j_1-2k)_{k-1}(i_1-\frac{1}{2})^p (j_1-\frac{1}{2})^q}{(i_1-1)!(j_1-1)!(k-i_1)!(k-j_1)!}.$$

If we define  $A_{i,j}^{\mathbb{P}^{1,+}}$  via

(108) 
$$\sum_{i,j\geq 0} A_{i,j}^{\mathbb{P}^{1},+} \lambda^{-j-1} \eta^{-i-1} := B(\lambda,\mu) - \frac{1}{\lambda-\mu}$$

and let  $Z^{\mathbb{P}^{1,+}}(\mathbf{m})$  be the KP tau-function associated to the point in  $Gr_0$  with the affine coordinates  $A_{i,j}^{\mathbb{P}^{1,+}}$  (i.e., using formula (13)), then we know that  $Z^{\mathbb{P}^1}(\mathbf{m})$  and  $Z^{\mathbb{P}^{1,+}}(\mathbf{m})$  can only differ by multiplication by the exponential of a particular linear function of  $\mathbf{m}$ .

**Proposition 5.1.** We have

(109)

$$\log Z^{\mathbb{P}^{1,+}}(\mathbf{m}) = \sum_{n \ge 1} \frac{1}{n!} \sum_{i_1, \dots, i_n \ge 0} \prod_{j=1}^n m_{i_j} \sum_{g \ge 0} \sum_{d > 0} q^d \left\langle \prod_{j=1}^n \tau_{i_j}(2) \right\rangle_{g,d}^{\mathbb{P}^1}.$$

*Proof.* We already know that  $\log Z^{\mathbb{P}^1}(\mathbf{m}) - \log Z^{\mathbb{P}^{1,+}}(\mathbf{m})$  is a linear function of  $\mathbf{m}$ . This linear function could be calculated by using the formulas (31), (32), (34) of [23] and by performing the asymptotic

expansion of the digamma function:

(110) 
$$\log Z^{\mathbb{P}^1}(\mathbf{m}) - \log Z^{\mathbb{P}^1,+}(\mathbf{m}) = \sum_{k\geq 1} (1-2^{-k})\zeta(-k)\frac{m_{k-1}}{(k-1)!}$$

The proposition is then proved by identifying this linear function with the degree zero part [44] of stationary  $\mathbb{P}^1$  free energy.

Proposition 5.1 and Proposition 2.2 implies the following theorem.

**Theorem 5.2.** The generating series the affine coordinates  $A_{i,j}^{\mathbb{P}^1}(\mathbf{0})$  is the asymptotic expansion of (111)

$$\frac{1}{x-y} - D(x,y) \cdot \exp\left[\left(\log\frac{\Gamma(x+\frac{1}{2})}{\sqrt{2\pi}} - x\log x + x\right) - \left(\log\frac{\Gamma(y+\frac{1}{2})}{\sqrt{2\pi}} - y\log y + y\right)\right]$$

*Remark* 5.3. We consider the computations of  $\log Z_0^{\mathbb{P}^1}(\mathbf{b})$  and  $A_{i,j}^{\mathbb{P}^1}(\mathbf{b})$  by using Virasoro constraints [26] and the ETH in a future publication.

5.2. Affine coordinates from the Okounkov–Pandharipande formula. We proceed with recalling the Okounkov–Pandharipande formula [44] for  $Z^{\mathbb{P}^1}(\mathbf{m})$ , which is derived from the GW/H (Gromov– Witten/Hurwitz) correspondence established by Okounkov and Pandharipande [44].

To consider the GW invariants of  $\mathbb{P}^1$ , one has to consider all the moduli spaces  $\overline{\mathcal{M}}_{g,n}(\mathbb{P}^1, d)$  of stable maps, where g is the genus of the domain curve, n is the number of marked point on the domain curve, d is the degree of the map. For the purpose of GW/H correspondence, it is necessary to consider the n = 0 case. The expected dimension of  $\overline{\mathcal{M}}_{g,0}(\mathbb{P}^1, d)$  is 2g - 2 + 2d. When

(112) 
$$2g - 2 + 2d = 0,$$

we have a contribution of  $q^d$  to the free energy. This happens only when (g, d) = (0, 1) and (g, d) = (1, 0), The first case corresponds to  $\overline{\mathcal{M}}_{0,0}(\mathbb{P}^1, 1)$ , and the second case is impossible.

With the contribution of zero point correlators, the free energy of the stationary GW invariants of  $\mathbb{P}^1$  is then

(113) 
$$\tilde{\mathcal{F}}^{\mathbb{P}^1}(\mathbf{m}) = q + \mathcal{F}^{\mathbb{P}^1}(\mathbf{m}).$$

Let  $\tilde{Z}^{\mathbb{P}^1}(\mathbf{m}) = \exp \tilde{F}^{\mathbb{P}^1}(\mathbf{m})$  be the corrected partition function. Then

(114) 
$$\tilde{Z}^{\mathbb{P}^1}(\mathbf{m}) = e^q \cdot Z^{\mathbb{P}^1}(\mathbf{m})$$

As a corollary to Corollary 1.3, we get

**Corollary 5.4.** The corrected partition function  $\tilde{Z}^{\mathbb{P}^1}(\mathbf{m})$  is a KP taufunction. The corrected partition function  $\tilde{Z}^{\mathbb{P}^1}$  has the following expansion:

(115) 
$$\tilde{Z}^{\mathbb{P}^1}(\mathbf{m}) = \sum_{d \ge 0} q^d \sum_{n \ge 0} \sum_{i_1, \dots, i_n \ge 0} \frac{m_{i_1} \cdots m_{i_n}}{n!} \left\langle \prod_{j=1}^n \tau_{i_j}(2) \right\rangle_d^{\bullet \mathbb{P}^1},$$

where  $\left\langle \prod_{j=1}^{n} \tau_{i_j}(2) \right\rangle_d^{\bullet \mathbb{P}^1}$  denotes the *not-necessarily connected* GW invariants of  $\mathbb{P}^1$  of degree d in the stationary sector. By using the GW/H correspondence Okounkov and Pandharipande obtained [44] the following formula:

(116) 
$$\left\langle \prod_{j=1}^{n} \tau_{i_j}(2) \right\rangle_d^{\bullet \mathbb{P}^1} = \sum_{\lambda \in \mathcal{P}_d} \left( \frac{\dim V^{\lambda}}{d!} \right)^2 \prod_{j=1}^{n} \frac{\mathbf{p}_{i_j+1}(\lambda)}{(i_j+1)!},$$

where  $V^{\lambda}$  is the irreducible representation of  $S_d$  indexed by  $\lambda$ , and  $\mathbf{p}_k(\lambda)$  is defined by

(117) 
$$\mathbf{p}_k(\lambda) = \sum_{i\geq 1} \left[ \left(\lambda_i - i + \frac{1}{2}\right)^k - \left(-i + \frac{1}{2}\right)^k \right] + (1 - 2^{-k})\zeta(-k),$$

with  $\zeta(-k)$  given by

(118) 
$$\zeta(-k) = (-1)^k \frac{B_{k+1}}{k+1}.$$

Here  $B_m$  denotes the *m*th Bernoulli number. Dubrovin [20] gave a different proof of formula (116) using Symplectic Field Theory.

We now have

$$\tilde{Z}^{\mathbb{P}^{1}}(\mathbf{m}) = \sum_{d\geq 0} q^{d} \sum_{n\geq 0} \sum_{i_{1},\dots,i_{n}\geq 0} \frac{m_{i_{1}}\cdots m_{i_{n}}}{n!} \sum_{|\lambda|=d} \left(\frac{\dim V^{\lambda}}{d!}\right)^{2} \prod_{j=1}^{n} \frac{\mathbf{p}_{i_{j}+1}(\lambda)}{(i_{j}+1)!}$$
$$= \sum_{d\geq 0} \frac{q^{d}}{d!} \sum_{\lambda\in\mathcal{P}_{d}} \frac{(\dim V^{\lambda})^{2}}{d!} \exp\sum_{i\geq 0} \frac{m_{i}}{(i+1)!} \mathbf{p}_{i+1}(\lambda).$$

When all  $m_i$ 's are taken to be 0, we get the vacuum expectation value

(119) 
$$\tilde{Z}^{\mathbb{P}^1}(\mathbf{m}=\mathbf{0}) = \sum_{d\geq 0} \frac{q^d}{d!} \sum_{\lambda\in\mathcal{P}_d} \frac{(\dim V^\lambda)^2}{d!} = e^q.$$

By the definition (114) we have

$$Z^{\mathbb{P}^{1}}(\mathbf{m}) = e^{-q} \tilde{Z}^{\mathbb{P}^{1}}(\mathbf{m})$$
  
= 
$$\sum_{d \ge 0} \frac{q^{d}}{d!} e^{-q} \sum_{\lambda \in \mathcal{P}_{d}} \frac{(\dim V^{\lambda})^{2}}{d!} \exp \sum_{i \ge 0} \frac{m_{i}}{(i+1)!} \mathbf{p}_{i+1}(\lambda).$$

Now note that  $\{\{\frac{q^d}{d!}e^{-q}\frac{(\dim V^{\lambda})^2}{d!}\}_{\lambda\in\mathcal{P}_d}\}_{d\geq 0}$  is the Poissonized Plancherel measure on the set of partitions and  $\exp\sum_{i\geq 0}\frac{m_i}{(i+1)!}\mathbf{p}_{i+1}$  is a random variable on the set of partitions, so the partition function  $Z^{\mathbb{P}^1}(\mathbf{m})$  is just the expectation value:

(120) 
$$Z^{\mathbb{P}^1}(\mathbf{m}) = \left\langle \exp \sum_{i\geq 0}^{\infty} \frac{m_i}{(i+1)!} \mathbf{p}_{i+1}(\lambda) \right\rangle_{\text{Plancherel}}.$$

Now we recall that  $p_k = \frac{m_{k-1}}{(k-1)!}, k \ge 1$ , so we get

$$Z^{\mathbb{P}^1}(\mathbf{m}) = \sum_{d\geq 0} \frac{q^d}{d!} e^{-q} \sum_{\lambda\in\mathcal{P}_d} \frac{(\dim V^\lambda)^2}{d!} \exp\sum_{k\geq 1} \frac{p_k}{k} \mathbf{p}_k(\lambda).$$

Recall

(121) 
$$\exp\sum_{k\geq 1}\frac{1}{k}p_k(x)p_k(y) = \sum_{\mu}s_{\mu}(x)s_{\mu}(y),$$

where  $s_{\mu}$  is the Schur function which is a polynomial in **p**:

(122) 
$$s_{\mu}(\mathbf{p}) = \sum_{\nu} \frac{\chi^{\mu}_{\nu}}{z_{\nu}} p_{\nu}$$

(see Macdonald [38] for the details). It follows that

$$Z^{\mathbb{P}^{1}}(\mathbf{m}) = \sum_{d \ge 0} \frac{q^{d}}{d!} e^{-q} \sum_{|\lambda|=d} \frac{(\dim V^{\lambda})^{2}}{d!} \sum_{\mu} s_{\mu}(\mathbf{p}) \mathbf{s}_{\mu}(\lambda)$$
$$= \sum_{\mu} \langle \mathbf{s}_{\mu} \rangle_{\text{Plancherel}} \cdot s_{\mu}(\mathbf{p}).$$

Together with Corollary 1.3 we arrive at

**Theorem 5.5.** The affine coordinates  $A_{i,j}^{\mathbb{P}^1}$  of  $Z^{\mathbb{P}^1}(\mathbf{m})$  are given by the Plancherel expectation value of  $\mathbf{s}_{(i+1,1^j)}$ :

(123) 
$$A_{i,j}^{\mathbb{P}^1} = (-1)^j \langle \mathbf{s}_{(i+1,1^j)} \rangle_{\text{Plancherel}},$$

or explicitly,

(124) 
$$A_{i,j}^{\mathbb{P}^1} = (-1)^j \sum_{d \ge 0} \frac{q^d}{d!} e^{-q} \sum_{\lambda \in \mathcal{P}_d} \frac{(\dim V^\lambda)^2}{d!} \mathbf{s}_{(i+1,1^j)}(\lambda).$$

Combining Theorem 5.2 and Theorem 5.5 we arrive at

Corollary 5.6. The following identity holds:

(125) 
$$\frac{1}{\xi - \eta} + \sum_{i,j \ge 0} (-1)^j \sum_{d \ge 0} \frac{q^d}{d!} e^{-q} \sum_{\lambda \in \mathcal{P}_d} \frac{(\dim V^\lambda)^2}{d!} \frac{\mathbf{s}_{(i+1,1^j)}(\lambda)}{\xi^{j+1} \eta^{i+1}}$$
$$= -B(\xi, \eta) \frac{e^{\log \frac{\Gamma(\xi + \frac{1}{2})}{\sqrt{2\pi}} - \xi \log \xi + \xi}}{e^{\log \frac{\Gamma(\eta + \frac{1}{2})}{\sqrt{2\pi}} - \eta \log \eta + \eta}},$$

where  $\Gamma(\xi + \frac{1}{2})$  is understood as its asymptotic expansion for  $\xi$  large with  $\arg \xi < \pi - \epsilon$ , and  $B(\xi, \eta)$  is given by (107).

For example,

$$A_{0,0}^{\mathbb{P}^{1}} = \sum_{d \ge 0} \frac{q^{d}}{d!} e^{-q} \sum_{\lambda \in \mathcal{P}_{d}} \frac{(\dim V^{\lambda})^{2}}{d!} \mathbf{s}_{(1)}(\lambda)$$
  
$$= e^{-q} \left[ -\frac{1}{24} + \frac{23}{24}q + \left( \left( \frac{1}{2!} \right)^{2} \frac{47}{24} + \left( \frac{1}{2!} \right)^{2} \frac{47}{24} \right) q^{2} + \left( \left( \left( \frac{1}{3!} \right)^{2} \frac{71}{24} + \left( \frac{2}{3!} \right)^{2} \frac{71}{24} + \left( \frac{1}{3!} \right)^{2} \frac{71}{24} \right) q^{3} + \cdots \right]$$
  
$$= -\frac{1}{24} + q,$$

$$\begin{split} A_{1,0}^{\mathbb{P}^{1}} &= \sum_{d \ge 0} \frac{q^{d}}{d!} e^{-q} \sum_{\lambda \in \mathcal{P}_{d}} \frac{(\dim V^{\lambda})^{2}}{d!} \mathbf{s}_{(2)}(\lambda) \\ &= \sum_{d \ge 0} \frac{q^{d}}{d!} e^{-q} \sum_{\lambda \in \mathcal{P}_{d}} \frac{(\dim V^{\lambda})^{2}}{d!} \left(\frac{1}{2} \mathbf{p}_{(2)}(\lambda) + \frac{1}{2} \mathbf{p}_{1}(\lambda)^{2}\right) \\ &= \frac{1}{2} e^{-q} \left[ 0 + 0 \cdot q + \left( \left(\frac{1}{2!}\right)^{2} \cdot 2 + \left(\frac{1}{2!}\right)^{2} \cdot (-2) \right) q^{2} \\ &+ \left( \left(\frac{1}{3!}\right)^{2} \cdot 6 + \left(\frac{2}{3!}\right)^{2} \cdot 0 + \left(\frac{1}{3!}\right)^{2} \cdot (-6) \right) q^{3} + \cdots \right. \\ &+ \left( -\frac{1}{24} \right)^{2} + \left(\frac{23}{24} \right)^{2} q + \left( \left(\frac{1}{2!}\right)^{2} \left(\frac{47}{24}\right)^{2} + \left(\frac{1}{2!}\right)^{2} \left(\frac{47}{24}\right)^{2} \right) q^{2} \\ &+ \left( \left(\frac{1}{3!}\right)^{2} \left(\frac{71}{24}\right)^{2} + \left(\frac{2}{3!}\right)^{2} \left(\frac{71}{24}\right)^{2} + \left(\frac{1}{3!}\right)^{2} \left(\frac{71}{24}\right)^{2} \right) q^{3} + \cdots \right] \\ &= \frac{1}{1152} + \frac{11q}{24} + \frac{q^{2}}{2}. \end{split}$$

These match with the right-hand side of (125).

By a result due to Stanley [48],

(126) 
$$\langle \mathbf{s}_{\mu} \rangle_{d} := \sum_{\lambda \in \mathcal{P}_{d}} \frac{(\dim V^{\lambda})^{2}}{d!} \sum_{\mu} \mathbf{s}_{\mu}(\lambda)$$

is a polynomial in d of degree  $|\mu|$ . For example,

$$\langle \mathbf{p}_{1} \rangle_{d} = d - \frac{1}{24},$$

$$\langle \mathbf{p}_{2} \rangle_{d} = 0,$$

$$\langle \mathbf{p}_{1}^{2} \rangle_{d} = (d - \frac{1}{24})^{2},$$

$$\langle \mathbf{p}_{3} \rangle_{d} = \frac{3d^{2}}{2} - \frac{5d}{4} + \frac{7}{960},$$

$$\langle \mathbf{p}_{2} \mathbf{p}_{1} \rangle_{d} = 0,$$

$$\langle \mathbf{p}_{1}^{3} \rangle_{d} = d^{3} - \frac{1}{8}d^{2} + \frac{1}{192}d - \frac{1}{13824},$$

and from these we get:

$$\langle \mathbf{s}_{(1)} \rangle_d = d - \frac{1}{24},$$

$$\langle \mathbf{s}_{(2)} \rangle_d = \langle \mathbf{s}_{(1^2)} \rangle_d = (d - \frac{1}{24})^2,$$

$$\langle \mathbf{s}_{(3)} \rangle_d = \langle \mathbf{s}_{(1^3)} \rangle = \frac{d^3}{6} + \frac{23}{48}d^2 - \frac{479}{1152}d + \frac{1003}{414720},$$

$$\langle \mathbf{s}_{(21)} \rangle_d = \frac{d^3}{3} - \frac{13}{24}d^2 + \frac{241}{576}d - \frac{509}{207360}.$$

Note that by (110) we have

(127) 
$$Z^{\mathbb{P}^{1}}(\mathbf{m}) = Z^{+,\mathbb{P}^{1}}(\mathbf{m}) \cdot \exp \sum_{k \ge 1} (1 - 2^{-k})\zeta(-k) \frac{m_{k-1}}{(k-1)!},$$

so we can also define:

(128) 
$$\hat{\mathbf{p}}_k(\lambda) = \sum_{i \ge 1} \left[ \left( \lambda_i - i + \frac{1}{2} \right)^k - \left( -i + \frac{1}{2} \right)^k \right].$$

Then we repeat everything we have done in this subsection by replacing every  $\mathbf{p}_k$  with  $\hat{\mathbf{p}}_k$  to get:

$$Z^{\mathbb{P}^{1,+}}(\mathbf{m}) = \sum_{d \ge 0} \frac{q^d}{d!} e^{-q} \sum_{|\lambda|=d} \frac{(\dim V^{\lambda})^2}{d!} \exp \sum_{i \ge 0} \frac{m_i}{(i+1)!} \hat{\mathbf{p}}_{i+1}(\lambda),$$

and

(129) 
$$Z^{\mathbb{P}^{1,+}}(\mathbf{m}) = \left\langle \exp \sum_{i \ge 0} \frac{m_i}{(i+1)!} \hat{\mathbf{p}}_{i+1}(\lambda) \right\rangle_{\text{Plancherel}}.$$

And if we set

(130) 
$$\hat{\mathbf{s}}_{\mu} = \sum_{\nu} \frac{\chi^{\mu}_{\nu}}{z_{\nu}} \hat{\mathbf{p}}_{\nu},$$

then we have

$$Z^{\mathbb{P}^{1,+}}(\mathbf{m}) = \sum_{d\geq 0} \frac{q^{d}}{d!} e^{-q} \sum_{|\lambda|=d} \frac{(\dim V^{\lambda})^{2}}{d!} \sum_{\mu} s_{\mu}(\mathbf{p}) \hat{\mathbf{s}}_{\mu}(\lambda)$$
$$= \sum_{\mu} \langle \hat{\mathbf{s}}_{\mu} \rangle_{\text{Plancherel}} \cdot s_{\mu}(\mathbf{p}).$$

**Theorem 5.7.** The affine coordinates  $A_{i,j}^{\mathbb{P}^{1,+}}$  of  $Z^{+,\mathbb{P}^{1}}(\mathbf{m})$  are given by the Plancherel expectation value of  $\hat{\mathbf{s}}_{(i+1,1^{j})}$ :

(131) 
$$A_{i,j}^{\mathbb{P}^{1,+}} = (-1)^n \langle \hat{\mathbf{s}}_{(i+1,1^j)} \rangle_{\text{Plancherel}},$$

or explicitly,

(132) 
$$A_{i,j}^{\mathbb{P}^{1,+}} = (-1)^{j} \sum_{d \ge 0} \frac{q^{d}}{d!} e^{-q} \sum_{|\lambda|=d} \frac{(\dim V^{\lambda})^{2}}{d!} \hat{\mathbf{s}}_{(i+1,1^{j})}(\lambda).$$

Combining (108) and Theorem 5.7 we arrive at

Corollary 5.8. The following identity holds:

(133) 
$$\sum_{i,j} (-1)^j \sum_{d \ge 0} \frac{q^d}{d!} e^{-q} \sum_{|\lambda|=d} \frac{(\dim V^{\lambda})^2}{d!} \frac{\hat{\mathbf{s}}_{(i+1,1^j)}(\lambda)}{\xi^{j+1} \eta^{i+1}} = B(\xi,\eta),$$

where  $B(\xi, \eta)$  is given by (107).

For example,

$$\begin{aligned} A_{0,0}^{\mathbb{P}^{1,+}} &= \sum_{d \ge 0} \frac{q^{d}}{d!} e^{-q} \sum_{|\lambda|=d} \frac{(\dim V^{\lambda})^{2}}{d!} \hat{\mathbf{s}}_{(1)}(\lambda) \\ &= e^{-q} \left[ 0 + 1 \cdot q + \left( \left(\frac{1}{2!}\right)^{2} \cdot 2 + \left(\frac{1}{2!}\right)^{2} \cdot 2 \right) q^{2} \\ &+ \left( \left(\left(\frac{1}{3!}\right)^{2} \cdot 3 + \left(\frac{2}{3!}\right)^{2} \cdot 3 + \left(\frac{1}{3!}\right)^{2} \cdot 3 \right) q^{3} + \cdots \right] \\ &= q, \end{aligned}$$

$$\begin{split} A_{1,0}^{\mathbb{P}^{1,+}} &= \sum_{d \ge 0} \frac{q^{d}}{d!} e^{-q} \sum_{|\lambda|=d} \frac{(\dim V^{\lambda})^{2}}{d!} \hat{\mathbf{s}}_{(2)}(\lambda) \\ &= \sum_{d \ge 0} \frac{q^{d}}{d!} e^{-q} \sum_{|\lambda|=d} \frac{(\dim V^{\lambda})^{2}}{d!} \left(\frac{1}{2} \hat{\mathbf{p}}_{(2)}(\lambda) + \frac{1}{2} \hat{\mathbf{p}}_{(1)}(\lambda)^{2}\right) \\ &= \frac{1}{2} e^{-q} \left[ 0 + 0 \cdot q + \left(\left(\frac{1}{2!}\right)^{2} \cdot 2 + \left(\frac{1}{2!}\right)^{2} \cdot (-2)\right) q^{2} \right. \\ &+ \left. \left(\left(\frac{1}{3!}\right)^{2} \cdot 6 + \left(\frac{2}{3!}\right)^{2} \cdot 0 + \left(\frac{1}{3!}\right)^{2} \cdot (-6)\right) q^{3} + \cdots \right. \\ &+ \left. 0^{2} q + 1^{2} q + \left(\left(\frac{1}{2!}\right)^{2} \cdot 2^{2} + \left(\frac{1}{2!}\right)^{2} \cdot 2^{2}\right) q^{2} \right. \\ &+ \left. \left(\left(\frac{1}{3!}\right)^{2} \cdot 3^{2} + \left(\frac{2}{3!}\right)^{2} \cdot 3^{2} + \left(\frac{1}{3!}\right)^{2} \cdot 3^{2}\right) q^{3} + \cdots \right] \\ &= \frac{q}{2} + \frac{q^{2}}{2}. \end{split}$$

These match with the right-hand side of (133). The Plancherel averages of  $\hat{\mathbf{p}}_{\mu}$  and  $\hat{\mathbf{s}}_{\mu}$  are again polynomials, but they are simpler that the averages of  $\mathbf{p}_{\mu}$  and  $\mathbf{s}_{\mu}$ . For example,

$$\langle \hat{\mathbf{p}}_1 \rangle_d = d, \qquad \langle \hat{\mathbf{p}}_2 \rangle_d = 0, \qquad \langle \hat{\mathbf{p}}_1^2 \rangle = d^2, \\ \langle \hat{\mathbf{p}}_3 \rangle_d = \frac{3}{2} d^2 - \frac{5d}{4}, \qquad \langle \hat{\mathbf{p}}_2 \hat{\mathbf{p}}_1 \rangle_d = 0, \qquad \langle \hat{\mathbf{p}}_1^3 \rangle_d = d^3,$$

and from these we get:

$$\begin{aligned} \langle \hat{\mathbf{s}}_{(1)} \rangle_d &= d, \\ \langle \hat{\mathbf{s}}_{(2)} \rangle_d &= \langle \hat{\mathbf{s}}_{(1^2)} \rangle = \frac{d^2}{2} = \frac{(d)_2}{2} + \frac{d}{2}, \\ \langle \hat{\mathbf{s}}_{(3)} \rangle_d &= \langle \hat{\mathbf{s}}_{(1^3)} \rangle = \frac{d^3}{6} + \frac{1}{2}d^2 - \frac{5}{12}d, \\ \langle \hat{\mathbf{s}}_{(21)} \rangle_d &= \frac{d^3}{3} - \frac{1}{2}d^2 + \frac{5}{12}d. \end{aligned}$$

## References

- Adler, M., van Moerbeke, P.: Matrix integrals, Toda symmetries, Virasoro constraints and orthogonal polynomials. Duke Math. J. 80 (1995), 863–911.
- [2] Alexandrov, A.: Matrix model for the stationary sector of Gromov-Witten theory of P<sup>1</sup>. Comm. Math. Phys. 382 (2021), 1935–1960.

#### DI YANG, JIAN ZHOU

- [3] Alexandrov, A., Bychkov, B., Dunin-Barkowski, P., Kazarian, M., Shadrin, S.: KP integrability through the x-y swap relation. arXiv:2309.12176.
- [4] Ambjørn, J., Chekhov, L.: The matrix model for dessins d'enfants. Ann. Inst. Henri Poincaré D 1 (2014), 337–361.
- [5] Balogh, F., Yang, D.: Geometric interpretation of Zhou's explicit formula for the Witten–Kontsevich tau function. Lett. Math. Phys. 107 (2017), 1837–1857.
- [6] Behrend, K., Fantechi, B.: The intrinsic normal cone. Invent. Math. 128 (1997), 45–88.
- [7] Bertola, M., Dubrovin, B., Yang, D.: Correlation functions of the KdV hierarchy and applications to intersection numbers over M
  <sub>g,n</sub>. Phys. D **327** (2016), 30–57.
- [8] Bertola M., Dubrovin B., Yang D.: Simple Lie algebras, Drinfeld–Sokolov hierarchies, and multi-point correlation functions. Mosc. Math. J. 21 (2021), 233–270.
- [9] Bertola, M., Ruzza, G.: Matrix models for stationary Gromov-Witten invariants of the Riemann sphere. Nonlinearity **34** (2021), 1168–1196.
- [10] Carlet, G.: Extended Toda hierarchy and its Hamiltonian structure. PhD Thesis. 2003.
- [11] Carlet, G., Dubrovin, B., Zhang, Y.: The extended Toda hierarchy. Mosc. Math. J. 4 (2004), 313–332.
- [12] Carlet, G., van de Leur, J.: Hirota equations for the extended bigraded Toda hierarchy and the total descendent potential of CP<sup>1</sup> orbifolds. J. Phys. A 46 (2013), 405205, 16 pp.
- [13] Carlet, G., van de Leur, J.; Posthuma, H.; Shadrin, S.: Higher genera Catalan numbers and Hirota equations for extended nonlinear Schrödinger hierarchy Lett. Math. Phys. **111** (2021), Paper No. 63, 67 pp.
- [14] Deift, P.: Orthogonal polynomials and random matrices: a Riemann-Hilbert approach. Courant Lecture Notes in Mathematics 3. American Mathematical Society, Providence, R.I., 1999.
- [15] Dickey, L. A.: Soliton Equations and Hamiltonian Systems, 2nd edn. World Scientific, 2003.
- [16] Dijkgraaf, R., Witten, E.: Mean field theory, topological field theory, and multi-matrix models. Nuclear Phys. B 342 (1990), 486–522.
- [17] Dubrovin, B.: Integrable systems and classification of 2-dimensional topological field theories. In "Integrable systems" (Luminy, 1991), Progr. Math., vol. 115, pp. 313–359., Birkhäuser Boston, Boston, MA, 1993.
- [18] Dubrovin, B.: Geometry of 2D topological field theories. In: Francaviglia, M., Greco, S. (eds.) "Integrable Systems and Quantum Groups" (Montecatini Terme, 1993), Lecture Notes in Math., vol. 1620, pp. 120–348. Springer, Berlin, 1996.
- [19] Dubrovin, B.: Hamiltonian perturbations of hyperbolic PDEs: from classification results to the properties of solutions. In: Sidoravičius, V. (ed.) "New Trends in Mathematical Physics", pp. 231–276, Springer, Dordrecht, 2009.
- [20] Dubrovin, B.: Symplectic field theory of a disk, quantum integrable systems, and Schur polynomials. Ann. Henri Poincaré 17 (2016), 1595–1613.
- [21] Dubrovin, B., Yang, D.: Generating series for GUE correlators. Lett. Math. Phys. 107 (2017), 1971–2012.

- [22] Dubrovin, B., Yang, D.: On Gromov-Witten invariants of P<sup>1</sup>. Math. Res. Lett. 26 (2019), 729–748.
- [23] Dubrovin, B., Yang, D., Zagier, D.: Gromov-Witten invariants of the Riemann sphere. Pure Appl. Math. Q. 16 (2020), 153–190.
- [24] Dubrovin, B., Yang, D., Zagier, D.: On tau-functions for the KdV hierarchy. Selecta Mathematica 27 (2021), paper No. 12.
- [25] Dubrovin, B., Zhang, Y.: Normal forms of hierarchies of integrable PDEs, Frobenius Manifolds and Gromov–Witten invariants. arXiv:math/0108160.
- [26] Dubrovin, B., Zhang, Y.: Virasoro symmetries of the extended Toda hierarchy. Comm. Math. Phys. 250 (2004), 161–193.
- [27] Eguchi, T., Yang, S.-K.: The topological CP<sup>1</sup> model and the large-N matrix integral. Modern Physics Letters A 9 (1994), 2893–2902.
- [28] Eguchi, T., Hori, K., Yang, S.-K.: Topological σ-models and large-N matrix integral Internat. J. Modern Phys. A 10 (1995), 4203–4224.
- [29] Ercolani, N.M., McLaughlin, K.D.T.-R., Pierce, V.U.: Random matrices, graphical enumeration and the continuum limit of the Toda lattices. Comm. Math. Phys. 278 (2008), 31–81.
- [30] Flaschka, H.: On the Toda lattice. II. Inverse-scattering solution. Progr. Theoret. Phys. 51 (1974), 703–716.
- [31] Gerasimov, A., Marshakov, A., Mironov, A., Morozov, A., Orlov, A.: Matrix models of two-dimensional gravity and Toda theory. Nucl. Phys. B 357 (1991), 565–618.
- [32] Getzler, E.: The Toda conjecture. In: "Symplectic Geometry and Mirror Symmetry" (Seoul, 2000), World Sci., River Edge, N.J., 51–79, 2001.
- [33] Gisonni, M., Grava, T., Ruzza, G.: Laguerre ensemble: correlators, Hurwitz numbers and Hodge integrals. Ann. Henri Poincaré 21 (2020), 3285–3339.
- [34] Harer, J., Zagier, D.: The Euler characteristic of the moduli space of curves. Invent. Math. 85 (1986), 457–485.
- [35] Kazakov, V., Kostov, I., Nekrasov, N.: D-particles, matrix integrals and KP hierarchy. Nucl. Phys. B 557 (1999), 413–442.
- [36] Kazarian, M., Zograf, P.: Virasoro constraints and topological recursion for Grothendieck's dessin counting. Lett. Math. Phys. 105 (2015), 1057–1084.
- [37] Li, J., Tian, G.: Virtual moduli cycles and Gromov-Witten invariants of algebraic varieties. J. Amer. Math. Soc. 11 (1998), 119–174.
- [38] Macdonald, I.G.: Symmetric Functions and Hall Polynomials, 2nd edn. Oxford Mathematical Monographs. Oxford University Press Inc., New York, 1995.
- [39] Manakov, S. V.: Complete integrability and stochastization of discrete dynamical systems. J. Experiment. Theoret. Phys. 67 (1974), 543–555 (in Russian). English translation in: Soviet Physics JETP 40 (1974), 269–274.
- [40] Marchal, O.: WKB solutions of difference equations and reconstruction by the topological recursion. Nonlinearity **31** (2018), 226–262.
- [41] Mehta, M.L.: Random Matrices, 2nd edn. Academic Press, New York, 1991.
- [42] Milanov, T.: Hirota quadratic equations for the extended Toda hierarchy. Duke Math. J. 138 (2007), 161–178.
- [43] Morozov, A.: Integrability and matrix models. Physics-Uspekhi 37 (1994), 1–55.
- [44] Okounkov, A., Pandharipande, R.: Gromov–Witten theory, Hurwitz theory, and completed cycles. Annals of Mathematics **163** (2006), 517–560.

#### DI YANG, JIAN ZHOU

- [45] Okounkov, A., Pandharipande, R.: The equivariant Gromov-Witten theory of P<sup>1</sup>. Annals of Mathematics 163 (2006), 561–605.
- [46] Ruan, Y., Tian, G.: A mathematical theory of quantum cohomology. J. Diff. Geom. 42 (1995), 259–367.
- [47] Shaw, J. C., Tu, M. H., Yen, H. C.: Matrix models at finite N and the KP hierarchy, Chinese Journ. Phys. Vol. 30, No. 4, 497–507, 1992.
- [48] Stanley, R.P.: Some combinatorial properties of hook lengths, contents, and parts of partitions, Ramanujan J (2010) 23: 91–105.
- [49] Ueno, K., Takasaki, K.: Toda lattice hierarchy. Adv. Stud. Pure Math. 4, pp. 1–95, North-Holland Publishing Co., Amsterdam, 1984.
- [50] Yang, D.: On tau-functions for the Toda lattice hierarchy. Lett. Math. Phys. 110 (2020), 555–583.
- [51] Yang, D.: GUE via Frobenius manifolds. I. From matrix gravity to topological gravity and back. to appear in Acta Mathematical Sinica (English Series). arXiv:2205.01618.
- [52] Yang, D., Zhou, J.: Grothendieck's dessins d'enfants in a web of dualities. III. J. Phys. A 56 (2023), Paper No. 055201, 34 pp.
- [53] Zhang, Y.: On the CP<sup>1</sup> topological sigma model and the Toda lattice hierarchy. J. Geom. Phys. 40 (2002), 215–232.
- [54] Zhou, J.: Explicit Formula for Witten-Kontsevich Tau-Function. arXiv:1306.5429.
- [55] Zhou, J.: Emergent geometry and mirror symmetry of a point. arXiv:1507.01679.
- [56] Zhou, J.: Hermitian one-matrix model and KP hierarchy. arXiv:1809.07951.
- [57] Zhou, J.: Grothendieck's dessins d'enfants in a web of dualities. arXiv:1905.10773.
- [58] Zhou, J.: Grothendieck's dessins d'enfants in a web of dualities. II. arXiv:1907.00357.
- [59] Zograf, P.: Enumeration of Grothendieck's dessins and KP hierarchy, IMRN 2015, 13533–13544.

Di Yang, School of Mathematical Sciences, University of Science and Technology of China Hefei 230026, P.R. China e-mail: diyang@ustc.edu.cn

Jian Zhou, Department of Mathematical Sciences, Tsinghua University Beijing 100084, P.R. China e-mail: jianzhou@mail.tsinghua.edu.cn