Finding the Fairest Voting System Using Likelihood Analysis

Virochan Pandit vpandit3@jhu.edu Johns Hopkins University

Joseph Cutrone jcutron2@jhu.edu Johns Hopkins University

This research was supported by Johns Hopkins' Provost's Undergraduate Research Award (PURA)

We would like to acknowledge Christopher Ratigan and Sergey Kushnarev for their insights and suggestions on this paper. We also thank Alexander Shumakovitch for guidance on operating the Rockfish computer cluster. Further, we thank the Hopkins Office for Undergraduate Research (HOUR) for their support and guidance.

Abstract

This paper takes a statistical approach to determine which of 10 voting systems, or preference aggregation rules, is the fairest based on their probabilistic likelihood of violating Arrow's five social choice criteria. The voting systems considered are: Plurality, Borda, Dowdall, Top Two (Plurality Runoff), Instant Runoff, Coombs, Baldwin, Copeland, Pairwise Majority, and Minimax. This paper builds upon the work of Dougherty and Heckelman (2020) by computing violation frequencies for elections with a greater number of alternatives. Elections with up to 50,000 voters and between three and six alternatives are simulated using both Impartial Culture and Impartial Anonymous Culture. The results of these simulations produce new IIA violation likelihoods for each method and show that Pairwise Majority is the most likely to jointly satisfy all five of Arrow's criteria. Furthermore, of the systems that satisfy transitivity, the Baldwin method is most likely to jointly satisfy all five of Arrow's criteria in elections with three alternatives. As the number of alternatives increase, the joint satisfaction frequencies decrease rapidly for all systems.

1 Introduction

Voting systems can be evaluated based on whether they satisfy or violate social choice criteria. This paper takes a statistical approach to determine which voting system is the fairest based on the probabilistic likelihood of violating Arrow's five social choice criteria.

In total, 10 voting systems are analyzed with respect to five social choice criteria. Specifically, the methods considered are Plurality, Borda, Dowdall, Top Two (Plurality Runoff), Instant Runoff, Coombs, Baldwin, Copeland, Pairwise Majority, and Minimax. Through simulations with up to 50,000 voters and six alternatives, the violation frequencies are computed for each method. To model voters' preferences, two cultures are used: Impartial Culture (IC) and Impartial Anonymous Culture (IAC). First, voting systems are tested with respect to each individual criteria. Then, for each voting system, Joint-Satisfaction (JS) frequencies, the probability that systems jointly satisfy all five conditions, are computed. These frequencies are used to determine the fairest system.

This paper builds upon the work of previous empirical studies in social choice theory. Compared to prior works, this study computes violation frequencies for elections with a greater number of voters and alternatives than most literature. For the case of three-alternatives elections, the majority of our results align with Dougherty and Heckelman (2020). This paper both extends their analysis for up to six alternatives and finds higher lower bounds for the frequency of IIA violations. The main findings are summarized below:

Result 1: Of the 10 voting systems analyzed, Pairwise Majority is the most likely to jointly satisfy Arrow's five social choice criteria. In elections with three alternatives and at least 100 voters, Pairwise Majority has Joint-Satisfaction frequencies of over 85%.

Result 2: Of the systems that satisfy transitivity, Baldwin's Method has the highest Joint-Satisfaction frequencies. The frequencies, however, are less than 20%.

Result 3: As the number of alternatives increases, the Joint-Satisfaction frequencies decrease for each voting method. For all systems except Pairwise Majority, the Joint-Satisfaction frequencies in elections with at least 30 voters and four alternatives are 0%.

The paper proceeds as follows: Sections 2 and 3 detail all required background knowledge. Section 4 contains this study's contributions and new results. Section 5 describes the experiment set-up and methodology. Section 6 presents the results, and Section 7 discusses the differences in results to past papers. Section 8 concludes the paper and highlights the key points.

2 Voting Systems

We follow the same assumptions as in Dougherty and Heckelman (2020). We define below the voting systems analyzed in this paper which are a subset of those considered in their paper. In Dougherty and Heckelman (2020), Plurality, Minimax, Top Two, and Instant Runoff are said to be part of **Group 1** due to their similar numerical results, and we adopt this convention here.

In terms of preference notation, we follow (Mihara, 1997). For two alternatives A and B, $A \succ B$ indicates society prefers A over B, $A \succeq B$ indicates society prefers A at least as much as B, and $A \sim B$ indicates society is indifferent between A and B. The analogous symbols with a subscript of i (for example $A \succ_i B$) indicate individual preferences.

Plurality - Alternatives are ranked based on their first-place vote tally.

- **Borda Count** Let the election have n alternatives. For each 1st place vote, alternatives receive n points, for each 2nd place vote they receive n 1 points, and so on until for an nth place vote, alternatives receive one point. To obtain a societal preference order, the alternatives are ranked based on their point total.
- **Dowdall** Let the election have *n* alternatives. For each first place vote, alternatives receive 1 point, for a second place vote they receive $\frac{1}{2}$ point, for a third place vote they receive $\frac{1}{3}$ point and so on until for an *n*th place vote, alternatives receive $\frac{1}{n}$ points. The voters are then ranked based on their point tally.

- **Top Two/Plurality Runoff** There are two rounds. The first round is a plurality election. If a alternative receives a majority, they win the election. If not, the voters vote again, but can only chose between the two alternatives who receive the most first-place votes in the first round. The alternative with the most first-place votes in the second round wins the election.¹
- **Instant Runoff/Hare -** In each round, the alternative(s) with the least first place votes is eliminated. The votes of the eliminated alternative(s) are transferred to the alternatives ranked below them. The process repeats until all alternatives are eliminated. Alternatives are ranked based on reverse order of the round they were eliminated.
- **Coombs** This follows the same procedure as Instant Runoff, but has a different elimination rule. Each round, the alternative with the most last place votes is eliminated.²
- **Baldwin** This system follows the same procedure as Instant Runoff, but has a different elimination rule. Each round, the alternative with the least points, as calculated by the Borda Count rule, is eliminated.
- **Pairwise Majority** All alternatives are compared head to head pairwise. For two alternatives A and B, if A wins the head to head then $A \succ B$. If B wins the head to head then $B \succ A$. If they tie then $A \sim B$. Based on the comparisons, a transitive ordering is formed if one exists.
- **Copeland/Pairwise Comparison** All alternatives are compared head to head pairwise. For each comparison, the alternative that wins head to head receives 1 point. If the two alternatives tie they receive $\frac{1}{2}$ point each. After all comparisons are performed, alternatives are ranked based on their point total.
- Minimax/Simpson-Kramer All alternatives are compared head to head. For each alternative, their score is the magnitude of their worst pairwise defeat. Alternatives are ranked in the reverse order of their score (with lower scores giving higher rank).

2.1 Arrow's Criteria of Fairness

Following (Dougherty & Heckelman, 2020; Morreau, 2019; Penn, 2015), below is a list of Arrow's voting criteria used in this paper.

- 1. Non-Dictatorship (ND) There is no individual whose preferences uniquely determine the social raking of all alternatives, regardless of how other individuals rank the alternatives.
- 2. Unrestricted Domain (UD) The domain of the voting system can take any preference schedule as input. This means that for every preference schedule, the voting system produces societal preferences between any two alternatives; there are no restrictions on how voters can vote. Note that, for the scope of this paper, this condition does not imply the rankings have to be transitive.
- 3. Transitivity (TR) The voting method always produces transitive preferences, i.e., the method always yields a societal preference order. Formally, for any three candidates A, B, and $C, A \succeq B$ and $B \succeq C$ implies $A \succeq C$.
- 4. (Pareto) Unanimity (U) If all voters prefer alternative A over alternative B, then in the societal preference order, $A \succ B$.
- 5. Independence of Irrelevant Alternatives (IIA) Society's preference between any two alternatives only depends on voter's preferences between these two alternatives (Börgers, 2010; Hodge & Kilma, 2018). Equivalently, if there are two preference schedules in which each voter has the same relative preference between alternatives A and B, the societal preference between A and B must be the same in both preference schedules.

¹If there is a majority winner in the first round, alternatives are ranked based on their first-place vote tally. If there is no majority winner, the top two alternatives are compared and ranked first and second (or tied first) depending on their relative vote totals in the second round. Then the other alternatives are ranked below them based on their first round vote totals.

²In some descriptions of Coombs' method (Nurmi, 1983), the election runs only until a alternative receives a majority. This approach, however, does not produce a clear ranking of alternatives.

3 Arrow's Theorem

Arrow's theorem shows that no voting system can satisfy all five criteria with at least two voters and at least three alternatives (Arrow, 1963). Table 1 below summarizes which criteria each voting system satisfies. A check-mark means that the voting system always satisfies the criteria while a x-mark indicates that it is possible for that voting system to violate that criteria. The theoretical properties of voting systems are well studied in literature (Hodge & Kilma, 2018; Niou, 1987; Nurmi, 1983; Patitsas, 2008).

Voting System	ND	UD	TR	U	IIA
Plurality	\checkmark	\checkmark	\checkmark	Х	X
Borda Count	\checkmark	\checkmark	\checkmark	\checkmark	×
Instant Runoff	\checkmark	\checkmark	\checkmark	×	×
Coombs Method	\checkmark	\checkmark	\checkmark	\checkmark	×
Copeland	\checkmark	\checkmark	\checkmark	\checkmark	×
Top Two	\checkmark	\checkmark	\checkmark	×	×
Dowdall	\checkmark	\checkmark	\checkmark	\checkmark	×
Pairwise Majority	\checkmark	\checkmark	Х	\checkmark	\checkmark
Minimax	\checkmark	\checkmark	\checkmark	×	×
Baldwin	\checkmark	\checkmark	\checkmark	\checkmark	×

Table 1: Criteria satisfied by 10 Voting Systems

Every voting system considered satisfies ND and UD since none are dictatorial and none restrict the way voters can vote. Note that Arrow did not originally include Transitivity as a separate criterion, but he did argue for transitive preferences (Morreau, 2019).

4 New and Expanded Results

The numerical probability of violating each of Arrow's five conditions is investigated in Dougherty and Heckelman (2020). The study also calculates the likelihood that voting systems violate at least one of Arrow's conditions. The new contributions of this paper that extend their prior results are:

- 1. This paper collects more empirical data, analyzing elections for up to 50,000 voters and six alternatives.
- 2. This paper confirms and collates the results of previous studies, providing extensive data of the frequencies at which voting systems violate criteria.
- 3. This paper develops a new methodology to test for IIA violations, which finds substantially more violations than previous algorithms, and thus creates a new greatest lower bound for IIA violations.
- 4. A new metric, Joint-Satisfaction frequencies, is introduced and formalized to test for the fairness of voting systems. This is similar to Dougherty and Heckelman (2020) regarding the likelihood of violating at least one criteria.

The data tables with the results showing percentages of violations for each criterion and Joint-Satisfaction frequencies are located in Appendix A.

5 Simulations

All computational models were implemented using C++ and Python. The voting systems (Plurality, Borda, Instant Runoff, Coombs, Copeland, Dowdall, Pairwise Majority, Minimax, Top Two, and Baldwin) were tested against Arrow's criterion (ND, UD, IIA, U, TR) for three to six alternatives.

Ties were handled as follows: For Top Two, there is a possibility of a tie in the first round, where more than two alternatives have equal number of votes. In this case, ties are broken randomly. In all other methods, if alternatives are tied, they have the same rank in the societal preference order. Two preference distributions, Impartial Culture and Impartial Anonymous Culture (Eğecioğlu & Giritligil, 2013) are considered. Impartial Culture (IC) assumes each voter is equally likely to select any of the possible preference orders. Impartial Anonymous Culture (IAC) assumes each preference schedule is equally likely to occur. All pseudo-random number generation was done using the Mersenne Twister engine (Matsumoto & Nishimura, 1998) in C++ random library.

For each setting (fixing the criterion, number of voters, number of alternatives, and culture) 10,000 preference schedules were generated. Then the violation frequency, the percentage of times a voting system violates the criterion, was calculated for each of the 10 voting systems. For JS Frequencies, this involved testing voting systems against multiple criteria consecutively. Due to the simulation process, the frequency of violations calculated are numerical probabilities, not theoretical.

All computations were carried out at the Advanced Research Computing at Hopkins (ARCH) core facility (rockfish.jhu.edu), which is supported by the National Science Foundation (NSF) grant number OAC1920103.

6 Results and Analysis

The following sections details the likelihood of voting systems violating Arrow's social choice criterion. Note that all voting methods considered satisfy Non-Dictatorship and Unrestricted Domain by definition and as such are excluded from the data.

6.1 Independence of Irrelevant Alternatives (IIA)

Key Observations in IIA Data

- 1. In elections with at least three alternatives and two voters, Group 1 systems (Plurality, Instant Runoff, Minimax, and Top Two) violate IIA more than 99.99% of the time.
- 2. As the number of voters increases, IIA violations generally increase under IC. For all systems that violate IIA, the frequency of violations approaches 100% as the number of voters increase. For elections with at least 100 voters and three alternatives, IIA is almost never satisfied. However, this trend is not always observed as many voting systems violate IIA at a lower frequency in elections with four voters than elections with three voters. The IAC data does not strictly abide by this trend.
- 3. As the number of alternatives increases, IIA violations increase. In elections with 4 alternatives and at least 100 voters, all systems violate IIA more than 99.7% of the time for both IC and IAC. For elections with at least five alternatives, IIA is almost never satisfied.

Tables 2 and 3 show percentage violations for three-alternative elections under IC and IAC. Figures 1 and 2 plot the IC and IAC data respectively. Note that Pairwise Majority cannot violate IIA, thus the violation frequencies are not computed.

		Number of Voters							
Voting System	3	4	5	10	30	50	100	1000	
Plurality	100	100	100	100	100	100	100	100	
Instant Runoff	100	100	100	100	100	100	100	100	
Top Two	100	100	100	100	100	100	100	100	
Minimax	100	100	100	100	100	100	100	100	
Copeland	97.35	73.63	99.95	99.99	100	100	100	100	
Dowdall	97.35	99.52	92.87	99.79	100	100	100	100	
Borda Count	97.35	73.63	92.87	94.79	99.97	99.99	100	100	
Coombs	97.35	73.63	92.87	94.79	99.99	100	100	100	
Baldwin	97.35	73.63	92.87	94.79	99.96	100	100	100	

Table 2: Percentage IIA Violations for 3 alternatives (IC Data)




Figure 1: IIA Violations in elections with three alternatives (IC)

Under IC, in elections with at least 30 voters, the frequency of violating IIA is 100%. In elections with three voters, Dowdall, Copeland, Coombs, Borda, and Baldwin show identical IIA violations.

		Number of Voters								
Voting System	3	4	5	10	30	50	100	1000		
Plurality	100	100	100	100	100	100	100	100		
Instant Runoff	100	100	100	100	100	100	100	100		
Top Two	100	100	100	100	100	100	100	100		
Minimax	100	100	100	100	100	100	100	100		
Copeland	89.08	56.66	97.3	97.75	99.8	99.99	100	99.99		
Dowdall	89.08	95.03	78.62	91.8	96.72	97.29	96.91	97.03		
Borda Count	89.08	56.66	78.62	74.29	84.52	84.54	84.14	83.64		
Coombs	89.08	56.66	78.62	74.29	87.4	86.41	85.7	85.52		
Baldwin	89.08	56.66	78.62	74.29	83.23	83.38	83.35	82.49		

Table 3: Percentage IIA Violations for 3 alternatives (IAC Data)

The IAC data has slightly more promising results and there is significant variation between systems. Coombs, Borda, and Baldwin all violate IIA less than 86% of the time, even for a sizable electorate. IIA violations appear to slightly decrease for Baldwin, Borda, and Coombs as the number of voters increases. The percentage violations for 1000 voters is less than that for 100 voters.

Our three-alternative data substantially differs from Dougherty and Heckelman (2020). For Dowdall, Coombs, Copeland, and Baldwin our algorithm reports greater percentage IIA violations. For Coombs and Copeland, the difference in the IAC data is around 20 percentage points. These differences are further explained in Section 7.




Figure 2: IIA Violations in elections with three alternatives (IAC)

6.2 Pareto Unanimity (U)

In the following, a *Pareto pair* is defined to be a pair of alternatives where all voters prefer one alternative in the pair over the other.

Only Group 1 systems can theoretically violate U. Even for these systems, U is nearly always satisfied. In elections with at least 30 voters, all systems have a 0% probability of violating U. U is almost always satisfied because encountering a Pareto pair is very rare, especially as the number of voters increases. Table 4 shows the frequency with which Instant Runoff violates U.

	(a)	IC Data				(b) I	AC Data		
Number of	Nu	umber of	alternativ	/es	Number of	Number of alternatives			
voters	3	4	5	6	voters	3	4	5	6
1	100.00	100.00	100.00	100.00	1	100.00	100.00	100.00	100.00
2	16.23	58.00	86.12	96.61	2	28.05	59.94	86.13	96.56
3	2.56	17.38	41.34	62.30	3	11.02	21.64	41.69	63.20
4	0.43	4.59	13.93	28.63	4	4.42	7.96	15.58	28.35
5	0.05	1.14	4.58	10.87	5	2.32	2.83	5.00	10.14
10	0.00	0.00	0.00	0.07	10	0.19	0.06	0.04	0.05
30	0.00	0.00	0.00	0.00	30	0.01	0.00	0.00	0.00
50	0.00	0.00	0.00	0.00	50	0.00	0.00	0.00	0.00
100	0.00	0.00	0.00	0.00	100	0.00	0.00	0.00	0.00
1000	0.00	0.00	0.00	0.00	1000	0.00	0.00	0.00	0.00

Table 4: Percentage Unanimity Violations Instant Runoff

For Instant Runoff, as the number of voters increases, the frequency of violations decreases because it is less likely that there exists a Pareto pair. As the number of alternatives increase, the frequency of U violations generally increases. Although percentage U violations are very low in most cases, they are common in elections with many alternatives and few voters. These trends are consistent with literature (Dougherty & Heckelman, 2020; Nurmi & Uusi-Heikkilä, 1986) and apply to all other systems that violate U. The results for Instant Runoff, Plurality, and Top Two are identical. Data for Minimax is identical for three alternatives, but for four or more alternatives, Minimax is less likely to violate U than other Group 1 systems. The differing performance of Minimax to other systems with greater number of alternatives is noted in literature (Dougherty & Heckelman, 2020).

Table 5 shows U violations for elections with four alternatives using IAC for Group 1 systems. Figure 3 is a graph of the data.

			Nu	mber o	of Voter	s			
Voting System	1	2	3	4	5	10	30	50	100
Plurality/InstantRunoff/TopTwo	100.00	59.94	21.64	7.96	2.83	0.06	0	0	0
Minimax	100.00	44.53	12.93	4.04	0.98	0.01	0	0	0

Table 5: Percentage U Violations 4 alternatives IAC




Figure 3: U Violations in elections with four alternatives (IAC)

While unanimity does not pose issues in large elections, for election with few voters and many candidates, this criterion is violated often.

6.3 Transitivity (TR)

Pairwise Majority is the only system studied that violates transitivity. Table 6 shows the percentage TR violations for Pairwise Majority. Figure 4 is a 3D plot of the IC data.

Key Observations

- 1. As the number of alternatives increases, the transitivity violations increase.
- 2. As the number of voters increase, transitivity violations decrease and appear to approach a constant. For six alternatives, the frequency of TR violations approaches around 67% and past 10,000 voters the frequencies fluctuate around that value. Studies have shown that as the number of voters tends to infinity, the probability of violating transitivity approaches a fixed value (Gehrlein, 1994). For three alternative elections using IAC, that value is 6.25%, which is consistent with our findings.

Table 6: Percentage Trans	itivity Violations	Pairwise Majority
---------------------------	--------------------	-------------------

71.18

69.16

68.60

68.11

67.61

(a) IC Data								
Number of	Nur	nber of	alternat	ives				
voters	3	4	5	6				
2	33.47	66.49	86.56	95.54				
3	5.54	17.12	32.60	48.93				
4	30.69	63.91	85.84	95.51				
5	6.87	21.04	39.59	57.39				
10	23.81	54.73	78.98	92.50				
30	17.31	43.78	68.77	86.03				
50	15.36	40.17	64.69	82.66				
100	13.08	35.99	60.08	78.92				

9.95

9.35

9.19

9.11

8.93

1000

5000

10000

30000

50000

29.11

27.45

27.36

26.48

26.23

51.63

49.01

48.90

47.95

47.96

(b) IAC Data									
Number of	Number of alternatives								
voters	3	4	5	6					
2	28.81	63.97	86.10	95.35					
3	3.57	14.97	31.66	48.87					
4	23.65	60.59	85.05	95.49					
5	4.77	19.17	38.71	57.56					
10	15.50	48.88	77.53	92.31					
30	9.58	35.85	66.47	85.58					
50	8.19	32.14	61.44	82.22					
100	7.23	28.51	56.13	78.15					
1000	6.46	24.54	47.83	69.68					
5000	6.24	24.03	46.76	67.73					
10000	6.34	24.13	46.72	67.08					
30000	6.45	24.06	46.92	66.98					
50000	6.34	23.90	46.49	67.21					




Figure 4: TR Violations in Pairwise Majority elections (IC)

Observations for three-alternative elections match the literature (Dougherty & Heckelman, 2020; Gehrlein, 1994; Niemi & Weisberg, 1968). For four-candidate elections, the limiting value for transitivity violations using IC in our study does not match the results in Niemi and Weisberg (1968). This is likely because we also include what they call Type 1 intransitivity, where there exists a candidates that wins head to head against all other candidates, but there are still cycles between the other candidates.

6.4 Joint Satisfaction of the Conditions

Simulations were run to find the percentage of times a voting system violates at least one criteria. The order in which criteria were tested was U, IIA, and TR. The other criterion (ND, UD) were not simulated as no system violates ND or UD. Joint-Satisfaction (JS) Frequencies, defined as the percentage of times systems satisfied all criteria simultaneously, serve as an indication of fairness of a voting system.

For every system that violates IIA, the JS Frequencies almost exactly align with the likelihood of satisfying IIA. In elections with more than three alternatives and two voters, no system had a JS Frequency of 100%, which is consistent with Arrow's Impossibility Theorem. Table 7 shows the results for Baldwin. All other data tables can be found in Appendix A.

Number of	Nun	nber of a	lternat	ives
voters	3	4	5	6
10	5.16	0.01	0.00	0.00
30	0.10	0.00	0.00	0.00
50	0.00	0.00	0.00	0.00
100	0.00	0.00	0.00	0.00
1000	0.00	0.00	0.00	0.00
5000	0.00	0.00	0.00	0.00
10000	0.00	0.00	0.00	0.00
30000	0.00	0.00	0.00	0.00
50000	0.00	0.00	0.00	0.00

(a) IC Data

Table 7: JS Frequencies	(%)	for	Baldwin
-------------------------	-----	-----	---------

Number of

10

30

50

100

1000

5000

10000

30000

50000

voters

Number of alternatives 3 4

0.25

0.27

0.28

0.16

0.22

0.27

0.29

0.22

0.33

25.16

16.87

16.79

17.14

17.27

16.63

16.68

17.07

16.88

5

0.01

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

6

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

(b) IAC Data

Table 8 shows a comparison	1 of all the voting systems for	or 3 alternative elections	under IAC and Figure
5 is a graph of the data.			

Table 8: Joint-Satisfaction Frequencies for 3 alternatives (IAC)

		Number of Voters						
Voting System	10	100	1000	5000	10000	50000		
Group 1	0.00	0.00	0.00	0.00	0.00	0.00		
Copeland	1.90	0.02	0.00	0.02	0.03	0.02		
Dowdall	7.41	3.16	3.13	3.11	2.97	3.10		
Borda Count	25.16	16.15	16.00	15.48	15.37	15.75		
Coombs	25.16	14.50	14.20	13.90	13.40	14.02		
Baldwin	25.16	17.14	17.27	16.63	16.68	16.88		
Pairwise Majority	84.50	92.77	93.54	93.76	93.67	93.66		

The data shows that Pairwise Majority is the most likely to satisfy all the criteria jointly for elections with three alternatives. As the number of voters increase in Pairwise Majority, the JS Frequencies increase, while for other systems the JS Frequencies either decrease or remain approximately constant. This is due to the fact that as the number of voters increase, transitivity violations become more rare, while, in general, IIA violations become more common.

There are also clear differences in performance in the other systems in three-alternative elections. Of the systems that satisfy transitivity, Borda Count, Baldwin, and Coombs have the highest JS Frequencies. Figure 6 shows a zoomed in version of Figure 5 for these three systems. For Borda, Baldwin, and Coombs the JS Frequencies also slightly increase as the number of voters becomes very large. This is because the IIA violations in these systems tend to marginally decrease as the number of voters increases from 30 to 50,000.

However, as the number of alternatives increases, the JS Frequencies decrease for every system. Figure 7 shows this trend clearly. In fact, for all systems excluding Pairwise Majority, the JS Frequencies for elections with at least 10 voters and 4 alternatives are less than 0.5%. This is because IIA is almost always violated in elections with at least four alternatives. Even for Pairwise Majority, the JS Frequencies sharply decrease as the number of alternatives increase, falling from 93.66% for three alternatives elections to 32.80% for six alternatives elections.




Figure 5: JS Frequencies in elections with three alternatives (IAC)




Figure 6: JS Frequencies for Black, Borda, Baldwin, Coombs

The conclusion for three-alternative elections is consistent with literature. In Dougherty and Heckelman (2020), the likelihood of violating at least one of Arrow's conditions is computed. Their study agrees that Pairwise Majority has the greatest JS Frequencies. However, due to differences in the IIA data as discussed in Section 7, the data for Copeland, Coombs, Dowdall, Borda, and Baldwin does not match exactly. Most systems have lower JS Frequencies than in the 2020 study.




Figure 7: JS Frequencies in elections with 50,000 voters (IAC)

7 Remarks about Differences in IIA Results

The IIA algorithm used in this paper is different than that of Dougherty and Heckelman (2020) and we describe those differences here. We start by describing our algorithm.

As IIA is an inter-profile condition, to check if a voting system violates IIA, multiple preference schedules have to be compared. Our IIA algorithm loops through all possible pairs of alternative. For each alternative pair A and B, the following test is done: First, find the societal preference between A and B. Then, let j equal the number of voters who prefer A over B and k the number of voters who prefer B over A. Then randomly generate 500 additional preference schedules in which j voters still prefer A over B and k voters prefer B over A. The number 500 was chosen taking various factors into consideration including computational resources and program run time. If any of these new preference schedules produce a different societal preference between A and B than the original, an IIA violation occurs. This is repeated for every possible pair of alternatives.

This algorithm can miss violations, which is more likely when the number of voters is high. As the number of voters increases, there are more possible preference schedules where j voters prefer A over B and k voters vote B over A. Thus, in the 500 new preference schedules generated, there may be no preference schedule that has a different societal order, even if one exists. Checking every single preference is impractical since there are $\left(\frac{c!}{2}\right)^v$ possible schedules such that each voter maintains their relative preference between a pair, where c is the number of alternatives and v is the number of voters.

In Dougherty and Heckelman (2020), a different IIA algorithm was used. Our algorithm finds substantially more IIA violations that the algorithm in Dougherty and Heckelman (2020), in particular with the IAC data. For Dowdall, Coombs, Copeland, and Baldwin our algorithm reports a greater percentage IIA violations. For Coombs and Copeland, the difference in the IAC data is around 20 percentage points.

7.0.1 Disagreement in IIA Conclusions under Copeland

Dougherty and Heckelman (2020) conclude that Copeland cannot violate IIA if a majority of voters have identical preferences. However, consider an election with three voters and three alternatives with the following preference schedule where a strict majority have preference order $A \succ B \succ C$:

						0	
1	st	Α	А	В	В	С	С
21	nd	В	С	Α	С	Α	В
3	rd	C	В	\mathbf{C}	А	C A B	А

Under Copeland, the societal preference order is $A \succ B \succ C$. Now, consider the pair of alternatives A and B. Suppose that one voter who has preference $A \succ_i B \succ_i C$ changes their preference to $C \succ_i A \succ_i B$. Now the preference schedule is:

	1	0	0	-	1	0
1st	Α	А	В	В	С	С
1st 2nd	В	\mathbf{C}	Α	\mathbf{C}	Α	В
3rd	C	В	\mathbf{C}	Α	В	Α

The societal preference order is now A = B = C. Even though all voters kept their relative preference between A and B, the societal ranking between them changed. Copeland can violate IIA even if a majority have identical preferences.

Dougherty and Heckelman (2020) also state that Copeland violates IIA at a greater frequency for even number of voters N than for the adjacent odd values N - 1 and N + 1. They claim that this is the case as there are more ties for elections with an even number of voters. While ties do make it easier to violate IIA, there are many cases for four voter elections where there are no ties and IIA is satisfied. Our analysis shows that for even number of voters, IIA violations in Copeland actually occur less frequently. For instance in three-alternative IC elections, Copeland violates IIA 97.35% of the time for three voters and only 73.63% of the time for four voters under IC.

Another, possibly related, disagreement was found while using their algorithm. For the purpose of comparing algorithms, we coded their IIA algorithm in our Python model for testing. Using their algorithm, we obtained the exact same data table that they did for both IC and IAC, with the exception of Copeland. For Copeland elections with three alternatives and at least four voters (in elections with three voters, the data is in agreement), their reported values are higher than what their algorithm produces. For instance, in elections with four voters and three alternatives under IC, simulations using their algorithm show a 73.7% chance of violating IIA, while the value reported in their paper is 88.9%. Notably, our IIA tables show 73.6% violations for identical elections.

This over-counting is also seen for IAC elections with more than four voters. For elections with 30 voters and three alternatives under IAC, our implementation of their algorithm yields 72% IIA violations, but the reported value is 78.5%. Note that as the number of voters increases, the extent of over-counting decreases. For IC, their data reaches 100% violations, and for IAC the difference between their values and our implementation of their algorithm becomes almost 0. The authors suspect an error in their Copeland method for greater than four voters.

8 Summary of Results

After comparing all systems, Pairwise Majority is the fairest system as it has the greatest likelihood of jointly satisfying all social choice criterion in elections with at least three alternatives and two voters. The other systems perform very poorly in comparison, primarily due to the extremely high frequency of IIA violations. Of the systems that satisfy transitivity, Baldwin, Borda, and Coombs perform the best, but have JS Frequencies of less than 17%.

Of all systems considered, Group 1 systems perform the worst. They violate IIA 100% of the time and as such, always have a 0% probability of jointly satisfying all criterion in elections with at least three alternatives.

Another key observation is that as the number of alternatives increases, the frequency with which voting systems violate criteria generally increases. For IIA, TR, and U, most systems increase in percentage violations as the number of alternatives increase. Thus, in elections with more alternatives, the JS Frequencies are significantly lower.

In the end, in alignment with Arrow's Theorem, no voting system is perfect. Each method gives rise to results which are arguably undesirable from the point of view of a society. However, not all voting systems are equally flawed. While some systems always violate social choice criteria, there are systems with a high likelihood of satisfying all criteria. Finding and implementing such systems can help achieve fairer and more representative electoral outcomes.

9 Declarations

Competing Interests The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data Availability Statement The raw data generated during this study is available upon request. All relevant processed data is included within this article and the appendix.

References

Arrow, K. J. (1963). Social choice and individual values. (2nd ed.) [First edition published in 1951]. Wiley.

- Börgers, C. (2010). *Mathematics of social choice*. Society for Industrial and Applied Mathematics. https://epubs.siam.org/doi/book/10.1137/1.9780898717624
- Dougherty, K. L., & Heckelman, J. C. (2020). The probability of violating arrow's conditions. European Journal of Political Economy, 65, 101936. https://doi.org/https://doi.org/10.1016/j.ejpoleco.2020. 101936
- Eğecioğlu, Ö., & Giritligil, A. E. (2013). The impartial, anonymous, and neutral culture model: A probability model for sampling public preference structures. *The Journal of Mathematical Sociology*, 37(4), 203– 222. https://doi.org/10.1080/0022250X.2011.597012
- Gehrlein, W. V. (1994). The expected likelihood of transitivity: A survey. Theory and Decision, 37(2), 175–209. https://doi.org/10.1007/bf01079265
- Hodge, J. K., & Kilma, R. E. (2018). The mathematics of voting and elections: A hands-on approach: Second edition (Vol. 30). American Mathematical Society.
- Matsumoto, M., & Nishimura, T. (1998). Mersenne twister: A 623-dimensionally equidistributed uniform pseudo-random number generator. ACM Trans. Model. Comput. Simul., 8(1), 3–30. https://doi. org/10.1145/272991.272995
- Mihara, H. R. (1997). Anonymity and neutrality in arrow's theorem with restricted coalition algebras. Social Choice and Welfare, 14(4), 503–512. Retrieved July 18, 2023, from http://www.jstor.org/stable/ 41106230
- Morreau, M. (2019). Arrow's Theorem. In E. N. Zalta (Ed.), *The Stanford encyclopedia of philosophy* (Winter 2019). Metaphysics Research Lab, Stanford University.
- Niemi, R. G., & Weisberg, H. F. (1968). A mathematical solution for the probability of the paradox of voting. Behavioral Science, 13(4), 317–323. https://doi.org/https://doi.org/10.1002/bs.3830130406
- Niou, E. M. S. (1987). A note on nanson's rule [ID: Niou1987]. Public Choice, 54(2), 191–193. https://doi. org/10.1007/BF00123006
- Nurmi, H. (1983). Voting procedures: A summary analysis. British Journal of Political Science, 13(2), 181– 208. Retrieved June 27, 2023, from http://www.jstor.org/stable/193949
- Nurmi, H., & Uusi-Heikkilä, Y. (1986). Computer simulations of approval and plurality voting: The frequency of weak pareto violations and condorcet loser choices in impartial cultures. *European Journal of Political Economy*, 2(1), 47–59. https://doi.org/https://doi.org/10.1016/S0176-2680(86)80004-0
- Patitsas, E. (2008). Can the borda count ever rank the condorcet loser ahead of the condorcet winner?
- Penn, E. M. (2015). Arrow's theorem and its descendants. In Handbook of social choice and voting (pp. 237– 262). Edward Elgar Publishing. https://EconPapers.repec.org/RePEc:elg:eechap:15584_14
- Xia, L. (2021). Semi-Random Impossibilities of Condorcet Criterion (Papers No. 2107.06435). arXiv.org. https://arxiv.org/abs/2107.06435
- Xia, L. (2020). The smoothed possibility of social choice. CoRR, abs/2006.06875. https://arxiv.org/abs/2006.06875

A Data Tables

A.1 IIA

Table 9: Percentage IIA Violations Plurality and Instant Runoff

(a)	IC	Data
-----	---------------------	------

(b) IAC Data

Number of	Number of alternatives							
voters	3	4	5	6				
3	100.00	100.00	100.00	100.00				
4	100.00	100.00	100.00	100.00				
5	100.00	100.00	100.00	100.00				
10	100.00	100.00	100.00	100.00				
30	100.00	100.00	100.00	100.00				
50	100.00	100.00	100.00	100.00				
100	100.00	100.00	100.00	100.00				
1000	100.00	100.00	100.00	100.00				
5000	100.00	100.00	100.00	100.00				
10000	100.00	100.00	100.00	100.00				
30000	100.00	100.00	100.00	100.00				
50000	100.00	100.00	100.00	100.00				

Number of	Number of alternatives							
voters	3	4	5	6				
3	100.00	100.00	100.00	100.00				
4	100.00	100.00	100.00	100.00				
5	100.00	100.00	100.00	100.00				
10	100.00	100.00	100.00	100.00				
30	100.00	100.00	100.00	100.00				
50	100.00	100.00	100.00	100.00				
100	100.00	100.00	100.00	100.00				
1000	100.00	100.00	100.00	100.00				
5000	100.00	100.00	100.00	100.00				
10000	100.00	100.00	100.00	100.00				
30000	100.00	100.00	100.00	100.00				
50000	99.99	100.00	100.00	100.00				

Table 10: Percentage IIA Violations Top Two

(a) IC Data

(b) IAC Data

Number of	Number of alternatives		Number of	Number of alternatives		ves			
voters	3	4	5	6	voters	3	4	5	6
3	100.00	100.00	100.00	100.00	3	100.00	100.00	100.00	100.00
4	100.00	100.00	100.00	100.00	4	100.00	100.00	100.00	100.00
5	100.00	100.00	100.00	100.00	5	100.00	100.00	100.00	100.00
10	100.00	100.00	100.00	100.00	10	100.00	100.00	100.00	100.00
30	100.00	100.00	100.00	100.00	30	100.00	100.00	100.00	100.00
50	100.00	100.00	100.00	100.00	50	100.00	100.00	100.00	100.00
100	100.00	100.00	100.00	100.00	100	100.00	100.00	100.00	100.00
1000	100.00	100.00	100.00	100.00	1000	100.00	100.00	100.00	100.00
5000	100.00	100.00	100.00	100.00	5000	100.00	100.00	100.00	100.00
10000	100.00	100.00	100.00	100.00	10000	100.00	100.00	100.00	100.00
30000	100.00	100.00	100.00	100.00	30000	100.00	100.00	100.00	100.00
50000	100.00	100.00	100.00	100.00	50000	100.00	100.00	100.00	100.00

(a) IC Data

(b) IAC Data

Number of	Nı	umber of	alternativ	ves		Number of	Nı	umber of	alternativ	ves
voters	3	4	5	6		voters	3	4	5	6
3	100.00	100.00	100.00	100.00]	3	100.00	100.00	100.00	100.00
4	100.00	100.00	100.00	100.00		4	100.00	100.00	100.00	100.00
5	100.00	100.00	100.00	100.00		5	100.00	100.00	100.00	100.00
10	100.00	100.00	100.00	100.00		10	100.00	100.00	100.00	100.00
30	100.00	100.00	100.00	100.00		30	100.00	100.00	100.00	100.00
50	100.00	100.00	100.00	100.00		50	100.00	100.00	100.00	100.00
100	100.00	100.00	100.00	100.00		100	100.00	100.00	100.00	100.00
1000	100.00	100.00	100.00	100.00		1000	100.00	100.00	100.00	100.00
5000	100.00	100.00	100.00	100.00		5000	99.99	100.00	100.00	100.00
10000	100.00	100.00	100.00	100.00		10000	100.00	100.00	100.00	100.00
30000	100.00	100.00	100.00	100.00		30000	100.00	100.00	100.00	100.00
50000	100.00	100.00	100.00	100.00		50000	99.99	100.00	100.00	100.00
L	1				1	L	1			

Table 12: Percentage IIA Violations Borda

(a) IC Data

Number of Number of alternatives voters 3 6 54 99.78 3 97.3599.98100.00100.00473.6399.96100.00592.87 99.39100.00 100.001094.79 100.00 100.00100.0030 100.00100.0099.97100.005099.99 100.00 100.00100.00100100.00100.00100.00100.001000 100.00 100.00100.00100.005000 100.00 100.00 100.00100.0010000100.00100.00100.00100.0030000 100.00100.00100.00 100.0050000100.00100.00100.00100.00

(b) IAC Data

Number of	Number of alternatives							
voters	3	4	5	6				
3	89.08	99.08	99.96	100.00				
4	56.66	99.81	100.00	100.00				
5	78.62	97.75	100.00	100.00				
10	74.29	99.90	100.00	100.00				
30	84.52	99.87	100.00	100.00				
50	84.54	99.90	100.00	100.00				
100	84.14	99.93	100.00	100.00				
1000	83.64	99.91	100.00	100.00				
5000	83.60	99.87	100.00	100.00				
10000	84.12	99.90	100.00	100.00				
30000	83.74	99.85	100.00	100.00				
50000	83.96	99.76	100.00	100.00				

Table 13: Percentage IIA	Violations (Coombs
--------------------------	--------------	--------

(a) IC Data

(b) IAC Data	ı
--------------	---

	()	ie Data					
Number of	Number of alternatives						
voters	3	4	5	6			
3	97.35	99.78	99.98	100.00			
4	73.63	99.96	100.00	100.00			
5	92.87	99.39	100.00	100.00			
10	94.79	100.00	100.00	100.00			
30	99.99	100.00	100.00	100.00			
50	100.00	100.00	100.00	100.00			
100	100.00	100.00	100.00	100.00			
1000	100.00	100.00	100.00	100.00			
5000	100.00	100.00	100.00	100.00			
10000	100.00	100.00	100.00	100.00			
30000	100.00	100.00	100.00	100.00			
50000	100.00	100.00	100.00	100.00			

Number of	Number of alternatives						
voters	3	4	5	6			
3	89.08	99.08	99.96	100.00			
4	56.66	99.81	100.00	100.00			
5	78.62	97.75	100.00	100.00			
10	74.29	99.91	100.00	100.00			
30	87.40	100.00	100.00	100.00			
50	86.41	99.97	100.00	100.00			
100	85.70	100.00	100.00	100.00			
1000	85.52	99.99	100.00	100.00			
5000	85.57	100.00	100.00	100.00			
10000	85.91	99.99	100.00	100.00			
30000	85.43	99.99	100.00	100.00			
50000	85.45	99.98	100.00	100.00			

Table 14: Percentage IIA Violations Dowdall

(a) IC Data

(b) IAC Data

Number of	Number of alternatives						
voters	3	4	5	6			
3	97.35	99.78	99.98	100.00			
4	99.52	99.96	100.00	100.00			
5	92.87	100.00	100.00	100.00			
10	99.79	100.00	100.00	100.00			
30	100.00	100.00	100.00	100.00			
50	100.00	100.00	100.00	100.00			
100	100.00	100.00	100.00	100.00			
1000	100.00	100.00	100.00	100.00			
5000	100.00	100.00	100.00	100.00			
10000	100.00	100.00	100.00	100.00			
30000	100.00	100.00	100.00	100.00			
50000	100.00	100.00	100.00	100.00			

Number of	Number of alternatives							
voters	3	4	5	6				
3	89.08	99.08	99.96	100.00				
4	95.03	99.81	100.00	100.00				
5	78.62	99.99	100.00	100.00				
10	91.80	100.00	100.00	100.00				
30	96.72	100.00	100.00	100.00				
50	97.29	100.00	100.00	100.00				
100	96.91	100.00	100.00	100.00				
1000	97.03	100.00	100.00	100.00				
5000	96.66	100.00	100.00	100.00				
10000	96.93	100.00	100.00	100.00				
30000	96.83	99.99	100.00	100.00				
50000	96.97	100.00	100.00	100.00				

Table 15: I	Percentage	IIA V	Violations	Copeland
-------------	------------	-------	------------	----------

(a) IC Data	(a)	IC	Data
-------------	-----	----	------

(b) IAC Data

Number of	Nι	umber of	alternativ	ves	Number of	Nι	umber of	alternativ	ves
voters	3	4	5	6	voters	3	4	5	6
3	97.35	99.78	99.98	100.00	3	89.08	99.08	99.96	100.00
4	73.63	99.96	100.00	100.00	4	56.66	99.81	100.00	100.00
5	99.95	100.00	100.00	100.00	5	97.30	99.99	100.00	100.00
10	99.99	100.00	100.00	100.00	10	97.75	100.00	100.00	100.00
30	100.00	100.00	100.00	100.00	30	99.80	100.00	100.00	100.00
50	100.00	100.00	100.00	100.00	50	99.99	100.00	100.00	100.00
100	100.00	100.00	100.00	100.00	100	100.00	100.00	100.00	100.00
1000	100.00	100.00	100.00	100.00	1000	99.99	100.00	100.00	100.00
5000	100.00	100.00	100.00	100.00	5000	99.99	100.00	100.00	100.00
10000	100.00	100.00	100.00	100.00	10000	99.98	100.00	100.00	100.00
30000	100.00	100.00	100.00	100.00	30000	99.97	100.00	100.00	100.00
50000	100.00	100.00	100.00	100.00	50000	99.98	100.00	100.00	100.00

Table 16: Percentage IIA Violations Baldwin

(a) IC Data

(b) IAC Data

Number of	Nı	umber of	alternativ	ves
voters	3	4	5	6
3	97.35	99.78	99.98	100.00
4	73.63	99.96	100.00	100.00
5	92.87	99.39	100.00	100.00
10	94.79	99.99	100.00	100.00
30	99.96	100.00	100.00	100.00
50	100.00	100.00	100.00	100.00
100	100.00	100.00	100.00	100.00
1000	100.00	100.00	100.00	100.00
5000	100.00	100.00	100.00	100.00
10000	100.00	100.00	100.00	100.00
30000	100.00	100.00	100.00	100.00
50000	100.00	100.00	100.00	100.00

Number of	Nι	Number of alternatives						
voters	3	4	5	6				
3	89.08	99.08	99.96	100.00				
4	56.66	99.81	100.00	100.00				
5	78.62	97.75	100.00	100.00				
10	74.29	99.76	100.00	100.00				
30	83.23	99.76	100.00	100.00				
50	83.38	99.78	100.00	100.00				
100	83.35	99.75	100.00	100.00				
1000	82.49	99.76	100.00	100.00				
5000	82.63	99.73	100.00	100.00				
10000	83.14	99.74	100.00	100.00				
30000	82.74	99.74	100.00	100.00				
50000	82.87	99.70	100.00	100.00				

(a) IC Data									
Number of	Nı	Number of alternatives							
voters	3	4	5	6					
1	100.00	100.00	100.00	100.00					
2	16.23	58.00	86.12	96.61					
3	2.56	17.38	41.34	62.30					
4	0.43	4.59	13.93	28.63					
5	0.05	1.14	4.58	10.87					
10	0.00	0.00	0.00	0.07					
30	0.00	0.00	0.00	0.00					
50	0.00	0.00	0.00	0.00					
100	0.00	0.00	0.00	0.00					
1000	0.00	0.00	0.00	0.00					

Table 17: Percentage U Violations Plurality and Top Two

(b) IAC Data									
Number of	Nı	Number of alternatives							
voters	3	4	5	6					
1	100.00	100.00	100.00	100.00					
2	28.05	59.94	86.13	96.56					
3	11.02	21.64	41.69	63.20					
4	4.42	7.96	15.58	28.35					
5	2.32	2.83	5.00	10.14					
10	0.19	0.06	0.04	0.05					
30	0.01	0.00	0.00	0.00					
50	0.00	0.00	0.00	0.00					
100	0.00	0.00	0.00	0.00					
1000	0.00	0.00	0.00	0.00					

Table 18: Percentage U Violations Minimax

(a) IC Data

(b) IAC Data

	· · ·						· · ·			
Number of	Nu Nu	umber of	alternativ	ves	Nun	nber of	Number of alternatives			ves
voters	3	4	5	6	vote	ers	3	4	5	6
1	100.00	100.00	100.00	100.00		1	100.00	100.00	100.00	100.00
2	16.23	42.04	65.70	81.92		2	28.05	44.53	65.78	82.17
3	2.56	8.90	18.46	29.53		3	11.02	12.93	19.29	30.03
4	0.43	1.61	3.72	6.99		4	4.42	4.04	4.56	6.42
5	0.05	0.34	0.73	1.21		5	2.32	0.98	0.89	1.23
10	0.00	0.00	0.00	0.00		10	0.19	0.01	0.00	0.00
30	0.00	0.00	0.00	0.00		30	0.01	0.00	0.00	0.00
50	0.00	0.00	0.00	0.00		50	0.00	0.00	0.00	0.00
100	0.00	0.00	0.00	0.00		100	0.00	0.00	0.00	0.00
1000	0.00	0.00	0.00	0.00		1000	0.00	0.00	0.00	0.00

A.3 JS Frequencies

Recall that the JS Frequencies for Plurality, Instant Runoff, Minimax, and Top Two are always less than 0.01% in elections with at least three alternatives and two voters. For the rest of the systems, the JS Frequency tables are below.

Table 19: JS	Frequencies	Borda	Count
--------------	-------------	-------	-------

(a)	IC	Data
-----	----	------

	. ,			
Number of	Num	ber of	alterna	tives
voters	3	4	5	6
10	5.16	0.01	0.00	0.00
30	0.11	0.00	0.00	0.00
50	0.00	0.00	0.00	0.00
100	0.00	0.00	0.00	0.00
1000	0.00	0.00	0.00	0.00
5000	0.00	0.00	0.00	0.00
10000	0.00	0.00	0.00	0.00
30000	0.00	0.00	0.00	0.00
50000	0.00	0.00	0.00	0.00

(b)	IAC	Data
(b)	IAC	Data

Number of Number of alternatives							
Number of	Num	oer of a	atterna	tives			
voters	3	4	5	6			
10	25.16	0.12	0.00	0.00			
30	15.48	0.07	0.00	0.00			
50	15.71	0.18	0.00	0.00			
100	16.15	0.06	0.00	0.00			
1000	16.00	0.15	0.00	0.00			
5000	15.48	0.14	0.00	0.00			
10000	15.37	0.15	0.00	0.00			
30000	15.89	0.09	0.00	0.00			
50000	15.75	0.14	0.00	0.00			

(a) IC Data

Number of	Number of alternatives				
voters	3	4	5	6	
10	5.16	0.01	0.00	0.00	
30	0.02	0.00	0.00	0.00	
50	0.00	0.00	0.00	0.00	
100	0.00	0.00	0.00	0.00	
1000	0.00	0.00	0.00	0.00	
5000	0.00	0.00	0.00	0.00	
10000	0.00	0.00	0.00	0.00	
30000	0.00	0.00	0.00	0.00	
50000	0.00	0.00	0.00	0.00	

(b) IAC Data

Number of	Number of alternatives				
voters	3	4	5	6	
10	25.16	0.10	0.00	0.00	
30	12.81	0.01	0.00	0.00	
50	13.96	0.00	0.00	0.00	
100	14.50	0.00	0.00	0.00	
1000	14.20	0.02	0.00	0.00	
5000	13.90	0.02	0.00	0.00	
10000	13.40	0.01	0.00	0.00	
30000	14.29	0.02	0.00	0.00	
50000	14.02	0.01	0.00	0.00	

Table 21: JS Frequencies Dowdall

(a) IC Data

Number of	Number of alternatives				
voters	3	4	5	6	
10	0.22	0.00	0.00	0.00	
30	0.00	0.00	0.00	0.00	
50	0.00	0.00	0.00	0.00	
100	0.00	0.00	0.00	0.00	
1000	0.00	0.00	0.00	0.00	
5000	0.00	0.00	0.00	0.00	
10000	0.00	0.00	0.00	0.00	
30000	0.00	0.00	0.00	0.00	
50000	0.00	0.00	0.00	0.00	

(b) IAC Data

Number of	Number of alternatives				
voters	3	4	5	6	
10	7.41	0.02	0.00	0.00	
30	3.30	0.00	0.00	0.00	
50	3.27	0.00	0.00	0.00	
100	3.16	0.00	0.00	0.00	
1000	3.13	0.01	0.00	0.00	
5000	3.11	0.00	0.00	0.00	
10000	2.97	0.00	0.00	0.00	
30000	2.98	0.00	0.00	0.00	
50000	3.10	0.00	0.00	0.00	

Table	22:	JS	Frequencies	Copeland
-------	-----	----	-------------	----------

(a) IC Data							
Number of	Nun	nber of a	alternat	ives			
voters	3	4	5	6			
10	0.00	0.00	0.00	0.00			
30	0.00	0.00	0.00	0.00			
50	0.00	0.00	0.00	0.00			
100	0.00	0.00	0.00	0.00			
1000	0.00	0.00	0.00	0.00			
5000	0.00	0.00	0.00	0.00			
10000	0.00	0.00	0.00	0.00			
30000	0.00	0.00	0.00	0.00			
50000	0.00	0.00	0.00	0.00			

(b) IAC Data							
Number of	Nun	nber of a	alternat	ives			
voters	3	4	5	6			
10	1.90	0.00	0.00	0.00			
30	0.15	0.00	0.00	0.00			
50	0.08	0.00	0.00	0.00			
100	0.02	0.00	0.00	0.00			
1000	0.00	0.00	0.00	0.00			
5000	0.02	0.00	0.00	0.00			
10000	0.03	0.00	0.00	0.00			
30000	0.01	0.00	0.00	0.00			
50000	0.02	0.00	0.00	0.00			

(a) IC Data

Number of	Number of alternatives			
voters	3	4	5	6
10	5.16	0.01	0.00	0.00
30	0.10	0.00	0.00	0.00
50	0.00	0.00	0.00	0.00
100	0.00	0.00	0.00	0.00
1000	0.00	0.00	0.00	0.00
5000	0.00	0.00	0.00	0.00
10000	0.00	0.00	0.00	0.00
30000	0.00	0.00	0.00	0.00
50000	0.00	0.00	0.00	0.00

(b) IAC Data

Number of	Number of alternatives			
voters	3	4	5	6
10	25.16	0.25	0.01	0.00
30	16.87	0.27	0.00	0.00
50	16.79	0.28	0.00	0.00
100	17.14	0.16	0.00	0.00
1000	17.27	0.22	0.00	0.00
5000	16.63	0.27	0.00	0.00
10000	16.68	0.29	0.00	0.00
30000	17.07	0.22	0.00	0.00
50000	16.88	0.33	0.00	0.00

Table 24: JS Frequencies Pairwise Majority

(a) IC Data

Number of	Number of alternatives			
voters	3	4	5	6
10	76.19	45.27	21.02	7.50
30	82.69	56.23	31.23	13.97
50	84.64	59.83	35.31	17.34
100	86.92	64.02	39.92	21.08
1000	90.05	70.89	48.37	28.82
5000	90.65	72.55	50.99	30.84
10000	90.81	72.64	51.11	31.40
30000	90.89	73.53	52.06	31.89
50000	91.08	73.77	52.04	32.39

(b) IAC Data

Number of	Number of alternatives			
voters	3	4	5	6
10	84.50	51.12	22.47	7.69
30	90.42	64.15	33.53	14.42
50	91.81	67.86	38.56	17.78
100	92.77	71.49	43.87	21.85
1000	93.54	75.46	52.17	30.32
5000	93.76	75.97	53.24	32.27
10000	93.67	75.87	53.28	32.92
30000	93.55	75.94	53.08	33.02
50000	93.66	76.10	53.51	32.80