Our Dark Matter Stopping in the Earth

H.B. Nielsen, Niels Bohr Institut, and Colin D. Froggatt, Glasgow University

"Bled", July, 2023

Abstract

We have worked for some time on a model for dark matter, in which dark matter consists of small bubbles of a new speculated type of vacuum, which are pumped up by some ordinary matter such as diamond, so as to resist the pressure of the domain wall separating the two vacua. Here we put forward thoughts on, how such macroscopic pearls would have their surrounding dust cleaned off passing through the atmosphere and the Earth, and what their distribution would be as a function of the depth of their stopping point and the distribution of the radiation emitted from them. In our model we assume that they radiate 3.5 keV electrons and photons, after having been excited during their passage into the Earth. The purpose of such an estimation of the radiation distribution is to explain the truly mysterious fact that, among all the underground experiments seeking dark matter colliding with the Earth material, only the DAMA-LIBRA experiment has seen any evidence of dark matter. This is an experiment based on solid NaI scintillators and is rather deep at 1400 m. It is our point that we can arrange the main radiation to appear in the relatively deep DAMA-LIBRA site, and explain that the dark matter pearls cannot stop in a fluid, such as xenon in the xenon based experiments.

1 Introduction

It is still a great *mystery* of what the *dark matter*, of which one mostly has seen the effect of its gravitation, *consists*. An exceptionally great mystery in this connection is that among the experiments looking for dark matter hitting the Earth and being detected deep underground - to avoid the cosmic radiation background - there is only one experiment, DAMA-LIBRA [1] which has seen any evidence for dark matter. The experiments based on the fluid scintillator, fluid xenon, typically claim direct disagreement with DAMA-LIBRA, by obtaining so low upper limits on the cross section for the dark matter - using a WIMP model - that the observations by DAMA could

^{*}Speaker at the Work Shop "What comes beyond the Standard Models" in Bled.

not avoid having been seen in e.g. LUX. Our model has dark matter, that is not as weakly interacting as the usual WIMP model assumes. Rather our dark matter pearls consist of bubbles of a new (speculated) type of vacuum, containing ordinary matter and compressed to an outrageously high density. So, although still being per kg much less interacting than ordinary matter, our dark matter is much more strongly interacting per kg than the WIMPs usually considered. Thus our pearls of dark matter should not be called WIMPs but rather only IMPs (Interacting Massive Particles). The essential point for the present paper is, what happens to our model dark matter when hitting the Earth. It is not so much that they consist of a new type of vacuum etc. but rather that they are macroscopic objects causing a much bigger interaction that matters. However they still must be so massive compared to their cross section, that they do not just function like ordinary matter. But that does not prevent them getting stopped in the Earth, although with an appreciably longer stopping length than ordinary matter.

It is rather easy in our model to adjust parameters, so as to obtain whatever stopping length one might want for our dark matter pearls. At least we can easily believe that we could fit them to have a stopping length of the order of the depth of the DAMA-LIBRA experiment. Then if they were arranged to emit most of their excitation energy where they get stopped, they would be appreciably more visible to experiments in the depth of DAMA-LIBRA than in other depths.

If really the stopping of dark matter is needed for their easy observation, then the experiments with a fluid scintillator would be severely disfavoured because a dark matter pearl cannot really fully stop in a fluid. A little piece of fluid around the pearl will at least by gravity follow the pearl as it falls down, and it would spend much less time in a liquid xenon experiment than in a NaI(Tl) one. Even other NaI(Tl) experiments, if having a lower depth under the Earth surface than DAMA, might see only a little of the dark matter, because it passes through such experiments too fast and too little of the radiation from its excitation would be detected at such higher up experiments.

So our model of dark matter consists of small but still macroscopic pearls, in the sense of each consisting of many atoms, then made effectively "darker" by having these atoms concentrated by the domain wall between the new vacuum and the ordinary vacuum, which we assume to be degenerate (i.e. with the same energy density). We shall first briefly mention the evidence from dwarf galaxies for the interaction of dark matter with itself.

2 Our dark matter, stopping in the Earth, etc.

The two most crucial properties of our model [2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14] are as follows.

Self-interacting d



Figure 7. Same as Fig. 6, but extended to cover the range of MW-(~200 km/s) and cluster-size (~1000 – 5000 km/s) haloes' velocities. The figure shows upper and lower limits for σ/m_{χ} taken for substructure abundance studies (e.g. Volgelsberger et al. 2012 and Zavala et al. 2013), as well as based on halo shape/ellipticity studies and cluster lensing surveys (see text).

Figure 1: Self-interaction cross section over mass of the particles (pearls in our model) obtained by fitting velocities of (of course) the ordinary matter in dwarf galaxies around our Milky Way.

• Our dark matter is not so dark as WIMPs. Our dark matter pearls interact so much that they get stopped in the Earth.

We actually speculate that our dark matter pearls get essentially stopped in the Earth, down from their ~ 300 km/s galactic velocity, at a depth around that of DAMA-LIBRA. Velocity dependent fits of their "inverse darkness" $\frac{\sigma}{M}$ by Correa [15] from studying dwarf Galaxies is seen in Figure 1, which was motivated by the failure of the model that dark matter has only gravitational interactions as seen in Figure 2.

From Figure 2 we obtain the "inverse darkness" values:

$$\frac{\sigma}{M} \approx 150 cm^2/g = 15m^2/kg \,(\text{low velocity})$$
 (1)

$$\frac{\sigma}{M} \approx 1.5 cm^2/g = 0.15 m^2/kg \,(200 \,\mathrm{km/s})$$
 (2)

But here we must admit that with these "inverse darknesses" the pearls would not go so deep as we would like to make them stop in the depth of the DAMA-LIBRA experiment. So we have to help our model a little



Figure 2: This figure should illustrate, that when Correa simulated the dark matter distribution, under the assumption of a purely gravity interaction, and regained a prediction for the star velocities it was *not* successful. So there is a phenomenological call for e.g. interaction of the dark matter with itself.

bit by declaring that the pearls have caught up dust around them so as to get less dark, but that in the atmosphere or the upper shielding of the Earth this dust is washed off. Washing off the dust makes the cross section smaller presumably without losing much weight, so that after such a cleaning the inverse darkness is decreased and thus the stopping length is increased.

• The dark matter particles can be excited to radiate 3.5 keV X rays or presumably also electrons of the same energy per particle.

3 A hard problem: How can ANAIS disagree with DAMA-LIBRA?

The two underground experiments, ANAIS [16] and DAMA-LIBRA [1], and also a third one COSINE-100 [17], which has not yet reached sufficient accuracy to properly disagree with either of the first two, looking for dark matter are *very similar*.

- Both use NaI(Tl= Thallium),
- Both look for seasonal variation,
- Depth is 1400 m ~ 4200 m.w.e. for DAMA-LIBRA, while ANAIS is at 2450m.w.e. (COSINE-100 has a similar depth to ANAIS)

But ANAIS "sees" no dark matter yet and claims to deviate by 3 σ from DAMA-LIBRA, which sees 0.0103 cpd/kg/keV. Our idea to resolve this problem is that:

Dark matter passes quickly by ANAIS, but slows down or stops at DAMA-LIBRA, see Figure 3. If the DAMA-LIBRA signal is due to radiating dark matter pearls then this could solve the problem.

The COSINE-100 group searched for an annual modulation amplitude in their data taken over 3 years in the 1-6 keV region with the phase fixed to the DAMA value of 152.5 days. They subtracted their calculated time dependent background and obtained a positive amplitude of 0.0067 ± 0.0042 cpd/kg/keV, which is to be compared with the DAMA-LIBRA amplitude of 0.0103 cpd/kg/keV. The large error on the COSINE-100 amplitude means that it is consistent with both the DAMA-LIBRA result and no seasonal variation at all. But then they analysed their own COSINE-100 data in the "same way as DAMA" [18] by subtracting a constant background taken to be the average rate over one year, and generated what they consider to be a spurious seasonal variation. In fact they found a modulation amplitude of the opposite phase -0.0441 ± 0.0057 cpd/kg/keV (the wrong season has over abundance). Interestingly the WIMP model could not obtain such a result, but it is possible with our dark matter pearls.

In our model an experiment at some given depth, would have just those dark matter particles stop in the instrument, which have just the right velocity.

3.1 How slowdown can help and a huge day/night effect

In Figure 3 we illustrate how one must think about dark matter particles coming in and at first having high speed but then slowing down due to the interaction with the earth or stone in the shielding. When the speed gets low even the Earth's gravity can make so much extra acceleration that the trajectories of the dark matter pearls become curved.



Signal from slowing down radiating particles

Figure 3: Under the assumption, that dark matter after excitation under the stopping radiates say electrons, and that it is these electrons, that have the best chance to be observed, the fast moving dark matter pearls in the region closer to the surface of the Earth leave only weak signals, while a much stronger signal will be left in the counters of DAMA-LIBRA in a depth, where the dark matter pearls have slowed down.(m. w. e. = meter water equivalent). Notice that we have drawn the trajectories as a bit curved, especially where the dark matter pearls have almost stopped, so much as to move slower. Then namely the effect of the Earth's gravity becomes relatively larger.

But now we have to call attention to the fact that our model would - if we do not as we shall do in a moment find a way of avoiding it - predict a huge variation of the counting rate between day and night. The point is that the solar system moves with about 200 km/s relative to the dark matter center of mass system and, although the spread in the dark matter velocity in the radial direction to and from the center of the galaxy is presumably large, the spread in the direction around the galaxy in which the solar system moves will be much smaller. It might be as small as say ± 90 km/s in which case the average of the dark matter velocity seen from the Earth or the Sun would be say 2 times as large as the spread. On the other hand with stopping dark matter, as in our model, the dark matter can only come into the Earth and to an underground experiment from the side of the Earth on which the experiment is located. It can only come from the direction conceived as from above as seen from the experiment. This means that there will be a huge difference in the rate of visible impacts depending on whether the experiment in on the forward or the backward side of the Earth relative to the motion of the Earth seen from the dark matter average rest system.

When the experiment in question is on the front side of the Earth in its motion relative to the dark matter rest system, most of the dark matter can be overtaken by the Earth and fall into the experiment from the sky side, and one should get a very large rate of impact in this situation. Twelve hours later the experiment will be on the back side and now only very few dark matter particles have such a high individual velocity compared to the average velocity of the dark matter relative to the Earth, that they can run in the opposite direction to the majority of the dark matter. So in this case only relatively few dark matter particles can be observed.

For an underground experiment only particles running down towards its site can hit it, except that the Earth can overtake some slow ones. See Figure 4.

In Figure 5 we see the situation of the Earth running relative to the center of mass for dark matter. With the experiment in front of a huge part of the dark matter, all except the white tip would hit the Earth, while in the case of the experiment being on the backside only a small tip, the green one, will hit the Earth.

In Figure 6 we have added some small displacement lines to illustrate how the borders between the amounts of dark matter hitting and not hitting the Earth is shifted by the relatively small velocity changes due to the season. When the experiment is in front in the motion towards the dark matter center of mass motion the hitting rate increases in the summer when the relative velocity is larger. However, when we are in the time of day when the experiment is on the backside the larger summer relative velocity will mean that the little tip of dark matter hitting the Earth gets even smaller. So in the time of day-night in which the dark matter mainly comes in from the side of the Earth opposite to that of the experiment, the seasonal effect



Figure 4: If the stopping length is so short as we speculate in our model, it is only dark matter that penetrates into the Earth from the side of the experiment, e.g. COSINE-100, that has any chance of being detected, while in WIMP-models dark matter particles coming through the earth-interior before interacting is not at all excluded. This figure concentrates on three examples of the velocity of particles having managed to come in on the front side of the Earth say, on which side the experiments lies. Whether dark matter particles now hit the earth, and how fast, depends on the relative velocity.



Figure 5: On this figure the Gaussian distribution is the distribution of the dark matter particle velocities in the rest frame for the average velocity of the dark matter, the center of mass. Let us imagine the Earth at first coming in from the right side of the figure with the site of the experiment, COSINE-100 say, on the place having the left direction in zenith. Then, if the velocity towards the dark matter of the Earth - moving to the left - is very high the dark matter in the velocity bands denoted on the figure as dark blue, red and green, i.e. all the ranges except the white one, would hit the Earth on the side of this experiment. But now if the Earth moved slower compared to the dark matter, say it followed the center of mass for the dark matter, then only the red and the green amounts of dark matter would hit the Earth. And if the Earth "moved away" in the sense that the experiment was on the backside of the Earth compared to its velocity in the dark matter center of mass frame, then say only the green part of the dark matter would hit the Earth in the region of the experiment. In this case, where the experiment is on the backside relative to the motion only the very fastest part, the green band say, would be able to overtake the Earth and hit the experiment.



Figure 6: Like in figure 5 we consider here a Gaussian dark matter velocity distribution, and look for how with different velocities of the Earth (with the experiment on the left side) moving in the dark matter sees more dark matter in the experiment the faster this Earth moves forward. E.g. if the Earth moves with the experiment on the left side to the left, only the white part moves too fast for the Earth to catch up. But now we are interested in how small variations in the Earth's velocity, exemplified by the vertical lines drawn to the right of the color separation places on the figure. They symbolize a little bit slower Earth. The interesting point is that the effect of such a small velocity lowering is not the same independent of how fast the Earth moves already compared to the dark matter average. Even relatively the extra velocity gives different extra contributions, different even taken relative to the original one.

is actually opposite to that when the dark matter comes in on the same side as the experiment, i.e there are most counts in the winter and fewer in the summer.

This is definitely a very interesting prediction of ours, if the data on the detection of the dark matter is sufficiently detailed that one can distinguish day and night so to say. But now in detail our own model has it that the dark matter pearls are counted by means of the radiation of say electrons which are sent out delayed compared to the time at which the pearls were excited. If the decay lifetime is several days, then the very big day night oscillation gets washed out in our model and would not be seen. Indeed if the day night effect was as strong as we predict - percentwise stronger than the seasonal one - at first, i.e. if we did not have the washout by the pearls being excited and decaying with a several day lifetime scale, then it should have been easy for DAMA-LIBRA to have observed the day-night variation!

(So it means that a model of our type, with only a limited penetration into the Earth, would not work unless we also have a washout due to the decay lifetime being of order of at least days.)

When our dark matter pearls slow down, they either stop completely or continue falling slowly under gravity.

- If they stop completely, we should find a lot of stationary dark matter to be dug out as heavy pearls (much like if it were gold dust and one should wash it out).
- If they sink slowly, they of course go deeper inside the Earth.

4 Crude estimate of stopping distance and mass of a dark matter pearl

For simplicity we shall only consider the motion in one direction, down, and ignore the rest. The depth, into which a dark matter particle will penetrate before (effectively) stopping, will be a smooth function of its velocity relative to the Earth. So "topologically" the *distribution of stopping-depths* will reflect the *initial velocity spectrum* in the downward direction, as illustrated in Figure 7.

We note that DAMA-LIBRA is about twice as deep down as ANAIS and COSINE-100, see Figure 8. This figure illustrates that in our model ANAIS and COSINE-100 could, in principle, have annual modulation amplitudes with the opposite phase to DAMA-LIBRA, but of course it is not necessary.

In order for our dark matter pearls to stop just at DAMA-LIBRA crudely, we need a penetration depth of the order of magnitude:

Penetration depth
$$L = \frac{M}{\sigma} * 30/\rho_{stone} \approx 1km$$
 (3)

$$\Rightarrow \frac{M}{\sigma} \approx 10^5 kg/m^2 \tag{4}$$

$$\Rightarrow \frac{\sigma}{M} \approx 10^{-5} m^2 / kg \tag{5}$$

The number 30 is a crude estimate involving a logarithm and factors of order unity roughly - see equation (42).

We may compare this value of $\frac{\sigma}{M}$ for reaching just the DAMA wanted order of magnitude with the numbers given in equations (1, 2). This means that from the low velocity value (1) we need a decrease by a factor 1.5×10^6 and from the 200 km/s value (2) a decrease by 1.5×10^4 . Thus we must take it that so much dust around the dark matter pearl is washed off during the impact, that the cross section goes down by a factor of the order of respectively a million and 10000. Taking it that the relevant velocity is of the order of 200 km/s, the factor ten thousand is enough.



Figure 7: If one has a definite formula for the drag force stopping the pearls as a function of the velocity, then this formula will give the effective stopping depth - or rather the penetration distance - as a function of the initial impact velocity (we consider only the vertical direction as an approximation). And then there will be simply a transformation from the impact velocity in the vertical direction to the depth at the point of stopping (only "effectively", if the particle really continues with a much lower fixed velocity, which is then neglected here). Here we have drawn symbolically the image under this transformation of the initial vertical velocity distribution. It still bears some similarity to the supposed Gaussian velocity distribution.



Figure 8: This is again the vertical density distribution transformed into the depth spectrum (ignoring the horizontal components of the velocity), but now we have shown three variants corresponding to the seasons in which the velocity of the dark matter relative to the Earth is different. In summer this relative velocity is higher and thus the stopping spectrum is for summer on average a bit deeper. On the contrary the winter spectrum lies a bit less deep. An average spectrum is also drawn, which we could consider to be that for autumn and spring. Rather than to be realistic, the drawing has been made with the purpose of illustrating that indeed there is the possibility of the relatively close to the surface lying experiments ANAIS and COSINE-100 getting *less counts in the summer than in winter* opposite to the DAMA-LIBRA-experiment, which is about twice as deep (1400 m) and should see most events in the summer.

4.1 Size of the bubble

The need for assuming the dirt to be washed off the pearl means that the inverse darkness $\frac{\sigma}{M} = 10^{-5} m^2/kg$, identified with the one needed for the penetration depth just reaching DAMA-LIBRA, must be that of the supposedly hard bubble making up the main and most heavy part of the dark matter pearls. Now we used to think that we could estimate the density of this bubble filled with ordinary matter under high pressure by using a dimensional argument [8] - of course rather uncertain - to obtain the Fermi-momentum from the HOMO-LUMO gap E_H identified with the $E_H = 3.5 \, keV$ X-ray photon energy line likely to be associated with the dark matter:

$$E_H = \sqrt{2} \left(\frac{\alpha}{c}\right)^{3/2} E_f. \tag{6}$$

Here E_f is the Fermi energy in the compressed ordinary matter inside the new vacuum bubble, and α is the fine structure constant, which for our a bit special dimensional argument has been taken to be of dimension velocity, so that it is α/c that is the usual fine structure constant 1/137.036... We actually even calculated the $\sqrt{2}$, but that would clearly be far outside the expected accuracy. From this dimensional argumentation we then got the density of the material inside the new vacuum bubble:

$$\rho_B = 5 * 10^{11} kg/m^3. \tag{7}$$

Of course the radius of the bubble with the compressed ordinary matter is given as

$$R = \sqrt[3]{\frac{3M}{\rho_B * 4\pi}} = \sqrt[3]{M} * 8 * 10^{-4} kg^{-1/3} m.$$
(8)

and cross section

$$\sigma = \pi R^2 = \pi^{1/3} 3^{2/3} 4^{-2/3} M^{2/3} \rho_B^{-2/3}$$
(9)

$$= 1.21 * (M/\rho_b)^{2/3} = 1.9 * 10^{-8} kg^{-2/3} m^2 * M^{2/3}.$$
 (10)

So we obtain

$$\frac{\sigma}{M} = 1.9 * 10^{-8} kg^{-2/3} m^2 (\sqrt[3]{M})^{-1}$$
(11)

To achieve the value of $\sigma/M = 10^{-5}m^2/kg$ to just reach DAMA we need

$$10^{-5}m^2/kg = 1.9 * 10^{-8}kg^{-2/3}m^2M^{-1/3}$$
(12)

giving
$$\sqrt[3]{M} = 1.9 * 10^{-3} kg^{1/3}$$
 (13)

i.e.
$$M = 7 * 10^{-9} kg.$$
 (14)

In a moment we shall see below, by considering the energy of impact of the dark matter, that we would for the purpose of what DAMA sees have preferred the value $9.4 * 10^{-17} kg$.

The density of dark matter in the region of the solar system is

$$D_{sun} = 0.3 \, GeV/cm^3. \tag{15}$$

The rate of impact energy is then

$$"Rate" = v * D_{sun} \tag{16}$$

$$= 300km/s * 0.3 * 1.79 * 10^{-27}kg/(10^{-6}m^3)$$
(17)

$$= 1.6 * 10^{-16} kg/m^2/s \tag{18}$$

$$= 1.4 * 10^{-11} kg/m^2/day \tag{19}$$

Now let us call the mass of the individual pearls M and then we can compute the number density of pearls of dark matter:

number density falling = $1.4 * 10^{-11} kg/m^2/day/M$ (20)

Spreading over a kilometer we obtain:

number density =
$$1.4 * 10^{-14} kg/m^3/day/M$$
 (21)

For an earth density $= 3000 kg/m^3$ we then have:

number per kg of earth =
$$4.7 * 10^{-18} / Mper \, day$$
 (22)

DAMA saw
$$S_M = 0.01 cpd/kg/keV$$
 (23)

Over say 5 keV:
$$5keVS_m = 0.05cpd/kg$$
 (24)

If there is only one count per particle, we get

$$4.7 * 10^{-18} kg/Mcpd = 0.05 cpd \tag{25}$$

giving
$$M = 9.4 * 10^{-17} kg$$
 (26)

$$= 5.3 * 10^{10} GeV \tag{27}$$

4.2 Discussion of mass value

The agreement of the mass $M = 9.4 * 10^{-17} kg$ needed to get the DAMA number observations with just one electron emission from each dark matter pearl and the number crudely estimated from the dimensional argument and the wish for the appropriate stopping length $7 * 10^{-9} kg$ is far from perfect. But we shall have in mind that the mass came in via a third root, so that a miscalculation of say ρ_B , the density of the compressed ordinary matter by just one order of magnitude would lead to a factor 1000 in the mass. This density ρ_B again depends on the third power of the Fermi energy E_f , so that indeed the mass we need to achieve the wanted inverse darkness $\frac{\sigma}{M} = 10^{-5} m^2 / kg$ will go as the ninth power, if there is a mistake in our E_f estimate. So our 8 orders of magnitude deviation in the mass M corresponds to E_f only being about one order of magnitude wrong.

But our model is tensioned in the direction that the density of the bubbles should go up and we should preferably get several counts out of the same dark matter pearl. It should sit in the NaI(Tl) and radiate for say several days. Then we could tolerate somewhat heavier pearls and still get the required number of counts.

So in reality we can consider our model to be successful, because the deviations were not more than about 8 orders of magnitude in the mass, corresponding to one order of magnitude in the Fermi energy.

5 Model

We now give a short review of our dark matter pearl model:

- Our "new physics": There are several different phases of the vacuum, but all with same energy density (*Multiple Point Criticality Principle* = *MPP* [19, 20, 21, 22, 23, 24, 25]). But even this "several vacuum phases" hypothesis is not truly new physics, if we believe the speculation that the quark masses have been adjusted so that two phases of the QCD-theory can be degenerate [13]: One phase with chiral symmetry spontaneously broken, and another one where the breaking should rather be said to be due to the quark masses.
- Our dark matter particles are macroscopic objects consisting of bubbles of a second vacuum filled with some ordinary matter e.g. carbon.
- There is a skin/wall separating the two vacuum phases with a tension of the order of

$$S^{1/3} \sim 10 \, MeV.$$
 (28)

• The nucleons have a lower potential in the inside-the-pearl-vacuum than in the outside vacuum, wherein we live, by a difference

$$\Delta V \approx 3 \, MeV. \tag{29}$$

The value $\Delta V \approx 3 \, MeV$ was fitted to the inside material having a gap - a homolumo gap - between the empty and the filled electronstates arranged to let the dark matter preferably emit X-rays with the observed energy of 3.5 keV.

- The dark matter pearls are so macroscopic and of such a size compared to mass that they get stopped in the Earth at a distance of the order of the depth of the DAMA-LIBRA experiment (1400 m stone).
- The pearls are excitable so as to radiate X-ray photons of just the energy needed to give the "mysterious 3.5 keV line".

5.1 The 3.5 keV line

The observation of a mysterious new X-ray emission line at 3.5 keV was reported [26, 27] in 2014. It was detected in the Andromeda galaxy, the Perseus cluster and different combinations of other galaxy clusters. Later the line was detected in the Milky Way Center [28]. This line has been suggested to originate from dark matter and our model has been adjusted to fit the observations. However this interpretation and even the very existence of the line is controversial. There are indeed several details in the observation of this 3.5 keV X-ray, which a priori does not look supportive for the hypothesis that this line really comes from dark matter: In fact Jeltema et al. [29] have seen it in the Kepler Supernova Remnant where there is far too little dark matter for them to be able to see it, unless dark matter interacts not only with dark matter but also with ordinary matter. Also the distribution seen from the Center of the Milky Way does not a priori look so much like dark matter produced X-rays. Furthermore there are some problems with the details of fitting the 3.5 keV radiation from the outskirts of the Perseus galaxy cluster. Assuming that the dark matter can be brought to radiate the X-rays by interaction not only with other dark matter particles, but also with ordinary matter, we believe we can improve the understanding of these mentioned three problems in fitting the 3.5 keV radiation observations with dark matter pearls.

We should point out that the observations of the 3.5 keV line are only seen as very small deviations of the spectra observed from what is understood from the various ions in the sources. They are on the borderline of being statistically significant (see Figure 9 for the Perseus Cluster and Figure 10 for Andromeda taken from from Iakubovskyi [30]):

We identify the emission of some particle or another as seen in DAMA-LIBRA and potentially in COSINE-100 etc. with the emission from the mysterious 3.5 keV line. So it becomes of course crucial that the energy seen in the events of the seasonal varying type in DAMA-LIBRA is indeed just 3.5 keV. In Figure 11 we show a fit (red) to the DAMA-LIBRA low-energy spectrum [31] consisting of a background model (grey/dashed) and a Gaussian distribution function (green). The parameters of the Gaussian are shown in the figure and the energy is given as 3.15 keV in remarkably good agreement!



Fig. 1: The combined MOS spectrum of Perseus cluster scaled to 3-4 keV energy range. On top of their best-fit model, the series of the single-bin residuals corresponding to the extra emission line at 3.57 keV is shown in red. (Adapted from Figure 7 in [70]).

Figure 9: Spectrum of X-rays from the galaxy cluster Perseus. The curve is fitted to the expected X-ray spectrum from known ions. The little deviation at 3.5 keV could be from dark matter?



Fig. 2: The same as in the previous Figure [] but for the combined spectrum of Andromeda galaxy. (Adapted from Figure 1 in 71).

Figure 10: X-ray spectrum from Andromeda.



Figure 11: Here the seasonally varying part of the DAMA-LIBRA data is fitted to a Gaussian with the average energy 3.15 keV. This is remarkably close to the line observed from Perseus Cluster and Andromeda and other galaxy clusters etc.

6 Distributions

Really to get an idea of how the radiation from excited dark matter pearls get distributed in the Earth, we must take into account that the distribution of their incoming direction is presumably very smoothly distributed over the sky. We shall now make a couple of somewhat special approximations and derive the distribution at the depth of the point under the surface of the Earth at which one detects the radiation.

We shall make 4 ideal calculations based on the following assumptions:

- What is counted is the emissions from the dark matter particles of some radiation (electrons and/or X-rays, actually with energy 3.5 keV), which is assumed to go on with a constant rate for a long time (we may think of weeks) after the impact of the pearl. This means that in order to obtain a distribution with respect to depth, we need to calculate the length of time during which a dark matter particle is in each little interval of depth.
- At first we assume that gravity is so weak that we can ignore it and assume that the pearls stop completely. Then in the second calculation we suppose that the pearls continue to fall directly downwards under gravity with a constant very low velocity. After the set in of this "terminal velocity" the pearls contribute the same amount of radiation in all depth layers below the point where the "terminal velocity" has set in. This means that we consider that there are two different stages in the motion of a dark matter particle:

- A fast stage from when the dark matter particle enters the Earth



Figure 12: We consider a model in which the incoming direction of the dark matter pearls has a constant distribution over the sky. At first we assume they stop completely after passing through a "stopping length" of earth and only then radiate significantly. Then we get a flat distribution of radiation delivery as a function of the depth down to a deepest depth equal to the "stopping length", beyond which there is no more probability for finding any radiation. This corresponds to the black rectangle on the figure. However if the pearls do not stop completely, but begin to fall with a constant velocity determined by a balance between the friction and the gravitational attraction by the Earth after reaching the "stopping length", then the distribution of the radiation delivered is given by the red curve as a function of the depth.

- through its surface until the particle has slowed down and we assume it moves so fast that it has no time to emit significant amounts of radiation. So during this stage we can neglect the radiation and the particle is effectively invisible.
- In the next stage we first consider the case where the dark matter particle stops and stays sitting in the Earth until its amount of excitation has burned out (after a long time, say a week, but taken at least as an average as a fixed constant). This case is illustrated by the *black* rectangle in Figure 12.
- Further, as a simplifying assumption, we assume that all the particles have the same stopping length (in the earth) and have the same incoming velocity. So all the dark matter particles hitting the Earth's surface at a certain point end up and stop (or in the second case considered below go into the next stage with a terminal velocity) at a point lying

on a half sphere around the point of impact with the Earth, having a radius equal to the stopping length.

- Again for simplicity, we assume that the distribution of the directions from which the dark matter particles come is evenly distributed over the sky. But it is of course in our model with a stopping length much shorter than the radius of the Earth, only the half sphere where the particles going downwards can come from that gets populated. Particles seeking to come upwards are stopped on the other side of the Earth.
- We also assume that the radiation (of electrons or X-rays of energy 3.5 keV) emitted during the first stage when the pearls pass quickly through the medium (essentially the earth, or the experimental apparatus scintillator) is negligible compared to the radiation emitted at the stopping point or, in the second case, when the dark matter sinks (before it runs out of excitation energy).

Let us now illustrate the rather trivial calculation of the radiation intensity observed:

First notice that if the dark matter particle comes in with a direction making an angle θ with the downward vertical, then the depth to which it has reached when the particle has stopped is

stopping depth = "stopping length"
$$*\cos(\theta)$$
 (30)

Now the area on the half sphere - which by the assumption of an even distribution on the sky is proportional to the fraction of the dark matter particles ending up there - corresponding to the depth z being in the infinitesimal interval dz is given as

$$dz = "stopping length" * d\cos\theta \tag{31}$$

$$= (-)"stopping length" * \sin \theta * d\theta \tag{32}$$

while

$$d``area'' = 2\pi * (``stopping length'')^2 * d\theta * \sin \theta.$$
(33)

so that

$$dz \propto d$$
"area" $\propto d$ "probability" up to $z =$ "stopping depth". (34)

This means that the radiation is present only down to the depth just equal to the stopping length, and in the interval with lower depth than that the radiation rate is quite constant. This is the black rectangular distribution given in Figure 12, which is namely the case in which the dark matter particles sit still on the surface of the half sphere and radiate out.



Figure 13: Here we take into account that the Earth's gravity acts on the dark matter pearl even while it is in the fast stage of its motion. Then even a pearl coming in almost horizontally will sink a bit before it stops or falls with its terminal velocity So in fact the black rectangle from figure 12 will essentially be lowered by a constant amount in depth. Also the distribution for the delivery of the radiation in the case of the pearl having a terminal velocity gets a corresponding lowering of the red curve from figure 12.

In the second case the dark matter particles almost stop and then start moving straight downwards with a constant terminal velocity, which is so slow that they contribute significant amounts of radiation at lower depths. Hence one gets the radiation rate for a depth z to be given by the integral of the rectangular distribution over all the higher depths:

$$"Rate"(z) \propto \int_0^z "rate"_{rectangular}(z')dz', \qquad (35)$$

It is easy to see that the resulting radiation rate "Rate''(z) has a non-zero slope in the uppermost stopping length, while it becomes constant under this domain.

Next we want to repeat the above two calculations including the effect of gravitation under the fast stage, which we have neglected so far. We perform this improvement very crudely assuming:

• We can ignore the variation as a function of the angle θ of the time the stopping in the fast stage takes. I.e. whatever the angle and thereby the depth in which the particle essentially stops it takes the same time, counted from the passage through the Earth's surface.

• Much like in an Einstein elevator the particle falls compared to the motion ignoring gravity by the amount that a free fall in the same time would have caused an object to fall.

With these assumptions the effect of introducing gravity on the curves of Figure 12 is simply to move them downward by the free fall distance. We here mean the free fall distance of a body falling for as long as the stopping time. The uppermost range of depth corresponding to this falling distance, becomes completely free of radiation in our approximations. So an experiment so close to the surface should see no dark matter at all.

These moved down curves are drawn in Figure 13.

If we could arrange say that ANAIS should lie so close to the surface as the falling distance during the stopping time then, under the rather simplified assumptions just presented, we could explain the lack of observation of dark matter.

7 Size

The dragging force stopping the dark matter pearl is

v

$$F_D = C_D \frac{Av^2 \rho_{material outside pearl}}{2}$$
(36)

where v is its velocity and $\rho_{material outside pearl}$ the density of the fluid or material through which the pearl falls. C_D is the drag coefficient and is of order unity at high speed. $A \approx \sigma$ is the area shown to the motion.

The equation of motion (Newton's second law) becomes:

$$M\dot{v} = -F_D = -\frac{C_D \sigma v^2 \rho_{material outside pearl}}{2}$$
 (37)

which can be rewritten and integrated to give the stopping length $L_{stopping}$ as follows:

$$\frac{\dot{v}}{v^2} = -\frac{\sigma}{M} * \frac{C_D}{2} \rho_{material outside pearl}$$
(38)

$$-\frac{1}{v} = -t * \frac{\sigma}{M} * \frac{C_D}{2} \rho_{material outside pearl} + const.$$
(39)

$$= \frac{1}{t * \frac{\sigma}{M} * \frac{C_D}{2}\rho_{material outside pearl} - const.}$$
(40)

$$L_{stopping} = \int v dt = \int_{v=300km/s}^{v \ small} \frac{dt}{t * \frac{\sigma}{M} * \frac{C_D}{2} \rho_{material \ outside \ pearl} - \dots} (41)$$
$$\approx -\left(\frac{\sigma}{M} * \frac{C_D}{2} \rho_{material \ outside \ pearl}\right)^{-1} \ln\left(\frac{"small"}{300km}\right)$$

Rewriting the estimate of the stopping length we have

$$L_{stopping} \approx \frac{1}{\frac{\sigma}{M} * \frac{C_D}{2} \rho_{material outside pearl} \ln(\frac{300 km/s}{"small"})}$$

where $M = \frac{1}{M} \ln(\frac{300 km/s}{small"})$ (42)

Now $L_{stopping}\sigma * \rho_{material outside pearl}$ is the amount of material pressed away by the passage of the pearl through the earth. So noticing that $\frac{2}{C_D} * \ln \frac{300 km/s}{"small"}$ is just of order unity, say 30, we see that the mass of the material pressed away by the the pearl is only an order of unity times bigger than the mass M of the pearl itself.

If the pearl essentially stops at the DAMA depth with 4200m w.e. then we have

$$4200m * 1000kg/m^3 = \frac{M}{\sigma} \frac{2}{C_D} * \ln \frac{300km/s}{"small"}$$
(43)

giving
$$\frac{\sigma}{M} = \frac{1}{4.2 * 10^6 kg/m^2 * 30}$$
 (44)

$$= 10^{-7} m^2 / kg. (45)$$

This should be compared with the inverse darkness obtained by Correa for velocities around 300 km/s from her analysis of dwarf galaxies [15]

$$\frac{\sigma}{M} = 1cm^2/g = 0.1m^2/kg.$$
 (46)

This means we need that so much dust washed off the pearls coming into the Earth that their cross section is diminished by a factor 1 million. So in area, they should have lost a factor 10^6 and in linear scale a factor 10^3 .

8 Physics of hoped for phase transition

By fitting to our dark matter model we found the order of magnitude of the tension S of the domain wall between the two phases should lie in the range

$$S = (\text{few } MeV)^3 \text{ to say } (30 MeV)^3 \tag{47}$$

This indicates that the physics involved in making this two vacua, if it is right, should be in an energy range which is *not at all new*.

The number $(30MeV)^3$ is what letting the domain walls replace the cosmological constant (the dark energy) would require [14].

We obtained the relationship between the mass M and surface tension S for our pearls mainly by adjusting the density of the strongly compressed material of ordinary matter inside the new type of vacuum to have a HO-MOLUMO gap suitable for emitting a 3.5 keV X-ray line. We further assumed that the pearls are not far from collapsing and spitting out the nuclei

inside, in spite of a potential keeping them in of $\Delta V \approx 3 MeV$. This relationship is illustrated here by giving examples of the mass M for a few values of the cubic root of the surface tension S.

$$S^{1/3} = 1 \, GeV \quad \Rightarrow \quad M = 24 \, ton \tag{48}$$

while
$$S^{1/3} = 100 \, MeV \Rightarrow M = 24 \, mg$$
 (49)

and
$$S^{1/3} = 10 \, MeV \Rightarrow M = 2.4 * 10^{-14} \, kg = 1.4 * 10^{13} \, GeV(50)$$

If we take the tension to be say $(10MeV)^3$, then we should look for physics at this scale to find out what could be used to cause the phase transition making the two vacua.

9 Conclusion

We have briefly reviewed our model for the dark matter being pieces of a new vacuum phase - which should though be understandable in terms of Standard Model physics, namely the Nambu JonaLasinio chiral symmetry breaking and QCD - filled under high pressure with ordinary matter.

But mainly we have searched for a way to resolve the seemingly very hard mystery that, while the DAMA-LIBRA experiment has collected more and more evidence for dark matter other experiments have not yet seen any. In particular the ANAIS and COSINE-100 experiments are very similar in that they also use NaI(Tl) scintillator but do not confirm the seasonal variation in their data which DAMA-LIBRA use to single out the dark matter. AMAIS and COSINE-100 claim a 3 σ disagreement with DAMA-LIBRA but COSINE-100 has less data. Even more, all the xenon scintillator based experiments having searched for dark matter have seen nothing.

The solution we have proposed to this disagreement is that dark matter is macroscopic and interacts more strongly than assumed in other models such as WIMP theories, but still sufficiently dark to fit observations. WIMPS are not significantly stopped by the earth shielding. However our dark matter pearls are stopped in a depth of order of that of the DAMA-LIBRA experiment 1400m, after passing through the Earth's surface with a high galactic velocity of the order of 200 km/s. In our model the dark matter can be excited and then emit electrons or photons with the preferred energy of 3.5 keV (corresponding to the controversial X-ray line associated with dark matter). So we can claim that the main effect seen by DAMA-LIBRA is this 3.5 keV radiation emitted by stopped or very much slowed down dark matter particles. But the dark matter particles pass through the experiments like ANAIS and COSINE-100 so fast that there is not sufficient time for them to give an observable signal. If really we take the dark matter to only be effectively observable when stopped and having time to emit its excitation energy, then the *fluid* xenon used by the majority of the underground experiments looking for dark matter cannot keep the dark matter stationary and thus, under such assumptions, have no chance to "see" the dark matter. At least if the Earth's gravitation is sufficient to drive the dark matter particles through the liquid xenon the liquid xenon experiments cannot observe them.

Very crucial for our speculation is that the signal DAMA-LIBRA sees is indeed 3.5 keV radiation emitted by stopped dark matter particles. It is a *remarkable fact* that the DAMA-LIBRA spectrum of the seasonally varying component is fitted by an essentially Gaussian distribution in energy with an average energy $3.15 \, keV$ close to the $3.5 \, keV$ line.

We remark that in our model we can even obtain a negative seasonal variation (i.e. with more events in winter than in summer). Interestingly COSINE-100 generated such an effect, which they consider as spurious, by using the DAMA-LIBRA background subtraction procedure on their data.

But if now dark matter is indeed, as we suggest, stopped in a depth under the Earth's surface of the order of the depth of DAMA, 1400 m, then one should - of course - seek to *dig dark matter out at this depth*. In water equivalent depth it would accidentally be very close to the bottom of the oceans, 5 km of water. The dark matter should be easily distinguished by having an abnormally high specific weight.

So finally we believe that the DAMA-LIBRA results, contrary to other experiments, point to the need for a macroscopic model of dark matter and that the mean free path in earth for galactic velocity dark matter must be of the order of 1400 m.

Acknowledgement

One of us H.B.N. acknowledges emeritus status at the Niels Bohr Institute, and support to the tour to Bled.

References

- [1] R. Bernabei et al, Prog. Part. Nucl. Phys. **114**, 103810 (2020).
- [2] C. D. Froggatt and H. B. Nielsen, Phys. Rev. Lett. 95 231301 (2005) [arXiv:astro-ph/0508513].
- C.D. Froggatt and H.B. Nielsen, Proceedings of Conference: C05-07-19.3 (Bled 2005); arXiv:astro-ph/0512454.
- [4] C. D. Froggatt and H. B. Nielsen, Int. J. Mod. Phys. A 30 no.13, 1550066 (2015) [arXiv:1403.7177].
- [5] C. D. Froggatt and H. B. Nielsen, Mod. Phys. Lett. A30 no.36, 1550195 (2015) [arXiv:1503.01089].

- [6] H.B. Nielsen, C.D. Froggatt and D. Jurman, PoS(CORFU2017)075.
- [7] H.B. Nielsen and C.D. Froggatt, PoS(CORFU2019)049.
- [8] C. D. Froggatt, H. B. Nielsen, "The 3.5 keV line from non-perturbative Standard Model dark matter balls", arXiv:2003.05018.
- [9] H. B. Nielsen (speaker) and C.D. Froggatt, "Dark Matter Macroscopic Pearls, 3.5 keV -ray Line, How Big?", 23rd Bled Workshop on What comes beyond the Standard Models (2020), arXiv:2012.00445.
- [10] C. D. Froggatt and H.B.Nielsen, "Atomic Size Dark Matter Pearls, Electron Signal", 24th Bled Workshop on What comes beyond the Standard Models (2021), arXiv:2111.10879.
- [11] C. D. Froggatt and H.B. Nielsen, "Atomic Size Pearls being Dark Matter giving Electron Signal", arXiv:2203.02779.
- [12] H. B. Nielsen (speaker) and C.D. Froggatt, "Dusty Dark Matter Pearls Developed", 24th Bled Workshop on What comes beyond the Standard Models (2022), arXiv:2303.06061.
- [13] C.D. Froggatt and H.B. Nielsen PoS(CORFU2022)003, arXiv:2305.12291
- [14] C.D. Froggatt and H.B. Nielsen PoS(CORFU2022)205, arXiv:2305.18645
- [15] C. A. Correa, MNRAS **503** 920 (2021) [arXiv:2007.02958].
- [16] J. Amaré et al, Phys. Rev. **D103**, 102005 (2021) [arXiv:2103.01175]
- [17] G. Adhikari et al. Phys. Rev. **D106**, 052005 (2022)
- [18] Adhikari, G. et al., Sci. Rep. **13**, 4676 (2023).
- [19] D. L. Bennett, C. D. Froggatt and H. B. Nielsen, NBI-HE-94-44, GUTPA-94-09-3, Presented at Conference: C94-07-20 (ICHEP 1994), p.0557-560.
- [20] D. L. Bennett, C. D. Froggatt and H. B. Nielsen, NBI-95-15, GUTPA-95-04-1, Presented at Conference: C94-09-13 (Adriatic Meeting 1994), p.0255-279 [arXiv:hep-ph/9504294].
- [21] D. L. Bennett and H. B. Nielsen, Int. J. Mod. Phys. A9 5155 (1994).
- [22] D. L. Bennett, C. D. Froggatt and H. B. Nielsen, NBI-HE-95-07, Presented at Conference: C94-08-30 (Wendisch-Rietz) p.394-412.

- [23] C. D. Froggatt and H. B. Nielsen, Phys. Lett. B368 96 (1996) [arXiv:hep-ph/9511371].
- [24] H.B. Nielsen (Speaker) and C.D. Froggatt, Presented at Conference: C95-09-03.1 (Corfu 1995); arXiv:hep-ph/9607375.
- [25] Gia Dvali, arXiv:1107.0956.
- [26] E. Bulbul, M. Markevitch, A. Foster et al., ApJ. 789, 13 (2014) [arXiv:1402.2301];
 E. Bulbul, M. Markevitch, A. Foster et al., ApJ. 831, 55 (2016) [arXiv:1605.02034].
- [27] A. Boyarsky, O. Ruchayskiy, D. Iakubovskyi and J. Franse, Phys. Rev. Lett. 113, 251301 (2014) [arXiv:1402.4119].
- [28] A. Boyarsky, J. Franse, D. Iakubovskyi and O. Ruchayskiy, Phys. Rev. Lett. 115, 161301 (2015) [arXiv:1408.2503].
- [29] T. Jeltema and S. Profumo, MNRAS **450**, 2143 (2015) [arXiv:1408.1699].
- [30] D.A. Iakubovskyi, Advances in Astronomy and Space Physics 6, 3-15 (2016) [arXiv:1510.00358]
- [31] Z. Ahmed et al, Phys. Rev. D81, 042002 (2010) [arXiv:0907.1438]