# SYMMETRIES OF QUANTIZATION OF FINITE GROUPOIDS

JACKSON VAN DYKE

ABSTRACT. This paper proves certain facts concerning the equivariance of quantization of  $\pi$ -finite spaces. We argue that these facts establish an analogy between this quantization procedure and the geometric quantization of a symplectic vector space.

Specifically, we observe that symmetries of a given polarization/Lagrangian always induce coherent symmetries of the quantization. On the other hand, symmetries of the entire phase space a priori only induce projective symmetries. For certain three-dimensional theories, this projectivity appears via a twice-categorified analogue of Blattner-Kostant-Sternberg kernels in geometric quantization and the associated integral transforms.

# CONTENTS

1. Introduction	2
1.1. Automorphisms of the phase space	3
1.2. Automorphisms of the base	4
1.3. Outline	5
1.4. Acknowledgements	6
2. Postnikov obstructions	6
2.1. Preliminaries	6
2.2. Higher groupoids attached to a braided fusion category	7
2.3. Pointed braided fusion categories	8
2.4. Obstruction theory	10
3. Topological symmetry	13
3.1. TQFTs associated to $\pi$ -finite spaces	13
3.2. Module structures	16
4. Semiclassical symmetry	17
4.1. Semiclassical group actions	17
4.2. Spectral sequence obstruction	18
5. Symmetries of the base	20
5.1. Action via bimodules up to isomorphism	21
5.2. System of products	21
5.3. Associators	23
5.4. Semiclassical action	25
6. Orthogonal symmetries	25
6.1. Analogue of BKS kernels	25
6.2. Analogue of the intertwining operators	27
Appendix A. Bicharacters and the Schur multiplier	28
Appendix B. TQFTs and the Cobordism Hypothesis	29
B.1. The Cobordism Hypothesis for X-theories	29
B.2. Relative theories	30
References	31

Date: December 4, 2023.

#### 1. INTRODUCTION

Geometric quantization relates (semi)classical states to quantum ones. Namely, the geometric quantization of a polarized symplectic vector space V consists of some sort of "functions on half of the variables". I.e. if we have a polarization  $V \simeq T^*\ell$ , one model for the Hilbert space of states is  $L^2(\ell)$ . This space of states has the curious property that, even though it is most naturally thought of as associated to the symplectic vector space V, it is only *projectively* equivariant for the natural symplectic group of symmetries.

The traditional way of identifying this projectivity is by constructing a natural integral transform  $T_g$  associated to each symplectic group element g, using what are called Blattner-Kostant-Sternberg (BKS) kernels [LV80, Pol03, BW97]. E.g. when g is the symplectic matrix sending the position coordinates in V to the momentum coordinates, the associated integral transform is the *Fourier transform*. The integral transforms don't quite compose coherently:

$$c(g,h) = T_{gh}^{-1} \circ gT_h \circ T_g$$

for some scalar  $c(g,h) \in U(1)$ . This defines a 2-cocycle, and the extension classified by  $[c] \in H^2(B \operatorname{Sp}(T^*\ell))$  is essentially the metaplectic group.

In summary: Geometric quantization is only *projectively* a functor out of the appropriate category of phase spaces. On the other hand, it is manifestly (linearly) functorial out of *polarized* phase spaces.

We study an analogue of this projective (resp. linear) equivariance with respect to automorphisms of the full phase space (resp. of the Lagrangian) in the setting of quantization of  $\pi$ -finite groupoids in dimension three. Quantization of  $\pi$ -finite groupoids was introduced in [Kon88] and studied further in [Qui95, Tur10]. We will consider the fully local case introduced in [Fre94] and studied in [FHLT10, §3,8] and [FMT22, §A]. We write the quantization of a  $\pi$ -finite space, possible decorated by a cocycle  $\tau$  representing a class in  $H^d(X)$ , as  $\sigma^d_{X,\tau}$ .

Under the analogy considered herein, the analogue of the Lagrangian  $\ell$  is a finite group L, and the analogue of the symplectic vector space  $V = T^*\ell$  is  $A = L \oplus L^{\vee}$ . I.e. we are considering a finite higher analogue of quantization, wherein vector spaces are replaced by (classifying spaces of) abelian groups. The symplectic form is replaced by a quadratic<sup>1</sup> form on A given by evaluation. The analogue of the geometric quantization itself is  $\sigma_{BL}^3$ .

In any dimension, several different avatars of a "projective action" on a TQFT are shown to be equivalent in [VD23, §C.4]. Restricting to dimension three, it is shown in [VD23, Theorem 3.5, 3.12] that orthogonal automorphisms of A a priori only induce projective automorphisms of the theory  $\sigma_{BL}^3$ . Equivalently, certain 3-group extensions (defined in [VD23, §2.5]) of O (A, ev) act linearly on  $\sigma_{BL}^3$ . This is analogous to the linear action of the extension Spin of SO on spinors, as is explained in [VD23, §2.7].

In Section 6, we identify a certain object which plays the role of the BKS kernels in this context. In particular, this object will be a class in  $H^2$  of A, which defines a morphism between any two polarizations in a sense which is made precise in Sections 3.1 and 6.1.

On the other hand, we show that automorphisms of (possibly nonabelian) L (preserving a twist  $\tau$ ) induce automorphisms of the theory  $\sigma_{BL,\tau}^3$ . A certain notion of projectivity can appear, however it necessarily appears already semiclassically.

Now we provide a more detailed summary of the main results of this paper.

<sup>&</sup>lt;sup>1</sup>There is a discrepancy between the two sides of the analogy: The symplectic structure is replaced by an orthogonal one. This is a version of the well-known analogy between between the Weil representation and spinors [Del99].

1.1. Automorphisms of the phase space. Let  $V \simeq L \oplus L^{\vee}$  be a polarized vector space over a finite field  $\mathbb{F}_p$ . In Section 6.1, we define a certain "higher integral transform" associated to a reflection in  $O(L \oplus L^{\vee}, ev)$ . This is the analogue of the BKS kernels in geometric quantization.

Namely, we identify a correspondence from L to its image under an orthogonal reflection g, i.e. this is a semiclassical defect in the sense of [FMT22]. The **Vect** [L]-**Vect** [gL] bimodule category, i.e. higher integral transform, is the quantization of this semiclassical defect, in the sense of [FMT22].

We have a canonical splitting

$$s: O(A,q) \to \pi_{\leq 1} \operatorname{Aut}_{\mathbf{EqBr}} (\mathcal{Z})$$

(see Section 2.4.2). In [ENO10, §5.2] an equivalence

$$\Psi \colon \operatorname{Aut}_{\mathbf{EqBr}}(\mathcal{Z}) \xrightarrow{\sim} \pi_{<1} \operatorname{Aut}_{\mathbf{Fus}}(\mathcal{C})$$

is provided. The following is stated as Theorem 6.8 in the body of this paper.

**Theorem.** Assume Hypothesis Q. Let  $A \simeq L \oplus L^{\vee}$  be a vector space over a finite field. Sending a reflection  $g \in O(A, q)$  to the intertwining bimodule f(g) from (6.7) defines the same equivalence as the composition:

$$O(A,q) \xrightarrow{s} \pi_0 \operatorname{Aut}_{\mathbf{EqBr}} (\mathcal{Z}) \xrightarrow{\Psi} \pi_0 \operatorname{Aut}_{\mathbf{Fus}} (\mathcal{C})$$

*Warning.* The theory of quantizing finite groupoids (and the higher correspondences/defects between them) has not received a full treatment in the literature, however this is the subject of a work in progress of Claudia Scheimbauer and Tashi Walde. See Hypothesis Q for the formal features of this construction which we assume in this work. (This is the same as [VD23, Hypothesis Q].)

Explicitly, this construction defines a map from O(A, ev) to the group of isomorphism classes of **Vect** [L]-bimodules:

$$f: O(A, ev) \to \pi_0 \operatorname{Aut}_{\mathbf{Fus}} (\mathbf{Vect}[L])$$
.

The projectivity/anomaly is the obstruction to lifting this to a functor:

$$\mathbf{f} \colon \mathcal{O}(A, \mathrm{ev}) \to \mathrm{Aut}_{\mathbf{Fus}} \left( \mathbf{Vect}\left[L\right] \right)$$

where O(A) now denotes the discrete 2-category with objects O(A).

Obtaining such a map **f** is one avatar of a group acting on a theory: Vect [L] classifies the theory  $\sigma_{BL}^3$  by the Cobordism Hypothesis.

This obstruction turns out to vanish when V is a vector space over  $\mathbb{F}_p$  by [EG18, Theorem 6.1]. Therefore one could explicitly write down trivializations of the obstructions to lifting f to **f**, as we do in Section 5 for automorphisms of the base, however we do not pursue this.

The fact that A is a vector space over  $\mathbb{F}_p$  is only used in the proof of Theorem 6.8 when we invoke the fact that O(A,q) is generated by reflections. The same argument would work for any orthogonal transformation g such that gL is transverse to L. In these more general cases, the obstruction may not vanish.

If the top obstruction does not vanish, it may be interpreted as a nontrivial anomaly as explained in [VD23]. Namely, there are multiple different avatars of a "projective action" of a group on a TQFT, which are shown to be equivalent in [VD23, Theorem C.16, C.18].

Remark 1.1. Recall isomorphism classes of indecomposable bimodule categories over Vect [L], for L a finite group, are classified by subgroups  $H \subset L \times L$  equipped with a cocycle  $\psi \in Z^2(H, \mathbf{k}^{\times})$ .

Under the interpretation of these bimodule categories as twice-categorified higher integral transforms, this classification is analogous to the Schwarz Kernel Theorem. Versions of the once-categorified version of this type of statement can be found in [Toë07, BZFN10].

1.2. Automorphisms of the base. Now we will consider an action of a finite group G on L preserving the class of  $\tau$ , which will turn out to define a canonical action of G on  $\sigma^d_{BL,\tau}$ . But first we will make some general considerations regarding actions of groups on semiclassical data.

1.2.1. Semiclassical action. Following ideas in [FMT22], we define a semiclassical action (Definition 4.1) of a finite group G on semiclassical data  $(X, \tau)$  (where  $\tau$  represents a class in  $H^d(X, \mathbf{k}^{\times})$ ) to be a correspondence (i.e. 1-morphism) between the trivial  $\pi$ -finite space to BG such that when we pull back along  $* \to BG$  we obtain  $(X, \tau)$ :



I.e. this consists of an X-bundle over BG and an extension of  $\tau$  to the total space  $C_G$ .

The semiclassical action is *projective* if BG is replaced by (BG, c) for some (d + 1)-cocycle on BG, and a trivialization defines a correspondence (1-morphism) from (BG, c) to BG (Definition 4.4). Composing the projective action with this correspondence "linearizes" the projective action to an ordinary one.

Assuming we have a semiclassical action of G on X, we would like to identify the obstruction to extending this to a semiclassical action of G on  $(X, \tau)$ . I.e. we have an X-bundle over BGwith total space  $C_G$ , and we seek the obstruction to extending  $\tau$  from X to  $C_G$ .

The Serre spectral sequence provides a general method for answering this question. In Theorem 4.6, under some assumptions, we show that the spectral sequence obstruction encodes an anomaly/anomalous symmetry of the theory  $\sigma_{X,\tau}^d$ . Theorem 4.6 generalizes the 't Hooft anomalies studied in [Mü20, MS20, KT14] to the fully-extended setting.

In Corollary 4.9, we note that the anomaly/projectivity agrees with (a restriction of) the top k-invariant of the higher automorphism groupoid. Corollary 4.9 generalizes part of the arguments made in [ENO10, §A] to arbitrary dimensions.

1.2.2. Three dimensions. Now we specialize to three dimensions and consider the particular space X = BL for L a finite group with  $\tau$  a group cocycle representing a class in  $H^3(BL, \mathbf{k}^{\times}) \simeq H^3_{\text{Grn}}(L, \mathbf{k}^{\times})$ . Let G be a finite group, and consider a group homomorphism

$$\varphi \colon G \to \operatorname{Aut}(L,\tau)$$

where Aut  $(L, \tau)$  denotes the group of group automorphisms of L which preserve the cohomology class of  $\tau$ .

The fully-extended TQFT

$$\sigma_{L,\tau} \colon \mathbf{Bord}_3^{\mathrm{fr}} \to \mathbf{Fus}$$

(see Section 3) sends the point to

$$\mathcal{C} = \mathbf{Vect} \left[ L \right]^{\tau} ,$$

the fusion category of vector spaces graded by L, with convolution product twisted by  $\tau$ .

Therefore, one avatar of a group acting on a category might be a map:

$$\mathbf{f}: G \to \operatorname{Aut}_{\mathbf{Fus}}(\mathcal{C})$$

In Section 5, we define such a map by explicitly defining the map on the underlying ordinary groups, and trivializing the obstructions in [ENO10]. Any functor

$$BG \to B\operatorname{Aut}_{\mathbf{Fus}}\left(\sigma_{BL,\tau}^{3}\left(*\right)\right)$$

is equivalent to a  $(\sigma_{BG}^4, \rho)$ -module structure on  $\sigma_{BL,\tau}^3$  by Hypothesis Q. This shows the following, which is stated as Theorem 5.15 in the body of this paper.

**Theorem.** Assume Hypothesis Q. Let L and G be finite groups. Let  $\tau \in Z^3_{Grp}(L, \mathbf{k}^{\times})$  be a 3-cocycle in the group cohomology of L. Then any group homomorphism

$$\varphi \colon G \to \operatorname{Aut}(L,\tau)$$

canonically defines a  $(\sigma_{BG}^4, \rho)$ -module structure on  $\sigma_{BL,\tau}^3$ .

This canonical module structure can be twisted by a triple  $(\gamma, \mu, \alpha)$  where

$$\gamma \in C^{1}_{Grp}\left(G, C^{2}_{Grp}\left(L, \mathbf{k}^{\times}\right)\right) \qquad \mu \in Z^{2}_{Grp}\left(G, C^{1}_{Grp}\left(L, \mathbf{k}^{\times}\right)\right) \qquad \alpha \in Z^{3}\left(G, \mathbf{k}^{\times}\right) \ .$$

satisfy:

$$d_L(\gamma(g)) = \varphi(g)(\tau)\tau^{-1} \qquad \qquad d_L\mu = d_G\gamma^{-1}$$

As is shown in [ENO10, §11], the existence of such a functor

$$(f, m, a) : BG \to B \operatorname{Aut}_{\mathbf{Fus}} \left( \sigma^3_{BL, \tau} \left( * \right) \right)$$

implies the existence of an extension of  $\tau$  to a group 3-cocycle  $\tau_G$  on  $L \rtimes_{\varphi} G$ . The quantization of the correspondence



defines a  $(\sigma_{BG}^4, \rho)$ -module structure on  $\sigma_{BL,\tau}^3$ . This module structure equivalently defines a functor

$$BG \to B\operatorname{Aut}_{\mathbf{Fus}}(\operatorname{\mathbf{Vect}}[L]^{\tau})$$
,

which provides an alternative description of the map (f, m, a). This is discussed in more detail in Section 5.4.

1.3. **Outline.** We give an overview of the obstruction theory in [ENO10] in Section 2. Preliminaries are developed in Sections 2.1 to 2.3, and the obstruction classes are discussed in Section 2.4. Section 3 is an overview of topological symmetry, mostly following [FHLT10, FMT22]. In Section 3.1 we discuss quantization of finite groupoids, and introduce Hypothesis Q. In Section 3.2 we review the notion of a module structure on a QFT from [FMT22].

Section 4 consists of a discussion of semiclassical symmetries. Namely, we define linear and projective semiclassical actions in Section 4.1, and discuss the anomaly/obstruction to extending a semiclassical action on a  $\pi$ -finite space X to a semiclassical action on the decorated space  $(X, \tau)$  for some cocycle  $\tau$ .

In Section 5, we show that automorphisms of a finite group L preserving a 3-cocycle  $\tau$  act canonically on the theory  $\sigma_{BL,\tau}^3$ . In Section 5.1 we define the action up to isomorphism f, in Section 5.2 we define a system of products m, and in Section 5.3 we define an associator. In Section 2.4.2 we discuss the obstruction  $O_3(f)$ , and in Section 2.4.3 we discuss the obstruction  $O_4(f,m)$ . In Section 5.3.2 we state and prove Theorem 5.15. The connection with semiclassical actions is discussed in Section 5.4.

In Section 6 we instead consider automorphisms of the full group  $L \oplus L^{\vee}$  which preserve the quadratic form ev. In Section 6.1 we define an analogue of the BKS kernel. The quantization of this is considered in Section 6.2, where this is also shown to agree with the equivalence in [ENO10] in Theorem 6.8.

Appendix A is dedicated to a discussion of some technical points regarding the Schur isomorphism, which are used in Section 6.

Appendix B establishes some general facts concerning fully-extended TQFTs. The particular flavor of the Cobordism Hypothesis which is most useful for this paper is stated in appendix B.1. Some conventions are fixed regarding relative/boundary theories in Appendix B.2.

1.4. Acknowledgements. I warmly thank my advisor David Ben-Zvi for his constant help and guidance, without which this project would not exist. I would also like to thank Dan Freed, David Jordan, Lukas Müller, German Stefanich, Rok Gregoric, and Will Stewart for helpful discussions.

Part of this project was completed while the author was supported by the Simons Collaboration on Global Categorical Symmetries.

This research was also partially completed while visiting the Perimeter Institute for Theoretical Physics. Research at Perimeter is supported by the Government of Canada through the Department of Innovation, Science, and Economic Development, and by the Province of Ontario through the Ministry of Colleges and Universities.

# 2. Postnikov obstructions

2.1. **Preliminaries.** Unless otherwise specified, all categories are k-linear, where k is algebraically closed of characteristic zero. Throughout the paper, we will write **Vect** for the k-linear category of finite-dimensional vector spaces over k. Let **Pr** be the symmetric monoidal 2-category of locally presentable categories with colimit preserving functors (and natural transformations). We will now sketch some definitions of the targets for our 3 and 4-dimensional TQFTs. See [BJS21, §3] for a more detailed discussion of the Morita theory of (braided) tensor categories, and in particular the relationship with [Sch14, Hau17].

The Morita 3-category of tensor categories,  $\mathbf{Tens} = \mathbf{Alg}_1(\mathbf{Pr})$ , consists of the following:

- the objects are tensor categories,
- the 1-morphisms are bimodule categories,
- the 2-morphisms are functors between bimodule categories, and
- the 3-morphisms are natural transformations.

We will focus on the subcategory **Fus** consisting of fusion categories, semisimple bimodule categories, compact-preserving cocontinuous bimodule functors, and natural transformations. It was shown in [BJS21] that this forms a subcategory.

Similarly, braided tensor categories form a 4-category  $\mathbf{BrTens} = \mathbf{Alg}_2(\mathbf{Pr})$ :

- the objects are braided tensor categories,
- the 1-morphisms are associative algebra objects in the category of bimodule categories,
- the 2-morphisms are bimodule categories between bimodule categories,
- the 3-morphisms are functors between bimodule categories, and
- the 4-morphisms are natural transformations.

The subcategory **BrFus** consists of braided fusion categories, fusion categories equipped with central structures, finite semisimple bimodule categories, compact-preserving cocontinuous bimodule functors, and bimodule natural transformations. It was shown in [BJS21] that this forms a subcategory.

Braided fusion categories also form a 2-category:

- the objects are braided fusion categories,
- the 1-morphisms are functors preserving the braided structure, and
- the 2-morphisms are natural transformations.

The 2-groupoid given by the invertible part of this 2-category is written **EqBr**, as in [ENO10].

2.2. Higher groupoids attached to a braided fusion category. Let  $\mathcal{A}$  be a braided fusion category. The classifying space of the 2-group of braided equivalences forms a 2-type  $B \operatorname{Aut}_{\mathbf{EqBr}}(\mathcal{A})$ .

To  $\mathcal{A}$  we can also attach the 2-category of  $\mathcal{A}$ -module categories. The braiding defines an embedding from the 2-category of  $\mathcal{A}$ -modules to the 2-category of  $\mathcal{A}$ -bimodules. Therefore the category of  $\mathcal{A}$ -modules inherits a monoidal structure from the natural one on the category of  $\mathcal{A}$ -bimodules.

**Definition.** The invertible  $\mathcal{A}$ -modules, written Pic ( $\mathcal{A}$ ), comprise the *Picard* 3-group of  $\mathcal{A}$ .

From any 3-group, we can construct its classifying space  $B \operatorname{Pic}(\mathcal{A})$ , which is a 3-type. I.e. it has three nontrivial homotopy groups.

*Remark* 2.1. Some authors write the full 3-group/type as  $B\underline{\text{Pic}}(\mathcal{A})$ , and write the truncation to a 2-type as  $B\underline{\text{Pic}}(\mathcal{A})$  and to a 1-type as  $B\underline{\text{Pic}}(\mathcal{A})$ .

Instead, we write  $B \operatorname{Pic}(\mathcal{A})$  for the full 3-group, and write  $\tau_{\leq 2} B \operatorname{Pic}(\mathcal{A})$  and  $\tau_{\leq 1} B \operatorname{Pic}(\mathcal{A})$  for the truncations.

**Theorem** ([ENO10]). For a nondegenerate braided fusion category  $\mathcal{A}$  we have an equivalence:

(2.2) 
$$\pi_{<2}B\operatorname{Pic}(\mathcal{A}) \simeq B\operatorname{Aut}_{\mathbf{EqBr}}(\mathcal{A})$$

The homotopy groups of  $B \operatorname{Pic}(\mathcal{A})$ , e.g. from [ENO10, Proposition 7.5] are as follows:  $\pi_1$  is given by the ordinary group of isomorphism classes of braided equivalences, and  $\pi_2$  is given by the group of tensor isomorphisms of the identity functor on  $\mathcal{A}$ . The top homotopy group, which is not involved in (2.2), is  $\pi_3 B \operatorname{Pic}(\mathcal{A}) = \mathbf{k}^{\times}$ .

2.2.1. Braided fusion categories which are Drinfeld centers. Let C be a fusion category. To this we can attach the monoidal 2-category  $\operatorname{End}_{\operatorname{Fus}}(C)$ .

**Definition.** The *Brauer-Picard* 3-group of  $\mathcal{C}$  is  $\operatorname{Aut}_{\mathbf{Fus}}(\mathcal{C})$ .

Recall the Drinfeld center of a monoidal category C is a braided monoidal category, written  $\mathcal{Z}(C)$ .

**Theorem** ([EO04]). *There is an equivalence of* 2*-categories:* 

$$\operatorname{End}_{\mathbf{Fus}}\left(\mathcal{C}\right)\xrightarrow{\sim}\mathcal{Z}\left(\mathcal{C}\right)$$
-mod.

Passing to the invertible part, we obtain:

(2.3) 
$$\operatorname{Aut}_{\mathbf{Fus}}(\mathcal{C}) \simeq \operatorname{Pic}\left(\mathcal{Z}\left(\mathcal{C}\right)\right)$$

Remark 2.4. In [ENO11], it is shown that two fusion categories are Morita equivalent if and only if their Drinfeld centers are braided equivalent. This was strengthened in [ENO10]: they prove that there is a fully-faithful embedding of groupoids  $\mathbf{EqBr} \to \mathbf{Fus}^{\times}$ . Indeed, combining (2.2) and (2.3) we obtain

(2.5) 
$$\pi_{\leq 2} B \operatorname{Aut}_{\mathbf{Fus}}(\mathcal{C}) \simeq B \operatorname{Aut}_{\mathbf{EqBr}}(\mathcal{Z}(\mathcal{C})) .$$

So the center determines the Morita class of the fusion category itself. However, since the equivalence (2.2) is with the truncated part of  $\operatorname{Aut}_{\mathbf{Fus}}(\mathcal{C})$ , there is a subtle lack of "coherence", captured by an obstruction class originally studied in [ENO10] and surveyed in Section 2.4.3.

Related results were shown in [KZ18] and [KK12, §5].

In summary: given a braided monoidal category  $\mathcal{A}$ , we have a 3-type  $B \operatorname{Pic}(\mathcal{A})$  attached to it. If it happens to be the case that  $\mathcal{A} = \mathcal{Z}(\mathcal{C})$  for some fusion category  $\mathcal{C}$ , then this 3-type is the classifying space of the Brauer-Picard 3-group of  $\mathcal{C}$ . All together we have:



2.3. **Pointed braided fusion categories.** A fusion category is *pointed* if all simple objects are invertible.

*Example* 2.6. Let L be a finite group. Consider the category of L-graded vector spaces, **Vect** [L], with convolution, i.e. for  $a, b, c \in L$ :

$$(W * W')_a = \bigoplus_{a=bc} W_b \otimes W_c$$
.

This category has simple objects  $\mathbf{k}_a$  for  $a \in L$ , and is therefore pointed. Given a 3-cocycle  $\tau$  on G, we can define a variant of this category **Vect**  $[L]^{\tau}$ , which is still pointed.

As it turns out, all pointed fusion categories are of the form  $\operatorname{Vect} [L]^{\tau}$ . (Note however that the twisted version may have no fiber functor. This is a higher analogue of the fact that not all algebras have an augmentation map.)

2.3.1. Metric groups. Metric groups [DGNO10, ENO10] play an important role in the theory of pointed braided fusion categories. Let  $\mathbf{k}$  be an algebraically closed field of characteristic zero, and let A be a finite abelian group. Write  $A^{\vee} := \text{Hom}(A, \mathbf{k}^{\times})$  for the character dual of A.

**Definition.** A biadditive map

$$\langle \cdot, \cdot \rangle : A \times A \to \mathbf{k}^{\times}$$

is a symmetric bicharacter if for all  $a, b \in A$  we have  $\langle a, b \rangle = \langle b, a \rangle$ . By acting on the first argument, such a bicharacter  $\langle \cdot, \cdot \rangle$  defines a homomorphism  $A \to A^{\vee}$ . If this is an isomorphism, then we say that  $\langle \cdot, \cdot \rangle$  is nondegenerate.

**Definition.** A function

$$q: A \to \mathbf{k}^{\times}$$

is a quadratic form on A if

- (a) for all  $a \in A$  we have q(a) = q(-a), and
- (b) the following expression defines a symmetric bicharacter:

$$\langle a,b\rangle_q \coloneqq \frac{q\,(a+b)}{q\,(a)\,q\,(b)}$$

We say q is nondegenerate if the bicharacter  $\langle\cdot,\cdot\rangle_q$  is nondegenerate.

*Remark* 2.7. If |A| is odd, then sending q to  $\langle \cdot, \cdot \rangle_q$  defines a bijection between quadratic forms and bicharacters.

**Definition.** Any pair (A, q) is a *pre-metric group*. A pair (A, q) is called a *metric group* if q is nondegenerate.

**Definition.** The orthogonal group of a metric group (A, q) is:

$$\mathcal{O}(A,q) \coloneqq \{f \in \operatorname{Hom}(A,A) \,|\, q \circ f = q\} \;\;.$$

**Definition.** Let (A, q) be a metric group. Define the *determinant* 

det: 
$$O(A,q) \to \mathbb{Q}^{\times}_{>0} / \left( \mathbb{Q}^{\times 2}_{>0} \right)$$

by sending  $g \in O(A, q)$  to the image of  $|(g-1)A| \in \mathbb{N}$ .

The special orthogonal group is

(2.8) 
$$\operatorname{SO}(A,q) \coloneqq \ker(\det)$$

The upshot of upgrading an ordinary group to a metric group (A, q) is that the category of A-graded vector spaces acquires a braiding. We can specify this braiding on simple objects:

(2.9) 
$$\mathbf{k}_a \ast \mathbf{k}_b = \mathbf{k}_{ab} \xrightarrow{\langle a,b \rangle \cdot \mathrm{id}_{\mathbf{k}_{ab}}} \mathbf{k}_{ab} = \mathbf{k}_b \ast \mathbf{k}_a \; .$$

It is shown in [JS93] that braided fusion categories are classified by finite abelian groups A (the group of simple objects) equipped with a quadratic form  $q: A \to \mathbf{k}^{\times}$  which is not necessarily nondegenerate. See also [EGNO15, §8.4].

*Example* 2.10. If L is any finite abelian group, then the finite abelian group  $L \oplus L^{\vee}$  has a nondegenerate quadratic form given by evaluation, so  $(L \oplus L^{\vee}, ev)$  is a metric group. The symmetric bicharacter corresponding to evaluation is:

$$\langle (\ell, \chi), (\ell', \chi') \rangle = \chi (\ell') \chi' (\ell)$$
.

We obtain a braiding on **Vect**  $[L \oplus L^{\vee}]$  via (2.9).

*Example* 2.11. Let  $C = \text{Vect}[L]^{\tau}$  for L a vector space over  $\mathbb{F}_p$  and  $\tau \in H^3(L, \mathbf{k}^{\times})$ . Then, by [MN18, Corollary 5.3],  $\mathcal{Z}(C)$  is pointed if and only if  $\tau$  has trivial alternating component.<sup>2</sup>

2.3.2. Lagrangian subgroups.

**Definition.** A subgroup  $L \subset A$  of a metric group (A, q) is *isotropic* if  $q(\ell) = 1$  for all  $\ell \in L$ . The subgroup is Lagrangian if  $|L|^2 = |A|$ .

*Example* 2.12. If  $A = L \oplus L^{\vee}$  and q = ev, then  $L \oplus \{0\}$  and  $\{0\} \oplus L^{\vee}$  are both Lagrangian subgroups.

Let L be a finite abelian group. The Drinfeld center of the fusion category (Vect [L], \*) is:

$$\mathcal{Z}\left(\mathbf{Vect}\left[L\right],*
ight)\cong\mathbf{Vect}\left[L\oplus L^{\vee}
ight]$$

The tensor structure is convolution, and the braiding is induced by evaluation as in Example 2.10. So if we started with the metric group  $(L \oplus L^{\vee}, ev)$ , and turn it into a braided fusion category  $\mathcal{A}$ , we see that Lagrangians in the metric group give rise to fusion categories  $\mathcal{C}$  such that  $\mathcal{A} \cong \mathcal{Z}(\mathcal{C})$ .

<sup>&</sup>lt;sup>2</sup>Recall  $H^3(L, \mathbf{k}^{\times})$  decomposes as a direct sum of the alternating factor  $\wedge^3 L^{\vee}$  and the symmetric factor  $\operatorname{Sym}^2 L^{\vee}$ .

2.3.3. Higher groupoids in the pointed case. When the (braided) fusion categories in question are pointed, the groupoids introduced in Section 2.2 have a more concrete description. Let (A, q) be an (abelian) metric group, and let  $\mathcal{A} = \mathbf{Vect} [A]$  be the associated braided category of A-graded vector spaces. The Picard 3-type of  $\mathcal{A}$  has the following homotopy groups:

(2.13)  $\pi_1 = O(A,q) \qquad \pi_2 = A \qquad \pi_3 = \mathbf{k}^{\times}$ .

E.g. if  $A = L \oplus L^{\vee}$  then, in light of (2.3), this tells us that  $B \operatorname{Aut}_{\mathbf{Fus}}(\mathbf{Vect}[L])$  has  $\pi_2 = L \oplus L^{\vee}$ . If L is not abelian, then  $B \operatorname{Aut}_{\mathbf{Fus}}(\mathbf{Vect}[L])$  has  $\pi_2$  given by  $Z(L) \oplus L^{\vee}$ .

*Remark* 2.14. As explained in [ENO10, Remark 10.7], we can understand these homotopy groups as follows. First of all, we know that  $\pi_2 = A$  and  $\pi_2 = \mathbf{k}^{\times}$ , since this is the underlying groupoid of  $\mathcal{A}$  itself (delooped twice).

The bottom homotopy group  $\pi_1$  is identified with O(A, q) as follows. The Whitehead halfsquare (see [ENO10, §7.3]) is a homotopy invariant, which is a map  $\pi_2 \to \pi_3$ . As it turns out, in this case it is given by the (square of the) braiding on  $\mathcal{A}$ , which is defined by q in this case. The action of  $\pi_1$  on  $\pi_2$  by conjugation must preserve this invariant, and hence this defines a map  $\pi_1 \to O(A, q)$ , which turns out to be an equivalence.

This conjugation action is a shadow of a more general way to understand these groupoids. The invertible part of any monoidal 2-category acts on the endomorphism of the identify object by conjugation. As is explained in [ENO10, Remark 5.4], by setting the monoidal 2-category to be  $\operatorname{End}_{\mathbf{Fus}}(\mathcal{C})$ , we obtain a map:

$$\operatorname{Aut}_{\operatorname{\mathbf{Fus}}}(\mathcal{C}) \to \operatorname{Aut}_{\operatorname{\mathbf{EqBr}}}(\mathcal{Z}(\mathcal{C}))$$

When we restrict this to the truncation  $\pi_{\leq 2}$ , we obtain the equivalence (2.5). When we set the 2-category to be the category of modules over  $\mathcal{Z}(\mathcal{C})$  we obtain the equivalence (2.2).

2.4. **Obstruction theory.** We give an executive summary of the obstruction theory developed in [ENO10].

2.4.1. *Postnikov and Whitehead towers*. Recall the definition of the Postnikov and Whitehead towers of a space, and the associated *k*-invariants. See [Pos55, Whi78], or [MP12, Chapter 3] for references.

The 3-type  $B \operatorname{Aut}_{\mathbf{Fus}}(\mathcal{C}) \simeq B \operatorname{Pic}(\mathcal{Z}(\mathcal{C}))$  has Postnikov and Whitehead towers:



Recall the homotopy groups  $\pi_i$  from (2.13) and Remark 2.14.

2.4.2. Degree three obstruction.

Question 1. Given a group G and a morphism of groups  $f: G \to \pi_1$ , what is the obstruction to performing the following lift?



This question is equivalent to asking if the k-invariant  $k_2 \in H^3(B\pi_1, \pi_2)$  of the 2-type  $B \operatorname{Aut}_{\mathbf{EqBr}}(\mathcal{A})$  pulls back to something trivializable on G. I.e. the pullback

(2.16)  $O_3(f) \coloneqq (Bf)^* k_2 \in H^3(G, \pi_2)$ 

is the obstruction to lifting:

$$B \operatorname{Aut}_{\mathbf{EqBr}} (\mathcal{A})$$

$$\downarrow$$

$$BG \xrightarrow[Bf]{} B\pi_1 = \pi_{\leq 1} B \operatorname{Aut}_{\mathbf{EqBr}} (\mathcal{A})$$

*Remark* 2.17. At the level of classifying spaces, (2.2) tells us that the truncation of  $B \operatorname{Pic}(\mathcal{A})$  to a 2-type is identified with  $B \operatorname{Aut}_{\mathbf{EqBr}}(\mathcal{A})$ . Therefore  $O_3$  is the same as the obstruction to lifting:

$$BG \xrightarrow{\mathcal{B}_{5}} B\pi_{1} \simeq \pi_{<1}B\operatorname{Pic}(\mathcal{A})$$

*Remark* 2.18. When  $\mathcal{A} = \mathcal{Z}(\mathcal{C})$  for some fusion category  $\mathcal{C}$ , (2.3) allows us to describe this obstruction in terms of the 3-type  $B \operatorname{Aut}_{\mathbf{Fus}}(\mathcal{C})$ . In particular, it is the same as the obstruction to lifting:

$$BG \xrightarrow{B_{f}} B\pi_{1} \simeq \pi_{\leq 1}B\operatorname{Aut}_{\mathbf{Fus}}(\mathcal{C})$$

2.4.3. Degree four obstruction in the split case. Assume that the obstruction  $O_3(f)$  is trivializable, and let s be the section of  $B\operatorname{Aut}_{\mathbf{EqBr}}(\mathcal{Z}(\mathcal{C}))$  over  $B\pi_1$  corresponding to a fixed trivialization of  $O_3(f)$ . Recall the trivializations of  $O_3(f)$  form a torsor over  $H^2(BG, \pi_2)$ . By Remark 2.17, the upshot of the existence of s is that we have an associated map  $Bf_s: BG \to \pi_{\leq 2}B\operatorname{Pic}(\mathcal{A})$ .

Question 2. What is the obstruction to performing the following lift?



Just as before, the obstruction comes from the k-invariant. Now the relevant k-invariant is  $(2.19) k_3 \in H^4(\pi_{\leq 2}B\operatorname{Pic}(\mathcal{A}), \pi_3) .$ 

The pullback

$$O_4(f,s) \coloneqq (Bf_s)^* k_3 \in H^4(BG,\pi_3)$$

is the obstruction to lifting.

Remark 2.20. Again, if  $\mathcal{A} \cong \mathcal{Z}(\mathcal{C})$  for some fusion category  $\mathcal{C}$  then, by Remark 2.18, we have an associated map  $Bf_s : BG \to \pi_{\leq 2}B$  Aut<sub>Fus</sub> ( $\mathcal{C}$ ). By (2.3), the above question is equivalent to the following.

Question 3. What is the obstruction to performing the following lift?

$$B\operatorname{Aut}_{\mathbf{Fus}}(\mathcal{C})$$

$$BG \xrightarrow{\check{\mathcal{B}}_{f_s}} \pi_{\leq 2}B\operatorname{Aut}_{\mathbf{Fus}}(\mathcal{C})$$

The obstruction to lifting this is still  $O_4(f,s) \in H^4(BG,\pi_3)$ .

As it turns out, e.g. from [ENO10, Proposition 7.2, 7.3]:

$$\pi_3 \left( B \operatorname{Aut}_{\mathbf{Fus}} \left( \mathcal{C} \right) \right) \cong \mathbf{k}^{\times} \cong \pi_3 \left( B \operatorname{Pic} \left( \mathcal{A} \right) \right)$$

Therefore the obstruction is classified by familiar  $\mathbf{k}^{\times}$ -cohomology:

$$(2.21) O_4(f,s) \in H^4(BG,\mathbf{k}^{\times})$$

2.4.4. Full degree four obstruction. In Section 2.4.3, we assume that the map into

$$\pi_{<2}B\operatorname{Pic}(\mathcal{A})\simeq B\operatorname{Aut}_{\mathbf{EqBr}}(\mathcal{A})$$

is given by a trivialization of  $O_3(f)$  for f some map from a group to  $\pi_1$ . There is a slightly more general obstruction one can consider.

Question 4. Given a 2-group  $G_2$ , and a map

$$f_2: BG_2 \to \pi_{\leq 2} B \operatorname{Pic}(\mathcal{A}) \simeq B \operatorname{Aut}_{\mathbf{EqBr}}(\mathcal{A}) ,$$

what is the obstruction to performing the following lift?

$$B\operatorname{Pic}(\mathcal{A})$$

$$\downarrow$$

$$BG_{2} \xrightarrow{f_{2}} \pi_{\leq 2}B\operatorname{Pic}(\mathcal{A})$$

This is equivalent to asking if the k-invariant  $k_3$  from (2.19) pulls back to something trivializable on  $G_2$ .

Remark 2.22. As in the preceding subsections, this is equivalent to the analogous lifting question for  $B \operatorname{Aut}_{\mathbf{Fus}}(\mathcal{C})$ , by (2.3). Furthermore, a combination of (2.2) and (2.3) tell us that we can consider the universal version of this question:  $G_2 = B \operatorname{Aut}_{\mathbf{EqBr}}(\mathcal{Z}(\mathcal{C}))$ . I.e.

(2.23) 
$$k_3 \colon B\operatorname{Aut}_{\mathbf{EqBr}}(\mathcal{Z}(\mathcal{C})) \to B^4 \mathbf{k}^{\times}$$

is the obstruction to lifting

$$\begin{array}{c} B\operatorname{Aut}_{\mathbf{Fus}}\left(\mathcal{C}\right)\\ \downarrow\\ B\operatorname{Aut}_{\mathbf{EqBr}}\left(\mathcal{Z}\left(\mathcal{C}\right)\right) \xrightarrow{\sim} \pi_{\leq 2}B\operatorname{Aut}_{\mathbf{Fus}}\left(\mathcal{C}\right)\end{array}$$

2.4.5. Obstruction theory in the pointed case. Recall the homotopy groups in (2.13). If we take G = O(A, q), we get a universal obstruction class:

$$O_3 \in H^3(B \operatorname{O}(A,q), A)$$

Let |A| be odd. Then, as discussed in [EG18, §6] (and [CGPW16, §5]), the obstruction  $O_3$  vanishes, and in fact has a canonical splitting:

$$s: B \mathcal{O}(A,q) \to \pi_{\leq 2} B \operatorname{Pic}(\mathcal{A})$$
.

Therefore we have a well-defined universal version of the obstruction class in (2.21):

$$(2.24) O_4(A,q) \in H^4(B \cup (A,q), \mathbf{k}^{\times})$$

Remark 2.25. In other words,  $s^*B\operatorname{Aut}_{\mathbf{Fus}}(\mathcal{C})$  defines a  $B^3\mathbf{k}^{\times}$ -bundle over B O(A, q). At the level of 3-groups, this is actually a *central* extension of O(A, q) by  $B^2\mathbf{k}^{\times}$  since, for Brauer–Picard 3-groups of fusion categories,  $\pi_1$  automatically acts trivially on  $\pi_3$ .

Example 2.26. Let  $G = \mathbb{Z}/2$ . Then  $H^2(BG, \mathbf{k}^{\times}) = 0$  so the class  $O_4$  necessarily vanishes, and so the anomaly vanishes. There are two nonequivalent trivializations classified by  $H^3(BG, \mathbf{k}^{\times}) = \mathbb{Z}/2$ .

This is used in [ENO10, Example 9.4] to reproduce the classification in [TY98] of  $\mathbb{Z}/2$ -graded fusion categories.

Remark 2.27. Recall that the trivializations of  $O_3(f)$  form a torsor over  $H^2(BG, \pi_2)$ . I.e. given two trivializations s and s' of  $O_3(f)$ , there exists  $L \in H^2(BG, \pi_2)$  such that s' = Ls. By [ENO10, Proposition 8.15] we have that

$$O_4(f, s') / O_4(f, s) = \operatorname{PW}(L) ,$$

where PW denotes the Pontrjagin-Whitehead quadratic function from [ENO10, §8.7].

Also see [Jen11, Proposition 7.3] and [CGPW16, Proposition 8] for a concrete formula for the Pontrjagin-Whitehead quadratic function. It is shown to vanish in some examples in [GJ19].

2.4.6. Vanishing of the obstruction over a finite field. Let p be an odd prime, and let A = V be a 2*n*-dimensional vector space over a finite field. Let  $q_{\text{split}}$  be the quadratic form of signature (n, n).

**Observation 2.28.** The orthogonal group  $O(V, q_{split})$  is the split orthogonal group over  $\mathbb{F}_p$ .

Note that, if L is a vector space over  $\mathbb{F}_p$  of dimension n, then

$$\mathcal{E}\left(\mathbf{Vect}\left[L
ight],*
ight)\simeq\left(\mathbf{Vect}\left[V
ight],*,eta_{q}
ight)$$
 .

The obstruction  $O_4(A, q_{\text{split}})$  vanishes in this case by [EG18, Theorem 6.1].

There is a well-known analogy between the Weil representation and the spin representation. Correspondingly, the categorical representation(s) of O(V, q) studied in this paper are analogous to the Weil representation as well. This vanishing can be thought as an analogue of the fact that the Weil representation splits over a finite field [GH09, GH11].

# 3. TOPOLOGICAL SYMMETRY

3.1. **TQFTs associated to**  $\pi$ -finite spaces. Let X be a space (i.e. higher groupoid) which is (connected, pointed, and)  $\pi$ -finite<sup>3</sup>. There is a recipe for constructing a TQFT using X, which was introduced in [Kon88] and studied further in [Qui95, Tur10]. We will consider the fully local case introduced in [Fre94] and studied in [FHLT10, §3,8] and [FMT22, §A]. A formal treatment of these theories is the subject of an upcoming work of Claudia Scheimbauer and Tashi Walde.

<sup>&</sup>lt;sup>3</sup>This means X has finitely many homotopy groups, each of which is finite.

3.1.1. The quantization map. We will proceed heuristically, following [FHLT10, §3] and [FMT22, §A.2], to fix notation and describe expectations which will eventually be stated and assumed in Hypothesis Q.

Let  $\mathbf{Fam}_{d+1}$  denote the category with objects finite (d+1)-groupoids, 1-morphisms given by correspondences of  $\pi$ -finite spaces, 2-morphisms given by correspondences of correspondences, and so on until level (d+1). (Two (d+1)-morphisms are regarded as identical if they are equivalent.)

Let  $\mathcal{T}$  be the arbitrary symmetric monoidal target with duals, fixed in Appendix B. Let Y be an object of  $\operatorname{Fam}_{d+1}$ . A local system on Y valued in  $\mathcal{T}$  is a functor  $Y \to \mathcal{T}$ . Write  $\operatorname{Fam}_{d+1}(\mathcal{T})$ for the category of  $\pi$ -finite spaces equipped with a local system valued in  $\mathcal{T}$ .

For example,  $\mathbf{Fam}_{d+1}(B^{d+1}\mathbf{k}^{\times})$  has objects given by pairs  $(Y, \tau)$ , where  $\tau$  is a cocycle

$$\tau \colon Y \to B^{d+1} \mathbf{k}^{2}$$

representing a class in  $H^{d+1}(Y, \mathbf{k}^{\times})$ . A morphism is a correspondence:

(3.1) 
$$(Y_1, \tau_1) \xrightarrow{(E, \mu)} (Y_2, \tau_2)$$

where  $\mu \colon E \to B^d \mathbf{k}^{\times}$  satisfies

(3.2) 
$$d\mu = (p_1^* \tau_1)^{-1} \cdot (p_2^* \tau_2)$$

2-morphisms are correspondences between correspondences with a similar condition on the cocycles, and so on to define morphisms up to level (d + 1).

Recall we assumed  $\Omega^{d+1}\mathcal{T} = \mathbf{k}$ . Therefore there is a natural functor:  $B^{d+1}\mathbf{k}^{\times} \to \mathcal{T}$  inducing a functor:

(3.3) 
$$\mathbf{Fam}_{d+1}\left(B^{d+1}\mathbf{k}^{\times}\right) \to \mathbf{Fam}_{d+1}\left(\mathcal{T}\right) .$$

We assume that the following holds.

**Hypothesis Q.** For  $\mathcal{T}$  appropriately additive<sup>4</sup>, there is a quantization<sup>5</sup> functor

(3.4) 
$$\operatorname{Sum}_{d+1} \colon \operatorname{Fam}_{d+1} \left( B^{d+1} \mathbf{k}^{\times} \right) \to \mathcal{T}$$

such that there is an invertible natural transformation between

$$(X, \tau) \mapsto \operatorname{Hom}_{\mathcal{T}} (1, \operatorname{Sum}_{d+1} (X, \tau))$$

and

$$(X, \tau) \mapsto \operatorname{Hom}(\operatorname{hofib}(\tau), \Omega \mathcal{T})$$

viewed as functors out of  $\operatorname{Fam}_{d+1}(B^{d+1}\mathbf{k}^{\times})$ .

*Remark* 3.5. Unpacking the existence of the natural transformation in Hypothesis Q, we see that it ensures that we have equivalences

(3.6) 
$$\operatorname{Hom}_{\mathcal{T}}(1, \operatorname{Sum}_{d+1}(X)) \simeq \operatorname{Hom}(\operatorname{hofib}(\tau), \Omega \mathcal{T})$$

<sup>&</sup>lt;sup>4</sup>As is explained in [FHLT10, §3], for X an ordinary groupoid, then  $\mathcal{T}$  must be additive in the sense that the colimit  $\varinjlim_{x \in X} \tau(x)$  in  $\mathcal{T}$  exists, and agrees with the limit  $\varprojlim_{x \in X} \tau(x)$ , where we are regarding  $\tau$  as defining a  $\mathcal{T}$ -valued local system on X by (3.3).

<sup>&</sup>lt;sup>5</sup>See [FMT22, Remark A.7.1] where it is explained how  $\operatorname{Sum}_{d+1}(X)$  (and therefore  $\sigma_X^{d+1}$ ) can be obtained by integrating over fluctuating fields.

for all objects  $(X, \tau)$  of  $\mathbf{Fam}_{d+1}(B^{d+1}\mathbf{k}^{\times})$ , and given a morphism from  $(X_1, \tau_1)$  to  $(X_2, \tau_2)$  in  $\mathbf{Fam}_{d+1}(B^{d+1}\mathbf{k}^{\times})$ , we have a commuting diagram:

I.e. higher representations of  $\Omega X$  are equivalent to morphisms from 1 to  $\operatorname{Sum}_{d+1}(X)$  in  $\mathcal{T}$ . This generalizes the classical fact about modules over the group algebra and *G*-representation, as in Example 3.9 (ii).

Remark 3.8. In [FHLT10, §8.2], the quantization map  $\text{Sum}_n$  is constructed at the level of objects and shown to be a functor up to 2-morphisms. In particular, the cases in Example 3.9 are worked out in [FHLT10, §8.1,8.3].

*Example* 3.9. (i) Let  $\mathcal{T} =$ **Vect**. Then  $\text{Sum}_1(X) = \mathbf{k}(\pi_0 X)$  is the vector space of **k**-valued functions on  $\pi_0 X$ . Hypothesis Q is satisfied, since the natural transformation (3.6) required in Hypothesis Q is:

 $\operatorname{Hom}_{\operatorname{Vect}}(1, \operatorname{Sum}_1(X)) \simeq \operatorname{Hom}(X, \mathbf{k}) \simeq \operatorname{Hom}(\pi_0 X, \mathbf{k})$ .

(ii) Set X = BG for a finite group G, and let  $\mathcal{T} = \mathbf{Alg}$  be the Morita 2-category of algebras. Let  $\operatorname{Sum}_2(BG) = \mathbf{k}[G]$  be the group algebra. Hypothesis Q is satisfied, since the natural transformation (3.6) required in Hypothesis Q is:

 $\operatorname{Hom}_{\operatorname{Alg}}(1, \operatorname{Sum}_{2}(BG)) = \mathbf{k}[G] \operatorname{-mod} \simeq \operatorname{Rep}(G) \simeq \operatorname{Hom}(BG, \operatorname{Vect})$ .

We can equip BG with a 2-cocycle  $\mu$  (i.e. a group 2-cocycle) which twists the convolution structure on the group algebra, resulting in Sum<sub>2</sub> ( $BG, \tau$ ).

- (iii) Set X = BG. Let d = 2, and take  $\mathcal{T}$  to be the Morita 3-category of monoidal categories. Then Sum<sub>3</sub> (BG) is **Vect** [G], the fusion category of vector bundles on G with convolution. We can equip BG with a 3-cocycle  $\alpha$  which twists the fusion structure on **Vect** [G], yielding  $\sigma_{BG,\alpha}^3$  (\*). This is [FMT22, Example A.65].
- (iv) Set  $X = B^2 G$ . Let d = 3 and take  $\mathcal{T}$  to be the Morita 4-category of braided monoidal categories, **BrFus** (see Section 2.1). Consider a cocycle  $\tau: B^2 G \to B^4 \mathbf{k}^{\times}$ . It is a classical theorem of Eilenberg-Maclane [EM46] that cohomology classes in  $H^4 (B^2 G, \mathbf{k}^{\times})$  correspond to quadratic forms  $G \to \mathbf{k}^{\times}$ . Write  $q_{\tau}$  for the form corresponding to  $[\tau]$ . This defines a symmetric bicharacter on G:

$$\langle g,h \rangle_{\tau} \coloneqq \frac{q_{\tau} \left(g+h\right)}{q_{\tau} \left(g\right) q_{\tau} \left(h\right)}$$

Then Sum<sub>4</sub> ( $B^2G$ ) is Vect [G] with convolution, and with braiding specified on simples by:

$$eta_{ au} \colon \mathbf{k}_g \ast \mathbf{k}_h = \mathbf{k}_{gh} \xrightarrow{\langle g,h \rangle \operatorname{id}_{\mathbf{k}_{gh}}} \mathbf{k}_{gh} = \mathbf{k}_{hg} = \mathbf{k}_h \ast \mathbf{k}_g \; .$$

- (v) Let X = BG and d = 3. If G is abelian, and we take  $\mathcal{T} = \mathbf{BrFus}$ , then this quantizes to the symmetric (so in particular braided) fusion category Vect [G].
- (vi) Let X = BG and d = 3, only now we would like to use the category of monoidal 2categories. Then BG can be quantized to the fusion 2-category (in the sense of [DR18]) of G-graded 2-vector spaces [DR18, Construction 2.1.13].

As far as the author is aware, dualizability in the Morita 4-category of monoidal 2-categories has not been extensively studied. However this fusion 2-category is expected define a fully extended 4-dimensional TQFT [DR18].

3.1.2. TQFTs from the quantization map. The upshot of assuming Hypothesis Q is that, for a fixed object  $(X, \tau)$  of  $\operatorname{Fam}_{d+1}(B^{d+1}\mathbf{k}^{\times})$ , (3.4) can be composed with the mapping space functor<sup>6</sup> to obtain the theory

$$\sigma_{X,\tau}^{d+1} \colon \mathbf{Bord}_{d+1}^{\mathrm{fr}} \xrightarrow{\pi_{\leq d+1} \operatorname{Map}(-,X)} \mathbf{Fam}_{d+1}\left(\mathcal{T}\right) \xrightarrow{\operatorname{Sum}_{d+1}} \mathcal{T} \xrightarrow{}$$

*Remark* 3.10. In [FHLT10, §3,8], the theories  $\sigma_X^{d+1}$  are studied for  $\mathcal{T} = \mathbf{Alg}[d]$  the Morita (d+1)-category of "d-algebras", discussed in [FHLT10, §7].

*Remark* 3.11. As is remarked in [FMT22, §A] and [FHLT10, §3,8],  $\sigma_{X,\tau}^{d+1}$  can be upgraded to an oriented theory, and if  $\tau$  is trivial then it can even be upgraded to an unoriented theory. We will work with the underlying framed theories in this paper.

The following follows directly from Hypothesis Q. See [FMT22] for more.

**Proposition 3.12.** Assuming Hypothesis Q, a morphism from  $(X_1, \tau_1)$  to  $(X_2, \tau_2)$  in

 $\operatorname{Fam}_{d+1}(B^{d+1}\mathbf{k}^{\times})$ 

(i.e. correspondence as in (3.1)) induces a bimodule (i.e. domain wall)

$$\sigma_{X_1,\tau_1}^{d+1} \to \sigma_{X_2,\tau_2}^{d+1}$$
.

Remark 3.13. Proposition 3.12 can be shown directly (i.e. without the Cobordism Hypothesis) since a correspondence of spaces  $X \leftarrow C \rightarrow Y$  defines a correspondences of mapping spaces  $\operatorname{Map}(M, X) \leftarrow \operatorname{Map}(M, C) \rightarrow \operatorname{Map}(M, Y)$ , for any bordism M. This is the perspective taken in [FMT22].

Recall the motivation in Remark 3.5 for the existence of the natural transformation in Hypothesis Q. Proposition 3.14, proved in [VD23, Proposition B.14], writes this in terms of the theories  $\sigma_X^{d+1}$ .

**Proposition 3.14.** Assuming Hypothesis Q, every boundary theory for  $\sigma_{X,\tau}^{d+1}$  is classified by a symmetric-monoidal functor:

$$\operatorname{Bord}_d^{\operatorname{hofib}(\tau)} \to \Omega \mathcal{T}$$

If  $\tau$  is trivial, then the boundary theories are classified by symmetric-monoidal functors:

$$\mathbf{Bord}_d^X o \Omega \mathcal{T}$$
 .

3.2. Module structures. We summarize the material used in our construction from [FMT22]. See [FMT22, §3] for a more detailed discussion of these definitions. Let

$$\sigma \colon \mathbf{Bord}_{d+1}^{\mathrm{fr}} o \mathcal{T}$$

be a (d+1)-dimensional TQFT valued in the fixed target  $\mathcal{T}$  from Appendix B. Recall the notion of a boundary theory from Remark B.12. Recall the following definition in [FMT22].

**Definition.** A *d*-dimensional quicke is a pair  $(\sigma, \rho)$  in which  $\rho$  is a right topological boundary theory (or right  $\sigma$ -module), which we will write as  $\rho: \sigma \to 1$ .

<sup>&</sup>lt;sup>6</sup>Note that (3.3) allows us to construct a  $\mathcal{T}$ -valued local system from the cocycle  $\tau$ .

*Remark* 3.15. All of the quiches considered in this paper will be of the form  $(\sigma_{X,\tau}, \rho_{X,\tau})$ , where  $\sigma_{X,\tau}$  is the theory associated to  $(X,\tau)$  as in Section 3.1, and the (right) boundary theory  $\rho_{X,\tau}$  is the natural transformation induced by the correspondence diagram:



as in Proposition 3.12. Given a pointed space X, we will always write  $\rho_{X,\tau}$  for this boundary theory.

A quiche is an abstract symmetry datum, in the same sense as an algebra. The following definition is the analogue of a module, i.e. a realization of the quiche as symmetries of a given theory.

**Definition.** Let  $(\sigma, \rho)$  be a *d*-dimensional quiche, and let *F* be a *d*-dimensional TQFT. A  $(\sigma, \rho)$ -module structure on *F* is the pair  $(F_{\sigma}, \theta)$  where  $F_{\sigma}$  is a (left) boundary theory  $F_{\sigma} : 1 \to \sigma$  which is equipped with an isomorphism of *d*-dimensional theories:

$$\theta \colon \rho \otimes_{\sigma} F_{\sigma} \xrightarrow{\sim} F$$
.

# 4. Semiclassical symmetry

4.1. Semiclassical group actions. Fix  $d \in \mathbb{Z}^{\geq 0}$ , and let X be a  $\pi$ -finite space with cocycle  $\tau \colon X \to B^d \mathbf{k}^{\times}$ .

**Definition 4.1.** A semiclassical action of a group G on  $(X, \tau)$  is a correspondence of spaces

as in Proposition 3.12 such that we have a pullback:



I.e. a semiclassical action of G on  $(X, \tau)$  is just an X-bundle over BG, and an extension of  $\tau$  to the total-space of this bundle.

*Example* 4.3. Let X = BL for a finite group L. For trivial  $\tau$ , there is always a canonical semiclassical action of Aut<sub>Grp</sub> (L) given by

$$C_{\text{Aut}} = B\left(L \rtimes \text{Aut}_{\mathbf{Grp}}\left(L\right)\right)$$

since the following is a pullback diagram:



The same holds for any  $\varphi \colon G \to \operatorname{Aut}_{\mathbf{Grp}}(L)$ . Namely,

$$C_G = B\left(L \rtimes_{\varphi} G\right)$$

is the pullback



and this defines a semiclassical action because we also have a pullback



**Definition 4.4.** A projective semiclassical action of a group G on  $(X, \tau)$  with projectivity  $c: BG \to B^{d+1}\mathbf{k}^{\times}$  is a correspondence of spaces



as in Proposition 3.12 such that we have a pullback:



4.2. Spectral sequence obstruction. Consider a semiclassical action  $C_G$  of G on X as in Definition 4.1 (i.e. the twist is trivial). Now, given some nontrivial  $\tau$ , we can pose the question of whether there is a semiclassical action  $(C_G, \tau_G)$  of G on  $(X, \tau)$ . I.e. we are asking if the cocycle  $\tau$  extends to a cocycle  $\tau_G$  on  $C_G$ :



The Serre spectral sequence provides a general method for answering this question. The first quadrant Serre spectral sequence corresponding to the fibration

$$\begin{array}{c} X \stackrel{\iota}{\longrightarrow} C_G \\ \downarrow \\ BG \end{array}$$

18

is

$$E_2^{p,q} = H^p\left(BG, H^q\left(X, \mathbf{k}^{\times}\right)\right) \Rightarrow H^{p+q}\left(C_G, \mathbf{k}^{\times}\right)$$

Recall the differentials:

$$d_r^{p,q}: E_r^{p,q} \to E_r^{p+r,q-r+1}$$
.

The group G acts on  $H^d(X, \mathbf{k}^{\times})$  (induced by conjugation in  $\Omega C_G$ ) and every cocycle on  $C_G$  is invariant under this action, so we insist that

$$au \in H^d\left(X, \mathbf{k}^{\times}\right)^G$$

Then we have that

$$\tau \in \operatorname{im}\left(\iota^{*}\right) = E_{d+2}^{0,d} \hookrightarrow H^{d}\left(X,\mathbf{k}^{\times}\right)^{d}$$

if and only if

(4.5) 
$$d_k^{0,d} \tau = d_k \tau = 0 \in E_k^{k,d+1-k}$$

**Theorem 4.6.** If the first (d-1) obstructions (4.5) vanish, i.e.  $d_k \tau$  for  $2 \le k \le d$ , then there is a projective semiclassical action of G on  $(X, \tau)$ , and this quantizes to an anomalous action of G on the theory  $\sigma_{X,\tau}^d$  with anomaly classified by  $d_{d+1}\tau$ .

*Proof.* Assuming the obstructions  $d_k \tau$  vanish for  $2 \le k \le d$ , there exists some class

$$\tau_d \colon C_G \to B^d \mathbf{k}^{\diamond}$$

satisfying

$$d\tau_d = d_{d+1}\tau_d$$

so that we have a projective semiclassical action (Definition 4.4):

$$(BG, d_{d+1}\tau) (C_G, \tau_G)$$

The quantization of this correspondence is a boundary theory:

$$1 \to \sigma^{d+1}_{BG,d_{d+1}\tau}$$

such that pairing with the regular boundary condition gives a theory canonically equivalent to  $\sigma_{X,\tau}^d$ .

By [VD23, Theorem C.16], this is equivalent to an anomalous theory defined on **Bord**<sup>BG</sup>, with anomaly theory classified by  $d_{d+1}\tau \colon BG \to B^{d+1}\mathbf{k}^{\times}$ .

Recall our d-dimensional theories  $\sigma^d_{X,\tau}$  are valued in  $\Omega \mathcal{T}$  from Appendix B. Consider the d-type

$$B\operatorname{Aut}_{\mathcal{T}}\left(\sigma_{X,\tau}^{d}\left(*\right)\right)$$

with top homotopy group  $\pi_3$ . Because we assumed  $\Omega^{d+1}\mathcal{T} \simeq \mathbf{k}^{\times}$ , we have a map:

(4.7) 
$$\iota \colon B^d \mathbf{k}^{\times} \to B^d \pi_3$$

Note that the target of  $\iota$  is the initial term of the Whitehead tower of the *d*-type

$$B\operatorname{Aut}_{\mathcal{T}}\left(\sigma_{X,\tau}^{d}\left(*\right)\right)$$
.

Write the k-invariant of  $B\operatorname{Aut}_{\mathcal{T}}\left(\sigma_{X,\tau}^{d}\left(*\right)\right)$  over the truncation  $\pi_{\leq d-1}$  as

(4.8) 
$$k \colon \pi_{\leq d-1} B \operatorname{Aut}_{\mathcal{T}} \left( \sigma_{X,\tau}^d \left( * \right) \right) \to B^{d+1} \pi_3 .$$

**Corollary 4.9.** If the first (d-1) obstructions (4.5) vanish, then there is a map

$$\mathbf{f}: BG \to \pi_{\leq d-1}B\operatorname{Aut}_{\mathcal{T}}(\sigma_{X,\tau}(*))$$

and the projectivity of the projective G-action from Theorem 4.6 agrees with the restriction of the k-invariant:

$$BG \xrightarrow{\mathbf{f}} \pi_{\leq d-1} B \operatorname{Aut}_{\mathcal{T}} (\sigma_{X,\tau} (*))$$
$$\downarrow^{d_{d+1\tau}} \qquad \qquad \qquad \downarrow^{k}$$
$$B^{d+1} \mathbf{k}^{\times} \xrightarrow{B\iota} B (B\Omega)^{d} B \operatorname{Aut}_{\mathcal{T}} (\sigma_{X,\tau} (*))$$

where  $\iota$  is from (4.7) and k is from (4.8).

*Proof.* Let  $(C_G, \tau_G)$  be the projective semiclassical action (Definition 4.4) from Theorem 4.6. The quantization of this is a boundary theory  $1 \to \sigma_{BG,c}^{d+1}$ .

By [VD23, Theorem C.16], this is equivalent to a TQFT defined on **Bord**<sup>hofb( $d_{d+1}\tau$ )</sup>, which is classified via the Cobordism Hypothesis (B.6) by a functor

hofib 
$$(d_{d+1}\tau) \to B\operatorname{Aut}_{\Omega \mathcal{T}} \left( \sigma^d_{X,\tau} \left( * \right) \right)$$

which truncates to a functor

$$BG \to \pi_{\leq d-1} B \operatorname{Aut} (\mathcal{C})$$

# 5. Symmetries of the base

Let *L* be a finite group, and  $\tau$  a group cocycle representing a class in  $H^3(L, \mathbf{k}^{\times})$ . Let  $\mathcal{C} = \mathbf{Vect} [L]^{\tau}$  be the fusion category of vector spaces graded by *L*, twisted by  $\tau$ .

In Section 5.1, we will construct an action by bimodules up to isomorphism:

 $f: G \to \pi_0 \operatorname{Aut}_{\mathbf{Fus}} (\mathcal{C})$ 

which is induced by an action of G on the base:

$$\varphi \colon G \to \operatorname{Aut}(L,\tau)$$

where Aut  $(L, \tau)$  consists of group automorphisms of L which preserve the cohomology class of  $\tau$ . This action can be twisted by a class  $\gamma$  satisfying a certain equation.

The map f will lift to a map to the full 3-group:

$$\mathbf{f}: G \to \operatorname{Aut}_{\mathbf{Fus}}(\mathcal{C})$$

if and only if certain obstructions identified in [ENO10] vanish, and are trivialized. In Section 5.2 and Section 5.3, we explicitly describe the data necessary to trivialize these obstructions, which will be written as  $\mu$  and  $\alpha$  respectively. All together, given L and  $\tau$ , the map **f** is determined by the map  $\varphi$  and the triple  $(\gamma, \mu, \alpha)$ .

Importantly, when the twist is trivial, the map  $\varphi$  canonically defines such a map **f**. This is the content of Theorem 5.15.

In Section 5.4, we describe a semiclassical action (Definition 4.1) associated to the same data  $(\varphi, \gamma, \mu, \alpha)$ . Furthermore, the quantization of this semiclassical action agrees with the map **f**.

# 5.1. Action via bimodules up to isomorphism. Consider an arbitrary map

$$\varphi \colon G \to \operatorname{Aut}(L,\tau)$$

and a group 1-cochain on G valued in group 2-cochains on L:

(5.1) 
$$\gamma \colon G \to C^2_{\rm grp}\left(L, \mathbf{k}^{\times}\right)$$

If this satisfies the equation:

(5.2) 
$$d_L(\gamma(g)) = \varphi(g)(\tau) \tau^{-1},$$

then this defines a morphism in  $\mathbf{Fam}_d$  (as in Proposition 3.12):

where  $d_L$  is the group differential for L. Note that such a  $\gamma$  exists because the action  $\varphi$  preserves the cohomology class of  $\tau$ .

Write  $C_{\gamma}(g)$  for this 1-morphism in  $\mathbf{Fam}_d$ . Then we define:

(5.4) 
$$f(g) \coloneqq \operatorname{Sum}_d(C_{\gamma}(g))$$

where  $\operatorname{Sum}_d$  is the quantization functor (3.4).

The C-bimodule f(g) has the following description. The underlying category is the same as C, and the left C-module structure is simply the tensor product in C. The right action, say by some simple  $\mathbf{k}_{\ell} \in C$ , is given by tensor product with  $\mathbf{k}_{\varphi(\ell)}$ .

The twist of this construction by the 2-cocycle  $\gamma(g)$  can be thought of as follows. Rather than defining f(g) to be the trivial  $\mathcal{C}$ -bimodule twisted by a group automorphism of L, as above, we might consider twisting by a tensor autoequivalence  $\Phi$  of  $\mathcal{C}$ . I.e. now an object  $X \in \mathcal{C}$  acts on the right by tensoring with  $\Phi(X)$ . The tensor autoequivalences form an extension:

$$1 \to H^2(L, \mathbf{k}^{\times}) \to \pi_0 \operatorname{Aut}_{\otimes}(\mathcal{C}) \to \operatorname{Aut}(L, \tau) \to 1$$
,

where Aut  $(L, \tau)$  denotes the group automorphisms of L preserving the cohomology class defined by the cocycle  $\tau$ . See [ENO10, §11.2] for more.

5.2. **System of products.** Now we would like to construct a system of products, i.e. equivalences:

$$m(g,h): f(h) \circ f(g) \xrightarrow{\sim} f(gh)$$

attached to all pairs  $g, h \in G$ .

We will obtain m(g,h) by quantizing a correspondence between correspondences as in Proposition 3.12. Note that for any  $g, h \in G$  we have

$$\Gamma\left(\varphi\left(g\right)\right)\circ\Gamma\left(\varphi\left(h\right)\right)=\Gamma\left(\varphi\left(g\right)\circ\varphi\left(h\right)\right)=\Gamma\left(\varphi\left(gh\right)\right)\ .$$

so  $f(h) \circ f(g)$  is the quantization of



and f(qh) is the quantization of

$$(\Gamma\left(\varphi\left(gh\right)\right),\gamma\left(gh\right))$$

$$(L,\tau)$$

$$(L,\varphi\left(gh\right)(\tau))$$

Therefore we require a group 2-cochain on G valued in 1-cochains on L:

(5.5) 
$$\mu \colon G^2 \to C^1_{\rm grp}\left(L, \mathbf{k}^{\times}\right) \ .$$

If  $\mu$  (and  $\gamma$ ) satisfy:

$$d_{L}(\mu(g,h)) = \gamma(gh)\gamma(g)^{-1}\gamma(h)^{-1}$$

in  $C^2(L, \mathbf{k}^{\times})$  (where  $d_G$  and  $d_L$  are the differentials for the group cohomology of G and L respectively) or more succinctly:

$$(5.6) d_L \mu = d_G \gamma^{-1}$$

then this defines a 2-morphism (correspondence of correspondences) in  $\mathbf{Fam}_d$ :

(5.7) 
$$(\Gamma (\varphi (gh)), \mu (g, h)) \longrightarrow (\Gamma (\varphi (gh)), \gamma (g) \gamma (h)) (\Gamma (\varphi (gh)), \gamma (gh))$$

Write  $C_{\mu}(g,h)$  for this 2-morphism in **Fam**<sub>d</sub>.

(5.8) 
$$m(g,h) \coloneqq \operatorname{Sum}_d(C_\mu(g,h))$$

5.2.1. The degree three obstruction. Recall the obstruction  $O_3(f) \in H^3_{\text{Grp}}(G, L \oplus L^{\vee})$  from Section 2.4.2. This is defined using f and m, but only depends on f [ENO10]. Recall  $\gamma$  from (5.1) satisfying (5.2). Insisting that the obstruction  $O_3(f)$  vanishes would put a further condition on  $\gamma$ , however in the setting considered in this section, it will always vanish.

Consider the cocycle:

$$d_G \mu \in Z^3_{\mathrm{Grp}}\left(G, C^1\left(L\right)\right)$$
.

In fact, this defines an element

$$d_G \mu \in Z^3_{\mathrm{Grp}}\left(G, L^{\vee}\right)$$

where

$$L^{\vee} = \operatorname{Hom}\left(L, \mathbf{k}^{\times}\right) = Z^{1}\left(L, \mathbf{k}^{\times}\right)$$

because

$$d_L \circ d_G(\mu) = d_G(d_L\mu) = d_G(d_G\gamma)^{-1} = 0$$

since  $\gamma$  and  $\mu$  satisfy (5.6).

*Remark* 5.9. This section is similar to the discussion of the same obstruction in [CGPW16, §5.1].

**Proposition 5.10.** If f is the group homomorphism defined by  $\varphi$  and  $\gamma$  in (5.4), then  $O_3(f)$  vanishes.

*Proof.* Recall from Section 2.3.3, in this case we have:

$$O_{3}(f) \in H^{3}\left(G, Z(L)_{\operatorname{Grp}} \oplus L^{\vee}\right)$$
.

As noted in [ENO10, §9.1], this lies in the image of:

$$H^{3}_{\mathrm{Grp}}\left(G, Z\left(L\right)\right) \to H^{3}_{\mathrm{Grp}}\left(G, Z\left(L\right) \oplus L^{\vee}\right)$$

The image of

$$[d_G\mu] \in H^3_{\mathrm{Grp}}\left(G, L^{\vee}\right)$$

under the map

 $(\Gamma (\varphi$ 

$$H^{3}_{\mathrm{Grp}}\left(G, L^{\vee}\right) \to H^{3}_{\mathrm{Grp}}\left(G, Z\left(L\right) \oplus L^{\vee}\right)$$

induced by the inclusion  $L^{\vee} \hookrightarrow Z(L) \oplus L^{\vee} = \pi_2$  is the obstruction  $O_3(f)$  for f defined in (5.4). I.e.  $O_3(f)$  is trivial.

5.3. Associators. Now we would like to construct a system of associators, i.e. equivalences:

$$a\left(g,h,k\right):m\left(gh,k\right)\circ\left(m\left(g,h\right)\otimes\mathrm{id}_{f(k)}\right)\xrightarrow{\sim}m\left(g,hk\right)\circ\left(\mathrm{id}_{f(g)}\otimes m\left(h,k\right)\right)\ .$$

The source of a(g, h, k) will be the quantization of:

$$(ghk)), \gamma (g) \gamma (h) \gamma (k)) \xrightarrow{(\Gamma (\varphi (ghk)), \mu (gh, k) \mu (gh))} (\Gamma (\varphi (ghk)), \gamma (ghk))$$

and the target will be the quantization of:

$$(\Gamma\left(\varphi\left(ghk\right)\right),\gamma\left(g\right)\gamma\left(h\right)\gamma\left(k\right)) \xrightarrow{\left(\Gamma\left(\varphi\left(ghk\right)\right),\mu\left(g,hk\right)\mu\left(hk\right)\right)} (\Gamma\left(\varphi\left(ghk\right)\right),\gamma\left(ghk\right))$$

Therefore we require a group 3-cochain on G valued in  $C^0(L, \mathbf{k}^{\times}) = \mathbf{k}^{\times}$ :

(5.11) 
$$\alpha \colon G^3 \to \mathbf{k}^{\times}$$
.

The analogue of (5.2) for  $\gamma$  (and similarly of (5.6) for  $\mu$ ) is

$$d_{L}(\alpha(g,h,k)) = \mu(g,hk) \mu(h,k) \mu(gh,k)^{-1} \mu(g,h)^{-1}$$

or more succinctly:

 $(5.12) d_L \alpha = d_G \mu$ 

in  $C^1(L, \mathbf{k}^{\times})$ . Since  $\alpha(g, h, k)$  is a 0-cocycle on L (in cohomology for the trivial action of L on  $\mathbf{k}^{\times}$ ), it is always closed. I.e. the condition reduces to

(5.13) 
$$d_G \mu = 0$$
.

The 3-cochain  $\alpha$  defines a correspondence (between correspondences of correspondences):

$$\overbrace{\left(\Gamma\left(\varphi\left(ghk\right)\right),\mu\left(gh,k\right)\mu\left(gh\right)\right)}^{\left(\Gamma\left(\varphi\left(ghk\right)\right),\alpha\left(g,h,k\right)\right)},\left(\Gamma\left(\varphi\left(ghk\right)\right),\mu\left(g,hk\right)\mu\left(hk\right)\right)}$$

i.e. a 3-morphism in  $\mathbf{Fam}_d$ , which we write as  $C_{\alpha}(g, h, k)$ . Finally, the associator is defined by: (5.14)  $a(g, h, k) \coloneqq \operatorname{Sum}_d(C_{\alpha}(g, h, k))$ .

5.3.1. Degree four obstruction. Recall the obstruction 
$$O_4(f, m) \in H^4_{\text{Grp}}(G, \mathbf{k}^{\times})$$
. This is defined using  $f, m$ , and  $a$ , but only depends on  $f$  and  $m$  [ENO10]. When  $f$  is the map defined in (5.4), the system of products  $m$  is the one defined in (5.8), and the associator  $a$  is the one defined in (5.14), we have that the obstruction class is simply:

$$O_4(f,m) = [d_G\alpha] \in H^4_{\mathrm{Grp}}(G,\mathbf{k}^{\times}) ,$$

which always vanishes.

5.3.2. Fully coherent action induced by an automorphism of the base. Let L and G be finite groups. Let  $\tau \in Z^3_{\text{Grp}}(L, \mathbf{k}^{\times})$  be a 3-cocycle in the group cohomology of L with coefficients in  $\mathbf{k}^{\times}$ , and consider an arbitrary group homomorphism

$$\varphi \colon G \to \operatorname{Aut}(L,\tau)$$
,

where Aut  $(L, \tau)$  denotes the group of automorphisms preserving the cohomology class of  $\tau$ .

**Theorem 5.15.** Assume Hypothesis Q. Let L and G be finite groups. Let  $\tau \in Z^3_{Grp}(L, \mathbf{k}^{\times})$  be a 3-cocycle in the group cohomology of L. Then any group homomorphism

$$\varphi \colon G \to \operatorname{Aut}(L,\tau)$$

canonically defines a  $(\sigma_{BG}^4, \rho)$ -module structure on  $\sigma_{BL,\tau}^3$ .

This canonical module structure can be twisted by a triple  $(\gamma, \mu, \alpha)$  where

$$\gamma \in C^{1}_{Grp}\left(G, C^{2}_{Grp}\left(L, \mathbf{k}^{\times}\right)\right) \qquad \mu \in Z^{2}_{Grp}\left(G, C^{1}_{Grp}\left(L, \mathbf{k}^{\times}\right)\right) \qquad \alpha \in Z^{3}\left(G, \mathbf{k}^{\times}\right)$$

satisfy:

$$d_L(\gamma(g)) = \varphi(g)(\tau)\tau^{-1} \qquad \qquad d_L\mu = d_G\gamma^{-1}$$

*Proof.* The canonical structure will correspond to trivial  $\gamma$ ,  $\mu$ , and  $\alpha$ . We will show that we obtain a module structure for arbitrary  $(\gamma, \mu, \alpha)$ .

 $\varphi(g)(\tau) \tau^{-1} \in Z^2_{\text{Grp}}(L, \mathbf{k}^{\times})$  defines the trivial cohomology class (since the action determined by  $\varphi$  preserves the cohomology class of  $\tau$ ). Therefore any  $\gamma$  satisfies (5.2). The fact that  $O_3(f)$ for the group homomorphism f defined by  $\varphi$  and  $\gamma$  follows from Proposition 5.10. As in Section 5.3.1, the obstruction  $O_4(f, m) = 0$  vanishes for all such  $\tau$  and  $\varphi$ , and for all  $\gamma$  and  $\mu$ .

Combining the group homomorphism f from (5.4) (defined by  $\varphi$  and  $\gamma$ ) the system of products m from (5.8) (defined by  $\mu$ ) and the associator a from (5.14) (defined by  $\alpha$ ), we obtain a map of 3-groups:<sup>7</sup>

(5.16) 
$$\mathbf{f} = (f, m, a) : G \to \operatorname{Aut}_{\mathbf{Fus}}(\mathcal{C})$$

This equivalently defines a  $(\sigma_{BG}^{d+1}, \rho)$ -module structure on  $\sigma_{X,\tau}^d$  by [VD23, Theorem C.16].  $\Box$ 

Remark 5.17. We consider semiclassical actions  $BG \leftarrow C_G \rightarrow *$  of G on BL (Definition 4.1) of a very special form in this section:

$$C_G = B\left(L \rtimes_\varphi G\right) \; .$$

Arbitrary such semiclassical actions  $C_G$  are equivalent to arbitrary group extension of G by L, which are classified by degree 2 nonabelian group cohomology of G with coefficients in L.

Even if L is abelian, extensions given by semidirect products are still quite restrictive: For every  $\varphi$ , there is an extension associated to every class in  $H^2_{\text{Grp},\varphi}(G,L)$ , where this is group cohomology with the action  $\varphi$  of G on L.

Such, more general, considerations are made in in [ENO10, §9.1,§11]. Namely,

$$\varphi(gh)^{-1} \circ \varphi(g) \circ \varphi(h)$$

might be conjugation by some element  $n(g,h) \in L$ . This defines a class  $[n] \in H^2(BG, L)$ .

Once we have this twist n, the extension  $L \rtimes_{\varphi} G$  is replaced by the group extension determined by the pair  $(\varphi, n)$ . This introduces a possible nonzero obstruction class  $O_4(f, m)$ .

<sup>&</sup>lt;sup>7</sup>Here G denotes the discrete 2-category with objects G.

5.4. Semiclassical action. In this section we will construct a semiclassical action of G on  $(BL, \tau)$ , from the same data from above: an action  $\varphi$  and a triple  $(\gamma, \mu, \alpha)$ . The quantization of this semiclassical action will be equivalent to the map (f, m, a) in (5.16).

Note that the map  $\varphi$  canonically defines a semiclassical action of G on BL (Example 4.3), and recall the anomaly/obstruction to the existence of a semiclassical action of G on  $(BL, \tau)$ studied in Section 4.2. In this case, the space X is BL, so the Serre spectral sequence considered in Section 4.2 reduces to the Lyndon-Hochschild-Serre (LHS) spectral sequence associated to the normal subgroup  $L \hookrightarrow L \rtimes_{\varphi} G$ :

$$E_{2}^{p,q} = H_{\mathrm{Grp}}^{p}\left(G, H^{q}\left(L, \mathbf{k}^{\times}\right)\right) \Rightarrow H_{\mathrm{Grp}}^{p+q}\left(L \rtimes_{\varphi} G, \mathbf{k}^{\times}\right)$$

See [Mü20, MS20, KT14] and [ENO10, §11.8] for more on the LHS spectral sequence as a tool for encoding anomalies.

The following follows from [ENO10, §11.8].

**Proposition 5.18.** A map (f, m, a) as in (5.16) equivalently defines an extension of the cocycle  $\tau$  to a cocycle  $\tau_G$  on  $L \rtimes_{\varphi} G$  which agrees with  $\tau$  when restricted to L.

Therefore, we have a morphism as in Proposition 3.12:

$$BG \xrightarrow{(B(L \rtimes_{\varphi} G), \tau_G)} *$$

The quantization of this correspondence is a boundary theory:

$$F_G: 1 \to \sigma_{BG}^{d+1}$$
,

such that when we pair with the regular boundary condition, we obtain  $\sigma_{BL}^{d+1}$ . By [VD23, Theorem C.16],  $F_G$  is equivalent to a functor

$$F_G \colon BG \to B\operatorname{Aut}_{\mathbf{Fus}}(\mathcal{C})$$
.

This functor  $F_G$  provides another description of he functor (f, m, a).

### 6. Orthogonal symmetries

6.1. Analogue of BKS kernels. Let (A, q) be a finite abelian metric group (i.e. a finite abelian group equipped with nondegenerate quadratic form  $q: A \to \mathbf{k}$ ) with two Lagrangian subgroup  $L, K \subset A$ . Then we have a correspondence of groups:

$$A/(L \cap K) \xrightarrow{/(L/L \cap K)} A/K \xrightarrow{/(K/L \cap K)} A/L$$

If  $L \cap K = \{1\}$ , i.e. they are transverse, then this takes the simpler form:



By the Schur isomorphism (A.2), which in this case is:

$$H^{2}(A, \mathbf{k}^{\times}) \simeq H^{2}(L, \mathbf{k}^{\times}) \oplus H^{2}(L^{\vee}, \mathbf{k}^{\times}) \oplus \operatorname{Hom}(A, \mathbf{k}^{\times})$$
,

the quadratic form q defines a class in  $H^2(A, \mathbf{k}^{\times})$ . Therefore, for any 2-cocycle  $\kappa_q$  representing this class, we have a morphism as in Proposition 3.12:



This cocycle  $\kappa_q$  should be thought of as an analogue of the classical BKS kernel.

Assume there exists a polarization  $A \simeq L \oplus L^{\vee}$ , and that q is given by evaluation:

$$\operatorname{ev} \colon A \simeq L \oplus L^{\vee} \to \mathbf{k}^{\times}$$

*Example* 6.2. Let  $K = L^{\vee}$ . The twice-categorified Fourier transform is the quantization (in the sense of Hypothesis Q) of the correspondence  $L \leftarrow (A, \kappa_{ev}) \rightarrow L^{\vee}$ .

If gL is transverse to L, then we have (6.1) for K = gL:

Fix some isomorphism

$$\iota \colon gL \simeq A/L = L^{\vee} \ .$$

This allows us to rewrite (6.3) as:

(6.4) 
$$(L^2, \kappa_{q,g})$$

where  $\kappa_{q,g}$  is a 2-cocycle on  $L^2$  representing the class in  $H^2_{\text{Grp}}(L^2, \mathbf{k}^{\times})$  determined (by virtue of the Schur isomorphism (A.2)) by the map  $L^2 \to \mathbf{k}^{\times}$  sending

$$(\ell, k) \mapsto \iota \circ g(\ell)(k)$$
.

Now we will specialize to the case when L is a vector space over a finite field  $\mathbb{F}_p$ . Recall the following theorem.

**Theorem** (Cartan–Dieudonné theorem). Let (A,q) be an 2*m*-dimensional, non-degenerate quadratic space over a field of characteristic  $\neq 2$ . Every element of O(A,q) must be the composition of at most 2*m* reflections.

**Lemma 6.5.** Let  $g \in O(L \oplus L^{\vee}, ev)$  be a reflection. The image gL of a Lagrangian under g is transverse to L.

*Proof.* By definition g preserves Lagrangians, so this follows from  $gL \neq L$ . If gL = L, then g is the reflection over either L or its orthogonal complement, neither of which are orthogonal transformations.

Summarizing, to each reflection  $g \in O(L \oplus L^{\vee}, ev)$  we have defined a morphism as in Proposition 3.12:



6.2. Analogue of the intertwining operators. Continue to assume  $A \simeq L \oplus L^{\vee}$  is a vector space over a finite field. Recall the quantization map Sum<sub>3</sub> from Hypothesis Q and write  $C_{\text{ev},g}$  for the correspondence in (6.6). Then define a map:

$$f: O(L \oplus L^{\vee}) \to \pi_0 \operatorname{Aut}_{\mathbf{Fus}} (\operatorname{Vect} [L])$$

by sending a reflection g to:

(6.7) 
$$f(g) \coloneqq \operatorname{Sum}_3(C_{\operatorname{ev},q})$$

The bimodule category f(g) has the following explicit description. The underlying category is **Vect** and the action of **Vect** [L] on either side is trivial. The only nontrivial data is the middle associator, which is determined by

$$[\kappa_{\mathrm{ev},g}] \in \mathrm{Hom}\left(L \oplus L, \mathbf{k}^{\times}\right) \subset H^2\left(L^2, \mathbf{k}^{\times}\right)$$

We have a canonical splitting

$$s: O(A,q) \to \pi_{\leq 1} \operatorname{Aut}_{\mathbf{EqBr}} (\mathcal{Z})$$

(see Section 2.4.2). In [ENO10, §5.2] an equivalence

$$\Psi \colon \operatorname{Aut}_{\mathbf{EqBr}}(\mathcal{Z}) \xrightarrow{\sim} \pi_{\leq 1} \operatorname{Aut}_{\mathbf{Fus}}(\mathcal{C})$$

is provided.

**Theorem 6.8.** Assume Hypothesis Q. Let  $A \simeq L \oplus L^{\vee}$  be a vector space over a finite field. Sending a reflection  $g \in O(A, q)$  to the intertwining bimodule f(g) from (6.7) defines the same equivalence as the composition:

(6.9) 
$$O(A,q) \xrightarrow{s} \pi_0 \operatorname{Aut}_{\mathbf{EqBr}}(\mathcal{Z}) \xrightarrow{\Psi} \pi_0 \operatorname{Aut}_{\mathbf{Fus}}(\mathcal{C})$$

*Proof.* The equivalence (6.9) is given a description in [ENO10, §10] via the hyperbolic part of the category of Lagrangian correspondences.

Namely, they assign the following Vect  $[L^2]$ -module category to  $g \in O(L \oplus L^{\vee}, ev)$ . Recall Vect  $[L^2]$ -module categories are classified by a subgroup  $H \subset L^2$  and a class in  $H^2(L^2, \mathbf{k}^{\times})$ , or equivalently (according to (A.1)) an alternating bicharacter on  $L^2$ . The subgroup H is the projection of the graph of g to  $L^2 \subset (L \oplus L^{\vee})^2$ . The alternating bicharacter  $\chi_g$  is defined by:

$$\chi_g((\ell_1, k_1), (\ell_2, k_2)) \left\langle (\widetilde{\ell_1, k_1}), (\ell_2, 1, k_2, 1) \right\rangle_{\text{ev}_{L^2}}$$

where  $(\widetilde{\ell_1, k_2})$  denotes an arbitrary lift of  $(\ell_1, k_2)$  to the graph of g in  $(L \oplus L^{\vee})^2$ .

In particular, this bicharacter works out to be:

$$\chi_g \left( (\ell_1, k_1), (\ell_2, k_2) \right) = \left\langle (\widetilde{\ell_1, k_1}), (\ell_2, 1, k_2, 1) \right\rangle$$
  
=  $\operatorname{ev}_{L^2} \left( (\ell_1 + \ell_2, \varphi, k_1 + k_2, \psi) \right)$   
=  $\varphi \left( \ell_2 \right) \psi \left( k_2 \right) ,$ 

where  $\varphi \in L^{\vee}$  is such that

$$-k_1 = p_3 g\left(\ell_1, \varphi\right)$$

where  $p_3$  denotes the projection from the graph of g to the third factor L, and

$$\psi \coloneqq p_4 g\left(\ell_1, \varphi\right)$$
.

Now using Lemma A.3, this defines a bimodule with precisely the same middle associator as c(g) defined above.

# APPENDIX A. BICHARACTERS AND THE SCHUR MULTIPLIER

Let  $\wedge^2(-)$  denote the group of alternating bicharacters, and let  $Z^2(-, \mathbf{k}^{\times})$  denote the group of degree two group cocycles. Let A be an abelian group. There is a map

alt: 
$$Z^{2}(A, \mathbf{k}^{\times}) \to \wedge^{2}(A)$$

given by sending a cocycle  $\alpha \in Z^2(A)$  to the alternating bicharacter sending:

$$(a_1, a_2) \mapsto \alpha (a_1, a_2) \alpha (a_2, a_1)^{-1}$$

It is a well-known fact that this induces an equivalence:

(A.1) 
$$\operatorname{alt}: H^2(A, \mathbf{k}^{\times}) \xrightarrow{\sim} \wedge^2 A$$
.

See e.g. [NN08, 2.4].

If  $A = A_1 \oplus A_2$ , recall the Schur isomorphism:

(A.2) 
$$H^{2}(A) \simeq H^{2}(A_{1}) \oplus H^{2}(A_{2}) \oplus \operatorname{Hom}\left(A_{1} \oplus A_{2}, \mathbf{k}^{\times}\right)$$

See e.g. [Kar87, Theorem 2.2.10].

There is an equivalence of Vect  $[L^2]$ -modules and Vect [L]-bimodules sending a module to the same underlying category, with bimodule structure:

$$\mathbf{k}_{\ell} \otimes (-) \otimes \mathbf{k}_{k} = \mathbf{k}_{(\ell,-k)} \otimes (-)$$
.

**Lemma A.3.** Under this equivalence, Vect with associator  $[\alpha] \in H^2(L^2)$  gets sent to Vect with middle associator

$$(\ell, k) \mapsto (\operatorname{alt} \alpha) ((\ell, 0), (0, k))$$

*Proof.* Let  $\alpha \in Z^2(L^2)$  be the associator for a Vect  $[L^2]$ -module category with underlying category Vect. The middle associator for the associated Vect [L]-bimodule category is an equivalence between

$$(\mathbf{k}_{\ell_1} \otimes -) \otimes \mathbf{k}_{k_2} = \mathbf{k}_{(0,-k_2)} \otimes (\mathbf{k}_{(\ell_1,0)} \otimes -)$$

and

$$\mathbf{k}_{\ell_1} \otimes (- \otimes \mathbf{k}_{k_2}) = \mathbf{k}_{(\ell_1,0)} \otimes \left( \mathbf{k}_{(0,-k_2)} \otimes - \right) \;.$$

Because A is abelian, this is provided by the following composition:

	-	-
L		
L		
-	-	

### APPENDIX B. TQFTS AND THE COBORDISM HYPOTHESIS

The axioms for functorial quantum field theory originated in [Ati88, Seg04]. In this work, we will use TQFT to refer to *fully-extended functorial topological quantum field theories*, which first appeared in [BD95]. These were further studied in [Lur09]. Other references include [Fre93, Wal06, Tel16, SP14, Kap10].

The Cobordism Hypothesis was formulated by Baez-Dolan in [BD95]. A detailed sketch of the proof was given in [Lur09], and an approach for a proof using factorization homology is in [AF17]. A proof in two dimensions is in [SP09]. See [Fre13, SP14] for other references.

Let  $d \in \mathbb{Z}^{\geq 0}$ . Given a symmetric-monoidal  $(\infty, d+1)$ -category  $\mathcal{T}$ , which will be the target for our theories, write  $\mathcal{T}^{\sim}$  for the groupoid obtained by discarding all non-invertible morphisms at all levels. We will assume  $\mathcal{T}$  has duals.<sup>8</sup> We will also assume that  $\Omega^{d+1}\mathcal{T} = \mathbf{k}$  for  $\mathbf{k}$  an algebraically closed field of characteristic zero. Throughout the paper, we will write **Vect** for the **k**-linear category of finite-dimensional vector spaces over  $\mathbf{k}$ .

Remark B.1. Besides having duals and satisfying  $\Omega^{d+1}\mathcal{T} = \mathbf{k}$ , we will occasionally assume that  $\mathcal{T}$  satisfies an additional hypothesis. In particular, when we discuss module structures in the sense of [FMT22], we will assume that the target  $\mathcal{T}$  is sufficiently additive to support quantization of  $\pi$ -finite spaces in the sense of [FHLT10, §8] and [FMT22, §A] (this is Hypothesis Q).

The (framed) Cobordism Hypothesis, [Lur09, Theorem 2.4.6], asserts that, since  $\mathcal{T}$  has duals, the functor given by evaluation on the point is an equivalence:

$$\operatorname{Fun}^{\otimes}\left(\operatorname{\mathbf{Bord}}_{d+1}^{\operatorname{fr}}, \mathcal{T}\right) \xrightarrow{\sim} \mathcal{T}^{\sim}$$
.

Remark B.2. The category Fun<sup> $\otimes$ </sup> (**Bord**,  $\mathcal{T}$ ) consists of symmetric monoidal functors with morphisms given by strong natural transformations (in the sense of [JFS17]). A priori this is an  $(\infty, d+1)$ -category, but the statement of the Cobordism Hypothesis is that it is actually an  $(\infty, 0)$ -category, and in particular it is equivalent to the  $(\infty, 0)$ -category  $\mathcal{T}^{\sim}$ .

We will occasionally use the variant of  $\operatorname{Fun}^{\otimes}(\operatorname{\mathbf{Bord}},\mathcal{T})$  which has the same objects, however the 1-morphisms are lax natural transformations. We will write this as  $\operatorname{Fun}^{\operatorname{lax}}(\operatorname{\mathbf{Bord}},\mathcal{T})$ . See Remark B.10.

B.1. The Cobordism Hypothesis for X-theories. There is a refined version of the Cobordism Hypothesis obtained as follows. We follow [Lur09]. Let X be a topological space with real rank d+1 vector bundle  $\zeta$ , and let M be a manifold of dimension  $m \leq d+1$ . An  $(X, \zeta)$ -structure on M consists of

- (i) a continuous map  $f: M \to X$ , and
- (ii) an isomorphism of bundles  $TM \oplus \mathbb{R}^{d+1-m} \simeq f^* \zeta$ .

Write  $\mathbf{Bord}_{d+1}^{(X,\zeta)}$  for the category of bordisms equipped with an  $(X,\zeta)$ -structure. This is [Lur09, Definition 2.4.17]. We will refer to symmetric monoidal functors out of  $\mathbf{Bord}_{d+1}^{(X,\zeta)}$  as (d+1)-dimensional  $(X,\zeta)$ -TQFTs, or just  $(X,\zeta)$ -theories. The Cobordism Hypothesis for  $(X,\zeta)$ -theories is [Lur09, Theorem 2.4.18]. It is the equivalence

(B.3) 
$$\operatorname{Fun}^{\otimes} \left( \operatorname{Bord}_{d+1}^{(X,\zeta)}, \mathcal{T} \right) \xrightarrow{\sim} \operatorname{Hom} \left( \widetilde{X}, \mathcal{T}^{\sim} \right) ,$$

where  $\widetilde{X}$  is the associated principal O (d+1)-bundle of orthonormal frames in  $\zeta$ .

Let G be a topological group with continuous group homomorphism to the orthogonal group  $\chi: G \to O(d+1)$ . Particularly important cases of  $(X, \zeta)$ -structures are given by X = BG and  $\zeta = (\mathbb{R}^{d+1} \times EG) / G$  defined by  $\chi$ . In this case, we sometimes call theories defined on  $\mathbf{Bord}_{d+1}^{BG}$  G-theories.

<sup>&</sup>lt;sup>8</sup>Otherwise replace  $\mathcal{T}$  with the subcategory consisting of the fully-dualizable objects of  $\mathcal{T}$ .

*Example* B.4. Let G be trivial and  $\chi = \iota$  be the inclusion. A  $(BG, \zeta_{\iota})$ -structure is a framing. *Example* B.5. Let G = SO(d+1) and  $\chi = \iota$  be the inclusion. A  $(B SO(d+1), \zeta_{\iota})$ -structure is an orientation.

We will write  $\mathbf{Bord}_{d+1}^X$  for the category of bordisms with  $(X, \zeta)$ -structure with  $\zeta$  trivial. E.g. X = BG and  $\chi: G \to O(d+1)$  trivial. In spite of fr being removed from the notation, one should think of  $\mathbf{Bord}_{d+1}^X$  as consisting of framed bordisms with a map to X. We will refer to symmetric monoidal functors  $\mathbf{Bord}_{d+1}^X \to \mathcal{T}$  as X-TQFTs (or just X-theories). These should be thought of as families of framed TQFTs over X. In this case, the Cobordism Hypothesis (B.3) reduces to:

(B.6) 
$$\operatorname{Fun}^{\otimes} \left( \operatorname{\mathbf{Bord}}_{d+1}^X, \mathcal{T} \right) \xrightarrow{\sim} \operatorname{Hom} \left( X, \mathcal{T}^{\sim} \right) \ .$$

*Example* B.7. When X = BG with  $\chi = \text{triv}: G \to O(d+1)$  trivial, then the Cobordism Hypothesis reduces to [Lur09, Theorem 2.4.26]:

$$\operatorname{Fun}^{\otimes} \left( \operatorname{\mathbf{Bord}}_{d+1}^{BG}, \mathcal{T} \right) \xrightarrow{\sim} \left( \mathcal{T}^{\sim} \right)^{hG}$$

where  $(\mathcal{T})^{hG}$  denotes the homotopy fixed points of  $\mathcal{T}$ .

We will sometimes call symmetric monoidal functors  $\mathbf{Bord}_{d+1}^{BG} \to \mathcal{T}$  BG-theories, or G-theories.

Endomorphisms of the trivial TQFT are equivalent to theories of one lower dimension. By the Cobordism Hypothesis this is equivalent to endomorphisms of the identity in the target being decategorification. Proposition B.9 is this result for X-theories.

**Lemma B.8.** If A and B are monoidal  $(\infty, d+1)$ -categories, then

$$\Omega \operatorname{Fun}^{\otimes}(\mathcal{A},\mathcal{B}) \simeq \operatorname{Fun}^{\otimes}(\mathcal{A},\Omega\mathcal{B})$$

*Proof.* The LHS consists of natural transformations:

$$egin{array}{cccc} \mathcal{A} & \longrightarrow 1 \ & & \downarrow \ & & \downarrow \ 1 & \longrightarrow \mathcal{B} \end{array}$$

where **1** denotes the trivial monoidal category. The category  $\Omega \mathcal{B}$  is the pullback of the diagram  $\mathbf{1} \rightarrow \mathcal{B} \leftarrow \mathbf{1}$ , so the result follows from the universal property.

**Proposition B.9.** If  $1_X^{d+1}$  is the trivial functor  $\mathbf{Bord}_{d+1}^X \to \mathcal{T}$ , then

$$\operatorname{Fun}^{\otimes}\left(\mathbf{Bord}_{d}^{X}, \Omega \mathcal{T}\right) \simeq \operatorname{End}\left(1_{X}^{d+1}\right) \;,$$

where  $\Omega \mathcal{T} \coloneqq \operatorname{End}_{\mathcal{T}}(1)$  denotes the looping of  $\mathcal{T}$ .

*Proof.* This follows from Lemma B.8 for  $\mathcal{A} = \mathbf{Bord}_d^X$  and  $\mathcal{B} = \mathcal{T}$ .

B.2. Relative theories. A symmetric monoidal functor

$$\alpha \colon \mathbf{Bord}_d^X \to \mathcal{T}$$

is a *once-categorified d-dimensional X-TQFT*, where  $\mathcal{T}$  is the fixed target from the beginning of Appendix B. Let

$$1_X \colon \mathbf{Bord}_d^X \to \mathcal{T}$$

denote the trivial once-categorified *d*-dimensional X-TQFT. Recall the notion of a relative theory [FT14]. These are also called *twisted theories* [ST11, JFS17].

**Definition.** A theory defined right-relative to  $\alpha$  is a lax natural transformation (in the sense of [JFS17])

$$\alpha \to 1$$
 .

A theory defined left-relative to  $\alpha$  is a lax natural transformation

$$1 \rightarrow \alpha$$
 .

Remark B.10. Recall the definition of lax (resp. oplax) natural transformations from [JFS17]. Consider the arrow categories  $\mathcal{T}^{\downarrow}$  and  $\mathcal{T}^{\rightarrow}$ , and the source and target functors  $s, t: \mathcal{T}^* \rightarrow \mathcal{T}$  for  $* = \downarrow, \rightarrow$ . Following [JFS17], a lax (resp. oplax) natural transformation  $\alpha \rightarrow 1$  is a functor

(B.11) 
$$F_{\alpha} \colon \mathbf{Bord}_{d}^{X} \to \mathcal{T}^{\downarrow}$$
  $\left( \operatorname{resp.} F_{\alpha} \colon \mathbf{Bord}_{d}^{X} \to \mathcal{T}^{\rightarrow} \right)$ 

such that  $s \circ F_{\alpha} = \alpha$ , and  $t \circ F_{\alpha} = 1$ .

Throughout, we will use the lax version, as written in the above definition of relative theories. The reason we use the lax version, as noted in [JFS17, Example 7.3], that the *lax* natural transformations from the trivial theory to itself consist of theories of dimension lower ([JFS17, Theorem 7.4]) whereas the same is not true when lax is replaced with oplax. We need the analogous result for X-theories (Proposition B.9), in particular to establish a trivialized anomalous theory  $1 \xrightarrow{\sim} \alpha \rightarrow 1$  as an honest theory of one dimension lower.

Also noted in [JFS17, Example 7.3], is the fact that oplax natural transformations are "elements"  $F_{\sigma}(M) : 1 \to \alpha(M)$  for M a *closed* bordism of any codimension, which is for example the point of view taken in [FT14]. Oplax natural transformations are also used in [FT21]. Besides Proposition B.9, and the results depending on it, the remaining constructions and facts in this paper hold for the oplax version, obtained my replacing all lax natural transformations with oplax ones, and replacing  $\mathcal{T}^{\downarrow}$  with  $\mathcal{T}^{\rightarrow}$ .

*Remark* B.12. Often the once-categorified d-dimensional theory  $\alpha$  extends to a (d + 1)-dimensional theory:



In this case, the relative theory  $\alpha \to 1$  is upgraded to what is called a *boundary theory*. Boundary theories are defined as functors out of the extended bordism category  $\mathbf{Bord}_{d+1}^{X,\partial}$ , described in [Lur09, §4.3]. The connection with the notion defined here is made in [JFS17, Theorem 7.15].

#### References

- [AF17] David Ayala and John Francis, *The cobordism hypothesis*, 2017. 29
- [Ati88] Michael Atiyah, Topological quantum field theories, Inst. Hautes Études Sci. Publ. Math. (1988), no. 68, 175–186 (1989). MR 1001453 29
- [BD95] John C. Baez and James Dolan, Higher-dimensional algebra and topological quantum field theory, J. Math. Phys. 36 (1995), no. 11, 6073–6105. MR 1355899 29
- [BJS21] Adrien Brochier, David Jordan, and Noah Snyder, On dualizability of braided tensor categories, Compos. Math. 157 (2021), no. 3, 435–483. MR 4228258 6
- [BW97] Sean Bates and Alan Weinstein, Lectures on the geometry of quantization, Berkeley Mathematics Lecture Notes, vol. 8, American Mathematical Society, Providence, RI; Berkeley Center for Pure and Applied Mathematics, Berkeley, CA, 1997. MR 1806388 2
- [BZFN10] David Ben-Zvi, John Francis, and David Nadler, Integral transforms and Drinfeld centers in derived algebraic geometry, J. Amer. Math. Soc. 23 (2010), no. 4, 909–966. MR 2669705 3
- [CGPW16] Shawn X. Cui, César Galindo, Julia Yael Plavnik, and Zhenghan Wang, On gauging symmetry of modular categories, Comm. Math. Phys. 348 (2016), no. 3, 1043–1064. MR 3555361 13, 22

32	JACKSON VAN DYKE
[Del99] [DGNO10]	Pierre Deligne, <i>Notes on spinors</i> , Quantum fields and strings: a course for mathematicians, Vol. 1, 2 (Princeton, NJ, 1996/1997), Amer. Math. Soc., Providence, RI, 1999, pp. 99–135. MR 1701598 2 Vladimir Drinfeld, Shlomo Gelaki, Dmitri Nikshych, and Victor Ostrik, <i>On braided fusion categories</i> .
	I, Selecta Math. (N.S.) 16 (2010), no. 1, 1–119. MR 2609644 8
[DR18]	Christopher L. Douglas and David J. Reutter, Fusion 2-categories and a state-sum invariant for 4-manifolds, 2018. 15
[EG18]	Pavel Etingof and César Galindo, <i>Reflection fusion categories</i> , J. Algebra <b>516</b> (2018), 172–196. MR 3863475 3, 13
[EGNO15]	Pavel Etingof, Shlomo Gelaki, Dmitri Nikshych, and Victor Ostrik, <i>Tensor categories</i> , Mathemat- ical Surveys and Monographs, vol. 205, American Mathematical Society, Providence, RI, 2015. MR 3242743 9
[EM46]	Samuel Eilenberg and Saunders MacLane, Determination of the second homology and cohomology groups of a space by means of homotopy invariants, Proc. Nat. Acad. Sci. U.S.A. <b>32</b> (1946), 277–280. MR 19307 15
[ENO10]	Pavel Etingof, Dmitri Nikshych, and Victor Ostrik, <i>Fusion categories and homotopy theory</i> , Quantum Topol. <b>1</b> (2010), no. 3, 209–273, With an appendix by Ehud Meir. MR 2677836 3, 4, 5, 6, 7, 8, 10, 12, 13, 20, 21, 22, 23, 24, 25, 27
[ENO11]	., Weakly group-theoretical and solvable fusion categories, Adv. Math. <b>226</b> (2011), no. 1, 176–205 MR 2735754 7
[EO04]	Pavel Etingof and Viktor Ostrik, <i>Finite tensor categories</i> , Mosc. Math. J. 4 (2004), no. 3, 627–654, 782–783. MB 2119143 7, 8
[FHLT10]	Daniel S. Freed, Michael J. Hopkins, Jacob Lurie, and Constantin Teleman, <i>Topological quantum field theories from compact Lie groups</i> , A celebration of the mathematical legacy of Raoul Bott, CRM Proc. Lecture Notes, vol. 50, Amer. Math. Soc., Providence, RI, 2010, pp. 367–403.
[FMT22]	MR 2648901 2, 5, 13, 14, 15, 16, 29 Daniel S. Freed, Gregory W. Moore, and Constantin Teleman, <i>Topological symmetry in quantum field theory</i> , 2022. 2, 3, 4, 5, 13, 14, 15, 16, 29
[Fre93]	Daniel S. Freed, <i>Lectures on topological quantum field theory</i> , Integrable systems, quantum groups, and quantum field theories (Salamanca, 1992), NATO Adv. Sci. Inst. Ser. C: Math. Phys. Sci., vol. 409, Kluwer Acad, Publ., Dordrecht, 1993, pp. 95–156, MR 1262837, 29
[Fre94]	, Higher algebraic structures and quantization, Comm. Math. Phys. <b>159</b> (1994), no. 2, 343– 398. MB 1256993 2, 13
[Fre13]	, <i>The cobordism hypothesis</i> , Bull. Amer. Math. Soc. (N.S.) <b>50</b> (2013), no. 1, 57–92. MR 2904995 29
[FT14]	Daniel S. Freed and Constantin Teleman, <i>Relative quantum field theory</i> , Comm. Math. Phys. <b>326</b> (2014) no. 2, 459–476, MB 3165462 30, 31
[FT21]	
[GH09]	Shamgar Gurevich and Ronny Hadani, <i>Quantization of symplectic vector spaces over finite fields</i> , I. Symplectic Geom 7 (2009) no. 4, 475–502 MR 2552002 13
[GH11]	Shamgar Gurevich and Ronny Hadani, <i>The categorical weil representation</i> , 2011. 13
[GJ19]	Terry Gannon and Corey Jones, Vanishing of categorical obstructions for permutation orbifolds, Comm. Math. Phys. <b>369</b> (2019), no. 1, 245–259. MR 3959559 13
[Hau17]	Rune Haugseng, The higher Morita category of $\mathbb{E}_n$ -algebras, Geom. Topol. <b>21</b> (2017), no. 3, 1631–1730. MR 3650080 6
[Jen11]	Evan Jenkins, Extensions of groups by braided 2-groups, 2011. 13
[JFS17]	Theo Johnson-Freyd and Claudia Scheimbauer, ( <i>Op</i> )lax natural transformations, twisted quantum field theories, and "even higher" Morita categories, Adv. Math. <b>307</b> (2017), 147–223. MR 3590516
[JS93]	André Joyal and Ross Street, Braided tensor categories, Adv. Math. <b>102</b> (1993), no. 1, 20–78. MR 1250465 0
[Kap10]	Anton Kapustin, <i>Topological field theory, higher categories, and their applications</i> , Proceedings of the International Congress of Mathematicians. Volume III, Hindustan Book Agency, New Delhi, 2010, pp. 2021–2043, MR 2827874 29
[Kar87]	Gregory Karpilovsky, <i>The Schur multiplier</i> , London Mathematical Society Monographs. New Series, vol. 2. The Clarendon Press, Oxford University Press, New York, 1987, MR 1200015, 28
[KK12]	Alexei Kitaev and Liang Kong, Models for gapped boundaries and domain walls, Comm. Math. Phys. <b>313</b> (2012), no. 2, 351–373. MR 2942952 7

- [Kon88] Maxim Kontsevich, Rational conformal field theory and invariants of 3-dimensional manifolds, CNRS Luminy preprint (1988). 2, 13
- [KT14] Anton Kapustin and Ryan Thorngren, Anomalies of discrete symmetries in various dimensions and group cohomology, 2014. 4, 25
- [KZ18] Liang Kong and Hao Zheng, The center functor is fully faithful, Adv. Math. 339 (2018), 749–779. MR 3866911 7
- [Lur09] Jacob Lurie, On the classification of topological field theories, Current developments in mathematics, 2008, Int. Press, Somerville, MA, 2009, pp. 129–280. MR 2555928 29, 30, 31
- [LV80] Gérard Lion and Michèle Vergne, The Weil representation, Maslov index and theta series, Progress in Mathematics, vol. 6, Birkhäuser, Boston, Mass., 1980. MR 573448 2
- [MN18] Ian Marshall and Dmitri Nikshych, On the Brauer-Picard groups of fusion categories, Math. Z. 288 (2018), no. 3-4, 689–712. MR 3778972 9
- [MP12] J. P. May and K. Ponto, More concise algebraic topology, Chicago Lectures in Mathematics, University of Chicago Press, Chicago, IL, 2012, Localization, completion, and model categories. MR 2884233 10
- [MS20] Lukas Müller and Richard J. Szabo, 't Hooft anomalies of discrete gauge theories and non-abelian group cohomology, Comm. Math. Phys. 375 (2020), no. 3, 1581–1627. MR 4091494 4, 25
- [Mü20] Lukas Müller, Extended functorial field theories and anomalies in quantum field theories, 2020. 4, 25
- [NN08] Deepak Naidu and Dmitri Nikshych, Lagrangian subcategories and braided tensor equivalences of twisted quantum doubles of finite groups, Comm. Math. Phys. 279 (2008), no. 3, 845–872. MR 2386730 28
- [Pol03] Alexander Polishchuk, Abelian varieties, theta functions and the Fourier transform, Cambridge Tracts in Mathematics, vol. 153, Cambridge University Press, Cambridge, 2003. MR 1987784 2
- [Pos55] M. M. Postnikov, Investigations in homotopy theory of continuous mappings, Izdat. Akad. Nauk SSSR, Moscow, 1955, Trudy Mat. Inst. Steklov. no. 46. MR 0074819 10
- [Qui95] Frank Quinn, Lectures on axiomatic topological quantum field theory, Geometry and quantum field theory (Park City, UT, 1991), IAS/Park City Math. Ser., vol. 1, Amer. Math. Soc., Providence, RI, 1995, pp. 323–453. MR 1338394 2, 13
- [Sch14] Claudia I Scheimbauer, Factorization homology as a fully extended topological field theory, Ph.D. thesis, ETH Zurich, 2014. 6
- [Seg04] Graeme Segal, The definition of conformal field theory, Topology, geometry and quantum field theory, London Math. Soc. Lecture Note Ser., vol. 308, Cambridge Univ. Press, Cambridge, 2004, pp. 421–577. MR 2079383 29
- [SP09] Christopher John Schommer-Pries, The classification of two-dimensional extended topological field theories, ProQuest LLC, Ann Arbor, MI, 2009, Thesis (Ph.D.)–University of California, Berkeley. MR 2713992 29
- [SP14] Christopher J. Schommer-Pries, Dualizability in low-dimensional higher category theory, Topology and field theories, Contemp. Math., vol. 613, Amer. Math. Soc., Providence, RI, 2014, pp. 111–176. MR 3221292 29
- [ST11] Stephan Stolz and Peter Teichner, Supersymmetric field theories and generalized cohomology, Mathematical foundations of quantum field theory and perturbative string theory, Proc. Sympos. Pure Math., vol. 83, Amer. Math. Soc., Providence, RI, 2011, pp. 279–340. MR 2742432 30
- [Tel16] Constantin Teleman, Five lectures on topological field theory, Geometry and quantization of moduli spaces, Adv. Courses Math. CRM Barcelona, Birkhäuser/Springer, Cham, 2016, pp. 109–164. MR 3675464 29
- [Toë07] Bertrand Toën, The homotopy theory of dg-categories and derived Morita theory, Invent. Math. 167 (2007), no. 3, 615–667. MR 2276263 3
- [Tur10] Vladimir Turaev, Homotopy quantum field theory, EMS Tracts in Mathematics, vol. 10, European Mathematical Society (EMS), Zürich, 2010, Appendix 5 by Michael Müger and Appendices 6 and 7 by Alexis Virelizier. MR 2674592 2, 13
- [TY98] Daisuke Tambara and Shigeru Yamagami, Tensor categories with fusion rules of self-duality for finite abelian groups, J. Algebra 209 (1998), no. 2, 692–707. MR 1659954 13
- [VD23] Jackson Van Dyke, Projective symmetries of three-dimensional TQFTs, 2023, arXiv: 2311.01637 [math.QA]. 2, 3, 16, 19, 20, 24, 25
- [Wal06] Kevin Walker, Tqfts, https://canyon23.net/math/tc.pdf, 2006. 29
- [Whi78] George W. Whitehead, Elements of homotopy theory, Graduate Texts in Mathematics, vol. 61, Springer-Verlag, New York-Berlin, 1978. MR 516508 10