

Subsystem eigenstate thermalization hypothesis for translation invariant systems

Zhiqiang Huang (黄志强)^{1,*} and Xiao-Kan Guo (郭肖侃)²

¹*State Key Laboratory of Magnetic Resonance and Atomic and Molecular Physics,
Innovation Academy for Precision Measurement Science and Technology,
Chinese Academy of Sciences, Wuhan 430071, China*

²*Department of Applied Mathematics, Yancheng Institute of Technology, Jiangsu 224051, China*
(Dated: December 4, 2023)

The eigenstate thermalization hypothesis for translation invariant quantum spin systems has been proved recently by using random matrices. In this paper, we study the stronger subsystem version of eigenstate thermalization hypothesis for translation invariant quantum systems without referring to random matrices. By showing the small upper bounds on the quantum variance or the Belavkin-Staszewski relative entropy, we prove the subsystem eigenstate thermalization hypothesis for translation invariant quantum systems in an elementary way.

I. INTRODUCTION

The equilibration and the thermalization of an isolated quantum system are fundamental for understanding the emergence of quantum statistical mechanics from unitary quantum mechanics. By thermalization, it means that either an isolated quantum system would evolve into a thermal state, or the observables would attain their values in a statistical ensemble, after a unitary quantum evolution of the isolated quantum system for a period of time that is long enough. Since a unitary quantum evolution preserves the pure state, it is not easy to understand how the statistical mixture emerges if the initial state of an isolated quantum system is a pure state. Numerous approaches have been proposed to understand various aspects of this problem, cf. the reviews [1, 2].

The eigenstate thermalization hypothesis (ETH) [3, 4], that the expectation values of quantum observables in an energy eigenstate should approximately coincide with the thermal expectation values, provides a possible mechanism for the thermalization of an isolated quantum system. Although the ETH has more and more numerical and experimental evidences in specific closed quantum models/systems, its physical origin and mathematical description are not completely understood by now. In the original proposal by Deutsch and Srednicki [5–7], a random perturbation is added to a closed quantum system, and the ETH holds if the perturbed system becomes chaotic. By modeling the random perturbations as random matrices, the ETH for deterministic observables with the Hamiltonians sampled from the Wigner random matrix ensemble without further unitary symmetry is mathematically proved in the recent work [8]. This scenario, however, is not universal. For one thing, if further unitary symmetries are present, the conserved quantities would obstruct the thermalization to Gibbs states and the original ETH would fail. More recently in [9], the ETH for translation invariant spin systems is proved using the same method from random matrices,

thereby generalizing its validity to various translation invariant lattice spin models.

In many studies of the “weak” ETH, for example, [10–13], one does not presume the random energy perturbations, or simply the random Hamiltonians, but tries to derive the statistical properties solely from quantum properties. From this perspective, the quantum entanglement inside a closed quantum system, together with its dynamics under the global unitary evolution, should play a crucial role for thermalization, which has indeed been experimentally observed in [14]. To quantify the entanglement in a closed quantum system, we need to work at the level of subsystems of the total system to compute the entanglement entropies and alike. This observation leads to the *subsystem* ETH [15, 16], which hypothesizes the convergence of the subsystem density matrices to the thermal Gibbs density matrix. In fact, the trace distance between two density matrices is bounded by the relative entropy between two density matrices. Since the entanglement entropies and relative entropies are calculable in many conformal field theories (CFT), the subsystem ETH and its violation have been tested in many CFTs [16–22]. Notice that the conformal symmetry forms an infinite-dimensional group, so the infinite number of conserved KdV charges make the generalized Gibbs states as the proper equilibrated states for CFTs [23, 24]. It is then natural to ask for a quantum system/model with a smaller symmetry group such that the subsystem ETH still holds.

For translation invariant quantum lattice systems, we already know that the strong ETH [9], the weak ETH [11, 12], and the canonical typicality [25] are true. In addition, a version of the generalized ETH, i.e. thermalization to the generalized Gibbs ensemble, for translation invariant quasi-free fermionic integrable models is also proved in [26]. We therefore see that the translation invariant quantum lattice systems are good tested for checking various versions of ETH. In this paper, we make an effort to prove the subsystem ETH for translation invariant systems without referring to random matrices.

We will work in the setting of translation invariant

* hzq@wipm.ac.cn

quantum lattice system in the sense of [12]. Unlike the considerations by Iyoda *et al.* [12], we find a relation between the quantum variance and the Belavkin-Staszewski relative entropy in an average sense, thereby establishing a connection of the scaling analysis on the variance given in [12] and the subsystem ETH formulated as the relative-entropic bounds on the trace distance between the subsystem state and the canonical thermal state. In fact, we are able to prove the following form of subsystem ETH

$$\|\rho_{\text{sub}} - \rho^c\| \leq 2\hat{S}(\rho_{\text{sub}}\|\rho^c) \sim \mathcal{O}(N^{-\alpha}), \quad (1)$$

$$\|\rho_{\text{sub}} - \rho'_{\text{sub}}\| \sim \mathcal{O}(N^{-\alpha}), \quad (2)$$

where ρ_{sub} is the state of a subsystem, ρ^c is the canonical thermal state, and $0 < \alpha < 1$, for translation invariant quantum lattice systems. Since the Belavkin-Staszewski relative entropy $\hat{S}(\rho_{\text{sub}}\|\rho^c)$ is greater than or equal to the quantum relative entropy, a small $\hat{S}(\rho_{\text{sub}}\|\rho^c)$ entails the relative-entropic bound on subsystem ETH considered in [15, 16].

We begin in Section II with some preliminary results about ETH, subsystem ETH, and in particular the setting of translation invariant quantum lattice system from [12]. In Section III, we introduce the main technical input, i.e. the relation between the quantum variance and the Belavkin-Staszewski relative entropy in an average sense. Using this relation, we analyze the scaling of both the variance and the Belavkin-Staszewski relative entropy and prove the subsystem ETH. In the final Section V we conclude this paper and discuss some related issues.

II. PRELIMINARIES

In this section, we recollect the basics of ETH and subsystem ETH, and the weak ETH with eigenstate typicality in the sense of [12].

A. ETH and subsystem ETH

Consider an isolated or closed quantum system B with Hamiltonian h . This Hamiltonian h could include a random perturbation $h_{\text{pert.}}$. Suppose h has eigenvectors $|E_i\rangle$, $i = 1, 2, \dots, N$ with energy eigenvalues E_i , i.e. $h|E_i\rangle = E_i|E_i\rangle$. For a few-body observable A , the local ETH can be formulated in terms of the expectation values of A in the energy eigenstates as

$$\langle E_i|A|E_j\rangle = \mathcal{A}(E)\delta_{ij} + e^{-S(E)/2}f(E, \omega)R_{ij} \quad (3)$$

where $E = \frac{1}{2}(E_i + E_j)$, $\omega = E_i - E_j$, and $e^{S(E)} = E \sum_i \delta(E - E_i)$ is the density of states of the system B . The $\mathcal{A}(E)$ and $f(E, \omega)$ are smooth functions, while the fluctuation factor R_{ij} is of order 1. Particularly the thermalization requires that $\mathcal{A}(E)$ should be approximately

the thermal average of A in the canonical ensemble, $\mathcal{A} = \langle A \rangle_c + \mathcal{O}(N^{-1}) + \mathcal{O}(e^{-S/2})$, in the large N limit.

This local form (3) of ETH can be derived based on Berry's chaotic conjecture [7]. If we sample the Hamiltonian h from a random matrix ensemble, the following form of inequality for ETH,

$$|\langle E_i|A|E_j\rangle - \langle A \rangle_{\text{mc}}(E)\delta_{ij}| \leq \mathcal{O}(e^{-S/2}), \quad (4)$$

where $\langle \rangle_{\text{mc}}$ denotes the thermal average in the micro-canonical ensemble, can be proved mathematically in several cases, including the translation invariant systems, by using properties of random matrices [8, 9].

Both(3) and (4) are local conditions, as the ETH are assumed for each energy eigenstate. Therefore, in analogy to the canonical typicality of a subsystem B_i , we can envision the *subsystem* ETH,

$$\|\rho_{B_i} - \rho^c(E_i)\| \sim \mathcal{O}(e^{-S/2}), \quad (5)$$

$$\|\rho_{ij}\| \sim \mathcal{O}(e^{-S/2}), \quad (6)$$

where $\rho_{B_i} = \text{Tr}_{\bar{B}_i} |E_i\rangle \langle E_i|$ is the reduced density matrix of the subsystem B_i , ρ^c is a universal density matrix that could be the thermal canonical one, and $\rho_{ij} = \text{Tr}_{\bar{B}_i} |E_i\rangle \langle E_j|$. The norm here refers to the trace distance $\|\rho_1 - \rho_2\| = \frac{1}{2}\text{Tr}\sqrt{(\rho_1 - \rho_2)^2}$. The subsystem ETH as given by (5) and (6) is in fact stronger than the local ETH as in (3), due to the following inequality [16]

$$|\langle A \rangle - \langle A \rangle_c| \leq \sqrt{\|\rho - \rho^c\| \text{Tr}[(\rho + \rho^c)A^2]} \quad (7)$$

where $\langle A \rangle = \text{Tr}(\rho A)$ and $\langle A \rangle_c = \text{Tr}(\rho^c A)$.

What is important in the following is that the trace distance in (5) can be bounded by the relative entropy between two density matrices,

$$\|\rho_{B_i} - \rho^c(E_i)\|^2 \leq 2S(\rho_{B_i}\|\rho^c), \quad (8)$$

where $S(\rho_1\|\rho_2) = \text{tr}(\rho_1 \log \rho_1) - \text{tr}(\rho_1 \log \rho_2)$ is the (Umegaki) quantum relative entropy. This inequality (8) is the so-called quantum Pinsker inequality in quantum information theory [27].

B. Weak ETH with eigenstate typicality

In proving the weak ETH for translation invariant quantum lattice systems [12], the quantum uncertainty of measuring an observable plays an important role. Conventionally, the uncertainties, either classical or quantum, can be quantified by the variance [28]. For instance, given a quantum state ρ , the quantum uncertainty of measuring an observable A in the state ρ can be quantified by the variance

$$V(\rho, A) = \text{Tr}(\rho A^2) - (\text{Tr}\rho A)^2 = \text{Tr}[\rho(A - \langle A \rangle)^2]. \quad (9)$$

Let $\rho_B = \sum_j p_j \Pi_B^j$ be the state of the total system B expanded in the orthonormal basis $\{\Pi_B^j\}$ of rank-1 projectors, then in terms of these projectors one can define

particularly the following quantity, which is called as *fluctuation* in [12],

$$\Delta(\rho, \mathcal{A}) = \sum_j p_j \text{Tr}(\Pi_B^j \mathcal{A})^2 - (\text{Tr} \rho \mathcal{A})^2. \quad (10)$$

We have

$$\Delta(\rho, \mathcal{A}) \leq \sum_{j,k} p_j \text{Tr}(\Pi_B^j \mathcal{A} \Pi_B^k \mathcal{A}) - (\text{Tr} \rho \mathcal{A})^2 = V(\rho, \mathcal{A}), \quad (11)$$

because the additional off-diagonal terms are positive, i.e. $\text{Tr}(\Pi_B^j \mathcal{A} \Pi_B^k \mathcal{A}) = |\langle j | \mathcal{A} | k \rangle|^2 \geq 0$. This $\Delta(\rho, \mathcal{A})$ is related to the following (in)distinguishability measure of quantum states:

$$d(\Pi_B^j, \rho; \mathcal{A}) = |\text{Tr}[(\Pi_B^j - \rho) \mathcal{A}]|^2. \quad (12)$$

Indeed, $\Delta(\rho, \mathcal{A})$ can be considered as the quantification of the probabilistic typicality or concentration with respect to the measure (12),

$$\Delta(\rho, \mathcal{A}) = \int d(\Pi_B^j, \rho; \mathcal{A}) p_d \quad (13)$$

where the probability distribution p_d is obtained from the p_j 's through a Radon-Nykodym derivative. By the Chebyshev inequality, we have that

$$P_\rho(|\text{Tr}(\Pi_B^j \mathcal{A}) - \text{Tr}(\rho \mathcal{A})| \geq \epsilon) \leq \frac{\Delta(\rho, \mathcal{A})^2}{\epsilon^2}, \quad \forall \epsilon \in \mathbb{R}^+. \quad (14)$$

Therefore, when $\Delta(\rho, \mathcal{A})$ is very small, the expectation of the projectively measured observable would concentrate on the expectation of observable calculated with respect to the state ρ . In other words, the indistinguishability of measurement outcomes induced a description by a mixed state.

In ETH, one considers the local energy eigenstates. So, let $\sigma_{B_1}^j = \text{Tr}_{\bar{B}_1} \Pi_B^j$ be the reduced projection/state on the subsystem B_1 . Then we should consider

$$d(\sigma_{B_1}^j, \rho; \mathcal{A}_{B_1}) = |\text{Tr}[(\sigma_{B_1}^j - \rho_{B_1}) \mathcal{A}_{B_1}]|^2 \leq \|\sigma_{B_1}^j - \rho_{B_1}\|_1^2 \|\mathcal{A}_{B_1}\|_\infty^2, \quad (15)$$

where $\|\cdot\|_k$ is the Schatten k -norm. Next, let ρ^{mc} be the density matrix for the microcanonical ensemble. According to eqs. (10), (11) and (14), if

$$\Delta(\rho^{\text{mc}}, \mathcal{A}_{B_1}) \sim \mathcal{O}(N^{-\alpha}), \quad (16)$$

with $0 < \alpha < 1$, i.e. the expectations of a local observable \mathcal{A}_{B_1} with respect to the results of local measurements concentrate the expectation of \mathcal{A}_{B_1} with respect to ρ^{mc} , then we know that each pure state $\sigma_{B_1}^j$ cannot be distinguished from the microcanonical ρ^{mc} in the large N limit. This is the *weak ETH with eigenstate typicality* [12]. Furthermore, by using the equivalence of the ensembles, one

also has a similar weak ETH on the concentration of $\sigma_{B_i}^j$ to the canonical-ensemble density matrix ρ_c .

In the proofs of the weak ETH with eigenstate typicality for quantum lattice systems [12], the translation invariance in the following sense is crucial. Let us partition the lattice of system B into $\mathcal{C} = |B|/|B_1|$ blocks with the same size, where $|B|$ means the number lattice points of in B . These \mathcal{C} blocks are identical copies of B_1 . Let us also define the which are translational copies A^{B_i} of A^{B_1} defined on B_i obtained by the translations from block to block. Then the translation invariance means

$$\text{Tr}[\Pi_B^j A^{B_i}] = \text{Tr}[\Pi_B^j A^{B_1}]. \quad (17)$$

We can introduce the average observable

$$A^B = \frac{1}{\mathcal{C}} \sum_i A^{B_i}, \quad (18)$$

then the translation invariance (17) entails $\Delta(\rho, A^{B_1}) = \Delta(\rho, A^B)$. Therefore, the weak ETH can be proved by bounding $\Delta(\rho, A^B)$.

III. RELATING VARIANCE TO RELATIVE ENTROPY

Eqs. (9) and (12) depend on the measured observable \mathcal{A} . In order to quantify the quantum uncertainty in a way that depends only on the quantum measurements but not on the measured observables, the following entropic uncertainty used in the entropic uncertainty relation [29] serves the purpose,

$$H_\Pi(\rho) = \sum_i p_i S(\rho_i \| \rho) \quad (19)$$

where $\rho_i = \Pi^i \rho \Pi^i / p_i$ with $p_i = \text{Tr}(\Pi^i \rho)$ and $\{\Pi^i\}$ being the (not necessarily rank-1) measurement operators.

In view of the frequent usages of the maps between the total system B and its subsystems B_i in the proofs in [12], we consider the Belavkin-Staszewski (BS) relative entropy [30, 31]

$$\hat{S}(\sigma \| \rho) = \text{Tr}[\sigma \ln(\mathcal{J}_\sigma^{1/2}(\rho^{-1}))] \quad (20)$$

$$= \text{Tr}[\sigma \ln(\rho^{-1} \sigma)] \quad (21)$$

$$= \text{Tr}[\rho \mathcal{J}_\rho^{-1/2}(\sigma) \ln(\mathcal{J}_\rho^{-1/2}(\sigma))] \quad (22)$$

where $\mathcal{J}_\rho^\alpha(\cdot) := \rho^\alpha(\cdot)\rho^\alpha$ is the rescaling map. Since different ρ_i are orthogonal to each other by definition, the entropic uncertainty can be generalized by using (20) as

$$\sum_i p_i \hat{S}(\rho_i \| \rho) = \text{Tr}[\rho \ln(\sum_i \mathcal{J}_{\rho_i}^{1/2}(\rho^{-1}))]. \quad (23)$$

By the power series of the matrix logarithm

$$\ln(X) = (X - I) - \frac{1}{2}(X - I)^2 + \dots, \quad (24)$$

we can obtain the following first-order relations

$$\begin{aligned} \sum_i \text{Tr} \rho [\mathcal{J}_{\rho_i}^{1/2}(\rho^{-1}) - 1] &= \sum_i p_i \left(\text{Tr} \rho_i [\mathcal{J}_{\rho_i}^{1/2}(\rho^{-1})] - 1 \right) \\ &= \sum_i p_i \left(\text{Tr} [\rho (\mathcal{J}_{\rho}^{-1/2}(\rho_i))^2] - 1 \right) = \sum_i p_i V(\rho, O_i) \end{aligned} \quad (25)$$

where in the first line we have used the $\text{Tr} \rho_i = 1$, the second line follows from (22), and $O_i \equiv \mathcal{J}_{\rho}^{-1/2}(\rho_i)$ with $\langle O_i \rangle = 1$. Eq. (25) establishes a link between the variance and the entropic uncertainty in the first-order sense. Since the quantum relative entropy encodes the closeness between two density matrices, the $V(\rho, O_i)$ is again a quantity measuring the (in)distinguishability between states but independent on the measured observable A .

The above relation (25) can also be considered for localized states on subsystems. Let $\rho_B = \sum_j p_j \Pi_B^j$ as before. For the pure state Π_B^j , its reduced density matrix on a subsystem, say B_1 , $\sigma_{B_1}^j = \text{Tr}_{\bar{B}_1} \Pi_B^j$ is no longer pure in general. So, the reduced density matrix of ρ_B on B_1 is $\rho_{B_1} = \sum_i p_i \sigma_{B_1}^i$. In this setting, we can consider the local observable

$$O_i^{B_1} = \mathcal{J}_{\rho_{B_1}}^{-1/2}(\sigma_{B_1}^i). \quad (26)$$

Again, we have $\langle O_i^{B_1} \rangle = 1$. Similar to (25), we also have

$$\begin{aligned} V(\rho_B, O_i^{B_1}) &= \text{Tr} \rho_B [(\mathcal{J}_{\rho_{B_1}}^{-1/2}(\sigma_{B_1}^i))^2 - 1] = \\ &= \text{Tr} [\sigma_{B_1}^i (\mathcal{J}_{\sigma_{B_1}^i}^{1/2}(\rho_{B_1}^{-1}) - 1)] \end{aligned} \quad (27)$$

as the first order expansions of $\hat{S}(\sigma_{B_1}^i \| \rho_{B_1})$. In (27) we have used the property that $\text{Tr} \rho_B = 1$ as a normalized density matrix. Eq. (27) relates the indistinguishability of local states and the measurement uncertainty (of $O_i^{B_1}$) in ρ , in an average sense.

Recall that the BS relative entropy and the quantum relative entropy satisfy $\hat{S}(\sigma \| \rho) \geq S(\sigma \| \rho)$ [31], thereby

$$\hat{S}(\sigma_{B_1}^i \| \rho_{B_1}) \geq \frac{1}{2} \|\sigma_{B_1}^i - \rho_{B_1}\|_1^2 \quad (28)$$

where the Schatten-1 norm $\|\cdot\|_1$ is just the trace distance introduced above. On the other hand, the variance $V(\rho_B, O_i^{B_1})$ before the series expansion is by definition a Schatten-2 norm,

$$\begin{aligned} V(\rho_B, O_i^{B_1}) &= \text{Tr}[(\sigma_{B_1}^i - \rho_{B_1}) \rho_{B_1}^{-1} (\sigma_{B_1}^i - \rho_{B_1})] = \\ &= \|(\sigma_{B_1}^i - \rho_{B_1}) \rho_{B_1}^{-1/2}\|_2^2. \end{aligned} \quad (29)$$

By the Hölder's inequality, we have

$$\begin{aligned} \|(\sigma_{B_1}^i - \rho_{B_1})\|_1 &= \|(\sigma_{B_1}^i - \rho_{B_1}) \rho_{B_1}^{-1/2} \rho_{B_1}^{1/2}\|_1 \\ &\leq \|(\sigma_{B_1}^i - \rho_{B_1}) \rho_{B_1}^{-1/2}\|_2 \|\rho_{B_1}^{1/2}\|_2 \\ &= \|(\sigma_{B_1}^i - \rho_{B_1}) \rho_{B_1}^{-1/2}\|_2 = V(\rho_B, O_i^{B_1}). \end{aligned} \quad (30)$$

Thus, we can approach the subsystem ETH by bounding either $V(\rho_B, O_i^{B_1})$ or $\hat{S}(\sigma_{B_1}^i \| \rho_{B_1})$.

With the definition (21), we can also rewrite the variance (27) as

$$\text{Tr}[(\mathcal{J}_{\rho_B}^{1/2} \circ \mathcal{J}_{\rho_{B_1}}^{-1/2}(\sigma_{B_1}^i))(\rho_B^{-1} \mathcal{J}_{\rho_B}^{1/2} \circ \mathcal{J}_{\rho_{B_1}}^{-1/2}(\sigma_{B_1}^i) - 1)], \quad (31)$$

which is the first-order expansion of $\hat{S}(\mathcal{J}_{\rho_B}^{1/2} \circ \mathcal{J}_{\rho_{B_1}}^{-1/2}(\sigma_{B_1}^i) \| \rho_B)$. In this form (31), we find that the map $\mathcal{J}_{\rho_B}^{1/2} \circ \mathcal{J}_{\rho_{B_1}}^{-1/2}$ is just the Petz recovery map $\mathcal{R}_{\rho}^{B_k \rightarrow B}$ of the CPTP map $\mathcal{N}_{B \rightarrow B_k} = \text{Tr}_{\bar{B}_k}$ with respect to the reference state ρ_B , cf. [32]. In this way, we can rewrite

$$\hat{S}(\sigma_{B_1}^i \| \rho_{B_1}) = \hat{S}(\mathcal{R}_{\rho}^{B_1 \rightarrow B}(\sigma_{B_1}^i) \| \rho_B). \quad (32)$$

IV. SUBSYSTEM ETH FOR TRANSLATION INVARIANT SYSTEMS

Now we can turn to the proof of the subsystem ETH. The strategy is to derive general bounds on the trace distance and show that they are small in the large N limit. In the light of the relations (28) and (30), we need to study the bounds on either the variance $V(\rho_B, O_i^{B_1})$ or the BS relative entropy $\hat{S}(\sigma_{B_1}^i \| \rho_{B_1})$.

We consider the translation invariant quantum lattice systems as in the last paragraph of section II B or [12]. As in (18), we consider the average observable

$$O_i^B = \frac{1}{\mathcal{C}} \sum_k \mathcal{J}_{\rho_{B_k}}^{-1/2}(\sigma_{B_k}^i). \quad (33)$$

We still have $\langle O_i^B \rangle = 1$. The quantum variance

$$\begin{aligned} V(\rho, O_i^B) &= \\ &= \text{Tr} \left[\left(\frac{1}{\mathcal{C}} \sum_k \mathcal{R}_{\rho}^{B_k \rightarrow B}(\sigma_{B_k}^i) \right) (\rho_B^{-1} \frac{1}{\mathcal{C}} \sum_l \mathcal{R}_{\rho}^{B_l \rightarrow B}(\sigma_{B_l}^i) - 1) \right], \end{aligned} \quad (34)$$

as given by eq. (31), is the first order expansion of the BS relative entropy

$$\hat{S} \left(\frac{1}{\mathcal{C}} \sum_k \mathcal{R}_{\rho}^{B_k \rightarrow B}(\sigma_{B_k}^i) \| \rho_B \right). \quad (35)$$

Since the Petz recovery map $\mathcal{R}_{\rho}^{B_k \rightarrow B}$ is also CPTP, we see that $\text{Tr}[\frac{1}{\mathcal{C}} \sum_k \mathcal{R}_{\rho}^{B_k \rightarrow B}(\sigma_{B_k}^i)]$ is also a legitimate density matrix. For example, consider that there is no correlation between the blocks $B_1, \dots, B_{\mathcal{C}}$, i.e. $\rho_B = \rho_{B_1} \otimes \dots \otimes \rho_{B_{\mathcal{C}}}$, then

$$\mathcal{R}_{\rho}^{B_k \rightarrow B}(\sigma_{B_k}^i) = \rho_{B_1} \otimes \dots \otimes \sigma_{B_k}^i \otimes \dots \otimes \rho_{B_{\mathcal{C}}}. \quad (36)$$

By the joint convexity of relative entropy, it is easy to

show that

$$\begin{aligned} \hat{S}\left(\frac{1}{\mathcal{C}} \sum_{\mathbf{k}} \mathcal{R}_{\rho}^{\mathbf{B}_k \rightarrow \mathbf{B}}(\sigma_{\mathbf{B}_k}^i) \parallel \rho_{\mathbf{B}}\right) &\leq \frac{1}{\mathcal{C}} \sum_{\mathbf{k}} \hat{S}(\mathcal{R}_{\rho}^{\mathbf{B}_k \rightarrow \mathbf{B}}(\sigma_{\mathbf{B}_k}^i) \parallel \rho_{\mathbf{B}}) \\ &= \frac{1}{\mathcal{C}} \sum_{\mathbf{k}} \hat{S}(\sigma_{\mathbf{B}_k}^i \parallel \rho_{\mathbf{B}_k}), \end{aligned} \quad (37)$$

the last expression of which is just the average of the local (in)distinguishability. As we can see from (37), if the $\hat{S}(\sigma_{\mathbf{B}_k}^i \parallel \rho_{\mathbf{B}_k})$ are small for all blocks \mathbf{B}_i , then $\hat{S}(\frac{1}{\mathcal{C}} \sum_{\mathbf{k}} \mathcal{R}_{\rho}^{\mathbf{B}_k \rightarrow \mathbf{B}}(\sigma_{\mathbf{B}_k}^i) \parallel \rho_{\mathbf{B}})$ must be small, but the converse is not true.

To prove the subsystem ETH, we need to show that

$$\hat{S}\left(\frac{1}{\mathcal{C}} \sum_{\mathbf{k}} \mathcal{R}_{\rho}^{\mathbf{B}_k \rightarrow \mathbf{B}}(\sigma_{\mathbf{B}_k}^i) \parallel \rho_{\mathbf{B}}^c\right) \sim \mathcal{O}(\mathbf{N}^{-\alpha}), \quad (38)$$

$$\text{or } V(\rho^c, \mathbf{O}_i^{\mathbf{B}}) \sim \mathcal{O}(\mathbf{N}^{-\alpha}), \quad (39)$$

for $0 < \alpha < 1$ as in (16). Here, the \mathbf{N} can be simply considered as \mathcal{C} .

Firstly, the quantum variance can be rewritten as

$$\begin{aligned} V(\rho, \mathbf{O}_i^{\mathbf{B}}) &= \frac{1}{\mathcal{C}^2} \sum_{\mathbf{k}} V(\rho, \mathbf{O}_i^{\mathbf{B}_k}) + \\ &+ \frac{1}{\mathcal{C}^2} \sum_{\mathbf{k} \neq \mathbf{l}} \text{Tr}[\mathbf{O}_i^{\mathbf{B}_k} \otimes \mathbf{O}_i^{\mathbf{B}_l} (\rho_{\mathbf{B}_k \mathbf{B}_l} - \rho_{\mathbf{B}_k} \otimes \rho_{\mathbf{B}_l})]. \end{aligned} \quad (40)$$

The first term in (40) is the local variance, in which the terms $V(\rho, \mathbf{O}_i^{\mathbf{B}_k})$ are defined locally on \mathbf{B}_k , so that they will not grow with \mathcal{C} . So, if $V(\rho, \mathbf{O}_i^{\mathbf{B}_k}) \leq C, \forall \mathbf{B}$, then

$$\frac{1}{\mathcal{C}^2} \sum_{\mathbf{k}} V(\rho, \mathbf{O}_i^{\mathbf{B}_k}) \leq \frac{1}{\mathcal{C}^2} \sum_{\mathbf{k}} C \sim \mathcal{O}(\mathcal{C}^{-1}). \quad (41)$$

The second term depends on the correlations between \mathbf{B}_k and \mathbf{B}_l . Suppose that the correlation is bounded by $\|\rho_{\mathbf{B}_k \mathbf{B}_l} - \rho_{\mathbf{B}_k} \otimes \rho_{\mathbf{B}_l}\| \leq C'$ for \mathcal{C}_{∂} blocks, then the second term in (40) is less than or equal to

$$\frac{1}{\mathcal{C}^2} \sum_{\mathbf{k}} \sum_{\mathbf{B}_l \in \partial} C' \sim \mathcal{O}(\mathcal{C}_{\partial} \mathcal{C}^{-1}), \quad (42)$$

where ∂ denotes the region in which the bounds on correlations hold. For short-range correlations such that

$$\mathcal{O}(\mathcal{C}_{\partial} \mathcal{C}^{-1}) \sim \mathcal{O}(\mathcal{C}^{-\alpha}), \quad (43)$$

we see that (39) holds in such cases.

Secondly, we study the bounds on the BS relative entropy (35). To this end, define the m -th moment of the (expanded logarithm) operator $\mathbf{O}_i^{\mathbf{B}} - \mathbf{I}$,

$$\mathbf{M}^{(m)} = \text{Tr}[\rho_{\mathbf{B}}(\mathbf{O}_i^{\mathbf{B}} - \mathbf{I})^m] \quad (44)$$

which is the higher-moment generalization of (27). Then, by the power series of the matrix logarithm, we have

$$\begin{aligned} \hat{S}\left(\frac{1}{\mathcal{C}} \sum_{\mathbf{k}} \mathcal{R}_{\rho}^{\mathbf{B}_k \rightarrow \mathbf{B}}(\sigma_{\mathbf{B}_k}^i) \parallel \rho_{\mathbf{B}}\right) &= \\ &= \frac{1}{\mathcal{C}} \sum_{\mathbf{k}} \text{Tr}[\rho_{\mathbf{B}} \mathbf{O}^{\mathbf{B}_k} \ln(\frac{1}{\mathcal{C}} \sum_{\mathbf{l}} \mathbf{O}^{\mathbf{B}_l})] = \\ &= V(\rho, \mathbf{O}_i^{\mathbf{B}}) + \dots + \frac{(-1)^n}{n-1} (\mathbf{M}^{(n)} + \mathbf{M}^{(n-1)}) + \dots \\ &= \frac{1}{2} V(\rho, \mathbf{O}_i^{\mathbf{B}}) + \sum_{n=3}^{\infty} \frac{(-1)^n}{(n-1)n} \mathbf{M}^{(n)}. \end{aligned} \quad (45)$$

The first term $V(\rho, \mathbf{O}_i^{\mathbf{B}})$ has been bounded as in (41). The other terms in (45) depend on the multipartite correlations, and the higher moments $\mathbf{M}^{(m)}$ in them can be bounded in the same way as in [13],

$$\mathbf{M}^{(m)} \leq \frac{1}{\mathcal{C}^m} \mathcal{O}(\mathcal{C}^{m/2}) \sim \mathcal{O}(\mathcal{C}^{-m/2}), \quad (46)$$

where we omit those parts that do not increase with \mathcal{C} . Thus, we obtain the overall bounds (38) on the BS relative entropy.

Thirdly, if we replace the canonical thermal state ρ^c by another local state $\sigma_{\mathbf{B}_l}$ in the above formulas, we will find that the scaling analysis still holds. In other words, for two local states, we have

$$\|\sigma_{\mathbf{B}_k}^i - \sigma_{\mathbf{B}_l}^i\| \sim \mathcal{O}(\mathbf{N}^{-\alpha}). \quad (47)$$

We have therefore successfully proved the subsystem ETH by showing the bounds or decaying behaviors (38) and (39).

Notice that, compared to the proofs given in (the appendix) of [12], we have changed the (in)distinguishability measure (12) by the variance or BS relative entropy to get the bounds on them. We can apply such a replacement back to the proofs of the weak ETH with eigenstate typicality as in [12] to see what happens. To this end, we consider a microcanonical energy shell $[E - \delta, E]$ with width δ with the index set

$$\mathcal{M}_{E, \delta} = \{i | E_i \in [E - \delta, E]\}. \quad (48)$$

We can discuss the (in)distinguishability of the microcanonical and canonical ensembles by using

$$\hat{S}(\rho_{\mathbf{B}_1}^{\text{mc}} \parallel \rho_{\mathbf{B}_1}^c) \leq \sum_{i \in \mathcal{M}_{E, \delta}} \frac{1}{\mathbf{D}} \hat{S}(\sigma_{\mathbf{B}_1}^i \parallel \rho_{\mathbf{B}_1}^c) \quad (49)$$

which is just a consequence of the joint convexity of relative entropy. In the large \mathbf{N} limit, we have from (38) the right hand side of (49) is very small, so we can conclude the equivalence between the microcanonical and canonical ensembles that in this case.

Before ending this section, let us look at several weaker conditions without proofs. By replacing d by $\hat{S}(\sigma_{\mathbf{B}_1}^i \parallel \rho_{\mathbf{B}_1}^c)$

and using (32), we see that the microcanonical average (13) becomes

$$\begin{aligned} & \sum_{i \in \mathcal{M}_{E,\delta}} \frac{1}{D} \hat{S}(\sigma_{B_1}^i, \|\rho_{B_1}^c\|) = \\ & = \hat{S}\left(\sum_{i \in \mathcal{M}_{E,\delta}} \frac{1}{D} \mathcal{R}_\rho^{B_1 \rightarrow B}(\sigma_{B_1}^i) \otimes \Pi_{B'}^i, \|\rho_B^c \otimes \rho_{B'}^{mc}\|\right), \end{aligned} \quad (50)$$

where D is number of block given by the index set $\mathcal{M}_{E,\delta}$ and $\rho_{B'}^{mc} = \sum_{i \in \mathcal{M}_{E,\delta}} \frac{1}{D} \Pi_{B'}^i$ is the microcanonical density matrix. In the second line of (50), we have extended the state of B by a microcanonical state of B' to a quantum-classical state; it is easy to see that added part in the tensor product contributes zero to the relative entropy, which is a conventional technique for treating the Holevo information [27]. With (50) at our disposal, we can propose the following quantity

$$\hat{S}\left(\sum_{i \in \mathcal{M}_{E,\delta}} \frac{1}{D} \mathcal{R}_\rho^{B_1 \rightarrow B}(\sigma_{B_1}^i) \otimes \Pi_{B'}^i, \|\rho_B^{mc} \otimes \rho_{B'}^{mc}\|\right) \quad (51)$$

which is just the analog of (13) with the $\mathbf{d}(\Pi_B^j, \rho^{MC}; \mathcal{A}^{B_1})$ replaced by $\hat{S}(\sigma_{B_1}^i, \|\rho_{B_1}^{mc}\|)$. And in a similar way, we have the Chebyshev-type inequality,

$$\begin{aligned} & P_\rho(\hat{S}(\sigma_{B_1}^i, \|\rho_{B_1}^{mc}\|) \geq \epsilon^2) \\ & \leq \frac{1}{\epsilon^2} \hat{S}\left(\sum_{i \in \mathcal{M}_{E,\delta}} \frac{1}{D} \mathcal{R}_\rho^{B_1 \rightarrow B}(\sigma_{B_1}^i) \otimes \Pi_{B'}^i, \|\rho_B^{mc} \otimes \rho_{B'}^{mc}\|\right) \end{aligned} \quad (52)$$

for $\epsilon > 0$. When eq. (50) is very small, we can conclude that the measurement results concentrate on the results predicted by the microcanonical ensemble ρ^{mc} . This hypothesis is similar to the weak ETH with eigenstate typicality. This hypothesis can also be translated to the canonical ensemble using the ensemble equivalence, e.g. (49).

V. CONCLUSION AND DISCUSSION

We have studied the subsystem ETH for translation invariant quantum systems. We develop upon the setting for translation invariant systems given in [12] by relating the quantum variance to the BS relative entropy. Surprisingly, with this technical input, we are able to prove the subsystem ETH for translation invariant systems using the similar scaling analysis as in [12]. The proof given above is elementary, without referring to the advanced techniques from random matrix theory. Since the subsystem is stronger than the local ETH, our results corroborate the previous results for local ETH for translation invariant systems [9, 11, 12].

It seems that the Hamiltonian of the system does not make an appearance in our proof, but in fact, the Hamiltonian is important in the condition of short-range correlations leading to (43). For long-range interacting systems, a deviation to some extent from the short-range result could be expected, cf. the recent numerical test [33]. Another point is that in the analysis of (45) the higher moments are relevant. The higher-moment versions of ETH can be related to many interesting structures, such as the out-of-time-ordered correlation functions indicating quantum chaos [34]. This could be a possible approach to relating the chaotic conjecture and the present analysis without referring to random matrices. Moreover, it is also interesting to study the eigenstate fluctuation theorems [12, 35] at the subsystem level, which might be a suitable situation for thermalized open/non-Hermitian quantum system. All these aspects are left to future investigations.

ACKNOWLEDGMENTS

This work is supported by the National Natural Science Foundation of China under Grants No. 12305035.

-
- [1] C. Gogolin and J. Eisert, Equilibration, thermalisation, and the emergence of statistical mechanics in closed quantum systems, *Rep. Prog. Phys.* **79**, 056001 (2016).
 - [2] T. Mori, T. N. Ikeda, E. Kaminishi, and M. Ueda, Thermalization and prethermalization in isolated quantum systems: A theoretical overview, *J. Phys. B: At. Mol. Opt. Phys.* **51**, 112001 (2018).
 - [3] L. D'Alessio, Y. Kafri, A. Polkovnikov, and M. Rigol, From quantum chaos and eigenstate thermalization to statistical mechanics and thermodynamics, *Adv. Phys.* **65**, 239 (2016).
 - [4] J. M. Deutsch, Eigenstate thermalization hypothesis, *Rep. Prog. Phys.* **81**, 082001 (2018).
 - [5] J. M. Deutsch, Quantum statistical mechanics in a closed system, *Phys. Rev. A* **43**, 2046 (1991).
 - [6] M. Srednicki, Chaos and quantum thermalization, *Phys. Rev. E* **50**, 888 (1994).
 - [7] M. Srednicki, The approach to thermal equilibrium in quantized chaotic systems, *J. Phys. A: Math. Gen.* **32**, 1163 (1999).
 - [8] G. Cipolloni, L. Erdos, and D. Schröder, Eigenstate thermalization hypothesis for Wigner matrices, *Commun. Math. Phys.* **388**, 1005 (2021).
 - [9] S. Sugimoto, J. Henheik, V. Riabov, and L. Erdos, Eigenstate thermalisation hypothesis for translation invariant spin systems, *J. Stat. Phys.* **190**, 128 (2023).
 - [10] G. Biroli, C. Kollath, and A. M. Läuchli, Effect of rare fluctuations on the thermalization of isolated quantum systems, *Phys. Rev. Lett.* **105**, 250401 (2010).
 - [11] T. Mori, Weak eigenstate thermalization with large deviation bound, arXiv:1609.09776.
 - [12] E. Iyoda, K. Kaneko, and T. Sagawa, Fluctuation theorem for many-body pure quantum states, *Phys. Rev. Lett.* **119**, 100601 (2017).

- [13] T. Kuwahara and K. Saito, Eigenstate thermalization from the clustering property of correlation, *Phys. Rev. Lett.* **124**, 200604 (2020) .
- [14] A. M. Kaufman, M. E. Tai, A. Lukin, M. Rispoli, R. Schittko, P. M. Preiss, and M. Greiner, Quantum thermalization through entanglement in an isolated many-body system, *Science* **353**, 794 (2016).
- [15] A. Dymarsky, N. Lashkari, and H. Liu, Subsystem eigenstate thermalization hypothesis, *Phys. Rev. E* **97**, 012140 (2018).
- [16] N. Lashkari, A. Dymarsky, and H. Liu, Eigenstate thermalization hypothesis in conformal field theory, *J. Stat. Mech.* (2018) 033101.
- [17] N. Lashkari, A. Dymarsky, and H. Liu, Universality of quantum information in chaotic CFTs, *JHEP03(2018)070*.
- [18] P. Basu, D. Das, S. Datta, and S. Pal, Thermality of eigenstates in conformal field theories, *Phys. Rev. E* **96**, 022149 (2017).
- [19] S. He, F.-L. Lin, and J.-J. Zhang, Subsystem eigenstate thermalization hypothesis for entanglement entropy in CFT, *JHEP08(2017)126*.
- [20] S. He, F.-L. Lin, and J.-J. Zhang, Dissimilarities of reduced density matrices and eigenstate thermalization hypothesis, *JHEP12(2017)073*.
- [21] T. Faulkner and H.-J. Wang, Probing beyond ETH at large c , *JHEP06(2018)123*.
- [22] W.-Z. Guo, F.-L. Lin, and J.-J. Zhang, Note on ETH of descendant states in 2D CFT, *JHEP01(2019)152*.
- [23] A. Dymarsky and K. Pavlenko, Generalized eigenstate thermalization hypothesis in 2D conformal field theories, *Phys. Rev. Lett.* **123**, 111602 (2019).
- [24] A. Dymarsky and K. Pavlenko, Generalized Gibbs ensemble of 2d CFTs at large central charge in the thermodynamic limit, *JHEP01(2019)098*.
- [25] M. P. Müller, E. Adlam, L. Masanes, and N. Wiebe, Thermalization and canonical typicality in translation-invariant quantum lattice systems, *Commun. Math. Phys.* **340**, 499 (2015).
- [26] J. Riddell and M. P. Müller, Generalized eigenstate typicality in translation-invariant quasifree fermionic models, *Phys. Rev. B* **97**, 035129 (2018).
- [27] J. Watrous, *The Theory of Quantum Information*, Cambridge University Press, 2018.
- [28] S.-L. Luo, Quantum versus classical uncertainty, *Theor. Math. Phys.* **143**, 681 (2005).
- [29] K. Korzekwa, M. Lostaglio, D. Jennings, and T. Rudolph, Quantum and classical entropic uncertainty relations, *Phys. Rev. A* **89**, 042122 (2014).
- [30] V. P. Belavkin and P. Staszewski, C^* -algebraic generalization of relative entropy and entropy, *Ann. Inst. Henri Poincaré A* **37**, 51 (1982).
- [31] A. Bluhm, A. Capel, and A. Pérez-Hernández, Exponential decay of mutual information for Gibbs states of local Hamiltonians, *Quantum* **6**, 650 (2022).
- [32] H. Kwon and M. S. Kim, Fluctuation theorems for a quantum channel, *Phys. Rev. X* **9**, 031029 (2019).
- [33] S. Sugimoto, R. Hamazaki, and M. Ueda, Eigenstate thermalization in long-range interacting systems, *Phys. Rev. Lett.* **129**, 030602 (2022).
- [34] L. Foini and J. Kurchan, Eigenstate thermalization hypothesis and out of time order correlators, *Phys. Rev. E* **99**, 042139 (2019).
- [35] E. Iyoda, K. Kaneko, and T. Sagawa, Eigenstate fluctuation theorem in the short- and long-time regimes, *Phys. Rev. E* **105**, 044106 (2022).