

Quantum Speed Limits Based on Schatten Norms: Universality and Tightness

Alberto J. B. Rosal,¹ Diogo O. Soares-Pinto,¹ and Diego Paiva Pires²

¹*Instituto de Física de São Carlos, Universidade de São Paulo, CP 369, 13560-970 São Carlos, Brazil*

²*Departamento de Física, Universidade Federal do Maranhão, Campus Universitário do Bacanga, 65080-805, São Luís, Maranhão, Brazil*

We present two families of quantum speed limits (QSLs) for finite-dimensional quantum systems undergoing a general physical process. These QSLs were obtained using Schatten α -norms, firstly exploiting the geometric features of the space of quantum states, and secondly employing the Holder's inequality for matrix norms. In particular, for the case of single-qubit states, we find that the geometric QSL is independent of the Schatten norm chosen, thus revealing a universality behavior of such quantifiers. Furthermore, we provide a comparison of these quantum speed limits with existing paradigmatic QSLs in literature, thus showing that the latter results represent particular cases of a general class of QSLs related to Schatten α -norms. Noteworthy, we address necessary and sufficient conditions for the tightness of the quantum speed limit that mostly depends on the populations and quantum coherences of the evolved single-qubit state, and also present a geometric interpretation for these set of conditions. Finally, we compare the two QSL obtained for the dynamics of single-qubit states, also presenting an inequality between them that has a clear geometrical meaning.

Introduction.— Quantum speed limits (QSLs) are of fundamental and practical interest in several branches of physics, ranging from non-equilibrium thermodynamics [1–4], to quantum many-body systems [5–9], also including the study of quantum metrology [10], communication and quantum computing [11, 12], quantum entanglement [13–16], quantum-to-classical transition [17–19], optimal control theory [20, 21], quantum batteries [22], and also non-Hermitian systems [23–25]. Recent studies include experimental investigations of the quantum speed limit in both platforms of trapped atoms [26], and Nuclear Magnetic Resonance systems [27, 28]. In short, the QSL sets the shortest time in which a quantum system undergoing a given dynamics can evolve from some initial to final state. We refer to Refs. [29, 30] for recent reviews covering the subject.

For closed quantum systems, Mandelstam and Tamm (MT) [31] proved that $\tau \geq \tau_{MT}$, with $\tau_{MT} := (\hbar/\Delta E) \arccos(|\langle\psi_0|\psi_\tau\rangle|)$ being the QSL time, while $\Delta E = \sqrt{\langle H^2 \rangle - \langle H \rangle^2}$ is the variance of the time-independent Hamiltonian H that generates the unitary dynamics, whereas $|\psi_0\rangle$ and $|\psi_\tau\rangle$ are the initial and final pure states of the quantum system, respectively. In particular, for two orthogonal states, the QSL time reduces to $\tau_{MT} = \pi\hbar/(2\Delta E)$. In turn, Margolus and Levitin [32] realized the bound $\tau \geq \tau_{ML}$ for closed quantum systems and maximally distinguishable states, with the QSL time $\tau_{ML} := \hbar\pi/[2(\langle\psi_0|H|\psi_0\rangle - E_0)]$, where E_0 is the ground state of H . We shall mention another restrictive QSL bound for closed systems related to pure states with a bounded energy spectrum that is dual to the ML bound [33]. Noteworthy, Levitin and Toffoli (LT) have shown that the tighter QSL for orthogonal states is achieved by combining the results by MT and ML as $\tau_{QSL} = \max\{\tau_{MT}, \tau_{ML}\}$ [34]. In addition, QSLs have also been addressed for driven closed quantum systems initialized in mixed states [35–38]. Recent investigations include the derivation of an extended MT and ML bounds for states with arbitrary fidelity [39, 40].

QSLs have been also investigated for open quantum systems. We point out that Taddei *et al.* [41] and del Campo *et al.* [42] originally addressed such issue for nonunitary evolu-

tions, thus deriving MT-like bounds for general physical processes. The former bound is based on the quantum Fisher information, while the latter QSL time is obtained from the relative purity between initial and final states. In turn, Deffner and Lutz [43] proposed a MT bound for the QSL time that is based on operator norms and the so-called Bures angle, also indicating that non-Markovianity signatures can speed up the quantum evolution. However, it has been shown that the QSL time is sensitive to the choice of the initial state of the system, also depending on the details of its dynamical evolution [44–49]. QSLs have been discussed within the perspective of information geometry [50]. This approach provides a family of QSLs based on contractive Riemannian metrics equipping the space of quantum states, and holds for both unitary and nonunitary dynamics, mixed and pure states [51–53]. We also mention the interplay between QSLs and the theory of Finsler metrics, which finds applications in quantum information processing [54]. In this setting, to define the QSL time, we look for an information-theoretic quantifier that defines a distance measure in the space of quantum states, i.e., a certain information metric [55].

QSLs have also been addressed by using families of matrix norms. For example, Ref. [56] discusses a QSL based on Schatten α -norms obtained by using an approach that is restricted to the Wigner phase space. In turn, Ref. [57] comment on the interplay between QSLs and statistical speeds for unitary dynamics with local Hamiltonians. In addition, Refs. [58–60] study speed limits for open quantum systems based on trace distance. Recently, Ref. [61] introduced a family of coherent QSL bounds based on Schatten α -norms that only apply to closed quantum systems. Based on the different QSLs present in the literature, each related to some measure of distinguishability of quantum states, would it be possible to generalize them to obtain an ultimate QSL that would be valid for any Schatten norm and general physical processes? If so, how could this be? How does a given QSL change when choosing another information metric to equip the space of quantum states? For example, to the best of our knowledge, there is no clear relationship in literature between the QSL obtained using the Bures angle [43] with those QSLs

obtained using the Hilbert-Schmidt norm [38]. We will seek to answer these questions in this work, bringing an idea of universality between different types of QSL. In this work, we aim to develop families of QSLs based on general Schatten α -norms defined in the space of quantum states that holds for arbitrary physical processes, thereby generalizing other types of QSLs that already exist in the literature. We also compare the QSL based on Schatten norms and the QSL developed from the geometrical approach with those existing in the literature [38, 42, 43]. Furthermore, we analyze the necessary and sufficient conditions under the dynamics for the QSL induced by Schatten norms to be tight.

Geometric QSL from a normed space of quantum states.— We consider a quantum system related to a finite-dimensional Hilbert space \mathcal{H} , with $d = \dim(\mathcal{H})$. The system is initialized at the probe state $\rho_0 \in \mathcal{D}(\mathcal{H})$, where $\mathcal{D}(\mathcal{H}) = \{\rho \in \mathcal{H} \mid \rho^\dagger = \rho, \rho \geq 0, \text{Tr}(\rho) = 1\}$ defines the convex set of quantum states. Let Γ_Ω be the length of a given path γ that connects two quantum states $\rho, \varrho \in \mathcal{D}(\mathcal{H})$ in the space of quantum states. The geodesic path is such that minimize the length of a curve connecting the states ρ and ϱ . This minimal length will be called geodesic distance $D^{\text{geo}}(\rho, \varrho)$. For two quantum states $\rho, \varrho \in \mathcal{D}(\mathcal{H})$, and for a given path γ that connects these two states, we have that the geodesic distance $D^{\text{geo}}(\rho, \varrho)$ is upper bounded by the length Γ_γ of the path γ as follows

$$D^{\text{geo}}(\rho, \varrho) \leq \Gamma_\gamma. \quad (1)$$

For example, if the space of quantum states $\mathcal{D}(\mathcal{H})$ is endowed with a given matrix norm $\|\cdot\|$, this space becomes a metric space, and through of the distance induced from this norm, we can take $D^{\text{geo}}(\rho, \varrho) = \|\rho - \varrho\|$.

Next, let $\gamma_\tau = \{\rho_t = \Lambda_{t,0}(\rho_0) : t \in [0, \tau]\}$ be a given dynamical path, where the system evolves from $\varrho = \rho_0$ to $\rho = \Lambda_{\tau,0}(\rho_0) = \rho_\tau$, while $\Lambda_{t,0}$ is a time-differentiable dynamical map. Note that for a given initial condition and a time-differentiable dynamical map, and for all $t \geq 0$, we have a point ρ_t in the space state $\mathcal{D}(\mathcal{H})$, and the result is a continuous and smooth path given by the curve $\rho_t \in \mathcal{D}(\mathcal{H})$. In this setting, the length Γ_{γ_τ} is given by

$$\Gamma_{\gamma_\tau} = \int_0^\tau dt \left\| \frac{d\rho_t}{dt} \right\|. \quad (2)$$

Hereafter, we set the geodesic distance $D^{\text{geo}}(\rho_0, \rho_\tau) = \|\rho_\tau - \rho_0\|$. Hence, combining Eqs. (1) and (2), one gets the result

$$\|\rho_\tau - \rho_0\| \leq \int_0^\tau dt \left\| \frac{d\rho_t}{dt} \right\|. \quad (3)$$

Therefore, from Eq. (3), we obtain a lower bound on the evolution time given by

$$\tau \geq \tau_{QSL}^{\|\cdot\|}, \quad (4)$$

where $\tau_{QSL}^{\|\cdot\|} \equiv \tau_{QSL}^{\|\cdot\|}(\tau)$ sets the geometric quantum speed limit induced from the norm $\|\cdot\|$ and it is defined as follows

$$\tau_{QSL}^{\|\cdot\|} := \frac{\|\rho_\tau - \rho_0\|}{\left\langle \left\| \frac{d\rho_t}{dt} \right\| \right\rangle_\tau}, \quad (5)$$

with $\langle \left\| \frac{d\rho_t}{dt} \right\| \rangle_\tau := (1/\tau) \int_0^\tau dt \left\| \frac{d\rho_t}{dt} \right\|$ being the mean length of the dynamical path. We note that the QSL bound in Eq. (5) holds for general physical processes and applies for any family matrix norms, and thus generalizes the results discussed in Refs. [56, 58–61]. In what follows, we set the norm $\|\cdot\| = \|\cdot\|_\alpha$ being the Schatten α -norm and discuss details on the QSL in Eq. (5).

a. Schatten α -QSL for qubits — Here we set the matrix norm $\|\cdot\|$ as the Schatten α -norm $\|\cdot\|_\alpha$, with $\alpha \in [1, \infty)$. In this setting, space of quantum states $\mathcal{D}(\mathcal{H})$ becomes a metric space equipped with the geodesic distance $D^{\text{geo}}(\rho, \varrho) := D_\alpha(\rho, \varrho) = \|\rho - \varrho\|_\alpha$, where $\|A\|_\alpha := (\text{Tr}\{(\sqrt{A^\dagger A})^\alpha\})^{\frac{1}{\alpha}}$ [see Supplemental Material, Sec. S.I], and thus from Eq. (5) the quantum speed limit induced from this norm, for a time of evolution τ , is given by

$$\tau_{QSL}^\alpha(\tau) := \frac{\|\rho_\tau - \rho_0\|_\alpha}{\int_0^\tau dt \left\| \frac{d\rho_t}{dt} \right\|_\alpha} \tau. \quad (6)$$

In the following, we consider the dynamics of single-qubit states whose the evolution is governed by a dynamical map that is time-differentiable. For such a dynamic, we can compute explicitly the distance $\|\rho_\tau - \rho_0\|_\alpha$ as well as we can get the norm $\left\| \frac{d\rho_t}{dt} \right\|_\alpha$ [see Supplemental Material, Sec. S.II]. Moreover, we will consider that $\varrho = \rho_0$ is the initial state and then $\rho_\tau = \Lambda_{\tau,0}(\rho_0)$ is the final state for an evolution time τ . We will demonstrate the following result:

Theorem 1 (Universality of the α -QSL for single-qubit states). *Let us consider a two-dimensional quantum system initially described by the single-qubit state $\rho_0 = (1/2)(\mathbb{I} + \vec{n}_0 \cdot \vec{\sigma})$. The dynamics of the system is given in terms of a time-differentiable, CPTP, dynamical map $\Lambda_{t,0}(\bullet)$, and its instantaneous state becomes $\rho_t = \Lambda_{t,0}(\rho_0) = (1/2)(\mathbb{I} + \vec{n}_t \cdot \vec{\sigma})$, for all $t \geq 0$, where $\vec{n}_t = (n_{x,t}, n_{y,t}, n_{z,t})$ is the Bloch vector for every $t \in [0, \tau]$. From Eq. (6), the geometric α -QSL that is induced from Schatten α -norm, for every $\alpha \in [1, \infty)$, yields*

$$\tau_{QSL}^\alpha(\tau) = \frac{d_{\text{euc}}(\vec{n}_0, \vec{n}_\tau)}{\int_0^\tau dt \left\| \frac{d\vec{n}_t}{dt} \right\|} \tau, \quad (7)$$

with

$$\left\| \frac{d\vec{n}_t}{dt} \right\| = \sqrt{\sum_{j=x,y,z} \left(\frac{dn_{j,t}}{dt} \right)^2}, \quad (8)$$

while

$$d_{\text{euc}}(\vec{n}_0, \vec{n}_\tau) = \sqrt{\sum_{j=x,y,z} (n_{j,\tau} - n_{j,0})^2} \quad (9)$$

stands for the Euclidean distance between the initial and the final Bloch vectors in the Bloch space.

Proof. See Supplemental Material, Sec. S.III. \square

Theorem 1 shows that there is some kind of universality of the geometric QSL induced through of a α -norm. That is,

does not matter what is the metric chosen, a unique quantifier is given by the Eq. (7). In principle, one expects that different values of α could generate a different value of the quantifier τ_{QSL}^α , but according to this theorem the QSL time remains invariant regardless the value of $1 \leq \alpha < \infty$.

b. Comparing α -QSL with others quantifiers — We shall begin by comparing the geometric QSL induced from an α -norm [see Eq. (7)] with the QSL developed by Deffner and Lutz (DL) [43]. By taking a pure initial state $\rho_0 = |\psi_0\rangle\langle\psi_0|$ undergoing a certain nonunitary dynamics, they showed the lower bound $\tau \geq \tau_{QSL}^{DL}$, with

$$\tau_{QSL}^{DL} = \frac{1 - \langle\psi_0|\rho_\tau|\psi_0\rangle}{\frac{1}{\tau} \int_0^\tau dt \left\| \frac{d\rho_t}{dt} \right\|_\infty}. \quad (10)$$

Since we are working with a two-dimensional Hilbert space \mathcal{H} , we can choose an orthonormal basis $\{|\psi_0\rangle, |\psi_0^\perp\rangle\}$, where $\langle\psi_0|\psi_0^\perp\rangle = 0$. In this basis, we can define the Pauli matrix $\sigma_z = |\psi_0\rangle\langle\psi_0| - |\psi_0^\perp\rangle\langle\psi_0^\perp|$. Therefore, in this basis, taking the Bloch representation for the final state, we can write

$$1 - \langle\psi_0|\rho_\tau|\psi_0\rangle = \frac{1 - n_{z,\tau}}{2} \leq \frac{1}{2} d_{\text{euc}}(\vec{n}_0, \vec{n}_\tau), \quad (11)$$

where we have used the fact that $n_{z,t} \leq 1$ for every $t \geq 0$, and $d_{\text{euc}}(\vec{n}_0, \vec{n}_\tau) = \sqrt{n_{x,\tau}^2 + n_{y,\tau}^2 + (1 - n_{z,\tau})^2}$ is the Euclidean distance between the time-dependent vector $\vec{n}_t = (n_{x,t}, n_{y,t}, n_{z,t})$, and the initial Bloch vector $\vec{n}_0 = (0, 0, 1)$ in the chosen basis $\{|\psi_0\rangle, |\psi_0^\perp\rangle\}$. Therefore, taking the Theorem 1, we can conclude that

$$\tau_{QSL}^{DL} \leq \tau_{QSL}^\alpha, \quad (12)$$

where we have used that $\|d\rho_t/dt\|_\infty = (1/2)\|d\vec{n}_t/dt\|$.

Importantly, Eq. (12) holds whenever we have an initial single-qubit pure state. For a mixed initial state, we can purify this one, finding again the previous case. Then, we can see that the geometric QSL induced from a Schatten α -norm [Eq. (6)], for every $\alpha \in [1, \infty)$, is tighter than the QSL obtained by Deffner and Lutz [Eq. (10)].

Next, we compare the geometric α -QSL [see Eq. (7)] with the speed limit developed by Campaioli *et al.* (CPM) [38], which in turn depends on the Hilbert-Schmidt norm. Namely, they showed that

$$\tau_{QSL}^{CPM} := \frac{\|\rho_\tau - \rho_0\|_2}{\int_0^\tau dt \left\| \frac{d\rho_t}{dt} \right\|_2} \tau. \quad (13)$$

That is, Eq. (13) is a particular case of the general geometric α -QSL [see Eq. (6)] for $\alpha = 2$. In particular, taking a qubit dynamics, surprisingly due to the Theorem 1 and the Eq. (7), we have that this two quantifiers are, in fact, equals for every value of $\alpha \in [1, \infty)$. That is, for a qubit dynamics and for every $\alpha \in [1, \infty)$, we have

$$\tau_{QSL}^{CPM} = \tau_{QSL}^{(\alpha=2)}(\tau) = \tau_{QSL}^\alpha(\tau) = \frac{d_{\text{euc}}(\vec{n}_0, \vec{n}_\tau)}{\int_0^\tau dt \left\| \frac{d\vec{n}_t}{dt} \right\|} \tau. \quad (14)$$

Although Eq. (14) shows that, on the one hand, the CPM QSL in Eq. (13) is a particular case of a more general QSL but, on the other hand, due to the Theorem 1, for a qubit dynamics the value assumed for the CPM QSL is the same for the general case.

c. Tightness condition of α -QSL for the dynamics of single-qubit states — The geometric QSL induced from a α -norm [Eq. (6)] is tight, that is $\tau_{QSL}^\alpha/\tau = 1$, if and only if the dynamical path is the geodesic path, which represents the optimal path in the state space $\mathcal{D}(\mathcal{H})$. Since the state space has an Euclidean metric induced from the Schatten norm, the geodesic path is, in fact, the straight line connecting the initial state ρ_0 to the final state ρ_τ . Next, let's consider a qubit dynamics. From Eq. (7), we can see that this condition also holds in the Bloch space. That is:

Lemma 1 (Tightness condition in the Bloch space for the α -QSL). *Let $\rho_t = (1/2)(\mathbb{I} + \vec{n}_t \cdot \vec{\sigma}) = \rho_{11,t}|0\rangle\langle 0| + (1 - \rho_{11,t})|1\rangle\langle 1| + \rho_{12,t}|0\rangle\langle 1| + \rho_{12,t}^*|1\rangle\langle 0|$ be the single-qubit state of a given two-level system, with $\rho_{jl,t}$ being time-dependent matrix elements of the state ρ_t respective to the computational basis $\{|0\rangle, |1\rangle\}$, and $\vec{n}_t = (n_{x,t}, n_{y,t}, n_{z,t})$. For the general dynamics of this state, the geometric QSL induced from the Schatten α -norm [see Eq. (7)] is tight, i.e., $\tau_{QSL}^\alpha/\tau = 1$, if one gets*

$$\frac{d\vec{n}_t}{dt} = \left\| \frac{d\vec{n}_t}{dt} \right\| \hat{r}_{0,t}, \quad (15)$$

which implies that

$$(\text{Re}(d\rho_{12,t}/dt), -\text{Im}(d\rho_{12,t}/dt), d\rho_{11,t}/dt) = \Omega \hat{r}_{0,t}, \quad (16)$$

where

$$\hat{r}_{0,t} = \frac{\vec{n}_t - \vec{n}_0}{\|\vec{n}_t - \vec{n}_0\|} \quad (17)$$

is the unit vector which define the direction of the line that connect the initial and the final Bloch vectors, with

$$\Omega := \sqrt{\left| \frac{d\rho_{11,t}}{dt} \right|^2 + \left| \frac{d\rho_{12,t}}{dt} \right|^2}, \quad (18)$$

and $\|d\vec{n}_t/dt\|$ is defined in Eq. (8).

Proof. See Supplemental Material, Sec. S.IV. \square

Therefore, due to the Theorem 1 the optimal dynamical path in the Bloch space is, such as in the state space $\mathcal{D}(\mathcal{H})$ of the density operators, the straight line connecting the initial Bloch vector \vec{n}_0 to the final one \vec{n}_τ . This condition can be seen as the geometrical interpretation of the Lemma 1. From the algebraic point of view, see that the unit vector $\hat{r}_{0,\tau}$, which depends only on the initial and final Bloch vectors, tell us what is the signal of the values of the rates of the populations and quantum coherences. For example, if $(\hat{r}_{0,\tau})_z = (1/\Omega)d\rho_{11,t}/dt \leq 0$, then the optimal physical process need be such that $d\rho_{11,t}/dt \leq 0$, that is, a dissipative process. Then, under this algebraic point of view, we can see

that the final and initial single-qubit states restrict the class of possible dynamical maps in which the QSL can be tight.

QSL from Holder's inequality.— Here we will present a family of quantum speed limits through the Holder's inequality. In addition, we will compare this one with others QSL, and analyze the necessary and sufficient condition for this QSL to be tightness. Let \mathcal{H} be a finite-dimensional Hilbert space and $\mathcal{D}(\mathcal{H})$ its associated space of quantum states. By hypothesis, this last set is a normed space as it is equipped with a family of Schatten α -norms through which distances between quantum states can be evaluated. In addition, we will consider that the dynamical map $\Lambda_{t,0}(\bullet)$ is time-differentiable, with $\rho_t = \Lambda_{t,0}(\rho_0)$ for some initial state $\rho_0 \in \mathcal{D}(\mathcal{H})$. Therefore, $d\rho_t/dt$ is supposed to be well defined for every $t \geq 0$ and the dynamical path is a continuous and smooth path in the state space.

We define the overlap between the initial and final states as $\xi(t) = \text{Tr}(\rho_0 \rho_t)$. In particular, when $\rho_0 = |\psi_0\rangle\langle\psi_0|$ is a pure initial state, the overlap $\xi(t) = \langle\psi_0|\rho_t|\psi_0\rangle$ becomes the expected fidelity. Due to the Holder's inequality $|\text{Tr}[Y^\dagger X]| \leq \|X\|_\alpha \|Y\|_\beta$, which holds for every two linear operators $X, Y \in L(\mathcal{H})$ and real numbers $\alpha, \beta \in [1, \infty)$ where $\frac{1}{\alpha} + \frac{1}{\beta} = 1$, one finds that the absolute value of rate of such overlap becomes

$$\left| \frac{d\xi(t)}{dt} \right| = \left| \text{Tr} \left(\rho_0 \frac{d\rho_t}{dt} \right) \right| \leq \|\rho_0\|_\beta \left\| \frac{d\rho_t}{dt} \right\|_\alpha, \quad (19)$$

where we have used that $\rho_0 = \rho_0^\dagger$ is Hermitian. Integrating this inequality in the interval $[0, \tau]$, we obtain

$$|\xi(\tau) - \xi(0)| \leq \|\rho_0\|_\beta \int_0^\tau dt \left\| \frac{d\rho_t}{dt} \right\|_\alpha, \quad (20)$$

where we are using $|\xi(\tau) - \xi(0)| = \left| \int_0^\tau dt (d\xi(t)/dt) \right| \leq \int_0^\tau dt |d\xi(t)/dt|$. However, note that $\xi(0) = \text{Tr}(\rho_0^2) \in [\frac{1}{d}, 1]$ is the purity of the initial state, while $\xi(\tau) = \text{Tr}(\rho_\tau \rho_0) \in [0, 1]$ is the overlap between the initial and final states. By using these results, one gets

$$\frac{1}{\|\rho_0\|_\beta} |\text{Tr}[(\rho_\tau - \rho_0)\rho_0]| \leq \int_0^\tau dt \left\| \frac{d\rho_t}{dt} \right\|_\alpha. \quad (21)$$

Therefore, for all $\alpha, \beta \in [1, \infty)$ such that $\frac{1}{\alpha} + \frac{1}{\beta} = 1$, Eq. (21) implies the lower bound on the time of evolution

$$\tau \geq \tau_{QSL}^{\alpha, \beta}, \quad (22)$$

where we have introduced the so-called (α, β) -QSL defined as follows

$$\tau_{QSL}^{\alpha, \beta} = \frac{|\text{Tr}[(\rho_\tau - \rho_0)\rho_0]|}{\|\rho_0\|_\beta \int_0^\tau dt \left\| \frac{d\rho_t}{dt} \right\|_\alpha} \tau. \quad (23)$$

Finally, we specialize the general QSL obtained in Eq. (23) for an initial pure state $\rho_0 = |\psi_0\rangle\langle\psi_0|$. In this case, we find that $\text{Tr}(\rho_0^2) = 1$ and $\text{Tr}(\rho_0 \rho_\tau) = \langle\psi_0|\rho_\tau|\psi_0\rangle \in [0, 1]$. Since ρ_0 is a projector, we have that $\rho_0^2 = \rho_0$, and therefore $\|\rho_0\|_\beta =$

1 for every $\beta \in [1, \infty)$. Taking all together, Eq. (23) readily becomes

$$\frac{\tau_{QSL}^{\alpha, \beta}}{\tau} = \frac{1 - \langle\psi_0|\rho_\tau|\psi_0\rangle}{\int_0^\tau dt \left\| \frac{d\rho_t}{dt} \right\|_\alpha}. \quad (24)$$

Noteworthy, Eq. (24) shows that, for initial pure states, there is no more dependence on the dual norm $\|\cdot\|_\beta$ of the initial state, and the bound only depends on the norm $\|\cdot\|_\alpha$. Until now, we are working with a general Hilbert space \mathcal{H} . In what follows, we will investigate the case of single-qubit states, that is, when the Hilbert space is two-dimensional.

a. (α, β) -QSL for the single-qubit dynamics — Here we address $\tau_{QSL}^{\alpha, \beta}$ in Eq. (23) for single-qubit states $\rho_t = (1/2)(\mathbb{I} + \vec{n}_t \cdot \vec{\sigma})$ that can be mixed or pure, undergoing a general physical processes over the time interval $0 \leq t \leq \tau$. In this setting, it can be verified that Eq. (23) becomes

$$\frac{\tau_{QSL}^{\alpha, \beta}}{\tau} = \frac{2^{1-\frac{1}{\alpha}} |(\vec{n}_\tau - \vec{n}_0) \cdot \vec{n}_0|}{[(1 + \|\vec{n}_0\|)^\beta + (1 - \|\vec{n}_0\|)^\beta]^{\frac{1}{\beta}} \int_0^\tau dt \left\| \frac{d\vec{n}_t}{dt} \right\|}. \quad (25)$$

In particular, let's see what happens when the initial state is pure. In this case, we have $\|\vec{n}_0\| = 1$, and the QSL given by the Eq. (25) can be written as

$$\frac{\tau_{QSL}^{\alpha, \beta}}{\tau} = 2^{-\frac{1}{\alpha}} \frac{(1 - \vec{n}_\tau \cdot \vec{n}_0)}{\int_0^\tau dt \left\| \frac{d\vec{n}_t}{dt} \right\|}. \quad (26)$$

We note that the QSL time in Eq. (26) is independent of the dual norm $\|\cdot\|_\beta$ of the Schatten norm $\|\cdot\|_\alpha$. In fact, we can obtain Eq. (26) directly from the Eq. (24).

b. Comparing (α, β) -QSL with others quantifiers — In the following, we will compare the general QSL obtained from the Holder's inequality [Eq. (23)] with others quantifiers existing in literature. In especial, we will show that the quantifier $\tau_{QSL}^{\alpha, \beta}$ is tighter than the del Campo *et al* [42]. QSL, it is equal to Deffner-Lutz QSL for $\alpha = \infty$ (operator norm) and $\beta = 1$ (dual trace norm), and in general it is proportional to the Deffner-Lutz QSL in the case of a qubit dynamics.

We address a comparison between the (α, β) -QSL in Eq. (23), and the quantum speed limit obtained by del Campo *et al.* (CEPH) [42]. To do so, we fix $\alpha = \beta = 2$, and thus define the Schatten 2-norm as a *bona fide* distance measure on the space of quantum states. In this case, the QSL in Eq. (23) becomes

$$\frac{\tau_{QSL}^{2,2}}{\tau} = \frac{\|\rho_0\|_2 |f(\tau) - f(0)|}{\int_0^\tau dt \left\| \frac{d\rho_t}{dt} \right\|_2}, \quad (27)$$

where we introduce the relative purity as follows

$$f(t) := \frac{\text{Tr}(\rho_0 \rho_t)}{\|\rho_0\|_2^2}, \quad (28)$$

with $0 \leq f(t) \leq 1$ for all $t \in [0, \tau]$, and $f(0) = 1$. In particular, introducing the parameterization $f(t) = \cos \theta_t$, with $0 \leq \theta_t \leq \pi/2$, and also taking into account the fact that

$|\cos \theta_\tau - 1| \geq 4\theta_\tau^2/\pi^2$, one finds that Eq. (27) implies the lower bound as

$$\frac{\tau_{QSL}^{2,2}}{\tau} = \frac{\|\rho_0\|_2 |\cos \theta_\tau - 1|}{\int_0^\tau dt \left\| \frac{d\rho_t}{dt} \right\|_2} \geq \frac{\|\rho_0\|_2}{\int_0^\tau dt \left\| \frac{d\rho_t}{dt} \right\|_2} \frac{4\theta_\tau^2}{\pi^2}. \quad (29)$$

The quantity $\tau_{QSL}^{CEPH}/\tau := (4\theta_\tau^2/\pi^2)\|\rho_0\|_2/\int_0^\tau dt \|d\rho_t/dt\|_2$ in the right-hand side of Eq. (29) defines the QSL obtained by del Campo *et al.* (CEPH) for the nonunitary evolution in open quantum systems [42]. Therefore, we can conclude that

$$\tau_{QSL}^{2,2} \geq \tau_{QSL}^{CEPH}. \quad (30)$$

Equation (30) shows that, for $\alpha = \beta = 2$, the QSL obtained from the Holder's inequality [see Eq. (23)] is expected to be tighter than the QSL developed by del Campo *et al.* We emphasize that the choice $\alpha = \beta = 2$ is necessary because τ_{QSL}^{CEPH} was originally derived using the Hilbert-Schmidt norm.

Next, we compare the general QSL in Eq. (23) developed from Holder's inequality, with the QSL obtained by Deffner-Lutz [43]. On the one hand, for an initial pure state $\rho_0 = |\psi_0\rangle\langle\psi_0|$, the Deffner-Lutz QSL is given in Eq. (10). On the other hand, for the same initial state, the $\tau_{QSL}^{\alpha,\beta}$ is given in Eq. (24). Note that we have

$$\lim_{\alpha \rightarrow \infty} \tau_{QSL}^{\alpha,\beta} = \tau_{QSL}^{DL}. \quad (31)$$

In particular, for a single-qubit dynamics, these two QSL can be written as

$$\frac{\tau_{QSL}^{DL}}{\tau} = \frac{1 - \vec{n}_\tau \cdot \vec{n}_0}{\int_0^\tau dt \left\| \frac{d\vec{n}_t}{dt} \right\|}, \quad (32)$$

and

$$\frac{\tau_{QSL}^{\alpha,\beta}}{\tau} = 2^{-\frac{1}{\alpha}} \frac{(1 - \vec{n}_\tau \cdot \vec{n}_0)}{\int_0^\tau dt \left\| \frac{d\vec{n}_t}{dt} \right\|}. \quad (33)$$

Therefore, we can conclude that for a qubit dynamics, considering a initial pure state and taking a Schatten norm with $\alpha \in [1, \infty)$, the quantifier $\tau_{QSL}^{\alpha,\beta}$ is related with τ_{QSL}^{DL} by

$$\tau_{QSL}^{\alpha,\beta} = 2^{-\frac{1}{\alpha}} \tau_{QSL}^{DL}. \quad (34)$$

In the next section, we will analyze the necessary and sufficient condition for the (α, β) -QSL to be tight, providing a geometrical interpretation in terms of a optimal path in the Bloch space, and an algebraic one in terms of the populations and coherences of the density operator.

c. Tightness condition of (α, β) -QSL for the dynamics of single-qubit states — Here we address the QSL in Eq. (26) and analyze the necessary and sufficient condition to obtain $\tau_{QSL}^{\alpha,\beta} = \tau$, that is, the necessary and sufficient condition to get the optimal dynamics. In order to develop this condition, we consider the general dynamics of an initial single-qubit pure state $\rho_0 = (1/2)(\mathbb{I} + \vec{n}_0 \cdot \vec{\sigma})$, where \vec{n}_0 is the Bloch vector.

Lemma 2 (Tightness condition in the Bloch space for the (α, β) -QSL). *Let us consider a two-level system with the initial pure state $\rho_0 = |\psi_0\rangle\langle\psi_0|$, while its evolved state becomes $\rho_t = (1/2)(\mathbb{I} + \vec{n}_t \cdot \vec{\sigma}) = \rho_{11,t}|\psi_0\rangle\langle\psi_0| + (1 - \rho_{11,t})|\psi_0^\perp\rangle\langle\psi_0^\perp| + \rho_{12,t}|\psi_0\rangle\langle\psi_0^\perp| + \rho_{12,t}^*|\psi_0^\perp\rangle\langle\psi_0|$. Then, in the orthonormal basis $\{|\psi_0\rangle, |\psi_0^\perp\rangle\}$ the (α, β) -QSL [see Eq. (26)] has the following necessary and sufficient condition for to be tight:*

$$\begin{aligned} \frac{\tau_{QSL}^{\alpha,\beta}}{\tau} = 1 &\Leftrightarrow \left\| \frac{d\vec{n}_t}{dt} \right\| = -2^{-\frac{1}{\alpha}} \left(\vec{n}_0 \cdot \frac{d\vec{n}_t}{dt} \right) \\ &= 2^{-\frac{1}{\alpha}} \left(-\frac{dn_{z,t}}{dt} \right), \end{aligned} \quad (35)$$

with $dn_{z,t}/dt \leq 0$, or equivalently

$$\frac{\tau_{QSL}^{\alpha,\beta}}{\tau} = 1 \Leftrightarrow \Omega = 2^{-\frac{1}{\alpha}} \left(-\frac{d\rho_{11,t}}{dt} \right), \quad (36)$$

where $d\rho_{11,t}/dt = d\langle\psi_0|\rho_t|\psi_0\rangle/dt \leq 0$, with $\|d\vec{n}_t/dt\|$ defined in Eq. (8), and Ω is given in Eq. (18).

Proof. See Supplemental Material, Sec. S.V. \square

Let's now interpret the result of the Lemma 2. From the algebraic point of view, we can see that a necessary condition for have $\tau_{QSL}^{\alpha,\beta} = \tau$ is that the dynamical map is dissipative, that is, the populations of the initial pure state $\rho_0 = |\psi_0\rangle\langle\psi_0|$, namely $\rho_{11,t} = \langle\psi_0|\rho_t|\psi_0\rangle$, needs to be a monotonically decreases function in the time. This result restricts the class of quantum channels for which the $\tau_{QSL}^{\alpha,\beta}$ is tight given a initial pure state.

Taking a geometric point of view, since the initial state is pure, we have $\|\vec{n}_0\| = 1$ and the initial Bloch vector is unit. Therefore, we find that the condition $\tau_{QSL}^{\alpha,\beta}/\tau = 1$ is achieved whenever one gets $\|d\vec{n}_t/dt\| = -2^{-1/\alpha} [\vec{n}_0 \cdot (d\vec{n}_t/dt)]$, which implies that the QSL is tight as long as the dynamics of the single-qubit state satisfies the following constraint

$$\frac{d\vec{n}_t}{dt} = -2^{\frac{1}{\alpha}} \left\| \frac{d\vec{n}_t}{dt} \right\| \vec{n}_0. \quad (37)$$

Equation (37) means that, if the dynamical path in the Bloch space is radial in the Bloch sphere, and it has opposite direction of the initial Bloch vector \vec{n}_0 , then the dynamic is tight, i.e., one finds that $\tau_{QSL}^{\alpha,\beta}/\tau = 1$.

Finally, we address the particular case where $\alpha = \infty$ and $\beta = 1$. As discussed above, in this case and taking a pure initial state, we have that

$$\lim_{\alpha \rightarrow \infty} \tau_{QSL}^{\alpha,\beta} = \tau_{QSL}^{DL}. \quad (38)$$

From the Lemma 2 and since $\|\vec{n}_0\| = 1$, we have that

$$\begin{aligned} \frac{\tau_{QSL}^{\alpha=\infty,\beta=1}}{\tau} = 1 &\Leftrightarrow \left\| \frac{d\vec{n}_t}{dt} \right\| = - \left(\vec{n}_0 \cdot \frac{d\vec{n}_t}{dt} \right) \\ &\Leftrightarrow \frac{d\vec{n}_t}{dt} = - \left\| \frac{d\vec{n}_t}{dt} \right\| \vec{n}_0. \end{aligned} \quad (39)$$

Therefore, for $\alpha = \infty$ and $\beta = 1$, and for a pure initial state, the QSL $\tau_{QSL}^{\infty,1} = \tau_{QSL}^{DL}$ is tight if, and only if, the dynamical path is radial in the Bloch sphere and it has the opposite direction of the initial unit Bloch vector \vec{n}_0 . Moreover, this condition can be seen as the geometric interpretation of tightness of the Deffner-Lutz QSL.

Comparing α -QSL and (α, β) -QSL.— Here we investigate the relationship between QSLs τ_{QSL}^α and $\tau_{QSL}^{\alpha,\beta}$. In order to do that, we state the following result:

Lemma 3 (Inequality between τ_{QSL}^α and $\tau_{QSL}^{\alpha,\beta}$). *For a qubit dynamics, if $\vec{n}_0 \cdot \vec{n}_\tau \leq 0$ and the initial state is pure, then*

$$\tau_{QSL}^\alpha \leq 2^{\frac{1}{2} + \frac{1}{\alpha}} \tau_{QSL}^{\alpha,\beta}. \quad (40)$$

Proof. See Supplemental Material, Sec. S.VI. \square

The inequality in Eq. (40) has an interesting geometrical interpretation. The initial pure state is represented by a point \vec{n}_0 in the surface of the Bloch sphere, since $\|\vec{n}_0\| = 1$. This unit vector divides the Bloch sphere into two hemispheres through the plane orthogonal to this vector. Let us call the northern hemisphere the half of the sphere that has the point \vec{n}_0 , and the southern hemisphere the other half, which has the point $-\vec{n}_0$. In this setting, we have $\vec{n}_\tau \cdot \vec{n}_0 \leq 0$ whenever the final vector \vec{n}_τ belongs to the southern hemisphere, and the result of the Lemma is true in these cases.

Conclusion.— In this work, we discuss two families of QSLs induced by general Schatten norms. The first was obtained through the geometric approach, while the second one from the Holder's inequality. Such QSLs were compared with others speed limits found in literature, and the tightness condition for these two quantifiers was also obtained.

On the one hand, we obtain the α -QSL within a geometric approach, which in turn provides a generalization of the QSL obtained by Campaioli *et al.* [38] for general Schatten norms. We find that the value of such a QSL does not depend on the specific Schatten norm when we are considering a single-qubit dynamics, highlighting a certain notion of universality. Furthermore, it was noted that when the initial state is pure, such

a QSL is tighter than the Deffner-Lutz QSL [43]. On the other hand, (α, β) -QSL was obtained from the Holders' inequality, and can be seen as a generalization of the del Campo *et al.* QSL [42] to general Schatten norms and also that it is tighter. Furthermore, (α, β) -QSL recovers the Deffner-Lutz QSL when we take the operator norm and that this is proportional to this one in the case of qubit dynamics.

We investigate the necessary and sufficient conditions for such QSL induced from Schatten norms to be tight under a general qubit dynamics. We obtain conditions with geometric interpretations in terms of the dynamic path in the Bloch space, just as we obtain the constraints on the behavior of the populations and coherences of the dynamic state of the qubit, thus reflecting an algebraic condition.

Finally, we compared the two QSL developed in this work for the case of a qubit system. We found an inequality constraint between them that holds whenever the initial state is pure. This link exhibits a geometric interpretation in terms of the initial and final vectors in Bloch space. In addition to obtaining a family of QSLs, these quantifiers generalizes previous QSLs discussed in the literature and which were developed under completely different approaches. This again reinforces the notion of universality between the different types of QSL.

Note added.— After completion of this work, a paper has appeared [61] reporting coherent QSL bounds based on Schatten α -norms. We note that their approach is different from ours, and their results only apply to closed quantum systems. In addition, α -QSL and (α, β) -QSL has been not achieved in that paper.

Acknowledgments.— This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior–Brasil (CAPES) (Finance Code 001). D. O. S. P acknowledges the support by the Brazilian funding agencies CNPq (Grant No. 304891/2022-3), FAPESP (Grant No. 2017/03727-0) and the Brazilian National Institute of Science and Technology of Quantum Information (INCT/IQ). D. P. P. also acknowledges Fundação de Amparo à Pesquisa e ao Desenvolvimento Científico e Tecnológico do Maranhão (FAPEMA).

-
- [1] S. Deffner and E. Lutz, Generalized Clausius Inequality for Nonequilibrium Quantum Processes, *Phys. Rev. Lett.* **105**, 170402 (2010).
 - [2] E. Aghion and J. R. Green, Thermodynamic speed limits for mechanical work, *J. Phys. A: Math. Theor.* **56**, 05LT01 (2023).
 - [3] B. Mohan and A. K. Pati, Reverse quantum speed limit: How slowly a quantum battery can discharge, *Phys. Rev. A* **104**, 042209 (2021).
 - [4] Y. Hasegawa, Unifying speed limit, thermodynamic uncertainty relation and Heisenberg principle via bulk-boundary correspondence, *Nat. Commun.* **14**, 2828 (2023).
 - [5] T. Fogarty, S. Deffner, T. Busch, and S. Campbell, Orthogonality Catastrophe as a Consequence of the Quantum Speed Limit, *Phys. Rev. Lett.* **124**, 110601 (2020).
 - [6] A. del Campo, Probing Quantum Speed Limits with Ultracold Gases, *Phys. Rev. Lett.* **126**, 180603 (2021).
 - [7] R. Puebla, S. Deffner, and S. Campbell, Kibble-Zurek scaling in quantum speed limits for shortcuts to adiabaticity, *Phys. Rev. Research* **2**, 032020 (2020).
 - [8] D. P. Pires and T. R. de Oliveira, Relative purity, speed of fluctuations, and bounds on equilibration times, *Phys. Rev. A* **104**, 052223 (2021).
 - [9] L. P. García-Pintos, S. B. Nicholson, J. R. Green, A. del Campo, and A. V. Gorshkov, Unifying Quantum and Classical Speed Limits on Observables, *Phys. Rev. X* **12**, 011038 (2022).
 - [10] V. Giovannetti, S. Lloyd, and L. Maccone, Advances in quantum metrology, *Nature Photon.* **5**, 222 (2011).
 - [11] J. D. Bekenstein, Energy Cost of Information Transfer, *Phys. Rev. Lett.* **46**, 623 (1981).
 - [12] S. Lloyd, Ultimate physical limits to computation, *Nature* **406**, 1047 (2000).
 - [13] V. Giovannetti, S. Lloyd, and L. Maccone, The role of entan-

- glement in dynamical evolution, *EPL* **62**, 615 (2003).
- [14] J. Batle, M. Casas, A. Plastino, and A. R. Plastino, Connection between entanglement and the speed of quantum evolution, *Phys. Rev. A* **72**, 032337 (2005).
- [15] J. Batle, M. Casas, A. Plastino, and A. R. Plastino, Erratum: Connection between entanglement and the speed of quantum evolution [Phys. Rev. A 72, 032337 (2005)], *Phys. Rev. A* **73**, 049904 (2006).
- [16] F. Centrone and M. Gessner, Breaking local quantum speed limits with steering, [arXiv:2401.04599](https://arxiv.org/abs/2401.04599) (2024).
- [17] B. Shanahan, A. Chenu, N. Margolus, and A. del Campo, Quantum Speed Limits across the Quantum-to-Classical Transition, *Phys. Rev. Lett.* **120**, 070401 (2018).
- [18] M. Okuyama and M. Ohzeki, Quantum Speed Limit is Not Quantum, *Phys. Rev. Lett.* **120**, 070402 (2018).
- [19] R. Hamazaki, Speed Limits for Macroscopic Transitions, *PRX Quantum* **3**, 020319 (2022).
- [20] T. Caneva, M. Murphy, T. Calarco, R. Fazio, S. Montangero, V. Giovannetti, and G. E. Santoro, Optimal Control at the Quantum Speed Limit, *Phys. Rev. Lett.* **103**, 240501 (2009).
- [21] K. Kobayashi and N. Yamamoto, Quantum speed limit for robust state characterization and engineering, *Phys. Rev. A* **102**, 042606 (2020).
- [22] J.-Y. Gyhm, D. Rosa, and D. Šafránek, The minimal time it takes to charge a quantum system, [arXiv:2308.16086](https://arxiv.org/abs/2308.16086) (2023).
- [23] F. Impens, F. M. D'Angelis, F. A. Pinheiro, and D. Guéry-Odelin, Time scaling and quantum speed limit in non-Hermitian Hamiltonians, *Phys. Rev. A* **104**, 052620 (2021).
- [24] Y.-Y. Wang and M.-F. Fang, Quantum speed limit time of a non-Hermitian two-level system, *Chinese Phys. B* **29**, 030304 (2020).
- [25] D. P. Pires, Unified entropies and quantum speed limits for nonunitary dynamics, *Phys. Rev. A* **106**, 012403 (2022).
- [26] G. Ness, M. R. Lam, W. Alt, D. Meschede, Y. Sagi, and A. Alberti, Observing crossover between quantum speed limits, *Sci. Adv.* **7**, 9119 (2021).
- [27] D. V. Villamizar, E. I. Duzzioni, A. C. S. Leal, and R. Auccaise, Estimating the time evolution of NMR systems via a quantum-speed-limit-like expression, *Phys. Rev. A* **97**, 052125 (2018).
- [28] D. P. Pires, E. R. deAzevedo, D. O. Soares-Pinto, F. Brito, and J. G. Filgueiras, Experimental Investigation of Geometric Quantum Speed Limits in an Open Quantum System, [arXiv:quant-ph/9705052](https://arxiv.org/abs/quant-ph/9705052) (2023).
- [29] S. Deffner and S. Campbell, Quantum speed limits: from Heisenberg's uncertainty principle to optimal quantum control, *J. Phys. A: Math. Theor.* **50**, 453001 (2017).
- [30] Z. Gong and R. Hamazaki, Bounds in nonequilibrium quantum dynamics, *Int. J. Mod. Phys. B* **36**, 2230007 (2022).
- [31] L. Mandelstam and I. Tamm, The uncertainty relation between energy and time in non-relativistic quantum mechanics, in *Selected Papers*, edited by B. M. Bolotovskii, V. Y. Frenkel, and R. Peierls (Springer, Berlin, Heidelberg, 1991) pp. 115–123.
- [32] N. Margolus and L. B. Levitin, The maximum speed of dynamical evolution, *Physica D* **120**, 188 (1998).
- [33] G. Ness, A. Alberti, and Y. Sagi, Quantum Speed Limit for States with a Bounded Energy Spectrum, *Phys. Rev. Lett.* **129**, 140403 (2022).
- [34] L. B. Levitin and T. Toffoli, Fundamental Limit on the Rate of Quantum Dynamics: The Unified Bound Is Tight, *Phys. Rev. Lett.* **103**, 160502 (2009).
- [35] V. Giovannetti, S. Lloyd, and L. Maccone, Quantum limits to dynamical evolution, *Phys. Rev. A* **67**, 052109 (2003).
- [36] D. Mondal, C. Datta, and S. Sazim, Quantum coherence sets the quantum speed limit for mixed states, *Phys. Lett. A* **380**, 689 (2016).
- [37] D. P. Pires, K. Modi, and L. C. Céleri, Bounding generalized relative entropies: Nonasymptotic quantum speed limits, *Phys. Rev. E* **103**, 032105 (2021).
- [38] F. Campaioli, F. A. Pollock, and K. Modi, Tight, robust, and feasible quantum speed limits for open dynamics, *Quantum* **3**, 168 (2019).
- [39] N. Hörnedal, D. Allan, and O. Sönnernborn, Extensions of the Mandelstam-Tamm quantum speed limit to systems in mixed states, *New J. Phys.* **24**, 055004 (2022).
- [40] N. Hörnedal and O. Sönnernborn, Margolus-Levitin quantum speed limit for an arbitrary fidelity, *Phys. Rev. Res.* **5**, 043234 (2023).
- [41] M. M. Taddei, B. M. Escher, L. Davidovich, and R. L. de Matos Filho, Quantum Speed Limit for Physical Processes, *Phys. Rev. Lett.* **110**, 050402 (2013).
- [42] A. del Campo, I. L. Egusquiza, M. B. Plenio, and S. F. Huelga, Quantum Speed Limits in Open System Dynamics, *Phys. Rev. Lett.* **110**, 050403 (2013).
- [43] S. Deffner and E. Lutz, Quantum Speed Limit for Non-Markovian Dynamics, *Phys. Rev. Lett.* **111**, 010402 (2013).
- [44] Z. Sun, J. Liu, J. Ma, and X. Wang, Quantum speed limits in open systems: Non-Markovian dynamics without rotating-wave approximation, *Sci. Rep.* **5**, 8444 (2015).
- [45] X. Meng, C. Wu, and H. Guo, Minimal evolution time and quantum speed limit of non-Markovian open systems, *Sci. Rep.* **5**, 16357 (2015).
- [46] N. Mirkin, F. Toscano, and D. A. Wisniacki, Quantum-speed-limit bounds in an open quantum evolution, *Phys. Rev. A* **94**, 052125 (2016).
- [47] M. Cianciaruso, S. Maniscalco, and G. Adesso, Role of non-Markovianity and backflow of information in the speed of quantum evolution, *Phys. Rev. A* **96**, 012105 (2017).
- [48] J. Teittinen, H. Lyyra, and S. Maniscalco, There is no general connection between the quantum speed limit and non-Markovianity, *New J. Phys.* **21**, 123041 (2019).
- [49] J. Teittinen and S. Maniscalco, Quantum Speed Limit and Divisibility of the Dynamical Map, *Entropy* **23**, 331 (2021).
- [50] I. Bengtsson and K. Życzkowski, *Geometry of Quantum States: An Introduction to Quantum Entanglement* (Cambridge University Press, Cambridge, 2006).
- [51] D. P. Pires, M. Cianciaruso, L. C. Céleri, G. Adesso, and D. O. Soares-Pinto, Generalized Geometric Quantum Speed Limits, *Phys. Rev. X* **6**, 021031 (2016).
- [52] E. O'Connor, G. Guarnieri, and S. Campbell, Action quantum speed limits, *Phys. Rev. A* **103**, 022210 (2021).
- [53] K. Lan, S. Xie, and X. Cai, Geometric quantum speed limits for Markovian dynamics in open quantum systems, *New J. Phys.* **24**, 055003 (2022).
- [54] B. Russell and S. Stepney, Applications of Finsler geometry to speed limits to quantum information processing, *Int. J. Found. Comput. Sci.* **25**, 489 (2014).
- [55] J. Anandan and Y. Aharonov, Geometry of quantum evolution, *Phys. Rev. Lett.* **65**, 1697 (1990).
- [56] S. Deffner, Geometric quantum speed limits: a case for Wigner phase space, *New J. Phys.* **19**, 103018 (2017).
- [57] M. Gessner and A. Smerzi, Statistical speed of quantum states: Generalized quantum Fisher information and Schatten speed, *Phys. Rev. A* **97**, 022109 (2018).
- [58] K. Funo, N. Shiraishi, and K. Saito, Speed limit for open quantum systems, *New J. Phys.* **21**, 013006 (2019).
- [59] T. Van Vu and Y. Hasegawa, Geometrical Bounds of the Irreversibility in Markovian Systems, *Phys. Rev. Lett.* **126**, 010601 (2021).

- [60] S. Nakajima and Y. Utsumi, Speed limits of the trace distance for open quantum system, [New J. Phys. **24**, 095004 \(2022\)](#).
- [61] H. Wang and X. Qiu, Generalized Coherent Quantum Speed Limits, [arXiv:2401.01746 \(2024\)](#).

Supplemental Material

Quantum Speed Limits Based on Schatten Norms: Universality and Tightness

Alberto J. B. Rosal, Diogo O. Soares-Pinto, and Diego Paiva Pires

S.I. PRELIMINARIES AND DEFINITIONS

Let us start with some definitions about Schatten α -norms, followed by the metric space structure for the state space, and the Bloch representation of quantum states. These mathematical tools are very important in the discussion developed in this work.

S.I.1. Normed spaces and Schatten norms

Let us consider the Hilbert space \mathcal{H} and we will denote $L(\mathcal{H})$ as the set of every linear transformations on this space. In fact, $L(\mathcal{H})$ is a complex vectorial space and this one can be transformed in a normed space with the help of the Schatten α -norms:

Definition S1 (Schatten norms). *For every linear operator $A \in L(\mathcal{H})$ acting on a Hilbert space \mathcal{H} , we define its Schatten α -norm by:*

$$\|A\|_\alpha := (\text{Tr}\{(\sqrt{A^\dagger A})^\alpha\})^{\frac{1}{\alpha}}, \quad (\text{S1})$$

where $\alpha \in [1, \infty)$.

Some values of α for the Schatten norms defined in the Definition S1 are very common in the study of the space of quantum states. In particular, for $\alpha = 1$ we have the trace norm (or L_1 -norm), for $\alpha = 2$ we have the Hilbert-Schmidt (or L_2 -norm) norm and finally for the limit $\alpha \rightarrow \infty$ we have the operator norm (or spectral norm). We can compute these Schatten norms through the singular values. We point out that the singular values of some linear operator $A \in L(\mathcal{H})$ are given by the eigenvalues of the positive operator $\sqrt{A^\dagger A}$. In this way, the Schatten α -norm of the operator A can be given by

$$\|A\|_\alpha = \left(\sum_{i=1}^r \sigma(A)_i^\alpha \right)^{\frac{1}{\alpha}}, \quad (\text{S2})$$

where $\sigma(A)_i$ are the singular values of the operator A , and $r = \text{rank}(A)$ is the rank of A . Through Eq. (S2), we have that the trace norm, Hilbert-Schmidt norm, and operator norm can be written as

$$\|A\|_1 = \sum_{j=1}^r \sigma(A)_j, \quad \|A\|_2 = \sqrt{\sum_{j=1}^r \sigma(A)_j^2}, \quad \|A\|_\infty = \max_j \{\sigma(A)_j\}, \quad (\text{S3})$$

respectively. Now, since the singular values are the eigenvalues of $\sqrt{A^\dagger A}$, if the operator A is Hermitian (self-adjoint), $A^\dagger = A$, then the eigenvalues of A , denoted by $\lambda(A)$, are related with the singular values $\{\sigma(A)_j\}_{j=1, \dots, r}$ as follows

$$\sigma(A)_j = |\lambda(A)_j|. \quad (\text{S4})$$

In particular, note that if the operator A is positive semi-definite, then the singular values are equals to the eigenvalues.

The Schatten α -norms have a large number of interesting properties. In particular, we have the follow result:

Theorem S1 (Duality of the Schatten norms). *For every linear operator A , the Schatten norm can be given as a supremum:*

$$\|A\|_\alpha = \sup_{\chi \neq 0} \left\{ |\text{Tr}[\chi^\dagger A]| : \|\chi\|_\beta \leq 1, \frac{1}{\alpha} + \frac{1}{\beta} = 1 \right\}. \quad (\text{S5})$$

In particular, due to the Theorem S1, we have the Holder's inequality, defined as

$$|\text{Tr}[Y^\dagger X]| \leq \|X\|_\alpha \|Y\|_\beta, \quad (\text{S6})$$

which holds for every two linear operators $X, Y \in L(\mathcal{H})$ and for every real numbers $\alpha, \beta \in [1, \infty)$ where $\frac{1}{\alpha} + \frac{1}{\beta} = 1$. In this sense, the norms $\|\cdot\|_\alpha$ and $\|\cdot\|_\beta$ are said to be *dual* to each other. The Holder's inequality in Eq. (S6) is the fundamental inequality that implied the (α, β) -QSL discussed in the main text.

S.I.2. Metric spaces and space of quantum states

Since $L(\mathcal{H})$ is in fact a complex vectorial space, we can write its elements in a particular chosen basis. This idea is the core for the Bloch representation of quantum states (density operators). In this way, we can see density operators as vectors in the Euclidean space, namely the Bloch space. Here, let us denote $\mathcal{D}(\mathcal{H})$ as the set of every quantum state ρ associated to the Hilbert space \mathcal{H} . Obviously $\mathcal{D}(\mathcal{H}) \subseteq L(\mathcal{H})$, since every density operator is also a linear operator acting on the Hilbert space \mathcal{H} . Then, the set $\mathcal{D}(\mathcal{H})$ will be called state space of the Hilbert space \mathcal{H} . In fact, the space of states equipped with a Schatten α -norm is not only a normed space, but it is also a metric space. Remember that a metric space is an ordered pair (M, D) where M is a set and D is a metric on M , where $D : M \times M \rightarrow \mathbb{R}$ is a function such that, for every $x, y, z \in M$, we have:

- **Positivity semi-definite:** $D(x, y) \geq 0$, with $D(x, y) = 0$ if and only if $x = y$.
- **Symmetry:** $D(x, y) = D(y, x)$.
- **Triangle inequality:** $D(x, z) \leq D(x, y) + D(y, z)$.

In this way, taking the set $M = \mathcal{D}(\mathcal{H})$, and considering the α -norm in the state space, we can define the metric

$$D_\alpha(\rho, \eta) = \|\rho - \eta\|_\alpha, \quad (\text{S7})$$

and then the state space becomes an Euclidean metric space. Therefore, an normed state space $\mathcal{D}(\mathcal{H})$ equipped with a Schatten α -norm becomes, in fact, an authentic metric space. We can work with elements of the state space in a very useful fashion, called Bloch representation. This approach provides a useful geometric interpretation for two-dimensional quantum systems.

S.I.3. General qudit systems and Bloch representation

Now, let us consider a finite dimensional Hilbert space \mathcal{H} with dimension d , and let $\{F_\alpha\}_{\alpha=0}^{d^2-1}$ be an orthonormal traceless Hermitian basis in $L(\mathcal{H})$, that is $\text{Tr}(F_\alpha F_\beta) = \delta_{\alpha,\beta}$, together with $F_0 = \mathbb{1}/\sqrt{d}$ and $F_\alpha^\dagger = F_\alpha$, as well as $\text{Tr}(F_\alpha) = 0$ for every $\alpha = 1, 2, \dots, d^2 - 1$. In this way, any quantum state (density operator) ρ of the Hilbert Space \mathcal{H} can be represented by

$$\rho_t = \frac{1}{d} \left(\mathbb{1} + \vec{x}_t \cdot \vec{F} \right), \quad (\text{S8})$$

where $\vec{x}_t \cdot \vec{F} = \sum_{i=1}^{d^2-1} x_{i,t} F_i$ sets for the standard inner product, while $x_{j,t} = \text{Tr}(F_j \rho_t)$, where $\vec{x}_t = (x_{1,t}, \dots, x_{d^2-1,t}) \in \mathbb{R}^{d^2-1}$, and $\vec{F} = (F_1, \dots, F_{d^2-1})$. In this setting, every density operator ρ can be represented by a vector \vec{x} belonging to a subset of the Euclidean space \mathbb{R}^{d^2-1} . The subset of \mathbb{R}^{d^2-1} of every accessible states is called Bloch space. For a qubit system, the set of accessible states has an spherical geometry, and it is called Bloch sphere. For systems in which the dimension d is greater than two, the geometry of the Bloch space can becomes very complicated. Actually, note that the operators F_α are the generators of the $SU(d)$ Lie algebra.

S.II. SINGLE-QUBIT SYSTEMS

S.II.1. General single-qubit state and the Bloch representation

For a two-dimensional Hilbert space, we have the Pauli matrix operators as a basis for $\mathcal{D}(\mathcal{H})$. In this case, we have $\rho_t = (1/2)(\mathbb{1} + \vec{n}_t \cdot \vec{\sigma})$, where $\vec{n}_t = (n_{x,t}, n_{y,t}, n_{z,t}) \in \mathbb{R}^3$ is a time-dependent three-dimensional vector, and $\vec{\sigma} = (\sigma_x, \sigma_y, \sigma_z)$ is the vector of Pauli matrices. We remember that the density matrix is Hermitian, $\rho_t^\dagger = \rho_t$, positive semi-definite, $\rho_t \geq 0$, with trace equal to the unit, $\text{Tr}(\rho_t) = 1$, for all $t \geq 0$. Here, we set the eigenbasis $\{|0\rangle, |1\rangle\}$ of the operator σ_z for the two-dimensional Hilbert space \mathcal{H} , with $\sigma_z |\ell\rangle = (-1)^\ell |\ell\rangle$, and $\ell = \{0, 1\}$. With respect to this eigenbasis, it follows that the aforementioned single-qubit state is written as follows

$$\rho_t = \frac{1}{2} \begin{pmatrix} 1 + n_{z,t} & n_{x,t} - in_{y,t} \\ n_{x,t} + in_{y,t} & 1 - n_{z,t} \end{pmatrix}. \quad (\text{S9})$$

The eigenvalues of the density matrix in Eq. (S9) can be written as

$$\lambda_\pm(t) = \frac{1}{2}(1 \pm \|\vec{n}_t\|), \quad (\text{S10})$$

where $\|\vec{n}_t\| = \sqrt{n_{x,t}^2 + n_{y,t}^2 + n_{z,t}^2}$ is the Euclidean norm of the vector \vec{n}_t , with $\|\vec{n}_t\| \leq 1$. The Bloch space is a subset of the three-dimensional sphere in the Euclidean space \mathbb{R}^3 . We note that the density matrix ρ_t exhibits the singular values $\sigma_{\pm} = (1/2)(1 \pm \|\vec{n}_t\|)$ [see Eq. (S4)]. Finally, since we know the singular values of ρ_t , we can compute its Schatten α -norm as [see Eq. (S2)]

$$\|\rho_t\|_{\alpha} = \frac{1}{2} [(1 - \|\vec{n}_t\|)^{\alpha} + (1 + \|\vec{n}_t\|)^{\alpha}]^{\frac{1}{\alpha}}. \quad (\text{S11})$$

Note that both populations and coherences of the density matrix ρ_t can be written in terms of the elements $\{n_{j,t}\}_{j=x,y,z}$ of the Bloch vector \vec{n}_t , and vice-versa. On the one hand, by writing the density matrix in the eigenbasis $\{|0\rangle, |1\rangle\}$, we have that $\rho_t = p_t|0\rangle\langle 0| + (1 - p_t)|1\rangle\langle 1| + q_t|0\rangle\langle 1| + q_t^*|1\rangle\langle 0|$, where $p_t = \langle 0|\rho_t|0\rangle \in [0, 1]$ is the population of the state $|0\rangle$, and $q_t = \langle 0|\rho_t|1\rangle$ is the coherence between vector states $|0\rangle$ and $|1\rangle$. On the other hand, from Eq. (S9), we find that $p_t = (1/2)(1 + n_{z,t})$, and $q_t = (1/2)(n_{x,t} - i n_{y,t})$, which readily implies the relations $n_{z,t} = 2p_t - 1$, $n_{x,t} = 2 \operatorname{Re}(q_t)$, and $n_{y,t} = -2 \operatorname{Im}(q_t)$.

Next, let us suppose that the qubit has a time-differentiable dynamical map $\Lambda_{(t,0)}$ whose the initial state is ρ_0 . In the Bloch representation, taking the eigenbasis of σ_z for the two-dimensional Hilbert space \mathcal{H} , we find that

$$\frac{d\rho_t}{dt} = \frac{1}{2} \frac{d\vec{n}_t}{dt} \cdot \vec{\sigma} = \frac{1}{2} \begin{pmatrix} dn_{z,t}/dt & dn_{x,t}/dt - i dn_{y,t}/dt \\ dn_{x,t}/dt + i dn_{y,t}/dt & -dn_{z,t}/dt \end{pmatrix}. \quad (\text{S12})$$

The eigenvalues of the operator $d\rho_t/dt$ are given by $\gamma_{\pm} = \pm(1/2) \|d\vec{n}_t/dt\|$, where we define the Euclidean norm $\|d\vec{n}_t/dt\|^2 = \sum_{j=x,y,z} (dn_{j,t}/dt)^2$. The singular values of this operator are given by $\omega_1 = \omega_2 = (1/2) \|d\vec{n}_t/dt\|$. Hence, Eq. (S2) implies the following result

$$\left\| \frac{d\rho_t}{dt} \right\|_{\alpha} = 2^{-1+\frac{1}{\alpha}} \left\| \frac{d\vec{n}_t}{dt} \right\|. \quad (\text{S13})$$

Equation (S13) means that the Schatten α -norm $\|d\rho_t/dt\|_{\alpha}$ in the state space is proportional to the Euclidean norm $\|d\vec{n}_t/dt\|$ in the Bloch space.

S.II.2. The space of single-qubit states as a metric space

Let us compute the distance $D_{\alpha}(\rho_t, \eta_t) = \|\rho_t - \eta_t\|_{\alpha}$ [see Eq. (S7)] between the single-qubit states $\rho_t, \eta_t \in \mathcal{D}(\mathcal{H})$, with $\rho_t = (1/2)(\mathbb{I} + \vec{n}_t \cdot \vec{\sigma})$, and $\eta_t = (1/2)(\mathbb{I} + \vec{m}_t \cdot \vec{\sigma})$. First, we note that

$$\rho_t - \eta_t = \frac{1}{2}(\vec{n}_t - \vec{m}_t) \cdot \vec{\sigma}, \quad (\text{S14})$$

where $\vec{n}_t = (n_{x,t}, n_{y,t}, n_{z,t})$, and $\vec{m}_t = (m_{x,t}, m_{y,t}, m_{z,t})$ are the Bloch vectors of ρ_t and η_t , respectively. We see that Eqs. (S12) and (S14) are similar each other for the single-qubit dynamics. Hence, the eigenvalues of the operator $\rho_t - \eta_t$ and the respective singular values can be obtained in the same way. In particular, the Schatten α -norm of the operator $\rho_t - \eta_t$ [that is, the distance $D_{\alpha}(\rho_t, \eta_t)$] is given by

$$D_{\alpha}(\rho_t, \eta_t) = \|\rho_t - \eta_t\|_{\alpha} = 2^{-1+\frac{1}{\alpha}} d_{\text{euc}}(\vec{n}_t, \vec{m}_t), \quad (\text{S15})$$

where

$$d_{\text{euc}}(\vec{n}_t, \vec{m}_t) = \sqrt{(n_{x,t} - m_{x,t})^2 + (n_{y,t} - m_{y,t})^2 + (n_{z,t} - m_{z,t})^2}. \quad (\text{S16})$$

Equation (S15) means that the distance D_{α} in the state space $\mathcal{D}(\mathcal{H})$ induced from the α -norm is proportional to the Euclidean distance d_{euc} in the Bloch space.

S.III. PROOF OF THEOREM 1

The quantum speed limit time induced by the α -Schatten norm, for a time of evolution τ , is given by

$$\tau_{QSL}^{\alpha}(\tau) := \frac{\|\rho_{\tau} - \rho_0\|_{\alpha}}{\int_0^{\tau} dt \left\| \frac{d\rho_t}{dt} \right\|_{\alpha}} \tau. \quad (\text{S17})$$

Given the single-qubit states $\rho_0 = (1/2)(\mathbb{I} + \vec{n}_0 \cdot \vec{\sigma})$ and $\rho_\tau = (1/2)(\mathbb{I} + \vec{n}_\tau \cdot \vec{\sigma})$, one can verify the identity [see Sec. S.II]

$$\|\rho_\tau - \rho_0\|_\alpha = 2^{-1+\frac{1}{\alpha}} d_{\text{euc}}(\vec{n}_0, \vec{n}_\tau), \quad (\text{S18})$$

with the Euclidean distance defined as

$$d_{\text{euc}}(\vec{n}_0, \vec{n}_\tau) = \sqrt{\sum_{j=x,y,z} (n_{j,\tau} - n_{j,0})^2}. \quad (\text{S19})$$

We also note that

$$\left\| \frac{d\rho_t}{dt} \right\|_\alpha = 2^{-1+\frac{1}{\alpha}} \left\| \frac{d\vec{n}_t}{dt} \right\|, \quad (\text{S20})$$

with

$$\left\| \frac{d\vec{n}_t}{dt} \right\| = \sqrt{\sum_{j=x,y,z} \left(\frac{dn_{j,t}}{dt} \right)^2}. \quad (\text{S21})$$

Hence, by combining Eqs. (S17), (S18), and (S20), one gets the result

$$\tau_{QSL}^\alpha(\tau) = \frac{d_{\text{euc}}(\vec{n}_0, \vec{n}_\tau)}{\int_0^\tau dt \left\| \frac{d\vec{n}_t}{dt} \right\|} \tau, \quad (\text{S22})$$

which concludes the proof of Theorem 1.

S.IV. PROOF OF LEMMA 1

Here we discuss the conditions in which the α -QSL in Eq. (S22) for single-qubit states is tight, i.e., $\tau_{QSL}^\alpha(\tau)/\tau = 1$. To do so, one should satisfies the constraint as follows

$$d_{\text{euc}}(\vec{n}_0, \vec{n}_\tau) = \int_0^\tau dt \left\| \frac{d\vec{n}_t}{dt} \right\|, \quad (\text{S23})$$

where the Euclidean distance $d_{\text{euc}}(\vec{n}_0, \vec{n}_\tau)$ is defined in Eq. (S19). By differentiating both sides of Eq. (S23) with respect to the parameter τ , one obtains the result

$$\left(\hat{r}_{0,t} \cdot \frac{d\vec{n}_t}{dt} \right)_{t=\tau} = \left(\left\| \frac{d\vec{n}_t}{dt} \right\| \right)_{t=\tau}, \quad (\text{S24})$$

where we have defined the unit vector

$$\hat{r}_{0,t} := \frac{\vec{n}_t - \vec{n}_0}{\|\vec{n}_t - \vec{n}_0\|}. \quad (\text{S25})$$

We note that Eq. (S24) holds for all $t \geq 0$ (or even $\tau \geq 0$), and from now on we deliberately work with such an equation written in terms of the parameter t . In addition, Eq. (S24) can be solved in terms of the quantity $\|d\vec{n}_t/dt\|$ as long as one chooses the vector $d\vec{n}_t/dt$ as follows

$$\frac{d\vec{n}_t}{dt} = \left\| \frac{d\vec{n}_t}{dt} \right\| \hat{r}_{0,t}, \quad (\text{S26})$$

where we used the fact that $\hat{r}_{0,t} \cdot \hat{r}_{0,t} = 1$, for all $t \geq 0$. Therefore, Eq. (S26) means that $\tau_{QSL}^\alpha = \tau$ if and only if the dynamical path in the Bloch space is given by the straight line connecting the initial Bloch vector \vec{n}_0 to the final one \vec{n}_τ . Next, we note that Eq. (S26) can be recasted in terms of the populations and quantum coherences of the evolved state ρ_t respective to the computational eigenbasis $\{|0\rangle, |1\rangle\}$. To do so, we use the fact that $n_{z,t} = 2\rho_{11,t} - 1$, $n_{x,t} = 2\text{Re}(\rho_{12,t})$, and $n_{y,t} = -2\text{Im}(\rho_{12,t})$ for single-qubit states. In this case, one readily obtains

$$\begin{aligned} \frac{d\vec{n}_t}{dt} &= (dn_{x,t}/dt, dn_{y,t}/dt, dn_{z,t}/dt) \\ &= 2(\text{Re}(d\rho_{12,t}/dt), -\text{Im}(d\rho_{12,t}/dt), d\rho_{11,t}/dt). \end{aligned} \quad (\text{S27})$$

Next, Eq. (S27) implies that the Euclidean norm $\|d\vec{n}_t/dt\|$ is written as follows

$$\left\| \frac{d\vec{n}_t}{dt} \right\| = \sqrt{\sum_{j=x,y,z} \left(\frac{dn_{j,t}}{dt} \right)^2} = 2\Omega, \quad (\text{S28})$$

where we have defined the quantity

$$\Omega := \sqrt{\left| \frac{d\rho_{11,t}}{dt} \right|^2 + \left| \frac{d\rho_{12,t}}{dt} \right|^2} \quad (\text{S29})$$

Therefore, by combining Eqs. (S26), (S27) and (S28), we find that

$$(\text{Re}(d\rho_{12,t}/dt), -\text{Im}(d\rho_{12,t}/dt), d\rho_{11,t}/dt) = \Omega \hat{r}_{0,t}, \quad (\text{S30})$$

and one concludes the proof of Lemma 1.

S.V. PROOF OF LEMMA 2

In order to achieve the tighter (α, β) -QSL, i.e., $\tau_{QSL}^{\alpha, \beta}/\tau = 1$, we require that

$$2^{-\frac{1}{\alpha}}(1 - \vec{n}_\tau \cdot \vec{n}_0) = \int_0^\tau dt \left\| \frac{d\vec{n}_t}{dt} \right\|. \quad (\text{S31})$$

We note that Eq. (S31) is satisfied as long as one sets the Euclidean norm as

$$\left\| \frac{d\vec{n}_t}{dt} \right\| = -2^{-\frac{1}{\alpha}} \left(\vec{n}_0 \cdot \frac{d\vec{n}_t}{dt} \right). \quad (\text{S32})$$

In particular, we hereafter fix the orthonormal basis $\{|\psi_0\rangle, |\psi_0^\perp\rangle\}$. In this setting, one gets the vector $\vec{n}_0 = (0, 0, 1)$, which in turn implies that $\vec{n}_0 \cdot (d\vec{n}_t/dt) = dn_{z,t}/dt$. Hence, Eq. (S32) becomes

$$\left\| \frac{d\vec{n}_t}{dt} \right\| = 2^{-\frac{1}{\alpha}} \left(-\frac{dn_{z,t}}{dt} \right). \quad (\text{S33})$$

We note that, since $\|d\vec{n}_t/dt\| \geq 0$, we thus necessarily have that $dn_{z,t}/dt \leq 0$ in Eq. (S33). From Sec. S.IV, we know that $dn_{z,t}/dt = 2 d\rho_{11,t}/dt = 2 d\langle \psi_0 | \rho_t | \psi_0 \rangle / dt$ [see Eq. (S27)], and also $\|d\vec{n}_t/dt\| = 2\Omega$ [see Eqs. (S28) and (S29)]. Finally, by substituting these results in Eq. (S33), one obtains that

$$\Omega = 2^{-\frac{1}{\alpha}} \left(-\frac{d\rho_{11,t}}{dt} \right), \quad \text{with} \quad \frac{d\rho_{11,t}}{dt} = \frac{d}{dt} \langle \psi_0 | \rho_t | \psi_0 \rangle \leq 0. \quad (\text{S34})$$

In conclusion, Eq. (S34) proved the Lemma 2.

S.VI. PROOF OF LEMMA 3

We consider a two-level system initialized in a single-qubit pure state that undergoes a general evolution. In this setting, given the dynamics between states $\rho_0 = (1/2)(\mathbb{I} + \vec{n}_0 \cdot \vec{\sigma})$ and $\rho_\tau = (1/2)(\mathbb{I} + \vec{n}_\tau \cdot \vec{\sigma})$, we proved the geometric α -QSL time as

$$\frac{\tau_{QSL}^\alpha}{\tau} = \frac{d_{\text{euc}}(\vec{n}_0, \vec{n}_\tau)}{\int_0^\tau dt \|d\vec{n}_t/dt\|}, \quad (\text{S35})$$

and also the (α, β) -QSL time as

$$\frac{\tau_{QSL}^{\alpha, \beta}}{\tau} = 2^{-\frac{1}{\alpha}} \frac{(1 - \vec{n}_\tau \cdot \vec{n}_0)}{\int_0^\tau dt \|d\vec{n}_t/dt\|} \quad (\text{S36})$$

that is based on Holder's inequality, where $d_{\text{euc}}(\vec{n}_0, \vec{n}_\tau) = \|\vec{n}_\tau - \vec{n}_0\|$ is the Euclidean distance between the initial and final states, with $\|\vec{x}\| = \sqrt{\vec{x} \cdot \vec{x}}$ for every vector \vec{x} in the Bloch space. We note that the Euclidean distance can be written as follows

$$d_{\text{euc}}(\vec{n}_0, \vec{n}_\tau) = \sqrt{\|\vec{n}_0\|^2 + \|\vec{n}_\tau\|^2 - 2\vec{n}_0 \cdot \vec{n}_\tau}. \quad (\text{S37})$$

Because $\|\vec{n}_t\| \leq 1$ for all $t \in [0, \tau]$, one gets $d_{\text{euc}}(\vec{n}_0, \vec{n}_\tau) \leq \sqrt{2}\sqrt{1 - \vec{n}_\tau \cdot \vec{n}_0}$. In particular, by considering the case $\vec{n}_\tau \cdot \vec{n}_0 \leq 0$, one gets that $1 - \vec{n}_\tau \cdot \vec{n}_0 \geq 1$. In this setting, by taking into account that $\sqrt{x} \leq x$ for every $x \geq 1$, we find that $\sqrt{1 - \vec{n}_\tau \cdot \vec{n}_0} \leq 1 - \vec{n}_\tau \cdot \vec{n}_0$. Hence, for $\vec{n}_\tau \cdot \vec{n}_0 \leq 0$, Eq. (S37) can be written as follows

$$d_{\text{euc}}(\vec{n}_0, \vec{n}_\tau) \leq \sqrt{2}(1 - \vec{n}_\tau \cdot \vec{n}_0). \quad (\text{S38})$$

Finally, by combining Eqs. (S35), (S36), and (S38), we obtain the result

$$\tau_{QSL}^\alpha \leq 2^{\frac{1}{2} + \frac{1}{\alpha}} \tau_{QSL}^{\alpha, \beta}, \quad (\text{S39})$$

which holds whenever $\vec{n}_\tau \cdot \vec{n}_0 \leq 0$. This proves Lemma 3.