Two Results in the Quantum Theory of Measurements^{*}

Simone Del Vecchio, Jürg Fröhlich, Alessandro Pizzo, Alessio Ranallo

December 4, 2023

Dedicated to the memory of our colleague, teacher and friend *Gianni Morchio*

Abstract

Two theorems with applications to the quantum theory of measurements are stated and proven. The first one clarifies and amends von Neumann's Measurement Postulate used in the Copenhagen interpretation of quantum mechanics. The second one clarifies the relationship between "events" and "measurements" and the meaning of measurements in the ETH-Approach to quantum mechanics.

Contents

1	Introduction and summary of contents	1
	Von Neumann's Measurement Postulate 2.1 An amended form of von Neumann's Postulate	2 3
	The Description of Measurements in the ETH -Approach to QM	6
4	Proof of the Main Result	10

1 Introduction and summary of contents

In this paper, we present two mathematical results of relevance to the quantum theory of measurements,¹ which we treat in a spirit close to the Copenhagen interpretation/heuristics of quantum mechanics (QM), as amended in [1, 2, 3].

Let \mathfrak{E} be a large ensemble of physical systems identical (isomorphic) to a specific system, S, of finitely many degrees of freedom to be described quantum-mechanically. We are interested in understanding the effect of measurements of a physical quantity, \hat{X} , characteristic of S for all systems in \mathfrak{E} . In text-book QM, one tends to invoke *von Neumann's measurement postulate* (see [4])

^{*}to appear in "Trails in Modern Theoretical Physics. A Volume in Tribute of Giovanni Morchio," Andrea Cintio and Alessandro Michelangeli (eds.), Springer-Verlag

¹As far as we remember, Gianni Morchio had an interest in the foundations of quantum mechanics; so he would probably have appreciated our results.

to predict properties of the resulting state, averaged over all systems in \mathfrak{E} , right after a successful completion of the measurements of \hat{X} . The standard formulation of this postulate appears to be afflicted with some problems, which we will discuss and attempt to clarify in the following.

We begin this paper by describing the systems $\simeq S$ we have in mind. A physical quantity, \hat{X} , characteristic of S is represented by a self-adjoint operator, $X = X^*$, acting on a separable Hilbert space, \mathcal{H} . An average over \mathfrak{E} of states of these systems is called an "ensemble state" and is given by a density matrix, i.e., by a positive, trace-class operator, Ω , on \mathcal{H} of trace tr $\Omega = 1$. To mention an example, a system $S \in \mathfrak{E}$ might consist of a particle, such as an electron, propagating in physical space \mathbb{E}^3 , \hat{X} might be a component of the spin or a bounded function of a component of the position- or the momentum of the particle, and

$$\mathcal{H} = L^2(\mathbb{R}^3, d^3x) \otimes \mathbb{C}^{2s+1},$$

where $x \in \mathbb{R}^3$ is the position and s the spin of the particle.

The purpose of this paper is to clarify what is meant by the statement that a measurement of the quantity \hat{X} has been completed successfully. Since we will try to follow the spirit of the Copenhagen Interpretation/heuristics of QM, where appropriate, we will usually adopt an *ensemble point of view*, emphasizing statements that are obtained by taking averages over all systems in the ensemble \mathfrak{E} . But when combined with results in [2, 3], our results have implications relevant for the theory of measurements carried out on individual systems.

Next, we outline the contents of this paper. In Sect. 2, we recall von Neumann's measurement postulate and point out some problems with it. We then formulate a revised version of this postulate and state the main result proven in this paper. In Sect. 3 we sketch how measurements are described in the ETH- Approach to quantum mechanics [1, 2, 3]. In Sect. 4, we present the proof of our main result.

2 Von Neumann's Measurement Postulate

We imagine that the initial ensemble state when measurements of \hat{X} set in, for all systems in \mathfrak{E} , is described by a density matrix Ω_{in} . In his book [4] on the foundations of QM, von Neumann postulated that, when averaging over \mathfrak{E} , the effect of measuring \hat{X} for all systems belonging to \mathfrak{E} amounts to replacing the state Ω_{in} by a certain ensemble state, Ω_{out} , describing the average of states of systems belonging to \mathfrak{E} right *after* the measurements of \hat{X} have been completed, where Ω_{out} satisfies the following postulate.

Von Neumann's Postulate:

Let $X = X^*$ be the self-adjoint operator on \mathcal{H} representing the physical quantity \hat{X} , and let

$$X = \int_{\mathbb{R}} \xi \, d\Pi(\xi) \tag{1}$$

be the spectral decomposition of X, with $\Pi(\Delta)$ its spectral projection associated with an arbitrary Borel set $\Delta \subset \mathbb{R}$. The ensemble state Ω_{out} right after completion of the measurements of \hat{X} has the properties that

$$[\Omega_{out}, X] = 0, \quad \text{and} \\ \operatorname{tr}(\Omega_{in} \cdot \Pi(\Delta)) = \operatorname{tr}(\Omega_{out} \cdot \Pi(\Delta)), \; \forall \; \text{Borel sets} \; \Delta \subset \mathbb{R} \; (Born's \; Rule) \quad \Box$$

$$(2)$$

<u>*Remark:*</u> We will see shortly that this formulation of von Neumann's postulate is inadequate, except if the operator X has pure point spectrum (for which case it was originally formulated) – but even then it is problematic, as will become apparent in Sect. 3.

The spectral decomposition of a density matrix Ω has the form

$$\Omega = \sum_{n=1}^{N} \omega_n \pi_n, \qquad 1 \ge \omega_1 > \omega_2 > \dots > \omega_N > 0,$$

$$\pi_n = \pi_n^*, \quad \pi_n \cdot \pi_m = \delta_{nm} \pi_n, \quad \forall \ n, m = 1, 2, \dots, N,$$
(3)

for some $N \leq \infty$. The operators π_n are disjoint orthogonal projections of finite rank (the eigenprojections of Ω), and

$$\operatorname{tr}(\Omega) = \sum_{n=1}^{N} p_n = 1$$
, where $p_n = \omega_n \cdot \dim \pi_n$, $n = 1, 2, \dots, N$

We set

$$\pi_{\infty} := \mathbf{1} - \sum_{n=1}^N \pi_n \,.$$

If, as in the formulation of von Neumann's postulate given in Eq. (2), the operator X is assumed to commute with Ω_{out} , then it satisfies the identity

$$X = \sum_{n=1}^{N} \pi_n X \pi_n + \pi_\infty X \pi_\infty, \quad \text{where } \pi_n X \pi_n \text{ is of finite rank, } \forall n = 1, 2, \dots, N.$$
(4)

We observe that, for every n = 1, 2, ..., N, $\pi_n X \pi_n$ is a selfadjoint, finite-dimensional matrix; hence its spectrum consists of finitely many (discrete) eigenvalues. Let \mathcal{H}^+ be the subspace of \mathcal{H} given by the range of $1 - \pi_\infty$. It follows that if X satisfies (2) then the operator $X|_{\mathcal{H}^+}$ has *pure-point spectrum*. (Of course, if the range of π_∞ is infinite-dimensional then $\pi_\infty X \pi_\infty$ may have continuous spectrum; but this is irrelevant for measurements of \hat{X} that result in states occupied by the systems in \mathfrak{E} whose average is given by Ω_{out} .) Thus, at best, von Neumann's postulate in the formulation of Eq. (2) can only be applied to measurements of physical quantities with pure-point spectrum. However, a component of the position or of the momentum of a quantum particle propagating in physical space \mathbb{E}^3 has simple continuous spectrum occupying the entire real line \mathbb{R} .

We conclude that Eqs. (2) *cannot* be valid verbatim when physical quantities represented by operators with *continuous spectrum* are measured, and we should find out how to modify them in such instances.

2.1 An amended form of von Neumann's Postulate

We imagine that measurements of a physical quantity \hat{X} are carried out for all systems belonging to a large ensemble \mathfrak{E} of systems identical to a system S, with the result that the average over \mathfrak{E} of the final states of these systems after completion of the measurements of \hat{X} is found to be *close* (but not necessarily equal) to an essemble state given by a density matrix Ω_{out} with the property that

$$\left\| \left[\Omega_{out}, X \right] \right\| < \varepsilon \,, \tag{5}$$

for some ε smaller than the error margin of the instrument used to measure \hat{X} . One may add the assumption that, for Ω_{out} , Born's Rule holds, as formulated in the second equation of (2). We will

establish the following

<u>Main Result</u>: If condition (5) holds for a sufficiently small $\varepsilon \ll 1$ then one may replace Ω_{out} by a modified density matrix Ω'_{out} and X by a modified operator X',

$$X' = \sum_{k=1}^{K} \xi_k \Pi_k, \quad \text{for some } K \le \infty, \qquad (6)$$

where $\xi_1 > \xi_2 > \cdots > \xi_K > -\infty$ are the eigenvalues of X' and Π_1, \ldots, Π_K the corresponding eigen-projections, with the properties that

- (i) the operator X' has pure-point spectrum and is close to the operator X representing \hat{X} in the operator norm;
- (ii) the density matrix Ω'_{out} is close to the density matrix Ω_{out} in the trace norm; and
- (iii) the operators X' and Ω'_{out} commute, i.e.,

$$\left[\Omega_{out}^{\prime}, X^{\prime}\right] = 0.$$
⁽⁷⁾

The closeness of X' to X and of Ω'_{out} to Ω_{out} depends on the size of the commutator of X with Ω_{out} : the smaller the norm, $\|[\Omega_{out}, X]\|$, of this commutator the closer are X' to X and Ω'_{out} to Ω_{out} . The size of $\|[\Omega_{out}, X]\|$ is thus a measure for the precision of the instrument used to measure \hat{X} – the smaller this norm, the higher the precision of the instrument.

The proof of the *Main Result* stated above is given in Sect. 4. At the end of the present section, we sketch the very easy proof in the special case where $\dim(\mathcal{H}) < \infty$.

<u>Remarks</u>:

(1) Another possible amendment of von Neumann's postulate can be formulated as follows. We cover the spectrum, $\operatorname{spec}(X)$, of the operator X with small closed intervals $\Delta_k \subset \mathbb{R}, k = 1, 2, \ldots, K$, for some $K < \infty$, with the properties that $\Delta_k \cap \Delta_{k'}$ is empty or consists of a single point (assumed not to be an eigenvalue of X) whenever $k \neq k'$, and $\bigcup_{k=1}^{K} \Delta_k \supseteq \operatorname{spec}(X)$. These intervals are assumed to be determined by properties of the instrument used to measure \hat{X} . One may then assume that the \mathfrak{E} -average of the states of the systems after completion of the measurements of \hat{X} is given by a density matrix Ω_{out} satisfying

$$\Omega_{out} = \sum_{k=1}^{K} \Pi_k \Omega_{out} \Pi_k, \quad \text{where} \quad \Pi_k = \Pi(\Delta_k), \ \forall \ k = 1, 2, \dots, K.$$
(8)

The operator X' is chosen to be given by

$$X' = \sum_{k=1}^{K} \xi_k \, \Pi_k,$$

where ξ_k is the midpoint of the interval $\Delta_k \subset \mathbb{R}$, for all k. Assuming that the length of all the intervals Δ_k is bounded above by 2ε , we conclude that

$$\left\| \left[\Omega_{out}, X \right] \right\| < \varepsilon, \quad \left[\Omega_{out}, X' \right] = 0, \quad \text{and} \quad \|X - X'\| < \varepsilon.$$

$$(9)$$

This amendment of von Neumann's postulate is somewhat arbitrary and involves assumptions on what is meant by a measurement of a physical quantity that are more detailed than condition (5).

- (2) The *Main Result* stated above is reminiscent of a theorem that says that if two bounded selfadjoint operators almost commute then there are two operators close in norm to the original ones that *do* commute; see [5, 6, 7].
- (3) We conjecture that our *Main Result* is a special case of the following more general statement: Let \mathfrak{A} be a von Neumann algebra with unit $\mathbf{1}$, and let ω be a normal state on \mathfrak{A} . For an operator $X \in \mathfrak{A}$, we define a bounded linear functional on \mathfrak{A} by

$$\operatorname{ad}_{X}[\omega](Y) := \omega([Y, X]), \quad \forall Y \in \mathfrak{A}.$$
 (10)

Suppose now that ω and X are such that

$$\left|\operatorname{ad}_{X}[\omega](Y)\right| < \varepsilon \|Y\|, \quad \forall Y \in \mathfrak{A}, \text{ for some } \varepsilon \ll 1.$$
 (11)

Then there exist a normal state ω' on \mathfrak{A} and an operator $X' \in \mathfrak{A}$, with $\|\omega' - \omega\| < \delta(\varepsilon)$ and $\|X' - X\| < \delta(\varepsilon)$, for some $\delta(\varepsilon) \searrow 0$, as $\varepsilon \searrow 0$, such that

$$\operatorname{ad}_{X'}[\omega'] = 0. \tag{12}$$

Our *Main Result* shows that this conjecture holds in the special case where \mathfrak{A} is isomorphic to the algebra of all bounded operators on a separable Hilbert space.

As a warm-up we prove the *Main Result* in the special case of a finite-dimensional Hilbert space \mathcal{H} , which is very easy. In items (i) through (iii), one may then set $\Omega'_{out} = \Omega_{out}$ and only slightly modify the operator X, or one may set X' = X and only slightly modify Ω_{out} , and end up with (7).

Let $\mathcal{H} = \mathbb{C}^M$, with $M < \infty$. Then

$$X = \sum_{k=1}^{K} \xi_k \Pi_k, \qquad K \le M \,, \quad \text{and}$$

$$\Omega = \sum_{n=1}^{N} \omega_n \pi_n, \qquad N \le M \,,$$
(13)

where $\xi_1 > \xi_2 > \cdots > \xi_K > -\infty$ are the eigenvalues of X and $\Pi_1, \Pi_2, \ldots, \Pi_K$ are the corresponding eigen-projections, and $\omega_1 > \omega_2 > \cdots > \omega_n > 0$ are the non-zero eigenvalues of Ω , with $\pi_1, \pi_2, \ldots, \pi_N$ the corresponding eigen-projections. We define $\pi_{N+1} := \mathbf{1} - \sum_{n=1}^N \pi_n$, and

$$\gamma_{\Omega} := \min_{1 \le n \le N} \left(\omega_n - \omega_{n+1} \right) > 0, \quad \text{with } \omega_{N+1} := 0, \quad (14)$$

to be the smallest gap between distinct eigenvalues of Ω . Let us assume that

$$\|[X,\Omega]\| \leq \varepsilon, \quad \text{for some } \varepsilon \ll \gamma_{\Omega}.$$
 (15)

We define an operator X' by setting

$$X' := \sum_{n=1}^{N+1} \pi_n X \pi_n .$$
 (16)

Obviously X' commutes with Ω , and we claim that

$$\|X' - X\| < \text{const.}\,\varepsilon\,. \tag{17}$$

Proof of (17). Clearly

$$X = \sum_{n,n'=1,2,\dots,N+1} \pi_n X \pi_{n'}.$$
 (18)

By (15) we have that

$$[\pi_n X \pi_{n'}, \Omega] \| = \|\pi_n [X, \Omega] \pi_{n'} \| \leq \varepsilon, \quad \forall \ n, n'.$$
⁽¹⁹⁾

Plainly $[\pi_n X \pi_n, \Omega] = 0$, $\forall n = 1, 2, \dots, N + 1$. If $n \neq n'$ then

$$[\pi_n X \pi_{n'}, \Omega] = (\omega_{n'} - \omega_n) \pi_n X \pi_{n'}.$$

By Eqs. (14) and (19), we have that

$$\|\pi_n X \pi_{n'}\| \leq \gamma_{\Omega}^{-1} \varepsilon$$
, for $n \neq n'$.

Thus, using (18) we find that

$$\|X - X'\| \le (N+1)N \,\gamma_{\Omega}^{-1} \,\varepsilon < M^2 \gamma_{\Omega}^{-1} \varepsilon \,, \tag{20}$$

as claimed in (17).

In the calculations just shown we can obviously exchange the roles of X and Ω . We set

$$\gamma_X := \min_{1 \le k < K} (\xi_k - \xi_{k+1}) > 0,$$

and we then replace the density matrix Ω by

$$\Omega' := \sum_{k=1}^K \Pi_k \,\Omega \,\Pi_k \,.$$

Clearly Ω' is a non-negative operator, and $tr(\Omega') = 1$, because $\sum_{k=1}^{K} \Pi_k = 1$; i.e., Ω' is a density matrix; and it obviously commutes with X. Repeating the arguments shown above, we find that

$$\operatorname{tr}(|\Omega - \Omega'|) \leq M \left(K - 1\right) K \gamma_X^{-1} \varepsilon < M^3 \gamma_X^{-1} \varepsilon.$$
(21)

Of course, the problem with the estimates in (20) and (21) is the dependence of the right sides on the dimension, M, of the Hilbert space \mathcal{H} . This problem is addressed in Sect. 4, where we state a result that is *uniform* in the dimension of the Hilbert space, but at the price that we have to slightly modify both, X and Ω . This result enables one to modify von Neumann's measurement postulate so as to avoid the shortcomings of the original version, as indicated above.

3 The Description of Measurements in the *ETH*-Approach to QM

In this section we sketch how measurements can be described in the formulation of QM proposed in [1, 2, 3] under the name of "*ETH*-Approach to QM" (assuming some familiarity with these papers).

We begin with the obvious observation that a successful measurement of a physical quantity \hat{X} characteristic of a system S (belonging to an experiment \mathfrak{E}) results in an *event*, namely the event that \hat{X} takes a – possibly somewhat imprecise – value belonging to some small interval contained in the real line whose length depends on the accuracy of the instrument used to measure \hat{X} . To

understand the significance of this statement it is necessary to clarify what, in the *ETH*-Approach to QM, is meant by an "event". We recall the definition proposed in [2, 3]. Abstractly, a "potential event", \mathfrak{e} , associated with a physical system $S \in \mathfrak{E}$ is a partition of unity, $\mathfrak{e} = \{\pi_n\}_{n=1}^{\infty}$, by orthogonal projections satisfying

$$\pi_n = \pi_n^*, \quad \pi_n \cdot \pi_{n'} = \delta_{nn'} \pi_n, \ \forall \ n, n' = 1, 2, \dots, \quad \sum_{n=1}^{\infty} \pi_n = \mathbf{1}.$$
(22)

An operator X representing a physical quantity \hat{X} characteristic of a system $S \in \mathfrak{E}$ at some time $\geq t$ and the projections $\pi \in \mathfrak{e}$ of an arbitrary potential event \mathfrak{e} that may occur in S at a time $\geq t$ are supposed to belong to some algebra $\mathfrak{A} = \mathcal{E}_{\geq t}$, which, in general, depends *non-trivially* on time t. For systems, S, with finitely many degrees of freedom, \mathfrak{A} is the algebra, $B(\mathcal{H})$, of all bounded operators on a separable Hilbert space \mathcal{H} and is independent of t. But, for systems with infinitely many degrees of freedom, including those describing the quantized electromagnetic field,² the time-dependence of $\mathfrak{A} = \mathcal{E}_{\geq t}$ tends to be *non-trivial*, and \mathfrak{A} is a more exotic (type-III₁) algebra. Our analysis in this section does not require any specific assumptions on \mathfrak{A} . (It is only assumed that the algebra \mathfrak{A} is weakly closed, i.e., that it is a von Neumann algebra; but it need not and usually will *not* be isomorphic to $B(\mathcal{H})$.) States at time t are states on $\mathfrak{A} = \mathcal{E}_{\geq t}$ (i.e., positive, normalized linear functionals on $\mathcal{E}_{\geq t}$). They are denoted by lower-case Greek letters, ω, \ldots .

In the following discussion we fix a time t and suppress explicit reference to time-dependence wherever possible. We suppose that a state, ω , on \mathfrak{A} is an *ensemble state*, i.e., that it has the meaning of being an average over the ensemble \mathfrak{E} of states of individual systems, all $\simeq S$. If a potential event $\mathfrak{e} = \{\pi_n\}_{n=1}^{\infty} \subset \mathfrak{A}$ is *actualizing* (i.e., is observed to happen) at some time t then, according to the *ETH*- Approach to QM, the state $\omega = \omega_t$ has the property that

$$\omega(X) = \sum_{\pi \in \mathfrak{e}} \omega(\pi \cdot X \cdot \pi), \qquad \forall \ X \in \mathfrak{A},$$
(23)

i.e., ω is a convex combination of states in the images of the projections $\pi \in \mathfrak{e}$. Potential events actualizing at some time are called "actualities". (For a more precise characterization of actualities, see, e.g., [3].) If $\mathfrak{A} = B(\mathcal{H})$ then

$$\omega(X) = \operatorname{tr}(\Omega \cdot X), \quad \forall \ X \in \mathfrak{A},$$

for some density matrix Ω on \mathcal{H} , and the projections π belonging to the event \mathfrak{e} that actualizes, given the state ω , are the spectral projections of the density matrix Ω .

If \mathfrak{e} is an event actualizing at some time t then the state at time t of an *individual* system in the ensemble \mathfrak{E} is expected to belong to the image of a projection $\pi \in \mathfrak{e}$, with a probability, $prob_{\omega}(\pi)$, given by *Born's Rule*, namely

$$prob_{\omega}(\pi) = \omega(\pi),$$

where ω is the ensemble state at time t.

We are interested in characterizing actualities $\mathfrak{e} = \{\pi_n\}_{n=1}^N \subset \mathfrak{A}, N \leq \infty$, that can be interpreted as corresponding to the completion of the measurement of a certain physical quantity \hat{X} . We thus consider a state ω satisfying Eq. (23). Given a non-negative number $\varepsilon \ll 1$, there exists an integer $N_0 < \infty$ such that

$$\sum_{n=1}^{N_0-1} \omega(\pi_n) > 1 - \varepsilon, \ i.e., \quad \omega(\pi^{(N_0)}) < \varepsilon, \quad \text{where} \quad \pi^{(N_0)} := \sum_{n=N_0}^N \pi_n.$$
(24)

²the only systems for which (in our opinion) the "measurement problem" has a satisfactory solution

It is then very unlikely that an individual system in \mathfrak{E} is found to occupy a state in the range of a projection $\pi \leq \pi^{(N_0)}$. If \mathfrak{e} is the potential event actualizing at a certain time t and ω is the ensemble state at time t satisfying (23) then the slightly coarser event $\mathfrak{e}_0 := \{\pi_1, \pi_2, \ldots, \pi_{N_0-1}, \pi^{(N_0)}\}$ can be viewed to be an actuality at time t, too. To avoid irrelevant complications, we henceforth replace \mathfrak{e} by \mathfrak{e}_0 throughout the following discussion, and we simplify our notations by writing \mathfrak{e} , instead of \mathfrak{e}_0 , and π_{N_0} , instead of $\pi^{(N_0)}$, with $N_0 < \infty$.

We assume that the operator X representing the physical quantity \hat{X} has the form

$$X = \sum_{k=1}^{K} \xi_k \Pi_k, \quad \text{for some} \quad K < \infty,$$
(25)

where the real numbers ξ_k are the eigenvalues of X and the operators Π_k are the corresponding eigen-projections, k = 1, 2, ..., K. (We should mention that the projections Π_k may be given by $\Pi_k = \Pi(\Delta_k)$, where the sets Δ_k are intervals of the real line of length $< 2\varepsilon$ whose union covers spec(X), and ξ_k may be (e.g.) the midpoint of the interval Δ_k , for all k, as discussed in Remark (1) of Subsect. 2.1.)

If the actuality \mathfrak{e} can be interpreted to correspond to the likely completion of a measurement of \hat{X} , with an accuracy measured by ε , then there must exist a decomposition of $\{1, 2, \ldots, N_0\}$ into disjoint subsets \mathcal{I}_k , $k = 1, 2, \ldots, K$, such that

$$\|[\pi_n, \Pi_k]\| < \mathcal{O}(N_0^{-2}\varepsilon), \quad \forall \ n \le N_0, \quad \text{and}$$
$$\sum_{n \notin \mathcal{I}_k, n < N_0} \|\pi_n \Pi_k \pi_n\| < \mathcal{O}(\varepsilon), \quad \forall \ k = 1, 2, \dots K.$$
(26)

The second equation tells us that if a system is found in a state in the range of a projection $\pi_n, n \notin \mathcal{I}_k, n < N_0$, then the quantity \hat{X} is very unlikely to have the measured value ξ_k . By (24), if the ensemble state is given by ω then it is very unlikely that an individual system in \mathfrak{E} is found in a state belonging to the range of the projection π_{N_0} .

Since $\sum_{n=1}^{N_0} \pi_n = 1$, one obviously has that

$$X = \sum_{n,n'=1,2,...,N_0} \pi_n X \, \pi_{n'}$$

Since $\pi_n \cdot \pi_{n'} = 0$, for $n \neq n'$, the first inequality in (26) then implies that the operator X is approximated in norm by

$$X' := \sum_{n=1}^{N_0} \pi_n X \pi_n , \qquad (27)$$

up to an error of $\mathcal{O}(\varepsilon)$; and (24) tells us that the Born probability of picking up a correction in determining the outcome of the measurement of \hat{X} that is due to the operator $\pi_{N_0} X \pi_{N_0}$ is bounded by $\mathcal{O}(\varepsilon)$, hence very small. One may then wonder whether the actuality \mathfrak{e} could occur as the result of a measurement of a *slightly different* physical quantity $\simeq \hat{X}$.

The second inequality in (26) implies that X' is well approximated by the operator

$$X'' := \sum_{k=1}^{K} \sum_{n \in \mathcal{I}_k} \xi_k \, \pi_n \, \Pi_k \, \pi_n + \pi_{N_0} \, X \, \pi_{N_0}$$
(28)

with

$$|X'' - X'|| < \mathcal{O}(K\varepsilon).$$
⁽²⁹⁾

Next, we note that the first inequality in (26) implies that

$$\left|\left|\left(\pi_n \Pi_k \pi_n\right)^2 - \pi_n \Pi_k \pi_n\right|\right| < \mathcal{O}\left(N_0^{-2}\varepsilon\right).$$

This estimate enables us to apply the following

<u>Lemma</u>. Let P be a self-adjoint operator in a von Neumann algebra \mathfrak{A} , and let $\delta < \frac{1}{2}$. If $||P^2 - P|| < \delta$ then there exists an orthogonal projection $\hat{P} \in \mathfrak{A}$ whose image belongs to the range of P such that

$$\|\widehat{P} - P\| < \delta.$$

See Lemma 8 and Appendix C of [8]. This lemma implies that if $N_0^{-2}\varepsilon$ is small enough then there exists an orthogonal projection $\pi_{k,n}$ with the property that the image of $\pi_{k,n}$ is contained in or equal to the image of π_n and such that

$$\|\pi_{k,n} - \pi_n \Pi_k \pi_n\| < \mathcal{O}\left(N_0^{-2}\varepsilon\right).$$

We define

$$X''' := \sum_{k=1}^{K} \xi_k \left(\sum_{n \in \mathcal{I}_k} \pi_{k,n} \right) + \pi_{N_0} X \pi_{N_0}, \quad \text{and} \quad X_{fin} := X''' - \pi_{N_0} X \pi_{N_0}.$$
(30)

We are ready to state a result in the theory of measurements, according to the ETH-Approach to QM.

Theorem 3.1 We assume that the bounds in (24) and (26) hold for some $\varepsilon \ll 1$. Then we have that

- (i) the Born probability of finding an individual system in the ensemble \mathfrak{E} in a state that belongs to the range of the projection $\pi_{N_0} = \pi^{(N_0)}$ is bounded above by ε ;
- (ii) the operator X''' defined in (30) is reduced by the projections π_{N_0} and $1 \pi_{N_0}$;
- (iii) the norm of X''' X is bounded by

$$\|X''' - X\| < \mathcal{O}(\varepsilon),$$

i.e., the physical quantity \hat{X} is well approximated by a slightly modified physical quantity represented by the operator X''';

(iv) the eigenvalues of $X_{fin} = X''' - \pi_{N_0} X \pi_{N_0}$ are contained in or equal to the spectrum, $\{\xi_k\}_{k=1}^K$, of the operator X and the eigen-projection of X_{fin} corresponding to ξ_k is given by the projection $\sum_{n \in \mathcal{I}_k} \pi_{k,n}$ (which is dominated by the projection $\sum_{n \in \mathcal{I}_k} \pi_n$), for k = 1, 2, ..., K; and

$$\left[X_{fin}, \pi_n\right] = 0, \quad \forall \ \pi_n \in \mathfrak{e} \,. \qquad \Box$$

We conclude that, under the hypotheses of Theorem 3.1, one may interpret the actualization of the event \mathfrak{e} as being accompanied by the completion of a measurement of a physical quantity $\hat{X}''' \approx \hat{X}$, where \hat{X}''' is represented by an operator X''' that is a tiny modification of the operator X representing \hat{X} .

In this section, we have not tried to optimize our results; we have attempted to outline the basic ideas of how measurements can be interpreted in the ETH-Approach described in [1, 2, 3].

4 Proof of the Main Result

In this section we prove the *Main Result* announced in Sect. 2. We consider a density matrix Ω on a separable Hilbert space \mathcal{H} with spectral decomposition

$$\Omega = \sum_{n=1}^{\infty} \omega_n \, \pi_n \,, \qquad \omega_1 > \omega_2 > \cdots \,. \tag{31}$$

as in Eq. (3) of Sect. 2. We define $p_n := \omega_n \cdot \dim \pi_n$, $n = 1, 2, \ldots$ Given a positive number $\varepsilon \ll 1$, we define Δ_{ε} by

$$\Delta_{\varepsilon} := \sum_{n : \omega_n \leqslant \varepsilon^{1/4}} p_n \,. \tag{32}$$

Clearly, $\Delta_{\varepsilon} \searrow 0$, as $\varepsilon \searrow 0$. The *Main Result* is a consequence of the following theorem.

Theorem 4.1 Let Ω and Δ_{ε} be as in (31) and (32), respectively, and let X be a self-adjoint operator on \mathcal{H} , with $||X|| \leq 1$. We assume that

$$\left\| \left[\Omega, X\right] \right\| \leqslant \varepsilon \,. \tag{33}$$

Then, for sufficiently small values of ε and Δ_{ε} , there exist a density matrix Ω' and a self-adjoint operator X' such that

$$||X - X'|| \leq \varepsilon^{1/4}, \quad and \quad tr|\Omega - \Omega'| \leq 2\Delta_{\varepsilon} + \mathcal{O}(\varepsilon^{1/4}).$$
 (34)

Proof.

As announced in the theorem, our goal is to construct a density matrix Ω' close to Ω in the trace norm and a self-adjoint operator X' close to X in the operator norm such that $[\Omega', X'] = 0$. We begin with the construction of Ω' .

In the following it is convenient to rewrite the spectral decomposition of Ω as follows:

$$\Omega = \sum_{j=1}^{\infty} \omega_j |u_j\rangle \langle u_j|, \qquad \omega_1 \ge \omega_2 \ge \cdots \ge 0, \qquad \sum_{j=1}^{\infty} \omega_j = 1, \qquad (35)$$

where $\{u_j\}_{j=1}^{\infty}$ is an orthonormal system of eigenvectors of Ω , and $|u_j\rangle\langle u_j|$ is the orthogonal projection onto u_j , for all j. Then assumption (33) implies that

$$\|[\Omega, X]u_i\|^2 = \sum_{j=1}^{\infty} (\omega_i - \omega_j)^2 |\langle u_i, Xu_j \rangle|^2 \leqslant \varepsilon^2, \quad \forall i.$$
(36)

In the following steps, we construct a positive trace-class operator $\widetilde{\Omega} \leq \Omega$, (hence tr $\widetilde{\Omega} \leq 1$).

1) We preserve the eigenvectors of the density operator Ω , but - where necessary - modify the corresponding eigenvalues in such a way that the spectrum of the modified operator $\tilde{\Omega}$ consists of (possibly degenerate) eigenvalues separated by gaps of specified size. To begin with we choose two exponents, δ and β (later set equal to 1/4 and 3/4, respectively), with

$$0 < \delta < \beta < 1 \quad \text{and} \quad \beta > 2\delta \,, \tag{37}$$

and we modify the spectrum of $\widetilde{\Omega}$ in such a way that the gaps between the non-coinciding modified eigenvalues, i.e., between the distinct eigenvalues of $\widetilde{\Omega}$, will be larger than ε^{β} .

1-i) We observe that, since $\Omega \ge 0$, with tr $\Omega = 1$, the dimension of the direct sum of the eigenspaces of Ω corresponding to eigenvalues larger than or equal to ε^{δ} is bounded above by $O(1/\varepsilon^{\delta})$.

1-ii) Next, we define ω_{i_1} to be the smallest eigenvalue of Ω of order ε^{δ} with the property that its separation from the previous (next larger) eigenvalue is bounded below by ε^{β} . It is not assumed that an eigenvalue with the properties of ω_{i_1} exists.

But if such an eigenvalue ω_{i_1} exists then we denote by $(\omega_{i_1})_-$ its precursor. By construction, we have that $(\omega_{i_1})_- \leq \mathcal{O}(\varepsilon^{\delta} + \varepsilon^{\beta-\delta})$, because there are at most $\mathcal{O}(\varepsilon^{-\delta})$ eigenvalues separated by gaps bounded by $\leq \varepsilon^{\beta}$ in between ω_{i_1} and $(\omega_{i_1})_-$, as follows from 1-i).

We define

- an interval I_0 by $I_0 := [0, (\omega_{i_1})_-],$
- and a subspace $\mathcal{H}_0 \subset \mathcal{H}$ as the direct sum of the eigenspaces of Ω corresponding to eigenvalues contained in the interval I_0 .

If an eigenvalue with the properties of ω_{i_1} does not exists then we conclude that the largest eigenvalue, ω_{max} , of Ω must be smaller than $\mathcal{O}(\varepsilon^{\delta} + \varepsilon^{\beta - \delta})$. In this case, we define $I_0 := [0, \omega_{max}]$.

We define $\widetilde{\Omega}$ to vanish on the subspace \mathcal{H}_0 .

1-iii) We next assume that $I_0 \neq [0, \omega_{max}]$, i.e., that an eigenvalue with the properties of ω_{i_1} exists. Then we consider the smallest eigenvalue of Ω larger than ω_{i_1} with the property that its separation from the previous eigenvalue is larger than ε^{β} .

If such an eigenvalue exists we denote it by ω_{i_2} and its precursor by $(\omega_{i_2})_-$, and we then have that $(\omega_{i_2})_- \leq \mathcal{O}(\omega_{i_1} + \varepsilon^{\beta - \delta})$. We also define

- $I_1 := [\omega_{i_1}, (\omega_{i_2})_-];$
- $n_1 :=$ number of eigenvalues (with multiplicity) of Ω contained in I_1 ;
- $\mathcal{H}_1 :=$ direct sum of the corresponding eigenspaces (notice that dim $\mathcal{H}_1 = n_1$).

If an eigenvalue with the properties of ω_{i_2} does not exists we conclude that the largest eigenvalue, ω_{max} , of Ω is smaller than $\mathcal{O}(\omega_{i_1} + \varepsilon^{\beta - \delta})$, and we define $I_1 := [\omega_{i_1}, \omega_{max}]$

On the subspace \mathcal{H}_1 we define

$$\widetilde{\Omega}\big|_{\mathcal{H}_1} := \widetilde{\omega}_1 \cdot \mathbf{1}\big|_{\mathcal{H}_1},$$

where $\widetilde{\omega}_1 := \omega_{i_1}$.

1-iv) We iterate these arguments: If $I_{m-1} \neq [\omega_{i_{m-1}}, \omega_{max}]$, then, starting from ω_{i_m} , we consider the eigenvalue of Ω with the property that its separation from the previous one is bounded below by ε^{β} .

If an eigenvalue of Ω with these properties exists we denote it by $\omega_{i_{m+1}}$ and the previous one by $(\omega_{i_{m+1}})_{-} (\leq \mathcal{O}(\omega_{i_m} + \varepsilon^{\beta - \delta}))$. We also define

- $I_m := [\omega_{i_m}, (\omega_{i_{m+1}})_-];$
- $\mathcal{H}_m :=$ direct sum of eigenspaces of Ω corresponding to eigenvalues contained in the interval I_m ; and $n_m := \dim \mathcal{H}_m$.

If this eigenvalue does not exists we conclude that the largest eigenvalue, ω_{max} , of Ω is bounded above by $\mathcal{O}(\omega_{i_m} + \varepsilon^{\beta - \delta})$, and we define $I_m := [\omega_{i_m}, \omega_{max}]$

On the subspace \mathcal{H}_m we define the operator $\widetilde{\Omega}$ by $\widetilde{\Omega}|_{\mathcal{H}_m} := \widetilde{\omega}_m \cdot \mathbf{1}|_{\mathcal{H}_m}$, where $\widetilde{\omega}_m := \omega_{i_m}$. 1-v) The construction described above must necessarily stop at some step $\overline{m} \ge 0$, because Ω is trace-class and $\varepsilon^{\beta} > 0$. The spectrum of the operator $\widetilde{\Omega}$ constructed above consists of the points

$$\{\omega_{i_0} := 0, \, \omega_{i_1}, \, \dots, \, \omega_{i_{\overline{m}}}\}. \tag{38}$$

1-vi) We note that $\tilde{\Omega}$ has been defined as the operator whose eigenspaces are the subspaces \mathcal{H}_m and the corresponding eigenvalues are given by $\tilde{\omega}_m$. (To avoid possible confusion we stress that the eigenvalues $\tilde{\omega}_m$ of $\tilde{\Omega}$ are *increasing* in *m* whereas the eigenvalues ω_i of Ω are *decreasing* in *i*.) The operator $\tilde{\Omega}$ enjoys the property

$$\operatorname{tr}|\Omega - \widetilde{\Omega}| \leq o(1) + \mathcal{O}\left(\varepsilon^{\beta - \delta}(n_1 + \dots + n_{\overline{m}})\right) \leq o(1) + O(\varepsilon^{\beta - 2\delta}), \qquad (39)$$

which holds, because

$$0 < \sum_{\omega_i \leqslant \varepsilon^{\delta}} \omega_i \leqslant o(1); \tag{40}$$

(recall that Ω is trace-class and that, in (32), we have noticed that $\sum_{i:\omega_i \leq \varepsilon^{1/4}} \omega_i =: \Delta_{\varepsilon} \ll 1$). Moreover, we use the facts that any eigenvalue of Ω corresponding to an eigenvector in \mathcal{H}_m

is included in the interval $[\tilde{\omega}_m, \tilde{\omega}_m + \epsilon^{\beta-\delta}]$, by construction of \mathcal{H}_m , and that

$$n_1 + \dots + n_{\overline{m}} \leqslant O(1/\varepsilon^{\circ}),$$
(41)

as shown in 1-i).

2) Next, we modify the operator X. The modified operator is denoted by X' and is defined by its matrix elements in the basis, $\{u_j\}_{j=1}^{\infty}$, of eigenvectors of Ω , which are given by

$$(X')_{i,j} := \langle u_i, X u_j \rangle, \tag{42}$$

provided that u_i and u_j belong to the same subspace \mathcal{H}_p , $p \leq \overline{m}$, and

$$(X')_{i,j} := 0, (43)$$

if u_i and u_j belong to *different* eigenspaces, \mathcal{H}_p , $\mathcal{H}_{p'}$, of Ω .

We thus have by construction that

$$\left[\widetilde{\Omega}, X'\right] = 0. \tag{44}$$

Next, we show that ||X - X'|| = o(1). This follows from

$$\sup_{i} \sum_{j=1}^{\infty} |\langle u_j, (X - X') u_i \rangle|^2 = \varepsilon^{2(1-\beta)}, \qquad (45)$$

where the summands are non-zero only if u_i and u_j belong to different eigenspaces \mathcal{H}_{p_i} , \mathcal{H}_{p_j} of $\tilde{\Omega}$, so that

$$\sum_{j=1}^{\infty} |\langle u_j, (X - X') u_i \rangle|^2 = \sum_{j: u_j \in \mathcal{H}_{p_j}, p_j \neq p_i} |\langle u_j, X u_i \rangle|^2.$$
(46)

But if $p_j \neq p_i$ then $|\omega_i - \omega_j| \ge \varepsilon^{\beta}$, where ω_i and ω_j are the eigenvalues of Ω on the vectors $u_j \in \mathcal{H}_{p_i}$ and $u_i \in \mathcal{H}_{p_i}$, respectively. Next, we exploit the bound assumed in (33), namely

$$\varepsilon^2 \ge \| \left[\,\Omega \,, \, X \,\right] u_i \, \|^2 \tag{47}$$

$$=\sum_{j=1}^{\infty} (\omega_i - \omega_j)^2 |\langle u_i, X u_j \rangle|^2$$
(48)

$$\geq \varepsilon^{2\beta} \sum_{j: u_j \in \mathcal{H}_{p_j}, p_j \neq p_i} |\langle u_i, X u_j \rangle|^2$$
(49)

To conclude the proof of the theorem, we normalize $\tilde{\Omega}$ by dividing by its trace, defining $\Omega' := \frac{\tilde{\Omega}}{\operatorname{tr} \tilde{\Omega}}$. Setting $\delta = \frac{1}{4}$ and $\beta = \frac{3}{4}$, and using that

$$\operatorname{tr}|\Omega - \widetilde{\Omega}| \leq \Delta_{\varepsilon} + \mathcal{O}(\varepsilon^{\frac{1}{4}}),$$

we conclude that

$$\operatorname{tr}|\Omega - \Omega'| \leq 2\Delta_{\varepsilon} + \mathcal{O}(\varepsilon^{\frac{1}{4}}).$$

References

- J. Fröhlich and B. Schubnel, Quantum probability theory and the foundations of quantum mechanics, in: The Message of Quantum Science. Springer, Ph. Blanchard and J. Fröhlich (eds.), Springer-Verlag, Berlin, 2015; and arXiv:1310.1484; see also: Ph. Blanchard, J. Fröhlich and B. Schubnel, A "garden of forking paths" The quantum mechanics of histories of events, Nucl. Phys. B 912, 463-484 (2016)
- [2] J. Fröhlich and A. Pizzo, The Time-Evolution of States in Quantum Mechanics according to the ETH-Approach, Commun. Math. Phys. 389, 1673-1715 (2022)
- [3] J. Fröhlich, Z. Gang and A. Pizzo, A Tentative Completion of Quantum Mechanics, arXiv:2303.11112v1 [math-ph] 20 Mar 2023
- [4] J. von Neumann, Mathematical foundations of quantum mechanics (with an introduction by Iván Abonyi), Akadémiai Kiadó (Publishing House of the Hungarian Academy of Sciences), Budapest, 1980; (translated from the 1964 Russian edition by Ákos Sebestyén); see also:
 G. Lüders, Über die Zustandsänderung durch den Messprozess, Annalen der Physik, 443(5-8) 322-328 (1950)
- [5] H. Lin, Almost commuting self-adjoint matrices and applications, Fields. Inst. Commun. 13, 193 (1995)
- [6] M. B. Hastings, Making Almost Commuting Matrices Commute, Commun. Math. Phys. 291, 321–345 (2009)
- [7] I. Kachkovskiy and Y. Safarov, Distance to Normal Elelemnts in C*-Algebras of Real Rank Zero, arXiv:1403.2021v3 [math.OA] 23 Feb 2015

[8] J. Fröhlich and B. Schubnel, Do we understand quantum mechanics – finally? in: Erwin Schrödinger – 50 Years After, Wolfgang L. Reiter and J. Yngvason (eds.), ESI Lectures in Mathematics and Physics, pages 37-84, European Mathematical Society Publ., 2013

Simone Del Vecchio, Dipartimento di Matematica, Università degli Studi di Bari, Italy, simone.delvecchio@uniba.it

Jürg Fröhlich, ETH Zürich, Institute for Theoretical Physics,

juerg@phys.ethz.ch

Alessandro Pizzo, Dipartimento di Matematica, Università di Roma "Tor Vergata", Italy, pizzo@mat.uniroma2.it

Alessio Ranallo, Dipartimento di Matematica, Università di Roma "Tor Vergata", Italy, ranallo@mat.uniroma2.it