# Motivic coaction and single-valued map of polylogarithms from zeta generators

Hadleigh Frost<sup>a</sup>, Martijn Hidding<sup>b,c</sup>, Deepak Kamlesh<sup>a</sup>, Carlos Rodriguez<sup>b</sup>, Oliver Schlotterer<sup>b</sup> and Bram Verbeek<sup>b</sup>

<sup>a</sup> Mathematical Institute, University of Oxford, OX2 6GG Oxford, United Kingdom

<sup>b</sup> Department of Physics and Astronomy, Uppsala University, Box 516, 75120 Uppsala, Sweden and

<sup>c</sup> Institute for Theoretical Physics, ETH Zurich, 8093 Zürich, Switzerland

We introduce a new Lie-algebraic approach to explicitly construct the motivic coaction and single-valued map of multiple polylogarithms in any number of variables. In both cases, the appearance of multiple zeta values is controlled by conjugating generating series of polylogarithms with Lie-algebra generators associated with odd zeta values. Our reformulation of earlier constructions of coactions and single-valued polylogarithms preserves choices of fibration bases, exposes the correlation between multiple zeta values of different depths and paves the way for generalizations beyond genus zero.

### INTRODUCTION

The advent of multiple polylogarithms (MPLs) [1–5] led to a wealth of new perturbative results and structural insights in quantum field theories [6–10], string theory [11, 12] and numerous other areas of theoretical physics where iterated integrals of rational functions are encountered. Apart from systematizing the integration over marked points on the sphere, MPLs enjoy Hopf-algebra structures including the motivic coaction and the single-valued map which exposed surprising unifying phenomena in the recent amplitudes literature.

The motivic coaction [4, 13–16] mapping MPLs to tensor products of similar iterated integrals turns out to stabilize and connect perturbative quantities in a variety of quantum field theories and string theories [17, 18]. Examples include periods in  $\phi^3$  and  $\phi^4$  theory [19, 20], the anomalous magnetic moment of the electron [21],  $\mathcal{N}=4$  super-Yang-Mills [22], various families of Feynman integrals [23–29], the disk integrals in string tree-level amplitudes [30, 31] and the antipodal duality between amplitudes and form factors [32–34]. In this broad spectrum of cases, the motivic coaction of MPLs in physical quantities hints at novel types of symmetries under the so-called cosmic Galois group [18, 35]. Such Galois symmetries extract an infinite pool of perturbative information from amplitude contributions at low loop orders.

The single-valued map [17, 36, 37] in turn solves the problem of eliminating the monodromies of MPLs by systematically adding complex conjugate MPLs and multiple zeta values (MZVs), while preserving holomorphic differential equations. The resulting single-valued MPLs [38] and their associated single-valued MZVs [17, 36] became key ingredients for studying the multi-Regge regime of  $\mathcal{N}=4$  super-Yang-Mills [39-42], the high-energy limit of more general gauge theories [43, 44] and numerous classes of Feynman integrals [17, 45-47]. Moreover, the single-valued map is a number-theoretic bridge between open- and closed-string interactions in flat spacetime, with a detailed understanding at tree level [30, 48–52] and first loop-level echoes in [53–57]. Finally, the singlevalued map has initiated bootstrap approaches to string amplitudes in AdS backgrounds [58–61].

In this work, we provide a new construction of both the motivic coaction and the single-valued map of MPLs in any number of variables. The main novelty is the use of Lie-algebra structures that efficiently reorganize the appearance of MZVs and lower-complexity MPLs. Apart from the well-known braid operators for marked points on a sphere, our construction makes essential use of Lie-algebra generators associated with odd Riemann zeta values [62, 63]. While state-of-the-art formulations of both the motivic coaction [4, 13, 15, 26] and single-valued MPLs [38, 40, 41] in principle cover cases in any number of variables and of arbitrary transcendental weight, the advantages of our alternative construction are

- (i) The results are in fully simplified form with respect to relations among MPLs and MZVs over Q, i.e. are automatically cast into a fibration basis of MPLs.
- (ii) The appearance of odd Riemann zeta values completely determines the coefficients of MZVs beyond depth one and products of arbitrary MZVs.
- (iii) The Lie-algebraic approach inspires generalizations beyond genus zero where additional generators are associated to all moduli of the surface.

While the first point (i) may be viewed as a computational benefit, the exposure of correlations (ii) between MZVs of different depths is a structural virtue of our approach to coaction formulas and single-valued periods. The last point (iii) already found an explicit genus-one manifestation in section 4.1.2 of [64]<sup>1</sup> for Brown's single-valued iterated Eisenstein integrals [70–72] which indicates that the series of zeta generators in this work are in fact a genus-agnostic building block. In this way, our results should unlock new perspectives on the appearance of MZVs and genus-zero MPLs in coaction formulas for elliptic MPLs [73–76] and ultimately their generalizations

<sup>&</sup>lt;sup>1</sup> In follow-up work [65], generating series of single-valued iterated Eisenstein integrals will be shown to exhibit even closer parallels to the genus-zero results of this work once the non-geometric parts  $z_{2k+1}$  of the zeta generators in [64] are completed to generators  $\sigma_{2k+1}$  including Tsunogai's derivations dual to holomorphic Eisenstein series [66–69].

to higher-genus surfaces or even higher-dimensional varieties. Hence, the results of this work are hoped to serve as a stepping stone towards an all-loop understanding of the role of the cosmic Galois group in quantum field theory and string theory, as well as the interplay of openand closed-string interactions.

# REVIEW OF MULTIPLE POLYLOGARITHMS

We follow the standard conventions in [4] and define multiple polylogarithms (MPLs) recursively by

$$G(a_1, a_2, \dots, a_w; z) = \int_0^z \frac{\mathrm{d}t}{t - a_1} G(a_2, \dots, a_w; t)$$
 (1)

with labels  $a_1, \ldots, a_w \in \mathbb{C}$ , argument  $z \in \mathbb{C}$ , (transcendental) weight  $w \in \mathbb{N}$  and convention  $G(\emptyset; z) = 1$ . Endpoint divergences are shuffle-regularized<sup>2</sup> with the assignment  $G(0; z) = \log(z)$  and  $G(z; z) = -\log(z)$  of regularized values at weight one (see e.g. [77]). Evaluation of MPLs with the restricted alphabet  $a_i \in \{0, 1\}$  at z = 1 yields multiple zeta values (MZVs)

$$\zeta_{n_1, n_2, \dots, n_r} = \sum_{0 < k_1 < k_2 < \dots < k_r}^{\infty} k_1^{-n_1} k_2^{-n_2} \dots k_r^{-n_r}$$

$$= (-1)^r G(\underbrace{0, \dots, 0}_{n_r - 1}, 1, \dots, \underbrace{0, \dots, 0}_{n_2 - 1}, 1, \underbrace{0, \dots, 0}_{n_1 - 1}, 1; 1)$$

of depth r and weight  $n_1 + \ldots + n_r$  with  $n_i \in \mathbb{N}$  and  $n_r \ge 2$ .

**A. Motivic coaction:** Goncharov's and Brown's work [4, 13, 14, 78] gives an explicit formula for the motivic coaction of MPLs (1) at arbitrary  $a_i, z \in \mathbb{C}$ . The outcome of the coaction is conventionally denoted by a tensor product, where the first and second entry are mathematically understood as motivic periods and de Rham periods, respectively; see [13, 14, 16, 79] for the deep algebraic-geometry background of these two types of periods. We will drop the tensor product and rely on superscripts to distinguish between the motivic and the de Rham entry: throughout this work, we will write  $G^{\mathfrak{m}}$  instead of  $G^{\mathfrak{m}} \otimes 1$  and  $G^{\mathfrak{dr}}$  instead of  $1 \otimes G^{\mathfrak{dr}}$  (and similarly for  $\zeta^{\mathfrak{m}}, \zeta^{\mathfrak{dr}}$ ). De Rham periods are only defined up to discontinuities such that powers of  $i\pi$  are set to zero in the second entry of the coaction, i.e. we have  $\zeta_{i}^{\mathfrak{dr}} = 0$ .

A simple class of terms in the motivic coaction of MPLs is obtained from deconcatenating its labels  $a_i$ 

$$\Delta G^{\mathfrak{m}}(a_{1}, a_{2}, \dots, a_{w}; z) = \sum_{j=0}^{w} G^{\mathfrak{m}}(a_{j+1}, \dots, a_{w}; z) \quad (3)$$

$$\times G^{\mathfrak{dr}}(a_{1}, \dots, a_{j}; z) + \dots$$

However, the coaction formula of [4, 13] typically yields numerous additional terms in the ellipsis which are more challenging to state in closed form and involve at least one factor of  $\zeta^{\mathfrak{dr}}$  or  $G^{\mathfrak{dr}}(\ldots; a_i)$  with  $a_i \neq z$ . The simplest example in the one-variable case, with  $a_i \in \{0, 1\}$ , is the last term  $G^{\mathfrak{m}}(1; z)\zeta_0^{\mathfrak{dr}}$  of

$$\Delta G^{\mathfrak{m}}(0,0,1,1;z) = G^{\mathfrak{dr}}(0,0,1,1;z) + G^{\mathfrak{m}}(1;z)G^{\mathfrak{dr}}(0,0,1;z) + G^{\mathfrak{m}}(1,1;z)G^{\mathfrak{dr}}(0,0;z) + G^{\mathfrak{m}}(0,1,1;z)G^{\mathfrak{dr}}(0;z) + G^{\mathfrak{m}}(0,0,1,1;z) + G^{\mathfrak{m}}(1;z)\zeta_{3}^{\mathfrak{dr}}$$
(4)

A main result of this work is a streamlined description of the general form of the ellipsis in (3). In particular, we will employ generating series to formulate the completion of (3) for MPLs in any number of variables  $a_i \in \{0, 1, y, \ldots\}$ . The rapid growth in complexity of this problem with the number of variables is illustrated by

$$\Delta G^{\mathfrak{m}}(1, y; z) = G^{\mathfrak{m}}(1, y; z) + G^{\mathfrak{m}}(y; z)G^{\mathfrak{dr}}(1; z) + G^{\mathfrak{dr}}(1, y; z) + G^{\mathfrak{m}}(1; z) [G^{\mathfrak{dr}}(1; y) - G^{\mathfrak{dr}}(0; y)] - G^{\mathfrak{m}}(y; z)G^{\mathfrak{dr}}(1; y)$$
(5)

**B.** f-alphabet: The so-called f-alphabet [14, 78] provides a useful description of MZVs that exposes the entirety of their currently known relations over  $\mathbb{Q}$ . By formally passing to motivic MZVs<sup>3</sup>,  $\zeta_{n_1,...,n_r} \to \zeta_{n_1,...,n_r}^{\mathfrak{m}}$ , we can isomorphically map MZVs to words in

- one commutative generator,  $f_2$ , whose  $n^{\text{th}}$  powers capture the even Riemann zeta values,  $\zeta_{2n} \in \mathbb{Q}\pi^{2n}$
- an infinite set of non-commutative generators,  $f_3, f_5, f_7, \ldots$ , in odd degrees  $\geq 3$ , which capture the odd Riemann zeta values  $\zeta_{2k+1}$

such that the counting of independent words  $f_2^n f_{i_1} f_{i_2} \dots f_{i_r}$  with  $n \in \mathbb{N}$  as well as  $i_1, \dots, i_r \in 2\mathbb{N}+1$  and  $f_{i_a} f_{i_b} \neq f_{i_b} f_{i_a}$  reproduces the expected number of  $\mathbb{Q}$ -independent MZVs at weight  $2n+i_1+\dots+i_r$  [80]. The isomorphism  $\phi$  into the f-alphabet preserves the shuffle product  $\phi(\zeta_{\vec{n}}^m \cdot \zeta_{\vec{u}}^m) = \phi(\zeta_{\vec{n}}^m) \sqcup \phi(\zeta_{\vec{u}}^m)$  (with  $\vec{n}, \vec{u} \in \mathbb{N}^{\times}$ ), e.g.  $f_i \sqcup f_j = f_i f_j + f_j f_i$ . The simplest examples read

$$\phi(\zeta_{2k+1}^{\mathfrak{m}}) = f_{2k+1}, \quad \phi(\zeta_{2}^{\mathfrak{m}}) = f_{2}, \quad \phi(\zeta_{3,5}^{\mathfrak{m}}) = -5f_{3}f_{5}$$

$$\phi(\zeta_{3,3,5}^{\mathfrak{m}}) = -5f_{3}f_{3}f_{5} - 45f_{9}f_{2} - \frac{6}{5}f_{7}f_{2}^{2} + \frac{4}{7}f_{5}f_{2}^{3}$$
 (6)

The images of conjecturally indecomposable higher-depth MZVs such as  $\phi(\zeta_{3,5}^{\mathfrak{m}})$  are determined by imposing  $\phi$  to preserve the motivic coaction. With the notation  $^{\mathfrak{m}}$  and  $^{\mathfrak{dr}}$  to distinguish the first and second entry of the tensor product, the coaction in the f-alphabet is by definition given by the simple deconcatenation formula

$$\Delta (f_2^n f_{i_1} \dots f_{i_r})^{\mathfrak{m}} = \sum_{j=0}^r (f_2^n f_{i_1} \dots f_{i_j})^{\mathfrak{m}} (f_{i_{j+1}} \dots f_{i_r})^{\mathfrak{dr}}$$
(7)

<sup>&</sup>lt;sup>2</sup> Shuffle-regularization amounts to imposing that the shuffle product  $G(\vec{a};z)G(\vec{b};z) = \sum_{\vec{c} \in \vec{a} \sqcup \vec{b}} G(\vec{c};z)$  (for ordered sets  $\vec{a}, \vec{b}, \vec{c}$  of labels) universal to iterated integrals extends to regularized values of formally divergent cases.

<sup>&</sup>lt;sup>3</sup> For the purposes of this work, the key point of the elaborate definition of motivic MZVs in algebraic-geometry literature [13, 14, 16, 79] is that they by definition obey no relations other than the currently known Q-relations among conventional MZVs.

However, since this requirement does not fix the coefficient of  $f_{n_1+\ldots+n_r}$  in  $\phi(\zeta_{n_1,\ldots,n_r}^{\mathfrak{m}})$  (with  $f_{2n}=\frac{\zeta_{2n}}{(\zeta_2)^n}f_2^n$  in case of even weight), the f-alphabet images beyond depth one are in general non-canonical. Following [30] and (6), our convention is to single out the higher-depth MZVs at weight  $w\geq 8$  that lead to the conjectural  $\mathbb{Q}$ -basis of [81],

basis MZVs 
$$\ni \zeta_{3,5}, \zeta_{3,7}, \zeta_{3,3,5}, \zeta_{3,9}, \zeta_{1,1,4,6}, \dots$$
 (8)

and to impose  $f_w$  to be absent in their  $\phi$  images, see (6).

C. Single-valued map: The MPLs of (1) are meromorphic and exhibit monodromies when the integration path from 0 to z is deformed to wind around the singular points  $a_1, \ldots, a_w$  of the integration kernels. Still, one can construct single-valued versions sv  $G(a_1, \ldots, a_w; z)$  that share the holomorphic differential equations of (1)

$$\partial_z \text{sv } G(a_1, a_2, \dots, a_w; z) = \frac{\text{sv } G(a_2, \dots, a_w; z)}{z - a_1}$$
 (9)

by combining meromorphic MPLs with their complex conjugates and MZVs. Explicit constructions of single-valued MPLs in one and multiple variables can be found in [38] and [40, 41], respectively. Generalizations of single-valued MPLs that also accommodate primitives of  $\frac{1}{z\bar{z}+az+b\bar{z}+c}$  with  $a,b,c\in\mathbb{C}$  are developed in [82] and go beyond the scope of this work (see [20, 83] for applications).

Similar to the contributions (3) to the motivic coaction of MPLs, the generating series in [38, 40, 41] show that the single-valued map includes the following simple class of terms:

$$\operatorname{sv} G(a_1, a_2, \dots, a_w; z) = \sum_{j=0}^{w} G(a_1, \dots, a_j; z)$$

$$\times \overline{G(a_w, \dots, a_{j+1}; z)} + \dots$$
(10)

A second main result of this work is a new streamlined formulation of the additional terms in the ellipsis. These terms rapidly grow in complexity with the number of variables  $a_i \in \{0, 1, y, \ldots\}$  and have as coefficients MZVs or single-valued MPLs in fewer variables.

Evaluation of single-valued MPLs in one variable at z=1 as in (2) defines single-valued MZVs [17, 36]

$$\zeta_{n_1,\dots,n_r}^{\text{sv}} = (-1)^r \text{sv } G(\underbrace{0,\dots,0}_{n_r-1},1,\dots,\underbrace{0,\dots,0}_{n_2-1},1,\underbrace{0,\dots,0}_{n_1-1},1;1)$$
(11)

Single-valued MZVs are expressible in terms of MZVs, but form a proper subset since for instance  $\zeta_{2n}^{\rm sv}=0$  and  $\zeta_{2n+1}^{\rm sv}=2\zeta_{2n+1}$ . Higher depth examples include

$$\zeta_{3,5}^{\text{sv}} = -10\zeta_3\zeta_5, \quad \zeta_{3,7}^{\text{sv}} = -28\zeta_3\zeta_7 - 12\zeta_5^2$$

$$\zeta_{3,3,5}^{\text{sv}} = 2\zeta_{3,3,5} - 5\zeta_3^2\zeta_5 + 90\zeta_2\zeta_9 + \frac{12}{5}\zeta_2^2\zeta_7 - \frac{8}{7}\zeta_2^3\zeta_5$$
(12)

The single-valued map, sv :  $\zeta_{n_1,n_2,\dots,n_r} \to \zeta_{n_1,n_2,\dots,n_r}^{\text{sv}}$ , takes the most convenient form in the f-alphabet:

sv 
$$f_2^n f_{i_1} f_{i_2} \dots f_{i_r} = \delta_{n,0} \sum_{j=0}^r f_{i_j} \dots f_{i_2} f_{i_1} \coprod f_{i_{j+1}} \dots f_{i_r}$$
(13)

This annihilates sv  $f_2 = 0$  and resembles the deconcatenation formula (7) for the coaction of odd-degree generators  $f_{2k+1}$ . Even though the single-valued map is only well-defined in a motivic setting [36], we omit superscripts  $\zeta^{\mathfrak{m}}$  and  $G^{\mathfrak{m}}$  in a slight abuse of notation except where they specify the entries of coaction formulas. Note that the single-valued map preserves the product structure  $\operatorname{sv}(A \cdot B) = \operatorname{sv}(A) \cdot \operatorname{sv}(B)$  for arbitrary combination A, B of MPLs and MZVs.

### KEY GENERATING SERIES

Our results for the motivic coaction and the single-valued map of MPLs are best expressed using generating series. In this section, we define the relevant generating series of MPLs and MZVs.

A. Polylogarithmic generating series: Let the alphabet A be a set of labels,  $a \in A$ , and introduce noncommuting formal variables  $e_a$  for each label. The meromorphic MPLs (1) are organised into the following generating series

$$\mathbb{G}_{A}(z; e_{a}) = \sum_{r=0}^{\infty} \sum_{a_{1}, \dots, a_{r} \in A} e_{a_{1}} \dots e_{a_{r}} G(a_{r}, \dots, a_{2}, a_{1}; z)$$
(14)

$$=1+\sum_{a_1\in A}e_{a_1}G(a_1;z)+\sum_{a_1,a_2\in A}e_{a_1}e_{a_2}G(a_2,a_1;z)+\dots$$

The generating series  $\mathbb{G}_A(z; e_a)$  is a solution to the following Knizhnik–Zamolodchikov (KZ) equation

$$\partial_z \mathbb{G}_A(z; e_a) = \mathbb{G}_A(z; e_a) \sum_{a_i \in A} \frac{e_{a_i}}{z - a_j}$$
 (15)

This is a special case of multivariable KZ equation

$$\partial_{z_i} \mathbb{F}(z_a; x_{bc}) = \mathbb{F}(z_a; x_{bc}) \sum_{j \neq i} \frac{x_{ij}}{z_i - z_j}$$
 (16)

where the braid operators  $x_{ij} = x_{ji}$  obey the Yang–Baxter relations [84, 85]

$$[x_{ij}, x_{ik} + x_{jk}] = 0, \quad [x_{ij}, x_{kl}] = 0$$
 (17)

for pairwise distinct i, j, k, l. The KZ equation (15) of the polylogarithmic series (14) is obtained by setting  $z_i \mapsto z$ ,  $z_j \mapsto a_j$ , and identifying  $x_{ij}$  with  $e_{a_j}$ . Since (17) implies no relations among the  $x_{ij}$  at fixed i, the universal enveloping algebra of the  $e_{a_j}$  in (14) is freely generated. An individual MPL,  $G(a_1, \ldots, a_r; z)$ , is uniquely specified by  $\mathbb{G}_A(z; e_a)$  as the coefficient of the word  $e_{a_r} \cdots e_{a_1}$ .

**B. Drinfeld associator:** Restricting (14) to the alphabet  $A = \{0,1\}$  gives the generating series,  $\mathbb{G}_A(z; e_0, e_1)$  of MPLs in one variable. Evaluating this at z = 1 yields the Drinfeld associator [86, 87]

$$\Phi(e_0, e_1) = \sum_{r=0}^{\infty} \sum_{a_1, \dots, a_r \in \{0, 1\}} e_{a_1} \dots e_{a_r} G(a_r, \dots, a_2, a_1; 1)$$

$$= 1 + \zeta_2[e_0, e_1] + \zeta_3[[e_0, e_1], e_0 + e_1] + \dots (18)$$

This is the generating series of shuffle-regularized MZVs, (2), with G(0;1) = G(1;1) = 0 [88]. Denote the coefficients in  $\Phi(e_0, e_1)$  of the odd zeta values,  $\zeta_{2k+1}$ , as

$$W_{2k+1}(e_0, e_1) \equiv \Phi(e_0, e_1) \Big|_{\zeta_{2k+1}}$$
 (19)

Because of the multitude of  $\mathbb{Q}$  relations among MZVs, coefficients of MZVs in  $\Phi(e_0, e_1)$  are non-canonical. Our convention is that all the MZVs at weight 2k+1 in  $\Phi(e_0, e_1)$  are to be decomposed into the conjectural  $\mathbb{Q}$ -basis with higher-depth elements in (8) and [81]. With this understood,  $W_{2k+1}$  is a unique Lie polynomial in  $e_i$ .

**C. Zeta generators:** For the results in this paper, it is important to introduce a second generating series of MZVs,  $\mathbb{M}$ , adapted to the f-alphabet. Introducing noncommuting formal generators  $M_3, M_5, \ldots$ , we define

$$\mathbb{M} = \sum_{r=0}^{\infty} \sum_{i_1, \dots, i_r \in 2\mathbb{N}+1} \phi^{-1}(f_{i_1} \dots f_{i_r}) M_{i_1} \dots M_{i_r}$$
 (20)

$$=1+\sum_{i_1\in 2\mathbb{N}+1}\zeta_{i_1}M_{i_1}+\sum_{i_1,i_2\in 2\mathbb{N}+1}\phi^{-1}(f_{i_1}f_{i_2})M_{i_1}M_{i_2}+\ldots$$

The zeta generators,  $M_{2k+1}$  [62, 63], have no relations among themselves and cannot be expressed in terms of braid operators  $e_{a_i}$ . However, we will later encounter an algebra on the  $M_{2k+1}$  and  $e_{a_i}$ , such that commutators  $[M_{2k+1}, e_{a_i}]$  are equal to nested commutators of the braid operators,  $e_{a_i}$ . In other words, the braid algebra is normalized by the  $M_{2k+1}$ .

However, recall that the f-alphabet of MZVs is noncanonical. For instance, departing from our convention specified in the discussion around (6) would redefine  $M_{11}$ by adding a rational multiple of  $[[M_3, M_5], M_3]$ . Similar ambiguities apply to  $M_{13}, M_{15}, \ldots$ , which can be redefined by nested brackets of an odd number of  $M_{2k+1}$ .

The generating series  $\mathbb{M}$  is related to similar series encountered in open-string tree-level amplitudes [30] and configuration-space integrals over marked points on the sphere [89]. However, in these contexts, one studies specific matrix representations of the Lie algebra satisfied by the  $M_{2k+1}$  and  $e_{a_i}$ . It is important for our results to refrain from specifying a matrix representation. In this way, we ensure that no terms in (20) drop out due to non-generic relations that can arise in specific matrix representations.

**D. Inverse series:** Both polylogarithms and the f-alphabet algebra obey the shuffle relations. This makes it straightforward to verify that the series inverses of  $\mathbb{G}_A$  and  $\mathbb{M}$  with respect to concatenation are

$$\mathbb{G}_A(z; e_a)^{-1} = \sum_{r=0}^{\infty} (-1)^r \sum_{a_1, \dots, a_r \in A} e_{a_1} \dots e_{a_r} G(a_1, \dots, a_r; z)$$

$$\mathbb{M}^{-1} = \sum_{r=0}^{\infty} (-1)^r \sum_{i_1, i_2, \dots, i_r \in 2\mathbb{N} + 1} \phi^{-1}(f_{i_1} f_{i_2} \dots f_{i_r}) M_{i_r} \dots M_{i_2} M_{i_1}$$
(21)

## POLYLOGARITHMS IN ONE VARIABLE

We begin by presenting our results as applied to MPLs in one variable. The generalisation to more than one variable is given in later sections. Recall that the generating series for MPLs in one variable is  $\mathbb{G}_{\{0,1\}}(z) = \mathbb{G}_{\{0,1\}}(z; e_0, e_1)$ , where the alphabet in (14) is specialized to  $A = \{0, 1\}$ .

**A.** The motivic coaction: Our first main result is a formula for the motivic coaction. We claim that the motivic coaction is given by

$$\Delta \mathbb{G}^{\mathfrak{m}}_{\{0,1\}}(z) = (\mathbb{M}^{\mathfrak{dr}})^{-1} \, \mathbb{G}^{\mathfrak{m}}_{\{0,1\}}(z) \, \mathbb{M}^{\mathfrak{dr}} \, \mathbb{G}^{\mathfrak{dr}}_{\{0,1\}}(z) \qquad (22)$$

where the coaction  $\Delta$  of individual MPLs in one variable is uniquely specified by the coefficient

$$\Delta G^{\mathfrak{m}}(a_1, a_2, \dots, a_r; z) = \Delta \mathbb{G}^{\mathfrak{m}}_{\{0,1\}}(z) \big|_{e_{a_r} \dots e_{a_2} e_{a_1}}$$
 (23)

of the generating series  $\Delta \mathbb{G}_{\{0,1\}}^{\mathfrak{m}}(z)$ .

The 'motivic' and 'de Rham' superscripts of  $\mathbb{G}^{\mathfrak{m}}$ ,  $\mathbb{G}^{\mathfrak{dr}}$ ,  $\mathbb{M}^{\mathfrak{dr}}$  apply to the MPLs and MZVs in the respective expansions (14), (20), and they distinguish the first and second entry of the series  $\Delta \mathbb{G}^{\mathfrak{m}}_{\{0,1\}}(z)$ . The multiplication of the generating series in (22) is just the concatenation product of the braid operators  $e_0$ ,  $e_1$  and the zeta generators  $M_{2k+1}$ .

As a first check of (22), note that the simple terms (3) in the coaction formula for MPLs are recovered by ignoring all the MZV terms in M (i.e. setting  $\mathbb{M}^{\mathfrak{dr}} \to 1$  in (22)). The additional terms involving MZVs arise in our formula (22) from commutators of the  $M_{2k+1}$  and  $e_a$ . This algebra is specified below in subsection  $\mathbf{C}$ .

**B.** The sv map: Our second main result is a formula for the single-valued map, sv. We claim that the sv map on MPLs in one variable is given by

$$\operatorname{sv} \mathbb{G}_{\{0,1\}}(z) = (\operatorname{sv} \mathbb{M})^{-1} \overline{\mathbb{G}_{\{0,1\}}(z)^t} (\operatorname{sv} \mathbb{M}) \mathbb{G}_{\{0,1\}}(z)$$
 (24)

where the action of sv on individual MPLs is uniquely specified by the coefficient

$$\operatorname{sv} G(a_1, a_2, \dots, a_r; z) = \operatorname{sv} \mathbb{G}_{\{0,1\}}(z) \big|_{e_{a_r} \dots e_{a_2} e_{a_1}}$$
 (25)

of the generating series sv  $\mathbb{G}_{\{0,1\}}(z)$ .

The factor  $\mathbb{G}_{\{0,1\}}(z)^t$  in (24) is defined from the series  $\mathbb{G}_{\{0,1\}}(z)$  by applying complex conjugation to the MPLs and word reversal on the braid operators:  $(e_1 \cdots e_j)^t = e_j \cdots e_1$ . The factor sv M in (24) is a generating series of single-valued MZVs. This is defined from the series expansion (20) of M by acting with the sv on the f-algebra, (13). For example, the first few terms are

$$\operatorname{sv} \mathbb{M} = 1 + 2 \sum_{i_1 \in 2\mathbb{N} + 1} \zeta_{i_1} M_{i_1} + 2 \sum_{i_1, i_2 \in 2\mathbb{N} + 1} \zeta_{i_1} \zeta_{i_2} M_{i_1} M_{i_2}$$
 (26)  
 
$$+ \sum_{i_1, i_2, i_3 \in 2\mathbb{N} + 1} \phi^{-1} \left( \operatorname{sv}(f_{i_1} f_{i_2} f_{i_3}) \right) M_{i_1} M_{i_2} M_{i_3} + \dots$$

The factors of 2 arise from  $sv(f_i) = 2f_i$  and  $sv(f_if_j) = 2f_i \sqcup f_j$ , and  $\phi^{-1}(sv(f_{i_1}f_{i_2}f_{i_3}))$  at weight  $i_1+i_2+i_3 \geq 11$  comprise (conjecturally) irreducible single-valued MZVs at depth  $\geq 3$ , see for instance (6) and (12).

As a first check of (24), note that the simple terms (10) in the expression for single-valued MPLs are recovered by setting sv  $\mathbb{M} \to 1$  in (24), i.e. ignoring all MZVs.

C. Commutators with zeta generators: Both of the new formulas, (22) and (24), for the motivic coaction and sv map of MPLs in one variable involve conjugations by a series in MZVs and zeta generators  $M_{2k+1}$ . In order to attain well-defined coefficients of  $e_{a_r} \dots e_{a_1}$  in (23) and (25), we must specify how conjugations with  $M_{2k+1}$  conspire with the braid operators to yield a series solely in  $e_i$ .

As a first step, we expand the conjugation of an arbitrary series  $\mathbb{G}$  in the braid operators  $e_0, e_1$  with  $\mathbb{M}$  in terms of nested commutators

$$\phi((\mathbb{M})^{-1}\mathbb{GM}) = \sum_{r=0}^{\infty} \sum_{i_1, i_2, \dots, i_r \in 2\mathbb{N}+1} f_{i_1} f_{i_2} \dots f_{i_r}$$

$$\times \left[ \left[ \dots \left[ \left[ \mathbb{G}, M_{i_1} \right], M_{i_2} \right], \dots, M_{i_{r-1}} \right], M_{i_r} \right]$$

$$= \mathbb{G} + \sum_{i_1 \in 2\mathbb{N}+1} f_{i_1} \left[ \mathbb{G}, M_{i_1} \right] + \sum_{i_1, i_2 \in 2\mathbb{N}+1} f_{i_1} f_{i_2} \left[ \left[ \mathbb{G}, M_{i_1} \right], M_{i_2} \right] + \dots$$

The second step is to specify the following commutation relations among the zeta generators  $M_{2k+1}$  that normalize the free algebra generated by  $e_0, e_1$ 

$$[e_0, M_{2k+1}] = 0, \quad [e_1, M_{2k+1}] = [W_{2k+1}(e_0, e_1), e_1]$$
(28)

where  $W_{2k+1}(e_0, e_1)$  are the Lie polynomials appearing in the expansion of the Drinfeld associator, (19). For example, identifying  $W_3(e_0, e_1) = [[e_0, e_1], e_0 + e_1]$  from (18),

$$[e_0, M_3] = 0, \quad [e_1, M_3] = [[e_0, e_1], e_0 + e_1], e_1]$$
 (29)

These relations featured in the context of the stable derivation algebra<sup>4</sup> [90, 91] and were later observed for specific matrix representations in (4.23) of [89].

Upon iterative use of the commutators (28) in (27), all the zeta generators entering (22) and (24) via  $(\mathbb{M}^{\mathfrak{dr}})^{-1}\mathbb{G}^{\mathfrak{m}}_{\{0,1\}}(z)\mathbb{M}^{\mathfrak{dr}}$  and  $(\operatorname{sv}\mathbb{M})^{-1}\overline{\mathbb{G}_{\{0,1\}}(z)^{t}}(\operatorname{sv}\mathbb{M})$  conspire to yield series  $\Delta\mathbb{G}^{\mathfrak{m}}_{\{0,1\}}(z)$  and  $\operatorname{sv}\mathbb{G}_{\{0,1\}}(z)$  solely in  $e_0$  and  $e_1$ . This is a crucial prerequisite to extract the motivic coaction and single-valued map of MPLs in one variable from the respective coefficients in (23) and (25).

For example, the simplest contribution to the motivic coaction (22) involving a MZV from M is

$$(\mathbb{M}^{\mathfrak{dr}})^{-1} \mathbb{G}_{\{0,1\}}(z) \mathbb{M}^{\mathfrak{dr}} \Big|_{\zeta_3^{\mathfrak{dr}}}$$

$$= G(1; z) \big[ \big[ [e_0, e_1], e_0 + e_1 \big], e_1 \big] + \dots$$
(30)

with MPLs of weight  $\geq 2$  and words involving  $\geq 5$  braid operators in the ellipsis. Isolating the word  $e_1e_1e_0e_0$  in (30), this reproduces the contribution  $G^{\mathfrak{m}}(1;z)\zeta_3^{\mathfrak{dr}}$  to  $\Delta G^{\mathfrak{m}}(0,0,1,1;z)$  in the introductory example (4).

Likewise, the simplest MZV contribution to (24) is

$$(\operatorname{sv} \mathbb{M})^{-1} \overline{\mathbb{G}_{\{0,1\}}(z)^{t}} (\operatorname{sv} \mathbb{M}) \Big|_{\zeta_{3}}$$

$$= 2\overline{G(1;z)} [[[e_{0},e_{1}],e_{0}+e_{1}],e_{1}] + \dots$$
(31)

where the ellipsis contains words with  $\geq 5$  braid operators. Isolating the word  $e_1e_1e_0e_0$ , this reproduces the contribution  $2\zeta_3\overline{G(1;z)}$  to sv G(0,0,1,1;z).

**D.** Equivalence with earlier results: We claim that the formula (22) for the motivic coaction on MPLs in one variable is equivalent to the Ihara formula [92]

$$\Delta \mathbb{G}_{\{0,1\}}^{\mathfrak{m}}(z; e_{0}, e_{1}) = \mathbb{G}_{\{0,1\}}^{\mathfrak{m}}(z; e_{0}, e_{1}') \mathbb{G}_{\{0,1\}}^{\mathfrak{dr}}(z; e_{0}, e_{1})$$
$$e_{1}' = \Phi^{\mathfrak{dr}}(e_{0}, e_{1}) e_{1} \Phi^{\mathfrak{dr}}(e_{0}, e_{1})^{-1} \qquad (32)$$

which is implicit in (6.6) of [36] and explicit in Proposition 8.3 of [93]. Similarly, we claim that the formula (24) for the generating series of single-valued MPLs is equivalent to the following formula of Brown [38]

$$\operatorname{sv} \mathbb{G}_{\{0,1\}}(z; e_0, e_1) = \overline{\mathbb{G}_{\{0,1\}}(z; e_0, \hat{e}_1)^t} \mathbb{G}_{\{0,1\}}(z; e_0, e_1)$$

$$\hat{e}_1 = \left(\operatorname{sv} \Phi(e_0, e_1)\right) e_1 \left(\operatorname{sv} \Phi(e_0, e_1)\right)^{-1}$$

$$= \operatorname{sv} e'_1 \tag{33}$$

Unlike our expressions, (22) and (24), the earlier formulas, (32) and (33), employ a change of alphabet  $e_1 \to e'_1$  or  $\hat{e}_1$  for one of the braid operators within the series  $\mathbb{G}^{\mathfrak{m}}_{\{0,1\}}(z)$  or  $\overline{\mathbb{G}_{\{0,1\}}(z)^t}$ .

A full proof that our results follow from these earlier formulas is to be given in [94]. Here we sketch the argument in brief. First, insert  $1 = \mathbb{M}^{\mathfrak{dr}}(\mathbb{M}^{\mathfrak{dr}})^{-1}$  and  $1 = \text{sv } \mathbb{M}(\text{sv } \mathbb{M})^{-1}$  between any pair of braid operators  $e_i$  in the series  $\mathbb{G}^{\mathfrak{m}}_{\{0,1\}}(z)$  in (22) and  $\overline{\mathbb{G}}_{\{0,1\}}(z)^t$  in (24), respectively. Given that  $e_0$  commutes with the  $M_{2k+1}$ ,

$$(\mathbb{M}^{\mathfrak{dr}})^{-1}e_0\,\mathbb{M}^{\mathfrak{dr}} = (\operatorname{sv}\,\mathbb{M})^{-1}e_0\,(\operatorname{sv}\,\mathbb{M}) = e_0 \tag{34}$$

our formulas, (22) and (24), can be rewritten as

$$\Delta \mathbb{G}^{\mathfrak{m}}_{\{0,1\}}(z; e_{0}, e_{1}) = \mathbb{G}^{\mathfrak{m}}_{\{0,1\}}(z; e_{0}, (\mathbb{M}^{\mathfrak{dr}})^{-1} e_{1} \mathbb{M}^{\mathfrak{dr}}) \times \mathbb{G}^{\mathfrak{dr}}_{\{0,1\}}(z; e_{0}, e_{1})$$

$$\operatorname{sv} \mathbb{G}_{\{0,1\}}(z; e_{0}, e_{1}) = \overline{\mathbb{G}_{\{0,1\}}(z; e_{0}, (\operatorname{sv} \mathbb{M})^{-1} e_{1} \operatorname{sv} \mathbb{M})^{t}} \times \mathbb{G}_{\{0,1\}}(z; e_{0}, e_{1})$$

<sup>&</sup>lt;sup>4</sup> For instance,  $W_3$  satisfies the equation for commutators 'f' in the formulation of standard derivation algebra as stated in section 2.2 of [90]. Taking the commutators in (29) with respect to  $M_3$  gives rise to the corresponding derivation  $D_{W_3}$  on the free Lie algebra generated by  $e_0, e_1$ .

Next, it remains to show that

$$(\mathbb{M}^{\mathfrak{dr}})^{-1} e_1 \mathbb{M}^{\mathfrak{dr}} = \Phi^{\mathfrak{dr}}(e_0, e_1) e_1 \Phi^{\mathfrak{dr}}(e_0, e_1)^{-1}$$
 (36)

As a first check, note that the coefficients of  $\zeta_{2k+1}^{\mathfrak{dr}}$  on both sides of this equation agree, as a consequence of the commutators (28). In the full proof of (36), to be given in [94], matching the appearance of  $f_{i_1} \dots f_{i_r}$  at  $r \geq 2$  relies on the coaction properties of  $\Phi(e_0, e_1)$  and  $\mathbb{M}$ .

Note that the changes of alphabet in the earlier formulas, (32) and (33), cannot be rewritten by conjugating  $\mathbb{G}^{\mathfrak{m}}_{\{0,1\}}(z)$  or  $\overline{\mathbb{G}_{\{0,1\}}(z)^t}$  with the Drinfeld associator,  $\Phi^{\mathfrak{dr}}(e_0,e_1)$  or sv  $\Phi(e_0,e_1)$ . This is because the Drinfeld associator does not commute with  $e_0$ . Our formulas are only made possible by introducing the zeta generators  $M_{2k+1}$  outside the universal enveloping algebra of  $e_0,e_1$ .

**E.** Correlations between different MZVs: Our formulas, (22) and (24), for the motivic coaction and the single-valued map, expose a key fact. In both cases, the coefficients of higher-depth MZVs,  $\phi^{-1}(f_{i_1} \dots f_{i_r})$  at  $r \geq 2$ , are completely determined by the coefficients of the odd zeta values,  $\zeta_i = \phi^{-1}(f_i)$ . This is closely related to the simple properties

$$\Delta \mathbb{M}^{\mathfrak{m}} = \mathbb{M}^{\mathfrak{m}} \mathbb{M}^{\mathfrak{dr}}, \quad \operatorname{sv} \mathbb{M} = \mathbb{M}^{t} \mathbb{M}$$
 (37)

of the series M. By contrast, the Drinfeld associator does not satisfy these properties. The expressions for  $\Delta \Phi^{\mathfrak{m}}(e_0, e_1)$  and sv  $\Phi(e_0, e_1)$  [31, 36] necessitate the above change of alphabet  $e_1 \to e'_1$  or  $\hat{e}_1$ .

The proof of (36) in [94] follows from understanding how these properties of  $\mathbb{M}$  and  $\Phi(e_0, e_1)$  are related. The Drinfeld associator enjoys an expansion similar to that of  $\mathbb{M}$  in (20), in which the concatenation product on the zeta generators  $M_{2k+1}$  is replaced by a certain product  $\circ$  on the Lie polynomials  $W_{2k+1}$  [31, 95].

# POLYLOGARITHMS IN TWO VARIABLES

The results for MPLs in one variable generalize to any number of variables. Before presenting the general result in the next section, we devote this section to the case of MPLs in two variables. In this way, the additional features of the Lie algebras governing the multivariable case (as opposed to MPLs in one variable) will be illustrated in a combinatorially simple setting.

We study MPLs in two variables, y and z, in the fibration basis involving products of MPLs  $G(a_1, \ldots, a_r; z)$   $(a_i \in \{0, 1, y\})$  and  $G(b_1, \ldots, b_s; y)$   $(b_i \in \{0, 1\})$ . The respective generating series for these MPLs are now defined using two sets of braid group generators. To avoid confusion, we use the superscript  $^{(2)}$  to distinguish the new generators from the braid operators  $e_0, e_1$  of the one-variable case. Hence, the protagonist of this section is the generating series (14) with  $A = \{0, 1, y\}$ ,

$$\mathbb{G}_{\{0,1,y\}}(z) = \mathbb{G}_{\{0,1,y\}}(z; e_0^{(2)}, e_1^{(2)}, e_y^{(2)})$$
 (38)

**A.** The results: For MPLs in two variables, we claim that the motivic coaction is generated by the formula

$$\Delta \mathbb{G}_{\{0,1,y\}}^{\mathfrak{m}}(z) = \left(\mathbb{G}_{\{0,1\}}^{\mathfrak{dr}}(y)\right)^{-1} (\mathbb{M}^{\mathfrak{dr}})^{-1} \mathbb{G}_{\{0,1,y\}}^{\mathfrak{m}}(z) \times \mathbb{M}^{\mathfrak{dr}} \mathbb{G}_{\{0,1\}}^{\mathfrak{dr}}(y) \mathbb{G}_{\{0,1,y\}}^{\mathfrak{dr}}(z)$$
(39)

Similarly, for MPLs in two variables, our formula for the single-valued map is

$$\operatorname{sv} \mathbb{G}_{\{0,1,y\}}(z) = \left(\operatorname{sv} \mathbb{G}_{\{0,1\}}(y)\right)^{-1} \left(\operatorname{sv} \mathbb{M}\right)^{-1} \overline{\mathbb{G}_{\{0,1,y\}}(z)^{t}} \times \left(\operatorname{sv} \mathbb{M}\right) \left(\operatorname{sv} \mathbb{G}_{\{0,1\}}(y)\right) \mathbb{G}_{\{0,1,y\}}(z)$$
(40)

see (26) and (24) for the generating series sv  $\mathbb{M}$  and sv  $\mathbb{G}_{\{0,1\}}(y)$  of single-valued MZVs and one-variable MPLs, respectively.

The action of  $\Delta$  and sv on a specific MPL  $G(a_1,\ldots,a_r;z)$  in two variables  $(a_i\in\{0,1,y\})$  is uniquely specified by extracting the coefficient of the word  $e_{a_r}^{(2)}\cdots e_{a_1}^{(2)}$  in (39) and (40), respectively. However, both (39) and (40) feature zeta generators  $M_{2k+1}$  and the distinct braid operators  $e_0,e_1$  of the one-variable case on the right-hand side. To use these formulas, one must eliminate all of  $M_{2k+1},e_0,e_1$  in favor of the generators  $e_0^{(2)},e_1^{(2)},e_y^{(2)}$ . This can be done by using the Yang–Baxter relations of the braid algebra, (17), and also extending the braid algebra by specifying the commutators of the  $M_{2k+1}$  with the braid generators.

**B.** The braid algebra: In the present notation, the braid-algebra relations for two variables are given by

$$[e_0^{(2)}, e_0] = 0, [e_0^{(2)}, e_1] = 0 (41)$$

$$[e_1^{(2)}, e_0] = [e_1^{(2)}, e_y^{(2)}], [e_1^{(2)}, e_1] = [e_y^{(2)}, e_1^{(2)}]$$

$$[e_y^{(2)}, e_0] = [e_0^{(2)}, e_y^{(2)}], [e_y^{(2)}, e_1] = [e_1^{(2)}, e_y^{(2)}]$$

These commutators can be derived from the Yang–Baxter relations (17) under the identifications

$$e_0^{(2)} = x_{z0}, e_1^{(2)} = x_{z1}, e_y^{(2)} = x_{yz}$$
  
 $e_0 = x_{y0} + x_{yz}, e_1 = x_{y1}$  (42)

which follow from the KZ equation (16) of the product,  $\mathbb{G}_{\{0,1\}}(y)\mathbb{G}_{\{0,1,y\}}(z)$ , as in [89].

The key point is that the braid-algebra relations (41) do not feature any operators  $e_0, e_1$  of the one-variable case on their right-hand side. One can therefore eliminate any appearance of  $e_0, e_1$  in favor of  $e_i^{(2)}$  on the right-hand sides of our main formulas (39) and (40). More specifically, iterative use of (41) ensures that the expansion of

$$\mathbb{G}_{\{0,1\}}(y)^{-1} \mathbb{X} \mathbb{G}_{\{0,1\}}(y) = \mathbb{X} + \sum_{a_1 \in \{0,1\}} G(a_1; y) [\mathbb{X}, e_{a_1}] + \sum_{a_1, a_2 \in \{0,1\}} G(a_1, a_2; y) [[\mathbb{X}, e_{a_2}], e_{a_1}] + \dots (43)$$

preserves the alphabet  $\{e_0^{(2)},e_1^{(2)},e_y^{(2)}\}$  for any series  $\mathbb X$  in these three variables.

C. The extension by zeta generators: Finally, to complete the presentation of the main results (39) and (40) in terms of  $e_0^{(2)}$ ,  $e_1^{(2)}$ ,  $e_y^{(2)}$ , we extend the braid algebra (41) by the following commutators for zeta generators:

$$\begin{split} [e_0^{(2)}, M_{2k+1}] &= 0 \tag{44} \\ [e_y^{(2)}, M_{2k+1}] &= [W_{2k+1}(e_0^{(2)}, e_y^{(2)}), e_y^{(2)}] \\ [e_1^{(2)}, M_{2k+1}] &= [W_{2k+1}(e_0^{(2)} + e_y^{(2)}, e_1^{(2)}) + W_{2k+1}(e_0^{(2)}, e_y^{(2)}) \\ &+ W_{2k+1}(e_y^{(2)}, e_0 - e_y^{(2)}), e_1^{(2)}] \end{split}$$

Again,  $W_{2k+1}$  are the Lie-polynomial coefficients of the series expansion of the Drinfeld associator (19). These commutators involving  $e_i^{(2)}$  generalize the Lie brackets (28) with zeta generators in the one-variable case. The derivation of (44) will be given in [94] and is based on analytic continuations of the series  $\mathbb{G}_{\{0,1,y\}}(z)$  in two-variable MPLs. Note that the braid-algebra relations (41) can be used to eliminate the  $e_0$  from the Lie polynomial  $W_{2k+1}(e_y^{(2)}, e_0 - e_y^{(2)})$  in the last line of (44).

The extended braid algebra in (44) and (41) can now be used to extract the coaction and sv map of individual MPLs in two variables from their generating series (39) and (40). The first step is to conjugate by  $\mathbb{M}^{\mathfrak{dr}}$  (or by sv  $\mathbb{M}$ ) via (27), and to then eliminate the  $M_{2k+1}$  from the series using the commutators (44). The second step is to conjugate by  $\mathbb{G}^{\mathfrak{dr}}_{\{0,1\}}(y)$  (or by sv  $\mathbb{G}_{\{0,1\}}(y)$ ) via (43), and to then eliminate the  $e_0, e_1$  using the braid relations (41).

The introductory example (5) for  $\Delta G^{\mathfrak{m}}(1,y;z)$  can be extracted from the coefficient of  $e_{y}^{(2)}e_{1}^{(2)}$  in the coaction formula (39). The weight-one terms of  $\mathbb{G}_{\{0,1,y\}}^{\mathfrak{m}}(z)$  on the right-hand side suffice to recover the second line of (5) from the conjugation (43) and the braid algebra (41).

**D. Equivalence with earlier results:** The proof of our results, (39) and (40), again follows by showing that they are equivalent to earlier formulas. In [94], it will be shown that our coaction formula (39) is equivalent to the multivariate generalization of the Ihara formula in [93], and that our formula for the sv map (40) is equivalent to the results in [40] for single valued MPLs in two variables.

As in the single-variable case, these earlier formulas introduce a change of alphabet, defining new letters such as  $\hat{e}_1^{(2)}, \hat{e}_y^{(2)}$  in case of the single-valued map [40]. The first step of the proof is to show that the conjugation by  $\mathbb{M}^{\mathfrak{dr}} \mathbb{G}^{\mathfrak{dr}}_{\{0,1\}}(y)$  in (39) and by  $\mathrm{sv}(\mathbb{MG}_{\{0,1\}}(y))$  in (40) can be rewritten as a change of alphabet from  $e_0^{(2)}, e_1^{(2)}, e_y^{(2)}$  to the new alphabet of [93] and [40]. Note that  $e_0^{(2)}$  is not changed in this substitution, and this is consistent with the fact that  $e_0^{(2)}$  commutes with  $M_{2k+1}, e_0, e_1$  (see for comparison the monodromy arguments of [40]).

### GENERAL MULTIVARIABLE RESULT

This section gives our results for the motivic coaction and the single-valued map of MPLs in full generality, for any number n of variables. The formulas are structurally analogous to those for MPLs in two variables presented in the previous section. As we will see, the additional challenges of the n-variable case are mostly of combinatorial nature, in particular to present the explicit form of the braid algebra and its extension by zeta generators.

At the end of this section, we comment on how our general results relate to the properties of period matrices, and explain how the new results relate to previous works on configuration-space integrals at genus zero subject to multivariable KZ equations.

**A. Notation:** We again specify a fibration basis for MPLs where the labels of  $G(a_1, \ldots, a_r; z_n)$  are restricted to  $a_i \in \{0, 1, z_1, \ldots, z_{n-1}\}$ . In order to state the general result, it is helpful to introduce the following shorthand for the generating series of n-variable MPLs

$$\mathbb{G}_{n} := \mathbb{G}_{A = \{0, 1, z_{1}, z_{2}, \dots, z_{n-1}\}}(z_{n}; e_{i}^{(n)})$$

$$= \sum_{r=0}^{\infty} \sum_{\substack{a_{1}, \dots, a_{r} \\ \in \{0, 1, z_{1}, z_{2}, \dots, z_{n-1}\}}} e_{a_{1}}^{(n)} \dots e_{a_{r}}^{(n)} G(a_{1}, \dots, a_{r}; z_{n})$$

In this notation, the n+1 letters in the alphabet A and the associated braid operators  $e_i^{(n)}$  are left implicit. In particular, for n=1 or n=2 variables, we now write

$$\mathbb{G}_{1} = \mathbb{G}_{\{0,1\}}(z_{1}; e_{0}^{(1)}, e_{1}^{(1)})$$

$$\mathbb{G}_{2} = \mathbb{G}_{\{0,1,z_{1}\}}(z_{2}; e_{0}^{(2)}, e_{1}^{(2)}, e_{z_{1}}^{(2)})$$
(46)

with  $e_i^{(1)} = e_i$  in earlier sections. In this notation, our formulas for n = 2, (39) and (40), become

$$\Delta \mathbb{G}_{2}^{\mathfrak{m}} = (\mathbb{M}^{\mathfrak{dr}} \mathbb{G}_{1}^{\mathfrak{dr}})^{-1} \, \mathbb{G}_{2}^{\mathfrak{m}} \, \mathbb{M}^{\mathfrak{dr}} \, \mathbb{G}_{1}^{\mathfrak{dr}} \mathbb{G}_{2}^{\mathfrak{dr}}$$

$$\operatorname{sv} \mathbb{G}_{2} = \operatorname{sv}(\mathbb{M} \, \mathbb{G}_{1})^{-1} \, \overline{\mathbb{G}_{2}^{t}} \, \operatorname{sv}(\mathbb{M} \, \mathbb{G}_{1}) \, \mathbb{G}_{2}$$

$$(47)$$

**B. The results:** For MPLs in any number n of variables, our main results for the motivic coaction and the single-valued map are generated by

$$\Delta \mathbb{G}_{n}^{\mathfrak{m}} = (\mathbb{M}^{\mathfrak{dr}} \, \mathbb{G}_{1}^{\mathfrak{dr}} \dots \mathbb{G}_{n-1}^{\mathfrak{dr}})^{-1} \, \mathbb{G}_{n}^{\mathfrak{m}} \, \mathbb{M}^{\mathfrak{dr}} \, \mathbb{G}_{1}^{\mathfrak{dr}} \dots \mathbb{G}_{n-1}^{\mathfrak{dr}} \, \mathbb{G}_{n}^{\mathfrak{dr}}$$

$$\operatorname{sv} \, \mathbb{G}_{n} = \operatorname{sv}(\mathbb{M} \, \mathbb{G}_{1} \dots \mathbb{G}_{n-1})^{-1} \, \overline{\mathbb{G}_{n}^{t}} \, \operatorname{sv}(\mathbb{M} \, \mathbb{G}_{1} \dots \mathbb{G}_{n-1}) \, \mathbb{G}_{n}$$

$$(48)$$

where the second line recursively determines sv  $\mathbb{G}_n$  in terms of generating series sv  $\mathbb{G}_i$  in fewer variables j < n.

As a key advantage of these formulas over earlier expressions for the coaction [4, 13, 14, 78] and the single-valued map [40, 41], the right-hand sides are already in fully simplified form. First, the right-hand side of (48) incorporates all known  $\mathbb{Q}$ -relations among MZVs by virtue of the f-alphabet representation (20) of  $\mathbb{M}$ . Second, all

MPLs  $G(b_1, \ldots, b_r; z_j)$  in  $j \leq n$  variables on the right-hand side of (48) are already in a uniform fibration basis with  $b_i \in \{0, 1, z_1, \ldots, z_{j-1}\}$ .

The proof of our main results (48) will be given in [94], by demonstrating that the coaction formula is equivalent to the multivariate Ihara formula [93], and that the sv formula is equivalent to the construction of single-valued polylogarithms in [40].

In order to read off the coaction and single-valued map of individual n-variable MPLs, it remains to understand the braid algebra obeyed by the operators  $e_i^{(j)}$  at  $j \leq n$  and its extension by the zeta generators  $M_{2k+1}$ . Using the commutation relations of this algebra, the series in (48) can be systematically expanded in terms of solely the braid operators  $e_i^{(n)}$  with  $i \in \{0, 1, z_1, \ldots, z_{n-1}\}$  at fixed n. Since the universal enveloping algebra of the  $e_i^{(n)}$  entering  $\mathbb{G}_n$  in (45) is freely generated, the coaction and single-valued map of n-variable MPLs  $G(a_1, \ldots, a_r; z_n)$  is uniquely specified by the coefficients of  $e_{a_r} \ldots e_{a_1}$  (48).

Following the treatment of the two-variable case, we will present the commutation relations for the extended braid algebra in two steps in the next subsections: the Yang–Baxter relations for the braid algebras in the *n*-variable case in subsection  $\mathbf{C}$ , followed by the brackets of  $e_i^{(n)}$  with zeta generators in subsection  $\mathbf{D}$ .

- **C. The braid algebra:** In the present notation, the braid algebra for the *n*-variables case is defined by the commutators  $[e_a^{(n)}, e_b^{(m)}]$ , with m < n. We group these commutators according to the following three cases:
  - let m < n and j < n, then:

$$\begin{split} &[e_0^{(n)},e_0^{(m)}]=0\,,\qquad [e_1^{(n)},e_0^{(m)}]=[e_1^{(n)},e_{z_m}^{(n)}] \qquad (49)\\ &[e_0^{(n)},e_1^{(m)}]=0\,,\qquad [e_1^{(n)},e_1^{(m)}]=[e_{z_m}^{(n)},e_1^{(n)}]\\ &[e_0^{(n)},e_{z_j}^{(m)}]=0\,,\qquad [e_1^{(n)},e_{z_j}^{(m)}]=0 \end{split}$$

• let m < n and i < n, then:

$$[e_{z_i}^{(n)}, e_0^{(m)}] = \begin{cases}
 [e_{z_i}^{(n)}, e_{z_m}^{(n)}], & i < m \\
 [e_0^{(n)} + \sum_{j=m+1}^{n-1} e_{z_j}^{(n)}, e_{z_m}^{(n)}], & i = m \\
 0, & i > m
 \end{cases}$$

$$[e_{z_i}^{(n)}, e_1^{(m)}] = \begin{cases}
 [e_1^{(n)}, e_{z_m}^{(n)}], & i = m \\
 0, & i \neq m
 \end{cases}$$
(50)

• let m < n, i < n, and j < m, then:

$$[e_{z_i}^{(n)}, e_{z_j}^{(m)}] = \begin{cases} [e_{z_m}^{(n)}, e_{z_i}^{(n)}], & i = j \\ [e_{z_j}^{(n)}, e_{z_m}^{(n)}], & i = m \\ 0, & i \neq j, m \end{cases}$$
(51)

The key point is that these commutation relations reduce any commutator  $[e_a^{(n)}, e_b^{(m)}]$  with m < n to the braid operators  $e_i^{(n)}$  that appear in the generating series  $\mathbb{G}_n$ .

The commutation relations (49) to (51) follow from the Yang–Baxter relations (17) under the identifications

$$e_0^{(n)} = x_{z_n 0}, e_1^{(n)} = x_{z_n 1}, e_{z_j}^{(n)} = x_{z_n z_j}$$

$$e_0^{(m)} = x_{z_m 0} + \sum_{j=m+1}^n x_{z_m z_j}, e_1^{(m)} = x_{z_m 1}$$

$$e_{z_j}^{(m)} = x_{z_m z_j}, m < n (52)$$

**D.** The extension by zeta generators: To complete the description of the extended braid algebra for n variables, we spell out the commutators of the zeta generators  $M_{2k+1}$  with the braid operators  $e_i^{(n)}$  entering  $\mathbb{G}_n$ . The combinatorially more involved generalization of the one- and two-variable cases (28) and (44) is given by

$$[e_{0}^{(n)}, M_{2k+1}] = 0$$

$$[e_{z_{j}}^{(n)}, M_{2k+1}] = \left[W_{2k+1}(e_{0}^{(n)} + \sum_{i=j+1}^{n-1} e_{z_{i}}^{(n)}, e_{z_{j}}^{(n)}) + \sum_{i=j+1}^{n-1} \left\{W_{2k+1}(e_{0}^{(n)} + \sum_{r=i+1}^{n-1} e_{z_{r}}^{(n)}, e_{z_{i}}^{(n)}) + W_{2k+1}(e_{z_{i}}^{(n)}, e_{0}^{(i)} - e_{z_{i}}^{(n)})\right\}, e_{z_{j}}^{(n)}]$$

with j = 0, 1, ..., n-1, where we define  $e_{z_0}^{(n)} = e_1^{(n)}$  to also specify  $[e_1^{(n)}, M_{2k+1}]$  through the j = 0 instance of the last three lines. The sums over i in the second and third line are absent when j = n-1. The Lie polynomials,  $W_{2k+1}$ , are defined by (19). A derivation of the commutators (53) will appear in [94], based on studying analytic continuations of the generating series for MPLs.

The key point for our present purposes is that the commutators (53), together with the Yang-Baxter relations, suffice to extract the motivic coaction and the single-valued map of individual MPLs from our main formulas in (48). First, the commutators (53) can be applied in our formulas to eliminate the zeta generators  $M_{2k+1}$  in favor of the braid operators  $e_i^{(j)}$ . The final step is to eliminate all braid operators  $e_i^{(m)}$  with m < n in favour of the n+1 generators  $e_i^{(n)}$  that occur in  $\mathbb{G}_n$  by means of the braid algebra (49) to (51).

**E. Relation to period matrices:** We briefly relate our results to earlier work, by mentioning an alternative way to present our formulas. Consider the product

$$\mathbb{F}_n = \mathbb{M} \, \mathbb{G}_1 \, \mathbb{G}_2 \dots \mathbb{G}_n \tag{54}$$

of the generating series  $\mathbb{G}_j$  at  $1 \leq j \leq n$ . This product  $\mathbb{F}_n$  solves the multivariable KZ equations (16). By induction on n, one can show that the main results (48) of this work can be written compactly as

$$\Delta \mathbb{F}_n^{\mathfrak{m}} = \mathbb{F}_n^{\mathfrak{m}} \, \mathbb{F}_n^{\mathfrak{dr}}, \quad \text{sv} \, \mathbb{F}_n = \overline{\mathbb{F}_n^t} \, \mathbb{F}_n$$
 (55)

In practice, these attractive formulas are not as computationally useful as the formulas (48) involving  $\mathbb{G}_n$ . This is because  $\mathbb{F}_n$  takes values in the universal enveloping algebra of all of  $M_{2k+1}$  and  $e_i^{(j)}$  with  $j \leq n$  subject to a multitude of relations. A meaningful comparison of coefficients in (55) is only possible after exhaustive use of the extended braid-algebra commutators (49) to (51) and (53) to line up the ordering of generators on the left- and right-hand sides. By contrast, the n+1 generators  $e_i^{(n)}$  in  $\mathbb{G}_n$  generate a free algebra, and there is no need to specify any ordering convention in comparing coefficients in (48).

However, the formulas (55) help to illustrate the relation of our results to earlier work. In [89], configurationspace integrals over marked points on a disk or sphere are shown to give rise to solutions to KZ equations with the same product structure as  $\mathbb{F}_n$  in (54). These solutions in the reference can be obtained from products  $\mathbb{PF}_n$  (with a generating series  $\mathbb{P}=1+\sum_{k=1}^{\infty}\zeta_2^kP_{2k}$  in even zeta values) by choosing particular matrix representations of the braid algebra, its extension by zeta generators  $M_{2k+1}$ , and the left-multiplicative  $P_{2k}$ . In fact, our main results were initially anticipated from matrix representations of (54) studied in [89]. However, in the present work, the Lie-algebra generators  $e_i^{(j)}$  and  $M_{2k+1}$  are studied without specialising to the matrix representations in the reference. This ensures that the generators satisfy only the extended braid-algebra relations (49) to (51) and (53), without obeying any additional relations that arise as an artifact of the choice of representation.

The products  $\mathbb{P} \mathbb{F}_n$  in [89] take the role of twisted period matrices for configuration spaces at genus zero with different numbers of integrated and unintegrated marked points. By the corollary  $\mathbb{P}^{\mathfrak{dr}} = 1$  of  $\zeta_{2k}^{\mathfrak{dr}} = 0$ , the period matrices  $\mathbb{PF}_n$  in the reference obey the same coaction properties as  $\mathbb{F}_n$  in (55). Hence, the specializations of our main coaction formula in (55) to matrix representations yield examples of the so-called "master formula" [25, 26] proposed for the motivic coaction of more general (twisted) period matrices. Moreover, the formula (55) for sv  $\mathbb{F}_n$  lines up with the general theory of singlevalued period matrices in [37]. When specializing to the matrix representations in [89] and taking the properties of  $\mathbb{P}$  into account, the expression for sv  $\mathbb{F}_n$  in (55) can be derived from the generalization of the genus-zero KLT relations to an arbitrary number of unintegrated marked points [51, 96], with further details to be given in [94].

# CONCLUSION AND OUTLOOK

In this work, we have provided a new formulation of the motivic coaction and single-valued map for multiple polylogarithms (MPLs) in any number of variables. Our results are based on Lie-algebra structures that intertwine braid operators with additional generators for odd Riemann zeta values. The main result (48) (which is specialised to the one- and two-variables cases in (22–24) and (39–40)) is expressed in terms of generating series that combine these two types of generators with MPLs and MZVs. The coaction and single-valued map of the individual MPLs can be read off after applying the Liebrackets of the generators that we determine explicitly from Yang–Baxter relations and the Drinfeld associator. Unlike earlier approaches, the MPLs and MZVs in our formulas are automatically fully simplified with respect to changes of fibration basis and MZV relations over  $\mathbb{Q}$ .

A longer companion paper [94] will give both heuristic derivations and proofs of the results reported here. In particular, the construction of the single-valued map from the coaction and the antipode [36, 40] will make manifest that all of these results are underpinned by the same extension of the braid algebra by zeta generators.

For the genus-zero MPLs studied in this work, the motivic coaction and the single-valued map already find a broad spectrum of applications to quantum-field-theory and string amplitudes. The coaction gives rise to smoking guns for new symmetries of physical theories under the so-called cosmic Galois group. The single-valued map in turn is a rich source of integration techniques and relations between open- and closed-string amplitudes. However, both of these intriguing structures have mostly been studied at the level of genus-zero contributions to scattering amplitudes, i.e. MPLs and MZVs. Accordingly, investigations of coactions and single-valued maps of elliptic polylogarithms and more general functions beyond MPLs encountered in amplitudes will have important implications for understanding the symmetries of and relations among fundamental interactions.

Our reformulation of coactions and single-valued maps at genus zero via zeta generators is tailored to facilitate generalizations to elliptic polylogarithms and beyond. Indeed, Brown's equivariant and single-valued iterated Eisenstein integrals [70–72] at genus one give another class of functions where zeta generators control the appearance of MZVs in non-holomorphic modular forms. More specifically, Brown's construction can be rewritten in terms of adjoint actions of generating series similar to the one sv M in this work [64, 65], where the zeta generators act on certain derivations dual to Eisenstein series [66–69], instead of on the braid operators at genus zero. The close analogy between functions of the modular parameter at genus one and MPLs in one variable at genus zero will be highlighted in upcoming work [65], and this connection calls for future investigations of multivariable cases and implications for coaction formulas.

In the long run, our unified approach to the motivic coaction and single-valued map via Lie-algebra structures including zeta generators might offer a systematic line of attack for MPLs on surfaces of arbitrary genus. It appears promising to look for similar results using these methods for more general classes of iterated integrals depending on increasing numbers of marked points and complex-structure moduli.

#### ACKNOWLEDGEMENTS

We are grateful to Ruth Britto, Francis Brown, Daniele Dorigoni, Mehregan Doroudiani, Joshua Drewitt, James Drummond, Axel Kleinschmidt, Carlos Mafra, Lionel Mason, Sebastian Mizera, Pierre Lochak, Franziska Porkert, Leila Schneps, Oliver Schnetz and Yoann Sohnle for combinations of valuable discussions and collaboration on related topics. The research of HF is supported by Merton College, Oxford, and received additional support from the European Research Council under ERC-COG-724638 GALOP. The research of MH is supported in part by the European Research Council under ERC-STG-804286 UNISCAMP and in part by the Knut and Alice Wallenberg Foundation under grant KAW 2018.0116. The research of DK was supported by the European Research Council under ERC-COG-724638 GALOP. The research of CR and OS is supported by the European Research Council under ERC-STG-804286 UNISCAMP. The research of BV was supported by the Knut and Alice Wallenberg Foundation under grant KAW2018.0162.

- A. B. Goncharov, Geometry of configurations, polylogarithms, and motivic cohomology, Advances in Mathematics 114 (1995) 197–318.
- A. B. Goncharov, Multiple polylogarithms, cyclotomy and modular complexes,
   Math. Res. Lett. 5 (1998) 497–516, [1105.2076].
- [3] E. Remiddi and J. A. M. Vermaseren, *Harmonic polylogarithms*, Int. J. Mod. Phys. A 15 (2000) 725–754, [hep-ph/9905237].
- [4] A. B. Goncharov, Multiple polylogarithms and mixed Tate motives, math/0103059.
- [5] J. Vollinga and S. Weinzierl, Numerical evaluation of multiple polylogarithms, Comput. Phys. Commun. 167 (2005) 177, [hep-ph/0410259].
- [6] C. Duhr, Mathematical aspects of scattering amplitudes, in Theoretical Advanced Study Institute in Elementary Particle Physics: Journeys Through the Precision Frontier: Amplitudes for Colliders, pp. 419–476, 2015. 1411.7538.
- [7] J. M. Henn, Lectures on differential equations for Feynman integrals, J. Phys. A 48 (2015) 153001, [1412.2296].
- [8] J. L. Bourjaily et al., Functions Beyond Multiple Polylogarithms for Precision Collider Physics, in Snowmass 2021, 3, 2022. 2203.07088.
- [9] S. Abreu, R. Britto and C. Duhr, The SAGEX review on scattering amplitudes Chapter 3: Mathematical structures in Feynman integrals, J. Phys. A 55 (2022) 443004, [2203.13014].
- [10] S. Weinzierl, Feynman Integrals: A Comprehensive Treatment for Students and Researchers. Cambridge Monographs on Mathematical Physics. Springer Cham, 2022.
- [11] N. Berkovits, E. D'Hoker, M. B. Green, H. Johansson and O. Schlotterer, *Snowmass White Paper: String*

- Perturbation Theory, in Snowmass 2021, 3, 2022. 2203.09099.
- [12] C. R. Mafra and O. Schlotterer, Tree-level amplitudes from the pure spinor superstring, Phys. Rept. 1020 (2023) 1–162, [2210.14241].
- [13] A. B. Goncharov, Galois symmetries of fundamental groupoids and noncommutative geometry, Duke Math. J. 128 (2005) 209, [math/0208144].
- [14] F. Brown, Mixed Tate motives over Z, Ann. of Math. (2) 175 (2012) 949–976
- [15] C. Duhr, Hopf algebras, coproducts and symbols: an application to Higgs boson amplitudes, JHEP 08 (2012) 043, [1203.0454].
- [16] F. Brown, Motivic periods and the projective line minus three points, in Proceedings of the ICM 2014, 2014. 1407.5165.
- [17] O. Schnetz, Graphical functions and single-valued multiple polylogarithms, Commun. Num. Theor. Phys. 08 (2014) 589–675, [1302.6445].
- [18] F. Brown, Feynman amplitudes, coaction principle, and cosmic Galois group, Commun. Num. Theor. Phys. 11 (2017) 453–556, [1512.06409].
- [19] E. Panzer and O. Schnetz, The Galois coaction on  $\phi^4$  periods, Commun. Num. Theor. Phys. 11 (2017) 657–705, [1603.04289].
- [20] M. Borinsky and O. Schnetz, Recursive computation of Feynman periods, JHEP 22 (2022) 291, [2206.10460].
- [21] O. Schnetz, The Galois coaction on the electron anomalous magnetic moment, Commun. Num. Theor. Phys. 12 (2018) 335–354, [1711.05118].
- [22] S. Caron-Huot, L. J. Dixon, F. Dulat, M. Von Hippel, A. J. McLeod and G. Papathanasiou, The Cosmic Galois Group and Extended Steinmann Relations for Planar N = 4 SYM Amplitudes, JHEP 09 (2019) 061, [1906.07116].
- [23] S. Abreu, R. Britto, C. Duhr and E. Gardi, From multiple unitarity cuts to the coproduct of Feynman integrals, JHEP 10 (2014) 125, [1401.3546].
- [24] S. Abreu, R. Britto and H. Grönqvist, Cuts and coproducts of massive triangle diagrams, JHEP 07 (2015) 111, [1504.00206].
- [25] S. Abreu, R. Britto, C. Duhr and E. Gardi, Algebraic Structure of Cut Feynman Integrals and the Diagrammatic Coaction, Phys. Rev. Lett. 119 (2017) 051601, [1703.05064].
- [26] S. Abreu, R. Britto, C. Duhr and E. Gardi, Diagrammatic Hopf algebra of cut Feynman integrals: the one-loop case, JHEP 12 (2017) 090, [1704.07931].
- [27] M. Tapušković, Motivic Galois coaction and one-loop Feynman graphs, Commun. Num. Theor. Phys. 15 (2021) 221–278, [1911.01540].
- [28] S. Abreu, R. Britto, C. Duhr, E. Gardi and J. Matthew, Generalized hypergeometric functions and intersection theory for Feynman integrals, PoS (2019) 067, [1912.03205].
- [29] O. Gürdoğan, From integrability to the Galois coaction on Feynman periods, Phys. Rev. D 103 (2021) L081703, [2011.04781].
- [30] O. Schlotterer and S. Stieberger, Motivic Multiple Zeta

- Values and Superstring Amplitudes, J. Phys. **A46** (2013) 475401, [1205.1516].
- [31] J. M. Drummond and E. Ragoucy, Superstring amplitudes and the associator, JHEP 08 (2013) 135, [1301.0794].
- [32] L. J. Dixon, O. Gurdogan, A. J. McLeod and M. Wilhelm, Folding Amplitudes into Form Factors: An Antipodal Duality, Phys. Rev. Lett. 128 (2022) 111602, [2112.06243].
- [33] L. J. Dixon, O. Gürdoğan, Y.-T. Liu, A. J. McLeod and M. Wilhelm, Antipodal Self-Duality for a Four-Particle Form Factor, Phys. Rev. Lett. 130 (2023) 111601, [2212.02410].
- [34] L. J. Dixon and Y.-T. Liu, An eight loop amplitude via antipodal duality, JHEP 09 (2023) 098, [2308.08199].
- [35] P. Cartier, La folle journée, de Grothendieck à Connes et Kontsevich. évolution des notions d'espace et de symétrie, Publications Mathématiques de l'IHÉS S88 (1998) 23-42.
- [36] F. Brown, Single-valued Motivic Periods and Multiple Zeta Values, SIGMA 2 (2014) e25, [1309.5309].
- [37] F. Brown and C. Dupont, Single-valued integration and double copy, J. Reine Angew. Math. 2021 (2021) 145–196, [1810.07682].
- [38] F. Brown, Polylogarithmes multiples uniformes en une variable, C. R. Acad. Sci. Paris Ser. I 338 (2004) 527–532.
- [39] L. J. Dixon, C. Duhr and J. Pennington, Single-valued harmonic polylogarithms and the multi-Regge limit, JHEP 10 (2012) 074, [1207.0186].
- [40] V. Del Duca, S. Druc, J. Drummond, C. Duhr, F. Dulat, R. Marzucca et al., Multi-Regge kinematics and the moduli space of Riemann spheres with marked points, JHEP 08 (2016) 152, [1606.08807].
- [41] J. Broedel, M. Sprenger and A. Torres Orjuela, Towards single-valued polylogarithms in two variables for the seven-point remainder function in multi-Regge-kinematics, Nucl. Phys. B 915 (2017) 394–413, [1606.08411].
- [42] V. Del Duca, S. Druc, J. M. Drummond, C. Duhr, F. Dulat, R. Marzucca et al., All-order amplitudes at any multiplicity in the multi-Regge limit, Phys. Rev. Lett. 124 (2020) 161602, [1912.00188].
- [43] V. Del Duca, L. J. Dixon, C. Duhr and J. Pennington, The BFKL equation, Mueller-Navelet jets and single-valued harmonic polylogarithms, JHEP 02 (2014) 086, [1309.6647].
- [44] V. Del Duca, C. Duhr, R. Marzucca and B. Verbeek, The analytic structure and the transcendental weight of the BFKL ladder at NLL accuracy, JHEP 10 (2017) 001, [1705.10163].
- [45] J. M. Drummond, Generalised ladders and single-valued polylogarithms, JHEP 02 (2013) 092, [1207.3824].
- [46] F. Brown and O. Schnetz, Single-valued multiple polylogarithms and a proof of the zig-zag conjecture, Journal of Number Theory 148 (2015) 478–506.
- [47] C. Duhr and F. Porkert, Feynman integrals in two dimensions and single-valued hypergeometric functions, 2309.12772.
- [48] S. Stieberger, Closed superstring amplitudes, single-valued multiple zeta values and the Deligne associator, J. Phys. A47 (2014) 155401, [1310.3259].
- [49] S. Stieberger and T. R. Taylor, Closed String

- Amplitudes as Single-Valued Open String Amplitudes, Nucl. Phys. **B881** (2014) 269–287, [1401.1218].
- [50] O. Schlotterer and O. Schnetz, Closed strings as single-valued open strings: A genus-zero derivation, J. Phys. A52 (2019) 045401, [1808.00713].
- [51] P. Vanhove and F. Zerbini, Single-valued hyperlogarithms, correlation functions and closed string amplitudes, Adv. Theor. Math. Phys. 26 (2022) 455–530, [1812.03018].
- [52] F. Brown and C. Dupont, Single-valued integration and superstring amplitudes in genus zero, Commun. Math. Phys. 382 (2021) 815–874, [1910.01107].
- [53] E. D'Hoker, M. B. Green, Ö. Gürdogan and P. Vanhove, *Modular graph functions*, *Commun. Num. Theor. Phys.* **11** (2017) 165–218, [1512.06779].
- [54] J. Broedel, O. Schlotterer and F. Zerbini, From elliptic multiple zeta values to modular graph functions: open and closed strings at one loop, JHEP 01 (2019) 155, [1803.00527].
- [55] J. E. Gerken, A. Kleinschmidt and O. Schlotterer, Heterotic-string amplitudes at one loop: modular graph forms and relations to open strings, JHEP 01 (2019) 052, [1811.02548].
- [56] D. Zagier and F. Zerbini, Genus-zero and genus-one string amplitudes and special multiple zeta values, Commun. Num. Theor. Phys. 14 (2020) 413–452, [1906.12339].
- [57] J. E. Gerken, A. Kleinschmidt, C. R. Mafra, O. Schlotterer and B. Verbeek, Towards closed strings as single-valued open strings at genus one, J. Phys. A 55 (2022) 025401, [2010.10558].
- [58] L. F. Alday, T. Hansen and J. A. Silva, AdS Virasoro-Shapiro from single-valued periods, JHEP 12 (2022) 010, [2209.06223].
- [59] L. F. Alday, T. Hansen and J. A. Silva, Emergent Worldsheet for the AdS Virasoro-Shapiro Amplitude, Phys. Rev. Lett. 131 (2023) 161603, [2305.03593].
- [60] L. F. Alday and T. Hansen, The AdS Virasoro-Shapiro amplitude, JHEP 10 (2023) 023, [2306.12786].
- [61] G. Fardelli, T. Hansen and J. A. Silva, AdS Virasoro-Shapiro amplitude with KK modes, JHEP 11 (2023) 064, [2308.03683].
- [62] P. Deligne and A. B. Goncharov, Groupes fondamentaux motiviques de Tate mixte, Ann. Sci. École Norm. Sup. (4) 38 (2005) 1–56, [math/0302267].
- [63] F. Brown, Zeta elements in depth 3 and the fundamental Lie algebra of the infinitesimal Tate curve, Forum Math. Sigma 5 (2017) Paper No. e1, 56, [1504.04737].
- [64] D. Dorigoni, M. Doroudiani, J. Drewitt, M. Hidding, A. Kleinschmidt, N. Matthes et al., Modular graph forms from equivariant iterated Eisenstein integrals, JHEP 12 (2022) 162, [2209.06772].
- [65] D. Dorigoni, M. Doroudiani, J. Drewitt, M. Hidding, A. Kleinschmidt, O. Schlotterer et al., Non-holomorphic modular forms from zeta generators, 2023.
- [66] Y. Ihara, Braids, Galois groups, and some arithmetic functions, in Proceedings of the International Congress of Mathematicians, Vol. I, II (Kyoto, 1990),

- pp. 99-120, Math. Soc. Japan, Tokyo, 1991.
- [67] Y. Ihara and N. Takao, Seminar talk, 1993.
- [68] H. Tsunogai, On some derivations of Lie algebras related to Galois representations, Publ. Res. Inst. Math. Sci. 31 (1995) 113-134.
- [69] A. Pollack, "Relations between derivations arising from modular forms." https://dukespace.lib.duke.edu/dspace/handle/10161/1281, M. Jimbo, Introduction to the Yang-Baxter equation,
- [70] F. Brown, Multiple modular values and the relative completion of the fundamental group of  $\mathcal{M}_{1,1}$ , 1407.5167.
- [71] F. Brown, A class of non-holomorphic modular forms I, Res. Math. Sci. 5 (2018) 5:7, [1707.01230].
- [72] F. Brown, A class of non-holomorphic modular forms II: equivariant iterated Eisenstein integrals, Forum of Mathematics, Sigma 8 (2020) 1, [1708.03354].
- [73] J. Broedel, C. Duhr, F. Dulat, B. Penante and L. Tancredi, Elliptic symbol calculus: from elliptic polylogarithms to iterated integrals of Eisenstein series, JHEP 08 (2018) 014, [1803.10256].
- [74] M. Wilhelm and C. Zhang, Symbology for elliptic multiple polylogarithms and the symbol prime, JHEP **01** (2023) 089, [2206.08378].
- [75] A. Forum and M. von Hippel, A Symbol and Coaction for Higher-Loop Sunrise Integrals, SciPost Phys. 6 (2023) 050, [2209.03922].
- [76] M. Tapušković, The cosmic Galois group, the sunrise Feynman integral, and the relative completion of  $\Gamma_1(6)$ , 2303.17534.
- [77] E. Panzer, Feynman integrals and hyperlogarithms. PhD thesis, Humboldt U., 2015. 1506.07243.
- [78] F. Brown, On the decomposition of motivic multiple zeta values, in Galois-Teichmüller theory and arithmetic geometry, vol. 63 of Adv. Stud. Pure Math., pp. 31–58. Math. Soc. Japan, Tokyo, 2012. 1102.1310.
- [79] F. Brown, Notes on motivic periods, Communications in Number Theory and Physics 11 (2015) 557-655, [1512.06410]
- [80] D. Zagier, Values of Zeta Functions and Their Applications, pp. 497–512. Birkhäuser Basel, Basel,
- [81] J. Blümlein, D. J. Broadhurst and J. A. M. Vermaseren, The Multiple Zeta Value Data Mine, Comput. Phys. Commun. 181 (2010) 582–625,

- [0907.2557].
- [82] O. Schnetz, Generalized single-valued hyperlogarithms, 2111.11246.
- [83] M. Borinsky and O. Schnetz, Graphical functions in even dimensions, Commun. Num. Theor. Phys. 16 (2022) 515-614, [2105.05015].
- International Journal of Modern Physics A 4 (1989) 3759-3777.
- [85] C. Kassel, Quantum groups, vol. 155. Springer Science & Business Media, 2012.
- [86] V. Drinfeld, Quasi Hopf algebras, Leningrad Math. J. 1 (1989) 1419–1457.
- [87] V. Drinfeld, On quasitriangular quasi-Hopf algebras and on a group that is closely connected with  $Gal(\overline{\mathbb{Q}}/\mathbb{Q})$ , Leningrad Math. J. 2 (4) (1991) 829–860.
- [88] T. Le and J. Murakami, Kontsevich's integral for the Kauffman polynomial, Nagoya Math J. 142 (1996) 93-65.
- [89] R. Britto, S. Mizera, C. Rodriguez and O. Schlotterer, Coaction and double-copy properties of configuration-space integrals at genus zero, JHEP 05 (2021) 053, [2102.06206].
- [90] H. Furusho, The multiple zeta value algebra and the stable derivation algebra, Publications of The Research Institute for Mathematical Sciences 39 (2000) 695–720. [math/0011261].
- [91] Y. Ihara, On the stable derivation algebra associated with some braid groups, Israel Journal of Mathematics **80** (1992) 35–153.
- [92] Y. Ihara, The Galois representation arising from  $\mathbb{P}^1 - \{0, 1, \infty\}$  and Tate twists of even degree, 1989.
- [93] F. Brown and C. Dupont, Lauricella hypergeometric functions, unipotent fundamental groups of the punctured Riemann sphere, and their motivic coactions, Nagoya Math. J. **249** (2023) 148–220, [1907.06603].
- [94] H. Frost, M. Hidding, D. Kamlesh, C. Rodriguez, O. Schlotterer and B. Verbeek, work in progress, 2024.
- [95] P. P. Deligne, Le groupe fondamental de la droite projective moins trois points, in Galois Groups over  $\mathbb{Q}$ , (New York, NY), pp. 79-297, Springer US, 1989.
- [96] H. Kawai, D. C. Lewellen and S. H. H. Tye, A Relation Between Tree Amplitudes of Closed and Open Strings, Nucl. Phys. **B269** (1986) 1–23.