

# Random Forest for Dynamic Risk Prediction of Recurrent Events: A Pseudo-Observation Approach

ABIGAIL LOE<sup>\*,1</sup>, SUSAN MURRAY<sup>1</sup>, ZHENKE WU<sup>1,2</sup>

<sup>1</sup>*Department of Biostatistics, University of Michigan, Ann Arbor, MI 48109, USA*

<sup>2</sup>*Michigan Institute for Data Science, University of Michigan, Ann Arbor, MI 48109, USA*

\*agloe@umich.edu

## SUMMARY

Recurrent events are common in clinical, healthcare, social and behavioral studies. To overcome some long-standing challenges in analyzing censored recurrent event data, a recent regression analysis framework constructs a censored longitudinal data set consisting of times to the first recurrent event in multiple pre-specified follow-up windows of length  $\tau$  (Xia, Murray and Tayob, 2020, XMT). Traditional regression models struggle with nonlinear and multi-way interactions, with success depending on the skill of the statistical programmer. With a staggering number of potential predictors being generated from genetic, -omic, and electronic health records sources, machine learning approaches such as the random forest regression are growing in popularity, as they can nonparametrically incorporate information from many predictors with nonlinear and multi-way interactions involved in prediction. In this paper, we (1) develop a random forest approach for dynamically predicting probabilities of remaining event-free during a subsequent  $\tau$ -duration follow-up period from a reconstructed censored longitudinal data set, (2) modify the XMT regression approach to predict these same probabilities, subject to the limitations that traditional regression models typically have, and (3) demonstrate how to incorporate patient-specific history of recurrent events for prediction in settings where this information may be partially missing. We show the increased ability of our random forest algorithm for predicting the probability of remaining event-free over a  $\tau$ -duration follow-up period when compared to our modified XMT method for prediction in settings where association between predictors and recurrent event outcomes is complex in nature. We also show the importance of incorporating past recurrent event history in prediction algorithms when event times are correlated within a subject. The proposed random forest algorithm is demonstrated using recurrent exacerbation data from the Azithromycin for the Prevention of Exacerbations of Chronic Obstructive Pulmonary Disease (Albert *and others*, 2011).

\*To whom correspondence should be addressed.

*Key words:* Censored data; Longitudinal data; Pseudo-observations; Random forest; Recurrent events

## 1. INTRODUCTION

Recurrent events are common in healthcare settings, such as exacerbations of pulmonary diseases, or bleeding events for chemotherapy patients. Clinicians are often interested in predicting the probability of remaining event-free until a patient's next check-up. Information from events over time helps improve precision of predictions relative to analyses of a single time-to-event. Prediction is also greatly enhanced by the large number of patient-specific predictors that are collected now, for instance, electronic health records data, gene expression levels, metabolomic data, etc. The very large dimensions of the predictors and the multivariate nature of recurrent event outcomes requires careful thought for analysis, particularly when censoring of recurrent events is a feature of the data. A method that can handle both highly correlated data and recurrent events would be of interest to modelers and practitioners.

The majority of literature in recurrent event analysis has been parametric or semi-parametric in nature, with specific modeled relationships between predictors and recurrent event outcomes. Models have been designed for the analysis of recurrent events, gap times, and times-to-first-event across different follow-up windows, each with different assumptions imposed by the modeling paradigms (e.g., Andersen and Gill, 1982; Miloslavsky *and others*, 2004; Ang *and others*, 2009; Mao and Lin, 2016; Hengelbrock *and others*, 2016; Kong *and others*, 2016; Chan and Wang, 2017*a*; Liang *and others*, 2017; Xu *and others*, 2017; Choi *and others*, 2017; Chan and Wang, 2017*b*; Xia, Murray and Tayob, 2020; Safari *and others*, 2023). Such models have interpretable parameter estimates that can inform medical literature, generate hypotheses, and to some degree, predict outcomes. Disadvantages include limits to estimating these same parameters when there are many predictors or co-linearity between important predictors. In addition, methods that incorrectly assume independence of gap times between recurrent events measured on an individual give biased results due to induced dependent censoring. A few of these papers use pseudo-observation approaches to model recurrent event data (Yokota and Matsuyama, 2019; Xia, Murray and Tayob, 2020), an attractive approach that we also consider.

As to the goal of modeling many potentially collinear predictors, perhaps with non-linear associations with recurrent event outcomes, we are drawn to the nonparametric nature of random forest algorithms.

Machine learning methods, and in particular random forest algorithms, have gained widespread acceptance as a method of building predictions based on complex relationships of biomarkers (first proposed by Breiman *and others* (1984), applied more recently in statistical literature such as Ishwaran *and others* (2004), Wager and Athey (2018), Zhao *and others* (2020), Gerber *and others* (2021a), Gerber *and others* (2021b), and reviewed in Hu and Szymczak (2023)). These algorithms naturally accommodate non-canonical relationships between covariates and outcomes, an attractive feature in an age of increasingly high-dimensional data where a traditional modeler cannot realistically find all the interactions, higher-order covariates and non-linear patterns that may be predictive of an outcome. By using information across different sets of covariates within each of the generated trees of a random forest, a modeler can side-step several traditional pitfalls of parametric and semi-parametric models: (1) multicollinearity of predictors and (2) limited ability to include a large number of covariates. In particular, Zhao *and others* (2020) showed that for a single event time, imposing a censored longitudinal data structure introduced by Tayob and Murray (2015) and incorporating ideas from pseudo-observation literature, random forest regression can address nonlinear relationships and interactions between covariates and outcomes for dynamic prediction of event-free periods.

In our manuscript we incorporate ideas from Xia, Murray and Tayob (2020), who propose a censored longitudinal data structure for the analysis of recurrent event data. The Xia, Murray and Tayob (XMT) model estimates restricted mean time-to-next-recurrent-event based on clinical characteristics. As part of this manuscript, we will develop a modified version of the XMT model that instead estimates event-free probabilities over  $\tau$ -duration follow-up windows. This more standard regression approach to prediction will be compared to a novel random forest dynamic prediction model for potentially censored recurrent event data that we introduce in this manuscript and recommend when prediction is the primary goal of analysis and when many covariates are available. For both of these modeling approaches, we explore the importance of patient-specific historical event-rate as a predictor. To date, we are not aware of any methods for dynamic prediction applicable to censored recurrent event data.

The rest of this paper is organized as follows. In Section 2, we describe how to transform recurrent event data to a censored longitudinal data framework for analysis using our proposed methods. In Section 3, we review a random forest algorithm applicable to uncensored longitudinal data. In Section 4, we describe how to construct pseudo-observations based on the censored longitudinal data structure, described in Section

2, that can be used with the random forest algorithm developed in Section 3. In Section 5, we compare via simulation our modified XMT regression approach versus our proposed random forest approach for the analysis of censored recurrent event data, and evaluate different methods for including historical event rate information in these analyses. We apply both the modified XMT and our recurrent event random forest algorithm to data from Azithromycin for the Prevention of Exacerbation of COPD (Albert *and others*, 2011) in Section 6. We conclude with a discussion in Section 7.

## 2. NOTATION AND CENSORED LONGITUDINAL DATA STRUCTURE

For individual  $i$ , denote recurrent event times  $T_{i,1} < T_{i,2} < \dots$ , that are potentially censored by the random variable  $C_i$ , where  $C_i$  is independent of  $T_{i,j}$ ,  $i = 1, \dots, n$ , and  $j \geq 1$ . Observed data for each individual is  $X_{i,j}^* = \min(T_{i,j}, C_i)$ ,  $i = 1, \dots, n$ , and  $j \geq 1$ . These event times are sometimes converted to gap times between recurrent events  $G_{i,j} = T_{i,j+1} - T_{i,j}$ ,  $i = 1, 2, \dots, n$ ,  $j \geq 1$ . A well-known challenge in the analysis of correlated censored gap times is the dependence between gap time random variables,  $G_{i,j}$ , and corresponding censoring times,  $C_i - G_{i,j-1}$  (Lin *and others*, 2000).

One approach to avoiding dependent censoring in analyses of recurrent events was proposed by Tayob and Murray (2015), who transformed correlated recurrent event times into a censored longitudinal data structure. First a series of regularly spaced follow-up window start times are spaced every  $a$  units apart, so that follow-up window start times are  $\mathcal{T} = \{t_0, t_1, t_2, \dots\}$ , with  $t_j - t_{j-1} = a$ ,  $j = 1, 2, \dots$ . Briefly,  $\mathcal{T}$  can include window start times that are as close as  $a = 1$  day apart, if desired. Computational speed of the analysis is usually the limiting factor in choosing  $a$ . Xia and Murray (2019) found that  $a$  equal to one-third of the mean gap time in the control group resulted in capturing 90% of the recurrent events from the original data set in at least one follow-up window. In practice, more frequent follow-up window start times have the potential for more precise estimation, but with diminishing returns once all recurrent events have been captured in at least one follow up window.

In this paper, at each of the pre-specified ‘‘check-in time points’’ in  $\mathcal{T}$ , we will focus on estimating the probability of remaining recurrent-event-free over a subsequent follow-up window of length  $\tau$ , where  $\tau$  is user-specified and ideally informed by clinical expertise. Although the check-in time points are shared across subjects by design, potential censoring may result in different numbers of effective check-ins; we

therefore denote the subject-specific sets of check-in time points by  $\{\mathcal{T}_i\}_{i=1}^n$ , where  $\mathcal{T}_i = \{t_0, t_1, \dots, t_{m_i}\}$ ,  $t_{m_i} < \min\{C_i, D - \tau\}$ , and  $D$  is the time of administrative censoring, e.g., indicating the end of study.

We follow Tayob and Murray (2015) and construct the censored longitudinal data that is essential for creating pseudo-observations in Section 4. We repeat the following two steps at all the check-in time points  $t \in \mathcal{T}_i$ ,  $i = 1, \dots, n$ :

- i) Identify the first recurrent event occurred after  $t$  and let  $\eta_i(t) = \min\{j : X_{i,j}^* \geq t, j = 1, \dots, n_i\}$  be its index among all the observed recurrent events for subject  $i$ ; also define the residual recurrent-event-free time by  $T_i(t) = T_{i,\eta_i(t)} - t$  with the corresponding censoring random variable  $C_i(t) = C_i - t$ , which remains independent of  $T_i(t)$ ;
- ii) Construct the observed censored longitudinal data set at time  $t$  by  $\{X_i(t) = X_{i,\eta_i(t)}^* - t, \delta_i(t) = I(X_i(t) \leq C_i(t)) : i = 1, 2, \dots, n\}$ .

Our prediction target can now be formally expressed as  $P(T_i(t) \geq \tau | Z_i(t))$ , i.e., at a pre-specified time point  $t$ , the probability of remaining recurrent-event-free over a subsequent follow-up period of length  $\tau$  given covariates  $Z_i(t)$ . In the following, we will occasionally suppress the subscript  $i$  and write  $P(T(t) \geq \tau | Z(t))$  for ease of presentation.

Figure 1 uses three hypothetical subjects to illustrate the above three steps of converting traditional recurrent event data  $\{X_{i,j}^*, \delta_{i,j}\}$  into the censored longitudinal data format  $\{X_i(t), \delta_i(t) : t \in \mathcal{T}_i\}$ . In Section 4, we will use the constructed censored longitudinal data to create longitudinal pseudo-observations that will be inputs of a random forest algorithm, which we now review.

### 3. RANDOM FOREST FOR UNCENSORED LONGITUDINAL OUTCOMES: A REVIEW

Regression trees are a popular approach to incorporating multi-way interactions among predictors by finding groups of observations that are similar. A tree is grown in a few steps where at each step a new branch sorts the data leftover from the preceding step into bins based on one of the predictors. The sequential branching slices the space of predictors into rectangular partitions and approximates the true outcome-predictor relationship with the average outcome within each partition. Therefore, to grow a tree is to find bins that best discriminate among the outcomes. The specific predictor and the value split at each branching are chosen to minimize prediction error.




Fig. 1: Visualization of recurrent events in both traditional format (black) and censored longitudinal format (blue). Start times of follow-up windows ( $\mathcal{T} = \{0, 60, 120, 180\}$  days, with  $a = 60$ ) are highlighted by vertical dashed lines. Traditional observed recurrent event data for individuals  $i = 1, 2, 3$  are  $\{\{X_{1,1}^* = 80, \delta_{1,1} = 1\}, \{X_{2,1}^* = 125, \delta_{2,1} = 0\}, \{X_{3,1}^* = 48, \delta_{1,1} = 1\}, \{X_{3,2}^* = 62, \delta_{1,2} = 1\}, \{X_{3,3}^* = 75, \delta_{1,3} = 1\}, \{X_{3,4}^* = 147, \delta_{1,4} = 1\}, \{X_{3,5}^* = 240, \delta_{1,5} = 0\}\}$ . For these same individuals, the observed censored longitudinal data visualized here is  $\{\{X_1(0) = 80, \delta_1(0) = 1\}, \{X_1(60) = 20, \delta_1(60) = 1\}, \{X_1(120) = 83, \delta_1(120) = 1\}, \{X_1(180) = 23, \delta_1(180) = 1\}, \{X_2(0) = 125, \delta_2(0) = 0\}, \{X_2(60) = 65, \delta_2(60) = 0\}, \{X_2(120) = 5, \delta_2(120) = 0\}, \{X_3(0) = 48, \delta_3(0) = 1\}, \{X_3(60) = 2, \delta_3(60) = 1\}, \{X_3(120) = 27, \delta_3(120) = 1\}, \{X_3(180) = 60, \delta_3(180) = 0\}\}$ . All participants were administratively censored after 8 months (240 days). Participant 2 has no recorded events before  $C_2$ . Additionally, while  $X_{33}^*$  is technically observed, it is not included in the censored longitudinal data set because it is the second observation in the window, and there is no  $t$  such that  $X_i(t) = X_{33}^*$ . Besides potential correlation between separate event times seen in the same individual, this data structure induces correlation due to the same event contributing to the final censored longitudinal values in more than one follow-up window. For example,  $X_{12}^*$  produces two measures in the longitudinal data structure,  $X_1(120) = 83$  and  $X_1(180) = 23$ .

The number of possible trees is combinatorically large and precludes efficient global optimization (the number of binary trees with  $K$  leaves is  $(2K - 3)!!$ ). Greedy algorithms have been developed to approximate the optimal global tree by myopically optimizing prediction error at the start of each branch. We will focus on binary trees in this paper for their popularity and effectiveness. The loss associated with the prediction error for a branch is often termed “impurity” which measures how similarly the observations behave on either side of the binary split. The branching procedure halts, for example, when the number of observations in a terminal node or the number of terminal nodes reaches respective thresholds. Advantages of a regression tree include invariance to monotonic transformation of predictors, flexible approximation to potentially severe nonlinearities, and the capacity to approximate  $L - 1$  way interactions for a tree with depth  $L$ .

To overcome potential overfitting, ensemble methods can be used to combine predictions from many trees into a single prediction. Random forest is such an ensemble regularizer based on Breiman’s bootstrap aggregation (Breiman, 2001), or “bagging,” which averages over multiple predictions obtained from trees grown from multiple bootstrap samples hence stabilizing the overall prediction. Random forest is a variation of bagging designed to further reduce the correlation among trees grown using different bootstrap samples by “dropout” which considers only a randomly drawn subset of predictors for splitting at each potential branch. Such a strategy ensures that early branches for some trees will not always split on predictors that offer the most gain in prediction accuracy. This reduces the correlation among predictions by multiple trees to further improve the variance reduction relative to the standard bagging along with reduced computational cost.

Among the many variants of random forest, we are particularly interested in random forests for longitudinal data, as recurrent event analysis uses data obtained from subjects over time. Adler, Potapov and Lausen (2011) proposed an algorithm for tree-based ensemble methods that takes into consideration the dependence structure inherent in longitudinal data analysis. We provide a primer on this specific variation of random forest.

Let  $(Y_i(t), \mathbf{Z}_i(t))$  be a pair of a continuous outcome and a set of  $p$ -dimensional covariates, measured at time  $t \in \mathcal{T}_i$  for  $i = 1, 2, \dots, n$ ; here we use  $\mathcal{T}_i$ ’s introduced in Section 2 with shared measurement timings to align more closely with our construction in Section 4, although the algorithm presented herein is applicable to scenarios with subject-specific measurement timings. In Section 4, we will substitute  $Y_i(t)$  here with a longitudinal outcome  $(\widehat{S}_i^T(t))$  that is constructed based on pseudo-observation technique to overcome censoring. Also note that the  $\mathbf{Z}_i(t)$  may include time  $t$ . The following algorithm assumes  $Y_i(t) = g\{\mathbf{Z}_i(t)\} + \epsilon_i(t)$  with  $\mathbb{E}[\epsilon_i(t)] = 0$  and is designed to estimate  $g(\cdot)$  which describes a potentially complex relationship between the outcome and the predictors.

For  $b = 1, \dots, B$ , we repeat Step 1 and 2 below ( $B = 500$  is used in this manuscript):

Step 1 (Bootstrap): Generate a bootstrap sample  $\{(Y_i(t), \mathbf{Z}_i(t)), (i, t) \in \text{Bootstrap}(b)\}$  from the original data set, where  $\text{Bootstrap}(b)$  is the  $b$ -th bootstrap sample;  $\text{Bootstrap}(b)$  is obtained by first sampling  $n$  subjects with replacement, followed by selecting all check-in time points  $t \in \{t_0, t_1, \dots, t_{m_i}\}$  for each bootstrapped subject  $i$  (Adler, Potapov and Lausen, 2011). The data corresponding to individuals in  $\text{Bootstrap}(b)$  is called “bagged”, and data not in the sample is called “out-of-bag.”

Step 2 (Grow a tree): Initialize the tree stump by grouping all observations together.

- 2a. (Form split variables) At a potential branching, randomly select  $m$  predictors from the  $p$  predictors with  $m \ll p$ ; in this paper we use the default choice of  $m = \sqrt{p}$  which works well in our simulation and data analysis. Let the randomly selected predictors be  $\{Z_{i,s_1}(t), \dots, Z_{i,s_m}(t)\}$  at a particular potential branching for bagged observation  $(i, t)$  where  $(s_1, \dots, s_m)$  are the indices for the subset of  $m$  predictors. For ease of presentation, we denote it by  $\mathbf{W} = (W_1, \dots, W_m)$ . For a continuous or categorical observed covariate  $W_k$ , construct a grid of all observed values of the covariate,  $\widetilde{W}_k = \{w_{kl}, l = 1, \dots, L_k\}$ . Consider all possible thresholds in  $\widetilde{W}_k$  and their corresponding indicators,  $\{I(Z_{i,s_k}(t) < w_{kl}), (i, t) \in \text{Bootstrap}(b), w_{kl} \in \widetilde{W}_k\}$  for subject  $i$  at check-in time  $t$  in the bootstrap sample  $\text{Bootstrap}(b)$ . These derived indicator functions for continuous and categorical covariates, along with covariates in  $\mathbf{W}$  that were already binary, are “potential split variables”.
- 2b. (Choose branching): Choosing a covariate among  $\mathbf{W}$  and the split value  $w_{kl}$  will result in two daughter nodes,  $\{N_{w_{kl},1}, N_{w_{kl},0}\}$ . Criteria for selecting the best variable and split vary by outcome type and metric of prediction accuracy. For continuous outcomes, the split with minimal sum of squared errors (SSE), is selected, though for other responses types, loss functions such as the Gini Index may be selected. For any particular node split under consideration using covariate  $w$ , that partitions individuals into two groups ( $\{N_{w,0}, N_{w,1}\}$ ) of sizes  $n_{w,0}$  and  $n_{w,1}$ , we minimize the criterion  $SSE(w) = \sum_{c=0,1} \sum_{q=1}^{n_{w,c}} (Y_q - \bar{Y}_c)^2$ , where  $\bar{Y}_c = \frac{1}{n_{w,c}} \sum_{q=1}^{n_{w,c}} Y_{q,c} = 0, 1$ .
- 2c. (Stopping criteria and prediction from the grown tree) Repeat Step 2a and 2b until one of the following hyperparameters has been met: minimum node size ( $\min\{n_{w,c}, c = 0, 1\}$ ) is set to 40 in this paper) is reached, or  $SSE(w)$  does not decrease any further with more branches. This results in a single binary decision tree. For each terminal node (“leaf”) of the tree  $C$ , predict  $Y_i(t)$  by the average of the outcome for observations in leaf  $C$ . Let the prediction be  $\widehat{Y}_b(\mathbf{Z}_i(t); \widehat{\boldsymbol{\theta}}_b)$ , where  $\widehat{\boldsymbol{\theta}}_b$  is the collection of terminal node predictions in the  $b$ -th tree.

Step 3 (Ensemble prediction by random forest): The final prediction for an individual  $i'$  with covariates  $\mathbf{Z}_{i'}(t)$  is given by the average of all the outputs from  $B$  binary decision trees above:  $\widehat{Y}_{i'}(t) = \frac{1}{B} \sum_{b=1}^B \widehat{Y}_b(\mathbf{Z}_{i'}(t); \widehat{\boldsymbol{\theta}}_b)$ .



REMARK 3.1 The two-stage bootstrapping scheme has a few statistical and computational advantages consistent with results from Adler, Potapov and Lausen (2011): (1) subjects with many measurements are as likely to be included in the bootstrap sample as subjects with few measurements, making it unlikely that observations from subjects with longer follow-up time overpower subjects with shorter follow-up times in the tree-growing stage and (2) out-of-bag data for each tree remains independent of the bagged data, making predictions from such a forest remarkably simple.

Permutation tests are applied to the out-of-bag sample to generate  $p$ -values assessing statistical significance for predictors used within the random forest algorithm while controlling for the remaining predictors. Each variable,  $W$ , is selected and the values of  $W$  in the out-of-bag sample are permuted  $d$  times, to obtain  $\mathcal{W}_s$ ,  $s = 1, 2, \dots, d$ . This breaks their association with  $Y(t)$ . The out-of-bag samples corresponding to both  $\mathcal{W}_s$  and  $W$  are then passed through the forest to obtain new predictions,  $\mathcal{P}_{\mathcal{W}_s}$  and  $\mathcal{P}_W$  respectively. Exploiting the fact that  $p_{w,\text{test}} = \frac{\frac{1}{d} \sum_{s=1}^d (\mathcal{P}_W - \mathcal{P}_{\mathcal{W}_s})}{\sigma_{\mathcal{W},W}}$  has an asymptotically standard normal distribution, one can then obtain a Wald-type test of variable significance ( $\sigma_{\mathcal{W},W}$  is estimated from the permutation distribution of prediction errors).

Random forests for survival data can be evaluated using Harrell’s C-statistic (also called “C-index”), a non-parametric measure of concordance (Harrell *and others*, 1982). Briefly, the C-statistic compares the predicted risk scores or fit values for any two observations with the actual time-to-event for those two observations. The proportion of observations that are correctly ordered is reported as the C-index. Any model which perfectly orders patients based on some risk score has a C-index of 1, while a model with no predictive power has C-index equal to 0.50.

In the next section we review pseudo-observation methodology that can convert the censored longitudinal data developed in Section 2 into a version that can be used with the above algorithm.

#### 4. CONSTRUCTION OF PSEUDO-OBSERVATIONS

Pseudo-observations are a variation of jack-knife methodology applicable to censored time-to-event outcomes (Andersen, Klein and Rosthøj, 2003). This technical device has been used successfully for a single time-to-event and recurrent events in manuscripts such as: Andrei and Murray (2007); Hengelbrock *and others* (2016); Xia, Murray and Tayob (2020); Zhao *and others* (2020). Essentially a pseudo-observation of a cen-

sored outcome, once created, can be used in regression contexts as if it was an uncensored outcome. In this section, we will first describe the mechanics of how to create a pseudo-observation for each potentially censored time-to-first event in a follow-up window of length  $\tau$  starting at  $t$ . We will then briefly describe properties of pseudo-observations in the context of the random forest described in Section 3.

In this manuscript, our parameter of interest is survival probability, but we recognize that pseudo-observation is a technique that can be applied to a wide variety of other targets, including  $\tau$ -restricted mean survival time and hazard estimation. To construct pseudo-observations to be used with our random forest algorithm, we return to the censored longitudinal data structure described in Section 2. We calculate two estimates of the probability of recurrent-event-free in a follow-up window of length  $\tau$  beginning at check-in time  $t$ :  $\widehat{P}(X_i(t) \geq \tau)$  and  $\widehat{P}^{(-i)}(X_i(t) \geq \tau)$  via the Kaplan-Meier estimator without conditioning on any covariate information for an individual  $i = 1, 2, \dots, n(t)$  at risk at time  $t$ , where  $\widehat{P}^{(-i)}(X_i(t) \geq \tau)$  is estimated excluding individual  $i$ . The pseudo-observation for individual  $i$  at time  $t$  is defined by:

$$\widehat{S}_i^\tau(t) = n(t)\widehat{P}(X_i(t) \geq \tau) - (n(t) - 1)\widehat{P}^{(-i)}(X_i(t) \geq \tau).$$

To gain intuition behind the use of pseudo-observations, consider the simplest case where there is no censoring. Then, algebraically,  $\widehat{P}(T_i(t) \geq \tau) = \frac{\sum_{j=1}^{n(t)} I(T_j(t) \geq \tau)}{n(t)}$  and  $\widehat{P}^{(-i)}(T_i(t) \geq \tau) = \frac{\sum_{j \neq i, j=1}^{n(t)} I(T_j(t) \geq \tau)}{n(t)-1}$ . Here  $\widehat{S}_i^\tau(t)$  reduces to  $I(T_i(t) \geq \tau)$ , so that  $E[\widehat{S}_i^\tau(t) \mid \mathbf{Z}_i(t)] = E[I(T_i(t) \geq \tau) \mid \mathbf{Z}_i(t)] = P(T_i(t) \geq \tau \mid \mathbf{Z}_i(t))$ . This property is approximately held when we use Kaplan-Meier estimates for  $\widehat{P}(T_i(t) \geq \tau)$  and  $\widehat{P}^{(-i)}(T_i(t) \geq \tau)$  in the independently right-censored case. Andersen, Klein and Rosthøj (2003) argued that this property enables consistent estimation by directly regressing the pseudo-observations upon covariates. When pseudo-observations of probability of time-to-first-recurrent-events are applied to random forests in Section 3 for prediction, we call the algorithm “RFRE.PO” (short for “Random Forest for Recurrent Events based on Pseudo-Observations”); we depict the analysis pipeline in Figure 2.

## 5. SIMULATION STUDY

As mentioned earlier in Section 1, random forest algorithms are known for nonparametrically capturing complex relationships between a large number of predictors and outcomes of interest, where these complex relationships might involve higher-order interactions, nonlinearity, and colinearity between predictors. In this section, we demonstrate the ability of RFRE.PO to predict  $P(T_i(t) \geq \tau \mid \mathbf{Z}_i(t))$  in a scenario where the true




Fig. 2: A flowchart depicting the analysis pipeline of RFRE.PO. Labels above arrows correspond to the section which describes the corresponding step in greater detail. An analyst begins with data recorded as recurrent event times before constructing an appropriate  $\mathcal{T}$ , and transforming data into the censored longitudinal framework. Upon creating such a structure, the analyst can select the preferred quantity for calculating pseudo-observations, which are ultimately fed to a historical random forest.

relationship between  $Z_i(t)$  and  $I(T_i(t) \geq \tau)$  is many times more complicated than a modeler who tends to depend on main effect terms would anticipate. For comparison, we also describe results from using a perfectly specified Xia, Murray and Tayob (XMT) model (Xia, Murray and Tayob, 2020) (Model A); and the XMT model that only uses main effect terms for the  $Z_i(t)$  in its regression model (Model B). Note that in Xia, Murray and Tayob (2020), the XMT model was for modeling the restricted mean rather than  $\hat{S}_i^\tau(t)$ ; however, their methods can be extended to our case. We did not include highly colinear predictors that would cause modified XMT model convergence issues in our simulations, although random forest algorithms handle colinear predictors well.

In our experience, one of the most important predictors of future recurrent events is past event rate history. This is particularly important in settings where  $T_i(t)$ ,  $t \in \mathcal{T}_i$ , are highly correlated within individual  $i$ . Studies do not always have access to recurrent event history at follow-up time  $t = t_0$ , although this information grows and is available for later follow-up windows. The goals of our simulation study are (1) to verify high performance of our random forest algorithm for prediction of  $P(T_i(t) \geq \tau | Z_i(t))$  in the recurrent

event setting and (2) to demonstrate the importance of event rate history in modeling  $P(T_i(t) \geq \tau | \mathbf{Z}_i(t))$  with suggestions on how this can be done. As part of goal (2), for each modeling strategy (RFRE.PO, Model A, Model B), we consider three different settings corresponding to availability of historical event rate information: (1) *none*, where there is no history covariate available (2) *full*, where the history covariate is available for each individual  $i$  in each follow-up window starting at  $t \in \mathcal{T}_i$  or (3) *partial*, where the history covariate is only recorded for windows beginning at  $t > t_0$ .

We now describe the details of how data were simulated for our study. The true relationship between recurrent events ( $T_{i,1} < T_{i,2} < \dots$ ) and patient characteristics (continuous covariates  $Z_1, Z_2, Z_4$ , and categorical covariates  $Z_3, Z_5, Z_6, Z_7$ ) is designed to be intentionally complex. We generate the recurrent events by simulating and transforming possibly dependent gap-times by Gaussian copula. The recurrent event gap-times marginally follow an exponential distribution with hazard:

$$\lambda_i = \exp\{Z_{i2} \sin(Z_{i1}/Z_{i6}) - Z_{i3}I(Z_{i2} > 2) + Z_{i3}I(Z_{i2} \leq 2) + Z_{i1}Z_{i6} + Z_{i2}^2Z_{i4}\}.$$

We simulate  $Z_1 \sim N(0, 5)$ ,  $Z_2 \sim N(2, 0.8)$ ,  $Z_3 + 2 \sim \text{Poisson}(4)$ ,  $Z_4 - 0.1 \sim \text{Beta}(7, 1)$ ,  $Z_5 \in \{0, 1, 2\}$  with probability  $(\frac{2}{3}, \frac{1}{6}, \frac{1}{6})$ ,  $Z_6 \in \{-5, -2, 2, 3\}$  with probability  $(\frac{1}{10}, \frac{1}{3}, \frac{11}{30}, \frac{1}{5})$ , and  $Z_7 \in \{0, 1, 2, 3\}$  with equal probability. Individuals with simulated hazards outside of the range of  $\frac{8}{15}$  and 15 years are removed from the sample and replaced. We induce correlation among exponentially distributed gap times for each subject  $i$  by utilizing the Gaussian copula method as described by Xia, Murray and Tayob (2020), for  $\rho = 0, 0.3, 0.6$  or 0.9. The resultant recurrent event times are converted to a censored longitudinal data format as described in Section 2. Censoring is independently generated from an exponential distribution so that 0%, 23%, 45%, or 63% of observations are censored before the end of the study.

The censored longitudinal data structure assumes  $\mathcal{T} = \{0, \frac{1}{12}, \frac{1}{6}, \dots, 2 \text{ years}\}$ , and  $\tau = \frac{1}{6}$  year. The modified XMT model estimates  $\text{logit}\{E(S_i^T(t))\}$  assuming the relationship  $\text{logit}\{E(S_i^T(t))\} = \beta^\top \mathbf{Z}_i^*(t)$ . For Model A, the ad hoc nonlinear model,  $\mathbf{Z}_i^*(t) = \{Z_{i2} \sin(Z_{i1}/Z_{i6}), (-1)^{I(Z_{i2} > 2)}Z_{i3}, Z_{i1}Z_{i6}, Z_{i2}^2Z_{i4}\}^\top$ . Model B (the main effects modified XMT model) has covariate vector  $\mathbf{Z}_i^*(t) = \{Z_{i1}, Z_{i2}, Z_{i3}, Z_{i4}, Z_{i5}, Z_{i6}, Z_{i7}\}^\top$ . A derived historical covariate, not already included in Model A will be added to augmented  $\mathbf{Z}_i(t)$  in the *full* and *partial* scenarios,  $H_{f,i}(t)$  and  $H_{p,i}(t)$  respectively. At a particular window start time  $t_k$ ,  $H_{f,i}(t_k) = \frac{\sum_{t \in \mathcal{T}^+, t < t_k} \min\{X_i(t), \frac{1}{12} \text{ years}\}}{\sum_{t \in \mathcal{T}^+} I(t < t_k)}$ , and  $H_{p,i}(t_k) = \frac{\sum_{t \in \mathcal{T}_i, t_0 < t < t_k} \min\{X_i(t), \frac{1}{12} \text{ years}\} + \widehat{H}_{p,i}(t_0)}{\sum_{t \in \mathcal{T}_i^+} I(t < t_k)}$ , where  $\mathcal{T}_i^+ = \{t_0 - \frac{1}{12} \text{ years}\} \cup \mathcal{T}_i$ ,  $\widehat{H}_{p,i}(t_0) = \min(\widetilde{X}_i(t_0), \frac{1}{12} \text{ years})$ .  $\widetilde{X}_i(t_0)$  is multiply imputed from an approximate

Exponential(7.5) distribution, consistent with clinician knowledge of recurrent event rates in the population. Hence results from partial scenarios will be combined across 10 imputed data sets in the partial history setting.

For each of the models described above, as well as `RFRE.PO`, Table 1 lists the mean and empirical standard deviation of the index of concordance of the algorithms discussed and Figure 2 shows the distribution of the index of concordance of the algorithms discussed. Increasing levels of correlation between recurrent event times are listed from left to right. As one moves down the table and figure, censoring levels increase.

For all levels of censoring, correlation and historical covariate use, excluding the highest correlation level with historical covariate use, Model A (the ad hoc nonlinear modified XMT model) has the best C-index with varying degrees of superiority to Model B (the main effects modified XMT model) and `RFRE.PO` method. The `RFRE.PO` approach outperforms Model B in all settings, except high correlation with use of covariate history, in which Model B competes with both `RFRE.PO` and Model A. This suggests that in highly correlated settings, patient history is more important than baseline covariate information.

In the absence of correlation, use of  $H_p(t)$  in Model B weakens predictive performance, whereas predictive performance is similar for the Model B with  $H_f(t)$  versus without use of  $H_f(t)$ . The `RFRE.PO` procedure is able to make large gains in predictive ability from use of  $H_p(t)$  and  $H_f(t)$  in the same case with no correlation. Generally, use of  $H_f(t)$  gives superior predictive performance to use of  $H_p(t)$  in settings with correlation between recurrent event times. Our summary of the `RFRE.PO` results is that any use of covariate history in settings with and without correlated recurrent event outcomes is quite important and assists in recovering information from the data as censoring increases.

## 6. PREDICTION OF COPD EXACERBATIONS

Chronic obstructive pulmonary disease (COPD) is most often attributed to a smoking history, although cases may also be caused by breathing polluted air such as biomass fuel or industrial chemicals (Safiri *and others*, 2022). COPD is characterized by periods of relative stability punctuated with episodic exacerbations of coughing, struggling for breath, and/or wheezing. Exacerbations may or may not result in hospitalization, in which case they are called “severe.” Such exacerbations increase overall disease progression, eventually leading to lung transplantation or death (Seemungal *and others*, 1998; Donaldson *and others*, 2002). Therefore, it




Fig. 3: A violin plot of the distributions of the C-statistic of the models under consideration in simulation. Measures of center and spread can be found in Table 1. Within a box, historical covariate information increases from left to right. For partial history information, the distribution for the average C-statistic from  $m = 10$  imputed data sets is displayed.

Table 1: Simulated C-statistics listed as Mean (ESD) for varying prediction algorithms, censoring percentages (none is 0%, light is 23%, moderate is 45%, and heavy is 63%), and gap-time correlations ( $\rho = 0, 0.3, 0.6, 0.9$ ). 300 replicates were performed with  $n = 500$  for each replicate.

	$\rho = 0$	$\rho = 0.3$	$\rho = 0.6$	$\rho = 0.9$
No Censoring				
Model A	0.677 (0.006)	0.672 (0.007)	0.664 (0.008)	0.651 (0.010)
RFRE.PO	0.598 (0.014)	0.589 (0.015)	0.576 (0.016)	0.558 (0.016)
Model B	0.530 (0.007)	0.531 (0.009)	0.531 (0.010)	0.529 (0.011)
Model A with $H_p(t)$	0.675 (0.006)	0.674 (0.007)	0.684 (0.007)	0.710 (0.007)
RFRE.PO with $H_p(t)$	0.629 (0.007)	0.651 (0.007)	0.683 (0.008)	0.710 (0.008)
Model B with $H_p(t)$	0.489 (0.023)	0.582 (0.019)	0.666 (0.011)	0.724 (0.009)
Model A with $H_f(t)$	0.676 (0.006)	0.677 (0.006)	0.691 (0.011)	0.712 (0.010)
RFRE.PO with $H_f(t)$	0.634 (0.008)	0.658 (0.007)	0.686 (0.009)	0.710 (0.007)
Model B with $H_f(t)$	0.519 (0.035)	0.645 (0.035)	0.685 (0.017)	0.731 (0.010)
Light Censoring				
Model A	0.676 (0.006)	0.672 (0.008)	0.665 (0.009)	0.650 (0.011)
RFRE.PO	0.594 (0.014)	0.585 (0.014)	0.574 (0.016)	0.557 (0.016)
Model B	0.530 (0.009)	0.531 (0.009)	0.532 (0.010)	0.530 (0.011)
Model A with $H_p(t)$	0.676 (0.007)	0.673 (0.008)	0.681 (0.007)	0.704 (0.008)
RFRE.PO with $H_p(t)$	0.628 (0.008)	0.649 (0.008)	0.680 (0.006)	0.707 (0.007)
Model B with $H_p(t)$	0.491 (0.025)	0.574 (0.019)	0.659 (0.013)	0.717 (0.011)
Model A with $H_f(t)$	0.675 (0.006)	0.677 (0.007)	0.689 (0.010)	0.705 (0.010)
RFRE.PO with $H_f(t)$	0.631 (0.008)	0.655 (0.008)	0.684 (0.008)	0.707 (0.008)
Model B with $H_f(t)$	0.523 (0.037)	0.642 (0.017)	0.680 (0.019)	0.724 (0.011)
Moderate Censoring				
Model A	0.676 (0.007)	0.672 (0.007)	0.664 (0.009)	0.649 (0.012)
RFRE.PO	0.591 (0.014)	0.584 (0.015)	0.570 (0.015)	0.555 (0.017)
Model B	0.531 (0.009)	0.532 (0.010)	0.531 (0.010)	0.530 (0.012)
Model A with $H_p(t)$	0.675 (0.007)	0.672 (0.008)	0.679 (0.008)	0.701 (0.008)
RFRE.PO with $H_p(t)$	0.623 (0.009)	0.644 (0.008)	0.676 (0.007)	0.705 (0.007)
Model B with $H_p(t)$	0.492 (0.023)	0.565 (0.021)	0.649 (0.015)	0.712 (0.012)
Model A with $H_f(t)$	0.675 (0.007)	0.676 (0.007)	0.686 (0.011)	0.700 (0.011)
RFRE.PO with $H_f(t)$	0.629 (0.009)	0.653 (0.008)	0.680 (0.009)	0.704 (0.008)
Model B with $H_f(t)$	0.528 (0.034)	0.639 (0.016)	0.674 (0.021)	0.718 (0.012)
Heavy Censoring				
Model A	0.676 (0.007)	0.672 (0.009)	0.664 (0.010)	0.649 (0.011)
RFRE.PO	0.586 (0.017)	0.577 (0.016)	0.568 (0.018)	0.551 (0.017)
Model B	0.532 (0.009)	0.532 (0.009)	0.533 (0.012)	0.530 (0.012)
Model A with $H_p(t)$	0.674 (0.007)	0.672 (0.009)	0.676 (0.009)	0.695 (0.009)
RFRE.PO with $H_p(t)$	0.618 (0.010)	0.640 (0.009)	0.671 (0.008)	0.700 (0.007)
Model B with $H_p(t)$	0.492 (0.025)	0.554 (0.022)	0.640 (0.016)	0.703 (0.013)
Model A with $H_f(t)$	0.675 (0.007)	0.675 (0.008)	0.684 (0.011)	0.695 (0.012)
RFRE.PO with $H_f(t)$	0.625 (0.010)	0.650 (0.009)	0.676 (0.009)	0.700 (0.009)
Model B with $H_f(t)$	0.531 (0.032)	0.636 (0.016)	0.668 (0.024)	0.711 (0.014)

is of clinical interest to dynamically predict the chances of being exacerbation-free over time.

We apply RFRE.PO to predict the probability of remaining exacerbation-free over a subsequent  $\tau = 180$  day follow-up period, using data from Azithromycin for the Prevention of COPD (Albert *and others*, 2011).

Azithromycin for the Prevention of COPD was a multi-center clinical trial, aiming to estimate the treatment effect 250mg of oral azithromycin had on reducing exacerbations in COPD patients. We pooled data from both arms of the clinical trial, and used treatment assignment as one of many covariates in our prediction model.

Our example is applied to study participants ( $n = 1035$ ) who have complete baseline data, as well as longitudinal sleep study variables. Recurrent exacerbations were monitored over the following year. There are 962 covariates measured, including demographic variables, social well-being surveys, sleep quality metrics and clinical characteristics such as forced vital capacity (FVC), and medication use. There is strong correlation between variables as some are derived from each other, making `RFRE.PO` an attractive alternative to model-based predictions (see Supplemental Figure ??). Censored longitudinal data is constructed with  $\mathcal{T} = \{0, 30, 60, \dots, 180\}$  and used for all analyses shown in this section.

Baseline information on the time since the most recent exacerbation was not available. Hence in this study we are in a scenario similar to the partial history scenario in Section 5. In this example, we create two partial history variables,  $H_{p_1}(t)$  and  $H_{p_2}(t)$ .  $H_{p_1}(t)$  was assumed to take a categorical form based on whether the participant had an exacerbation in the past zero to 30 days, 31-92 days, 93-182 days, 183 days to 365 days, or finally, if that time was more than 365 days or never.  $H_{p_2}(t)$  is 100 times the estimated 30-day event rate using all follow-up prior to time  $t$  (so as to provide point estimates on a reasonable scale). For the first  $H_{p_1}(0)$  and  $H_{p_2}(0)$ , multiple imputation was based on event rates in the placebo group for all participants, and Rubin’s rule (Rubin, 1987) was used to combine inference across multiply imputed datasets as appropriate.

For comparison purposes, we present `RFRE.PO` results alongside results from two modified XMT models, where modified XMT models are fit via the `geese` command from the `geepack` library in R. The first modified XMT model (Model 1) uses the same predictors identified in Xia, Murray and Tayob (2020) for this cohort, where these predictors were based on clinical judgement for estimating treatment effect, adjusted for known confounders. The second modified XMT model (Model 2) is based on Wald forward selection applied to the first multiply imputed data set with  $p < 0.05$  required for entry; covariates that introduce model instability are discarded from consideration. Prediction algorithms for `RFRE.PO` and Model 2 were applied to a training cohort (70% of original cohort) and all algorithms were evaluated via the C-statistic in a validation cohort



(remaining 30% of original cohort). For any particular multiply imputed validation dataset, standard errors of the C-statistic are calculated via the bootstrap with  $b = 100$  bootstrap samples. C-statistics for models involving multiply imputed history variables are combined across multiply imputed datasets using Rubin's rule. Parameter estimates and standard errors for Model 2, which involves the imputed partial history covariates, are combined across multiply imputed validation datasets via Rubin's Rule. Permutation tests associated with the RFRE.PO algorithm predictors are calculated within the training dataset, using out-of-bag samples as described in Section 3.

For Model 1, where estimation of treatment effect, and not necessarily prediction was the goal, predictors that were identified as important in Xia, Murray and Tayob (2020) were baseline forced expiratory volume in liters (FEV), age in decades (Age10), gender (0 for female, 1 for male), and smoking status at baseline (Smoker):

$$\text{logit}\{E(S_i^T(t))\} = \beta_0 + \beta_1 I(\text{Treatment})_i + \beta_2 (\text{Baseline FEV})_i + \beta_3 (\text{Age10})_i + \beta_4 I(\text{Male})_i + \beta_5 I(\text{Smoker})_i,$$

where parameter estimates for the modified XMT model are displayed in Supplemental Table ??.

Model 2, the Wald forward selection model, is:

$$\begin{aligned} \text{logit}\{E(S_i^T(t))\} = & \beta_0 + \beta_1 I(H_{p_1} = \text{Exacerbation in 0 -30 days prior to time } t)_i + \\ & \beta_2 I(\text{Bronchiectasis diagnosis before time } t)_i + \beta_3 I(\text{Male gender})_i + \beta_4 I(\text{Azithromycin group})_i + \\ & \beta_5 I(\text{Non-selective beta-blocker use at time } t)_i + \beta_6 I(\text{Mixed beta-blocker use at time } t)_i + \\ & \beta_7 (\text{FEV}_1/\text{FVC } \% \text{ predicted at time } t)_i + \beta_8 (\text{SGRQ symptoms at time } t)_i + \\ & \beta_9 I(\text{Supplemental oxygen use at time } t)_i + \beta_{10} I(\text{Hospitalization one year before study start})_i + \\ & \beta_{11} H_{p_2}(t)_i + \beta_{12} I(\text{Inhaled corticosteroid and long-acting muscarinic antagonist use at time } t)_i + \\ & \beta_{13} I(\text{Inhaled corticosteroid and long-acting beta-agonist use at time } t)_i, \end{aligned}$$

with parameter estimates displayed in Supplemental Table ??.

For the RFRE.PO algorithm, Supplemental Table ?? displays the corresponding p-values of the permutation tests, combined across multiply imputed datasets via Rubin's rule. Predictors in Supplemental Table ?? had a statistically significant permutation test z-score for at least one of the multiply imputed training datasets. Additional predictors that did not achieve statistical significance via the permutation test, but were used in the RFRE.PO algorithm in some form, are not shown.

Figure 4 displays the Z-scores for predictors included in Models 1 and 2 (validation dataset) or Table ?? (out-of-bag samples from training dataset). As seen in Section 5, history variables were among the most important variables in the RFRE.PO approaches as well as the modified XMT model selected through Wald Z-score forward selection. While there is overlap between predictors included in the various algorithms, in terms of the C-statistic evaluated in the validation cohort, the RFRE.PO algorithm outperformed both the algorithm based on clinical input (Model 1) and the Wald forward selection model (Model 2) (C-statistic 0.616 compared to 0.565 and 0.550, respectively, see Table 2). A large number of variables that were included in the random forest algorithm had statistical significance at the level of 0.05 via the permutation test in at least one of the multiply imputed datasets (see Figure 4 for the z-statistic and p-values in Supplemental Table ?? for the predictors deemed important). Statistically significant variables from Model 1 are similarly listed as important RFRE.PO predictors. Variables identified via Wald forward selection in Model 2 are for the most part identified as important variables in the RFRE.PO algorithm. The version of the predictors may vary between these two approaches. For example, variables related to the long-acting muscarinic or bronchodilator, and/or steroid treatment are incorporated in different forms into these models. Similarly, different heart medications are featured in Model 2 (beta-blockers) versus the RFRE.PO algorithm (anti-coagulants). The RFRE.PO algorithm highlights a few variables that were not used in Model 2, including pain level at time  $t$  and leukocytosis present at time  $t$ . The RFRE.PO algorithm has no problem including information from correlated predictors; for instance two included variables measure pain (pain level at time  $t$ , standardized pain level at time  $t$ ) and two included variables measure general health (general health score at time  $t$  standardized general health score at time  $t$ ).

The C-statistic corresponding to Model 1, which benefits from prior clinical knowledge of the causal pathway, validated well. The RFRE.PO random forest algorithm includes predictors, with corresponding split decisions, that minimize squared error of the predictions relative to the observed data. This results in predictor information that is included without rigorous checks for statistical significance, so validation in an independent data set is quite important when using this type of algorithm. Because Model 2 requires statistical significance of all included predictors, its individual predictors already have some theoretical validity built into the selection mechanism. However, with variable power to detect statistical significance of useful predictors, the comparisons with methods that do not require statistical significance will leave Model

Table 2: Comparison of the C-statistics of models fit to the validation set of the Azithromycin in the Prevention of COPD cohort. Models 1 and 2 are fit using the modified XMT approach. Model 1 is the model selected using clinician input from Xia, Murray and Tayob (2020). Model 2 includes predictors through automated forward selection requiring  $p < 0.05$  for entry into the model. Missing values in Model 2 and `RFRE.PO` for time since most recent exacerbation are multiply imputed using the coding scheme described in Section 6. `RFRE.PO` processes our random forest approach using the same predictors considered in Model 2. C-statistic estimates and standard errors for all methods and models are averaged across results from multiply imputed data with Rubin’s rule applied to bootstrapped ( $b = 100$ ) C-statistic standard errors.

Model Name	C-Statistic	
	Mean	Standard Error
<code>RFRE.PO</code>	0.616	0.009
Model 1	0.565	0.009
Model 2	0.550	0.010

2-style strategies at a disadvantage.

The `RFRE.PO` algorithm is completely agnostic to statistical significance and scientific experience, so long as the squared error within nodes along cut-points decreases. And although it may occasionally include a predictor that is not useful in the validation set (Type I Error analog), it has a much better chance of including predictors that are useful but would not have met statistical significance (akin to a Type II error) in traditional statistical model building environments. If the scientific goal of analysis is prediction, the `RFRE.PO` approach seems remarkably simple, effective, and reproducible.

## 7. DISCUSSION

In this paper we presented a novel method for dynamically estimating probability of event-free survival during different follow-up windows. Using `RFRE.PO`, we showed the utility of our method in simulation, comparing its performance to traditional longitudinal model fits with an modified XMT model. In all instances, except for the most highly correlated, `RFRE.PO` outperformed the main effects modified XMT model that a traditional modeler might first attempt to fit. In instances of high correlation, `RFRE.PO` and the main effects model competed with the ad hoc nonlinear modified XMT model- that is, the perfect model that exactly reflected how the data were simulated.

An attractive feature of random forest applied to longitudinal follow-up windows is that a split based on the follow-up window times can occur and predictors that are more helpful in later follow up times can emerge naturally in importance. We explored recent history prior to the beginning of each follow-up window




Fig. 4: Wald Statistic Z-scores derived from the validation set for predictors that were included in Models 1 (modified XMT Xia, Murray, Tayob 2020 Model) and 2 (modified XMT Wald Forward Selection) after adjusting for multiple imputation using Rubin’s Rule, where appropriate, and permutation test Z-scores derived from out-of-bag samples as part of the RFRE.PO algorithm, adjusting for multiple imputation using Rubin’s Rule where appropriate. For the RFRE.PO method, 100 permutations were used in calculating the permutation Z-scores. Note that some Wald forward selection variables lost significance as additional terms were added.

as a covariate in the algorithm, and found that it was one of the most useful predictors in settings with high correlation between recurrent events. The use of complex scoring algorithms has increased with the development of machine learning predictors such as RFRE.PO. For example, the COPD research community has agreed that algorithmic prediction models need to be user-friendly to be used in clinical practice. To this end, the COPDGene Study <sup>®</sup> has created an app to calculate various risk scores based on patient characteristics that a user can input. Apps that simplify risk score calculations along these lines are one such way to bring complicated scores to clinical practitioners.

It is always important when introducing a new piece of methodology to clearly identify research questions that it can and cannot address. Random forest technology in general is aimed towards prediction, and for **RFRE.PO**, predicting the probability of remaining event-free during subsequent follow-up windows after patient check-ins. In this case, all available predictors that the treating clinician can access are valid and important to use. There is no goal, per sé, of establishing which predictors are statistically significant or have causal relationships with the outcomes. It is likely that some, if not most, of the predictors with high variable importance scores have these features, but metrics assessing statistical significance and causality are not necessarily valid for these algorithms.

In contrast, models such as the modified XMT model for recurrent event outcomes often have the goal of measuring the impact of a particular predictor or set of predictors on the outcome via an effect size, confidence interval and p-value. In the Azithromycin for Prevention of COPD cohort, the azithromycin effect was the key research question, requiring a measure of statistical significance as part of the drug approval process. Some predictors that would be very helpful in prediction are not allowed in these models assessing treatment effect. For instance, it is well-known that one should not include updates to patient characteristics after treatment has been initiated if these characteristics could have been improved by treatment; doing so would adjust away part of the treatment effect under study. Similarly, clinical trial analyses often require careful adjustment for confounders. If the goal is prediction and not assessment of treatment effect, these same confounders might not be included in the algorithm and in fact may not be helpful relative to other available predictors.

The ability of flexible algorithms like **RFRE.PO** to sift through large numbers of correlated predictors with complicated relationships to outcomes is an argument for their use over traditional reliance on models such as the modified XMT model. The growing popularity of machine learning tools is likely to dominate analyses of high-dimensional data, and the **RFRE.PO** algorithm contributes to this literature as one of the first machine learning methods to be able to make dynamic predictions from recurrent event data.

#### SUPPLEMENTARY MATERIAL

Supplementary material is available online at <http://biostatistics.oxfordjournals.org>.

## ACKNOWLEDGMENT

The research is supported in part by a Precision Health Investigator Award from University of Michigan, Ann Arbor (AL, ZW). *Conflict of Interest*: None.

## DATA AVAILABILITY STATEMENT

Code for simulation and data analysis is available at <https://github.com/AbigailLoe/RFRE.PO>. The data that support the findings in this paper are available from the corresponding author upon reasonable request.

## REFERENCES

- ADLER, W., POTAPOV, S. AND LAUSEN, B. (2011, Mar). Classification of repeated measurements data using tree-based ensemble methods. *Computational Statistics* **26**, 355–369.
- ALBERT, R. K., CONNETT, J., BAILEY, W., CASABURI, R., COOPER, A., CRINER, G., CURTIS, J., DRANSFIELD, M., HAN, M., LAZARUS, S., MAKE, B., MARCHETTI, N., MARTINEZ, F., MADINGER, N., McEVOY, C., NIEWOEHNER, D., PORSASZ, J., PRICE, C., REILY, J., SCANLON, P., SCIURBA, F., SCHARF, S., WASHKO, G., WOODRUFF, P. *and others.* (2011). Azithromycin for prevention of exacerbations of copd. *New England Journal of Medicine* **365**(8), 689–698.
- ANDERSEN, P. K. AND GILL, R. D. (1982). Cox’s regression model for counting processes: A large sample study. *The Annals of Statistics* **10**(4), 1100–1120.
- ANDERSEN, P. K., KLEIN, J. P. AND ROSTHØJ, S. (2003). Generalised linear models for correlated pseudo-observations, with applications to multi-state models. *Biometrika* **90**(1), 15–27.
- ANDREI, A.-C. AND MURRAY, S. (2007, Jun). Regression models for the mean of the quality-of-life-adjusted restricted survival time using pseudo-observations. *Biometrics* **63**, 398–404.
- ANG, Y., MEYERSON, L., TANG, Y. AND QIAN, N. (2009, Aug). Statistical methods for the analysis of relapse data in ms clinical trials. *Journal of Neurological Sciences* **285**(1-2), 206–211.
- BREIMAN, L. (2001). Random forests. *Machine Learning* **45**, 5–32.
- BREIMAN, L., FREIDMAN, J. H., OLSHEN, R. A. AND STONE, C. J. (1984). *CART: Classification and Regression Trees*.

- CHAN, K. C. G. AND WANG, M.-C. (2017a). Semiparametric modeling and estimation of the terminal behavior of recurrent marker processes before failure events. *Journal of the American Statistical Association* **112**(517), 351–362.
- CHAN, K. C. G. AND WANG, M.-C. (2017b, May). Semiparametric modeling and estimation of the terminal behavior of recurrent marker processes before failure events. *U.S. National Library of Medicine*.
- CHOI, S., HUANG, X., JU, H. AND NING, J. (2017). Semiparametric accelerated intensity models for correlated recurrent and terminal events. *Statistica Sinica* **27**(2), 625–643.
- DONALDSON, G., SEEMUNGAL, T., BHOWMIK, A. AND WEDZICHA, J. (2002, October). Relationship between exacerbation frequency and lung function in chronic obstructive pulmonary disease. *Thorax* **57**(10), 847–852.
- GERBER, G., FAOU, Y. L., LOPEZ, O. AND TRUPIN, M. (2021a). The impact of churn on client value in health insurance, evaluation using a random forest under various censoring mechanisms. *Journal of the American Statistical Association* **116**(536), 2053–2064.
- GERBER, G., FAOU, Y. L., LOPEZ, O. AND TRUPIN, M. (2021b). The impact of churn on client value in health insurance, evaluation using a random forest under various censoring mechanisms. *Journal of the American Statistical Association* **116**(536), 2053–2064.
- HARRELL, J. FRANK E., CALIFF, R. M., PRYOR, D. B., LEE, K. L. AND ROSATI, R. A. (1982, 05). Evaluating the Yield of Medical Tests. *Journal of the American Medical Association* **247**(18), 2543–2546.
- HENGELBROCK, J., GILLHAUS, J., KLOSS, S. AND LEVERKUS, F. (2016). Safety data from randomized controlled trials: applying models for recurrent events. *Pharmaceutical Statistics* **15**(4), 315–323.
- HU, J. AND SZYMCZAK, S. (2023). A review on longitudinal data analysis with random forest. *Briefings in Bioinformatics* **2**(24), 1–11.
- ISHWARAN, H., BLACKSTONE, E. H., POTHIER, C. E. AND LAUER, M. S. (2004). Relative risk forests for exercise heart rate recovery as a predictor of mortality. *Journal of the American Statistical Association* **99**(467), 591–600.

- KONG, M., XU, S., LEVY, S. AND DATTA, S. (2016, May). Gee type inference for clustered sero-inflated negative binomial regression with application to dental caries. *Computational Statistics and Data Analysis* **85**, 54–66.
- LIANG, B., TONG, X., ZENG, D. AND WANG, Y. (2017). Semiparametric regression analysis of repeated current status data. *Statistica Sinica* **27**(3), 1079–1100.
- LIN, D. Y., WEI, L.-J., YANG, I. AND YING, Z. (2000). Semiparametric regression for the mean and rate functions of recurrent events. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* **62**(4), 711–730.
- MAO, L. AND LIN, D. Y. (2016, Apr). Semiparametric regression for the weighted composite endpoint of recurrent and terminal events. *U.S. National Library of Medicine*.
- MILOSLAVSKY, M., KELEŞ, S., VAN DER LAAN, M. J. AND BUTLER, S. (2004). Recurrent events analysis in the presence of time-dependent covariates and dependent censoring. *Journal of the Royal Statistical Society. Series B (Statistical Methodology)* **66**(1), 239–257.
- RUBIN, D. B. (1987). Multiple imputation for nonresponse in surveys.
- SAFARI, A., PETKAU, J., FITZGERALD, M. J. AND SADATSAFAVI, M. (2023). A parametric model to jointly characterize rate, duration, and severity of exacerbations in episodic diseases. *BMC Medical Informatics and Decision Making* **23**(1), 1–11.
- SAFIRI, S., CARSON-CHAHHOUD, K., NOORI, M., NEJADGHADERI, S. A., SULLMAN, M. J. M., HERIS, J. A., ANSARIN, K., MANSOURNIA, M. A., COLLINS, G. S., KOLAHI, A.-A. and others. (2022). Burden of chronic obstructive pulmonary disease and its attributable risk factors in 204 countries and territories, 1990-2019: results from the global burden of disease study 2019. *BMJ* **378**, e069679.
- SEEMUNGAL, T., DONALDSON, G., PAUL, E., JEFFRIES, D. AND WEDZICHA, J. (1998, May). Effect of Exacerbation on quality of life in patients with chronic obstructive pulmonary disease. *American Journal of Critical Care Medicine* **157**(5), 1418–22.
- TAYOB, N. AND MURRAY, S. (2015, January). Nonparametric tests of treatment effect based on combined endpoints for mortality and recurrent events. *Biostatistics* **16**(1), 73–83.



- WAGER, S. AND ATHEY, S. (2018). Estimation and inference of heterogeneous treatment effects using random forests. *Journal of the American Statistical Association* **113**(523), 1228–1242.
- XIA, M. AND MURRAY, S. (2019, July). Commentary on Tayob and Murray (2014) with a useful update pertaining to study design. *Biostatistics* **20**(3), 542—545.
- XIA, M., MURRAY, S. AND TAYOB, N. (2020, January). Regression analysis of recurrent-event-free time from multiple follow-up windows. *Statistics in Medicine* **39**, 1–15.
- XU, G., CHIOU, S. H., HUANG, C.-Y., WANG, M.-C. AND YAN, J. (2017). Joint scale-change models for recurrent events and failure time. *Journal of the American Statistical Association* **112**(518), 794–805.
- YOKOTA, I. AND MATSUYAMA, Y. (2019, Feb). Dynamic prediction of repeated events data based on leanmaking model: application to colorectal liver metastases data. *BMC Medical Research Methodology* **19**(31), 1–11.
- ZHAO, L., MURRAY, S., MARIANI, L. H. AND JU, W. (2020). Incorporating longitudinal biomarkers for dynamic risk prediction in the era of big data: A pseudo-observation approach. *Statistics in Medicine* **39**(26), 3685–3699.

[Received XX XX, XXXX; revised XX XX, XXXX; accepted for publication XXX XX, XXXX]

# Supporting Information for “Random Forest for Dynamic Risk Prediction of Recurrent Events: A Pseudo-Observation Approach”

Abigail Loe<sup>1,\*</sup>, Susan Murray<sup>1,+</sup>, and Zhenke Wu<sup>1,+</sup>

<sup>1</sup>Department of Biostatistics, University of Michigan, Ann Arbor, MI 48109, USA; E-mail: \*agloe@umich.edu; +: co-senior authors

## A1 Tables and Figures

Table S1: XMT Model 1 parameter estimates, based on covariates selected by Xia et al. (2020), applied to censored longitudinal data with  $\tau = 180$ ,  $a = 30$ .

Covariate	$e^{\hat{\beta}^\dagger}$	95% CI	p
Azithromycin (versus placebo)	1.43	(1.14, 1.79)	< <b>0.01</b>
Smoker at baseline (versus non-smoker)	1.07	(0.80, 1.43)	0.66
FEV <sub>1</sub> in liters at baseline	1.32	(1.05, 1.66)	<b>0.02</b>
Male gender (versus female)	1.36	(1.07, 1.74)	<b>0.01</b>
Age in decades	1.14	(0.99, 1.32)	0.07

<sup>†</sup>  $e^{\hat{\beta}}$  is the estimated odds ratio for being event-free during a 180-day follow-up period corresponding to a 1-unit increase in the (continuous) predictor, or the presence of a binary covariate, adjusted for other predictors.

FEV<sub>1</sub> is forced expiratory volume in one second, a clinical measure of lung function.




Figure S1: A correlation heat map of continuous and ordinal variables in the Azithromycin for the Prevention of COPD Cohort, measured at baseline. Variables derived from each other have instances of particularly high positive or negative correlation. These correlation patterns make random forest a more attractive modeling alternative compared to semi-parametric and parametric methods.

Table S2: Parameter Estimates from Model 2, or the model selected through Wald forward-selection on the first of  $m = 10$  imputed datasets. A very recent exacerbation, bronchitis, gender, treatment, the ratio of FEV to FVC %predicted, SGRQ symptoms, oxygen use, recent hospitalization, rate of exacerbations, and use of ICS medication remain significant across the  $m = 10$  imputed datasets. Exacerbation history indicators and rate of exacerbations are multiply imputed.

Covariate	$e^{\hat{\beta}}$ †	95% CI	p
Exacerbation in the 0-30 days prior to time $t$ (versus more than 30 days)	2.18	(1.91, 2.48)	< <b>0.01</b>
Bronchiectasis diagnosis at time $t$ (versus no diagnosis)	0.60	(0.44, 0.81)	< <b>0.01</b>
Male gender (versus female gender)	1.94	(1.43, 2.64)	< <b>0.01</b>
Azithromycin (versus placebo)	1.62	(1.20, 2.18)	< <b>0.01</b>
Non-selective beta-blocker use at time $t$ (versus selective and no beta-blocker)	0.14	(0.00, 5.71)	0.30
Mixed beta-blocker use at time $t$ (versus selective and no beta-blocker)	2.54	(0.98, 6.59)	<b>0.05</b>
FEV <sub>1</sub> /FVC % predicted at time $t$	1.01	(1.00, 1.02)	<b>0.03</b>
SGRQ symptoms at time $t$	0.99	(0.99, 1.00)	<b>0.03</b>
Supplemental oxygen use at time $t$ (versus no supplemental oxygen)	0.74	(0.55, 0.99)	<b>0.04</b>
Hospitalization in the year before study time (versus not hospitalized)	0.60	(0.45, 0.80)	< <b>0.01</b>
Rate of exacerbation prior to time $t$	1.25	(1.05, 1.50)	<b>0.01</b>
Corticosteroid and long-acting muscarinic antagonist use at time $t$ (versus not)	0.72	(0.55, 0.95)	<b>0.02</b>
Corticosteroid and long-acting beta-agonist use at time $t$ (versus not)	1.30	(1.03, 1.63)	<b>0.03</b>

†:  $e^{\hat{\beta}}$  is the estimated odds ratio for being event-free during a 180-day follow-up period corresponding to a 1-unit increase in the (continuous) predictor, or the presence of a binary covariate, adjusted for other predictors.

FEV<sub>1</sub> is forced expiratory volume in one second and FVC is forced vital capacity, clinical measures of lung function.

SGRQ is the Saint George's Respiratory Questionnaire, a patient reported outcome measuring quality of life.

Table S3: P-values from the RFRE.PO permutation test applied to the out-of-bag-samples combined across  $m = 10$  multiply imputed datasets. Included predictors had a statistically significant permutation test z-score for at least one of the multiply imputed training datasets. Additional predictors that did not achieve statistical significance via the permutation test, but were used in the RFRE.PO algorithm in some form, are not shown.

Covariate	p
Exacerbation in the 0-30 days prior to time $t$	<b>0.04</b>
Exacerbation in the 31-92 days prior to time $t$	0.06
Exacerbation in the 366+ days (or none) prior to time $t$	<b>0.01</b>
Categorical time since most recent exacerbation prior to time $t$ , or $H_{p_1}(t)$	0.79
Rate of Exacerbation	0.61
Had a severe exacerbation on study time prior to time $t$	0.05
General health score at time $t$	0.90
Standardized general health z-score at time $t$	< <b>0.01</b>
Male gender	< <b>0.01</b>
Anti-coagulant use at time $t$	<b>0.01</b>
Pain score at time $t$	< <b>0.01</b>
Standardized pain z-score at time $t$	0.26
FEV <sub>1</sub> in liters at time $t$	<b>0.01</b>
Treatment group	0.42
SGRQ symptoms at time $t$	< <b>0.01</b>
Long-acting beta-agonist use at time $t$	0.90
Long-acting muscarinic antagonist use at time $t$	<b>0.02</b>
Inhaled corticosteroids use in the year before study time	0.07
Hospitalization in the year before study time	0.38
Phlegm at time $t$	0.51
Baseline smoking	0.05
Leukocytosis at time $t$	0.05
Oxygen use at time $t$	0.07
Bronchiectasis diagnosis at time $t$	0.50
FEV <sub>1</sub> /FVC % Predicted at time $t$	0.40
Saw a mental health provider for depression or anxiety at time $t$	0.53
Anxiety HADS score at time $t$	0.21

FEV<sub>1</sub> is forced expiratory volume in one second and FVC is forced vital capacity, clinical measures of lung function.

SGRQ is the Saint George's Respiratory Questionnaire, a patient reported outcome measuring quality of life.

HADS Anxiety score Hospital Anxiety and Depression Scale is a clinical measure of mental well-being.

## References

Xia, M., Murray, S., and Tayob, N. (2020). Regression analysis of recurrent-event-free time from multiple follow-up windows. *Statistics in Medicine*, 39:1–15.