

M_{W_R} dependence of leptogenesis in Minimal Left-Right Symmetric Model with different strengths of Type-II seesaw mass.

Ankita Kakoti^{1,*} and Mrinal Kumar Das^{1,†}

¹*Department of Physics, Tezpur University, Tezpur 784028, India*

Abstract

Left Right Symmetric Model (LRSM) being an extension of the Standard model of particle physics incorporates within itself Type-I and Type-II seesaw mass terms naturally. Both the mass terms can have significant amount of contribution to the resulting light neutrino mass within the model and hence on the different phenomenology associated within. In this paper, we have thoroughly analyzed and discussed the implications of specifying different weightages to both the mass terms and also the study has been carried out for different values of M_{W_R} which is mass of the right-handed gauge boson. This paper also gives a deeper insight into the new physics contributions of Neutrinoless Double Beta Decay ($0\nu\beta\beta$) and their variations with the net baryon asymmetry arising out of the model. Therefore, the main objective of the present paper rests on investigating the implications of imposing different weightage to the type-I and type-II seesaw terms and different values of M_{W_R} on the new physics contributions of $0\nu\beta\beta$ and net baryon asymmetry arising out as a result of resonant leptogenesis. LRSM in this work has been realized using modular group of level 3, $\Gamma(3)$ which is isomorphic to non-abelian discrete symmetry group A_4 , the advantage being the non-requirement of flavons within the model and hence maintaining the minimality of the model.

* ankitak@tezu.ernet.in

† mkdas@tezu.ernet.in

I. INTRODUCTION

Considering the fact that neutrinos are massive and mixing occur between their flavors which has been proved by the neutrino oscillation experiments, we are compelled to move beyond the Standard paradigm. Although Standard Model (SM) has been enormously victorious in explaining the fundamental particles and their interactions, it fails to address some issues like the origin of tiny neutrino masses, baryon asymmetry of the universe (BAU) [1], lepton flavor violation (LFV) [2], origin of dark matter etc [3, 4].

In this context, we can state that the seesaw mechanism [5–11] has proven to be a very straightforward way to explain the tiny neutrino masses, the two main components of the mechanism being, addition of right-handed (RH) neutrinos to the SM and endowing the RH neutrinos with a Majorana mass which breaks the accidental global (B-L) symmetry of the SM. However, in the context of electroweak gauge theory of the SM described by $SU(3)_C \otimes SU(2)_L \otimes U(1)_Y$, there is no underlying principle that explains this mechanism, but rather they are put in by hand. In this context, we have two ultraviolet-complete theories where both the components of seesaw arise in a natural manner and these are,

- The left-right (LR) symmetric theories of weak interactions based on the gauge group $SU(3)_C \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$ [7, 12–18].
- $SO(10)$ grand-unified theory for all interactions [19].

In this work, our focus is on low-scale Left-Right symmetric model (LRSM) where the seesaw scale can be in a few TeV range, such that it is accessible to the LHC, while satisfying the observed charge-lepton and neutrino mass spectra.

One important question that arises here is that why the seesaw mechanism is so favourable for the explanation of tiny neutrino masses. This is because, when we take seesaw mechanism into consideration, the same Yukawa that are responsible for generating small neutrino masses may also be helpful in resolving one of the outstanding puzzles of cosmology, namely, the origin of matter-antimatter asymmetry, via leptogenesis. Leptogenesis can be picturized by the out-of equilibrium decays of the RH neutrinos via the modes $N \rightarrow L_l \phi$ and $N \rightarrow L_l^c \phi^c$, where, $L_l = (\nu_L l)_L^T$ are the $SU(2)_L$ lepton doublets, ϕ is the Higgs doublet and the superscript c denotes the CP conjugate. In the presence of CP violation in the Yukawa sector, the decays mentioned above can lead to lepton

asymmetry in the early universe satisfying the three Sakharov conditions. This asymmetry then undergoes thermodynamic evolution with the expansion of the universe and different reactions present in the model have their impact on washing out part of the asymmetry. The remaining final lepton asymmetry is converted to baryon asymmetry via sphaleron processes before the electroweak phase transition.

For TeV scale seesaw models, we require resonant leptogenesis (RL) for the generation of adequate lepton asymmetry, where atleast two of the heavy Majorana neutrinos have a small difference comparable to their decay widths. The TeV scale new particles in LRSM also leads to interesting collider signatures.

In this work, we have also taken into account the study of neutrinoless double beta decay (NDBD) [20], [21], because the possible observation of NDBD would play an important role in understanding the origin of BAU as NDBD would imply that lepton number is indeed not conserved, which is one of the essential conditions for leptogenesis. In LRSM, there are several contributions to NDBD that involve left-right sectors individually and also some contributions that involves both the sector through left-right mixing accompanied by both light and heavy neutrinos. Left-Right mixing is a ratio of the Dirac and Majorana mass scales $[M_D M_{RR}^{-1}]$, appearing in the type-I seesaw mass formula as we will discuss later. While taking to account large left-right mixing we have to focus on the momentum dependent contributions of NDBD, that is, the contribution that arises as a result of the mixed diagrams with simultaneous mediation of W_L and W_R accompanied by light left-handed and heavy right-handed neutrino known as λ and η contributions.

In this work, we have firstly realized LRSM with the help of modular A_4 symmetry and thus obtained the mass matrices in terms of the modular forms (Y_1, Y_2, Y_3) , and then we move forward to study the linkage between the different contributions of NDBD and baryogenesis. We have considered different strengths of the type-I and type-II seesaw masses with different values of the $SU(2)_R$ breaking scale, namely, type-II seesaw mass being 30%, 50%, 70% dominant for $M_{W_R} = 1.5, 2.2, 2.5, 3, 5, 10, 18$ TeV. We have not considered any particular texture for the Dirac and Majorana mass matrices and in order to account for the neutrino mass of the order of sub eV, the Dirac mass that we have determined is in the scale of MeV, as the heavy right handed neutrino is kept in the TeV scale. This leads to not so large left-right mixing and hence we get non-negligible effects of the momentum dependent contributions of NDBD. Although we have

studied all the contributions of NDBD, however to relate baryogenesis to the phenomena, we have considered only the contributions appearing from the mediation of heavy right-handed neutrino and the momentum dependent λ and η contributions. Since the effective mass governing NDBD is dependent on the Majorana phases, α and β , it would be compelling to study if there is a link between NDBD and BAU. So, this paper has been organized by firstly discussing about LRSM and the origin of neutrino mass in section II, section III describing the realization of LRSM with A_4 modular symmetry and section IV describes the phenomenon of resonant leptogenesis and NDBD within LRSM, section V discusses the numerical analysis and results and we conclude our work by discussion and conclusion in section VI and VII. The appendices discussed the minimisation of the Higgs potential for the model and also introduces the concept of modular symmetry which has been used for the realisation of the model.

II. LEFT-RIGHT SYMMETRIC MODEL AND NEUTRINO MASS.

Left-Right Symmetric Model (LRSM) is described by the gauge group $SU(3)_C \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$. As is evident from the gauge group that unlike SM, both the left-handed and right handed particles are taken as doublets in the said model. The scalar sector of the model consists of one Higgs bidoublet $\phi(1, 2, 2, 0)$ and two scalar triplets given by $\Delta_L(1, 3, 1, 2)$ and $\Delta_R(1, 1, 3, 2)$. Because of the presence of right-handed (RH) neutrinos and scalar triplets, LRSM incorporates within itself *Type – I* and *Type – II* seesaw masses naturally.

The fermions can attain mass only when the necessary Yukawa Lagrangian is constructed which shows coupling of particles to the bidoublet ϕ . The Yukawa Lagrangian for fermions incorporating the bidoublet is given by,

$$\mathcal{L}_{\mathcal{D}} = \bar{l}_{iL}(Y_{ij}^l \phi + \widetilde{Y}_{ij}^l \widetilde{\phi}) l_{jR} + h.c \quad (2.1)$$

where, l_L and l_R are the left-handed and right-handed lepton fields. Y^l being the Yukawa coupling corresponding to leptons and Y^q being the Yukawa coupling for the quarks. The Yukawa Lagrangian where the scalar triplets play a role in providing Majorana mass to the neutrinos is given by,

$$\mathcal{L}_{\mathcal{M}} = f_{L,ij} \Psi_{L,i}^T C i \sigma_2 \Delta_L \Psi_{L,j} + f_{R,ij} \Psi_{R,i}^T C i \sigma_2 \Delta_R \Psi_{R,j} + h.c \quad (2.2)$$

f_L and f_R are the Majorana Yukawa couplings and are equal because of the discrete left-right symmetry. The family indices i, j runs from 1 to 3 representing the three generations of the fermions. $C = i\gamma_2\gamma_0$ is the charge conjugation operator, where γ_μ are the Dirac matrices and $\tilde{\phi} = \tau_2\phi^*\tau_2$.

The symmetry breaking of the gauge group describing the model takes place in two steps. The model gauge group is first broken down to the Standard Model gauge group by the VEV of the scalar triplet Δ_R , and then the Standard Model gauge group is broken down to the electromagnetic gauge group i.e., $U(1)_{em}$ by the VEV of the bidoublet and a tiny vev of the scalar triplet Δ_L .

So the resultant light neutrino mass of LRSM is expressed as a sum of the type-I and type-II seesaw mass terms, given as,

$$M_\nu = M_\nu^I + M_\nu^{II} \quad (2.3)$$

where,

$$M_\nu^I = M_D M_R^{-1} M_D^T \quad (2.4)$$

is the type-I seesaw mass, and type-II seesaw mass is given by

$$M_\nu^{II} = M_{LL} \quad (2.5)$$

M_D is the Dirac mass matrix and M_R is the right-handed Majorana mass matrix, where, $M_R = \sqrt{2}v_R f_R$ and $M_L = \sqrt{2}v_L f_L$. v_R and v_L are the respective VEVs of Δ_R and Δ_L . The magnitudes of the VEVs follows the relation, $|v_L|^2 < |k^2 + k'^2| < |v_R|^2$. In LRSM however, the type-I and type-II mass terms can be expressed in terms of the heavy right-handed Majorana mass matrix, so Eq.(2.3) will follow,

$$M_\nu = M_D M_R^{-1} M_D^T + \gamma \left(\frac{M_W}{v_R} \right)^2 M_{RR} \quad (2.6)$$

where, γ is a dimensionless parameter which is a function of various couplings, appearing in the VEV of the triplet Higgs Δ_L , i.e., $v_L = \gamma \left(\frac{v^2}{v_R} \right)$ and here, $v = \sqrt{k^2 + k'^2}$, and

$$\gamma = \frac{\beta_1 k k' + \beta_2 k^2 + \beta_3 k'^2}{(2\rho_1 - \rho_3)(k^2 + k'^2)} \quad (2.7)$$

In our model, the dimensionless parameter γ has been fine tuned to $\gamma \approx 10^{-13}$ and v_R is of the order of TeV . β_i and ρ_i are dimensionless parameters which appear in the potential of the model

which has been thoroughly elaborated in the appendix.

The light neutrino mass obtained can be expressed in terms of a matrix given as,

$$M_\nu = \begin{pmatrix} M_{LL} & M_D \\ M_D^T & M_{RR} \end{pmatrix} \quad (2.8)$$

This matrix is a 6×6 matrix which can be diagonalized by a unitary matrix as follows,

$$\nu^T M_\nu \nu = \begin{pmatrix} \hat{M}_\nu & 0 \\ 0 & \hat{M}_{RR} \end{pmatrix} \quad (2.9)$$

where, ν represents the diagonalizing matrix of the full neutrino mass matrix, $M_\nu, \hat{M}_\nu = \text{diag}(m_1, m_2, m_3)$, with m_i being the light neutrino masses and $\hat{M}_{RR} = \text{diag}(M_1, M_2, M_3)$, with M_i being the heavy right-handed neutrino masses.

The diagonalizing matrix can be represented as,

$$\nu = \begin{pmatrix} U & S \\ T & V \end{pmatrix} \approx \begin{pmatrix} 1 - \frac{1}{2}RR^\dagger & R \\ -R^\dagger & 1 - \frac{1}{2}R^\dagger R \end{pmatrix} \begin{pmatrix} V_\nu & 0 \\ 0 & V_R \end{pmatrix} \quad (2.10)$$

where, R describes the left-right mixing and is given by,

$$R = M_D M_{RR}^{-1} + O(M_D^3 (M_{RR}^{-1})). \quad (2.11)$$

The matrices U, V, S and T are as follows,

$$U = [1 - \frac{1}{2}M_D M_{RR}^{-1} (M_D M_{RR}^{-1})^\dagger] V_\nu \quad (2.12)$$

$$V = [1 - \frac{1}{2}(M_D M_{RR}^{-1})^\dagger M_D M_{RR}^{-1}] V_\nu \quad (2.13)$$

$$S = M_D M_{RR}^{-1} v_R f_R \quad (2.14)$$

$$T = -(M_D M_{RR}^{-1})^\dagger V_\nu \quad (2.15)$$

In this work, realization of LRSM has been done with the help of A_4 modular symmetry, which has been discussed thoroughly in the next section.

III. REALIZATION OF LRSM WITH MODULAR GROUP $\Gamma(3)$ HAVING WEIGHT 2 .

It has been realized since a long course of time that symmetries do hold an important position in the context of particle physics. LRSM has been constructed with the help discrete flavor symmetries in several earlier works like [22], [23],[24],[25]. In our present work, we have realized the model using A_4 modular symmetry (non-SUSY case). The advantage of using modular symmetry rather than flavor symmetry is that we do not require the use of any flavons within the model and hence it helps in keeping the model minimal. The model contains usual particle content of LRSM [24]. The lepton doublets transform as triplets under A_4 and the bidoublet and scalar triplets transform as 1 under A_4 [26]. As we have considered modular symmetry, we assign modular weights to the particles, keeping in mind that matter multiplets corresponding to the model can have negative modular weights, but the modular forms cannot be assigned negative weights. We assign modular weights to the particles in a manner that the sum of the modular weights of the particles in each term of the Lagrangian sum up to be zero. Modular weights corresponding to each particle is shown in table I. The Yukawa Lagrangian for the leptonic sector in LRSM is given by Eq.(2.1),(2.2) from which we can write the Yukawa Lagrangian of our A_4 modular symmetric LRSM, for the lepton sector, by introducing Yukawa coupling in the form of modular forms Y is given as,

$$\mathcal{L}_y = \bar{l}_L \phi l_R Y + \bar{l}_L \tilde{\phi} l_R Y + l_R^T C i\tau_2 \Delta_R l_R Y + l_L^T C i\tau_2 \Delta_L l_L Y \quad (3.1)$$

Gauge group	l_L	l_R	ϕ	Δ_L	Δ_R
$SU(3)_C$	1	1	1	1	1
$SU(2)_L$	2	1	2	3	1
$SU(2)_R$	1	2	2	1	3
$U(1)_{B-L}$	-1	-1	0	2	2
A_4	3	3	1	1	1
k_I	0	-2	0	-2	2

TABLE I: Charge assignments for the particle content of the model.

The Yukawa couplings $Y = (Y_1, Y_2, Y_3)$ are expressed as modular forms of level 3.

	Y (modular forms)
A_4	3
k_I	2

TABLE II: Charge assignment and modular weight for the corresponding modular Yukawa form for the model.

Table II shows the charge assignment for the modular form of level 3 and its corresponding modular weight. In our work, we are concerned with the mass of the neutrinos and as such, using A_4 modular symmetry and using the multiplication rules for A_4 group [27], we construct mass matrices as given below.

The Dirac mass matrix from the Yukawa Lagrangian has been determined in terms of (Y_1, Y_2, Y_3) as,

$$M_D = v \begin{pmatrix} 2Y_1 & -Y_3 & -Y_2 \\ -Y_2 & -Y_1 & 2Y_3 \\ -Y_3 & 2Y_2 & -Y_1 \end{pmatrix} \quad (3.2)$$

where, v is the VEV of the Higgs bidoublet ϕ .

It is to be noted here that the textures of the Dirac mass matrix and also the right-handed mass matrix has been determined, so it is a fact that the type-I seesaw mass texture can be determined. But, the pivotal investigation in the present study revolves around the assignment of different strengths to the type-II seesaw mass term and hence from there, the value of the type-I seesaw mass has been calculated in order to determine the light neutrino mass.

The right-handed Majorana mass matrix is given by,

$$M_R = v_R \begin{pmatrix} 2Y_1 & -Y_3 & -Y_2 \\ -Y_3 & 2Y_2 & -Y_1 \\ -Y_2 & -Y_1 & 2Y_3 \end{pmatrix} \quad (3.3)$$

where, v_R is the VEV for the scalar triplet Δ_R . As it is seen that the Majorana mass matrix for our model is found to be symmetric in nature as it should be. Under these assumptions for modular symmetric LRSM and the basis that we have considered, our charged lepton mass matrix is also

found to be diagonal.

The type-II seesaw mass is given by,

$$M_\nu^{II} = M_{LL} \quad (3.4)$$

where,

$$M_\nu^{II} = v_L \begin{pmatrix} 2Y_1 & -Y_3 & -Y_2 \\ -Y_3 & 2Y_2 & -Y_1 \\ -Y_2 & -Y_1 & 2Y_3 \end{pmatrix} \quad (3.5)$$

where, v_L is the vev of the scalar triplet Δ_L .

The determination of the values of the modular forms (Y_1, Y_2, Y_3) has been done using the relations as mentioned in the section of numerical analysis and results.

IV. RESONANT LEPTOGENESIS AND NEUTRINOLESS DOUBLE BETA DECAY ($0\nu\beta\beta$) IN TEV SCALE LRSM.

In TeV scale seesaw models, an adequate amount of lepton asymmetry can be generated by the phenomenon of resonant leptogenesis (RL) [28–31]. For RL, two of the right-handed Majorana neutrinos need to be degenerate which has been found to be so in the present work. The presence of RH neutrinos and scalar triplets in the framework of LRSM suggests that their decays can give rise to lepton asymmetry. The net lepton asymmetry will therefore be generated due to the two seesaw terms. So, in this work, we investigate the contribution of both the seesaw terms towards the lepton asymmetry, assigning them different weightages (e.g. 30%, 50% and 70%).

For RL however there is a basic requirement that the a pair of Majorana neutrinos must have a mass difference comparable to their decay widths which will be illustrated later in the work.

The rate at which the RH neutrinos decay is governed by the Yukawa couplings, and is hence given by [32],

$$\Gamma_i = (Y_\nu^\dagger Y_\nu)_{ii} \frac{M_i}{8\pi} \quad (4.1)$$

As already mentioned previously that one important condition for RL is that the mass difference of the two heavy RH neutrinos must be comparable to their decay width, i.e., $M_i - M_j = \Gamma$. In such a case, the CP asymmetry may become very large. The CP violating asymmetry is thus given

by [33],

$$\epsilon_i = \frac{\text{Im}[(Y_\nu^\dagger Y_\nu)_{ij}^2]}{(Y_\nu^\dagger Y_\nu)_{11}(Y_\nu^\dagger Y_\nu)_{22}} \cdot \frac{(M_i^2 - M_j^2)M_i\Gamma_j}{(M_i^2 - M_j^2)^2 + M_i^2\Gamma_j^2} \quad (4.2)$$

The variables i, j run over 1 and 2 and $i \neq j$.

The CP asymmetries ϵ_1 and ϵ_2 can give rise to a net lepton asymmetry, provided the expansion rate of the universe is larger than Γ_1 and Γ_2 . This can further be converted into baryon asymmetry of the universe by $B + L$ violating sphaleron processes.

Now, due to the presence of several new heavy particles in LRSM, along with the standard light neutrino contribution to $0\nu\beta\beta$, several new physics contributions also come into the picture. There are total eight contributions of the phenomenon to LRSM. They are mentioned as under,

- Standard Model contribution to NDBD where the intermediate particles are the W_L bosons and light neutrinos, the process in which the amplitude depends upon the leptonic mixing matrix elements and light neutrino masses.
- Heavy right-handed neutrino contribution in which the mediator particles are the W_L bosons and the amplitude depends upon the mixing between light and heavy neutrinos as well as the mass of the heavy neutrino.
- Light neutrino contribution to NDBD where the intermediate particles are W_R bosons and the amplitude depends upon the mixing between light and heavy neutrinos as well as mass of the right-handed gauge boson W_R .
- Heavy right-handed neutrino contribution where the mediator particles are the W_R bosons. The amplitude of this process is dependent on the elements of the right handed leptonic mixing matrix and mass of the right-handed gauge boson, W_R as well as the mass of the heavy right handed Majorana neutrino.
- Light neutrino contribution from the Feynman diagram mediated by both W_L and W_R , and the amplitude of the process depends upon the mixing between light and heavy neutrinos, leptonic mixing matrix elements, light neutrino masses and the mass of the gauge bosons, W_L and W_R (λ contribution).
- Heavy neutrino contribution from the Feynman diagram mediated by both W_L and W_R , and the amplitude of the process depends upon the right handed leptonic mixing matrix

elements, mixing between the light and heavy neutrinos, also the mass of the gauge bosons, W_L and W_R and the mass of the heavy right handed neutrino (η contribution).

- Scalar triplet contribution (Δ_L) in which the mediator particles are W_L bosons, and the amplitude for the process depends upon the masses of the W_L bosons, left-handed triplet Higgs, as well as their coupling to leptons.
- Right-handed scalar triplet contribution (Δ_R) contribution to NDBD in which the mediator particles are W_R bosons, and the amplitude for the process depends upon the masses of the W_R bosons, right-handed triplet Higgs, Δ_R as well as their coupling to leptons.

In this work, the momentum dependent, that is, λ and η contributions of $0\nu\beta\beta$ have been considered. The standard light neutrino contribution and new physics contributions by the heavy RH neutrino and scalar triplet contribution in modular symmetric LRSM have already been done in one of our previous work [34].

V. NUMERICAL ANALYSIS AND RESULTS

The light neutrino mass in LRSM is given by,

$$M_\nu = M_\nu^I + M_\nu^{II} \quad (5.1)$$

where, the type-I seesaw mass term is,

$$M_\nu^I = M_D M_R^{-1} M_D^T \quad (5.2)$$

and,

$$M_\nu^{II} = M_{LL} \quad (5.3)$$

In this work, we aim to study the strengths of each seesaw terms that contribute towards the light neutrino mass and hence the associated phenomenology. So, we have considered,

$$M_\nu^{II(diag)} = X M_\nu^{(diag)} \quad (5.4)$$

where, the parameter X decides the relative strength of the type-II seesaw mass term towards the resulting light neutrino mass. For example, considering that the type-II mass term contributes

10% towards the light neutrino mass, then,

$$X = 10\% = \frac{10}{100} = 0.1 \quad (5.5)$$

In this work, we have considered the contributions of type-II seesaw mass terms towards the light neutrino mass as, $X = 0.3, 0.5, 0.7$ that is, contribution from type-II seesaw term is considered to be 30%, 50%, 70% respectively.

As we are using modular symmetry in LRSM, so our mass matrices are expressed in terms of the modular forms (Y_1, Y_2, Y_3) . So, the type-II mass matrix is expressed as,

$$M_\nu^{II} = v_L \begin{pmatrix} 2Y_1 & -Y_3 & -Y_2 \\ -Y_3 & 2Y_2 & -Y_1 \\ -Y_2 & -Y_1 & 2Y_3 \end{pmatrix} \quad (5.6)$$

where, v_L is of the order of eV .

Using (5.4), we can determine the values of the modular forms and hence further we can calculate the values of the neutrino masses and take into consideration effective masses of different contributions of $0\nu\beta\beta$ in the present work. In LRSM, from type-II seesaw term, M_R can be written as,

$$M_R = \frac{1}{\gamma} \left(\frac{v_R}{M_{W_L}} \right)^2 M_\nu^{II} \quad (5.7)$$

As already mentioned that γ in this work has been fine tuned to 10^{-13} . We have considered different values of the $SU(2)_R$ breaking scale v_R for our further analysis, namely 1.5, 2.2, 2.5, 3, 5, 10 and 18 TeV. These values will be useful for us to determine the common parameter space of the phenomena being considered like, baryon asymmetry of universe (BAU), contributions of $0\nu\beta\beta$ etc. The mass of the left-handed gauge boson is taken as, $M_{W_L} = 80GeV$.

To proceed further into our study and for the required analysis, firstly we need to determine the values of Yukawa couplings expressed as modular forms (Y_1, Y_2, Y_3) . In order to determine their values, we make use of Eq.(5.4). For that, we first check whether the type-II mass matrix M_ν^{II} is diagonalizable. In the present work, M_ν^{II} is indeed diagonalizable and so we diagonalize the matrix to obtain $M_\nu^{II(diag)}$ in terms of (Y_1, Y_2, Y_3) . The RHS (right-hand side) of Eq.(5.4) contains the parameters whose values are known, that is X is taken to be 0.3, 0.5 and 0.7, also $M_\nu^{diag} = diag(m_1, m_2, m_3)$ is the diagonal light neutrino mass matrix. Equating each of the elements

corresponding to the derived matrices on the RHS and LHS of Eq.(5.4) has helped us obtain the values of (Y_1, Y_2, Y_3) . Now, we substitute these values into the obtained mass matrices so as to determine the required parameters associated with the model. The range of the Yukawa couplings for both normal and inverted hierarchies have been found to be in the order of $O(10^{-4})$ to $O(1)$. We then determine the values of the leptonic CP asymmetry ϵ_1 and ϵ_2 , by using (4.2), where $Y_v = \frac{M_D}{v}$, M_D being the Dirac mass and v is the VEV of the Higgs bidoublet and is in GeV . The decay rates in (4.2) can be calculated using the relation (4.1).

The CP violating asymmetries ϵ_1 and ϵ_2 can give rise to net lepton number asymmetry, provided the expansion rate of the universe is larger than Γ_1 and Γ_2 . The net baryon asymmetry is then calculated using the following relation [35],

$$\eta_B \approx -0.96 \times 10^{-2} \sum_i (k_i \epsilon_i) \quad (5.8)$$

k_i being the efficiency factors measuring the washout effects. Some parameters are needed to defined as,

$$K_i \equiv \frac{\Gamma_i}{H} \quad (5.9)$$

Equation (5.9) is defined at temperature $T = M_i$. The Hubble's constant is given by, $H \equiv \frac{1.66\sqrt{g_*}T^2}{M_{Planck}}$, where, $g_* = 107$ and $M_{Planck} = 1.2 \times 10^{19}$ GeV is the Planck mass. decay width is estimated using (4.1). The efficiency factors k_i can be calculated using the formula [36],

$$k_1 \equiv k_2 \equiv \frac{1}{2}(\sum_i K_i)^{-1.2} \quad (5.10)$$

Equation (5.10) holds valid for two nearly degenerate heavy Majorana masses and in most general cases $5 \leq K_i \leq 100$. To calculate the baryon asymmetry parameter, we have used formula (5.8). The results are shown as a variation of the effective neutrino masses calculated for momentum dependent contributions and heavy right handed neutrino contribution as described below. The variations are shown for different values of M_{W_R} and also variations are shown for different strengths of the seesaw mass terms.

Before delving into the results showing the connection between lepton asymmetry and $0\nu\beta\beta$, we have first determined the value of η_B corresponding to a range of values of M_{W_R} for both normal and inverted hierarchy and also for $X = 0.3, 0.5, 0.7$ as shown in figures 1 and 2.





FIG. 1: Variation of η_B with M_{W_R} for normal hierarchy(NH) for $X = 0.3, 0.5, 0.7$. The black line indicates the Planck limit on η_B .






FIG. 2: Variation of η_B with M_{W_R} for inverted hierarchy(IH) for $X = 0.3, 0.5, 0.7$. The black line indicates the Planck limit on η_B .

Talking about neutrinoless double beta decay ($0\nu\beta\beta$), in this work, the momentum dependent mechanisms namely, λ and η contributions have been taken into consideration, which are given by the following formulae,

- For λ contribution, the dimensionless parameter η_λ is given by,

$$|\eta_\lambda| = \left(\frac{M_{W_L}}{M_{W_R}} \right)^2 |\Sigma_i U_{ei} T_{ei}^*| \quad (5.11)$$

- For η contribution, the dimensionless parameter describing $0\nu\beta\beta$ is given by,

$$|\eta_\eta| = \tan \xi |\Sigma_i U_{ei} T_{ei}^*| \quad (5.12)$$

In the above equations, U_{ei} represents the first row of the neutrino mixing matrix. $|\Sigma_i U_{ei} T_{ei}^*|$ can be simplified to the form $-[M_D M_{RR}^{-1}]_{ee}$ as described in [37]. T is represented by the equation (2.15) and,


$$\tan 2\xi = -\frac{2k_1 k_2}{v_R^2 - v_L^2} \quad (5.13)$$

The heavy right-handed (RH) neutrino contribution for $0\nu\beta\beta$ is given by,

$$M_{eff}^N = p^2 \left(\frac{M_{W_L}^4}{M_{W_R}^4} \right) \frac{U_{Rei}^2}{M_i} \quad (5.14)$$

where, $\langle p^2 \rangle = m_e m_p \frac{M_N}{M_\nu}$ is the typical momentum exchange of the process. m_e and m_p are the masses of the electron and proton respectively and M_N is the nuclear matrix element (NME) for

the right-handed neutrino exchange. The allowed value of p is in the range (100 – 200) MeV. But for our analysis we have used $p = 180\text{MeV}$ [38]. Figure 3,4 and 5 below shows the variation of baryon asymmetry parameter with the effective mass for heavy right-handed neutrino contribution for right-handed gauge boson masses 3,5,10 and 18TeV for type-II seesaw mass term contributing 30%, 50% and 70% to light neutrino mass respectively.







FIG. 3: Variation of η_B with effective masses corresponding to heavy right-handed neutrino contribution for $M_{W_R} = 3, 5, 10$ and 18TeV respectively for normal hierarchy(NH) when $X = 0.3$.







FIG. 4: Variation of η_B with effective masses corresponding to heavy right-handed neutrino contribution for $M_{W_R} = 3, 5, 10$ and 18 TeV respectively for NH and IH when $X = 0.5$.






FIG. 5: Variation of η_B with effective masses corresponding to heavy right-handed neutrino contribution for $M_{W_R} = 3, 5, 10$ and 18TeV respectively for NH and IH when $X = 0.7$.

The effective mass for λ contribution, that is, $|m_{eff}^\lambda|$ is expressed in terms of the dimensionless parameter η_λ as,

$$|\eta_\lambda| = \frac{|m_{eff}^\lambda|}{m_e} \quad (5.15)$$

The variations of baryon asymmetry parameter with the effective masses corresponding to the aforementioned contribution are shown in figures 6 to 8.




FIG. 6: Variation of effective mass with baryon asymmetry parameter for NH and IH and $X=0.3$.




FIG. 7: Variation of effective mass with baryon asymmetry parameter for NH and IH and $X=0.5$.




FIG. 8: Variation of effective mass with baryon asymmetry parameter for NH and IH and $X=0.7$.

The effective mass for η contribution, that is, $|m_{eff}^\eta|$ is expressed in terms of the dimensionless

parameter η_η as,

$$|\eta_\eta| = \frac{|m_{eff}^\eta|}{m_e} \quad (5.16)$$




FIG. 9: Variation of effective mass with baryon asymmetry parameter for NH and IH and $X=0.3$.




FIG. 10: Variation of effective mass with baryon asymmetry parameter for NH and IH and $X=0.5$.




FIG. 11: Variation of effective mass with baryon asymmetry parameter for NH and IH and $X=0.7$.

Figures 9 to 11 shows the variation of baryon asymmetry parameter with the effective masses for η contribution of $0\nu\beta\beta$.





FIG. 12: Variation of effective mass with baryon asymmetry parameter for NH and $X=0.3, 0.5$ and 0.7 .

Figure 12 shows the variation of baryon asymmetry parameter with the effective mass corresponding to right-handed neutrino contribution for 1.5, 2.2 and 2.5 TeV for normal hierarchy.







FIG. 13: Variation of effective mass for heavy right-handed neutrino contribution with baryon asymmetry parameter for IH and $X=0.3, 0.5$ and 0.7 .

Figure 13 shows the variation of baryon asymmetry parameter with the effective mass corresponding to right-handed neutrino contribution for $1.5, 2.2$ and 2.5 TeV for inverted hierarchy. It has been observed that the heavy right-handed neutrino corresponding to $X = 0.3$ and $X = 0.5$ is very less in case of $1.5, 2.2$ and 2.5 TeV . But when $X = 0.7$ that is, when the type-II seesaw mass term is dominant, there are some amount of data obtained within the experimental limits. Same has been observed in case of lambda and eta contributions of $0\nu\beta\beta$ as well and as such, we are showing only the plots corresponding to $X = 0.7$.






FIG. 14: Variation of effective mass for η contribution with baryon asymmetry parameter for NH and $X=0.7$.

Figure 14 shows the variation of effective mass corresponding to η contribution with the baryon asymmetry parameter for the type-II seesaw mass contributing 70% towards the neutrino mass. The plots are for normal hierarchy. The figure 15 shows the variations for inverted hierarchy.




FIG. 15: Variation of effective mass for η contribution with baryon asymmetry parameter for IH and $X=0.7$.

Variations of baryon asymmetry parameter with effective mass corresponding to λ contribution have also been shown in figures 16 and 17.





FIG. 16: Variation of effective mass for λ contribution with baryon asymmetry parameter for NH and $X=0.7$.






FIG. 17: Variation of effective mass for λ contribution with baryon asymmetry parameter for IH and $X=0.7$.

VI. DISCUSSION

The following points describe the complete study and analysis of the present work and the results obtained hereafter,

- Left-Right Symmetric model has been realized with the help of modular group of level 3 and weight 2 ($\Gamma(3)$), isomorphic to the A_4 non-abelian discrete symmetry group, as discussed in the appendix.
- The aim of the study focusses on describing the variation of baryogenesis (via leptogenesis) with variations in the value of the right-handed gauge boson mass and strength of the type-II seesaw mass.
- In several previous literatures, it was stated that while taking into consideration resonant leptogenesis, the dominant contribution was given by type-I seesaw mass, however, we have also taken into consideration the strength of the type-II seesaw mass term and it has been found that the analysis gives considerable results for the same.
- We have considered strength of the type-II seesaw mass to be 30% , 50% and 70%, and masses of the right-handed gauge bosons are considered to be $M_{W_R} = 1.5, 2.2, 2.5, 3, 5, 10, 18$ TeV. Here, the condition $M_N \leq M_{W_R}$ has been used.
- From the present analysis, it can be stated that a lower bound of $M_{W_R} \geq 1.5$ TeV has been

obtained for the case $g_L = g_R$, that is, when the gauge couplings corresponding to left-handed and right-handed components are equal. In the present work, the Yukawa couplings have been expressed in terms of modular forms (Y_1, Y_2, Y_3) and the values determined for the same have been found to be much larger than the canonical value of $O(10^{-5.5})$ [31, 39]. Due to the enhanced Yukawa couplings, the reaction rates of the two body decay becomes more dominant over the W_R mediated scatterings as well as the three body decays [39]. In the present model, the CP asymmetry parameter has also been numerically investigated rather than considering it as one.

As stated in [40], we can take into consideration two different ways of deriving a lower bound on the value of M_{W_R} . The first one being the careful scanning of the parameter space of the model and the second one being the use of the approximate expression 6.1 [39, 40],

$$|\eta^{\Delta L}(z_c)| = \frac{9}{4\zeta(3)} \frac{z_c^2 K_1(z_c)}{S_R(z_c)} \frac{r_s r_d^2}{(3 + r_s r_d)(3 + r_s)} \epsilon_{total}^Y \quad (6.1)$$

where, r_s and r_d are the relative washout strengths of the two-body decays, W_R mediated decays and scatterings. $\zeta(x)$ is the Riemann function, $K_1(z_c)$ is the first order Bessel function and $z = \frac{m_{N_1}}{T}$, T being the temperature of the Universe, is a dimensionless variable, and z_c is the critical value beyond which the sphaleron processes become ineffective [39, 40]. The value of these strengths depend upon the values of M_N , M_{W_R} and most importantly the Yukawa couplings. we have used the first method to determine the lower bound on M_{W_R} , which was found to be $M_{W_R} \geq 1.5 TeV$. As stated in the paper, [39], a dependence on the Yukawa couplings of the decay rates can be mathematically expressed as $D \equiv Y^2$ which suggests that with larger value of Yukawa couplings, the decay rate will be larger as compared to that of the scatterings induced due to RH currents.

- The analysis shows that the weaker bounds on the M_{W_R} mass is applicable only in the case where the type-II term contributes dominantly. This might be the result of the values of the Yukawa couplings and textures of the mass matrices obtained as a result of implementing modular symmetry in LRSM rather than flavor symmetry.
- As already stated that, a lower bound has been found through the present study on M_{W_R} , but we have also calculated the baryon asymmetry parameter for values less than and also greater than $1.5 TeV$, but the points are more specific only when $M_{W_R} \geq 1.5 TeV$. This lower

bound be a result of the larger Yukawa couplings obtained in the present work ranging from 10^{-4} to 10^{-1} and some maximum values are also greater than 1.

- We have also shown the variations of η_B with M_{W_R} to emphasize on the fact that even though the range of M_{W_R} is taken to be quite high, but successful explanation of leptogenesis and its connection with the low energy phenomena, that is, Neutrinoless Double Beta Decay can be put forward, only for certain values of X (most preferably $X > 0.5$) that is when the type-II seesaw mass term is dominant and only at specific values of M_{W_R} . The results are found to be more reliable for inverted hierarchy as compared to that of normal hierarchy and for the range $1.5 \leq M_{W_R} \leq 2.5$, the values predicted by the model for baryon asymmetry are found to be consistent with the observed value for both normal and inverted hierarchy.

Several previous works have been done taking into consideration the flavor symmetric realization of LRSM as studied in [23–25, 41] etc. The authors in the mentioned literatures have also carried out phenomenological studies related to the model. However, in the present work we have not considered flavor symmetry but, we have taken into consideration modular symmetry. The use of modular symmetry provides the advantage of maintaining the minimality in construction of the model without any flavons. In the present work, we have considered modular group of level 3, $\Gamma(3)$ and weight 2, which is isomorphic to A_4 non-abelian discrete symmetry group. Here, it is to be noted that the present model has been designed in the non-SUSY framework and as such, Left-Right Symmetric Model does not restrict the use of infinite number of modular forms, but as we are using $\Gamma(3)$ modular group having weight 2, as stated in the appendix in table III, we will have three number of modular forms represented as (Y_1, Y_2, Y_3) . As such, the mass matrices corresponding to the present model have been expressed in terms of these three modular forms, discussed in section (III) of the manuscript.

Some previous work as discussed in [32, 42] have also taken into account the study of several phenomenology like $0\nu\beta\beta$, Lepton Flavor Violation (LFV) with changing values of M_{W_R} . In the present work, we have also taken into account the different values of M_{W_R} to study the relation between $0\nu\beta\beta$ and leptogenesis, however it has been done taking into account different strengths of the type-II seesaw mass term. One of the most significant results of the present work can be stated to be the observation of considerable amount of leptogenesis phenomena taking place when the leading contribution of neutrino mass is given by the type-II seesaw mass term (70%).

As stated by the authors in [32] that while taking into consideration the phenomena of resonant leptogenesis, type-I seesaw term plays the dominant role and the contribution of Δ_L is negligible. But, in the present work, type-II seesaw mass term shows considerable contribution towards the said phenomenology. The exact reason although not known, this may be thought of as a result of emerging modular forms in the picture which changes the structure of the mass matrices within the model.

VII. CONCLUSION

The experimental verification of neutrino oscillations paved the gateway for physics beyond the Standard Model. In the present work, LRSM has been realized with the help of modular A_4 symmetry, the advantage being, no extra particles called 'flavons'. The Yukawa couplings are represented as modular forms expressed as expansions of q . The values of the Yukawa couplings (Y_1, Y_2, Y_3) are calculated using the relation $M_\nu^{II(diag)} = X M_\nu^{(diag)}$. The mass matrix for the type-II seesaw dominance has been determined using the multiplication rules for A_4 group stated in the Appendix. The Majorana mass matrix is found to be symmetric and under the considered basis, the charged lepton mass matrix is also diagonal. To summarize our work, some results are stated as under,

- In the present work we have considered the arbitrary constant that can be multiplied with the Yukawa term to be one. This is to reduce the number of free parameters in the model and to determine the data more accurately.
- The model has been realized using A_4 modular symmetry, the advantage being no use of extra particles or flavons. We can also use other symmetry groups depending upon the level of modular form used and as such, the number of modular forms will also change, as per the table III in the manuscript. As we have used A_4 modular symmetry, there are three modular forms (Y_1, Y_2, Y_3) and hence it became easier for us to construct the 3×3 mass matrices in terms of the aforementioned modular forms which are consistent with the neutrino data.
- The Yukawa couplings determined for the cases when $X = 0.3, 0.5, 0.7$ has been found to satisfy the condition that its modulus is greater than unity.

- Effective masses corresponding to heavy right-handed neutrino contribution, and momentum dependent contributions of $0\nu\beta\beta$ are determined and their variations with the baryon asymmetry parameter η_B has been studied. The results are found to satisfy the experimental constraints, for both normal and inverted hierarchies.

The effective masses for the $0\nu\beta\beta$ contributions are calculated and it has been found that the values for the effective mass corresponding to each contribution is well within the experimental bounds, which in fact makes us clearly state that the building of the model with modular symmetry is advantageous to that of flavor symmetries. In this model, no extra particles have been used and the calculations are carried out taking into picture the usual particle content of the model and hence conclusively it brings to our attention that different phenomenology in the Beyond Standard Model framework can be studied using modular symmetry that helps in keeping the model minimal.

APPENDIX A : MINIMISATION OF THE POTENTIAL TERM.

Let us consider the Higgs potential of our model that has quadratic and quartic coupling terms given by [43],

$$\begin{aligned}
V_{\phi,\Delta_L,\Delta_R} = & -\mu_{ij}^2 Tr[\phi_i^\dagger \phi_j] + \lambda_{ijkl} Tr[\phi_i^\dagger \phi_j] Tr[\phi_k^\dagger \phi_l] + \lambda'_{ijkl} Tr[\phi_i^\dagger \phi_j \phi_k^\dagger \phi_l] - \mu_{ij}^2 Tr[\Delta_L^\dagger \Delta_L + \Delta_R^\dagger \Delta_R] \\
& \rho_1 [(Tr[\Delta_L^\dagger \Delta_L])^2 + (Tr[\Delta_R^\dagger \Delta_R])^2] + \rho_2 (Tr[\Delta_L^\dagger \Delta_L \Delta_L^\dagger \Delta_L] + Tr[\Delta_R^\dagger \Delta_R \Delta_R^\dagger \Delta_R]) + \rho_3 (Tr[\Delta_L^\dagger \Delta_L] Tr[\Delta_R^\dagger \Delta_R]) + \\
& \alpha_{ij} Tr[\phi_i^\dagger \phi_j] (Tr[\Delta_L^\dagger \Delta_L] + Tr[\Delta_R^\dagger \Delta_R]) + \beta_{ij} (Tr[\Delta_L^\dagger \Delta_L \phi_i \phi_j^\dagger] + (Tr[\Delta_R^\dagger \Delta_R \phi_i \phi_j^\dagger])) + \gamma_{ij} (Tr[\Delta_L^\dagger \phi_i \Delta_R \phi_j^\dagger] + h.c)
\end{aligned} \tag{7.1}$$

where, i,j,k,l runs from 1 to 2 with $\phi_1 = \phi$ and $\phi_2 = \tilde{\phi}$. As mentioned above after SSB, the scalar sector obtains VEV. So after the substitution of the respective VEVs and determining the traces, so after simplification the potential can be written as,

$$V = -\mu^2(v_L^2 + v_R^2) + \frac{\rho}{4}(v_L^4 + v_R^4) + \frac{\rho'}{2} + \frac{\alpha}{2}(v_L^2 + v_R^2)k^2 + \gamma v_L v_R k^2 \tag{7.2}$$

where, we have used the approximation $k' \ll k$, and $\rho' = 2\rho_3$. Our minimization conditions are, $\frac{\delta V}{\delta v_L} = \frac{\delta V}{\delta v_R} = \frac{\delta V}{\delta k} = \frac{\delta V}{\delta k'} = 0$

Therefore, we get,

$$\frac{\delta V}{\delta v_L} = -2\mu^2 v_L + \rho v_L^3 + \rho' v_L k^2 + \gamma v_R k^2 \tag{7.3}$$

Here, it is evident that the Majorana mass of the left-handed neutrino M_{LL} is dependent on the vev v_L as already defined above. Again, we have

$$\frac{\delta V}{\delta v_R} = -2\mu^2 v_R + \rho v_R^3 + \rho' v_R k^2 + \gamma v_L k^2 \quad (7.4)$$

So, the right handed Majorana mass M_{RR} is dependent on the vev v_R . Similarly, the calculations for the same can be carried out and it can be found out the Dirac mass term M_D can be expressed in terms of the vev for the Higgs bidoublet as also defined previously.

Now, we are to determine a relation between the VEVs for the scalars and so after using the minimization conditions and simplifying the equations, we come to a relation given by,

$$v_L v_R = \frac{\gamma}{\xi} k \quad (7.5)$$

where, $\xi = \rho - \rho'$.

The neutrino mass for LRSM is given as a summation of the type-I and type-II term as already mentioned above. So, in the approximation that $k' \ll k$, and if we consider that our Yukawa coupling Y^l corresponding to the neutrino masses is y_D and the coupling \widetilde{Y}^l for the charged fermion masses is denoted by y_L , so considering $y_D k \gg y_L k'$ we can write,

$$M_\nu = \frac{k^2}{v_R} y_D f^{-1} y_D^T + f_L v_L \quad (7.6)$$

Since, for due to left-right symmetry, we can consider $f_L = f_R = f$, so the above equation can be written as,

$$M_\nu = \frac{k^2}{v_R} y_D f^{-1} y_D^T + f v_L \quad (7.7)$$

So, from this equation we can come to a relation given by,

$$M_\nu = \left(f \frac{\gamma}{\xi} + y_D f^{-1} y_D^T \right) \frac{k^2}{v_R} \quad (7.8)$$

Here, we can consider two situations, namely

- If $f \frac{\gamma}{\xi} \ll y_D f^{-1} y_D^T$, the light neutrino mass is given by the type-I term $M_D M_{RR}^{-1} M_D^T$. That is, here type-I is dominant and the light neutrino mass is from the suppression of heavy ν_R .
- If $f \frac{\gamma}{\xi} \gg y_D f^{-1} y_D^T$, the light neutrino mass is given by the type-II term $f v_L$. That is, in this case type-II mass term is dominant and the light neutrino mass is because of the tiny value of ν_L .

APPENDIX B : PROPERTIES OF A_4 DISCRETE SYMMETRY GROUP.

A_4 is a non-abelian discrete symmetry group which represents even permutations of four objects. It has four irreducible representations, three out of which are singlets ($1, 1', 1''$) and one triplet 3 (3_A represents the anti-symmetric part and 3_S the symmetric part). Products of the singlets and triplets are given by,

$$1 \otimes 1 = 1$$

$$1' \otimes 1' = 1''$$

$$1' \otimes 1'' = 1$$

$$1'' \otimes 1'' = 1'$$

$$3 \otimes 3 = 1 \oplus 1' \oplus 1'' \oplus 3_A \oplus 3_S$$

If we have two triplets under A_4 say, (a_1, a_2, a_3) and (b_1, b_2, b_3) , then their multiplication rules are given by,

$$1 \approx a_1 b_1 + a_2 b_3 + a_3 b_2$$

$$1' \approx a_3 b_3 + a_1 b_2 + a_2 b_1$$

$$1'' \approx a_2 b_2 + a_3 b_1 + a_1 b_3$$

$$3_S \approx \begin{pmatrix} 2a_1b_1 - a_2b_3 - a_3b_2 \\ 2a_3b_3 - a_1b_2 - a_2b_1 \\ 2a_2b_2 - a_1b_3 - a_3b_1 \end{pmatrix}$$

$$3_A \approx \begin{pmatrix} a_2b_3 - a_3b_2 \\ a_1b_2 - a_2b_1 \\ a_3b_1 - a_1b_3 \end{pmatrix}$$

APPENDIX C : MODULAR SYMMETRY.

Modular symmetry has gained much importance in aspects of model building [44], [45]. This is because it can minimize the extra particle called 'flavons' while analyzing a model with respect to a particular symmetry group. An element q of the modular group acts on a complex variable τ which belongs to the upper-half of the complex plane given as [45] [46]

$$q\tau = \frac{a\tau + b}{c\tau + d} \quad (7.9)$$

where a, b, c, d are integers and $ad - bc = 1$, $\text{Im}\tau > 0$.

The modular group is isomorphic to the projective special linear group $\text{PSL}(2, \mathbb{Z}) = \text{SL}(2, \mathbb{Z}) / Z_2$ where, $\text{SL}(2, \mathbb{Z})$ is the special linear group of integer 2×2 matrices having determinant unity and $Z_2 = (I, -I)$ is the centre, I being the identity element. The modular group can be represented in terms of two generators S and T which satisfies $S^2 = (ST)^3 = I$. S and T satisfies the following matrix representations:

$$S = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad (7.10)$$

$$T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \quad (7.11)$$

corresponding to the transformations,

$$S : \tau \rightarrow -\frac{1}{\tau}; T : \tau \rightarrow \tau + 1 \quad (7.12)$$

Finite modular groups ($N \leq 5$) are isomorphic to non-abelian discrete groups, for example, $\Gamma(3) \approx A_4$, $\Gamma(2) \approx S_3$, $\Gamma(4) \approx S_4$. While using modular symmetry, the Yukawa couplings can be expressed

in terms of modular forms, and the number of modular forms present depends upon the level and weight of the modular form. For a modular form of level N and weight $2k$, the table below shows the number of modular forms associated within and the non-abelian discrete symmetry group to which it is isomorphic [46].

N	No. of modular forms	$\Gamma(N)$
2	$k + 1$	S_3
3	$2k + 1$	A_4
4	$4k + 1$	S_4
5	$10k + 1$	A_5
6	$12k$	
7	$28k - 2$	

TABLE III: No. of modular forms corresponding to modular weight $2k$.

In our work, we will be using modular form of level 3, that is, $\Gamma(3)$ which is isomorphic to A_4 discrete symmetry group. The weight of the modular form is taken to be 2, and hence it will have three modular forms (Y_1, Y_2, Y_3) which can be expressed as expansions of q given by,

$$Y_1 = 1 + 12q + 36q^2 + 12q^3 + 84q^4 + 72q^5 + 36q^6 + 96q^7 + 180q^8 + 12q^9 + 216q^{10} \quad (7.13)$$

$$Y_2 = -6q^{1/3}(1 + 7q + 8q^2 + 18q^3 + 14q^4 + 31q^5 + 20q^6 + 36q^7 + 31q^8 + 56q^9) \quad (7.14)$$

$$Y_3 = -18q^{2/3}(1 + 2q + 5q^2 + 4q^3 + 8q^4 + 6q^5 + 14q^6 + 8q^7 + 14q^8 + 10q^9) \quad (7.15)$$

where, $q = \exp(2\pi i\tau)$.

ACKNOWLEDGEMENTS

Ankita Kakoti acknowledges Department of Science and Technology (DST), India (grant DST/INSPIRE Fellowship/2019/IF190900) for the financial assistantship.

REFERENCES

- [1] E. W. Kolb and S. Wolfram, “The Development of Baryon Asymmetry in the Early Universe,” *Phys. Lett. B* **91** (1980) 217–221.
- [2] G. Altarelli, F. Feruglio, L. Merlo, and E. Stamou, “Discrete Flavour Groups, θ_{13} and Lepton Flavour Violation,” *JHEP* **08** (2012) 021, [arXiv:1205.4670 \[hep-ph\]](#).
- [3] M. Taoso, G. Bertone, and A. Masiero, “Dark Matter Candidates: A Ten-Point Test,” *JCAP* **03** (2008) 022, [arXiv:0711.4996 \[astro-ph\]](#).
- [4] M. S. Turner, “Dark matter: Theoretical perspectives,” *Proc. Nat. Acad. Sci.* **90** (1993) 4827, [arXiv:astro-ph/9302003](#).
- [5] P. Minkowski, “ $\mu \rightarrow e\gamma$ at a Rate of One Out of 10^9 Muon Decays?,” *Phys. Lett. B* **67** (1977) 421–428.
- [6] J. Schechter and J. W. F. Valle, “Neutrino Masses in $SU(2) \times U(1)$ Theories,” *Phys. Rev. D* **22** (1980) 2227.
- [7] R. N. Mohapatra and G. Senjanovic, “Neutrino Mass and Spontaneous Parity Nonconservation,” *Phys. Rev. Lett.* **44** (1980) 912.
- [8] C. Wetterich, “Neutrino Masses and the Scale of B-L Violation,” *Nucl. Phys. B* **187** (1981) 343–375.
- [9] S. Antusch and S. F. King, “Type II Leptogenesis and the neutrino mass scale,” *Phys. Lett. B* **597** (2004) 199–207, [arXiv:hep-ph/0405093](#).
- [10] R. Foot, H. Lew, X. G. He, and G. C. Joshi, “Seesaw Neutrino Masses Induced by a Triplet of Leptons,” *Z. Phys. C* **44** (1989) 441.
- [11] N. Haba, H. Ishida, and Y. Yamaguchi, “Naturalness and lepton number/ flavor violation in inverse seesaw models,” *JHEP* **11** (2016) 003, [arXiv:1608.07447 \[hep-ph\]](#).
- [12] G. Senjanovic, “Spontaneous Breakdown of Parity in a Class of Gauge Theories,” *Nucl. Phys. B* **153** (1979) 334–364.
- [13] G. Senjanovic and V. Tello, “Disentangling the seesaw mechanism in the minimal left-right

- symmetric model,” *Phys. Rev. D* **100** no. 11, (2019) 115031, [arXiv:1812.03790 \[hep-ph\]](#).
- [14] P. S. Bhupal Dev, R. N. Mohapatra, W. Rodejohann, and X.-J. Xu, “Vacuum structure of the left-right symmetric model,” *JHEP* **02** (2019) 154, [arXiv:1811.06869 \[hep-ph\]](#).
- [15] W. Grimus, “Introduction to left-right symmetric models,” in *4th Hellenic School on Elementary Particle Physics*, pp. 619–632. 3, 1993.
- [16] R. N. Mohapatra and J. C. Pati, “A Natural Left-Right Symmetry,” *Phys. Rev. D* **11** (1975) 2558.
- [17] J. C. Pati and A. Salam, “Lepton Number as the Fourth Color,” *Phys. Rev. D* **10** (1974) 275–289. [Erratum: *Phys.Rev.D* 11, 703–703 (1975)].
- [18] G. Senjanovic and R. N. Mohapatra, “Exact Left-Right Symmetry and Spontaneous Violation of Parity,” *Phys. Rev. D* **12** (1975) 1502.
- [19] M. Pernow, *Phenomenology of $SO(10)$ Grand Unified Theories*. PhD thesis, Royal Inst. Tech., Stockholm, 2019.
- [20] J. D. Vergados, H. Ejiri, and F. Simkovic, “Theory of Neutrinoless Double Beta Decay,” *Rept. Prog. Phys.* **75** (2012) 106301, [arXiv:1205.0649 \[hep-ph\]](#).
- [21] L. Cardani, “Neutrinoless Double Beta Decay Overview,” *SciPost Phys. Proc.* **1** (2019) 024, [arXiv:1810.12828 \[nucl-ex\]](#).
- [22] P. Duka, J. Gluza, and M. Zralek, “Quantization and renormalization of the manifest left-right symmetric model of electroweak interactions,” *Annals Phys.* **280** (2000) 336–408, [arXiv:hep-ph/9910279](#).
- [23] B. B. Boruah and M. K. Das, “Realization of Left-Right Symmetric Model by Discrete Flavor Symmetries,” *Springer Proc. Phys.* **265** (2022) 285–291.
- [24] P. Sahu, S. Patra, and P. Pritimita, “ A_4 realization of left-right symmetric linear seesaw,” [arXiv:2002.06846 \[hep-ph\]](#).
- [25] W. Rodejohann and X.-J. Xu, “A left–right symmetric flavor symmetry model,” *Eur. Phys. J. C* **76** no. 3, (2016) 138, [arXiv:1509.03265 \[hep-ph\]](#).
- [26] M. Abbas, “Fermion masses and mixing in modular A_4 Symmetry,” *Phys. Rev. D* **103** no. 5, (2021) 056016, [arXiv:2002.01929 \[hep-ph\]](#).
- [27] G.-J. Ding, S. F. King, and A. J. Stuart, “Generalised CP and A_4 Family Symmetry,” *JHEP* **12** (2013) 006, [arXiv:1307.4212 \[hep-ph\]](#).

- [28] T. Asaka and T. Yoshida, “Resonant leptogenesis at TeV-scale and neutrinoless double beta decay,” *JHEP* **09** (2019) 089, [arXiv:1812.11323 \[hep-ph\]](#).
- [29] S. Blanchet, Z. Chacko, S. S. Granor, and R. N. Mohapatra, “Probing Resonant Leptogenesis at the LHC,” *Phys. Rev. D* **82** (2010) 076008, [arXiv:0904.2174 \[hep-ph\]](#).
- [30] M. Flanz, E. A. Paschos, U. Sarkar, and J. Weiss, “Baryogenesis through mixing of heavy Majorana neutrinos,” *Phys. Lett. B* **389** (1996) 693–699, [arXiv:hep-ph/9607310](#).
- [31] P. S. B. Dev, “TeV Scale Leptogenesis,” *Springer Proc. Phys.* **174** (2016) 245–253, [arXiv:1506.00837 \[hep-ph\]](#).
- [32] H. Borgohain and M. K. Das, “Lepton number violation, lepton flavor violation, and baryogenesis in left-right symmetric model,” *Phys. Rev. D* **96** no. 7, (2017) 075021, [arXiv:1709.09542 \[hep-ph\]](#).
- [33] Z.-z. Xing and Z.-h. Zhao, “A review of μ - τ flavor symmetry in neutrino physics,” *Rept. Prog. Phys.* **79** no. 7, (2016) 076201, [arXiv:1512.04207 \[hep-ph\]](#).
- [34] A. Kakoti, B. B. Boruah, and M. K. Das, “Minimal left–right symmetric model with A4 modular symmetry,” *Int. J. Mod. Phys. A* **38** no. 28, (2023) 2350150, [arXiv:2301.13552 \[hep-ph\]](#).
- [35] W. Buchmuller, P. Di Bari, and M. Plumacher, “Some aspects of thermal leptogenesis,” *New J. Phys.* **6** (2004) 105, [arXiv:hep-ph/0406014](#).
- [36] S. Blanchet and P. Di Bari, “New aspects of leptogenesis bounds,” *Nucl. Phys. B* **807** (2009) 155–187, [arXiv:0807.0743 \[hep-ph\]](#).
- [37] J. Barry and W. Rodejohann, “Lepton number and flavour violation in TeV-scale left-right symmetric theories with large left-right mixing,” *JHEP* **09** (2013) 153, [arXiv:1303.6324 \[hep-ph\]](#).
- [38] J. Chakraborty, H. Z. Devi, S. Goswami, and S. Patra, “Neutrinoless double- β decay in TeV scale Left-Right symmetric models,” *JHEP* **08** (2012) 008, [arXiv:1204.2527 \[hep-ph\]](#).
- [39] P. S. Bhupal Dev, C.-H. Lee, and R. N. Mohapatra, “Leptogenesis Constraints on the Mass of Right-handed Gauge Bosons,” *Phys. Rev. D* **90** no. 9, (2014) 095012, [arXiv:1408.2820 \[hep-ph\]](#).
- [40] P. S. Bhupal Dev, C.-H. Lee, and R. N. Mohapatra, “TeV Scale Lepton Number Violation and Baryogenesis,” *J. Phys. Conf. Ser.* **631** no. 1, (2015) 012007, [arXiv:1503.04970 \[hep-ph\]](#).
- [41] B. B. Boruah and M. K. Das, “Neutrinoless double-beta decay and lepton flavor violation in discrete flavor symmetric left–right symmetric model,”

- Int. J. Mod. Phys. A* **37** no. 06, (2022) 2250026, [arXiv:2111.10341 \[hep-ph\]](#).
- [42] H. Borgohain and M. K. Das, “Neutrinoless double beta decay and lepton flavour violation in broken $\mu - \tau$ symmetric neutrino mass models,” *Int. J. Theor. Phys.* **56** no. 9, (2017) 2911–2934, [arXiv:1705.00922 \[hep-ph\]](#).
- [43] M. J. Luo and Q. Y. Liu, “Small Neutrino Masses From Structural cancellation In Left-Right Symmetric Model,” *JHEP* **12** (2008) 061, [arXiv:0812.3453 \[hep-ph\]](#).
- [44] S. J. D. King and S. F. King, “Fermion mass hierarchies from modular symmetry,” *JHEP* **09** (2020) 043, [arXiv:2002.00969 \[hep-ph\]](#).
- [45] P. P. Novichkov, J. T. Penedo, S. T. Petcov, and A. V. Titov, “Generalised CP Symmetry in Modular-Invariant Models of Flavour,” *JHEP* **07** (2019) 165, [arXiv:1905.11970 \[hep-ph\]](#).
- [46] F. Feruglio, *Are neutrino masses modular forms?*, pp. 227–266. 2019. [arXiv:1706.08749 \[hep-ph\]](#).