The electroweak magnetic monopole in the presence of KSVZ axion

Tong Li^{1,*} and Rui-Jia Zhang^{1,†}

¹School of Physics, Nankai University, Tianjin 300071, China

Abstract

The Witten effect implies the dynamics of axion and magnetic monopole. The Cho-Maison monopole is a realistic electroweak monopole arisen in the Weinberg-Salam theory. This monopole of TeV scale mass motivates the dedicated search for electroweak monopole at colliders. In this work we investigate the implication of KSVZ axion to the electroweak magnetic monopole. We use the spherically symmetric ansatz for the electroweak dyon and introduce the spherically symmetric function for the axion field. The effective Lagrangian is then shown in terms of the electroweak monopole part, the axion kinetic energy as well as the axion interaction term. We derive the consequent equations of motion in the presence of the axion-photon coupling and show the numerical results of the topological solutions. We then calculate the changed characteristics of the electroweak monopole such as the monopole mass and the electromagnetic charges, as well as the axion potential energy.

^{*} litong@nankai.edu.cn

[†] zhangruijia@mail.nankai.edu.cn

CONTENTS

I. Introduction	2
II. The KSVZ axion model and the electroweak monopole	4
III. The solution of electroweak monopole in the presence of axion	6
IV. Characteristics of electroweak monopole and axion	8
A. The monopole energy	8
B. The electromagnetic charges	10
C. The axion potential energy	11
V. Conclusion	13
Acknowledgments	13
References	13

I. INTRODUCTION

The axion and magnetic monopole are two of the longstanding and interesting candidates of physics beyond the standard model (SM). The pseudo-Goldstone boson axion was proposed to solve the strong CP problem in quantum chromodynamics (QCD), as a result of the spontaneous breaking of a QCD anomalous Peccei-Quinn (PQ) global symmetry $U(1)_{PQ}$ [1–4]. The chiral transformation of the quark fields with PQ charges leads to the anomaly under quantum electro-dynamics (QED) and the coupling between the axion field *a* and electromagnetic fields

$$\frac{1}{4}g_{a\gamma\gamma}aF_{\mu\nu}\tilde{F}^{\mu\nu} = g_{a\gamma\gamma}a\vec{E}\cdot\vec{B} .$$
(1)

The QCD axion and axion-like particles (ALPs) can also play as dark matter through the misalignment mechanism [5, 6]. The theory of axion and the detection of axion couplings have received a wide interest in both theoretical and experimental aspects (see Refs. [7, 8] for recent reviews).

P. Dirac first suggested the existence of magnetic monopole in quantum theory in 1931 [9]. It can be constructed from a non-singular Abelian theory [10] or can arise from the spontaneous breaking of non-Abelian gauge symmetries, resulting to monopole in grand unification theory (GUT) [11, 12]. Besides GUT monopole, the existence of electroweak monopole in particular gains much interest and attention [13–17]. Cho and Maison showed that the Weinberg-Salam model has a topology of magnetic monopole and there does exist a new type of dyon solutions in the SM [13] (see Ref. [18] for a recent review). This electroweak generalization of the Dirac monopole comes from the non-trivial topology of the SM. The SU(2) of the Weinberg-Salam model has the same non-Abelian monopole topology as the Georgi-Glashow model. The $U(1)_Y$

part has the Abelian monopole topology. The Cho-Maison monopole (CMM) is thus a hybrid of 't Hooft-Polyakov monopole and Dirac monopole. Its magnetic charge is twice that of the Dirac monopole and the monopole mass is of the order of

$$\frac{4\pi}{e^2}M_W \simeq 11 \text{ TeV} , \qquad (2)$$

where e is the elementary electric charge and M_W is the W boson mass. This low mass motives the detection of such realistic monopole at the ATLAS [19] or MoEDAL [20] detector of LHC or future high-energy upgrades. The electroweak monopole also leads to various impacts on cosmology, such as the primordial magnetic black holes and dark matter.

In 1979, E. Witten pointed out that a CP violating term in the non-Abelian SO(3) theory provides an additional electric charge for the 't Hooft-Polyakov monopoles in this theory [21]. The generic value of electric charge q_e becomes

$$q_e = e\left(n_e + \frac{\theta}{2\pi}n_m\right),\tag{3}$$

where n_e $(n_m) \in \mathbb{Z}$ is the electric (magnetic) quantum number, and θ denotes a CP violating parameter. When substituting the parameter θ by the axion field, this so-called Witten effect built the close relationship between axion and magnetic monopole. W. Fischler et al. then derived the dyon-axion dynamics under the classical electromagnetism in 1983 [22]. There are many recent studies discussing the possible modification of standard axion electrodynamics inspired by Witten effect in both theory [23–25] and phenomenology [26–32]. There also exist some open questions in the axion-magnetic monopole aspect [33].

What is then the implication of axion to the electroweak monopole? In this work we try to make the first attempt. We explore the solution of Cho-Maison electroweak monopole in the benchmark model of axion, i.e. the Kim-Shifman-Vainshtein-Zakharov (KSVZ) model [34, 35]. Besides axion interactions at low energies, the content of KSVZ model is exactly the same as the SM. We are able to construct the topological solution of the electroweak monopole in the SM. More importantly, inspired by the Witten effect, the characteristic features of electroweak monopole would be changed in light of the interaction of axion and electromagnetic field. For instance, the axion-photon coupling would induce additional electric charge for the monopole and modify the monopole mass. The monopole background would also confine the axion potential. We will concentrate on the solution of spherically symmetrical function of axion field in a background magnetic field. The contributions of axion to the properties of electroweak monopole will be studied in details. These changes have impacts on both the existence of axion and the search for electroweak monopole.

This paper is organized as follows. In Sec. II we overview the benchmark axion model and the electroweak monopole called Cho-Maison monopole. The topological solutions of electroweak monopole and axion field will be explored in Sec. III. In Sec. IV we discuss the impact of axion on the properties of electroweak monopole. Our conclusions are drawn in Sec. V.

II. THE KSVZ AXION MODEL AND THE ELECTROWEAK MONOPOLE

We first overview the KSVZ axion model and the electroweak monopole in the SM called Cho-Maison monopole. The Lagrangian for the KSVZ model is

$$\mathcal{L}_{\rm KSVZ} = -\frac{1}{4} W^a_{\mu\nu} W^{a\mu\nu} - \frac{1}{4} B_{\mu\nu} B^{\mu\nu} + |D_\mu \Phi|^2 + |\partial_\mu \phi|^2 + \mathcal{L}_{\rm VLF} - V_{\rm KSVZ}(\Phi, \phi) , \qquad (4)$$

where $W^a_{\mu\nu}$ (a = 1, 2, 3) and $B_{\mu\nu}$ respectively denote the field strength tensors of $SU(2)_L$ and $U(1)_Y$ gauge fields, Φ is the SM Higgs doublet, and ϕ is a complex scalar singlet. The covariant derivative to the Higgs doublet is

$$D_{\mu}\Phi = \left(\partial_{\mu} + \frac{ig}{2}\sigma^{a}W_{\mu}^{a} + \frac{ig'}{2}B_{\mu}\right)\Phi, \qquad (5)$$

where σ^a denotes the Pauli matrix, and g and g' are the gauge couplings of $SU(2)_L$ and $U(1)_Y$, respectively. The KSVZ model introduces the kinetic term and Yukawa term for a vector-like fermion (VLF) $Q \sim (3, 1, 0)$

$$\mathcal{L}_{\rm VLF} = \overline{\mathcal{Q}}i\not\!\!{D}\mathcal{Q} - y_{\mathcal{Q}}\overline{\mathcal{Q}_L}\mathcal{Q}_R\phi + h.c. , \qquad (6)$$

which emerges a $U(1)_{PQ}$ symmetry and the scalar potential

$$V_{\rm KSVZ}(\Phi,\phi) = \lambda_{\Phi} \left(|\Phi|^2 - \frac{v^2}{2} \right)^2 + \lambda_{\phi} \left(|\phi|^2 - \frac{v_a^2}{2} \right)^2 \,. \tag{7}$$

Here v is the vacuum expectation value (vev) of the Higgs field Φ and the $U(1)_{PQ}$ symmetry is spontaneously broken with the vev parameter v_a . The complex scalar field is decomposed as

$$\phi = \frac{1}{\sqrt{2}} (v_a + \sigma_a) e^{ia/v_a} , \qquad (8)$$

where a denotes the axion field and σ_a is the radial mode. After performing the axial rotation of the VLF field

$$\mathcal{Q} \to e^{-i\gamma_5 \frac{a}{2v_a}} \mathcal{Q} , \qquad (9)$$

and removing the axion field in the Yukawa term, one obtains the following anomalous Lagrangian

$$\delta \mathcal{L}_{\text{KSVZ}} = \frac{\alpha_s N}{4\pi} \frac{a}{v_a} G^a_{\mu\nu} \tilde{G}^{a\mu\nu} + \frac{\alpha E}{4\pi} \frac{a}{v_a} F_{\mu\nu} \tilde{F}^{\mu\nu} , \qquad (10)$$

where $G^a_{\mu\nu}$ $(a = 1, \dots, 8)$ and $F_{\mu\nu}$ are the field strength tensors of $SU(3)_c$ and $U(1)_{\rm em}$, respectively, the dual field strengths are defined as $\tilde{X}_{\mu\nu} \equiv \frac{1}{2} \epsilon_{\mu\nu\alpha\beta} X^{\alpha\beta}$ with $\epsilon_{0123} = 1^{-1}$, and E and $N = N_{\rm DW}/2$ are the anomaly coefficients with $N_{\rm DW}$ being the domain wall (DW) number ².

¹ We follow the convention in Ref. [7] and adopt the metric tensor as diag(+1, -1, -1, -1) throughout the paper.

² The $N_{\rm DW} = 1$ model is taken throughout the paper.

Throughout the paper, we only consider the second coupling between axion and the physical photon. Taking into account both the quark kinetic term and the quark mass operator, the axion-photon coupling can be given as

$$\frac{1}{4}g_{a\gamma\gamma}aF_{\mu\nu}\tilde{F}^{\mu\nu} \tag{11}$$

with

$$g_{a\gamma\gamma} = \frac{\alpha}{\pi v_a} \left[E - N \left(\frac{2}{3} \frac{4m_d + m_u}{m_u + m_d} \right) \right] \simeq \frac{\alpha}{\pi v_a} (E - 1.92N) . \tag{12}$$

We take $g_{a\gamma\gamma} \simeq -0.0024/v_a$ in the minimal KSVZ model.

Next we show the ansatz for the spherically symmetry solution of the Cho-Maison monopole in the KSVZ model with the effective Lagrangian

$$\mathcal{L}_{eff} = -\frac{1}{4} W^{a}_{\mu\nu} W^{a\mu\nu} - \frac{1}{4} B_{\mu\nu} B^{\mu\nu} + |D_{\mu}\Phi|^{2} + |\partial_{\mu}\phi|^{2} -\lambda_{\Phi} \left(|\Phi|^{2} - \frac{v^{2}}{2} \right)^{2} - \lambda_{\phi} \left(|\phi|^{2} - \frac{v^{2}_{a}}{2} \right)^{2} + \frac{1}{4} g_{a\gamma\gamma} a F_{\mu\nu} \tilde{F}^{\mu\nu} .$$
(13)

The authors of Refs. [13, 18] make use of the gauge-independent Abelian decomposition to Abelianize the non-Abelian gauge theory. The Abelianized dual potential has both a nontopological electric potential and a topological magnetic potential. They verify that the Abelian decomposition of the Weinberg-Salam model is similar to that of the Georgi-Glashow model and they have the same SU(2) monopole topology. Inspired by this implication, one can parameterize the Higgs doublet in terms of the spherical coordinates (t, r, θ, φ) as [13, 18]

$$\Phi = \frac{1}{\sqrt{2}}\rho(r)\xi(\theta,\varphi), \qquad \xi^{\dagger}\xi = 1, \qquad (14)$$

with $\rho(r)$ being a real function, and $\xi(\theta, \varphi) = i(\sin \frac{\theta}{2}e^{-i\varphi}, -\cos \frac{\theta}{2})^T$ being complex unit doublet. The ansatz for the spherically symmetry solution of the electroweak dyon is given as [13, 18]

$$g\dot{W}_{\mu} = A(r)(\partial_{\mu}t)\hat{r} + (f(r) - 1)\hat{r} \times \partial_{\mu}\hat{r},$$

$$g'B_{\mu} = B(r)\partial_{\mu}t - (1 - \cos\theta)\partial_{\mu}\varphi,$$
(15)

with $\hat{r} = -\xi^{\dagger} \vec{\sigma} \xi = (\sin \theta \cos \varphi, \sin \theta \sin \varphi, \cos \theta)$. Either using the unitary gauge or using the gauge independent Abelian decomposition, the physical fields are expressed as [13, 18]

$$A_{\mu} = e \left(\frac{A(r)}{g^2} + \frac{B(r)}{g'^2} \right) \partial_{\mu} t - \frac{1}{e} (1 - \cos \theta) \partial_{\mu} \varphi , \qquad (16)$$

$$W_{\mu} = \frac{if(r)}{\sqrt{2}g} e^{+i\varphi} (\partial_{\mu}\theta + i\sin\theta\partial_{\mu}\varphi) , \qquad (17)$$

$$W^*_{\mu} = \frac{if(r)}{\sqrt{2}g} e^{-i\varphi} \left(-\partial_{\mu}\theta + i\sin\theta\partial_{\mu}\varphi\right), \qquad (18)$$

$$Z_{\mu} = \frac{e}{gg'} (A(r) - B(r))\partial_{\mu}t .$$
⁽¹⁹⁾

The complex scalar singlet is parameterized as

$$\phi = \frac{1}{\sqrt{2}} (v_a + \sigma_a) e^{i\hat{a}(r)} , \qquad (20)$$

where σ_a as the radial component is supposed to be stabilized and will not be considered below, and $\hat{a}(r) \equiv a(r)/v_a$ is the dimensionless axion (angular) component defined in the domain $[0, 2\pi v_a)$. The axion-photon coupling in Eq. (11) leads to the modified equations of motion (EoM)

$$\partial_{\mu}(F^{\mu\nu} - g_{a\gamma\gamma}a\tilde{F}^{\mu\nu}) = 0 , \qquad (21)$$

and the relevant modified Gauss's law for the electric field

$$\vec{\nabla} \cdot (\vec{E} + g_{a\gamma\gamma} a \vec{B}) = 0 .$$
⁽²²⁾

(23)

Suppose there is a magnetic monopole with a magnetic charge q_m , the above Gauss's law changes the usual electric charge quantization $q_e/e = n \in \mathbb{Z}$ to

$$\frac{q_e}{e} + \frac{q_m}{e} g_{a\gamma\gamma} a(\infty) = n \in \mathbb{Z} .$$
(24)

This turns out to be the generic feature of Witten effect in U(1) gauge group.

III. THE SOLUTION OF ELECTROWEAK MONOPOLE IN THE PRESENCE OF AXION

Using the spherically symmetric ansatz in Eqs. (14) and (15), we can rewrite the above effective Lagrangian in Eq. (13) in terms of five radial functions $\rho(r)$, f(r), A(r), B(r) and a(r). The complete expansion of the Lagrangian contains three parts: the Cho-Maison monopole part, the axion kinetic energy and the axion interaction

$$\mathcal{L}_{eff} = \mathcal{L}_{\rm CMM} + \mathcal{L}_{\rm axion \ kin.} + \mathcal{L}_{\rm axion \ int.}$$
(25)

with

$$\mathcal{L}_{\text{CMM}} = \frac{1}{2} (\partial_{\mu} \rho)^{2} - \frac{\lambda_{\Phi}}{4} (\rho^{2} - v^{2})^{2} - \frac{1}{4} F_{\mu\nu}^{2} - \frac{1}{4} Z_{\mu\nu}^{2} + \frac{g^{2}}{4} \rho^{2} W_{\mu}^{*} W^{\mu} + \frac{g^{2} + {g'}^{2}}{8} \rho^{2} Z_{\mu}^{2} - \frac{1}{2} \left| (D_{\mu} W_{\nu} - D_{\nu} W_{\mu}) + ie \frac{g}{g'} (Z_{\mu} W_{\nu} - Z_{\nu} W_{\mu}) \right|^{2} + ie F_{\mu\nu} W^{*\mu} W^{\nu} + ie \frac{g}{g'} Z_{\mu\nu} W^{*\mu} W^{\nu} + \frac{g^{2}}{4} (W_{\mu}^{*} W_{\nu} - W_{\nu}^{*} W_{\mu})^{2} = \frac{1}{2g^{2}} \left(\frac{dA}{dr} \right)^{2} + \frac{1}{8} (A - B)^{2} \rho^{2} + \frac{f^{2} A^{2}}{g^{2} r^{2}} - \frac{f^{2} \rho^{2}}{4r^{2}} + \frac{1}{2g'^{2}} \left(\frac{dB}{dr} \right)^{2} - \frac{1}{g^{2} r^{2}} \left(\frac{df}{dr} \right)^{2} - \frac{1}{2g'^{2} r^{4}} - \frac{1}{2} \left(\frac{d\rho}{dr} \right)^{2} - \frac{\lambda_{\Phi}}{4} (\rho^{2} - v^{2})^{2} - \frac{(f^{2} - 1)^{2}}{2g^{2} r^{4}} ,$$
(26)

$$\mathcal{L}_{\text{axion kin.}} = \frac{1}{2} (\partial_{\mu} a)^2 = -\frac{1}{2} \left(\frac{da}{dr} \right)^2 , \qquad (27)$$

$$\mathcal{L}_{\text{axion int.}} = \frac{1}{4} g_{a\gamma\gamma} a F_{\mu\nu} \tilde{F}^{\mu\nu} = -g_{a\gamma\gamma} \frac{a}{r^2} \left[\left(\frac{dA}{dr} \right) \left(\frac{\sin^2 \theta_W}{g^2} (1 - f^2) + \frac{\cos^2 \theta_W}{g^2} \right) + \left(\frac{dB}{dr} \right) \left(\frac{\sin^2 \theta_W}{g'^2} (1 - f^2) + \frac{\cos^2 \theta_W}{g'^2} \right) \right] , \qquad (28)$$

where $D_{\mu} = \partial_{\mu} + ieA_{\mu}$ and $Z_{\mu\nu} = \partial_{\mu}Z_{\nu} - \partial_{\nu}Z_{\mu}$. Then, the equations of motion for the above radial functions can be obtained as

$$\frac{d^2\rho}{dr^2} + \frac{2}{r}\frac{d\rho}{dr} - \frac{f^2}{2r^2}\rho + \frac{1}{4}(A-B)^2\rho = \lambda_{\Phi}(\rho^2 - v^2)\rho , \qquad (29)$$

$$\frac{d^2f}{dr^2} - \frac{(f^2 - 1)f}{r^2} - T_f = \left(\frac{g^2\rho^2}{4} - A^2\right)f, \qquad (30)$$

$$\frac{d^2A}{dr^2} + \frac{2}{r}\frac{dA}{dr} - \frac{2f^2A}{r^2} + T_A = \frac{g^2}{4}(A-B)\rho^2 , \qquad (31)$$

$$\frac{d^2B}{dr^2} + \frac{2}{r}\frac{dB}{dr} + T_B = -\frac{g'^2}{4}(A-B)\rho^2 , \qquad (32)$$

$$\frac{d^2a}{dr^2} + \frac{2}{r}\frac{da}{dr} + T_a = 0 , \qquad (33)$$

where

$$T_f = -g_{a\gamma\gamma}af\left[\sin^2\theta_W\left(\frac{dA}{dr}\right) + \cos^2\theta_W\left(\frac{dB}{dr}\right)\right] , \qquad (34)$$

$$T_A = T_B = -g_{a\gamma\gamma} \frac{1}{r^2} \left[\left(\frac{da}{dr} - \frac{2a}{r} \right) \left(1 - \sin^2 \theta_W f^2 \right) - 2\sin^2 \theta_W a f \left(\frac{df}{dr} \right) \right] , \qquad (35)$$

$$T_a = -g_{a\gamma\gamma} \frac{1}{r^2} \left[\left(\frac{dA}{dr} \right) \left(\frac{1}{g^2} - \frac{\sin^2 \theta_W}{g^2} f^2 \right) + \left(\frac{dB}{dr} \right) \left(\frac{1}{g'^2} - \frac{\sin^2 \theta_W}{g'^2} f^2 \right) \right].$$
(36)

When $g_{a\gamma\gamma}$ is absent, the first four equations of motion would restore to those for pure Cho-Maison monopole [18]. The last equation becomes the equation of motion for axion field.

As stated in Refs. [13, 18], the energy of pure Cho-Maison monopole is infinite according to the Lagrangian \mathcal{L}_{CMM} . To make the electroweak monopole finite, one has to include the quantum correction and regularize the Cho-Maison monopole [36–38]. The ultraviolet regularization introduces the scale-dependent parameters α , β and changes the CMM Lagrangian to

$$\mathcal{L}_{CMM}' = \frac{1}{2g^2} \left(\frac{dA}{dr}\right)^2 + \frac{1}{8}(A-B)^2\rho^2 + \frac{f^2A^2}{g^2r^2} - \frac{f^2\rho^2}{4r^2} + \frac{1}{2g'^2} \left(\frac{dB}{dr}\right)^2 - \frac{1}{g^2r^2} \left(\frac{df}{dr}\right)^2 - \frac{1}{2g'^2r^4} - \frac{1}{2} \left(\frac{d\rho}{dr}\right)^2 - \frac{\lambda_{\Phi}}{4}(\rho^2 - v^2)^2 - \frac{1}{2g^2r^4} \left[1 - 2(1+\alpha)f^2 + (1+\beta)f^4\right] (37)$$

Then, the equation of motion in Eq. (30) is modified as

$$\frac{d^2f}{dr^2} - \frac{1}{r^2} \Big[(1+\beta)f^2 - (1+\alpha) \Big] f - T_f = \Big(\frac{g^2\rho^2}{4} - A^2\Big) f .$$
(38)

To make the monopole energy finite, the constants α and β should be given by

$$1 + \alpha = \frac{1}{f(0)^2} \frac{g^2}{e^2} , \quad 1 + \beta = \frac{1}{f(0)^4} \frac{g^2}{e^2} . \tag{39}$$

Under appropriate boundary conditions, we can numerically integrate the above differential equations to obtain the electroweak dyon solution with KSVZ axion. Following Ref. [18], we choose the following boundary conditions

$$\rho(0) = 1 , \ \rho(\infty) = v , \ f(0) = c , \ f(\infty) = 0 ,$$

$$A(0) = 0 , \ B(0) = 0.2gv , \ A(\infty) = B(\infty) = gv/4 = M_W/2 ,$$
(40)

for the Higgs boson, W boson and Z boson represented by ρ , f and A - B, respectively. Here we also choose c = 1 for the Cho-Maison monopole and $c = g/e = 1/\sin\theta_W$ for UV regularized electroweak monopole. The latter choice fixes the correction constants as $\alpha = 0$ and $\beta = -\cos^2\theta_W$. As the axion is defined as an angular variable in $[0, 2\pi v_a)$ and the QCD axion potential is also periodic in this range, we set the boundary conditions for axion as

$$a(0) = 0, \ a(\infty) = 2\pi v_a$$
 (41)

In order to solve the above differential equations in Eqs. (29,30,31,32,33), we use the relaxation method for numerical iterative solution [39]. Besides, for the region of $x = M_W \cdot r < 0.5$, we take the following analytical approximation in each iteration

Cho-Maison monopole:
$$\rho \simeq c_0 r^{\delta_-}, \quad f \simeq 1 + c_1 r^2, \quad A \simeq c_2 r^3, \quad B \simeq b_0 + c_3 r^{2\delta_+}, \quad a \simeq c_4 r^3,$$

regularized monopole: $\rho \simeq c_0 r^{\delta_1}, \quad \frac{f}{f(0)} \simeq 1 + c_1 r^{\delta_2}, \quad A \simeq c_2 r^3, \quad B \simeq b_0 + c_3 r^{\delta_4}, \quad a \simeq c_4 r^3,$
(42)

where $\delta_{\pm} = (\sqrt{3}\pm 1)/2$, $\delta_1 = \frac{1}{2}(\sqrt{1+2f^2(0)}-1)$, $\delta_2 = \frac{1}{2}(1+\sqrt{8\alpha+9})$, $\delta_3 = \frac{1}{2}(\sqrt{1+8f^2(0)}-1)$, $\delta_4 = \sqrt{1+2f^2(0)}+1$. The $c_{0,1,2,3,4}$ and b_0 coefficients are chosen to match the numerical solution smoothly. In Fig. 1, we show the solutions of Cho-Maison monopole with energy divergence at r = 0 (dashed lines), the finite electroweak monopole (solid lines) as well as axion field. The anomaly coefficients of axion in KSVZ model are E = 0 and N = 1/2. The PQ scale is set as $v_a = 10^7$ GeV for illustration. In the presence of small axion-photon coupling, the monopole solutions remain the same as those of pure Cho-Maison electroweak monopole in Ref. [18].

IV. CHARACTERISTICS OF ELECTROWEAK MONOPOLE AND AXION

A. The monopole energy

We first show the monopole energy in the presence of axion. The 00 component of the energymomentum tensor T^{00} in terms of the gauge fields is

$$T^{00} = 2(D^0\Phi^{\dagger})(D^0\Phi) - (W^a)^{0i}(W^a)^0_{\ i} - B^{0i}B^0_{\ i} - \frac{1}{2}g_{a\gamma\gamma}a \left[2\sin^2\theta_W(\tilde{W}^3)^{0i}(W^3)^0_{\ i}\right]$$



FIG. 1. The solutions of Cho-Maison monopole with energy divergence at r = 0 (dashed lines), the finite electroweak monopole (solid lines) as well as axion field. The monopole with finite energy is obtained with $f(0) = 1/\sin \theta_W$. The anomaly coefficients of axion in KSVZ model are E = 0 and N = 1/2. The PQ scale is set as $v_a = 10^7$ GeV for illustration.

$$+2\cos^{2}\theta_{W}\tilde{B}^{0i}B^{0}_{i} + \sin\theta_{W}\cos\theta_{W}\left(\tilde{B}^{0i}(W^{3})^{0}_{i} + (\tilde{W}^{3})^{0i}B^{0}_{i}\right)\right] - \mathcal{L}_{eff} .$$
(43)

The total energy of monopole in the KSVZ model can be obtained as

$$E = E_{\text{CMM}} + E_{\text{axion int.}}, \qquad (44)$$

$$E_{\text{CMM}} = \frac{4\pi}{g'^2} \int_0^\infty \frac{dr}{2r^2} + 4\pi \int_0^\infty dr \left\{ \frac{r^2}{2} \left(\frac{d\rho}{dr}\right)^2 + \frac{1}{g^2} \left[\left(\frac{df}{dr}\right)^2 + \frac{r^2}{2} \left(\frac{dA}{dr}\right)^2 + \frac{1}{2} \frac{(f^2 - 1)^2}{r^2} + A^2 f^2 \right] + \frac{r^2}{2} \left(\frac{dB}{dr}\right)^2 + \frac{\lambda_{\Phi}}{2} \left(\frac{2}{r^2} - \frac{2}{2}\right)^2 + \frac{1}{r^2} \left(\frac{r^2}{r^2} - \frac{r^2}{r^2}\right)^2 + \frac{1}{2} \left(\frac{r^2}{r^2} - \frac{1}{r^2}\right)^2 + \frac{1}{r^2} \left(\frac{r^2}{$$

$$+\frac{r^{2}}{2g^{\prime 2}}\left(\frac{dB}{dr}\right)^{2} + \frac{\lambda_{\Phi}}{4}r^{2}(\rho^{2} - v^{2})^{2} + \frac{1}{4}f^{2}\rho^{2} + \frac{r^{2}}{8}\rho^{2}(A - B)^{2}\right\},$$
(45)

$$E_{\text{axion int.}} = 2\pi g_{a\gamma\gamma} \left(\frac{\sin^2 \theta_W}{g'^2}\right) \int_0^\infty dr a(r) \left[\left(\frac{dA}{dr}\right) + (1 - f^2)\left(\frac{dB}{dr}\right)\right], \tag{46}$$

where the first term of E_{CMM} is apparently infinite. The other terms in E_{CMM} and $E_{\text{axion int.}}$ are all finite. The origin of this infinite energy comes from the singularity of the point-like magnetic monopole of $U(1)_Y$. In Refs. [18, 36–38], the author utilized the UV regularization to include the quantum correction and proved the existence of electroweak monopole with a finite energy. According to the Lagrangian with scale-dependent parameters in Eq. (37), the modified energy E_{CMM} becomes

$$E'_{\rm CMM} = \frac{4\pi}{g^2} \int_0^\infty \frac{dr}{2r^2} \left\{ \frac{g^2}{e^2} - \frac{(1+\alpha)^2}{1+\beta} + (1+\beta) \left(f^2 - \frac{1+\alpha}{1+\beta} \right)^2 \right\}$$

$$+4\pi \int_{0}^{\infty} dr \left\{ \frac{r^{2}}{2} \left(\frac{d\rho}{dr} \right)^{2} + \frac{1}{g^{2}} \left[\left(\frac{df}{dr} \right)^{2} + \frac{r^{2}}{2} \left(\frac{dA}{dr} \right)^{2} + A^{2} f^{2} \right] + \frac{r^{2}}{2g'^{2}} \left(\frac{dB}{dr} \right)^{2} + \frac{\lambda_{\Phi} r^{2}}{4} (\rho^{2} - v^{2})^{2} + \frac{1}{4} f^{2} \rho^{2} + \frac{r^{2}}{8} \rho^{2} (A - B)^{2} \right\}.$$
 (47)

Given the fixed α and β in Eq. (39), the energy becomes finite at the origin. We evaluate the above integrals to obtain the energy E for the electroweak monopole (with $g_{a\gamma\gamma} = 0$ and a(r) = 0) and that in the presence of axion. The obtained monopole masses in units of TeV are shown in the second column in Table I. The axion contribution to the mass is from the axion interaction in Eq. (46) and is governed by the $g_{a\gamma\gamma}$ coupling. One can see that the monopole mass is slightly decreased by 0.1% (~ 10 GeV) and actually the change does not depend on the PQ scale v_a . Suppose the electroweak monopole will be observed in future, its property may confine the nature of axion. On the other hand, the observation of axion would also have impact on the monopole of TeV scale.

Characteristics of monopole	mass (in units of TeV)	electric charge (in units of $4\pi/e$)
w/o axion	13.67	-0.23
w/ axion	13.66	-0.28

TABLE I. The characteristics of monopole without or with the presence of KSVZ axion, under the choice of $f(0) = 1/\sin \theta_W$.

B. The electromagnetic charges

We next discuss the electromagnetic properties of electroweak monopole in the presence of KSVZ axion. According to the ansatz of neutral gauge components in Eq. (15), we can obtain the electromagnetic tensor $F_{\mu\nu}$ given by the electromagnetic field A_{μ} in Eq. (16). The corresponding electric field \vec{E} and magnetic field \vec{B} can be then written as

$$\vec{E} = -e\left(\frac{\dot{A}(r)}{g^2} + \frac{\dot{B}(r)}{g'^2}\right)\hat{r}$$
, (48)

$$\vec{B} = -\frac{\hat{r}}{er^2} \,. \tag{49}$$

where the extra minus sign on the right-handed side of \vec{B} is due to our choice of metric tensor diag(+1, -1, -1, -1). These electromagnetic fields are purely generated by the electroweak monopole in vacuum. However, if we take into account the axion which couples to the physical photon, the axion acts as a kind of source to excite another electromagnetic field. It changes the electric field as

$$\vec{E} \to \vec{E} + \vec{E}_a = \vec{E} + g_{a\gamma\gamma} a \vec{B}$$
 (50)

For the electric and magnetic charges of the electroweak monopole, they are defined as the integrals of electromagnetic fields over a closed curve. We have the following charges

$$\begin{split} q_e &= \oint_{r=\infty} d\vec{S} \cdot (\vec{E} + g_{a\gamma\gamma} a\vec{B}) = -4\pi e \int_0^\infty dr \left[r^2 \left(\frac{\dot{A}(r)}{g^2} + \frac{\dot{B}(r)}{g'^2} \right) \right]' + g_{a\gamma\gamma} \oint_{r=\infty} d\vec{S} \cdot a\vec{B} \\ &= -4\pi e \left[r^2 \left(\frac{\dot{A}(r)}{g^2} + \frac{\dot{B}(r)}{g'^2} \right) \right] \left|_{r=\infty} + q_m g_{a\gamma\gamma} a(r) \right|_{r=\infty} \\ &= -\frac{8\pi}{e} \sin^2 \theta_W \int_0^\infty dr f^2(r) A(r) + \frac{4\pi}{e} \int_0^\infty dr r^2 T_A + q_m g_{a\gamma\gamma} 2\pi v_a , \\ q_m &= \oint_{r=\infty} d\vec{S} \cdot \vec{B} = -\frac{4\pi}{e} . \end{split}$$

The magnetic charge q_m is exactly the same as that of the Cho-Maison monopole. The first term of electric charge q_e is that of the pure Cho-Maison monopole and the last two terms are induced by the presence of axion-photon interaction. We show the modified electric charge in the third column of Table I in units of $4\pi/e$. One can see that the KSVZ axion changes the electric charge of electroweak monopole by about 20%. The change does not depend on the PQ scale v_a either.

C. The axion potential energy

Finally, we reexamine the axion potential in the background field of a monopole which was first studied by Fischler et al. in Ref. [22]. The axion potential energy can be divided into two parts. One part is provided by the kinetic energy of axion. The axion itself has a lot of kinetic energy. The other part comes from the electrostatic field energy $1/2\vec{E}_a^2$. The induced electric field \vec{E}_a is proportion to the magnetic field of monopole \vec{B} . Thus, there exists a non-negligible energy cost $1/2\vec{E}_a^2$ in the vicinity of the axion. In other words, the axion is strongly repelled by the monopole in electromagnetic properties and needs to pay a price of large energy if it wants to be combined with monopole. The axion potential energy is given by

$$V_{a} = \frac{1}{2} \int dV \left[(\nabla a)^{2} + \vec{E}_{a}^{2} \right]$$

$$= E_{\text{axion kin.}} + \frac{1}{2} \int dV \left[g_{a\gamma\gamma}^{2} \frac{a(r)^{2}}{e^{2}r^{4}} \right]$$

$$= 2\pi \int_{0}^{\infty} dr r^{2} \left(\frac{da}{dr} \right)^{2} + 2\pi \int_{0}^{\infty} dr \left(\frac{g_{a\gamma\gamma}a(r)}{er} \right)^{2} \equiv \int_{0}^{\infty} dr K(r) .$$
(51)

We can then follow the substitution of integral variable in Ref. [22] to minimize V_a . The axion potential can be rewritten as

$$V_{a} = -2\pi \left[r_{0} \int_{0}^{\infty} dz \left(\frac{da}{dz} \right)^{2} + \frac{g_{a\gamma\gamma}^{2}}{e^{2}} \frac{1}{r_{0}} \int_{0}^{\infty} dz a^{2} \right] , \qquad (52)$$

where the radius r is replaced with a dimensionless variable $z = r_0/r$ with $r_0 = |g_{a\gamma\gamma}/e|$. The axion in the configuration of $a(r) = 2\pi v_a e^{-r_0/r}$ approaches $2\pi v_a$ when $z \to 0$ and can minimize V_a [22]. The minimum potential energy is then given by

$$V_a^{\min} = -\pi (2\pi v_a)^2 \left(r_0 + \frac{g_{a\gamma\gamma}^2}{e^2 r_0} \right) = -8\pi^3 v_a^2 \frac{g_{a\gamma\gamma}}{e} .$$
(53)

Note that the calculation of this minimum value is only reliable in the classical limit. As seen from the left panels of Fig. 2, the above a(r) configuration (red line) agrees with our numerical solution of axion field (blue line) only for $r \gg r_0$. For small radial distance, there exhibits significant distinction between them. As v_a decreases, the analytical result approaches the numerical solution. The integrand K(r) is shown in the right panels of Fig. 2. One can see that there is sizable contribution in addition to this minimum value from the effect of axion radial mode. In Table II, we show the comparison of the axion potential energy V_a between the minimum value calculated as above and the one by integrating the numerical solution of the axion field. Indeed, the completely numerical result exhibits a much larger axion potential energy than the minimum value.



FIG. 2. The normalized axion field a/v_a (left) and the integrand K(r) in Eq. (51) (right), as a function of r in units of 1/v. We compare the result of the analytical form $a(r) \sim e^{-r_0/r}$ (red line) with that of our numerical solution (blue line). Both cases of $v_a = 10^7$ GeV (top) and 10^3 GeV (bottom) are shown.

V_a	minimum value V_a^{\min}	numerical result V_a
$v_a = 10^7 \text{ GeV}$	$1.89\times 10^7~{\rm GeV}$	$3.32 imes 10^{13} { m GeV}$
$v_a = 10^3 \ {\rm GeV}$	$1.89\times 10^3{\rm GeV}$	$3.32 imes 10^5 { m ~GeV}$

TABLE II. The comparison of the axion potential in Eq. (51) between the minimum value calculated from the analytical formula of axion solution in Eq. (53) and the one by integrating the numerical solution of the axion field from EoM.

V. CONCLUSION

The Witten effect implies the dynamics of axion and magnetic monopole. The Cho-Maison monopole is a realistic electroweak monopole arisen in the Weinberg-Salam theory. This monopole of TeV scale mass motivates the dedicated search for electroweak monopole at colliders. It is plausible to explore the implication of axion to the electroweak magnetic monopole.

In this work we investigate the topological solutions of Cho-Maison electroweak monopole in the presence of KSVZ axion. We use the spherically symmetric ansatz for the electroweak dyon and introduce the spherically symmetric function for the axion field. The effective Lagrangian is then showed in terms of the radial functions. It includes the electroweak monopole part, the axion kinetic energy as well as the axion interaction term. We derive the consequent equations of motion in the presence of the axion-photon coupling and show the numerical results. Given the above topological solutions, we calculate the characteristics of the electroweak monopole such as the monopole mass and the electromagnetic charges.

We find that the monopole mass can be slightly changed by the axion-photon interaction. The presence of KSVZ axion changes the electric charge of the monopole by 20%. These changes would have impacts on either the existence of axion or the testability of electroweak monopole. We also calculate the axion potential energy in terms of the axion solution from EoM, and compare with the result in the classical limit.

ACKNOWLEDGMENTS

We would like to thank Yongcheng Wu and Ke-Pan Xie for very useful discussions. T. L. is supported by the National Natural Science Foundation of China (Grant No. 12375096, 12035008, 11975129) and "the Fundamental Research Funds for the Central Universities", Nankai University (Grants No. 63196013).

^[1] R. D. Peccei and H. R. Quinn, Phys. Rev. Lett. 38, 1440 (1977).

^[2] R. D. Peccei and H. R. Quinn, Phys. Rev. D 16, 1791 (1977).

^[3] S. Weinberg, Phys. Rev. Lett. 40, 223 (1978).

- [4] F. Wilczek, Phys. Rev. Lett. 40, 279 (1978).
- [5] J. Preskill, M. B. Wise, and F. Wilczek, Phys. Lett. B 120, 127 (1983).
- [6] M. Dine and W. Fischler, Phys. Lett. B 120, 137 (1983).
- [7] L. Di Luzio, M. Giannotti, E. Nardi, and L. Visinelli, Phys. Rept. 870, 1 (2020), arXiv:2003.01100 [hep-ph].
- [8] P. Sikivie, Reviews of Modern Physics 93 (2021), 10.1103/revmodphys.93.015004.
- [9] P. A. M. Dirac, Proc. Roy. Soc. Lond. A 133, 60 (1931).
- [10] T. T. Wu and C. N. Yang, Phys. Rev. D 12, 3845 (1975).
- [11] G. 't Hooft, Nucl. Phys. B 79, 276 (1974).
- [12] A. M. Polyakov, JETP Lett. 20, 194 (1974).
- [13] Y. M. Cho and D. Maison, Phys. Lett. B 391, 360 (1997), arXiv:hep-th/9601028.
- [14] P. Q. Hung, Nucl. Phys. B 962, 115278 (2021), arXiv:2003.02794 [hep-ph].
- [15] J. Alexandre and N. E. Mavromatos, Phys. Rev. D 100, 096005 (2019), arXiv:1906.08738 [hep-ph].
- [16] J. Ellis, N. E. Mavromatos, and T. You, Phys. Lett. B 756, 29 (2016), arXiv:1602.01745 [hep-ph].
- [17] G. Lazarides and Q. Shafi, Phys. Rev. D 103, 095021 (2021), arXiv:2102.07124 [hep-ph].
- [18] Y. M. Cho, Phil. Trans. Roy. Soc. Lond. A 377, 20190038 (2019).
- [19] G. Aad et al. (ATLAS), Phys. Rev. Lett. 124, 031802 (2020), arXiv:1905.10130 [hep-ex].
- [20] B. Acharya et al. (MoEDAL), Phys. Rev. Lett. 123, 021802 (2019), arXiv:1903.08491 [hep-ex].
- [21] E. Witten, Phys. Lett. B 86, 283 (1979).
- [22] W. Fischler and J. Preskill, Phys. Lett. B 125, 165 (1983).
- [23] A. V. Sokolov and A. Ringwald, (2022), arXiv:2205.02605 [hep-ph].
- [24] A. V. Sokolov and A. Ringwald, (2023), 10.1002/andp.202300112, arXiv:2303.10170 [hep-ph].
- [25] B. Heidenreich, J. McNamara, and M. Reece, (2023), arXiv:2309.07951 [hep-ph].
- [26] T. Li, R.-J. Zhang, and C.-J. Dai, JHEP 03, 088 (2023), arXiv:2211.06847 [hep-ph].
- [27] M. E. Tobar, C. A. Thomson, B. T. McAllister, M. Goryachev, A. Sokolov, and A. Ringwald, Annalen Phys. 2023, 2200594 (2022), arXiv:2211.09637 [hep-ph].
- [28] B. T. McAllister, A. Quiskamp, C. O'Hare, P. Altin, E. Ivanov, M. Goryachev, and M. Tobar, Annalen Phys. 2023, 2200622 (2022), arXiv:2212.01971 [hep-ph].
- [29] T. Li, C.-J. Dai, and R.-J. Zhang, (2023), arXiv:2304.12525 [hep-ph].
- [30] T. Li and R.-J. Zhang, (2023), arXiv:2305.01344 [hep-ph].
- [31] M. E. Tobar, A. V. Sokolov, A. Ringwald, and M. Goryachev, Phys. Rev. D **108**, 035024 (2023), arXiv:2306.13320 [hep-ph].
- [32] A. Patkos, (2023), arXiv:2309.05523 [hep-ph].
- [33] P. Agrawal, K. V. Berghaus, J. Fan, A. Hook, G. Marques-Tavares, and T. Rudelius, in <u>Snowmass 2021</u> (2022) arXiv:2203.08026 [hep-ph].
- [34] J. E. Kim, Phys. Rev. Lett. 43, 103 (1979).
- [35] M. A. Shifman, A. I. Vainshtein, and V. I. Zakharov, Nucl. Phys. B 166, 493 (1980).
- [36] W. S. Bae and Y. M. Cho, J. Korean Phys. Soc. 46, 791 (2005), arXiv:hep-th/0210299.

- [37] Y. M. Cho, K. Kimm, and J. H. Yoon, Mod. Phys. Lett. A 31, 1650053 (2016), arXiv:1212.3885 [hep-ph].
- [38] Y. M. Cho, K. Kim, and J. H. Yoon, Eur. Phys. J. C 75, 67 (2015), arXiv:1305.1699 [hep-ph].
- [39] W. H. Press, S. A. Teukolsky, W. T. Vetterling, and B. P. Flannery, <u>Numerical Recipes 3rd Edition: The Art of Scientific Computing</u>, 3rd ed. (Cambridge University Press, 2007).