# **Relation between PLS and OLS regression in terms of the eigenvalue distribution of the regressor covariance matrix**

**David del Val<sup>1</sup> | José R. Berrendero<sup>2</sup> | Alberto Suárez<sup>3</sup>**

<sup>1</sup>Departamento de Matemáticas and Departamento de Ingeniería Informática, Universidad Autónoma de Madrid <sup>2</sup>Departamento de Matemáticas, Universidad Autónoma de Madrid, and Instituto de Ciencias Matemáticas ICMAT (CSIC-UAM-UCM-UC3M) <sup>3</sup>Departamento de Ingeniería Informática, Escuela Politécnica Superior, Universidad Autónoma de Madrid

#### **Correspondence**

Email: delvaldavid1@gmail.com (D. del Val) Email: joser.berrendero@uam.es (J.R. Berrendero) Email: alberto.suarez@uam.es (A. Suárez)

## **Summary**

Partial least squares (PLS) is a dimensionality reduction technique introduced in the field of chemometrics and successfully employed in many other areas. The PLS components are obtained by maximizing the covariance between linear combinations of the regressors and of the target variables. In this work, we focus on its application to scalar regression problems. PLS regression consists in finding the least squares predictor that is a linear combination of a subset of the PLS components. Alternatively, PLS regression can be formulated as a least squares problem restricted to a Krylov subspace. This equivalent formulation is employed to analyze the distance between  $\hat{\beta}_\text{PLS}^{(L)}$ , the PLS estimator of the vector of coefficients of the linear regression model based on  $L$ PLS components, and  $\hat{\beta}_{\text{OLS}}$ , the one obtained by ordinary least squares (OLS), as a function of L. Specifically,  $\hat{\beta}_{\text{PLS}}^{(L)}$  is the vector of coefficients in the aforementioned Krylov subspace that is closest to  $\hat{\beta}_{\text{OLS}}$  in terms of the Mahalanobis distance with respect to the covariance matrix of the OLS estimate. We provide a bound on this distance that depends only on the distribution of the eigenvalues of the regressor covariance matrix. Numerical examples on synthetic and real-world data are used to illustrate how the distance between  $\hat{\beta}_{\rm PLS}^{(L)}$  and  $\hat{\beta}_{\rm OLS}$  depends on the number of clusters in which the eigenvalues of the regressor covariance matrix are grouped.

#### **KEYWORDS:**

dimensionality reduction, linear regression, partial least squares, Krylov subspaces

## **1 INTRODUCTION**

Partial least squares (PLS) is a family of dimensionality reduction methods introduced in the field of chemometrics [\(Noonan & Wold 1977\)](#page-10-0), where it is extensively used [\(Frank & Friedman 1993;](#page-10-1) [S. Wold, Sjöström, & Eriksson 2001\)](#page-11-0). Its success in this discipline has led to its adoption in other scientific areas such as medicine [\(Nguyen & Rocke 2002;](#page-10-2) [Worsley 1997\)](#page-11-1), physiology [\(Lobaugh, West, & McIntosh 2001\)](#page-10-3), and pharmacology [\(Nilsson,](#page-10-4) [de Jong, & Smilde 1997\)](#page-10-4). In PLS, two blocks of random variables  $X$  and  $Y$  are considered. The PLS components are built in a stepwise manner by maximizing the covariance between linear combinations of the components of  $X$  and of  $Y$  and imposing orthogonality to the previously identified components. In variants of PLS used only for dimensionality reduction,  $X$  and  $Y$  are handled in a symmetric manner. As a result, the maximum number of components is limited by the block that has the lowest dimension. When PLS is employed for regression, non-symmetric variants are used to account for the distinct roles played by the two blocks: predictor  $(X)$  and response  $(Y)$  variables. In contrast to the symmetric case, the number of components is limited only by the dimensionality of  $X$ , a feature that is of particular relevance in scalar regression problems.

In this paper, we apply PLS to a random vector  $X$  with  $D$  components and a scalar random variable  $Y$ . In this setting, PLS extracts a sequence of  $L\leq D$  (random) components  $\{t_l\}_{l=1}^L$ , each of which is a linear combination of the coordinates of  $X$ . The PLS components are orthogonal directions along which the covariance with the response variable, Y, is maximal. This optimization criteria is closely related to the ones used in other linear dimensionality reduction techniques such as principal component analysis (PCA) and canonical correlation analysis (CCA). The CCA components are identified by maximizing the correlation with the response variable, instead of the covariance. In PCA, the principal components are defined solely in terms of the regressor variables: they are linear combinations of the coordinates of X, obtained sequentially by maximizing the variance in the space orthogonal to the one spanned by the previously identified components. The covariance of  $t_l$ , the  $l$ -th PLS component,

<span id="page-1-1"></span>and the response variable can be expressed in terms of the correlation and the corresponding variances:

$$
cov(t_l, Y)^2 = var(t_l)corr(t_l, Y)^2 var(Y).
$$

This expression makes it clear that the PLS objective  $(\mathrm{cov}(t_l,Y))$  combines the optimization objective of PCA  $(\mathrm{var}(t_l))$  and CCA  $(\mathrm{corr}(t_l,Y))$ .

PLS was originally introduced in [Noonan and Wold](#page-10-0) [\(1977\)](#page-10-0). In that work, the PLS components are defined computationally as the result of applying the NIPALS (non-linear iterative partial least squares) algorithm. For scalar  $Y$ , the components identified by NIPALS are the solution of a constrained optimization problem. This problem consists in finding orthogonal linear combinations of the coordinates of  $X$  that maximize the covariance with the response variable [\(de Jong 1993\)](#page-10-5). Also in the case of scalar response, PLS regression with  $L$  components can be formulated as a least squares problem restricted to the Krylov subspace of order L generated by  $cov(X, X)$ , the covariance matrix of X, and  $cov(X, Y)$ , the vector of cross-covariances between X and Y [\(Helland 1988\)](#page-10-6). As shown in Section [4,](#page-4-0)  $\hat{\beta}_{\rm PLS}^{(L)}$  is the estimator in the Krylov subspace of order  $L$  that is closest to  $\hat{\beta}_{\text{OLS}}$ , the ordinary least squares (OLS) estimator, in terms of the Mahalanobis distance with the covariance matrix of the OLS estimator. Moreover, the PLS estimator is equal to the OLS one when L is the number of distinct eigenvalues of  $cov(X, X)$ . Given this equivalency, it is also possible to show that the conjugate gradient method [\(S. Wold, Ruhe, Wold, & Dunn 1984\)](#page-11-2) and the Lanczos bidiagonzalization algorithm [\(Eldén](#page-10-7) [2004\)](#page-10-7) can be used as an alternative to NIPALS for PLS regression. Another contribution of this work is to utilize the properties of the conjugate gradient method [\(Hestenes & Stiefel 1952;](#page-10-8) [Nocedal & Wright 1999\)](#page-10-9) to quantify the differences between the PLS and OLS estimators of the vector of regression coefficients, by establishing an equivalency between PLS and a polynomial fitting problem. From this reformulation of the problem it is possible to derive an upper bound for the distance between these estimators that depends only on the spectrum of  $cov(X, X)$ . In light of this analysis, we explore the relation between these estimators in terms of the characteristics of the distribution of eigenvalues. In particular, if these eigenvalues are grouped into k tight clusters, PLS with  $k$  components provides a good approximation to OLS.

Finally, we carry out an empirical comparison of PCR (principal components regression) and PLS regression. This comparison shows that PLS and PCR are optimal for different types of eigenvalue distributions. PLS performs best when the eigenvalues are clustered around a few values. In contrast, PCA works best in the presence of a few dominant eigenvalues.

The article is organized as follows: In Section [2,](#page-1-0) PLS is introduced as a dimensionality reduction method. The NIPALS algorithm for PLS regression is also detailed, and its properties are analyzed. The use of PLS in regression is described in Section [3.](#page-2-0) Specifically, we provide a novel derivation of the equivalence between the standard formulation, as a least squares regression problem in the space spanned by the first PLS components, and one based on solving a least squares problem restricted to a Krylov space. In Section [4,](#page-4-0) the differences between the PLS estimator of the vector of regressor coefficients and the OLS one are quantified in terms of the Mahalanobis distance with the covariance matrix of the OLS estimator. Proofs of some of these relations are given in the Appendix. In Section [5,](#page-6-0) the results of a numerical investigation of the performance of PLS are presented for different scenarios using synthetic and real-world regression problems. Finally, the conclusions of this work are presented in Section [6.](#page-8-0)

## <span id="page-1-0"></span>**2 DIMENSIONALITY REDUCTION WITH PARTIAL LEAST SQUARES**

Consider the sample  $\{(\mathbf{x}_i, y_i)\}_{i=1}^N,$  whose  $i$ -th element is characterized by  $\mathbf{x}\in\mathbb{R}^D$  and  $\mathbf{y}_i\in\mathbb{R}.$  The goal of PLS is to identify a set of components  $\{t_l\}_{l=1}^L$  that capture linear relations between X and Y. Originally, the PLS components were introduced as the output of the NIPALS algorithm, which will be described later in this section. In [de Jong \(1993\)](#page-10-5), it is shown that the  $l$ -th PLS component is the solution of the following optimization problem:

$$
\mathbf{t}_{l} = \underset{\mathbf{t}}{\operatorname{argmax}} \quad \text{cov}(\mathbf{t}, \mathbf{y}) \quad \text{subject to} \quad \mathbf{t} = \mathbf{X}\mathbf{r}, \ \mathbf{r} \in \mathbb{R}^{D}, \ \|\mathbf{r}\| = 1; \\
\mathbf{t}^{\top}\mathbf{t}_{i} = 0 \quad i = 1, \dots, \ l - 1,
$$
\n(1)

where  $\mathbf{X}=(\mathbf{x}_1,\dots\mathbf{x}_N)^\top\in\mathbb{R}^{N\times D}$  and  $\mathbf{y}=(y_1,\dots y_N)^\top.$  Associated with the components, the weight vectors  $\{\mathbf{r}_l\}_{l=1}^L$  are defined so that  $\|\mathbf{r}_l\|=1$  and  $\mathbf{t}_l=\mathbf{X}\mathbf{r}_l$ , for  $l=1,\ldots,L$ .

The PLS components can be computed using different algorithms. The NIPALS algorithm, which we will describe in the remainder of this section, was the first one introduced and is still widely used today. This algorithm was first introduced in [Noonan and Wold](#page-10-0) [\(1977\)](#page-10-0), and has been the object of successive refinements [\(Wegelin 2000\)](#page-10-10). In this work we focus on the NIPALS algorithm for scalar PLS regression [\(Eldén 2004;](#page-10-7) [Rosipal &](#page-10-11) [Krämer 2005\)](#page-10-11). NIPALS follows an iterative approach. At the end of each iteration, X, the data matrix, is modified by removing the projection on the component computed in that iteration (line 7). As a result, a sequence of projections of the data matrix can be considered:  $\{{\bf X}_l\}_{l=1}^L.$  This deflation step ensures that subsequent components computed by the algorithm are orthogonal to the ones extracted up to that point. In particular, from lines 6 and 7,  $\mathbf{X}_l=\Big(\mathbf{I}-\frac{\mathbf{t}_l\mathbf{t}_l^\top}{\mathbf{t}_l^\top\mathbf{t}_l}\Big)\mathbf{X}_{l-1},$  for  $l=1,\ldots L.$ 

The *l*-th component can be computed as  $\mathbf{t}_l = \mathbf{X}\mathbf{r}_l$ , the projection of the original data onto the direction defined by the vector of weights  $\mathbf{r}_l$ . Alternatively, it is  ${\bf t}_l = {\bf X}_{l-1}{\bf w}_l$ , where  ${\bf w}_l$  is the  $l$ -th weight vector extracted by NIPALS. Finally, the regressor and response loadings are defined as  $\mathbf{p}_l = \mathbf{X}_{l-1}^{\top}\mathbf{t}_l/\|\mathbf{t}_l\|^2$  and  $q_l = \mathbf{y}^{\top}\mathbf{t}_l/\|\mathbf{t}_l\|^2$ , for  $l = 1, \ldots, L$ , respectively.

<span id="page-2-5"></span>

From this algorithm, we can derive a series of properties. The ones relevant for the rest of this paper are included in the following propositions, whose proofs are given in the appendix.

**Proposition 1.** From the NIPALS algorithm, the following properties can be derived:

1. In terms of the PLS components, the original data can be expressed as

<span id="page-2-4"></span>
$$
\mathbf{X} = \mathbf{T}_L \mathbf{P}_L^\top + \mathbf{X}_L, \qquad \mathbf{y} = \mathbf{T}_L \mathbf{Q}_L^\top + \mathbf{y}_L,\tag{2}
$$

3

where  $\mathbf{X}_L \in \mathbb{R}^{N \times D}$  and  $\mathbf{y}_L \in \mathbb{R}^N$  are defined as

<span id="page-2-6"></span>
$$
\mathbf{X}_L = \prod_{i=1}^L \left( \mathbf{I} - \frac{\mathbf{t}_i \mathbf{t}_i^\top}{\mathbf{t}_i^\top \mathbf{t}_i} \right) \mathbf{X}, \qquad \mathbf{y}_L = \prod_{i=1}^L \left( \mathbf{I} - \frac{\mathbf{t}_i \mathbf{t}_i^\top}{\mathbf{t}_i^\top \mathbf{t}_i} \right) \mathbf{y}.
$$

Additionally,  $\mathbf{T}_L$ ,  $\mathbf{P}_L$  and  $\mathbf{Q}_L$  are defined as  $\mathbf{T}_L = (\mathbf{t}_1, \ldots, \mathbf{t}_L) \in \mathbb{R}^{N \times L}$ ,  $\mathbf{P}_L = (\mathbf{p}_1, \ldots, \mathbf{p}_L) \in \mathbb{R}^{D \times L}$ , and  $\mathbf{Q}_L = (q_1, \ldots, q_L) \in \mathbb{R}^{1 \times L}$ .

- 2. The Frobenius norms of  $\mathbf{X}_L$  and  $\mathbf{y}_L$  decrease as  $L$  increases.
- 3. After L iterations,  $X_L$  is orthogonal to the weights:  $X_L \mathbf{W}_L = 0$ , where  $\mathbf{W}_L = (\mathbf{w}_1, \dots \mathbf{w}_L) \in \mathbb{R}^{M \times L}$
- 4. The loading matrices  ${\bf P}_L$  and  ${\bf Q}_L$  can be expressed in terms of the components and the original data as  ${\bf P}_L$   $=$   $\, {\bf X}^\top {\bf T}_L {\bf D}_L^{-2}$  and  $\mathbf{Q}_L = \mathbf{y}^\top \mathbf{T}_L \mathbf{D}_L^{-2}$ , with  $\mathbf{D}_L = \mathsf{diag}\left(\|\mathbf{t}_1\|,\ldots,\|\mathbf{t}_L\|\right) \in \mathbb{R}^{L \times L}.$

NIPALS calculates both the components and the weights needed to express the components as projections of the deflated  $X_L$  data matrices. However, it is advantageous to express the components as a projection of the original data  $X$  to simplify the resulting expressions. The following proposition, whose proof can be found in the appendix, provides an expression for these projection directions

<span id="page-2-1"></span>**Proposition 2.** The matrix  $\mathbf{R}_L \in \mathbb{R}^{D \times L}$  that fulfills  $\mathbf{T}_L = \mathbf{X} \mathbf{R}_L$  is  $\mathbf{R}_L = \mathbf{W}_L (\mathbf{P}_L^\top \mathbf{W}_L)^{-1}$ .

## <span id="page-2-0"></span>**3 PARTIAL LEAST SQUARES REGRESSION WITH SCALAR RESPONSE**

Consider the linear regression model  $Y=\bm{\beta}^\top X + \epsilon,$  where  $X$  is the regressor vector with  $D$  components,  $\bm{\beta}$  is the vector of coefficients, which needs to be estimated,  $\epsilon$  is random noise independent of X, and Y is the scalar response. For the sake of simplicity, and without loss of generality, both  $X$  and  $Y$  are assumed to have zero mean. To fit this model,  $N$  independent observations drawn from this model are available:  $\{(\mathbf{x}_i, y_i)\}_{i=1}^N$ We further assume that  $\{\epsilon_i\}_{i=1}^N$  are iid with variance  $\sigma^2.$  In this setting, we seek to estimate a vector of coefficients  $\bm{\beta}$  such that  $y_i=\bm{\beta}^\top\mathbf{x}_i+\epsilon_i,$ where  $i = 1, \ldots, N$ . These equations can be grouped row-wise into the matrix equation

<span id="page-2-3"></span>
$$
y = X\beta + \epsilon, \tag{4}
$$

where  $\mathbf{y} = (y_1, \ldots, y_N)^\top \in \mathbb{R}^N$ ,  $\mathbf{X} = (\mathbf{x}_1, \ldots, \mathbf{x}_N)^\top \in \mathbb{R}^{N \times D}$  and  $\boldsymbol{\epsilon} = (\epsilon_1, \ldots, \epsilon_N)^\top \in \mathbb{R}^N$ .

One possible estimator for  $\beta$  is the ordinary least squares estimator (OLS), given by

<span id="page-2-2"></span>
$$
\hat{\boldsymbol{\beta}}_{\text{OLS}} = \underset{\boldsymbol{\beta} \in \mathbb{R}^D}{\text{argmin}} \|\mathbf{y} - \mathbf{X}\boldsymbol{\beta}\|^2 = (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y} = \boldsymbol{\Sigma}_{\text{XX}}^{-1} \boldsymbol{\Sigma}_{\text{XY}},
$$
\n(5)

where  $\|\cdot\|$  is the euclidean norm,  $\Sigma_{XX} = X^\top X$  is the empirical estimate of the covariance matrix of X scaled by the number of observations, and  $\Sigma_{XY} = \mathbf{X}^\top \mathbf{y}$  is the empirical estimate of the covariance matrix of X and Y scaled by the number of observations as well.

A different estimator of  $\beta$  is obtained using PLS regression. The first step is to extract L PLS components as described in the previous section. Then, a linear prediction is made in terms of these components:  $\sum_{l=1}^L\hat\gamma_l{}^{(L)}\mathbf{t}_l=\mathbf{T}_L\hat\gamma^{(L)},$  with  $\hat\gamma^{(L)}=\left(\hat\gamma_1^{(L)},\ldots,\hat\gamma_L^{(L)}\right)^\top$  determined by least squares as

$$
\hat{\boldsymbol{\gamma}}^{(L)} = \underset{\boldsymbol{\gamma} \in \mathbb{R}^L}{\operatorname{argmin}} \|\mathbf{y} - \mathbf{T}_L \boldsymbol{\gamma}\|^2 = (\mathbf{T}_L^\top \mathbf{T}_L)^{-1} \mathbf{T}_L^\top \mathbf{y} = \mathbf{D}_L^{-2} \mathbf{T}_L^\top \mathbf{y}.
$$
 (6)

The PLS estimator of  $\beta$  is obtained by expressing this linear predictor in terms of the original variables  $\texttt{T}_L\hat{\gamma}^{(L)}=\textbf{X}\hat{\beta}_{\text{PLS}}^{(L)}.$  Using the definition of  ${\bf R}_L$  from Proposition [2:](#page-2-1)  ${\bf X}\hat{\bf \beta}_{\rm PLS}^{(L)}={\bf T}_L\hat{\bf \gamma}^{(L)}={\bf X}{\bf R_L}\hat{\bf \gamma}^{(L)}$  . Therefore, the PLS estimator of the vector of regression coefficients is

<span id="page-3-0"></span>
$$
\hat{\boldsymbol{\beta}}_{\text{PLS}}^{(L)} = \mathbf{R}_L \hat{\boldsymbol{\gamma}}^{(L)} = \mathbf{R}_L \mathbf{D}_L^{-2} \mathbf{T}_L^{\top} \mathbf{y}.
$$
\n<sup>(7)</sup>

This estimator is then used to yield the linear prediction  $\mathbf{X}\hat{\bm{\beta}}_\text{PLS}^{(L)}$ , in terms of the original variables.

Alternatively,  $\hat{\beta}_{\rm PLS}^{(L)}$  can be viewed as the least squares estimator of  $\beta$  when the optimization is constrained to a Krylov subspace. Krylov subspaces are defined as follows:

**Definition 1.** The Krylov subspace of order  $L \leq D$  generated by the matrix  $A \in \mathbb{R}^{D \times D}$  and the vector  $b \in \mathbb{R}^D$ ,  $b \neq 0$  is

$$
\mathcal{K}_L(\mathbf{A}, \mathbf{b}) = \text{span}\{\mathbf{b}, \mathbf{A}\mathbf{b}, \dots, \mathbf{A}^{L-1}\mathbf{b}\}.
$$
 (8)

<span id="page-3-3"></span>**Theorem 1.** The PLS estimator with L components defined in [\(7\)](#page-3-0) is the solution to the least squares problem

<span id="page-3-1"></span>
$$
\hat{\boldsymbol{\beta}}_{\text{PLS}}^{(L)} = \underset{\boldsymbol{\beta} \in \mathcal{K}_L(\boldsymbol{\Sigma}_{\text{XX}}, \boldsymbol{\Sigma}_{\text{XY}})}{\operatorname{argmin}} ||\mathbf{y} - \mathbf{X}\boldsymbol{\beta}||^2, \tag{9}
$$

where  $\mathcal{K}_L(\Sigma_{XX}, \Sigma_{XY})$  is the Krylov subspace of order L generated by the matrix  $\Sigma_{XX}$  and the vector  $\Sigma_{XY}$ .

*Proof.* Assume that the columns of  $B_L \in \mathbb{R}^{D \times L}$  constitute a basis of the Krylov subspace  $\mathcal{K}_L(\Sigma_{XX}, \Sigma_{XY})$ . Then any  $\beta \in \mathcal{K}_L(\Sigma_{XX}, \Sigma_{XY})$  can be expressed as  $\beta = B_L \alpha$  for some  $\alpha \in \mathbb{R}^L$ . Thus, the constrained optimization problem given by [\(9\)](#page-3-1) can be transformed into an unconstrained optimization problem in  $\mathbb{R}^L$ :

<span id="page-3-2"></span>
$$
\underset{\beta \in \mathcal{K}_L(\Sigma_{\text{XX}}, \Sigma_{\text{XY}})}{\operatorname{argmin}} ||\mathbf{y} - \mathbf{X}\beta||^2 = \underset{\mathbf{\alpha} \in \mathbb{R}^L}{\operatorname{argmin}} ||\mathbf{y} - \mathbf{X}\mathbf{B}_L \mathbf{\alpha}||^2 = \mathbf{B}_L (\mathbf{B}_L^\top \mathbf{X}^\top \mathbf{X} \mathbf{B}_L)^{-1} \mathbf{B}_L^\top \mathbf{X}^\top \mathbf{y}.
$$
\n(10)

As shown in [Eldén](#page-10-7) [\(2004\)](#page-10-7), the columns of the matrix  $W_L$  obtained after L iterations of NIPALS constitute a basis of  $\mathcal{K}_L(\Sigma_{XX},\Sigma_{XY})$ . Therefore, [\(10\)](#page-3-2) holds for  ${\bf B}_L={\bf W}_L.$  It is then possible to show that  $\hat{\beta}_{\rm PLS}^{(L)}$  can be expressed in the form given by the rhs of (10) with  ${\bf B}_L={\bf W}_L.$  To this end, Propositions [1](#page-1-1) and [2](#page-2-1) are applied repeatedly to [\(7\)](#page-3-0):

$$
\hat{\beta}_{\text{PLS}}^{(L)} = \mathbf{R}_{L} \mathbf{D}_{L}^{-2} \mathbf{T}_{L}^{\top} \mathbf{y} = \mathbf{W}_{L} (\mathbf{P}_{L}^{\top} \mathbf{W}_{L}^{-1}) \mathbf{D}_{L}^{-2} \mathbf{R}_{L}^{\top} \mathbf{X}^{\top} \mathbf{y} = \mathbf{W}_{L} (\mathbf{P}_{L}^{\top} \mathbf{W}_{L})^{-1} \mathbf{D}_{L}^{-2} (\mathbf{W}_{L} (\mathbf{P}_{L}^{\top} \mathbf{W}_{L})^{-1})^{\top} \mathbf{X}^{\top} \mathbf{y} =
$$
\n
$$
= \mathbf{W}_{L} (\mathbf{W}_{L}^{\top} \mathbf{P}_{L} \mathbf{D}_{L}^{2} \mathbf{P}_{L}^{\top} \mathbf{W}_{L})^{-1} \mathbf{W}_{L}^{\top} \mathbf{X}^{\top} \mathbf{y} = \mathbf{W}_{L} (\mathbf{W}_{L}^{\top} \mathbf{X}^{\top} \mathbf{T}_{L} \mathbf{P}_{L}^{\top} \mathbf{W}_{L})^{-1} \mathbf{W}_{L}^{\top} \mathbf{X}^{\top} \mathbf{y} =
$$
\n
$$
= \mathbf{W}_{L} (\mathbf{W}_{L}^{\top} \mathbf{X}^{\top} (\mathbf{X} - \mathbf{X}_{L}) \mathbf{W}_{L})^{-1} \mathbf{W}_{L}^{\top} \mathbf{X}^{\top} \mathbf{y} = \mathbf{W}_{L} (\mathbf{W}_{L}^{\top} \mathbf{X}^{\top} \mathbf{X} \mathbf{W}_{L})^{-1} \mathbf{W}_{L}^{\top} \mathbf{X}^{\top} \mathbf{y},
$$
\nthen holds because of the orthogonality between  $\mathbf{X}_{L}$  and  $\mathbf{W}_{L}$  (Proposition 1)

where the last step holds because of the orthogonality between  $X_L$  and  $W_L$  (Proposition [1\)](#page-1-1).

Other approaches can be adopted to prove this theorem. In [Eldén](#page-10-7) [\(2004\)](#page-10-7), the proof is based on the relation of PLS with the Lanczos bidiagonalization algorithm. An alternative derivation is given in [Takane and Loisel](#page-10-12) [\(2016\)](#page-10-12), leveraging the properties of some bidiagonal and tridiagonal matrices in the NIPALS algorithm [\(Noonan & Wold 1977\)](#page-10-0). However, in this proof, only simpler relationship between the matrices that are defined in the NIPALS algorithm are needed.

The expression of the vector of PLS regression coefficients given by [\(9\)](#page-3-1) opens up the possibility of using numerical optimization algorithms that accept linear constraints to compute  $\hat{\beta}_{\rm PLS}^{(L)}$ . It suffices to minimize  $\|{\bf y}-{\bf X}\beta\|^2$  subject to  $\beta$  belonging to  $\mathcal{K}_L(\pmb{\Sigma}_{\sf XX},\pmb{\Sigma}_{\sf XY})$ . In particular, the conjugate gradient algorithm is an iterative algorithm that minimizes a quadratic form  $\psi(z) = \mathbf{z}^\top \mathbf{A} \mathbf{z} - \mathbf{b}^\top \mathbf{z}$  while exploring  $\mathcal{K}_L(\mathbf{A}, \mathbf{b})$  in the L-th iteration [\(Nocedal & Wright 1999\)](#page-10-9). Thus, the optimization problem in Theorem [1](#page-3-3) can be solved using the conjugate gradient algorithm with  $A = \Sigma_{XX}$  and  $\mathbf{b} = \Sigma_{XY}$ . In the next section, we take advantage of this observation to study how the PLS estimator approximates the OLS one.

Additionally, the following theorem establishes an important link between the OLS and the PLS estimators.

<span id="page-3-4"></span>**Theorem 2.** The OLS estimator is contained in  $K_M(\Sigma_{XX}, \Sigma_{XY})$ , where M is the number of distinct eigenvalues of  $\Sigma_{XX}$ .

*Proof.* As a consequence of the Cayley-Hamilton theorem (e.g. [Bronson & Costa 2009,](#page-10-13) p.220), since Σ<sub>XX</sub> is a non-singular symmetric matrix, there exists a polynomial  $P_{\Sigma_{XX}}$  of degree  $M-1$  such that  $P_{\Sigma_{XX}}(\Sigma_{XX})\Sigma_{XX} = I$ , where M is the number of different eigenvalues of  $\Sigma_{XX}$ . Applying this result to the usual formula of OLS, we obtain  $\hat{\beta}_{\text{OLS}} = (\mathbf{X}^{\top} \mathbf{X})^{-1} \mathbf{X}^{\top} \mathbf{y} = P_{\Sigma_{XX}}(\Sigma_{XX}) \Sigma_{XY} \in \mathcal{K}_M(\Sigma_{XX}, \Sigma_{XY}).$  $\Box$ 

 $\Box$ 

 $\Box$ 

**Corollary 1.** The PLS estimator coincides with the OLS estimator after M iterations, where M is the number of different eigenvalues of  $\Sigma_{XX}$ :

$$
\hat{\boldsymbol{\beta}}_{\text{PLS}}^{(M)} = \hat{\boldsymbol{\beta}}_{\text{OLS}}.
$$
\n(11)

*Proof.* It is a direct consequence of Theorem [2](#page-3-4) and the definition of  $\hat{\beta}_{\rm PLS}^{(L)}$  as a restricted least squared estimator in Theorem [1.](#page-3-3)

## <span id="page-4-0"></span>**4 RELATION BETWEEN PARTIAL LEAST SQUARES AND ORDINARY LEAST SQUARES**

As described in the previous section, the PLS estimator of the vector of coefficients of a linear regression model with L components converges to the ordinarly least squares estimator as L increases. Furthermore, they coincide when  $L \geq M$ , the number of distinct eigenvalues of  $\Sigma_{XX}$ . The goal of this section is to provide an upper bound for the distance between  $\hat{\beta}_{\rm PLS}^{(L)}$  and  $\hat{\beta}_{\rm OLS}$ . In order to do so, we take advantage of the formulation of PLS in Theorem [1,](#page-3-3) as a constrained optimization problem that can be solved using conjugate gradients. The first part of this section follows the convergence analysis for the conjugate gradient method in [Nocedal and Wright](#page-10-9) [\(1999\)](#page-10-9). First, the PLS estimator is defined as the solution of yet another optimization problem in which a distance to the OLS estimator is minimized subject to some constrains.

<span id="page-4-1"></span>Proposition 3. The PLS estimator of the vector of coefficients of a linear regression model with L components is the solution to the optimization problem

<span id="page-4-2"></span>
$$
\hat{\boldsymbol{\beta}}_{\mathrm{PLS}}^{(L)} = \underset{\boldsymbol{\beta} \in \mathcal{K}_{L}(\boldsymbol{\Sigma}_{\mathrm{XX}}, \boldsymbol{\Sigma}_{\mathrm{XY}})}{\operatorname{argmin}} \left\| \boldsymbol{\beta} - \hat{\boldsymbol{\beta}}_{\mathrm{OLS}} \right\|_{\boldsymbol{\Sigma}_{\mathrm{XX}}}^{2},\tag{12}
$$

where  $\|{\bf z}\|_{\bf \Sigma_{XX}}^2 = {\bf z}^\top \bf \Sigma_{XX} {\bf z}$ , the square of the quadratic-form norm with the positive definite matrix  $\bf \Sigma_{XX}$ .

*Proof.* This result is a consequence of the definition of the PLS estimator with L components provided in Theorem [1:](#page-3-3)

$$
\hat{\beta}_{\text{PLS}}^{(L)} = \underset{\beta \in \mathcal{K}_L(\Sigma_{\text{XX}}, \Sigma_{\text{XY}})}{\operatorname{argmin}} ||\mathbf{y} - \mathbf{X}\beta||^2 = \underset{\beta \in \mathcal{K}_L(\Sigma_{\text{XX}}, \Sigma_{\text{XY}})}{\operatorname{argmin}} \left( \mathbf{y}^\top \mathbf{y} - 2\beta^\top \mathbf{X}^\top \mathbf{y} + \beta^\top \mathbf{X}^\top \mathbf{X}\beta \right) = \underset{\beta \in \mathcal{K}_L(\Sigma_{\text{XX}}, \Sigma_{\text{XY}})}{\operatorname{argmin}} \left( \beta^\top \mathbf{X}^\top \mathbf{X}\beta - 2\beta^\top \mathbf{X}^\top \mathbf{y} \right) =
$$
\n
$$
= \underset{\beta \in \mathcal{K}_L(\Sigma_{\text{XX}}, \Sigma_{\text{XY}})}{\operatorname{argmin}} \left( \beta^\top \Sigma_{\text{XX}} \beta - 2\beta^\top \Sigma_{\text{XX}} \hat{\beta}_{\text{OLS}} + \hat{\beta}_{\text{OLS}}^\top \Sigma_{\text{XX}} \hat{\beta}_{\text{OLS}} \right) = \underset{\beta \in \mathcal{K}_L(\Sigma_{\text{XX}}, \Sigma_{\text{XY}})}{\operatorname{argmin}} ||\beta - \hat{\beta}_{\text{OLS}}||_{\Sigma_{\text{XX}}}^2,
$$
\nhere we have used that  $\mathbf{X}^\top \mathbf{v} = \mathbf{X}^\top \mathbf{X} \hat{\beta}_{\text{OLS}} = \Sigma_{\text{XX}} \hat{\beta}_{\text{OLS}}$ ,

where we have used that  $X'y = X'X\beta_{OLS} = \Sigma_{XX}\beta_{OLS}$ .

The quadratic-form norm  $\|\cdot\|_{\Sigma_{XX}}$  is related to the Mahalanobis distance with the covariance matrix of the OLS estimator of  $\beta$  The following observation motivates the use of this norm as a natural way to quantify the differences between  $\hat\beta_{\rm PLS}^{(L)}$  and  $\hat\beta_{\rm OLS}$ .

**Corollary 2.** The PLS estimator of the vector of coefficients of a linear regression model with L components is the solution of the optimization problem

$$
\hat{\beta}_{\rm PLS}^{(L)} = \underset{\beta \in \mathcal{K}_L(\Sigma_{\rm XX}, \Sigma_{\rm XY})}{\text{argmin}} d_M(\beta, \hat{\beta}_{\rm OLS}),\tag{13}
$$

where  $d_M$  is the Mahalanobis distance with respect to the matrix  $\frac{1}{\sigma^2}\mathbf{\Sigma}_{\text{XX}}{}^{-1}$ , which is the covariance matrix of the OLS estimator of the regression coefficients conditioned to the observations of X.

*Proof.* From [\(5\)](#page-2-2), the variance of the OLS estimator conditioned to  $x_1,...x_N$  is  $C_{\text{OLS}} = \text{var}(\hat{\beta}_{\text{OLS}}|x_1,...x_N) = \sigma^2(\mathbf{X}^T\mathbf{X})^{-1}$ , where we have used that the  $\text{var}(\mathbf{y}|\mathbf{x}_1,\dots\mathbf{x}_N)=\text{var}(\bm{\epsilon})$ , and that the observations of  $\epsilon$  are iid random variables with variance  $\sigma^2$ . As a result, the squared Mahalanobis distance between the  $\hat{\beta}_{\text{OLS}}$  estimator and some other estimator  $\hat{\beta}$  can be expressed as

$$
d_M(\hat{\boldsymbol{\beta}}, \hat{\boldsymbol{\beta}}_{\text{OLS}})^2 = (\hat{\boldsymbol{\beta}} - \hat{\boldsymbol{\beta}}_{\text{OLS}})^\top \mathbf{C}_{\text{OLS}}^{-1} (\hat{\boldsymbol{\beta}} - \hat{\boldsymbol{\beta}}_{\text{OLS}}) = \frac{1}{\sigma^2} (\hat{\boldsymbol{\beta}} - \hat{\boldsymbol{\beta}}_{\text{OLS}})^\top (\mathbf{X}^\top \mathbf{X}) (\hat{\boldsymbol{\beta}} - \hat{\boldsymbol{\beta}}_{\text{OLS}}) = \frac{1}{\sigma^2} ||\hat{\boldsymbol{\beta}} - \hat{\boldsymbol{\beta}}_{\text{OLS}}||_{\mathbf{\Sigma}_{\text{XX}}}^2.
$$

Thus, the distance induced by the quadratic form norm  $\|\cdot\|_{\mathbf{\Sigma}_{\mathsf{XX}}}$  is proportional to the Mahalanobis distance with  $\sigma^2\mathbf{\Sigma}_{\mathsf{XX}}{}^{-1}$ , the covariance matrix of the OLS estimator.  $\Box$ 

Therefore, with L components, PLS finds the closest estimator to  $\hat{\beta}_{OLS}$  with respect to the Mahalanobis distance with the covariance matrix of the OLS estimator in the Krylov subspace characterized by  $\Sigma_{XX}$  and  $\Sigma_{XY}$  of order L. The Mahalanobis distance provides a natural measure of differences in the space of estimators, one that captures its geometry better than the Euclidean distance. For once, the Mahalanobis distance between the estimators is deeply related to the euclidean distance between the predictions:

$$
d_M(\hat{\boldsymbol{\beta}}_{\mathrm{PLS}}^{(L)},\hat{\boldsymbol{\beta}}_{\mathrm{OLS}})^2 = \frac{1}{\sigma^2} (\hat{\boldsymbol{\beta}}_{\mathrm{PLS}}^{(L)} - \hat{\boldsymbol{\beta}}_{\mathrm{OLS}})^{\top} (\mathbf{X}^{\top} \mathbf{X}) (\hat{\boldsymbol{\beta}}_{\mathrm{PLS}}^{(L)} - \hat{\boldsymbol{\beta}}_{\mathrm{OLS}}) = \frac{1}{\sigma^2} ||\hat{\mathbf{y}}_{\mathrm{OLS}} - \hat{\mathbf{y}}_{\mathrm{PLS}}||^2.
$$

Additionally, the structure of Krylov subspaces makes it possible to identify each element in a Krylov subspace of order  $L$  with a polynomial of order  $L - 1$ . As a result, the optimization problem in Proposition [3,](#page-4-1) is equivalent to the optimization problem given in the following Corollary:

<span id="page-5-5"></span>**Corollary 3.** The PLS estimator with  $L$  component is  $\hat{{\boldsymbol \beta}}_{\rm PLS}^{(L)} = P_{L-1}^*(\boldsymbol{\Sigma}_{\rm XX})\boldsymbol{\Sigma}_{\rm XY}$ , where

<span id="page-5-0"></span>
$$
P_L^* = \underset{P \in \mathcal{P}_L}{\text{argmin}} \left\| P(\Sigma_{XX}) \Sigma_{XY} - \hat{\boldsymbol{\beta}}_{\text{OLS}} \right\|_{\Sigma_{XX}}^2, \tag{14}
$$

and  $P_L$  is the space of polynomials of degree lower or equal to  $L$ .

*Proof.* Since  $\hat{\bm{\beta}}_{\rm PLS}^{(L)} \in \mathcal{K}_L(\bm{\Sigma}_{\rm XX},\bm{\Sigma}_{\rm XY})$ , it can be expressed as  $\hat{\bm{\beta}}_{\rm PLS}^{(L)}=P_{L-1}^*(\bm{\Sigma}_{\rm XX})\bm{\Sigma}_{\rm XY}$ . By substituting this expression into [\(12\)](#page-4-2), we obtain [\(14\)](#page-5-0).

As stated in the following theorem, whose proof is given in the appendix, the difference between the PLS estimator and the OLS estimator can be expressed as an optimization problem in a space of polynomials:

<span id="page-5-1"></span>**Theorem 3.** The distance between the PLS estimator and the OLS estimator fulfills

$$
\left\|\hat{\beta}_{\rm PLS}^{(L)} - \hat{\beta}_{\rm OLS}\right\|_{\Sigma_{\rm XX}}^2 = \min_{Q_L \in \Omega_L} \sum_{d=1}^D Q_L(\lambda_d)^2 \lambda_d \xi_d^2,
$$
\n(15)

where  $\{\lambda_d\}_{d=1}^D$  are the eigenvalues of  $\pmb{\Sigma}_{\sf XX}$ ,  $\{\xi_d\}_{d=1}^D$  are the coefficients of the expansion of  $\hat{\bm{\beta}}_{\sf OLS}$  in  $\{\mathbf{u}_d\}_{d=1}^D,$  the basis of eigenvectors of  $\pmb{\Sigma}_{\sf XX}$ , and  $\Omega_L = \{Q_L \in \mathcal{P}_L : Q_L(0) = -1\}$ . Additionally, for each  $L$ , the minimum is reached for  $Q_L^*(t) = tP_L^*(t) - 1$ .

This theorem implies that any polynomial  $Q_L \in \Omega_L$  can be used to provide an upper bound for the distance between the OLS and PLS estimators:

$$
\left\|\hat{\boldsymbol{\beta}}_{\text{PLS}}^{(L)} - \hat{\boldsymbol{\beta}}_{\text{OLS}}\right\|_{\boldsymbol{\Sigma}_{\text{XX}}}^2 \le \sum_{d=1}^D Q_L(\lambda_d)^2 \lambda_d \xi_d^2, \quad \text{for all } Q_L \in \Omega_L.
$$
 (16)

Furthermore, it is possible to obtain an upper bound also in terms of the norm of the OLS estimator and of  $Q_L$  evaluated at the eigenvalues of  $\Sigma_{\rm XX}$ . **Corollary 4.** Given a function  $H: \Omega_L \to \mathbb{R}$  that, for any polynomial  $R \in \Omega_L$ , fulfills  $R(\lambda_d)^2 \leq H(R)$  over all  $d = 1, \ldots, D$ , and given a particular polynomial  $Q_L \in \Omega_L$ ,

$$
\left\|\hat{\beta}_{\rm PLS}^{(L)} - \hat{\beta}_{\rm OLS}\right\|_{\Sigma_{\rm XX}}^2 \le H(Q_L)\left\|\hat{\beta}_{\rm OLS}\right\|_{\Sigma_{\rm XX}}^2, \quad \text{for all } Q_L \in \Omega_L. \tag{17}
$$

*Proof.* From Theorem [3,](#page-5-1) and the condition  $R(\lambda_d)^2 \le H(R)$  for  $d = 1, ..., D$ ,

$$
\left\|\hat{\beta}_{\text{PLS}}^{(L)} - \hat{\beta}_{\text{OLS}}\right\|_{\boldsymbol{\Sigma}_{\text{XX}}}^2 = \min_{R \in \Omega_L} \sum_{d=1}^D R(\lambda_d)^2 \lambda_d \xi_d^2 \le \min_{R \in \Omega_L} H(R) \sum_{d=1}^D \lambda_d \xi_d^2 = \min_{R \in \Omega_L} H(R) \left\|\hat{\beta}_{\text{OLS}}\right\|_{\boldsymbol{\Sigma}_{\text{XX}}}^2 \le H(Q_L) \left\|\hat{\beta}_{\text{OLS}}\right\|_{\boldsymbol{\Sigma}_{\text{XX}}}^2.
$$

Therefore, by choosing an H function and a specific polynomial  $Q_L$ , an upper bound on the PLS error can be obtained. There are different choices for  $H.$  In [Nocedal and Wright](#page-10-9) [\(1999\)](#page-10-9), a number of results are given using the upper bound  $H_1(Q_L)=\max_d Q_L(\lambda_d)^2.$  However, this bound has a major disadvantage: it is not straightforward to calculate the polynomial  $Q_L$  that minimizes  $H_1$ . In the remainder of this section, the simpler upper bound  $H_2(Q_L)=\sum_{d=1}^D Q_L(\lambda_d)^2$  is considered. The following theorem provides an uper bound on the PLS error by calculating the polynomial in  $\Omega_L$  that minimizes  $H_2$ .

<span id="page-5-3"></span>**Theorem 4.** The following bound for the squared norm of the difference between the L-th PLS estimator and the OLS estimator holds:

<span id="page-5-2"></span>
$$
\left\| \hat{\boldsymbol{\beta}}_{\text{PLS}}^{(L)} - \hat{\boldsymbol{\beta}}_{\text{OLS}} \right\|_{\boldsymbol{\Sigma}_{\text{XX}}}^2 \le C_L \left\| \hat{\boldsymbol{\beta}}_{\text{OLS}} \right\|_{\boldsymbol{\Sigma}_{\text{XX}}}^2, \tag{18}
$$

where

$$
C_L = D(1 - \mathbf{c}_L^{\top} \mathbf{H}_L^{-1} \mathbf{c}_L), \qquad \mathbf{H}_L = \begin{pmatrix} \mu'_2 & \cdots & \mu'_{L+1} \\ \vdots & \ddots & \vdots \\ \mu'_{L+1} & \cdots & \mu'_{2L} \end{pmatrix}, \qquad \mathbf{c}_L = \begin{pmatrix} \mu'_1 \\ \vdots \\ \mu'_L \end{pmatrix},
$$
(19)

and  $\mu'_l$  is the  $l$ -th raw moment of the distribution of the eigenvalues of  $\boldsymbol{\Sigma}_{\mathsf{XX}}$ .

This result provides an upper bound for the distance between  $\hat{\beta}_{\rm PLS}^{(L)}$  and  $\hat{\beta}_{\rm OLS}$  that depends only on the distribution of the eigenvalues of the regressor covariance matrix. Explicit expressions of this bound can be derived for PLS regression with one and two components.

<span id="page-5-4"></span>**Corollary 5.** The bounds given in [\(18\)](#page-5-2) for  $L = 1$  and  $L = 2$  can be expressed as a function of the coefficient of variation  $(c_v = \sigma/\mu)$ , the coefficient of asymmetry ( $\gamma$ ) and the kurtosis ( $\kappa$ ) of the eigenvalues of  $\Sigma_{XX}$ .

$$
C_1 = D \frac{c_v^2}{1 + c_v^2}, \qquad C_2 = D \frac{c_v^4 (\kappa - \gamma^2 - 1)}{(\kappa - \gamma^2) c_v^4 + (\kappa - 3 - 2\gamma) c_v^3 - 2\gamma c_v + 1}.
$$
\n(20)

*Proof.* These identities are obtained by expressing the raw moments that appear in  $C_L$  in terms of  $\mu$ ,  $\sigma$ ,  $\gamma$  and  $\kappa$ , and then simplifying the resulting formulas. $\Box$ 

Additionally, from Pearson's inequality ( $\kappa\geq 1+\gamma^2$ ), the quantity  $\kappa-\gamma^2-1$  is non-negative [\(Sharma & Bhandari 2015\)](#page-10-14). This quantity is zero for dichotomous distributions. Thus,  $C_2$  should be small when the eigenvalues are distributed in two tightly packed clusters. In the next section, we provide numerical illustrations of the dependence of distance between  $\hat{\beta}_{\rm PLS}^{(L)}$  and  $\hat{\beta}_{\rm OLS}$  as a function of  $L$ , for different distributions of the eigenvalues of  $\Sigma_{XX}$ .

### <span id="page-6-0"></span>**5 EMPIRICAL STUDY**

In this section, an empirical study is carried out to investigate the effect of the eigenvalue distribution of the regressor covariance matrix on PLS. Specifically, we analyze the dependence of the quadratic-form distance between  $\hat{\beta}_{\rm PLS}^{(L)}$  and  $\hat{\beta}_{\rm OLS}$ , the upper bound established in Theorem [4](#page-5-3) for this distance, and the accuracy of the linear predictor as a function of the number of PLS components considered. The analysis is first performed in regression problems with synthetic data for different forms of the distribution of eigenvalues. The corresponding analysis is then performed for the California Housing dataset [\(Kelley Pace & Barry 1997\)](#page-10-15).

### **5.1 Synthetic data**

In this section, synthetic data are used to illustrate the behavior of the PLS method depending on the eigenvalue distribution of the regressor covariance matrix. Five regression problems are considered. In these problems, X is modelled as a multivariate normal vector  $X \sim N(0, \Sigma)$ . The eigenvalues of the covariance matrix  $\bm\Sigma,$   $\{\lambda_d\}_{d=1}^D,$  are sampled from different distributions with specific characteristics. Specifically,  $D=30$ eigenvalues are selected with the following characteristics:

- 1. 30 equally spaced eigenvalues from 2.5 to 7.5.
- 2. One cluster of 30 eigenvalues sampled from  $N(5, 0.1)$ .
- 3. Two clusters of 15 eigenvalues, each sampled from  $N(2.5, 0.1)$  and  $N(7.5, 0.1)$ .
- 4. Three clusters of 10 eigenvalues sampled from  $N(2.5, 0.1)$ ,  $N(5, 0.1)$ , and  $N(7.5, 0.1)$ .
- 5. Three clusters of 10 eigenvalues sampled from  $N(0.2, 0.1)$ ,  $N(5, 0.1)$ , and  $N(7.5, 0.1)$ ; so that one of the clusters is very close to zero.

These eigenvalue distributions are displayed in Figure [1.](#page-12-0) The actual covariace matrix is generated by a random rotation of the diagonal eigenvalue matrix:  $\Sigma = \mathbf{Q}^\top \text{diag}(\lambda_1, \dots, \lambda_D) \mathbf{Q}$ , where  $\mathbf{Q}$  is a uniformly-distributed orthogonal random matrix. The rotation matrix  $\mathbf{Q}$  is obtained from the QR decomposition of a random matrix whose entries are sampled from a standard normal distribution [\(Mezzadri 2007\)](#page-10-16). Finally, the data matrix,  $\mathbf{X} = (\mathbf{x}_1, \dots, \mathbf{x}_N)^\top$  is obtained by stacking  $N = 1000$  samples from this random vector.

To generate the response data, the linear model with additive noise presented in [\(4\)](#page-2-3) is used. The  $\beta$  parameter is a random vector whose entries are sampled from a uniform distribution in  $[0,1].$  The noise  $\bm{\epsilon}$  is sampled from a  $N(0,\sigma^2)$  distribution, where  $\sigma=0.1\,\mathrm{std}(\mathbf{X}\bm{\beta}),$  so that the model is not dominated by the noise. Finally, the response vector is computed as  $\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}$ .

In the experiments carried out, the closeness between  $\hat\beta_{\rm PLS}^{(L)}$  and  $\hat\beta_{\rm OLS}$  is quantified in terms of the normalized estimator difference:

$$
NED_L = \frac{\|\hat{\beta}_{\rm PLS}^{(L)} - \hat{\beta}_{\rm OLS}\|_{\Sigma_{\rm XX}}^2}{\|\hat{\beta}_{\rm OLS}\|_{\Sigma_{\rm XX}}^2},\tag{21}
$$

From [\(18\)](#page-5-2), it is apparent that  $C_L$  is a bound on  $NED_L$ . The results reported are averages over 20 realizations of the data.

The plots in the left column of Figure [2](#page-13-0) display the dependence of the normalized differences between the estimators,  $NED<sub>L</sub>$ , and of the corresponding upper-bound,  $C_L$ , on L, the number of PLS components considered. These plots show how, as L increases, the decrease of the bound introduced in Theorem [4](#page-5-3) parallels that of the difference between the estimators. As discussed in the previous section, PLS can be formulated as a polynomial fitting problem. In particular, Theorem [3](#page-5-1) provides a way of expressing the error of the estimator with  $L$  iterations as a function of the values of some polynomial  $Q_L$ , of degree lower or equal to  $L$  that fulfills  $Q_L(0)=-1$ . The optimal polynomials  $Q_L^*$  defined in Theorem [3](#page-5-1) are plotted in the right column of Figure [2](#page-13-0) .

It is possible to interpret the features of the curves displayed in the left column of Figure [2](#page-13-0) from the characteristics of the polynomials plotted in the right column of this figure. In the first scenario, in which the eigenvalues are uniformly distributed in an interval separate from zero, considering

more components allows to find polynomials that fulfill  $Q_L(0) = -1$  and take small values for all the eigenvalues. In the second one, the decrease of  $NED<sub>L</sub>$  is much steeper because having the eigenvalues closely packed in a single cluster makes the polynomial fitting problem much simpler. For a given numbers of components, the corresponding polynomials take smaller values on the eigenvalues in the second scenario than in the first one. This result is consistent with the dependency of  $C_1$  and  $C_2$  on  $c_v$  given in Corollary [5.](#page-5-4)

Figure [2](#page-13-0) also shows that the decrease of NED<sub>L</sub> with L follows different patterns depending on the number of clusters in which the eigenvalues are grouped. In particular, the decrease is sharper for specific numbers of components. When the eigenvalues are grouped in two clusters, the first abrupt decrease of NED<sub>L</sub> occurs between  $L = 1$  and  $L = 2$  components. This observation can be explained by noting it is not possible to find a polynomial of degree one (i.e., a straight line) that takes small values on both clusters and passes through the point  $(0, -1)$ . However, a polynomial of degree two (i.e., a parabola) provides a reasonable fit. Significant improvements are observed also for  $L = 4$  and  $L = 6$  components. This is due to the fact that, in those cases, it is possible to find polynomials that pass through (0,-1) with equal numbers of roots located in the vicinity of each of the clusters. A similar analysis can be carried out for the fourth scenario, in which the eigenvalues are clustered in 3 groups. In this case, sharper improvements are found for  $L = 3$  and  $L = 6$ .

To complete the analysis, we consider a case in which one of the clusters of eigenvalues is close to 0. From the plot in the bottom left of Figure [2](#page-13-0) it is apparent that the decrease of NED<sub>L</sub> with L is rather slow. The reason for this is that, since the fitted polynomial has to go through (0, -1), large values of  $L$  are needed so that the polynomial can take simultaneously small values for the eigenvalues in the vicinity of 0 and in the other clusters.

We now compare the performance of PCA and PLS regression as a function of  $L$ , the number of components considered. The quality of the predictions is measured in terms of the coefficient of determination ( $R^2$  score), which represents the proportion of explained variance. In most regression problems PLS is expected to outperform PCA because, in the definition of the components, the correlations between the regressor and response variables are taken into account in the former, but not in the latter [\(Frank & Friedman 1993\)](#page-10-1). Since the properties of PLS depend on the distribution of the eigenvalues of  $\Sigma_{XX}$ , the regressor covariance matrix, we carry out the analysis for the five scenarios described earlier. In Figure [3](#page-14-0) we compare the curves that trace the dependence  $R^2$  on L, for PLS (left plots) and PCA (right plots) in the first two synthetic datasets. This comparison illustrates the differences between problems in which the eigenvalues of the regressor covariance matrix are uniformly distributed and problems in which they are clustered around a particular value, different from zero. As expected, PLS obtains better results when the eigenvalues concentrate around a single value. In fact, when they are clustered in a single tight group, the PLS regression model with only one component provides a very accurate prediction of the response. By contrast, when the eigenvalues are uniformly distributed, more components are needed. The behavior of PCA is markedly different. In the case of clustered eigenvalues, the  $R^2$  score of PCA increases linearly with the number of eigenvalues considered. This is to be expected since the increment in explained variance is proportional to the eigenvalue that corresponds to the eigenvector considered by PCA at each step. If the eigenvalues are spread out uniformly, PCA considers first the components that correspond to the larger eigenvalues. Therefore, the magnitude of the eigenvalues decreases as more components are considered, which leads to a reduction of the rate at which  $R^2$  increases for larger  $L$ . Additionally, Figure [3](#page-14-0) also shows that PCA needs many more components to achieve the same  $R^2$  scores that PLS.

The plots displayed in Figure [4](#page-14-1) illustrate the properties of the curves that trace the evolution of the  $R^2$  score as a function of  $L$ , depending on the number of clusters in which the eigenvalues are grouped. From these results we conclude that, in this case, the number of PLS components necessary to obtain a value of  $R^2$  close to 1 (perfect prediction) coincides with the number of eigenvalue clusters. This is consistent with the analysis of the differences between  $\hat{\beta}_{\rm PLS}^{(L)}$  and  $\hat{\beta}_{\rm OLS}$  for these datasets. Regarding PCA, we can see how the number of clusters of eigenvalues has only minor effects in dependence of the  $R^2$  scores with  $L.$  For example, with two clusters, the  $R^2$  increases faster during the first 15 iterations, which corresponds to the cluster with the largest 15 eigenvalues. For the scenario with three clusters of eigenvalues, the rate of increase drops after 10 and 20 components have been considered. These correspond to having included in the model all the components in the first, and in the first and second largest clusters, respectively.

Finally, we use the last two scenarios to investigate the impact of having a cluster of eigenvalues close to zero. Figure [5](#page-14-2) shows how that for  $L > 1$ , the performance of PLS deteriorates when there is a cluster of small eigenvalues. This is again to be expected from the theoretical analysis carried out in the previous section because of the difficulties of fitting a polynomial that goes throgh  $(0, 1)$  and takes small values at the locations of the eigenvalues in the clusters. By contrast, PCA achieves better results when a sizeable fraction of the eigenvalues are close to zero. In fact, the maximum value of  $R^2$  is attained for  $L=20,$  once all the components that correspond to eigenvalues significantly larger than zero have been selected. Nonetheless, PLS outperforms PCA regression also in these scenarios.

#### **5.2 The Californian Housing dataset**

In this section we analyze the properties of PLS regression for the California Housing dataset [Kelley Pace and Barry](#page-10-15) [\(1997\)](#page-10-15) In this problem, the goal is to predict the median house value in a particular block group in a California district using 8 attributes ( $D = 8$ ): the median house age, the average number of rooms, the average number of bedrooms, the number of people residing within the block, the average number of household members, and the latitude and longitude of the block group. As a preprocessing step, both the regressor vector and the response variable are centered so that they have zero mean. Each column of X is scaled so that it has unit variance. In the original dataset, a median house value of  $500,000\$ \$ is assigned to instances whose actual value is above that threshold. To avoid distortions associated to this thresholding, these examples have been discarded.

Figure [6](#page-15-0) shows the eigenvalue distribution of the regressor covariance matrix for the California Housing dataset. The eigenvalues are roughly grouped in three clusters, one of them close to zero. This pattern is similar to the last synthetic dataset analyzed in the previous section. However, the eigenvalues in the central cluster are more spread out. This dispersion hinders somewhat the performance of PLS, which is nonetheless fairly good. The differences between the PLS and the OLS estimators as a function of  $L$  are analyzed in Figur[e7.](#page-15-1) The left plot displays the dependence of these differences, quantified by  $NED<sub>L</sub>$ , and of  $C<sub>L</sub>$ , the upper bound of these differences derived in this work, as a funtion of L, the number of PLS components considered. Note that, for  $L = 8$  the PLS coincides with the OLS estimator. As expected, the distance between the estimators decreases slowly, because of the presence of the small eigenvalues and, to a lesser extent, the dispersion of the medium-sized eigenvalues.

Figure [8](#page-15-2) presents the results of a comparison between PCA and PLS regression. The left plot displays the curves that trace the dependence of the  $R^2$  score, a measure of the quality of the predictions, with  $L$ , the number of components considered. From these results one concludes that PLS obtains better results than PCA, and needs fewer components to achieve an accuracy comparable to OLS. The evolution of  $C<sub>L</sub>$  as a function of  $L$  is displayed in the right plot of this figure. Note that the descent of the bound mirrors the increase of the  $R^2$  score as  $L$  increases. This illustrates that the upper bound defined in Theorem [4](#page-5-3) provides an effective way to monitor the performance of PLS.

## <span id="page-8-0"></span>**6 CONCLUSIONS**

In this work, the relation between ordinary least squares (OLS) and partial least squares (PLS) regression has been established by analyzing a number of different but equivalent optimization problems. In the context of scalar regression, the PLS components are orthogonal linear combinations of the regressor variables  $(X)$  that maximize the covariance with the response variable  $(Y)$ . A linear predictor is then built by taking a linear combination of a subset of size  $L$  of the PLS components. The coefficients in this linear combination are obtained by least squares. The PLS predictor can be expressed also as a linear combination of the original regression variables. The estimate of the vector of regression coefficients in the original variables given by PLS with  $L$  components,  $\hat{\beta}_{\rm PLS}^{(L)}$ , is the solution of a restricted least squares problem in a Krylov subspace of order  $L$ generated by  $cov(X, X)$ , the covariance matrix of X, and  $cov(X, Y)$ , the vector of cross-covariances between X and Y. An important contribution of this work is to show that  $\hat{\beta}_{\rm PLS}^{(L)}$  is the vector of regression coefficients that is closest to the OLS estimator in this Krylov subspace. Closeness is measured in terms of the Mahalanobis distance with the covariance matrix of the OLS estimator, which is a natural measure of differences in the space of estimators. Finally, leveraging the connection between optimization in Krylov subspaces and conjugate gradients, PLS regression is related to a polynomial optimization problem. From this reformulation, we derive an upper bound for the differences between the PLS and the OLS estimators of the vector of regression coefficients. This bound depends only on the eigenvalue distribution of the covariance matrix of the regressor variables. In particular, PLS is expected to be most effective when the eigenvalues are not close to zero and appear tightly grouped in a few clusters. Furthermore, if the regressor covariance matrix has only  $M$  distinct eigenvalues, convergence would be reached after  $M$  steps. An empirical study using simulated data is carried out to analyze the effect of different types of eigenvalue distributions on the effectiveness of PLS. Finally, a similar studied is performed for the California Housing dataset. The results obtained illustrate the relevance of the theoretical analysis presented in this work and the advantages of PLS regression in real-world applications.

## **ACKNOWLEDGMENTS**

J.R.B. acknowledges financial support from Grant CEX2019-000904-S funded by MCIN/AEI/ 10.13039/501100011033, and Spanish Ministry of Education and Innovation project PID2019-109387GB-I00.

A.S. acknowledges financial support from project PID2022-139856NB-I00 funded by MCIN/ AEI / 10.13039/501100011033 / FEDER, UE and project PID2019-106827GB-I00 / AEI / 10.13039/501100011033 and from the Autonomous Community of Madrid (ELLIS Unit Madrid).

**How to cite this article:** del Val D. Berrendero JR. Suárez A. (2023), Relation between PLS and OLS regression in terms of the eigenvalue distribution of the regressor covariance matrix .

# **APPENDIX**

# **A PROOFS**

**Proof of Proposition [1.](#page-1-1)** The identity for X in [\(2\)](#page-2-4) is a direct consequence of substituting line 6 into line 7 of Algorithm [1.](#page-2-5) The corresponding identity for y can be derived in a similar manner once one notices that adding a deflation step for y at the end of each iteration would not affect the results of NIPALS. This holds true since  ${\bf y}$  is used only in the calculation of  ${\bf w}_l$ . After including the deflation step,  ${\bf w}_l = {\bf X}_{l-1}^\top {\bf y}_{l-1}/\|{\bf X}_{l-1}^\top {\bf y}_{l-1}\|$ . However,  $\mathbf{X}_{l-1}^\top \mathbf{y}_{l-1} = \mathbf{X}_{l-1}^\top \mathbf{y}$  since  $\mathbf{y}_{l-1} = \mathbf{y} - \sum_{i=1}^{l-1} \mathbf{t}_i q_i$ , and  $\mathbf{X}_{l-1}^\top \mathbf{t}_i = \mathbf{0}$  as long as  $i < l$ . Regarding the decrease of the Frobenius norm, this is a consequence of the expressions for  ${\bf X}_l$  and  ${\bf y}_l$  in [\(3\)](#page-2-6). We will prove the result for  ${\bf X}_l$  . From (3), one obtains  ${\bf X}_l = {\bf \Pi}_l {\bf X}_{l-1},$  where  ${\bf \Pi}_l = \left( {\bf I} - \frac{{\bf t}_l {\bf t}_l^\top}{{\bf t}_l^\top {\bf t}_l} \right)$ .

Then, to show the decrement of the norms, We need only show that  $||\mathbf{X}_l||_F \le ||\mathbf{X}_{l-1}||_F$  for  $1 \le l \le L$ :

$$
\|\mathbf{X}_{l}\|_{F} = \|\mathbf{\Pi}_{l}\mathbf{X}_{l-1}\|_{F} = \|\mathbf{U}_{l}\mathbf{S}_{l}\mathbf{U}_{l}^{\top}\mathbf{X}_{l-1}\|_{F} = \|\mathbf{S}_{l}\mathbf{U}_{l}^{\top}\mathbf{X}_{l-1}\|_{F} \leq \|\mathbf{U}_{l}^{\top}\mathbf{X}_{l-1}\|_{F} = \|\mathbf{X}_{l-1}\|_{F},
$$

where  $\bm{\Pi}_l=\textbf{U}_l\textbf{S}_l\textbf{U}_l^\top$  is the eigenvector decomposition of  $\bm{\Pi}_l$ . Since  $\bm{\Pi}_l$  is a real symmetric matrix,  $\textbf{U}_l$  is a unitary matrix and we can apply that the Frobenius norm is invariant under unitary operations. Additionally, since  $\bm{\Pi}_l$  is positive-definite and idempotent, its eigenvalues are either 0 or 1. Therefore,  $S_i$  has only 0s or 1s in the diagonal. As a result, multiplying by it can only reduce the Frobenius norm.

The orthogonality between  $X_L$  and W can be proven showing that  $X_Lw_l = 0$  if  $l \leq L$ . From [\(3\)](#page-2-6),

$$
\mathbf{X}_L \mathbf{w}_l = \left(\mathbf{I} - \frac{\mathbf{t}_L \mathbf{t}_L^\top}{\mathbf{t}_L^\top \mathbf{t}_L}\right) \dots \left(\mathbf{I} - \frac{\mathbf{t}_l \mathbf{t}_l^\top}{\mathbf{t}_l^\top \mathbf{t}_l}\right) \mathbf{X}_{l-1} \mathbf{w}_l = \left(\mathbf{I} - \frac{\mathbf{t}_L \mathbf{t}_L^\top}{\mathbf{t}_L^\top \mathbf{t}_L}\right) \dots \left(\mathbf{I} - \frac{\mathbf{t}_l \mathbf{t}_l^\top}{\mathbf{t}_l^\top \mathbf{t}_l}\right) \mathbf{t}_l = \mathbf{0}
$$

Regarding the expressions for the loadings, both identities can be proven in the same way. We will prove the identity for  $P$ , the  $X$  loadings, showing the equality for each column of both sides of the equation. This equality is, in turn, a consequence of the expression for  $X$  in [\(3\)](#page-2-6).

$$
\mathbf{X}^{\top}\mathbf{t}_{l}\|\mathbf{t}_{l}\|^{-2} = (\mathbf{T}_{l-1}\mathbf{P}_{l-1})^{\top}\mathbf{t}_{l}\|\mathbf{t}_{l}\|^{-2} + \mathbf{X}_{l-1}^{\top}\mathbf{t}_{l}\|\mathbf{t}_{l}\|^{-2} = \mathbf{P}_{l-1}^{\top}\mathbf{T}_{l-1}^{\top}\mathbf{t}_{l}\|\mathbf{t}_{l}\|^{-2} + \mathbf{X}_{l-1}^{\top}\mathbf{t}_{l}\|\mathbf{t}_{l}\|^{-2} = \mathbf{p}_{l},
$$

where  $\mathbf{T}_{l-1}^\top \mathbf{t}_l = \mathbf{0}$  because the extracted components are orthogonal.  $\qquad \qquad \Box$ 

**Proof of Proposition [2.](#page-2-1)** From Proposition [1,](#page-1-1)  $X_L W_L = 0$ . Applying this to the decomposition for X in [\(2\)](#page-2-4), we obtain:

$$
\mathbf{X}\mathbf{R}_L = (\mathbf{T}_L\mathbf{P}_L^\top + \mathbf{X}_L)(\mathbf{W}_L(\mathbf{P}_L^\top\mathbf{W}_L)^{-1}) = \mathbf{T}_L\mathbf{P}_L^\top\mathbf{W}_L(\mathbf{P}_L^\top\mathbf{W}_L)^{-1} + \mathbf{X}_L\mathbf{W}_L(\mathbf{P}_L^\top\mathbf{W}_L)^{-1} = \mathbf{T}_L.
$$

**Proof of Theorem [3.](#page-5-1)** Since  $\Sigma_{XX} = \mathbf{X}^\top \mathbf{X}$  is a real, symmetric matrix, it is possible to find a sequence of non-negative eigenvalues  $\{\lambda_1, \dots, \lambda_D\}$ and orthonormal eigenvectors:  $\{{\bf u}_1,\ldots,{\bf u}_D\}$  such that  $\bm\Sigma_\textsf{XX}=\sum_{d=1}^D\lambda_d{\bf u}_d{\bf u}_d^\top.$  This eigenvalue decomposition has three properties. First, the eigenvectors span the entire  $\R^D$  space. Therefore,  $D$  scalars  $\{\xi_d\}_{d=1}^D$  can be found such that  $\hat\beta_{\rm OLS}=\sum_{d=1}^D \xi_d\mathbf{u}_d.$  Second, the norm of a vector can be calculated as  $\|\mathbf{z}\|_{\mathbf{\Sigma}_{\mathsf{XX}}}^2 = \mathbf{z}^\top \mathbf{\Sigma}_{\mathsf{XX}} \mathbf{z} = \sum_{d=1}^D \lambda_d (\mathbf{u}_d^\top \mathbf{z})^2$  Third, for any polynomial  $P$ , it holds that  $P(\mathbf{\Sigma}_{\mathsf{XX}})\mathbf{u}_d = P(\lambda_d)\mathbf{u}_d$ , for  $d=1,\ldots,D$ . Using these properties we can now find an expression to calculate  $\left\|\hat{\boldsymbol{\beta}}_L-\hat{\boldsymbol{\beta}}_{\rm OLS}\right\|$ 2  $\sum_{\mathbf{\Sigma}_{\mathsf{XX}}}$  in terms of the polynomials  $P_L^*$ .

<span id="page-9-0"></span>
$$
\left\|\hat{\boldsymbol{\beta}}_L - \hat{\boldsymbol{\beta}}_{\text{OLS}}\right\|_{\boldsymbol{\Sigma}_{\text{XX}}}^2 = \left\|(P_{L-1}^*(\boldsymbol{\Sigma}_{\text{XX}})\boldsymbol{\Sigma}_{\text{XX}} - \mathbf{I})\hat{\boldsymbol{\beta}}_{\text{OLS}}\right\|_{\boldsymbol{\Sigma}_{\text{XX}}}^2 = \left\|\sum_{d=1}^D (P_{L-1}^*(\boldsymbol{\Sigma}_{\text{XX}})\boldsymbol{\Sigma}_{\text{XX}} - \mathbf{I})\xi_d\mathbf{u}_d\right\|_{\boldsymbol{\Sigma}_{\text{XX}}}^2 = \sum_{d=1}^D Q_L^*(\lambda_d)^2\lambda_d\xi_d^2,
$$
 (A1)

where  $Q_L^*(t)=tP_{L-1}^*(t)-1$ , a polynomial of degree lower or equal to  $L$  that fulfills  $Q_L^*(0)=-1$ . Additionally, Corollary [3](#page-5-5) shows that  $P_{L-1}^*$  is the polynomial that minimizes the RHS of [\(A1\)](#page-9-0) over all polynomials of degree lower or equal to  $L-1.$  Therefore,  $Q^*_L$  minimizes that same quantity over all the polynomials  $Q_L$  of degree at most L such that  $Q_L(0) = -1$ . That is to say, over  $\Omega_L$ .

**Proof of Theorem [4.](#page-5-3)** All polynomials in  $\Omega_L$  can be expressed as  $R_L(t) = -1 + a_1t + \cdots + a_Lt^L$  for some coefficients  $a_1,\ldots,a_L$ . Therefore, as a function of the coefficients of the polynomials, the bound can be expressed as  $h_L(a_1,\dots,a_L)=\sum_{d=1}^D\big(-1+a_1\lambda_d+\dots+a_L\lambda_d^L\big)^2.$  To minimize this function, we calculate its gradient and determine the coefficients for which it is zero:

$$
\frac{\partial h_L}{\partial a_l} = 2 \sum_{d=1}^D (-1 + a_1 \lambda_d + \dots + a_L \lambda_d^L) \lambda_d^l = -2 \sum_{d=1}^D \lambda_d^l + 2a_1 \sum_{d=1}^D \lambda_d^{l+1} + \dots + 2a_L \sum_{d=1}^D \lambda_d^{l+L} = 0, \qquad l = 1, \dots, L.
$$
 (A2)

By rewritting these equations in terms of the sample raw moments of the eigenvalues, we obtain  $a_1\mu'_{l+1}+\cdots+a_L\mu'_{l+l}=\mu'_l,$  for  $l=1,\ldots,L$ These equations can be expressed as the system  ${\bf H}_L {\bf a}_L = {\bf c}_L.$  Therefore, the coefficients that minimize  $h_L$  are  ${\bf a}_L^* = {\bf H}_L^{-1} {\bf c}_L.$  Additionally, we express  $h_L$  as

$$
h_L(a_1,\ldots,a_L) = (-1,\mathbf{a}_L) \begin{pmatrix} 1 & \ldots & 1 \\ \vdots & \ddots & \vdots \\ \lambda_1^L & \ldots & \lambda_D^L \end{pmatrix} \begin{pmatrix} 1 & \ldots & \lambda_1^L \\ \vdots & \ddots & \vdots \\ 1 & \ldots & \lambda_D^L \end{pmatrix} \begin{pmatrix} -1 \\ \mathbf{a}_L \end{pmatrix} = (-1,\mathbf{a}_L) \begin{pmatrix} D & D\mathbf{c}_L^\top \\ D\mathbf{c}_L & D\mathbf{H}_L \end{pmatrix} \begin{pmatrix} -1 \\ \mathbf{a}_L \end{pmatrix}.
$$
 (A3)

Substituting the expression for  $\mathbf{a}_L^*$  in the previous formula shows that  $h_L(\mathbf{a}_L^*)=(-1,\mathbf{H}_L^{-1}\mathbf{c}_L)=D(1-\mathbf{c}_L^\top\mathbf{H}_L^{-1}\mathbf{c}_L).$  Finally, note that the obtained coefficients  $\mathbf{a}^*_L$  define the polynomial  $R^*_L(t) = -1 + a_1^*t + \cdots + a_L^*t^L$ , which minimizes  $H_2$ .

#### **References**

Bro, R., & Eldén, L. (2009). PLS works. *Journal of Chemometrics*, *23*(2), 69–71. doi: 10.1002/cem.1177

- <span id="page-10-13"></span>Bronson, R., & Costa, G. B. (2009). 7 - Matrix Calculus. In R. Bronson & G. B. Costa (Eds.), *Matrix Methods (Third Edition)* (pp. 213–255). Boston: Academic Press. doi: 10.1016/B978-0-08-092225-6.50013-9
- <span id="page-10-5"></span>de Jong, S. (1993). SIMPLS: An alternative approach to partial least squares regression. *Chemometrics and Intelligent Laboratory Systems*, *18*(3), 251–263. doi: 10.1016/0169-7439(93)85002-X

Devore, J. (1987). *Probability and Statistics for Engineering and the Sciences*. Brooks/Cole Publishing Company.

- <span id="page-10-7"></span>Eldén, L. (2004). Partial least-squares vs. Lanczos bidiagonalization—I: Analysis of a projection method for multiple regression. *Computational Statistics and Data Analysis*, *46*(1), 11–31. doi: https://doi.org/10.1016/S0167-9473(03)00138-5
- <span id="page-10-1"></span>Frank, I. E., & Friedman, J. H. (1993). A Statistical View of Some Chemometrics Regression Tools. *Technometrics*, *35*(2), 109–135. doi: 10.2307/1269656
- Geladi, P. (1988). Notes on the history and nature of partial least squares (PLS) modelling. *Journal of Chemometrics*, *2*(4), 231–246. doi: 10.1002/cem.1180020403
- Geladi, P., & Kowalski, B. R. (1986). Partial least-squares regression: A tutorial. *Analytica Chimica Acta*, *185*, 1–17. doi: 10.1016/0003- 2670(86)80028-9
- <span id="page-10-6"></span>Helland, I. S. (1988). On the structure of partial least squares regression. *Communications in Statistics - Simulation and Computation*, *17*(2), 581–607. doi: 10.1080/03610918808812681
- <span id="page-10-8"></span>Hestenes, M. R., & Stiefel, E. (1952). Methods of conjugate gradients for solving linear systems. *Journal of research of the National Bureau of Standards*, *49*, 409–435. doi: 10.6028/jres.049.044
- Höskuldsson, A. (1988). PLS regression methods. *Journal of Chemometrics*, *2*(3), 211–228. doi: https://doi.org/10.1002/cem.1180020306
- <span id="page-10-15"></span>Kelley Pace, R., & Barry, R. (1997). Sparse spatial autoregressions. *Statistics & Probability Letters*, *33*(3), 291–297. doi: 10.1016/S0167- 7152(96)00140-X
- <span id="page-10-3"></span>Lobaugh, N. J., West, R., & McIntosh, A. R. (2001). Spatiotemporal analysis of experimental differences in event-related potential data with partial least squares. *Psychophysiology*, *38*(3), 517–530. doi: 10.1017/s0048577201991681
- Lyttkens, E. (1972). Regression aspects of canonical correlation. *Journal of Multivariate Analysis*, *2*(4), 418–439. doi: https://doi.org/10.1016/0047- 259X(72)90036-X
- <span id="page-10-16"></span>Mezzadri, F. (2007). How to generate random matrices from the classical compact groups. *Notices of the American Mathematical Society*, *54*(5), 592–604. doi: 10.48550/arXiv.math-ph/0609050
- Naes, T., & Martens, H. (1985). Comparison of prediction methods for multicollinear data. *Communications in Statistics Simulation and Computation*, *14*(3), 545–576. doi: 10.1080/03610918508812458
- <span id="page-10-2"></span>Nguyen, D., & Rocke, D. (2002). Tumor Classification by Partial Least Squares Using Microarray Gene Expression Data. *Bioinformatics (Oxford, England)*, *18*, 39–50. doi: 10.1093/bioinformatics/18.1.39
- <span id="page-10-4"></span>Nilsson, J., de Jong, S., & Smilde, A. K. (1997). Multiway calibration in 3D QSAR. *Journal of Chemometrics*, *11*(6), 511–524. doi: 10.1002/(SICI)1099- 128X(199711/12)11:6<511::AID-CEM488>3.0.CO;2-W
- <span id="page-10-9"></span><span id="page-10-0"></span>Nocedal, J., & Wright, S. J. (Eds.). (1999). *Numerical Optimization*. New York: Springer-Verlag. doi: 10.1007/b98874
- Noonan, R., & Wold, H. (1977). NIPALS Path Modelling with Latent Variables. *Scandinavian Journal of Educational Research SCAND J EDUC RES*, *21*, 33–61. doi: 10.1080/0031383770210103
- Phatak, A., & Hoog, F. (2002). Exploiting the connection between PLS, Lanczos methods and conjugate gradients: Alternative proofs of some properties of PLS. *Journal of Chemometrics*, *16*, 361–367. doi: 10.1002/cem.728
- <span id="page-10-11"></span>Rosipal, R., & Krämer, N. (2005). Overview and Recent Advances in Partial Least Squares. *Lecture Notes in Computer Science*, *3940*, 34–51. doi: 10.1007/11752790\_2
- <span id="page-10-14"></span>Sharma, R., & Bhandari, R. (2015). Skewness, kurtosis and Newton's inequality. *Rocky Mountain Journal of Mathematics*, *45*(5), 1639–1643. doi: 10.1216/RMJ-2015-45-5-1639
- <span id="page-10-12"></span>Takane, Y., & Loisel, S. (2016). On the PLS Algorithm for Multiple Regression (PLS1). In H. Abdi, V. Esposito Vinzi, G. Russolillo, G. Saporta, & L. Trinchera (Eds.), *The Multiple Facets of Partial Least Squares and Related Methods* (pp. 17–28). Cham: Springer International Publishing. doi: 10.1007/978-3-319-40643-5\_2
- <span id="page-10-10"></span>Wegelin, J. (2000). A Survey of Partial Least Squares (PLS) Methods, with Emphasis on the Two-Block Case. In *Technical Report.* Department of Statistics, University of Washington, Seattle.
- Wold, H. (1980). Model Construction and Evaluation When Theoretical Knowledge Is Scarce. In *Evaluation of Econometric Models* (pp. 47–74). Academic Press.

- <span id="page-11-2"></span>Wold, S., Ruhe, A., Wold, H., & Dunn, W. J., III. (1984). The Collinearity Problem in Linear Regression. The Partial Least Squares (PLS) Approach to Generalized Inverses. *SIAM Journal on Scientific and Statistical Computing*, *5*(3), 735–743. doi: 10.1137/0905052
- <span id="page-11-0"></span>Wold, S., Sjöström, M., & Eriksson, L. (2001). PLS-regression: A basic tool of chemometrics. *Chemometrics and Intelligent Laboratory Systems*, *58*(2), 109–130. doi: 10.1016/S0169-7439(01)00155-1
- <span id="page-11-1"></span>Worsley, K. J. (1997). An overview and some new developments in the statistical analysis of PET and fMRI data. *Human Brain Mapping*, *5*(4), 254–258. doi: 10.1002/(SICI)1097-0193(1997)5:4<254::AID-HBM9>3.0.CO;2-2

<span id="page-12-0"></span>

**Figure 1** Eigenvalue distributions in the synthetic regression problems

 $\overline{1}$ 

<span id="page-13-0"></span>

**Figure 2** PLS estimator distance analysis with different distributions of eigenvalues of the regressor covariance matrix

<span id="page-14-0"></span>

**Figure 3** Accuracy of the predictions of PCA and PLS regression measured in terms of the  $R^2$  score depending on whether the eigenvalues are concentrated or spread out uniformly.

<span id="page-14-1"></span>

**Figure 4** Accuracy of the predictions of PCA and PLS regression measured in terms of the  $R^2$  score depending on the number of clusters in which the eigenvalues are grouped

<span id="page-14-2"></span>

**Figure 5** Accuracy of the predictions of PCA and PLS regression measured in terms of the  $R^2$  score, depending on whether there is a cluster of eigenvalues near zero

<span id="page-15-0"></span>

**Figure 6** Eigenvalue distribution in the California Housing dataset

<span id="page-15-1"></span>

**Figure 7** PLS estimator distance analysis in the California Housing dataset

<span id="page-15-2"></span>

**Figure 8** Accuracy of the predictions of PCA and PLS regression measured in terms of the  $R^2$  score in the California Housing dataset