

Differential $e\mu bb$ Cross-Sections and New Higgses at the
Electroweak ScaleSUMIT BANIK * ¹, GUGLIELMO COLORETTI², ANDREAS CRIVELLIN³*Physik-Institut, Universität Zürich, Winterthurerstrasse 190, CH-8057 Zürich,
Switzerland**Paul Scherrer Institut, CH-5232 Villigen PSI, Switzerland*BRUCE MELLADO⁴*School of Physics and Institute for Collider Particle Physics, University of the
Witwatersrand, Johannesburg, Wits 2050, South Africa**iThemba LABS, National Research Foundation, PO Box 722, Somerset West 7129,
South Africa*

ATLAS found that none of their Standard Model simulations can describe the measured differential lepton distributions in their $t\bar{t}$ analysis reasonably well. Therefore, we study the possibility that this measurement has a new physics contamination. We consider a benchmark model motivated by the indications for di-photon resonances: A heavy scalar decays into two lighter Higgs bosons with masses of 152 GeV and 95 GeV, with subsequent decay to WW and bb , respectively. In this setup, the description of data is improved by at least 5.6σ .

PRESENTED AT

16th International Workshop on Top Quark Physics
(Top2023), 24–29 September, 2023

¹sumit.banik@physik.uzh.ch (*speaker)²guglielmo.coloretti@physik.uzh.ch³andreas.crivellin@cern.ch⁴bmellado@mail.cern.ch

1 Introduction

With the discovery of the Brout-Englert-Higgs boson [1–4] at the Large Hadron Collider (LHC) in 2012 [5, 6], the Standard Model (SM) has been completed. However, the minimality of the SM scalar sector is not guaranteed by any symmetry principle and a plethora of extensions, including the addition of $SU(2)_L$ singlets [7–9], doublets [10–14] and triplets [15–20] have been proposed.

In fact, there are indications of new Higgses at the electroweak (EW) scale with masses of ≈ 95 GeV [21–25] and ≈ 152 GeV [26] with global significances of 3.8σ and 4.9σ [27, 28], respectively. Furthermore, the “multi-lepton anomalies” [29–32], processes involving multiple leptons and missing energy, with and without (b-)jets, where significant deviations from the SM expectations have been observed over the last years, including WW , WWW , Wh , tW , $t\bar{t}$, $t\bar{t}W$ and $t\bar{t}t\bar{t}$ signatures (see Ref. [32] and references therein) are compatible with, or even suggest, a new scalar with a mass of ≈ 150 GeV [33, 34].

In these proceedings, we will summarize the statistically most significant multi-lepton excess encoded in the latest ATLAS analysis [35] of the $t\bar{t}$ differential cross-sections as analyzed in Ref. [36]. We focus on the invariant di-lepton mass ($m^{e\mu}$) and the angle between the leptons ($\Delta\phi^{e\mu}$) for various combinations of Monte Carlo (MC) simulators.* For this, we consider a simplified model in which a new scalar H is produced via gluon fusion and decays to S and S' , with masses of 152 GeV and 95 GeV, respectively, as suggested by the hints for narrow resonances.

2 Analysis and Results

ATLAS [35] considered six different SM simulations, as given in Table 2, for the differential $t\bar{t}$ cross-section, including the invariant mass of the electron-muon pair ($m^{e\mu}$) and the angle between the leptons ($|\Delta\phi^{e\mu}|$). From the dashed lines in Fig. 1 one can see the poor agreement of the SM predictions with data.

In our simplified model, where the masses are fixed by the di-photon resonances to $m_S = 152$ GeV and $m_{S'} = 95$ GeV, we assume that the dominant decay of S is into pairs of W boson, while S' decays dominantly to $b\bar{b}$. Note that this is naturally the case if S is a SM-like Higgs and S' is the neutral component of an $SU(2)_L$ triplet with hypercharge 0.

We simulated in this setup the process $pp \rightarrow H \rightarrow SS' \rightarrow (WW^{(*)} \rightarrow \ell^+\nu\ell^-\nu)b\bar{b}$, with $\ell = e, \mu, \tau$, using **MadGraph5aMC@NLO** [38], **Pythia8.3** [39], and **Delphes** [40]. The results are shown as solid lines in Fig. 1. One can see that for all SM simulations used by ATLAS, the agreement with data is significantly improved. In fact, as given

*Note that the CMS analysis of $t + W$ [37] is less precise but consistent with the ATLAS findings [29].




Figure 1: The dashed coloured lines show six different SM predictions (MC) normalized to data given in Ref. [35]. The solid lines include the NP contribution from our benchmark model obtained by a combined global fit to $m^{e\mu}$ and $\Delta\phi^{e\mu}$ data including its correlations. The black lines are obtained by averaging the six predictions and the grey band shows the total uncertainty (systematic and statistical). One can see that the agreement between theory and experiment is significantly increased over the whole range of the distributions by adding an NP effect.

in Table. 2, the χ^2 is reduced between $\Delta\chi^2 = 34$ and $\Delta\chi^2 = 158$, corresponding to a preference of the NP model over the SM hypothesis by 5.8σ – 13.0σ while the average of the SM predictions results in 9.6σ .

So far, we fixed the masses of S and S' by using the hints for narrow di-photon resonances and took $m_H = 270$ GeV as a benchmark point which avoids constraints from SUSY and non-resonant di-Higgs searches. Let us now discuss the dependence on the masses. First of all, we checked that the effect of changing either the mass of H or of S' is very small. This means that one can pick any value for m_H so that this does not need to be counted as a degree of freedom in the statistical analysis. Now we can find the best-fit value in the m_S - ϵ_{NP} plane by averaging the x_i values of the six different MC simulations. For this, we generated 500k events each for m_S between 140 GeV and 160 GeV in 2 GeV steps and interpolated. As one can see from Fig. 3 the result is perfectly consistent with a mass of 152 GeV as suggested by the invariant mass of the di-photon system [26, 27].

Monte Carlo	χ_{SM}^2	χ_{NP}^2	σ_{NP}	Sig.	m_S [GeV]
Powheg+Pythia8	213	102	9pb	10.5 σ	143–156
aMC@NLO+Herwig7.1.3	102	68	5pb	5.8 σ	--
aMC@NLO+Pythia8	291	163	10pb	11.3 σ	148-157
Powheg+Herwig7.1.3	261	126	10pb	11.6 σ	149-156
Powheg+Pythia8 (rew)	69	35	5pb	5.8 σ	--
Powheg+Herwig7.0.4	294	126	12pb	13.0 σ	149-156
Average	182	88	9pb	9.6 σ	143-157

Figure 2: χ^2 values, preferred cross-section (σ_{NP}) significance (Sig.) etc. for the combined fit to $m^{e\mu}$ and $|\Delta\phi^{e\mu}|$ distributions for the six different SM simulations and their average. The χ_{NP}^2 is for our benchmark scenario with $m_S \approx 152$ GeV while the m_S gives the preferred range of it from the fit.

Finally, let us consider the consistency of the preferred signal strength for $pp \rightarrow H \rightarrow SS'$ with the di-photon excess at 95 GeV. The branching ratio to photons of an SM-like Higgs with 95 GeV is $\approx 1.4 \times 10^{-3}$, $\text{Br}[S' \rightarrow b\bar{b}] \approx 0.86$ [41] and the production cross section at 13 TeV via gluon fusion is ≈ 68 pb [41]. Given that the preferred $\gamma\gamma$ signal at 95 GeV, with respect to a (hypothetical) SM-like Higgs with a mass of 95 GeV, is $\mu_{\gamma\gamma} = 0.24_{-0.08}^{+0.09}$ [42] we can see in Fig. 3 that, under the assumption that S' is SM-like and $\text{Br}[S \rightarrow WW^*] \approx 100\%$, the associated production of S' , determined by the preferred signal strength from $t\bar{t}$, perfectly explains the excess.

3 Conclusions

ATLAS observed that the measured differential lepton distributions in its analysis of $t\bar{t}$ significantly differ from the predictions of the SM, which were obtained through various combinations of Monte Carlo simulators: “No model (simulation) can describe all measured distributions within their uncertainties.” This suggests that the measurement may be influenced by new physics contributions.

We suggest that the $pp \rightarrow H \rightarrow SS' \rightarrow WWb\bar{b}$ could constitute an NP background for the measurement. In fact, motivated by hints of di-photon resonances at around 95 GeV and 152 GeV, we have examined a benchmark scenario where S' and S have corresponding masses and decay to $b\bar{b}$ and WW , respectively, with $m_H = 270$ GeV. This NP hypothesis is preferred over the SM hypothesis by at least 5.8σ .

Furthermore, by varying m_S we found that the preferred range is compatible with $m_S \approx 152$ GeV, as motivated by the $\gamma\gamma$ and WW excesses. Since the latter analysis employs a jet veto, this further disfavours the possibility that higher-order QCD corrections are the origin of the tension in the $t\bar{t}$ differential distributions. Assuming




Figure 3: Preferred regions from the $t\bar{t}$ differential distributions (red) as a function of m_S and the total cross section $pp \rightarrow H \rightarrow SS' \rightarrow WWbb$ assuming S' to be SM-like and $\text{Br}[S \rightarrow WW] = 100\%$. The blue region is preferred by the 95 GeV $\gamma\gamma$ signal strength.

that S' is SM-like, this results in a di-photon signal strength in agreement with the $\gamma\gamma$ excess at 95 GeV. Since for S a dominant decay to W bosons is needed while the ZZ rate should be low to respect the limits from inclusive 4ℓ searches, this suggests that it could be the neutral component of the $SU(2)_L$ triplet with hypercharge 0.

ACKNOWLEDGEMENTS

The work of A.C., S.B. and G.C. is supported by a professorship grant from the Swiss National Science Foundation (No. PP00P2_211002). B.M. gratefully acknowledges the South African Department of Science and Innovation through the SA-CERN program, the National Research Foundation, and the Research Office of the University of the Witwatersrand for various forms of support.

References

- [1] Peter W. Higgs. Broken symmetries, massless particles and gauge fields. *Phys. Lett.*, 12:132–133, 1964.
- [2] F. Englert and R. Brout. Broken Symmetry and the Mass of Gauge Vector Mesons. *Phys. Rev. Lett.*, 13:321–323, 1964.
- [3] Peter W. Higgs. Broken Symmetries and the Masses of Gauge Bosons. *Phys. Rev. Lett.*, 13:508–509, 1964.
- [4] G. S. Guralnik, C. R. Hagen, and T. W. B. Kibble. Global Conservation Laws and Massless Particles. *Phys. Rev. Lett.*, 13:585–587, 1964.
- [5] Georges Aad et al. Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC. *Phys. Lett. B*, 716:1–29, 2012.
- [6] Serguei Chatrchyan et al. Observation of a New Boson at a Mass of 125 GeV with the CMS Experiment at the LHC. *Phys. Lett. B*, 716:30–61, 2012.
- [7] Vanda Silveira and A. Zee. SCALAR PHANTOMS. *Phys. Lett. B*, 161:136–140, 1985.
- [8] Massimo Pietroni. The Electroweak phase transition in a nonminimal supersymmetric model. *Nucl. Phys. B*, 402:27–45, 1993.
- [9] John McDonald. Gauge singlet scalars as cold dark matter. *Phys. Rev. D*, 50:3637–3649, 1994.
- [10] T. D. Lee. A Theory of Spontaneous T Violation. *Phys. Rev. D*, 8:1226–1239, 1973.
- [11] Howard E. Haber and Gordon L. Kane. The Search for Supersymmetry: Probing Physics Beyond the Standard Model. *Phys. Rept.*, 117:75–263, 1985.
- [12] Jihn E. Kim. Light Pseudoscalars, Particle Physics and Cosmology. *Phys. Rept.*, 150:1–177, 1987.
- [13] R. D. Peccei and Helen R. Quinn. CP Conservation in the Presence of Instantons. *Phys. Rev. Lett.*, 38:1440–1443, 1977.
- [14] Neil Turok and John Zadrozny. Electroweak baryogenesis in the two doublet model. *Nucl. Phys. B*, 358:471–493, 1991.

- [15] W. Konetschny and W. Kummer. Nonconservation of Total Lepton Number with Scalar Bosons. *Phys. Lett. B*, 70:433–435, 1977.
- [16] T. P. Cheng and Ling-Fong Li. Neutrino Masses, Mixings and Oscillations in SU(2) x U(1) Models of Electroweak Interactions. *Phys. Rev. D*, 22:2860, 1980.
- [17] George Lazarides, Q. Shafi, and C. Wetterich. Proton Lifetime and Fermion Masses in an SO(10) Model. *Nucl. Phys. B*, 181:287–300, 1981.
- [18] J. Schechter and J. W. F. Valle. Neutrino Masses in SU(2) x U(1) Theories. *Phys. Rev. D*, 22:2227, 1980.
- [19] M. Magg and C. Wetterich. Neutrino Mass Problem and Gauge Hierarchy. *Phys. Lett. B*, 94:61–64, 1980.
- [20] Rabindra N. Mohapatra and Goran Senjanovic. Neutrino Masses and Mixings in Gauge Models with Spontaneous Parity Violation. *Phys. Rev. D*, 23:165, 1981.
- [21] R. Barate et al. Search for the standard model Higgs boson at LEP. *Phys. Lett. B*, 565:61–75, 2003.
- [22] Albert M Sirunyan et al. Search for a standard model-like Higgs boson in the mass range between 70 and 110 GeV in the diphoton final state in proton-proton collisions at $\sqrt{s} = 8$ and 13 TeV. *Phys. Lett. B*, 793:320–347, 2019.
- [23] Searches for additional Higgs bosons and vector leptoquarks in $\tau\tau$ final states in proton-proton collisions at $\sqrt{s} = 13$ TeV. 2022.
- [24] Search for a new resonance decaying to two scalars in the final state with two bottom quarks and two photons in proton-proton collisions at $\sqrt{s} = 13$ TeV. 2022.
- [25] Search for diphoton resonances in the 66 to 110 GeV mass range using 140 fb⁻¹ of 13 TeV *pp* collisions collected with the ATLAS detector. 2023.
- [26] Georges Aad et al. Search for dark matter in events with missing transverse momentum and a Higgs boson decaying into two photons in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. *JHEP*, 10:013, 2021.
- [27] Andreas Crivellin, Yaquan Fang, Oliver Fischer, Abhaya Kumar, Mukesh Kumar, Elias Malwa, Bruce Mellado, Ntsoko Rapheeha, Xifeng Ruan, and Qiyu Sha. Accumulating Evidence for the Associate Production of a Neutral Scalar with Mass around 151 GeV. 9 2021.

- [28] Srimoy Bhattacharya, Guglielmo Coloretti, Andreas Crivellin, Salah-Eddine Dahbi, Yaquan Fang, Mukesh Kumar, and Bruce Mellado. Growing Excesses of New Scalars at the Electroweak Scale. 6 2023.
- [29] Stefan Buddenbrock, Alan S. Cornell, Yaquan Fang, Abdualazem Fadol Mohammed, Mukesh Kumar, Bruce Mellado, and Kehinde G. Tomiwa. The emergence of multi-lepton anomalies at the LHC and their compatibility with new physics at the EW scale. *JHEP*, 10:157, 2019.
- [30] Stefan von Buddenbrock, Richard Ruiz, and Bruce Mellado. Anatomy of inclusive $t\bar{t}W$ production at hadron colliders. *Phys. Lett. B*, 811:135964, 2020.
- [31] Yesenia Hernandez, Mukesh Kumar, Alan S. Cornell, Salah-Eddine Dahbi, Yaquan Fang, Benjamin Lieberman, Bruce Mellado, Kgomotso Monnagotla, Xifeng Ruan, and Shuiting Xin. The anomalous production of multi-lepton and its impact on the measurement of Wh production at the LHC. *Eur. Phys. J. C*, 81(4):365, 2021.
- [32] Oliver Fischer et al. Unveiling hidden physics at the LHC. *Eur. Phys. J. C*, 82(8):665, 2022.
- [33] Stefan von Buddenbrock, Alan S. Cornell, Abdualazem Fadol, Mukesh Kumar, Bruce Mellado, and Xifeng Ruan. Multi-lepton signatures of additional scalar bosons beyond the Standard Model at the LHC. *J. Phys. G*, 45(11):115003, 2018.
- [34] Guglielmo Coloretti, Andreas Crivellin, Srimoy Bhattacharya, and Bruce Mellado. Searching for low-mass resonances decaying into W bosons. *Phys. Rev. D*, 108(3):035026, 2023.
- [35] Georges Aad et al. Inclusive and differential cross-sections for dilepton $t\bar{t}$ production measured in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector. *JHEP*, 07:141, 2023.
- [36] Sumit Banik, Guglielmo Coloretti, Andreas Crivellin, and Bruce Mellado. Uncovering New Higgses in the LHC Analyses of Differential $t\bar{t}$ Cross Sections. 8 2023.
- [37] Albert M Sirunyan et al. Measurement of the production cross section for single top quarks in association with W bosons in proton-proton collisions at $\sqrt{s} = 13$ TeV. *JHEP*, 10:117, 2018.

- [38] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H. S. Shao, T. Stelzer, P. Torrielli, and M. Zaro. The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations. *JHEP*, 07:079, 2014.
- [39] Torbjörn Sjöstrand, Stefan Ask, Jesper R. Christiansen, Richard Corke, Nishita Desai, Philip Ilten, Stephen Mrenna, Stefan Prestel, Christine O. Rasmussen, and Peter Z. Skands. An introduction to PYTHIA 8.2. *Comput. Phys. Commun.*, 191:159–177, 2015.
- [40] J. de Favereau, C. Delaere, P. Demin, A. Giammanco, V. Lemaître, A. Mertens, and M. Selvaggi. DELPHES 3, A modular framework for fast simulation of a generic collider experiment. *JHEP*, 02:057, 2014.
- [41] D. de Florian et al. Handbook of LHC Higgs Cross Sections: 4. Deciphering the Nature of the Higgs Sector. 2/2017, 10 2016.
- [42] T. Biekötter, S. Heinemeyer, and G. Weiglein. The 95.4 GeV di-photon excess at ATLAS and CMS. 6 2023.