Z_3 symmetry of the CKM and PMNS matrices

Piotr Kielanowski^a, S. Rebeca Juárez Wysozka^b, Liliana Vázquez Mercado^c

^aDepartamento de Física, Centro de Investigación y Estudios Avanzados, Av. Instituto Politécnico Nacional 2508, Ciudad de México, 07000, Mexico ^bDepartamento de Física, Escuela Superior de Física y Matemáticas, Instituto Politécnico Nacional. U.P Adolfo López Mateos, Ciudad de México, 07739, Mexico ^cDepartamento de Física, Centro Universitario de Ciencias Exactas e Ingenierías, Universidad de Guadalajara, Av. Revolución 1500, Colonia Olímpica, Guadalajara, 44430, Jalisco, Mexico

Abstract

We develop for the CKM and PMNS matrices a new representation with special properties. It is obtained by splitting each of these matrices into two rotations by the angle $\sim 2\pi/3$ and a universal diagonal matrix with elements, which are cubic roots of 1. Such a representation of the CKM and PMNS matrices may indicate the Z_3 symmetry to be present in the Yukawa sector of the SM. Identical mathematical structure of the CKM and PMNS matrices is also an extension of the quark-lepton universality. In this approach the CP violation is a natural consequence of the structure of the Yukawa couplings. The CP violating phase is not a fitted parameter and its value is governed by the parameters of two rotations. The parameters of the diagonalizing matrices of the bi-unitary transformation do not exhibit a hierarchy, which means that the origins of the hierarchy of quark masses and of the CKM

Keywords: CKM matrix, PMNS matrix, CP violation, Z_3 symmetry, Yukawa couplings

Abstract
We develop for the CKM and PMN
matrices into two rotations by the
a representation of the CKM and I
Identical mathematical structure o
approach the CP violation is a nat
fitted parameter and its value is go
bi-unitary transformation do not ex
matrix elements are not the same. *Keywords:* CKM matrix, PMNS n
In classical physics the invaria
the discrete symmetries *P*, *T* and *C*considered certain that it should a
physics at any level. This belief
experimental confirmation [1] of t
violation [2] in weak interactions
violation of the charge conjugatio
in the experiment measuring the
decays of charged pions [3]. Nex
bined CP invariance was experime
neutral *K* meson decays [4]. Th
sal invariance *T* was observed for
system [5].
The discussion of the violatic
violation as new experimental results v
oretical attempt of the descriptio
by Fermi [6, 7] with only vector In classical physics the invariance of all phenomena under the discrete symmetries P, T and C was beyond doubt, so it was considered certain that it should also be extended to quantum physics at any level. This belief was unexpectedly broken by experimental confirmation [1] of the prediction of space parity violation [2] in weak interactions. Almost simultaneously the violation of the charge conjugation invariance C was observed in the experiment measuring the polarization of muons in the decays of charged pions [3]. Next the hypothesis of the combined CP invariance was experimentally proven to be broken in neutral K meson decays [4]. The violation of the time reversal invariance T was observed for the first time in the neutral K

The discussion of the violation of discrete symmetries invariance was gradually evolving in the theory of weak interaction as new experimental results were emerging. The first theoretical attempt of the description of β -decay was formulated by Fermi [6, 7] with only vector charged currents interactions and it did not contain the breaking of discrete symmetries. The parity violation through V-A weak current was introduced into the Fermi's theoretical description of weak interactions by Sudarshan and Marshak in Refs. [8, 9] and Feynman and Gell-Mann in Ref. [10]. The next step in the theoretical development was the reconciliation of universality of weak interactions of weak decays of strange particles with β decays of muons by expressing the weak hadronic current as a mixture described by

one angle θ of strangeness conserving current and strangeness changing current [11]. CP violation can only be consistently described in the theory with three generations of quarks and it was done in the framework of the Standard Model (SM) by Kobayashi and Maskawa by introducing the 3×3 unitary mixing matrix for quark interactions with charged vector bosons [12]. We thus see that description of charged current weak interactions by the Cabibbo-Kobayashi-Maskawa (CKM) matrix is the result of the 40 years study of the properties of elementary particles, with emphasis on the lepton-hadron universality.

For massive neutrinos in the lepton sector the mixing matrix, called the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) matrix [13-17], plays an identical role and has similar mathematical properties as the CKM matrix for quarks. The only difference is the number of parameters that appear for the Majorana neutrinos.

The least understood part of the SM is the Yukawa sector. In the classical SM (with masless neutrinos) it contains 13 phenomenological parameters: 9 quark and lepton masses and 4 parameters of the CKM matrix. Better understanding of the Yukawa sector may lead to a significant reduction of the number of parameters of the SM.

2. General remarks about the CKM and PMNS matrices

The three most important results of our analysis are: 1. The CP violation appears naturally and the CP violating phase is not a free parameter. 2. The CKM and PMNS matrices have a common algebraic structure and this extends the quark lepton universality. 3. The experimental data for the CKM and PMNS matrices suggest the Z_3 symmetry to be present in the Yukawa sector of the Standard Model.

Email addresses: piotr.kielanowski@cinvestav.mx (Piotr Kielanowski), rebecajw@gmail.com (S. Rebeca Juárez Wysozka), liliana.vmercado@academicos.udg.mx (Liliana Vázquez Mercado)

The CKM matrix V, when derived from the Yukawa couplings of quarks with the Higgs field is equal to

$$V = V_L^u V_L^{d^{\dagger}},\tag{1}$$

where V_L^u and V_L^d are the unitary matrices that are determined by the biunitary diagonalization of the Yukawa interactions matrices of the *up* and *down* quarks.

It is not simple to obtain properties of the Yukawa couplings from the experimental values of the CKM matrix [18], because there is an excessive arbitrariness in the splitting of the matrix V in Eq. (1) into two unitary matrices. The conventional way to derive those properties is to assume a discrete flavor symmetry or some other properties, like textures for the Yukawa couplings and then compare the predictions with experimental values for the CKM or PMNS matrix elements. For a recent illustration of the first method see Ref. [19] and for the recent paper on textures see [20].

In phenomenological applications there are two mostly used parameterizations of the CKM matrix: the standard one [21], which is a superposition of three rotations around the x, y and z axes with a complex rotation around the y axis and a second one is the Wolfenstein parameterization [22], which highlights the hierarchy between the CKM matrix elements.

3. New representation of the CKM and PMNS matrices

In this paper we will use a special representation of a general unitary matrix as a product of three matrices

$$U = O_1 D O_2^T. (2)$$

Here O_1 and O_2 are orthogonal (rotation) matrices ¹ and *D* is the diagonal matrix whose elements are imaginary exponents. A sketch of a proof that a general unitary matrix can be expressed in a form (2) is given in Ref. [23].

Now using the representation in Eq. (2) we will analyze the CKM and PMNS (Dirac case) matrices ². Our aim is not to find a new parameterization of those matrices, but to show that the matrices O_1 , O_2 and D have a special form, which can be identified with the symmetry present in the Yukawa interactions of quarks and leptons. For this purpose we will determine the matrices O_1 , O_2 and D from the experimental input for the CKM and PMNS matrices using the method described below.

The CKM and PMNS matrices have dimension 3×3 and are unitary. The rephasing freedom of the quark and lepton fields results in the reduction to 4 of the number of significant parameters of each matrix. A general 3×3 unitary matrix has 9 free parameters. This means that we have some freedom in the choice of the values of the parameters in representation (2) for the reproduction of the experimental values of the CKM and PMNS matrices. In order to uncover the Z_3 symmetry of the Yukawa sector we will make the following assumptions about the structure of the matrices O_1 , O_2 and D in Eq. (2) 1. The diagonal matrix *D* for *both* CKM and PMNS matrices comprises of the third order roots of 1 (*Z*₃ symmetry) and is equal

$$D_{Z_3} = \text{diag}(e^{\frac{4\pi i}{3}}, e^{\frac{2\pi i}{3}}, 1).$$
(3)

matrix The choice of such a form is also motivated by the successful two-angle parameterization of the CKM matrix [24], which uses a similar form of this matrix. This ansatz for the matrix D reduces the number of free parameters for each matrix by 3.

2. We choose the angles α of the rotation of the real orthogonal matrices O_1 and O_2 to be *fixed* and equal to each other, so these matrices differ only by the axis of rotation and thus we remain with 4 free parameters for each matrix CKM and PMNS.

These assumptions lead the following representation of the CKM and PMNS matrices

$$V = \mathcal{R}(\alpha; \theta_u, \phi_u) D_{Z_3} \mathcal{R}(\alpha; \theta_d, \phi_d)^T.$$
(4)

Here $\mathcal{R}(\alpha; \theta, \phi)$ is the matrix of rotation, α is the angle of rotation and θ and ϕ are the spherical angles of the axis of rotation.

4. Parameters of the CKM and PMNS matrices from experiment

Our goal is to investigate the compatibility of the representation (4) with the measured values of the elements of the CKM and PMNS matrices, by determination of the parameters of the CKM matrix in representation (4) from the absolute values of the matrix elements of the CKM and PMNS matrices generated by the fit of these matrices to the experimental values. In such a way the parameters of the unitary matrix in Eq. (4) will be determined exactly. As our input we use the central values of the parameters of each matrix given by PDG [18], which are

for the CKM matrix

$$\lambda = 0.22500, A = 0.826, \bar{\rho} = 0.159, \bar{\eta} = 0.348,$$

for the PMNS matrix (normal order)
 $\sin^2(\theta_{12}) = 0.307, \sin^2(\theta_{23}) = 0.547,$
 $\sin^2(\theta_{13}) = 0.0220, \ \delta_{CP} = 1.23\pi \text{ radians}$
for the PMNS matrix (inverted order)
 $\sin^2(\theta_{12}) = 0.307, \ \sin^2(\theta_{23}) = 0.534,$
 $\sin^2(\theta_{13}) = 0.0220, \ \delta_{CP} = 1.23\pi \text{ radians}$
(5)

and calculate the absolute values of the elements of the CKM and PMNS matrices. Next we determine *exactly* the spherical angles θ and ϕ for both matrices by solving the set of corresponding equations for these absolute values.

As the next step in the search of the Z_3 symmetry we test the assumption that the angle of rotation α is equal *exactly* to $2*\pi/3$. Such an assumption is *not compatible* with the experimental information for the CKM and PMNS matrices. Subsequently we tested a small deviation of angle α from $2\pi/3$ for both matrices. The positive result of our search is shown in Table 1 where

¹For the 3-dimensional matrices we parameterize them by three angles, which are: two angles θ and ϕ which define the direction of the axis of rotation and the rotation angle α around the given axis.

²The Majorana case will be discussed elsewhere.

we give the values of the angle α and spherical angles θ and ϕ which *exactly* reproduce the CKM and PMNS matrices generated from the data given in Eq. (5) and we see that the deviation

| Matrix | α | θ_u | ϕ_u | $	heta_d$ | ϕ_d |
|----------|-------------------------|------------|----------|-----------|----------|
| CKM | $\frac{2\pi}{3} * 1.04$ | 52.92° | 46.35° | 52.70° | 50.25° |
| PMNS (A) | $\frac{2\pi}{3} * 1.01$ | 64.24° | 138.66° | 156.96° | 33.02° |
| PMNS (B) | $\frac{2\pi}{3} * 1.01$ | 64.18° | 138.98° | 156.55° | 32.52° |

Table 1: The angles α , θ and ϕ of the CKM and PMNS matrices. PMNS (A) stands for normal order and PMNS (B) stands for the inverted order. Angle α is in radians and the remaining angles are in degrees.

from the value $2\pi/3$ is small: for the CKM matrix it is 4% and for the PMNS matrix it is only 1%. The striking fact is that none of the angles in Table 1 are small, which suggests a non perturbative nature of the CKM and PMNS matrices.

5. Discussion of the results

The data in Table 1 show that the spherical angles of the axes of rotation in the CKM matrix for the *up* and *down* quarks are almost equal and this is a consequence of the fact that the matrix of the absolute values CKM matrix is *almost* symmetric. The spherical angles of the axes of rotation of the CKM and PMNS matrices are different, but this was to be expected, because these matrices are not equal.

The structure of the matrix D_{Z_3} and the values of the angle α for the CKM and PMNS matrices give a strong support for the idea of the presence of the Z_3 symmetry for the matrices CKM and PMNS and thus also for the Yukawa couplings of quarks and leptons. A deviation of the angle α from the value $2 * \pi/3$ may be caused by a small Z_3 symmetry breaking.

Our representation of the CKM and PMNS matrices allows the determination of the *left* bi-unitary diagonalizing matrices V_L^u and V_L^d

$$V_L^f = \mathcal{R}(\alpha, \theta_f, \phi_f) D_f, \quad f = u, d, \tag{6}$$

where D_u and D_d are the diagonal matrices

$$D_u = \text{diag}(1, e^{2\pi i/3}, e^{4\pi i/3}), \quad D_d = \text{diag}(e^{2\pi i/3}, 1, e^{4\pi i/3})$$
 (7)

and

$$D_{Z_3} = D_{\mu} D_{\mu}^{\dagger}. \tag{8}$$

It follows from the biunitary diagonalization that the quark Yukawa couplings matrices Y^u and Y^d are equal

$$\frac{\nu}{\sqrt{2}}Y^{f} = V_{L}^{f^{\dagger}}M^{f}V_{R}^{f}$$
$$= D_{f}^{\dagger}\mathcal{R}(\alpha,\theta_{f},\phi_{f})^{T}M^{f}V_{R}^{f}, \quad f = u, d. \quad (9)$$

Here M^f is the diagonal quark or lepton mass matrix and v is the vacuum expectation value of the Higgs field. The *right* biunitary diagonalizing matrices do not enter into any observables and we will consider two cases: $V_R^f = I$ or $V_R^f = V_L^f$. These choices lead to different results for the matrices of the quark and lepton Yukawa couplings:

$$\frac{v}{\sqrt{2}}Y^{f} = \begin{cases} \text{case A} & D_{f}^{\dagger}\mathcal{R}(\alpha,\theta_{f},\phi_{f})^{T}M^{f} \text{ for } V_{R}^{u} = I\\ \text{case B} & D_{f}^{\dagger}\mathcal{R}(\alpha,\theta_{f},\phi_{f})^{T}M^{f}V_{L}^{f}D_{f} \text{ for } V_{R}^{f} = V_{L}^{f} \end{cases}$$
(10)

Below we give the explicit numerical values of the V_l^J and Y^f matrices for quarks obtained from the values of the angles in Table 1. For clarity we did not substitute the phases $e^{2\pi i/3}$ and $e^{4\pi i/3}$ by its numerical values, because in such a way the mathematical structure of these matrices becomes clear.

$$V_L^u = \begin{pmatrix} -0.094 & 0.0043 \cdot e^{2\pi i/3} & 1.00 \cdot e^{4\pi i/3} \\ 0.99 & -0.047 \cdot e^{2\pi i/3} & 0.094 \cdot e^{4\pi i/3} \\ 0.048 & 1.00 \cdot e^{2\pi i/3} & 0.00021 \cdot e^{4\pi i/3} \end{pmatrix}$$
(11)

$$V_L^d = \begin{pmatrix} -0.16 \cdot e^{2\pi i/3} & -0.0088 & 0.99 \cdot e^{4\pi i/3} \\ 0.99 \cdot e^{2\pi i/3} & 0.017 & 0.16 \cdot e^{4\pi i/3} \\ -0.018 \cdot e^{2\pi i/3} & 1.00 & 0.0069 \cdot e^{4\pi i/3} \end{pmatrix}$$
(12)

These matrices have the following properties

- The numerical structure of the matrices V_L^u and V_L^d is similar.
- The phases of the matrix elements in each column are equal.
- There is a hierarchy between the matrix elements of the matrices: the matrix elements: in each row (or column) one matrix element is of the order ~ 1 and the remaining matrix elements are small.
- The hierarchy of the matrix elements is *independent* of the masses of quarks.
- There are no exact textures.

Next, we calculate from Eq. (9) the matrices of the Yukawa couplings Y^f for two cases explained in Eq. (10). In the equations below we provide the numerical values of the matrices $vY^{u,d}/(m_{t,b}\sqrt{2})$ for the *up* and *down* quarks. For Y^u and Y^d we normalize the values of these matrices by the masses m_t and m_b , respectively.

case A:
$$\frac{v}{m_t \sqrt{2}} Y^u$$

= $\begin{pmatrix} -1.18 \cdot 10^{-6} & 0.0073 & 0.048 \\ 5.35 \cdot 10^{-8} \cdot e^{4\pi i/3} & -0.00035 \cdot e^{4\pi i/3} & 1.00 \cdot e^{4\pi i/3} \\ 0.000012 \cdot e^{2\pi i/3} & 0.00069 \cdot e^{2\pi i/3} & 0.00021 \cdot e^{2\pi i/3} \end{pmatrix}$ (13a)

case B:
$$\frac{v}{m_t \sqrt{2}} Y^u$$

= $\begin{pmatrix} 0.0095 & 0.047 \cdot e^{2\pi i/3} & 0.00070 \cdot e^{4\pi i/3} \\ 0.047 \cdot e^{4\pi i/3} & 1.00 & 0.00018 \cdot e^{2\pi i/3} \\ 0.00070 \cdot e^{2\pi i/3} & 0.00018 \cdot e^{4\pi i/3} & 0.000078 \end{pmatrix}$ (13b)

case A:
$$\frac{v}{m_b \sqrt{2}} Y^d$$

= $\begin{pmatrix} -0.00018 \cdot e^{4\pi i/3} & 0.022 \cdot e^{4\pi i/3} & -0.018 \cdot e^{4\pi i/3} \\ -9.87 \cdot 10^{-6} & 0.00038 & 1.00 \\ 0.0011 \cdot e^{2\pi i/3} & 0.0037 \cdot e^{2\pi i/3} & 0.0060 \cdot e^{2\pi i/3} \end{pmatrix}$ (14a)

case B:
$$\frac{v}{m_b \sqrt{2}} Y^d$$

= $\begin{pmatrix} 0.022 & -0.018 \cdot e^{4\pi i/3} & 0.0033 \cdot e^{2\pi i/3} \\ -0.018 \cdot e^{2\pi i/3} & 1.00 & 0.0060 \cdot e^{4\pi i/3} \\ 0.0033 \cdot e^{4\pi i/3} & 0.0060 \cdot e^{2\pi i/3} & 0.0017 \end{pmatrix}$ (14b)

We observe that the Yukawa couplings Y^f have the following properties

- For case A the phases of the matrix elements in each row are equal and for case B the matrices are hermitian.
- There are no *exact* textures, because all matrix elements of $\mathcal{R}(\alpha, \theta_f, \phi_f)$ are non-vanishing.
- The absolute value of only one matrix element of each matrix is close to 1, the remaining ones are several order of magnitude smaller. For case A, the matrix element (2, 3) dominates, and for case B, the element (2, 2) dominates.
- The hierarchy in the *Y*^f matrices is stronger than in case of the *V*^f_L matrices which is caused by the strong hierarchy of the quark masses.

6. Conclusions

Let us summarize the obtained results by stating that we formulated a new representation of the CKM and PMNS matrices with the following properties

1. *Both* matrices CKM and PMNS have the following identical mathematical structure

$$V_{\text{CKM,PMNS}} = \mathcal{R}(\alpha, \theta_u, \phi_u) D_{Z_3} \mathcal{R}(\alpha, \theta_d, \phi_d)^T$$

with D_{Z_3} fixed, given in Eq. (3) and the angle α is close to $2\pi/3$.

- 2. The structure of the diagonal matrix D_{Z_3} , whose elements are cubic roots of 1 and the value of the angle α close to $2\pi/3$ support the idea of the presence of the Z_3 symmetry in the Yukawa sector of the Standard Model.
- 3. The CP violation phase in the CKM matrix *is not* a fitted parameter, but is the consequence of the structure of the phases of the Yukawa couplings matrices (see Eq. 9). This may be a starting point of a future theory of CP violation.

- 4. The angles describing the diagonalizing matrices V_L^u and V_L^d are not small (see Table 1). This means that the hierarchy present in the CKM matrix seems to be accidental and has a different origin than the hierarchy of the quark and lepton masses. This point of view is supported by the lack of the hierarchy in the PMNS matrix.
- 5. The Yukawa couplings do not have *exact* textures.

Points 1., 2. and 3. provide a description of CP violation without free parameters in the SM and reestablish the quark-lepton universality, which seemed missing in the Yukawa sector. Points 4. and 5. may be a good testing ground in search of the quark flavor symmetry.

Acknowledgements

This work is supported in part by Proyecto SIP20231470, Secretaría de Investigación y Posgrado, Beca EDI y Comisión de Operación y Fomento de Actividades Académicas (COFAA) del Instituto Politécnico Nacional (IPN), Mexico. We also thank professor Xing Zhi Zhong for constructive comments.

References

- C. S. Wu, E. Ambler, R. W. Hayward, D. D. Hoppes, R. P. Hudson, Experimental Test of Parity Conservation in Beta Decay, Phys. Rev. 105 (1957) 1413-1415. doi:10.1103/PhysRev.105.1413. URL https://link.aps.org/doi/10.1103/PhysRev.105.1413
- [2] T. D. Lee, C. N. Yang, Question of Parity Conservation in Weak Interactions, Phys. Rev. 104 (1956) 254–258. doi:10.1103/PhysRev.104.254. URL https://link.aps.org/doi/10.1103/PhysRev.104.254
- [3] R. L. Garwin, L. M. Lederman, M. Weinrich, Observations of the Failure of Conservation of Parity and Charge Conjugation in Me Phys. Rev. 105 (1957) 1415–1417. doi:10.1103/PhysRev.105.1415. URL https://link.aps.org/doi/10.1103/PhysRev.105.1415
- [4] J. H. Christenson, J. W. Cronin, V. L. Fitch, R. Turlay, Evidence for the 2π Decay of the K_2^0 Meson, Phys. Rev. Lett. 13 (1964) 138–140. doi:10.1103/PhysRevLett.13.138. URL https://link.aps.org/doi/10.1103/PhysRevLett.13.138
- [5] A. Angelopoulos et al., First direct observation of time-reversal non-invariance in the Physics Letters B 444 (1) (1998) 43-51. doi:https://doi.org/10.1016/S0370-2693(98)01356-2. URL https://www.sciencedirect.com/science/article/pii/S037026939
- [6] E. Fermi, Versuch einer Theorie der β-Strahlen. I, Zeitschrift für Physik
 88 (3) (1934) 161–177. doi:10.1007/BF01351864.
 URL https://doi.org/10.1007/BF01351864
- [7] F. L. Wilson, Fermi's Theory of Beta Decay, American Journal of Physics 36 (12) (1968) 1150–1160, a complete English translation is given of the classic Enrico Fermi paper on beta decay published in Zeitschrift für Physik in 1934. arXiv:https://pubs.aip.org/aapt/ajp/article-pdf/36/12/1150/11891 doi:10.1119/1.1974382.
- URL https://doi.org/10.1119/1.1974382 [8] E. C. G. Sudarshan, R. E. Marshak, The Nature of the Four Fermion Inter-
- [6] E. C. O. Sudarshan, K. E. Marshak, The Nature of the Four Fermion Interaction, in: N. Zanichelli (Ed.), Proc. of the Padua-Venice Conference on Mesons and Newly-Discovered Particles, September 1957, Società Italiana di Fisica, 1958, pp. 508–515.
- [9] E. C. G. Sudarshan, R. E. Marshak, Chirality Invariance and the Universal Fermi Interaction, Phys. Rev. 109 (1958) 1860–1862. doi:10.1103/PhysRev.109.1860.2. URL https://link.aps.org/doi/10.1103/PhysRev.109.1860.2
- [10] R. P. Feynman, M. Gell-Mann, Theory of the Fermi Interaction, Phys. Rev. 109 (1958) 193-198. doi:10.1103/PhysRev.109.193.
 URL https://link.aps.org/doi/10.1103/PhysRev.109.193

- [11] N. Cabibbo, Unitary Symmetry and Leptonic Decays, Phys. Rev. Lett. 10 (1963) 531-533. doi:10.1103/PhysRevLett.10.531. URL https://link.aps.org/doi/10.1103/PhysRevLett.10.531
- [12] M. Kobayashi, T. Maskawa, CP-Violation in the Renormalizable Theory of Weak Interaction, Progress of Theoretical Physics 49 (2) (1973) 652-657. arXiv:https://academic.oup.com/ptp/article-pdf/49/2/652/5257692/49-2-652.pdf, doi:10.1143/PTP.49.652. URL https://doi.org/10.1143/PTP.49.652
- B. Pontecorvo, Mesonium and anti-mesonium, Sov. Phys. JETP 6 (1957) 429, [Zh.Eksp.Teor.Fiz. 33 (1957) 549-551].
- [14] B. Pontecorvo, Inverse beta processes and nonconservation of lepton charge, Sov. Phys. JETP 7 (1958) 172–173, [Zh.Eksp.Teor.Fiz. 34 (1957) 247].
- [15] Z. Maki, M. Nakagawa, S. Sakata, Remarks on the Unified Model of Elementary Particles, Progress of Theoretical Physics 28 (1962) 870–880.
- [16] B. Pontecorvo, Neutrino Experiments and the Problem of Conservation of Leptonic Charge, Sov. Phys. JETP 26 (1968) 984–988, [Zh.Eksp.Teor.Fiz. 53 (1967) 1717–1725].
- [17] V. N. Gribov, B. Pontecorvo, Neutrino astronomy and lepton charge, Phys. Lett. B 28 (1969) 493–496. doi:10.1016/0370-2693(69)90525-5.
- [18] R. L. Workman, Others, Review of Particle Physics, PTEP 2022 (2022) 083C01. doi:10.1093/ptep/ptac097.
- [19] A. E. C. Hernández, C. Espinoza, J. C. Gómez-Izquierdo, J. M. González, M. Mondragón, Predictive extended 3HDM with S₄ family symmetry (2022). arXiv:2212.12000.
- [20] B. Belfatto, Z. Berezhiani, Minimally modified Fritzsch texture for quark masses and CKM mixing (2023). arXiv:2305.00069.
- [21] L.-L. Chau, W.-Y. Keung, Comments on the parametrization of the kobayashi-maskawa matrix, Phys. Rev. Lett. 53 (1984) 1802–1805. doi:10.1103/PhysRevLett.53.1802. URL https://link.aps.org/doi/10.1103/PhysRevLett.53.1802
- [22] L. Wolfenstein, Parametrization of the kobayashi-maskawa matrix, Phys. Rev. Lett. 51 (1983) 1945–1947. doi:10.1103/PhysRevLett.51.1945.
- URL https://link.aps.org/doi/10.1103/PhysRevLett.51.1945
 [23] Mathematics Stack Exchange: orangeskid, Unitary matrix decomposition using orthogonal matrices, https://math.stackexchange.com/questions/1127779/unitary-matrix-decomposition-using-orthogonal-matrices, Accessed: 2023-08-05 (2015).
- [24] P. Kielanowski, Two-angle parametrization of the Kobayashi-Maskawa matrix: A relation between CP violation and the Cabibbo-type angles, Phys. Rev. Lett. 63 (1989) 2189–2191. doi:10.1103/PhysRevLett.63.2189. URL https://link.aps.org/doi/10.1103/PhysRevLett.63.2189