Allowed region for the (second) lightest mass $M_{1,2}$ of right-handed neutrino $\nu_{R1,2}$ with $SO(10)$ -inspired relations and sequential dominance

Masaki J. S. Yang^{1,*}

¹Department of Physics, Saitama University, Shimo-okubo, Sakura-ku, Saitama, 338-8570, Japan

In this paper, we evaluate the (second) lightest mass $M_{1,2}$ of right-handed neutrino $\nu_{R1,2}$ in the type-I seesaw mechanism with $SO(10)$ -inspired relations that predicts an almost massless neutrino $m_{1 \text{ or } 3} \sim 0$. By chiral perturbative treatment, the masses $M_{1,2}$ are expressed as $M_1 = m_{D1}^2/m_{11}, M_2 = m_{D2}^2m_{11}/(m_{11}m_{22} - m_{12}^2)$ with the mass matrix of left-handed neutrinos m in the diagonal basis of the Dirac mass matrix m_D .

Assuming m_{Di} and the unitary matrix V in the singular value decomposition $(m_D)_{ij} =$ $\sum_{k} V_{ik} m_{Dk} U^{\dagger}_{kj}$ are close to observed fermion masses and the CKM matrix, we obtain expressions of $M_{1,2}$ by parameters in the low energy and unknown phases. As a result, for $m_{D1} \simeq 0.5$ MeV and $m_{D2} \simeq 100$ MeV, their allowed regions are found to be $M_1^{\rm NH} \simeq 3 \times 10^{4-6}$ GeV and $M_2^{\text{NH}} \simeq 3 \times 10^{6-8}$ GeV in the NH, $M_1^{\text{IH}} \simeq 5 \times 10^{3-4}$ GeV and $M_2^{\text{IH}} \simeq 4 \times 10^{8-9}$ GeV in the IH. These upper and lower bounds are proportional to m_{Di}^2 or $m_{D1}m_{D2}$.

I. INTRODUCTION

The type-I seesaw mechanism[[1](#page-13-0)–[4](#page-13-0)] remains one of the most important subjects in particle physics. For the three mass matrices in this mechanism, texture studies have been extensively investigated[[5](#page-13-0)–[9](#page-13-0)]. For example, the sequential dominance (SD) [\[10](#page-13-0)–[22](#page-14-0)] attempt to explain the large mixings of the neutrino oscillation by a very large mass M_3 of the heaviest right-handed neutrino ν_{R3} . In the limit of $M_3 \to \infty$, a neutrino approaches to be massless. It has also been argued that such a light neutrino is associated with an approximate lepton number symmetry [\[23](#page-14-0)–[27](#page-14-0)]. These lepton number symmetries $U(1)_L³ \times U(1)_R³$ can be reinterpreted as kind of chiral symmetries.

Meanwhile, chiral perturbative treatment by singular values m_{Di} of the Dirac mass matrix m_D has made it possible to calculate three right-handed neutrino masses M_i [[28–30](#page-14-0)]. These

[∗]Electronic address: mjsyang@mail.saitama-u.ac.jp

results are used in the evaluation of N_2 leptogenesis [\[31](#page-14-0), [32\]](#page-14-0) based on $SO(10)$ -inspired mass relations[[33–40](#page-14-0)] (and for a review,[[41\]](#page-14-0)). In this treatment, input parameters are the mass matrix of light neutrinos m_{ν} , the singular values m_{Di} of the Dirac mass matrix m_D , and the left-handed unitary matrix V in the singular value decomposition $m_D = V m_D^{\text{diag}} U^{\dagger}$. Thus, by assuming m_{Di} and V to be about the observed fermion masses and the CKM matrix V_{CKM} , we can understand analytic dependences of M_i on the low energy observables in some grand unified theories.

In this paper, as a simplified problem, we evaluate the lighter masses $M_{1,2}$ when the lightest neutrino mass $m_{1 \text{ or } 3}$ is very small. In the parameter region where the chiral perturbative expansion is valid, the limit $m_{1 \text{ or } 3} \to 0$ corresponds to the sequential dominance with $M_3 \to \infty$. We will analyze the allowed regions for the lighter masses $M_{1,2}$ in such a model. This result would be valid for discussions of cosmology, such as leptogenesis, because it is encompassed by the results of some grand unified theories with seesaw mechanisms.

This paper is organized as follows. The next section gives a review of the chiral perturbative diagonalization of M_R and the sequential dominance in the type-I seesaw mechanism. In Sec. 3, we evaluate $M_{1,2}$ with $SO(10)$ -inspired relations and the sequential dominance. The final section is devoted to a summary.

II. CHIRAL PERTURBATIVE DIAGONALIZATION OF M_R

In this section, we review the chiral perturbative diagonalization of the mass matrix of righthanded neutrinos M_R [\[29](#page-14-0), [38](#page-14-0), [39](#page-14-0), [42, 43\]](#page-14-0). The Dirac mass matrix m_D and the mass matrix of neutrinos $m_{\nu} = U_{\text{MNS}} m_{\nu}^{\text{diag}} U_{\text{MNS}}^{T}$ are defined in a basis where the Yukawa matrix of charged leptons Y_e and M_R are diagonal. Here, U_{MNS} is the MNS matrix. By the singular value decomposition $m_D = V m_D^{\text{diag}} U^{\dagger}$ with unitary matrices V and U, the type-I seesaw mechanism in the diagonal basis of m_D is represented as

$$
m_D^{\text{diag}} V^T m_\nu^{-1} V m_D^{\text{diag}} = U^T M_R^{\text{diag}} U. \tag{1}
$$

By defining the mass matrix $V^T m_{\nu}^{-1} V \equiv m^{-1}$ (and $V^{\dagger} m_{\nu} V^* = m$) in this basis, matrix elements of M_R are simply written by $(M_R)_{ij} \equiv (U^T M_R^{\text{diag}} U)_{ij} = m_{Di}(m^{-1})_{ij} m_{Dj}$. Here, singular values m_{Di} of m_D are assumed to be sufficiently hierarchical $(m_{D3} \gg m_{D2} \gg m_{D1})$.

For an invertible m (i.e., the lightest mass $m_{1 \text{ or } 3}$ is finite), M_R is diagonalized chiral perturbatively. Once the heaviest right-handed neutrino ν_{R3} is integrated out, the mass matrix M_{R0} of the remaining two lighter generations is

$$
M_{R0} = \begin{pmatrix} m_{D1}^2 (m^{-1})_{11} & m_{D1} m_{D2} (m^{-1})_{12} \\ m_{D1} m_{D2} (m^{-1})_{12} & m_{D2}^2 (m^{-1})_{22} \end{pmatrix}
$$
 (2)

$$
-\frac{1}{m_{D3}^2(m^{-1})_{33}}m_{D3}^2\left(\frac{m_{D1}(m^{-1})_{13}}{m_{D2}(m^{-1})_{23}}\right)\otimes\left(m_{D1}(m^{-1})_{13} \ m_{D2}(m^{-1})_{23}\right) \tag{3}
$$

$$
= \frac{1}{m_{11}m_{22} - m_{12}^2} \begin{pmatrix} m_{D1} & 0 \\ 0 & m_{D2} \end{pmatrix} \begin{pmatrix} m_{22} & -m_{12} \\ -m_{12} & m_{11} \end{pmatrix} \begin{pmatrix} m_{D1} & 0 \\ 0 & m_{D2} \end{pmatrix} . \tag{4}
$$

This means that to integrate out ν_{R3} simply yields the seesaw mechanism of the remaining two generations.

By integrating out the second heaviest neutrino ν_{R2} again, m^{-1} becomes the "inverse of 1×1 matrix" $1/m_{11}$, and the diagonalized elements M_i of the mass matrix M_R are found to be

$$
(M_1, M_2, M_3) \simeq \left(\frac{m_{D1}^2}{m_{11}}, \frac{m_{D2}^2 m_{11}}{m_{11} m_{22} - m_{12}^2}, m_{D3}^2 (m^{-1})_{33}\right). \tag{5}
$$

Strictly the singular values of M_R are the absolute values of Eq. (5). This is a kind of chiral perturbative expansion because M_i becomes zero in the limit of $m_{Di} \rightarrow 0$.

The diagonalization matrix of M_R associated with the integration out of heavy states is ¹

$$
U \simeq \begin{pmatrix} 1 & \frac{m_{D1}}{m_{D2}} \frac{m_{12}^*}{m_{11}^*} & 0 \\ -\frac{m_{D1}}{m_{D2}} \frac{m_{12}}{m_{11}} & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & -\frac{m_{D1}}{m_{D3}} \frac{(m^{-1})_{13}^*}{(m^{-1})_{23}^*} \\ 0 & 1 & -\frac{m_{D2}}{m_{D3}} \frac{(m^{-1})_{23}^*}{(m^{-1})_{33}^*} \\ \frac{m_{D1}}{m_{D3}} \frac{(m^{-1})_{13}^*}{(m^{-1})_{33}} & \frac{m_{D2}}{m_{D3}} \frac{(m^{-1})_{23}^*}{(m^{-1})_{33}} \end{pmatrix}
$$
(6)

$$
\simeq \begin{pmatrix} 1 & \frac{m_{D1}}{m_{D2}} \frac{m_{12}^*}{m_{11}} & \frac{m_{D1}}{m_{D3}} \frac{m_{13}^*}{m_{11}^*} \\ -\frac{m_{D1}}{m_{D2}} \frac{m_{12}}{m_{11}} & 1 & -\frac{m_{D2}}{m_{D3}} \frac{(m^{-1})_{23}^*}{(m^{-1})_{33}^*} \\ \frac{m_{D1}}{m_{D3}} \frac{(m^{-1})_{13}^*}{(m^{-1})_{33}} & \frac{m_{D2}}{m_{D3}} \frac{(m^{-1})_{23}^*}{(m^{-1})_{33}} & 1 \end{pmatrix} .
$$
(7)

This procedure perturbatively "solves" all six constraints of the type-I seesaw mechanism [\[38](#page-14-0), [39\]](#page-14-0). In other words, U and M_i are determined by input parameters m_{ν}, m_{Di} and V [[44\]](#page-14-0).

¹ The 1-3 element of U is not a misprint, but due to the sum of two terms. This makes U not look like a unitary matrix, because m_{D1}/m_{D3} behaves as a second-order perturbation.

For the chiral perturbative diagonalization to be a good description, U must not have a large mixing and the following conditions are required;

$$
\frac{m_{D1}}{m_{D2}} \left| \frac{m_{12}}{m_{11}} \right| , \frac{m_{D1}}{m_{D3}} \left| \frac{(m^{-1})_{13}}{(m^{-1})_{33}} \right| , \frac{m_{D2}}{m_{D3}} \left| \frac{(m^{-1})_{23}}{(m^{-1})_{33}} \right| \lesssim 0.1 . \tag{8}
$$

Since elements m_{ij} are expected to be comparable unless m_{ν} and V have special forms, this perturbative treatment is valid in a wide parameter region.

A. Sequential dominance by the limit of $m_{1 \text{ or } 3} \rightarrow 0$

The above discussion requires that m_{ν} have an inverse matrix. Let us examine the behavior of the limit $m_{1 \text{ or } 3} \rightarrow 0$ (This is corresponds to performing a chiral perturbative expansion by $m_{1 \text{ or } 3}$). There are two limits in which the matrix $m_{\nu} = m_D M_R^{-1} m_D^T$ becomes singular;

- 1. The rank of m_D approaches two.
- 2. $M_3 \to \infty$.

In the case of 1, (almost) zero singular values of m_D and m_{ν} yields a chiral symmetry associated with the lepton number[[25–27](#page-14-0), [45\]](#page-14-0);

$$
R m_D = m_D
$$
, $R m_{\nu} = m_{\nu} R = m_{\nu}$, $R = \text{diag}(e^{i\alpha}, 1, 1)$. (9)

The following shows that R for m_{ν} and m_D is identical. For each singular value decomposition $m_{\nu} \equiv U_{\rm MNS} m_{\nu}^{\rm diag} U_{\rm MNS}^{T}$ and $m_{D} = V m_{D}^{\rm diag} U^{\dagger}$, the seesaw mechanism is written by

$$
\begin{pmatrix} 0 & 0 & 0 \ 0 & m_2 & 0 \ 0 & 0 & m_3 \end{pmatrix} = U_{\text{MNS}}^{\dagger} V \begin{pmatrix} 0 & 0 & 0 \ 0 & m_{D2} & 0 \ 0 & 0 & m_{D3} \end{pmatrix} U^{\dagger} (M_R^{\text{diag}})^{-1} U^* \begin{pmatrix} 0 & 0 & 0 \ 0 & m_{D2} & 0 \ 0 & 0 & m_{D3} \end{pmatrix} V^T U_{\text{MNS}}^*.
$$
 (10)

The first column of U_{MNS} and V must match in order for the singular value decomposition (by $U_{\text{MNS}}^{\dagger}V$ to close in the 2-3 subspace. Thus, the eigenvectors of U_{MNS} and V in the massless direction must coincide, and Eq. (9) holds simultaneously in a proper basis.

For the normal hierarchy (NH) $m_1 = 0$ and the inverted hierarchy (IH) $m_3 = 0$, the form of $m = V^{\dagger} m_{\nu} V^*$ in this basis is respectively

$$
m = \begin{pmatrix} 0 & 0 & 0 \\ 0 & * & * \\ 0 & * & * \end{pmatrix} \quad \text{or} \quad \begin{pmatrix} * & * & 0 \\ * & * & 0 \\ 0 & 0 & 0 \end{pmatrix}, \tag{11}
$$

where $*$ denotes appropriate matrix elements. That is, m_{1i} or $m_{3i} = 0$ is a consequence of rank $m_D = 2$ (for det $M_R \neq 0$). In this case the perturbative diagonalization for M_R breaks down [\[43](#page-14-0)], because the cancellation of the approximate chiral symmetry between m_D and m_{ν}^{-1} leads to a matrix M_R without the chiral symmetry. Chiral perturbative treatment can be possible in the remaining two generations. However, in this situation, the unitary matrix $V \sim U_{MNS}$ of the Dirac mass matrix is far from the CKM matrix. Specifically,

$$
V^{\rm NH} \sim \begin{pmatrix} -2/\sqrt{6} & * & * \\ 1/\sqrt{6} & * & * \\ 1/\sqrt{6} & * & * \end{pmatrix}, \quad V^{\rm IH} \sim \begin{pmatrix} * & * & 0 \\ * & * & -1/\sqrt{2} \\ * & * & 1/\sqrt{2} \end{pmatrix}, \tag{12}
$$

for the case of NH and IH. Thus, there is no guideline for m_{Di} and V from unified theories.

On the other hand, the second case $M_3 \to \infty$ can be treated by chiral perturbative expansion and it corresponds to the sequential dominance[[10–12](#page-13-0)]. Since several parameters become unphysical as the heaviest neutrino ν_{R3} is decoupled, the seesaw mechanism is simplified and analysis becomes easier in this limit.

B. Behavior of M_R in the limit of $m_{1 \text{ or } 3} \rightarrow 0$

Here, we explore the chiral perturbative behavior of M_R in the limit of $m_{1 \text{ or } 3} \to 0$. A misalignment of diagonalization W between m_{ν} and m_D is defined as $V^{\dagger}U_{\rm MNS} \equiv W \equiv (w_1, w_2, w_3)$ by 3-dimensional vectors w_i . The spectral decomposition $m^{-1} = V^T m_{\nu}^{-1} V = W^* m_{\nu}^{\text{diag}-1} W^{\dagger}$ is written by

$$
(m^{-1})^* = \frac{1}{m_1} w_1 \otimes w_1^T + \frac{1}{m_2} w_2 \otimes w_2^T + \frac{1}{m_3} w_3 \otimes w_3^T.
$$
 (13)

In Eq. (13), the rank one matrix diverging in the limit of $m_{1 \text{ or } 3} \rightarrow 0$ contributes to M_3 . In particular,

$$
(m^{-1})_{33}^* = \frac{W_{31}^2}{m_1} + \frac{W_{32}^2}{m_2} + \frac{W_{33}^2}{m_3} \simeq \frac{W_{31}^2}{m_1} \text{ or } \frac{W_{33}^2}{m_3} \,. \tag{14}
$$

If the matrix V is similar to V_{CKM} , W has large mixings and matrix elements of W are comparable. Then this approximation holds with $m_{1 \text{ or } 3}/m_2 \lesssim 0.1$, and it is valid when $m_1 \lesssim 1 \text{ meV}$ in NH and $m_3 \lesssim 5$ meV in IH. In this case the heaviest singular value M_3 is evaluated as

$$
M_3 \simeq m_{D3}^2 (m^{-1})_{33} \simeq \frac{m_{D3}^2}{m_1} |W_{31}^*|^2 \quad \text{or} \quad \frac{m_{D3}^2}{m_3} |W_{33}^*|^2 \; . \tag{15}
$$

This situation will be realized by a sufficiently heavy M_3 in some GUT models with strong hierarchy.

These chiral relations of M_i are expected to hold at a higher energy scale because the Yukawa interactions of the first and second generations hardy receive the renormalization. Only when m_{D3} is about as large as the top quark mass m_t , it receives a quantum correction of about 10 % [\[46](#page-14-0)]. Furthermore, the renormalization of gauge interactions in the Standard Model cancels out in the numerator and denominator, the expressions for $M_{1,2}$ are rather insensitive to quantum corrections.

III. THE (SECOND) LIGHTEST MASS $M_{1,2}$ OF RIGHT-HANDED NEUTRINO $\nu_{R1,2}$ WITH $SO(10)$ -INSPIRED RELATIONS AND SEQUENTIAL DOMINANCE

Here we evaluate the lighter masses $M_{1,2}$ and its allowed regions with $SO(10)$ -inspired relations and sequential dominance. First of all, the singular value decompositions of the Yukawa matrices of leptons in a general basis are defined as

$$
Y_{\nu}^{0} \equiv V_{\nu} Y_{\nu}^{\text{diag}} U_{\nu}^{\dagger}, \quad Y_{e}^{0} \equiv V_{e} Y_{e}^{\text{diag}} U_{e}^{\dagger}.
$$
 (16)

On the other hand, the mass matrices m_D and m_{ν} are defined in the diagonal basis of Y_e ;

$$
Y_e = Y_e^{\text{diag}}, \quad m_D = V m_D^{\text{diag}} U^{\dagger}, \quad m_{\nu} = U_{\text{MNS}} m_{\nu}^{\text{diag}} U_{\text{MNS}}^T. \tag{17}
$$

From this, $V = V_e^{\dagger} V_{\nu}$ and U_{MNS} implicitly involve V_e .

With some grand unified theories in mind, the following $SO(10)$ -inspired relations are assumed.

- 1. $m_{D1} \simeq m_u(\Lambda_{\text{GUT}}) \simeq m_e(\Lambda_{\text{GUT}}) \simeq 0.5 \text{ MeV}, m_{D2} \simeq m_c(\Lambda_{\text{GUT}}) \sim m_u(\Lambda_{\text{GUT}}) \sim 100 \text{ MeV}$ [\[46](#page-14-0)].
- 2. The unitary matrix V of m_D is dominated by 1-2 mixing, and its mixing angle is at most about the Cabibbo angle θ_C .

Specifically, V is expressed by,

$$
V \simeq \begin{pmatrix} e^{i\sigma_1} & 0 & 0 \\ 0 & e^{i\sigma_2} & 0 \\ 0 & 0 & e^{i\sigma_3} \end{pmatrix} \begin{pmatrix} c_\theta & s_\theta e^{i\phi} & O(10^{-3}) \\ -s_\theta e^{-i\phi} & c_\theta & O(10^{-2}) \\ O(10^{-3}) & O(10^{-2}) & 1 \end{pmatrix},
$$
(18)

where $s_{\theta} \equiv \sin \theta \lesssim 0.2$, $c_{\theta} \equiv \cos \theta \sim 1$ and ϕ , σ_i are unknown phases. The signs of s_{θ} , c_{θ} are absorbed into the phases, and θ is restricted to the first quadrant $0 \leq \theta \leq \pi/2$. Unknown phases ρ_i from V_{ν} are omitted because their contributions $m_{ij} \to e^{i\rho_i} m_{ij} e^{i\rho_j}$ are unphysical.

Since these assumptions eliminate the possibility of identical chiral symmetry such as Eq.([11](#page-3-0)), we consider only SD in this situation. From $|W_{3i}| \simeq |U_{\tau i}|$, the absolute value of M_3 [\(15\)](#page-4-0) is hardly affected by V ;

$$
M_3 \simeq m_{D3}^2 (m^{-1})_{33} \simeq \frac{m_{D3}^2}{m_1} |U_{\tau 1}|^2 \quad \text{or} \quad \frac{m_{D3}^2}{m_3} |U_{\tau 3}|^2 \; . \tag{19}
$$

To express $M_{1,2}$ by low energy parameters, m_{11} is written by matrix elements of m_{ν} with $m_{\alpha\beta}=\sum_i m_i U_{\alpha i} U_{\beta i}$ as

$$
m_{11} \simeq (V_{11}^*)^2 m_{ee} + 2V_{11}^* V_{21}^* m_{e\mu} + (V_{21}^*)^2 m_{\mu\mu} , \qquad (20)
$$

$$
= c_{\theta}^{2} e^{-2i\sigma_{1}} m_{ee} - 2s_{\theta} c_{\theta} e^{i(\phi - \sigma_{1} - \sigma_{2})} m_{e\mu} + s_{\theta}^{2} e^{2i(\phi - \sigma_{2})} m_{\mu\mu}.
$$
\n(21)

Since the final result is an absolute value, the extra phase can be removed by multiplying by $e^{2i\sigma_1}$;

$$
|m_{11}| \simeq |c_{\theta}^2 m_{ee} - 2s_{\theta} c_{\theta} e^{i\beta} m_{e\mu} + s_{\theta}^2 e^{2i\beta} m_{\mu\mu}|, \qquad (22)
$$

where $\beta \equiv \phi + \sigma_1 - \sigma_2$ is the only unknown phase that cannot be determined from phenomenology.

Besides, using elements W_{ij} and m_i , we obtain

$$
m_{11}^{\text{NH}} \simeq W_{12}^2 m_2 + W_{13}^2 m_3 \,, \quad m_{11}^{\text{IH}} \simeq W_{11}^2 m_1 + W_{12}^2 m_2 \,. \tag{23}
$$

By uniform redefinition of phases $e^{i\sigma_1}W_{1i} = c_\theta U_{ei} - s_\theta e^{i\beta}U_{\mu i}$ and PDG parameterization of U_{MNS} , these matrix elements are rewritten as

$$
e^{i\sigma_1}W_{11} \simeq c_{\theta}c_{12}c_{13} + s_{\theta}e^{i\beta}(c_{23}s_{12} + c_{12}s_{13}s_{23}e^{i\delta}), \qquad (24)
$$

$$
e^{-i\alpha/2}e^{i\sigma_1}W_{12} \simeq c_\theta s_{12}c_{13} - s_\theta e^{i\beta}(c_{12}c_{23} - s_{12}s_{13}s_{23}e^{i\delta}), \qquad (25)
$$

$$
e^{i\sigma_1} W_{13} \simeq c_\theta s_{13} e^{-i\delta} - s_\theta e^{i\beta} c_{13} s_{23} , \qquad (26)
$$

where $c_{ij} \equiv \cos \theta_{ij}$ and $s_{ij} \equiv \sin \theta_{ij}$. In the limit of $M_3 \to \infty$, the only Majorana phase α is defined as associated with m_2 .

The latest global fit (without Super-Kamiokande) is used for input parameters[[47\]](#page-14-0). The mass differences are

$$
\Delta m_{21}^2 = 74.2 \,\text{meV}^2 \,, \quad \Delta m_{31}^2 = 2515 \,\text{meV}^2 \,, \quad \Delta m_{32}^2 = -2498 \,\text{meV}^2 \,, \tag{27}
$$

where $\Delta m_{31}^2 > 0$ for NH and $\Delta m_{32}^2 < 0$ for IH. The three mixing angles are

$$
\sin^2 \theta_{12}^{\text{NH}} = 0.304 \,, \quad \sin^2 \theta_{23}^{\text{NH}} = 0.573 \,, \quad \sin^2 \theta_{13}^{\text{NH}} = 0.0222 \,, \tag{28}
$$

$$
\sin^2 \theta_{12}^{\text{NH}} = 0.304 \,, \quad \sin^2 \theta_{23}^{\text{NH}} = 0.578 \,, \quad \sin^2 \theta_{13}^{\text{NH}} = 0.0224 \,. \tag{29}
$$

In this paper, only best-fit values are used to grasp behaviors of $M_{1,2}$. The errors in chiral perturbative expansion of m_{Di} are $O(m_{Di}^2/m_{Dj}^2) \lesssim 1$ % [\[48](#page-15-0)], which is smaller than that of neutrino oscillation experiments. There are also another errors of about 4 % due to the neglect of V_{cb} , $V_{ts} \sim 0.04$. However, the largest source of errors come from the neglect of $m_{1 \text{ or } 3}/m_2$.

A. Inverted hierarchy

We start the analysis of IH first because it is easier than NH. From the triangular inequality $|z_1| - |z_2| \le |z_1 + e^{i\alpha}z_2| \le |z_1| + |z_2|$ for complex numbers $z_{1,2}$,

$$
m_1|W_{11}|^2 - m_2|W_{12}|^2 \le |m_{11}^{\text{IH}}| \le m_1|W_{11}|^2 + m_2|W_{12}|^2. \tag{30}
$$

Since W_{11} and W_{12} are functions of β and δ , inequality is not used anymore.

The upper bound is evaluated as

$$
m_1|W_{11}|^2 + m_2|W_{12}|^2
$$
\n
$$
\simeq m_1 \left| c_\theta c_{12} c_{13} + s_\theta e^{i\beta} (c_{23} s_{12} + c_{12} s_{13} s_{23} e^{i\delta}) \right|^2 + m_2 \left| c_\theta s_{12} c_{13} - s_\theta e^{i\beta} (c_{12} c_{23} - s_{12} s_{13} s_{23} e^{i\delta}) \right|^2
$$
\n
$$
= c_{13}^2 c_\theta^2 (m_1 c_{12}^2 + m_2 s_{12}^2) + 2c_{13} c_\theta s_\theta [s_{13} s_{23} (m_1 c_{12}^2 + m_2 s_{12}^2) \cos(\beta + \delta) + c_{12} c_{23} (m_1 - m_2) s_{12} \cos \beta)]
$$
\n
$$
+ s_\theta^2 [2c_{12} c_{23} (m_1 - m_2) s_{12} s_{13} s_{23} \cos \delta + c_{23}^2 (m_1 s_{12}^2 + m_2 c_{12}^2) + s_{13}^2 s_{23}^2 (m_1 c_{12}^2 + m_2 s_{12}^2)].
$$
\n(32)

The maximum value $|m_{11}^{\text{IH}}|^{\text{max}}$ is exist near the point $\beta = \delta = \pi$ because of $m_2 \gtrsim m_1$ and terms proportional to s_{θ}^2 are negligble second-order perturbations. Thus, the Majorana phase is obviously chosen to be $\alpha \simeq 0$.

Substituting $m_1 \simeq m_2$, we obtain

$$
|m_{11}^{\text{IH}}|^{\text{max}} \simeq m_1 (c_\theta^2 c_{13}^2 + 2 s_\theta c_\theta s_{13} c_{13} s_{23} + s_\theta^2 (c_{23}^2 + s_{13}^2 s_{23}^2)) \simeq 50 \,\text{meV} \,. \tag{33}
$$

The dependence of θ is small because the terms in the first order of s_{θ} cancel out by $m_1 \simeq m_2$ and the lowest order is proportional to $s_{\theta}s_{13}$.

The minimum value is evaluated in the same manner.

$$
\frac{m_1|W_{11}|^2 - m_2|W_{12}|^2}{m_1} \tag{34}
$$

$$
\simeq \left| c_{\theta} c_{12} c_{13} + s_{\theta} e^{i\beta} (c_{23} s_{12} + c_{12} s_{13} s_{23} e^{i\delta}) \right|^2 - \left| c_{\theta} s_{12} c_{13} - s_{\theta} e^{i\beta} (c_{12} c_{23} - s_{12} s_{13} s_{23} e^{i\delta}) \right|^2 \tag{35}
$$

$$
= c_{13}^2 c_{\theta}^2 (c_{12}^2 - s_{12}^2) + 2c_{13} c_{\theta} s_{\theta} [2c_{12} c_{23} s_{12} \cos \beta + s_{13} s_{23} (c_{12}^2 - s_{12}^2) \cos(\beta + \delta)] - s_{\theta}^2 [(c_{12}^2 - s_{12}^2)(c_{23}^2 - s_{13}^2 s_{23}^2) - 4c_{12} c_{23} s_{12} s_{13} s_{23} \cos \delta].
$$
\n(36)

Among terms with phase dependences, the dominant one is the term proportional to $\cos \beta$. The rests are suppressed by s_{13} and/or s_{θ} , and the minimum exists at a point near $\beta = \pi, \delta = 0$. We have confirmed these behaviors numerically. From this, the minimum value $|m_{11}^{\text{IH}}|^{\text{min}}$ exists around $\alpha = \pi$,

$$
|m_{11}^{\text{IH}}|^{\text{min}} \simeq m_1(|c_{\theta}c_{12}c_{13} - s_{\theta}(c_{23}s_{12} + c_{12}s_{13}s_{23})|^2 - |c_{\theta}s_{12}c_{13} + s_{\theta}(c_{12}c_{23} - s_{12}s_{13}s_{23})|^2). \tag{37}
$$

By ignoring the second-order perturbations for s_{θ} and s_{13} ,

$$
|m_{11}^{\text{IH}}|^{\text{min}} \simeq m_1(c_{13}^2 c_{\theta}^2(c_{12}^2 - s_{12}^2) - 4s_{\theta} c_{\theta} s_{12} c_{13} c_{12} c_{23}) \simeq 19(1 - 3s_{\theta}) \,\text{meV}. \tag{38}
$$

The numerical values are approximately

$$
|m_{11}^{\text{IH}}|_{\theta=0}^{\text{min}} \simeq 18.6 \,\text{meV} \,, \quad |m_{11}^{\text{IH}}|_{\theta=0.1}^{\text{min}} = 12.2 \,\text{meV} \,, \quad |m_{11}^{\text{IH}}|_{\theta=0.2}^{\text{min}} = 5.53 \,\text{meV} \,. \tag{39}
$$

These expressions coincide with the upper and lower bounds of $|m_{ee}|$ in the limit of $\theta \to 0$.

Therefore, in grand unified theories where the chiral perturbative expansion is valid, the range of $M_1^{\text{IH}} = m_{D1}^2 / |m_{11}^{\text{IH}}|$ in IH with $m_3 \to 0$ is

$$
5 \,\mathrm{TeV} \left(\frac{m_{D1}}{0.5 \,\mathrm{MeV}}\right)^2 \lesssim M_1^{\mathrm{IH}} \lesssim 50 \,\mathrm{TeV} \left(\frac{m_{D1}}{0.5 \,\mathrm{MeV}}\right)^2 \,. \tag{40}
$$

Around these upper and lower bounds, all CP phases are almost trivial and some CP symmetry is expected in the lepton sector. In addition, the latest limit of double beta decay by the KamLAND-Zen collaboration $|m_{ee}| \lesssim 36 - 156 \,\text{meV}$ [\[49](#page-15-0)] is roughly corresponds to about 7 – $1.6 \,\text{TeV} \gtrsim M_1^{\text{IH}}.$

The parameter m_{11} also dominates M_2 . From Eq. [\(5\)](#page-2-0) and the behavior [\(14](#page-4-0)) of $(m^{-1})_{33}$ in SD,

$$
M_2^{\rm IH} \simeq m_{D2}^2 \left| \frac{m_{11}}{\text{Det } m(m^{-1})_{33}} \right| \simeq \frac{m_{D2}^2}{m_1 m_2} \left| \frac{m_{11}}{(W_{33}^*)^2} \right| \simeq \frac{m_{D2}^2}{m_1 m_2} \left| \frac{m_{11}}{U_{73}^2} \right| \,. \tag{41}
$$

The fermion masses of the second generation at the GUT scale are roughly[[46](#page-14-0)]

$$
m_c \simeq 200 \,\text{MeV}, \quad m_s \simeq 20 \,\text{MeV}, \quad m_\mu \simeq 100 \,\text{MeV}.\tag{42}
$$

By normalizing for $m_{D2} \sim 100 \, \text{MeV},$

$$
M_2^{\rm IH} \simeq \left(\frac{m_{D2}}{100 \text{ MeV}}\right)^2 \left|\frac{m_{11}}{m_1}\right| \left(\frac{10 \text{ meV}}{m_2 |U_{\tau 3}^2|}\right) 10^9 \text{ GeV}.
$$
 (43)

Then the allowed region of M_2^{IH} of IH is determined as

$$
0.4 \times 10^9 \,\text{GeV} \left(\frac{m_{D2}}{100 \,\text{MeV}}\right)^2 \lesssim M_2^{\text{IH}} \lesssim 4.0 \times 10^9 \,\text{GeV} \left(\frac{m_{D2}}{100 \,\text{MeV}}\right)^2. \tag{44}
$$

B. Normal hierarchy

We proceed to the analysis of NH. As in the case of IH, there exists the triangular inequality,

$$
m_2|W_{12}|^2 - m_3|W_{13}|^2 \le |m_{11}^{\text{NH}}| \le m_2|W_{12}|^2 + m_3|W_{13}|^2. \tag{45}
$$

For the upper limit,

$$
|W_{12}|^2 m_2 + |W_{13}|^2 m_3 \tag{46}
$$

$$
\simeq m_2 \left| c_\theta s_{12} c_{13} - s_\theta e^{i\beta} (c_{12} c_{23} - s_{12} s_{13} s_{23} e^{i\delta}) \right|^2 + m_3 \left| c_\theta s_{13} e^{-i\delta} - s_\theta e^{i\beta} c_{13} s_{23} \right|^2 \tag{47}
$$

$$
= c_{\theta}^{2}(m_{2}c_{13}^{2}s_{12}^{2}+m_{3}s_{13}^{2}) - 2s_{\theta}[c_{13}c_{\theta}\{m_{2}c_{12}c_{23}s_{12}\cos\beta + (m_{3}-m_{2}s_{12}^{2})s_{13}s_{23}\cos(\beta+\delta)\}] + s_{\theta}^{2}[m_{2}(c_{12}^{2}c_{23}^{2}+s_{12}^{2}s_{13}^{2}s_{23}^{2})+m_{3}s_{23}^{2}c_{13}^{2} - 2m_{2}c_{12}c_{23}s_{12}s_{13}s_{23}\cos\delta].
$$
\n(48)

Due to phase-dependent terms of the first-order in s_{θ} , it reaches the maximum near $\beta = \pi, \delta = 0$ and $\alpha = 0$. The maximum value $|m_{11}^{\text{NH}}|^{\text{max}}$ is expressed as

$$
|m_{11}^{\text{NH}}|^{\text{max}} \simeq m_2[c_{\theta}s_{12}c_{13} + s_{\theta}(c_{12}c_{23} - s_{12}s_{13}s_{23})]^2 + m_3(c_{\theta}s_{13} + s_{\theta}c_{13}s_{23})^2 \tag{49}
$$

$$
\simeq (m_2 s_{12}^2 + m_3 s_{13}^2) + 2s_\theta (s_{13} s_{23} (m_3 - m_2 s_{12}^2) + m_2 c_{12} c_{23} s_{12}), \tag{50}
$$

where the second order terms of s_{θ} are ignored in the second equality. Numerically, it is approximately evaluated as

$$
|m_{11}^{\text{NH}}|_{\theta=0}^{\text{max}} = 3.67 \,\text{meV} \,, \quad |m_{11}^{\text{NH}}|_{\theta=0.1}^{\text{max}} = 5.50 \,\text{meV} \,, \quad |m_{11}^{\text{NH}}|_{\theta=0.2}^{\text{max}} = 7.81 \,\text{meV} \,. \tag{51}
$$

For M_1^{NH} , the same equation as IH holds. For M_2^{NH} , we obtain

$$
M_2 \simeq m_{D2}^2 \left| \frac{m_{11}}{\text{Det } m(m^{-1})_{33}} \right| \simeq \frac{m_{D2}^2}{m_2 m_3} \left| \frac{m_{11}}{U_{\tau_1}^2} \right| \,. \tag{52}
$$

By a proper normalization,

$$
M_2 \simeq \left(\frac{m_{D2}}{100 \,\text{MeV}}\right)^2 \left(\frac{100 \,\text{meV}^2}{m_2 m_3 |U_{\tau 1}|^2}\right) \left(\frac{|m_{11}|}{1 \,\text{meV}}\right) 10^8 \,\text{GeV} \,. \tag{53}
$$

Therefore, the minimum value of M_1^{NH} and the maximum value of M_2^{NH} are given by the maxmum value of $|m_{11}|$ with $\theta = 0.2$,

$$
M_1^{\min} \simeq 30 \,\text{TeV} \left(\frac{m_{D1}}{0.5 \,\text{MeV}} \right)^2 \,, \quad M_2^{\max} \simeq 3 \times 10^8 \,\text{GeV} \left(\frac{m_{D2}}{100 \,\text{MeV}} \right)^2 \,. \tag{54}
$$

On the other hand, the minimum of m_{11}^{NH} is

$$
|W_{12}|^2 m_2 - |W_{13}|^2 m_3 \tag{55}
$$

$$
\simeq m_2 \left| c_\theta s_{12} c_{13} - s_\theta e^{i\beta} (c_{12} c_{23} - s_{12} s_{13} s_{23} e^{i\delta}) \right|^2 - m_3 \left| c_\theta s_{13} e^{-i\delta} - s_\theta e^{i\beta} c_{13} s_{23} \right|^2 \tag{56}
$$

$$
= m_2 s_{12}^2 c_{13}^2 c_{\theta}^2 - m_3 (c_{13}^2 s_{23}^2 s_{\theta}^2 + s_{13}^2 c_{\theta}^2) + 2c_{13} c_{\theta} s_{\theta} (s_{13} s_{23} (m_2 s_{12}^2 + m_3) \cos(\beta + \delta) - c_{12} c_{23} m_2 s_{12} \cos \beta) + m_2 s_{\theta}^2 (c_{12}^2 c_{23}^2 + s_{12}^2 s_{13}^2 s_{23}^2 - 2c_{12} c_{23} s_{12} s_{13} s_{23} \cos \delta).
$$
\n(57)

This is minimized near $\beta = 0$ and $\delta = \pi$. Thus the minimum value $(m_{11}^{\text{NH}})^{\text{min}}$ of m_{11}^{NH} exists near $\alpha \simeq \pi$, and a rough evaluation of the first order of s_{θ} is

$$
(m_{11}^{\text{NH}})^{\text{min}} \simeq m_2(c_{\theta}s_{12}c_{13} - s_{\theta}(c_{12}c_{23} + s_{12}s_{13}s_{23}))^2 - m_3(c_{\theta}s_{13} + s_{\theta}c_{13}s_{23})^2 \tag{58}
$$

$$
\simeq (m_2 s_{12}^2 - m_3 s_{13}^2) - 2s_\theta (s_{13} s_{23} (m_3 + m_2 s_{12}^2) + m_2 c_{12} c_{23} s_{12}). \tag{59}
$$

The numerical values are

$$
(m_{11}^{\text{NH}})_{\theta=0}^{\text{min}} \simeq 1.44 \,\text{meV} \,, \quad (m_{11}^{\text{NH}})_{\theta=0.1}^{\text{min}} = -0.50 \,\text{meV} \,, \quad (m_{11}^{\text{NH}})_{\theta=0.2}^{\text{min}} = -2.92 \,\text{meV} \,. \tag{60}
$$

For a finite s_{θ} , m_{11}^{NH} can have a negative value. This corresponds to the region where $|m_{11}^{\text{NH}}|^{min} = 0$ is possible for appropriate complex phases. The condition for such a cancellation to occur is $\sqrt{m_2}W_{12} \simeq \pm i\sqrt{m_3}W_{13}$, and the minimum θ that satisfies the condition is

$$
\sqrt{m_2}(c_{\theta}s_{12}c_{13} - s_{\theta}(c_{12}c_{23} + s_{12}s_{13}s_{23})) = \pm \sqrt{m_3}(c_{\theta}s_{13} + s_{\theta}c_{13}s_{23}). \tag{61}
$$

By choosing the solution with the smaller θ ,

$$
s_{\theta} = \frac{c_{13}s_{12} - \sqrt{\frac{m_3}{m_2}}s_{13}}{c_{13}\sqrt{\frac{m_3}{m_2}}s_{23} + c_{12}c_{23} + s_{12}s_{13}s_{23}} \simeq 0.077. \tag{62}
$$

For a larger θ than this value, there is a canceling region where $(m_{11}^{\text{NH}})^{\text{min}} \approx 0$. The phase dependence of the minimum is shown in Fig. 1. We can see that such cancellation is likely to occur in the region $\delta + \beta \sim \pi$.

FIG. 1: Plots of $|W_{12}|^2 m_2 - |W_{13}|^2 m_3$ [\(48\)](#page-9-0) for δ and β in NH. The left and right ones represent the case of $s_{\theta} = 0.1, 0.2$ respectively.

C. $M_{1,2}$ in the level crossing point $(m_{11}^{\text{NH}})^{\text{min}} \simeq 0$

The chiral perturbative expansion breaks down in the limit of $m_{11} \rightarrow 0$. This is clear from Eq. [\(4](#page-2-0)) because the diagonalization of the mass matrix M_{R0} for the lighter generations has a large mixing. In this *level crossing point* [[29\]](#page-14-0), the masses M_1 and M_2 are of equal magnitude, and the relation $M_1^{-1} \propto m_{11} \propto M_2$ no longer holds. Let us examine how the mass matrices m and M_{R0} behave in the region where m_{11} is small. From Eq. [\(14](#page-4-0)), the minor determinant is evaluated as

$$
|m_{11}m_{22} - m_{12}^2| = |\text{Det }m(m^{-1})_{33}| \simeq m_2 m_3 |(W_{31}^*)^2| \simeq m_2 m_3 |U_{\tau 1}|^2. \tag{63}
$$

Thus in the limit of $m_{11} \rightarrow 0$, $|m_{12}| \simeq \sqrt{m_2 m_3} |U_{\tau_1}| \simeq 15 \text{ meV}$ holds. Since the value $|m_{22}| \simeq$ $|m_{\mu\mu}| \simeq m_3/2 \simeq 25 \,\text{meV}$ is a roughly constant under θ , the behaviors of m and M_{R0} [\(4\)](#page-2-0) in the limit of $m_{11} \rightarrow 0$ are

$$
M_{R0} = \frac{1}{m_{11}m_{22} - m_{12}^2} \begin{pmatrix} m_{D1} & 0 \\ 0 & m_{D2} \end{pmatrix} \begin{pmatrix} m_{22} & -m_{12} \\ -m_{12} & m_{11} \end{pmatrix} \begin{pmatrix} m_{D1} & 0 \\ 0 & m_{D2} \end{pmatrix}, \quad (64)
$$

$$
|M_{R0}| \simeq \frac{1}{m_2 m_3 |U_{\tau 1}|^2} \begin{pmatrix} m_{D1}^2 m_3 / 2 & m_{D1} m_{D2} \sqrt{m_2 m_3} |U_{\tau 1}| \\ m_{D1} m_{D2} \sqrt{m_2 m_3} |U_{\tau 1}| & 0 \end{pmatrix}.
$$
 (65)

From $|m_{12}| \sim |m_{22}|$, the off-diagonal element is dominant in unified theories with $m_{D2} \gtrsim 20 m_{D1}$, and the two mass singular values are almost same $M_1 \simeq M_2$.

Such a condition is also evaluated from the relation of determinants and Eq. [\(15](#page-4-0));

$$
M_1 M_2 M_3 = \frac{m_{D1}^2 m_{D2}^2 m_{D3}^2}{m_1 m_2 m_3}, \quad M_1 M_2 = \frac{m_{D1}^2 m_{D2}^2}{m_2 m_3 |U_{\tau 1}|^2}.
$$
 (66)

When M_1 and M_2 are approximately equal,

$$
M_1 \simeq M_2 \simeq \frac{m_{D1}m_{D2}}{\sqrt{m_2 m_3}|U_{\tau 1}|}.
$$
\n(67)

It gives an upper bound on M_1^{NH} and a lower bound on M_2^{NH} simultaneously,

$$
M_1^{\max} \simeq M_2^{\min} \simeq \left(\frac{m_{D1}}{1 \,\text{MeV}}\right) \left(\frac{m_{D2}}{100 \,\text{MeV}}\right) \left(\frac{10 \,\text{meV}}{|U_{\tau 1}|\sqrt{m_2 m_3}}\right) 10^7 \text{ GeV}. \tag{68}
$$

Therefore, the bounds will be about 10 PeV. Unlike with the case of IH, these upper and lower bounds are proportional to $m_{D1}m_{D2}$. To summarize these results,

$$
3 \times 10^4 \,\text{GeV} \left(\frac{m_{D1}}{0.5 \text{ MeV}}\right)^2 \lesssim M_1^{\text{NH}} \lesssim 3 \times 10^6 \,\text{GeV} \left(\frac{m_{D1} m_{D2}}{50 \text{ MeV}^2}\right),\tag{69}
$$

$$
3 \times 10^6 \,\text{GeV} \left(\frac{m_{D1}m_{D2}}{50 \text{ MeV}^2}\right) \lesssim M_2^{\text{NH}} \lesssim 3 \times 10^8 \,\text{GeV} \left(\frac{m_{D2}}{100 \text{ MeV}}\right)^2. \tag{70}
$$

Since M_1M_2 is approximately constant under θ , the widths of these allowed regions are comparable.

IV. SUMMARY

In this paper, we perform a chiral perturbative evaluation of the (second) lightest mass $M_{1,2}$ of right-handed neutrino $\nu_{R1,2}$ in the type-I seesaw mechanism with $SO(10)$ -inspired relations

and an almost massless neutrino $m_{1 \text{ or } 3} \sim 0$. By chiral perturbative treatment, the masses $M_{1,2}$ are expressed as $M_1 = m_{D1}^2/m_{11}$, $M_2 = m_{D2}^2 m_{11}/(m_{11}m_{22} - m_{12}^2)$ with the mass matrix of left-handed neutrinos m in the diagonal basis of the Dirac mass matrix m_D . Assuming m_{Di} and the unitary matrix V in the singular value decomposition $(m_D)_{ij} = \sum_k V_{ik} m_{Dk} U^{\dagger}_{kj}$ are close to observed fermion masses and the CKM matrix, we obtained expressions for $M_{1,2}$ by parameters in the low energy and unknown phases.

As a result, the upper and lower bounds of $M_{1,2}$ are obtained as

$$
3 \times 10^4 \,\text{GeV} \left(\frac{m_{D1}}{0.5 \text{ MeV}}\right)^2 \lesssim M_1^{\text{NH}} \lesssim 3 \times 10^6 \,\text{GeV} \left(\frac{m_{D1} m_{D2}}{50 \text{ MeV}^2}\right),\tag{71}
$$

$$
3 \times 10^6 \,\mathrm{GeV} \left(\frac{m_{D1}m_{D2}}{50 \,\mathrm{MeV}^2}\right) \lesssim M_2^{\mathrm{NH}} \lesssim 3 \times 10^8 \,\mathrm{GeV} \left(\frac{m_{D2}}{100 \,\mathrm{MeV}}\right)^2. \tag{72}
$$

for the NH, and

$$
5 \times 10^3 \,\text{GeV} \left(\frac{m_{D1}}{0.5 \text{ MeV}}\right)^2 \lesssim M_1^{\text{IH}} \lesssim 5 \times 10^4 \,\text{GeV} \left(\frac{m_{D1}}{0.5 \text{ MeV}}\right)^2 ,\tag{73}
$$

$$
4 \times 10^8 \,\text{GeV} \left(\frac{m_{D2}}{100 \,\text{MeV}}\right)^2 \lesssim M_2^{\text{IH}} \lesssim 4 \times 10^9 \,\text{GeV} \left(\frac{m_{D2}}{100 \,\text{MeV}}\right)^2 ,\tag{74}
$$

for the IH. The difference in the dependence of m_{Di} is due to a cancellation of m_{11} in NH. This result would be valid for discussions of cosmology, such as leptogenesis, because it is encompassed by the results of some grand unified theories with seesaw mechanisms.

- [1] P. Minkowski, Phys. Lett. 67B, 421 (1977).
- [2] M. Gell-Mann, P. Ramond, and R. Slansky, Conf. Proc. C790927, 315 (1979).
- [3] T. Yanagida, Conf. Proc. C7902131, 95 (1979).
- [4] R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. 44, 912 (1980).
- [5] H. Fritzsch and Z.-z. Xing, Prog. Part. Nucl. Phys. 45, 1 (2000), arXiv:hep-ph/9912358.
- [6] Z.-z. Xing and Z.-h. Zhao, Rept. Prog. Phys. 79, 076201 (2016), arXiv:1512.04207.
- [7] Z.-z. Xing, Phys. Rept. 854, 1 (2020), arXiv:1909.09610.
- [8] Z.-z. Xing and Z.-h. Zhao, Rept. Prog. Phys. 84, 066201 (2021), arXiv:2008.12090.
- [9] Z.-z. Xing, (2022), arXiv:2210.11922.
- [10] S. F. King, Phys. Lett. B 439, 350 (1998), arXiv:hep-ph/9806440.
- [11] S. F. King, Nucl. Phys. B 562, 57 (1999), arXiv:hep-ph/9904210.
- [12] S. F. King, Nucl. Phys. B 576, 85 (2000), arXiv:hep-ph/9912492.
- [13] S. F. King, JHEP 09, 011 (2002), arXiv:hep-ph/0204360.
- [14] S. Antusch and S. F. King, New J. Phys. 6, 110 (2004), arXiv:hep-ph/0405272.
- [15] S. Antusch and S. F. King, Phys. Lett. B 591, 104 (2004), arXiv:hep-ph/0403053.
- [16] S. Antusch, S. F. King, and A. Riotto, JCAP 11, 011 (2006), arXiv:hep-ph/0609038.
- [17] S. Antusch, S. Boudjemaa, and S. F. King, JHEP 09, 096 (2010), arXiv:1003.5498.
- [18] S. F. King, JHEP 08, 105 (2005), arXiv:hep-ph/0506297.
- [19] S. Antusch and S. F. King, Phys. Lett. B 659, 640 (2008), arXiv:0709.0666.
- [20] S. Antusch, S. F. King, C. Luhn, and M. Spinrath, Nucl. Phys. B 856, 328 (2012), arXiv:1108.4278.
- [21] S. Antusch, S. F. King, and M. Spinrath, Phys. Rev. D 87, 096018 (2013), arXiv:1301.6764.
- [22] F. Björkeroth and S. F. King, J. Phys. G 42, 125002 (2015), arXiv:1412.6996.
- [23] D. Wyler and L. Wolfenstein, Nucl. Phys. B 218, 205 (1983).
- [24] S. T. Petcov and S. T. Toshev, Phys. Lett. B 143, 175 (1984).
- [25] G. C. Branco, W. Grimus, and L. Lavoura, Nucl. Phys. B 312, 492 (1989).
- [26] J. Kersten and A. Y. Smirnov, Phys. Rev. D 76, 073005 (2007), arXiv:0705.3221.
- [27] R. Adhikari and A. Raychaudhuri, Phys. Rev. D 84, 033002 (2011), arXiv:1004.5111.
- [28] G. C. Branco, R. Gonzalez Felipe, F. R. Joaquim, and M. N. Rebelo, Nucl. Phys. B 640, 202 (2002), arXiv:hep-ph/0202030.
- [29] E. K. Akhmedov, M. Frigerio, and A. Y. Smirnov, JHEP 09, 021 (2003), arXiv:hep-ph/0305322.
- [30] S. Davidson, JHEP 03, 037 (2003), arXiv:hep-ph/0302075.
- [31] P. Di Bari, Nucl. Phys. B 727, 318 (2005), arXiv:hep-ph/0502082.
- [32] O. Vives, Phys. Rev. D73, 073006 (2006), arXiv:hep-ph/0512160.
- [33] P. Di Bari and A. Riotto, Phys. Lett. B 671, 462 (2009), arXiv:0809.2285.
- [34] E. Bertuzzo, P. Di Bari, and L. Marzola, Nucl. Phys. B 849, 521 (2011), arXiv:1007.1641.
- [35] P. Di Bari and A. Riotto, JCAP 04, 037 (2011), arXiv:1012.2343.
- [36] P. Di Bari and L. Marzola, Nucl. Phys. B 877, 719 (2013), arXiv:1308.1107.
- [37] P. Di Bari, S. King, and M. Re Fiorentin, JCAP 03, 050 (2014), arXiv:1401.6185.
- [38] P. Di Bari, L. Marzola, and M. Re Fiorentin, Nucl. Phys. B 893, 122 (2015), arXiv:1411.5478.
- [39] P. Di Bari and M. Re Fiorentin, JHEP 10, 029 (2017), arXiv:1705.01935.
- [40] M. Chianese and P. Di Bari, JHEP 05, 073 (2018), arXiv:1802.07690.
- [41] P. Di Bari, Prog. Part. Nucl. Phys. **122**, 103913 (2022), arXiv:2107.13750.
- [42] M. J. S. Yang, Nucl. Phys. B **994**, 116318 (2023), arXiv:2211.15101.
- [43] M. J. S. Yang, Phys. Lett. B 843, 138061 (2023), arXiv:2304.04513.
- [44] S. Davidson and A. Ibarra, JHEP 09, 013 (2001), arXiv:hep-ph/0104076.
- [45] M. J. S. Yang, Nucl. Phys. B **982**, 115893 (2022), arXiv:2204.08607.
- [46] Z.-z. Xing, H. Zhang, and S. Zhou, Phys. Rev. D77, 113016 (2008), arXiv:0712.1419.
- [47] M. C. Gonzalez-Garcia, M. Maltoni, and T. Schwetz, Universe 7, 459 (2021), arXiv:2111.03086.
- [48] M. J. S. Yang, (2024), arXiv:2402.02767.
- [49] KamLAND-Zen, S. Abe et al., Phys. Rev. Lett. 130, 051801 (2023), arXiv:2203.02139.
- [50] L. Covi, E. Roulet, and F. Vissani, Phys. Lett. B 384, 169 (1996), arXiv:hep-ph/9605319.
- [51] S. Blanchet and P. Di Bari, JCAP 03, 018 (2007), arXiv:hep-ph/0607330.