Type-II Seesaw Leptogenesis along the Ridge

Chengcheng Han^{1,2}, Zhanhong Lei¹, Jin Min Yang^{3,4}

¹School of Physics, Sun Yat-Sen University, Guangzhou 510275, P. R. China
²Asia Pacific Center for Theoretical Physics, Pohang 37673, Korea
³CAS Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, P. R. China
⁴Department of Physics, Henan Normal University, Xinxiang 453007, P. R. China
E-mail: hanchch@mail.sysu.edu.cn, leizhh3@mail2.sysu.edu.cn, jmyang@itp.ac.cn

ABSTRACT: Type-II seesaw leptogenesis is a model that integrates inflation, baryon number asymmetry, and neutrino mass simultaneously. It employs the Affleck-Dine mechanism to generate lepton asymmetry, with the Higgs bosons serving as the inflaton. Previous studies assumed inflation to occur in a valley of the potential, employing the single-field approximation. In this work, we explore an alternative scenario for the type-II seesaw leptogenesis, where the inflation takes place along a ridge of the potential. Firstly, we conduct a comprehensive numerical calculation in the canonical scenario, where inflation occurs in a valley, confirming the effectiveness of the single-field approximation. Then, we introduce a novel scenario wherein inflation initiates along the potential's ridge and transitions to the valley in the late stages. In this case, the single-field inflation approximation is no longer valid, yet leptogenesis is still successfully achieved. We find that this scenario can generate a significant non-Gaussianity signature, offering testable predictions for future experiments.

Contents

1	Introduction	2	
2	 Type-II Seesaw and Inflation 2.1 Type-II seesaw 2.2 Inflation from Higgs 2.3 The potential and inflation trajectory 	3 3 4 5	
3	The Affleck-Dine Leptogenesis3.1Along the valley3.2Along the ridge	6 8 9	
4	Non-Gaussianity	12	
5	Conclusions	14	
A	cknowledgments	15	
References			

1 Introduction

Many open problems remain to be solved in particle physics and cosmology, including the three typical ones: the origin of neutrino masses, the matter-antimatter asymmetry, and the nature of inflation. To solve these problems, various models have been proposed, among which the seesaw mechanisms [1-10] are rather popular since they can interpret the tiny neutrino masses and the baryon asymmetry through the leptogenesis [11]. There are also attempts to simultaneously solve these three problems, usually through considering some extensions to the seesaw models [12-37].

Interestingly, recent studies [38, 39] have shown that the minimal Type-II seesaw may address the above three problems simultaneously. In this framework, the inflaton is provided by the mixing of the SM Higgs H with the triplet scalar Δ in the Type-II seesaw model. The nonminimal coupling of these two scalars with gravity induces a flat direction, along which the slow-roll inflation can be achieved. Considering the Affleck-Dine mechanism [40], as the inflaton evolves along the inflationary trajectory, the phase of Δ carrying lepton number also evolves to give rise to a net lepton number density. Then, after the inflation ends and the universe thermalizes, the asymmetry of lepton number is converted into the asymmetry of baryon number through the sphaleron process [41]. Finally, after spontaneous electroweak symmetry breaking, the neutral component of Δ obtains a nonzero vev, which is responsible for the neutrino masses.

In [38, 39] the authors considered the large field approximation and the inflation can be effectively described as a single field. In this work we conduct the calculation numerically and show that under certain conditions, the inflaton indeed evolve along a single flat direction, which is the valley of the potential. Even if the initial values of the fields are chosen arbitrarily, the inflaton will fall quickly into the valley and the subsequent evolution is essentially analogous to single-field slow-roll inflation, known as the attractor solution. In addition, we observe that there exist other cases where the evolution of inflaton deviates from the standard single-field slow-roll inflation. Particularly, if there exist a local minimum of the potential at the top of the ridge, the inflaton initially on the ridge will evolve along the ridge for a period, and then roll off and evolve along the valley until the inflation ends. Our calculations show that as long as the inflaton rolls off the ridge late enough, then the produced net lepton number density will not be diluted too much by the subsequent inflation and the baryon asymmetry of the universe can be interpreted. Moreover, the deviation from the single-field approximation may give rise to large primordial non-Gaussianity, which is an important feature to distinguish inflation models. Using the δN formalism, we calculate the non-Gaussianity and find that the successful Type-II seesaw leptogenesis along the ridge can give rise to a sizable $|f_{NL}|$, which can be tested by the upcoming CMB measurements. Other phenomenology studies on the type-II seesaw leptogenesis can be found in [42-44].

This paper is organized as follows. In Sec. 2 we briefly review the inflation dynamics under the Type-II seesaw framework. In Sec. 3 we analyse the Affleck-Dine mechanism along the inflationary trajectory. We conduct the calculation in the cases where the inflaton evolves along the valley and along the ridge respectively, and show that successful leptogenesis can be achieved in both cases. After that, we calculate the non-Gaussianity produced in the second case in Sec. 4. Finally, in Sec. 5 we give a summary and draw conclusions.

2 Type-II Seesaw and Inflation

In this section, we briefly overview the inflation dynamics in type-II seesaw model. The nonminimal couplings of the triplet Higgs in type-II seesaw model and the SM Higgs to gravity induces a flat potential for large fields and induce a Starobinsky type inflation in the early universe [45–58]. The multifield inflation dynamics is analysed in the Einstein frame, under which the inflation trajectory can be deduced from the equation of motion of the inflaton.

2.1 Type-II seesaw

The type-II seesaw mechanism attempts to explain neutrino masses through introducing an $SU(2)_L$ triplet scalar Δ to the SM, which carries a hypercharge Y = 1. The triplet and SM Higgs are parameterized as

$$H = \begin{pmatrix} h^+ \\ h \end{pmatrix}, \quad \Delta = \begin{pmatrix} \Delta^+ / \sqrt{2} & \Delta^{++} \\ \Delta^0 & -\Delta^+ / \sqrt{2} \end{pmatrix}, \quad (2.1)$$

where h and Δ^0 are the neutral components. The triplet Δ induces a gauge invariant, renormalizable Yukawa term,

$$\mathcal{L}_{\text{Yukawa}} = \mathcal{L}_{\text{Yukawa}}^{\text{SM}} - \frac{1}{2} y_{jk} \overline{L}_{j}^{c} i \sigma^{2} \Delta L_{k} + \text{ h.c.} , \qquad (2.2)$$

where L_i represents a SM left-handed lepton doublet. Thus we can assign a lepton charge of $Q_L = -2$ to Δ . After electroweak symmetry breaking, this term generates a neutrino mass matrix $m_{ij}^{\nu} = y_{ij}v_{\Delta}$ when Δ^0 obtains a non-zero vev. Diagonalizing m_{ij}^{ν} by the PMNS matrix gives the Majorana neutrino masses.

The field Δ also induces new terms to the scalar potential, given by

$$V(H,\Delta) = -m_H^2 H^{\dagger} H + m_{\Delta}^2 \text{Tr}(\Delta^{\dagger} \Delta) + \lambda_H (H^{\dagger} H)^2 + \lambda_1 (H^{\dagger} H) \text{Tr}(\Delta^{\dagger} \Delta) + \lambda_2 \left(\text{Tr}(\Delta^{\dagger} \Delta) \right)^2 + \lambda_3 \text{Tr}(\Delta^{\dagger} \Delta)^2 + \lambda_4 H^{\dagger} \Delta \Delta^{\dagger} H + \left[\mu (H^T i \sigma^2 \Delta^{\dagger} H) \right] + \frac{\lambda_5}{M_P} (H^T i \sigma^2 \Delta^{\dagger} H) (H^{\dagger} H) + \frac{\lambda_5'}{M_P} (H^T i \sigma^2 \Delta^{\dagger} H) (\Delta^{\dagger} \Delta) + h.c. \right] .$$
(2.3)

Here the terms in the bracket lead to lepton number violation, which is necessary for successful leptogenesis. Two dimension-5 operators are also included, which are suppressed at low energy but would be important during the early universe.

2.2 Inflation from Higgs

The CMB observation indicates that the inflaton should evolve along a flat potential. As is known in the scenario of Higgs inflation, the nonminimal coupling of Higgs scalars to the Ricci scalar R induces a flat direction in the large field limit. In our framework, the Lagrangian in the Jordan frame with nonminimal coupling is given by

$$\frac{\mathcal{L}}{\sqrt{-g}} = \frac{1}{2} M_P^2 R + \xi_H H^{\dagger} H R + \xi_\Delta \operatorname{Tr}(\Delta^{\dagger} \Delta) R - g^{\mu\nu} (D_{\mu} H)^{\dagger} (D_{\nu} H) - g^{\mu\nu} \operatorname{Tr}(D_{\mu} \Delta)^{\dagger} (D_{\nu} \Delta) - V(H, \Delta) + \mathcal{L}_{Yukawa} .$$
(2.4)

The following analysis will focus on the neutral components of the scalars which have nonzero vev, thus the cosmological relevant Lagrangian is given by

$$\frac{\mathcal{L}}{\sqrt{-g}} = \frac{1}{2}M_P^2 R + \xi_H |h|^2 R + \xi_\Delta |\Delta^0|^2 R - (\partial_\mu h)^2 - (\partial_\mu \Delta^0)^2 - V(h, \Delta^0) = \frac{1}{2}(M_P^2 + \xi_H \rho_H^2 + \xi_\Delta \rho_\Delta^2) R - \frac{1}{2}(\partial_\mu \rho_H)^2 - \frac{1}{2}(\partial_\mu \rho_\Delta)^2 - V(h, \Delta^0)$$
(2.5)

and the scalar potential can be simplified as

$$V(h, \Delta^{0}) = -m_{H}^{2}|h|^{2} + m_{\Delta}^{2}|\Delta^{0}|^{2} + \lambda_{H}|h|^{4} + \lambda_{\Delta}|\Delta^{0}|^{4} + \lambda_{H\Delta}|h|^{2}|\Delta^{0}|^{2} - \left(\mu h^{2}\Delta^{0^{*}} + \frac{\lambda_{5}}{M_{P}}|h|^{2}h^{2}\Delta^{0^{*}} + \frac{\lambda_{5}'}{M_{P}}|\Delta^{0}|^{2}h^{2}\Delta^{0^{*}} + h.c.\right) + \dots, \qquad (2.6)$$

where $\lambda_{\Delta} = \lambda_2 + \lambda_3$ and $\lambda_{H\Delta} = \lambda_1 + \lambda_4$. We have introduced the polar coordinate parameterizations

$$h \equiv \frac{1}{\sqrt{2}} \rho_H e^{i\eta}, \quad \Delta^0 \equiv \frac{1}{\sqrt{2}} \rho_\Delta e^{i\theta}.$$
 (2.7)

Since the potential is dominated by the radial directions ρ_H and ρ_{Δ} , the angular motion of η and θ can be ignored in the analysis of inflationary dynamics.

After a Weyl transformation,

$$g_{E\mu\nu} = \Omega^2 g_{\mu\nu}, \quad \Omega^2 = 1 + \frac{\xi_H \rho_H^2}{M_P^2} + \frac{\xi_\Delta \rho_\Delta^2}{M_P^2} ,$$
 (2.8)

we obtain the Lagrangian in the Einstein frame,

$$\frac{\mathcal{L}}{\sqrt{-g_E}} = \frac{1}{2} M_P^2 R_E - \frac{1}{2\Omega^2} \left((\partial_\mu \rho_H)^2 + (\partial_\mu \rho_\Delta)^2 \right) - 3M_P^2 (\partial_\mu \log \Omega)^2 - \frac{V(h, \Delta^0)}{\Omega^4} .$$
(2.9)

In the following we analyse the inflation dynamics by considering the action of multifield inflation. We denote $\phi^I = (\rho_H, \rho_{\Delta})$, then from Eq. 2.9, the action in the Einstein frame is

$$S = \int d^4x \sqrt{-g_E} \times \left[\frac{M_P^2}{2}R - \frac{1}{2}G_{IJ}g_E^{\mu\nu}\partial_\mu\phi^I\partial_\nu\phi^J - V_E(\phi)\right] , \qquad (2.10)$$

where $V_E(\phi) = V(h, \Delta^0)/\Omega^4$ and G_{IJ} is the field metric given by

$$G_{IJ} = \frac{M_P^2}{2f(\phi^I)} \left[\delta_{IJ} + \frac{3}{f(\phi^I)} f_{,I} f_{,J} \right] , \qquad (2.11)$$

with $f(\phi^I) = \frac{1}{2}(M_P^2 + \xi_H \rho_H^2 + \xi_\Delta \rho_\Delta^2)$ being the nonminimal coupling function and $f_{,I} = \partial f / \partial \phi^I$.

Varying the inflation action with respect to $g_{E\mu\nu}$ and ϕ^I , and focusing on the background part of inflaton, $\phi^I = \varphi^I + \delta \phi^I$, we obtain the Friedmann equations and the Klein-Gordon equation,

$$H^{2} = \frac{1}{3M_{P}^{2}} \left[\frac{1}{2} G_{IJ} \dot{\varphi}^{I} \dot{\varphi}^{J} + V \right] ,$$

$$\dot{H} = -\frac{1}{2M_{P}^{2}} G_{IJ} \dot{\varphi}^{I} \dot{\varphi}^{J} ,$$

(2.12)

$$\mathcal{D}_t \dot{\varphi}^I + 3H \dot{\varphi}^I + G^{IK} V_{,K} = 0 , \qquad (2.13)$$

where $\mathcal{D}_t A^I \equiv \dot{\varphi}^J \mathcal{D}_J A^I$, and the covariant derivative in the field space is $\mathcal{D}_J A^I = \partial_J A^I + \Gamma^I_{JK} A^K$. By numerically solving Eq. 2.13, we can obtain the evolution of the background trajectory.

In the case of multifield inflation, the evolution of the inflaton may be complicated, which, however, can be drastically simplified in the so-called kinematical basis [59–61]. We can define the speed of the field vector as

$$\dot{\sigma} = \left| \dot{\varphi}^I \right| = \sqrt{G_{IJ} \dot{\varphi}^I \dot{\varphi}^J} , \qquad (2.14)$$

then the Friedmann equations simplify to

$$H^{2} = \frac{1}{3M_{P}^{2}} \left[\frac{1}{2} \dot{\sigma}^{2} + V \right], \quad \dot{H} = -\frac{1}{2M_{P}^{2}} \dot{\sigma}^{2} .$$
 (2.15)

It can be seen that $\dot{\sigma}$ captures the evolution characteristics along the field trajectory, which is similar to the single-field case. Thus the slow-roll parameters can be defined as in single-field inflation,

$$\begin{aligned} \epsilon &\equiv -\frac{\dot{H}}{H^2} = \frac{3\dot{\sigma}^2}{(\dot{\sigma}^2 + 2V)} ,\\ \eta_{\sigma\sigma} &\equiv M_P^2 \frac{\mathcal{M}_{\sigma\sigma}}{V} , \end{aligned}$$
(2.16)

where $\mathcal{M}_{\sigma\sigma} = \hat{\sigma}^K \hat{\sigma}^J (\mathcal{D}_K \mathcal{D}_J V)$, and $\hat{\sigma}^I = \dot{\varphi}^I / \dot{\sigma}$.

The deviation from single-field inflation is captured by the turn rate $\omega^I = \mathcal{D}_t \hat{\sigma}^I$, which represents how quickly the field trajectory is changing direction. If the so-called 'slow-turn limit' $|\omega^I| \ll 1$ is satisfied, then the inflation dynamics is effectively single-field inflation.

2.3 The potential and inflation trajectory

In the following we adopt the natural unit $M_P = 1$ with M_P being the Planck scale. Considering $\xi_H \rho_H^2 + \xi_\Delta \rho_\Delta^2 \gg 1$ during inflation, the kinetic Lagrangian of the scalar fields is

$$\mathcal{L}_{kin} = -\frac{3}{4} \left(\partial_{\mu} \log(\xi_H \rho_H^2 + \xi_\Delta \rho_\Delta^2) \right)^2 - \frac{1}{2(\xi_H \rho_H^2 + \xi_\Delta \rho_\Delta^2)} \left((\partial_{\mu} \rho_H)^2 + (\partial_{\mu} \rho_\Delta)^2 \right) , \quad (2.17)$$

which is noncanonical. Thus we redefine the fields as

$$\chi = \sqrt{\frac{3}{2}} \log(\xi_H \rho_H^2 + \xi_\Delta \rho_\Delta^2), \quad \kappa = \frac{\rho_H}{\rho_\Delta} . \tag{2.18}$$

For large nonminimal couplings ξ_H , $\xi_{\Delta} \gg 1$, the mixing term $(\partial_{\mu}\chi)(\partial^{\mu}\kappa)$ is suppressed, and the kinetic Lagrangian reduces to

$$\mathcal{L}_{kin} = -\frac{1}{2} (\partial_{\mu} \chi)^2 - \frac{1}{2} \frac{\xi_H^2 \kappa^2 + \xi_{\Delta}^2}{(\xi_H \kappa^2 + \xi_{\Delta})^3} (\partial_{\mu} \kappa)^2 , \qquad (2.19)$$

in which χ is canonically normalised. A canonically normalized field κ' can also be defined in different regimes for κ .

Taking the large field limit, the quartic terms in the potential dominate, and the scalar potential in the Einstein frame reduces to

$$U(\rho_H, \rho_\Delta) = \frac{V}{\Omega^4} = \frac{\lambda_H \kappa^4 + \lambda_{H\Delta} \kappa^2 + \lambda_\Delta}{4(\xi_H \kappa^2 + \xi_\Delta)^2} .$$
(2.20)

It can be shown that the mass of κ' is of order $1/\sqrt{\xi}$, always larger than the Hubble parameter during inflation which is of order $1/\xi$. Therefore, the field κ can be integrated out, and the inflation dynamics is determined by χ .

As pointed by [55], the potential $U(\rho_H, \rho_\Delta)$ has the following minima,

(1)
$$2\lambda_H \xi_\Delta - \lambda_{H\Delta} \xi_H > 0$$
, $2\lambda_\Delta \xi_H - \lambda_{H\Delta} \xi_\Delta > 0$, $\kappa = \sqrt{\frac{2\lambda_\Delta \xi_H - \lambda_{H\Delta} \xi_\Delta}{2\lambda_H \xi_\Delta - \lambda_{H\Delta} \xi_H}}$, (2.21)

(2)
$$2\lambda_H \xi_\Delta - \lambda_{H\Delta} \xi_H > 0, \ 2\lambda_\Delta \xi_H - \lambda_{H\Delta} \xi_\Delta < 0, \ \kappa = 0$$
, (2.22)

(3)
$$2\lambda_H\xi_\Delta - \lambda_{H\Delta}\xi_H < 0, \ 2\lambda_\Delta\xi_H - \lambda_{H\Delta}\xi_\Delta > 0, \ \kappa = \infty$$
, (2.23)

(4)
$$2\lambda_H \xi_\Delta - \lambda_{H\Delta} \xi_H < 0, \ 2\lambda_\Delta \xi_H - \lambda_{H\Delta} \xi_\Delta < 0, \ \kappa = 0, \infty$$
 (2.24)

As shown in Fig. 1, in case (1) the minimum is in the direction of the mixing of h and Δ^0 , while in other cases the valley extends in the direction of h or Δ^0 . If the inflaton evolves along the valley, as will be discussed in Sec. 3.1, the inflation dynamics is essentially equivalent to the single-field inflation. In this situation, successful leptogenesis can only occur in case (1). However, as will be discussed in Sec. 3.2, if a local minimum exhibits in the ridge in case (2), then successful leptogenesis can also be achieved. Moreover, in this case the single-field inflation approximation is invalid, and some interesting features of multifield inflation arise.

3 The Affleck-Dine Leptogenesis

In the Affleck-Dine mechanism, the baryon asymmetry is generated through a nonzero angular motion of phases of the scalar fields carrying $U(1)_B$ charge. In our model the lepton asymmetry is generated analogously by the angular rotation of the phase of Δ^0 when it evolves along the inflation trajectory. Thus, to analyse leptogenesis, we should keep track of the evolution



Figure 1. The potential $U(\rho_H, \rho_{\Delta})$ with different minima.

of the phases by solving the Klein-Gordon equation Eq. 2.13 with $\varphi^{I} = (\rho_{H}, \eta, \rho_{\Delta}, \theta)$. The generated lepton number density during this process can be calculated through the Noether current,

$$n_L = j_L^0 = \frac{Q_L \rho_\Delta^2 \dot{\theta}}{1 + \xi_H \rho_H^2 + \xi_\Delta \rho_\Delta^2}.$$
 (3.1)

After reheating [47, 62–70], the generated lepton number density will be present as SM particles. Then, through the equilibrium electroweak sphaleron process before electroweak symmetry breaking, part of lepton asymmetry is converted into baryon asymmetry, $n_B = -\frac{28}{79}n_L$ [41, 71–73]. As pointed by [38, 39], a lepton number density $n_L \sim 10^{-16}M_P^{-3}$ at the end of the inflation could just explain the baryon asymmetry of our universe.

To have non-trivial motion of θ during inflation, we need the presence of $U(1)_L$ breaking term during inflation. This condition requires the motion of the inflation to occur along a path where both h and Δ^0 are non-vanishing. One obvious choice is that if the conditions in case (1) are satisfied, the inflation will follow a direction with fixed κ and the $U(1)_L$ breaking condition is satisfied. This scenario is pointed out by [38, 39] and we denote it as the "canonical scenario" of type-II seesaw leptogenesis. Note that in the previous studies [38, 39], the estimation of the lepton asymmetry is calculated via the single-field approximation. Instead, in this work we adopt a full calculation without any approximation, and we will show that the single-field approximation indeed work very well, supporting the result of [38, 39]. For the other three cases, it seems that the inflation will occur in a direction where either h or Δ^0 is nearly zero and thus the motion of θ is strongly suppressed because the $U(1)_L$ breaking term is too small. However, we will show that there exist another scenario that the lepton asymmetry can be generated if the inflaton evolves along the ridge. In this scenario, the single field approximation breaks down and a sizable non-Gaussianity could be generated.

3.1 Along the valley

As discussed in the preceding section, the field κ can be effectively integrated out, and the minima of the potential extend in some specific directions with a fixed κ . Then the inflaton evolves along the valley and a single-field approximation can be used during the inflation. In order to achieve successful leptogenesis, the valley should extend in the direction of the mixing of h^0 and Δ^0 , as in case (1). In this work, instead of using the single-field approximation, we present a full numerical calculation to solve the evolution of the fields.

Table 1. A benchmark point in the parameter region satisfying case (1).

ξ_H	ξ_{Δ}	λ_H	λ_Δ	$\lambda_{H\Delta}$	μ	λ_5	λ_5'
300	300	4×10^{-5}	6×10^{-5}	4×10^{-5}	0	0	-10^{-9}

In Table. 1 we show one benchmark point in the parameter space satisfying the conditions in case (1). With the initial conditions chosen as $\rho_H = 0.4, \rho_\Delta = 0.4, \eta = 0, \theta = 0.1, \dot{\rho}_H = \dot{\rho}_\Delta = \dot{\eta} = \dot{\theta} = 0$, the evolution of the inflaton is solved and depicted in Fig. 2. We have parameterized the evolution with respect to $H_0 t$, with H_0 defined as the Hubble constant when the inflation starts. The inflation ends at $H_0 t = 74.44$, leading to the number of efoldings of expansion N = 70.22.

The slow-roll parameters and the turn rate are depicted in Fig. 3. It is obvious that, after a short period of rolling down to the valley, the slow-roll parameters and the turn rate quickly approach the slow-roll slow-turn limit ϵ , $|\eta_{\sigma\sigma}|$, $|\omega^{I}| \ll 1$. Then, as the inflaton evolves along the valley, the single-field approximation is valid and the inflation is effectively standard single-field inflation.

The spectral index of the power spectrum at the horizon crossing $N_{\star} = N_{end} - 60$ is also calculated, leading to $n_s(N_{\star}) = 0.967$ which is in excellent agreement with current CMB



Figure 2. The evolution of the inflaton along the valley.



Figure 3. The slow-roll parameters when the inflaton evolves along the valley.

observations.

Since the evolution of inflaton is solved, the lepton number generated during inflation can be determined, which is depicted in Fig. 4. We see that at this benchmark point an adequate lepton asymmetry is generated. By simply setting $\lambda'_5 = -10^{-12}$, we can obtain $|n_L(t_{end})| \sim 10^{-16} M_P^{-3}$, which is required to explain the baryon asymmetry observed today.

3.2 Along the ridge

As demonstrated above, typically the inflaton rapidly descends to the valley, evolving as a single field. However, exceptions exist. If a local minimum exists along the ridge and the inflaton initially resides there, it may traverse the ridge before ultimately transitioning to the valley. Should this transition occur in the late stages, the CMB data remains explicable, and successful leptogenesis is attainable. In such cases, the single-field approximation becomes



Figure 4. The generated lepton number density during inflation along the valley.

invalid.

Here we take the case (2) as an example, where the ridge extends in the direction of ρ_H . A local minimum on the ridge requires the potential

$$\frac{\partial^2 U}{\partial \rho_{\Delta}^2}\Big|_{\rho_{\Delta}=0} = \frac{\rho_H^2}{2(1+\xi_H \rho_H^2)^3} \left[\lambda_{H\Delta} + (\lambda_{H\Delta}\xi_H - 2\lambda_H\xi_{\Delta})\rho_H^2\right] > 0.$$
(3.2)

Table 2. A benchmark point in the parameter region satisfying case (2) and Eq. 3.2.

ξ_H	ξ_{Δ}	λ_H	λ_{Δ}	$\lambda_{H\Delta}$	μ	λ_5	λ_5'
100	100.02	5×10^{-6}	3.5×10^{-6}	1×10^{-5}	0	0	-10^{-10}

In Table 2 we present a benchmark point in the parameter region satisfying the conditions in case (2) and Eq. 3.2. With the initial condition chosen as $\rho_H = 1$, $\rho_{\Delta} = 8.7 \times 10^{-5}$, $\eta = 0$, $\theta = 0.1$, $\dot{\rho}_H = \dot{\rho}_{\Delta} = \dot{\eta} = \dot{\theta} = 0$, the evolution of the inflaton is solved and depicted in Fig. 5. The inflation ends at $H_0 t = 75.95$, leading to the number of efoldings of expansion N = 73.31. The spectral index in this case, $n_s(N_{\star}) = 0.967$, is also in excellent agreement with current CMB observations.

The slow-roll parameters and the turn rate are depicted in Fig. 6. One can observe that when the inflaton evolves along the ridge, the slow-roll slow-turn limit ϵ , $|\eta_{\sigma\sigma}|$, $|\omega^I| \ll 1$ is satisfied. However, when the inflaton rolls down the ridge, the deviation from the single-field inflation leads to something interesting. As depicted in Fig. 7, when the inflaton rolls off, the rapid turn in the field space leads to a rapid motion of θ , and thus a large amount of lepton number is generated at this time. To match the observed baryon asymmetry today, we can simply set $\lambda'_5 = -3 \times 10^{-13}$ in this case.

For the aforementioned benchmark point, we deliberately selected a very small value for ρ_{Δ} to ensure that the inflaton remains on the ridge for an extended duration. However,



Figure 5. The evolution of the inflaton along the ridge.



Figure 6. The slow-roll parameters when the inflaton evolves along the valley.

varying this parameter can significantly alter the trajectory of inflation. In Fig. 8, we depict the inflationary paths for different initial conditions of ρ_{Δ} , specifically 4×10^{-5} , 8.7×10^{-5} , and 2×10^{-4} . The results indicate that deviating the initial ρ_{Δ} significantly from the local minima causes the inflaton to descend to the valley much earlier. Subsequently, it follows the valley where $\rho_h = 0$, resulting in a substantial suppression of lepton number generation. Simultaneously, the previously generated lepton asymmetry is diluted during the late-stage inflation. This combined effect leads to an exceedingly small lepton asymmetry in such instances. The right panel of Fig. 8 illustrates this, with the purple line indicating a rapid decrease in the generated lepton number during the late stage of inflation.

Conversely, choosing the initial ρ_{Δ} too close to the local minima results in a continuous stay at this point without any turning. In such cases, the generated lepton asymmetry remains consistently small, as depicted by the red line on the right panel of Fig. 8. In this case, a larger λ_5 seems needed to get the correct value of baryon asymmetry in our universe.



Figure 7. The generated lepton number density during inflation along the ridge.



Figure 8. The type-II seesaw leptogenesis along the ridge with different initial conditions. The red, blue, and purple lines correspond to initial field value $\rho_{\Delta} = 4 \times 10^{-5}$, 8.7×10^{-5} , 2×10^{-4} respectively.

4 Non-Gaussianity

In multifield inflation, an important prediction is the primordial non-Gaussianity of cosmological perturbations [60, 61, 74–92]. The non-Gaussianity leads to nonzero three-point or higher order correlation functions. In the following, we calculate the primordial bispectra produced by the inflation processes discussed above.

In order to handle field perturbations in the models with nontrivial field-space manifolds, we adopt the covariant formalism [93–95]. The field perturbation $\delta \phi^I = \phi^I - \varphi^I$ does not transform covariantly. However, after parametrizing the geodesic connecting φ^I and ϕ^I by λ such that $\phi^I(\lambda = 0) = \varphi^I$ and $\phi^I(\lambda = 1) = \phi^I$, $\delta \phi^I$ can be expanded as

$$\delta\phi^{I} = \mathcal{Q}^{I} - \frac{1}{2}\Gamma^{I}_{JK}\mathcal{Q}^{J}\mathcal{Q}^{K} + \frac{1}{6}\left(\Gamma^{I}_{LM}\Gamma^{M}_{JK} - \Gamma^{I}_{JK;L}\right)\mathcal{Q}^{J}\mathcal{Q}^{K}\mathcal{Q}^{L} + \cdots, \qquad (4.1)$$

where $Q^I = \frac{d\phi^I}{d\lambda}\Big|_{\lambda=0}$ is the tangent vector. We also adopt the gauge invariant Mukhanov-Sasaki variable $Q^I = Q^I + \psi \frac{\dot{\varphi}^I}{H}$, where ψ is the scalar component of spatial metric perturbation. In the spatially flat gauge up to first order, $Q^I = Q^I = \delta \phi^I$.

The quantity of interest is the bispectrum of the curvature perturbation ζ , parametrized as

$$\langle \zeta(\mathbf{k}_1)\zeta(\mathbf{k}_2)\zeta(\mathbf{k}_3)\rangle = S(k_1, k_2, k_3) \frac{1}{(k_1 k_2 k_3)^2} \mathcal{P}_{\zeta}^2(2\pi)^7 \delta^3(\mathbf{k}_1 + \mathbf{k}_2 + \mathbf{k}_3),$$
(4.2)

where \mathcal{P}_{ζ} is the power spectrum of ζ and the $S(k_1, k_2, k_3)$ is the shape function.

Adopting the δN formalism [96–98], ζ on super-Hubble scales can be expanded as a function of the field perturbations on the initial flat hypersurface,

$$\zeta(x) = N_{,I}Q^{I}(x) + \frac{1}{2}N_{,IJ}Q^{I}(x)Q^{J}(x) + \cdots, \qquad (4.3)$$

where $N_{,I} = \mathcal{D}_I N|_{\varphi_{\star}^I}$, $N_{,IJ} = \mathcal{D}_I \mathcal{D}_J N|_{\varphi_{\star}^I, \varphi_{\star}^J}$ and N is the e-fold number between the initial flat hypersurface and the final uniform energy density hypersurface. Then the power spectrum and the bispectrum can be expanded as

$$\langle \zeta \left(\boldsymbol{k}_{1} \right) \zeta \left(\boldsymbol{k}_{2} \right) \rangle = (2\pi)^{5} \frac{\mathcal{P}_{\zeta}}{2k_{1}^{3}} \delta^{3} \left(\boldsymbol{k}_{1} + \boldsymbol{k}_{2} \right), \quad \mathcal{P}_{\zeta} = \left(\frac{H_{\star}}{2\pi} \right)^{2} N_{,I} N^{,I}, \tag{4.4}$$

$$\langle \zeta(\mathbf{k}_1)\zeta(\mathbf{k}_2)\zeta(\mathbf{k}_3)\rangle = N_{,I}N_{,J}N_{,K}\langle Q^I(\mathbf{k}_1)Q^J(\mathbf{k}_2)Q^K(\mathbf{k}_3)\rangle_* + \frac{1}{2}N_{,IJ}N_{,K}N_{,L}\int \frac{d^3q}{(2\pi)^3}\langle Q^I(\mathbf{k}_1 - \mathbf{q})Q^K(\mathbf{k}_2)\rangle_*\langle Q^J(\mathbf{q})Q^L(\mathbf{k}_3)\rangle_*$$
(4.5)
 + cyclic permutations.

We can see that there are two contributions to the bispectrum of ζ . One results from the linear transfer of the intrinsic bispectra of the field perturbations at horizon crossing, corresponding to the first term at the right side. The other results from the non-linear relation between the Gaussian field perturbations ζ , corresponding to the other terms. The in-in formalism is needed to calculate $\langle Q^{I}(\mathbf{k}_{1})Q^{J}(\mathbf{k}_{2})Q^{K}(\mathbf{k}_{3})\rangle_{\star}$. Luckily, it was shown that the first contribution remains considerably smaller than the latter contribution for the family of models of interest [94], thus we can reasonably assume that at horizon crossing, the perturbation is Gaussian. Then the bispectrum of ζ reduces to

$$\left\langle \zeta\left(\mathbf{k}_{1}\right)\zeta\left(\mathbf{k}_{2}\right)\zeta\left(\mathbf{k}_{3}\right)\right\rangle = N_{,IJ}N^{,I}N^{,J}\frac{H_{*}^{4}}{4}\left(\frac{1}{k_{1}^{3}k_{2}^{3}} + \frac{1}{k_{2}^{3}k_{3}^{3}} + \frac{1}{k_{3}^{3}k_{1}^{3}}\right)(2\pi)^{3}\delta^{3}\left(\mathbf{k}_{1} + \mathbf{k}_{2} + \mathbf{k}_{3}\right).$$
(4.6)

According to Eq. 4.2, the shape function is

$$S(k_1, k_2, k_3) = \frac{1}{4} \frac{N_{,IJ} N^{,I} N^{,J}}{(N_{,K} N^{,K})^2} \left(\frac{k_1^2}{k_2 k_3} + \text{cyclic permutations}\right).$$
(4.7)

Shape function of this type is called local shape, which peaks at the squeezed triangle limit. Conventionally, the amplitude of bispectrum is denoted as f_{NL} by matching to the shape function,

$$S(k_1, k_2, k_3) \xrightarrow[\text{limit}]{k_1 = k_2 = k_3} \frac{9}{10} f_{NL}.$$
 (4.8)

In this case, f_{NL} is given by

$$f_{NL} = -\frac{5}{6} \frac{N^{,I} N^{,J} \mathcal{D}_I \mathcal{D}_J N}{\left(N_{,K} N^{,K}\right)^2}.$$
(4.9)

In general multifield models, it is difficult to find the analytical form of δN formalism, thus we use finite difference method to perform the calculation numerically. For example, $N_{,\rho_H}$ can be obtained by

$$N_{,\rho_H} = \frac{N(\rho_H + \Delta \rho_H, \rho_\Delta) - N(\rho_H - \Delta \rho_H, \rho_\Delta)}{2\Delta \rho_H},\tag{4.10}$$

where $N(\rho_H, \rho_{\Delta})$ is the number of e-folds from horizon-crossing to the end of inflation, and ρ_H, ρ_{Δ} are the initial values at horizon crossing. We give a perturbation $\Delta \phi^I$ to the initial value, and numerically solve the classical field equation of motion to obtain $N(\phi^I + \Delta \phi^I)$. Then, after tuning $\Delta \rho_H$ or $\Delta \rho_{\Delta}$ to sufficiently small, the numerical result can converge, as depicted in Fig. 9.



Figure 9. The non-Gaussianity produced in the inflation processes along the valley and the ridge.

We see that when the inflaton evolves along the valley, the generated non-Gaussianity is negligible. However, when the inflaton evolves along the ridge, the generated non-Gaussianity can be as large as $|f_{NL}| \sim 2.1$ for fiducial k_{\star} which first crossed the horizon $N_{\star} = 60$ e-folds before inflation ends. Given current limit f_{NL} is -0.9 ± 5.1 for local shape [99], such a large non-Gaussianity may be observed and tested in the upcoming CMB experiments.

5 Conclusions

Type-II seesaw leptogenesis is a model that simultaneously explains inflation, baryon number asymmetry, and neutrino mass, employing the Affleck-Dine mechanism to generate lepton asymmetry and using the Higgs bosons as the inflaton. Previous studies assumed inflation to occur in a valley of the potential, taking the single-field approximation. In this work, we explored an alternative scenario for such type-II seesaw leptogenesis, where the inflation takes place along a ridge of the potential. Firstly we conducted a comprehensive numerical calculation in the canonical scenario, where the inflation occurs in a valley, confirming the effectiveness of the single-field approximation. We also introduced a novel scenario wherein the inflation initiates along the potential's ridge and transitions to the valley in the late stages. During the transition, we find the generation of the lepton asymmetry can be enhanced. In this case, the single-field inflation approximation is no longer valid, yet leptogenesis is still successfully achieved. Furthermore, we demonstrated that this scenario can generate a significant non-Gaussianity signature, offering testable predictions for future experiments.

Acknowledgements

C. H. is supported by the Guangzhou Basic and Applied Basic Research Foundation under Grant No. 202102020885, the Sun Yat-Sen University Science Foundation and the Fundamental Research Funds for the Central Universities under Grant No. 22qntd3007. This work was also supported by the National Natural Science Foundation of China (NSFC) under grant Nos. 11821505, 12075300 and 1233500, and by Peng-Huan-Wu Theoretical Physics Innovation Center (12047503).

References

- [1] P. Minkowski, $\mu \to e\gamma$ at a Rate of One Out of 10⁹ Muon Decays?, Phys. Lett. B 67 (1977) 421.
- [2] T. Yanagida, Horizontal gauge symmetry and masses of neutrinos, Conf. Proc. C 7902131 (1979) 95.
- [3] S. L. Glashow, The Future of Elementary Particle Physics, NATO Sci. Ser. B 61 (1980) 687.
- [4] M. Gell-Mann, P. Ramond and R. Slansky, Complex Spinors and Unified Theories, Conf. Proc. C 790927 (1979) 315 [1306.4669].
- [5] M. Magg and C. Wetterich, Neutrino Mass Problem and Gauge Hierarchy, Phys. Lett. B 94 (1980) 61.
- [6] T. P. Cheng and L.-F. Li, Neutrino Masses, Mixings and Oscillations in SU(2) x U(1) Models of Electroweak Interactions, Phys. Rev. D 22 (1980) 2860.
- [7] G. Lazarides, Q. Shafi and C. Wetterich, Proton Lifetime and Fermion Masses in an SO(10) Model, Nucl. Phys. B 181 (1981) 287.
- [8] R. N. Mohapatra and G. Senjanovic, Neutrino Masses and Mixings in Gauge Models with Spontaneous Parity Violation, Phys. Rev. D 23 (1981) 165.
- [9] R. Foot, H. Lew, X. G. He and G. C. Joshi, Seesaw Neutrino Masses Induced by a Triplet of Leptons, Z. Phys. C 44 (1989) 441.
- [10] C. H. Albright and S. M. Barr, Leptogenesis in the type III seesaw mechanism, Phys. Rev. D 69 (2004) 073010 [hep-ph/0312224].

- [11] M. Fukugita and T. Yanagida, Baryogenesis Without Grand Unification, Phys. Lett. B 174 (1986) 45.
- [12] H. Murayama, H. Suzuki, T. Yanagida and J. Yokoyama, *Chaotic inflation and baryogenesis by right-handed sneutrinos*, *Phys. Rev. Lett.* **70** (1993) 1912.
- [13] H. Murayama, H. Suzuki, T. Yanagida and J. Yokoyama, *Chaotic inflation and baryogenesis in supergravity*, *Phys. Rev. D* 50 (1994) R2356 [hep-ph/9311326].
- [14] M. Senami and K. Yamamoto, Affleck-Dine leptogenesis with triplet Higgs, Phys. Lett. B 524 (2002) 332 [hep-ph/0105054].
- [15] J. L. Evans, T. Gherghetta and M. Peloso, Affleck-Dine Sneutrino Inflation, Phys. Rev. D 92 (2015) 021303 [1501.06560].
- [16] R. N. Mohapatra and N. Okada, Unified model for inflation, pseudo-Goldstone dark matter, neutrino mass, and baryogenesis, Phys. Rev. D 105 (2022) 035024 [2112.02069].
- [17] R. N. Mohapatra and N. Okada, Affleck-Dine Leptogenesis with One Loop Neutrino Mass and strong CP, 2207.10619.
- [18] M. P. Hertzberg and J. Karouby, Generating the Observed Baryon Asymmetry from the Inflaton Field, Phys. Rev. D 89 (2014) 063523 [1309.0010].
- [19] K. D. Lozanov and M. A. Amin, End of inflation, oscillons, and matter-antimatter asymmetry, Phys. Rev. D 90 (2014) 083528 [1408.1811].
- [20] M. Yamada, Affleck-Dine baryogenesis just after inflation, Phys. Rev. D 93 (2016) 083516 [1511.05974].
- [21] K. Bamba, N. D. Barrie, A. Sugamoto, T. Takeuchi and K. Yamashita, Ratchet baryogenesis and an analogy with the forced pendulum, Mod. Phys. Lett. A33 (2018) 1850097 [1610.03268].
- [22] K. Bamba, N. D. Barrie, A. Sugamoto, T. Takeuchi and K. Yamashita, *Pendulum Leptogenesis*, *Phys. Lett. B* 785 (2018) 184 [1805.04826].
- [23] J. M. Cline, M. Puel and T. Toma, Affleck-Dine inflation, Phys. Rev. D 101 (2020) 043014 [1909.12300].
- [24] N. D. Barrie, A. Sugamoto, T. Takeuchi and K. Yamashita, *Higgs Inflation, Vacuum Stability*, and Leptogenesis, JHEP 08 (2020) 072 [2001.07032].
- [25] C.-M. Lin and K. Kohri, Inflaton as the Affleck-Dine Baryogenesis Field in Hilltop Supernatural Inflation, Phys. Rev. D 102 (2020) 043511 [2003.13963].
- [26] M. Kawasaki and S. Ueda, Affleck-Dine inflation in supergravity, JCAP 04 (2021) 049 [2011.10397].
- [27] A. Kusenko, L. Pearce and L. Yang, Postinflationary Higgs relaxation and the origin of matter-antimatter asymmetry, Phys. Rev. Lett. 114 (2015) 061302 [1410.0722].
- [28] Y.-P. Wu, L. Yang and A. Kusenko, Leptogenesis from spontaneous symmetry breaking during inflation, JHEP 12 (2019) 088 [1905.10537].
- [29] Y.-Y. Charng, D.-S. Lee, C. N. Leung and K.-W. Ng, Affleck-Dine Baryogenesis, Split Supersymmetry, and Inflation, Phys. Rev. D 80 (2009) 063519 [0802.1328].

- [30] J. G. Ferreira, C. A. de S. Pires, J. G. Rodrigues and P. S. Rodrigues da Silva, Inflation scenario driven by a low energy physics inflaton, Phys. Rev. D 96 (2017) 103504 [1707.01049].
- [31] E. Babichev, D. Gorbunov and S. Ramazanov, Affleck–Dine baryogenesis via mass splitting, Phys. Lett. B **792** (2019) 228 [1809.08108].
- [32] J. G. Rodrigues, M. Benetti, M. Campista and J. Alcaniz, Probing the Seesaw Mechanism with Cosmological data, JCAP 07 (2020) 007 [2002.05154].
- [33] S. M. Lee, K.-y. Oda and S. C. Park, Spontaneous Leptogenesis in Higgs Inflation, JHEP 03 (2021) 083 [2010.07563].
- [34] S. Enomoto, C. Cai, Z.-H. Yu and H.-H. Zhang, Leptogenesis due to oscillating Higgs field, Eur. Phys. J. C 80 (2020) 1098 [2005.08037].
- [35] A. Lloyd-Stubbs and J. McDonald, A Minimal Approach to Baryogenesis via Affleck-Dine and Inflaton Mass Terms, Phys. Rev. D 103 (2021) 123514 [2008.04339].
- [36] R. N. Mohapatra and N. Okada, Affleck-Dine baryogenesis with observable neutron-antineutron oscillation, Phys. Rev. D 104 (2021) 055030 [2107.01514].
- [37] R. N. Mohapatra and N. Okada, Neutrino mass from Affleck-Dine leptogenesis and WIMP dark matter, JHEP 03 (2022) 092 [2201.06151].
- [38] N. D. Barrie, C. Han and H. Murayama, Affleck-Dine Leptogenesis from Higgs Inflation, Phys. Rev. Lett. 128 (2022) 141801 [2106.03381].
- [39] N. D. Barrie, C. Han and H. Murayama, *Type II Seesaw leptogenesis*, *JHEP* 05 (2022) 160 [2204.08202].
- [40] I. Affleck and M. Dine, A New Mechanism for Baryogenesis, Nucl. Phys. B 249 (1985) 361.
- [41] V. A. Kuzmin, V. A. Rubakov and M. E. Shaposhnikov, On the Anomalous Electroweak Baryon Number Nonconservation in the Early Universe, Phys. Lett. B 155 (1985) 36.
- [42] C. Han, Z. Lei and W. Liao, Testing type II seesaw leptogenesis at the LHC*, Chin. Phys. C 47 (2023) 093104 [2303.15709].
- [43] C. Han, S. Huang and Z. Lei, Vacuum stability of the type II seesaw leptogenesis from inflation, Phys. Rev. D 107 (2023) 015021 [2208.11336].
- [44] N. D. Barrie and S. T. Petcov, Lepton Flavour Violation tests of Type II Seesaw Leptogenesis, JHEP 01 (2023) 001 [2210.02110].
- [45] A. A. Starobinsky, A New Type of Isotropic Cosmological Models Without Singularity, Phys. Lett. B 91 (1980) 99.
- [46] F. L. Bezrukov and M. Shaposhnikov, The Standard Model Higgs boson as the inflaton, Phys. Lett. B659 (2008) 703 [0710.3755].
- [47] F. Bezrukov, D. Gorbunov and M. Shaposhnikov, On initial conditions for the Hot Big Bang, JCAP 0906 (2009) 029 [0812.3622].
- [48] J. Garcia-Bellido, D. G. Figueroa and J. Rubio, Preheating in the Standard Model with the Higgs-Inflaton coupled to gravity, Phys. Rev. D 79 (2009) 063531 [0812.4624].
- [49] J. L. F. Barbon and J. R. Espinosa, On the Naturalness of Higgs Inflation, Phys. Rev. D79 (2009) 081302 [0903.0355].

- [50] A. O. Barvinsky, A. Yu. Kamenshchik, C. Kiefer, A. A. Starobinsky and C. Steinwachs, Asymptotic freedom in inflationary cosmology with a non-minimally coupled Higgs field, JCAP 0912 (2009) 003 [0904.1698].
- [51] F. Bezrukov and M. Shaposhnikov, Standard Model Higgs boson mass from inflation: Two loop analysis, JHEP 07 (2009) 089 [0904.1537].
- [52] G. F. Giudice and H. M. Lee, Unitarizing Higgs Inflation, Phys. Lett. B 694 (2011) 294 [1010.1417].
- [53] F. Bezrukov, A. Magnin, M. Shaposhnikov and S. Sibiryakov, *Higgs inflation: consistency and generalisations*, *JHEP* 01 (2011) 016 [1008.5157].
- [54] C. P. Burgess, H. M. Lee and M. Trott, Comment on Higgs Inflation and Naturalness, JHEP 07 (2010) 007 [1002.2730].
- [55] O. Lebedev and H. M. Lee, *Higgs Portal Inflation, Eur. Phys. J. C* **71** (2011) 1821 [1105.2284].
- [56] Y. Hamada, H. Kawai, K.-y. Oda and S. C. Park, Higgs Inflation is Still Alive after the Results from BICEP2, Phys. Rev. Lett. 112 (2014) 241301 [1403.5043].
- [57] H. M. Lee, Light inflaton completing Higgs inflation, Phys. Rev. D 98 (2018) 015020 [1802.06174].
- [58] S.-M. Choi, Y.-J. Kang, H. M. Lee and K. Yamashita, Unitary inflaton as decaying dark matter, JHEP 05 (2019) 060 [1902.03781].
- [59] C. Gordon, D. Wands, B. A. Bassett and R. Maartens, Adiabatic and entropy perturbations from inflation, Phys. Rev. D 63 (2000) 023506 [astro-ph/0009131].
- [60] C. M. Peterson and M. Tegmark, Testing Two-Field Inflation, Phys. Rev. D 83 (2011) 023522 [1005.4056].
- [61] C. M. Peterson and M. Tegmark, Testing multifield inflation: A geometric approach, Phys. Rev. D 87 (2013) 103507 [1111.0927].
- [62] J. Garcia-Bellido, D. G. Figueroa and J. Rubio, Preheating in the Standard Model with the Higgs-Inflaton coupled to gravity, Phys. Rev. D 79 (2009) 063531 [0812.4624].
- [63] Y. Ema, R. Jinno, K. Mukaida and K. Nakayama, Violent Preheating in Inflation with Nonminimal Coupling, JCAP 02 (2017) 045 [1609.05209].
- [64] M. P. DeCross, D. I. Kaiser, A. Prabhu, C. Prescod-Weinstein and E. I. Sfakianakis, Preheating after Multifield Inflation with Nonminimal Couplings, I: Covariant Formalism and Attractor Behavior, Phys. Rev. D 97 (2018) 023526 [1510.08553].
- [65] M. P. DeCross, D. I. Kaiser, A. Prabhu, C. Prescod-Weinstein and E. I. Sfakianakis, Preheating after multifield inflation with nonminimal couplings, III: Dynamical spacetime results, Phys. Rev. D 97 (2018) 023528 [1610.08916].
- [66] M. P. DeCross, D. I. Kaiser, A. Prabhu, C. Prescod-Weinstein and E. I. Sfakianakis, Preheating after multifield inflation with nonminimal couplings, II: Resonance Structure, Phys. Rev. D 97 (2018) 023527 [1610.08868].
- [67] M. He, R. Jinno, K. Kamada, S. C. Park, A. A. Starobinsky and J. Yokoyama, On the violent preheating in the mixed Higgs-R² inflationary model, Phys. Lett. B 791 (2019) 36 [1812.10099].

- [68] M. He, R. Jinno, K. Kamada, A. A. Starobinsky and J. Yokoyama, Occurrence of tachyonic preheating in the mixed Higgs-R² model, JCAP 01 (2021) 066 [2007.10369].
- [69] M. He, Perturbative Reheating in the Mixed Higgs-R² Model, JCAP 05 (2021) 021 [2010.11717].
- [70] E. I. Sfakianakis and J. van de Vis, Preheating after Higgs Inflation: Self-Resonance and Gauge boson production, Phys. Rev. D 99 (2019) 083519 [1810.01304].
- [71] F. R. Klinkhamer and N. S. Manton, A Saddle Point Solution in the Weinberg-Salam Theory, Phys. Rev. D30 (1984) 2212.
- [72] M. Trodden, Electroweak baryogenesis, Rev. Mod. Phys. 71 (1999) 1463 [hep-ph/9803479].
- [73] A. Sugamoto, The neutrino mass and the monopole-Anti-monopole dumb-bell system in the SO(10) grand unified model, Phys. Lett. 127B (1983) 75.
- [74] K. A. Malik and D. Wands, Cosmological perturbations, Phys. Rept. 475 (2009) 1 [0809.4944].
- [75] D. Wands, Multiple field inflation, Lect. Notes Phys. 738 (2008) 275 [astro-ph/0702187].
- [76] N. Bartolo, E. Komatsu, S. Matarrese and A. Riotto, Non-Gaussianity from inflation: Theory and observations, Phys. Rept. 402 (2004) 103 [astro-ph/0406398].
- [77] X. Chen, Primordial Non-Gaussianities from Inflation Models, Adv. Astron. 2010 (2010) 638979 [1002.1416].
- [78] C. T. Byrnes and K.-Y. Choi, Review of local non-Gaussianity from multi-field inflation, Adv. Astron. 2010 (2010) 724525 [1002.3110].
- [79] E. Komatsu et al., Non-Gaussianity as a Probe of the Physics of the Primordial Universe and the Astrophysics of the Low Redshift Universe, 0902.4759.
- [80] T. Qiu and K.-C. Yang, Non-Gaussianities of Single Field Inflation with Non-minimal Coupling, Phys. Rev. D 83 (2011) 084022 [1012.1697].
- [81] F. Bernardeau and J.-P. Uzan, NonGaussianity in multifield inflation, Phys. Rev. D 66 (2002) 103506 [hep-ph/0207295].
- [82] F. Bernardeau and J.-P. Uzan, Inflationary models inducing non-Gaussian metric fluctuations, Phys. Rev. D 67 (2003) 121301 [astro-ph/0209330].
- [83] D. Langlois and S. Renaux-Petel, Perturbations in generalized multi-field inflation, JCAP 04 (2008) 017 [0801.1085].
- [84] D. Seery and J. E. Lidsey, Primordial non-Gaussianities from multiple-field inflation, JCAP 09 (2005) 011 [astro-ph/0506056].
- [85] S. Yokoyama, T. Suyama and T. Tanaka, Primordial Non-Gaussianity in Multi-Scalar Slow-Roll Inflation, JCAP 07 (2007) 013 [0705.3178].
- [86] S. Yokoyama, T. Suyama and T. Tanaka, Primordial Non-Gaussianity in Multi-Scalar Inflation, Phys. Rev. D 77 (2008) 083511 [0711.2920].
- [87] C. T. Byrnes, K.-Y. Choi and L. M. H. Hall, Conditions for large non-Gaussianity in two-field slow-roll inflation, JCAP 10 (2008) 008 [0807.1101].
- [88] J. Elliston, D. J. Mulryne, D. Seery and R. Tavakol, Evolution of fNL to the adiabatic limit, JCAP 11 (2011) 005 [1106.2153].

- [89] D. Seery, D. J. Mulryne, J. Frazer and R. H. Ribeiro, Inflationary perturbation theory is geometrical optics in phase space, JCAP 09 (2012) 010 [1203.2635].
- [90] C. M. Peterson and M. Tegmark, Non-Gaussianity in Two-Field Inflation, Phys. Rev. D 84 (2011) 023520 [1011.6675].
- [91] J.-O. Gong and H. M. Lee, Large non-Gaussianity in non-minimal inflation, JCAP 11 (2011) 040 [1105.0073].
- [92] A. Mazumdar and L.-F. Wang, Separable and non-separable multi-field inflation and large non-Gaussianity, JCAP 09 (2012) 005 [1203.3558].
- [93] J.-O. Gong and T. Tanaka, A covariant approach to general field space metric in multi-field inflation, JCAP 03 (2011) 015 [1101.4809].
- [94] D. I. Kaiser, E. A. Mazenc and E. I. Sfakianakis, Primordial Bispectrum from Multifield Inflation with Nonminimal Couplings, Phys. Rev. D 87 (2013) 064004 [1210.7487].
- [95] J. Elliston, D. Seery and R. Tavakol, The inflationary bispectrum with curved field-space, Journal of Cosmology and Astroparticle Physics 2012 (2012) 060.
- [96] M. Sasaki and E. D. Stewart, A General analytic formula for the spectral index of the density perturbations produced during inflation, Prog. Theor. Phys. 95 (1996) 71 [astro-ph/9507001].
- [97] D. H. Lyth, K. A. Malik and M. Sasaki, A General proof of the conservation of the curvature perturbation, JCAP 05 (2005) 004 [astro-ph/0411220].
- [98] H.-C. Lee, M. Sasaki, E. D. Stewart, T. Tanaka and S. Yokoyama, A New delta N formalism for multi-component inflation, JCAP 10 (2005) 004 [astro-ph/0506262].
- [99] PLANCK collaboration, Y. Akrami et al., Planck 2018 results. IX. Constraints on primordial non-Gaussianity, Astron. Astrophys. 641 (2020) A9 [1905.05697].