

LIE THEORETIC APPROACH TO UNITARY GROUPS OF C^* -ALGEBRAS

HIROSHI ANDO AND MICHAL DOUCHA

ABSTRACT. Following Robert's [23], we study the structure of unitary groups and groups of approximately inner automorphisms of unital C^* -algebras, taking advantage of the former being Banach-Lie groups. For a given unital C^* -algebra A , we provide a description of the closed normal subgroup structure of the connected component of the identity of the unitary group, denoted by U_A , resp. of the subgroup of approximately inner automorphisms induced by the connected component of the identity of the unitary group, denoted by V_A , in terms of perfect ideals, i.e. ideals admitting no characters. When the unital algebra is locally AF, we show that there is a one-to-one correspondence between closed normal subgroups of V_A and perfect ideals of the algebra, which can be in the separable case conveniently described using Bratteli diagrams; in particular showing that every closed normal subgroup of V_A is perfect. We also characterize unital C^* -algebras A such that U_A , resp. V_A are topologically simple, generalizing the main results from [23]. In the other way round, under certain conditions, we characterize simplicity of the algebra in terms of the structure of the unitary group. This in particular applies to reduced group C^* -algebras of discrete groups and we show that when A is a reduced group C^* -algebra of a non-amenable countable discrete group, then A is simple if and only if U_A/\mathbb{T} is topologically simple.

1. INTRODUCTION

Unitary groups of unital C^* -algebras and von Neumann algebras are objects that have been studied from several different perspectives in various subjects such as operator algebras, infinite dimensional differential geometry, mathematical physics, and topological group theory. Even within the C^* -algebra theory, the interest is both in the intrinsic structure of these groups, such as the normal subgroup structure when the corresponding algebra is simple, or in their relation to areas such as K-theory (see e.g. the monograph [24]) and exponential rank and length (see [21] and references therein). It was rarely used though that when equipped with the norm topology these groups are linear Banach-Lie groups, i.e. very well behaved, in general, infinite-dimensional Lie groups for which the powerful machinery of Lie theoretic techniques is available. A rare innovative exception, from which also our inspiration comes, was the paper [23] of Robert which shows that the closed commutator subgroup modulo its center of the connected component of the unit of a unitary group of any simple unital C^* -algebra is topologically simple using Lie theoretic methods. Robert also showed that the group of approximately inner automorphisms induced by unitaries from the connected component of the unit of any simple unital C^* -algebra is topologically simple resolving a long standing open problem from [11]. For another recent work related to unitary groups of C^* -algebras from the Lie perspective, we refer e.g. to [7, 12, 2].

We remark that Robert himself was inspired by a result [6] on Lie ideals in C^* -algebras, another area where Lie theory meets C^* -algebras and which has been already extensively studied (see e.g. [19, 18, 16, 22, 13]).

It is the goal of this paper to continue in this line of research and study the connected component of the unit of unitary groups, and their closely related cousins: the groups of approximately inner automorphisms induced by unitaries from such components, using Lie group techniques.

2020 *Mathematics Subject Classification.* 22E65, 22F50, 46L05.

Key words and phrases. unitary groups, C^* -algebras, approximately inner automorphisms, Banach-Lie groups, C^* -simplicity.

H. Ando was supported by Japan Society for the Promotion of Sciences (20K03647). M. Doucha was supported by the GAČR project 22-07833K and by the Czech Academy of Sciences (RVO 67985840).

We refine Robert's description of normal subgroup structure of these groups from [23] and relate them to *perfect ideals*, i.e. closed two-sided ideals admitting no characters. For a unital C^* -algebra A , denote by U_A the connected component of the unit of the unitary group of A (equipped with the norm topology) and by V_A the closure, in $\text{Aut}(A)$, of the inner automorphisms coming from U_A .

Theorem A. *Let A be a unital C^* -algebra. There is a one-to-one correspondence between perfect ideals of A and perfect closed normal subgroups of U_A and V_A , and an 'almost' one-to-one correspondence between perfect ideals and general closed normal subgroups of U_A and V_A .*

The vague wording 'almost one-to-one' will be made precise in Theorems 4.5 and 5.2 where these results are proved.

For some C^* -algebras A we obtain precise description of the closed normal subgroups of V_A .

Theorem B. *Let A be a unital locally AF algebra. Then there is a one-to-one correspondence between perfect ideals of A and closed normal subgroups of V_A . Moreover, when A is separable, as a consequence, there is a one-to-one correspondence between closed normal subgroups of V_A and directed and hereditary subsets of a Bratteli diagram of A that do not contain an infinite path of 1s.*

Denoting by $V(A)$ the group of all approximately inner automorphisms of a unital C^* -algebra A , we have the normal series $V_A \trianglelefteq V(A) \trianglelefteq \text{Aut}(A)$ and the structure of the quotients $\text{Aut}(A)/V(A)$, resp. $V(A)/V_A$ can be understood in some cases (see e.g. [11]), we emphasize that it is indeed important to understand the structure of V_A when one is interested in $\text{Aut}(A)$.

The rest of the paper concerns the following problems. First we note that the converse to Robert's theorem, i.e. that when the algebra A is simple, then the closed commutator subgroup of U_A modulo the center and V_A are topologically simple, does not hold. We provide a characterization of those C^* -algebras for which these groups are topologically simple. Then we turn to the issue whether the simplicity of the algebra can be detected from the corresponding unitary group structure. We provide several partial results in this direction that in particular apply to the reduced group C^* -algebras of discrete groups, and among other things we prove the following.

Theorem C. *Let G be a countable discrete non-amenable group. Then $A := C_r^*(G)$ is simple if and only if U_A/\mathbb{T} is topologically simple.*

The paper is organized as follows. The following Section 2 introduces all the necessary notions and notations. Section 3 contains first preliminary results that are used in the sequel and Sections 4 and 5 concern the normal subgroup structure of U_A and V_A respectively. In particular, Theorems A and B are proved there. Section 6 contains some additional information about the Lie structure of certain normal subgroups of U_A which in particular shows that subalgebras that are V_A -invariant are also $V(A)$ -invariant. Section 7 focuses on the simplicity of C^* -algebras and Theorem C is proved there.

2. PRELIMINARIES AND NOTATION

In this short section, we recall basic facts from (infinite-dimensional) Lie theory and C^* -algebras. We also introduce here all the main notation that we shall work with throughout the whole paper.

2.1. Infinite-dimensional Lie groups. A *Banach-Lie group* is a group that is also a Banach manifold, i.e. an infinite dimensional smooth manifold modelled on a Banach space, and such that the group operations are smooth. In this paper we solely work with a much more concrete and tractable subclass of Banach-Lie groups that we define below.

Definition 2.1. Let A be a unital Banach algebra and denote by $GL(A)$ the group of invertible elements of A equipped with the norm topology. A closed subgroup $G \subseteq GL(A)$ is a *linear Lie subgroup* if the map $\exp : A \rightarrow GL(A)$, $a \rightarrow \sum_{k=0}^{\infty} \frac{a^k}{k!}$, is a local homeomorphism around 0 between $\mathfrak{L}(G) := \{a \in A \mid \exp(ta) \in G \forall t \in \mathbb{R}\}$ and G .

It follows from [15, Definition 5.32 and Corollary 5.34] that G is then a Banach-Lie group modelled on $\mathfrak{L}(G)$, which is a Banach-Lie algebra of G . That is, a Banach subspace of A closed under the bracket operation

$$[x, y] = xy - yx, \quad x, y \in \mathfrak{L}(G).$$

In particular, the following properties hold true for G and $\mathfrak{L}(G)$ that we shall use without further reference in the sequel.

- The connected component G_0 of the unit 1_G of G is also a linear Lie group and $\mathfrak{L}(G_0) = \mathfrak{L}(G)$.
- For each $a \in \mathfrak{L}(G)$, $\exp(a) \in G_0$. Conversely, for each $g \in G$ there are $a_1, \dots, a_n \in \mathfrak{L}(G)$ such that $g = \prod_{i=1}^n \exp(a_i)$.
- (The Trotter product formula)

$$\exp(a + b) = \lim_{n \rightarrow \infty} \left(\exp\left(\frac{1}{n}a\right) \exp\left(\frac{1}{n}b\right) \right)^n,$$

for every $a, b \in \mathfrak{L}(G)$.

- (The commutator formula)

$$\exp([a, b]) = \lim_{n \rightarrow \infty} \left(\exp\left(\frac{1}{n}a\right) \exp\left(\frac{1}{n}b\right) \exp\left(-\frac{1}{n}a\right) \exp\left(-\frac{1}{n}b\right) \right)^{n^2},$$

for every $a, b \in \mathfrak{L}(G)$.

- (The Baker-Campbell-Hausdorff formula)

$$\exp(a) \exp(b) = \exp \left(a + b + \frac{1}{2}[a, b] + \frac{1}{12}[a, [a, b]] - \frac{1}{12}[b, [a, b]] + \dots \right),$$

for all sufficiently small $a, b \in \mathfrak{L}(G)$. In fact, by [4, Theorem 5.56], the right hand side converges whenever $a, b \in \mathfrak{L}(G)$ satisfy $\|a\| + \|b\| < \frac{1}{2} \log 2$.

Notice that in particular, $GL(A)$ is a linear Banach-Lie group and $\mathfrak{L}(GL(A)) = A$.

2.2. Unitary groups of C^* -algebras. Let A be a unital C^* -algebra. By $\mathcal{Z}(A)$ we shall denote the center of A . Similarly, if G is a (topological) group, by $\mathcal{Z}(G)$ we shall denote the center of G . For two subsets $I, J \subseteq A$ we denote by $[I, J]$ the linear span of the commutators $\{xy - yx \mid x \in I, y \in J\}$ in A . Analogously, for subgroups H_1, H_2 of G the subgroup of G generated by commutators $(h_1, h_2) = h_1 h_2 h_1^{-1} h_2^{-1}$, $h_1 \in H_1, h_2 \in H_2$ is denoted by (H_1, H_2) . Denote by $U(A)$ the unitary group, i.e. $U(A) = \{u \in A \mid uu^* = u^*u = 1\} \subseteq GL(A)$. Then $U(A)$ is a linear Lie group and its Lie algebra $\mathfrak{L}(U(A))$ is by definition the set of skew-adjoint elements, i.e. the set $\{a \in A \mid a^* = -a\}$. However, here we follow the standard practice from C^* -algebra theory and mathematical physics and we identify $\mathfrak{L}(U(A))$ with the set $A_{\text{sa}} = \{a \in A \mid a^* = a\}$ of self-adjoint elements. Note that as a result we work with the exponential map $a \in A_{\text{sa}} \rightarrow \exp(ia) \in U(A)$ and the bracket operation $i[x, y]$, for $x, y \in A_{\text{sa}}$. We also emphasize that we shall interchangeably write both $\exp(a)$ and e^a when using the exponential map as the latter is sometimes notationally more convenient when writing down formulas.

For $S \subseteq A$, we denote by $\text{Id}_A(S)$ the closed two-sided ideal of A generated by $S \subseteq A$.

The connected component of the unit in $U(A)$ will be denoted by U_A . It is therefore again a linear Lie subgroup with A_{sa} as a Lie algebra. Moreover, if $I \subseteq A$ is a closed two-sided ideal, we define $U(I) := \{u \in U(A) \mid u - 1 \in I\}$ and then U_I to be the connected component of the unit of $U(A)$. We have that U_I is a closed connected normal linear Lie subgroup of U_A and $\mathfrak{L}(U_I) = I_{\text{sa}} := A_{\text{sa}} \cap I$. This follows immediately from Definition 2.1 and the fact that the exponential map $a \rightarrow e^{ia}$ maps I into $1 + I$ and the logarithm map, whenever defined, maps $1 + I$ into I . The closure of the commutator subgroup (U_I, U_A) (resp. (U_I, U_I)) is denoted by $CU_{I,A}$ (resp. CU_I). These two groups are actually equal and topologically perfect. The latter is proved, for CU_I , in [23, Theorem 3.6], the former is implicit in [23] and explicitly proved in Lemma 3.2.

Denote by $\text{Aut}(A)$ the group of all automorphisms of A equipped with the pointwise convergence topology. For each $u \in U(A)$, $\text{Ad}(u) : A \rightarrow A$, defined by $a \rightarrow uau^*$, is an automorphism called an *inner automorphism*. Clearly, $\{\text{Ad}(u) \mid u \in U(A)\} \subseteq \text{Aut}(A)$ is a subgroup, which is

not closed in general. We denote by $V(A)$ the closure $\overline{\{\text{Ad}(u) \mid u \in U(A)\}}$ in $\text{Aut}(A)$, which is called the group of *approximately inner automorphisms*. We shall also denote by V_A the closure $\overline{\{\text{Ad}(u) \mid u \in U_A\}}$. Notice that we have a normal series $V_A \trianglelefteq V(A) \trianglelefteq \text{Aut}(A)$.

For any closed two-sided ideal $I \subseteq A$, the groups $V(I)$ and V_I are defined analogously.

3. FIRST RESULTS

In this section, we collect and prove several basic results that will become handy in the sequel.

3.1. Ideals and commutator subgroups. The next lemma is proved in [6, Proposition 5.25], by an argument using W^* -algebras and a result from [20]. As is mentioned in the last paragraph to [22, Lemma 1.4], it can be shown using quasi-central approximate units. Here we include such a proof.

Lemma 3.1. *Let A be a unital C^* -algebra and I be a closed two-sided ideal of A . Then $\overline{[A, A]} \cap I = \overline{[I, I]}$ holds.*

Proof. To ease notation for $x, y \in A$ and $\varepsilon > 0$, we denote $x \overset{\varepsilon}{\sim} y$ to mean $\|x - y\| < \varepsilon$. It is clear that $\overline{[I, I]} \subseteq \overline{[A, A]} \cap I$. Let $x \in \overline{[A, A]} \cap I$ and $\varepsilon > 0$. Then there exists $x_0 \in [A, A]$ such that $x \overset{\varepsilon}{\sim} x_0$. Let $(e_\lambda)_{\lambda \in \Lambda}$ be an approximate units for I that is quasicentral in A , meaning that $\lim_\lambda \|ae_\lambda - e_\lambda a\| = 0$ for every $a \in A$ (see e.g., [9, Theorem I.9.16]). We will use for functions $f, g: \Lambda \rightarrow A$ the notation $f(\lambda) \overset{\lambda \rightarrow \infty}{\sim} g(\lambda)$ to mean $\lim_\lambda \|f(\lambda) - g(\lambda)\| = 0$. By $x \in I$, there exists $\lambda_0 \in \Lambda$ such that $e_\lambda x e_\lambda \overset{\varepsilon}{\sim} x$ for every $\lambda \geq \lambda_0$. Because e_λ 's are positive contractions, we also have $e_\lambda x e_\lambda \overset{\varepsilon}{\sim} e_\lambda x_0 e_\lambda$ ($\lambda \geq \lambda_0$). Write

$$x_0 = \sum_{k=1}^n (a_k b_k - b_k a_k), \quad a_k, b_k \in A \quad (1 \leq k \leq n).$$

Because $(e_\lambda)_\lambda$ is quasicentral in A , we have

$$e_\lambda x_0 e_\lambda \overset{\lambda \rightarrow \infty}{\sim} \sum_{k=1}^n (e_\lambda a_k e_\lambda b_k - e_\lambda b_k e_\lambda a_k),$$

so that there exists $\lambda_1 \geq \lambda_0$ such that

$$e_\lambda x_0 e_\lambda \overset{\varepsilon}{\sim} \sum_{k=1}^n [e_\lambda a_k, e_\lambda b_k] =: x_1(\lambda) \in [I, I], \quad (\lambda \geq \lambda_1).$$

Then for $\lambda \geq \lambda_1$,

$$x \overset{\varepsilon}{\sim} e_\lambda x e_\lambda \overset{\varepsilon}{\sim} e_\lambda x_0 e_\lambda \overset{\varepsilon}{\sim} x_1(\lambda) \in [I, I].$$

Since ε is arbitrary, we obtain $x \in \overline{[I, I]}$. □

Lemma 3.2. *Let A be a unital C^* -algebra and I be a closed two-sided ideal of A . Then $CU_I = CU_{I,A}$.*

Proof. The inclusion $CU_I \subseteq CU_{I,A}$ is obvious. To show the converse inclusion, note that by [23, Lemma 3.1], the set $\{e^{ic} \mid c \in [I, A] \cap A_{\text{sa}}\}$ generates $CU_{I,A}$ as a topological group. Therefore it suffices to show that $e^{ic} \in CU_I$ for every $c \in [I, A] \cap A_{\text{sa}}$. But in this case it holds by Lemma 3.1 that $c \in \overline{[A, A]} \cap I_{\text{sa}} = \overline{[I, I]} \cap A_{\text{sa}}$, which by [23, Lemma 3.1] again implies that $e^{ic} \in CU_I$. □

The next lemma is essentially [23, Lemma 3.1] and the same proof works for it.

Lemma 3.3. *Let I be a closed two-sided ideal of a unital C^* -algebra A . Then CU_I is generated by $\{e^{ix} \mid x \in [I, I] \cap A_{\text{sa}}\}$ as a topological group.*

The next lemma improves Robert's lemma above.

Lemma 3.4. *Let A be a unital C^* -algebra, I be a closed two-sided ideal of A . Then the set $E_{I,A} = \{e^{[a,b]} \mid a \in I_{\text{sa}}, b \in A_{\text{sa}}\}$ generates $CU_{I,A}$ as a topological group.*

Proof. Let H be the closed subgroup of U_A generated by $E_{I,A}$. Then H is a normal subgroup of U_A , since $ue^{[a,b]}u^* = e^{[uau^*,ubu^*]}$ with $uau^* \in I_{\text{sa}}$, $ubu^* \in A_{\text{sa}}$ if $a \in I_{\text{sa}}$, $b \in A_{\text{sa}}$ and $u \in U_A$. By [23, Lemma 3.1], the set $E'_{I,A} = \{e^{ix} \mid x \in [I, A] \cap A_{\text{sa}}\}$ generates $CU_{I,A}$ as a topological group. Because $E_{I,A} \subseteq E'_{I,A}$, this implies that $H \subseteq CU_{I,A}$. Thus we only have to show that $CU_{I,A} \subseteq H$. Let $u \in U_I$ and $v \in U_A$, and we show that $(u, v) \in H$. For this purpose, we recall the formula (essentially the same as [23, (2.1)]) that for elements x, y, z in a group G ,

$$(1) \quad (z, xy) = (z, x)x(z, y)x^{-1}.$$

Since for every $r > 0$ the exponentials of skew-adjoint operators of norm at most $r > 0$ in I (resp. A) algebraically generates U_I (resp. U_A), by the formulas [23, (2.1)] and (1), we may assume that u, v are of the form $u = e^{ia}$, $v = e^{ib}$ for $a \in I_{\text{sa}}$ and $b \in A_{\text{sa}}$ such that their norms are so small so that the Campbell–Baker–Hausdorff formula converges.

This implies that

$$\begin{aligned} (u, v) &= e^{ia}e^{-ivav^*} \\ &= \exp \left(i(a - vav^*) - \frac{i}{2}[a, vav^*] + \frac{1}{12}([a, i[a, vav^*] + [vav^*, i[a, vav^*]]) \cdots \right), \end{aligned}$$

Moreover, note that $f(t) = e^{itb}ae^{-itb}$ is a norm-differentiable function in $t \in \mathbb{R}$ with $f'(t) = e^{itb}i[b, a]e^{-itb}$ for every $t \in \mathbb{R}$. Thus we can perform the vector-valued Riemann integration and see that

$$\begin{aligned} i(a - vav^*) &= i(a - f(1)) = ia - i \left(f(0) + \int_0^1 f'(t)dt \right) \\ &= - \int_0^1 \underbrace{[e^{itb}ae^{-itb}, b]}_{\in I_{\text{sa}}} dt. \end{aligned}$$

This in particular implies, by the standard approximation of vector-valued Riemann integrals by vector-valued step functions, that there exists sequences x_n, y_n all of which are finite sums of elements from the set $\{[c, d] \mid c \in I_{\text{sa}}, d \in A_{\text{sa}}\}$, such that $i(a - vav^*) = \lim_{n \rightarrow \infty} x_n$, and that (by the continuity of \exp) $(u, v) = \lim_{n \rightarrow \infty} \exp(x_n + y_n)$. By a repeated application of the Trotter product formula, we see that $\exp(x_n + y_n) \in H$ for every $n \in \mathbb{N}$, whence $(u, v) \in H$ as well. This shows the claim. \square

Proposition 3.5. *Let I be a closed two-sided ideal of a unital C^* -algebra A . Then $\mathcal{Z}(CU_I) = \mathcal{Z}(CU_A) \cap CU_I$ holds.*

Proof. It is clear that $CU_I \cap \mathcal{Z}(CU_A) \subseteq \mathcal{Z}(CU_I)$, so we prove the converse inclusion. Let $g \in \mathcal{Z}(CU_I)$ and we show that g commutes with every $h \in CU_A$. By Lemma 3.4, the set $\{e^{[a,b]} \mid a, b \in A_{\text{sa}}\}$ generates CU_A as a topological group, we may assume that $h = e^{[a,b]}$ where $a, b \in A_{\text{sa}}$. Let $(e_\lambda)_{\lambda \in \Lambda}$ be an approximate unit in I that is quasi-central in A .

In order to prove $gh = hg$, it is sufficient to show that $(g - 1)x = x(g - 1)$, where $x = [a, b]$. Because $g - 1 \in I$ and (e_λ) is an approximate unit for I that is quasiceutral in A , we have (use

also $g \in \mathcal{Z}(CU_I)$

$$\begin{aligned}
(g-1)(ab-ba) &= \lim_{\lambda} (g-1)e_{\lambda}^4(ab-ba) = \lim_{\lambda} (g-1)(e_{\lambda}ae_{\lambda}^2be_{\lambda} - e_{\lambda}be_{\lambda}^2ae_{\lambda}) \\
&= \lim_{\lambda} \frac{d}{dt} \Big|_{t=0} (g-1) \underbrace{\exp(t[e_{\lambda}ae_{\lambda}, e_{\lambda}be_{\lambda}])}_{\in CU_I} \\
&= \lim_{\lambda} \frac{d}{dt} \Big|_{t=0} \exp(t[e_{\lambda}ae_{\lambda}, e_{\lambda}be_{\lambda}]) (g-1) \\
&= \lim_{\lambda} [e_{\lambda}ae_{\lambda}, e_{\lambda}be_{\lambda}] (g-1) \\
&= \lim_{\lambda} (ab-ba)e_{\lambda}^4 (g-1) \\
&= (ab-ba)(g-1).
\end{aligned}$$

Thus $gx = xg$, whence $g \in \mathcal{Z}(CU_A)$ holds. \square

Lemma 3.6. *Let A be a unital C^* -algebra H, K be normal subgroups of U_A . Then $\text{Id}_A([H, K]) = \text{Id}_A((H, K) - 1)$ holds.*

Proof. Let $h \in H, k \in K$. Then

$$hk - kh = (hkh^{-1}k^{-1} - 1)kh,$$

which shows that $[H, K] \subseteq \text{Id}_A((H, K) - 1)$. Therefore $\text{Id}_A([H, K]) \subseteq \text{Id}_A((H, K) - 1)$ holds. From the identity

$$hkh^{-1}k^{-1} - 1 = (hk - kh)h^{-1}k^{-1}$$

we see that $(H, K) - 1 \subseteq \text{Id}_A([H, K])$, whence $\text{Id}_A((H, K) - 1) \subseteq \text{Id}_A([H, K])$. \square

Lemma 3.7. *Let A be a unital C^* -algebra and I be a closed two-sided ideal of A . Then $\text{Id}_A([I, I]) = \text{Id}_A([CU_I, CU_I])$.*

Proof. By Lemma 3.6 and since CU_I is topologically perfect, we have

$$\begin{aligned}
\text{Id}_A([CU_I, CU_I]) &= \text{Id}_A((CU_I, CU_I) - 1) \\
&= \text{Id}_A(\overline{(CU_I, CU_I)} - 1) \\
&= \text{Id}_A(CU_I - 1) = \text{Id}_A((U_I, U_I) - 1) \\
&= \text{Id}_A([U_I, U_I]) = \text{Id}_A([I, I]).
\end{aligned}$$

\square

3.2. Perfect Ideals.

Definition 3.8. Let I be a closed two-sided ideal of a unital C^* -algebra A . We call I *perfect* if $I = \text{Id}_A([I, I])$ holds.

Remark 3.9. Let A be a C^* -algebra, and I be a closed two-sided ideal of A . Then $\text{Id}_A([I, I]) = \text{Id}_I([I, I])$ holds. Indeed, it is clear that $\text{Id}_I([I, I]) \subseteq \text{Id}_A([I, I])$. On the other hand, for all $a, b \in A$ and $x, y \in I$, we have

$$a[x, y]b = \lim_{j \rightarrow \infty} ae_j[x, y]e_jb \in \text{Id}_I([I, I]),$$

where $(e_j)_{j \in \mathbb{N}}$ is an approximate unit for I . Since $\text{Id}_A([I, I])$ is the closed linear span of the elements of this type, we obtain $\text{Id}_A([I, I]) = \text{Id}_I([I, I])$. From this observation it is safe to write $\text{Id}([I, I])$ instead of $\text{Id}_A([I, I])$. We will follow this convention in the sequel.

Proposition 3.10. *Let I be a closed two-sided ideal of a unital C^* -algebra A . Then $I' := \text{Id}([I, I])$ is perfect. Moreover, it is the biggest perfect ideal contained in I .*

Lemma 3.11. *Let A be a (not necessarily unital) C^* -algebra. Then the following two conditions are equivalent.*

- (1) A is perfect in itself: $A = \text{Id}([A, A])$.

(2) *There is no nonzero $*$ -homomorphism (i.e., a character) $A \rightarrow \mathbb{C}$.*

Proof. (1) \implies (2) Any $*$ -homomorphism $\chi: A \rightarrow \mathbb{C}$ must vanish on $[A, A]$, hence on $\text{Id}([A, A]) = A$.

(2) \implies (1) If (1) does not hold, then $I = \text{Id}([A, A])$ is a proper closed two-sided ideal of A , and $B = A/I$ is then a nonzero abelian C^* -algebra. Thus any pure state $\chi: B \rightarrow \mathbb{C}$ is a nonzero $*$ -homomorphism. By composing χ with the quotient map $A \rightarrow B$, we obtain a nonzero $*$ -homomorphism $A \rightarrow \mathbb{C}$, whence (2) fails. \square

Proof of Proposition 3.10. It is clear that the only $*$ -homomorphism $\text{Id}([I, I]) \rightarrow \mathbb{C}$ is the trivial homomorphism, so I' is perfect by Lemma 3.11.

The ‘moreover’ part is clear. \square

Remark 3.12. Even though the set $\{e^{ix} \mid x \in [A, A]_{\text{sa}}\}$ generates CU_A as a topological group, it is not in general true that the Lie algebra of CU_A is $\overline{[A, A]_{\text{sa}}}$. For example, let A be the hyperfinite II_1 factor, and fix $\theta \in [0, 1] \setminus \mathbb{Q}$ and consider unitaries $u, v \in U(A) = U_A$ satisfying $(u, v) = e^{2\pi i \theta} 1$. Then because θ is irrational, CU_A contains \mathbb{T} . If $\overline{[A, A]_{\text{sa}}}$ were the Lie algebra of CU_A , then for each $s \in \mathbb{R}$, the condition $e^{ist} \in CU_A$ ($t \in \mathbb{R}$) implies that $s = i^{-1} \frac{d}{dt} \big|_{t=0} e^{ist} \in \overline{[A, A]_{\text{sa}}}$, whence $\overline{[A, A]_{\text{sa}}} \supset \mathbb{R}$, which is a contradiction (the unique tracial state on A vanishes on $\overline{[A, A]_{\text{sa}}}$, but not on \mathbb{R} . Note that by [8, Proposition 2.7], $A_{\text{sa}} = \overline{[A, A]_{\text{sa}}} + \mathbb{R}1$ holds).

4. NORMAL SUBGROUP STRUCTURE OF U_A

Here the core of the paper starts. In this section we prove the results concerning the normal subgroup structure of U_A , hereby proving half of Theorem A.

Proposition 4.1. *Let H be a closed normal subgroup of U_A , for a unital C^* -algebra A . Then $\overline{(H, H)}$ is a topologically perfect closed normal subgroup of U_A - the biggest topologically perfect subgroup contained in H - and it is of the form CU_I , where $I \subseteq A$ is a closed two-sided ideal.*

Proof. Let H be a closed normal subgroup of U_A . We prove that $\overline{(H, H)}$ is topologically perfect. It will be automatically normal, since it is normal in H and H is normal in U_A , so by [23, Theorem 3.7] it is normal in U_A . It will be also the biggest subgroup with such a property, since every closed topologically perfect subgroup $H' \leq H$ must be contained in $\overline{(H, H)}$, which follows from the equality $H' = \overline{(H', H')}$.

Set $I := \text{Id}([H, A])$. We show

$$\overline{(H, H)} = \overline{(H, U_A)} = CU_I,$$

and this will finish the proof since CU_I is topologically perfect. In fact, it is enough to show that $\overline{(H, U_A)} = CU_I$ since the former will be then a topologically perfect subgroup of H which thus must be contained in $\overline{(H, H)}$, and obviously contains $\overline{(H, H)}$. However, by [23, Theorem 3.3] we have $\overline{(H, U_A)} = \overline{(U_I, U_A)}$ and the latter is equal to CU_I by Lemma 3.2. \square

Lemma 4.2. *Let I be a closed two-sided ideal of a unital C^* -algebra A . Then $CU_I = CU_{I'}$, where $I' = \text{Id}([I, I])$.*

Proof. By Lemma 3.3, the set $\{e^{ix} \mid x \in [I, I] \cap A_{\text{sa}}\}$ generates CU_I as a topological group and if $x \in [I, I] \cap A_{\text{sa}} \subseteq I'$ then clearly $e^{ix} \in U_{I'}$. Therefore we have $CU_I \subseteq U_{I'}$. By Proposition 4.1, $CU_{I'} = \overline{(U_{I'}, U_{I'})}$ is the biggest topologically perfect normal subgroup of $U_{I'}$. On the other hand, CU_I is a topologically perfect normal subgroup of $U_{I'}$, thus $CU_I \subseteq CU_{I'}$. Since $I' \subseteq I$, we have also $CU_{I'} \subseteq CU_I$, and so $CU_I = CU_{I'}$ holds. \square

We will need the next well-known result. We include the proof for completeness.

Lemma 4.3. *Let A be a unital C^* -algebra. Then U_A spans A .*

Proof. Since A_{sa} spans A , it suffices to show that any self-adjoint contraction $x \in A$ is in the linear span of U_A , and note that the standard argument produces the unitary $u(x) = x + i(1 - x^2)^{\frac{1}{2}}$

such that $x = \frac{1}{2}(u(x) + u(x)^*)$, and $u(x)$ is homotopic to 1 by the continuous path $[0, 1] \ni t \mapsto e^{-\frac{\pi i}{2}(1-t)}u(tx) \in U(A)$. \square

Lemma 4.4. *Let I and J be perfect ideals of a unital C^* -algebra A . If $CU_I = CU_J$, then $I = J$.*

Proof. Let I and J be perfect ideals such that $CU_I = CU_J$. We show that $I = J$. Assume by contradiction that $I \not\subseteq J$. By the perfectness of I and $I \not\subseteq J$, we have $[I, I] \not\subseteq J$. Note that since U_I spans \tilde{I} by Lemma 4.3, the set $\{xu - ux \mid x \in I_{\text{sa}}, u \in U_I\}$ spans $[I, I]$. Thus there exists $x \in I_{\text{sa}}$ and $u \in U_I$ such that $ux - xu \in [I, I] \setminus J$, whence $uxu^* - x = u(xu^*) - (xu^*)u \in [I, I] \setminus J$. We may also assume that $\|x\|$ is small enough so that $uxu^* - x$ lies in a neighborhood of 0 in A where \exp is a local homeomorphism. Then by Lemma 3.3, we have $v := \exp(i(uxu^* - x)) \in CU_I$, however $v \notin U_J$, thus $v \notin CU_J$, which is a contradiction. Therefore, $I \subseteq J$ holds, and by symmetry, the same argument shows $J \subseteq I$ as well. \square

Theorem 4.5. *There is a one-to-one bijection between perfect ideals $I \subseteq A$ and topologically perfect closed normal subgroups of U_A of the form $I \rightarrow CU_I$.*

Proof. For every two-sided ideal $I \subseteq A$, CU_I is a topologically perfect closed normal subgroup of U_A .

By Lemma 4.4, the map $I \rightarrow CU_I$ is injective when restricted to perfect ideals. We check that the map $I \rightarrow CU_I$ is onto the set of all topologically perfect closed normal subgroups of U_A . Let H be a topologically perfect closed normal subgroup of U_A . Then $H = \overline{(H, H)}$, where the right hand side is by Proposition 4.1 equal to CU_I for some ideal $I \subseteq A$. But then for $I' := \text{Id}([I, I])$ we have that I' is perfect by Proposition 3.10 and $CU_I = CU_{I'}$ by Lemma 4.2. \square

Definition 4.6. For a closed two-sided ideal I of a unital C^* -algebra A , we define $N_I = \{u \in U_A \mid (u, U_A) \subseteq \tilde{I}\}$.

Lemma 4.7. *Let A be a unital C^* -algebra, I be a closed two-sided ideal of A . Then $N_I = \{u \in U_A \mid (u, U_A) \subseteq CU_I\}$ holds.*

Proof. Denote the set on the right hand side by M_I . Clearly, $M_I \subseteq N_I$. The converse follows immediately from [23, Lemma 3.2] and Lemma 3.2. \square

As an immediate consequence of Lemmas 4.2 and 4.7 we get the following corollary.

Corollary 4.8. *Let A be a unital C^* -algebra, I be a closed two-sided ideal of A . Then $N_I = N_{I'}$ holds.*

Theorem 4.9. *Let A be a unital C^* -algebra. The set of all closed normal subgroups of U_A is equal to the disjoint union*

$$\bigsqcup_{I \subseteq A \text{ perfect ideal}} \{H \leq U_A \mid CU_I \leq H \leq N_I\}.$$

Proof. Let $H \subseteq U_A$ be a closed normal subgroup. By [23, Theorem 3.3] the ideal $I = \text{Id}([H, A])$ satisfies $CU_{I,A} = \overline{(H, U_A)} \subseteq H$. By Lemmas 3.2 and 4.2, we have $CU_{I,A} = CU_I = CU_{I'}$, and I' is a perfect ideal by Proposition 3.10. Moreover, the equality $\overline{(H, U_A)} = CU_I$, together with Lemmas 4.7 and 4.8 implies that $H \subseteq N_I = N_{I'}$.

Conversely, suppose that H is a closed subgroup of U_A such that for some perfect ideal I of A , we have $CU_I \leq H \leq N_I$. We claim that H is normal in U_A . Pick $h \in H$ and $u \in U_A$. Since $H \subseteq N_I$ we have $uhu^*h^* \in CU_I$, thus $uhu^* \in hCU_I \subseteq H$.

It remains to show that the union from the statement is taken of disjoint sets. Suppose that $H \subseteq U_A$ is a closed normal subgroup and there are two perfect ideals I and J such that $CU_I \leq H \leq N_I$ and $CU_J \leq H \leq N_J$. Then since $(N_I, N_I) \subseteq CU_I = \overline{(CU_I, CU_I)}$ by the definition of N_I and since CU_I is topologically perfect, we have $\overline{(N_I, N_I)} = CU_I = \overline{(CU_I, CU_I)}$ and $\overline{(N_J, N_J)} = CU_J = \overline{(CU_J, CU_J)}$. Therefore, $\overline{(H, H)}$ must be equal to both CU_I and CU_J , showing that $CU_I = CU_J$, which implies, by Lemma 4.4, that $I = J$. \square

Robert proves in [23, Corollary 3.8] that when a unital C^* -algebra is simple, then $CU_A/\mathcal{Z}(CU_A)$ is topologically simple. The converse is not true though. Theorem 4.9 however allows us to completely characterize when $CU_A/\mathcal{Z}(CU_A)$ is topologically simple. We provide this characterization in Theorem 5.10 in the next section jointly with the characterization when V_A is topologically simple.

5. NORMAL SUBGROUP STRUCTURE OF V_A

This section is parallel to the previous one. Here we prove the results concerning the normal subgroup structure of V_A , having then proved the second half of Theorem A. We also provide a characterization of those unital C^* -algebras satisfying that $CU_A/\mathcal{Z}(CU_A)$ and V_A are topologically simple.

We start with the following lemma that appears in [1, Lemma 2.1].

Lemma 5.1. *Let A be a unital C^* -algebra. For every $\phi \in V_A$ and every closed two-sided ideal I we have $\phi[I] \subseteq I$, thus ϕ naturally induces $\pi_I \circ \phi =: P_I(\phi) \in V_{A/I}$, where $\pi_I : A \rightarrow A/I$ is the quotient map. The map $P_I : V_A \rightarrow V_{A/I}$ is a continuous homomorphism satisfying $P_I(\text{Ad}(u)) = \text{Ad}(\pi_I(u))$.*

Notice that $V_I \subseteq \ker(P_I)$. The converse is in general not clear.

The following theorem, concerning the normal subgroup structure of V_A , is analogous to Theorem 4.9 that concerns the normal subgroup structure of U_A .

Theorem 5.2. *Let A be a unital C^* -algebra. The set of all closed normal subgroups of V_A is equal to the disjoint union*

$$\bigsqcup_{I \subseteq A \text{ perfect ideal}} \{H \leq V_A \mid V_I \leq H \leq \ker(P_I)\}.$$

We need a few lemmas for the proof. The first one is an immediate corollary to [23, Corollary 3.8 and Lemma 3.9] (it is mentioned in [23, Theorem 3.10]).

Lemma 5.3 ([23]). *Let A be a unital C^* -algebra I be a closed two-sided ideal. Then V_I is topologically perfect and the equality $V_I = \overline{\text{Ad}(CU_I)}$ holds.*

Proof. By [23, Lemma 3.9] and Lemma 3.2, $V_I = \overline{\text{Ad}(CU_{I,A})} = \overline{\text{Ad}(CU_I)}$, and by [23, Theorem 3.6], CU_I is topologically perfect. Thus

$$\overline{(V_I, V_I)} = \overline{(\text{Ad}(CU_I), \text{Ad}(CU_I))} = \overline{\text{Ad}((CU_I, CU_I))} = \overline{\text{Ad}(CU_I)} = V_I.$$

□

Lemma 5.4. *Let A be a unital C^* -algebra. Then V_A has trivial center.*

Proof. Suppose that $\phi \in \mathcal{Z}(V_A)$. Then for each $u \in \mathcal{U}_A$ and $x \in A$,

$$\text{Ad}(u^*) \circ \phi \circ \text{Ad}(u)(x) = \phi(x).$$

The left hand side of the above equality is $u^*\phi(u)\phi(x)\phi(u^*)u = \gamma(u)\phi(x)\gamma(u)^*$, where $\gamma(u) = u^*\phi(u)$. Since ϕ is onto and x is arbitrary, it shows that $\gamma(u)$ is in the center of A . By [23, Lemma 3.11], ϕ is trivial. □

The next lemma is analogous to Lemma 4.4.

Lemma 5.5. *Let I and J be perfect ideals of a unital C^* -algebra A . If $V_I = V_J$, then $I = J$.*

Proof. Set

$$\tilde{I} := \text{Id}(\{\phi(x) - x \mid \phi \in V_I = V_J, x \in A\}) = \text{Id}(\{uxu^* - x \mid \text{Ad}(u) \in V_I = V_J, x \in A\}) = \text{Id}([CU_I, A]),$$

where the first equality between the ideals is straightforward and the second follows from the fact that $\text{Ad}[CU_I]$ is dense in $V_I = V_J$ by Lemma 5.3. Analogously, we set

$$\tilde{J} := \text{Id}(\{\phi(x) - x \mid \phi \in V_I = V_J, x \in A\}) = \text{Id}(\{uxu^* - x \mid \text{Ad}(u) \in V_I = V_J, x \in A\}) = \text{Id}([CU_J, A]).$$

Notice that we have $\tilde{I} = \tilde{J}$, so it suffices to check that $I = \tilde{I}$ and $J = \tilde{J}$. We show the former equality. By Lemma 3.7, we have

$$\tilde{I} = \text{Id}([CU_I, A]) \supset \text{Id}([CU_I, CU_I]) = \text{Id}([I, I]) = I' = I,$$

since I is perfect. On the other hand, for every $u \in U_I$ and $x \in A$, we have $uxu^* - x \in I$, whence $\tilde{I} \subseteq I$. Therefore $I = \tilde{I}$ and $I = J$ holds. \square

Proof of Theorem 5.2. Let H be a closed normal subgroup of V_A . Set

$$I := \text{Id}(\{\phi(x) - x \mid \phi \in H, x \in A\}).$$

By [23, Theorem 3.10], we have $\overline{(H, V_A)} = V_I$. Without loss of generality, we may assume that I is perfect. Indeed, otherwise we replace I by $I' := \text{Id}([I, I])$ which is perfect by Proposition 3.10. Then by Lemma 4.2 we have $CU_I = CU_{I'}$ and so by Lemma 5.3, we get

$$V_{I'} = \overline{\{\text{Ad}(u) \mid u \in CU_{I'}\}} = \overline{\{\text{Ad}(u) \mid u \in CU_I\}} = V_I.$$

Since H is normal and closed we have $\overline{(H, V_A)} \subseteq H$, thus we get $V_I \leq H \leq NV_I$, where

$$NV_I := \{\phi \in V_A \mid (\phi, V_A) \subseteq V_I\}.$$

We now claim that $NV_I = \ker(P_I)$. First, we show that $\ker(P_I) \subseteq NV_I$. By repeating the argument from the paragraph above with the closed normal subgroup $H := \ker(P_I)$ we get that $\ker(P_I) \subseteq NV_J$, where

$$J := \text{Id}(\{\phi(x) - x \mid \phi \in \ker(P_I), x \in A\}).$$

However, $J \subseteq I$, so $\ker(P_I) \subseteq NV_J \subseteq NV_I$. Indeed, for every $\phi \in \ker(P_I)$ and $x \in A$ we have (where $\pi_I : A \rightarrow A/I$ is the quotient map)

$$\pi_I(\phi(x) - x) = P_I(\phi)(\pi_I(x)) - \pi_I(x) = 0,$$

implying that $\phi(x) - x \in I$, which in turn implies that $J \subseteq I$.

Next we prove the other inclusion $NV_I \subseteq \ker(P_I)$. Pick $\phi \in NV_I$ and $u' \in U_{A/I}$, and let $u \in U_A$ be such that $u' = \pi_I(u)$. Since $(\phi, \text{Ad}(u)) \in V_I$ we have that $P_I(\phi)$ and $\text{Ad}(u')$ commute. As $u' \in U_{A/I}$ was arbitrary and $\{\text{Ad}(u) \mid u \in U_{A/I}\}$ is dense in $V_{A/I}$, we get $P_I(\phi) \in Z(V_{A/I})$. By Lemma 5.4, it follows that $P_I(\phi)$ is trivial, in other words, $\phi \in \ker(P_I)$. Therefore, $\ker(P_I) = NV_I$ holds. We thus obtain $V_I \leq H \leq \ker(P_I)$.

Conversely, if I is a perfect ideal and H is a closed subgroup of V_A such that $V_I \leq H \leq \ker(P_I)$, then we claim that H is normal in V_A . Indeed, pick any $\phi \in H$ and $\psi \in V_A$. Then since $\phi \in NV_I$, we have $\psi\phi\psi^{-1}\phi^{-1} \in V_I$, thus $\psi\phi\psi^{-1} \in V_I\phi \subseteq H$, showing that H is normal in V_A .

By Lemma 5.5, we have that if I and J are perfect ideals, and $V_I = V_J$, then $I = J$.

Finally, we show that the union from the statement consists of disjoint sets. This is done similarly as in the proof of Theorem 4.9. Suppose that $H \subseteq V_A$ is a closed normal subgroup and there are perfect ideals I and J such that $V_I \leq H \leq \ker(P_I)$ and $V_J \leq H \leq \ker(P_J)$. Since $\ker(P_I) = NV_I$ and $\ker(P_J) = NV_J$, the quotients $\ker(P_I)/V_I$ and $\ker(P_J)/V_J$ are abelian and we have that $\overline{(H, H)}$ must be equal to both V_I and V_J . Therefore, by the argument from the previous paragraph, $I = J$ holds. \square

Corollary 5.6. *There is a one-to-one bijection between perfect ideals $I \subseteq A$ and perfect closed normal subgroups of V_A of the form $I \rightarrow V_I$.*

Proof. The proof that the map $I \rightarrow V_I$ is injective, when restricted to perfect ideals, is given in the proof of Theorem 5.2. For each perfect ideal I of A , V_I is topologically perfect by Lemma 5.3. Now if $H \leq V_A$ is any topologically perfect closed normal subgroup, then by Theorem 5.2 there is a perfect ideal $I \subseteq A$ such that $V_I \subseteq H \subseteq \ker(P_I) = NV_I$. Since NV_I/V_I is abelian, it follows that $(NV_I, NV_I)/V_I$ is trivial, so $V_I = \overline{(\ker(P_I), \ker(P_I))} = \overline{(V_I, V_I)}$, thus because H is topologically perfect, we have $H = \overline{(H, H)} = V_I$. \square

In view of Theorem 5.2, it is important to know under which conditions the two groups V_I and $\ker(P_I) = NV_I$ coincide, in which case we have a more precise description of all closed normal subgroups of V_A .

A step in this direction is a part of the main result from [1] (see [1, Theorem 2.4]) that in particular answers in positive this question for unital locally AF algebras, and the following is therefore an immediate corollary of Theorem 5.2 and [1]. We note that in [1], U_I , for an ideal I of a unital C^* -algebra A , is defined to be the connected component of the unit of unitary group of the unitization \tilde{I} rather than $\{u \in U_A \mid u - 1 \in I\}$ as in this paper. The reader can readily verify that these two definitions differ only by the presence of \mathbb{T} and in particular induce the same group V_I .

Theorem 5.7. *Let A be a unital locally AF algebra. Then there is a one-to-one correspondence between perfect ideals of A and closed normal subgroups of V_A , which is of the form*

$$I \rightarrow V_I.$$

In particular, every closed normal subgroup of V_A is topologically perfect.

Observe that Theorem 5.7 proves half of Theorem B from Introduction. We now focus on the other half.

Recall that a unital separable AF algebra A which is a direct limit of a sequence $(A_n)_{n \in \mathbb{N}}$ of finite-dimensional algebras admits a convenient description using a Bratteli diagram \mathcal{D} . To fix a notation, vertices of \mathcal{D} are denoted by pairs (p, i) which correspond to i -th direct summand of A_p . We denote by $d(p, i)$ the dimension of this summand. Recall that there may be multiple arrows between vertices (p, i) and $(p+1, j)$ and the number of such arrows corresponds to the multiplicity of the partial embedding between the i -th direct summand of A_p and j -th direct summand of A_{p+1} . Recall, e.g. from [9, III.4], that there is a one-to-one correspondence between ideals of A and directed and hereditary subsets of \mathcal{D} , and that if $\mathcal{S} \subseteq \mathcal{D}$ is a directed and hereditary subset corresponding to an ideal $I \subseteq A$, then $\mathcal{D} \setminus \mathcal{S}$ is a Bratteli diagram of the quotient A/I .

Theorem 5.8. *Let A be a separable unital AF algebra with Bratteli diagram \mathcal{D} . There is a one-to-one correspondence between closed normal subgroups of V_A and directed hereditary subsets of \mathcal{D} that do not contain a maximal infinite path $(p, n_0) \rightarrow (p+1, n_1) \rightarrow (p+2, n_2) \rightarrow \dots$ such that for all $q \geq p$ we have $d(q, n_{q-p}) = 1$.*

Proof. We view A as a direct limit of a sequence $(A_n)_{n \in \mathbb{N}}$ of finite-dimensional algebras. We retain the notation from the previous paragraph. Applying Theorem 5.7, it is equivalent to prove a one-to-one correspondence between perfect ideals of A and directed hereditary subsets of \mathcal{D} that do not contain a maximal infinite path $(p, n_0) \rightarrow (p+1, n_1) \rightarrow (p+2, n_2) \rightarrow \dots$ such that for all $q \geq p$ we have $d(q, n_{q-p}) = 1$. Clearly, the multiplicity of each such partial embedding is necessarily 1.

We claim that a directed and a hereditary subset $\mathcal{S} \subseteq \mathcal{D}$ corresponding to an ideal $I \subseteq A$ defines a perfect ideal if and only if \mathcal{S} does not contain a maximal infinite path $(p, n_0) \rightarrow (p+1, n_1) \rightarrow (p+2, n_2) \rightarrow \dots$ such that for all $q \geq p$ we have $d(q, n_{q-p}) = 1$. Suppose that \mathcal{S} contains such a maximal path $(p, n_0) \rightarrow (p+1, n_1) \rightarrow (p+2, n_2) \rightarrow \dots$ and denote by \mathcal{S}' the vertices of this path. We claim that $\mathcal{S} \setminus \mathcal{S}'$ is directed and hereditary. To see this, notice that for every $(q, n_{q-p}) \in \mathcal{S}'$ there is exactly one arrow (inside \mathcal{D}) from (q, n_{q-p}) (necessarily to $(q+1, n_{q+1-p})$) and at most one arrow to (q, n_{q-p}) . Therefore if $(r, n_{r-p}) \in \mathcal{S} \setminus \mathcal{S}'$ and there is an arrow $(r, n_{r-p}) \rightarrow (r+1, n_{r+1-p})$ within \mathcal{S} , then $(r+1, n_{r+1-p}) \notin \mathcal{S}'$ as otherwise $d(r+1, n_{r+1-p}) = 1$, thus also $d(r, n_{r-p}) = 1$ and so $(r, n_{r-p}) \in \mathcal{S}'$ because of maximality \mathcal{S}' , a contradiction. This verifies directedness. Now if $(r, n_{r-p}) \in \mathcal{S}$ is such that for every arrow $(r, n_{r-p}) \rightarrow (r+1, n_{r+1-p})$ we have $(r+1, n_{r+1-p}) \notin \mathcal{S}'$, then necessarily also $(r, n_{r-p}) \notin \mathcal{S}'$ since otherwise, if $(r, n_{r-p}) \in \mathcal{S}'$, there is only one arrow from (r, n_{r-p}) and by definition it is into a vertex from \mathcal{S}' . This verifies the hereditary property. Therefore $\mathcal{S} \setminus \mathcal{S}'$ defines a subideal $J \subseteq I$ and I/J corresponds to \mathcal{S}' which clearly defines a one-dimensional C^* -algebra, i.e. \mathbb{C} . Therefore, I admits a non-trivial character, so it is not perfect by Lemma 3.11.

Conversely, suppose that I is not perfect. Then by Lemma 3.11, I admits a non-trivial character, so there exists a subideal $J \subseteq I$ such that $I/J = \mathbb{C}$. Let $\mathcal{F} \subseteq \mathcal{S}$ be a directed and hereditary subset that defines J . It follows that the quotient $\mathbb{C} = I/J$ corresponds to the Bratteli subdiagram $\mathcal{S} \setminus \mathcal{F}$. However, $\mathcal{S} \setminus \mathcal{F}$ may correspond to \mathbb{C} only if for every $(p, n) \in \mathcal{S} \setminus \mathcal{F}$ we have $d(p, n) = 1$. Since $\mathcal{S} \setminus \mathcal{F}$ is infinite, by Zorn's lemma we can extract a maximal infinite path $\mathcal{S}' \subseteq \mathcal{S} \setminus \mathcal{F}$. In particular, \mathcal{S} contains a maximal infinite path $(p, n_0) \rightarrow (p+1, n_1) \rightarrow (p+2, n_2) \rightarrow \dots$ such that for all $q \geq p$ we have $d(q, n_{q-p}) = 1$. This finishes the proof of the claim. \square

Remark 5.9. We remark that such a precise description of closed normal subgroups as it is done for V_A in Theorem 5.7 is rather impossible for U_A or CU_A . Indeed, notice that by Theorem 4.9, for every proper non-trivial ideal I of a unital C^* -algebra A we get the closed normal subgroups $CU_I \leq U_I \leq N_I$ of U_A and by Lemma 6.1, $U_I \subsetneq N_I$ if $\mathcal{Z}(A/I)$ is non-trivial. In particular, picking B any unital simple AF algebra and setting $A := B \oplus B \oplus B$ and $I := B \oplus \{0\} \oplus \{0\}$, we get $\mathcal{Z}(A/I)$ is two-dimensional, so non-trivial. Therefore there is no one-to-one correspondence between (perfect) ideals and closed normal subgroups of U_A even when A is AF.

We conclude this section with the following result that generalizes the main results from [23] that if A is a simple C^* -algebra, then $CU_A/\mathcal{Z}(CU_A)$ and V_A are topologically simple, and employs Theorems 4.9 and 5.2.

Theorem 5.10. *Let A be a unital C^* -algebra. Then the following are equivalent:*

- (1) *For every closed two-sided ideal I of A , either $I \subseteq \mathcal{Z}(A)$ or A/I is abelian.*
- (2) *The only perfect ideals of A are the trivial ideal $\{0\}$ and $\text{Id}([A, A])$.*
- (3) *V_A is topologically simple.*
- (4) *$CU_A/\mathcal{Z}(CU_A)$ is topologically simple.*
- (5) *For every closed two-sided ideal I of A , V_I is either trivial or equal to V_A .*
- (6) *For every closed two-sided ideal I of A , either CU_I is trivial, or $CU_I = CU_A$.*

Proof. (1) \implies (2): let $I \subseteq A$ be a perfect ideal. If $I \subseteq \mathcal{Z}(A)$, then $[I, I] = \{0\}$ and so since $I = \text{Id}([I, I])$ we get that $I = \{0\}$. If A/I is abelian, then necessarily $[A, A] \subseteq I$. It follows from

$$I \subseteq \text{Id}([I, I]) \subseteq \text{Id}([A, A]) \subseteq I$$

that $I = \text{Id}([A, A])$. Therefore, (2) holds.

(2) \implies (1): let $I \subseteq A$ be a closed two-sided ideal and suppose that $I \not\subseteq \mathcal{Z}(A)$. Since by Lemma 3.1, $\overline{[I, I]} = \overline{[A, A]}$ and the latter is nonzero as $I \not\subseteq \mathcal{Z}(A)$, we get that the perfect ideal $I' = \text{Id}([I, I])$ is nonzero. By the assumption we must have that $I' = \text{Id}([A, A])$ but then $[A, A] \subseteq I$, so A/I is abelian. Therefore (1) holds.

(2) \implies (3): since $V_{\{0\}} = \{\text{Id}\}$ and $\ker(P_{\{0\}}) = \{\text{Id}\}$, and also $V_{\text{Id}([A, A])} = V_A$ and $\ker(P_{\text{Id}([A, A])}) = V_A$, it immediately follows from Theorem 5.2 that the only closed normal subgroups of V_A are the trivial subgroup and V_A itself, thus V_A is topologically simple.

(3) \implies (2): Let $I \subseteq A$ be a perfect ideal. Then $V_I \leq V_A$ is a perfect closed normal subgroup and thus V_I is either trivial or equal to V_A which by Corollary 5.6 gives that I is either trivial or equal to $\text{Id}([A, A])$.

(2) \implies (4): let $H' \leq CU_A/\mathcal{Z}(CU_A)$ be a closed normal subgroup. By the fourth isomorphism theorem, there is a closed normal subgroup $\mathcal{Z}(CU_A) \leq H \leq CU_A$ such that $H' = H/\mathcal{Z}(CU_A)$. By [23, Theorem 3.7], H is normal in U_A , so by Theorem 4.9, there is a perfect ideal $I \subseteq A$ such that $CU_I \leq H \leq N_I$. If $I = \{0\}$ then $N_I = \mathcal{Z}(U_A)$ and so $H \subseteq CU_A \cap \mathcal{Z}(U_A) \subseteq \mathcal{Z}(CU_A)$, thus $H = \mathcal{Z}(CU_A)$. If $I = \text{Id}([A, A])$, then by Lemma 4.2, $CU_I = CU_A$ and since $CU_I \leq H \leq CU_A$ we get that $H = CU_A$. Therefore, $CU_A/\mathcal{Z}(CU_A)$ is topologically simple.

(4) \implies (2): we prove the contraposition. Suppose there is a perfect ideal $I \subseteq A$ that is neither equal to $\{0\}$ nor equal to $\text{Id}([A, A])$. Then by Lemma 4.4, CU_I is neither equal to $CU_{\{0\}} = \{1\}$ nor equal to $CU_{\text{Id}([A, A])} = CU_A$, and thus CU_I is a non-trivial proper closed normal subgroup of CU_A . Let H_I be the closed normal subgroup of CU_A generated by CU_I and $\mathcal{Z}(CU_A)$, which is the closure of $CU_I \cdot \mathcal{Z}(CU_A)$. Set $H'_I = H_I/\mathcal{Z}(CU_A)$, which is by

the second isomorphism theorem topologically isomorphic to $CU_I/(CU_I \cap \mathcal{Z}(CU_A))$, which is equal to $CU_I/\mathcal{Z}(CU_I)$ by Proposition 3.5. We claim that H'_I is a proper non-trivial closed normal subgroup of $CU_A/\mathcal{Z}(CU_A)$. H'_I must be non-trivial as otherwise, $CU_I = \mathcal{Z}(CU_I)$, which means that CU_I is abelian, and this contradicts that CU_I is non-trivial and topologically perfect. It remains to show that $H'_I \neq CU_A/\mathcal{Z}(CU_A)$, for which it suffices to show that $H_I \neq CU_A$. However, $\overline{(H_I, H_I)} = \overline{(CU_I, CU_I)} = CU_I$, while $\overline{(CU_A, CU_A)} = CU_A$, and $CU_I \subsetneq CU_A$. Therefore, (4) does not hold.

(5) \implies (3): let H be a closed normal subgroup of V_A . Then by Theorem 5.2, there exists a perfect ideal $I \subseteq A$ such that $V_I \leq H \leq \ker(P_I)$. If $V_I = V_A$ then clearly $H = V_A$, and if $V_I = \{\text{Id}\}$, then $I = \{0\}$, so $\ker(P_I) = \{\text{Id}\}$, and thus $H = \{\text{Id}\}$. Thus V_A is topologically simple.

(3) \implies (5): let $I \subseteq A$ be any closed two-sided ideal. Then V_I is a closed normal subgroup of V_A . Since V_A is topologically simple, V_I must be either trivial or equal to V_A , proving (5).

(4) \implies (6): let $I \subseteq A$ be a closed two-sided ideal of A . Let H_I be the closed subgroup of CU_A generated by CU_I and $\mathcal{Z}(CU_A)$, which is the closure of $CU_I \cdot \mathcal{Z}(CU_A)$. Then $H'_I = H_I/\mathcal{Z}(CU_A)$ is a closed normal subgroup of the topologically simple group $CU_A/\mathcal{Z}(CU_A)$, which is, as showed in the proof of (4) \implies (2), topologically isomorphic to $CU_I/\mathcal{Z}(CU_I)$. If $H'_I = \{1\}$, then $CU_I \subseteq \mathcal{Z}(CU_I)$, so CU_I is abelian and topologically perfect, whence $CU_I = \{1\}$. If $H'_I = CU_A/\mathcal{Z}(CU_A)$, then $H_I = CU_A$, so

$$CU_A = \overline{(H_I, H_I)} = \overline{(CU_I \cdot \mathcal{Z}(CU_A), CU_I \cdot \mathcal{Z}(CU_A))} = \overline{(CU_I, CU_I)} = CU_I,$$

so (6) holds.

(6) \implies (4): let $H' \leq CU_A/\mathcal{Z}(CU_A)$ be a closed normal subgroup. By the third isomorphism theorem, there exists a closed normal subgroup $\mathcal{Z}(CU_A) \leq H \leq CU_A$ such that $H' = H/\mathcal{Z}(CU_A)$. By [23, Theorem 3.7] H is normal in U_A so by Theorem 4.9, there exists a perfect ideal $I \subseteq A$ such that $CU_I \leq H \leq N_I$. Now we use the assumption. If $CU_I = CU_A$, then $H = CU_A$ and $H' = CU_A/\mathcal{Z}(CU_A)$, so assume that CU_I is trivial $= CU_{\{0\}}$. However, then $I = \{0\}$ by the perfectness of I and Corollary 5.6, and arguing as in the implication (2) \implies (4) we get that $H \subseteq \mathcal{Z}(CU_A)$, thus $H = \mathcal{Z}(CU_A)$ and so H' is trivial. Therefore, $CU_A/\mathcal{Z}(CU_A)$ is topologically simple. \square

6. LIE STRUCTURE OF U_A AND APPLICATIONS

In this section, we shall further investigate the Lie structure of U_A . The main application, proved in Corollary 6.5, is that C^* -subalgebras that are invariant under unitaries from the connected component of the identity are invariant under all unitaries.

The following lemma is already implicit in [23] and it immediately follows from [23, Lemma 3.2] and Lemma 3.2.

Lemma 6.1. *Let A be a unital C^* -algebra and $I \subseteq A$ be a two-sided closed ideal. Let $P_I : A \rightarrow A/I$ be the canonical quotient map. Then $N_I = P_I^{-1}(\mathcal{Z}(U_{A/I})) \cap U_A$.*

Let us define \mathfrak{n}_I to be the set

$$\{a \in A_{\text{sa}} \mid [a, A] \subseteq \overline{[I, I]}\}.$$

Lemma 6.2. *We have*

$$\mathfrak{n}_I = \{a \in A_{\text{sa}} \mid \exp(ita) \in N_I \ \forall t \in \mathbb{R}\};$$

i.e. \mathfrak{n}_I is the Lie algebra of N_I .

Proof. Denote the set on the right hand side from the statement as X .

First, let us show $\mathfrak{n}_I \subseteq X$. Pick $a \in \mathfrak{n}_I$ and $u \in U_A$. We may suppose that $\|a\|$ is small enough so that $\log((u, e^{ia})) = \log(\exp(iuau^*) \exp(-ia))$ is defined and equal to ic for some $c \in A_{\text{sa}}$, which is moreover a converging series given by the Baker-Campbell-Hausdorff formula.

By definition, we have $ua - au \in \overline{[I, I]} \subseteq I$, and thus also $uau^* - a \in I$. Since also, by the Baker-Campbell-Hausdorff formula, and since by definition $[a, A] \in \overline{[I, I]}$, we have

$$\log(\exp(iuau^*)\exp(-ia)) - i(uau^* - a) \in I,$$

we obtain $\log(\exp(iuau^*)\exp(-ia)) \in I$, so $c \in I$. It follows that $(\exp(ia), u) \in \tilde{I}$, so $\exp(ia) \in N_I$. Clearly, $\exp(ita) \in N_I$, for any $t \in \mathbb{R}$ with $|t| \leq 1$. Since N_I is a group, $\exp(ita) \in N_I$ for any $t \in \mathbb{R}$, and $\mathfrak{n}_I \subseteq X$ holds.

Now we show that $X \subseteq \mathfrak{n}_I$. Pick some $a \in X$ and $u \in A$. We check that $[a, u] \in \overline{[I, I]}$. Since the linear span of U_A is A by Lemma 4.3, we may suppose that $u \in U_A$. We may also suppose that $\|a\|$ is small enough so that $\log(\exp(iuau^* - a))$ is defined and equal to $i(uau^* - a)$.

Since $a \in X$, by applying the Trotter product formula, we obtain

$$\exp(i(uau^* - a)) = \lim_{n \rightarrow \infty} ((\exp(ia/n), u))^n \in CU_I.$$

It follows that $uau^* - a \in I$, and thus also $ua - au = [u, a] \in I$. Since $[A, A] \cap I \subseteq \overline{[I, I]}$ by Lemma 3.1, we are done. \square

Proposition 6.3. N_I is a Banach-Lie group with \mathfrak{n}_I as the Lie algebra.

Proof. We show that for every $a \in A_{\text{sa}}$ such that $\|a\|$ is sufficiently small we have $a \in \mathfrak{n}_I$ if and only if $\exp(ia) \in N_I$ since then we will have that \exp induces a local homeomorphism between \mathfrak{n}_I and N_I and we can conclude by [15, Definition 5.32 and Corollary 5.34].

By Lemma 6.2, if $a \in \mathfrak{n}_I$ then $\exp(ia) \in N_I$. So conversely assume that $e^{ia} \in N_I$ and that $\|a\|$ is small enough so that $\log(e^{ia})$ and $\log(e^{P(ia)})$ are defined (and equal to ia and $P(ia)$ respectively). We show that $a \in \mathfrak{n}_I$. Again by Lemma 6.2, it suffices to show that $e^{ita} \in N_I$, for all $t \in \mathbb{R}$. It is clear that in fact it is enough to prove that for t such that $|t| \leq 1$. Let $P_I : A \rightarrow A/I$ be the quotient map. Then by Lemma 6.1 we have

$$P_I(e^{ia}) \in \mathcal{Z}(U_{A/I}) = \mathcal{Z}(A/I) \cap U_{A/I},$$

where the last equality holds since by Lemma 4.3, $U_{A/I}$ spans A/I .

Consequently,

$$P_I(a) = P_I(\log(e^{ia})/i) = \log(P_I(e^{ia}))/i \in \mathcal{Z}(A/I)$$

and thus also $P_I(ta) \in \mathcal{Z}(A/I)$, for any t such that $|t| \leq 1$.

Therefore $\exp(iP_I(ta)) \in \mathcal{Z}(A/I) \cap U_{A/I} = \mathcal{Z}(U_{A/I})$ and since $P_I(\exp(ita)) = \exp(iP_I(ta))$ we get that $\exp(ita) \in N_I$ by Lemma 6.1. \square

Lemma 6.4. Every closed connected Lie normal subgroup of U_A is normal in $U(A)$.

Proof. Let H be a closed connected Lie normal subgroup of U_A . By Theorem 4.9 there is a perfect ideal I of A such that $CU_I \leq H \leq N_I$. Recall from the proof of Theorem 4.9 that we have $[H, A] \subseteq I$. One can check directly from their respective definitions that both CU_I and N_I are normal in $U(A)$. It suffices to verify that for every $u \in U(A)$ and $g \in H$ we have $(u, g) \in CU_I$ since then $ugu^* \in CU_I \cdot H \subseteq H$.

Let us first check $(u, g) \in \tilde{I}$. This follows since $ug - gu \in I$ as $[H, A] \subseteq I$, thus for $x = ug - gu \in I$ we have $ugu^*g^* = (x + gu)u^*g^* = xu^*g^* + 1 \in \tilde{I}$.

Now since H is connected and Lie, $g = e^{ib_1} \dots e^{ib_n}$, for some self-adjoint elements b_1, \dots, b_n from the Lie algebra of H . Without loss of generality we can suppose that $g = e^{ib}$, for $\|b\|$ small enough, so that $c := \log(u, g)$ is defined and the Baker-Campbell-Hausdorff formula for $e^{iubu^*}e^{-ib}$ converges. Thus $(u, e^{ib}) = e^{ic}$. The rest of the argument is the same as [23, Lemma 3.2]. We include the argument for completeness. For each $t \in \mathbb{R}$, $e^{itb} \in H$, so we differentiate $(u, e^{itb}) - 1$ at $t = 0$ to obtain $ubu^* - b \in \tilde{I}$, which implies $[u, b] \in \tilde{I}$ and thus $[u, b] \in I$. Then

$$(u, e^{ib}) - (ubu^* - b) = e^{iubu^*}e^{-ib} - \underbrace{(ubu^* - b)}_{\in I} \in \overline{[A, H]} \subset I,$$

whence $e^{ic} \in I \cap \overline{[A, A]} = \overline{[I, I]}$ and thus $e^{ic} \in CU_I$ by Lemma 3.3. \square

Corollary 6.5. *Let A be a unital C^* -algebra and let $B \subseteq A$ be a C^* -subalgebra that is Ad_{U_A} -invariant. Then B is $\text{Ad}_{U(A)}$ -invariant.*

Proof. Let A and B be as in the statement. Then U_B is a closed normal subgroup of U_A , which is moreover connected and Lie. Thus by Lemma 6.4, U_B is normal in $U(A)$, i.e. it is $\text{Ad}_{U(A)}$ -invariant. Since U_B spans B by Lemma 4.3, also B is $\text{Ad}_{U(A)}$ -invariant. \square

7. GROUP CHARACTERIZATIONS OF SIMPLICITY OF C^* -ALGEBRAS

In this final section, we investigate conditions on the structure of U_A that guarantee simplicity of A . That is, in a way, a reverse problem to what we consider in Theorem 5.10. We find applications to reduced group C^* -algebras.

7.1. General conditions on simplicity of C^* -algebras.

Lemma 7.1. *(well-known) Let A be a unital simple C^* -algebra. Then $\mathcal{Z}(A) = \mathbb{C}$.*

Proof. Let $a \in \mathcal{Z}(A) \setminus \{0\}$. Consider $I = \overline{Aa} = \overline{aA}$, which is a closed two-sided ideal of A . Since A is unital, $a \in I$, so $I \neq \{0\}$. Therefore by the simplicity of A , $I = A$ holds. Thus there exists $b \in A$ such that $\|ba - 1\| < 1$. Then ba , hence a is invertible. This shows that any nonzero element in $\mathcal{Z}(A)$ is invertible, whence $\mathcal{Z}(A) = \mathbb{C}$ by Banach–Mazur theorem. \square

The following result of Cuntz and Pedersen from [8, Proposition 2.7] will be used often, so we state it here. For a unital C^* -algebra A , we denote by $T(A)$ the state of all tracial states on A .

Theorem 7.2 ([8]). *Let A be a C^* -algebra. If A does not admit a tracial state, then $\overline{[A, A]} = A$ holds. If A admits a tracial state, then $\overline{[A, A]} = \bigcap_{\tau \in T(A)} \ker(\tau)$ holds.*

Lemma 7.3. *Let A be a unital C^* -algebra. If A does not admit a tracial state, then $U_A = CU_A$, and if A admits exactly one tracial state, then U_A/CU_A is trivial or equal to \mathbb{T} .*

Proof. If A does not admit a tracial state, then by Theorem 7.2, $\overline{[A, A]} = A$, thus by the combination of Lemma 3.1 and [23, Lemma 3.1] we get that $CU_A = U_A$. Suppose that A admits exactly one tracial state. Then again by Theorem 7.2 we get $A = \overline{[A, A]} \oplus \mathbb{C}$. Denote by Q the quotient map $U_A \rightarrow U_A/CU_A$. Let $u \in U_A$. Write u as $\exp(ia_1) \cdots \exp(ia_n)$, where $a_1, \dots, a_n \in A_{\text{sa}}$. Write $a_k = b_k + r_k 1$, $b_k \in \overline{[A, A]}_{\text{sa}}$, $r_k \in \mathbb{R}$ for each $k = 1, \dots, n$. Then

$$u = e^{ib_1} \cdots e^{ib_n} \exp(ir)$$

where $r = \sum_{k=1}^n r_k$, so $Q(u) = Q(e^{ir})$ because $e^{ib_k} \in CU_A$ by [23, Lemma 3.1]. Consequently, $Q[U_A] = Q[\mathbb{T}]$. Since Q is continuous and U_A is connected, U_A/CU_A is a connected quotient of \mathbb{T} , thus either trivial, or isomorphic to \mathbb{T} . \square

Theorem 7.4. *Let A be a unital C^* -algebra with at most one tracial state which we then moreover require not to be a character. Then A is simple if and only if $CU_A/\mathcal{Z}(CU_A)$ is topologically simple and $\mathcal{Z}(CU_A) \subseteq \mathbb{T}$.*

Proof. Suppose first that A is simple. By [23, Corollary 3.8], $CU_A/\mathcal{Z}(CU_A)$ is topologically simple. Set $H := \mathcal{Z}(CU_A)$. It is a closed normal subgroup of CU_A and by [23, Theorem 3.7], it is also normal in U_A . Therefore, by Theorem 4.9, there exists a perfect ideal $I \subseteq A$ such that $CU_I \leq H \leq N_I$. If $I = \text{Id}([A, A])$, then $H = CU_A$, implying that CU_A is abelian and topologically perfect, whence trivial.

So suppose that $I = \{0\}$. Then $H \leq N_I = \mathcal{Z}(U_A)$. Since A is simple, by Lemma 7.1, $\mathcal{Z}(A) = \mathbb{C}$, so $\mathcal{Z}(U_A) = \mathbb{T}$ (cf. Lemma 4.3) and we have $\mathcal{Z}(CU_A) \subseteq \mathbb{T}$.

Now we prove the converse. Let $I \subseteq A$ be a non-zero closed two-sided ideal. By Theorem 5.10, either $I \subseteq \mathcal{Z}(A)$ or A/I is abelian.

Case 1. $I \subseteq \mathcal{Z}(A)$. Assume first that A admits a unique tracial state τ . Pick a non-zero $b \in I_{\text{sa}}$. By Theorem 7.2, we can write $b = a + r1$, where $a \in \overline{[A, A]}$ and $r \in \mathbb{R}$. Since $b \in \mathcal{Z}(A)$, also $a \in \mathcal{Z}(A)$. Then by [23, Lemma 3.1], for every $t \in \mathbb{R}$,

$$\exp(ita) \in CU_A \cap \mathcal{Z}(U_A) \subseteq \mathcal{Z}(CU_A) \subseteq \mathbb{T}.$$

By differentiating the left hand side at $t = 0$, we obtain $a \in \mathbb{R}$, which implies $a \in \mathbb{R} \cap \overline{[A, A]} = \{0\}$. Therefore, for every $b \in I_{\text{sa}}$ we have $b = r \in I \cap \mathbb{C}^\times$. Thus $I = A \subseteq \mathcal{Z}(A)$, so A is an abelian C^* -algebra with a unique tracial state. Thus $A = \mathbb{C}$, which admits a character, a contradiction.

Next, assume that A does not admit a tracial state. Then by Theorem 7.2 and Lemma 7.3, we have $A = \overline{[A, A]}$, $U_A = CU_A$, and $\mathcal{Z}(U_A) = \mathcal{Z}(CU_A) \subseteq \mathbb{T}$. Thus $\mathcal{Z}(U_A) = \mathbb{T}$. This implies that $\mathcal{Z}(A) = \mathbb{C}$. Since A does not admit a tracial state, $A \neq \mathbb{C}$ and thus I is a proper ideal of A contained in \mathbb{C} , proving that $I = \{0\}$, which is a contradiction.

Case 2. A/I is abelian. It follows that $\overline{[A, A]} \subseteq I \subseteq A$. Assume that A admits a unique tracial state τ .

Since again by Theorem 7.2, $A = \overline{[A, A]} \oplus \mathbb{C}$ and if $I \neq A$ we get that $I = \overline{[A, A]}$. Thus it follows that $A/I = \mathbb{C}$ and so A admits a non-trivial character, a contradiction. Thus $I = A$.

Next assume that A does not have a tracial state. Then by Theorem 7.2, $A = \overline{[A, A]} = I$.

This concludes that A is simple. \square

7.2. Simplicity of reduced group C^* -algebras. Let G be a countable discrete group, $C_r^*(G)$ be the reduced group C^* -algebra of G . Denote by $\text{Rep}(G)$ the class of all unitary representations of G , and $\text{Rep}_\lambda(G)$ the subclass of all $\pi \in \text{Rep}(G)$ with $\pi \prec \lambda$. Here, λ is the left regular representation of G and \prec denotes the weak containment (we refer to [3, Part II] for a background on unitary representations of groups, weak containment, and its relations with group C^* -algebras). For $\pi \in \text{Rep}(G)$ acting on a Hilbert space H and for any $x \in \mathbb{C}G$ (the group algebra of G), we define

$$\pi(x) = \sum_{g \in G} a_g \pi(g), \quad x = \sum_{g \in G} a_g \cdot g \quad a_g \in \mathbb{C}.$$

We will be using the following characterization of the weak containment of representations.

Proposition 7.5. *For $\pi_1, \pi_2 \in \text{Rep}(G)$, the following conditions are equivalent.*

- (1) $\pi_1 \prec \pi_2$.
- (2) $\|\pi_1(x)\| \leq \|\pi_2(x)\|$ for every $x \in \mathbb{C}G$.
- (3) There exists a unital $*$ -homomorphism $\Phi: C_{\pi_2}^*(G) \rightarrow C_{\pi_1}^*(G)$ such that $\Phi(\pi_2(g)) = \pi_1(g)$ for every $g \in G$.

See [3, Theorem F.4.4] for a proof. The next result is well-known. We include the proof for completeness.

Proposition 7.6. *Let G be a countable discrete group. The following conditions are equivalent.*

- (1) There exists a unital surjective $*$ -homomorphism from $C_r^*(G)$ onto a unital abelian C^* -algebra.
- (2) There exists a unital surjective $*$ -homomorphism from $C_r^*(G)$ onto \mathbb{C} .
- (3) G is amenable.

Proof. (3) \implies (2) Assume that G is amenable and let ι_G be the trivial representation of G . Then $\iota_G \prec \lambda_G$, whence there exists a unital surjective $*$ -homomorphism from $C_r^*(G)$ onto $\mathbb{C} = C_{\iota_G}^*(G)$.

(2) \implies (3) Assume that there exists a unital surjective $*$ -homomorphism $\Phi: C_r^*(G) \rightarrow \mathbb{C}$. Then $\pi(g) = \Phi(\lambda(g))$, ($g \in G$) defines a unitary representation $\pi: G \rightarrow \mathbb{T}$. Let $N = \ker(\pi)$. Then the restriction $\lambda_G|_N$ of the left regular representation λ_G of G to the subgroup N is unitarily equivalent to a multiple of λ_N and $\pi|_N$ is unitarily equivalent to a multiple of ι_N . Thus for every $x \in \mathbb{C}N$, we have

$$\|\iota_N(x)\| = \|\pi(x)\| \leq \|\lambda_G(x)\| = \|\lambda_N(x)\|,$$

which shows that $\iota_N \prec \lambda_N$, whence N is amenable. Since subgroups of \mathbb{T} are abelian hence amenable, and amenability passes to extensions, it follows that G is also amenable.

(2) \implies (1) is trivial.

(1) \implies (2) Unital abelian C^* -algebras have pure states, and they are nonzero $*$ -homomorphisms onto \mathbb{C} . \square

Recall that an infinite group is called *ICC* (infinite conjugacy classes) if all the non-trivial conjugacy classes are infinite. Recall also that every locally compact group contains a largest normal amenable subgroup called the *amenable radical*. It is known for a discrete group G that if $C_r^*(G)$ is simple, then the amenable radical of G is trivial (see [10, Proposition 3] and that if G has trivial amenable radical (which by Breuillard–Kalantar–Kennedy–Ozawa theorem from [5] is equivalent to $C_r^*(G)$ having the unique tracial state), then G is ICC (see e.g. [14, Theorem 4.3]). We summarize the above results.

Theorem 7.7 ([5, 14, 10]). *Let G be a countable discrete group. Consider the following conditions.*

- (1) $C_r^*(G)$ is simple.
- (2) $C_r^*(G)$ has unique tracial state.
- (3) G has trivial amenable radical.
- (4) G is ICC.

Then $(1) \implies (2) \iff (3) \implies (4)$.

Proposition 7.8. *Let G be a countable discrete ICC group. Then for $A := C_r^*(G)$ we have that all proper quotients of A are abelian if and only if $CU_A/\mathcal{Z}(CU_A)$ is topologically simple. Moreover, in that case $\mathcal{Z}(CU_A) \subseteq \mathcal{Z}(U_A) = \mathbb{T}$.*

We will need the following well-known lemma.

Lemma 7.9. *Let G be a countable discrete ICC group, $A = C_r^*(G)$ be the reduced group C^* -algebra of G . Then $\mathcal{Z}(A) = \mathbb{C}$.*

Proof. Note that A is a C^* -subalgebra of the group von Neumann algebra $M = L(G)$ acting on $\ell^2(G)$. Thus if $z \in \mathcal{Z}(A)$, then write $z = \sum_{g \in G} z(g)\lambda(g)$ in the 2-norm expansion inside M . Exactly as in the proof that M is a (type II₁) factor, the condition $\lambda(h)z\lambda(h)^* = z$ ($h \in G$) implies that for any nontrivial element $g \in G \setminus \{1\}$, $z(h^{-1}gh) = z(g)$ for all $h \in G$, which by the condition $\sum_{g \in G} |z(g)|^2 < \infty$ and the ICC property imply that $z(g) = 0$, whence $z = z(1)1 \in \mathbb{C}$ holds. \square

Proof of Proposition 7.8. First, notice that by Lemma 7.9, $\mathcal{Z}(A) = \mathbb{C}$ which is equivalent with $\mathcal{Z}(U_A) = \mathbb{T}$.

Suppose first that all proper quotients of A are abelian. Let $H' \leq CU_A/\mathcal{Z}(CU_A)$ be a closed normal subgroup and, by the fourth isomorphism theorem, let $\mathcal{Z}(CU_A) \leq H \leq CU_A$ be a closed normal subgroup such that $H' = H/\mathcal{Z}(CU_A)$. By [23, Theorem 3.7], H is also normal in U_A . Therefore by Theorem 4.9 there exists a perfect ideal $I \subseteq A$ such that $CU_I \leq H \leq N_I$.

Suppose that $I \neq \{0\}$. Because A/I is abelian, $\overline{[A, A]} \subseteq I$ holds. It follows, by [23, Lemma 3.1], that $CU_A \leq U_I$ and thus, since by Proposition 4.1, CU_I is the largest closed perfect normal subgroup contained in U_I , that $CU_A = CU_I$. So since $CU_I \leq H \leq CU_A$ we get $H = CU_A$.

Suppose that $I = \{0\}$. Then $\mathcal{Z}(CU_A) \leq H \leq N_I = \mathcal{Z}(U_A) = \mathbb{T}$. Since $\mathcal{Z}(U_A) \cap CU_A \subseteq \mathcal{Z}(CU_A)$ we get $H = \mathcal{Z}(CU_A)$ and so H' is trivial.

Therefore, $CU_A/\mathcal{Z}(CU_A)$ is topologically simple.

Now we conversely assume that $CU_A/\mathcal{Z}(CU_A)$ is topologically simple. By Theorem 5.10 we get that for every nonzero proper ideal $I \subseteq A$ we have that either $I \subseteq \mathcal{Z}(A)$ or A/I is abelian. However, the former is not possible since $\mathcal{Z}(A) = \mathbb{C}$ by Lemma 7.9.

Notice that by the first part of the proof, with H' being the trivial subgroup, we get that $\mathcal{Z}(CU_A) \subseteq \mathcal{Z}(U_A) = \mathbb{T}$. \square

Lemma 7.10. *Let G be a countable discrete group and suppose that $C_r^*(G)$ is simple. Then $U_A/\mathbb{T} = CU_A/\mathcal{Z}(CU_A)$.*

Proof. By Theorem 7.7, G is necessarily ICC and so by Proposition 7.8, $\mathcal{Z}(CU_A) \subseteq \mathcal{Z}(U_A) = \mathbb{T}$. Moreover, by Theorem 7.7 again, $C_r^*(G)$ has a unique tracial state and so by Theorem 7.2, $A = \overline{[A, A]} \oplus \mathbb{C}$.

We define a topological isomorphism $T : U_A/\mathbb{T} \rightarrow CU_A/\mathcal{Z}(CU_A)$. For $u \in U_A$, we denote by $[u]_1$ its equivalence class in U_A/\mathbb{T} , and for $u \in CU_A$, we denote by $[u]_2$ its equivalence class in $CU_A/\mathcal{Z}(CU_A)$. Let $[u]_1 \in U_A/\mathbb{T}$ be an element with a representative $u \in U_A$. Then $u = \exp(i(a_1 + r_1)) \cdots \exp(i(a_n + r_n))$, where $a_1, \dots, a_n \in \overline{[A, A]}$ and $r_1, \dots, r_n \in \mathbb{R}$. We set

$$T([u]_1) := [\exp(ia_1) \cdots \exp(ia_n)]_2.$$

Then T is well-defined. Indeed, if $u = e^{i(a_1+r_1)} \cdots e^{i(a_n+r_n)}$ and $u' = e^{i(a'_1+r'_1)} \cdots e^{i(a'_m+r'_m)}$ are elements in U_A which defines the same class in U_A/\mathbb{T} where $a_1, \dots, a_n, a'_1, \dots, a'_m \in \overline{[A, A]}_{\text{sa}}$ and $r_1, \dots, r_n, r'_1, \dots, r'_m \in \mathbb{R}$, then there exists $z \in \mathbb{T}$ such that $u' = zu$, whence

$$e^{ia'_1} \cdots e^{ia'_m} = we^{ia_1} \cdots e^{ia_n},$$

where $w = \frac{z\lambda}{\lambda}$, $\lambda = \exp(i(r_1 + \cdots + r_n))$, $\lambda' = \exp(i(r'_1 + \cdots + r'_m)) \in \mathbb{T}$. In particular, $w \in CU_A \cap \mathbb{T} = \mathcal{Z}(CU_A)$ holds. Therefore,

$$[e^{ia_1} \cdots e^{ia_n}]_2 = [e^{ia'_1} \cdots e^{ia'_m}]_2,$$

whence T is well-defined.

It is clear that T is a surjective group homomorphism. In order to check that T is a homeomorphism, it is enough to check that T is injective and continuous, since then T^{-1} is a Borel homomorphism between Polish groups, thus continuous (see [17, Theorem 9.10]).

To show that T is injective, pick $[u]_1 \neq 1 \in U_A/\mathbb{T}$, where $u \in U_A$. Then $u \notin \mathbb{T}$, so writing $u = \exp(i(a_1 + r_1)) \cdots \exp(i(a_n + r_n))$, where $a_1, \dots, a_n \in \overline{[A, A]}$ and $r_1, \dots, r_n \in \mathbb{R}$, and then defining $u' := \exp(ia_1) \cdots \exp(ia_n)$, we get that $u' \neq 1$. Moreover, $u' \notin \mathbb{T}$ and since $\mathcal{Z}(CU_A) \subseteq \mathbb{T}$ we get

$$T([u]_1) = [u']_2 \neq 1.$$

Since $[u]_1 \in U_A/\mathbb{T}$ was arbitrary, we get that T is injective. Finally, we check the continuity. Suppose $u_n \in U_A$ satisfies $[u_n]_1 \rightarrow [1]$ in U_A/\mathbb{T} . Without loss of generality, we may assume that $u_n \rightarrow 1_{U_A}$. For each n , write u_n as $e^{i(a_1+r_1)} \cdots e^{i(a_m+r_m)}$ and denote by $u'_n \in CU_A$ the corresponding element $e^{ia_1} \cdots e^{ia_m}$. Notice that for each n , we have $u_n = u'_n t_n$, where $t_n \in \mathbb{T}$. By compactness of \mathbb{T} and passing to a subsequence if necessary we may suppose that $t_n \rightarrow t \in \mathbb{T}$. Since $u_n \rightarrow 1_{U_A}$ we get $u'_n \rightarrow t \in \mathbb{T}$. Then $t \in CU_A \cap \mathbb{T} = \mathcal{Z}(CU_A)$ and since $T([u_n]_1) = [u'_n]_2$ for each n , we get $T([u_n]_1) \rightarrow CU_A/\mathcal{Z}(CU_A)$. This finishes the proof. \square

Theorem 7.11. *Let G be a countable discrete non-amenable ICC group. Then $A := C_r^*(G)$ is simple if and only if U_A/\mathbb{T} is topologically simple.*

Proof. If A is simple, then by Proposition 7.8, $CU_A/\mathcal{Z}(CU_A)$ is topologically simple and so by Lemma 7.10, U_A/\mathbb{T} is topologically simple.

Conversely, suppose that U_A/\mathbb{T} is topologically simple. We claim that $CU_A/\mathcal{Z}(CU_A)$ is then topologically simple. Indeed, let H' be a closed normal subgroup of $CU_A/\mathcal{Z}(CU_A)$ and let $\mathcal{Z}(CU_A) \leq H \leq CU_A$ be a closed normal subgroup such that $H' = H/\mathcal{Z}(CU_A)$. Again by [23, Theorem 3.7], H is normal in U_A , so either $H \subseteq \mathbb{T}$ or $H = U_A$. Notice that this argument, applied to $H = \mathcal{Z}(CU_A)$, in particular implies that $\mathcal{Z}(CU_A) \subseteq \mathbb{T}$ as the case $\mathcal{Z}(CU_A) = U_A$ is impossible.

Case 1. $H = U_A$: this implies that $CU_A = U_A$, and thus $CU_A/\mathcal{Z}(CU_A) = CU_A/T = U_A/\mathbb{T}$ is topologically simple.

Case 2. $H \subseteq \mathbb{T}$: It follows that H is central in CU_A and thus $H = \mathcal{Z}(CU_A)$, so H' is trivial.

We have therefore proved that $CU_A/\mathcal{Z}(CU_A)$ is topologically simple. So by Proposition 7.8, all the proper quotients of $C_r^*(G)$ are abelian. However, since G is not amenable, by Proposition 7.6, $C_r^*(G)$ is simple. \square

REFERENCES

- [1] H. ANDO AND M. DOUCHA, *An application of Kirchberg's lemma on central sequence algebras to groups of approximately inner automorphisms*, arXiv preprint arXiv:2310.11922, (2023).
- [2] H. ANDO, M. DOUCHA, AND Y. MATSUZAWA, *Large scale geometry of Banach-Lie groups*, Trans. Am. Math. Soc., 375 (2022), pp. 2827–2881.
- [3] B. BEKKA, P. DE LA HARPE, AND A. VALETTE, *Kazhdan's property (T)*, vol. 11 of New Mathematical Monographs, Cambridge University Press, Cambridge, 2008.
- [4] A. BONFIGLIOLI AND R. FULCI, *Topics in noncommutative algebra*, vol. 2034 of Lecture Notes in Mathematics, Springer, Heidelberg, 2012. The theorem of Campbell, Baker, Hausdorff and Dynkin.
- [5] E. BREUILLARD, M. KALANTAR, M. KENNEDY, AND N. OZAWA, *C^* -simplicity and the unique trace property for discrete groups*, Publ. Math. Inst. Hautes Études Sci., 126 (2017), pp. 35–71.
- [6] M. BREŠAR, E. KISSIN, AND V. S. SHULMAN, *Lie ideals: from pure algebra to C^* -algebras*, J. Reine Angew. Math., 623 (2008), pp. 73–121.
- [7] A. CHAND AND L. ROBERT, *Simplicity, bounded normal generation, and automatic continuity of groups of unitaries*, Adv. Math., 415 (2023), p. 52. Id/No 108894.
- [8] J. CUNTZ AND G. K. PEDERSEN, *Equivalence and traces on C^* -algebras*, J. Functional Analysis, 33 (1979), pp. 135–164.
- [9] K. R. DAVIDSON, *C^* -algebras by example*, vol. 6 of Fields Institute Monographs, American Mathematical Society, Providence, RI, 1996.
- [10] P. DE LA HARPE, *On simplicity of reduced C^* -algebras of groups*, Bull. Lond. Math. Soc., 39 (2007), pp. 1–26.
- [11] G. A. ELLIOT AND M. RØRDAM, *The automorphism group of the irrational rotation C^* -algebra*, Commun. Math. Phys., 155 (1993), pp. 3–26.
- [12] L. FOGANG TAKOUTSING AND L. ROBERT, *Distinguishing C^* -algebras by their unitary groups*, arXiv preprint arXiv:2306.15825, (2023).
- [13] E. GARDELLA AND H. THIEL, *Prime ideals in C^* -algebras and applications to lie theory*, arXiv preprint arXiv:2306.16510, (2023).
- [14] U. HAAGERUP, *A new look at C^* -simplicity and the unique trace property of a group*, in Operator algebras and applications—the Abel Symposium 2015, vol. 12 of Abel Symp., Springer, [Cham], 2017, pp. 167–176.
- [15] K. H. HOFMANN AND S. A. MORRIS, *The structure of compact groups—a primer for the student—a handbook for the expert*, vol. 25 of De Gruyter Studies in Mathematics, De Gruyter, Berlin, fourth ed., [2020] ©2020.
- [16] A. HOPENWASSER AND V. PAULSEN, *Lie ideals in operator algebras.*, J. Oper. Theory, 52 (2004), pp. 325–340.
- [17] A. S. KECHRIS, *Classical descriptive set theory*, vol. 156 of Grad. Texts Math., Berlin: Springer-Verlag, 1995.
- [18] L. W. MARCOUX, *Projections, commutators and Lie ideals in C^* -algebras*, Math. Proc. R. Ir. Acad., 110A (2010), pp. 31–55.
- [19] L. W. MARCOUX AND G. J. MURPHY, *Unitarily-invariant linear spaces in C^* -algebras*, Proc. Am. Math. Soc., 126 (1998), pp. 3597–3605.
- [20] C. R. MIERS, *Closed Lie ideals in operator algebras*, Canadian J. Math., 33 (1981), pp. 1271–1278.
- [21] N. C. PHILLIPS, *A survey of exponential rank*, in C^* -algebras: 1943–1993 (San Antonio, TX, 1993), vol. 167 of Contemp. Math., Amer. Math. Soc., Providence, RI, 1994, pp. 352–399.
- [22] L. ROBERT, *On the Lie ideals of C^* -algebras*, J. Operator Theory, 75 (2016), pp. 387–408.
- [23] ———, *Normal subgroups of invertibles and of unitaries in a C^* -algebra*, J. Reine Angew. Math., 756 (2019), pp. 285–319.
- [24] M. RØRDAM, F. LARSEN, AND N. LAUSTSEN, *An introduction to K -theory for C^* -algebras*, vol. 49 of London Mathematical Society Student Texts, Cambridge University Press, Cambridge, 2000.

HIROSHI ANDO, DEPARTMENT OF MATHEMATICS AND INFORMATICS, CHIBA UNIVERSITY, 1-33 YAYOI-CHO, INAGE, CHIBA, 263- 8522, JAPAN

Email address: hiroando@math.s.chiba-u.ac.jp

INSTITUTE OF MATHEMATICS, CZECH ACADEMY OF SCIENCES, ŽITNÁ 25, 115 67 PRAHA 1, CZECH REPUBLIC

Email address: doucha@math.cas.cz

URL: <https://users.math.cas.cz/~doucha/>