On the One-Dimensional Contact Process with Enhancements

Enrique Andjel

Leonardo T. Rolla

March 25, 2025

Abstract

We study a one-dimensional contact process with two infection parameters, one giving the infection rates at the boundaries of a finite infected region and the other one the rates within that region. We prove that the critical value of each of these parameters is a strictly monotone continuous function of the other parameter. We also show that if one of these parameters is equal to the critical value of the standard contact process and the other parameter is strictly larger, then the infection starting from a single point has positive probability of surviving. This is in contrast with another result also obtained here, that the critical contact process on the half line with enhanced infection rate at finitely many sites also dies out.

1 Introduction

In this paper we consider a one-dimensional contact process with modified boundaries. This model was introduced by Durrett and Schinazi in [5]. It differs from the standard one-dimensional contact process only in the way the rightmost (leftmost) infected site transmits its infection to its right (left) nearest neighbor. These infection rates are given by a parameter λ_e while infection everywhere else occurs at rate λ_i , where e and i refer to external and internal (as usual, we assume that the recovery rate is 1). When $\lambda_e = \lambda_i = \lambda$, this process reduces to the case of standard contact process, and we refer the reader to [7, 11, 15] for its basic properties. (At the end of this introduction, we also consider another variant of the process.)

Let $\theta(\lambda_i, \lambda_e)$ denote the probability that, starting with only the origin infected, the infection survives throughout time.

Although comparisons between different pairs (λ_i, λ_e) on the parameter space are not immediate, they can be compared with points (λ_i, λ_i) and (λ_e, λ_e) through any standard graphical construction (we describe one in the next section). Using this fact, that the critical contact process dies out [3], and comparing the boundary of the process with a simple random walk, one immediately gets the picture of the phase space shown in Figure 1.1a. It turns out that $\theta(\lambda_i, \lambda_e)$ is non-decreasing in both parameters. We note that although this property is very intuitive, its proof is not immediate [5, Proposition 1]. So we define

$$\lambda_c := \inf\{\lambda : \theta(\lambda, \lambda) > 0\} = \sup\{\lambda : \theta(\lambda, \lambda) = 0\},\\ \lambda_*^i(\lambda_e) := \inf\{\lambda : \theta(\lambda, \lambda_e) > 0\} = \sup\{\lambda : \theta(\lambda, \lambda_e) = 0\},\\ \lambda_*^e(\lambda_i) := \inf\{\lambda : \theta(\lambda_i, \lambda) > 0\} = \sup\{\lambda : \theta(\lambda_i, \lambda) = 0\}.$$

The research program reported here started with the question of whether $\lambda_*^i(\lambda_e)$ would be strictly larger than λ_c for every $\lambda_e < \lambda_c$. This question resembles a typical question of enhancement: does a decrease in λ_e , however small, have a strong enough effect on the dynamics so as to cause the critical value of λ_i to increase? For percolation systems with two parameters, there are well established techniques that can often be used to prove that the critical value for one of the parameters is a strictly monotone function of the other parameter (see [1] and [9, Section 3.3]). However, these techniques are much better adapted to the non-oriented case, and will usually break down for the contact process and other oriented models. For instance, in the classic contact process, we are not aware of a proof that the critical parameter is strictly decreasing as we increase the spatial dimension (although $\lambda_c \to 0$ as $d \to \infty$ [8, 10]). Enhancement arguments being of no help, we have to rely on other methods which depend on the one-dimensional properties of our processes.

The following were proved in [5] and are illustrated in Figure 1.1b.

Theorem 1. For $\lambda_e > \lambda_c$, we have $\lambda^i_*(\lambda_e) = \lambda_c$.

Theorem 2. For $\lambda_e > 1$, we have $\lambda_*^i(\lambda_e) < \infty$.

Corollary 1.1. $\lim_{\lambda_i \to \infty} \lambda^e_*(\lambda_i) = 1.$

We now state the main results of this paper, illustrated in Figure 1.1c.



Figure 1.1: (a) Simple properties of the phase space: $\theta(\lambda_i, \lambda_e) > 0$ if $\min\{\lambda_i, \lambda_e\} > \lambda_c$ and $\theta(\lambda_i, \lambda_e) = 0$ if $\max\{\lambda_i, \lambda_e\} \leq \lambda_c$ or $\lambda_e \leq 1$. (b) Results of [5]. (c) Results of this paper. (color online)

The first and second theorems below state that the critical value of λ_i is a strictly decreasing function of λ_e in the relevant region, and vice versa. As a consequence, the critical value of either parameter is a continuous function of the other parameter.

Theorem 3. The function $\lambda_*^i(\lambda_e)$ is strictly decreasing for $\lambda_e \in (1, \lambda_c]$.

In particular, we answer the original question affirmatively:

Corollary 1.2. For $\lambda_e < \lambda_c$, we have $\lambda^i_*(\lambda_e) > \lambda_c$.

Theorem 4. The function $\lambda_*^e(\lambda_i)$ is strictly decreasing on $[\lambda_c, +\infty)$.

Corollary 1.3. For all $\lambda_i < \infty$ we have $\lambda_*^e(\lambda_i) > 1$.

Corollary 1.4. For $\lambda_e = \lambda_c$, we have $\lambda_*^i(\lambda_e) = \lambda_c$.

The following corollary is a consequences of Theorems 3 and 4. For the first two parts note that if one of the functions λ_*^e and λ_*^i is strictly monotone the other one must be continuous. Then these functions are inverse of each other and the last two parts of the corollary follow.

Corollary 1.5. The critical curve has the following properties.

- (i) The function $\lambda_*^e : [\lambda_c, \infty) \to (1, \lambda_c]$ is continuous.
- (ii) The function $\lambda_*^i: (1, \lambda_c] \to [\lambda_c, \infty)$ is continuous.
- (*iii*) $\lim_{\lambda_e \to 1} \lambda^i_*(\lambda_e) = +\infty.$
- (iv) For every $a \in [\lambda_c, \infty)$ and $b \in (1, \lambda_c]$, $\lambda^e_*(a) = b$ if and only if $\lambda^i_*(b) = a$.

Remark 1.6. In the course of proving Theorems 3 and 4, we show existence and uniqueness of a stationary state for the process seen from the rightmost infected site, as well as convergence to such state, for supercritical pairs of parameters (Proposition 4.1).

Corollary 1.4 is somehow expected, it says that a process with supercritical parameter in the bulk and critical parameter at the boundary will survive with positive probability. Perhaps more surprising is its counterpart that we state now.

Theorem 5. For $\lambda_i = \lambda_c$, we have $\lambda^e_*(\lambda_i) = \lambda_c$.

We make a few remarks about the above theorem.

Remark 1.7. We will in fact prove something slightly stronger: if the underlying graph is \mathbb{Z}_+ instead of \mathbb{Z} , and the infection rate is $\lambda_c + \varepsilon$ at the right-hand side boundary and λ_c everywhere else, then the process survives.

Remark 1.8. Even though the theorem says that the process survives for $\lambda_i = \lambda_c$ and $\lambda_e > \lambda_c$, the process does not have a non-trivial invariant measure. Indeed, if it had such a measure, it would be supported on configurations that have infinitely many occupied sites in both directions and would thus be invariant for the standard contact process with critical parameter, contradicting a.s. extinction established in [3].

Remark 1.9. The critical contact process seen from the rightmost infected site has a unique stationary state [4]. In the course of proving Theorem 5, we prove that the critical contact process started from this distribution has zero asymptotic speed (Proposition 4.7).

Remark 1.10. It is conjectured that the critical contact process dies out even if we change the infection rate to an arbitrarily large value at a finite number of edges. This conjecture is supported by an analogous result in non-oriented percolation [16]. In contrast, the above theorem says that, if at each time we increase the infection rate by $\varepsilon > 0$ at a specific, dynamically chosen, pair of edges (those at the boundary of the infected interval), then the process survives with positive probability.

Regarding the conjecture mentioned in Remark 1.10, we obtain a partial result to be contrasted with Theorem 5 and Remark 1.7.

Theorem 6. Consider the standard contact process on \mathbb{Z}_+ , with recovery rate 1 and infection rate λ_c at all but finitely many sites, and having at finitely many sites a recovery rate $\delta > 0$ and an infection rate $\lambda < \infty$ arbitrary but fixed. This process dies out a.s.

Remark 1.11. Our proof relies strongly on the underlying graph being \mathbb{Z}_+ rather than \mathbb{Z} . This creates one of those situations where, although the full conjecture cannot be mathematically established, partial results make its negation more and more implausible (as is the case for the $\theta(p_c) = 0$ conjecture in Bernoulli percolation). In the present context, the above theorem says that, for the analogous process on \mathbb{Z} , if the infection survives then *every* infinite infection path has to visit sites that are arbitrarily far from the origin *in both directions*.

We briefly ention some challenges faced in proving Theorem 5. Since this is no longer in the regime $\lambda_e \leq \lambda_i$, process under consideration is not attractive. So there is no simple way to compare different configurations. As an anecdote, we had previously obtained several "proofs" for Theorem 5 that turned out to have subtle flaws in them. In order to prove that the process with parameters $\lambda_i = \lambda_c$ and $\lambda_e = \lambda_c + \varepsilon$ survives, we compare it with a process with parameters $\lambda_e = \lambda_i = \lambda_c$, which is attractive and has zero speed (Proposition 4.7). Due to lack of attractiveness, we cannot use subadditivity to study growth speed either, and, in fact, it is not known whether the process has an asymptotic speed at all. To overcome these problems, we perform a kind of "restart" each time an extra infected site is added to the former process. At each such restart, we perform a thinning of the previous process, obtaining a configuration distributed as the invariant measure for the critical process seen from the rightmost infected site [4], the domination between the concerned measures being provided by Proposition 4.6. This sequence of restarted critical processes is carefully constructed so as make it stationary on the one hand, and in some sense comparable with the invariant measure on the other hand. From stationarity, we can conclude that this auxiliary process

has a speed a.s., and that the speed is positive with non-zero probability. From there we conclude that the non-restarted process has positive lower speed with positive probability, which fortunately is enough to conclude that the process survives.

The remaining sections are organized as follows. Section 2 introduces the notation and some basic definitions. Section 3 contains results concerning the asymptotic speed in the attractive regime $\lambda_e \leq \lambda_i$. In Section 4 we obtain a series of results for the process starting from semi-infinite configurations seen from the edge, which may be of independent interest. In Section 5 we show that the asymptotic speed is 0 when the pair (λ_i, λ_e) is critical. In Sections 6 and 7, we prove Theorems 3 and 4. Section 8 is devoted to the proof of Theorem 5, it uses results from Section 4 but it is otherwise independent of the other sections as it considers the non-attractive regime. Section 9 expands the arguments of Section 8 to prove Remark 1.7. Section 10 is independent of all the others and gives a proof of Theorem 6. Section 11 states some open problems.

2 Terminology, definitions and notation

Configurations are subsets of \mathbb{Z} represented by capital letters A and B if they are fixed, or η, ζ, ξ if they are random. The set of configurations is denoted $\Sigma = \mathcal{P}(\mathbb{Z})$. The set of semi-infinite configurations (to the left) is denoted Σ^{\ominus} . The shift to the right is defined as $TA = A + 1 \subseteq \mathbb{Z}$. For $A \in \Sigma$, we define $\mathcal{R}A = \sup A$ and $\mathcal{L}A = \inf A$.

Throughout this paper we use an enlarged version of the standard graphical construction of the contact process, which can handle all values of λ_i , λ_e and ε simultaneously. The relevant features of this graphical construction are that each space-time point (x, t) can have a recovery mark and each oriented edge from (x, t) to $(x \pm 1, t)$ can be λ -open for a given value of λ , in a way that λ -open edges are λ' -open for all $\lambda' \ge \lambda$. We also have an extra clock which is not attached to any specific site (it can be used to account for an increase in one of the parameters). We let \mathbb{P} denote an underlying probability measure in a space where these elements are defined.

Here is a possible implementation. Sample for each $x \in \mathbb{Z}$ a PPP (Poisson Point Process) $\omega_x \subseteq (0, +\infty)$ of intensity 1. We say that there is a recovery mark at (x, t) if $t \in \omega_x$. For each oriented edge from (x, t) to $(x \pm 1, t)$, sample a PPP $\omega_{x,x\pm 1} \subseteq (0, +\infty) \times (0, +\infty)$ with intensity 1, and say that this edge is λ -open if $\omega_{x,x\pm 1} \cap (\{t\} \times (0, \lambda]) \neq \emptyset$. Finally sample a PPP $\omega_{\varepsilon} \subseteq (0, +\infty)$ of intensity ε , and say that there is an ε -mark at time t if $t \in \omega_{\varepsilon}$.

Given $A \subseteq \mathbb{Z}$, a pair (λ_i, λ_e) , and $s \ge 0$, we construct the process $(\eta_{s,t}^A)_{t\ge s}$ as follows. At time $s, \eta_{s,s}^A = A$. The process jumps from B to $B \setminus \{x\}$ at time t if $x \in B$ and $t \in \omega_x$; it jumps to $B \cup \{x+1\}$ if $x \in B$ and either $x \ne \mathcal{R}B$ and the edge from (x,t) to (x+1,t) is λ_i -open or $x = \mathcal{R}B$ and the edge from (x,t) to (x+1,t) is λ_i -open or $x = \mathcal{R}B$ and the edge from (x,t) to (x+1,t) is λ_i -open or $x = \mathcal{L}B$ and the edge from (x,t) to (x-1,t) is λ_i -open or $x = \mathcal{L}B$ and the edge from (x,t) to (x-1,t) is λ_i -open or $x = \mathcal{L}B$ and the edge from (x,t) to (x-1,t) is λ_i -open or $x = \mathcal{L}B$ and the edge from (x,t) to (x-1,t) is λ_i -open or $x = \mathcal{L}B$ and the edge from (x,t) to (x-1,t) is λ_i -open or $x = \mathcal{L}B$ and the edge from (x,t) to (x-1,t) is λ_i -open or $x = \mathcal{L}B$ and the edge from (x,t) to (x-1,t) is λ_i -open or $x = \mathcal{L}B$ and the edge from (x,t) to (x-1,t) is λ_i -open or $x = \mathcal{L}B$ and the edge from (x,t) to (x-1,t) is λ_i -open or $x = \mathcal{L}B$ and the edge from (x,t) to (x-1,t) is λ_i -open or $x = \mathcal{L}B$ and the edge from (x,t) to (x-1,t) is λ_i -open. These rules a.s. define $\eta_{s,t}^A$ for all $t \ge s$ when A is finite. For infinite A, we define $\eta_{s,t}^A$

by taking a limit $A_n \uparrow A$. When the parameters need to be specified, we write $\eta^A_{s,t,\lambda_i,\lambda_e}$. Finally, if s = 0, we may omit it and write η^A_t .

For given $A \in \Sigma$, this defines a process $(\eta_t^A)_{t \ge 0}$ started from configuration $\eta_0 = A$, informally described in the previous section. Process $(\eta_t^-)_t$ starts from $\eta_0 = \mathbb{Z}_-$, $(\eta_t^+)_t$ starts from $\eta_0 = \mathbb{Z}_+$, $(\eta_t^x)_t$ starts from $\eta_0 = \{x\}, (\eta_t^0)_t$ starts from $\eta_0 = \{0\}, (\eta_t^\nu)_t$ starts from random η_0 with distribution ν . We reserve the letter μ for the distribution provided by Proposition 4.1.

Processes denoted by the letter η always evolve with parameters (λ_i, λ_e) and always use the same graphical construction, differing only in the starting time and initial configuration. Processes with different rules will be denoted with letters ξ or ζ , depending on the section. In a slight abuse of notation, the superscript in $\xi_t^1, \xi_t^2, \ldots, \xi_t^n, \ldots$ may be used to index a sequence of processes rather than specify the initial condition.

Remark 2.1. $\mathbb{P}(\eta_t^A \neq \emptyset \ \forall t)$ is either positive for all finite non-empty A or zero for all finite non-empty A.

A path is a discrete càdlàg function $\gamma : [t_0, t_1] \to \mathbb{Z}$ whose jumps are of length 1. We say that a path γ defined in the time interval [0, t] is (λ_i, λ_e) -active for an initial configuration A if its jumps coincide with λ_i or λ_e -open edges and $\gamma_s \in \eta_s^A$ for all $s \in [0, t]$. Due to the nearest neighbor property of the jumps, if there is a (λ_i, λ_e) -active path from $B \subset \mathbb{Z}$ to $C \subset \mathbb{Z}$ for a given initial configuration B, then there exits a rightmost (λ_i, λ_e) -active path joining B to C in the same time interval.

3 Attractiveness and asymptotic speed

We start this section with a basic lemma:

Lemma 3.1. On the octant $\{\lambda_i \ge \lambda_e \ge 0\}$, the configuration $\eta^A_{s,t,\lambda_i,\lambda_e}$ is increasing in A, λ_i and λ_e for fixed $t \ge s \ge 0$.

The lemma follows immediately from the construction described in §2, since infecting more sites can only affect the infection rates positively, by turning some external edges into internal ones.

Remark 3.2. For the same reason, the process is not attractive when $\lambda_e > \lambda_i$. To see this, consider $A = \{0\}$ and $B = \{3\}$, then for small values of t we have $\mathbb{P}(1 \in \eta_t^A) > \mathbb{P}(1 \in \eta_t^{A \cup B})$.

Remark 3.3. When $\lambda_e < \lambda_i$, the process is attractive but not additive. Indeed, with the same A and B as above, we have $\mathbb{P}(1 \in \eta_t^{A \cup B}) > \mathbb{P}(1 \in \eta_t^A) + \mathbb{P}(1 \in \eta_t^B)$ for small t, hence there is no coupling such that $\eta_t^{A \cup B} = \eta_t^A \cup \eta_t^B$, even though $\eta_t^{A \cup B} \supseteq \eta_t^A \cup \eta_t^B$.

Since many properties of the contact process are usually derived from its additivity, we have to modify or reinvent some of their proofs. Fortunately, for $\lambda_i \ge \lambda_e$, attractiveness is enough to use the Subadditive Ergodic Theorem as in [11, Chapter VI]. This allows us to work with the asymptotic

speed of the boundary of the process. With this in mind, we define

$$\alpha_t(\lambda_i, \lambda_e) = \mathbb{E}[\mathcal{R}\eta_t^-]$$

and

$$\alpha(\lambda_i, \lambda_e) = \inf_{t>0} \frac{\alpha_t(\lambda_i, \lambda_e)}{t}.$$

Lemma 3.4. Let $0 < \lambda_e \leq \lambda_i < \infty$ be fixed. Then,

$$\frac{\mathcal{R}\eta_t^-}{t} \xrightarrow{a.s.} \alpha(\lambda_i, \lambda_e) = \lim_{t \to \infty} \frac{\alpha_t(\lambda_i, \lambda_e)}{t}.$$

Thanks to Lemma 3.1, the proof is the same as in [11, VI.2.19].

Lemma 3.5. Suppose $\lambda_e \leq \lambda_i$ and $\theta(\lambda_i, \lambda_e) > 0$. Let $B \in \Sigma^{\ominus}$. Then,

$$\frac{\mathcal{R}\eta^B_t}{t} \xrightarrow{a.s.} \alpha(\lambda_i, \lambda_e).$$

Proof. Write $\alpha = \alpha(\lambda_i, \lambda_e)$. First note that, if $\eta_t^0 \neq \emptyset$, then $\mathcal{R}\eta_t^- = \mathcal{R}\eta_t^0$ (this holds because both processes are defined from the same graphical construction and infections are to nearest-neighbor only). Hence, a.s., if $\eta_t^x \neq \emptyset$ for all $t \ge 0$, then $\lim_t t^{-1}\mathcal{R}\eta_t^x = \alpha$ by the previous lemma. Now since $\theta(\lambda_i, \lambda_e) > 0$ and B has infinitely many points, a.s. there exists $x \in B$ such that $\eta_t^x \neq \emptyset$ for all $t \ge 0$, which implies that $\liminf_t t^{-1}\mathcal{R}\eta_t^B \ge \lim_t t^{-1}\mathcal{R}\eta_t^x = \alpha$ because $\{x\} \subseteq B$.

On the other hand, we can assume $B \subseteq \mathbb{Z}_{-}$ without loss of generality, and from this we get $\limsup_{t} t^{-1} \mathcal{R} \eta_t^B \leq \lim_{t} t^{-1} \mathcal{R} \eta_t^- = \alpha$, a.s.

Lemma 3.6. The function $\alpha(\cdot, \lambda_e)$ is right continuous on $[\lambda_e, \infty)$, and the function $\alpha(\lambda_i, \cdot)$ is right continuous on $[0, \lambda_i]$.

Proof. For all t > 1, the function $t^{-1}\alpha_t(\cdot, \lambda_e)$ is continuous. Hence, $\alpha(\cdot, \lambda_e)$ is upper semicontinuous. By Lemma 3.1, it is non-decreasing, and therefore it is right-continuous. The right continuity of $\alpha(\lambda_i, \cdot)$ is proved in exactly the same way. (Cf. [11, VI.2.27b].)

Lemma 3.7. Suppose $\lambda_e \leq \lambda_i$ and $\theta(\lambda_i, \lambda_e) > 0$. Then $\alpha(\lambda_i, \lambda_e) \geq 0$.

Proof. By Lemma 3.1, $\eta_t^0 \subseteq \eta_t^-$ and $\eta_t^0 \subseteq \eta_t^+$ for all t. If $\alpha(\lambda_i, \lambda_e) < 0$, by Lemma 3.5, we have $\eta_t^- \cap \eta_t^+ = \emptyset$ for all large t, almost surely, which by the previous sentence implies that $\eta_t^0 = \emptyset$ for large t, and therefore $\theta(\lambda_i, \lambda_e) = 0$. (Cf. [11, VI.2.27a].)

Theorem 7. For $1 < \lambda_e \leq \lambda_c$, we have that $\alpha(\lambda_*^i(\lambda_e), \lambda_e) = 0$. For $\lambda_i \geq \lambda_c$, we have $\alpha(\lambda_i, \lambda_*^e(\lambda_i)) = 0$.

We give the proof in §5.

Proposition 3.8. Suppose $\theta(\lambda_i, \lambda_e) > 0$, and $\lambda_e + \varepsilon \leq \lambda_i$. Then $\alpha(\lambda_i, \lambda_e + \varepsilon) \geq \alpha(\lambda_i, \lambda_e) + \varepsilon$.

We give the proof in §6.

Recall Theorem 2 which says that $\lambda^i_*(\lambda_e) < \infty$ for $\lambda_e > 1$.

Corollary 3.9. Let $1 < \lambda_e < \lambda'_e \leq \lambda_c$. Then $\alpha(\lambda^i_*(\lambda_e), \lambda'_e) > 0$.

Proof. Let $1 < \lambda_e < \lambda'_e < \lambda_c$ and write $\varepsilon = \lambda'_e - \lambda_e$. Take $\lambda_i = \lambda^i_*(\lambda_e) \ge \lambda_c > \lambda'_e$. For $\delta > 0$, we have $\theta(\lambda_i + \delta, \lambda_e) > 0$. By Proposition 3.8 and Lemma 3.7, $\alpha(\lambda_i + \delta, \lambda'_e) \ge \varepsilon$. Letting $\delta \downarrow 0$, by Lemma 3.6 we get $\alpha(\lambda_i, \lambda'_e) \ge \varepsilon$.

Proposition 3.10. Let $\lambda'_i > \lambda_i > \lambda_c$. Then $\alpha(\lambda'_i, \lambda^e_*(\lambda_i)) > 0$.

We give the proof in 7.

Proof of Theorems 3 and 4. Suppose that $1 < \lambda_e < \lambda'_e \leq \lambda_c$ and take $\lambda_i = \lambda^i_*(\lambda_e)$ (note that $\lambda_i \geq \lambda_c > 0$ and, by Theorem 2, $\lambda_i < \infty$). By Corollary 3.9, $\alpha(\lambda_i, \lambda'_e) > 0$. By Theorem 7, $\lambda_i > \lambda^i_*(\lambda'_e)$, since otherwise we would have $\alpha(\lambda_i, \lambda'_e) = 0$. So $\lambda^i_*(\lambda_e) > \lambda^i_*(\lambda'_e)$, proving Theorem 3. Note that $\lambda^e_*(\cdot)$ is non-increasing on $[\lambda_c, +\infty)$ by attractiveness, since $\lambda^e_*(\cdot) \leq \lambda_c$ on this interval. Let $\lambda'_i > \lambda_i > \lambda_c$ and take $\lambda_e = \lambda^e_*(\lambda_i)$ (note that $1 \leq \lambda_e \leq \lambda_c$). By Proposition 3.10, $\alpha(\lambda'_i, \lambda_e) > 0$. By Theorem 7, $\lambda_e > \lambda^e_*(\lambda'_i)$, since otherwise we would have $\alpha(\lambda'_i, \lambda_e) = 0$. So $\lambda^e_*(\lambda_i) > \lambda^e_*(\lambda'_i)$, proving Theorem 4.

4 The process seen from the edge

In this section we obtain a series of results for the process starting from semi-infinite configurations, seen from the edge. These results will be used in subsequent sections and may be of independent interest.

We start with a little bit of notation. Define $\Sigma^{\odot} = \{A \in \Sigma^{\ominus} : \mathcal{R}A = 0\}$. We also define the map $\Psi : \Sigma^{\ominus} \to \Sigma^{\odot}$ by $\Psi A = T^{-\mathcal{R}A}A$, and it is convenient to define $\Psi A = \emptyset$ for $A = \emptyset$. We define the addition of a site $\Xi : \Sigma^{\ominus} \to \Sigma^{\ominus}$ by $\Xi \eta = \eta \cup \{\mathcal{R}\eta + 1\}$. A real-valued function f on Σ will be called *increasing* if $f(A) \leq f(B)$ whenever $A \subseteq B$. If η and ξ are random elements of Σ , we say that η stochastically dominates ξ , denoted $\eta \succeq \xi$, if $\mathbb{E}[f(\eta)] \ge \mathbb{E}[f(\xi)]$ for every bounded increasing measurable function f.

In subsequent sections, we will use the propositions below, which can be of independent interest.

Proposition 4.1. Suppose $\lambda_e \leq \lambda_i$ and $\theta(\lambda_i, \lambda_e) > 0$, or $\lambda_i = \lambda_e = \lambda_c$. Then there exists a measure μ supported on Σ^{\odot} such that $\Psi \eta_t^{\mu} \sim \mu$ for every $t \geq 0$, and $\Psi \eta_t^A \rightarrow \mu$ weakly for every fixed $A \in \Sigma^{\ominus}$.

Throughout this paper, the letter μ refers to the above measure. The case $\lambda_e = \lambda_i = \lambda_c$ is [4, Theorem 1]. So we have prove Proposition 4.1 for $\lambda_e \leq \lambda_i$ and $\theta(\lambda_i, \lambda_e) > 0$. We will use the following lemma.

Lemma 4.2. Let $\lambda_i > \lambda_c$ and $\lambda_e \ge 0$. Then for all $t \ge 0$,

 $\Psi\eta_t^- \succcurlyeq \zeta \cap \mathbb{Z}_-,$

where ζ denotes a random configuration distributed as the unique non-trivial invariant measure for the standard contact process with parameter $\lambda_i > \lambda_c$.

Proof. Fix $t \ge 0$, and let Γ denote the rightmost active path connecting η_0^- at time 0 to η_t^- at time t, similar to the path used in the proof of Proposition 4.6. For a fixed path γ , the event $\Gamma = \gamma$ is determined by the graphical construction ω on γ and to the right of γ . So, given that $\Gamma = \gamma$, the conditional distribution of graphical construction to the left of γ remains the same, and the configuration η_t^- is given by the set of sites $x \le \gamma_t$ such that there is a backwards λ_i -open path from (x, t) to either γ or $\mathbb{Z} \times \{0\}$. This dominates the set of sites $x \le \gamma_t$ that are connected to $-\infty$ by backwards λ_i -open path, and translating this set by $-\gamma_t$ we get a set distributed exactly like $\zeta \cap \mathbb{Z}_-$. On the other hand, translating η_t^- by the same amount we get $\Psi \eta_t^-$, which shows the lemma.

Proof of Proposition 4.1. Note that the transition probabilities of the process $(\Psi \eta_t)_t$ are discontinuous at finite configurations. However, it is still true that, for every $t \ge 0$ and every continuous bounded $f: \Sigma \to \mathbb{R}$, the functional $S_t f$ given by $A \mapsto \mathbb{E}[f(\Psi \eta_t^A)]$ is continuous at points $A \in \Sigma^{\odot}$.

Consider the law of $\Psi \eta_t^-$ at time $t \in \mathbb{N}$. Since Σ is a compact metric space there is a subsequence $(t_k)_k$ such that $t_k \to +\infty$ and $\Psi \eta_{t_k}^- \to \mu$ weakly as $k \to \infty$, for some probability measure μ on Σ . The construction in [6, §3] then gives

$$\Psi \eta_t^A \to \mu$$
 weakly as $t \to \infty$, for every $A \in \Sigma^{\odot}$. (4.3)

It is thus enough to show that

$$\mu(\Sigma^{\odot}) = 1 , \qquad (4.4)$$

since these two properties, together with the fact that $S_t f$ is continuous on Σ^{\odot} for every continuous $f: \Sigma \to \mathbb{R}$, will imply that $\Psi \eta_t^{\mu} \sim \mu$ for every $t \ge 0$.

But (4.4) is a consequence of (4.3) and Lemma 4.2, and this concludes the proof of Proposition 4.1 for $\lambda_e \leq \lambda_i$ and $\theta(\lambda_i, \lambda_e) > 0$.

Proposition 4.5. Suppose $\lambda_e \leq \lambda_i$ and $\theta(\lambda_i, \lambda_e) > 0$. Then

$$\mathbb{E}[\mathcal{R}\eta_t^{\mu}] = \alpha(\lambda_i, \lambda_e) \cdot t$$

for all $t \ge 0$, where μ is given by Proposition 4.1.

Proof. Note that $\mathcal{R}\eta_t^{\mu}$ has stationary increments. Thus $t^{-1}\mathbb{E}[\mathcal{R}\eta_t^{\mu}] = \mathbb{E}[\mathcal{R}\eta_1^{\mu}] \in [-\infty, \lambda_e]$. Moreover, by the Ergodic Theorem, $n^{-1}\mathcal{R}\eta_n^{\mu} \to V$ a.s., for (possibly random) V with $\mathbb{E}V = \mathbb{E}[\mathcal{R}\eta_1^{\mu}]$. On the other hand, by Lemma 3.5, $n^{-1}\mathcal{R}\eta_n^{\mu} \to \alpha(\lambda_i, \lambda_e)$ a.s., concluding the proof.

Proposition 4.6. Let $\lambda_e \leq \lambda_i$ and suppose $\theta(\lambda_i, \lambda_e) > 0$, or that $\lambda_e = \lambda_i = \lambda_c$. If ζ denotes a random configuration in Σ^{\odot} with law μ , then $\Xi \zeta \geq T \zeta$.

Proof. It follows from Proposition 4.1 that $\Psi \eta_t^- \to \mu$ in distribution as $t \to \infty$. Since Ξ and T are continuous, it suffices to show that $\Xi \Psi \eta_t^- \geq T \Psi \eta_t^-$ for all $t \geq 0$. So let t be fixed. Let Γ denote the rightmost active path for the initial configuration \mathbb{Z}_- in the time interval [0, t].

For a deterministic path $(\gamma_s)_{s\in[0,t]}$, let $D_{\gamma}^{\ominus} = \{(x,s): s\in[0,t], x<\gamma_s\wedge\gamma_{s-}\}, D_{\gamma}^{\odot} = \{(x,s)\in D_{\gamma}^{\ominus}: s=0 \text{ or } x=\gamma_s\wedge\gamma_{s-}\}$ and $D_{\gamma}^{\oplus} = \{(x,s): s\in[0,t], x\geqslant\gamma_s\wedge\gamma_{s-}\}.$

On the event $\Gamma = \gamma$, the configuration η_t is given by the set of sites that can be reached from D_{γ}^{\odot} within D_{γ}^{\ominus} via a λ_i -open path. On the other hand, the event $\Gamma = \gamma$ is determined by the graphical construction ω on the region D_{γ}^{\oplus} . Hence, the conditional law of η_t given that $\Gamma = \gamma$ is the law of the set of sites that can be reached from D_{γ}^{\odot} within D_{γ}^{\ominus} , without conditioning. We denote this law by ν^{γ} .

Define the re-centered curve $(\bar{\gamma}_s)_{s\in[0,t]}$ by $\bar{\gamma}_s = \gamma_s - \gamma_t$. Since $\Gamma_t = \mathcal{R}\eta_t^-$, by translation invariance, the conditional distribution of $\Psi\eta_t^-$ given that $\Gamma = \gamma$ equals $\nu^{\bar{\gamma}}$.

We now consider the configuration $\Xi \eta_t^-$, which equals $\eta_t^- \cup \{\mathcal{R}\eta_t^- + 1\}$. To give a similar description as before, we define the path $(\gamma_s^+)_{s \in [0,t]}$ given by

$$\gamma_s^+ = \begin{cases} \gamma_s, & s \in [0, t), \\ \gamma_t + 1, & s = t. \end{cases}$$

As before, the conditional law of $\Xi \eta_t^-$ given that $\Gamma = \gamma$ is the law of the set of sites that can be reached from $D_{\gamma^+}^{\odot}$ within $D_{\gamma^+}^{\ominus}$, without conditioning. Also as before, we define the re-centered curve $(\bar{\gamma}_s^+)_{s \in [0,t]}$ by $\bar{\gamma}_s^+ = \gamma_s^+ - \gamma_t^+$. Since $\gamma_t^+ = \mathcal{R}(\Xi \eta_t^-)$, by translation invariance, the conditional distribution of $\Psi \Xi \eta_t^-$ given that $\Gamma = \gamma$ equals $\nu^{\bar{\gamma}^+}$.

To conclude the proof, we just need to notice that $\nu^{\bar{\gamma}^+} \succeq \nu^{\bar{\gamma}}$. This is true because $\bar{\gamma}^+ \leqslant \bar{\gamma}$. Indeed, $\bar{\gamma}_t^+ = \bar{\gamma}_t = 0$ and $\bar{\gamma}_s^+ = \bar{\gamma}_s - 1$ for $s \in [0, t)$.

Proposition 4.7. For
$$\lambda_i = \lambda_e = \lambda_c$$
, $\lim_{t \to \infty} \frac{\mathcal{R}\eta_t^{\mu}}{t} = \mathbb{E}[\mathcal{R}\eta_1^{\mu}] = \alpha(\lambda_c, \lambda_c) = 0$ a.s.

Proof. As in the proof of Proposition 4.5, there is $\beta \in [-\infty, \lambda_c]$ such that $\mathbb{E}[\mathcal{R}\eta_t^{\mu}] = \beta t$ for all $t \ge 0$ and $\frac{\mathcal{R}\eta_t^{\mu}}{t} \to V$ a.s., where V is such that $\mathbb{E}V = \beta$. However, here we cannot use Lemma 3.5. Below we will show that $V = \beta$ a.s. and that $\beta = \alpha$, where $\alpha = \alpha(\lambda_c, \lambda_c)$. Since $\eta_t^{\mu} \subseteq \eta_t^-$, we have

$$V = \lim_{t} \frac{\mathcal{R}\eta_t^{\mu}}{t} \leqslant \lim_{t} \frac{\mathcal{R}\eta_t^{-}}{t} = \alpha \quad \text{a.s.},$$

where the last limit holds by Lemma 3.4. So it is enough to prove that $\beta \ge \alpha$.

For $A \in \Sigma^{\ominus}$, let SA denote the distance between the rightmost and second rightmost site in A. We first note that

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathbb{E}[\mathcal{R}\eta_t^\nu] = \lambda_c - \mathbb{E}[\mathcal{S}\eta_t^\nu]$$

for every measure ν on Σ^{\ominus} . More precisely, if $\mathbb{E}|\mathcal{R}\eta_t^{\nu}| < \infty$ and $\mathbb{E}[\mathcal{S}\eta_t^{\nu}] < \infty$, then the above equation holds, and if $\mathbb{E}[\mathcal{R}\eta_t^{\nu}] = -\infty$ then $\mathbb{E}[\mathcal{R}\eta_s^{\nu}] = -\infty$ for all s > t.

Recall that, by Lemma 3.4, $\frac{\mathbb{E}[\mathcal{R}\eta_t^-]}{t} \to \alpha$. Since $\Psi\eta^- \to \mu$ in distribution as $t \to \infty$ by Proposition 4.1, and since $\mathbb{E}|\mathcal{R}\eta_t^-| < \infty$ for all $t \ge 0$, we have

$$\limsup_{t} \frac{\mathrm{d}}{\mathrm{d}t} \mathbb{E}[\mathcal{R}\eta_t^-] = \lambda_c - \liminf_{t} \mathbb{E}[\mathcal{S}\eta_t^-] \leqslant \lambda_c - \mathbb{E}[\mathcal{S}\eta_t^\mu] = \beta,$$

using Fatou's Lemma. In particular,

$$\alpha = \lim_t \frac{\mathbb{E}[\mathcal{R}\eta_t^-]}{t} \leqslant \beta$$

concluding the proof.

5 Zero speed at criticality

In this section we prove Theorem 7. It is possible to prove this theorem rather easily once it is known that $\theta(\lambda_i, \lambda_*^e(\lambda_i)) = 0$ and $\theta(\lambda_*^i(\lambda_e), \lambda_e) = 0$. This is proved in Chapter 2 of [14], where the proof is an adaptation of [3]. A more general result of this nature is available in [2]. However all these references rely on the rather involved dynamic renormalization technique. For this reason we provide a different proof.

The general structure of the proof is a classical block argument. However, the specifics of this model bring a number of complications. In particular, the building block is more subtle to define and analyze because whether the infection spreads through a given path may depend on the configuration outside this path.

Assuming that $\lambda_e \leq \lambda_i$ and $\alpha(\lambda_i, \lambda_e) > 0$, we will derive a finite condition (a condition that depends on a bounded space-time box in the graphical construction) which in turn implies that $\theta(\lambda_i, \lambda_e) > 0$. Since said condition is finite, it is still satisfied for slightly smaller values of the parameters. This way we can conclude that $\lambda_e > \lambda_*^e(\lambda_i)$ and $\lambda_i > \lambda_*^i(\lambda_e)$, and this chain of implications combined with Lemmas 3.6 and 3.7 proves the theorem.

We call *domain* any subset $D \subseteq \mathbb{Z} \times \mathbb{R}$ of the form

$$\{(x,t): t_0 \leqslant t \leqslant t_1, L(t) \leqslant x \leqslant R(t)\}$$

for some $t_0 \leq t_1 \in \mathbb{R}$ and $L, R : [t_0, t_1] \to \mathbb{Z}$ such that $L(t) \leq R(t)$ for all $t \in (t_0, t_1]$. The bottom of D is given by the region $\{(x, t_0) : L(t_0) \leq x \leq R(t_0)\}$, the top is defined analogously.

Definition 5.1 (Open branch). A (λ_i, λ_e) -open branch \mathcal{O} in a domain D is a collection of paths γ contained in D such that each path $\gamma \in \mathcal{O}$ satisfies:

- (i) γ starts at the bottom of D,
- (*ii*) γ contains no recovery marks,
- (*iii*) there is a λ_i -mark at each jump of γ ,
- (*iv*) there is a λ_e -mark at each jump of γ that points in a direction not occupied by other paths in \mathcal{O} .

For each $t \in (t_0, t_1]$ denote $\mathcal{O}_t = \{x : \exists \gamma \in \mathcal{O}, \gamma(t) = x\}$. The points \mathcal{RO}_t and \mathcal{LO}_t denote the rightmost and leftmost points attained by this branch at time t. With this notation, condition (iv) above reads as follows: for each $\gamma \in \mathcal{O}$, if a given jump of γ corresponds to a jump of \mathcal{LO}_t to the left or a jump of \mathcal{RO}_t to the right, then there is a λ_e -mark corresponding to that jump of γ .

An open branch is *complete* if it has a path which starts at the bottom of D and reaches the top of D.

As a side remark, for $\lambda_e \leq \lambda_i$, a subset of an open branch need not be an open branch, but the



Figure 5.1: Open branches in D (left), D' (center), and the grafting of D' onto D (right). Blue paths only jump to the right, and are λ_e -open. Red paths only jump to the left, and are λ_e -open. Green paths jump in both directions and are λ_i -open. For each open branch, at each instant in time, the rightmost site of the branch is occupied by a blue path and the leftmost site of the branch is occupied by a red path. Thin light gray lines indicate t_0, t'_0, t_1, t'_1 . A grayed version of D behind D' illustrates their relative position, and the four gray dots illustrate conditions $R(t'_0) \leq L'(t'_0)$, and $R'(t_1) \leq L(t_1)$. (color online)

union of two open branches in D is an open branch.

We now describe a way to concatenate open branches, see Figure 5.1.

Definition 5.2 (Crossing domains and grafting). Given two domains D and D' (with associated t_0, t_1, L, R and t'_0, t'_1, L', R' , respectively), we say that D' crosses D from right to left if $t_0 \leq t'_0 \leq t_1 \leq t'_1, R(t'_0) \leq L'(t'_0)$, and $R'(t_1) \leq L(t_1)$. If D' crosses D from right to left, we define the grafting of D' onto D as the domain D'' defined as follows. Take $t''_0 = t_0, t''_1 = t'_1$,

$$L''(t) = \begin{cases} L(t), & t_0 \leqslant t \leqslant t'_0, \\ L(t) \land L'(t), & t'_0 < t < t_1, \\ L'(t), & t_1 \leqslant t \leqslant t'_1, \end{cases}$$

and

$$R''(t) = \begin{cases} R(t), & t_0 \le t < t'_0 \\ R(t) \lor R'(t), & t'_0 \le t \le t_1 \\ R'(t), & t_1 < t \le t'_1 \end{cases}$$

See Figure 5.1.

Lemma 5.3 (Grafting lemma). Suppose a domain D' crosses a domain D from right to left. For $\lambda_e \leq \lambda_i$, if there are complete open branches in D and in D', then there is a complete open branch in the grafting of D' onto D.

Proof. Let \mathcal{O} and \mathcal{O}' be complete open branches in D and D'. We will describe how to construct a complete open branch \mathcal{O}'' in the grafting D'' of D' onto D. A look at Figure 5.1 may help understand how \mathcal{O}'' is constructed.

Let $t_0 \leq t'_0 \leq t_1 \leq t'_1$ denote the start and end times of D and D'.

Fix some path $\gamma^* \in \mathcal{O}$ that starts at the bottom of D and ends at the top of D. For each $\gamma' \in \mathcal{O}'$, write I for its time span and define γ'' by taking, for each $t \in [t_0, t'_0] \cup I$,

$$\gamma''(t) = \begin{cases} \gamma^*(t), & t \leq t'_0, \\ \gamma^*(t) \land \gamma'(t), & t'_0 < t \leq t_1, \\ \gamma'(t), & t > t_1. \end{cases}$$

Let \mathcal{O}'' be the collection of all γ'' obtained this way, as well as all $\gamma \in \mathcal{O}$.

To prove the lemma, it is enough to check that \mathcal{O}'' is a complete open branch in D''. The key observation is that $\mathcal{LO}''_t = \mathcal{LO}_t \wedge \mathcal{LO}'_t$ and $\mathcal{RO}''_t = \mathcal{RO}_t$ for every $t \in [t'_0, t_1]$. Hence, each time \mathcal{LO}''_t jumps to the left, it corresponds to \mathcal{LO}_t or \mathcal{LO}'_t jumping to the left, and in either case there is a corresponding λ_e -open edge in the graphical construction. Likewise, jumps of \mathcal{RO}''_t to the right correspond to a λ_e -open edge. Other jumps correspond to a jump of a path in \mathcal{O} or \mathcal{O}' , which in turn occur at a λ_i -open edge. For times $t \leq t'_0$, we have $\mathcal{O}''_t = \mathcal{O}_t$, and for times $t > t_1$ we have $\mathcal{O}''_t = \mathcal{O}'_t$. It follows that \mathcal{O}'' satisfies condition (*iv*) above. The other conditions are more immediate to verify, and we omit the tedious details.

Given $\alpha \neq 0, k > 0$ and $L \in \mathbb{N}$, define the domain

$$D_{L,k,\alpha} := \left\{ (x,t) \in \mathbb{Z} \times \mathbb{R} : t \in [0, \frac{kL}{|\alpha|}], |x - \alpha t| \leq L \right\}.$$

Denote by $\mathscr{E}_{L,k,\alpha,\lambda_i,\lambda_e}$ the event that $D_{L,k,\alpha}$ has a complete (λ_i,λ_e) -open branch. Our building block is the following.

We leave k as a free parameter because the next lemma is already hard to visualize with $k = \frac{5}{2}$, but later on we will use it with k = 20.

Lemma 5.4. Suppose $\lambda_e \leq \lambda_i$ and $\alpha = \alpha(\lambda_i, \lambda_e) > 0$. Then, for each $k \in \mathbb{N}$ fixed, $\mathbb{P}(\mathscr{E}_{L,k,\alpha,\lambda_i,\lambda_e}) \to 1$ as $L \to \infty$, $L \in 4\mathbb{N}$.

Proof. Given $\varepsilon > 0$ and T > 0, let $\mathscr{E}'_{\varepsilon,T}$ denote the event that

$$|\mathcal{R}\eta_t^- - \alpha t| \leq \varepsilon T$$
 for all $t \in [0, T]$.

By Lemma 3.4, almost surely $|\mathcal{R}\eta_t^- - \alpha t| \leq \varepsilon t$ for large enough t. Hence, $\mathbb{P}(\mathscr{E}'_{\varepsilon,T}) \to 1$ as $T \to \infty$ for every $\varepsilon > 0$.



Figure 5.2: Illustration for n = 5 of how simultaneous occurrence of n + 1 similar events result in a complete open branch in $D_{4\ell,\frac{n}{2},\alpha}$. Green paths are λ_i -open, red and blue paths are λ_e -open. Existence of paths in the gray areas ensure that λ_i -open edges in between can be used. The green paths may be needed as they connect and lead to blue and red paths. (high resolution, color online)

Remark the following about $\mathscr{E}'_{\varepsilon,T}$. If this event occurs, then, for each $t \in [0,T]$, there exists a λ_i -open path γ from $\mathbb{Z}^- \times \{0\}$ to $(\mathcal{R}\eta^-_t, t)$ such that $\gamma_s \leq \mathcal{R}\eta^-_s$ for all $s \in [0,T]$ and each jump of γ to the right that coincides with a jump of $\mathcal{R}\eta^-_s$ to the right happens at a λ_e -open edge.

For convenience, let $\mathscr{E}'_{\varepsilon,T,x}$ denote the event that $|\mathcal{R}\eta^{(-\infty,x]}_t - \alpha t - x| \leq \varepsilon T$ for all $t \in [0,T]$. In words, $\mathscr{E}'_{\varepsilon,T,x}$ is the same event as $\mathscr{E}'_{\varepsilon,T}$ except that the condition is translated by x units in space to the right. Also, let $\mathscr{E}'_{\varepsilon,T}$ denote the event that $|\mathcal{L}\eta^+_t + \alpha t| \leq \varepsilon T$ for all $t \in [0,T]$. In words, $\mathscr{E}'_{\varepsilon,T}$ is the same event as $\mathscr{E}'_{\varepsilon,T}$ except that the condition is mirrored in space. Finally, let $\mathscr{E}'_{\varepsilon,T,x}$ denote the translation of $\mathscr{E}''_{\varepsilon,T}$ by x units in space to the right.

Given positive integers n and ℓ , take $\varepsilon = \frac{\alpha}{2n}$ and $T = \frac{2n\ell}{\alpha}$. Note that the event $\mathscr{E}'_{\varepsilon,T}$ implies that the random points $(\mathcal{R}\eta^-_t, t)$ are in $D_{\ell,2n,\alpha}$ for all $t \in [0,T]$. Combining several similar events together, we have

$$\mathscr{E}_{\varepsilon,T,3\ell}' \cap \mathscr{E}_{\varepsilon,T,\ell}'' \cap \mathscr{E}_{\varepsilon,T,5\ell}'' \cap \mathscr{E}_{\varepsilon,T,9\ell}'' \cap \cdots \cap \mathscr{E}_{\varepsilon,T,(4n-3)\ell}'' \subseteq \mathscr{E}_{4\ell,\frac{n}{2},\alpha,\lambda_i,\lambda_e}$$

That is, simultaneous occurrence of n+1 events of the form \mathscr{E}' and \mathscr{E}'' imply existence of a complete open branch in $D_{4\ell,\frac{n}{2},\alpha}$, as shown in Figure 5.2. In particular, $\mathbb{P}(\mathscr{E}_{4\ell,\frac{n}{2},\alpha,\lambda_i,\lambda_e}) \to 1$ as $\ell \to \infty$ for each n fixed.

Turning back to the statement of the lemma, we can take n = 2k and let $L = 4\ell \to \infty$, so we get $\mathbb{P}(\mathscr{E}_{L,k,\alpha,\lambda_i,\lambda_e}) \to 1$ as $L \to \infty$ in $4\mathbb{N}$, which proves the lemma.

Proof of Theorem 7. We are assuming that $\lambda_e \leq \lambda_i$ and $\alpha = \alpha(\lambda_i, \lambda_e) > 0$. Fix k = 20 and, given $L \in \mathbb{N}$, consider a grid made of copies of scissor-shaped regions, as in Figure 5.3a. Each scissor consists of a translation of $D_{L,k,\alpha}$ and a translation of $D_{L,k,-\alpha}$. (The choice of k = 20 is not optimal, but it makes the picture more clear, especially considering that the figure was produced on a discrete space-time grid.)

We say that a scissor is (λ_i, λ_e) -good if its blades contain complete (λ_i, λ_e) -open branches. Note that each scissor only overlaps with its 6 nearest-neighboring scissors (two below, two above, one on each side), and the goodness of scissors in any collection not having nearest-neighbors is distributed as



Figure 5.3: Good blocks and long paths.

a product measure. By [12], there is $\delta > 0$ such that, if $\mathbb{P}(\mathscr{E}_{L,k,\alpha,\lambda_i,\lambda_e}) > 1 - \delta$, then the collection of (λ_i, λ_e) -good scissors stochastically dominates an i.i.d. configuration which is supercritical for oriented site percolation, illustrated in Figure 5.3b. By Lemma 5.4, we can fix $L \in \mathbb{N}$ such that $\mathbb{P}(\mathscr{E}_{L,k,\alpha,\lambda_i,\lambda_e}) > 1 - \delta$. On the other hand, since $D_{L,k,\alpha}$ is finite, there exist $\lambda'_i < \lambda_i$ and $\lambda'_e < \lambda_e$ such that $\lambda'_e \leq \lambda'_i$ and $\mathbb{P}(\mathscr{E}_{L,k,\alpha,\lambda'_i,\lambda'_e}) > 1 - \delta$ (note that we keep the same α).

So with positive probability there is an infinite oriented path of (λ'_i, λ'_e) -good scissors, such that the bottom of the first scissor in this path is contained in $A \times \{0\}$, where $A := \{-6L, \ldots, 6L\} \subseteq \mathbb{Z}$. We now argue that, when this event occurs, $\eta^A_{0,t,\lambda'_i,\lambda'_e} \neq \emptyset$ for all $t \ge 0$. Indeed, suppose it occurs and let t > 0. Consider a scissor whose time span includes t and which belongs to this infinite oriented path of (λ'_i, λ'_e) -good scissors. By following this good path and applying Lemma 5.3 a finite number of times, we get a complete (λ'_i, λ'_e) -open branch in a domain whose bottom is contained in $A \times \{0\}$ and whose time span includes t. Hence, $\eta^A_{0,t,\lambda'_i,\lambda'_e} \neq \emptyset$ as claimed.

By Remark 2.1, $\theta(\lambda'_i, \lambda'_e) > 0$. Thus, $\lambda_e > \lambda^e_*(\lambda_i)$ and $\lambda_i > \lambda^i_*(\lambda_e)$ whenever $\lambda_e \leq \lambda_i$ and $\alpha(\lambda_i, \lambda_e) > 0$. 1. Hence, for $\lambda_e \leq \lambda_c$, since $\lambda^i_*(\lambda_e) \ge \lambda_c \ge \lambda_e$, we have $\alpha(\lambda^i_*(\lambda_e), \lambda_e) \leq 0$. Likewise, for $\lambda_i \ge \lambda_c$, 1. since $\lambda^e_*(\lambda_i) \le \lambda_c \le \lambda_i$, we have $\alpha(\lambda_i, \lambda^e_*(\lambda_i)) \le 0$. On the other hand, by Lemmas 3.6 and 3.7 we 1. have $\alpha(\lambda^i_*(\lambda_e), \lambda_e) \ge 0$ and $\alpha(\lambda_i, \lambda^e_*(\lambda_i)) \ge 0$, proving the theorem.

6 Speed increase with external infection rate

In the section we prove Proposition 3.8. A small increase in the external infection rate can be seen as a forced infection of the site immediately to the right of the current configuration, with some small but positive frequency. We want to show that this amounts to an increase in the growth speed. To take advantage of this extra infected site, we compare the effect of infecting one site with the effect of shifting the configuration to the right (and the latter obviously amounts to a positive increase in the speed). To achieve such a comparison (Proposition 4.6), we start from the invariant measure for the process seen from the rightmost infected site, provided by Proposition 4.1.

Notation. Throughout this section, letter η means that parameters (λ_i, λ_e) are being used, letter ξ means that parameters $(\lambda_i, \lambda_e + \varepsilon)$ are being used, via an extra Poisson clock of rate ε to account for the increase in λ_e . The values of $\varepsilon > 0$ and (λ_i, λ_e) such that $\theta(\lambda_i, \lambda_e) > 0$ and $\lambda_e + \varepsilon \leq \lambda_i$ are arbitrary but fixed.

Proof of Proposition 3.8. We will consider a sequence of coupled processes $(\eta_t^n, \xi_t^n)_{t \ge 0}$ indexed by $n \in \mathbb{N}$. Denote by $0 < \tau_1 < \tau_2 < \ldots$ the times of the external Poisson clock.

Start by defining the pair $(\eta_t, \xi_t)_{t \ge 0}$ starting from $\eta_0 = \xi_0$ sampled from μ , which is supported on Σ^{\odot} . For all $t \in [0, \tau_1)$ we have $\xi_t = \eta_t$ and at time τ_1 we have

$$\xi_{\tau_1} = \Xi \eta_{\tau_1} = \eta_{\tau_1} \cup \{ \mathcal{R} \eta_{\tau_1} + 1 \}_{\tau_1}$$

where Xi was defined in Section 4. Since the process $(\eta_t)_{t \ge 0}$ is independent of τ_1 , the distribution of $\Psi\eta_{\tau_1}$ is μ . Thus, by Proposition 4.6, $\Psi\xi_{\tau_1} \ge \mu$. Hence, enlarging the underlying probability space if necessary, there exists a random configuration ζ^1 such that $\Psi\zeta^1$ is μ -distributed, $\mathcal{R}\zeta^1 = \mathcal{R}\xi_{\tau_1}$ and $\zeta^1 \subseteq \xi_{\tau_1}$. Moreover, ζ^1 is independent of τ_1 and of the graphical construction ω after time τ_1 . We now consider a coupled process $(\eta_t^1, \xi_t^1)_{t \ge \tau_1}$ which evolves using the same graphical construction and whose initial condition is $\eta_{\tau_1}^1 = \xi_{\tau_1}^1 = \zeta^1$. By Lemma 3.1, $\eta_{\tau_1+t}^1 \subseteq \xi_{\tau_1+t}^1 \subseteq \xi_{\tau_1+t}$ for all $t \ge 0$. Moreover, since the process $(\eta_{\tau_1+t}^1)_{t\ge 0}$ is equal in distribution to the process $(T\eta_{\tau_1+t})_{t\ge 0}$ we have $\mathbb{E}[\mathcal{R}\eta_{\tau_1+t}^1] = \mathbb{E}[\mathcal{R}\eta_{\tau_1+t}] + 1$. Proceeding recursively, for each n we obtain processes $(\eta_t^n, \xi_t^n)_{t\ge \tau_n}$ such that $\eta_{\tau_n+t}^n \subseteq \xi_{\tau_n+t}^n \subseteq \xi_{\tau_n+t}$ and $\mathbb{E}[\mathcal{R}\eta_{\tau_n+t}^n] = \mathbb{E}[\mathcal{R}\eta_{\tau_n+t}] + n$ for all $t \ge 0$. In particular, for all $n \in \mathbb{N}$,

$$\mathbb{E}[\mathcal{R}\xi_{\tau_n}] \geqslant \mathbb{E}[\mathcal{R}\eta_{\tau_n}] + n.$$

We now show from this inequality that

$$\alpha(\lambda_i, \lambda_e + \varepsilon) \ge \alpha(\lambda_i, \lambda_e) + \varepsilon.$$

Since $\mathbb{E}[\mathcal{R}\eta_t] = \alpha(\lambda_i, \lambda_e) \cdot t$ by Proposition 4.5, and $(\eta_t)_t$ is independent of $(\tau_k)_k$, we have

$$\mathbb{E}[\mathcal{R}\eta_{\tau_n}] = \frac{\alpha(\lambda_i, \lambda_e)}{\varepsilon} \, n,$$

as τ_n is the *n*-th arrival of Poisson clock of rate ε .

This and the previous inequality imply that, for all $n \in \mathbb{N}$,

$$\mathbb{E}\Big[\frac{\mathcal{R}\xi_{\tau_n}}{n}\Big] \geqslant \frac{\alpha(\lambda_i,\lambda_e) + \varepsilon}{\varepsilon}.$$

On the other hand, by Lemma 3.5, $\frac{\mathcal{R}\xi_{\tau_n}}{\tau_n} \to \alpha(\lambda_i, \lambda_e + \varepsilon)$ a.s., and therefore

$$\frac{\mathcal{R}\xi_{\tau_n}}{n} \to \frac{\alpha(\lambda_i, \lambda_e + \varepsilon)}{\varepsilon} \quad \text{a.s}$$

To finish the proof, we want to combine the above equations, and for this we need to relate the a.s. limit of $\frac{\mathcal{R}\xi_{\tau_n}}{n}$ to its expectation.

Let $(Z_t)_{t\geq 0}$ be the process on \mathbb{Z} that starts at the origin and jumps one unit to the right whenever the process $(\mathcal{R}\xi_t)_{t\geq 0}$ jumps to the right. Then $\mathcal{R}\xi_t \leq Z_t$ a.s. for all $t \geq 0$. Also, the sequence $\frac{Z_{\tau_n}}{n}$ converges a.s. and in mean to a constant v. Applying Fatou's lemma to $\frac{Z_{\tau_n} - \mathcal{R}\xi_{\tau_n}}{n}$, we get

$$v - \mathbb{E}\left[\limsup_{n} \frac{\mathcal{R}\xi_{\tau_n}}{n}\right] \leq v - \limsup_{n} \mathbb{E}\left[\frac{\mathcal{R}\xi_{\tau_n}}{n}\right]$$

Hence,

$$\frac{\alpha(\lambda_i,\lambda_e)+\varepsilon}{\varepsilon} \leqslant \limsup_n \mathbb{E}\Big[\frac{\mathcal{R}\xi_{\tau_n}}{n}\Big] \leqslant \mathbb{E}\Big[\limsup_n \frac{\mathcal{R}\xi_{\tau_n}}{n}\Big] = \frac{\alpha(\lambda_i,\lambda_e+\varepsilon)}{\varepsilon},$$

proving the proposition.

7 Speed increase with internal infection rates

In this section we prove Proposition 3.10. We want to show that a small increase in the internal infection rate amounts to an increase in the speed. The argument here rather indirect if compared to the previous section. It considers an auxiliary process that has infection rate λ_e only at the rightmost site but still λ_i at the leftmost one, shows that this process survives with probability $\theta_* > 0$, then shows that adding a site to a given configuration at a certain time increases the position of the rightmost infected site by at least θ_* on average, at any future time. This effect is cumulative, and from there we conclude that a process with larger λ_i has larger speed, on average. Since speed can be defined as expected speed (Lemma 3.4), this is enough to derive Theorem 4.

Notation. Throughout this section, letter η means that parameters (λ_i, λ_e) are being used, and ξ means that parameters (λ'_i, λ_e) are being used. An auxiliary process, denoted with ζ , has infections to the right of the rightmost infected site occurring at rate λ_e and other infections occurring at rate λ_i . The values of $\lambda_i > \lambda_c$ and $\lambda'_i > \lambda_i$ are fixed, and $\lambda_e = \lambda_e^*(\lambda_i) \leq \lambda_c$.

Let

$$\theta_* := \mathbb{P}(\zeta_t^0 \neq \emptyset, \forall t \ge 0)$$

Lemma 7.1. We have $\theta_* > 0$.

Proof. By Lemma 3.4 and Theorem 7, $\frac{\mathcal{R}\eta_t^-}{t} \to 0$ a.s. Hence, for any $\delta > 0$

$$\lim_{n} \mathbb{P}(\mathcal{R}\eta_t^- \ge -n - \delta t \text{ for all } t \ge 0) = 1.$$

Similarly, $\frac{\mathcal{L}\zeta_t^+}{t} \to -\alpha(\lambda_i, \lambda_i)$, so, for every $\delta' < \alpha(\lambda_i, \lambda_i)$,

$$\lim_{n} \mathbb{P}(\mathcal{L}\zeta_t^+ \leqslant n - \delta' t \text{ for all } t \ge 0) = 1$$

Now take some δ and δ' such that $0 < \delta < \delta' < \alpha(\lambda_i, \lambda_i)$, and take n such that the above probabilities are larger than $\frac{1}{2}$. Translating the latter event horizontally by -3n, we get the event $\{\mathcal{L}\zeta^{\{-3n,\ldots,0\}\cup\mathbb{Z}_+} \leq -2n - \delta't \text{ for all } t \geq 0\}$. Simultaneous occurrence of this event and $\{\mathcal{R}\eta_t^- \geq -n - \delta t \text{ for all } t \geq 0\}$ implies $\{\zeta_t^{\{-3n,\ldots,0\}} \neq \emptyset \text{ for all } t \geq 0\}$. By Remark 2.1, $\theta^* > 0$.

Lemma 7.2. Let $A \in \Sigma^{\ominus}$ and let $A' = A \cup \{x\}$ for some $x > \mathcal{R}A$. Then,

$$\mathbb{E}[\mathcal{R}\eta_t^{A'}] \geqslant \mathbb{E}[\mathcal{R}\eta_t^A] + \theta_*$$

for all $t \ge 0$.

Proof. We will construct a coupling of two contact processes η^A and $\eta^{A'}$ starting from A and A', respectively. Let $y = \mathcal{R}A$.

For the process $\eta^{A'}$, we first explore the (λ_i, λ_e) -active paths for configuration A' starting at (x, 0). We then translate this collection of active paths by (y - x, 0) and use them for the process η^A . Call $\tau \in (0, +\infty]$ the global extinction time of these active paths.

Then construct the paths starting to the left of x with the same graphical construction up to the time they reach an already constructed active path, i.e., an active path staring at (x, 0) for $\eta^{A'}$ and at (y, 0) for η^{A} .

We now observe that, for $t < \tau$, we have $\mathcal{R}\eta_t^{A'} = \mathcal{R}\eta_t^A + (x - y)$ and, for $t \ge \tau$ we have $\eta_t^{A'} \supseteq \eta_t^A$ because for any active path for A' which did not meet the descendants of x there is a corresponding active path for A which did not meet the descendants of y. Therefore,

$$\mathcal{R}\eta_t^{A'} - \mathcal{R}\eta_t^A \ge \mathbb{1}_{\{\tau > t\}}.$$

This proves the desired inequality and thus the lemma.

Proof of Proposition 3.10. First note that there exists $\delta > 0$ such that, for all $A \in \Sigma^{\ominus}$, at t = 1 we have $\mathbb{P}(\mathcal{R}\xi_1^A \ge \mathcal{R}\eta_1^A + 1) \ge \delta$ (proof omitted).

Now for each $n \in \mathbb{N}$, consider the process $(\xi_t^n)_{t \ge n}$ starting from $\xi_n^n = \eta_n^-$. Using Lemma 7.2, one gets $\mathbb{E}[\mathcal{R}\xi_t^-] \ge \mathbb{E}[\mathcal{R}\xi_t^1] + \delta\theta_*$ for all $t \ge 1$.

Likewise, $\mathbb{P}(\mathcal{R}\xi_2^1 \ge \mathcal{R}\eta_2^- + 1) \ge \delta$, hence for $t \ge 2$

$$\mathbb{E}[\mathcal{R}\xi_t^-] \geqslant \mathbb{E}[\mathcal{R}\xi_t^1] + \delta\theta_* \geqslant \mathbb{E}[\mathcal{R}\xi_t^2] + 2\delta\theta_* \geqslant \mathbb{E}[\mathcal{R}\eta_t^-] + 2\delta\theta_*.$$

The induction is clear and gives

$$\mathbb{E}[\mathcal{R}\xi_t^-] \geqslant \mathbb{E}[\mathcal{R}\eta_t^-] + n\delta\theta,$$

for all $n \in \mathbb{N}$ and $t \ge n$. Therefore,

$$\alpha(\lambda_i', \lambda_*^e(\lambda_i)) \ge \alpha(\lambda_i, \lambda_*^e(\lambda_i)) + \delta\theta_* = \delta\theta_* > 0$$

by Theorem 7 and Lemma 7.1, concluding the proof.

8 Domination in the non-attractive case

In this section we prove Theorem 5.

Notation. Throughout this section, $\lambda_i = \lambda_e = \lambda_c$. Letter ξ means that parameters $(\lambda_i, \lambda_e + \varepsilon)$ are being used, via an external Poisson clock to account for the ε increase. The value of $\varepsilon > 0$ is arbitrary but fixed. As usual, letter η means that parameters (λ_i, λ_e) are being used.

Denote by $0 = \tau_0 < \tau_1 < \tau_2 < \dots$ the times of the external Poisson clock (which has intensity ε). We will construct a sequence of processes denoted $(\xi_t^i)_{t \ge \tau_i}$, for $i \in \mathbb{N}_0$. Their initial distribution will

be the measure μ provided by Proposition 4.1 for $\lambda_i = \lambda_e = \lambda_c$, and the superscripts will be used to enumerate them. The process $(\xi_t^0)_{t\geq 0}$ is given by $(\xi_t)_{t\geq 0}$ which starts form a random configuration $\xi_0^0 \sim \mu.$

Suppose the process $(\xi^{i-1})_{t \ge \tau_{i-1}}$ has been constructed. We will show how to construct $(\xi^i)_{t \ge \tau_i}$.

We start by describing how we sample $\xi_{\tau_i}^i$, the initial configuration at time τ_i for $(\xi_t^i)_{t \ge \tau_i}$, from the configuration $\xi_{\tau_i}^{i-1}$. This configuration $\xi_{\tau_i}^i$ will satisfy $\xi_{\tau_i}^i \subseteq \xi_{\tau_i}^{i-1}$, $\mathcal{R}\xi_{\tau_i}^i = \mathcal{R}\xi_{\tau_i}^{i-1}$ and $\Psi\xi_{\tau_i}^i \sim \mu$. We will describe it rather explicitly, because later on we will need to argue that a certain sequence is stationary, and this will be key to prove Theorem 5. First note that, by Proposition 4.1, $\Psi \xi_{\tau_i}^{i-1} \sim \mu$. Also note that $\xi_{\tau_i}^{i-1} = \Xi(\xi_{\tau_i}^{i-1})$. By Proposition 4.6, $\Psi \xi_{\tau_i}^{i-1} \succeq \mu$. We will show how to use $\xi_{\tau_i}^{i-1}$ and some extra randomness to sample the configuration $\xi^i_{\tau_i}$ having the above properties. Without loss of generality, we assume that $\mathcal{R}\xi_{\tau_i}^{i-1} = 0$, otherwise just use $\Psi\xi_{\tau_i}^{i-1}$ and translate the resulting $\xi^i_{\tau_i}$ accordingly. Let $(V^i_k)_{k\in\mathbb{N}}$ be an infinite sequence of independent random variables uniformly distributed on [0,1] which are also independent of all the Poisson processes used previously. Also let ν be a distribution on $\Sigma^{\odot} \times \Sigma^{\odot}$ such that its first marginal is μ , its second marginal is the distribution of $\xi_{\tau_i}^{i-1}$ and $\nu(\{(\eta,\xi):\eta\subseteq\xi\})=1$. This measure ν has a conditional distribution of η given ξ , given by a regular conditional distribution. We start by declaring that $0 \in \xi_{\tau_i}^i$. Once $\xi_{\tau_i}^i \cap \{-k, \dots, 0\}$ has been determined, we declare that $-k - 1 \in \xi_{\tau_i}^i$ if $V_{k+1}^i \leq \nu(-k - 1 \in \eta \,|\, \xi = 1)$ $\xi_{\tau_i}^{i-1}$, $\eta = \xi_{\tau_i}^i$ on $\{-k, \ldots, 0\}$), and that $-k - 1 \notin \xi_{\tau_i}^i$ otherwise. This determines $\xi_{\tau_i}^i$ with the claimed properties.

To describe the evolution of $(\xi_i^i)_{t \ge \tau_i}$, we start looking at the active paths starting from the spacetime point $(\mathcal{R}\xi_{\tau_i}^{i-1}, \tau_i)$ for the configuration $\xi_{\tau_i}^{i-1}$. These are the paths whose jumps are supported by a λ_c -open edge, plus an extra jump to the right at the rightmost active path at times $\tau_{i+1}, \tau_{i+2}, \ldots$ For simplicity we assume this set of paths is finite in time (the opposite already implies Theorem 5 anyway). Let $A_{i,0}$ be the set of points in these paths, let $\tau_{i,0} = \tau_i$ and let

$$\tau_{i,1} = \sup\{s : A_{i,0} \cap \{(x,s), x \in \mathbb{Z}\} \neq \emptyset\}.$$

Then proceed by induction on j: first let $A_{i,j-1}$ be the set of points in the open paths starting at $(\mathcal{R}\xi_{\tau_{i,j-1}}^{i-1}, \tau_{i,j-1})$ and after that let

$$\tau_{i,j} = \sup\{s : A_{i,j-1} \cap \{(x,s), x \in \mathbb{Z}\} \neq \emptyset\}.$$

The construction of the process ξ^i is now done as follows:

- 1. In the time interval $[\tau_{i,j}, \tau_{i,j+1})$ the open paths starting at $(\mathcal{R}\xi^i_{\tau_{i,j}}, \tau_{i,j})$ are given by $T^{\mathcal{R}\xi^i_{\tau_{i,j}}-\mathcal{R}\xi^{i-1}_{\tau_{i,j}}}A_{i,j}$. That is, we translate the open paths starting from the rightmost point of $\xi_{\tau_i}^{i-1}$ to the rightmost point of $\xi^i_{\tau_{i,i}}$
- 2. the open paths in the same time interval starting at the other points of $\xi^i_{\tau_{i,j}}$ are given by the Poisson processes of parameters λ_i and 1 until they merge with the open paths starting from the rightmost point of $\xi_{\tau_{i,i}}^i$ (if this occurs).

The point of this perhaps intricate construction is that it simultaneously satisfies two properties.

First, the sequence $(\mathcal{R}\xi_{\tau_n}^n - \mathcal{R}\xi_{\tau_{n-1}}^{n-1})_{n\in\mathbb{N}}$ is stationary. Second, stated below, it allows a comparison between ξ^{i+1} and ξ^i .

Lemma 8.1. With the above construction of the processes ξ^i , we have $\mathcal{R}\xi_t^{i+1} \leq \mathcal{R}\xi_t^i$ for all $i \in \mathbb{N}_0$ and all $t \geq \tau_{i+1}$.

Proof. We prove by induction that, for all $j \in \mathbb{N} \cup \{0\}$,

$$\mathcal{R}(\xi_t^{i+1}) \leqslant \mathcal{R}(\xi_t^i)$$

for all $t \in [\tau_{i+1,j}, \tau_{i+1,j+1})$ and $\xi_{\tau_{i+1,j}}^{i+1} \subseteq \xi_{\tau_{i+1,j}}^i$. For j = 0 we obviously have $\xi_{\tau_{i+1,0}}^{i+1} = \xi_{\tau_{i+1,0}}^{i+1} \subseteq \xi_{\tau_{i+1,0}}^{i+1} = \xi_{\tau_{i+1,0}}^{i+1}$ and $\mathcal{R}(\xi_t^{i+1}) = \mathcal{R}(\xi_t^i)$ for all $t \in [\tau_{i+1,0}, \tau_{i+1,1})$. For the inductive step assume that $\mathcal{R}(\xi_t^{i+1}) \leq \mathcal{R}(\xi_t^i)$ for all $t \in [\tau_{i+1,k}, \tau_{i+1,k+1})$ and that $\xi_{\tau_{i+1,k}}^{i+1} \subseteq \xi_{\tau_{i,+1k}}^i$. At time $\tau_{i+1,k+1}$ the open paths of ξ^i starting at $(\mathcal{R}(\xi_{\tau_{i+1,k}}^{i+1}), \tau_{i+1,k})$ and the open paths of ξ^{i+1} starting at $(\mathcal{R}(\xi_{\tau_{i+1,k}}^{i+1}), \tau_{i+1,k}), \tau_{i+1,k})$ have vanished. Hence, the open paths starting at the left of these points which have merged with the paths starting at these points have also vanished. Since the open paths of ξ^{i+1} starting at $(\mathcal{R}(\xi_{\tau_{i+1,k}}^{i+1}), \tau_{i+1,k})$ are to the left of the open paths of ξ^i starting at $(\mathcal{R}(\xi_{\tau_{i+1,k}}^{i+1}), \tau_{i+1,k})$ we have $\xi_{\tau_{i+1,k+1}}^{i+1} \subseteq \xi_{\tau_{i+1,k+1}}^i$ and this implies that $\mathcal{R}(\xi_t^{i+1}) \leq \mathcal{R}(\xi_t^i)$ for all $t \in [\tau_{i+1,k+1}, \tau_{i+1,k+2})$ completing the inductive step and the proof of the lemma.

To simplify the notation, let $Y_n = \mathcal{R}\xi_{\tau_n}^n - \mathcal{R}\xi_{\tau_{n-1}}^{n-1}$.

Lemma 8.2. The random variables Y_n are integrable and $\mathbb{E}Y_n = 1$.

Proof. Write $Y_n = X_n + 1$, so X_n is distributed as the increment of the position of the rightmost point of $(\eta^{\mu})_{t \ge 0}$ in the time interval $[0, \tau]$ where τ is an exponential random variable that is independent of the process $(\eta^{\mu})_{t \ge 0}$. That $\mathbb{E}X_n = 0$ follows from the proposition below.

Proof of Theorem 5. We will prove that

$$\mathbb{P}\left(\liminf_{t} \frac{\mathcal{R}\xi_{t}^{\mu}}{t} > 0\right) > 0, \tag{8.3}$$

and then discuss how to get Theorem 5 therefrom. Remark 1.7 requires extra work and is handled in the next section.

From the definition of Y_k , we have

$$\mathcal{R}\xi_{\tau_n}^n = \sum_{k=1}^n Y_k$$

Recall that $(Y_n)_n$ is stationary. By the Ergodic Theorem, $\frac{\mathcal{R}(\xi_{\tau_n}^n)}{n} \to Y$ a.s., for some (possibly random) Y. Moreover, by Lemma 8.2, $\mathbb{E}Y = 1$. Hence,

$$\mathbb{P}\Big(\lim_{n}\frac{\mathcal{R}(\xi_{\tau_n}^n)}{n}>0\Big)>0.$$

On the other hand, by Lemma 8.1, we have $\mathcal{R}\xi^{\mu}_{\tau_n} \ge \mathcal{R}\xi^n_{\tau_n}$, whence

$$\mathbb{P}\big(\liminf_n \frac{\mathcal{R}(\xi_{\tau_n}^{\mu})}{n} > 0\big) > 0.$$

Since $\sup_{t \in [\tau_n, \tau_{n+1}]} (\mathcal{R}(\xi_{\tau_{n+1}}^{\mu}) - \mathcal{R}(\xi_t^{\mu}))$ is bounded above by a geometric random variable with parameter $\frac{\varepsilon}{\lambda_c + \varepsilon}$, using a simple Borel-Cantelli argument and $\frac{\tau_n}{n} \to \varepsilon^{-1}$ a.s., we can conclude (8.3). Now, consider a process $(\zeta_t)_t$ that has rate $\lambda_e + \varepsilon$ for infections to the right of the rightmost infected site and λ_i everywhere else. We claim that $\mathbb{P}(\zeta_t^0 \neq \emptyset \forall t \ge 0) > 0$. Indeed, if this probability were zero then there would exist a diverging sequence of random times t_n such that $\xi_{t_n}^{\mu} = \eta_{t_n}^{\mu}$ where η^{μ} is the process with parameters (λ_c, λ_c) , and, since $\lim \frac{\mathcal{R}\eta_t^{\mu}}{t} = 0$ a.s. by Proposition 4.7, this would imply that $\liminf_t \frac{\mathcal{R}\xi_t^{\mu}}{t} = 0$ a.s.

To get Theorem 5 from the above claim, note that, although a direct comparison between $(\zeta_t^0)_t$ and $(\eta_t^0)_t$ using the same graphical construction may break down due to lack of attractiveness, we can compare the probability that these processes survive using Proposition 1 of [5], thus proving Theorem 5.

9 Survival of the critical contact process on N with enhancement at the boundary

Continuing from the previous section, we now bootstrap from the intermediate results used in the proof of Theorem 5 so as to finally justify Remark 1.7.

Note that survival of the process $(\zeta_t)_t$ on \mathbb{Z} is a weaker version of Remark 1.7, differing only on the underlying physical space being \mathbb{Z} instead of \mathbb{Z}_+ . So it remains to show that, with positive probability, the process survives even if the negative half-line is shut down.

Let us make this approach more precise. On the event $\{\zeta_t^0 \neq \emptyset \ \forall t \ge 0\}$, define the random path $(\Gamma(t))_{t\ge 0}$ as the rightmost infinite path started from (0,0). That is, take

$$\Gamma(t) = \max\{x \in \mathbb{Z} : (0,0) \text{ infects } (t,x) \text{ and } (t,x) \text{ infects } \infty\},\$$

which corresponds to an infinite infection path in the graphical construction, and is such that any other infinite infection path γ starting from $\gamma(0) = 0$ satisfies $\gamma(t) \leq \Gamma(t)$ for all $t \geq 0$. Then our goal is to show that $\mathbb{P}(\Gamma(t) \geq 0 \forall t) > 0$, since on this event the process survives regardless of the infections taking place on the negative line, which can be completely suppressed.

In the previous section we actually showed that $\{\zeta_t^0 \neq \emptyset \ \forall t \ge 0\} \cap \{\liminf_t \frac{\mathcal{R}\xi_t^{\mu}}{t} > 0\}$ has positive probability. Now notice that, on the event $\{\zeta_t^0 \neq \emptyset \ \forall t \ge 0\}$, we have $\mathcal{R}\zeta_t^0 = \mathcal{R}\xi_t^{\mu} \ \forall t \ge 0$, so combining these events we get

$$\mathbb{P}(\mathcal{R}\zeta_t^0 \ge \delta t \ \forall t \ge T_0) > 0$$

for some $T_0 < \infty$ and some $\delta > 0$.

After some some thinking and drawing some pictures, the reader will conclude that, on the event $\{\mathcal{R}\zeta_t^0 \ge \delta t \ \forall t \ge T_0\} \cap \{\Gamma(t) \le 0 \text{ i.o.}\}$, there exists an increasing diverging sequence $(T_n)_n$ such that there are λ_c -open paths from $(0, T_n)$ to the line $\{(\lceil \delta t \rceil, t) : t \ge 0\}$. A detailed proof of this implication is very tedious and will be omitted.

To finish the proof of Remark 1.7, it remains to show that the latter event (infinitely many crossings of a cone by λ_c -open paths) has probability zero.

Lemma 9.1. Consider the critical contact process on \mathbb{Z} and let $\delta > 0$. The probability that there are infinitely many paths from $\{(0,t):t \ge 0\}$ to $\{(\lceil \delta t \rceil, t):t \ge 0\}$ is zero.

Proof. From Lemma 2 of [13], we get $0 < c_1 < C_1 < \infty$ such that

$$\mathbb{P}(\sup_{s\in[0,t]}\mathcal{R}\eta_t^- \ge \delta t) \leqslant C_1 e^{-c_1 t}$$
(9.2)

for all $t \ge 0$. Indeed, take $\lambda > \lambda_c$ such that $\alpha(\lambda, \lambda) < \delta$ and apply [13, Lemma 2] directly.

From (9.2), we get

$$\mathbb{P}(\sup_{s\in[t,2t]}\frac{\mathcal{R}\eta_s^-}{s} \ge \delta) \leqslant C_2 e^{-c_2 t}$$

for all t > 0 for new constants $0 < c_2 < C_2 < \infty$ (all constants depend on δ).

Fix t > 0. Taking a union bound, the previous estimate gives

$$\mathbb{P}(\sup_{s \ge t} \frac{\mathcal{R}\eta_s^-}{s} \ge \delta) \leqslant \sum_{k \ge 1} \mathbb{P}(\sup_{s \in [kt, 2kt]} \frac{\mathcal{R}\eta_s^-}{s} \ge \delta) \leqslant \sum_{k \ge 1} C_2 e^{-c_2 kt} \leqslant C_3 e^{-c_3 t}$$

Combining the previous estimates, we get

$$\mathbb{P}(\sup_{s \ge 0} \frac{\mathcal{R}\eta_s^-}{s+t} \ge \delta) \leqslant C_4 e^{-c_4 t},\tag{9.3}$$

which is valid for all t > 0. We only need another layer of tedious arguments the proof will be finished.

Fix T > 0. And let A_T denote the event that there is a path from $\{0\} \times [T, +\infty)$ to $\{(\lceil \delta t \rceil, t) : t \ge 0\}$. Denote the recovery marks on $\{0\} \times [T, +\infty)$ by $T + \tau_1, T + \tau_1 + \tau_2, T + \tau_1 + \tau_2 + \tau_3, \ldots$, and notice that $(\tau_n)_{n \in \mathbb{N}}$ are i.i.d. standard exponential variables. Observe that $A_T \subseteq \bigcup_{n \in \mathbb{N}_0} A_{T,n}$, where $A_{T,n}$ is the event that there is a λ_c -open path from $T + \sum_{k=1}^n \tau_k$ to some (x, t) with $x \ge \delta t$. Let $\beta = \mathbb{E}[e^{-c_4\tau_1}] < 1$. Using (9.3), we get

$$\mathbb{P}(A_{T,n}) = \mathbb{E}[\mathbb{P}(A_{T_n}|\tau_1,\ldots,\tau_n)] \leqslant \mathbb{E}[C_4 \exp(-c_4(T+\tau_1+\cdots+\tau_n))] \leqslant C_4 e^{-c_4 T} \beta^n,$$

and thus

$$\mathbb{P}(A_T) \leqslant \sum_{n=0}^{\infty} \mathbb{P}(A_{T,n}) \leqslant \frac{C_4 e^{-c_4 T}}{1-\beta}.$$

Since this vanishes as $T \to \infty$, the lemma is proved.

Using the above lemma, we conclude that $\mathbb{P}(\Gamma(t) \ge 0 \ \forall t) > 0$, proving Remark 1.7.

10 No survival for the critical contact process on \mathbb{N} with finite enhancements

In this section we prove Theorem 6.

The proof is by contradiction. Let $k \in \mathbb{N}$, $\lambda < \infty$ and $\delta > 0$. Let $\zeta_{t;k,\lambda,\delta}^0$ denote the contact process on \mathbb{Z}_+ started from $\zeta_0 = \{0\}$, having infection rate λ and recovery rate δ at sites $\{0, 1, \ldots, k\}$, and infection rate λ_c and recovery rate 1 at sites $\{k + 1, k + 2, \ldots\}$. Suppose by contradiction that $\mathbb{P}(\zeta_{t;k,\lambda,\delta}^0 \neq \emptyset \forall t \ge 0) > 0$. Fix some $0 < \delta' < \delta'' < \delta$ and $\lambda' = \lambda + (\delta - \delta'')/2 < \infty$. By defining both processes on the same graphical construction, we have $\zeta_{t;k,\lambda',\delta'}^0 \supseteq \zeta_{t;k,\lambda,\delta}^0$, and therefore $\mathbb{P}(\zeta_{t;k,\lambda',\delta'}^0 \neq \emptyset \forall t \ge 0) > 0$.

We will now produce a new graphical construction with parameters δ and λ from the graphical construction with parameters δ' and λ' , using extra randomness to resample some of the marks. On this new graphical construction, we will define a process $(\xi_t)_{t \ge 0}$, which is thus distributed as $(\zeta_{t;k,\lambda,\delta}^0)_{t \ge 0}$. At each site $x \in \{0,\ldots,k\}$, each λ' -infection mark from x to $x \pm 1$ is converted into a recovery mark at x with probability $1 - \lambda/\lambda'$, independently for each mark; moreover, an independent Poisson clock of rate $\delta'' - \delta'$ is added as new recovery marks. This new graphical construction has the same distribution as the graphical construction with parameters k, λ and δ , and we use it to define the process $(\xi_t)_t$. Below we show that $\mathbb{P}(\xi_t \neq \emptyset \forall t \ge 0) = 0$, contradicting the assumption that $\mathbb{P}(\zeta_{t;k,\lambda,\delta}^0 \neq \emptyset \forall t \ge 0) > 0$.

On the event $\{\zeta_{t;k,\lambda',\delta'}^0 \neq \emptyset \,\forall t \ge 0\}$, let Γ be the rightmost infinite open path started from (0,0). More precisely, and explicitly, take

$$\Gamma(t) = \max\{x \in \mathbb{Z}_+ : (0,0) \underset{(k,\lambda',\delta')}{\leadsto} (t,x) \text{ and } (t,x) \underset{(k,\lambda',\delta')}{\leadsto} \infty\},\$$

which corresponds to an infinite open path in the graphical construction, and is such that any other infinite open path γ starting from $\gamma(0) = 0$ satisfies $\gamma(t) \leq \Gamma(t)$ for all $t \geq 0$. On the above definition we are using the notation $(x, t) \rightsquigarrow_{(k,\lambda',\delta')}(y, s)$ to denote existence of a path from (x, t) to (y, s) in the model that uses parameters (k, λ', δ') .

Now observe that, almost surely, $\Gamma(t) \leq k$ for arbitrarily large times, since otherwise we would have an infinite open path in the region $\{k + 1, k + 2, ...\} \times [0, +\infty)$, contradicting [3]. So there is a divergent sequence of times $(t_n)_n$ such that $\Gamma(t_n) \leq k$ for all n. We can assume that $t_{n+1} > t_n + 1$ for all n.

Finally, let \mathcal{A}_n denote the event that, for each site $x \in \{0, \ldots, k\}$, during the time interval $[t_n, t_{n+1}]$ a recovery mark is added and the first infection mark (if there is one) is converted into a recovery mark. The events $(\mathcal{A}_n)_n$ are conditionally independent given the original graphical construction, and have (conditional) probability at least $((1 - \lambda/\lambda')(1 - e^{\delta'-\delta''}))^{k+1} > 0$. In particular, a.s. at least one such event will occur. Now observe that the occurrence of \mathcal{A}_n for some n is enough to simultaneously break every infinite open path that starts from (0,0). We have thus proved that $\mathbb{P}(\xi_t \neq \emptyset \forall t \ge 0) = 0$ as announced above, contradicting the assumption that this probability is positive. This proves Theorem 6.

11 Open problems

- 1. As already said in Remark 1.11, it is not known whether the contact process on \mathbb{Z} with parameter λ_c dies out if we increase the rate of infection emanating from a fixed site.
- 2. The proofs of the asymptotic speed results given in Section 3 depend on the Subadditive Ergodic Theorem. Do they still hold when $\lambda_i < \lambda_e$? The difficulty is due to the loss of attractiveness of the process.
- 3. Does the conclusion of Proposition 4.1 still holds when $\lambda_i = \lambda_c < \lambda_e$?

Acknowledgment

This paper was written while the first author was visiting IMPA in Rio de Janeiro. Thanks are given to its hospitality.

References

- M. AIZENMAN, G. GRIMMETT. Strict monotonicity for critical points in percolation and ferromagnetic models. J Statist Phys 63:817–835, 1991. doi.
- [2] C. BEZUIDENHOUT, L. GRAY. Critical attractive spin systems. Ann Probab 22:1160–1194, 1994. doi.
- [3] C. BEZUIDENHOUT, G. GRIMMETT. The critical contact process dies out. Ann Probab 18:1462– 1482, 1990. doi.
- [4] J. T. COX, R. DURRETT, R. SCHINAZI. The critical contact process seen from the right edge. Probab Theory Related Fields 87:325–332, 1991. doi.
- [5] R. DURRETT, R. B. SCHINAZI. Boundary modified contact processes. J Theoret Probab 13:575– 594, 2000. doi.

- [6] A. GALVES, E. PRESUTTI. Edge fluctuations for the one-dimensional supercritical contact process. Ann Probab 15:1131–1145, 1987. doi.
- [7] D. GRIFFEATH. The basic contact processes. Stochastic Process Appl 11:151–185, 1981. doi.
- [8] ———. The binary contact path process. Ann Probab 11:692–705, 1983. doi.
- [9] G. R. GRIMMETT. Percolation, vol. 321 of Grundlehren der mathematischen Wissenschaften. Springer-Verlag, 2 edn., 1999. doi.
- [10] R. HOLLEY, T. M. LIGGETT. Generalized potlatch and smoothing processes. Z Wahrsch Verw Gebiete 55:165–195, 1981. doi.
- [11] T. M. LIGGETT. Interacting particle systems. Classics in Mathematics. Springer-Verlag, Berlin, 2005. doi.
- [12] T. M. LIGGETT, R. H. SCHONMANN, A. M. STACEY. Domination by product measures. Ann Probab 25:71–95, 1997. doi.
- [13] T. S. MOUNTFORD, T. D. SWEET. An extension of Kuczek's argument to nonnearest neighbor contact processes. J Theoret Probab 13:1061–1081, 2000. doi.
- [14] C. TERRA. Dynamic phenomena in interacting particle systems: Phase transitions and equilibrium, 2024. doi. Preprint.
- [15] D. VALESIN. The contact processon on random graphs. Ensaios Matemáticos 40:1–115, 2024. doi.
- [16] Y. ZHANG. A note on inhomogeneous percolation. Ann Probab 22:803-819, 1994. doi.