# Direct and Inverse Problems in Special Geometry

Sergio Cecotti<sup>∗</sup>

Yanqi Lake Beijing Institute of Mathematical Sciences and Applications (BIMSA) Huaibei Town, Huairou District, Beijing 101408, China

and

Qiuzhen College, Tsinghua University, Beijing, China, SISSA, via Bonomea 265, Trieste, Italy

#### Abstract

The *inverse problem of special geometry* (Seiberg-Witten geometry of 4d  $\mathcal{N} = 2$  SCFT) asks for a recursive construction of all such geometries in rank r by assembling together known lower-rank "strata". This leads to a program to understand/construct/classify all special geometries which looks surprising effective. After reviewing some advanced topics in special geometry, in this long note we define the inverse problem and introduce the basic tools of the trade. The program is essentially completed in rank 2, and we pave the way to proceed to higher ranks. A central role is played by various notions of geometric rigidity: in addition to the obvious one (triviality of the conformal manifold), Falting-Saito-Peters rigidity and Deligne-Simpson rigidity also enter in the story.

December 6, 2023

<sup>∗</sup> e-mail: cecotti@sissa.it, cecotti@bimsa.cn

## Contents









## <span id="page-5-0"></span>1 Introduction: the Inverse Problem

A first fundamental step in any program to understand/construct/classify all 4d  $\mathcal{N} = 2$ SCFTs is to understand/construct/classify their special geometries (a.k.a. Seiberg-Witten geometries [\[1–](#page-148-0)[4\]](#page-148-1)). Such a geometric program has been especially pursued by Argyres and coworkers  $[5-20]^1$  $[5-20]^1$  $[5-20]^1$ ; for other work in the same line see e.g.  $[22-26]$  $[22-26]$ . This paper aims to introduce new powerful techniques and results in this direction.

A special geometry<sup>[2](#page-5-2)</sup> is a very nice mathematical gadget, with multiple physical interpretations and deep relations with several other topics, which may be studied from diverse math perspectives. A special geometry may be seen as the (complexification) of a classical integrable mechanical system [\[3,](#page-148-3)[4\]](#page-148-1) or as a non-perturbative description of a four-dimensional SUSY QFT  $[1,2]$  $[1,2]$ : we have two physical intuitions about them – one classical and one quantum – and the interplay between the two yields new unexpected insights. Special geometry in our SCFT sense looks very similar to the special geometry of  $\mathcal{N} = 2$  supergravity [\[27–](#page-150-2)[33\]](#page-150-3): formally the two are related by a simple "Wick rotation" in the signature of the fiber metric. The supergravity geometry describes the variation of Hodge structure (VHS [\[34–](#page-150-4)[42\]](#page-151-0)) of a family of Calabi-Yau 3-folds (3-CY) [\[29–](#page-150-5)[33\]](#page-150-3), and all ideas and techniques introduced in the study of moduli geometries of 3-CY can be equally well applied to the special geometries of  $\mathcal{N} = 2$  SCFT. Strangely enough, it seems that the previous SCFT literature did not leverage on the deep extensive work done in the Calabi-Yau contest. The 3-CY methods will be instead central in our discussion. Morally speaking, the program of the geometric classification of all 4d  $\mathcal{N} = 2$  SCFT may be seen as a simpler analogue of classifying all Calabi-Yau 3-folds, a basic problem with fundamental implications for quantum gravity [\[32\]](#page-150-6).

The main focus of this note is the *inverse method* in special geometry. While the paper is somehow written in a review format, essentially all results in this direction are novel, including the very notion of "inverse problem". In the rest of this introduction we outline what we mean by "inverse problem": a more precise description of the method is given in section [3](#page-44-0) in the proper geometric context.

Special geometry. In this paper a *special geometry* is a holomorphic integrable sys-tem<sup>[3](#page-5-3)</sup> associated to the Seiberg-Witten geometry of a 4d  $\mathcal{N}=2$  SCFT with all its implied structures [\[3,](#page-148-3) [4\]](#page-148-1). Thus a special geometry is a (smooth) complex symplectic  $\mathscr X$  variety together with a holomorphic fibration  $\pi: \mathscr{X} \to \mathscr{C}$  over the Coulomb branch  $\mathscr{C}$  with Lagrangian fibers which generically are polarized Abelian varieties. The Coulomb branch  $\mathscr C$  is the spectrum of the chiral ring  $\mathscr{R}$ , hence an affine complex variety. Most of our arguments work in the broader context of 4d  $\mathcal{N} = 2$  QFT, but we focus on the superconformal case where the geometry carries an additional structure: a  $\mathbb{C}^{\times}$ -action on  $\mathscr{X}$  by automorphisms of all the

<span id="page-5-1"></span> $1$  For a recent review see [\[21\]](#page-149-1).

<span id="page-5-2"></span><sup>2</sup> See section [2](#page-10-0) for precise definitions.

<span id="page-5-3"></span><sup>&</sup>lt;sup>3</sup> The integral systems of interest are actually *algebraic* not just *holomorphic*.

relevant geometric structures.<sup>[4](#page-6-0)</sup> The  $\mathbb{C}^{\times}$ -action reflects the existence of a conformal  $U(1)_R$ symmetry which is spontaneously broken to a discrete subgroup everywhere in  $\mathscr C$  but at a single closed point 0 (the origin). When the Coulomb branch is *smooth* at 0 (as a  $\mathbb{C}\text{-affine}$ variety)  $\mathscr{C} \simeq \mathbb{C}^r$ . We shall be concerned mainly, but not exclusively, with this case.

"Stratification". The quantum field theoretic intuition points out that a special geometry  $\mathscr{X} \to \mathscr{C}$  comes with a natural "stratification" in lower-dimension special geometries [\[16,](#page-149-2) [17\]](#page-149-3). We write the word "stratification" between quotes since the QFT intuition leads to a fairly new<sup>[5](#page-6-1)</sup> notion in symplectic geometry, which is *not* a stratification in any usual sense, and lays well outside the horizon of classical Liouvillian physical intuition. In math terms the "stratification" is a generalization of the Marsden-Weinstein-Meyer symplectic quotient [\[43,](#page-151-1)[44\]](#page-151-2) which applies to singular loci with suitable physical/geometric properties where the usual symplectic quotient becomes meaningless. The "stratification" of  $\mathscr X$  arises from an ordinary stratification of the Coulomb branch  $\mathscr{C}$  [\[16,](#page-149-2)[17\]](#page-149-3).

We outline the physical heuristics beyond "stratification". At the generic point of  $\mathscr C$ the IR physics, while not necessarily boring, is relatively simple, but there are special loci  $\mathscr{S}_a \subset \mathscr{C}$  where something "more interesting" happens. At a special point  $u \in \mathscr{C}$  three new phenomena may happen:

- (1) additional degrees of freedom become massless;
- (2) the unbroken subgroup  $R_u \subset U(1)_R$  of R-symmetry enhances;
- (3) we have both extra light degrees of freedom and R-symmetry enhancement.

In an interacting theory the special locus  $D \subset \mathscr{C}$  where we have additional massless degrees of freedom (called the discriminant) is a complex analytic subspace of pure codimension 1. We identify D with a simple effective divisor and write  $\mathcal{D} = \sum_a \mathcal{D}_a$  for its decomposition into irreducible components. In the SCFT case the  $\mathcal{D}_a$ 's are preserved by the  $\mathbb{C}^{\times}$ -action.

We focus on a point  $u \in \mathscr{C} \setminus \mathcal{D}$  which is at distance  $\epsilon$  from a generic point  $u_0 \in \mathcal{D}_a$  of the a-th discriminant component. The BPS states that are massless at  $u_0$  are parametrically light at u; since  $\mathcal{D}_a$  has codimension-1 in  $\mathscr C$  these states are charged under a rank-1 gauge group. Thus, asymptotically as  $u \to u_0$  the light degrees of freedom are described by an effective IR  $\mathcal{N} = 2$  theory which looks like a system composed of  $(r - 1)$  IR-free vectormultiplets and a rank-1 model which can be either IR-free or an interacting SCFT in its own right. This suggests that the special geometry in an infinitesimal neighborhood of  $\mathcal{D}_a$ should look as a rank-1 special geometry describing the light particles fibered over the local rank- $(r-1)$  geometry which describes the heavy states. While the actual situation may be a bit more intricate,<sup>[6](#page-6-2)</sup> this physical intuition yields a first *rough* cartoon of the geometry

<span id="page-6-0"></span><sup>&</sup>lt;sup>4</sup> Special geometries with this additional structure will be called  $\mathbb{C}^{\times}$ -isoinvariant.

<span id="page-6-1"></span> $5$  Fairly new  $\equiv$  the author is not aware of any previous mention of this idea in the math literature.

<span id="page-6-2"></span> $6$  See [\[26\]](#page-150-1).

in the vicinity of  $u_0$ . Making the idea a little more precise, one defines a rank- $(r - 1)$ special geometry  $\mathscr{X}_a \to \mathcal{D}_a$  over the codimension-1 locus  $\mathcal{D}_a \subset \mathscr{C}$ . The local geometry in an infinitesimal neighborhood of  $\mathcal{D}_a$  is fibered over this rank- $(r-1)$  special geometry. Iterating the construction, one gets a rank- $(r - s)$  special geometry  $\mathscr{X}_{a_s} \to \mathscr{C}_{a_s}$  over each subvariety in a finite set  $\{\mathscr{C}_{a_s}: a_s \in A_s\}$  of codimension-s subvarieties of  $\mathscr{C}$  for all  $0 \leq s \leq r$ . By the "stratification" of the special geometry  $\mathscr{X} \to \mathscr{C}$  we mean the full collection of special geometries

$$
\{\mathscr{X}_{a_s} \to \mathscr{C}_{a_s} : a_s \in A_s, \ 0 \le s \le r\}.\tag{1.1}
$$

The open sets  $\mathscr{C}_{a_s} \equiv \mathscr{C}_{a_s} \setminus (\cup_{a_{s-1} \in A_{s-1}} \mathscr{C}_{a_{s-1}}) \cap \mathscr{C}_{a_s}$  form a stratification of the Coulomb branch in the usual sense. The closures of the codimension- $(s + 1)$  strata,  $\mathscr{C}_{a_{s+1}} \subset \mathscr{C}$ , are the irreducible components of the discriminants of the special geometries  $\mathscr{X}_{a_s} \to \mathscr{C}_{a_s}$  over the (closed) codimension-s strata. In particular the  $\mathscr{C}_{a_1}$ 's coincide with the discriminant component  $\mathcal{D}_{a_1}$  of  $\mathscr{X}$ . The geometry along a codimension-1 stratum  $\mathcal{D}_a$  is determined by the singular fiber  $\mathscr{X}_{u_{\star}}$  at a generic point  $u_{\star} \in \mathcal{D}_a$ . The most important invariant of a divisor  $\mathcal{D}_a$  is the conjugacy class  $[\varrho_a]$  of the local monodromy  $\varrho_a$  around it; the type of the fiber  $\mathscr{X}_{u_\star}$ is determined by  $[\varrho_a]$  together with some subtler invariants [\[45–](#page-151-3)[50\]](#page-151-4). The local monodromy classes  $[\varrho_a]$  follows a Kodaira-like ADE classification [\[51–](#page-151-5)[54\]](#page-151-6). Thus to each component  $\mathcal{D}_a$  we can associate the (unique) rank-1 *asymptotic* special geometry over the (small) disk  $\Delta$  with the same monodromy class. Roughly speaking, the effective IR field theory which governs the degrees of freedom which are light along  $\mathcal{D}_a$  is described by this rank-1 geometry defined by the local monodromy  $\gamma_a$ . The detailed geometry/physics is a bit trickier [\[26\]](#page-150-1) and depends on the particular point  $u<sub>\star</sub>$ , not just on the discriminant component  $\mathcal{D}_a$ , but the most important aspects are already captured by  $|\gamma_a|$ .

The Coulomb branch  $\mathscr C$  carries a second stratification by unbroken  $U(1)_R$  symmetry. At a generic point in C the unbroken subgroup is  $\mathbb{Z}_k$  with  $k \in \{1, 2, 3, 4, 6\}$  [\[25\]](#page-150-7).<sup>[7](#page-7-0)</sup> The codimension-1 loci in  $\mathscr C$  where the unbroken R-symmetry enhances to a bigger group form a divisor  $E = \sum_{f} E_f$ . The unbroken R-symmetry at the generic point of  $E_f$  is  $\mathbb{Z}_{\ell_f}$  where  $\ell_f$ is a proper multiple of k. On multiple intersections  $\cap_j E_{f_i}$  we have further enhancement to  $\mathbb{Z}_{\text{lcm}(\ell_{f_j})}$ , etc. Some  $D_a$  may coincide with some  $E_f$  (case (3)).

By the *stratification data* of a  $\mathbb{C}^{\times}$ -isoinvariant special geometry we mean

<span id="page-7-1"></span>
$$
\{\mathcal{S}_a, [\varrho_a], \ell_a\}_{a \in A} \tag{1.2}
$$

where  $\{S_a\}_{a\in A}$  is the finite set of (C<sup>×</sup>-invariant) special divisors in  $\mathscr C$  where something interesting happens, and  $[\varrho_a]$  (resp.  $\ell_a$ ) is the conjugacy class of the local monodromy around  $\mathcal{S}_a$  (resp. the order of the unbroken R-symmetry along  $\mathcal{S}_a$ ).

<span id="page-7-0"></span><sup>&</sup>lt;sup>7</sup> If  $k \notin \{1,2\}$  the model is isotrivial and the geometry and physics can be determined by group-theoretical methods [\[25,](#page-150-7) [55\]](#page-151-7).

Inverse problem. The very fact that all special geometry has a "stratification" whose non-trivial strata are special geometries of strictly smaller dimension, suggests an inverse engineering strategy to construct all special geometries which, in particular, could lead to a complete classification of all 4d  $\mathcal{N}=2$  SCFT. The idea is that, once we have constructed all  $\mathbb{C}^{\times}$ -isoinvariant special geometries up to rank r, we can construct the rank  $r + 1$  ones by "gluing together" a finite set of rank-r geometries together with a rank- $(r + 1)$  open stratum  $\mathscr{C} = \mathscr{C} \setminus \mathscr{S}$  of "boring" generic vacua whose geometry is well understood. In addition to the stratification data [\(1.2\)](#page-7-1), the problem then requires to specify the generic vacuum which is encoded in two data: the action of the R-symmetry  $U(1)_R$  on  $\check{\mathscr{C}}$  and the generic Higgs branch. For smooth Coulomb branches the first datum is more conveniently replaced by the r-tuple of the Coulomb dimensions  $\{\Delta_i\}$ . For simplicity we assume that the generic Higgs branch is trivial, but this is not an essential assumption for our analysis.

This inverse engineering program is what we call "the inverse problem of special geometry". It requires to answer two main questions:

Question 1. Characterize the sets of the form  $\{(\Delta_i)_{i=1}^r,(\mathcal{S}_a,[\varrho_a],\ell_a)_{a=1}^s\}$  where  $\{\Delta_i\}$  are positive rational numbers,  $S_a \in \mathbb{C}^r$  are homogeneous irreducible divisors,  $[\varrho_a]$  Kodaira types, and  $\ell_a$  positive integers, which are the stratification data of indecomposable,<sup>[8](#page-8-0)</sup> rank-r, global special geometries.

Question 2. When  $\{(\Delta_i)_{i=1}^r, (\mathcal{S}_a, [\varrho_a], \ell_a)_{a=1}^s\}$  is an allowed stratification datum, construct all global special geometries that have this "stratification".

The answer to Question 1 may be given at different levels. There are necessary conditions on  $\{(\Delta_i)_{i=1}^r,(\mathcal{S}_a,[\varrho_a],\ell_a)_{a=1}^s\}$  for the existence of a global special geometry and also sufficient conditions. Necessary conditions are physical properties which all  $4d \mathcal{N} = 2 \text{ SCFT}$ should enjoy, and hence are statements of direct physical interest. Sufficient conditions produce examples which may lead to new insights. It would be desirable to have a precise necessary and sufficient condition. In this paper we shall come "close" to this goal but some work is still needed to get the complete story. The trickier part of **Question 1** is the classification of the special divisors  $\sum_a \mathcal{S}_a$  which may arise in a special geometry. This aspect is the main topic of section [9.](#page-122-1) When the base  $\mathscr C$  of the (algebraic) Lagrangian fibration is compact, i.e.  $\mathbb{P}^r$ , the possible discriminants are well-understood [\[56\]](#page-151-8), but in our set-up  $\mathscr{C}$  is affine, not projective, and a priori the discriminant is much less constrained.

Answers. In this long note we shall give essentially complete answers to the two Questions in rank-2. For higher rank we give some necessary conditions, discuss general aspects and properties, provide examples, and pave the way for a detailed study (which should be doable at least for low-ranks such as 3 or 4).

<span id="page-8-0"></span><sup>&</sup>lt;sup>8</sup> A geometry is indecomposable if it is not the product of lower-rank geometries, i.e. if the corresponding SCFT is does not split in two non-interacting sectors. The notion of "indecomposable" should not be confused with the notion of "irreducible".

**Rigidity.** Our basic tool to answer the two Questions is *rigidity* of various geometric structures implied by the definition of special geometry. Several inequivalent notions of rigidity are relevant in the present context: the "weaker one" (Faltings-Peter rigidity [\[57–](#page-151-9)[62\]](#page-152-0)) applies to general (irreducible, non-isotrivial) special geometries, while the stronger Deligne-Simpson rigidity [\[63](#page-152-1)[–65\]](#page-152-2) refers to the underlying monodromy representation and yields a typically sufficient condition. In addition we have (non)-rigidity in the special geometric sense, see section [5.](#page-54-0)

Classification scheme. The arguments in this paper suggest the classifying scheme:



Class I can be explored by elementary means for all rank  $r$  [\[55\]](#page-151-7). Classes II and III are severely constrained, "few" such geometries exist, and a systematic analysis may be possible also in higher rank. These geometries look rather "bizarre", but there is strong evidence that a small number of them do exist. In view of the **Folk-theorem**  $(\S, 5.2)$  $(\S, 5.2)$  class IV is also essentially charted for all ranks. For classes V and VI one falls back to the rank by rank strategy *via* the *inverse method*. One tries to construct them by induction on the rank by "gluing together" known lower rank "strata". Class VI is the less amenable to analysis and a priori the largest one, but not a single example of class VI geometry is known to the author, so there is hope that this hardest class is actually empty or at least very sparse.

Each one of the classes I-V is naturally decomposed into subclasses which are conveniently studied separately.

Organization. The paper is organized as follows. Section 2 contains a review of the basic geometric structures with new results and complements perhaps known to the experts but without a good reference in print. In section 3 we state the inverse problem and explain its relation to the classification program. In section 4 we discuss the families of Abelian varieties which arise from a  $\mathbb{C}^{\times}$ -isoinvariant special geometry. In section 5 we discuss the deformation theory of special geometry, the conformal manifold, and the Folk-theorem. To illustrate the main issues here we also discuss in detail the case of rank 2. In section 6 we consider the rigidity of implied geometric structures such as the Abelian family and the monodromy representation. Here we state the Deligne-Simpson problem. In section 7 we discuss how rigidity can be used to understand/construct/classify  $\mathbb{C}^{\times}$ -isoinvariant special geometries. In section 8 the program is carried on in rank 2 in detail. Here several tricky points are analyzed such as reducible representations. Finally in section 9 we lay the groundwork for an extension of the program to higher rank, focusing on the problem of understanding the class of allowed discriminant divisors. Some technicality is deferred to the appendices.

## <span id="page-10-0"></span>2 Special Geometry: review with complements

In this section we review special geometry to fix notation and terminology. We also present some complements which, while probably known to experts, are not spelled out in the literature (to the best of our knowledge). Some results are novel, at least in the present generality.

## <span id="page-10-1"></span>2.1 Defining structures

We first review the geometric structures implied by special geometry. We start with a rough sketch of the physical foundations of the topic. We omit all details and gloss over all subtle points since our purpose here is merely to motivate the precise definitions given in the next subsection  $(\S, 2.2)$  $(\S, 2.2)$  which we shall take as the geometric "axioms" for the rest of the paper.

An Euclidean 4d  $\mathcal{N} = 2$  SQFT admits a topological twisting *á la* Witten [\[66,](#page-152-3)[67\]](#page-152-4) by modifying the couplings of the various fields to the holonomy connection of the 4-dimensional spacetime manifold  $M$ . The twisted model is a Topological Field Theory (TFT) invariant under all oriented diffeomorphisms  $M \to M$  (see [\[68\]](#page-152-5) for an extensive review). By construction the twisting does not change the functional measure when  $M$  is locally flat.

#### <span id="page-10-2"></span>2.1.1 Chiral ring, Coulomb branch, Coulomb dimensions

The algebra of local operator of the twisted TFT is a subquotient  $\mathscr R$  of the algebra  $\mathfrak A$  of local operators of the parent  $\mathcal{N} = 2$  SQFT called the *chiral ring*. The correlation functions of operators in  $\mathcal R$  are independent of the choice of their representatives in  $\mathfrak A$ .  $\mathcal R$  is a commutative, associative C-algebra with unit, which in a sensible  $SQFT<sup>9</sup>$  $SQFT<sup>9</sup>$  $SQFT<sup>9</sup>$  must be a finitely-generated integral domain. Its spectrum  $\mathscr{C} = \mathsf{Spec} \mathscr{R}$  is then an *affine variety* over  $\mathbb C$  called the *Coulomb* branch. Its complex dimension  $r = \dim_{\mathbb{C}} \mathscr{C}$  is the rank. The basic observation is that when the Riemannian spacetime M is flat the correlation functions of operators in  $\mathscr R$  computed in the SQFT coincide with the ones computed in the corresponding twisted TFT.[10](#page-10-4)

By replacing  $\mathscr R$  with its normalization [\[70\]](#page-152-6), if necessary, we assume  $\mathscr C$  to be normal. In particular, the Coulomb branch is regular (as an affine variety<sup>[11](#page-10-5)</sup>) in codimension 1.

<span id="page-10-3"></span> $9 \text{ A}$  non finitely-generated chiral ring will violate the entropy bound any good QFT should obey.

<span id="page-10-4"></span><sup>&</sup>lt;sup>10</sup> This equality was the starting point of  $tt^*$  [\[69\]](#page-152-7).

<span id="page-10-5"></span><sup>&</sup>lt;sup>11</sup> We stress that some *other* relevant geometric structures on  $\mathscr{C}$ , such as the special Kähler metric, are

We are mainly concerned with the situation where the original  $\mathcal{N} = 2$  SQFT was a SCFT. In this special case the C-algebra  $\mathscr R$  is graded by the scaling dimension  $\Delta$ , that is:

<span id="page-11-1"></span>
$$
\mathcal{R} = \bigoplus_{\Delta \in \frac{1}{m}\mathbb{Z}_{\geq 0}} \mathcal{R}_{\Delta}, \qquad \mathcal{R}_{0} = \mathbb{C}, \qquad \mathcal{R}_{\Delta} \cdot \mathcal{R}_{\Delta'} \subseteq \mathcal{R}_{\Delta + \Delta'}, \tag{2.1}
$$

for some minimal positive integer m.

**Definition 1.** In this paper<sup>[12](#page-11-0)</sup> by a *chiral ring*  $\mathcal{R}$  of rank r we shall always mean a normal affine C-algebra of dimension r with a grading as in [\(2.1\)](#page-11-1). By its Coulomb branch  $\mathscr C$  we mean the normal affine r-variety  $\text{Spec } \mathcal{R}$  (or its underlying complex r-fold).

By definition the Coulomb branch  $\mathscr C$  is an affine quasi-cone with an algebraic  $\mathbb C^{\times}$ -action. The maximal ideal  $\mathscr{R}_+ \equiv \oplus_{\Delta>0}\mathscr{R}_\Delta$  defines a closed point  $0 \in \mathscr{C}$ , the tip of the quasi-cone, which physically represents the unique vacuum where the superconformal symmetry is not spontaneously broken.  $\{0\}$  is the only *closed*  $\mathbb{C}^{\times}$ -orbit, hence it is contained in the closure of all  $\mathbb{C}^{\times}$ -orbits.

**Smooth vs. non-smooth Coulomb branches.** The Coulomb branch  $\mathscr C$  is smooth at 0 (hence everywhere) iff the chiral ring is regular at the tip i.e.

$$
\dim(\mathcal{R}_+/\mathcal{R}_+^2) = \dim \mathcal{C} \equiv r,\tag{2.2}
$$

in which case  $\mathscr{R} = \mathbb{C}[u_1, \dots, u_r]$  where  $u_i$  has dimension  $\Delta_i \in \frac{1}{m}$  $\frac{1}{m}\mathbb{Z}_{>0}$ .

In the physical applications one usually assumes the affine quasi-cone  $\mathscr C$  to be smooth because the typical examples of Coulomb branches which arise in physics are smooth affine varieties. However, as it will be clear in the following, to construct a deep theory (i.e. to really understand the smooth case) one needs to allow for non-smooth Coulomb branches as well. The situation leads to two questions: (1) Which Coulomb branch singularities are allowed? (2) What is the physical meaning of a singular Coulomb branch? A plausible physical mechanism for a singularity is a "discrete gauging" which leads to Coulomb branches of the form  $\mathbb{C}^r/G$  with G a finite group. In this paper we shall assume the Coulomb branch to be smooth, except when explicitly stated otherwise.

**Coulomb dimensions.** When *C* is smooth, hence a copy of  $\mathbb{C}^r \simeq$  Spec  $\mathbb{C}[u_1, \ldots, u_r]$ with  $u_i \in \mathscr{R}_{\Delta_i}$ , the r-tuple of rational numbers  $\{\Delta_1, \ldots, \Delta_r\}$  is called the *Coulomb dimen*sions. We shall deduce various restrictions on the allowed Coulomb dimensions below: only finitely many r-tuples of Coulomb dimensions are allowed for a given  $r$  [\[22\]](#page-150-0).

typically non-regular along a divisor in the geometries we are interested in. This caveat applies to the question of smoothness in the next paragraph.

<span id="page-11-0"></span><sup>12</sup> Unless explicitly stated otherwise !!

The projective Coulomb branch  $\mathscr{P}$ . In the SCFT case it is natural to replace the ordinary Coulomb branch  $\mathscr{C} \equiv \text{Spec } \mathscr{R}$  by its projective counterpart

$$
\mathscr{P} = \text{Proj } \mathscr{R}.
$$
\n
$$
(2.3)
$$

When the Coulomb branch is smooth,  $\mathscr P$  is a weighted projective space [\[71\]](#page-152-8)

$$
\mathscr{P} \equiv \mathbb{P}(d_1, \dots, d_r), \qquad d_i = \frac{m \,\Delta_i}{\gcd\{m \,\Delta_j\}}.\tag{2.4}
$$

In particular  $\mathscr P$  is simply-connected  $\pi_1(\mathscr P)=1$ .  $\pi_1(\mathscr P)$  is at most finite even when  $\mathscr C$  is not smooth: indeed it is a quotient of the finite group F. In the smooth case  $\mathbb{P}(d_1, \dots, d_r)$ is isomorphic to a unique well-formed weighted projective space  $\mathbb{P}(q_1, \dots, q_r)$  such that  $gcd(q_1, ..., \hat{q}_j, ..., q_r) = 1$  for all j [\[71\]](#page-152-8).

#### <span id="page-12-0"></span>2.1.2 The Abelian fibration

The arguments in this subsection are heuristics to motivate the definitions in §. [2.2](#page-13-0) below.

By their very definition in all TFT the RG flow is trivial. Hence all topological correlation functions can be exactly computed using the low-energy effective theory [\[72,](#page-152-9)[73\]](#page-152-10) which, roughly speaking, is a supersymmetric  $\sigma$ -model with target space the set of SUSY vacua. We say "roughly speaking" because the vacuum space is typically singular, and one should be careful with the precise definition of the low-energy theory. For us the main point is that there are several distinct "IR effective theories" and we may use any one of them to compute TFT quantities. Independence of the TFT amplitudes from the choice of effective description implies compatibility conditions between these effective theories which we realize as additional geometric structures on  $\mathscr{C}$ .

Consider, for instance, the 4-manifold  $M \equiv \mathbb{R}^3 \times S^1$  where the circle has radius R. The local TFT observables are equal to their SQFT siblings (since M is flat) and independent of R by topological invariance. As  $R \to \infty$  they may be computed using the 4d IR effective  $\mathcal{N} = 2$ SQFT [\[72,](#page-152-9)[73\]](#page-152-10), while as  $R \to 0$  the very same local TFT observables may be computed using the 3d IR effective SQFT which now is a 3d  $\mathcal{N} = 4$  SUSY theory.<sup>[13](#page-12-1)</sup> We take for granted that the 3d Coulomb branch  $\mathscr X$  satisfies the same "regularity" conditions as its 4d counterpart  $\mathscr{C}$ , and hence that  $\mathscr{X}$  is a smooth<sup>[14](#page-12-2)</sup> scheme of finite type over  $\mathbb{C}$  (in facts over  $\mathscr{C}$ ). This follows from the physical heuristics sketched below, as well as from the usual special geometry folklore in the physical literature.<sup>[15](#page-12-3)</sup> On SUSY grounds [\[33\]](#page-150-3), the 3d Coulomb branch  $\mathscr X$  is a holomorphic symplectic manifold of quaternionic dimension  $r$  (complex dimension  $2r$ ):

<span id="page-12-1"></span><sup>&</sup>lt;sup>13</sup>  $\hat{\mathcal{L}}$  The physical low-energy effective theory depends in an intricate way from R [\[74\]](#page-152-11). Since the topological theory is independent of R, we are free to fix R in any convenient way. We use the asymptotic limit  $R \to \infty$ where things simplify dramatically.

<span id="page-12-3"></span><span id="page-12-2"></span><sup>&</sup>lt;sup>14</sup> Just as for the Coulomb branch, the condition that  $\mathscr X$  is smooth may be relaxed.

<sup>&</sup>lt;sup>15</sup> Of course, if  $\mathscr X$  is not an algebraic scheme (but still smooth), essentially everything we say will remain true in the analytic category.

indeed on the *real* tangent bundle  $T_{\mathbb{R}}\mathscr{X}$  we have a H-action induced by the 3d  $\mathcal{N}=4$  Rsymmetry which endows  $\mathscr X$  with a torsionless  $Sp(r, \mathbb C)$ -structure. We write  $\Omega$  for its  $(2, 0)$ holomorphic symplectic form. The compatibility of the TFT computation from the 4d and 3d viewpoints requires the algebra isomorphism

$$
\mathcal{R} \equiv \Gamma(\mathcal{C}, \mathcal{O}_{\mathcal{C}}) \simeq \Gamma(\mathcal{X}, \mathcal{O}_{\mathcal{X}}),\tag{2.5}
$$

which produces the canonical scheme-theoretic fibration

$$
\pi \colon \mathscr{X} \to \operatorname{Spec} \Gamma(\mathscr{X}, \mathcal{O}_{\mathscr{X}}) \simeq \mathscr{C}
$$
\n
$$
(2.6)
$$

over the Coulomb branch  $\mathscr C$ . The compatibility of the actions of the 3d and 4d R-symmetries implies that the fibers of  $\pi$  are Lagrangian submanifolds, i.e.  $\Omega|_{\mathscr{X}_u} = 0$  for all  $u \in \mathscr{C}$ . The map  $\pi$  has an obvious physical meaning in terms of the RG flow from  $R = 0$  to  $R = \infty$ . We claim that X is a commutative group-scheme over  $\mathscr{C}$ . In particular the fiber  $\mathscr{X}_u$  over the *generic* point u is a commutative algebraic group. Then  $\pi$  has a section s given by the zero of the group,  $s(u) = 0_u \in \mathscr{X}_u$ , and  $\mathscr{C}$  can be identified with the submanifold  $s(\mathscr{C}) \subset \mathscr{X}$  which should also be Lagrangian by comparison of R-symmetry actions in 3d and 4d. The claim can be easily checked in Lagrangian SCFTs and hence in full generality at generic points in  $\mathscr{C}$ : this suffices to establish that  $\mathscr{X}$  is a (smooth) birational model of a commutative group-scheme over  $\mathscr{C}$ . In the usual physical situation where the electromagnetic charges obey a Dirac quantization, the general fiber is actually a polarized Abelian variety. These physical requirements lead to the "conventional" special geometries which are the object of these notes. More exotic possibilities will be explored elsewhere.

## <span id="page-13-0"></span>2.2 Basic definitions

We codify the above informal discussion in a definition.

**Definition 2.** A rank-r special geometry is a holomorphic fibration  $\pi$  with section s

$$
\mathscr{X} \xrightarrow[s]{\pi} \mathscr{C}, \qquad \pi s = \text{Id}_{\mathscr{C}}, \tag{2.7}
$$

over a complex (normal) affine variety  $\mathscr C$  of dimension r, such that:

- (a) the total space  $\mathscr X$  has a holomorphic symplectic structure  $\Omega$ , and both the fibers  $\mathscr X_u$ of  $\pi$  and the section s are Lagrangian submanifolds:  $\Omega|_{\mathscr{X}_u} = \Omega|_s = 0;$
- (b) the generic fiber  $\mathscr{X}_u$   $(u \in \mathscr{C})$  is smooth and a polarized Abelian variety of dimension r with zero element  $s(u)$ .

The locus  $\mathcal{D} \subset \mathscr{C}$  of points u with non-smooth fiber  $\mathscr{X}_u$  is called the *discriminant*.

Although it is not part of the definition, we shall assume the polarization of the fiber to be principal for definiteness. Physically this is the most natural situation.

With this definition, a special geometry is (in particular) a birational model of an Abelian scheme over a (quasi-)projective variety, that is, of an Abelian variety defined over the function field  $\mathbb{C}(\mathscr{C})$  (e.g. in rank-1  $\mathscr{X}$  is the *Néron model* [\[75\]](#page-152-12) of an elliptic curve over the field of rational functions on a curve  $\overline{\mathscr{C}} \simeq \mathbb{P}^1$ .  $\mathscr{X}$  is also an Algebraic Integrable System where the Hamiltonians in involution are the holomorphic functions on  $\mathscr{C}$ , i.e. the chiral ring<sup>[16](#page-14-0)</sup>  $\mathscr{R}$ , while the conjugate angle variables are the doubly periodic coordinates along the fibers. Two weaker notions are often used:

- a non UV-complete special geometry is a fibration  $\mathscr{X} \to \mathscr{C}$  which satisfies all the properties of a special geometry except that  $\mathscr C$  is not quasi-projective. A non UV-complete geometry may or may not have a UV completion, i.e. an analytic extension over some affine base. Non UV-complete special geometries describe low-energy effective  $\mathcal{N}=2$ QFTs which are not UV-complete, and may or may not have a completion;
- a special Kähler geometry satisfies all the local conditions predicted by the  $\mathcal{N} = 2$ superspace approach to supersymmetry: existence of a local pre-potential  $\mathcal{F}$ , existence of a special Kähler metric on the Coulomb branch, etc., but whose underlying global structures (such as the VHS<sup>[17](#page-14-1)</sup>) are not defined over  $\mathbb{Q}$ , or some other number field  $\mathbb{F}$ , but over a local field such as  $\mathbb R$  or  $\mathbb C$ .

We stress that the results in this paper refer only to special geometries in the *strict* sense. In particular for us it will be absolutely essential that the Coulomb branch  $\mathscr C$  is quasi-projective. Most of our statements are *just false* when this condition does not hold.

Definition 3. A special geometry is called:

- decomposable if it is the product of two lower-rank special geometries. Decomposable geometries describe QFTs which decompose in two non-interacting sub-systems;
- free if the fibration is trivial i.e.  $\mathscr{X} = A \times \mathscr{C}$  for a fixed Abelian variety A. Free geometries describe free  $\mathcal{N} = 2$  QFT;
- *isotrivial* if all *smooth* fibers are isomorphic to a fixed polarized Abelian variety  $A$ . All  $\mathcal{N} \geq 3$  QFTs have isotrivial geometries, but there are also genuine  $\mathcal{N} = 2$  interacting SCFT with isotrivial geometries. The structure of isotrivial geometries is discussed in §. [2.10.4](#page-41-0) below;
- quasi-isotrivial if all smooth fibers are isogeneous to a product  $A \times Y_u$  where the first factor, A, is a fixed polarized Abelian variety while  $Y_u$  depends on  $u \in \mathscr{C}$ ;

<span id="page-14-0"></span><sup>16</sup> More precisely its Frechét completion.

<span id="page-14-1"></span><sup>&</sup>lt;sup>17</sup> VHS=variation of Hodge structure.

• unitary if it is complete as a Hamiltonian system, that is, iff for all Hamiltonian  $h \in \mathscr{R}$ the corresponding Hamiltonian vector field  $v(h)$ 

$$
\iota_{v(h)}\Omega \stackrel{\text{def}}{=} dh \tag{2.8}
$$

is complete, i.e. iff the solution of the (holomorphic) Hamilton equations

$$
\frac{dx^i}{dt} = v(h)^i \tag{2.9}
$$

exists for all times  $t \in \mathbb{C}$  and initial condition. The special geometries of unitary QFTs are unitary in this geometric sense. It is amusing that quantum unitarity of the  $\mathcal{N}=2$ QFT amounts to classical "unitarity" of the underlying Hamiltonian system.

 $\hat{\mathcal{L}}$  Caveat 1. In §.[2.8](#page-26-0) we introduce the notion of *reducible* special geometry. While decomposable implies reducible, there exist indecomposable geometries which are not irreducible. However their structure is very constrained, and their number rather small.

**Fact 1.** The discriminant D is either empty or of pure codimension 1. When  $D = \emptyset$  the geometry is free. Otherwise there are finitely-many elements  $p_a \in \mathcal{R}$  with  $(p_a)$  prime, such that  $\mathcal{D} = \{p_1p_2\cdots p_s = 0\} \subset \mathscr{C}$ .

*Argument.* The first assertion is e.g. **Proposition 3.1(2)** in [\[45\]](#page-151-3). The second one follows from the well-known fact that all families of smooth projective varieties parametrized by C are trivial (cf. §. 13.7 of [\[40\]](#page-151-10)). The third statement is well known.  $\Box$ 

The above definitions refer to the geometry of a general 4d  $\mathcal{N}=2$  SQFT. The geometries of  $\mathcal{N} = 2$  SCFTs have an extra geometric structure.

<span id="page-15-1"></span>**Definition 4.** A  $\mathbb{C}^{\times}$ -isoinvariant special geometry is a special geometry with a complete holomorphic vector field  $\mathcal{E} \in \Gamma(\mathcal{X}, T\mathcal{X})$ , called the *Euler vector*, such that for all  $z \in \mathbb{C}$ the exponential map  $\exp(z \mathcal{E})$ :  $\mathcal{X} \to \mathcal{X}$  is a complex automorphism preserving all special geometric structures: i.e.  $\exp(z \mathcal{E})$  preserves set-wise the section s and maps fibers into fibers,  $while<sup>18</sup>$  $while<sup>18</sup>$  $while<sup>18</sup>$ 

<span id="page-15-2"></span>
$$
\mathcal{L}_{\mathcal{E}}\Omega = \Omega, \quad \text{that is,} \quad \exp(z\,\mathcal{E})^*\Omega = e^z\,\Omega. \tag{2.10}
$$

The Euler vector  $\mathcal E$  induces a grading of the chiral ring  $\Gamma(\mathcal X, \mathcal O_{\mathcal X}) \simeq \mathcal R = \bigoplus_{\Delta} \mathcal R_{\Delta}$  by

$$
h \in \mathcal{R}_{\Delta} \quad \Leftrightarrow \quad \mathcal{L}_{\mathcal{E}} h = \Delta h. \tag{2.11}
$$

A Hamiltonian  $h \in \mathcal{R}_{\Delta}$  is said to have *dimension*  $\Delta$ . The identity has dimension zero. We shall see momentarily that in a unitary special geometry the identity 1 is the only element of  $\mathscr R$  with dimension zero, while all other Hamiltonians have  $\Delta \geq 1$ . We shall show below that

<span id="page-15-0"></span><sup>&</sup>lt;sup>18</sup> Here and below  $\mathscr{L}_v$  stands for the Lie derivative in the direction of the vector field v.

the  $\Delta$ 's are rational numbers and that only a short list of rational numbers may appear at a given rank r. Then  $\mathscr R$  is a finitely generated local graded C-algebra of the expected form [\(2.1\)](#page-11-1). When the Coulomb branch is smooth  $\mathscr{R} = \mathbb{C}[u_1, \dots, u_r]$  and the coordinates have definite dimension  $\mathscr{L}_{\mathcal{E}} u_i = \Delta_i u_i$ . In particular  $\exp(z \mathcal{E})$  induces a complex automorphism of the Coulomb branch,  $\exp(z\Delta): \mathscr{C} \to \mathscr{C}$ , such that

$$
\pi \exp(z \mathcal{E}) = \exp(z \Delta) \pi, \qquad (2.12)
$$

which for smooth  $\mathscr C$  is simply  $u_i \mapsto e^{z\Delta_i}u_i$ . It follows from **Definition [4](#page-15-1)** that the isoclass of the fiber, as a polarized Abelian variety, is invariant along an orbit of the group of automorphisms  $\exp(z \Delta)$ 

for all 
$$
z \in \mathbb{C}
$$
,  $u \in \mathscr{C} \setminus \mathcal{D}$ :  $\mathscr{X}_{\exp(z\Delta)u} \simeq \mathscr{X}_u$  as polarized Abelian varieties. (2.13)

The kernel of group homomorphism  $exp(\cdot \mathcal{E}) : \mathbb{C} \to Aut(\mathcal{X})$  is  $2\pi im\mathbb{Z}$  where m is the integer in eq.[\(2.1\)](#page-11-1). Indeed the induced map  $\exp(2\pi i m \Delta)$  is the identity on  $\mathscr{C}$ , so  $\exp(2\pi i m \mathscr{E})$ fixes the zero element  $0_u$  of each Abelian fiber  $\mathscr{X}_u$ , while it acts as the identity on its Lie algebra because of the isomorphism  $\Omega: T_{0u} \mathscr{X}_u \simeq T_u^* \mathscr{C}$ , so it is the identity in each fiber. The automorphism group acting effectively on  $\mathscr X$  is  $\mathbb C^{\times} \simeq \mathbb C/2\pi im\mathbb Z$ . The prime divisors  $p_a$  in the discriminant D are irreducible homogeneous elements of R, i.e.  $\mathscr{L}_{\mathcal{E}} p_a = \Delta(p_a) p_a$  for some  $\Delta(p_a) \in \frac{1}{m} \mathbb{N}$ .

Remark 1. In the physical literature it is more common to require the existence of a Seiberg-Witten (SW) differential  $\lambda$  such that  $\Omega = d\lambda$ .  $\lambda$  is simply the image of the vector  $\mathcal E$  under the natural isomorphism  $\Omega\colon T\mathscr{X}\to T^*\mathscr{X}$ , and the two data  $\lambda$ ,  $\mathscr{E}$  are equivalent in the  $\mathbb{C}^{\times}$ isoinvariant situation. It is more convenient to work with  $\mathcal E$  since it makes manifest that  $\lambda$  corresponds to an automorphisms ( $\equiv$  a symmetry) of the special geometry which is fully determined by its symplectic structure.

#### <span id="page-16-0"></span>2.2.1 Geometric version of the unitarity bound

Fact 2. (1) In a unitary  $\mathbb{C}^{\times}$ -isoinvariant special geometry all homogeneous elements of  $\mathcal R$ except the identity, have dimension  $\Delta \geq 1$ . (2) In a unitary, indecomposable, non-free  $\mathbb{C}^{\times}$ -isoinvariant geometry the inequality is strict  $\Delta > 1$ .

Argument. (1) Let h be a non-constant Hamiltonian of dimension  $\Delta$ . dh is not zero and  $\mathscr{L}_{\mathcal{E}} dh = d\mathscr{L}_{\mathcal{E}} h = \Delta dh$ . The corresponding Hamiltonian vector  $v(h)$  is non-zero and satisfies

$$
\Delta i_{v(h)}\Omega = \Delta dh = \mathcal{L}_{\mathcal{E}} dh = \mathcal{L}_{\mathcal{E}}(i_{v(h)}\Omega) = i_{\mathcal{L}_{\mathcal{E}}v(h)}\Omega + i_{v(h)}\mathcal{L}_{\mathcal{E}}\Omega = i_{\mathcal{L}_{\mathcal{E}}v(h)}\Omega + i_{v(h)}\Omega \quad (2.14)
$$

or  $\mathscr{L}_{\mathcal{E}} v(h) = (\Delta - 1)v(h)$ . Therefore

$$
\exp(z\mathcal{E})_* v(h) = z^{\Delta - 1} v(h). \tag{2.15}
$$

If  $\Delta$  < 1 the Hamiltonian vector field  $v(h)$  diverges at the tip of the quasi-cone and so cannot be complete. (2) follows from the following result whose proof is sketched in appendix [A.](#page-139-1)

<span id="page-17-1"></span>**Lemma 1.** The connected component  $(\mathscr{X}_0^{\text{reg}})$ <sup>reg</sup> $0 \text{ of the smooth locus } \mathscr{X}_0^{\text{reg}} \subset \mathscr{X} \text{ of the fiber}$ over the origin 0 has the structure

$$
(\mathcal{X}_0^{\text{reg}})^0 = A_0 \times \mathbb{C}^{r-\ell}, \quad \ell \equiv \dim A_0 = \dim \mathcal{R}_{\Delta=1},\tag{2.16}
$$

where  $A_0$  is a polarized Abelian variety, while the smooth fibers have the form  $\mathscr{X}_u = A_0 \times \mathscr{Y}_u$ with  $\mathscr{Y}_u$  a polarized Abelian variety of dimension  $r - \ell$ .

Clearly the geometry is the product of the free geometry  $(A_0 \times \mathbb{C}^{\ell}) \to \mathbb{C}^{\ell}$  with a special geometry  $\mathscr{Y} \to \mathbb{C}^{r-\ell}$  with fiber  $\mathscr{Y}_u$  and no dimension 1 coordinate.  $\Box$ 

Eq.[\(A.1\)](#page-139-2) says that the fiber over the origin has *additive* (as contrasted to *multiplica*tive) degeneration, thus in a non-free  $\mathbb{C}^{\times}$ -isoinvariant geometry  $\mathscr{X}_{0}$  is always an unstable degeneration of an Abelian variety.

**Remark 2.** The structure of the fiber over the origin, eq.  $(A.1)$  is one of the many properties of the special geometry which, taken together, guarantee that the corresponding  $\mathcal{N} = 2 \text{ QFT}$ is free of Landau poles. For instance, in rank-1 it rules out singular fibers of Kodaira type  $I_n$ , which would correspond to  $\mathcal{N} = 2$  SQCD a formal QFT affected by Landau poles.

#### <span id="page-17-0"></span>2.2.2 Implied structures: metric, SW periods, prepotential

Let L be the fibers' polarization seen as a integral  $(1, 1)$  class which restricts to a Kähler class in each fiber. On the complement  $\mathscr{E} = \mathscr{C} \setminus \mathcal{D}$  of the discriminant we have the function

$$
K: u \mapsto i \int_{\mathcal{X}_u} L^{r-1} \lambda \wedge \lambda^* = i \sum_i \left( \int_{A_i} \lambda \int_{B^i} \bar{\lambda} - \int_{A_i} \bar{\lambda} \int_{B^i} \lambda \right) \tag{2.17}
$$

where  $\{A_i, B^j\}$  is a symplectic basis of 1-cycles on the fibers  $\mathscr{X}_u$  defined locally in some  $U \subset \mathscr{C}$  with respect to the polarization L and  $\lambda \equiv \iota_{\mathcal{E}} \Omega$ . The function K is smooth and pluri-subharmonic in  $\mathscr{C} \setminus \mathcal{D}$ , so K is the potential of a Kähler metric with Kähler form

$$
\omega = i \int_{\mathcal{X}_u} L^{r-1} \wedge \Omega \wedge \bar{\Omega}.
$$
 (2.18)

Except in the free case (when  $\mathcal{D} = \varnothing$ ) this "special Kähler" metric is not complete since it has singularities along the discriminant  $D$  which are at finite distance in the special metric.

We define the SW period  $\Pi = (a_i^D, a^j)^t$  to be the *local* functions in U

$$
a_i^D(u) = \int_{B^i} \iota_{\mathcal{E}} \Omega \big|_{\mathcal{X}_u}, \qquad a^j(u) = \int_{A_i} \iota_{\mathcal{E}} \Omega \big|_{\mathcal{X}_u}, \quad u \in U, \ i, j = 1, \dots, r \tag{2.19}
$$

so that

$$
K = i \Pi^{\dagger} \Omega \Pi, \quad \text{where} \quad \Omega = \begin{pmatrix} 0 & \mathbf{1}_r \\ -\mathbf{1}_r & 0 \end{pmatrix} . \tag{2.20}
$$

The periods are locally-defined holomorphic functions on  $\mathscr{C}$ . Locally on  $\pi^{-1}(\mathscr{C})$  we may find Darboux coordinates such that  $\Omega = da^i \wedge dz_i$  where  $dz_i$  are holomorphic differentials on the fiber normalized as

$$
\int_{A_i} dz_j = \delta^i{}_j, \qquad \int_{B^i} dz_j = \tau_{ij}.
$$
\n(2.21)

Then  $\iota_{\mathcal{E}}\Omega = a^idz_i$  and

$$
a_i^D = \int_{B^i} \iota_{\mathcal{E}} \Omega \big|_{\mathcal{X}_u} = \tau_{ij} \, a^j. \tag{2.22}
$$

In particular the A-periods  $(a^1, \ldots, a^r)$  are local complex coordinates in  $\mathscr{C}$ . Clearly

$$
\mathcal{L}_{\mathcal{E}} \Pi = \Pi \tag{2.23}
$$

so that on local functions on  $\mathscr{E}$  we have  $\mathcal{E} = a^i \partial_{a^i}$  and  $\mathscr{L}_{\mathcal{E}} \tau_{ij} = 0$  gives  $a^i \partial_{a^i} \tau_{jl} = 0$  so that

<span id="page-18-2"></span>
$$
da_i^D = d(\tau_{ij} a^j) = \tau_{ij} da^j \quad \Rightarrow \quad da_i^D \wedge da^i = 0. \tag{2.24}
$$

The last equation says that, locally in  $U \subset \mathscr{C}$  there exists a holomorphic function  $\mathcal{F}(a^j)$ , called the prepotential, such that

$$
\mathcal{L}_{\mathcal{E}}\mathcal{F} = 2\mathcal{F} \quad \text{and} \quad a_i^D = \frac{\partial \mathcal{F}}{\partial a^i}, \quad \tau_{ij} = \frac{\partial^2 \mathcal{F}}{\partial a^i \partial a^j}.
$$
 (2.25)

**Remark 3.** The special coordinates yield an affine structure to  $\mathscr{C}$ . This is the holomorphic version of the integral affine structure on the base of a Liouville integrable system [\[76\]](#page-153-0).

## <span id="page-18-0"></span>2.3 Unbroken R-symmetry and the "characteristic dimension"

In the smooth case it is convenient to see the r-tuple of rational numbers<sup>[19](#page-18-1)</sup> { $\Delta_1, \cdots, \Delta_r$ } as a point in the projective space over Q and write

$$
(\Delta_1, \Delta_2, \dots, \Delta_r) = \lambda(d_1, d_2, \dots, d_r) \tag{2.26}
$$

where  $(d_1, d_2, \ldots, d_r)$  are the unique positive integers with  $gcd(d_1, \ldots, d_r) = 1$  which represent them same point in  $\mathbb{P}^{r-1}(\mathbb{Q})$ . We shall refer to  $d_i$  as the *degree* of  $u_i$ , while  $\Delta_i \equiv \lambda d_i$  is its dimension.  $\lambda$  is a positive rational number which we write in minimal terms as

$$
\lambda = \frac{n}{m} \quad \text{with } \gcd(m, n) = 1. \tag{2.27}
$$

<span id="page-18-1"></span><sup>&</sup>lt;sup>19</sup> We assume the  $\Delta_i$  to be ordered in a non-decreasing sequence, i.e.  $\Delta_{i+1} \geq \Delta_i$ .

 $\lambda$  is a basic invariant of the  $\mathbb{C}^{\times}$ -isoinvariant special geometry. It is well-defined also for non-smooth Coulomb branches: if  $\{\phi_1, \ldots, \phi_\ell\}$  is a set of generators of  $\mathscr{R}$ , one sets

$$
(\Delta(\phi_1), \ldots, \Delta(\phi_\ell)) = \lambda \left( d(\phi_1), \ldots, d(\phi_\ell) \right) \tag{2.28}
$$

with the  $d(\phi_i)$  integral and coprime. By construction the automorphism  $\exp(2\pi i \mathcal{E}/\lambda)$  acts trivially on  $\mathscr{R}$ , so it fixes all points of  $\mathscr{C}$ , while it multiplies  $\Omega$  by a *n*-th root of unity

$$
\exp(2\pi i \mathcal{E}/\lambda)^* \Omega = e^{2\pi i m/n} \Omega,
$$
\n(2.29)

which depends only on m mod n. The root  $\zeta \equiv e^{2\pi im/n}$ , or equivalently the pair  $(n, m \mod n)$ with gcd(m, n) = 1, was called the *characteristic dimension* in [\[25\]](#page-150-7) and written  $\varkappa = n/(m)_n$ where  $\langle m \rangle_n$  is the unique positive integer  $\leq n$  congruent to m mod n. The physical meanings of  $m$  and  $n$  are obvious:

- (1) the group which acts effectively on the geometry is  $\mathbb{C}^{\times} = \exp[\mathbb{C}/(z \sim z + 2\pi im)];$
- (2) at the *generic* point in  $\mathscr{C}$  the  $\mathbb{C}^{\times}$  symmetry is spontaneously broken to  $\mathbb{Z}_n$ ;
- (3) at any point  $u \in \mathscr{C}$  the  $U(1)_R$  symmetry is broken to a group which contains  $\mathbb{Z}_n$ . Hence, a part for the tip 0, to a discrete group of the form  $\mathbb{Z}_{nk}$ .

One shows  $|25|$  that n can take only five values

<span id="page-19-1"></span>
$$
n \in \{1, 2, 3, 4, 6\},\tag{2.30}
$$

hence we have only 8 possible  $\varkappa$ 

<span id="page-19-2"></span>
$$
\varkappa \in \left\{1, \frac{6}{5}, \frac{4}{3}, \frac{3}{2}, 2, 3, 4, 6\right\} \tag{2.31}
$$

in one-to-one correspondence with the conjugacy classes of torsion elements of  $SL(2, \mathbb{Z})$ , or equivalently with the Kodaira elliptic fibers with semisimple monodromy [\[51,](#page-151-5) [52\]](#page-151-11). The possible values of  $\varkappa$  are just the allowed dimensions of the Coulomb dimension  $\Delta$  in a  $r = 1$ special geometry: they are in 1-to-1 correspondence with the semisimple Kodaira fibers, see table [1.](#page-20-0) The basic result of [\[25\]](#page-150-7) was that the Abelian fiber over the generic point  $u \in \mathscr{C}$ admits complex multiplication by the cyclotomic field  $\mathbb{Q}(\zeta) \equiv \mathbb{Q}(e^{2\pi i/\varkappa})$  and hence its isoclass is uniquely determined unless  $\mathbb{Q}(\zeta) \equiv \mathbb{Q}$ , i.e. unless  $\varkappa \in \{1, 2\}$ .

**Fact 3.** When  $\varkappa \notin \{1,2\}$  all smooth fibers are isomorphic as polarized Abelian varieties, and the full geometry is isotrivial, hence easily constructed and studied by simple group theory [\[25\]](#page-150-7). The statement holds independently of the assumptions that  $\mathscr C$  is smooth and that the polarization is principal.

For later reference we sketch a proof of these statements. Preliminary we recall a few well know facts about polarized Abelian varieties.<sup>[20](#page-19-0)</sup> Let  $A$  be a polarized Abelian variety

<span id="page-19-0"></span><sup>&</sup>lt;sup>20</sup> For the facts about complex Abelian varieties used here and below see [\[77\]](#page-153-1) expecially chapter 13.

Kodaira fiber	$\mathbf{I}_0$	$\mathbf{I}^*_0$	п	$II^*$	ш	$III^*$	TV	$\mathbf{IV}^*$
κ			$\frac{6}{5}$	6	흫	4	$\frac{5}{2}$	3
monodromy	$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$	$\begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \qquad \begin{bmatrix} 1 & 1 \\ -1 & 0 \end{bmatrix} \qquad \begin{bmatrix} 0 & -1 \\ 1 & 1 \end{bmatrix} \qquad \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \qquad \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \qquad \begin{bmatrix} 0 & 1 \\ -1 & -1 \end{bmatrix} \qquad \begin{bmatrix} -1 & -1 \\ 1 & 0 \end{bmatrix}$						
Euler no.		6		10	3	9		8
Lie algebra		$\mathfrak{so}(8)$		$E_8$	$\mathfrak{su}(2)$	$E_7\,$	$\mathfrak{su}(3)$	$E_{6}$

<span id="page-20-0"></span>Table 1: Semi-simple Kodaira fibers

and Aut(A) the group of automorphisms  $A \to A$  which preserve the polarization. Aut(A) is a finite group with two natural representations. First we have the Q-defined rational representation  $\chi$  acting on  $H_1(A,\mathbb{Z})\otimes_{\mathbb{Z}}\mathbb{Q}$ . For principal polarizations  $\chi$  is a group embedding  $\chi:$  Aut $(A) \to Sp(2r, \mathbb{Z})$ ; for a general polarization see e.g. chapter 2 of [\[78\]](#page-153-2). Then we have the analytic representation  $\sigma: \text{Aut}(A) \to GL(r, \mathbb{C})$  which acts on the holomorphic tangent space at 0,  $T_0A \equiv \text{Lie}(A)$ . We have the isomorphism of complex representations

$$
\chi \otimes_{\mathbb{Q}} \mathbb{C} \simeq \sigma \oplus \bar{\sigma}.\tag{2.32}
$$

Since  $\chi$  preserves the lattice  $H_1(A,\mathbb{Z}) \subset H_1(A,\mathbb{Q})$ , this implies that for all  $\eta \in \text{Aut}(A)$  the characteristic polynomial det $[z - \chi(\eta)]$  is a product of cyclotomic polynomials whose roots are the eigenvalues of  $\sigma(\eta)$  and their conjugates. The automorphism

$$
\xi \stackrel{\text{def}}{=} \exp(2\pi i \mathcal{E}/\lambda) : \mathcal{X} \to \mathcal{X} \tag{2.33}
$$

generates a  $\mathbb{Z}_n$  group of automorphisms of  $\mathscr X$  which fixes point-wise the Coulomb branch  $\mathscr C$  and hence acts trivially on its holomorphic cotangent bundle  $T^*\mathscr C$ .  $\xi$  then restricts to an automorphism  $\xi_u \colon \mathscr{X}_u \to \mathscr{X}_u$  of each fiber (smooth or not).

Let  $u \in \mathscr{C}$  be a generic point, and  $\mathscr{X}_u$  its smooth Abelian fiber. The symplectic structure gives a linear isomorphism

$$
\Omega \colon T_{s(u)} \mathcal{X}_u \to T_u^* \mathcal{C}.\tag{2.34}
$$

The automorphism  $\xi$  acts as the identity on  $T^*_{u}\mathscr{C}$ , hence by eq.[\(2.10\)](#page-15-2) as multiplication by  $\zeta^{-1}$  on  $T_{s(u)}\mathscr{X}_u$ , that is,

$$
\sigma(\xi_u) = \zeta^{-1} \cdot \mathbf{1} \in GL(r, \mathbb{C}) \quad \Rightarrow \quad \det[z - \chi(\xi_u)] = \left( (z - \zeta^{-1})(z - \zeta) \right)^r \in \mathbb{Z}[z] \tag{2.35}
$$

which says that the root of unity  $\zeta$  is either  $\pm 1$  or an imaginary quadratic integer. This proves eqs.[\(2.30\)](#page-19-1),[\(2.31\)](#page-19-2). The final step is to recall<sup>[21](#page-20-1)</sup> the fact that when multiplication by

<span id="page-20-1"></span><sup>&</sup>lt;sup>21</sup> See Corollary 13.3.5 in [\[77\]](#page-153-1).

 $\zeta = i$  or  $e^{2\pi i/3}$  is an automorphism of the Abelian variety A we have  $2^2$ 

$$
A = \overbrace{E_{\zeta} \times E_{\zeta} \times \cdots \times E_{\zeta}}^{r \text{ factors}} \tag{2.36}
$$

and the geometry is (in particular) isotrivial, hence well understood [\[25,](#page-150-7) [55\]](#page-151-7). It remains to study the two cases  $\varkappa = 1$  and  $\varkappa = 2$  where the geometry may or may not be isotrivial.

## <span id="page-21-0"></span>2.4 The index of an irreducible homogeneous divisor

By an irreducible homogeneous divisor in  $\mathscr C$  we mean an irreducible codimension-1 subvariety preserved by the  $\mathbb{C}^{\times}$ -action (the closure of the  $\mathbb{C}^{\times}$ -orbits it contains). An irreducible homogeneous divisor has the form  $\{h = 0\} \subset \mathscr{C}$  for some irreducible quasi-homogeneous<sup>[23](#page-21-2)</sup> Hamiltonian  $h \in \mathcal{R}$  of degree d, i.e.

$$
\mathcal{L}_{\mathcal{E}/\lambda}h = d h. \tag{2.37}
$$

E.g. for a smooth geometry  $h = h(u_i)$  is a polynomial in the  $u_i$ 's of degree d, i.e.

$$
h(\lambda^{d_i} u_i) = \lambda^d h(u_i) \quad \forall \ \lambda \in \mathbb{C} \text{ and some } d \in \mathbb{N}.
$$
 (2.38)

Consider the Hilbert series  $P_h(t)$  of the algebra  $\mathcal{R}/(h)$  (graded by degree not dimension!). E.g. for  $\mathbb{C}[u_1,\ldots,u_r]/(h)$ 

$$
P_h(t) = \frac{(1 - t^d)}{\prod_i (1 - t^{d_i})}.
$$
\n(2.39)

The enhancement index  $\nu(h)$  of the divisor h is the largest positive integer such that

$$
P_h(t) \in \mathbb{C}[[t^{\nu(h)}]]. \tag{2.40}
$$

The physical meaning of  $\nu(h)$  is:

$$
\nu(h) = \left[ \left( \begin{array}{c} \text{unbroken subgroup of } \mathbb{C}^{\times} \text{ at the} \\ \text{generic point in the } h = 0 \text{ locus} \end{array} \right) : \left( \begin{array}{c} \text{unbroken subgroup of } \mathbb{C}^{\times} \\ \text{at the generic point in } \mathscr{C} \end{array} \right) \right]
$$
(2.41)

i.e.  $\nu(h)$  measures the enhancement of the unbroken R-symmetry along the divisor h. A divisor h is enhanced (resp. non-enhanced) if  $\nu(h) > 1$  (resp.  $\nu(h) = 1$ ).

Remark 4. The authors of [\[17\]](#page-149-3) distinguish the divisors in unknotted vs. knotted ones. Often, but not always, this coincides with the distinction enhanced vs. non-enhanced.

**Definition 5.** An irreducible homogeneous divisor  $(h)$  is *knotted* iff the fundamental group of its complement  $\pi_1(\mathbb{C}^r \setminus (h))$  is non-Abelian.

<span id="page-21-1"></span><sup>&</sup>lt;sup>22</sup> Through out this paper  $E_{\tau}$  stands for the elliptic curve of period  $\tau$ .

<span id="page-21-2"></span><sup>23</sup> In the sequel we simplify 'quasi-homogenous' in 'homogeneous'.

Smooth divisors are automatically unknotted.

**Example 2.1.** For  $r = 2$  the irreducible homogeneous divisors have one of the two forms:

$$
u_i
$$
 for  $i = 1, 2$  or  $u_1^{d_2} - z u_2^{d_1}$  with  $z \in \mathbb{C}^\times$ . (2.42)

For divisors of the second kind  $\nu = 1$ , while  $\nu(u_i) = d_1 d_2 / d_i$ . The divisors of the second kind are knotted unless  $d_1 = 1$ , while  $u_i$  is never knotted.

In a general rank-r smooth geometry, the irreducible divisors of degree  $d_i \in \{d_1, \ldots, d_r\}$  have  $\nu = \gcd\{d_1, \ldots, d_i, \ldots, d_r\}$  while divisors with  $d \notin \{d_1, \ldots, d_r\}$  have  $\nu = 1$ .

#### <span id="page-22-0"></span>2.5 Coulomb dimensions I: CM-types along submanifolds

Suppose we have an irreducible reduced total intersection  $H = \{h_1 = \cdots = h_\ell = 0\}$  with an enhancement index  $\nu(H) \equiv \nu > 1$  which is not contained in the discriminant  $\mathcal{D}$ . The automorphism  $\eta \equiv \exp(2\pi i \mathcal{E}/(\nu \lambda))$  fixes all points of H, hence its acts as the identity on its cotangent bundle  $T^*H \subset T^*\mathscr{C}|_H$  while it acts by polarized automorphisms on the fibers  $\mathscr{X}_u$ for  $u \in H$ . We write  $N^*H \to H$  for the rank- $\ell$  conormal bundle to H; it is spanned by the differentials  $dh_a$   $(a = 1, ..., \ell)$ . Let  $u \in H$  be a generic point; by assumption the fiber  $\mathscr{X}_u$  is smooth. One has

$$
T_u^* \mathscr{C} = T_u^* H \oplus N^* H_u \tag{2.43}
$$

 $\eta_u$  acts on the first summand as the identity and on the second as multiplication by the  $\ell \times \ell$ matrix

$$
\text{diag}\{\exp(2\pi i m\Delta(h_k))/(n\nu))\}.
$$
\n(2.44)

Using the isomorphims  $\Omega: T_{s(u)}\mathscr{X}_u \to T_u^*\mathscr{C}$  we get the eigenvalues of its analytic representation  $\sigma(\eta_u)$  are

<span id="page-22-2"></span>
$$
\overbrace{(e^{-2\pi i m/(n\nu)}, \cdots, e^{-2\pi i m/(n\nu)} )}^{(r-\ell) \text{ times}}, e^{-2\pi i m(1-\Delta(h_1))/(n\nu)}, \cdots, e^{-2\pi i m(1-\Delta(h_\ell)/(n\nu)})
$$
(2.45)

This r-tuple of roots of unity, together with their conjugates, are the roots (with multiplicities) of a product of cyclotomic polynomials. Enforcing these conditions on all total intersections of elements of  $\mathscr R$  yield a large web of constraints on the possible n, m and  $\nu$  and in turn on the Coulomb dimensions  $\Delta_i$ . We stress that nothing depends on the assumption that  $\mathscr C$  is smooth (except for the  $\Delta_i$ 's).

We give some examples.

#### <span id="page-22-1"></span>2.5.1 Example: dimension formulae along regular axes [\[22\]](#page-150-0)

For simplicity from now on we assume  $\mathscr C$  smooth. Suppose that the *i*-th coordinate axis  $H_i = \{u_j = 0$  for  $j \neq i\}$  is not contained in the discriminant: we say that  $H_i$  is a regular

axis. Then  $\nu = d_i$  and the list of roots in [\(2.45\)](#page-22-2) becomes

<span id="page-23-1"></span>
$$
\left\{ \exp[2\pi i(\Delta_1 - 1)/\Delta_i], \exp[2\pi i(\Delta_2 - 1)/\Delta_i], \cdots, \exp[2\pi i(\Delta_r - 1)/\Delta_i] \right\}
$$
 (2.46)

Note that a priori once we know this ordered list of roots of 1,  $1/\Delta_i$  is determined only mod 1, but since in a unitary geometry  $0 < 1/\Delta_i \leq 1$  knowing the *i*-th root in the list suffices to determine  $\Delta_i$  uniquely. Instead  $\Delta_j$  for  $j \neq i$  is determined by [\(2.46\)](#page-23-1) only up to addition of an integral multiple of  $\Delta_i$ . The ambiguity<sup>[24](#page-23-2)</sup> is however small since only a finite list of dimensions are allowed for any given  $r$  (see §. [2.12\)](#page-43-0). Moreover often there are several regular axes  $H_j \not\subset \mathcal{D}$  and we get one such formula for the dimensions for each one of these axes, getting an overdetermined system which gets rid of all ambiguities.

When the *i*-th coordinate axis  $H_i \not\subset \mathcal{D}$  the list of r  $(n d_i)$ -th roots of unity in eq.[\(2.46\)](#page-23-1) is a subset of the 2r roots of a product of cyclotomic polynomials with the property that the union of this subset with its conjugate yields the full set of all 2r roots. This entails that

$$
\exp(-2\pi i/\Delta_i) \tag{2.47}
$$

is a root in a cyclotomic field of degree at most 2r. That is, the dimension  $\Delta_i$  of a regular axis has the form

$$
\Delta_i = \frac{n_i}{m_i} \text{ where } \phi(n_i) \le 2r \text{ and } \gcd(n_i, m_i) = 1 \quad 1 \le m_i < n_i \tag{2.48}
$$

so that there are only finitely many allowed dimensions at a given rank  $r$  [\[22\]](#page-150-0). We shall show in §. [2.12](#page-43-0) that this conclusion remains valid even when  $H_i \subset \mathcal{D}$ ; in facts in that case we get the stronger condition  $\phi(n_i) \leq 2(r-1)$ .

#### <span id="page-23-0"></span>2.5.2 New dimensions and CM-type

The above results lead to the

**Definition 6.** A rational number  $\Delta$  is a new dimension in rank r if it is an allowed Coulomb dimension in rank r but not in rank  $(r-1)$ . The set of new dimensions in rank-r is

$$
\Xi(r) = \begin{cases} \left\{ \frac{n}{m} : \phi(n) = 2r, \text{ gcd}(n, m) = 1, 1 \le m < n \right\} & \text{for } r \ge 2\\ \left\{ 1, 6/5, 4/3, 3/2, 2, 3, 4, 6 \right\} & \text{for } r = 1 \end{cases}
$$
(2.49)

see [\[22\]](#page-150-0) for the cardinality  $|\Xi(r)|$  and other interesting properties of the set  $\Xi(r)$ .

We shall see in §. [2.12](#page-43-0) that an axis  $H_i$  with  $\Delta_i$  a new dimension in rank r is automatically regular. In facts, we have more:

<span id="page-23-2"></span> $24$  The role of this ambiguity was overlooked in [\[22\]](#page-150-0).

**Fact 4.** If the axis  $H_i$  has a new dimension  $\Delta_i$ , the fiber  $\mathscr{X}_u$  at the generic point  $u \in H_i$ has complex multiplication by the CM field  $\mathbb{Q}(e^{2\pi i/\Delta_i})$  of degree 2r and CM-type<sup>[25](#page-24-2)</sup> given by  $eq.(2.46).$  $eq.(2.46).$  $eq.(2.46).$ 

It is well-known that the number of isomorphism classes of polarized Abelian varieties of given CM-type is equal to the class number h of the corresponding number field  $\mathbb{Q}(e^{2\pi i/\Delta_i})$ . In particular for ranks  $r \leq 9$  all new dimensions correspond to fields with  $h = 1$  and the r-tuple of dimensions  $\{\Delta_1,\ldots,\Delta_r\}$  uniquely determines the fiber  $\mathscr{X}_u$  along a new dimension axis.

#### <span id="page-24-0"></span>2.5.3 Sample results

We list a few simple consequences of the above **Facts** for rank-2 that we shall use later in the examples. The geometries are assumed to be smooth. Here we use that an axis with a new dimension is automatically regular (see §. [2.12\)](#page-43-0).

- (1) A rank-2 special geometry having one of its dimensions in the list  $\{12, 12/5, 12/7, 12/11\}$ is automatically isotrivial;
- (2) if both axes are regular of new dimension then the dimensions are  $\{8, 12\}$  or  $\{\frac{8}{7}\}$  $\frac{8}{7}, \frac{10}{7}$  $\frac{10}{7}$ . A geometry with dimensions  $\{8/7, 10/7\}$  cannot be isotrivial;
- (3) the rank-2 geometries where both axes are regular must have dimensions  $\{5/4, 3/2\}$ ,  $\{8/7, 10/7\}$ ,  $\{8, 12\}$  or  $\{5/3, 4\}$ . The first two correspond to the Argyres-Douglas theories of types  $A_5$  and  $A_4$ , respectively, and the third one is the  $G_8$  isotrivial geometry constructed in [\[25\]](#page-150-7);
- (4) a rank-2 geometry with a new dimension of the type  $5/k$  or  $10/l$  is the Jacobian fibration of a fibration in generically smooth genus-2 curves.

*Proof.* (1) the dimension formulae yields pairs of dimensions with  $\varkappa \notin \{1,2\}$ . (2)(3) if both axes are regular we can compute  $\{\Delta_1, \Delta_2\}$  in two ways; consistency of the two computations yield the list. Second statement in (2): the fibers over the two axes have multiplication by  $\mathbb{Q}(e^{2\pi i/8})$  and respectively  $\mathbb{Q}(e^{2\pi i/10})$ , so are not isomorphic. (4) the fiber over the regular axis with the new dimension has muliplication by  $\mathbb{Q}(e^{2\pi i/5})$  hence is simple and not the product of two elliptic curves (cf. Corollary 13.3.6. of [\[77\]](#page-153-1)).  $\Box$ 

## <span id="page-24-1"></span>2.6 SW differential on enhanced axes

Suppose that the *i*-th axis  $H_i$  is regular. The R-symmetry which is unbroken along  $H_i$  is generated by  $\xi_i \equiv \exp(2\pi i \mathcal{E}/\Delta_i)$ .  $\xi_i$  acts by an automorphism of the generic (smooth) fiber  $\mathscr{X}_u$  with analytic representation

$$
\sigma(\xi_i) = \text{diag}\left\{\exp[2\pi i(\Delta_1 - 1)/\Delta_i], \exp[2\pi i(\Delta_2 - 1)/\Delta_i], \dots, \exp[2\pi i(\Delta_r - 1)/\Delta_i]\right\}
$$
 (2.50)

<span id="page-24-2"></span> $25$  For the notion of CM-type see e.g. chapter 13 of [\[77\]](#page-153-1).

and rational representation  $\chi(\xi_i) \in Sp(2r, \mathbb{Z})$  acting on  $H_1(\mathscr{X}_u, \mathbb{Z})$ , and

$$
\chi(\xi_i) \simeq \sigma(\xi_i) \oplus \overline{\sigma(\xi_i)}
$$
\n(2.51)

as complex representations. The polarization gives an isomorphism

$$
\text{Lie}(\mathcal{X}_u) \simeq H^{0,1}(\mathcal{X}_u) \simeq \overline{H^0(\mathcal{X}_u, \Omega^1_{\mathcal{X}_u})}
$$
(2.52)

Let  $\lambda \equiv \iota_{\mathcal{E}} \Omega$  be the SW differential and  $\lambda_u \equiv \lambda |_{\mathcal{X}_u}$  be its restriction to the fiber. We have

$$
\xi_i^* \lambda_u = e^{2\pi i/\Delta_i} \lambda_u. \tag{2.53}
$$

Let  $\{e_a\}$  be a symplectic basis of  $H_1(\mathscr{X}_u, \mathbb{Z})$ . The SW periods transform as

$$
\langle e_a, \lambda_u \rangle = \xi_i \cdot \langle e_a, \lambda_u \rangle = \langle \chi(\xi_i)_{ab} \, e_b \cdot \alpha, \xi_i^* \lambda_u \rangle = e^{2\pi i/\Delta_i} \, \chi(\xi_i)_{ab} \, \langle e_b, \lambda_u \rangle \tag{2.54}
$$

so, setting

$$
\Pi(u)_a \equiv \langle e_a, \lambda_u \rangle = \begin{pmatrix} a_i^D \\ a^j \end{pmatrix} \tag{2.55}
$$

$$
\chi(\xi_i)_{ab} \Pi(u)_b = e^{-2\pi i/\Delta_i} \Pi(u)_a \tag{2.56}
$$

i.e. the SW periods along  $H_i$  are eigenvectors of the rational representation of  $\xi_i$  of eigenvalue  $\exp(-2\pi i/\Delta_i)$  (up to overall scale the period is independent of  $u \in H_i$ , since the axis minus the origin is a  $\mathbb{C}^{\times}$ -orbit).

## <span id="page-25-0"></span>2.7  $\varkappa = 1$  vs.  $\varkappa = 2$  special geometries

The characteristic dimension  $\varkappa$  is a basic invariant of a  $\mathbb{C}^{\times}$ -isoinvariant special geometry. When  $\varkappa$  is not 1 or 2 the geometry is isotrivial, all smooth fibers are the product of r copies of  $E_{e^{2\pi i/\varkappa}}$ , and the possible geometries are essentially known globally.<sup>[26](#page-25-1)</sup> We remain with two cases:  $\varkappa = 1$  and  $\varkappa = 2$ . What is the difference between these two (potentially) non-isotrivial situations?

The automorphism  $\xi \equiv \exp(2\pi i \mathcal{E}/\lambda)$  which fixes point-wise the Coulomb branch  $\mathscr C$  acts on the SW differential  $\lambda \equiv \iota_{\mathcal{E}} \Omega$  as

$$
\xi^* \lambda = \begin{cases} \lambda & \varkappa = 1 \\ -\lambda & \varkappa = 2 \end{cases}
$$
 (2.57)

and in facts

$$
\xi^*|_{H^1(\mathcal{X}_u,\mathbb{C})} = (-1)^{\varkappa+1}.\tag{2.58}
$$

<span id="page-25-1"></span><sup>&</sup>lt;sup>26</sup> We stress that a geometry may be isotrivial even when  $\varkappa \in \{1,2\}.$ 

This means that when  $\varkappa = 2$  the SW periods  $(a_i^D, a^j)^t$  are well-defined only up to overall sign. In Lagrangian SCFTs with  $\varkappa = 2$  the sign flip is part of the gauge group (it is the longest word in the gauge Weyl group), so the two signs are obviously physically equivalent and indistinguishable. This property extends to special geometries with  $\varkappa = 2$  which do not describe Lagrangian SCFTs.

We can state the difference between  $\varkappa = 1$  and 2 in another (related) way. When  $\varkappa = 1$ there is one element h of the Coulomb branch function field<sup>[27](#page-26-2)</sup>  $\mathbb{C}(\mathscr{C})$  of dimension 1, while there is no such object when  $\varkappa = 2$ . When  $\varkappa = 2$  we have instead an element  $h^2 \in \mathbb{C}(\mathscr{C})$  of dimension 2.

This difference between the two situations will have fundamental consequences for the global aspects of a non-isotrivial special geometry.

## <span id="page-26-0"></span>2.8 Discriminant complement and monodromy representation

As before  $\mathcal{D} \subset \mathscr{C}$  is the discriminant i.e. the divisor of points with singular fibers; we write  $\sum_i \mathcal{D}_i$  for its decomposition into irreducible components.  $\mathcal{D}_i = (p_i)$  for irreducible quasi-homogeneous polynomials  $p_i$  of degree  $g_i$  and index  $\nu_i$ . We write  $\mathscr{C} \equiv \mathscr{C} \setminus \mathcal{D}$  for the complement of the discriminant, i.e. the open domain of "good" points with smooth Abelian fiber. We have a holomorphic family of polarized Abelian varieties

$$
\mathring{\pi} : \mathring{\mathscr{X}} \equiv \mathscr{X}|_{\mathring{\mathscr{C}}} \to \mathring{\mathscr{C}},\tag{2.59}
$$

parametrized by the complement  $\hat{\mathscr{C}}$ . We stress that  $\hat{\mathscr{C}}$  is quasi-projective over  $\mathbb{C}$ . Then we may apply the standard methods and results of the theory of Variations of Hodge Structure (VHS) over a quasi-projective base [\[34](#page-150-4)[–42\]](#page-151-0).

All fibers of  $\pi$  are diffeomorphic. This entails that over  $\mathscr{C}$  we have a flat Gauss-Manin local system with fiber  $H_1(\mathscr{X}_u, \mathbb{Z})$  [\[35,](#page-150-8) [40\]](#page-151-10). Going along a non-trivial closed curve  $\ell \subset \mathscr{C}$ the fiber of the local system gets back to itself up to a rotation by an element  $\gamma(\ell)$  of its polarized automorphism group: for principal polarizations this is the Siegel modular group  $Sp(2r, \mathbb{Z})$  (see [\[78\]](#page-153-2) for the general case). This yields a group homomorphism

$$
\varrho \colon \pi_1(\mathring{\mathscr{C}}) \to Sp(2r, \mathbb{Z}) \tag{2.60}
$$

called the monodromy representation [\[34](#page-150-4)[–42,](#page-151-0) [79\]](#page-153-3), a crucial invariant of the special geometry which controls many physical properties of the associated SCFT. The image  $\Gamma \subset Sp(2r, \mathbb{Z})$ of  $\rho$  is the monodromy group.

#### <span id="page-26-1"></span>2.8.1 Topology of the complement

A basic ingredient of special geometry is then the topology of the complement of the discriminant  $\mathscr{C}$ , in particular its fundamental group  $\pi_1(\mathscr{C})$ . While the Abelianization of  $\pi_1(\mathscr{C})$ ,

<span id="page-26-2"></span><sup>&</sup>lt;sup>27</sup>  $\mathbb{C}(\mathscr{C})$  is the quotient field of the integral domain  $\mathscr{R}$ .

i.e.  $H_1(\hat{\mathscr{C}}, \mathbb{Z})$ , is a pretty simple group [\[80\]](#page-153-4)

$$
H_1(\hat{\mathscr{C}}, \mathbb{Z}) \simeq \mathbb{Z}^k
$$
, where k is the number of components of  $\mathcal{D}$ ,  $(2.61)$ 

the group  $\pi_1(\hat{\mathscr{C}})$  is typically rather complicated. Roughly speaking, its derived group

$$
D\pi_1(\mathscr{C}) \stackrel{\text{def}}{=} \ker\left(\pi_1(\mathscr{C}) \to H_1(\mathscr{C}, \mathbb{Z})\right) \tag{2.62}
$$

is a measure of the interactions in the corresponding SCFT. A simple example will illustrate the issue.

**Example 2.2.** Consider the special geometry of 4d  $\mathcal{N} = 4$  SYM with  $G = SU(3)$  (seen as a special instance of  $\mathcal{N} = 2$  theory). The Coulomb dimensions  $\{\Delta_1, \Delta_2\}$  are the degrees of Weyl $(SU(3))$ ,  $\{2,3\}$ , while physically we expect the monodromy group to be the  $SU(3)$ Weyl group  $\Gamma = \text{Weyl}(SU(3)) \simeq \mathfrak{S}_3$ . Let us see how this comes about topologically. The discriminant has a unique irreducible component corresponding to the locus in  $\mathscr C$  where a  $SU(2)$  gauge symmetry is restored and the corresponding non-Abelian d.o.f. get light. Thus  $D = (u_1^3 - u_2^2)$ . One has [\[81\]](#page-153-5)

$$
\pi_1(\hat{\mathscr{C}}) \equiv \pi_1(\mathbb{C}^2 \setminus \{u_1^3 - u_2^2 = 0\}) = \langle \alpha, \beta : \alpha^3 = \beta^2 \rangle \tag{2.63}
$$

which is the Artin braid group  $\mathcal{B}_3$  of type  $A_2$ . Since the geometry is isotrivial with dimensions  $\{2,3\}$ , the monodromy representation  $\varrho$  factors through the finite quotient Weyl( $SU(3)$ )  $\simeq$  $\mathcal{B}_3/\mathcal{P}_3$  producing the physically expected results. ( $\mathcal{P}_3$  is the pure braid group in 3 strands). A similar story holds for  $\mathcal{N} = 4$  with any gauge group.

We list a few useful facts about the topology of hypersurface complements [\[80\]](#page-153-4). When the divisor D is normal crossing with k irreducible components the  $\pi_1$  of the complement is an Abelian group, generated by the  $k$  lassos circling the irreducible components. By the Zariski conjecture (proved in [\[82\]](#page-153-6)) the same holds for all nodal planar curves. In the general case we have an explicit description of  $\pi_1(\check{\mathscr{C}})$  given by the Zariski-van Kampen theorem [\[80,](#page-153-4) [83,](#page-153-7) [84\]](#page-153-8) that we shall review in §. [9.2.1.](#page-127-0)

#### <span id="page-27-0"></span>2.8.2 Subtleties when  $x = 2$

We assume our special geometry not to be isotrivial. At a generic point  $* \in \mathscr{C}$  the Rsymmetry is totally broken for  $\varkappa = 1$  while we have a residual unbroken  $\mathbb{Z}_2$  symmetry when  $\varkappa = 2$ . We define the SW periods  $\Pi \equiv (a_i^D, a^j)^t$  in a small neighborhood of \*, and analytically continue them along a non-trivial loop  $\ell \subset \mathscr{C}$  based at  $*$ . One would expect that the analytically continued periods Π′ satisfy

$$
\Pi' = \varrho(\ell) \,\Pi. \tag{2.64}
$$

However when  $\varkappa = 2$  the periods  $\Pi$  are well defined only up to sign (cf. §[.2.7\)](#page-25-0) and only the quotient representation

$$
\check{\varrho} \colon \pi_1(\mathring{\mathscr{C}}) \to PSp(2r, \mathbb{Z}) \equiv Sp(2r, \mathbb{Z})/\{\pm 1\}
$$
\n(2.65)

has a well-defined physical meaning.

#### <span id="page-28-0"></span>2.8.3 Algebraic invariants of Γ: the Hodge ring

We have a vector bundle  $\mathscr V$  over the complement  $\mathscr{\dot{C}}$  with fiber

$$
H_1(\mathscr{X}_u, \mathbb{C}) \equiv H_1(\mathscr{X}_u, \mathbb{Z}) \otimes_{\mathbb{Z}} \mathbb{C}
$$
\n(2.66)

equipped with the flat Gauss-Manin connection. We fix a reference point  $u<sub>star</sub>$  and write V for the fiber  $\mathcal{V}_{u_{\star}}$ . V carries the representation  $\varrho$  of  $\Gamma$  (defined over  $\mathbb{Q}$ ) while<sup>[28](#page-28-1)</sup>  $V \simeq V^{\vee}$ . This induces representations (over  $\mathbb{Q}$ ) of  $\Gamma$  on all tensor spaces  $V^{\otimes k}$ . Since  $\overset{\circ}{\mathscr{C}}$  is quasi-projective, one has:

**Fact 5** (See [\[79\]](#page-153-3)). The monodromy group  $\Gamma$  is finitely generated and semisimple, i.e. all its finite-dimensional representations split in direct sums of irreducible ones.

The subtle point about the previous statement is that we must work with representations defined over  $\mathbb Q$  instead of the algebraically closed field  $\mathbb C$ . However there are some simple situations.

**Fact 6.** Let  $V = V_0 \oplus V_1$  where  $V_0$  is the largest C-subspace where  $\Gamma$  acts trivially. Then:

- (1) The chiral ring contains precisely dim  $V_0/2$  elements with  $\Delta = 1$ .
- (2) The special geometry is a product with one factor a free geometry of rank dim  $V_0/2$ .

Argument. Let  $\{\gamma_I\} \equiv \{B^i, A_j\}$   $(I = 1, \ldots, 2r, i, j = 1, \ldots, r)$  be multi-valued sections of the local system with section  $H_1(\mathscr{X}_u, \mathbb{Z})$  which form a symplectic basis in each fiber. The SW periods

$$
\Pi_I(u) \stackrel{\text{def}}{=} \oint_{\gamma_I} \iota_{\mathcal{E}} \Omega \big|_{\mathcal{X}_u}, \qquad \Pi_I(u) \equiv \begin{pmatrix} a_i^D(u) \\ a^j(u) \end{pmatrix} \tag{2.67}
$$

are a set of 2r multi-valued holomorphic sections of  $\mathcal{V}^{\vee} \simeq \mathcal{V} \to \mathcal{C}$  which span the fibers and satisfy

$$
\mathcal{L}_{\mathcal{E}} \Pi_I = \Pi_I. \tag{2.68}
$$

Only half the periods are functionally independent: indeed  $da_i^D(u) \wedge da_i(u) = 0$  (cf. eq.[\(2.24\)](#page-18-2)) and  $a_i^D(u) = \tau_{ij}(u) a^j(u)$  where  $\tau_{ij}(u)$  is the (multivalued) period matrix of  $\mathscr{X}_u$  in the given basis  $\{B^j, A_i\}$ , Let  $P_I^J$  be the projector  $V \to V_0$  written in the basis  $\{\gamma_I\}$ . By construction

<span id="page-28-1"></span> $\frac{28}{28}$  The isomorphism is induced by the polarization.

 $P_I^J a_J$  are now global holomorphic functions on  $\mathscr{C}$ . It is easy to check that they are regular along the discriminant,<sup>[29](#page-29-1)</sup> and hence are defined and regular everywhere on  $\mathscr{C}$ . By definition, these holomorphic global functions belong to  $\mathscr R$  and have  $\Delta = 1$ . It is easy to see that exactly half of them are functionally independent.  $\Box$ 

We can generalize the story. Suppose that the symmetric k-th power  $V^{\odot k}$  contains a  $\Gamma$ -invariant tensor  $T^{I_1...I_k}$  (automatically defined over Q); the function

$$
T^{I_1\cdots I_k} \Pi_{I_1} \Pi_{I_2} \cdots \Pi_{I_k},\tag{2.69}
$$

(if not zero) is an element of R with  $\Delta = k$ . The invariants of  $\Gamma$  form a subalgebra of the chiral ring R called the Hodge ring  $\mathscr H$  [\[38,](#page-150-9) [39\]](#page-150-10), a fundamental invariant of special geometry. For instance, for a  $\mathcal{N} = 4$  special geometry  $\mathcal{H} = \mathcal{R}$ . More generally, for an isotrivial geometry  $\mathscr R$  is a finitely-generated  $\mathscr H$ -module. The ring  $\mathscr H$  is non-trivial for all quasi-isotrivial geometries as well as for the ones having a non-generic Mumford-Tate group.

More generally, if we have an invariant tensor  $T^{I_1...I_k;J_1...J_l} \in V^{\odot k} \otimes V^{\wedge l}$  we can form the global *l*-form of scaling dimension  $k + l$ 

$$
T^{I_1\ldots I_k;J_1\ldots J_l} \Pi_{I_1}\cdots \Pi_{I_k} d\Pi_{J_1} \wedge \cdots \wedge d\Pi_{J_l}
$$
\n
$$
(2.70)
$$

When  $T^{J_1J_2} \equiv \Omega^{J_1J_2}$  is the polarization, the corresponding invariant 2-form is trivial,  $\Omega^{IJ} d\Pi_I \wedge d\Pi_J = 0$ , in facts we have the stronger condition  $\Omega^{IJ} \Pi_I \wedge d\Pi_J = 0$ .

#### <span id="page-29-0"></span>2.8.4  $\textcolor{black}{\widehat{\mathfrak{D}}}$  The reducible case: subtleties

We consider now the case of a monodromy representation V which decomposes over  $\mathbb Q$  as  $V_1 \oplus V_2$  where the summands are non-trivial and non-isomorphic.

We consider two different scenarios. In the fist one the image of  $\Gamma$  in  $GL(V<sub>s</sub>, \mathbb{Q})$  is infinite for  $s = 1, 2$ . Working in a sufficiently small domain  $U \subset \mathscr{C}$ , we define the SW periods and use half of them as special coordinates in  $U$ . Taking  $\mathbb Q$ -linear combinations, we may split the periods in two sets,  $\Pi_I^{(1)}$  $\mathop{\mathrm{I}}\nolimits^{(1)}_I$  and  $\mathop{\mathrm{II}}\nolimits^{(2)}_J$  $J_J^{(2)}$  taking value respectively in  $V_1$  and  $V_2$ . When we go around a non-trivial path in  $\mathscr{C}$  and return back to U, each Q-period gets transformed into linear combinations of Q-periods from the same set. The two sets never mix, and the splitting of the local special coordinates in two sets  $a_{(1)}^i$  and  $a_{(2)}^j$  is globally defined. The cotangent bundle of  $\mathscr C$  splits into the subbundle spanned by the differentials  $da^i_{(1)}$  and the one spanned by the  $da_{(2)}^j$ 's. Indeed, up to isogeny, the fibers  $\mathscr{X}_u$  split into a product  $A_u^{(1)} \times A_u^{(2)}$  of Abelian varieties with  $H_1(A_u^{(s)}, \mathbb{Q}) \simeq V_s$ . Going to a finite cover we may assume the fiber is a product. We have

<span id="page-29-2"></span>
$$
T_0 A_u^{(1)} \oplus T_0 A_u^{(2)} \simeq T_u^* \mathscr{C}.
$$
\n
$$
(2.71)
$$

<span id="page-29-1"></span> $29$  By Hartogs theorem it suffice to check regularity in codimension 1.

The decomposition [\(2.71\)](#page-29-2) defines two complementary integrable distributions whose leaves have local equations in  $U$  of the form

$$
a_{(s)}^i = \text{const.} \quad s = 1, 2,
$$
\n(2.72)

so that (restricting U if necessary) locally the Coulomb branch can be written as  $U =$  $U_{(1)} \times U_{(2)}$  with  $a^i_{(s)}$  local coordinates on the factor  $U_{(s)}$   $(s = 1, 2)$ . The local pre-potential is a sum

$$
\mathcal{F}(a_{(1)}^i, a_{(2)}^j) = \mathcal{F}_{(1)}(a_{(1)}^i) + \mathcal{F}_{(2)}(a_{(2)}^j)
$$
\n(2.73)

where the period matrix of  $A_u^{(s)}$  is given by  $\partial^2 \mathcal{F}_{(s)}(a_{(s)}) \partial a_{(s)}^i \partial a_{(s)}^j$ , and the analytic continuation of  $\mathcal{F}(a_{(1)}^i, a_{(2)}^j)$  (as a *multivalued* holomorphic function in  $\hat{\mathscr{C}}$ ) is the sum of the analytic continuations of the two multivalued functions in the RHS.

Thus *locally* in  $U$  we have a product of two special geometries with Coulomb branches  $\mathscr{C}^{(s)}$  and fibers  $A^{(s)}$   $(s=1,2)$ . This local product structure is preserved by the monodromy so it is a kind of "global" feature.

At first sight it may seem that we have got a decomposable geometry. Is this really so? Not quite. A decomposable geometry has some peculiar properties which are not implied by the above discussion of reducible monodromy.

The  $r = 2$  situation. For definiteness we discuss the case  $r = 2$  which already presents all essential phenomena. All decomposable  $r = 2$  geometries are isotrivial and their discriminant is contained in the union of the two axes  $H_1 \cup H_2$  (in particular, it is normal crossing). Neither properties follow from reducibility of the monodromy representation. Indeed the geometry cannot be isotrivial and the discriminant cannot be normal crossing when  $\Gamma$  is infinite (as we are assuming). The point is that while the generic fiber is indeed the product of two elliptic curves  $E_{\tau_{(1)}} \times E_{\tau_{(2)}}$ , the periods of these curves need not to be constant: we have just  $\tau_{(s)} = \partial^2 \mathcal{F}_{(s)}(a_{(s)})/\partial^2 a_{(s)}$  for  $s = 1, 2$ . The only conclusion that we may draw is that the monodromy representation factors through

$$
SL(2,\mathbb{Q}) \times SL(2,\mathbb{Q}) \hookrightarrow Sp(4,\mathbb{Q})
$$
\n
$$
(2.74)
$$

while (by assumption) its image is infinite in both factors, in fact the Q-Zariski closure of the image in each factor is the full  $SL(2, \mathbb{Q})$  modulo finite groups. In particular there are no invariant symmetric elements in the tensor algebra  $T^{\bullet,\bullet}V_s$  of each summand  $V_s$ , so while the splitting of the periods in the two sets  $\Pi^{(1)}$  and  $\Pi^{(2)}$  is globally defined, we cannot use each set of special coordinates *separately* to define global functions on  $\mathscr C$  i.e. elements in  $\mathscr R$ which may play the role of coordinates  $u_i$  in the factor Coulomb branches of a decomposable geometry. Therefore we expect that the Coulomb coordinates  $u_1, u_2$  are non-trivial functions of both  $a_{(1)}$  and  $a_{(2)}$ . However, in order for the functions  $u_i = u(a_{(1)}, a_{(2)})$ , initially defined locally in U, to have a *univalued* analytic continuation everywhere in  $\mathscr{C}$ , the monodromy

<span id="page-31-0"></span>

Kodaira fiber	$\mathbf{I}_n$	$\mathbf{I}_n^*$
monodromy	$\left[\begin{matrix}1 & n\\0 & 1\end{matrix}\right]$	$\begin{bmatrix} -1 & -n \\ 0 & -1 \end{bmatrix}$
Euler number	$n_{\cdot}$	$n+6$
Lie algebra g	$\mathfrak{su}(n)$	$\mathfrak{so}(2n+8)$

Table 2: Non-semisimple Kodaira fibers. Here  $n \geq 1$ 

transformations of the two summands should be correlated, and given the Zariski density of Γ the only possibility which comes to mind is a further factorization

$$
\pi_1(\mathscr{C}) \to SL(2,\mathbb{Q}) \xrightarrow{\text{diag}} SL(2,\mathbb{Q}) \times SL(2,\mathbb{Q}) \hookrightarrow Sp(4,\mathbb{Q}).
$$
\n(2.75)

If this was the case, we would have an element of  $\mathscr R$  with  $\Delta = 2$  given by  $\epsilon^{ij} \Pi_i^{(1)} \Pi_j^{(2)}$ j .

In the second scenario (for  $r = 2$ ) the image of Γ in the first  $SL(2, \mathbb{Q})$  factor is infinite but is finite in the second one. (If both images are finite the geometry is isotrivial, see §. [2.10.4\)](#page-41-0). When pulled back to a finite cover  $\mathscr{C}^\sharp$  of  $\mathscr{C}$  the family takes the form

$$
E \times \mathcal{Y} \to \mathscr{C}^\sharp \tag{2.76}
$$

with E a constant elliptic curve and  $\mathscr{Y} \to \mathscr{C}^{\sharp}$  a non-isotrivial family of elliptic curves. This is what we call a quasi-isotrivial situation. If such a special geometry exists, it cannot be decomposable. However it looks pretty hard to cook up such a quasi-isotrivial family in a way which is consistent with all the other required structures to produce a regular C <sup>×</sup>-isoinvariant special geometry. A quasi-isotrivial geometry is a highly over-constrained problem which may have a solution only under very specific conditions: peculiar Coulomb dimensions, specific form of the discriminant, etc. A priori no one will bet a cent on their existence. However, wait for §§. [8.1,](#page-81-0) [8.9.1,](#page-108-1) [8.13](#page-119-0) and see.

This discussion motivates the following

**Definition 7.** An indecomposable geometry is *reducible* iff it is not isotrivial and its underlying monodromy representation is reducible.

Later we shall show that

<span id="page-31-1"></span>**Fact 7.** An indecomposable but reducible geometry is a very rare, bizzarre and ugly animal. They exist only for some special "troublesome" dimension r-tuples  $\{\Delta_1, \ldots, \Delta_r\}$ . In rank 2 they are either quasi-isotrivial or have a dimension 2 Coulomb coordinate.

## <span id="page-32-0"></span>2.9 Singular fibers along the discriminant

In this technical subsection we quickly sketch the structure of singular fibers as described in the math literature [\[45–](#page-151-3)[50\]](#page-151-4) and reviewed in [\[26\]](#page-150-1). We stress that our assumption that there is a section  $s: \mathscr{C} \to \mathscr{X}$  implies that only simple singular fibers may be present, so the phenomena associated with more general kinds of fibers are not present in this restricted setting (for "special geometries" with multiple fibers see [\[26\]](#page-150-1)).

In rank 1 the classification of the singular elliptic fibers is given by Kodaira [\[51–](#page-151-5)[54\]](#page-151-6); the local monodromies of simple exceptional elliptic fibers with semisimple monodromy are listed in table  $1^{30}$  $1^{30}$  $1^{30}$  $1^{30}$  and the ones with non-semisimple monodromy in table [2.](#page-31-0) In rank-r the possible conjugacy classes of the local monodromy around a discriminant component  $\mathcal{D}_a$  are again classified in Kodaira types but new phenomena appear. The local monodromy is not sufficient to determine the singular fiber in general [\[45](#page-151-3)[–50\]](#page-151-4).

#### <span id="page-32-1"></span>2.9.1 Generalities

We stress that the following analysis applies under *genericity* assumptions. For the highly non-generic situations in Fact [7](#page-31-1) one needs to introduce the appropriate modifications.

We consider a generic point u in the *i*-th discriminant component. The fiber  $\mathscr{X}_u$  is a union of components  $F_{u,a}$ ; the normalization of each component  $F_{u,a}^{\text{nor}}$  is a fibration over an Abelian variety  $A_u$  of dimension  $(r-1)$  (the Albanese variety of the fiber) which is the same for all components. For *simple* fibers (i.e. fibers of multiplicity 1) the fiber of  $F_{u,a}^{\text{nor}} \to A_u$ is a copy of  $\mathbb{P}^1$ . However it is not always true that  $\mathscr{X}_u \to A_u$  is globally a fibration: two components  $F_{u,a}, F_{u,b}$  are glued together by identifying a section  $\simeq A_u$  of the first component with a section  $\simeq A_u$  of the second one, but the identification does not necessarily carry the zero to the zero. When the fiber is not simply-connected this may result in an obstruction to the fiber being a global fibration over  $A_u$ . One may classify the fibers type in terms of the characteristic cycle: i.e. a maximal connected sequence of fibers. The possible (simple) characteristic cycles are classified by the Kodaira types plus an extra type  $I_{\infty}$  [\[45](#page-151-3)[–47\]](#page-151-12). If the type is semi-simple  $(I_0^*, II, II^*, III, III^*, IV,$  and  $IV^*$ ) we have a fibration over  $A_u$  with fiber the Kodaira singularity. In these cases by some blow-up/down and a finite cover we can get a fibration with no singular fiber (as we see from the period map) [\[46,](#page-151-13) [47\]](#page-151-12). These same operations allow to reduce the case  $I_b^*$  to the case  $I_{2b}$ . The case  $I_b$  (including  $b = \infty$ ) is the most subtle one, but (at least when the polarization is principal) it is described in great detail in [\[47\]](#page-151-12). The rough picture, is as follows: each irreducible component  $F_{u,a}$  is a fibration over  $A_u$  with fiber a  $\mathbb{P}^1$ . Two components  $F_{u,a}$ ,  $F_{u,b}$  cross along finitely many sections. In facts their intersection matrix  $F_{u,a} \# F_{u,b}$  (suitably defined) is equal to the intersection matrix  $\Theta_a \cdot \Theta_b$  of the components a Kodaira fiber K. If K has type  $I_b$ , the type of the characteristic cycle can be  $I_{mb}$  where m is a positive integer or  $\infty$  (in this case the characteristic cycle

<span id="page-32-2"></span> $30$  In the following we omit type  $I_0$  which is non-singular.

warps m times around the fiber). When the fiber is simply-connected,  $K$  coincides with the type of the characteristic cycle.

We stress that, while the type K is locally constant as we move the general point  $u$  in the discriminant locus, the type of the characteristic cycle may change abruptly: as shown in the examples of [\[47\]](#page-151-12), points whose fibers are of type  $I_{\infty}$  or type  $I_k$  (k finite) may be both dense in the discriminant.<sup>[31](#page-33-1)</sup> The physical meaning of the characteristic cycle was discussed in [\[26\]](#page-150-1); we refer the interested reader to that paper.

The type K controls the local monodromy  $\varrho_a$  around the a-th divisor component. It is clear from the above discussion that  $\varrho_a$  is conjugate in  $Sp(2r, \mathbb{Q})$  to the direct sum of the Kodaira monodromy of type K,  $\varrho_K \in SL(2,\mathbb{Z})$ , and the  $(2r-2) \times (2r-2)$  identity matrix. In particular, its characteristic polynomial has the form

$$
\det[z - \varrho_a] = (z - 1)^{2(r-1)} P_K(z) \tag{2.77}
$$

where  $P_K(z)$  is the characteristic polynomial of  $\varrho_K$ . Notice that the conjugacy class in  $Sp(2r, \mathbb{Q})$  distinguishes types II, III, IV from the ones with the same characteristic polynomial  $II^*, III^*, IV^*$  (respectively). A priori we would have an ambiguity for types  $I_b^*$  and  $I<sub>b</sub>$ . However this is precisely the case where the detailed analysis of [\[46,](#page-151-13)[47\]](#page-151-12) applies, and the relation of the monodromy class to the fiber geometry is established:

**Proposition 1** (See [\[47\]](#page-151-12)). Assume principal polarization. The monodromy  $\varrho_a$  around a semi-stable divisor  $\mathcal{D}_a$  (characteristic cycle of type  $I_b$ ) is conjugate over in  $Sp(2r, \mathbb{Z})$  to

<span id="page-33-2"></span>
$$
\begin{pmatrix} 1 & \ell \\ 0 & 1 \end{pmatrix} \oplus \mathbf{1}_{r-1} \tag{2.78}
$$

where  $\ell$  is the number of components of the fiber. As a consequence if  $\rho_a$  is a local monodromy around a divisor of type  $I_b^*$ ,  $\varrho_a^2$  is conjugate over  $Sp(2r, \mathbb{Z})$  to the matrix  $(2.78)$  with  $\ell = 2b$ .

The geometry also depends on the characteristic cycle, see [\[45–](#page-151-3)[47\]](#page-151-12) or [\[26\]](#page-150-1) for details.

#### <span id="page-33-0"></span>2.9.2 More details on the stable reduction

This subsection may be skipped in a first reading. Having constructed a putative special geometry, we have to check that it is a genuine one, that is, analytically regular. One important step is to check its behavior along the discriminant which is the locus where things may go wrong. This is done by comparing the near discriminant geometry with the model local geometry which describes the singular fiber type at the generic point in the discriminant. For the present purposes the important invariant of the singular fiber is its local monodromy class. As always we assume principal polarization. The following local models are borrowed from refs. [\[45–](#page-151-3)[47\]](#page-151-12).

<span id="page-33-1"></span> $31$  See Proposition 5.3 of [\[47\]](#page-151-12).

A. Transformation of the fiber type. Just as in the Kodaira situation (rank-1), locally in a neighborhood of  $y \in \mathcal{D}_i$  we can always reduce to the semi-stable case by a sequence of blow up/down of components of the fiber together with a (branched) cyclic base change, that is, by making  $\tilde{z} \to \tilde{z}^m = z$ , where z is the local coordinate in  $\mathscr{C}$  such that  $\mathcal{D}_i$ is given by  $z = 0$  and m is the order of the semi-simple part of the monodromy around  $\mathcal{D}_i$ :

model fiber	m	semi-simple reduction
$I_n^*, n \geq 1$		$I_{2n}$
$II, II^*$		$E(e^{2\pi i/3})$
$III, III^*$		E(i)
$IV, IV^*$	3	$E(e^{2\pi i/3})$

where  $E(\tau)$  stands for the elliptic curve of period  $\tau$ , and fiber type  $E(\tau)$  means that the fiber over the point  $\tilde{y}$  covering y has the form

$$
E(\tau) \times A \tag{2.80}
$$

for an Abelian variety A of dimension  $r - 1$ .

**B.** Transformation of the symplectic form.  $y$  is a generic point in a component  $\mathcal{D}_i$  of the discriminant, and  $U \ni y$  a sufficiently small neighborhood. Let  $\tilde{U} \to U$  be the branched cover of the (local) semi-stable reduction. We write  $V = \pi^{-1}(U)$  and  $\tilde{V} = \tilde{\pi}^{-1}(\tilde{U}),$ where tilded symbols refer to the semi-stable reduction. We can find local coordinates  $(x_i, y^i)$ in V (resp.  $(\tilde{x}_i, \tilde{y}^i)$  in  $\tilde{V}$ ) such that

- $x_i$ ,  $\tilde{x}_i$  are coordinates in the bases U and  $\tilde{U}$ ;
- the local equation of  $\mathcal{D}_i$  is  $x_r = \tilde{x}_r = 0$ .

There are three cases [\[46,](#page-151-13) [47\]](#page-151-12):

(I) Type  $I_k^*, k \geq 1$ . In this case

$$
x_r = \tilde{x}_r^2
$$
,  $y^r = \frac{\tilde{y}^r}{\tilde{x}_r}$ ,  $x_i = \tilde{x}_i$   $y^i = \tilde{y}^i$  for  $i = 1, ..., r - 1$  (2.81)

$$
\sum_{i} dx_{i} \wedge dy^{i} = \sum_{k=1}^{r-1} d\tilde{x}_{i} \wedge d\tilde{y}^{i} + 2 d\tilde{x}_{r} \wedge d\tilde{y}^{r}
$$
\n(2.82)

(II) Types  $I_0^*$ ,  $II^*$ ,  $III^*$  and  $IV^*$ . One has

$$
x_r = \tilde{x}_r^m
$$
,  $y^r = \frac{\tilde{y}^r}{\tilde{x}_r^{m-1}}$ ,  $x_i = \tilde{x}_i$   $y^i = \tilde{y}^i$  for  $i = 1, ..., r - 1$  (2.83)

$$
\sum_{i} dx_{i} \wedge dy^{i} = \sum_{k=1}^{r-1} d\tilde{x}_{i} \wedge d\tilde{y}^{i} + m d\tilde{x}_{r} \wedge d\tilde{y}^{r}
$$
\n(2.84)

where  $m = 2, 6, 4, 3$ , respectively.

(III) Types  $II$ ,  $III$  and  $IV$ . One has

$$
x_r = \tilde{x}_r^m, \quad y^r = \frac{\tilde{y}^r}{\tilde{x}_r}, \qquad x_i = \tilde{x}_i \quad y^i = \tilde{y}^i \quad \text{for } i = 1, \dots, r - 1 \tag{2.85}
$$

$$
\sum_{i} dx_{i} \wedge dy^{i} = \sum_{k=1}^{r-1} d\tilde{x}_{i} \wedge d\tilde{y}^{i} + m \tilde{x}^{m-2} d\tilde{x}_{r} \wedge d\tilde{y}^{r}
$$
\n(2.86)

where  $m = 6, 4, 3$ , respectively.

The monodromy in the cover geometry as  $\tilde{x}_r \to e^{2\pi i} \tilde{x}_r$  is equal to the monodromy in the original space as  $x_r \to e^{2\pi i m} x_r$  i.e. to the m-th power of the original monodromy which is trivial for all semisimple types and unipotent of type  $I_{2n}$  in the  $I_n^*$  case.

C. Local pre-potential in the semi-stable case. We consider a (sufficiently small) neighborhood U of a generic point x on a semi-stable divisor  $\mathcal{D}_i$ . Locally we can always reduce to this case.

**Proposition 2** ([\[47\]](#page-151-12)). In U we can find special coordinates  $(a^1, \ldots, a^r)$  such that

- the local equation of  $\mathcal{D}_i$  is  $a^r = 0$ ;
- the prepotential has the form

<span id="page-35-1"></span>
$$
\mathcal{F}(a^1, \cdots, a^r) = \tilde{\mathcal{F}}(a^1, \cdots, a^r) + \frac{\ell}{4\pi i} (a^r)^2 \log a^r \qquad (2.87)
$$

where  $\tilde{\mathcal{F}}(a^1, \dots, a^r)$  is holomorphic in U and l is a positive integer;

• the period matrix is

$$
\boldsymbol{\tau}_{ij} = \frac{\partial^2 \mathcal{F}}{\partial a_i \partial a_j}.
$$
\n(2.88)

• the local monodromy around the semi-stable discriminant with pre-potential  $(2.87)$  is unipotent of type  $I_{\ell}$ .

Indeed under the transformation  $a_r \to e^{2\pi i} a_r$  the periods will transform as

$$
a^i \to a^i, \qquad a^D_j \equiv \frac{\partial \mathcal{F}}{\partial a^j} \to a^D_j \quad j \neq r, \qquad a^D_r \equiv \frac{\partial \mathcal{F}}{\partial a^r} \to a^D_r + \ell a^r. \tag{2.89}
$$

#### <span id="page-35-0"></span>2.10 Period map and structure theorem

We saw above that the restriction of the fibration  $\pi$  to the complement  $\mathscr{C}$  is in particular a family  $\mathring{\mathscr{X}}$  of polarized Abelian varieties parametrized by the quasi-projective space  $\mathring{\mathscr{C}}$ . The
moduli space parametrizing isomorphism classes of principally polarized Abelian varieties of dimension  $r$  is the *Siegel variety* 

$$
S_r \equiv Sp(2r, \mathbb{Z}) \backslash Sp(2r, \mathbb{R}) / U(r).
$$
\n(2.90)

We then have a *holomorphic period map*  $\check{p}$ 

$$
\check{p} \colon \mathring{\mathscr{C}} \to \mathsf{S}_r \tag{2.91}
$$

which sends a point  $u \in \mathscr{C}$  into the isoclass  $[\mathscr{X}_u] \in \mathsf{S}_r$  of its fiber. The geometry is isotrivial iff  $\check{p}$  is the constant map.

We stress that the family  $\mathcal{X}$  of Abelian varieties over  $\mathcal{C}$  is not fully determined by the period map.<sup>[32](#page-36-0)</sup> Since  $\{\pm 1\} \subset Sp(2r, \mathbb{Z})$  acts trivially on the symmetric space  $Sp(2r, \mathbb{R})/U(r)$ , the period map  $\check{p}$  only determines the quotient representation

$$
\check{\varrho} \colon \pi_1(\mathring{\mathscr{C}}) \to PSp(2r, \mathbb{Z}) \equiv Sp(2r, \mathbb{Z})/\{\pm 1\},\tag{2.92}
$$

while the family of Abelian varieties over  $\mathscr{C}$  depends on the actual monodromy i.e. on the specific lift  $\rho$  of  $\check{\rho}$ 

$$
\pi_1(\stackrel{\circ}{\mathscr{E}}) \xrightarrow{\qquad \varrho \qquad} \longrightarrow Sp(2r, \mathbb{Z})
$$
\n
$$
\downarrow^{\text{can}} \qquad (2.93)
$$
\n
$$
\stackrel{\circ}{\longrightarrow} \text{Sp}(2r, \mathbb{Z})/\{\pm 1\}
$$

which is a *finer* invariant of the family  $\mathcal{X}^{\dagger}$  than  $\check{\rho}$ . There may be at most finitely many special geometries with the same period map [\[87\]](#page-153-0). When  $\varkappa = 2$  only  $\check{\varrho}$  is a a priori well-defined (cf. §. [2.8.2\)](#page-27-0), and one has to be careful with for the proper lift  $\rho$  which yields the correct family of Abelian varieties arising from the particular special geometry.

# 2.10.1 The Abelian family over  $\mathscr{\dot{P}}$

The holomorphic map  $\exp(z \mathcal{E})$  acts by automorphisms, hence

$$
\mathscr{X}_{\exp(z\mathcal{E})u} \simeq \mathscr{X}_u \quad \text{for all } z \in \mathbb{C}/2\pi im\mathbb{Z},\tag{2.94}
$$

i.e. the fibers over all points in a  $\mathbb{C}^{\times}$ -orbit are isomorphic (as polarized Abelian varieties). Then the map  $\check{p}$  factors through a holomorphic period map

<span id="page-36-1"></span>
$$
p: \mathscr{E}/\mathbb{C}^{\times} \simeq \text{Proj} \mathscr{R} \setminus Y \to \mathsf{S}_{r}
$$
\n
$$
(2.95)
$$

<span id="page-36-0"></span> $32$  This is well-known in the case of families of elliptic curves over a curve [\[51–](#page-151-0)[53,](#page-151-1) [85\]](#page-153-1). There are several families of elliptic curves over a curve with the same period map: they are related by the so-called *quadratic* transform [\[85,](#page-153-1) [86\]](#page-153-2).

where  $Y \subset \text{Proj } \mathscr{R} \equiv \mathscr{P}$  is the hypersurface  $\mathcal{D}/\mathbb{C}^{\times}$ . When  $\mathscr{C}$  is smooth

$$
\text{Proj } \mathscr{R} = \mathbb{P}(d_1, \cdots, d_r) \simeq \mathbb{P}(q_1, \ldots, q_r) \quad \text{(well-formed)}, \tag{2.96}
$$

and Y is the weighted projective hypersurface with quasi-cone the reduced discriminant  $\mathcal{D}$ .

However there is a subtlety.  $\mathbb{P}(d_1, \dots, d_r)$  is not projectively<sup>[33](#page-37-0)</sup> smooth on points which corresponds to  $\mathbb{C}^{\times}$ -orbits with enhanced R-symmetry, since the quotient of  $U(1)_R$  which acts effectively on  $\mathscr C$  does not act freely there. We are mainly interested in singularities in codimension one: over an enhanced divisor of index  $\nu$  we have  $\mathbb{Z}_{\nu}$  quotient singularities.

It is convenient to replace the divisor Y by the special locus divisor  $\mathscr{S} \subset \mathscr{P}$  whose irreducible components  $S_i$  are either projective images  $\pi(\mathcal{D}_i)$  of the discriminant components or divisors of  $\mathscr P$  which are the projective images of R-enhanced irreducible divisors. Note that a special divisor  $S_i$  can belong to both the discriminant and the R-enhancement locus.

We consider the projective monodromy representation

$$
\mu: \pi_1(\mathscr{P} \setminus \mathscr{S}) \to \begin{cases} Sp(2r, \mathbb{Z}) & \mathscr{U} = 1 \\ PSp(2r, \mathbb{Z}) & \mathscr{U} = 2 \end{cases} \tag{2.97}
$$

which we call the  $\mu$ -monodromy to distinguish it from the  $\varrho$ -monodromy representation of  $\pi_1(\mathscr{C} \setminus \mathcal{D})$ . The two monodromies are related and determine each other.

**Fact 8.** For  $\varkappa = 1$  (resp.  $\varkappa = 2$ ) the group  $\mu(\pi_1(\mathscr{P}\setminus\mathscr{S}))$  is equal to the Coulomb branch monodromy group  $\Gamma$  (resp. the image  $\tilde{\Gamma}$  of  $\Gamma \subset Sp(2r,\mathbb{Z})$  in the quotient group  $Sp(2r,\mathbb{Z})/\{\pm 1\}$ ).

Argument. Let  $*\in \mathscr{C} \setminus \pi^{-1}(\mathscr{S})$  be a generic point in the Coulomb branch. A closed path  $\ell \in \mathscr{C} \subseteq \mathscr{C} \setminus \mathcal{D}$ , based at \*, may be continuously deformed to lay in  $\mathscr{C} \setminus \pi^{-1}(\mathscr{S})$ . Then  $\ell^{\sharp} \equiv \pi(\ell)$  is a closed path in  $\mathscr{P} \setminus \mathscr{S}$  based at  $\pi(*)$ . On the other hand if  $\gamma(t)$  is a closed path in  $\mathscr{P}\setminus\mathscr{S}$  based at  $\pi(*)$  we can lift it into a path  $\tilde{\gamma}(t)$  starting at \* and ending at some point  $\tilde{\gamma}(1)$  in the same  $\mathbb{C}^{\times}$ -orbit as  $*$ . Hence

<span id="page-37-1"></span>
$$
* = \exp(a\mathcal{E})\tilde{\gamma}(1), \qquad \text{for some } a \in \mathbb{C}/2\pi im\mathbb{Z}
$$
 (2.98)

and we can define a closed loop  $\gamma^{\flat}$  in  $\mathscr{C} \setminus \pi^{-1}(\mathscr{S})$  based at  $*$ 

$$
\gamma^{\flat}(t) = \begin{cases}\n\tilde{\gamma}(2t) & 0 \le t \le 1/2 \\
\exp[(2t-1)a \mathcal{E}]\tilde{\gamma}(1) & 1/2 \le t \le 1\n\end{cases}
$$
\n(2.99)

Note that the homotopy class of  $\gamma^{\flat}$  is independent of the lift. When  $\varkappa = 1$  the two maps

$$
\pi_1(\mathscr{C}\setminus\pi^{-1}(\mathscr{S}))\xleftarrow{\sharp} \pi_1(\mathscr{P}\setminus\mathscr{S})
$$
\n(2.100)

<span id="page-37-0"></span><sup>&</sup>lt;sup>33</sup> The underlying complex space may be nevertheless smooth (this happens when the associated wellformed weighted projective space is  $\mathbb{P}^{r-1} \equiv \mathbb{P}(1,\ldots,1)$  [\[71\]](#page-152-0)). See [\[44\]](#page-151-2) for a discussion.

are clearly inverse of each other and  $\mu(\ell) = \varrho(\ell^{\flat})$ . However when  $\varkappa = 2$  the base point  $∗$  is fixed by an automorphism of the form  $exp(2πim E/2)$  (*m* odd) and *a* in eq.[\(2.98\)](#page-37-1) is ambiguous by the shift  $a \mapsto a + i\pi m$ . Hence  $\mu(\ell) = \pm \varrho(\ell^{\flat})$ .  $\Box$ 

**Example 2.3.** We return to **Example [2.2](#page-27-1)**:  $\mathcal{N} = 4$  with  $G = SU(3)$ . As complex manifolds, one has  $\mathbb{P}(2,3) \simeq \mathbb{P}^1$  via the map  $(u_1, u_2) \mapsto (u_1^3 : u_2^2)$  [\[71\]](#page-152-0). The only component of the discriminant is mapped to the point  $(1 : 1) \in \mathbb{P}^1$ . The sphere less one point is simplyconnected, so all monodromy groups are trivial, while we expect it to be the non-Abelian (but solvable) group Weyl( $SU(3)$ ). The local monodromy  $\mu_1$  around the discriminant point  $1 \equiv (1:1)$  is physically expected to generate Weyl $(SU(2)) \simeq \mathbb{Z}_2$ , while the full monodromy group is Weyl $(SU(3))$  (cf. **Example [2.2](#page-27-1)**). Taking into account the monodromy around the two R-enhanced divisors we get the correct answer: the local  $\mu$ -monodromy around 0 is of order  $\nu_0 = 2$  and the local monodromy around  $\infty$  is of order  $\nu_\infty = 3$ ; hence the  $\mu$ -monodromy group has the presentation

$$
\Gamma = \{\mu_0, \mu_1, \mu_\infty \colon \mu_0^2 = \mu_1^2 = \mu_\infty^3 = \mu_1 \mu_2 \mu_3 = 1\}
$$
\n(2.101)

which is the standard Coxeter presentation of  $Weyl(SU(3))$ .

In conclusion: the family of polarized Abelian r-varieties  $\mathring{\mathscr{X}} \to \mathring{\mathscr{C}}$  defines a family of polarized Abelian r-varieties

$$
\mathscr{A} \to \mathring{\mathscr{P}} \equiv \mathscr{P} \setminus \mathscr{S} \tag{2.102}
$$

which is our main object of study in this paper.

A basic property of  $\Gamma$  is that it contains finite-index normal subgroups which are *neat* (so, in particular, torsionless) [\[79\]](#page-153-3). We write  $\Upsilon \triangleleft \Gamma$  for such a subgroup and  $\mathsf{G} = \Gamma/\Upsilon$  for the quotient *finite* group. By general theory we have a finite holomorphic cover  $f: \mathcal{Q} \to \mathcal{P}$ , branched over  $\mathscr{S}$ , with deck group G such that the pulled back family

$$
f^*\mathscr{A} \to \mathscr{Q} \equiv \mathscr{Q} \setminus f^{-1}(\mathscr{S}) \tag{2.103}
$$

has neat monodromy group  $\Upsilon$ . For many purposes it is technically more convenient to consider the G-twisted period map

$$
P: \mathcal{Q} \to \Upsilon \backslash Sp(2r, \mathbb{R})/U(r), \tag{2.104}
$$

which contains the same information but is simpler since we got rid of the finite groups.

### 2.10.2 Local monodromies at special divisors

The special divisor  $S_i$  may be of three kinds:

(1) projective discriminant non-enhanced: the local monodromies satisfy  $[\mu_i] = [\rho_i];$ 

(2) R-enhanced non-discriminant. In this case the period map  $p$  may be extended holomorphically on the divisor, as it is evident from eq.  $(2.95)$ . Let  $(u, z)$  be local coordinates along a regular enhanced divisor with local homogeneous equation  $z = 0$ . If its index is  $\nu_i$ , we have that the points  $p_k \equiv (u, e^{2\pi i k/\nu_i} \epsilon)$  are all in the same  $\mathbb{C}^\times$  orbit, indeed  $p_{k+1} = \exp[2\pi i \mathcal{E}/(\Delta(z)\nu_i)]p_k$ . A curve in  $\mathscr{E}$  from  $p_k$  to  $p_{k+1}$  will project to a closed loop  $\ell$  in  $\mathscr{P}$ . Going around  $\ell$  for  $\nu_i$  times we return to the original point  $p_0 \equiv p_{\nu_i}$  in  $\hat{\mathscr{C}}$ ; since the locus  $z = 0$  is not in the determinant, there is no  $\varrho$ -monodromy. Then the  $\mu$ -monodromy along the loop  $\ell \subset \mathring{\mathscr{P}}$  satisfies

$$
\mu_i^{\nu_i} = 1 \in \begin{cases} Sp(2r, \mathbb{Z}) & \mathcal{U} = 1 \\ PSp(2r, \mathbb{Z}) & \mathcal{U} = 2. \end{cases} \tag{2.105}
$$

One can try to be more precise. By continuity as  $\epsilon \to 0$ ,  $\mu_i$  is (up to conjugacy) the same as the rational representation of the automorphism  $\exp[2\pi i \mathcal{E}/(\Delta(z)\nu_i)]$  of the generic fiber over the R-enhanced discriminant. Hence  $\mu_i$  is a  $\nu_i$ -root of  $1 \in PSp(2r, \mathbb{Z})$  which lifts in  $Sp(2r, \mathbb{Z})$  to a matrix with eigenvalues

$$
\left\{\exp[\pm 2\pi i(1-\Delta_1)/(\Delta(z)\nu_i)],\ldots,\exp[\pm 2\pi i(1-\Delta_r)/(\Delta(z)\nu_i)]\right\}\tag{2.106}
$$

(3) R-enhanced and discriminant. The same argument as before yields

$$
\mu_i^{\nu_i} = \varrho_i \in \begin{cases} Sp(2r, \mathbb{Z}) & \mathcal{H} = 1 \\ PSp(2r, \mathbb{Z}) & \mathcal{H} = 2. \end{cases} \tag{2.107}
$$

To find the proper root of  $\varrho_i$  (more precisely: to determine the minimal polynomial of  $\mu_i$ ) one compares with the case in which the geometry is locally the product of a special geometry with fiber an Abelian variety of dimension  $(r - 1)$  – the Albanese variety (cf. §. [2.9\)](#page-32-0) – on which  $\exp[2\pi i \mathcal{E}/(\Delta(z) \nu_i)]$  acts by polarized automorphisms times an elliptic fibration over a small disk with a Kodaira central fiber of the appropriate type. Up to conjugacy in  $Sp(2r, \mathbb{Q})$  (and overall sign for  $\varkappa = 2$ )  $\mu_i$  is the block diagonal matrix having a  $(2r-2) \times (2r-2)$  block of eigenvalues

$$
\left\{ \exp[\pm 2\pi i(1-\Delta_1)/(\Delta(z)\nu_i)], \ldots, \exp[\pm 2\pi i(1-\widehat{\Delta(z)})/(\Delta(z)\nu_i)], \ldots \right\}
$$
 (2.108)

and a  $2 \times 2$  block which is the monodromy obtained from the Kodaira monodromy of the fiber type in  $\mathscr X$  by the base change  $z \mapsto z^{\nu_i} = w$  (see e.g. **table 5.2** of [\[86\]](#page-153-2)).

### <span id="page-39-0"></span>2.10.3 The structure theorem

The structure theorem of VHS gives the period map in terms of the neat monodromy group  $\Upsilon$  [\[37–](#page-150-0)[40\]](#page-151-3). By definition  $\Upsilon$  is a discrete subgroup of the algebraic group  $Sp(2r, \mathbb{Q})$  defined over Q. The Q-Zariski closure  $\overline{\Upsilon}^{\mathbb{Q}}$  of  $\Upsilon$  is the smallest Q-algebraic group containing  $\Upsilon$ .  $\Upsilon$  and  $\overline{\Upsilon}^{\mathbb{Q}}$ 

are algebraically indistinguishable, in the sense that no algebraic invariant can discriminate between them.  $\overline{\Upsilon}^{\mathbb{Q}}$  is also called the algebraic monodromy group. Let  $\overline{\Upsilon}^{\mathbb{Q}}(\mathbb{R}) \subset Sp(2r, \mathbb{R})$ be the *real Lie group* of R-valued points of the Q-algebraic group  $\overline{\Upsilon}^{\mathbb{Q}}$ : it is a semi-simple Lie group with finitely many connected components.<sup>[34](#page-40-0)</sup> We write  $H$  for a real Lie group such that  $\overline{\Upsilon}^{\mathbb{Q}}(\mathbb{R}) \times H$  is a maximal subgroup of  $Sp(2r, \mathbb{R})$  and write K, L for maximal compact subgroups of the real Lie groups  $\overline{\Gamma}^{\mathbb{Q}}(\mathbb{R})$  and H, respectively. The structure theorem says that the period map factorizes through

$$
\mathring{\mathscr{B}} \to \left(\Upsilon \backslash \overline{\Gamma}^{\mathbb{Q}}(\mathbb{R})/K\right) \times H/L \to Sp(2r, \mathbb{Z}) \backslash Sp(2r, \mathbb{R})/U(r) \equiv \mathscr{S}_r,\tag{2.109}
$$

and is the constant map in the factor  $H/L$ .

Corollary 1. In special geometry

- (1) if  $\Gamma$  is finite p is the constant map;
- (2) if  $\Gamma$  is Abelian p is the constant map;
- (3) if  $\Gamma = \Gamma_1 \times \Gamma_2$  the period map decomposes into two distinct period maps  $p_1$  and  $p_2$ factoring, respectively, through  $\Gamma_1 \backslash \overline{\Gamma_1}^{\mathbb{Q}}(\mathbb{R})/K_1$  and  $\Gamma_2 \backslash \overline{\Gamma_2}^{\mathbb{Q}}(\mathbb{R})/K_2$ .

A special geometry where  $p$  is the constant map is called *isotrivial*. All its smooth fibers are isomorphic as polarized Abelian varieties. In the situation (3) if one of the two maps is constant we say that the special geometry is quasi-isotrivial.

**Corollary 2.** The R-algebraic monodromy group  $M \equiv \overline{\Upsilon}^{\mathbb{Q}}(\mathbb{R})$  of a non-isotrivial special geometry is isogeneous to

$$
M_1 \times M_2 \times \cdots \times M_r \tag{2.110}
$$

where  $M_i$  are **simple, non-compact** real Lie subgroup of  $Sp(2r, \mathbb{R})$  of odd Mumford-Tate type and  $2r = \bigoplus_i V_i$  for a real symplectic representation  $V_i$  of  $M_i$ .

For the list of allowed pairs  $(M_i, V_i)$  see e.g. the table in [\[88\]](#page-153-4) or [\[38\]](#page-150-1). For a given rank  $r$  there are finitely many allowed groups  $M$ , and we may classify the special geometries group by group. The simpler R-algebraic monodromy  $M$  the simpler the corresponding classification program. The case  $M = \{1\}$  is specially easy.

This has a consequence already noted in [\[17\]](#page-149-0):

**Corollary 3.** (1) If the discriminant  $\mathcal F$  is a simple normal crossing divisor, the geometry is a product of r rank-1 geometries; (2) let  $\mathscr{R} = \mathbb{C}[u_1, \dots, u_r, v_1, \dots, v_s]$  and  $\mathcal{D} = (f_1) + (f_2)$ where  $f_1 \in \mathbb{C}[u_1, \dots, u_r]$ ,  $f_2 \in \mathbb{C}[v_1, \dots, v_s]$ . Then the special geometry is the product of two special geometries with chiral rings  $\mathscr{R}_1 \in \mathbb{C}[u_1, \dots, u_r], \mathscr{R}_2 \in \mathbb{C}[v_1, \dots, v_s].$ 

<span id="page-40-0"></span><sup>&</sup>lt;sup>34</sup> To show the last statement one uses the fact that the Coulomb branch is - by definition - quasi-projective as a complex manifold hence Liouvillic for the plurisubharmonic functions.

**Example 2.4.** The discriminant of an *indecomposable* rank-2 geometry contains at least:

- (1) one knotted component;
- (2)  $or$  two unknotted components crossing non-transversely;
- (3) or three unknotted components with pairwise transverse crossings at the origin.

The geometries of the SCFT numbered  $#20, #21, #60$  in the tables of [\[89\]](#page-153-5) have a minimal discriminant of type (1); those numbered  $\#1, \#7, \#11, \#14, \#16, \#18, \#51-\#53, \#61-\$  $\#65$ ,  $\#68$  have a minimal discriminant of type (2); the ones numbered  $\#24$ ,  $\#27$ ,  $\#30$  a minimal discriminant of type (3). All other SCFT in the tables of [\[89\]](#page-153-5) have non-minimal discriminants with more components. Model #32 is not indecomposable.

The case of higher r will be sketched in section [9.](#page-122-0)

#### <span id="page-41-0"></span>2.10.4  $\Gamma$  finite: isotrivial special geometries

The isotrivial special geometries have several (equivalent) characterizations:

Fact 9. For a  $\mathbb{C}^{\times}$ -isoinvariant special geometry the following are equivalent:

- (1)  $\Gamma$  is finite or, equivalently,  $M = \{1\}$ ;
- (2) all smooth fibers of the special geometry  $\mathscr{X} \to \mathscr{C}$  are isomorphic as polarized Abelian varieties to a fixed variety A;
- (3) there is a finite cover  $f: \check{\mathscr{C}} \to \mathscr{C}$ , branched only over the discriminant, such that the pulled-back fibration is trivial, i.e.  $f^*\mathscr{X} = A \times \check{\mathscr{C}}$  for a fixed polarized Abelian variety A. In particular the special geometry  $\pi: \mathscr{X} \to \mathscr{C}$  is a global quotient [\[37–](#page-150-0)[40\]](#page-151-3)

$$
A \times \check{\mathscr{C}} \xrightarrow{\phantom{X}p_2} \check{\mathscr{C}}
$$
  
\n
$$
\downarrow \qquad \qquad \downarrow
$$
  
\n
$$
\mathscr{X} \equiv (A \times \check{\mathscr{C}})/\Gamma \xrightarrow{\pi} \mathscr{C} = \check{\mathscr{C}}/\Gamma
$$
  
\n(2.111)

(4) all local monodromies around the discriminant components  $\mathcal{D}_i$  have finite order.

 $(1) \Leftrightarrow (2) \Leftrightarrow (3)$  is very well-know in VHS and valid for all holomorphic fibration in polarized Abelian varieties over a quasi-projective base. We summarize the arguments: (1)  $\Leftrightarrow$  (2) is the structure theorem together with the Godemen criterion [\[90\]](#page-153-6). (1) + (2)  $\Rightarrow$  (3) follows from the obvious Galois cover of the complement  $\check{\check{\mathscr{C}}} = \check{\check{\mathscr{C}}}/\text{ker }\varrho$  with finite deck group Γ, together with Riemann's existence theorem [\[91\]](#page-153-7) which extends it to a finite branched cover  $\check{\mathscr{C}} \to \mathscr{C}$ . (3)  $\Rightarrow$  (1) follows from the fact that  $\Gamma$  is a subgroup of the finite group of polarized automorphisms of A.

The tricky (and important for non-perturbative physical considerations) part of the statement is the equivalence of (4) with one (hence all) of the first 3 conditions. It is the only point where one uses that  $\mathscr{X} \to \mathscr{C}$  is a  $\mathbb{C}^{\times}$ -isoinvariant special geometry and not merely a fibration in polarized Abelian varieties. (1)  $\Rightarrow$  (4) is trivial, and one needs to prove (4)  $\Rightarrow$  (1). There is a *finite* cover  $\mathscr{B} \to \mathscr{C}$ , branched only on the discriminant, such that the pull-back of the Abelian family to  $\mathscr B$  has trivial local monodromies. The pull-back of the special Kähler metric extends to a regular metric over the inverse image of the discriminant components of types  $I_0^*$ ,  $II^*$ ,  $III^*$ ,  $IV^*$ . If we have only local monodromies of these types, the pulled back special Kähler metric is conic and regular, hence it is the flat metric, which implies that the period  $\tau_{ij}$  is constant. Otherwise we have a covering  $\mathscr{B} \to \mathscr{A}$ , branched only on discriminant components of types II, II and IV , such that the singular special Kähler metric on  $\mathscr B$  is the pull back of a regular one in  $\mathscr A$  which then must be flat. In all cases the period matrix  $\tau_{ij}$  is locally constant.

Many  $\mathcal{N} = 2$  SCFT have isotrivial geometries: *(i)* all  $\mathcal{N} \geq 3$  models (regarded as special  $\mathcal{N} = 2$  SCFT); *(ii)* all models with  $\varkappa \notin \{1,2\}$ ; *(iii)* all rank-1 models. In principle the isotrivial special geometries can be constructed and classified in all ranks  $r$  [\[55\]](#page-151-4). More details will be given elsewhere.

## 2.11 "Stratification" of the special geometry

The Coulomb branch  $\mathscr C$  is naturally stratified [\[16,](#page-149-1) [17\]](#page-149-0). The closure of a codimension-s stratum is a connected component of a multiple intersection of discriminant components  $\mathcal{D}_{1_1} \cap \cdots \cap \mathcal{D}_{i_s}$  of codimension s. To this stratification there corresponds a "stratification" of the special geometry in the sense that the special geometry  $\mathscr{X} \to \mathscr{C}$  induces a Lagrangian fibration  $\mathscr{X}_{a_s} \to \mathscr{C}_{a_s}$  over each codimension-s stratum  $\mathscr{C}_{a_s}$  whose generic fibers are polarized Abelian varieties of dimension  $\dim \mathscr{C}_{a_s}$ .

Let  $\mathcal{D}_{a_1}$  be an irreducible component of the discriminant and  $u \in \mathcal{D}_{a_1}$  a general point. To u we may associate a polarized Abelian variety  $A<sub>u</sub>$ , namely the Albanese variety of the normalization of a connected component of the fiber  $\mathscr{X}_u$ .  $A_u$  is well-defined since it does not depend on the chosen component [\[45,](#page-151-5) [46,](#page-151-6) [50\]](#page-151-7). We can glue the various  $A_u$ 's along  $\mathcal{D}_{a_1}$  to get a fibration  $\mathscr{X}_{a_1} \to \mathcal{D}_{a_1} \equiv \mathscr{C}_{a_1}$  in polarized Abelian varieties which we claim is a special geometry in its own right. The geometry  $\mathscr{X}_{a_1} \to \mathcal{D}_{a_1} \equiv \mathscr{C}_{a_1}$  may be seen as a symplectic reduction of  $\mathscr{X} \to \mathscr{C}$  in a sense which is more general than the usual Marsden-Weinstein-Meyer (MWM) one [\[43,](#page-151-8)[44\]](#page-151-2) which applies when the usual regularity assumptions of the MWM theorem do not hold, but the Lagrangian fibration is holomorphic.

### 2.11.1 A generalization of Marsden-Weinstein-Meyer symplectic reduction

We recall how it works the usual MWM symplectic reduction of a phase space  $X$  with symplectic form  $\Omega$ : first one considers a level subset  $\iota: h^{-1}(c) \hookrightarrow X$  of a Hamiltonian function  $h: X \to \mathbb{C}$ , and then one constructs the space Y which parametrizes the orbits of the flow generated by the Hamiltonian vector  $v(h)$  dual to dh. In general Y may be quite weird. In the lucky case when the orbit space  $Y$  happens to be a nice manifold, it inherits a symplectic structure  $\omega$  from the one of X such that

<span id="page-43-0"></span>
$$
\iota^* \Omega = \pi^* \omega \tag{2.112}
$$

where  $\pi: h^{-1}(c) \to Y$  is the canonical projection sending a point to the orbit it belongs. In this regular case Y is a phase space in its own right, called the *symplectic reduction* of X along the hypersurface  $h = c$ . The typical application is when h is the momentum map of a  $U(1)$  action on X. Unfortunately, most often Y is not a nice manifold, indeed it can be a very ugly space, and the symplectic reduction does not make sense unless the necessary regularity conditions are met.

To get the special geometric "stratification" one may try to apply the MWM strategy to the zero level set of the prime homogeneous Hamiltonian  $p_a \in \mathcal{R}$  such that  $\mathcal{D}_a = (p_a)$ . However now the level subset  $p_a^{-1}(0) \subset \mathcal{X}$  is non-smooth by definition. The divisor  $p_a^{-1}(0)$ contains several non-reduced irreducible components intersecting each other in a Kodaira pattern. The zero-section s is contained in one component of multiplicity 1 of  $p_a^{-1}(0)$ . We focus on the normalization of this zero component, and consider the space parametrizing its orbits under the flow generated by the vector  $v(p_a)$ . Away from special points in  $\mathcal{D}_a$ , this orbit space  $\mathscr{X}_a$  is smooth. In facts the normalization of the zero component of the fiber over the generic point  $u \in \mathcal{D}_a$  is a fibration over the Albanese polarized Abelian variety  $A_u$  and the vector  $v(p_a)$  is tangent to the fiber [\[45\]](#page-151-5), so we may identify  $A_u$  with the fiber at  $u \in \mathcal{D}_a$ of the orbit space  $\mathscr{X}_a \to \mathcal{D}_a$ . In other words  $\mathscr{X}_a \to \mathcal{D}_a$  is the fibration with generic fiber  $A_u$ . Note that if we start with the normalization of any other irreducible component we would get the same orbit space  $\mathscr{X}_a$ , which then is an 'intrinsic' property of the divisor  $\mathcal{D}_a$ . It is easy to check tha  $\mathscr{X}_a$  carries a natural symplectic form such that [\(2.112\)](#page-43-0) holds.<sup>[35](#page-43-1)</sup>  $\mathscr{X}_a$  may (in facts, should) have singularities over a codimension-1 locus in  $\mathcal{D}_a$ . If these singularities admit a crepant resolution, the resolved  $X_a$  is a *bona fide* special geometry of rank  $(r-1)$ and we can consider recursively its own "stratification". We assume that this can be done.

## 2.12 "Stratification" vs. Coulomb dimensions

"Stratification" gives new information on the allowed Coulomb dimensions. The formulae of [\[22\]](#page-150-2) reviewed in §. [2.5.1](#page-22-0) assumed the existence of a *regular* coordinate axis  $H_i$  not contained in the discriminant. While this is often the case, there are many geometries where no such "good" axis is present, and we need a more general analysis.

We may assume without loss that  $\Delta_j > 1$  for all j. Fix an axis  $H_i$  and let  $\ell$  be the smallest integer such that  $H_i$  is contained in a codimension- $\ell$  (open) stratum; clearly  $1 \leq \ell \leq r$ . Let

<span id="page-43-1"></span><sup>&</sup>lt;sup>35</sup> As mentioned before, in the stable case (monodromy type  $I_m$ ) there is some additional subtlety related to the type of the characteristic cycle. This does not affect the conclusion, just requires some more care in the precise definitions of the various entities.

 $u \in H_i$  be a generic point. Let  $\mathscr{X}_{u,0}$  be the unique component of the fiber which contains the smooth point  $s(u)$ . The normalization of  $\mathscr{X}_{u,0}$  is a fibration over an Albanese variety  $A_u$  of dimension  $\ell$ .  $T_{s(u)}\mathscr{X}_{u,a}$  is isomorphic to  $T_u^*\mathscr{C}$ . The automorphism  $\xi_u = \exp(2\pi i \mathscr{E}/\Delta_i)$ fixes u and acts linearly on  $T_{s(u)}\mathscr{X}_{u,a}$  with eigenvalues

<span id="page-44-0"></span>
$$
\exp(2\pi i(\Delta_j - 1)/\Delta_i), \quad j = 1, 2, \cdots, r \tag{2.113}
$$

Since  $\xi_u$  is an automorphism of the fiber, it preserves its Albanese variety  $A_u$  and acts on it as a polarized automorphism for some *polarization*. Hence  $\ell$  out of the r eigenvalues [\(2.113\)](#page-44-0) are the eigenvalues of the analytic representation  $\sigma$  of a polarized automorphism of the Abelian variety  $A_u$  of dimension  $\ell$ . Moreover  $\xi_u$  acts trivially on  $\mathbb{C} du_i|_u \subset T_u\mathscr{C}$  which corresponds to a vector field tangent to the Albanese variety. Hence  $\exp(-2\pi i/\Delta_i)$  is one of the eigenvalues of  $\sigma$ . Writing  $\Delta_i = m/n$  with  $gcd(m, n) = 1$  and  $n < m$ , we get  $\phi(m) \leq 2\ell$ .

**Fact 10.** Let  $H_i$  be an axis of dimension  $\Delta_i$  which is new in rank  $\ell$ . The axis  $H_i$  is not contained in a stratum of  $\mathscr C$  of dimension  $\lt \ell$ . In particular, if  $\Delta_i$  is a new dimension  $H_i$ does not belong to the discriminant. If  $\Delta_i$  is new in rank r, the full r-tuple  $\{\Delta_1,\ldots,\Delta_r\}$  is given by the formulae of §.[2.5.1](#page-22-0). If  $\Delta_i$  is new in rank  $\ell$ , at least a subset of  $\ell$  dimensions are given by those formulae with r replaced by  $\ell$ .

In particular, when the rank is 2 and  $\Delta_i$  is new rank-2 the corresponding axis  $H_i$  does not belong to the discriminant, and the dimension formulae of §. [2.5.1](#page-22-0) can be used giving a small set of possibilities for the second dimension.

# 3 Classification: the inverse problem

One possible strategy to construct/classify all  $\mathcal{N} = 2$  SCFTs is to construct first all possible C <sup>×</sup>-isoinvariant special geometries and then work out the (finitely many) SCFT associated with each one of them. This program was initiated and carried on to a certain extend by Argyres and coworkers (see [\[5–](#page-148-0)[17\]](#page-149-0) and references therein). The first step in such a program requires to construct all  $\mathbb{C}^{\times}$ -isoinvariant special geometries, or at least to chart their "geography/zoology", by following either one of two natural strategies:

- (A) rank after rank;
- (B) or general subclass after general subclass in increasing order of "complexity".

Subprogram  $(A)$  is completed in rank  $1 \left[ 7-9, 23, 24 \right]$  and in good shape in rank  $2 \left[ 18-20 \right]$ , while subprogram (B) was initiated in [\[25\]](#page-150-5) with the introduction of a first numerical invariant,  $\kappa$ , which partially characterize the complexity subclasses. The subclasses with  $\kappa \notin \{1,2\}$  can be worked out explicitly for arbitrary r [\[55\]](#page-151-4) (at least when  $\mathscr C$  is smooth). The Lagrangian SCFT are also classified for arbitrary  $r$  [\[95\]](#page-154-0), but not all their global special geometries are explicitly known. In our view the best course is to follow a "'mixed" strategy where we describe the simplest subclasses for arbitrary ranks and then discuss the residual "hard" geometries rank by rank. We propose the following coarse classification of the indecomposable, non-free, unitary geometries:



Classes I and IV are essentially charted for all ranks. Classes II and III are very constrained, and a systematic analysis may be doable. In these cases one aims to produce strong necessary conditions which rule out such geometries except under "exceptional" circumstances such as peculiar values of the dimensions  $\{\Delta_1, \ldots, \Delta_r\}$  and of the local monodromies determining their "zoology". For classes V and VI one falls back to the rank by rank strategy. Class VI is the less amenable to analysis and a priori the largest one, but

Hope (Supported by some experimental evidence). Class VI is hard but empty.

Each class is further divided into subclasses.

## 3.1 The inverse problem

For the last two classes of geometries at the moment there is no better plan than studying one rank at the time. In this direction a natural idea, already put forward in [\[16,](#page-149-1) [17\]](#page-149-0), is to exploit the "stratification" of the special geometry in lower rank ones, and try to construct recursively all possible geometries in rank r by "gluing" together local rank- $(r - 1)$  special geometries around codimension-1 loci  $\mathcal{D}_a$  and a "boring" rank-r open stratum of generic vacua in the complement  $\mathbb{C}^r \setminus \cup_a \mathcal{D}_a$  in "all consistent ways". This strategy leads to what we call the Inverse problem which now we state in concrete terms:

Inverse Problem. Given the following data:

- (i) the Coulomb dimensions  $\{\Delta_1, \ldots, \Delta_r\},\$
- (*ii*) the discriminant  $\mathcal{D}$ ,

(iii) the conjugacy class of the local monodromy at each irreducible component  $\mathcal{D}_a$  of  $\mathcal{D}_a$ , which we label by the corresponding Kodaira type  $C_a$ ;

construct the indecomposable special geometries of rank r with these data or conclude that no such geometry exists.

We can restrict the set of allowed data. First of all we may consider only  $r$ -tuple  $\{\Delta_1,\ldots,\Delta_r\} \equiv \lambda \{d_1,\ldots,d_r\}$  with  $\Delta_i$ 's in the finite set  $\cup_{s=1}^r \Xi(s)$  of allowed dimensions in rank r, since otherwise the answer is negative for obvious reasons. We may also assume that  $1/\lambda$  is integral  $(\varkappa = 1)$  or half-integral  $(\varkappa = 2)$  since otherwise the geometry is automatically isotrivial and hence can be constructed/classified by group-theoretical means. If some of the  $\Delta_i$  is equal 2 the geometry is *non-rigid* (see §.[5\)](#page-54-0). A **folk-theorem** states that a non-rigid special geometry describes a Lagrangian SCFT which is more conveniently studied by different (physical) means. There are many other restrictions in the r-tuple from the dimension formulae and their generalizations. We can limit ourselves to admissible  $r$ tuples  $\{\Delta_1,\ldots,\Delta_r\}$  with  $\Delta_i\neq 1,2$  and  $2\lambda\in\mathbb{N}$  without essential loss. We know that when all Kodaira types  $C_a$  are semi-simple the geometry is isotrivial, therefore we may assume that at least one monodromy class  $C_a$  is  $I_n$  or  $I_n^*$  with  $n > 0$ . This, in particular, requires the discriminant  $\mathcal D$  not to be normal crossing; more generally we have the condition that  $\pi_1(\mathscr{C} \setminus \mathcal{D})$  has an infinite quotient  $\Gamma$  which is semisimple with a symplectic action on  $\mathbb{Q}^{2r}$ . When focusing on the "harder" classes IV, VI we may assume the action to be irreducible.

In physical terms the Inverse problem asks to find a common UV completion to the several low energy effective theories describing locally the degrees of freedom which are light along each discriminant component, if this (geometric) global UV completion exists.

### **Fact 11.** There are at most finitely many geometries for a given datum  $(\{\Delta_i\}, \mathcal{D}, \{C_a\})$ .

The general expectation is that the special geometry associated to an inverse datum, when it exists, is actually unique.<sup>[36](#page-46-0)</sup> The idea is that the existence of a global special geometry is already a very overdetermined problem, which can be solved positively only for a tiny set of data  $(\{\Delta_i\}, \mathcal{D}, \{C_a\})$ . It looks extremely unlikely that we have more than one solution for the same datum. The finiteness in the Fact follows from the fact that there are finitely many fibrations in Abelian r-varieties over  $\mathscr{C} \setminus \mathcal{D}$  for fixed  $\mathcal{D}$ . For almost all choices of  $D$ , the monodromy of all these finitely-many fibrations would not be compatible with any Kodaira-type/dimension data, so that *only a "short list" of discriminants are allowed in a* given rank r and for a given discriminant  $\mathcal D$  only a "small set" of Kodaira-type/dimension datum is consistent. We shall discuss in  $\S$ . [9](#page-122-0) the conditions on  $\mathcal{D}$ . They turn out to be very restrictive.

For each consistent set of "stratification" data there exist only finitely many Abelian fibrations, but this is just one of the many geometric structures required for a special geometry.

<span id="page-46-0"></span><sup>&</sup>lt;sup>36</sup> In this statement is important that  $C_a$  is defined as a Kodaira type, i.e. as a conjugacy class in  $SL(2,\mathbb{Z})$ ; if  $C_a$  is seen as a conjugacy class in a larger group, then multiple geometries do exist.

To be a special geometry the total space of an Abelian family over  $\mathscr{C} \setminus \mathcal{D}$  should have a symplectic structure which extends over the "bad" fibers such that all fibers are Lagrangian. This is quite a severe restriction to impose on the few surviving fibrations, and for most consistent-looking data no one will pass the test. The existence of more than one geometry for the same datum looks totally unlikely.

The inverse problem may be studied with different aims. In decreasing order of ambition:

- (1) to construct all  $\mathbb{C}^{\times}$ -isoinvariant geometries recursively in the rank r;
- (2) to discover interesting properties which hold for all such geometries;
- (3) to get necessary conditions for the existence of a geometry;
- (4) to construct examples with prescribed/desired properties.

# 3.2 Taking "stratification" seriously

At first sight the weak aspect of the **Inverse problem** strategy is that we need to specify a *priori* a discriminant  $\mathcal{D} \subset \mathscr{C}$  (or, equivalently, a special divisor  $\mathscr{S} \subset \mathscr{P}$ ). While the datum of a dimension r-tuple  $\{\Delta_1, \ldots, \Delta_r\}$  is to be chosen from a known *finite* set, and the Kodaira classes  $C_i$  from a known discrete set which becomes finite when we consider conjugacy classes in  $Sp(2r, \mathbb{R})$  instead of  $Sp(2r, \mathbb{Z})$ , the discriminant  $\mathcal{D} = (P)$  a priori is just a square-free homogeneous element  $P \in \mathcal{R}$  which may have (and indeed has) several components, with arbitrarily bad singularities, intersecting each other is all ugly non-transverse ways. At face value, describing the set of meaningful initial data for the problem looks harder than finding needles in a haystack.

Luckily it is not so. Consider (the normalization of) an irreducible component  $\mathcal{D}_a$  of the discriminant. If we take "stratification" seriously we conclude that it is the Coulomb branch of a  $\mathbb{C}^{\times}$ -isoinvariant special geometry on its own right. Thus  $\mathcal{D}_{a}$  cannot be an arbitrary hypersurface in  $\mathscr{C}$ , it should satisfy the conditions of special geometry.  $\mathcal{D}_a$  cannot be, say, a quasi-cone over a smooth Calabi-Yau. The affine variety  $\mathcal{D}_a$  needs not to be smooth, so its normalization may not be just a copy of  $\mathbb{C}^{r-1}$  (although this is rather common in examples), but at the very least it should be a quasi-cone over a Fano variety or (according to physical intuition) satisfy the even more restrictive condition of admitting a polynomial parametrization of the form  $u_i = u_i(v_i)$ .

Going on, consider the divisor

$$
\mathcal{D}'_a \equiv \mathcal{D}_a \cap (\mathcal{D} \setminus \mathcal{D}_a) \subset \mathcal{D}_a \tag{3.1}
$$

(or rather the respective normalizations). The pair  $(\mathcal{D}_a, \mathcal{D}'_a)$  should be in the list of allowed pairs (Coulomb branch, Discriminant) in rank  $(r - 1)$ , a condition that we call *hereditarity*. This sets a strong condition on D.

Then there are two other sets of constraints on  $D$ . The first one comes from topology: the fundamental group of the complement  $\pi_1(\mathscr{C} \setminus \mathcal{D})$  should have the right properties (not to be finite, or solvable, or a product, etc.). Thus D cannot be normal crossing, nor  $\mathcal{D}_1 + \mathcal{D}_2$ with all components of  $\mathcal{D}_1$  crossing transversely the ones of  $\mathcal{D}_2$ , etc.

However the most strong condition on  $\mathcal D$  comes from the hero of the present paper: rigidity which will be the central topic of the rest of the note.

We introduce a coarse equivalence relation between discriminants or rather special divisors ("commensurability").  $\mathscr{S} \sim \mathscr{S}'$  iff their complements  $\mathscr{P} \setminus \mathscr{S}$  and  $\mathscr{P} \setminus \mathscr{S}'$  admit a common finite unbranched cover, so that their fundamental groups have an isomorphic finite-index normal subgroup.

**Hope** (Well supported). In rank-r there are only finitely many equivalence classes of (deformations classes of ) allowed special divisor.

The validity of **Hope** is a tautological consequence of the (physically expected) fact that there are only finitely many special geometries at a given rank r. But what we really hope for is that the allowed divisor classes can be listed a priori without working out the classification of the geometries themselves. In the rest of the paper the hope will be illustrated by several examples in various situations.

## 4 Abelian families arising from a special geometry

We adopt a viewpoint (and language) which is more familiar in the context of Calabi-Yau moduli spaces and mirror symmetry. As stated in the introduction, the two problems are formally similar.

Suppose we are given an  $\mathbb{C}^{\times}$ -isoinvariant Abelian family  $\mathscr{X} \to \mathscr{C}$ , or equivalently a family

<span id="page-48-1"></span>
$$
\mathscr{A} \to \mathring{\mathscr{P}} \tag{4.1}
$$

where  $\mathscr{P} \equiv \mathscr{P} \setminus \mathscr{S}$  with  $\mathscr{S}$  the special divisor in  $\mathscr{P}$ . The Abelian family  $\mathscr{A}$  defines a monodromy representation  $\mu: \pi_1(\tilde{\mathscr{P}}) \to Sp(2r, \mathbb{Z})$  and also a flat Gauss-Manin (GM) differential equation of rank 2r of the form

<span id="page-48-0"></span>
$$
\nabla \Phi \equiv d\Phi - \Phi A = 0 \tag{4.2}
$$

where the connection A is locally a  $2r \times 2r$  matrix of holomorphic differentials and  $\Phi$  is the fundamental solution (a  $2r \times 2r$  matrix normalized to be the identity at a reference point  $* \in \mathscr{P}$ ). The differential equation has only *regular singularities* along  $\mathscr{S}$  [\[96\]](#page-154-1). When  $\dim \mathscr{P} = 1$  it reduces to a classical Picard-Fuchs equation, and we use the terms Gauss-Manin connection and Picard-Fuchs system interchangeably. For simplicity we consider the smooth case where  $\mathscr{P} = \mathbb{P}(q_1, \ldots, q_r)$  and we work in the affine patch  $z_r \neq 0$ . In this patch  $($   $\cong$   $\mathbb{C}^{r-1}$  the connection can be written as

$$
\nabla = d + \sum_{i} \frac{A_i(u)}{p_i(u)} \tag{4.3}
$$

where the  $p_i(u)$  are the equations of the irreducible components of  $\mathscr S$  and the  $A_i(u)$  are  $2r \times 2r$  matrices of holomorphic differentials. The connection should be flat

$$
\nabla^2 = dA + A^2 = 0, \qquad A \stackrel{\text{def}}{=} \sum_i \frac{A_i(u)}{p_i(u)} \tag{4.4}
$$

The Riemann-Hilbert correspondence states that there is a one-to-one equivalence between the GM PDEs and the monodromy representation up to the obvious equivalences. In our case the monodromy takes values in  $Sp(2r, \mathbb{Z})$ . Since we are mainly interested in irreducible non-isotrivial geometries, we may assume that the monodromy group  $\Gamma$  acts irreducibly and contains non-trivial unipotent elements.<sup>[37](#page-49-0)</sup> Since  $\pi_1(\mathscr{P})$  is trivial, the monodromy is generated by the images of loops encircling the irreducible components  $S_i$  of the special divisor  $\mathscr{S} = \sum_i \mathcal{S}_i$ . The monodromy group acts on the solution as

$$
\mu([\ell]) \cdot \Phi = \Phi \Big|_{\substack{\text{analytically} \\ \text{conjug } \ell \\ \text{along } \ell}} \tag{4.5}
$$

where  $\ell$  is a loop based at a reference point  $* \in \mathcal{P}$  which represents a given class in  $\pi_1(\mathcal{P}, *)$ . The Zariski closure over C of the group  $\Gamma \subset Sp(2r, \mathbb{Z})$  is the *differential Galois group of the* PDE [\(4.2\)](#page-48-0). We use the symbol  $\Pi$  for a column 2r-vector given by a linear combination of the columns of  $\Phi$  (with complex coefficients). The symplectic  $2r \times 2r$  matrix will be written  $\Omega$  (so  $\gamma \in \Gamma \Rightarrow \gamma^t \Omega \gamma = \Omega$ ). In a basis where the monodromy elements are integral symplectic matrices, the entries of  $\Pi$  are the periods of some holomorphic differential  $\eta$  with respect to a symplectic basis  $\{A^i, B_j\}$  of 1-cycles on the fibers

$$
\Pi = \begin{pmatrix} \int_{B^i} \eta \\ \int_{A_i} \eta \end{pmatrix} \tag{4.6}
$$

The period vector  $\Pi$  is multivalued on  $\mathscr{\dot{P}}$ . We see  $\Pi$  as a map defined on the universal cover  $\mathscr{\mathscr{P}}$  of  $\mathscr{\mathscr{P}}$ 

$$
\Pi \colon \widetilde{\mathscr{P}} \to \mathbb{P}^{2r-1} \tag{4.7}
$$

equivariant with respect to the deck group  $\pi_1(\tilde{\mathscr{P}})$ 

$$
\sigma^* \Pi = \mu(\sigma) \Pi, \qquad \sigma \in \pi_1(\hat{\mathscr{P}}). \tag{4.8}
$$

<span id="page-49-0"></span><sup>37</sup> If Γ does not contain non-trivial unipotent elements it is automatically finite.

Fact 12. The family [\(4.1\)](#page-48-1) of Abelian varieties arises from a  $\mathbb{C}^{\times}$ -isoinvariant special geometry, iff its GM equation has a preferred period  $\Pi$  such that its image in  $\mathbb{P}^{2r-1}$  is a Legendre submanifold of the canonical contact structure of  $\mathbb{P}^{2r-1}$ , that is,

<span id="page-50-0"></span>
$$
\Pi^t \,\Omega \,d\Pi = 0 \tag{4.9}
$$

while the  $(1, 1)$ -form i d $\Pi^{\dagger} \Omega d\Pi$  is positive definite.

*Proof.* (Adapted from [\[41\]](#page-151-9)) Write  $\Pi = (a_i^D, a^j)^t$   $(i, j = 1, ..., r)$ . For a Legendre submanifold in generic position, locally there exists a holomorphic function  $\mathcal{F}(a^j)$ , homogeneous of degree 2, such that

$$
a_i^D = \frac{\partial \mathcal{F}(a)}{\partial a^i}.
$$
\n(4.10)

On the other hand  $a^i = \int_{A_i} \eta$  implies  $\eta = \sum_i a^i \omega_i$  where  $\{\omega_i\}$  are holomorphic differential normalized as  $\int_{A_i} \omega_j = \delta^i{}_j$ . Then the period matrix of the Abelian variety is

$$
\tau_{ij} = \int_{B^i} \omega_j = \frac{\partial}{\partial a^j} \int_{B^i} \eta = \frac{\partial}{\partial a^j} a_i^D = \frac{\partial^2 \mathcal{F}(a)}{\partial a^i \partial a^j}.
$$
\n(4.11)

In particular Im  $\tau_{ij} > 0$  by the second Riemann relation for the Abelian fibers. A posteriori the differential  $\eta$  defined by the Legendre condition is identified (when it exists) with the Seiberg-Witten differential  $\lambda$ . In other words, the Legendre-manifold condition is equivalent to the existence of a Seiberg-Witten differential on the pull-back of the Abelian family from  $\mathscr{\hat{P}}$  to  $\mathscr{\hat{C}}$ . When such differential exists, it pulls back to a Lagrangian fibration  $\mathscr{X}$  over  $\mathscr{\hat{C}}$ by taking as local pre-potential  $\mathcal F$  the one defining (locally) the Legendre submanifold. The Legendre condition is also called the "integrability" condition.  $\Box$ 

**Comparison with the Calabi-Yau case.** A family of Calabi-Yau 3-folds with  $h^{2,1}$  =  $r-1$  is also described by a GM connection with a period  $\Pi$  satisfying the Legendre condition [\[41\]](#page-151-9). The difference is that in the CY case the matrix Im  $\tau_{ij}$  has signature  $(r-1, 1)$  instead of being positive-definite. A part for this crucial sign condition, all equations look formally identical, so we can borrow methods developed in the context of Calabi-Yau periods and mirror symmetry.

### 4.1 The Legendre condition

A priori the Legendre condition may look a severe constraint on the family of Abelian varieties. However often it is not hard to satisfy. Consider the 1-form  $\Pi^t \Omega d\Pi$  for a generic period. If it extends over the special divisor  $\mathscr{S}$ , it is a global holomorphic 1-form on  $\mathscr{P}$  hence zero since the latter has  $h^{1,0} = 0$ . So the question boils down to the existence of a period Π which is "sufficiently regular" along the special divisors  $\mathcal{S}_a$ . Since the local monodromy around  $S_a$  corresponds (by construction) to the IR physics of the degrees of freedom which

purportedly becomes light there, "sufficiently regular" means that the period  $\Pi$  approaches along  $\mathcal{S}_a$  the ones describing the local physics of the light degrees of freedom, i.e. that  $\Pi$ agrees near  $S_a$  with one particular local period, the "good" one. The "good" local period  $\Pi$  exists because we choose the exponents of the monodromy (encoded in the datum  $C_a$  of the inverse problem) precisely to guarantee that the local geometry follows the appropriate model (cf. §. [2.9.2\)](#page-33-0).

Seen in this way, the Legendre condition is a requirement on the connection formula for the PDE: the "good" local period near  $S_a$ , when analytically continued to a neighborhood of  $\mathcal{S}_b$ , should agree with the "good" local solution there for all b. In the language of the **inverse** problem: the several local geometries along the strata should "glue well" together. Alternatively one has to show that the holomorphic one-form  $\Pi^t \Omega d\Pi$  vanishes in a neighborhood of  $S_a$  and not just asymptotically when we approach it. This point of view clarifies that the special period  $\Pi$  (when it exists) is unique.

In the classical case of rank-2 we may be very precise. Here we use the analogy with mirror symmetry for one-modulus Calabi-Yau's [\[97,](#page-154-2) [98\]](#page-154-3).

## 4.2 The classical case  $r = 2$

In the case  $r = 2$  the normalization of  $\mathscr P$  is just the Riemann sphere  $\mathbb P^1$  for all pairs of Coulomb dimensions  $\{\Delta_1, \Delta_2\}$ , while  $\mathring{\mathscr{P}} \simeq \mathbb{P}^1 \setminus \{z_1, \ldots, z_k\}$  where  $\{z_i\}$  are k distinct points on the sphere. The Gauss-Manin PDE reduces to a Fuchsian ODE. Being primarily interested in irreducible geometries, we assume the monodromy to be irreducible. Then, by a "choice of gauge" the ODE can always be written in the form [\[99,](#page-154-4) [100\]](#page-154-5)

$$
\frac{d}{dz}\Pi = \Pi A, \qquad A \equiv \sum_{i} \frac{A_{\ell}}{z - z_{\ell}} \tag{4.12}
$$

for constant  $4 \times 4$  matrices  $A_{\ell}$  satisfying the non-resonant condition.<sup>[38](#page-51-0)</sup> The conjugacy class in  $GL(4,\mathbb{C})$  of the local monodromy  $\mu_{\ell}$  at  $z_{\ell}$  is

<span id="page-51-1"></span>
$$
\[\mu_{\ell}\] = \[\exp(2\pi i A_{\ell})\].\tag{4.13}
$$

The equation [\(4.13\)](#page-51-1) can be recast in the form of the 4-th order Fuchsian linear equation for a scalar function y:

<span id="page-51-2"></span>
$$
Ly = 0, \quad \text{where } L = \frac{d^4}{dz^4} + a_3(z)\frac{d^3}{dz^3} + a_2(z)\frac{d^2}{dz^2} + a_1(z)\frac{d}{dz} + a_0(z) \tag{4.14}
$$

whose coefficients are rational functions on  $\mathbb{P}^1$  determined by the points  $z_i$  and the monodromy representation of the original equation. In [\[98\]](#page-154-3) (see also [\[97\]](#page-154-2)) it is shown that the coefficients  $a_i(z)$  of a 4-th order ODE [\(4.14\)](#page-51-2) with differential Galois group contained in

<span id="page-51-0"></span><sup>&</sup>lt;sup>38</sup> That is: if two eigenvalues of  $A_{\ell}$  are equal mod 1, then they must be equal.

 $Sp(4,\mathbb{C})$  satisfy the differential relation<sup>[39](#page-52-0)</sup>

<span id="page-52-1"></span>
$$
4a_2a_3 - 8a_1 - a_3^3 + 8a_2' - 6a_3a_3' - 4a_3'' = 0.
$$
\n(4.15)

Let  $y_1, y_2, y_3, y_4$  be four linearly independent solutions to the ODE [\(4.14\)](#page-51-2). Out of these 4 functions we may produce 6 independent  $2 \times 2$  Wronskians

$$
W_{ab} \equiv \begin{vmatrix} y_a & y_b \\ y'_a & y'_b \end{vmatrix}, \quad 1 \le a < b \le 4.
$$
 (4.16)

In a general 4-th order Fuchsian ODE the Wronskians satisfy a linear ODE of order 6. The Almkvist-Zudilin theorem [\[98\]](#page-154-3) states that when the coefficients obey the condition [\(4.15\)](#page-52-1) the Wronskians  $W_{ab}$  solve an ODE of just order 5. In other words, a linear combination of the  $W_{ab}$  vanishes, or equivalently there exists a constant, non-degenerate, antisymmetric,  $4 \times 4$ matrix  $M_{ab}$  (unique up to overall normalization) such that

<span id="page-52-2"></span>
$$
M_{ab} y_a y'_b = 0. \t\t(4.17)
$$

We can find a constant matrix  $S \in GL(4,\mathbb{C})$  such that  $S^tMS = \Omega$ , with  $\Omega$  the standard symplectic  $4 \times 4$  matrix. Define

<span id="page-52-3"></span>
$$
\Pi_a \stackrel{\text{def}}{=} (S^{-1})_a^{\ b} y_b \tag{4.18}
$$

 $Eq.(4.17)$  $Eq.(4.17)$  becomes

<span id="page-52-4"></span>
$$
(\Pi^t \Omega \partial_z \Pi) dz = 0, \tag{4.19}
$$

which is the Legendre condition [\(4.9\)](#page-50-0). We see that the Legendre condition is an automatic consequence of having the "right" algebraic monodromy/differential Galois group. More precisely here there are two ingredients: the "right" differential Galois group, and the choice of the "good" solution for the 4 periods  $\Pi_a$ . Since all solutions are linear combinations of any one basis of solutions, the "good" ones are special linear combinations and the matrix  $S_a{}^b$  specifies the one we need to keep, cf. eq. [\(4.18\)](#page-52-3). Since  $\Pi$  are the periods of the so-called Seiberg-Witten differential, this shows that the monodromy representation also fixes which linear combination of the differentials on the fiber is the Seiberg-Witten differential. Note that we have still some freedom: making  $S \to t S g$  with  $g \in Sp(4,\mathbb{C})$  and  $t \in \mathbb{C}^\times$  we get a new Π which still satisfy eq.[\(4.19\)](#page-52-4). The overall scale freedom just reflects the fact that we consider the image of the periods in projective space. The freedom by a symplectic rotation is more interesting: for the moment Π are periods defined over some complex sympletic basis of  $H_1(\mathscr{A}_u, \mathbb{Z}) \otimes_{\mathbb{Z}} \mathbb{C}$ . Using  $g \in Sp(4, \mathbb{C})$  we transform this basis into an integral symplectic basis of 1-cycles (which exists since, by assumption, our ODE arises from an actual family of Abelian surfaces) characterized by the fact that the monodromy elements becomes integral matrices in  $Sp(4, \mathbb{Z})$  when written in terms of such a basis. Then  $\Pi \in \mathbb{P}^{2r-1}$  becomes unique

<span id="page-52-0"></span><sup>&</sup>lt;sup>39</sup> The inverse statement holds only modulo finite groups (equivalently modulo finite covers).

modulo the action of the Siegel modular group  $Sp(2r, \mathbb{Z})$ . Finally the condition  $\text{Im } \tau_{ij} > 0$ holds since the ODE arises from a family of Abelian varieties. We conclude

**Fact 13.** In the classical case  $r = 2$  all non-isotrivial, irreducible families of Abelian surfaces  $\mathscr{A} \to \mathbb{P}^1 \setminus \{z_1, \ldots, z_s\}$ , having a monodromy representation satisfying the appropriate conditions, admit an essentially unique SW differential  $\lambda$  making them into  $\mathbb{C}^{\times}$ -isoinvariant rank-2 special geometries.

In the "appropriate conditions" we include the properties which insures that the special geometry is regular at the special points  $z_i$ , i.e. along the discriminant components and the enhanced divisors. To insure these regularity conditions requires some easy extra checks (see below).

The message is that all that matters is the monodromy representation  $\mu$ . If the monodromy representation has the right properties, the integrability conditions will follow. This is the same statement as in the Calabi-Yau case, but of course, the monodromy representations and other important aspects are rather different in the two cases. From an abstract viewpoint we may state that special geometry is a topic in Representation Theory (RT), and the main source of difficulty is that the representation is defined over  $\mathbb{Z}$  (or another ring of integers) which makes the analysis harder and subtler than plain RT over (say) C.

The corollary is that to classify all irreducible rank-2 geometries (which are not isotrivial) it suffices to classify all non-isotrivial families of Abelian surfaces with the appropriate properties. We expect the higher rank case to work morally in the same way.

### 4.2.1 Recovering the special coordinates  $(r = 2)$

In terms of the solution  $\Pi(z)$  to the Picard-Fuch equation the SW periods on  $\mathscr{C} \simeq \mathbb{C}^2$  read

<span id="page-53-0"></span>
$$
\Pi_{\rm SW}(u_1, u_2) = \begin{cases} h(u_1, u_2) \, \Pi(u_1^{d_2}/u_2^{d_1}) & \varkappa = 1\\ \sqrt{h(u_1, u_2)^2} \, \Pi(u_1^{d_2}/u_2^{d_1}) & \varkappa = 2 \end{cases} \tag{4.20}
$$

where  $h(u_1, u_2)$  (resp.  $h(u_1, u_2)^2$ ) is an homogeneous element of the field  $\mathbb{C}(\mathscr{C})$  of dimension 1 (resp. 2). Note that in the  $\varkappa = 2$  case the periods are defined up to sign.

In order for a solution Π of the Picard-Fuchs equation to represent a regular special geometry we need in particular that the SW periods  $\Pi_{SW}$  along the special divisors  $S_i$ (whose leading behavior we can read from the local solutions defined by the conjugacy class  $C_i$ ) agree asymptotically with the local models of the singular fibers of type  $C_i$  which we reviewed in §. [2.9.2.](#page-33-0) Later we shall use eq.  $(4.20)$  to check the regularity of the candidate geometries along the special loci.

## 4.3 "Stratification" vs. Picard-Fuchs systems

Let us summarize the logic of our work. The starting point is that specifying a special geometry is the same as giving a Gauss-Manin connection  $\nabla$  on the projective Coulomb

branch  $\mathscr P$  with the appropriate properties (monodromy in  $Sp(2r, \mathbb{Z})$ , a "good" solution with the Legendre property, local regularity, and so on). The idea is to construct the global connection  $\nabla$  from the local data around the special divisors  $\mathcal{S}_a = (p_a)$ . The obvious local datum is the conjugacy class  $C_a$  of the local monodromy along  $S_a$ . The class of the local monodromy is just the class of the exponential of the residue matrix  $A_a(u)|_{\mathcal{S}_a}$  of the connection  $\nabla = d + \sum_a A_a(u) d \log p_a(u)$ :

$$
C_a = \left[ \exp(-2\pi i A_a(u)|_{\mathcal{S}_a}) \right]. \tag{4.21}
$$

It is a basic fact that the integrability condition  $\nabla^2 = 0$  implies that the class  $C_a$  is independent of the point along the special divisor  $S_a$  [\[100\]](#page-154-5). The monodromy representation (which fully determines the geometry) is generated by the local monodromies  $\mu_a$  around  $\mathcal{S}_a$  whose conjugacy classes is a datum of the inverse problem.

"Stratification" yields another strong constraint on  $\nabla$ . The restriction of the rank 2r Gauss-Manin connection on a special divisor (or rather to a special divisor which originates as a irreducible component of the discriminant) should reproduce the rank  $(2r-2)$  Gauss-Manin connection of the special geometry along the strata. Since we are working recursively on  $r$ , this GM connection is supposed to be know when running the inverse problem program (but it may be reducible or trivial, corresponding to an isotrivial geometry along the discriminant component).

The technical details of the procedure to restrict the GM PDE on a special divisor are described, for instance, in §.12.2 of the book [\[100\]](#page-154-5). For simplicity we consider the case when  $\mathcal{S}_a$  is a non-enhanced discriminant (we can always reduce to this case by a change of base and the elimination of apparent singularities). In this case the local monodromy  $\mu_a$  leaves invariant a subspace (defined over  $\mathbb{Q}$ )  $V_{\mathbb{Q}} \subset \mathbb{Q}^{2r}$  of dimension  $2r - 2$ . The solutions taking value in  $V_{\mathbb{Q}}$  have trivial monodromy around the divisor, so can be taken to be holomorphic. The connection restricted to  $\mathcal{S}_a$  and  $V_{\mathbb{Q}}$  should reproduce one of the allowed GM connection in one less rank, that we have already classified in the previous step of our recursive procedure. This fixes a  $(2r-2) \times (2r-2)$  diagonal block of the residue matrix  $A_a(u)$ . The complementary 2 × 2 block should be the  $(2\pi i)^{-1}$  times the log of the matrix giving the known Kodaira monodromy around the discriminant (see §. [2.9\)](#page-32-0), up to conjugacy. The restriction relations for the GM connection following from "stratification" is a defining property of special geometry.

# <span id="page-54-0"></span>5 Rigidity I: the conformal manifold

To get necessary/sufficient conditions for the solvability of the inverse problem, and to construct explicit solutions in lower rank, we use *rigidity theorems*. Three different notions of rigidity are relevant for our story:

(a) rigidity as a  $\mathbb{C}^{\times}$ -isotrivial special geometry;

- (b) rigidity of the underlying family of Abelian varieties over the quasi-projective manifold  $\mathscr{\tilde{P}}$  in the sense of Arakelov [\[57\]](#page-151-10), Faltings [\[58\]](#page-152-1), Saito [\[59\]](#page-152-2), and Peters [\[60–](#page-152-3)[62\]](#page-152-4);
- (c) rigidity of the underlying monodromy representation in the sense of Deligne [\[101\]](#page-154-6), Simpson [\[63–](#page-152-5)[65\]](#page-152-6), and Katz [\[102\]](#page-154-7).

A fourth rigidity notion, the VHS one in the sense of Griffiths [\[34–](#page-150-6)[37\]](#page-150-0) and Schmidt [\[103\]](#page-154-8), will not be discussed here since in the present setting it is subsumed in (b).

# 5.1 Rigidity as a  $\mathbb{C}^{\times}$ -isoinvariant special geometry

The first notion is rigidity of the  $\mathbb{C}^{\times}$ -isoinvariant special geometry in its full glory.

**Definition 8.** A family of  $\mathbb{C}^{\times}$ -isoinvariant special geometries over the Coulomb branch  $\mathscr{C}$ parametrized by the complex manifold  $\mathscr B$  is a holomorphic fibration

<span id="page-55-1"></span>
$$
\varpi: \mathscr{Z} \to \mathscr{C} \times \mathscr{B}, \qquad \mathscr{B} \text{ connected}
$$
\n
$$
(5.1)
$$

such that for all  $b \in \mathcal{B}$  the restricted fibration

$$
\varpi|_{\varpi^{-1}(\mathscr{C}\times\{b\})}\colon \mathscr{Z}|_{\varpi^{-1}(\mathscr{C}\times\{b\})}\to \mathscr{C}
$$
\n
$$
(5.2)
$$

is a C <sup>×</sup>-isoinvariant special geometry.

We may assume the manifold  $\mathscr{B}$  to be quasi-projective with no loss. Then, as always when working with families [\[104\]](#page-154-9), we may (or rather should) introduce a moduli functor.

**Definition 9.** The *(fine)* conformal manifold  $\mathcal{M}$  of a  $\mathbb{C}^{\times}$ -isoinvariant special geometry is the scheme which represents the moduli functor. Typically  $\mathscr M$  does not exist as a *fine* moduli space and one must content himself with a *coarse* conformal manifold. This happens (for instance) when points in the conformal manifold are fixed by finite subgroups of the S-duality group. Alternatively on considers a fine extended conformal manifold which parametrizes suitable pairs (SCFT, duality frames/ $\sim$ ). $^{40}$  $^{40}$  $^{40}$ 

Roughly speaking,  $M$  parametrizes continuous deformations of the geometry which keep it  $\mathbb{C}^{\times}$ -isoinvariant and special. The dimensions  $\{\Delta_1, \ldots, \Delta_r\}$ , being rational, are constant along the conformal manifold  $\mathcal{M}$ , as is the monodromy representation.

**Definition 10.** A  $\mathbb{C}^{\times}$ -isoinvariant special geometry is *rigid* if its conformal manifold M is zero dimensional (i.e. a point).

Fact 14. One has dim  $\mathscr{M} = \dim \mathscr{R}_{\Delta=2}$ .

<span id="page-55-0"></span><sup>&</sup>lt;sup>40</sup> A typical example (SU(2) with  $N_f = 4$ ) works as follows: the coarse conformal manifold is  $H/SL(2, \mathbb{Z})$ while taking ∼ to be the equivalence relation that two frames are identified if they differ by a rotation in  $\Gamma(2) \subset SL(2, \mathbb{Z})$  leads to a fine conformal manifold for the pair, namely  $H/\Gamma(2)$ .

*Proof.* Using the periods  $\Pi(u, m)$   $(u \in \mathscr{C}, m \in \mathscr{M})$  we construct the intrinsic linear map<sup>[41](#page-56-0)</sup>

$$
\phi \colon T_m \mathscr{M} \to \mathscr{R}_{\Delta=2}, \qquad \phi \colon \partial_{m^a} \mapsto \frac{1}{2} \Pi(u, m)^t \Omega \partial_{m^a} \Pi(u, m) \tag{5.3}
$$

A priori the function in the RHS is only defined in  $\mathscr{C}$ , but it is easy to check<sup>[42](#page-56-1)</sup> that it extends everywhere on C and hence is an element of the chiral ring  $\mathscr R$  of scaling dimension  $\Delta = 2$ . We have to show that the map  $\phi$  is an isomorphism. Consider a small coordinate patch  $U \subset \mathscr{C}$  and let  $\mathcal{F}(a,m)_U$  be the local pre-potential in U as a function of the conformal parameters  $m^a$ . Locally in U one has

$$
\frac{1}{2}\Pi(u,m)^t\Omega\partial_{m^a}\Pi(u,m) = \frac{1}{2}a^i\partial_{m^a}\partial_{a^i}\mathcal{F}_U = \partial_{m^a}\left(\frac{1}{2}a^i\partial_{a^i}\mathcal{F}_U\right) = \partial_{m^a}\mathcal{F}_U,\tag{5.4}
$$

which shows that  $\phi$  is injective since  $\partial_{m^a}F_U=0$  in the open set U implies that the infinitesimal deformation is trivial by uniqueness of the analytic continuation. On the other hand, consider an open cover  $\mathscr{C} = \cup_{\alpha} U_{\alpha}$  and locally modify the pre-potential  $\mathcal{F}_{\alpha}$  in each  $U_{\alpha}$  as

$$
\mathcal{F}_{\alpha} \to \mathcal{F}_{\alpha} + \epsilon h + O(\epsilon^2) \quad \text{for } h \in \mathcal{R}_{\Delta=2}.
$$
 (5.5)

It is easy to check that in the intersection  $U_{\alpha} \cap U_{\beta}$  the deformed local special geometries on  $U_{\alpha}$  and  $U_{\beta}$  agree to the first order in  $\epsilon$ . Hence the infinitesimal deformation makes sense at the global level on  $\mathscr C$  and defines a non-trivial infinitesimally deformed special geometry which corresponds to an element of the tangent space  $T_m\mathscr{M}$ . Hence  $\phi$  is also surjective.  $\Box$ 

Recall our standing assumption that  $\Delta_i > 1$  for all i. In this case:

**Fact 15.** When the Coulomb branch  $\mathscr C$  is smooth the dimension of the conformal manifold  $M$  is simply the multiplicity of 2 as a Coulomb dimension. In particular when 2 is not a Coulomb dimension the special geometry is rigid.

◯ Caveat 2. When the Coulomb branch is non-smooth we may have dim  $\mathcal{M}$  > dim  $\mathcal{C}$   $\equiv$  r.

## 5.2 Non-rigid geometries: the Folk-theorem

Everybody believes that the following is true:

Folk-theorem. A non-rigid, smooth, indecomposable  $\mathbb{C}^{\times}$ -isoinvariant special geometry with  $\dim_{\mathbb{C}} \mathscr{M} = s$  is the Seiberg-Witten geometry of a coupled  $\mathcal{N} = 2$  SYM + "matter" model, where the gauge group  $G$  has s simple factors, and the "matter" is a  $SCFT$  – which may be non-Lagrangian – such that the YM beta-functions vanish for all factor gauge groups.<sup>[43](#page-56-2)</sup>

<span id="page-56-1"></span><span id="page-56-0"></span><sup>&</sup>lt;sup>41</sup> Here  $m^a$  are local coordinates around the point  $m \in \mathcal{M}$ .

<sup>&</sup>lt;sup>42</sup> Use the classification of the possible singular fibers along the discriminant components and check the statement on a case by case basis.

<span id="page-56-2"></span> $^{43}$  This is equivalent to requiring that the central charge of the  $G_i$  current algebra of the "matter" SCFT has the correct value  $h(G_i)^\vee$  for all simple factor  $G_i$  of  $G$ .

The matter sector decouples into a set of SCFTs (possibly free) whose special geometries are rigid.

The main relevance for us of the **Folk-theorem** is that it says that all non-rigid special geometries can be constructed by "gauging" global symmetries of the rigid ones; therefore for the classification purpose we may (and do!) assume the geometry under consideration is rigid. This restriction will simplify our discussion. In this subsection we spend some words about the Folk-theorem for the benefit of the skeptical reader, sketching a proof of it in some special situations, and giving some applications to the construction of Seiberg-Witten "curves" including ones that were not known before.

The **Folk-theorem** is known to be true for *isotrivial* geometries with a smooth Coulomb branch [\[22\]](#page-150-2). In this subsection we present some general considerations which apply also to the non-isotrivial case.

 $\hat{\mathcal{L}}$  Caveat 3. The Folk-theorem is *false* in the non-smooth case.

### 5.2.1 Folk-theorem vs. classical special geometries

The precise statement of the Folk-theorem is that (under the assumptions) the coarse conformal manifold is a quasi-projective space  $\mathscr{M} = \overline{\mathscr{M}} \setminus \sum_i W_i$  while along some divisor at infinity  $W_{i_0}$  the gauge coupling gets parametrically small almost everywhere in  $\mathscr{C}$ . This means that (in an appropriate  $Sp(2r, \mathbb{Z})$  frame)  $r_G \leq r$  eigenvalues<sup>[44](#page-57-0)</sup> of the matrix Im  $\tau_{ij}(u)$ go to infinity for "most" points  $u \in \mathscr{C}$ . Physically we expect that the SCFT dynamics in this low-couplings regime has a weakly-coupled Lagrangian description,<sup>[45](#page-57-1)</sup> and moreover that the semiclassical treatment of the gauge sector gets (asymptotically) exact near  $W_i$ . Going from the dynamics of the SCFT to its special geometry, one is led to think that the special geometry should also approach a semiclassical geometry. This is morally right, but the actual story is subtler. We start our discussion of the **Folk-theorem** by briefly reviewing the geometry of a SCFT with a weakly-coupled Lagrangian description (hence with a Yang-Mills subsector).

The special geometry of an  $\mathcal{N}=2$  model containing a Yang-Mills sector with the gauge group  $G = G_1 \times \cdots \times G_s$  ( $G_a$  simple Lie groups) has a Coulomb branch of the form<sup>[46](#page-57-2)</sup>

<span id="page-57-4"></span>
$$
\mathscr{C}_{G_1} \times \mathscr{C}_{G_2} \times \cdots \mathscr{C}_{G_s} \times \mathscr{C}_{\text{matter}}^{\text{inter}} \tag{5.6}
$$

where<sup>[47](#page-57-3)</sup>

$$
\mathcal{C}_{G_a} = \mathbb{C}^{r_{G_a}} / \text{Weyl}(G_a) = \text{Spec } \mathcal{R}_{G_a} \simeq \mathbb{C}^{r_{G_a}} \tag{5.7}
$$

<span id="page-57-0"></span><sup>&</sup>lt;sup>44</sup> The eigenvalues of Im  $\tau_{ij}(u)$  may be identified with  $4\pi/g(u)^2_i$  where  $g(u)_i$  are the effective gauge couplings in vacuum u.

<span id="page-57-1"></span><sup>45</sup> In this description the "matter" is treated as a "black box": only the gauge degrees of freedom are described by weakly coupled fields.

<span id="page-57-2"></span><sup>&</sup>lt;sup>46</sup> The sector called "matter" may contain further Yang-Mills sectors.

<span id="page-57-3"></span> $^{47}$  As always  $\mathsf{Weyl}(G_a)$  is the Weyl group of the Lie group  $G_a.$ 

with  $r_{G_a}$  the rank of the simple Lie group  $G_a$  while  $\mathscr{R}_{G_a} \simeq \mathbb{C}[x_1,\ldots,x_{r_{G_a}}]^{\text{Weyl}(G_a)}$  is the graded polynomial ring in  $r_{G_a}$  variables  $u_i$  whose degrees are the degrees of  $G_a$ .

<span id="page-58-2"></span>**Definition 11.** A classical special geometry with gauge group  $G = G_1 \times \cdots \times G_s$  is a  $\mathbb{C}^{\times}$ isoinvariant special geometry over a Coulomb branch of the form [\(5.6\)](#page-57-4) whose general fiber is an Abelian variety containing a *fixed* Abelian subvariety  $A_G$  of rank  $r_G \equiv \sum_a r_{G_a}$  whose automorphism group  $\mathsf{Aut}(A_G)$  contains  $\mathsf{Weyl}(G) \equiv \prod_a \mathsf{Weyl}(G_a)$ .

In particular when  $r = r_G$ , that is, when "matter" is a system of free hypermultiplets, a classical geometry is automatically isotrivial. In this case the Abelian varieties  $A_G$  are as described in [\[106,](#page-154-10) [107\]](#page-154-11) (see also [\[22\]](#page-150-2)). In rank-2 all classical geometries are isotrivial, since the matter has at most rank-1 and hence is automatically isotrivial. The following statement is very well known:[48](#page-58-0)

**Fact 16.** When our  $\mathcal{N} = 2$  SCFT is actually  $\mathcal{N} = 4$  invariant, the non-renormalization theorems of extended supersymmetry imply that the special geometry is classical.

When  $r_G \leq r$  eigenvalues of Im  $\tau_{ij}$  go to infinity the Abelian generic fiber of the classical geometry  $\mathscr X$  degenerates.  $\mathscr X_u$  (u generic) remains a commutative algebraic group (hence still smooth) but is no longer compact:  $\mathscr{X}_u$  is a fibration over an Abelian variety of dimension  $(r - r_G)$  with fiber the algebraic torus  $(\mathbb{C}^\times)^{r_G}$  (again the Chevalley-Barsotti theorem<sup>[49](#page-58-1)</sup>). In particular in the (much simpler) case where  $r = r_G$ , the classical special geometry reduces in this weak-coupling limit to the classical geometry of the Coulomb branch of the 3d  $\mathcal{N}=4$ model with the same gauge group and matter content. However in 3d QFT quantum effects modify the geometry of the Coulomb branch, so that the *quantum* 3d  $\mathcal{N} = 4$  Coulomb branch is not the classical one: see [\[108\]](#page-154-12) and references therein for a rigorous mathematical definition of the 3d Coulomb branch. It is reasonable to expect that on the divisor  $W_i \subset \overline{\mathscr{M}}$  at infinity in the conformal manifold the special geometry reduces (in this simpler situation) to the quantum 3d one. Since the "proof" of the **Folk-theorem** asserts that "almost everywhere" on the Coulomb branch we get the semiclassical situation, we deduce two geometric properties of the 3d  $\mathcal{N} = 4$  Coulomb branch:

**Fact 17.** (1) the 3d  $\mathcal{N} = 4$  Coulomb branch of a gauge theory coupled to hypermultplets is a Lagrangian fibration over the 4d Coulomb branch of the corresponding model with fibers the dual of the maximal torus in the gauge group G, i.e. the fiber  $\simeq (\mathbb{C}^{\times})^{r_{G}}$ . (2) the special geometry on the divisor  $W_i \subset \overline{\mathcal{M}}$  is birational to the classical 3d  $\mathcal{N}=4$  Coulomb branch.

This statement is in fact a math theorem, see eq.(3.17), **Proposition 5.19**, and **Corollary** 5.21 of [\[108\]](#page-154-12). This Fact prompts the

<span id="page-58-0"></span><sup>&</sup>lt;sup>48</sup> A mathematical proof goes as follows.  $\mathcal{N} = 4$  SUSY implies that the scalar metric is flat [\[33\]](#page-150-7), hence the special geometry is isotrivial for the reasons explained in §. [2.10.4.](#page-41-0)

<span id="page-58-1"></span><sup>49</sup> A more general and deeper treatment will be presented elsewhere.

<span id="page-59-0"></span>**Definition 12.** By a *weakly-coupled geometry* we mean a *generalized* special geometry  $\pi: \mathscr{W} \to \mathscr{C}$  with Lagrangian fibers and section whose general fiber  $\mathscr{W}_u$  is a commutative algebraic group which is an extension of an Abelian variety  $A_u$  (possibly of dimension 0)

$$
1 \to T^{\vee} \to \mathscr{W}_u \to A_u \to 0 \tag{5.8}
$$

by the dual  $T^{\vee} \simeq (\mathbb{C}^{\times})^{r_G}$  of the maximal torus of a (compact, semisimple) Lie group G which is the product of s simple factors. The set of degrees of the group  $G$  are contained, with the appropriate multiplicities, in the r-tuple  $\{\Delta_i\}$  of Coulomb dimensions.

Remark 5. In Remark [2](#page-17-0) we noticed that the degeneration of the fiber as we approach the origin in  $\mathscr C$  is always additive. On the contrary, the degeneration at infinity in  $\mathscr M$  is always multiplicative.

Remark 6. The theory of generalized special geometries, their number-theoretical flavor, and their applications will be discussed elsewhere [\[109\]](#page-154-13).

In view of these remarks, we restate the **Folk-theorem** as follows

Folk-theorem. A smooth indecomposable  $\mathbb{C}^{\times}$ -isoinvariant special geometry where 2 has multiplicity s as a Coulomb dimension has a (coarse) conformal manifold  $\mathcal M$  which is a quasiprojective variety i.e. can be compactified in a projective variety  $\overline{\mathscr{M}}$  while  $\mathscr{M} = \overline{\mathscr{M}} \setminus W$ for a simple divisor  $W = \sum_i W_i$  at infinity. There is (at least one) component  $W_i$  of  $W$ such that as we go close to it the special geometry asymptotically approaches a classical (weakly-coupled) geometry for the appropriate Lie group G.

Remark 7. The Folk-theorem implicitly contains the idea that the conformal manifold  $\mathscr M$  has a *canonical* compactification. In the known examples this is true. For instance, for class S theories [\[110,](#page-154-14) [111\]](#page-154-15) (which form a sizable portion of all  $\mathcal{N}=2$  SCFTs) the conformal manifold is the moduli space  $\mathcal{M}_{g,n}$  of genus g curves with n punctures which has the canonical Deligne-Mumford compactification [\[112,](#page-155-0) [113\]](#page-155-1).

Remark 8. Typically the divisor W has several irreducible components at which we have inequivalent weakly-coupled geometries. When this happens we say that there is duality of the Argyres-Seiberg type [\[114\]](#page-155-2) between the corresponding weakly-coupled QFTs. The basic example  $[114]$  is  $SU(3)$  SYM with 6 fundamentals whose conformal manifold has a second divisor at infinity where it describes  $SU(2)$  coupled to "matter" consisting of one fundamental hyper plus one Minahan-Nemeshanski interacting SCFT of type  $E_6$  [\[115,](#page-155-3) [116\]](#page-155-4).

**Disclaimer.** Proving (or disproving) the **Folk-theorem** is not a purpose of this paper. We limit ourselves to sketch a proof in the special case of rank-2 where the necessary mathematical analysis has already been done in the math literature for different purposes. As a preparation we consider a slightly more general situation which is of independent interest.

### 5.2.2 A special class of dimension r-tuples  $\{\Delta_i\}$

We focus on the following:

**Special situation.** A non-isotrivial  $\mathbb{C}^{\times}$ -isoinvariant special geometry with r even, having a r-tuple of dimensions  $\{\Delta_1, \ldots, \Delta_r\}$  which contains 2 with multiplicity 1 and in addition a dimension  $\Delta_{i_1}$  which is new in rank  $r/2$ , while the axis spanned by  $u_{i_1}$  (called the "good" axis henceforth) is not contained in the discriminant.

**Remark 9.** This is not a too "special" situation. In particular all  $r = 2$  models with precisely one dimension 2 have this form with the following **qualification** (see §.[6.1\)](#page-69-0): as we move in the conformal manifold  $\mathscr M$  the discriminant divisor  $\mathcal D \subset \mathbb C^2$  will also move. In particular there is a point at infinity in  $\mathscr M$  where the "good" axis ends up being inside the discriminant: our arguments here apply only as long as we keep away from that particular corner of the conformal manifold.

**Remark 10.** The **Special situation** covers all Lagrangian  $\mathcal{N} = 2$  SCFT with a simple gauge group G of rank r which has a "good"<sup>[50](#page-60-0)</sup> degree d such that<sup>[51](#page-60-1)</sup>  $\phi(d) = r$ . Their list is

<span id="page-60-3"></span>
$$
SU(p) \text{ (p prime)}, \quad SO(2^k+1), \quad Sp(2^{k+1}), \quad G_2, \quad F_4, \quad E_6, \quad E_8 \tag{5.9}
$$

which contains all simple Lie groups of rank-2.

In the **Special situation**  $\mathcal{M}$  is a non-compact complex curve which we may write as a projective curve  $\overline{\mathscr{M}}$  minus a few points ("cusps at infinity"). In the known examples  $\mathscr{M}$ is a (non-compact) modular curve [\[117\]](#page-155-5): one expects this to be true in general when 2 has multiplicity 1. When  $r = 2$  one proves that  $\mathscr M$  is a non-compact modular curve of genus zero (see below).

We fix a point u on the "good" axis with  $u \neq 0$ . By restricting [\(5.1\)](#page-55-1) to u we get a one-parameter family of Abelian varieties over the conformal manifold  $52$ 

<span id="page-60-4"></span>
$$
\mathscr{Z}_u \to \mathscr{M} \tag{5.10}
$$

whose fibers are principally polarized Abelian varieties of dimension  $r$  which are left invariant by the (polarized) automorphism  $\xi \equiv \exp(2\pi i \mathcal{E}/\Delta_{i_1})$  which acts on the fibers via the analytic representation

<span id="page-60-5"></span>
$$
\sigma(\xi) = \text{diag}(e^{2\pi i (1 - \Delta_1)/\Delta_{i_1}}, \dots, e^{2\pi i (1 - \Delta_r)/\Delta_{i_1}}). \tag{5.11}
$$

<span id="page-60-0"></span><sup>&</sup>lt;sup>50</sup> For the groups in the list [\(5.9\)](#page-60-3) the Coxeter number h is always a "good" degree except for  $E_6$  which has the single "good" degree 9.  $F_4$  has 2 "good" degrees 8 and 12, while  $E_8$  has 3 "good" degrees 20, 24 and 30. <sup>51</sup> Throughout this paper  $\phi: \mathbb{N} \to \mathbb{N}$  is Euler's totient (multiplicative) function.

<span id="page-60-2"></span><span id="page-60-1"></span>

 $52$  Here we are oversimplifying since  $\mathcal M$  is typically coarse. Implicitly we add some extra structure to the geometry to make some finite cover of  $\mathscr M$  into a fine conformal manifold. This is implicitly done all the time in the physical literature about dualities of  $\mathcal{N} = 2$  SCFTs.

Since 2 is a dimension, the family  $(5.10)$  is non-isotrivial while both conjugate roots

$$
\exp(2\pi i/\Delta_{i_1}) \quad \text{and} \quad \exp(-2\pi i/\Delta_{i_1}) \tag{5.12}
$$

are eigenvalues of  $\sigma(\xi)$  and hence both must have multiplicity at least 2 as eigenvalues of the rational representation  $\chi(\xi)$ . By assumption the algebraic number  $\exp(2\pi i/\Delta_{i_1})$  has degree r, and hence

$$
\det[z - \chi(\xi)] = \Phi_m(z)^2 \tag{5.13}
$$

where  $\Phi_m(z)$  is a cyclotomic polynomial with  $\phi(m) = r$ . Then  $\det[z - \sigma(\xi)] = \Phi_m(z)$  and the r phases in eq.[\(5.11\)](#page-60-5) are precisely the primitive  $m$ -th roots of 1.

The classification of all **Special situation** geometries is then reduced to the construction of all non-isotrivial one-parameter families of principally polarized Abelian  $r$ -folds with a (polarization preserving)  $\mathbb{Z}_m$  automorphism  $\xi$  (with  $\phi(m) = r$ ) having the analytic repre-sentation in eq.[\(5.11\)](#page-60-5). We **Claim** that a sufficient (but not necessary!) condition for the validity of the Folk-theorem in the Special situation is that all one-parameter families of Abelian varietes with the stated properties have an automorphism group containing a non-trivial Weyl group of the proper kind. Next we show that this is the case in rank 2.

### 5.2.3 Folk-theorem: proof in rank 2

We return to the Folk-theorem in rank 2. We focus on the case with dimensions  $\{2, \Delta\}$ where  $\Delta$  is a rank-1 dimension  $\neq 1, 2$ . Since the discriminant locus  $\mathcal{D}_q \subset \mathscr{C}$  will move with the point  $q \in \mathcal{M}$ , at generic points in  $\mathcal{M}$  it stays away from the "good" axis of dimension  $\Delta$ and we are exactly in the **Special situation**. In particular  $m = 3, 4, 6$ , so either  $\Delta = m$  or  $\Delta = m/(m-1)$ , and the fibers  $X_u$  along the axis of dimension  $\Delta$  have  $\mathbb{Z}_m \subset \text{Aut}(X_u)$ .

**Proposition 3** (Cf. Theorem 4.8 of  $[118]^{53}$  $[118]^{53}$  $[118]^{53}$  $[118]^{53}$ ). The one-dimensional non-isotrivial families of principally polarized Abelian surfaces  $A_{\tau}$  with a non-trivial automorphism group  $\mathfrak A$  are:

Abelian surface $A_{\tau}$	$\mathfrak A$	normal subgroup	$\tau_{ij}$	analytic representation	Jacobian of $q=2$ curve
$E_{\tau} \times E_{\tau}$	$D_8\equiv \mathsf{Weyl}(C_2)$	$\mathbb{Z}_4$		$\begin{pmatrix} 7 & 0 \\ 0 & 7 \end{pmatrix} \qquad \left\langle \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right\rangle$	
	$(E_{\tau} \times E_{\tau})/\mathbb{Z}_2^2$ $D_8 \equiv \text{Weyl}(C_2)$				$\mathbb{Z}_4$ $\left(\begin{array}{cc} \tau^{-1/2} \\ 1/2 & \tau \end{array}\right)$ $\left\langle \left(\begin{array}{cc} 0 & 1 \\ -1 & 0 \end{array}\right), \left(\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array}\right) \right\rangle$ $y^2 = x(x^4 + \alpha x^2 + 1)$
	$(E_{\tau} \times E_{\tau})/\mathbb{Z}_3$ $D_{12} \equiv \text{Weyl}(G_2)$				$\mathbb{Z}_6$ $\begin{pmatrix} \tau & \tau/2 \\ \tau/2 & \tau \end{pmatrix}$ $\langle \begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \rangle$ $y^2 = x^6 + \alpha x^3 + 1$

where  $E_{\tau}$  is the elliptic curve of period  $\tau \in H \equiv SL(2,\mathbb{R})/U(1)$ ,  $\alpha \equiv \alpha(\tau)$  is a certain

<span id="page-61-1"></span><span id="page-61-0"></span><sup>&</sup>lt;sup>53</sup> A slightly weaker result is in **Theorem 13.4.5** of [\[77\]](#page-153-8). For our purposes we need the strong version of the theorem.

complex coordinate on the curve  $\mathcal{M} = H/\Gamma$ ,  $\Gamma$  a suitable congruence subgroup of  $SL(2, \mathbb{Z})$ , and  $D_{2k}$  is the dihedral group with  $2k$  elements.

Note that modulo isogeny we have  $A_{\tau} \sim E_{\tau} \times E_{\tau}$  in all three cases (cf. footnote [53\)](#page-61-0).

From the **Proposition** we see that the invariance of the one-parameter family under the cyclic automorphism group  $\langle \xi \rangle \simeq \mathbb{Z}_m$  implies its invariance under a full Weyl group which is Weyl $(G_2)$   $\supset$  Weyl $(A_2)$ , Weyl $(C_2)$ , and Weyl $(G_2)$  for respectively  $m = 3, 4, 6$ . Note that the derivative of the gauge coupling  $\tau_{ij}$  along the "good" axis with respect to the coordinate  $\alpha$ on  $\mathscr{M}$  is<sup>[54](#page-62-0)</sup>

$$
\frac{\partial \tau_{ij}}{\partial \alpha}\Big|_{\text{``good''}} \propto \left(\frac{\text{invariant scalar product}}{C_{ij} \text{ on the weight lattice}}\right). \tag{5.14}
$$

This is precisely the condition which we claimed above to be sufficient for the validity of the Folk-theorem. However we have still to explain why the *Claim* holds.

**Justification of the Claim for**  $r = 2$ . The first observation is that – for each choice of  $m \in \{3, 4, 6\}$  and family  $\{A_{\tau}\}_{{\tau \in M}}$  with  $\mathbb{Z}_m \subset \mathfrak{A}$  – we have a one-parameter family of isotrivial special geometries

$$
(A_{\tau} \times \mathbb{C}^2) / \text{Weyl}(G) \to \mathbb{C}^2 / \text{Weyl}(G), \qquad \tau \in \mathcal{M}
$$
 (5.15)

where  $G = A_2, C_2, G_2$  for respectively  $m = 3, 4, 6$ . The Weyl symmetry acts on the factor  $A_{\tau}$  via the embedding Weyl $(G) \subset \mathfrak{A}$  and on the factor  $\mathbb{C}^2$  via its degree 2 representation as a reflection group. In this way we get four rank-2 special geometries which are classical in the sense of **Definition [11](#page-58-2)**. For  $G = SU(3)$  or  $G_2$  we have just one classical geometry, but for  $Sp(4)$  we get two inequivalent classical geometries with  $A_{\tau} = (E_{\tau} \times E_{\tau})/\mathbb{Z}_{2}^{2}$  and  $A_{\tau} = E_{\tau} \times E_{\tau}$  respectively.

By the non-renormalization theorems of extended SUSY, the quantum special geometry of  $\mathcal{N} = 4$  SYM coincides with its semiclassical one. Hence for  $G = SU(3)$  or  $G_2$  the only semiclassical geometry must be the  $\mathcal{N} = 4$  one. In the case of  $Sp(4)$ , on the other hand, in addition to the  $\mathcal{N} = 4$  geometry, we have a second "exotic" semiclassical geometry. We stress than the "exotic" semiclassical geometry has the peculiarity that its fibers are not the Jacobians of genus 2 curves, i.e. there is no (smooth) Seiberg-Witten curve.

Roughly speaking the **Folk-theorem** asserts that in the limit  $\tau \to i\infty$  the geometry approaches a classical one; however the classical geometry itself is degenerating in the limit to a weakly-coupled one, so the actual statement is that as  $\tau \to i\infty$  we get a generalized geometry as in Definition [12](#page-59-0). Physically this is merely the statement that at extreme weak coupling the semiclassical approximation becomes exact. On the other hand Proposition **[3](#page-61-1)** states that in a sufficiently narrow  $\mathbb{C}^{\times}$ -stable neighborhood  $U_{\tau} \subset \mathscr{C}$  of the "good" axis the semiclassical description is a good approximation for all  $\tau$ . The size of the neighborhood  $U_{\tau}$ where the semiclassical approximation is reliable depends on the coupling  $\tau$ . Heuristically

<span id="page-62-0"></span><sup>54</sup> For the analogue statement in higher rank see [\[107\]](#page-154-11).

the *Claim* says that in the limit  $\tau \to i\infty$  the semiclassical domain  $U_{\tau} \subset \mathscr{C}$  will enlarge to cover "almost all" the Coulomb branch  $\mathscr C$  so that semiclassical physics in the "zero coupling" limit  $\tau \to i\infty$  gives the correct prediction for "almost all" observables and almost all vacual  $u \in \mathscr{C}$ . We formalize this heuristic picture as follow. Fix a huge constant  $\Lambda \gg 1$  and consider the  $\mathbb{C}^{\times}$ -stable open domain  $\mathscr{W}_{\Lambda,\tau} \subset \mathscr{C}$  consisting of Coulomb vacua where the gauge couplings are "very small" in the sense that all eigenvalues of the real matrix  $\text{Im } \tau_{ij}$  are larger than the fixed huge quantity  $\Lambda$ . The local physics at these vacua is essentially semiclassical, and we call (somehow abusively<sup>[55](#page-63-0)</sup>)  $\mathscr{W}_{\Lambda,\tau}$  the "semiclassical domain".

The *Claim* states that as  $\tau \to i\infty$  in *M* the size of the semiclassical domain  $\mathscr{W}_{\Lambda,\tau} \subset \mathscr{C}$ gets infinite for all Λ however large. That this is true follows from Ahlfors' lemma applied to the period map  $\tilde{p}: \tilde{\mathscr{Q}} \to Sp(2r,\mathbb{R})/U(r)$  where  $\mathscr{Q}$  is  $\mathscr{P}$  with all special points omitted but the one corresponding to the "good" axis. This makes sense since the "good" axis does not belong to the discriminant, so it has a local monodromy of finite order, and hence the covering period map  $\tilde{p}$  extends holomorphically over that special point.<sup>[56](#page-63-1)</sup> Since the geometry is not rigid,  $\widetilde{\mathscr{Q}}$  is the universal cover of  $\mathbb{P}^1$  less a number  $\geq 3$  of points (see §.[6.1](#page-69-0) below), hence  $\mathscr{Q}$  is the upper half-plane H with its canonical Poincaré distance function  $d_H(\cdot, \cdot)$ . We write  $B_H(*, R) \subset H$  for the Poincaré ball of radius R centered in a point  $* \in H$  which is (a lift of) the projection of the "good axis" on  $\mathscr P$  and set  $U(R) \equiv \varpi^{-1}(B_H(*, R)) \subset \mathscr C$ . The size of the semiclassical domain is the largest radius  $R_{\Lambda,\tau}$  such that  $U(R_{\Lambda,\tau}) \subset \mathscr{W}_{\Lambda,\tau}$ .

The precise version of the *Claim* is the statement that  $R_{\Lambda,\tau} \to \infty$  as  $\tau \to i\infty$  for all  $\Lambda$ . For  $\tau \gg \Lambda$  very large but finite  $\tilde{p}(*) = \tau C_{ij} + O(1)$  and  $\mathscr{W}_{\Lambda,\tau}$  contains the preimage  $U_{\Lambda,\tau}$  of the open ball  $B(\widetilde{p}(*), r_{\Lambda,\tau})_{\text{Siegel}}$  in Siegel space  $Sp(2r,\mathbb{R})/U(r)$  of radius

$$
r_{\Lambda,\tau} \equiv \inf_{M_{ij}} d_S(M_{ij}, \tilde{p}(*)),\tag{5.16}
$$

where  $d_S(\cdot, \cdot)$  is the canonical distance in the Siegel space and the infimum is taken over the set of complex symmetric matrices  $M_{ij}$  such that at least one of the eigenvalues of Im  $M_{ij}$  is  $\leq \Lambda$ :

$$
\mathscr{W}_{\Lambda,\tau} \supset U_{\Lambda,\tau} \equiv \{ x \in H \colon d_S(\widetilde{p}(x), \widetilde{p}(*)) < r_{\Lambda,\tau} \} \tag{5.17}
$$

and  $R_{\Lambda,\tau}$  is not smaller than the largest R such that  $U(R) \subset U_{\Lambda,\tau}$ . By Ahlfors' lemma<sup>[57](#page-63-2)</sup>  $U_{\Lambda,\tau}$ contains  $U(r_{\Lambda,\tau})$ , that is,

$$
R_{\Lambda,\tau} \ge r_{\Lambda,\tau} \tag{5.18}
$$

As  $\tau \to i\infty$ ,  $r_{\Lambda,\tau} \to \infty$  giving the *Claim*.

<span id="page-63-0"></span> $55$  The terminology may be justified as follows: the period map p extends holomorphically [\[119\]](#page-155-7) to the Barley-Borel compactification  $\overline{S}_r$  of the Siegel variety  $S_r \equiv Sp(2r,\mathbb{Z})\backslash Sp(2r,\mathbb{R})/U(r)$  [\[120\]](#page-155-8).  $\mathscr{W}_{\Lambda,\tau}$  gets mapped by p in a small neighborhood of a point at infinity in  $\overline{S}_r$  which shrinks to zero as  $\Lambda \to \infty$ , i.e. for large  $\tau$  the period map is "essentially constant" which is the trademark of a semiclassical geometry. However, this holds after replacing the period map by its Borel extension and should be taken with a pitch of salt.

<span id="page-63-1"></span><sup>56</sup> See e.g. Theorem 13.7.5 of [\[40\]](#page-151-3).

<span id="page-63-2"></span> $57$  See e.g. Corollary 13.7.2 in [\[40\]](#page-151-3).

### 5.2.4 Argyres-Seiberg dualities

The above discussion refers only to limits in the conformal manifold which are semiclassical of the special situation kind, i.e. which correspond to a weakly coupled gauge theory of rank 2. The boundary of the conformal manifold  $\mathcal{M}$  typically contains other components where a different kind of semiclassical geometry appears whose fixed Abelian variety  $A_{\tau}$  (in the sense of **Definition [11](#page-58-2)**) has rank  $\lt 2$ . This happens on a component of the boundary of  $\mathcal{M}$ where the axis parametrized by the coordinate  $v \equiv u_2$  with  $\Delta_2 = m$  lays in the discriminant. In this case  $\exp(2\pi i \mathcal{E}/m)$  acts by automorphisms of the singular fiber, hence of its Albanese variety which then has complex multiplication by  $\mathbb{Q}(e^{2\pi i/m})$  and hence is rigid. The onedimensional stratum with fiber this Albanese curve is the special geometry of a rank-1 SCFT with  $\Delta = m$ . Following [\[114\]](#page-155-2) we can describe this Albanese stratum using the genus-2 curve whose Jacobian yields the fiber over the v-axis. For, say,  $m = 3$  this curve is

$$
y^2 = x^6 + \alpha v x^3 + v^2. \tag{5.19}
$$

We rescale  $\alpha^{1/3}x = \tilde{x}/v$   $y = \tilde{y}/v$  getting

$$
\tilde{y}^2 = \frac{\tilde{x}^6}{\alpha^2 v^4} + \tilde{x}^3 + v^4
$$
\n(5.20)

which as  $\alpha \to \infty$  reduces to the elliptic curve for the  $E_6$  Minahan-Nemeshanski model. On the contrary, the divisor at infinity where the weakly coupled  $SU(3)$  SYM appears is  $\alpha^2 \to 4$ ; in this limit the hyperelliptic curve degenerates to [\[114\]](#page-155-2)

$$
y^2 = (x^3 + v)^2. \tag{5.21}
$$

### 5.2.5 Examples and checks

The above analysis gives predictions for the geometry of the rank-2 models with one dimension equal 2 which we wish to check against the known models. We stress that the same geometry may be shared by several inequivalent SCFTs. We write  $u, v$  for the Coulomb coordinate of dimension 2 and  $\Delta$ , respectively. The first consequence of the validity of the Folk-theorem in rank-2 is that  $\Delta$  is an integer  $\Delta \equiv m = 3, 4, 6$ . We consider each case in turn.

**Dimensions**  $\{2,3\}$  $\{2,3\}$  $\{2,3\}$ : generalities. From the table in Proposition 3 we see that all geometries with these dimensions have a genus-2 SW curve which when restricted to the locus  $u = 0$  is isomorphic to the one written in the last column of that table. Taking into account the scaling properties, we get a SW curve of the form

$$
y^{2} = x^{6} + \alpha(\tau) x^{3} v + v^{2} + z P(x, v; z)
$$
\n(5.22)

where  $P(x, v; z)$  is a polynomial in the variables x, v, of degree at most 4 in x, with coefficients in the ring of power series  $\mathbb{C}[[z]]$  in the local uniformizing parameter z of the normalization of the projective Coulomb branch  $\mathscr P$  at the special point associated with the "good" axis  $u = 0$ , that is,  $z = u^3/v^2$ . It satisfies the homogeneity condition

$$
P(t x, t^{3} v; z) = t^{4} P(x, v; z), \quad t \in \mathbb{C}.
$$
 (5.23)

The SW differential  $\lambda$  is uniquely determined by the analytic representation of  $\xi$  to be

<span id="page-65-2"></span>
$$
\lambda = v \frac{dx}{y} + u \frac{x \, dx}{y}.\tag{5.24}
$$

The simplest possibility is that  $z P(x, y; z)$  is a polynomial of the form  $u Q(x, y, u)$ ; by its scaling properties should have the form  $u Q(x, v, u) = c_1 x^4 u + c_2 x^2 u^2 + c_3 x u v + c_4 u^3$ . The coefficients  $c_i$  are further restricted by the condition that the discriminant has precisely 3 irreducible components. These predictions from the automorphism  $\xi$  are consistent with the known SW curve which yields the *non-isotrivial* geometry common to  $SU(3)$  with 6 fundamentals (#12 of [\[89\]](#page-153-5)) and  $SU(3)$  with one fundamental and one symmetric (SCFT  $\#38$  in [\[89\]](#page-153-5)) given by the equation [\[18\]](#page-149-4)

<span id="page-65-0"></span>
$$
y^{2} = x^{6} + \alpha(\tau) x^{3} (v - ux) + (v - ux)^{2}
$$
\n(5.25)

whose discriminant is proportional to

$$
v^{6} \left(27 v^{2} - 2(\alpha - \sqrt{\alpha^{2} - 4}) u^{3}\right) \left(27 v^{2} - 2(\alpha + \sqrt{\alpha^{2} - 4}) u^{3}\right),\tag{5.26}
$$

corresponding to two  $I_1$  component and one  $I_6$  component along the enhanced divisor  $v = 0$ .

Dimensions  $\{2,3\}$ : the isotrivial geometry. In addition to  $(5.25)$  there must be (at least) another SW curve corresponding to  $SU(3)$  with one adjoint. In this case we know from §. [2.10.4](#page-41-0) that the geometry is isotrivial (all smooth fibers are the Jacobian of the genus 2 curve in the table) and that the discriminant is irreducible (hence knotted). We also know that the monodromy around the discriminant is an automorphism of the fixed Abelian fiber of order 2 whose analytic representation has eigenvalues  $\{+1, -1\}$ . Since the cotangent space to the fiber (the dual of its Lie algebra) is spanned by the two forms

$$
\frac{dx}{y}, \qquad \frac{x \, dx}{y} \tag{5.27}
$$

this requires x to flip sign when we go around the unique component of the discriminant. Modulo trivial redefinitions, there is precisely one equation which satisfies all these requirements

<span id="page-65-1"></span>
$$
y^{2} = x^{6} + \alpha(\tau) x^{3} (v^{2} - u^{3})^{1/2} + (v^{2} - u^{3})
$$
\n(5.28)

whose discriminant is a bona fide algebraic divisor in the Coulomb branch

$$
729\left(\alpha(\tau)^2 - 4\right)^3 \left(v^2 - u^3\right)^5\tag{5.29}
$$

with only one irreducible component  $v^2 - u^3 = 0$ . The "odd-looking" square-root in [\(5.28\)](#page-65-1) is actually required to compensate for the flip of sign of  $x^3$  when we go around the discriminant. As always with isotrivial geometries, the SW curve is not a good way to describe the situation.

Dimensions  $\{2, 6\}$ : generalities. Again Proposition [3](#page-61-1) fixes the fiber along the "good" axis to be (up to isomorphism) the Jacobian of the same curve as in the {2, 3} case for all special geometries with dimensions  $\{2,3\}$  while the SW differential is still given by eq.[\(5.24\)](#page-65-2). The scaling argument yields a SW curve of the form

$$
v y^{2} = x^{6} + \alpha(\tau) x^{3} v^{2} + v^{4} + z P(x, v; z)
$$
\n(5.30)

where now  $z = u^3/v$ . There should be an isotrivial geometry corresponding to  $\mathcal{N} = 4$  with gauge group  $G_2$  and also a non-isotrivial one describing  $G_2$  SYM coupled to 4 fundamentals (model #49). The last model has 3 smooth discriminant components [\[89\]](#page-153-5).

For the non-isotrivial geometry analogy with the {2, 3} case (which is strictly related) suggests making the following replacement in the  $z = 0$  curve

<span id="page-66-0"></span>
$$
v \rightsquigarrow v - u \, x \tag{5.31}
$$

getting the SW curve

$$
v y^{2} = x^{6} + \alpha(\tau) x^{3} (v - u x)^{2} + (v - u x)^{4}
$$
 (5.32)

whose (properly normalized!!) discriminant is (up to an overall constant depending on  $\tau$ )

$$
v^{8}(v - \beta_{+}(\tau)u^{3})(v - \beta_{-}(\tau)u^{3})
$$
\n(5.33)

where  $\beta_{\pm}(\tau)$  are known functions of  $\alpha(\tau)$ . This expression says that we have two Kodaira fibers of Euler characteristic  $e = 1$ , i.e. of type  $I_1$ , while the third one has  $e = 8$ , i.e.  $I_8$ ,  $I_2^*$  or  $II^*$ . This is consistent with  $G_2$  with 4 fundamental which is known [\[89\]](#page-153-5) to have two discriminants of type  $I_1$  and one of type  $I_2^*$ .

**Dimensions**  $\{2, 6\}$ : the isotrivial geometry. The discriminant has two *smooth* components meeting non-transversally

$$
(v - 2u3)(v + 2u3) = 0
$$
\n(5.34)

Consider the curve

$$
v y^2 = x^6 + \alpha(\tau) x^3 (v^2 - 4u^6) + (v^2 - 4u^6)^2
$$
\n(5.35)



<span id="page-67-0"></span>Table 3: Known SCFTs with dimensions {2, 4} and their data from [\[18,](#page-149-4) [89\]](#page-153-5). "Types" refers to the Kodaira type of each irreducible component of the discriminant which are all smooth curves which intersect non-transversely. When the number of components is less than 3 the geometry is necessarily isotrivial (see  $\S$ , [2.10.3\)](#page-39-0). Models  $\#10$  and  $\#43$  share the same SW curve. For model  $\#11$  ref. [\[18\]](#page-149-4) determines that the geometry is not a Jacobian family arising from a family of smooth genus-2 curves.

which is isotrivial with the appropriate fixed fiber and scaling properties for dimensions  $\{2, 6\}$ . The discriminant of the polynomial in the RHS is (up to an overall constant) is

$$
(v - 2u3)10(v + 2u3)10
$$
 (5.36)

with the correct components. As for all isotrivial geometries, the SW curve is not a good description.

### 5.2.6 The subtle case: dimensions  $\{2,4\}$

**Generalities.** There are several known SCFTs with dimensions  $\{2,4\}$  we list them in table [3](#page-67-0) together with the relevant information on them taken from refs. [\[18,](#page-149-4) [89\]](#page-153-5). As always # refers to the number of the model in Martone's tables [\[89\]](#page-153-5).

From **Proposition [3](#page-61-1)** we know that there are *two distinct classes* of  $\{2, 4\}$  geometries:

(A) geometries whose fiber<sup>[58](#page-67-1)</sup> over the "good" axis  $u = 0$  is isomorphic to  $(E \times E)/\mathbb{Z}_2^2$  i.e. to the Jacobian of the hyperelliptic curve  $y^2 = x(x^4 + \alpha x^2 + 1)$ 

<span id="page-67-1"></span> $\frac{58}{100}$  At a general point in the conformal manifold  $\mathcal{M}$ .

(B) geometries whose fiber over the "good" axis  $u = 0$  is isomorphic to the non-Jacobian Abelian surface  $E \times E$ .

Class (A) geometries are expected to have a SW curve which is "nice" when the geometry is non-isotrivial. Class (B) geometries are not expected to have a SW curve neither nice nor ugly. The authors of ref. [\[18\]](#page-149-4) conclude that the SCFT  $\#11$  has a non-Jacobian geometry, i.e. model  $\#11$  is a first example of a  $\{2,4\}$  SCFT of class (B).

**The two isotrivial geometries.** We have two isotrivial  $\{2, 4\}$  geometries one of class (A) and one of class (B); they have the form

$$
\pi: (A \times \mathbb{C}^2) / \text{Weyl}(C_2) \to \mathbb{C}^2 / \text{Weyl}(C_2) \simeq \mathbb{C}^2 \equiv \mathscr{C}
$$
\n(5.37)

where the fixed Abelian variety  $\tilde{A}$  is, respectively, the Jacobian and non-Jacobian principally polarized Abelian surface with automorphism group  $Weyl(C_2)$ , cf. the table attached to Proposition [3](#page-61-1). This math result fits with the observation in [\[18\]](#page-149-4) that there are two inequivalent  $\{2, 4\}$  isotrivial geometries. Thus the SCFT #56 has the class (A) isotrivial geometry and the SCFT  $#11$  the class  $(B)$  isotrivial geometry.

Class (A) non-isotrivial. These geometries are the Jacobian of a fibration in genus-2 curves which on the locus  $u = 0$  are isomorphic to the curve in the last column of the table attached to Proposition [3](#page-61-1) and the SW differential in eq.[\(5.24\)](#page-65-2). Taking into account the proper scaling we have

$$
y^{2} = x(x^{4} + \alpha(\tau) x^{2} v + v^{2}) \text{ along } u = 0 \qquad (5.38)
$$

which is promoted by the replacement [\(5.31\)](#page-66-0) to the SW curve

$$
y^{2} = x(x^{4} + \alpha(\tau) x^{2} (v - x u) + (v - x u)^{2})
$$
\n(5.39)

which is the curve given in [\[18\]](#page-149-4) for models  $\#10$  and  $\#43$ .

The geometry of the model #48 looks "of the same kind" as the ones for the models  $#10, #43.$  Indeed the discriminant types suggest that these two geometries share the same C-VHS (of the Appell class). If this is the case, the two geometries are related by a Number-Theoretical phenomenon that we shall discuss in §. [8.7.1.](#page-102-0)

The geometry of  $Sp(4)$  with  $\frac{1}{2}$  16 is very subtle. It will be discussed in §.[8.1.](#page-81-0)

### 5.2.7 A few words about  $r > 2$

To use the above methods in arbitrary rank  $r$  we need the explicit classification of polarized Abelian varieties of dimension r corresponding to **Proposition [3](#page-61-1)** for  $r = 2$ . Without working out that classification we may only make some general consideration.

In the **Special situation** we expect – at least when the class number of the number field  $\mathbb{Q}(e^{2\pi i/\Delta_{i_1}})$  is 1 – that the fiber along the "good" axes has the form

$$
A_{\tau} = \left(\overbrace{E_{\tau} \times E_{\tau} \times \cdots \times E_{\tau}}^{r \text{ copies}}\right) / (\text{finite group}) \tag{5.40}
$$

where  $E_{\tau}$  is an elliptic curve of arbitrary period  $\tau$ , and

$$
Aut(A_{\tau}) \supseteq Weyl(G) \tag{5.41}
$$

for a simple Lie group G of rank  $r$ . Now everything boils down to the classification of the allowed (finite groups). Just as in the case of dimensions  $\{2, 4\}$  we have two possible groups leading to the geometry classes (A) and (B), in higher rank we expect a short list of finite group for each dimension  $r$ -tuple consistent with the **Special situation** condition. This will lead to a finite list of special geometry classes for each dimension  $r$ -tuple. Situations more general than the Special one can be studied along similar lines.

# 6 Rigidity II

## <span id="page-69-0"></span>6.1 Rigidity of the underlying Abelian family

A different notion of rigidity refers to the underlying Abelian fibration stripped of all other special-geometric structures (cf. Faltings [\[58\]](#page-152-1), Saito [\[59\]](#page-152-2), and Peters [\[60–](#page-152-3)[62\]](#page-152-4)). In this context we ask if we can continuously deform the holomorphic Abelian fibration  $\mathscr{A} \to \mathscr{P} \equiv \mathscr{P} \setminus \mathscr{S}$ while keeping fixed its basis (i.e. the special divisor  $\mathscr{S} \subset \mathscr{P}$ ). Here the crucial fact is that the base  $\mathscr{\tilde{P}}$  of the family is quasi-projective.

It is known that all non-isotrivial families of Abelian r-varieties over a quasi-projective base B are rigid if one of the following situations occur<sup>[59](#page-69-1)</sup> (cf. **Example 2.4(4)** of [\[62\]](#page-152-4) and references therein):

- $r < 7$ ;
- $\bullet$  the monodromy is irreducible and r is prime;
- $\bullet$  the base  $B$  is non-compact, the underlying variation of Hodge structure is irreducible and some local monodromy operator at the boundary has infinite order.

We have seen in §. [2.10.4](#page-41-0) that a non-isotrivial geometry has at least one local monodromy of infinite order, while our main interest is in geometries with irreducible monodromy. Then

**Fact 18.** The Abelian family  $\mathscr{A} \to \mathscr{\mathscr{P}}$  arising from a non-isotrivial, irreducible  $\mathbb{C}^{\times}$ -isoinvariant special geometry is rigid.

<span id="page-69-1"></span><sup>59</sup> I thank Chris Peters for clarifications on this issue.

Note that this holds independently of the assumption that  $\mathscr C$  is smooth. In practice we shall work only in lower rank, so rigidity in the non-isotrivial case is guaranteed by the condition that the rank is less than 8, even if the special geometry is reducible.

**Corollary 4.** The deformations of a non-isotrivial, irreducible,  $\mathbb{C}^{\times}$ -isoinvariant special geometry arise from isomonodromic deformations of the discriminant divisor D. The same holds in general for non-isotrivial geometries of rank less than 8.

In particular when  $\hat{\mathcal{P}}$  is rigid, the non-isotrivial special geometry is also rigid. Conversely:

<span id="page-70-1"></span>**Corollary 5.** If 2 is a Coulomb dimension and  $\mathcal{P}$  is rigid then the special geometry must be isotrivial.

### 6.2 The case of rank 2

In rank-2 all irreducible components of the special divisor  $\mathscr{S}$  are points in  $\mathscr{P} \simeq \mathbb{P}^1$ . Let s be the degree of  $\mathscr{S}$ . The moduli of allowed divisors is then a subspace of  $\mathcal{M}_{0,s}$ , the moduli space of curves of genus zero with s punctures. Then

Corollary 6. For a rank 2 non-isotrivial special geometry

<span id="page-70-0"></span>
$$
\dim \mathcal{M} \le s - 3. \tag{6.1}
$$

In particular the special geometry of a rank-2 SCFT with one (resp. two) Coulomb dimensions equal 2 either has an isotrivial geometry or at least 4 (resp. 5) special points in  $\mathbb{P}^1$ . An irreducible rank-2 SCFT has at least 3 special points.

Next we argue that the inequality is generically saturated in  $r = 2$ . Indeed, suppose we have a special geometry with s special points. Its  $\mu$ -monodromy be generated by the s local monodromies  $\mu_i \in Sp(4, \mathbb{Z})$  with  $\mu_1 \cdots \mu_s = 1$ . Now move the special points a little bit to new positions nearby. By the Riemann-Hilbert correspondence there is a Picard-Fuchs connection with the monodromies  $\mu_i \in Sp(4,\mathbb{Z})$  around these points. By the Almkvist-Zudilin theorem this defines a special geometry (the positivity of Im  $\tau_{ij}$  will be not destroyed by small deformation, nor it will spoil regularity). The deformed geometry is isomorphic to the original one if we can undo the deformation by a automorphism of  $\mathbb{P}^1 \simeq \mathscr{P}$ . Thus – if we are allowed to move the points freely – the number of proper deformations is at least  $s - 3$ . Since it is at most  $s - 3$ , we get that it is precisely  $s - 3$ . More generally, the number of deformations is at least the number of points which we may move freely less 3. Usually we have no constraint on the positions, and the inequality is saturated. The only mechanism that may freeze some degrees of freedom of the special discriminant  $\mathscr{S} = \sum_i z_i$ is the condition that the family should preserve some extra symmetry. This can only happen under exceptional circumstances. In this case fixing the positions of a suitable subset of  $\dim M + 3$  points of  $\mathscr S$  determines the other  $s - 3 - \dim M$ .

Corollary 7. The conformal manifold of geometries with dimensions  $\{2, \Delta\}, \Delta \neq 2$  is (modulo finite covers) the moduli space of the sphere less 4 points, i.e. the modular curve  $H/\Gamma_0(2)$ , while the moduli space of geometries with dimensions  $\{2,2\}$  is the moduli of the sphere less 5 points.

### 6.2.1 Comparison with known models: non-rigid geometries

Let us compare the last statement with the list of known rank-2 SCFTs (table 1 of [\[89\]](#page-153-5)): the models with one  $\Delta_i = 2$  are the numbers  $\#10, \#11, \#12, \#38, \#43, \#49, \#56, \#60, \#68,$ #69 where we have underlined the isotrivial ones. All isotrivial geometries have 3 special points,[60](#page-71-0) and all the non-isotrivial ones are known to have 4 special points except the last one which looks to have 7 special points, namely, 6 divisors of type  $I_1$  and the enhanced axis of dimension 4. The models with  $\Delta_1 = \Delta_2 = 2$  are the numbers #13, #31 and #32. The last one is reducible and so has only 2 special points, the other two have 5 special points. See e.g. [\[121\]](#page-155-9) where it is shown explicitly that the conformal manifold of the  $\{2, 2\}$  geometries is  $\mathcal{M}_{0.5}$  (the moduli space of the 5-punctured sphere) in agreement with Gaiotto's class- $\mathcal S$ picture  $[110]$ . Thus, with the single exception of model  $\#69$ , all other saturate the bound [\(6.1\)](#page-70-0). Note that in the non-rigid isotrivial special geometries (which correspond to  $\mathcal{N}=4$ SCFTs) the continuous deformations of the geometry leave fixed the discriminant  $D$ . This is not possible in the non-isotrivial case – that is, when at least along one component  $\mathcal{D}_i$  the light degrees of freedom are IR-free – because of Falting-Saito-Peters rigidity.

### 6.2.2 A scenario for the mysterious  $\#69$

 $Sp(4)$  SYM coupled to  $\frac{1}{2}$  16 has several peculiar properties which single it out from all other SCFT. It has no known construction in string theory so we have no clue on its geometry from physics. It has dimensions  $\{2, 4\}$ , six discriminant components of the form  $u_2 - z_a u_1^2 = 0$ ,  $z_a \in \mathbb{C}$   $(a = 1, \ldots, 6)$  and one regular enhanced axis  $u_1 = 0$  [\[89\]](#page-153-5). The conformal manifold is one-dimensional, so we have to freeze 3 out of the 7 special points. A scenario which comes to mind is the following one. Consider the  $\mathbb{Z}_2$ -quotient

$$
s: (u_1, u_2) \to (v_1, v_2) = (u_1, u_2^2) \tag{6.2}
$$

and suppose that on the base we have an Abelian family  $\mathscr{B} \to \mathbb{P}(2,8) \setminus \mathscr{S}$  with the 4 special points  $v_1 = 0$ ,  $v_2 - \alpha_k^2 v_1^4 = 0$  ( $k = 1, 2, 3$ ). We may fix  $\alpha_1$ ,  $\alpha_2$  using the residual automorphisms of  $\mathbb{P}^1$ , remaining with one essential parameter  $\alpha_3$ . The pull-back of this family via s would be an Abelian family over  $\mathscr P$  with 7 special points  $u_1 = 0$  and  $u_2 \pm \alpha_k u_1^2 = 0$  $(k = 1, 2, 3)$ , which depend on the single parameter  $\alpha_3$ . One should validate (or reject!) this scenario by constructing the Abelian family on the quotient. We shall leave the hard analysis to future work. A possibility to be explored is that the local monodromy  $\gamma$  of  $\mathscr{B}$  around the  $v_2$  axis is such that  $\gamma^2$  is the automorphism of the fiber of  $\mathscr A$  on the  $u_2$  axis (described in

<span id="page-71-0"></span><sup>60</sup> Cf. Corollary [5](#page-70-1).
**Proposition [3](#page-61-0)**) with analytic representation  $\left(\begin{smallmatrix} 1 & 0 \\ 0 & -1 \end{smallmatrix}\right) \in \text{Weyl}(Sp(4))$ . Note that the family  $\mathscr B$ cannot describe a special geometry on its own right since  $\{2, 8\}$  is not an allowed dimension pair (it existence is also ruled out by the Folk-theorem).

#### 6.2.3 Comparison with known models: rigid geometries

One expects that a rank-2 *rigid* special geometry has *exactly* 3 special points. In the table of known SCFTs in [\[89\]](#page-153-0) there is only one apparent counterexample to this statement, namely model  $#22$  with dimensions  $\{4, 6\}$ . However in §. 4.2 of that paper it is said:

Unfortunately there is another possible solution which is compatible with everything that we know, i.e. a single knotted stratum with  $\mathfrak{T}_{u^3+v^2} = [II, \emptyset]$ . [...] we pick the former but for not better reason than symmetries with the other unknotted stratum, also of  $I_n$  type. It is important to clarify, that these two choices are indistinguishable as far as the analysis we are performing here goes [...] we expect that a more careful analysis of the global structure of the CB could very likely distinguish the two cases.

Our guess is that the second choice is the correct one, thus eliminating the counterexample to equality. However another geometric scenario does exist, a covering scenario similar to the one suggested for  $\#69$ . Consider the  $\mathbb{Z}_2$  cover

$$
f: (u_1, u_2) \to (v_1, v_2) = (u_1^2, u_2)
$$
\n
$$
(6.3)
$$

over a plane with coordinates of dimensions  $\{8, 6\}$ . There is a known geometry with these dimensions, the one of the SCFT  $#2$  of [\[89\]](#page-153-0), which has a discriminant component  $v_1 = 0$ of type  $I_6^*$  and a divisor  $v_1^3 - v_2^4 = 0$  of type  $I_1$ . The pull-back of the Abelian family  $\mathscr{A} \to \mathbb{P}(8,6)$  to an Abelian family  $f^*\mathscr{A} \to \mathbb{P}(4,6)$  will have a discriminant component  $u_1 = 0$ with monodromy the square of the monodromy at  $v_1 = 0$ , i.e. Kodaira type  $I_{12}$ . In addition we have the two discriminant components  $u_1^3 \pm u_2^2 = 0$  both of type  $I_1$ . This is precisely the "stratification" proposed in [\[89\]](#page-153-0) for the  $#22$  model. For these precise monodromies, the would-be  $\{4, 6\}$  geometry passes the text of local regularity of the periods around the axis of dimension 4.

Thus the present analysis of the global structure of the Coulomb branch cannot decisively support one possibility over the other. It is even possible that both geometries exist: this will explain why both pass all possible checks geometrical as well as physical.

#### 6.2.4 Covering and primitive geometries

The discussion following **Corollary [6](#page-70-0)** shows that whenever  $s - 3 > \dim M$  there must be relations between the positions of the special points and we suggested above that this very rare phenomenon may arise when there is a cyclic cover  $f: \mathscr{P} \to \mathbb{Q}$  – branched at most over the points  $(1,0)$  and  $(0,1)$  – and the special divisor is invariant under the deck group. On

 $\mathscr{Q} \setminus f(\mathscr{S})$  we have an Abelian family  $\mathscr{B} \to \mathscr{Q}$  whose pull-back to  $\mathscr{\hat{P}}$  is the Abelian family defining the original special geometry.

**Definition 13.** A special geometry is *primitive* iff its underlying Abelian family  $\mathscr{A} \to \mathscr{P}$ cannot be written as the pull-back of a family on a non-trivial quotient of  $\mathscr{P}$ .

**Fact 19.** An indecomposable non-isotrivial rank-2 geometry with  $s-3 = \dim M$  is primitive.

Indeed the quotient geometry has dim  $\mathcal{M}$ +3−t  $\geq 1$  special points  $\neq$  (1 : 0), (0 : 1) where  $1 \le t \le 2$  is the number of non-trivial monodromies around the points  $(1:0),(0:1)$  in the quotient. The number of special points of the covering geometry is  $s = d(\dim \mathcal{M} + 3 - t) + \tilde{t}$ , where d is the degree of the cyclic cover and  $\hat{t}$  the the number of non-trivial monodromies around the points  $(1:0),(0:1)$  in the quotient. For a non-isotrivial geometry  $t \geq 2-\dim M$ . When  $t = \tilde{t}$  we have

$$
0 = s - 3 - \dim \mathcal{M} = (d - 1)(\dim \mathcal{M} + 3 - t)
$$
\n(6.4)

which can be satisfied only if  $d = 1$ , i.e. if the geometry is primitive. Otherwise we have dim  $M = 1, 2$  and

$$
\dim \mathcal{M} = 1 \qquad 4 = s = d(4 - t) + \tilde{t} \ge 2d + 1
$$
\n
$$
\dim \mathcal{M} = 2 \qquad 5 = s = d(5 - t) + \tilde{t} \ge 3d \tag{6.5}
$$

both of which can be satisfied only for  $d = 1$ . Thus for a rank-2 indecomposable nonisotrivial geometry  $s - 3 = \dim M$  if and only if the geometry is primitive. Most geometries are primitive. We shall focus on primitive geometries. The non-primitive ones (if they exist) can be studied by the same methods going to the quotient.

### 6.3 Finiteness theorems

These rigidity properties, together with a bound on the degree, give Falting-Peters finiteness theorems (cf. **Theorem 4.2** of  $|60|$ ):

**Theorem** (Falting, Peters). For fixed  $\mathscr{S} \subset \mathscr{P}$  there are only finitely many non-isotrivial Abelian families which can arise from an irreducible non-isotrivial special geometry with special divisor  $\mathscr{S}$ .

Remark 11. Since all simple Abelian surfaces are Jacobians of genus 2 curves, in rank 2 the above essentially reduces to Arakelov rigidity [\[57\]](#page-151-0).

This finiteness result parallel another deep result from the VHS side:

**Theorem** (Deligne finiteness theorem [\[124\]](#page-155-0)). Fix a connected quasi-projective manifold  $\mathscr{P}$ and an integer N. There are at most finitely many conjugacy classes of rational representations of  $\pi_1(\tilde{\mathscr{P}})$  of dimension N giving local systems that occur as a direct factor of a polarized variation of Hodge structure on  $\tilde{\mathscr{P}}$  (of any weight).

To get the expected result that only finitely many special geometries exist for each rank r it remains to show that only finitely many divisors  $\mathscr S$  are possible for rigid irreducible geometries. In rank-2 we have the stronger statement that there is a unique  $\mathscr S$  modulo automorphisms of  $\mathscr{P} \simeq \mathbb{P}^1$  namely  $\mathscr{S}$  should consists of three distinct points. In the general case we already said some words on this issue in §. [3.2](#page-47-0) and we shall return to the question in sect. [9.1.](#page-123-0)

## 6.4 Rigidity in the Deligne-Simpson sense

For the third notion of rigidity we strip the special geometry of all its structures but for the  $\mu$ -monodromy representation<sup>[61](#page-74-0)</sup>

$$
\mu \colon \pi_1(\mathscr{P} \setminus \mathscr{S}) \to Sp(2r, \mathbb{Z}) \subset GL(2r, \mathbb{C}), \tag{6.6}
$$

and we forget that the monodromy was defined over  $\mathbb Q$  (and in fact integral), seeing it as a mere representation in  $GL(2r, \mathbb{C})$ . In particular we consider two representations to be the same if they are conjugated in  $GL(2r,\mathbb{C})$ .

Then  $\mu$  is a complex representation of dimension  $2r$ , which in the interesting situations is irreducible, with the property that the local monodromy  $\mu_i$  around the *i*-th component of the divisor  $\mathscr S$  belongs to a prescribed quasi-unipotent conjugacy class  $C_i \subset GL(2r, \mathbb C)$ .

**Definition 14.** An irreducible representation  $\mu: \pi_1(\tilde{\mathscr{P}}) \to GL(2r, \mathbb{C})$  with  $\mu_i \in C_i$  is called *rigid* if all nearby irreducible representations with local monodromies  $\mu'_i \in C_i$  are conjugated in  $GL(2r, \mathbb{C})$  to it. It is called *absolutely rigid* if all such irreducible representations are conjugate.

When dim  $\mathscr{\dot{P}} = 1$  rigidity implies absolute rigidity. The Simpson celebrated conjecture [\[63\]](#page-152-1)  $states<sup>62</sup>$  $states<sup>62</sup>$  $states<sup>62</sup>$ 

Conjecture (Simpson). Every rigid irreducible representation of the fundamental group of a smooth complex algebraic variety  $\mathscr{P} = \mathscr{P} \setminus \mathscr{S}$ , with quasi-unipotent monodromies around the components of  $\mathscr S$  is motivic *i.e.* arises from a VHS.

The **Conjecture** is a theorem when dim  $\mathcal{P} = 1$ ; this follows from the work of many distinguished mathematicians culminating in the explicit construction of the rigid monodromies by Katz [\[102\]](#page-154-0) later reinterpreted in the representation theoretical language by Crawley-Boevey<sup>[63](#page-74-2)</sup> [\[128](#page-155-1)[–132\]](#page-156-0). In ref. [\[134\]](#page-156-1) it was shown how these results can be rephrased in terms of Seiberg duality for SUSY gauge theories.

The **Conjecture** is proven in general in the category of  $\mathbb{C}$ -VHS, that is, it is proven that a rigid irreducible representation (with the appropriate  $C_i$ 's) yields a *weak special geometry* 

<span id="page-74-1"></span><span id="page-74-0"></span><sup>&</sup>lt;sup>61</sup> For  $\varkappa = 2$  we need to choose a suitable lift of the monodromy from  $PSp(2r, \mathbb{Z})$  to  $Sp(2r, \mathbb{Z})$ .

 $62$  A somehow related statement is the Bombieri-Dwork conjectures about G-functions [\[125,](#page-155-2) [126\]](#page-155-3). The Laplace transform of G-functions are the BPS brane amplitudes in the asymmetric limit [\[127\]](#page-155-4).

<span id="page-74-2"></span><sup>63</sup> For a textbook account see [\[100\]](#page-154-1).

(see Lemma 4.5 of  $[63]$ ). The aspect of the **Conjecture** which is still open is the numbertheoretic one. When the monodromy is rigid, the representation  $\mu$  automatically takes value in  $GL(2r, \mathbb{F})$  for some number field  $\mathbb{F}$ , that is, it corresponds to a direct factor of a Q-variation of Hodge structure (Theorem 5 of [\[63\]](#page-152-1)). The Conjecture claims that this representation is also *integral*, i.e.  $\mu$  takes values in  $GL(2r, \mathcal{O}_{\mathbb{F}})$  where  $\mathcal{O}_{\mathbb{F}}$  is the ring of algebraic integers in **F.** Equivalently: the traces tr  $\gamma$  ( $\gamma \in \Gamma$ ) are rational integers.

#### 6.4.1 Properties of rigid monodromy representations

Let us explain in elementary terms a special (and simpler) instance of the 'surprising' result of [\[63\]](#page-152-1), that is, why an absolutely rigid, irreducible, complex, monodromy representation  $\mu$ with local conjugacy classes  $C_i \subset GL(2r, \mathbb{C})$  such that:

- (1) for all  $i$  the elements of  $C_i$  are quasi-unipotent with at most one non-trivial Jordan block of size 2;
- (2) for all *i*:  $C_i \cap Sp(2r, \mathbb{Z}) \neq \emptyset$ ;
- (3) the elements of at least one  $C_i$  have infinite order i.e. some power of the elements of  $C_i$  is unipotent with a single Jordan 2-block

is automatically contained in a real form of  $Sp(2r, \mathbb{C})$ , so yields a R-VHS which, under some "mild" regularity condition (cf. §. [4\)](#page-48-0) describes a *weak* special geometry. By assumption the  $C_i$ 's are invariant under  $g \mapsto (g^t)^{-1}$  and  $g \mapsto g^*$ . Hence by rigidity

$$
\mu = M^{-1}(\mu^t)^{-1}M, \qquad \mu = R^{-1}\mu^*R \tag{6.7}
$$

for some invertible matrices  $M$ ,  $R$  with

<span id="page-75-2"></span>
$$
(Mt)-1M \mu = \mu (Mt)-1M, \qquad RR^* \mu = \mu RR^*.
$$
 (6.8)

These equations imply

<span id="page-75-1"></span>
$$
\mu^t M \mu = M \quad \text{and} \quad \mu^\dagger H \mu = H \quad \text{where} \quad H \equiv M^* R. \tag{6.9}
$$

Since the representation is irreducible, the invariant bilinear form  $M$  must be either symmetric or antisymmetric while, being invertible,  $M$  is non-degenerate. A non-degenerate symmetric M is not consistent<sup>[64](#page-75-0)</sup> with condition (3) and should be ruled out. Changing basis, we may assume that M is the standard (real) symplectic matrix  $\Omega$ . This shows that the rigid monodromy group  $\Gamma \subset Sp(2r, \mathbb{C})$ . The second eq.[\(6.9\)](#page-75-1) then implies that  $\Gamma$  is in a real form of  $Sp(2r,\mathbb{C})$ . More specifically: by Schur's lemma and eq.[\(6.8\)](#page-75-2)  $RR^*$  and  $H^{\dagger}H^{-1}$ 

<span id="page-75-0"></span><sup>&</sup>lt;sup>64</sup> A matrix  $\mu$  with rank $(\mu - 1) = 1$  which satisfies the first eq.[\(6.9\)](#page-75-1) for  $M = M^t$  must have the form  $\mu = 1 + v \otimes v^t M$  with  $v^t M v = -2$ , while the condition  $(\mu - 1)^2 = 0$  requires  $v^t M v = 0$ .

are multiples of the identity, so  $H^{\dagger}H^{-1} =$  (phase). We can set  $RR^* = 1$  as a choice of normalization, fixing  $R$  modulo an overall phase. Now

$$
(\Omega R)^{\dagger} = H^{\dagger} = (\text{phase}) H = (\text{phase}) \Omega R \tag{6.10}
$$

and we may fix the phase of R so that H is anti-Hermitian i.e.  $R^{\dagger} \Omega = \Omega R$  or

$$
R^t \Omega R \equiv (R^\dagger)^{-1} \Omega R = \Omega,\tag{6.11}
$$

that is,  $R \in Sp(2r, \mathbb{C})$ . Since  $Sp(2r, \mathbb{C})$  is simply connected and  $R^* = R^{-1}$ , we can find  $L \in \mathfrak{sp}(2r, \mathbb{C})$  with  $L^* = -L$  such that  $R = \exp(L)$ . Let  $S \equiv \exp(L/2) \in Sp(2r, \mathbb{C})$ ; one has  $S^2 = R$  and  $SS^* = 1$ , hence

$$
(S\mu S^{-1})^* = S^* \mu^* (S^*)^{-1} = S^{-1} \mu^* S = S^{-1} (S^2 \mu^* S^{-2}) S = S \mu S^{-1}
$$
(6.12)

that is, after conjugating with S, the monodromy  $\mu$  takes value in  $Sp(2r, \mathbb{R})$ .

#### 6.4.2 Obstruction to be rational: the Brauer group

By rigidity, the traces  $tr(\gamma)$  ( $\gamma \in \Gamma$ ) are complex numbers invariant under all automorphisms of  $\mathbb{C}$ , so are algebraic numbers. Since the Galois group  $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$  fixes the traces, they take value in  $\mathbb Q$ . Then there exists a finite extension  $\mathbb F$  of  $\mathbb Q$  such that (after a suitable overall conjugation)  $\Gamma \subset Sp(2r, \mathbb{F})$ . The number field  $\mathbb F$  is either totally real or a purely imaginary quadratic extension of a totally real field [\[63\]](#page-152-1). As already mentioned, the Deligne-Simpson conjecture requires Γ to be integral:<sup>[65](#page-76-0)</sup> Γ ⊂  $Sp(2r, \mathcal{O}_{\mathbb{F}})$ . In particular the traces are ordinary integers. However to be the monodromy of an actual family of Abelian r-folds  $\mu$  should be integral and *rational* i.e.  $\Gamma \subset Sp(2r,\mathcal{O}_\mathbb{Q}) \equiv Sp(2r,\mathbb{Z})$ .

The Brauer group of the number field  $\mathbb F$  gives a potential obstruction to the rationality of the monodromy representation. Let  $\sigma \in \text{Gal}(\mathbb{F}/\mathbb{Q})$ ; by rigidity we have

<span id="page-76-1"></span>
$$
\gamma_i^{\sigma} = M_{\sigma}^{-1} \gamma_i M_{\sigma}, \quad \forall \ i \tag{6.13}
$$

for some matrix  $M_{\sigma} \in GL(2r, \mathbb{F})$  which is unique up to multiplication by an element of  $\mathbb{F}^{\times}$ by irreducibility. Now

$$
M_{\sigma\tau}^{-1} \gamma_i M_{\sigma\tau} = \gamma_i^{\sigma\tau} = (\gamma_i^{\tau})^{\sigma} = (M_{\tau}^{-1} \gamma_i M_{\tau})^{\sigma} =
$$
  
= 
$$
(M_{\tau}^{\sigma})^{-1} \gamma_i^{\sigma} (M_{\tau})^{\sigma} = (M_{\tau}^{\sigma})^{-1} M_{\sigma}^{-1} \gamma_i M_{\sigma} (M_{\tau})^{\sigma},
$$
 (6.14)

that is,

$$
C_{\sigma\tau} \equiv M_{\sigma\tau} (M_{\sigma} (M_{\tau})^{\sigma})^{-1} \in \mathbb{F}^{\times}, \qquad \sigma, \tau \in \text{Gal}(\mathbb{F}/\mathbb{Q}). \tag{6.15}
$$

is a 2-cocycle on the Galois group with values in  $\mathbb{F}^{\times}$ . When the cohomology class  $[C_{\sigma\tau}]$  in the

<span id="page-76-0"></span><sup>&</sup>lt;sup>65</sup> Recall that the symbol  $\mathcal{O}_{\mathbb{F}}$  stands for the ring of integers in the number field  $\mathbb{F}$ .

Brauer group  $H^2(\mathbb{F}, \mathbb{F}^\times)$  vanishes, we can chose  $M_{\sigma}$  so that  $M_{\sigma\tau} = M_{\sigma}(M_{\tau})^{\sigma}$ , and  $M_{\sigma}$  is a Galois 1-cocycle with values in  $GL(2r, \mathbb{F})$ . The non-Abelian extension of Hilbert's **Theorem** 90 states [\[135\]](#page-156-2)

$$
H^1(\text{Gal}(\mathbb{F}/\mathbb{Q}), GL(2r, \mathbb{F})) = \{1\},\tag{6.16}
$$

that is, there exists a matrix  $N \in GL(2r, \mathbb{F})$  such that

$$
M_{\sigma} = N^{-1} N^{\sigma} \qquad \forall \sigma \in \text{Gal}(\mathbb{K}/\mathbb{Q}). \tag{6.17}
$$

Substituting this equation in [\(6.13\)](#page-76-1) we get

$$
(N\gamma_i N^{-1})^{\sigma} \equiv N^{\sigma} \gamma_i^{\sigma} (N^{\sigma})^{-1} = N\gamma_i N^{-1} \quad \forall \ i, \quad \forall \ \sigma,
$$
\n(6.18)

that is, the matrices  $\eta_i = N \gamma_i N^{-1}$  yield a conjugate representation defined over Q which is integral by the Simpson conjecture.

However the Brauer class  $[C_{\sigma\tau}]$  is not zero in general. Therefore to conclude that a particular rigid irreducible representation satisfying  $(1)(2)(3)$  yields a viable monodromy group for special geometry we have to check that its Brauer class is zero. Otherwise we can only conclude that the monodromy is a direct summand in some higher degree VHS [\[63\]](#page-152-1), and hence not directly relevant for special geometry.

## 6.5 Use of rigidity to construct special geometries

It may happen that a special geometry has a rigid monodromy representation  $\mu$ .

**Definition 15.** A special geometry is  $\mu$ -rigid if it is rigid (i.e. its conformal manifold  $\mathcal{M}$ is a point), and its monodromy representation  $\mu: \pi_1(\tilde{\mathscr{P}}) \to GL(2r, \mathbb{C})$  is either irreducible and rigid (as a monodromy representation) or it decomposes in the direct sum of rigid representations.

By the Riemann-Hilbert correspondence being  $\mu$ -rigid is equivalent to saying that the associated Gauss-Manin (GM) connection (Picard-Fuchs ODE for  $r = 2$ ) is rigid (modulo gauge equivalence). In  $\S$ . [4](#page-48-0) we suggested that "under some mild conditions" an irreducible Gauss-Manin connection over  $\mathring{\mathscr{P}}$  with the appropriate local monodromies will yield a special geometry; in  $r = 2$  this follows<sup>[66](#page-77-0)</sup> from the Almkvist-Zudilin theorem [\[98\]](#page-154-2) and we expect that a similar story holds for all  $r$ . The basic idea is that one can construct the GM connection of the  $\mu$ -rigid special geometries by exploiting their rigidity. This is a method pioneered by Riemann in his 1851 lecture notes on the hypergeometric ODE: he was able to construct integral representations for the solutions and compute the connection coefficients using rigidity of the monodromy. Since Riemann's times a quantity of important results have been obtained using this method.

<span id="page-77-0"></span><sup>&</sup>lt;sup>66</sup> Subject to the checks of Im  $\tau_{ij} > 0$  and local regularity.

Our interest in the class of  $\mu$ -rigid special geometries stems from the fact that all rigid special geometries tend to be  $\mu$ -rigid.

## Conjecture. Most, and possibly all, rigid  $\mathbb{C}^{\times}$ -isoinvariant special geometries are  $\mu$ -rigid.

Evidence. In rank 1 the statement is trivially true. The seventy or so known rank 2 special geometries [\[89\]](#page-153-0) have rigid monodromy representation, including the non-rigid (i.e. Lagrangian) ones. For all but (at most) two indecomposable rank-2 geometries the monodromy representation is in addition irreducible. As emphasized on page 360 of [\[100\]](#page-154-1), rigidity becomes a milder and milder requirement on the monodromy as we increase the rank  $r$ . The point is that the fundamental group of the complement  $\pi_1(\mathscr{P} \setminus \mathscr{S})$  is finitely-presented for all geometries in any  $r$ . But while the number of generators grows slowly with  $r$  the number of independent relations grows quite rapidly, producing a typically overdetermined set of algebraic equations for the moduli space of monodromy representations which then is typically empty for most choices of the  ${C_a}$ 's. Since already for  $r = 2$  all (or at least the vast majority) of the special geometries are  $\mu$ -rigid, this suggests that this pattern would hold for all r.  $\Box$ 

Regardless for the validity of the **Conjecture**, the  $\mu$ -rigid geometries form a big chunk of the finite set of all rigid special geometries of given  $r$ . Then in our classification scheme (cf.  $\S. 3$ ) we introduce class V of special geometries which are  $\mu$ -rigid and do not belong to simpler classes I-IV which can be addressed with simpler techniques. Then we proceed to study, and hopefully solve, the inverse problem for this important class of geometries by exploiting the wonders of rigid monodromies (with the hope that the Conjecture is true and class VI is small or empty). We shall pursue this program for the rest of the paper.

The crucial question is when a rigid monodromy leads to a non-isotrivial special geometry. A necessary condition is that the local monodromies are quasi-unipotent, real symplectic, with at most one non-trivial Jordan block of dimension 2, and at least one local monodromy non-semisimple. The Deligne-Simpson conjecture (theorem for  $r = 2$ ) then says that the monodromy arises from a VHS. In particular the monodromy group is a subgroup of  $Sp(2r, \mathcal{O}_{\mathbb{F}})$  where  $\mathcal{O}_{\mathbb{F}}$  is the ring of algebraic integers in some number field  $\mathbb{F}$  (a finite extension of  $\mathbb{Q}$ ). Given the rigid monodromy representation  $\mu$  we can write the corresponding rigid GM connection  $\nabla$  on  $\mathscr{\dot{\mathscr{P}}}$  and solve the flat-section equation  $\nabla \Phi = 0$ . To declare that we have constructed a  $\mathbb{C}^{\times}$ -isoinvariant special geometry in the sense of **Definition [4](#page-15-0)** it remains only to check:

- (i) the existence of a particular solution  $\Pi$  of the flat-section equation which satisfies the Legendre condition. This is automatically true for  $r = 2$  (cf. [4\)](#page-48-0) and plausibly also for general r under some (yet to be fully understood) "mild" conditions;
- (ii) the Brauer class of  $\mu$  is trivial so that  $\Gamma \subset Sp(2r, \mathbb{Z});$
- (iii) Im  $\tau_{ij} > 0$ , i.e. the VHS implied by the Deligne-Simpson conjecture has weight 1;

(iv) regularity of the geometry, that is, that the SW periods pulled back on  $\mathscr{C}$  (as in §. [4.2.1](#page-53-0)) for  $r = 2$ ) have the behavior along the irreducible components of the discriminant dictated by the local models of the corresponding singular fiber types, cf. §. [2.9,](#page-32-0) and no singularity along the enhanced divisors, where they should satisfy the conditions in §. [2.6.](#page-24-0)

When the monodromy representation is an appropriate one, we expect that all these conditions are automatically satisfied, without imposing any new one. Since it is still not fully obvious what we shall mean by "appropriate" monodromy representation, when in doubt we shall perform the checks ex post.

# 7 The inverse problem vs. rigidity

We now return to our inverse problem in rank  $r$  which we state in more precise terms:

Inverse Problem. Suppose we are given the following set of data:

- (a) the Coulomb dimensions  $\{\Delta_1, \ldots, \Delta_r\}$  from which we construct the (normal projective) manifold  $\mathscr{P} \equiv \mathbb{P}(d_1, d_2, \ldots, d_r)$ , and identify its cyclic quotient orbifold divisors;<sup>[67](#page-79-0)</sup>
- (b) a finite list  $\{p_1, \dots, p_\ell\}$  of irreducible quasi-homogeneous polynomials in  $\mathbb{C}[u_1, \dots, u_r]$  $(\text{deg } u_i = d_i)$  as candidate irreducible components of the discriminant. The  $\ell$ -tuple of polynomials is severely restricted by the conditions briefly discussed in §. [3.2](#page-47-0); we shall return to them in §. [9.1](#page-123-0);
- (c) a conjugacy class  $C_a \subset Sp(2r, \mathbb{Z})$  of Kodaira type per each component  $(p_a)$  of the discriminant;

and ask:

- (1) there exists an indecomposable special geometry with these data?
- (2) how many are they? What are the degrees of the allowed polarizations?
- (3) What can we say about the geometry (when it exists)?

We consider only data  $(\{\Delta_i\}, \{p_a\}, C_a)$  for which the answer to  $(1)$  is not "no" for trivial reasons. So the  $\{\Delta_i\}$  is an allowed r-tuple of dimensions, the discriminant satisfies the various necessary conditions, and in particular the group  $\pi_1(\mathscr{C} \setminus \sum_a(p_a))$  is not solvable, finite, or a non-trivial product, and when an axis (or a coordinate plane) is not part of the discriminant the corresponding restrictions on the dimensions are satisfied, etc. Moreover we do not loose nothing in assuming that the data correspond to a geometry of a non-isotrivial rigid geometry, that is,

<span id="page-79-0"></span> $67$  Here we mean "orbifolds" in the projective sense, the underlying manifold/variety may be smooth along the divisor (this happens when the quotient is by a reflection group [\[71\]](#page-152-2)). For the distinction between the two notions we refer to the discussion in [\[44\]](#page-151-1).

- (d) neither 1 nor 2 belong to the list  $\{\Delta_i\}$
- (e) at least one local monodromy is not semisimple.

The first step to give a positive answer to  $(1)$  is to show that there exists a monodromy representation of  $\pi_1(\tilde{\mathscr{P}})$  of dimension 2r which has the given local monodromies  $\mu_a \in C_a$ . Again, we may set a part the reducible indecomposable case (which carries additional constraints) for a separate discussion, and focus on the typical case of *irreducible* monodromy representations. The irreducible monodromies with the appropriate local classes  ${C_a}$  are parametrized by a moduli space  $\mathcal{M}(\{\Delta_i\}, \{p_a\}, C_a)$ . The monodromy is rigid (resp. properly rigid) iff  $\mathcal{M}(\{\Delta_i\}, \{p_a\}, C_a)$  is non empty of dimension zero (resp. a point). Then a necessary condition for a positive answer to  $(1)$  is:

Necessary condition. The moduli space of irreducible representations of  $\pi_1(\tilde{\mathscr{P}})$  with the fixed local monodromy classes should be non-empty  $\mathcal{M}(\{\Delta_i\}, \{p_a\}, \{C_a\}) \neq \emptyset$ .

For "most" data  $(\{\Delta_i\}, \{p_a\}, \{C_a\})$  all representations with the required local monodromies turn out to be reducible, i.e. the moduli space  $\mathcal{M}(\{\Delta_i\}, \{p_a\}, \{C_a\})$  is empty. Thus the necessary condition is already very stringent. Indeed, for large  $r$  the set of representations with given  ${C_a}$  is empty even before imposing irreducibility.

Our main point is that there is also an "almost sufficient" condition:

## "Almost" sufficient condition. dim  $\mathcal{M}({\{\Delta_i\}}, \{p_a\}, \{C_a\}) = 0.$

This is "almost" sufficient because<sup>[68](#page-80-0)</sup> when it holds we have still to make three checks before we can declare that we have a bona fide special geometry:

- (i) the vanishing of the Brauer class  $[C_{\sigma\rho\tau}] \in H^2(\mathbb{F}, \mathbb{F}^\times);$
- $(ii)$  the regularity of the solution (which is expected to take care of itself most of the time);
- (*iii*) the positivity of  $\text{Im } \tau_{ij}$  (also expected to be typically true).

"Experimentally" the criterion looks pretty good as a "sufficient" condition in the sense that the checks have the tendency to go well. By definition the special geometries satisfying the "almost sufficient" condition are of class V.

## 8 The inverse problem in rank-2

In this section we run the inverse problem program in rank-2 in more or less full detail. We suppose the geometry to be indecomposable (thus  $\mathcal{R}_1 = 0$ ).

<span id="page-80-0"></span><sup>&</sup>lt;sup>68</sup> Assuming the validity of the Deligne-Simpson conjecture (which is a theorem in many special cases).

When the geometry is primitive and irreducible the situation is pretty clear. Let  $s =$  $\dim_{\mathbb{C}} \mathcal{R}_2$  be the dimension of the conformal manifold  $\mathcal{M}$ . The complement  $\mathcal{P}$  is the Riemann sphere  $\mathbb{P}^1$  minus exactly  $s+3$  special points  $p_a$ . Then

$$
\pi_1(\tilde{\mathscr{P}}) = \langle l_1, \dots, l_{s+3} : l_1 l_2 \cdots l_{s+3} = 1 \rangle. \tag{8.1}
$$

When  $x = 1$  we are interested in monodromy representations of  $\pi_1(\hat{S})$  taking value in the arithmetic group  $Sp(4, \mathbb{Z})$ , but let start by considering the underlying complex representations valued in  $GL(4,\mathbb{C})$ . An irreducible representation of  $\pi_1(\mathscr{P})$  consists of  $s+3$  complex  $4 \times 4$  matrices  $\mu_a \equiv \mu(\ell_a)$   $(a = 1, \ldots, s + 3)$  such that  $\mu_1 \cdots \mu_{s+3} = 1$  which have no nontrivial common invariant subspace in  $\mathbb{C}^4$ . In addition, the conjugacy classes of the  $s+3$  local monodromies  $\mu_a$  are prescribed. The space of all such complex representations is

$$
\{\mu_1, \dots, \mu_{s+3} \in GL(4, \mathbb{C}) : \mu_1 \cdots \mu_{s+3} = 1, \ \mu_a \in C_a \ a = 1, \dots, s+3, \text{ irreducible}\}\
$$
 (8.2)

Two representations are considered to be the same (as complex representations) iff they are related by an overall conjugation  $\mu'_{a} = U \mu_{a} U^{-1}$  with  $U \in GL(4, \mathbb{C})$ .

When  $\varkappa = 2$  the  $\mu$ -monodromy representation is well defined in  $PSp(4, \mathbb{Z})$ . Suppose we lift the local monodromies  $\mu_a$  to elements of  $Sp(4, \mathbb{Z})$ . Now the product  $\mu_1 \cdots \mu_{s+3}$ , which is the identity in  $PSp(4, \mathbb{Z})$ , may lift to either +1 or −1 in  $Sp(4, \mathbb{Z})$ . In the second case we may add a dummy special point  $p_{s+4}$  with local monodromy  $\mu_{s+4} = -1$  and reduce back to the standard case

<span id="page-81-1"></span>
$$
\mu_1 \cdots \mu_{s+3} \mu_{s+4} = 1. \tag{8.3}
$$

Alternatively we may flip the sign  $\mu_{s+3} \to -\mu_{s+3}$  to get back to the usual story.

The case of main interest is the rigid one,  $s = 0$ . In this case  $\tilde{\mathscr{P}}$  is the sphere less three points, which we call 0, 1 and  $\infty$ , while  $\pi_1(\tilde{\mathscr{P}}) \simeq F_2$ . The three local monodromies will be written as  $\mu_0$ ,  $\mu_1$ ,  $\mu_\infty$ , satisfying  $\mu_0\mu_1\mu_\infty = \pm 1$  (the sign is always + when  $\varkappa = 1$ ).

### <span id="page-81-2"></span>8.1 Reducible geometries in rank 2

Indecomposable but reducible special geometries are rare and quite subtle.

Suppose our  $\mu$ -monodromy representation M is reducible,  $M = V \oplus W$ , while it contains non-trivial unipotents: then at least one summand, say  $V$ , should be irreducible of dimension 2.<sup>[69](#page-81-0)</sup> Since  $\Omega: M \xrightarrow{\sim} M^{\vee}$  we have three possibilities:

$$
L^{\perp} \stackrel{\text{def}}{=} \{ v \in M : v^t \Omega \, l = 0 \ \forall l \in L \} \supset L
$$

<span id="page-81-0"></span><sup>&</sup>lt;sup>69</sup> Proof. Suppose  $L \subset M$  is an invariant subspace of dimension 1. Then

is also invariant of dimension 3. Since the monodromy is totally reducible (Deligne theorem [\[136\]](#page-156-3))  $L^{\perp} = V \oplus L$ for  $V$  an invariant subspace of dimension 2. We claim that  $V$  is irreducible. Indeed, otherwise, it would be the direct sum of two invariant spaces of dimension 1, and the monodromy representation would be the direct sum of 4 one-dimensional ones, hence Abelian, which is a contradiction since the monodromy group

- 1.  $V^{\vee} \simeq W$  and  $\Omega: V \xrightarrow{\sim} W^{\vee}$ ;
- 2.  $V^{\vee} \not\cong W$  and  $\Omega: V \xrightarrow{\sim} V^{\vee}$ ,  $\Omega: W \xrightarrow{\sim} W^{\vee}$ ;
- 3.  $V^{\vee} \simeq V \simeq W$  and  $\Omega: V \xrightarrow{\sim} V^{\vee}$ .

In the first case the monodromy is contained in  $GL(2, \mathbb{Q}) \subset Sp(4, \mathbb{Q})$ , in the second one is a subgroup of  $SL(2,\mathbb{Q}) \times SL(2,\mathbb{Q})$ , and in the third situation in the diagonal  $SL(2,\mathbb{Q})$ . Case 3. is a special instance of both 1. and 2.

#### 8.1.1 Case 1

In case 1. (so also in 3.) we have a symmetric Hodge 2-tensor, hence an element of the chiral ring of dimension 2, i.e. the special geometry *cannot be rigid*. According to the **Folk**theorem, the geometry should describe a Lagrangian SCFT whose list is known. We shall see later in this section that all geometries describing rank-2 SCFTs with  $s+3 = \dim M$  are irreducible. It remains the *mysterious* model  $Sp(4)$  with  $\frac{1}{2}$ **16** (#69 of [\[89\]](#page-153-0)) whose geometry requires further analysis.

#### 8.1.2 Case 2

In case 2. the monodromy is contained in  $SL(2,\mathbb{Q}) \times SL(2,\mathbb{Q})$  so, going to a finite cover if necessary, we have a fibration with fiber the product of two elliptic curves  $E_1 \times E_2$ , that is, the fibered product of two elliptic fibrations over  $\mathbb{P}^1$  less some points which we may replace by the smooth surfaces given by their respective Kodaira-Néron models<sup>[70](#page-82-0)</sup>  $\pi_a: \mathcal{E}_a \to \mathbb{P}^1$   $(a = 1, 2)$ 

<span id="page-82-1"></span>

If  $\mathcal{E}_a$  are non-trivial fibrations, the monodromy group has Zariski-dense images in each factor of  $SL(2,\mathbb{Q}) \times SL(2,\mathbb{Q})$ , so either we are back to case 3. or there is no operator of dimension 2, and the geometry is rigid, so we have exactly 3 special points on  $\mathbb{P}^1$ . Since 1, 2 are not Coulomb dimensions,  $d_1, d_2 \neq 1$ , and hence the axes in  $\mathscr C$  gives two special points  $(1:0)$  and  $(0:1)$ . We remain with only another special point, around which – generically – only the monodromy in one of the two  $SL(2,\mathbb{Q})$  factors is non-trivial. Therefore the other elliptic surface is isotrivial. An isotrivial family is not rigid but in the present case it may be rigidified by the extension on a special point if the monodromy around that point is

contains non-trivial unipotents.

<span id="page-82-0"></span><sup>70</sup> See [\[86\]](#page-153-1).

constrained to be  $\neq \pm 1$  say because the special point is regular enhanced or with a non-trivial semisimple monodromy. Then either:

- $(i)$  we are not in the generic situation, i.e. we are in case 3 which is a particular case of 1;
- $(ii)$  the geometry is quasi-isotrivial.

#### 8.1.3 Scenario  $(ii)$ : quasi-isotrivial geometries

In this case in the diagram  $(8.4)$  the second rational elliptic surface  $\mathcal{E}_2$  is isotrivial while the first one  $\mathcal{E}_1$  is non-isotrivial.

The non-rigid geometries are known not to be quasi-isotrivial except for the still *myste*rious one which deserves a separate analysis. Here we assume the quasi-isotrivial geometry to be rigid. Both dimensions  $\Delta_1, \Delta_2$  should be old and not 1 or 2 and  $\varkappa \in \{1,2\}$ . Thus the order of  $1/\Delta_1$  and  $1/\Delta_2$  should be in  $\{3,4,6\}$  and not both divisible by 4 or 3. Then one dimension should be either 4 or  $4/3$ , but the second possibility is ruled out by the values of the  $d_i$ . We remain with  $\{\Delta_1, \Delta_2\} = \{4, 3\}$  or  $\{4, 6\}$  depending on  $\varkappa = 1$  or 2.

We have 3 special points in  $\mathbb{P}^1$ . Two special points are  $p_0 \equiv (0:1)$  and  $p_{\infty} \equiv (1:0)$ corresponding to the axes and then we have a third 'knotted' special point with monodromy of the form  $\alpha_1 \oplus \mathbf{1}_2$ . The formulae for local regularity along the axes, yield *one* exponent of the local  $\mu$ -monodromy at each of the points  $p_0$ ,  $p_\infty$  regardless of the Kodaira type of the axes as discriminant components. For the present two allowed dimension pairs the statement is simply that  $1/\Delta_i$  is an exponent for the local monodromy around the special point corresponding to an axis of dimension  $\Delta_i = 3, 4, 6$ .

Since a non-isotrivial family should have at least 3 non-trivial local holonomies,  $\alpha_1$  is a local monodromy for  $\mathcal{E}_1$ , while  $\mathcal{E}_2$  has only two special fibers over  $(1:0)$  and  $(0:1)$  and their local  $\mu$ -monodromy,  $\beta$  and  $\beta^{-1}$ , are the inverse of each others so  $\mathcal{E}_2$  is an isotrivial rational elliptic surface with a singular fibers' configuration of the forms  $\{F, F^*\}$  where  $F = II, III,$ or IV. The local monodromies of  $\mathcal{E}_1$  are  $\alpha_0$ ,  $\alpha_1$  and  $\alpha_\infty$  with  $e(\alpha_0) + e(\alpha_1) + e(\alpha_\infty) = 12$ where  $e(\alpha)$  stands for the Euler characteristics of a Kodaira fiber with local monodromy  $[\alpha] \in SL(2, \mathbb{Z})$ . The *ρ*-monodromy

$$
(\alpha_0 \oplus \beta)^{d_1} = \begin{cases} \epsilon_0 (\mathbf{1} \oplus K_0) \\ \epsilon_0 (K_0 \oplus \mathbf{1}), \end{cases} \qquad (\alpha_\infty \oplus \beta^{-1})^{d_2} = \begin{cases} \epsilon_\infty (\mathbf{1} \oplus K_\infty) \\ \epsilon_\infty (K_\infty \oplus \mathbf{1}) \end{cases} \tag{8.5}
$$

where  $K_0$ ,  $K_\infty$  are the local Kodaira monodromies along the axes in  $\mathscr C$  and  $\epsilon_0$ ,  $\epsilon_\infty$  signs from the lift  $(\epsilon_0 = \epsilon_\infty = +1$  when  $\varkappa = 1)$ . A simple very preliminary "regularity" argument leads to the following

Fact 20. All rigid non-isotrivial quasi-isotrivial rank-2 geometries – IF THEY EXIST – should be one of the following: $71$ 

<span id="page-83-0"></span><sup>&</sup>lt;sup>71</sup> Not to confuse the reader, we recall that we are assuming that the Coulomb branch  $\mathscr C$  is smooth.

- (1) Coulomb dimensions  $\{3, 4\}$ , the axis of dimension 3 is a discriminant of type  $I_0^*$  and the axis of dimension  $\lambda$  is a discriminant of type IV<sup>\*</sup>. In addition we have a knotted discriminant of type  $I_m$   $(m > 0)$ ;
- (2) Coulomb dimensions  $\{3,4\}$ , the axis of dimension 3 is a discriminant of type III<sup>\*</sup>, the axis of dimension 4 is a discriminant of type  $I_n$   $(n > 0)$ , and the knotted discriminant has type  $I_m$   $(m > 0)$ ;
- (3) Coulomb dimensions  $\{4, 6\}$ , the axis of dimension 4 is a discriminant of type  $II^*$ , the axis of dimension 6 is a discriminant of Kodaira type  $I_n$  with  $n > 0$ , and we have a knotted discriminant of type  $I_m$   $(m > 0)$ ;
- (4) Coulomb dimensions  $\{4, 6\}$ , the axis of dimension 4 is a discriminant of type  $II^*$ , the axis of dimension 6 is a discriminant of type  $I_0^*$ , and we have a knotted discriminant of type  $I_m$   $(m > 0)$ .

All statements are shown in appendix [F](#page-146-0) except for the claim about the knotted discriminant that follows from considerations about the hypergeometric Picard-Fuchs ODE (see below). The 4 geometries are very similar. For future reference we list the eigenvalues of the  $\mu$ -monodromy at the points  $(1:0)$  and  $(0:1)$  (cf. appendix [F\)](#page-146-0):

	(2)	(3)	$\left(4\right)$	
			(1 : 0) $\left\{e^{\pm 2\pi i/3}, e^{\pm 2\pi i/6}\right\}$ $\left\{e^{\pm 2\pi i/3}, e^{\pm 2\pi i/4}\right\}$ $\left\{e^{\pm 2\pi i/4}, e^{\pm 2\pi i/6}\right\}$ $\left\{e^{\pm 2\pi i/4}, e^{\pm 2\pi i/6}\right\}$ (8.6)	
$(0:1)$ $\left\{e^{\pm 2\pi i/4}, e^{\pm 2\pi i/3}\right\}$ $\left\{e^{\pm 2\pi i/4}, e^{\pm 2\pi i/2}\right\}$ $\left\{e^{\pm 2\pi i/6}, e^{\pm 2\pi i/2}\right\}$ $\left\{e^{\pm 2\pi i/6}, e^{\pm 2\pi i/3}\right\}$				

Remark 12. In this section we shall return again on the intriguing possibility of the existence of quasi-isotrivial geometries from different viewpoints. We shall get further restrictions on them. There is evidence from physics [\[89\]](#page-153-0) that the geometry (3) indeed exists and represents a known SCFT.

From now on we shall (mainly) focus on *irreducible* (principal) rank-2 special geometries.

## 8.2 Review of the Deligne-Simpson problem

The Deligne-Simpson problem [\[137–](#page-156-4)[141\]](#page-156-5) asks for the determination of the irreducible degreen representations of the fundamental group of the sphere  $\mathbb{P}^1$  less  $\ell$  points  $\{p_1, \ldots, p_\ell\}$  with prescribed conjugacy classes  $C_a \subset GL(n, \mathbb{C})$  of the local monodromies at the marked points. The moduli space of such representations is given by

$$
\boldsymbol{m} = \left\{ \mu_1, \ldots, \mu_\ell \in GL(n, \mathbb{C}) : \mu_1 \cdots \mu_\ell = 1, \ \mu_a \in C_a \ \forall a, \ \lim_{\text{common invariant subspace}} \right\} / \sim \ (8.7)
$$

where  $\sim$  stands for equivalence up to overall conjugacy in  $GL(n,\mathbb{C})$ . The representation is rigid iff  $m$  is non-empty and zero dimensional.

The Deligne-Simpson problem was solved by W. Crawley-Boevey by relating it to the representation theory of quivers, see  $[128-133]$  $[128-133]$ .<sup>[72](#page-85-0)</sup> The theory is reviewed in the physical language of Seiberg duality in [\[134\]](#page-156-1). To summarize the story, we start by describing a convenient way to encode the conjugacy classes  $C_a \subset GL(n,\mathbb{C})$  of a  $\ell$ -tuple  $\{\mu_a\}_{a=1}^{\ell}$  of  $n \times n$  matrices. For each marked point  $p_a$  we choose a polynomial  $P_a(z)$  which is solved by the matrix  $\mu_a$ , i.e. such that  $P_a(\mu_a) = 0$ . Note that the polynomial depends only on the conjugacy class of  $\mu_a$ . While we can choose any polynomial with this property, for definiteness we assume  $P_a(z)$  to be *minimal*. Then we factorize  $P(z)$  into linear factors  $P_a(z) = \prod_{i=1}^{m_a} (z - \xi_{a,i})$  choosing an order between them. Now we set

$$
E_{a,k} \stackrel{\text{def}}{=} \prod_{i=1}^{k} (\mu_a - \xi_{a,i}) \mathbb{C}^n, \quad N_{a,k} \stackrel{\text{def}}{=} \dim E_{a,k} \quad a = 1, \dots, \ell, \quad k = 0, \dots, m_a. \tag{8.8}
$$

When the polynomial  $P_a(z)$  is minimal the non-negative integers are strictly-decreasing

$$
N_{a,k} < N_{a,k-1},\tag{8.9}
$$

and we choose the order of the factors in  $P_a(z)$  such that the integers  $N_{a,k}$  take their smallest possible values (we call this the minimal form). This choice is not necessary, but convenient. Then the conjugacy class  $C_a$  of  $\mu_a$  in  $GL(n,\mathbb{C})$  is encoded in the two sequences

$$
0 \equiv N_{a,m_a} < N_{a,m_a-1} < \cdots < N_{a,1} < N_{a,0} \equiv n, \quad \text{and} \quad \xi_{a,1}, \ldots, \xi_{a,m_a} \tag{8.10}
$$

indeed from these data we recover the Jordan-block structure of  $\mu_a$ . Henceforth we write  $\{(N_{a,i}, \xi_{a,i})\}\$ for the datum  $\{C_a\}$ . A conjugacy class  $C_a$  consisting only the element  $\lambda$  1  $(\lambda \in \mathbb{C}^{\times})$  will be called "dummy".  $m_a = 1$  if and only if  $C_a$  is dummy. In our application the local monodromy belongs to a dummy class only if it corresponds to the extra special point with local monodromy −1 arising from the lift of the monodromy representation from  $PSp(4, \mathbb{Z})$  to  $Sp(4, \mathbb{Z})$  in rank-2 geometries with  $\varkappa = 2$  (cf. eq.[\(8.3\)](#page-81-1)).

To the Deligne-Simpson datum  $\{(N_{a,i}, \xi_{a,i})\}$  one associates a star-shaped graph with one

<span id="page-85-0"></span><sup>72</sup> See [\[100\]](#page-154-1) for a textbook account.

branch for each class  $C_a$  containing  $m_a-1$  nodes (not counting the central one  $\star$ ) [\[128–](#page-155-1)[133\]](#page-156-6):



which we see as the Dynkin graph of a simply-laced Kac-Moody Lie algebra  $\mathfrak{G}$ . We write  $\alpha_{\star}$ and  $\alpha_{\bullet_{a,i}}$  for (respectively) the simple root associated to the node  $\star$  and  $\bullet_{a,i}$  while  $\Delta^+(\mathfrak{G})$ stands for the set of positive roots of  $\mathfrak{G}$ . Note that a "dummy" class yields a zero-lenght branch which has no effect on the definition of  $\Delta^+(\mathfrak{G})$  but will enter in the definition of the sub-sets  $R_+$  and  $\Sigma$  below. For  $\alpha$  a positive root

<span id="page-86-0"></span>
$$
\alpha = k_{\star} \alpha_{\star} + \sum_{a=1}^{\ell} \sum_{i=1}^{m_a - 1} k_{a,i} \alpha_{\bullet_{a,i}} \in \Delta^+(\mathfrak{G}), \qquad k_{\star}, k_{a,i} \in \mathbb{Z}_{\geq 0}
$$
(8.12)

we set

$$
d(\alpha) \stackrel{\text{def}}{=} 1 - \frac{1}{2} \langle \alpha, \alpha \rangle \equiv 1 - k_{\star}^2 + \sum_{a} \left( \sum_{i=1}^{m_a - 1} k_{a,i}^2 + \sum_{i=1}^{m_a - 1} k_{a,i} k_{a,i-1} \right), \tag{8.13}
$$

where  $\langle -, - \rangle$  is the integral quadratic form on the root lattice of  $\mathfrak{G}$  given by the Cartan matrix, and we use the convention  $k_{a,0} \equiv k_{\star}$  for all a. The integer  $d(\alpha)$  is non-negative for all roots and zero iff the root  $\alpha$  is real. We write  $R_+ \subset \Delta^+(\mathfrak{G})$  for the sub-set of positive roots [\(8.12\)](#page-86-0) such that the complex number

$$
\xi(\alpha) \stackrel{\text{def}}{=} \prod_{a=1}^{\ell} \xi_{a,1}^{k_{\star}} \prod_{i=1}^{m_{a}-1} \left( \xi_{a,i}^{k_{a,i-1}} / \xi_{a,i}^{k_{a,i}} \right)
$$
(8.14)

is equal 1, that is,

$$
R_{+} \stackrel{\text{def}}{=} \left\{ \alpha \in \Delta^{+}(\mathfrak{G}) : \xi(\alpha) = 1 \right\}.
$$
 (8.15)

Note that adding the "dummy" class  $\lambda \mathbf{1}$  has the effect  $\xi(\alpha) \rightsquigarrow \lambda^{k_{\star}} \xi(\alpha)$ . For convenience we rewrite  $\xi(\alpha)$  in the form

$$
\xi(\alpha) = \lambda_{\star}^{k_{\star}} \prod_{a} \left( \prod_{i=1}^{m_{a}-1} \lambda_{a,i}^{k_{a,i}} \right)
$$
\n(8.16)

and call  $\lambda_{\star} = \prod_a \xi_{a,1}$  and  $\lambda_{a,i} = \xi_{a,i+1}/\xi_{a,i}$  the *fugacity* of the quiver node  $\star$  and respectively  $\bullet_{a,i}$ . We specify the function  $\xi(\cdot)$  by drawing the quiver with the values of the fugacities written over the respective nodes.

**Definition 16.**  $\Sigma \subset R_+$  is the subset of positive roots  $\alpha$  such that for all nontrivial decomposition  $\alpha = \beta_1 + \cdots + \beta_s$  with  $\beta_j \in R_+$  we have

$$
d(\alpha) > \sum_{i} d(\beta_i). \tag{8.17}
$$

Theorem (Crawley-Boevey Theorems 1.1, 1.9 of [\[130\]](#page-156-7)). There is an irreducible representation of degree n with local monodromies in the given conjugacy classes  $\{(N_{a,i}, \xi_{a,i})\subset$  $GL(n,\mathbb{C})\},\;{\it iff}$ 

$$
\mathbf{N} = n \alpha_{\star} + \sum_{a} \sum_{i} N_{a,i} \alpha_{\bullet_{a,i}} \in \Sigma.
$$
 (8.18)

In this case dim  $m = 2 d(N)$  and hence an irreducible representation is rigid iff  $N \in \Sigma$  is a real *root* of the Kac-Moody algebra  $\mathfrak{G}$ .

**Remark 13.** When  $\alpha \in \Sigma$  admits a non-trivial decomposition in  $R_+$  we have both reducible and irreducible representations with the given local classes  $C_a$ . When  $\alpha$  is a real root in  $R_+$ either all representations are irreducible or all reducible.

**Remark 14.** If the graph  $\mathfrak{G}$  is the Dynkin graph of a finite-dimensional, affine, or hyperbolic Kac-Moody algebra, the condition  $d(\alpha) = 0$  (resp.  $d(\alpha) \geq 1$ ) implies that  $\alpha$  is a real root (resp. imaginary root).

## 8.3 Irreducible monodromies in  $Sp(4,\mathbb{R})$

With the possible exceptions discussed in  $\S$ . [8.1,](#page-81-2) all smooth  $\mathbb{C}^{\times}$ -isoinvariant, indecomposable, non-isotrivial, rank-2 geometries are irreducible with 3, 4 or 5 special points for, respectively, rigid geometries, geometries with one dimension 2, and geometries with two dimensions 2. The associated Dynkin graphs are stars with 3, 4, or 5 branches of the following kinds

<span id="page-87-0"></span>

where the circled number on the  $(a, i)$ -th node is  $N_{a,i}$ . Note that a  $\delta = 1$  branch can correspond only to a non-enhanced divisor of type  $I_n$   $(n > 0)$ .

The "virtual" dimension of the moduli space  $m$  of irreducible representations whose graph has s branches of kinds  $\{\delta_1, \ldots, \delta_s\}$  is  $2 d(\delta_1, \ldots, \delta_s)$  where

$$
d(\delta_1, ..., \delta_s) = 2s + \sum_{\ell=1}^s \delta_{s} - 15
$$
 (8.20)

and this quantity should be non-negative (zero in the rigid case). In the addition we have the other conditions in the Theorem which depend on the eigenvalues of the local monodromies.

#### 8.3.1 The conjugacy classes which may appear

We list the conjugacy classes in  $GL(4,\mathbb{C})$  which may appear. To the a-th special point in  $\mathscr{P} \simeq \mathbb{P}^1$  there corresponds a branch of the quiver of one of the four kinds [\(8.19\)](#page-87-0) with their fugacities.

In the case  $\varkappa = 2$  only the image of the monodromy in  $PSp(4, \mathbb{Z})$  is non-ambiguous, so its conjugacy class is determined up to overall sign. However the integers  $N_{a,i}$  are independent of the sign, as are the fugacities  $\lambda_{a,i} \equiv \xi_{a,i+1}/\xi_{a,i}$  of all nodes except the central one. Thus the only ambiguity is the sign of  $\lambda_{\star}$ . Therefore when  $\varkappa = 2$  we have to explore the two possible signs of the central fugacity  $\lambda_{\star}$ .

Discriminant component non-enhanced. In this case  $[\mu_i] = [\varrho_i] = [\sigma_i \oplus \mathbf{1}_2]$  with  $\sigma_i \in SL(2,\mathbb{Z})$  a Kodaira monodromy, so

<span id="page-88-0"></span>

**Enhanced divisor non-discriminant.** Suppose that  $u_1 = 0$  is not part of the discriminant. The local coordinate around the corresponding special point in  $\mathbb{P}^1$  is  $z = u_1^{d_2}/u_2^{d_1}$ . A path from  $(\epsilon, u_2)$  to  $(e^{2\pi i/d_2}\epsilon, u_2)$  covers a closed loop encircling  $u_1 = 0$ . Hence the local  $\mu$ -monodromy  $\mu$  satisfies

<span id="page-89-0"></span>
$$
\mu^{d_2} = \begin{cases} 1 & \text{for } \varkappa = 1 \\ \pm 1 & \text{for } \varkappa = 2. \end{cases} \tag{8.22}
$$

Since the central fiber in the covering family  $\mathscr X$  is a smooth Abelian fiber  $A_{u_1=0}$ , the monodromomy  $\mu$  should be the rational representation of an element of the automorphism group of  $A_{u_1=0}$ . The element  $\eta = \exp(2\pi i \mathcal{E}/\Delta_2) \in \text{Aut}(A_{u_1=0})$  satisfies

<span id="page-89-2"></span>
$$
\sigma(\eta)^{d_2} = \sigma(\exp(2\pi i \mathcal{E}/\lambda)) = \exp(2\pi i/\varkappa) \cdot \mathbf{1},\tag{8.23}
$$

so that, modulo the sign when  $\varkappa = 2$ , a solution to [\(8.22\)](#page-89-0) is  $\mu = \chi(\eta)$  whose eigenvalues are

<span id="page-89-1"></span>
$$
\left\{ \exp\left[\pm 2\pi i(1-\Delta_1)/\Delta_2\right], \exp\left[\pm 2\pi i(1-\Delta_2)/\Delta_2\right] \right\}.
$$
 (8.24)

When  $\Delta_2$  is a new dimension this is the only solution to [\(8.22\)](#page-89-0) up to equivalence. In this case the 4 eigenvalues are distinct and we get a  $\delta = 4$  branch of  $\mathfrak{G}$ . Otherwise one can find in principle other automorphisms of order  $d_2$  which may play the role of  $\mu$ . This seems unlikely, and we shall assume the equality  $\mu = \chi(\eta)$  in general. If the axis  $u_2 = 0$  does not belong to the discriminant, the same formulae apply with  $1 \leftrightarrow 2$ .

 $\mu$  is semisimple and integral, so a part for the dummies  $\pm 1$ , it may have

- 2 distinct eigenvalues both of multiplicity 2 of the form  $(+1, -1)$  or  $(e^{2\pi i/m}, e^{-2\pi i/m})$  $(m = 3, 4, 6)$  which lead to a  $\delta = 2$  branch,
- 3 distinct eigenvalues of multiplicites  $(2,1,1)$  of the form  $(\pm 1, \epsilon^{2\pi i/m}, e^{-2\pi i/m})$  (m = 3, 4, 6) which lead to a  $\delta = 3$  branch with fugacities

$$
\cdots \longrightarrow e^{2\pi i/m} \longrightarrow e^{-4\pi i/m} \tag{8.25}
$$

• in all other cases we have a  $\delta = 4$  branch.

Eigenvalues  $\pm 1$  are expected only when we have a regular axis with  $\Delta = 2$  which is rather uncommon. Hence (possibly with a few exceptions) we expect that only  $\delta = 2$  and  $\delta = 4$ branches appear at an enhanced non-discriminant orbifold point.  $\delta = 2$  means that the four phases [\(8.24\)](#page-89-1) are equal in pairs. This requires

$$
\frac{1-\Delta_1}{\Delta_2} = \pm \frac{1}{\Delta_2} \mod 1\tag{8.26}
$$

i.e. either (i)  $\Delta_1 = m\Delta_2$  which is possible only for  $\Delta_2 = 2$  and  $m > 1$ , or (ii)  $\Delta_1 = 2 + m\Delta_2$ where  $\Delta_2$  is an old dimension. In all other cases we have a  $\delta = 4$  branch.

**Remark 15.** The argument also shows that  $\phi(d_i) \leq 4$  for a rank-2 theory.

**Enhanced divisor in the discriminant.** We suppose that  $u_1 = 0$  is both enhanced and part of the discriminant. Arguing as around eq.[\(8.23\)](#page-89-2), we conclude that the monodromy  $\mu$  around the corresponding special point  $z = 0$  in  $\mathbb{P}^1$  is related to the monodromy  $\varrho$  around the divisor  $(u_1) \subset \mathscr{C}$  by

$$
\mu^{d_2} = \begin{cases} \varrho & \mathcal{U} = 1 \\ \pm \varrho & \mathcal{U} = 2. \end{cases} \tag{8.27}
$$

We need to determine  $\mu$  up to conjugacy in  $GL(4,\mathbb{C})$ . As argued in §. [2.9,](#page-32-0)  $\mu$  induces an action on the homology of the Albanese curve of the singular fiber which must be the rational representation of an automorphism, which (as in the previous paragraph) has eigenvalues  $\exp(\pm 2\pi i/\Delta_2)$  (up to the usual sign ambiguity when  $\varkappa = 2$ ). This is consistent since  $\Delta_2$ is necessarily a rank-1 dimension. The other two eigenvalues are determined by  $d_2$  and the Kodaira type of the singular fiber. If  $\rho$  is non-semisimple, so should be  $\mu$ , and hence these two eigenvalues must be equal, hence  $\pm 1$  since the determinant of  $\mu$  is 1. In particular

**Fact 21.** Assume  $\varkappa = 1$ . (1) if  $d_2$  is even the divisor  $u_1 = 0$  cannot be of type  $I_n^*$  with  $n > 0$ . (2) the only allowed semisimple types are:

 $d_2 = 2$ :  $I_0^*, IV, IV^*$  $d_2 = 3$ :  $I_0^*$ ,  $III$ ,  $III^*$  $d_2 = 4$ : IV, IV\*  $d_2 = 6$ :  $I_0^*$ 

When  $x = 2$  no a priori restriction on the non-semisimple fibers, while for the semisimple fibers we have the same table as above with IV,  $IV^*$  replaced by IV,  $IV^*$ , II and  $II^*$ .

No counterexample to these statements appears in the tables of [\[89\]](#page-153-0). The statement eliminates an ambiguity in the analysis of model #37 in that reference.

When the fiber type is additive, the other two eigenvalues besides  $\exp(\pm 2\pi i/\Delta_2)$  are related to the Kodaira type of the  $\varrho$ -monodromy by the base change formula (modulo the sign ambiguity when  $\varkappa = 2$ ) which may be read e.g. in table 5.2 of [\[86\]](#page-153-1). Since  $d_2 > 1$  for an enhanced discriminant, when the fiber is semisimple the two eigenvalues are distinct.

In conclusion: an enhanced discriminant  $u_1 = 0$  produces always a  $\delta = 4$  branch of the Dynkin graph except when  $\Delta_2 = 2$  which produces a  $\delta = 3$  graph.

## 8.4 Rank-2 special geometries with rigid  $\mu$ -monodromy

The classification/construction of non-isotrivial irreducible special geometries with rigid monodromy is relatively straightforward in rank-2. This would produce the full list of nonisotrivial irreducible geometries if the  $\mu$ -rigidity conjecture is true, and a large portion of them otherwise. The results below will strongly support the conjecture.

#### 8.4.1 Goursat rigid monodromies

In order 4 there are six Fuchsian ODEs with rigid monodromy. They were already discussed by Goursat in 1886 [\[142,](#page-156-8) [143\]](#page-156-9). Writing all conjugacy classes in their minimal form, the corresponding pairs (Dynkin graph, real root) are as in table [1.](#page-92-0) Some of these graphs describe classically well-known Fuchsian OPEs:

- (a) is the order-4 hypergeometric ODE of Thomae;
- (d) is a reduction to one-variable of the 2-variable Appell hypergeometric PDE, aka as the Okubo equation  $II^*$
- (f) is the order-4 (Jordan-)Pochhammer ODE.

Remark 16. The graph corresponding to 3 unipotent monodromies and one with minimal polynomial of degree 4



has  $d(\alpha) = 0$ , so superficially seems to be a rigid solution to the Deligne-Simpson problem, but it is not so since the Kac-Moody algebra is not hyperbolic and  $\alpha$  is not a root. We conclude that these conjugacy classes cannot appear.

#### 8.4.2 Graph selection rules

We note that  $\xi(\alpha)$  remains invariant if we cancel one or more  $\delta = 1$  branches. This fact eliminates some possibilities thus further restricting the allowed geometries.

<span id="page-91-0"></span>Fact 22. In a non-isotrivial, irreducible, rank-2 geometry:

- $(1)$  graph (c) is excluded;
- (2) graph (e) is excluded;
- (3) graph (f) is excluded when  $\varkappa = 1$ .

*Argument.* Part  $(1)$ . At least one branch is non-semisimple, so the fugacities are as follows





<span id="page-92-0"></span>Figure 1: The six order 4 rigid ODEs: graphs and roots in minimal form. The first graph is the order-4 hypergeometric, the 4-th one is the Appell equation (an ODE truncation of the 2-variable Appell PDE), the fifth one is a particular instance of Simpson equation, and the last one is the order-4 Pochhammer equation.

where  $\zeta$  and  $\xi$  are either  $-1$  (if the branch is non-semisimple) or a root of 1 of degree 2 (if it is semisimple). As before  $\pm$  corresponds to the two possible lifts in  $Sp(4)$  when  $\varkappa = 2$ . If the sign is + the simple root at the central node  $\alpha_{\star} \in R_+$  and  $\alpha - \alpha_{\star}$  is the minimal imaginary root of  $\widehat{E}_6$  which also belongs to  $R_+$ . Since  $d(\alpha) = 0$  the representation is reducible. Suppose now that the sign is −: we have the decomposition

$$
\begin{array}{cccc}\n0 & 1 & 1 \\
1 & 2 & 3 & 2 & 1\n\end{array} + \begin{array}{cccc}\n1 & 1 & 0 & 0 \\
0 & 0 & 1 & 0 & 0\n\end{array}
$$
\n(8.29)

into roots of the  $E_6$  and  $A_3$  subgraphs which also belong to  $R_+$  and again the representation is reducible.

**Part (2).** The reader may find in appendix [B](#page-141-0) the proof of the following

**Lemma 2.** A monodromy representation associated to the graph (e), whose local monodromies are in the allowed lists, is irreducible only when  $\varkappa = 2$  and the three semisimple monodromies have eigenvalues

<span id="page-93-1"></span>
$$
(1, 1, -1, -1), \quad (i, i, -i, -i), \quad (\pm \zeta_3, \pm \zeta_3, \pm \zeta_3^{-1}, \pm \zeta_3^{-1}), \tag{8.30}
$$

where  $\zeta_3$  is a primitive 3-rd root of 1.<sup>[73](#page-93-0)</sup>

If it exists, an irreducible geometry based on graph (e) should have a conformal manifold of dimension 1 hence dimensions  $\{2, \Delta\}$ , with  $\Delta$  an *old* dimension  $\neq 2$ . To have  $\varkappa = 2$  we must have  $\Delta \in \{6/5, 4/3, 4, 6\}$ . When  $\Delta$  is an integer = 4,6 there is only one enhanced divisor, and the spectra [\(8.30\)](#page-93-1) are inconsistent. When  $\Delta = 6/5$  we have  $(d_1, d_2) = (5, 3)$ and the eigenvalues [\(8.30\)](#page-93-1) are inconsistent. We remain with  $\{\Delta_1, \Delta_2\} = \{2, 4/3\}$  which has  $(d_1, d_2) = (3, 2)$ . But then the axis of dimension  $\Delta_1$  has a monodromy  $\rho = \pm \mu_1^3 = \pm 1$  and hence is not part of the discriminant, so

$$
\exp(\pm 2\pi i/\Delta_1), \qquad \exp(\pm 2\pi i(1-\Delta_2)/\Delta_1), \tag{8.31}
$$

should be all square roots of 1. This is not the case.

**Part (3).** When  $\varkappa = 1$  there is no extra sign from the lift from  $PSp(4, \mathbb{Z})$  to  $Sp(4, \mathbb{Z})$ while all local monodromies in graph (f) are unipotent, so that the group homomorphism

$$
\xi(\alpha): \bigoplus_{a \in \mathfrak{G}} \mathbb{Z} \alpha_a \to \mathbb{C}^\times \tag{8.32}
$$

is the trivial one, and the only roots in  $\Sigma_{\xi}$  are the ones in the fundamental domain (which are never real!) and the simple roots (cf. §. 5 of [\[128\]](#page-155-1)). When  $\varkappa = 2$  we have the central node  $\star$ may have fugacity  $\zeta_{\star} = -1$ , so that now  $R_{+}$  is the set of positive roots of the form  $2k\alpha_{\star} + \cdots$ . It is easy to check that no decomposition of the root in (f) into elements of  $R_+$  exists. Alternatively, since the ODE associated to graph (f) is the classical order-4 Pochhmmer

<span id="page-93-0"></span><sup>73</sup> This is the first example in [\[144\]](#page-156-10). I thank V. Kostov for clarifications.

Remark 17. The argument for part (2) is greatly simplified if one assumes the Folktheorem (which we argued is true in rank-2). A non-isotrivial geometry based on graph (e) should be the geometry of a Lagrangian SCFT, which were classified [\[95\]](#page-154-3), and no SCFT in that list corresponds to graph (e). Indeed a Lagrangian SCFT has integral  $\Delta_i$  which is incompatible with (e) on the nose.

**Remark 18.** According to the **Folk-theorem** graph (f) corresponds to dimensions  $\{2, 2\}$ which has automatically  $\varkappa = 2$  as predicted by rigidity and irreducibility.

**Fact 23.** In graph (a) we assume the relation  $\mu_1\mu_2\mu_3 = 1$  ( $\mu_1$  corresponds to the  $\delta = 1$ branch) by absorbing the possible extra sign from the lift from  $PSp(4, \mathbb{Z})$  to  $Sp(4, \mathbb{Z})$  when  $\varkappa = 2$  in the definition of  $\mu_3$ . With this convention, the matrices  $\mu_2$  and  $\mu_3$  cannot have any common eigenvalue.

Argument. (1) Suppose that they have the eigenvalue  $\eta$  in common. Since the monodromies are symplectic, also  $\eta^{-1}$  is an eigenvalue and by a suitable ordering of the factors in their respective minimal polynomial we get the graph fugacities

$$
\begin{array}{c}\n1 \\
\xi_1^{-2} - \xi_1 \eta^{-1} - \eta^2 - 1 - \eta^{-2} - \xi_2 \eta - \xi_2^{-2}\n\end{array} \tag{8.33}
$$

 $\Box$ 

and the simple root  $\alpha_{\star}$  and the imaginary root  $\alpha - \alpha_{\star}$  belong to  $\Sigma_{+}$ .

## 8.5 From rigid monodromy to special geometry

All non-isotrivial, irreducible,  $\mu$ -rigid rank-2 geometries are associated to one of four Kac-Moody graphs  $(a)$ ,  $(b)$ ,  $(d)$ ,  $(f)$  according to the rule:

- Graph (a): rigid geometries (2 not a dimension) with one "knotted" discriminant of type  $I_n$  for some  $n > 0$ ;
- Graph (b): rigid geometries (2 not a dimension) with one "knotted" discriminant not of type  $I_n$ ;
- Graph (d) geometries with dimensions  $\Delta_1 = 2, \Delta_2 \neq 2;$
- Graph (f) geometries with  $\{\Delta_1, \Delta_2\} = \{2, 2\}.$

We shall say that a  $\mu$ -rigid special geometry is of class  $\mathfrak{G}$  iff its monodromy corresponds to a real root of the Kac-Moody algebra G.

### 8.5.1 The inverse algorithm

The inverse algorithm for rank-2 works through the following steps:

- 1. If the datum  $(\{\Delta_i\}, \{\varrho_a\})$  corresponds to an isotrivial, a decomposable, a reducible, or a covering geometry, the special geometry can either be constructed by direct means or easily proven not to exist;
- 2. Otherwise (assuming the  $\mu$ -rigidity conjecture) either the datum  $(\{\Delta_i\}, \{\varrho_a\})$  corresponds to a graph  $(a)$ ,  $(b)$ ,  $(c)$ , or  $(d)$  or no geometry with the datum exists;
- 3. From the datum one reads the fugacities of the nodes of the graph  $\mathfrak{G}$ . Sometimes there may be an ambiguity, but with only few alternatives. One checks whether one or more of these alternative fugacities define an irreducible monodromy using the characterization of the root set  $\Sigma(\xi)$ . At this point we have a rigid  $GL(4,\mathbb{C})$  monodromy which is known to be in  $Sp(4,\mathbb{C})$  in facts it can be conjugated to lay in  $Sp(4,\mathbb{R})$  or  $Sp(4,\mathfrak{o}_F);$
- 4. We have to check that the Brauer class of the resulting monodromy is trivial, so that the monodromy can be defined over Q;
- 5. At this point we know that the monodromy is integral, rational, and symplectic for a unique skew-symmetric form  $\Omega$ . However we need the symplectic matrix  $\Omega$  to be principal, that is, with integral entries and determinant 1. Thus we have to check that by a simultaneous transformation

$$
\mu_a \leadsto U^{-1} \mu U, \qquad \Omega \leadsto U^t \Omega U, \qquad U \in GL(4, \mathbb{R}) \tag{8.34}
$$

we can set all matrices  $\mu_a$ ,  $\Omega$  to be integral while det  $\Omega = 1$ .

- 6.  $\hat{\mathcal{L}}$  the matrix U with the properties in 5. (if it exists) may be non-unique. Two such matrices U and U' yield equivalent integral monodromies iff  $U'U^{-1} \in Sp(4, \mathbb{Z})$ , otherwise they produce *inequivalent* integral monodromies (all equivalent in the  $GL(4,\mathbb{C})$ -sense).<sup>[74](#page-95-0)</sup> Thus we get *a list* of inequivalent integral monodromies associated to each fugacity assignment. In particular, it is quite frequent to have distinct special geometries arising from the same graph and fugacities which differs because the types  $I_n$  of the semistable discriminants have different values of  $n$  (see below for examples). We stress that the list is finite (an typically quite short) because of Deligne finiteness (or Faltings finiteness in view of 7.);
- 7. For each (rigid) integral monodromy in the list of item 6. we get a candidate  $\mathbb{C}^{\times}$ isoinvariant special geometry. To write it explicitly, we use Riemann's observation that the solutions to all Fuchsian ODE with rigid monodromy have a known explicit

<span id="page-95-0"></span><sup>&</sup>lt;sup>74</sup> This subtle phenomenon is already present in rank-1 where geometries with dimensions m and  $m/(m-1)$ (with  $m = 3, 4, 6$ ) have monodromies conjugates over  $GL(2, \mathbb{R})$  but not over  $SL(2, \mathbb{Z})$ .

integral representation (see [\[102\]](#page-154-0), for a textbook treatment [\[100\]](#page-154-1), for a review in physical language see §. 4 of [\[134\]](#page-156-1)). The Simpson-Deligne integrality implies that these integral representations are motivic i.e. the integrand is given in terms of algebraic functions. The periods of the SW differential  $(a_i^D, a^j)$  are linear combinations of these explicit transcendents, with coefficients uniquely fixed by the Almkvist-Zudilin theorem (§. [4.2\)](#page-51-0). The SW periods of the various geometries associated to a single set of fugacities are linear combinations of each other with coefficients dictated by the matrices U. We stress that, up to equivalence, only finitely many such U may exist by Faltings finiteness. Having constructed the SW periods as multivalued functions of the Coulomb branch coordinates we may check their regularity (which typically gives no problem). If this final text is passed, we declare to have constructed an explicit special geometry. If one wishes to write its equations as a variety, one may read them from the algebraic functions in the integrand of the integral representation of the periods.

8. the special geometry with the original datum  $(\{\Delta_i\}, \{\varrho_a\})$  is either an exceptional one in step 1, or one of the finite list produced in step 7, or it does not exist.

The classification program. Since only finitely many fugacity assignments are possible, we have only a finite list of cases to analyze along the lines of items 3.-7. above. This will produce all non-isotrivial, irreducible,  $\mu$ -rigid geometries thus completing the classification in rank-2 unless – contrary to expectations  $!$  – non-trivial non- $\mu$ -rigid geometries exist.

How to proceed in practice. To work out items 3.-7. one has to construct explicitly the matrices  $\mu_i$ 's realizing the rigid monodromy of the Abelian family over  $\mathscr{\dot{P}}$ . This has been an active field of research for almost two centuries, so a lot is known. Unfortunately, for some of the graphs the results in the literature refer to the simpler case of generic eigenvalues/fugacities whereas we are interested in a highly non-generic case. Thus for some graphs we cannot limit ourselves to search the answer in the existing math literature and novel computations are needed. Then we have to determine the list of relevant matrices U in step 6, and this gives us a "Diophantine" problem which may be not easy to solve. Anyhow a very large portion of the program can be carried on rather explicitly without going into detailed computations as we are going to show.

## 8.6 Checking regularity

Having found a rigid monodromy representation with the prescribed properties we have to check that it describes a regular special geometry. In principle the check splits in two parts:

(1) local checks around the special divisors. One has to check that there exist a local solution to the ODE which are (multivalued) local special coordinates which behave as predicted by the local models of the Lagrangian fibration near a singular fiber of the appropriate Kodaira type, cf. §. [2.9](#page-32-0) . Since the leading behavior of the local solutions is determined by the exponents of the ODE, i.e. the inverse data, these local check are easy and quick (see §. [8.6.1](#page-97-0) below);

(2) if the local checks are satisfied, in principle one has still to check that the "good" solution at one special divisor is the same one which is "good" at all other divisors. This requires the use of the connection formulae which are explicitly known for rigid ODEs [\[100\]](#page-154-1) so the check may be time-consuming but has no fundamental difficulty.

We shall not do checks of type (2) in this paper. We do not expect them to be a major issue: we know from the Almkvist-Zudilin theorem [\[98\]](#page-154-2) that there is a global solution which embeds  $\mathscr{\dot{P}}$  as a Legendre submanifold of  $\mathbb{P}^3$ . The "good" local solutions should agree with this global "good" solution.

#### <span id="page-97-0"></span>8.6.1 Local regularity along the axes

We consider a non-isotrivial rank-2 geometry with dimensions  $\{\Delta_1, \Delta_2\} = \lambda \{d_1, d_2\}$  where  $\lambda$ is written in minimal terms is  $k/\varkappa$  ( $k \in \mathbb{N}$ ) and  $\varkappa \in \{1,2\}$ . When  $\varkappa = 1$  we have an element  $h \in \mathbb{C}(\mathscr{C})$  with  $\Delta(h) = 1$  while for  $\varkappa = 2$  we have an element  $h^2 \in \mathbb{C}(\mathscr{C})$  with  $\Delta(h^2) = 2$ . The standard coordinate on the normalization  $\mathbb{P}^1$  of  $\mathscr{P}$  is  $z = u_1^{d_2}/u_2^{d_1}$  [\[71\]](#page-152-2).

The SW periods have the form

<span id="page-97-1"></span>
$$
a^i = h \cdot f_i(z), \qquad a_j^D = h \cdot f_j^D(z) \tag{8.35}
$$

where  $(f_j^D(z), f_i(z))$  is the solution to the Picard-Fuchs equation selected by the Almkvistwhere  $(f_j(z), f_i(z))$  is the solution to the ricard-ruchs equation selected by the Almitylst-<br>Zudilin theorem. In [\(8.35\)](#page-97-1) h is an abusive notation which stands for h when  $\varkappa = 1$  and  $\sqrt{h^2}$ when  $\varkappa = 2$ . In the latter case the periods are defined only up to overall sign.

We specialize the above expression along the coordinate axes. They correspond to the two special points  $z = 0$  and  $z = \infty$  in  $\mathscr{P}$ . To get the behavior along the axes we use the local solutions of the Picard-Fuchs equations around these special points [\[100\]](#page-154-1). If the monodromy is semisimple, the leading term has the form  $z^{\alpha_{\ell,i}}(1+O(z))$  with  $\alpha_{\ell,i} = \log \xi_{\ell,i}/2\pi i$  and exponent of the local monodromy  $\mu_{\ell}$  at these special points ( $\xi_{\ell,i}$  are the eigenvalues of  $\mu_{\ell}$ as before). If  $\mu_{\ell}$  has non-trivial Jordan blocks the leading term contains logarithms of z; however for a suitable choice of the symplectic basis only the "dual" special coordinates  $a_i^D$ contain logarithms, while the special coordinates  $a^j$  have the local form of a fractional power of z times a holomorphic function.

We consider regularity along the first axis  $(u_2 = 0)$   $(z = \infty)$ . The same discussion applies to the second axis with  $1 \leftrightarrow 2$ . Two (C-linear combination of the) periods,  $a^{\parallel}$  and  $a^{\perp}$  correspond to the special coordinates associated respectively to the first axis and its orthogonal bundle [\[17,](#page-149-0)[22\]](#page-150-0): these two particular periods do not contain logarithms. Comparing with the local models in §. [2.9.2](#page-33-0) we see that regularity along the first axis requires the existence of two exponents  $\alpha_{\parallel}$  and  $\alpha_{\perp}$  of  $\mu_{\infty}$  such that

<span id="page-98-1"></span>
$$
a^{\parallel} = h \cdot f_{\parallel}(z) \sim h \cdot z^{\alpha_{\parallel}} = u_{\ell}^{1/\Delta_1}
$$
  
\n
$$
a^{\perp} = h \cdot f_{\perp}(z) \sim h \cdot z^{\alpha_{\perp}} = u_2^{1/\Delta_K} \cdot u_1^{(1-\Delta_K)/\Delta_1}
$$
\n(8.36)

where  $\Delta_K$  is the dimension of the rank-1 (weak) geometry with the same Kodaira type as the  $\varrho$ -monodromy  $\varrho_1 = \pm \mu_{\infty}^{d_1}$  around the first axis, i.e.

$$
\begin{array}{c|ccccccccc}\n\text{type} & I_n & I_n^* & II & II^* & III & III^* & IV & IV^* \\
\hline\n\Delta_K & 1 & 2 & 6/5 & 6 & 4/3 & 4 & 3/2 & 3\n\end{array} \tag{8.37}
$$

with the convention that an enhanced regular axis is of type  $I_0$ .

 $\overline{\phantom{a}}$ 

#### 8.6.2 Check of regularity: an example

We start by illustrating how regularity of the SW periods works in an non-isotrivial example where we have an enhanced discriminant of semisimple type, we consider the geometry of the SCFT #3 of [\[89\]](#page-153-0) which has dimensions  $\{4, 10\}$ ,  $\varkappa = 2$ , and a class- $\widehat{E}_7$  geometry where the axis of dimension 4 is a component of the discriminant with Kodaira type  $II^*$ . The second axis is regular since it has the new dimension 10. Given that this geometry describes an actual SCFT, we know in advance that the regularity conditions are satisfied: we just want to illustrate how they work. The monodromies of the family over  $\mathring{\mathscr{P}}$  are explicitly

$$
\mu_1 = \begin{pmatrix} 1 & -1 & 0 & 0 \\ 2 & 0 & 1 & 1 \\ 0 & 0 & 0 & -1 \\ -1 & 0 & 0 & 0 \end{pmatrix}, \quad \mu_2 = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & -1 & -1 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \quad \mu_3 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \tag{8.38}
$$

which satisfy

<span id="page-98-0"></span>
$$
(\mu_1^2 + 1)(\mu_1^2 - \mu_1 + 1) = \mu_2^4 + \mu_2^3 + \mu_2^2 + \mu_2 + 1 = 0, \quad \mu_1 \mu_2 \mu_3 = -1,
$$
  

$$
\mu_a^t \Omega \mu_a = \Omega, \quad a = 1, 2, 3 \quad \text{where} \quad \Omega \equiv \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}
$$
 (8.39)

while from the form of  $\mu_3$  we see that there is a knotted discriminant of type  $I_1$ . The  $\varrho$ -monodromy around the axis of dimension 4,  $\varrho = -\mu_1^2$ , satisfies the minimal equation

$$
(\varrho - 1)(\varrho^2 - \varrho + 1) = 0 \tag{8.40}
$$

which is consistent with type  $II^*$  (or  $II$ ). Let us look at the special coordinates as multivalued functions in a neighborhood of the axes. Since  $\varkappa = 2$  they have the form

$$
a^i = \sqrt{h^2} f_i(z) \tag{8.41}
$$

where  $h^2$  is an element of the quotient ring of dimension 2, z is the coordinate on  $\mathbb{P}^1$  and  $f_i$ are solutions to the Picard-Fuchs ODE whose local expressions along the axes scale with the appropriate local exponents. In the present case  $h^2 = u_2/u_1^2$  and  $z = u_1^5/u_2^2$ .

The local exponents along the  $u_2$ -axis  $(u_1 = 0)$  for the special coordinates are  $1/5$  and  $2/5$  (as one sees using  $\mu_2^5 = 1$  and the dimension formulae along a regular axis) so that

$$
a_{\parallel} \sim \sqrt{h} z^{1/5} = \frac{u_2^{1/2}}{u_1} \cdot \frac{u_1}{u_2^{2/5}} = u_2^{1/10}
$$
  
\n
$$
a_{\perp} \sim \sqrt{h} z^{2/5} = \frac{u_2^{1/2}}{u_1} \cdot \frac{u_1^2}{u_2^{4/5}} = u_1 u_2^{-3/10}
$$
  
\nas  $u_1 \to 0$  (8.42)

which are precisely the regular behaviors expected along a regular axis, see [\[22\]](#page-150-0)[\[17\]](#page-149-0). More interesting are the periods along the  $u_1$ -axis which belong to a semisimple component of the discriminant. The local exponents are read from the minimal polynomial of  $\mu_1$ , eq.[\(8.39\)](#page-98-0): they are  $1/4$  and  $1/6$ . Then as  $u_2 \rightarrow 0$ 

$$
a_{\parallel} \sim \sqrt{h} z^{1/4} = \frac{u_2^{1/2}}{u_1} \cdot \frac{u_1^{5/4}}{u_2^{1/2}} = u_1^{1/4}
$$
  
\n
$$
a_{\perp} \sim \sqrt{h} z^{1/6} = \frac{u_2^{1/2}}{u_1} \cdot \frac{u_1^{5/6}}{u_2^{1/3}} = u_2^{1/6} u_1^{-1/6}
$$
\n(8.43)

which is the expected result for an axis of dimension 4 which is a discriminant of type  $II^*$ , cf. eq.[\(8.36\)](#page-98-1). Note that regularity rules out type II.

# 8.7 Geometries with  $\{\Delta_1, \Delta_2\} = \{2, 2\}$

The  $\{2, 2\}$  geometries are identified by the **Folk-theorem** with the geometries of Lagrangians SCFTs with gauge group  $SU(2) \times SU(2)$  and hence are known. However it is instructive to recover their classification from the solution of the corresponding Deligne-Simpson problem and Diophantine considerations. The perfect agreement of our Number-Theoretic analysis with the physical expectations yields a highly non-trivial check on our methodology and provides additional evidence for the  $\mu$ -rigidity conjecture.

The only *isotrivial* geometry with dimensions  $\{2, 2\}$  is  $\mathcal{N} = 4$  with the gauge group  $SU(2) \times SU(2)$ . Therefore we may assume the geometry to be non-isotrivial. We have  $x = 2$ , the conformal manifold has dimension 2, there are no enhanced divisors, and there must be 5 special points on  $\mathscr{P} \simeq \mathbb{P}^1$ . Moreover from the discussion in §.[8.1](#page-81-2) we have that an indecomposable  $\{2, 2\}$  geometry should be irreducible. Then  $\mu$ -rigidity yields

**Fact 24.** A non-isotrivial  $\{2, 2\}$  geometry has 5 discriminant components of type  $I_{n_j}$   $(n_j > 0, 1)$  $j = 1, \ldots, 5)$  and its Picard-Fuch equation is the 4-order Pochhammer ODE (i.e. graph (f)) with fugacities $75$ 

<span id="page-100-2"></span>
$$
\xi(\alpha_{\star}) = -1, \qquad \xi(\alpha_a) = 1 \text{ for } a \neq \star. \tag{8.44}
$$

The solutions of the Pochhmmer ODE have a well-known explicit expression in terms of contour integrals (see e.g. [\[146\]](#page-157-0)) from which one reads the monodromy representation [\[146\]](#page-157-0). Consider the following six explicit  $4 \times 4$  matrices:

<span id="page-100-3"></span>
$$
\mu_1 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ -2 & -2 & -2 & 1 \end{pmatrix} \quad \mu_2 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ -2 & -2 & 1 & 2 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \mu_3 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -2 & 1 & 2 & 2 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}
$$

$$
\mu_4 = \begin{pmatrix} 1 & 2 & 2 & 2 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \qquad \mu_5 = \begin{pmatrix} 3 & 2 & 2 & 2 \\ -2 & -1 & -2 & -2 \\ 2 & 2 & 3 & 2 \\ -2 & -2 & -2 & -1 \end{pmatrix} \quad M = \begin{pmatrix} 0 & 1 & 1 & 1 \\ -1 & 0 & 1 & 1 \\ -1 & -1 & 0 & 1 \\ -1 & -1 & -1 & 0 \end{pmatrix}
$$
(8.45)

The first four may be written as  $(i = 1, 2, 3, 4)$ 

$$
\mu_i = 1 + 2 v_i \otimes v_i^t M \tag{8.46}
$$

where  $\{v_4, v_3, v_2, v_1\}$  is the standard basis<sup>[76](#page-100-1)</sup> of  $\mathbb{Z}^4$ . The  $\mu_a$ 's satisfy (now  $a = 1, 2, ..., 5$ )

$$
(\mu_a - 1)^2 = 0 \qquad \text{rank}(\mu_a - 1) = 1, \qquad \mu_a - 1 \equiv 0 \mod 2
$$
  

$$
\mu_1 \mu_2 \mu_3 \mu_4 \mu_5 = -1 \qquad \text{det } M = 1 \qquad \qquad \mu_a^t M \mu_a = M,
$$
 (8.47)

so that  $\{\mu_1, \ldots, \mu_5\}$  is (a lift to  $Sp(4, \mathbb{Z})$  of) a monodromy representation in  $PSp(4, \mathbb{Z})$  with five local monodromies of type  $I_2$  where  $Sp(4, \mathbb{Z})$  is seen as the integral matrix group which leaves invariant the principal polarization M. In particular

**Corollary 8.** The Brauer class of the rigid representation associated to graph  $(f)$  with fugacities [\(8.44\)](#page-100-2) is trivial.

The change of basis which puts the polarization  $M$  in the standard form is

$$
S = \begin{pmatrix} 1 & 0 & -1 & 1 \\ 0 & 0 & 1 & -1 \\ 0 & 1 & 0 & 0 \\ 0 & -1 & 0 & 1 \end{pmatrix}, \qquad \Omega \stackrel{\text{def}}{=} S^t MS = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}.
$$
 (8.48)

In the standard symplectic basis the integral monodromies read

$$
\ell_a = S^{-1} \mu_a S \quad a = 1, 2, \dots, 5. \tag{8.49}
$$

In particular  $\ell_4$  becomes

$$
\ell_4 = \begin{pmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \tag{8.50}
$$

<span id="page-100-1"></span><span id="page-100-0"></span><sup>&</sup>lt;sup>75</sup> Recall from **Fact [22](#page-91-0)** that there is no irreducible lift of the  $PSp(4, \mathbb{Z})$  monodromy with  $\xi(\alpha_{\star}) = 1$ .

<sup>76</sup> Note the inversion of the order of the basis elements: it is chosen to agree with the conventions in the classical literature on the Pochhammer ODE.

while the five local monodromies are cyclically permuted by the  $\mathbb{Z}_5$  symmetry:

$$
\ell_{a+1 \bmod 5} = R \, \ell_a \, R^{-1},\tag{8.51}
$$

where

$$
R = \begin{pmatrix} 0 & 1 & 1 & -1 \\ 1 & -1 & -1 & 2 \\ -1 & 0 & 1 & 0 \\ 0 & -1 & 0 & 1 \end{pmatrix} \in Sp(4, \mathbb{Z}), \qquad R^5 = -1,
$$
 (8.52)

which makes manifest that all five local monodromies  $\ell_a$  are of type  $I_2$ . We conclude

**Corollary 9.** There is a  $\{2, 2\}$  geometry (with principal polarization) having 5 irreducible discriminant components of type  $I_2$ . All other (inequivalent)  $\{2, 2\}$  (principally polarized) geometries arise from monodromy representations in  $Sp(4, \mathbb{Z})$  which are conjugate over  $GL(4, \mathbb{C})$ to the  $\{\ell_1,\ldots,\ell_5\}$  one, but not conjugate in  $Sp(4,\mathbb{Z})$ .

Consider the matrix  $Q \in Sp(4,\mathbb{R})$ 

$$
Q = \left(\begin{array}{c|c}\n\sqrt{2} \mathbf{1}_2 & \cdots & \mathbf{1}_2 \\
\hline\n\frac{1}{\sqrt{2}} \mathbf{1}_2 & \cdots & \mathbf{1}_2\n\end{array}\right) \in Sp(4, \mathbb{R})
$$
\n(8.53)

which has the property that the five matrices  $Q\ell_aQ^{-1}$  have integral entries. One has

$$
\gcd_{ij} (Q\ell_1 Q^{-1} - 1)_{ij} = 4
$$
\n
$$
\gcd_{ij} (Q\ell_a Q^{-1} - 1)_{ij} = 1 \quad \text{for } a = 2, 3, 4, 5
$$
\n(8.54)

for instance, say,

$$
Q\ell_4 Q^{-1} = \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \tag{8.55}
$$

We interpret the monodromy representation  $\{\tilde{\mu}_a \equiv Q \ell_a Q^{-1}\}$  as a second *inequivalent* special geometry with  $\{\Delta_1, \Delta_2\} = \{2, 2\}$  having one  $I_4$  and four  $I_1$ .<sup>[77](#page-101-0)</sup>

<span id="page-101-1"></span>Fact 25. There are two inequivalent non-isotrivial, irreducible special geometries with dimensions  $\{2, 2\}$  with five discriminants of type  $I_2$  and, respectively, four  $I_1$  and one  $I_4$ .

According to the **Folk-theorem**, the above two special geometries should describe two inequivalent non-isotrivial Lagrangian SCFTs with gauge group  $SU(2) \times SU(2)$ . Indeed in the table of [\[89\]](#page-153-0) there are precisely two such models:  $\#13$  with five  $I_2$ 's, and  $\#31$  with four  $I_1$  and one  $I_4$ , in perfect agreement with our findings. These two  $\mathcal{N}=2$  models are Gaiotto

<span id="page-101-0"></span><sup>&</sup>lt;sup>77</sup> The matrix  $QRQ^{-1}$  which permutes the 5 local monodromies is now valued in  $Sp(4, \frac{1}{2}\mathbb{Z})$  so the five local monodromies, while conjugate in  $Sp(4,\mathbb{Q})$  are not longer conjugate in the Siegel modular group  $Sp(4,\mathbb{Z})$ .

class- $S[A_1]$  SCFTs [\[110\]](#page-154-4) as well as quiver gauge theories with quivers

<span id="page-102-1"></span>

From the physical side we do not expect other solutions, since there are no other Lagrangian SCFT with these properties. We shall give a math proof of this fact in the next subsection.

Remark 19. As a byproduct of the analysis we get explicit expressions for the periods of all {2, 2} SCFTs in terms of Pochhammer transcendents which have well-known integral representations and connection coefficients.

## <span id="page-102-2"></span>8.7.1 No other  $\{2, 2\}$  special geometries

We present a direct math proof that indeed there are no other integral symplectic monodromy representations conjugate in  $GL(4,\mathbb{C})$  to the (unique) rigid complex representation given by the solution of the Deligne-Simpson problem associated to graph (f). The reader may prefer to skip this technical subsection.

**Fact 26.** (Up to isomorphism) there are no other irreducible, non-isotrival special geometries with Coulomb dimensions  $\{2, 2\}$  besides the two listed in **Fact [25](#page-101-1)**.

Proof. We work in the Haraoka basis where the local monodromies are given by the matrices  $\mu_i$  and the symplectic form by M, cf. eq.[\(8.45\)](#page-100-3). Let  $L \in Sp(M, \mathbb{C})$  be a complex  $4 \times 4$  matrix such that  $L^tML = M$  while the five matrices  $\tilde{\mu}_i \equiv L\mu_i L^{-1}$   $(i = 1, ..., 5)$  have integral entries. For  $i = 1, \ldots, 4$  we have

<span id="page-102-0"></span>
$$
\tilde{\mu}_i - \mathbf{1} = 2 \, Lv_i \otimes v_i^t L^t M \quad \text{(not summed over } i! \text{)} \tag{8.57}
$$

where  $\{v_4, v_3, v_2, v_1\}$  is the standard column-vector basis of  $\mathbb{Z}^4$  (written in reverse order). We extend eq.[\(8.57\)](#page-102-0) to  $i = 5$  by setting  $v_5 = (-1, 1, -1, 1)^t$ . It is easy to check that

$$
v_i^t M v_j = 1 \quad \text{for} \quad 1 \le i < j \le 5. \tag{8.58}
$$

If the first four matrices in the LHS of  $(8.57)$  have to be integral,  $Lv_i$  must be a scalar multiple

of a 4-vector with integral coefficients, so there are 4 complex numbers  $\rho_a$  such that

$$
L = \begin{bmatrix} \rho_1 n_{1,1} & \rho_2 n_{1,2} & \rho_3 n_{1,3} & \rho_4 n_{1,4} \\ \rho_1 n_{2,1} & \rho_2 n_{2,2} & \rho_3 n_{2,3} & \rho_4 n_{2,4} \\ \rho_1 n_{3,1} & \rho_2 n_{3,2} & \rho_3 n_{3,3} & \rho_4 n_{3,4} \\ \rho_1 n_{4,1} & \rho_2 n_{4,2} & \rho_3 n_{4,3} & \rho_4 n_{4,4} \end{bmatrix} \equiv N \operatorname{diag}(\rho_1, \dots, \rho_4) \in Sp(4, \mathbb{C}) \tag{8.59}
$$

where  $n_{a,i}$  are integers,  $gcd\{n_{1,i}, n_{2,i}, n_{3,i}, n_{4,i}\} = 1$  for  $i = 1, \ldots, 4$ , and the four columns vectors  $n_i \equiv (n_{a,i})$  are linear independent over C. N is the integral matrix with entries  $n_{a,i}$ . One has

<span id="page-103-1"></span>
$$
1 = \det L = (\rho_1 \rho_2 \rho_3 \rho_4) \det N \qquad \det N \in \mathbb{Z}.
$$
 (8.60)

Moreover the integrality of  $\tilde{\mu}_i$  implies

<span id="page-103-0"></span>
$$
2\rho_i^2 \in \mathbb{Z}, \quad \text{and} \quad (\rho_i \rho_j)^{-1} \in \mathbb{Z} \qquad \text{for } 1 \le i < j \le 4,\tag{8.61}
$$

where the second condition follows from  $(Lv_i)^t M(Lv_j) = 1$  for  $0 \le i \le j \le 4$ . Requiring that  $\tilde{\mu}_5$  is also integral extends [\(8.61\)](#page-103-0) to  $i, j = 1, \ldots, 5$  where  $\rho_5$  is the complex number (unique when it exists) such that

$$
\rho_5 n_{i,5} \equiv (Lv_5)_i = -\rho_1 n_{i,1} + \rho_2 n_{i,2} - \rho_3 n_{i,3} + \rho_4 n_{i,4}
$$
\n(8.62)

where  $\{n_{1,5}, n_{2,5}, n_{3,5}, n_{4,5}\}$  are coprime integers. Dividing out by  $\rho_5$  we see that  $\rho_i/\rho_5 \in \mathbb{Q}$ , so  $\rho_i = q_i \rho_5$  with  $q_i \in \mathbb{Q}$   $(q_5 = 1)$ . Since  $2\rho_i^2 = 2\rho_5^2 q_i^2$ , from the first eq.[\(8.61\)](#page-103-0) we conclude that there is a square-free integer d such that

$$
\rho_i = \sqrt{\frac{d}{2}} \, k_i, \qquad k_i \in \mathbb{N}.\tag{8.63}
$$

Then

<span id="page-103-2"></span>
$$
(\rho_i \rho_j)^{-1} = \frac{2}{d k_i k_j} \in \mathbb{Z} \qquad \text{for } 1 \le i < j \le 5. \tag{8.64}
$$

From eq.[\(8.60\)](#page-103-1) we get

$$
1 = \frac{d^2}{4} k_1 k_2 k_3 k_4 \det N = \frac{\det N}{\frac{2}{d k_1 k_2} \frac{2}{d k_3 k_4}}
$$
(8.65)

Now we have two possibilities:

 $d = 2$  then  $k_i = 1$  and  $\rho_i = 1$  for  $i = 1, \ldots, 5$  while det  $N = 1$ , so that  $N \equiv L \in Sp(4, \mathbb{Z})$ and  $2\rho_i^2 = 2$  for  $i = 1, 2, 3, 4, 5$ . In this solution we have five  $I_2$ 's. It is our first integral monodromy corresponding to the upper quiver in  $(8.56)$ ;

 $d=1$  so that  $2\rho_i^2 = k_i^2$  is a perfect square for all i. Eq.[\(8.64\)](#page-103-2) shows that at most one  $k_i$ 

 $(i = 1, \ldots, 5)$  can be 2 all others should be 1. By the  $\mathbb{Z}_5$  symmetry [\(8.61\)](#page-103-0) we may assume that  $k_1 = k_2 = k_3 = k_4 = 1$  $k_1 = k_2 = k_3 = k_4 = 1$  $k_1 = k_2 = k_3 = k_4 = 1$ . Then one gets  $k_5 = 2$  (cf. **Lemma 4** in appendix [C\)](#page-142-1). Therefore we get a monodromy with one  $I_4$  and four  $I_1$ 's which is the second special geometry in Fact [25](#page-101-1).

 $\Box$ 

# 8.8 Irreducible geometries with  $\{2, \Delta\}, \Delta \neq 2$

We already discussed these geometries when discussing the **Folk-theorem**, now we look at them from the present  $\mu$ -rigidity viewpoint. The are 4 special points on  $\mathscr P$  and the Picard-Fuchs ODE has a graph  $\mathfrak{G}$  with 4 branches. Graph (e) was ruled out, so all  $\mu$ -rigid special geometries should arise from graph (d) which corresponds to an ODE reduction of the Appell PDE aka Okubo II<sup>∗</sup> ODE [\[147\]](#page-157-1).

<span id="page-104-1"></span>Fact 27. Assume *u*-rigidity. (1) There are no non-isotrivial, irreducible geometries with dimensions  $\{2, \Delta\}$  and  $\Delta \notin \{2, 3, 4, 6\}$ . (2) A non-isotrivial, irreducible geometry with dimensions  $\{2,3\}$  should have two knotted discriminant of types  $I_{n_1}$ ,  $I_{n_2}$  (with  $n_1$ ,  $n_2$  satisfying the conditions in appendix [D](#page-143-0)), while the third discriminant is  $u_2 = 0$  of type  $I_{2n_3}$   $(n_3 > 0)$ . (3) A non-isotrivial, irreducible geometry with dimensions  $\{2, 2m\}$  ( $m \geq 2$ ) should have two non-enhanced discriminant of type  $I_{n_1}$ ,  $I_{n_2}$  (with  $n_1, n_2$  as in appendix [D](#page-143-0)), while the third discriminant is  $u_2 = 0$  of type  $I_{n_3}^*$   $(n_3 > 0)$ .

These results can also be inferred from the **Folk-theorem**. The obvious solution to the conditions in appendix [D](#page-143-0) is  $n_1 = n_2 \in \{1, 2\}.$ 

Argument. (1)  $\Delta$  is a dimension in rank-1 so if the statement does not hold  $\Delta = m/(m-1)$ for  $m \in \{3, 4, 6\}$ . Then (say) for  $m = 4, 6$ 

$$
(\Delta_1, \Delta_2) = \frac{2}{m-1} \left( m - 1, \frac{m}{2} \right) = \lambda(d_1, d_2)
$$
 (8.66)

so  $(d_1, d_2)$  are coprime  $\neq 1$  and both axes are enhanced. Since  $(m - 1) > 2$  and  $\delta < 4$  both are regular. The  $\varrho$ -monodromy along the  $u_2 = 0$  axis then has eigenvalues

$$
\exp(\pm 2\pi i/2), \qquad \exp(\pm 2\pi i(1-\Delta)/2) = \exp(\pm 2\pi i/(2(m-1)) \tag{8.67}
$$

while the eigenvalues of  $\mu = \pm \varrho^{1/(m-1)}$  are not algebraic numbers of degree  $\leq 2$ .

Statement (2) reduces to (3) for  $2m = 6$  by the following trick. The family over  $\mathscr{\tilde{P}}$ of a  $\{2,3\}$  geometry has local monodromies  $\mu_1$ ,  $\mu_2$ ,  $\mu_3$ ,  $\mu_4$  with with  $\mu_1\mu_2\mu_3\mu_4 = 1$ ,  $\mu_1$ ,  $\mu_2$ (knotted) monodromies of type  $I_1$ , while  $(\mu_3)^3 = 1$  and  $(\mu_4)^2 = \varrho_4$  where  $\varrho_4$  is the monodromy around the axis  $u_2 = 0$  of dimension  $2 \equiv d_1$ . The local monodromies<sup>[78](#page-104-0)</sup> { $\mu_1, \mu_2, -\mu_3, -\mu_4$ }

<span id="page-104-0"></span><sup>&</sup>lt;sup>78</sup> The change of sign in  $\mu_3$  is due to the different sign of the phases  $\exp(\pm 2\pi i/\Delta_2)$  for  $\Delta_2 = 3$  and  $\Delta_2 = 6$ .

can also be seen (as a lift) of the local monodromy for a  $\{2,6\}$  geometry where now  $d_1 = 1$ so  $-\mu_4 = \varrho_4$ . Thus

$$
\varrho_4 \Big|_{\{2,3\}} = (\varrho_4)^2 \Big|_{\{2,6\}} \tag{8.68}
$$

and so  $(3) \Rightarrow (2)$ .

(3) the two  $\delta = 1$  branches in graph (d) correspond to  $I_{n_1}$   $(n_1 > 0)$  discriminants. The  $\delta = 2$  branch then corresponds to the enhanced axis of dimension  $2m$  whose local  $\mu$ -monodromy has eigenvalues  $\exp(\pm 2\pi i/2m)$  both with multiplicity 2 (and trivial Jordan blocks). The forth branch has then  $\delta = 3$ , and since the  $u_2 = 0$  axis is not enhanced, by eq.[\(8.21\)](#page-88-0) it must be a discriminant of type  $I_{n_2}^*$   $(n_2 > 0)$  or the local monodromy must satisfy  $\mu^{\ell} = \pm 1$  for  $\ell \in \{3, 4, 6\}$  (that is, a semisimple Kodaira type  $\neq I_0^*$ ). The second possibility looks inconsistent for the following reason. The period matrix  $\tau_{ij}$  along the  $u_2 = 0$  axis must have the form  $\text{diag}(\zeta, \tau')$  with  $\zeta^{\ell} = 1$  at all points  $\tau \in \mathcal{M}$  which is not consistent with the discussion in §. [5.2.3.](#page-61-1)

When the  $u_2 = 0$  axis is of type  $I_{n_2}^*$ , or  $\Delta \in \{3, 4, 6\}$  fugacities are

<span id="page-105-0"></span>
$$
e^{-2\pi i/\Delta}
$$
\n
$$
1 \longrightarrow \pm e^{\pi i/\Delta} \longrightarrow -1 \longrightarrow 1
$$
\n(8.69)

An element  $\lambda$  of the root lattice with  $\xi(\lambda) \in \mathbb{R}$  should be even at the central node; the only decomposition into a sum of roots with this property is

$$
\beta_1 + \beta_2 = \frac{1}{1 \cdot 2 \cdot 1 \cdot 1} + \frac{1}{1 \cdot 1} \tag{8.70}
$$

with  $\xi(\beta_a) = -1$ , so that the representation is irreducible hence rigid.

 $\Box$ 

**Remark 20.** All the known models with dimensions  $\{2, \Delta \neq 2\}$  (a part for the subtle #69 discussed in §. [5.2.3\)](#page-61-1) obey Fact [27](#page-104-1) with  $n_1 = n_2 \in \{1, 2\}$ . According to the Folk theorem the known models exhaust the geometries with these dimensions. Indeed a non-systematic search produced no integral representation corresponding to a semisimple discriminant along the  $u_2 = 0$  axis.

**Remark 21.** One can get the possible values of the integers  $n_1$  and  $n_3$  by constructing the explicit monodromy representation as we did in the Pochhammer case. The explicit construction of the monodromy also allows to check the triviality of the Brauer class (as we did in the previous case) and the positivity condition. Of course, both statements hold true since we know that these geometries exists (for suitable  $\{n_1, n_3\} \in \mathbb{N}^2$ ) because we may construct the corresponding SCFT by Lagrangian methods.

We give some examples.

### 8.8.1 Example: irreducible special geometries with  $\{\Delta_1, \Delta_2\} = \{2, 4\}$

While there is a substantial literature on the explicit monodromy representation for the Okubo  $II^*$  ODE, see e.g. [\[148,](#page-157-2)[149\]](#page-157-3), to the best of our knowledge all published results apply only under the simplifying assumption that the local monodromies are semisimple, which is not the case of interest for the present applications. Then, to construct explicitly the geometries, we need to proceed by learned guess-work. The nice aspect of the story is that if we manage to guess one monodromy representation, all others are conjugate to it by rigidity, so a single happy guess is good enough. We got the following one:

<span id="page-106-0"></span>**Fact 28.** The following four explicit integral  $4 \times 4$  matrices

matrix	minimal equation	branch of conjugacy class	
	$M_1 = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}$ $(M_1 + 1)^2(M_1 - 1) = 0$ $\underline{\qquad \qquad -1}$		
$M_2 = \left(\begin{array}{rrr} 1 & 0 & 0 & 0 \\ -1 & 1 & 0 & 1 \\ -1 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 \end{array}\right)$	$(M_2-1)^2=0$	$-1$	(8.71)
$M_3 = \begin{pmatrix} 0 & 1 & 1 & 0 \\ -1 & 2 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 1 & -1 & -1 & 1 \end{pmatrix}$ $(M_3 - 1)^2 = 0$		$-11$	
$M_4 = \left( \begin{array}{rrr} 0 & 1 & -1 & 2 \\ 0 & 0 & -1 & 1 \\ -1 & 1 & 1 & -1 \\ 1 & 1 & -1 & 1 \end{array} \right)$ $M_4^2 + 1 = 0$		$-12$	

satisfy

$$
M_1 M_2 M_3 M_4 = 1
$$
 and  $M_a^t \Omega M_a = \Omega$   $(a = 1, ..., 4)$  (8.72)

where

$$
\Omega = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix} \tag{8.73}
$$

In particular the Brauer class of the order-4 Okubo II<sup>\*</sup> ODE with the fugacities in eq.[\(8.69\)](#page-105-0) (with  $\Delta = 4$ ) is trivial.

Corollary 10. There exists a (non-isotrivial, irreducible) special geometry with dimensions  $\{2,4\}$  two knotted divisors of type  $I_1$  while the axis of dimension 2 is a component of the discriminant of type  $I_2^*$ .

In the list of [\[89\]](#page-153-0) there are two SCFT with precisely this geometry  $\#10$  (SO(5) with six 4's) and  $\#43$  (SO(5) with three 5's). There is another SCFT with dimensions  $\{2,4\}$  whose geometry is non-isotrivial irreducible:  $\#48$  (SO(5) with two 4 and two 5) which has two knotted discriminants of type  $I_2$  while the axis of dimension 2 has type  $I_1^*$ . Again this means

that we have a second integral-symplectic representation which is conjugated to the one in **Fact [28](#page-106-0)** in  $GL(4,\mathbb{C})$  but not in  $Sp(4,\mathbb{Z})$ . It follows from **Fact** (or the **Folk-theorem**) that there are no other geometries of this kind.

Going through the same (elementary) systematic procedure for the search of integral symplectic monodromies conjugate to the one we used in §. [8.7.1](#page-102-2) one gets

Fact 29. Besides the one in Fact [28](#page-106-0) there is precisely one other irreducible geometry with dimensions {2, 4} obtained by conjugating the previous one

$$
\tilde{M}_a = U M_a U^{-1} \quad a = 1, \dots, 4, \qquad U = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 & 0 & -1 \\ 0 & 2 & 0 & 0 \\ 0 & 2 & -2 & 0 \\ 0 & -1 & 1 & -1 \end{pmatrix} . \tag{8.74}
$$

Explicitly

$$
\tilde{M}_1 = \begin{pmatrix}\n-1 & -1 & 1 & 2 \\
0 & -1 & 0 & 0 \\
0 & -2 & 1 & 0 \\
0 & 1 & 0 & 1\n\end{pmatrix} \qquad \tilde{M}_2 = \begin{pmatrix}\n1 & 0 & 0 & 0 \\
-2 & 1 & 0 & 0 \\
0 & 0 & 1 & 0 \\
0 & 0 & 0 & 1\n\end{pmatrix}
$$
\n
$$
\tilde{M}_3 = \begin{pmatrix}\n-1 & 2 & 0 & 2 \\
-2 & 3 & 0 & 2 \\
-2 & 2 & 1 & 2 \\
0 & 0 & 0 & 1\n\end{pmatrix} \qquad \tilde{M}_4 = \begin{pmatrix}\n1 & -1 & -1 & -4 \\
0 & -1 & 0 & -2 \\
2 & -4 & -1 & -6 \\
0 & 1 & 0 & 1\n\end{pmatrix}
$$
\n(8.75)

which confirms the physical expectation that the second geometry has one discriminant of type  $I_1^*$  and two of type  $I_2$  as required for  $SO(5)$  with two 4 and two 5.

Again our findings agree with the **Folk-theorem**.

## 8.8.2 Example: irreducible special geometry with  $\{\Delta_1, \Delta_2\} = \{2, 3\}$

The Diophantine problem looks harder. However we may restrict the search by the following considerations: the monodromy  $\mu_1$  associated to the regular axis is known by **Proposition [3](#page-61-0)** (see also the more detailed analysis in [\[106,](#page-154-5)[107\]](#page-154-6)). So we can fix two of the required matrices

$$
\Omega = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix} \qquad \mu_1 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & -1 & 0 & 0 \\ 0 & 0 & -1 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix} \tag{8.76}
$$

which satisfy

$$
\mu_1^t \Omega \mu_1 = \Omega, \qquad \mu_1^3 = 1. \tag{8.77}
$$

Then  $\mu_2$ ,  $\mu_3$  must be symplectic with respect to the principal polarization  $\Omega$  and  $\mu_a - 1$ should have rank 1 for  $a = 2, 3$  i.e.

$$
\mu_a = \mathbf{1} + v_a \otimes v_a^t \Omega. \tag{8.78}
$$

for some integral 4-vectors  $v_a$ . By some try and error we find  $v_2 = (1, 0, -1, -1)^t$  and  $v_3 = (0, 1, 1, 1)^t$  so that

$$
\mu_2 = \begin{pmatrix} 2 & 1 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & -1 & 0 & 0 \\ -1 & -1 & -1 & 1 \end{pmatrix}, \qquad \mu_3 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 1 \\ -1 & -1 & 1 & 1 \\ -1 & -1 & 0 & 2 \end{pmatrix}
$$
 (8.79)
since  $v_1$  and  $v_2$  are integral and primitive, these are monodromies of type  $I_1$ . Then the relation  $\mu_1\mu_2\mu_3\mu_4 = 1$  yields

$$
\mu_4 = \begin{pmatrix} -1 & 0 & 0 & 1 \\ 1 & 1 & -1 & 3 \\ 0 & 0 & -1 & 1 \\ 0 & 0 & -1 & 1 \end{pmatrix} \tag{8.80}
$$

and the monodromy around the discriminant component  $u_2 = 0$  in the Coulomb branch is

$$
\varrho_4 \equiv \mu_4^2 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 6 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \tag{8.81}
$$

We conclude

**Fact 30.** There is a special geometry with dimensions  $\{2, 3\}$ , two knotted discriminant components of type  $I_1$ , and a third component  $u_2 = 0$  of type  $I_6$ .

Of course this is just the well-known geometry of  $SU(3)$  with six fundamentals. Here we reconstructed it from the  $\mu$ -rigidity perspective as a further check of the general philosophy. A (non-systematic) search has produced no other solution as physically expected.

### 8.9 Rigid, non-isotrivial, rank-2 special geometries

Rank-2 non-isotrivial, primitive, *rigid* geometries have three special points on  $\mathscr P$  and hence Dynkin graphs with 3 branches. Moreover  $\Delta_2$  cannot be an integral multiple of  $\Delta_1$  (since in this case one has  $\varkappa = \Delta_1$  while  $\Delta_1 \neq 1, 2$ , so the geometry is automatically isotrivial [\[25\]](#page-150-0)). Thus two special points are enhanced (they may or may not be part of the discriminant) while the third one arises from a non-enhanced discriminant component (which now should be knotted). Assuming  $\mu$ -rigidity, the graph selection rules of §.[8.4.2](#page-91-0) allow two classes of geometries corresponding, respectively, to the Dynkin graphs of the Kac-Moody algebras  $E_7$ and  $E_7$ .

### 8.9.1 Rigid geometries with a knotted discriminant of type  $I_n$

The vast majority of known special geometries of rank-2 SCFT with  $\mathcal{R}_2 = 0$  have a unique knotted discriminant which is of Kodaira type  $I_1$ . To shed light on some puzzles we shall encounter later, we begin by considering the implications of this fact per se, without making any other assumption (however plausible) nor using any restriction arising from our previous considerations. In this subsection we assume only that the  $\mu$ -monodromy is given by three quasi-unipotent matrices  $\mu_0, \mu_1, \mu_\infty \in Sp(4, \mathbb{Z})$ , satisfying  $\mu_0\mu_1\mu_\infty = \pm 1$ , while  $(\mu_1 - 1)$  is nilpotent of rank 1. The double sign  $\pm$  applies only when  $\varkappa = 2$ . The presence of a knotted discriminant guarantees that the geometry is indecomposable, but we do not rule out the possibility that  $\mu$  is reducible.

If the minimal polynomials of both  $\mu_0$  and  $\mu_\infty$  are of degree 4 we get an  $\widehat{E}_7$  graph and the representation generated by  $\mu_0$  and  $\mu_\infty$  is irreducible precisely when their minimal polynomials  $P_0(z)$ ,  $P_\infty(z)$  have no common zero i.e.  $gcd(P_0(z), P_\infty(z)) = 1$  or equivalently iff their resultant Res $(P_0(z), P_1(z)) \neq 0$ . When this is the case the representation is automatically irreducible and rigid. This "nice" case will be discussed in great detail in §. [8.10](#page-112-0) below. We ask what happens in the two "less nice" cases:

- (I) the minimal polynomial of, say,  $\mu_0$  has degree  $\leq 3$ ;
- (II)  $P_0(z), P_\infty(z)$  have degree 4 while the polynomial  $gcd(P_0(z), P_\infty(z))$  has positive degree.

In either case the monodromy representation cannot be irreducible and decomposes (over C) into smaller ones. Since there is a non-trivial unipotent monodromy associated to the knotted discriminant, the geometry cannot be isotrivial, and since 2 is not a dimension, it is rigid. Therefore we end up in the scenario discussed in §. [8.1.3.](#page-83-0)

<span id="page-109-0"></span>**Fact 31.** A rigid rank-2 geometry with a single knotted discriminant of type  $I_n$  ( $n > 0$ ) is either irreducible of class  $\widehat{E}_7$  or it is reducible quasi-isotrivial with Coulomb dimensions  ${3, 4}$  resp.  ${4, 6}$  and with the discriminant types listed in **Fact [20](#page-83-1)**. The two non-trivial periods can be expressed in terms of Gauss' hypergeometric functions.

#### 8.9.2 Rigid non-isotrivial geometries of class  $E_7$

Class- $E_7$  contains the non-isotrivial, primitive, rigid and  $\mu$ -rigid geometries where the knotted discriminant is not of type  $I_n$ . The  $E_7$  class is quite small. Indeed these geometries are severely constrained. The point is that class- $E_7$  rigid monodromies behave quite differently from their  $\widehat{E}_7$  cousins which are hypergeometric monodromies. In the  $\widehat{E}_7$  case the Brauer class is always trivial by the Levelt theorem [\[150,](#page-157-0) [151\]](#page-157-1), while "most"  $E_7$ -class rigid monodromies are known to have non-trivial Brauer classes [\[143\]](#page-156-0), so they do not describe special geometries: this is one of the reasons why class- $E_7$  special geometries are relatively rare. The computation of the Brauer class is difficult, so this subclass of special geometries is the hardest one to classify and construct in explicit terms. In principle we can settle the question with a finite amount of work since we do have explicit integral representation of the solutions [\[100\]](#page-154-0), but we shall not attempt to do it here.

The following statement is proven in appendix [E:](#page-145-0)

**Lemma 3.** In a (non-isotrivial) class- $E_7$  special geometry the short  $\delta = 2$  branch cannot correspond to a non-enhanced divisor.

Thus the  $\delta = 2$  branch represents a coordinate axis in the Coulomb branch  $\mathscr{C}$  which should be enhanced but cannot be part of the discriminant because the  $\delta = 2$  branch is too short to describe an enhanced discriminant. Working mod 2 we get:

**Fact 32.** In a class- $E_7$  special geometry the Kodaira types  $I_n$  and  $I_{2k}^*$  ( $k \in \mathbb{N}$ ) are excluded for the non-enhanced (i.e. knotted) discriminant associated to the  $\delta = 3$  branch.

Now we are ready to show:

<span id="page-110-0"></span>**Fact 33.** A rigid non-isotrivial (irreducible) geometry of class  $E_7$  whose two dimensions  $\{\Delta_1, \Delta_2\}$  are both **non new**, must have Coulomb dimensions  $\{4, 6\}.$ 

 $\hat{\mathcal{L}}$  Caveat 4. There are plenty of rigid non-isotrivial irreducible special geometries with dimensions  $\{4, 6\}$  which are not of class  $E_7$  but rather of class  $\widehat{E}_7$ . Note that  $\{4, 6\}$  is the same "troublesome" dimension pair which allows for quasi-isotrivial geometries.

**Remark 22.** There is one known class- $E_7$  SCFTs with dimensions  $\{4, 6\}$  and a non-enhanced divisor of type  $I_1^*$  (model  $\#66$ ); in addition one of the two possible geometries for the model #22 (the non-cover one) would be of class- $E_7$  with dimensions  $\{4, 6\}$  and a non-enhanced divisor of type II. Both are consistent with the above Facts.

Argument. We call  $\Delta_1$  the dimension of the regular axis corresponding to the  $\delta = 2$  branch, and  $\Delta_2$  the other one (beware:  $\Delta_1$  may be actually larger than  $\Delta_2$ ).  $\Delta_1$  is an old dimension  $\neq$  2. Applying the dimension formulae of §. [2.5.1](#page-22-0) to the regular axis of dimension  $\Delta_1$  we get

$$
\Delta_2 = 2 + k \Delta_1 \quad k \in \mathbb{Z} \setminus \{0\}.
$$
\n(8.82)

When  $\Delta_2$  is a new dimension the other axis is also regular, so we must have a non-enhanced discriminant which must be non-semisimple for the geometry not to be isotrivial. The nonenhanced discriminant should be associated with the  $\delta = 3$  branch and hence be of type  $I_n^*$  $(n > 0)$ . In this case we can apply the dimension formulae of §. [2.5.1](#page-22-0) also to the axis of dimension  $\Delta_2$  getting a list of allowed values for  $\Delta_1$  which should agree with the original one. One gets the table

sequence	rank-2 dimensions in the sequence	old dimensions $\Delta_1$ consistent with $\Delta_2$	notes	
$2+\frac{6}{5}\mathbb{Z}$	8	4, 6	$\Delta_1 = \frac{6}{5}$ ruled out	
$2+\frac{4}{3}\mathbb{Z}$	6		$\{\frac{4}{3}, 6\}$ allowed	
	10	4	$\{\frac{4}{3}, 10\}$ ruled out	
$2+\frac{3}{2}\mathbb{Z}$	8	4, 6	$\Delta_1 = \frac{3}{2}$ ruled out	(8.83)
$2+3\mathbb{Z}$	5	3, 4	$\{3,5\}$ <i>not</i> ruled out	
	8	4, 6	$\{3, 8\}$ ruled out	
$2+4\mathbb{Z}$	6		$\{4,6\}$ allowed	
	10	4	$\{4, 10\}$ <i>not</i> ruled out	
$2+6\mathbb{Z}$	8	6, 4	$\{6,8\}$ <i>not</i> ruled out	

Under the assumptions in **Fact [33](#page-110-0)**,  $\Delta_2$  is an old dimension and there are only two allowed dimension pairs with these properties

$$
\left\{\frac{4}{3}, 6\right\} \quad \text{and} \quad \{4, 6\}. \tag{8.84}
$$

The first one has  $d_2 = 9$  which is not acceptable in rank-2. Thus we remain with  $\{4, 6\}$ .  $\Box$ 

#### <span id="page-111-1"></span>8.9.3 Class- $E_7$  geometries with new dimensions?

The proof of **Fact [33](#page-110-0)** leaves open the possibility of non-isotrivial class  $E_7$ -geometries of with an *irreducible* non-enhanced discriminant of type  $I_{2k+1}^*$  and dimensions in the list<sup>[79](#page-111-0)</sup>  $\{3,5\}$ ,  $\{4, 10\}$ ,  $\{6, 8\}$ . All known SCFTs with dimensions  $\{3, 5\}$ ,  $\{4, 10\}$  and  $\{6, 8\}$  have geometries of class  $E_7$  and one wonders if any class- $E_7$  geometry with a new dimension exists.

However we can write an explicit integral, rigid, irreducible monodromy representation of class- $E_7$  which is appropriate to dimensions  $\{3, 5\}$  without making any new computation. Just take as  $\mu_1 = -R$  where R is the matrix defined in eq.[\(8.52\)](#page-101-0) and as  $\mu_2$  the second matrix in eq.[\(8.76\)](#page-107-0). Now

<span id="page-111-2"></span>
$$
\mu_1^t \Omega \mu_1 = \mu_2^t \Omega \mu_2 = \Omega, \quad \mu_1^5 = \mu_2^3 = \mathbf{1}
$$
 (8.85)

while

$$
\mu_3 \stackrel{\text{def}}{=} (\mu_1 \mu_2)^{-1} = \begin{pmatrix} 1 & 1 & 0 & -1 \\ -1 & 0 & 1 & -1 \\ 0 & 1 & 1 & -1 \\ -1 & 1 & 1 & -2 \end{pmatrix} \tag{8.86}
$$

which has a Jordan block of size 2 associated to the eigenvalue −1 and 2 trivial Jordan blocks associated to the eigenvalue  $+1$ . Moreover

<span id="page-111-3"></span>
$$
\mu_3^2 = \mathbf{1} + 2 w \otimes w^t \Omega \quad \text{where} \quad w^t = (0 \ 1 \ 0 \ 1), \tag{8.87}
$$

so that  $\mu_3$  is an integral symplectic monodromy of Kodaira type  $I_1^*$ . Thus a special geometry of dimensions  $\{3, 5\}$  with a single discriminant component of type  $I_1^*$  potentially may exist.

**Regularity?** Local regularity along the axes poses no problem for the putative  $\{3, 5\}$ geometry of class- $E_7$ . We have  $h = u_1^2/u_2$  and  $z = u_2^5/u_2^3$ . Along the first axis

$$
a^{\parallel} \sim h \cdot z^{1/3} = u_1^{1/3}, \qquad a^{\perp} \sim h \cdot z^{2/3} = u_2 \cdot u_1^{-4/3}
$$
 (8.88)

along the second axis

$$
a^{\parallel} \sim h \cdot z^{2/5} = u_2^{1/5}, \qquad a^{\perp} \sim h \cdot z^{1/5} = u_1 \cdot u_2^{-2/5}.
$$
 (8.89)

The only mechanism which may prevent this geometry to exist is a failure of the connection formula i.e. the locally good solution around one axis is not the analytic continuation of the one which is good at the other axis. It looks a very unlikely possibility, so here we have a promising candidate for a novel special geometry which was not previously known.

<span id="page-111-0"></span><sup>79</sup> Recall that when a rank-2 geometry with one dimension equal 12 is automatically isotrivial.

## <span id="page-112-0"></span>8.10 Rigid special geometries of class  $\widehat{E}_7$

The Fuchsian ODE with rigid monodromy and graph  $\widehat{E}_7$  is Thomae's order-4 hypergeometric equation whose monodromy is well understood thanks to Levelt [\[150\]](#page-157-0) and Beukers-Heckman [\[151\]](#page-157-1). In particular we know that the Brauer class is always trivial. We write  $\mu_0$ , and  $\mu_\infty$  for the local  $\mu$ -monodromies associated to the two  $\delta = 4$  branches and  $\mu_1$  for the one associated to the non-enhanced divisor i.e. to the  $\delta = 1$  branch of  $\mathfrak{G}$ . We absorb the possible sign in the lift to  $Sp(4, \mathbb{Z})$  in  $\mu_{\infty}$ , so that the relation  $\mu_0\mu_1\mu_{\infty} = 1$  holds.

We write  $X_a$   $(a = 0, \infty)$  for, respectively, the matrices  $\mu_0$  and  $\mu_{\infty}^{-1}$ . The fact that the conjugacy classes of the matrices  $X_0$  and  $X_\infty$  are described by  $\delta = 4$  branches is equivalent to the statement that their minimal polynomials have degree 4. In this case, up to conjugacy, they may be set in the form

<span id="page-112-2"></span>
$$
X_a = \begin{pmatrix} 0 & 0 & 0 & c_{0,a} \\ 1 & 0 & 0 & c_{1,a} \\ 0 & 1 & 0 & c_{2,a} \\ 0 & 0 & 1 & c_{3,a} \end{pmatrix} \quad a = 0, \infty
$$
 (8.90)

where the  $c_{i,a}$ 's are the coefficients in the characteristic polynomial of  $X_a$ :

<span id="page-112-1"></span>
$$
P_a(z) \equiv \det[z - X_a] = z^4 - c_{3,a} z^3 - c_{2,a} z^2 - c_{1,a} z - c_{0,a}.
$$
 (8.91)

Since  $X_a \in Sp(4,\mathbb{Z})$  the coefficients are integral and  $c_{0,a} = -1$  and  $c_{3,a} = c_{1,a}$ . Further restrictions on the  $c_{i,a}$  are obtained by the condition that the RHS of  $(8.91)$  is a product of cyclotomic polynomials. From now on we assume that the characteristic polynomials of  $X_0$ ,  $X_{\infty}$  satisfy these conditions.

Suppose both  $X_0, X_\infty$  are in the form [\(8.90\)](#page-112-2) up to a common conjugation. In this case

$$
X_1 \equiv X_0^{-1} X_{\infty} = \mathbf{1}_4 + V \otimes W^t, \quad \text{with } V = \begin{pmatrix} A \\ B \\ C \end{pmatrix}, \quad W = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}
$$
  
where  $A = c_{1,\infty} - c_{1,0}, \quad B = c_{2,\infty} - c_{2,0}$  (8.92)

is automatically in the conjugacy class described by a  $\delta = 1$  with all its eigenvalues are equal 1. A  $\mu$ -representation with these local monodromy classes is automatically rigid, and the three matrices  $\mu_0 \equiv X_0$ ,  $\mu_1 \equiv X_1$  and  $\mu_\infty = X_\infty^{-1}$  should generate the *unique* monodromy representation up to conjugacy in  $GL(4,\mathbb{C})$ . This statement is the Levelt theorem [\[150\]](#page-157-0). Since in our case the coefficients of the characteristic polynomials are rational integers, the Levelt representation is automatically integral and the Brauer class is trivial. The Levelt representation is  $Sp(4, \mathbb{Z})$ -valued and irreducible iff the two characteristic polynomials,  $P_0(z)$ and  $P_{\infty}(z)$ , satisfy the following conditions:

- (a)  $(P_0(z), P_\infty(z))$  is a pair of degree 4 products of cyclotomic polynomials  $\prod_n \Phi_n(z)^{m_n}$ with the property  $P_a(z) = z^4 P_a(1/z)$  (for  $a = 0, \infty)$ ;
- (b) each polynomial has at most one double root (corresponding to at most one non-trivial

Jordan block of size 2) which then must be  $\pm 1$ ;

- (c)  $P_0(z)$  and  $P_{\infty}(z)$  have no common root by **Fact [23](#page-94-0)**. In other words, their resultant must be non-zero,  $\text{Res}(P_0(z), P_\infty(z)) \neq 0;$
- (d) there should exist a pair of *coprime* integers  $(d_1, d_2)$  such that a pair  $(X_0, X_\infty)$  of  $4 \times 4$ matrices with minimal polynomials  $(P_0(z), P_\infty(z))$  have the property that  $X_0^{d_1}$ ,  $X_\infty^{d_2}$ have two trivial Jordan blocks of eigenvalue  $\pm 1$ . This condition eliminates, say, the troublesome polynomial pair  $(\Phi_1^2 \Phi_6, \Phi_2^2 \Phi_3)$ ;
- (e) we can disregard  $\Phi_{12}(z)$  since it is consistent only with isotrivial geometries and two Z-irreducible polynomials (i.e. two new dimensions) are possible only if they are  $\Phi_8$ and  $\Phi_{10}$  or  $\Phi_5$  (cf. §. [2.5.3\)](#page-24-0).

**Definition 17.** The three integral matrices  $X_0$ ,  $X_1$  and  $X_\infty$  will be called the *Levelt mon*odromy (of class- $E_7$ ) associated to the pair of polynomials  $(P_0(z), P_\infty(z))$  which satisfies the above restrictions.

The Kodaira type of  $X_1$  is  $I_f$  where the integer f is

$$
f \equiv \gcd(A, B). \tag{8.93}
$$

In the list of pairs  $(P_0(z), P_\infty(z))$  satisfying conditions  $(a), (b), (c)$  only  $(\Phi_1^2 \Phi_6, \Phi_2^2 \Phi_3)$  has  $f \notin \{1,2\}$  in facts  $f = 6$ . But this "bad" pair is ruled out by condition (d).

### 8.10.1 Symplectic polarization

With our restrictions on the characteristic polynomials  $(P_0(z), P_\infty(z))$ , the Levelt matrices  $X_a$  preserve the symplectic form

$$
\check{\Omega} = \begin{pmatrix}\n0 & A & -B & C \\
-A & 0 & A & -B \\
B & -A & 0 & A \\
-C & B & -A & 0\n\end{pmatrix} \text{ where } C = A(1 + c_{2,0}) - B c_{1,0} \quad (8.94)
$$

with

$$
\mathsf{Paf}(\check{\Omega}) = A^2 - B^2 + AC, \qquad \det(\check{\Omega}) \equiv \mathsf{Paf}(\check{\Omega})^2 = \mathsf{Res}\big(P_0(z), P_\infty(z)\big). \tag{8.95}
$$

Note that  $\check{\Omega}$  is f times an integral matrix with the GCD of its entries 1. In particular

$$
f^2 | \text{Paf}(\check{\Omega}), \text{ and } \check{\Omega} V = -\text{Paf}(\check{\Omega}) W.
$$
 (8.96)

If  $\tilde{\Omega}$  is principal (more generally Paf( $\tilde{\Omega}$ ) is square-free), f is automatically 1 and the knotted discriminant has Kodaira type  $I_1$ . In general  $\Omega$  is Z-equivalent to a symplectic matrix in the normal form [\[152\]](#page-157-2)

$$
f\begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & g \\ 0 & 0 & -g & 0 \end{pmatrix}, \qquad g \in \mathbb{N}, \quad \text{Paf}(\check{\Omega}) = f^2 g. \tag{8.97}
$$

The values of  $A, B, C, f$  and g for each pair of polynomials satisfying the three conditions  $(a), (b), (c)$  are listed in table [4.](#page-115-0)

When the invariant  $g = 1$ , the Levelt monodromy takes values in  $Sp(4, \mathbb{Z})$  and is a candidate for the  $\mu$ -monodromy associated to a special geometry. Otherwise one should look for other monodromies, conjugate to the Levelt one in  $GL(4,\mathbb{C})$ , which do take value in  $Sp(4, \mathbb{Z})$ . Moreover, as in §.[8.7,](#page-99-0) there may be several  $GL(4, \mathbb{C})$ -conjugate monodromies all taking value in  $Sp(4, \mathbb{Z})$ , but not conjugate in  $Sp(4, \mathbb{Z})$ , which may correspond to inequivalent special geometries. Before going to the general case, we present a few simple examples with  $f = g = 1.$ 

### 8.10.2 Example 1: the {8/7, 10/7} geometry

We know from  $\S$ . [2.5.3](#page-24-0) that a non-isotrivial geometry in rank-2 with two new dimensions must have dimensions  $\{8/7, 10/7\}$  (this is the geometry of the SCFT #21 in the list [\[89\]](#page-153-0)). The axes are regular enhanced with 4 distinct eigenvalues so the geometry must be of class  $\widehat{E}_7$ . The polynomial pair is<sup>[80](#page-114-0)</sup> ( $\Phi_8$ ,  $\Phi_{10}$ ). We have

$$
c_{1,0} = c_{2,0} = 0, \quad c_{1,\infty} = -c_{2\infty} = 1. \tag{8.98}
$$

and we get

$$
\Omega = \begin{pmatrix} 0 & -1 & -1 & -1 \\ 1 & 0 & -1 & -1 \\ 1 & 1 & 0 & -1 \\ 1 & 1 & 1 & 0 \end{pmatrix} \tag{8.99}
$$

which has  $f = g = 1$  so that the Levelt monodromy is valued in  $Sp(4, \mathbb{Z})$ . By a change of basis we put the polarization in the standard form

$$
\mu_0 = \begin{pmatrix} 0 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 \\ -2 & 0 & -1 & -1 \\ 1 & -1 & 0 & 1 \end{pmatrix} \quad \mu_1 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \mu_\infty = \begin{pmatrix} -1 & 0 & -1 & -1 \\ -1 & 1 & 0 & -1 \\ 1 & -1 & -1 & 0 \\ 0 & 1 & 1 & 1 \end{pmatrix} \quad \Omega = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix} \quad (8.100)
$$

which satisfy

$$
\mu_0^4 - \mu_0^3 + \mu_0^2 - \mu_0 + \mathbf{1} = 0, \qquad (\mu_1 - \mathbf{1})^2 = 0, \qquad \mu_\infty^4 = -\mathbf{1},
$$
  
\n
$$
\mu_0 \mu_1 \mu_\infty = \mathbf{1}, \qquad \mu_a^t \Omega \mu_a = \Omega \qquad \text{for } a = 0, 1, \infty.
$$
\n(8.101)

It is clear that the (knotted) discriminant is of type  $I_1$ .

Local regularity along the axes. As a further illustration we check local regularity (which must hold since we know that this geometry exists). We have  $\varkappa = 2$  with  $h^2 =$ 

<span id="page-114-0"></span><sup>&</sup>lt;sup>80</sup> ( $\Phi_8$ ,  $\Phi_5$ ) yields a different lift of the same  $PSp(4, \mathbb{Z})$  monodromy.



<span id="page-115-0"></span>Table 4: Pairs of degree-4 symmetric products of cyclotomic polynomials which have at most one non-simple root of multiplicity 2 and no common zero.

 $(u_2/u_1)^7$  and  $z = u_1^5/u_2^4$ . Along the first axis  $(u_1 = 0)$  the exponents are 7/8, 5/8 so:

$$
a^{\parallel} \sim \sqrt{h^2} \, z^{7/8} = u_1^{7/8} \equiv u_1^{1/\Delta_1}, \qquad a^{\perp} \sim \sqrt{h^2} \, z^{5/8} = u_2 \cdot u_1^{-3/8} \tag{8.102}
$$

along the second axis  $(u_1 = 0)$  the exponents are  $7/10$ ,  $9/10$  and

$$
a^{\parallel} \sim \sqrt{h^2} z^{7/10} = u_2^{7/10} \equiv u_2^{1/\Delta_2}, \qquad a^{\perp} \sim \sqrt{h^2} z^{9/10} = u_1 \cdot u_2^{-1/10}.
$$
 (8.103)

## 8.10.3 Example 2:  $\{3, 5\}$  geometries with one  $I_n^*$  discriminant

This geometry corresponds to SCFTs  $#8, #36, #42$  and  $#46$ . The polynomials are  $(\Phi_2^2 \Phi_3, \Phi_5)$ which give  $f = g = 1$ , so the Levelt monodromy is in  $Sp(4, \mathbb{Z})$  and the knotted discriminant is of type  $I_1$ . The other component of the discriminant is along the axis of dimension 3 and is associated with the  $\mu$ -monodromy

$$
X_0 = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & -3 \\ 0 & 1 & 0 & -4 \\ 0 & 0 & 1 & -3 \end{pmatrix}, \qquad (X_0 + 1)^2 (X_0^2 + X_0 + 1) = 0.
$$
 (8.104)

The corresponding  $\varrho$ -monodromy in  $\mathscr{C}$ ,  $\varrho_0 \equiv X_0^3$  has one Jordan block of size 2 associated to the eigenvalue −1 and two trivial Jordan blocks with eigenvalues +1. It satisfies

$$
\varrho_0^2 \equiv (X_0^3)^2 = \mathbf{1} + 6 v \otimes v^t \, \tilde{\Omega}, \qquad v \equiv (1 \ 2 \ 2 \ 1)^t,\tag{8.105}
$$

which shows that this enhanced discriminant has Kodaira type  $I_3^*$  in agreement with physical expectations.

**Local regularity.** This geometry has  $\varkappa = 1$ ,  $h = u_1^2/u_2$  and  $z = u_2^3/u_1^5$ . Along the second axis the exponents are 2/5 and 1/5

$$
a^{\parallel} \sim h \cdot z^{2/5} = u_2^{1/5}, \qquad a^{\perp} \sim h \cdot z^{1/5} = u_1 \cdot u_2^{-2/5}
$$
 (8.106)

along the first axis  $(u_2 = 0)$  the exponents are  $1/3$  and  $1/2$  so that

$$
a^{\parallel} \sim h \cdot z^{1/3} = u_1^{1/5}, \qquad a^{\perp} \sim h \cdot z^{1/2} \sim u_2^{1/2} \cdot u_1^{-1/2}
$$
 (8.107)

which is the expected behaviour for a discriminant of type  $I_n^*$   $(n \geq 0)$ .

### 8.10.4 Example 3: {4, 10} geometry with a semisimple enhanced discriminant

This is the geometry of the SCFT #3. This geometry has  $\varkappa = 2$  so we have a little ambiguity in the  $Sp(4,\mathbb{Z})$  lift. The pair of polynomials may be  $(\Phi_4\Phi_6,\Phi_{10})$  or  $(\Phi_4\Phi_6,\Phi_5)$ . Both pairs have  $f = g = 1$  so that the Levelt monodromy is already in  $Sp(4, \mathbb{Z})$ . For, say, the first pair we put the polarization in the standard form

$$
\Omega \equiv \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix} = -S^t \check{\Omega} S, \text{ where } S = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 1 \end{pmatrix}
$$
 (8.108)

In this canonical basis the monodromy around the axis of dimension 4 reads

$$
\varrho_0 = -S^{-1}X_0^2 S = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & -1 & 1 & 0 \\ 0 & -1 & -1 & -1 \end{pmatrix} \tag{8.109}
$$

which is semisimple with eigenvalues  $(\zeta_6, \zeta_6^{-1}, 1, 1)$  i.e. a discriminant of type  $II^*$  or  $II$ . The regularity of this example was already discussed in §[.8.6.2;](#page-98-0) as shown there regularity fixes the signs in the lift to  $Sp(4, \mathbb{Z})$  while the axis of dimension 4 must be a discriminant of type  $II^*$ , as physically expected.

### **8.10.5** Example 4: the  ${5/4, 3/2}$  geometry

This is the only other non-isotrivial geometry in the list of examples [\[89\]](#page-153-0) with an irreducible discriminant. It corresponds to the polynomials  $(\Phi_3 \Phi_6, \Phi_5)$  which have  $f = g = 1$  so that the Levelt monodromy is in  $Sp(4, \mathbb{Z})$ . This geometry has  $\varkappa = 1$ ,  $h = u_2^4/u_1^4$ ,  $z = u_1^6/u_2^5$ . The exponents for the first axis are 4/5 and 3/5 so that

$$
a^{\parallel} \sim h \cdot z^{4/5} = u_1^{4/5}, \qquad a^{\perp} \sim h \cdot z^{3/5} = u_2 \cdot u_1^{-2/5}
$$
 (8.110)

while around the second axis the exponents are  $2/3$  and  $5/6$  which gives

$$
a^{\parallel} \sim h \cdot z^{2/3} = u_2^{2/3}, \qquad a^{\perp} \sim h \cdot z^{h/6} = u_1 \cdot u_2^{-1/6}.
$$
 (8.111)

# 8.11  $\widehat{E}_7$  class: non-Levelt principal polarizations

The Levelt integral monodromy is a solution to our inverse problem only when  $\Omega$  is an integral multiple of a principal integral matrix, that is, when  $q = 1$ . Otherwise we have to search for representations conjugate to the Levelt one in  $GL(4,\mathbb{C})$  which take value in  $Sp(4,\mathbb{Z})$  much as we did in §. [8.7](#page-99-0) for the Pochhammer ODE monodromy. As in that discussion, there may be more than one inequivalent  $Sp(4, \mathbb{Z})$ -valued representations for a given rigid  $GL(4, \mathbb{C})$ monodromy. Even when the Levelt monodromy is principal it may be not the only one valued in  $Sp(4, \mathbb{Z})$  for the given polynomial pair.

### 8.11.1 Example: a  $\{6, 8\}$  geometry

Consider the following two pairs of polynomials  $(\Phi_2^2 \Phi_6, \Phi_8)$  and  $(\Phi_1^2 \Phi_3, \Phi_8)$  which have  $f = 1$ and  $g = 2$ . The first one is naturally interpreted as describing  $\varkappa = 2$  geometry with dimensions  $\{6, 8\}$  where the axis of dimension 6 has type  $I_{n>0}^*$ . The monodromy of the

second pair is obtained from the first one by a sign flip  $\mu_0 \to -\mu_0$ ,  $\mu_\infty \to -\mu_\infty$  so it gives a different lift of the same  $PSp(4, \mathbb{Z})$  monodromy. We have

$$
-2\Omega \equiv \begin{pmatrix} 0 & -2 & 0 & 0 \\ 2 & 0 & 0 & 0 \\ 0 & 0 & 0 & -2 \\ 0 & 0 & 2 & 0 \end{pmatrix} = S^t \check{\Omega} S, \quad \text{where} \quad S = \begin{pmatrix} 2 & 1 & 1 & 1 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}
$$
 (8.112)

so that conjugating the Levelt representation with  $S^{-1}$  we get the three matrices

$$
\mu_0 = \begin{pmatrix} -1 & -1 & -1 & 0 \\ 2 & 1 & 2 & 0 \\ 0 & 1 & -1 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix} \qquad \mu_1 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix} \qquad \mu_\infty = \begin{pmatrix} 1 & 1 & 0 & -1 \\ -2 & -1 & 0 & 0 \\ 2 & 1 & 1 & 2 \\ -2 & -1 & -1 & -1 \end{pmatrix} \tag{8.113}
$$

which satisfy

$$
\mu_0 \mu_1 \mu_\infty = \mathbf{1}, \quad \mu_a^t \Omega \mu_a = \Omega \quad (a = 0, 1, \infty)
$$
  

$$
\mu_\infty^4 + \mathbf{1} = (\mu_0^2 + 1)^2 (\mu_0^2 - \mu_0 + \mathbf{1}) = 0
$$
\n(8.114)

One has

$$
\mu_0^6 = \mathbf{1} + 4v \otimes v^t \Omega \qquad v = \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} \tag{8.115}
$$

which means that the axis of dimension 6 is a discriminant of type  $I_2^*$  as expected for model #44 of [\[89\]](#page-153-0). There should be another  $GL(4,\mathbb{C})$  equivalent but not  $Sp(4,\mathbb{Z})$ -equivalent solution which describes the geometries of the models  $#2, #33, #39$  where the type is  $I_6^*$ . Presumably they correspond to another  $Sp(4, \mathbb{Z})$  inequivalent integral realization of the same  $GL(4,\mathbb{C})$  monodromy.

# 8.12  $\widehat{E}_7$  class: Kodaira type of knotted discriminants

**Fact 34.** All knotted discriminants of class  $\widehat{E}_7$  geometries are of Kodaira type  $I_1$ .

This is confirmed from the known examples [\[89\]](#page-153-0).

*Argument.* Any integral symplectic monodromy which is conjugate in  $GL(4,\mathbb{C})$  to the Levelt one has the local monodromomy  $\mu_1$  associated to the  $\delta = 1$  branch of type  $I_n$  where  $n \mid f$ . Indeed  $\mu_1 = 1 \text{ mod } n$  and hence

$$
\mu_{\infty} = (\mu_0)^{-1} \bmod n \tag{8.116}
$$

and  $P_{\infty}(z) = P_0(z) \mod n$ , while f is the largest integer such that  $P_{\infty}(z) = P_0(z) \mod f$ . Looking to the table [4](#page-115-0) we see that  $f = 1$  except for the pair  $(\Phi_1^2 \Phi_6, \Phi_2^2 \Phi_3)$  which has  $f = 6$ and seven pairs which have  $f = 2$ . The pair  $(\Phi_1^2 \Phi_6, \Phi_2^2 \Phi_3)$  cannot correspond to any regular special geometry since both axes would be non-semisimple, and all consistent assignments of dimensions would lead to  $3 \times \infty$  which implies an isotrivial geometry which is inconsistent with the presence of 3 non-semisimple discriminants. Equivalently this pair is ruled out by requirement (d). Three of the pairs with  $f = 2$ , namely  $(\Phi_1^2 \Phi_3, \Phi_2^2 \Phi_6)$ ,  $(\Phi_1^2 \Phi_4, \Phi_8)$ ,  $(\Phi_2^2 \Phi_4, \Phi_8)$  should also be excluded because all consistent dimension assignments lead to

 $x > 2$  (requirement (d)). The four remaining pairs with  $f = 2$  are

<span id="page-119-0"></span>
$$
(\Phi_1^2 \Phi_6, \Phi_3 \Phi_4) \quad (\Phi_2^2 \Phi_6, \Phi_3 \Phi_4) \quad (\Phi_1^2 \Phi_3, \Phi_6 \Phi_4) \quad (\Phi_2^2 \Phi_3, \Phi_6 \Phi_4). \tag{8.117}
$$

In a would-be special geometry associated with one of these pairs the first axis is a nonsemisimple discriminant of dimension in the list  $\{3/2, 6/5, 3, 6\}$ . The second axis is a semisimple discriminant: a priori the Albanese variety may have an automorphim of degree 3 or 4, but the first possibility would lead to  $3 \mid \varkappa$ , hence to a contradiction. Let us focus, say, on the first pair. The dimensions pairs with  $1/\Delta_i$  of the correct order in  $\mathbb{Q}/\mathbb{Z}$  are  $\{\Delta_1, \Delta_2\} = \{6, 4\},$  $\{6/5, 4\}, \{6, 4/3\}, \text{ and } \{6/5, 4/3\}$  but the fractional dimensions may be ruled out by the values of the degree  $(d_1, d_2)$ . We remain with  $\{6, 4\}$  where the second axis of type  $IV^*$ ,  $II^*$  (or, less likely, IV, II). Then  $\varkappa = 2$ ,  $h^2 = u_1/u_2$ , and  $z = u_2^3/u_1^2 \text{ mod } 1$ . Along the second axis the exponents are  $\pm 1/4$  and  $\pm 1/3$ , then as  $u_1 \rightarrow 0$ 

$$
a_{\parallel} \sim \sqrt{h^2} \, z^{1/4} = \frac{u_1^{1/2}}{u_2^{1/2}} \cdot \frac{u_2^{3/4}}{u_1^{1/2}} = u_2^{1/4}
$$
\n
$$
\text{either} \quad a_{\perp} \sim \sqrt{h^2} \, z^{1/3} = \frac{u_1^{1/2}}{u_2^{1/2}} \cdot \frac{u_2}{u_1^{2/3}} = u_2^{1/2} \, u_1^{-1/6}
$$
\n
$$
\text{or} \quad a_{\perp} \sim \sqrt{h^2} \, z^{-1/3} = \frac{u_1^{1/2}}{u_2^{1/2}} \cdot \frac{u_1^{2/3}}{u_2} = u_2^{-3/2} \, u_1^{7/6}
$$
\n
$$
(8.118)
$$

and neither expression of  $a_{\perp}$  is consistent with a regular total space with a singular fiber of type  $IV^*$  or  $IV$  (nor  $II^*, II$ ). The same argument applies to the second pair which has the same putative dimensions. For the other two pair one has dimensions  $\{3, 4\}, \, \varkappa = 1$ ,  $h = u_2/u_1, z = u_1^4/u_2^3$  and as  $u_1 \to 0$ 

$$
a_{\parallel} \sim h z^{1/4} = u_2^{1/4}, \qquad a_{\perp} \sim \begin{cases} h z^{1/6} = u_1^{-1/3} u_2^{1/2} \\ h z^{-1/6} = u_1^{-5/2} u_2^{3/2} \end{cases}
$$
(8.119)

Neither expression of  $a_{\perp}$  is consistent with a bona fide special geometry. Therefore all consistent pairs have  $f = 1$  and a fortiori  $n = 1$ .  $\Box$ 

### 8.13 The puzzling SCFT  $\#45$

The SCFT #45 in the list [\[89\]](#page-153-0) is puzzling on the physical side for reasons explained in. that reference. Its most plausible (or less implausible) properties are listed in [\[89\]](#page-153-0): Coulomb dimensions  $\{4, 6\}$ , a knotted discriminant of type  $I_1$ , the axis of dimension 6 is a discriminant of type  $I_2$ , and the one of dimension 4 is a semisimple discriminant of (possibly) type  $II^*$ . This list of properties from physical considerations look a bit puzzling from the geometrical side too. Let us summarize the situation.

If we assume irreducibility, the presence of a knotted  $I_1$  discriminant suggests a class- $\widehat{E}_7$ 

geometry. This requires the characteristic polynomials of  $\mu_0$  and  $\mu_\infty$  to be coprime, hence in the list of table [4.](#page-115-0) From the dimensions, the fact that  $\mu_{\infty}$  has a non-trivial Jordan block while  $\mu_0$  is semisimple will suggest as candidate polynomials pairs (taking into account the various  $Sp(4, \mathbb{Z})$  lifts) precisely the ones in eq.[\(8.117\)](#page-119-0) which are not consistent with regularity. Moreover the most natural pair  $(\Phi_2^2 \Phi_6, \Phi_3 \Phi_4)$  has a Levelt monodromy in  $Sp(4, \mathbb{Z})$  which, however, has a knotted discriminant of the wrong type:  $I_2$  instead of  $I_1$ .

To solve the conundrum we use regularity backwards: we assume the minimal polynomial of  $\mu_{\infty}$  to be  $\Phi_2^2 \Phi_6$ , since this identification is not problematic, and ask for which characteristic polynomial of  $\mu_0$  we would have both regularity and the physically expected properties. In particular we already know that the polynomial  $\Phi_0(z)$  should have the form  $\Phi_4(z) Q(z)$  for some degree-2 *cyclotomic* polynomial  $Q(z)$ . The purported geometry has dimensions  $\{4, 6\}$ , so  $\varkappa = 2$ ,  $h^2 = u_2/u_1$  and  $z = u_1^3/u_2^2$ . Assuming the first axis  $(u_2 = 0)$  is a discriminant of type  $II^*$  we must have

$$
a^{\parallel} \sim \sqrt{h^2} \cdot z^{\alpha_{\parallel}} = u_1^{1/4}, \qquad a^{\perp} \sim \sqrt{h^2} \cdot z^{\alpha_{\perp}} = u_2^{1/6}
$$
 (8.120)

for some exponents  $\alpha_{\parallel}$  and  $\alpha_{\perp}$  of  $\mu_0$ . The first relation yields  $\alpha_{\parallel} = 1/4$  consistently with the factor  $\Phi_4(z)$  of  $P_0(z)$ . The second relation yields  $\alpha_{\perp} = 1/6$  which implies  $Q(z) = \Phi_6$ . We conclude

$$
P_0 = \Phi_4 \Phi_6
$$
,  $P_\infty = \Phi_2^2 \Phi_6$ ,  $gcd(P_0, P_\infty) = \Phi_6$ , (8.121)

which leads to a reducible geometry. We have studied them in §.[8.1.](#page-81-0) Since this geometry is rigid, it should be one of the quasi-isotrivial possibilities listed in §. [8.1.3](#page-83-0) (see also appendix [F\)](#page-146-0). Indeed the geometric data of model  $#45$  exactly matches with **Fact [20\(](#page-83-1)3)**: same dimensions, same Kodaira types of the three special divisors. The same conclusion will follows from Fact [31](#page-109-0).

However unlikely this situation appears, it is not ruled out by our analysis since #45 does have all the characteristics expected for a quasi-isotrivial geometry. While this is certainly not the last word about the geometry of model #45, it gives a plausible geometric scenario consistent with the physical intuition.

**Scenario.** The special geometry of the SCFT  $#45$  is quasi-isotrivial.

# <span id="page-120-0"></span>8.14  $\widehat{E}_7$  class: allowed polynomial pairs

A part for the case of the troublesome dimension pair  $\{4, 6\}$  (and perhaps its close cousin  $\{3,4\}$ , assuming  $\mu$ -rigidity the classification of rigid (non-cover) non-isotrivial geometries is reduced to: (i) listing the allowed pairs of coprime polynomials  $(P_0, P_\infty)$  (ii) checking local regularity, then *(iii)* for each such pair solve the Diophantine problem of finding all  $S \in \mathbb{C}^{\times} \backslash GL(4,\mathbb{C})/Sp(4,\mathbb{Z})$  such that  $S<sup>t</sup> \check{\Omega} S$  is a multiple of an integral skew-symmetric matrix of determinant 1 while  $\mu_a = S^{-1} X_a S$   $(a = 0, 1, \infty)$  are integral matrices (here  $X_a$  are the Levelt matrices).

The results of part *(i)* of the program are listed in the last column of table [4.](#page-115-0) There are various obvious reasons to rule out a polynomial pair. The first one is the non-existence of a consistent pair of dimensions  $\{\Delta_1, \Delta_2\}$  potentially associated to the given pair. If, say,  $\Delta_2$  is a new dimension the list of allowed  $\Delta_1$  is short and this allows to eliminate all pairs which do not correspond to an allowed set of dimensions. Then one eliminates all pairs which would correspond to dimensions with  $\varkappa \notin \{1,2\}$  since the corresponding geometries would be isotrivial which is in contradiction with the presence of a  $I_1$  discriminant. For instance,  $\Phi_{12}$ cannot appear since it would automatically imply that the geometry is isotrivial. The pairs which are eliminated for dimensional reasons have no:dim in the last column of the table. Then we can eliminate some pairs based on local regularity. We have already eliminated the four pairs in eq.[\(8.117\)](#page-119-0). On the other extremum there is the list of pairs which are obviously allowed, namely the ones associated to known SCFTs. The corresponding SCFT are written in the table (up to a small ambiguity).

After this exercise, we remain with two pairs which are neither obviously ruled out nor obviously allowed:

<span id="page-121-1"></span>
$$
(\Phi_3 \Phi_6, \Phi_8) \qquad (\Phi_4 \Phi_6, \Phi_8) \tag{8.122}
$$

In both cases the second axis is regular of dimension  $\Delta_2 = 8/k$  where  $gcd(k, 8) = 1$ . The first axis has dimension  $\Delta_1$  (written in minimal form)  $3/\ell$  or  $6/\ell$  for the first pair and  $4/\ell$ or  $6/\ell$  for the second one. The only allowed dimension pairs with these properties are

<span id="page-121-2"></span>
$$
\{4/3,8/3\}, \{8/3,4\}, \{4,8\}, \{6/5,8/5\}, \{6,8\}. \tag{8.123}
$$

All but the first pair are realized in the geometries of known  $SCFT^{81}$  $SCFT^{81}$  $SCFT^{81}$ . The first three pairs have  $x = 4$  or  $4/3$  and so are isotrivial. We claim that the last two sets of dimensions cannot be realized by monodromies with the polynomials in eq.[\(8.122\)](#page-121-1). Note that in the known geometries with these dimensions the axis of dimension  $\neq 8$  is a discriminant of non-semisimple type, while the polynomials  $(8.122)$  are satisfied by semisimple matrices.

Let us justify the claim. The last two dimension pairs in [\(8.123\)](#page-121-2) have both  $\varkappa = 2$  and  ${d_1, d_2} = {3, 4}.$  Hence

$$
\varrho_0 = \pm \mu_0^3 \text{ with eigenvalues } \begin{cases} \pm (1, 1, -1, -1) & \text{for } (\Phi_3 \Phi_6, \Phi_8) \\ \pm (-1, -1, i, -i) & \text{for } (\Phi_4 \Phi_6, \Phi_8) \end{cases} \tag{8.124}
$$

so the discriminant along the first axis has Kodaira type  $I_0^*$  and, respectively,  $III$  or  $III^*$ .

For both dimension pairs  $z = u_1^4/u_2^3$ , while for the first (resp. second) dimension pair  $h^2 = (u_2/u_1)^5$  (resp.  $h^2 = u_2/u_1$ ). We look for the exponents  $\alpha_{\perp}$  required to get a local solution along the first axis  $(u_2 = 2)$  which matches with the Kodaira type above in the

<span id="page-121-0"></span><sup>&</sup>lt;sup>81</sup> The dimension pairs are realized in the following SCFTs:  $\{8/3, 4\}$  in #58,  $\{4, 8\}$  in #4, 52, 62, 65,  $\{6/5, 8/5\}$  in #19, and  $\{6, 8\}$  in #2, 33, 39, 44.

several cases:



The only meaningful exponent is  $2/3$  in the third row. Hence  $(\Phi_4 \Phi_6, \Phi_8)$  is ruled out, while the pair  $(\Phi_3 \Phi_6, \Phi_8)$  may correspond to a special geometry with an enhanced discriminant of type  $I_0^*$  provided it has the exotic-looking dimension pair  $\{6/5, 8/5\}.$ 

## 8.15 Summary of rank-2

The main result of this section is that the list of known rank-2 special geometries ( $\equiv$  the SW geometries of known rank-2 SCFTs) is "essentially" complete in the sense that the only unknown geometries whose existence we have not yet ruled out are:

- geometries which are weakly equivalent to a known one, i.e. they have a monodromy which is  $GL(4,\mathbb{C})$ -equivalent but not  $Sp(4,\mathbb{Z})$ -equivalent to a known one;
- class- $E_7$  geometries with new dimensions not ruled out in §. [\(8.9.3\)](#page-111-1). A natural candidate follows from the explicit monodromy in eqs.[\(8.85\)](#page-111-2)-[\(8.87\)](#page-111-3).
- other reducible geometries with dimensions  $\{3, 4\}$  or  $\{4, 6\}$  besides #45. (This class deserves a more detailed study);
- additional cover geometries (looks unlikely);
- rigid geometries with a non rigid monodromy (which conjecturally do not exist).

# 9 The inverse problem in rank  $r \geq 3$

At first sight the inverse problem may seen impossibly hard when  $r > 2$ . Even the specification of the inverse data  $(\{\Delta_i\}, \{\mathcal{D}_a\}, \{\varrho_a\})$  may look bewildering because now the discriminant components  $\mathcal{D}_a$  are quasi-cones over weighted projective hypersurfaces  $\mathcal{S}_a \subset \mathscr{P}$  which a priori may be very complicated whereas for  $r = 2$  they were mere points in  $\mathbb{P}^1$ .

Luckily the story is much better than it may appear. The two basic ingredients that we used in  $r = 2$ , the rigidity of the Abelian family over  $\mathscr{S} \equiv \mathscr{P} \setminus \mathscr{S}$  and the special properties of rigid representations of the fundamental group  $\pi_1(\hat{\mathscr{S}})$  apply (the second one a bit conjecturally) to all quasi-projective bases  $\mathscr S$  of any dimension. In the rank-2 case we used the first property to conclude that when the geometry is not decomposable, or a cover, and 2 is not a Coulomb dimension (the only case which remains open assuming the Folk-theorem) the special divisor  $\mathscr S$  must consist of precisely 3 points in  $\mathbb P^1$ . One has to extend that argument to higher rank; this leads to:

**Question 3.** Which hypersurfaces complements  $\mathbb{P}(q_1, \ldots, q_r) \setminus \mathscr{S}$  in a weighted projective space are the higher dimensional analogues of the sphere with three punctures?

If we can give an a priori bound on how bad the singularities of the special divisor  $\mathscr S$ may be, the set of such special divisors should be finite for each  $r$ -tuple of well-posed weights  $(q_1, \ldots, q_r)$  which form themselves a finite set since only finitely many dimensions r-tuple  ${\{\Delta_1,\ldots,\Delta_r\}}$  are allowed in a given rank r.

We start by making some simple considerations about the **Question**.

## 9.1 The special divisor  $\mathscr S$

### 9.1.1 Conditions on  $\mathscr S$  from rigidity and "stratification"

We assume the Coulomb branch to be smooth and the geometry to be primary. Just as in rank-2 we have the

**Principle** (Rigidity). The space M of deformations of the (reduced) divisor  $\mathscr{S} \subset \mathscr{P}$  which preserve the group  $\pi_1(\mathscr{P} \setminus \mathscr{S})$  should have dimension equal to the multiplicity of 2 as a Coulomb dimension. In facts,  $M$  should coincide with the conformal manifold of the special geometry.

<span id="page-123-1"></span>Example 9.1. We are mainly interested in rigid special geometries, but below we shall consider also examples coming from the opposite extremum, namely a dimension  $r$ -tuple of the form  $\{2, 2, \ldots, 2\}$ . In this instance we expect finitely-many r-parameter families  $\{\mathscr{S}_m\}_{m\in\mathscr{M}}$  of special divisors<sup>[82](#page-123-0)</sup>. In view of the Gaiotto construction [\[110\]](#page-154-1), the **Folk-theorem** yields an explicit description of them: there is one family per pair  $(q, n)$  of non-negative integers such that  $3g - 3 + n = r$ . The conformal manifold of the  $(g, n)$ -family is the moduli space  $\mathcal{M}_{g,n}$  of Riemann surfaces of genus g with n-punctures  $p_i$ , while  $\mathscr P$  is the linear system of the divisor  $2K + \sum_i p_i$  which is a copy of  $\mathbb{P}^{r-1}$ . Fix  $m \in \mathcal{M}_{g,n}$  and consider the corresponding marked curve  $\Sigma_m$ . A point  $z \in \mathscr{P}$  corresponds to a quadratic differential  $\phi_{m,z}$  on  $\Sigma_m$  with at most simple poles at the punctures (modulo overall normalization). Consider the curve  $C_{m,z}$  in the (holomorphic) cotangent space  $T^*\Sigma_m$  (with fiber coordinate y) of equation  $y^2 = \phi_{m,z}$ .  $\mathscr{S}_m \subset \mathbb{P}^{r-1}$  is the locus of z's with singular curve  $C_{m,z}$ .

<span id="page-123-0"></span> $82$  In this situation all components  $S_a$  arise from the discriminant.

Rigidity is not the only restriction on  $\mathscr{S}$ . We have already discussed in §.[3.2](#page-47-0) other conditions following from the physical intuition of "stratification" and in particular that each irreducible component of the discriminant admits a polynomial parametrization. Some examples of divisor satisfying these constraints are in order.

**Example 9.2.** Consider the rank-3 special geometries of dimensions  $\{2, 2, 2\}$ . As described in **Example [9.1](#page-123-1)** we have three families of them with  $(q, n) = (0, 6), (1, 3), (2, 0)$ . We focus on the simpler  $(0, 6)$  family. We have a quadratic differential with simple poles at  $(say)$  $z_1 \equiv 0, z_2 = 1, z_3 = -1, z_4, z_5, z_6 \ (z_4, z_5, z_6 \text{ being local coordinates in } \mathcal{M}_{0,6})$ 

<span id="page-124-1"></span>
$$
\phi_{m,u} = \frac{u_1 z^2 + u_2 z + u_3}{z(z^2 - 1)(z - z_4)(z - z_5)(z - z_6)} dz^2
$$
\n(9.1)

where  $(u_1 : u_2 : u_3)$  are homogeneous coordinates on  $\mathscr{P}$ . The discriminant is the union of the loci where a zero of the quadratic polynomial in the numerator collides with a zero of the denominator and the locus where the two zeros of the numerators coalesce

$$
\mathcal{D}: \quad (u_2^2 - 4u_1u_3) \prod_{a=1}^6 (u_1 z_a^2 + u_2 z_a + u_3) = 0. \tag{9.2}
$$

Let us check that this divisor  $\mathcal D$  has the properties expected on physical grounds. We start with "rigidity".  $\mathcal D$  is the union of a smooth planar conic  $C$  and six lines

$$
L_a \equiv \{u_1 z_a^2 + u_2 z_a + u_3 = 0\} \subset \mathbb{C}^3, \quad a = 1, \dots, 6 \tag{9.3}
$$

each one of which is isomorphic to a copy of  $\mathbb{P}^1$ . Modulo projective equivalence there is a unique smooth conic. The six lines are all tangent to the conic  $C$ , and each line  $L_a$  is uniquely determined by its point of tangency  $p_a \in C$ . Thus  $\mathscr{S}_m$  is determined by the six point  $\{p_a\}$ . However the automorphism group  $O(3,\mathbb{C})$  of C acts on the points  $\{p_a\}$ ; since the action of  $O(3, \mathbb{C})$  on C is 3-transitive<sup>[83](#page-124-0)</sup>, we can fix 3 points, and the actual deformation space of  $\mathscr{S}_m$  is 3-dimensional as required by physics.

<span id="page-124-2"></span>**Remark 23.** A cheap curve  $C \subset \mathbb{P}^2$  with the property that the deformation space  $\mathcal{M}$ preserving the topology of its complement has dimension s is a smooth conic together with  $s + 3$  tangent lines.

Let us check "stratification" on this example. Restricting on the discriminant line  $L_a$ , the numerator has a zero at  $z = z_a$  which cancels a zero in the denominator, so that  $\phi_2$  becomes the quadratic differential of the model with  $(g, n) = (0, 5)$ ; hence the restriction on  $L_a$  yields back the first  $\{2, 2\}$  special geometry in eq.[\(8.56\)](#page-102-0). The restriction on the conic C produces the non-smooth special geometry discussed in §. ??, i.e. a  $\mathbb{Z}_2$  quotient of the free geometry.

<span id="page-124-0"></span><sup>&</sup>lt;sup>83</sup> This is the complexified version of the well-known *holography* statement that the Euclidean conformal group acts on the *d*-sphere  $S^d$  in a 3-transitive fashion.

Indeed

$$
\phi_{m,u}\Big|_{C} = \frac{u_1(z - u_2/2u_1)^2}{\prod_{a=1}^6 (z - z_a)} dz^2
$$
\n(9.4)

and setting  $y = \sqrt{u_1}(z - u_2/2u_1)\tilde{y}/\prod_a(z - z_a)$ , we get the equation

$$
\tilde{y}^2 = \prod_{a=1}^6 (z - z_a), \qquad \lambda = \sqrt{\phi_2} = (\sqrt{u_1} \, z - \sqrt{u_3}) \frac{dz}{\tilde{y}}.
$$
\n(9.5)

**Example 9.3.** Consider the rank-4 SCFT with dimensions  $\{2, 2, 2, 2\}$  associated to the sphere with 7 punctures. Eq. $(9.1)$  gets replaced by

<span id="page-125-0"></span>
$$
\phi_{m,u} = \frac{u_1 z^3 + u_2 z^2 + u_3 z + u_4}{\prod_{a=1}^7 (z - z_a)} dz^2, \qquad z_0, z_1, z_2 = 0, +1, -1 \tag{9.6}
$$

with discriminant  $\mathscr{S} \subset \mathbb{P}^3$  of equation

$$
\left(-27u_1^2u_4^2+18u_1u_2u_3u_4-4u_1u_3^3-4u_2^3u_4+u_2^2u_3^2\right)\prod_{a=1}^7\left(u_1z_a^3+u_2z_a^2+u_3z_a+u_4\right)\tag{9.7}
$$

the union of 7 planes  $L_a$  and a *singular* quartic  $Q_4$ , with the polynomial parametrization

$$
(u_1: u_2: u_3: u_4) = (t_1^3: (2t_2 + t_3)t_1^2: (2t_2t_3 + t_2^2)t_1: t_2^2t_3).
$$
\n(9.8)

 $Q_4$  and the a-th plane are tangent along the the image of the line  $t_2 + z_a t_1 = 0$ . If we restrict to a plane  $L_a$ , a zero in the numerator of  $(9.6)$  cancels a zero in the denominator and we get back the quadratic differential in one rank less, eq.[\(9.1\)](#page-124-1), as expected from stratification. We may look at the discriminant restricted on a line  $L_a$ , say to  $u_4 = 0$  we get a reducible curve: one component is the correct curve  $u_2^2 - 4u_1u_3 = 0$  in one rank less and the other is a "spurious" extra double line  $u_3^2 = 0$  tangent to the conic. This reflects the fact that we are (wrongly!) enforcing that the curve depends essentially on 4 parameters, and so it reduction on  $u_4 = 0$  must contain 7 tangent lines, whereas the actual discriminant depends only on 3 conformal moduli and hence should contain only 6 tangent line. When both the original geometry and the reduced geometry along a discriminant component are rigid we do not expect this mismatch to show up. We expect that for a rigid geometry the restriction of the special locus to a discriminant component should be a legitimate special locus in one less rank. We call this the *hereditary* principle.

<span id="page-125-1"></span>**Remark 24.** A priori the hereditary principle applies only to the components of  $\mathscr S$  which arise from the discriminant, not necessarily to the ones associated with regular R-enhanced divisors. However the only conditions that may not hold are the algebraic properties of the fundamental group of the complement which now may be Abelian etc. corresponding to (say) an isotrivial geometry along the R-enhanced divisor. For instance, in  $r = 3$  the

normalization of a R-enhanced divisor may be  $\mathbb{P}^1$  with just two punctures not three as in the "legitimate" rank-2 case.

**Example 9.4.** More generally, for the rank- $r$  theory with all dimensions equal 2 whose Gaiotto surface is a sphere with  $r + 3$  punctures  $\mathscr S$  is the union of  $r + 3$  hyperplanes

$$
H_a = \left\{ \sum_{s=1}^r u_s z_a^{r-s} = 0 \right\} \subset \mathbb{P}^{r-1}, \qquad a = 1, \dots, r+3 \tag{9.9}
$$

plus a singular hypersurface  $Y_{2r-4}$  of degree  $2r-4$  given by the discriminant of the degree r − 1 polynomial  $P(z) = \sum_{s=1}^{r} u_s z^{r-s} = 0$  which admits the rational parametrization

$$
(u_1: u_2: \dots: u_r) = (t_0^{r-1}, t_0^{r-2}e_1, t_0^{r-2}e_2, \dots, t_0e_{r-2}, e_{r-1}), \quad (t_0, t_1, \dots, t_{r-2}) \in \mathbb{P}^1 \quad (9.10)
$$

where  $e_k$  is the k-th elementary polynomials in the  $(r-1)$  variables  $(t_1, t_1, t_2, \ldots, t_{r-2})$  with  $t_1$  repeated twice. The validity of the hereditary principle is obvious.

## <span id="page-126-1"></span>9.1.2 Conditions on  $\mathscr S$  from  $\pi_1(\mathscr P \setminus \mathscr S)$

The above constraints from rigidity and "stratification" are not the only conditions on the divisor  $\mathscr{S} \subset \mathscr{P}$ . The requirement that the geometry is indecomposable and non-isotrivial (nor quasi-isotrivial) sets additional conditions on  $\mathscr{S}$ . Indeed the complement  $\mathscr{P}\setminus\mathscr{S}$  cannot be a product of lower dimensional quasi-projective space and its fundamental group  $\pi_1(\mathscr{P} \setminus$  $\mathscr{S}$  cannot be finite, Abelian, solvable, or a non-trivial product, etc. To see what this implies we review the fundamental group of the complement of the hypersurface  $\mathscr{S}$  [\[80\]](#page-153-1).

## <span id="page-126-2"></span>9.2 The fundamental group of a hypersurface complement

For special geometry in rank-r we are interested in the fundamental group  $\pi_1(\mathscr{P} \setminus \mathscr{S})$  of a weighted projective hypersurface  $\mathscr S$  in a weighted projective space  $\mathscr P = \mathbb P(q_1, \ldots, q_r)$  which we assume well-formed.

The first observation is that we can always reduce ourselves to hypersurface complements in the *ordinary* projective space<sup>[84](#page-126-0)</sup>  $\mathbb{P}^{r-1}$ . In our situation the hypersurface  $\mathscr S$  can be assumed to have form

$$
\mathcal{S} = \{u_1 = 0\} \cup \{u_2 = 0\} \cup \cdots \cup \{u_r = 0\} \cup \{h(u_1, \ldots, u_r) = 0\}
$$
(9.11)

where  $\{u_i = 0\}$  are (typically) the R-enhanced divisors and  $\mathscr{K} \equiv \{h = 0\}$  is a divisor (typically reducible) which does not lie in any linear sub-space spanned by the coordinate axes. We use the notation  $\mathscr{\tilde{P}} \equiv \mathscr{P} \setminus (u_1u_2\cdots u_r)$ ,  $\mathbb{P}^{r-1} \equiv \mathbb{P}^{r-1} \setminus (x_1x_2\cdots x_r)$ , so that

<span id="page-126-0"></span><sup>&</sup>lt;sup>84</sup> Below we shall use the symbols  $u_1, \ldots, u_r$  and  $x_1, \ldots, x_r$  for, respectively, the homogeneous coordinates in the weighted and ordinary projective space.

 $\mathscr{P} \setminus \mathscr{S} \equiv \tilde{\mathscr{P}} \setminus \mathscr{K}$ . By construction we have an unbranched finite cover

$$
\varpi: \ddot{\mathbb{P}}^{r-1} \to \ddot{\mathscr{P}}, \qquad \varpi: x_i \mapsto u_i = x_i^{q_i}, \qquad (9.12)
$$

with deck group  $\mathbb{Z}_{q_1} \times \mathbb{Z}_{q_2} \times \cdots \times \mathbb{Z}_{q_r}$  so we have an exact sequence of groups

$$
1 \to \prod_{a} \mathbb{Z}_{q_a} \to \pi_1(\tilde{\mathscr{P}} \setminus \mathscr{K}) \to \pi_1(\tilde{\mathbb{P}}^{r-1} \setminus \varpi^{-1}(\mathscr{K}))) \to 1
$$
\n(9.13)

so that, modulo a finite group, we can replace  $\pi_1(\mathscr{P} \setminus \mathscr{S})$  by the fundamental group of the complement in  $\mathbb{P}^{r-1}$  of the hypersurface  $\mathscr W$  of equation

$$
\mathscr{W}: \quad x_1 \, x_2 \, \cdots x_r \, h(x_1^{d_1}, x_2^{d_2}, \ldots, x_r^{d_r}) = 0. \tag{9.14}
$$

Note that some arguments, e.g. the structure theorem of VHS, are insensitive to finite groups.

#### 9.2.1 Zariski and Kempen-Zariski theorems

The fundamental group  $\pi_1(\mathbb{P}^{r-1} \setminus \mathscr{W})$  for  $r \geq 3$  is described by two theorems due to Zariski and Kempen-Zariski respectively (see e.g.[\[80\]](#page-153-1) or [\[100\]](#page-154-0)). The first theorem states that we can effectively reduce the problem to two-dimensions (i.e.  $r = 3$ ).

**Theorem** (Zariski). W is a hypersurface in  $\mathbb{P}^{r-1}$  with complement  $Y \equiv \mathbb{P}^{r-1} \setminus W$  and  $E_k \subset \mathbb{P}^{r-1}$  a generic k-dimensional linear subspace. The morphism induced by inclusion

$$
\iota_k \colon \pi_1(E_k \cap Y) \to \pi_1(Y) \tag{9.15}
$$

is surjective for  $k = 1$  and an isomorphism for  $k > 2$ . In particular:

- (a) The computation of  $\pi_1(Y)$  is reduced to the case  $r = 3$  i.e. to the complement of a planar curve  $E_2 \setminus E_2 \cap \mathscr{W}$ ;
- (b) The group  $\pi_1(Y)$  has a natural set of  $d = \deg(W) \equiv \#\{E_1 \cap W\}$  generators  $\gamma_j \equiv \iota_1(\ell_j)$ satisfying the relation  $\gamma_1 \cdots \gamma_d = 1$ .

In view of (b), to get a full description of the fundamental group  $\pi_1(Y)$  we have to complete its presentation by adding the relations between the generators  $\gamma_i$  which generate the kernel of the homomorphism  $\iota_1$ . This is precisely what the Kempen-Zariski theorem does, see e.g. [\[80,](#page-153-1) [100\]](#page-154-0) for details. E.g. if  $\mathscr W$  is a smooth planar curve of degree d,  $\pi_1(Y)$  =  $\mathbb{Z}/d\mathbb{Z}$ . The Kempen-Zariski relations take a particular simple form when the planar curve  $W \subset \mathbb{P}^2$  is the complexification of a real arrangement of lines. By (b) we have one generator  $\gamma_i$  per line (satisfying  $\gamma_1 \cdots \gamma_d = 1$ ); The Randell theorem [\[153\]](#page-157-3) states that we have one additional relation per intersection point between these lines. At a point where the  $\ell$  lines  $\gamma_{j_s}$   $(s = 1, \ldots, \ell)$  meet (ordered in cyclic order) we have the 'rotation' relations

$$
\gamma_{i_1}\gamma_{i_2}\cdots\gamma_{i_{s-1}}\gamma_{i_s} = \gamma_{i_2}\gamma_{i_3}\cdots\gamma_{i_s}\gamma_{i_1} = \gamma_{i_3}\gamma_{i_4}\cdots\gamma_{i_1}\gamma_{i_2} = \cdots = \gamma_{i_s}\gamma_{i_1}\cdots\gamma_{i_{s-2}}\gamma_{i_{s-1}} \qquad (9.16)
$$

In particular if we have only pairwise intersections the fundamental group is commutative. This is a particular case of the more general result

**Proposition 4.** If  $X \subset \mathbb{P}^2$  is a curve with only nodal singularities  $\pi_1(\mathbb{P}^2 \setminus X)$  is Abelian.<sup>[85](#page-128-0)</sup>

### <span id="page-128-1"></span>9.2.2 Rational covers

We have already noticed that for many purposes we are interested in the monodromy group only up to commensurability, i.e. up to finite groups. For  $r \geq 3$  there is a cheap way of getting new good special divisors  $\mathscr S$  out of old ones such that their complements have commensurable fundamental groups. For easy of exposition, we assume that  $\mathscr P$  is the ordinary projective space  $\mathbb{P}^{r-1}$ . Suppose we have a rational map  $\psi: \mathbb{P}^{r-1} \dashrightarrow \mathbb{P}^{r-1}$  of degree d and a good special divisor  $\mathscr S$  with the property that the hypersurface  $\mathscr S' \equiv \overline{\psi^{-1}(\mathscr S)}$  contains all the points of indeterminacy of  $\psi$  as well as the branching locus of  $\psi$ . Then the restriction of the map between the complements

$$
\psi \mid : \mathbb{P}^{r-1} \setminus \mathscr{S}' \to \mathbb{P}^{r-1} \setminus \mathscr{S} \tag{9.17}
$$

is an unbranched cover with a deck group  $F \subset \mathfrak{S}_d$  (hence finite). One has the exact sequence of groups

$$
1 \to F \to \pi_1(\mathbb{P}^{r-1} \setminus \mathscr{S}) \to \pi_1(\mathbb{P}^{r-1} \setminus \mathscr{S}') \to 1
$$
\n(9.18)

so that the fundamental groups of the two complements differ by a finite group. When  $d = 1$ , i.e.  $\psi$  is a Cremona transformation, the complements are isomorphic and in particular have the same fundamental groups (evident from the above exact sequence since  $F = 1$  in this case). The special divisor  $\mathscr{S}, \mathscr{S}'$  are different but essentially equivalent from the viewpoint of the monodromy analysis. For examples, see below.

More generally we may exploit the theory of Cremona transformations for weighted pro-jective space. See [\[160\]](#page-157-4) for examples when  $r = 3$  with applications to the computation of fundamental groups of complements of curves in weighted projective planes.

## 9.3 First "economical" examples

We focus on the basic case  $r = 3$ , i.e.  $\mathscr P$  of complex dimension 2, and assume the special geometry to be irreducible, rigid (2 is not a Coulomb dimension), and non-isotrivial. Under these conditions in rank-2 the complement  $\mathscr{P} \setminus \mathscr{S}$  was the sphere less 3 points; we look for complements  $\mathscr{P} \setminus \mathscr{S}$  which are "the higher dimensional analogues of the thrice-punctured sphere" in the sense that satisfy the "same" rigidity and non-triviality conditions. The curve

<span id="page-128-0"></span><sup>85</sup> For line arrangements the converse is also true.

 $\mathscr S$  should posses rather restrictive properties, in particular it should have no equisingular  $deformation<sup>86</sup>$  $deformation<sup>86</sup>$  $deformation<sup>86</sup>$  since the complements of equisingular curves are diffeomorphic [\[80\]](#page-153-1) so have the same fundamental group. To give the flavor of a planar curve satisfying all the relevant constraints, we present a simple example which may actually arise from a fully-fledged rank-3 special geometry. Later we shall discuss a deeper interpretation of the same example.

To keep the example as simple as possible, we consider the special case where  $\{\Delta_1, \Delta_2, \Delta_3\}$ are such:  $(i)$  1 and 2 are not dimensions,  $(ii)$  the geometry is not automatically isotrivial  $(\varkappa \in \{1,2\})$ , and  $(iii) \mathscr{P} \simeq \mathbb{P}^2$ . The typical example is  $\{\Delta_1, \Delta_2, \Delta_3\} = \{4, 6, 12\}$  where the isomorphism  $\mathscr{P} \simeq \mathbb{P}^2$  is given in homogeneous coordinates  $(x_1 : x_2 : x_3) = (u_1^6 : u_2^4 : u_3^2)$ . The reduced  $\mathscr S$  has the form

$$
\mathcal{S} = (x_1 \, x_2 \, x_3 \, h(x_1, x_2, x_3)) \tag{9.19}
$$

for some square-free homogeneous polynomial  $h \equiv h(x_1, x_2, x_3)$  of degree d which is not divisible by any  $x_i$ .  $\mathscr S$  consists of three lines in general position,  $\{x_a = 0\}$ , plus the planar curve  $h = 0$  which is reduced but typically reducible. h should satisfy a number of conditions. First,  $\mathscr S$  must have no equisingular deformations, that is, all modifications of the polynomial h which preserve the Milnor numbers of the singularities of  $\mathscr S$  may be undone by rescalings of the variables  $x_i$  ('rigidity'). If the curve h is smooth in general position the rigidity condition is satisfied only when  $d = 1$ . But then the curve  $\mathscr S$  has only simple nodes,  $\pi_1(\mathscr P \setminus \mathscr S)$  is Abelian, so the period map is constant, and the geometry (if it exists) is isotrivial. Next the line  $h = 0$  may be in a special position, i.e. it may pass through the intersection point of two coordinate lines, say through  $(x_2 = 0) \cap (x_3 = 0)$ , so  $h = \alpha x_2 + \beta x_3$ . But in this case

$$
\pi_1(\mathscr{C}\setminus(x_1x_2x_3(\alpha x_2+\beta x_3))) = \pi_1(\mathbb{C}\setminus(x_1)) \times \pi_1(\mathbb{C}^2\setminus(x_2x_3(\alpha x_2+\beta x_3)))
$$
\n(9.20)

and the special geometry decomposes in a rank-1 special geometry with  $\Delta = 4$  and a rank-2 geometry with  $\{\Delta_1, \Delta_2\} = \{6, 12\}$  (again isotrivial). We conclude that the minimal degree of h is 2, in which case the reduced curve  $h = 0$  is either two distinct lines or a smooth conic. We consider the two possibilities in turn.

 $h = 0$  are two distinct lines. Neither line can be in general position because of rigidity; then either both lines pass through a crossing point  $p_{ij} \equiv (x_i = 0) \cap (x_j = 0)$  of two coordinate lines, or they meet on a coordinate line (say on  $x_3 = 0$ ). They cannot pass through the same crossing point  $p_{ij}$  because this will lead to a decomposable geometry. Modulo permutation of the coordinates, we remain with the two possibilities

$$
\begin{cases}\n\alpha x_1 + \beta x_3 = 0 \\
\gamma x_2 + \delta x_3 = 0\n\end{cases} \n\text{ or } \n\begin{cases}\n\alpha x_1 + \beta x_2 + \gamma x_3 = 0 \\
\alpha x_1 + \beta x_2 + \gamma' x_3 = 0\n\end{cases} (9.21)
$$

<span id="page-129-0"></span> $86$  A deformation of a curve is *equisingular* if it preserves the Milnor number of each of its singularity, see [\[80\]](#page-153-1).

In the second case the parameter  $\gamma/\gamma'$  cannot be absorbed by a rescaling, so this configuration is excluded. We remain with the first possibility. We absorb the parameters by a suitable rescaling of the coordinates, and focus on the affine patch  $x_3 \neq 0$  of  $\mathbb{P}^2$ ;  $x_3 = 0$  is then the line at infinity  $L_{\infty}$ , and we write  $\mathscr{S} = X \cup L_{\infty}$ . Now  $\mathbb{P}^2 \setminus \mathscr{S} = \mathbb{C}^2 \setminus X$  where  $X \subset \mathbb{C}^2$  is the complexification of the arrangement in  $\mathbb{R}^2$  of the 4 real lines drawn solid in the figure

<span id="page-130-0"></span>

Adding the dashed line  $L_5 \equiv \{x_1 = x_2\}$  to X will not spoil rigidity nor any other required property. We claim that we should add  $L_5$  in order to satisfy the "natural" conditions of  $\pi_1(\mathscr{P} \setminus \mathscr{S})$  listed in §.[9.1.2.](#page-126-1) Indeed in eq.[\(9.33\)](#page-133-0) we will show that the fundamental group of the complement in  $\mathbb{C}^2$  of the 4 solid lines is a non-trivial product. Thus

Example 9.5. Our first "economic" higher dimensional analogue of the sphere less 3 points is  $\mathbb{P}^2$  less the six lines

$$
\{x_1 = 0\}, \ \{x_2 = 0\}, \ \{x_3 = 0\}, \ \{x_1 = x_3\}, \ \{x_2 = x_3\}, \ \{x_1 = x_2\}.\tag{9.23}
$$

i.e. the complement in  $\mathbb{C}^2$  of the five lines in the figure [\(9.22\)](#page-130-0). We check the validity of the "hereditary principle". Each special line contains precisely 3 special points, namely the two intersections with other lines visible in figure [\(9.22\)](#page-130-0) plus the intersection with the line at infinity which also has 3 special points associated to each set of parallel lines in [\(9.22\)](#page-130-0). We shall consider the reduction of the natural Picard-Fuchs equation with singularity locus  $(9.22)$  to one discriminant line (in the sense of §.[4.3\)](#page-53-0), in §.  $(9.3.1)$  below.

h is a smooth conic. Let us consider the case where  $h = 0$  is a conic. Remark [23](#page-124-2) gives us a first good configuration: h is the (unique) smooth conic in  $\mathbb{P}^2$  which is tangent to all 3 coordinate lines  $\{x_a = 0\}$  i.e.

<span id="page-130-1"></span>
$$
h: \quad x_1^2 + x_2^2 + x_3^2 - 2(x_1x_2 + x_2x_3 + x_3x_1) = 0. \tag{9.24}
$$

Each one of the 4 components of  $\mathscr S$  is a copy of  $\mathbb P^1$  and the (reduced) restriction of  $\mathscr S$  to it consists of 3 distinct points in perfect accordance with the hereditary principle.

Another rigid configuration is a conic which goes through all 3 intersection points  $p_{ij}$  i.e.

$$
h: \quad x_1 x_2 + x_2 x_3 + x_3 x_1 = 0 \tag{9.25}
$$

The three lines  $x_i + x_j = 0$   $(i < j)$  are tangent to  $h = 0$  at the crossing point  $p_{ij}$ ; adding some of them would not spoil rigidity. If we add exactly two of them, say the tangents to  $h = 0$  in  $p_{12}$  and  $p_{13}$ , all components have 3 special points, but for the line  $x_1 = 0$  which has only 2 such points and then must be a regular R-enhanced discriminant (cf. **Remark** [24](#page-125-1)). If we draw all 3 tangents we get a special divisor that, while rigid, is not hereditary: some components have 4 special points. This configuration should be disregarded. Thus to construct a new "economic" example we add just two tangents and draw them dashed meaning that the monodromy along them is allowed to be trivial (cf. the dashed line in [\(9.22\)](#page-130-0)). If we set the local monodromy along 0,1, or dashed lines to 1 we get a configuration with (respectively) 1, 2 or 3 lines  $\{x_a = 0\}$  which are regular R-enhanced. Note that this is consistent with the fact that only coordinate planes can be regular special divisors.

Finally we have the less symmetric situations where the conic is tangent (say) to the line  $x_3 = 0$  in the crossing point  $(0:1:0)$  and also passes through another crossing point, say  $(0:0:1)$ . While this curve is rigid, the conic component has only two special points: since it is not a coordinate line it cannot be a regular component and so, having only 2 special points, violates hereditary. This configuration should also be disregarded.

Example 9.6. Our two next "economic" examples of higher dimensional analogues of the 3 punctured sphere  $\mathbb{P}^1 \setminus \{0, 1, \infty\}$  are: (1) the complement in  $\mathbb{C}^2$  of a conic with three tangent lines and (2) the complement of three lines plus the conic passing through their crossing points and two out of the three lines tangent to the conic which pass through a crossing point of the original lines.

**Higher degree**  $d \geq 3$ . We can increase the degree d of h above 2 and rigidity will imply stronger and stronger non-genericity conditions and wilder singularities making the configuration of curves more and more "unlikely" and "ugly". Since our purpose is merely to present nice economical examples, we shall not pursue them here. Many other examples (possibly all) can be produced using the cover construction outlined in §. [9.2.2](#page-128-1) and so are essentially equivalent to the one described before. It is conceivable that all consistent special divisors arise in this way from the previous "economic" divisors (see below for some preliminary evidence).

**Summary of**  $r = 3$  examples. In conclusion, for our illustrative purposes we focus on the three "cheap" examples of higher dimensional analogues of the thrice-punctured sphere  $\mathbb{P}^1 \setminus \{0, 1, \infty\}$  given by the complement  $\mathbb{P}^2 \setminus \mathscr{S}_{(a)}$   $(a = 1, 2, 3)$  where

(1)  $\mathscr{S}_{(1)}$ :  $x_1 x_2 x_3 (x_3 - x_1) (x_2 - x_3) (x_1 - x_2) = 0$ 

- (2)  $\mathcal{S}_{(2)}$ :  $x_1 x_2 x_3 (x_1 + x_2) (x_2 + x_3) (x_1 x_2 + x_2 x_3 + x_1 x_3) = 0$
- (3)  $\mathscr{S}_{(3)}$ :  $x_1 x_2 x_3 (x_1^2 + x_2^2 + x_3^2 2x_1 x_2 2x_2 x_3 2x_3 x_1) = 0.$

**Remark 25.** The coordinate lines  $x_i = 0$  may correspond to R-enhanced loci which are or not part of the discriminant. On the contrary, all components of  $h = 0$  should be part of the discriminant. Hence a *necessary condition* for the coordinate axis  $\{u_i = 0\} \cap \{u_j = 0\}$ not to lay in the discriminant is that h does not cross the point  $p_{ij} \in \mathbb{P}^2$ . Therefore, when the three axes are all regular (and the degree of  $h \leq 2$ ), the only possibilities are (3) or (2) with the two "dashed" lines  $x_1 + x_2 = 0$ ,  $x_2 + x_3 = 0$  deleted. In rank-2 there are only 2 non-isotrivial geometries with all axes regular  $(\#20 \text{ and } \#21)$  and they both correspond to Argyres-Douglas SCFTs. Non-isotrivial geometries with all axes regular and an irreducible discriminant are quite rare.

Rational covers. In the case of the thrice-punctured sphere we had to study the representations of a single group  $\pi_1(\mathbb{P}^1 \setminus \{0, 1, \infty\}) \simeq F_2$  (the free group in two generators). In the present situation it may seem that we have to study three different groups already at the level of our "economic" examples. However it is not so [\[154\]](#page-157-5)

**Fact 35.** (1) The complements  $\mathbb{P}^2 \setminus \mathscr{S}_{(1)}$  and  $\mathbb{P}^2 \setminus \mathscr{S}_{(2)}$  are biholomorphic so, in particular, have isomorphic fundamental groups  $\pi_1(\mathbb{P}^2 \setminus \mathscr{S}_{(1)}) \simeq \pi_1(\mathbb{P}^2 \setminus \mathscr{S}_{(2)})$ . (2) We have an unbranched double cover  $\mathbb{P}^2 \setminus \mathscr{S}_{(1)} \to \mathbb{P}^2 \setminus \mathscr{S}_{(3)}$  with deck group  $\mathbb{Z}_2$  generated by  $x_1 \leftrightarrow x_2$ , so that

$$
1 \to \mathbb{Z}_2 \to \pi_1(\mathbb{P}^2 \setminus \mathcal{S}_{(3)}) \to \pi_1(\mathbb{P}^2 \setminus \mathcal{S}_{(1)}) \to 1 \tag{9.26}
$$

so the fundamental groups agree modulo a finite group.

Proof. (1) The Cremona transformation

$$
c: \mathbb{P}^2 \dashrightarrow \mathbb{P}^2, \qquad (x_1, x_2, x_3) \mapsto (y_1, y_2, y_3) \equiv (x_1 x_2 + x_2 x_3, -x_1 x_3, x_1 x_2) \tag{9.27}
$$

(projectively equivalent to a standard quadratic transform [\[155\]](#page-157-6)) has the property that the *reduced* divisor  $(c^*\mathscr{S}_{(1)})_{\text{red}}$  is equal to  $\mathscr{S}_{(2)}$ . Indeed

$$
c^*(y_1y_2y_3(y_3-y_1)(y_2-y_3)(y_1-y_2)) =
$$
  
= 
$$
-x_1^3x_2^3x_3^2(x_1+x_3)(x_2+x_3)(x_1x_2+x_1x_3+x_2x_3),
$$
 (9.28)

while the points of indeterminacy of  $c((1:0:0), (0:1:0), (0:0:1))$  and of  $c^{-1}((1:0:0),$  $(0:1:0), (1:0:1)$  belong to the excised curves so that c is an isomorphism when restricted to the complements. (2) Same story, except that the rational map

$$
r: (x_1, x_2, x_3) \mapsto (y_1, y_2, y_3) \equiv (x_1 x_2, (x_3 - x_1)(x_3 - x_2), x_3^2)
$$

$$
r^* \left(y_1 y_2 y_3 \left(\sum_a y_a^2 - 2 \sum_{a < b} y_a y_b\right)\right) = x_1 x_2 x_3^4 (x_1 - x_2)^2 (x_1 - x_3)(x_2 - x_3)
$$
(9.29)

has now degree 2. The Jacobian of the map is  $2(x_1 - x_2)x_3^2$ , so the map is an unbranched cover when restricted to the complement.  $\Box$ 

**Fundamental groups.** It is convenient to think of the "good" locus  $\mathcal{\dot{P}}$  as the complement of the arrangement of lines  $\mathscr{S}_{(1)}$  which allows us to write a simple presentation of its fundamental group using the Randell theorem [\[153\]](#page-157-3). We number the lines in [\(9.22\)](#page-130-0) (i.e. the component of  $\mathscr{S}_{(1)}$  distinct of the line at infinity  $L_{\infty} = \{x_3 = 0\}$  as

$$
L_1 = \{x_1 = 0\}, \ L_2 = \{x_2 = x_3\}, \ L_3 = \{x_1 = x_2\}, \ L_4 = \{x_1 = x_3\}, \ L_5 = \{x_2 = 0\} \tag{9.30}
$$

and write  $\mu_a$  for the monodromy of a loop going once around  $L_a$  in the positive direction. The Randell theorem then gives  $[100, 153]^{87}$  $[100, 153]^{87}$  $[100, 153]^{87}$  $[100, 153]^{87}$  $[100, 153]^{87}$  $[100, 153]^{87}$ 

<span id="page-133-3"></span>
$$
\pi_1(\mathbb{P}^2 \setminus \mathscr{S}_{(1)}) = \pi_1(\mathbb{C} \setminus \bigcup_{a=1}^5 L_a) = \left\langle \mu_1, \mu_2, \mu_3, \mu_4, \mu_5 \middle| \begin{array}{l} \mu_1 \mu_2 = \mu_2 \mu_1, & \mu_4 \mu_5 = \mu_5 \mu_4, \\ \mu_5 \mu_3 \mu_1 = \mu_3 \mu_1 \mu_5 = \mu_1 \mu_5 \mu_3, \\ \mu_4 \mu_3 \mu_2 = \mu_3 \mu_2 \mu_4 = \mu_2 \mu_4 \mu_3 \end{array} \right\} (9.31)
$$

and we have

$$
\mu_{\infty} = (\mu_1 \mu_2 \mu_3 \mu_4 \mu_5)^{-1}.
$$
\n(9.32)

When the dashed line  $L_3$  has trivial local monodromy,  $\mu_3 = 1$ , (the "*most* economical" example) all relations become commutations and we get  $88$ 

<span id="page-133-0"></span>
$$
\pi_1(\mathbb{P}^2 \setminus \mathscr{S}_{\text{m.e.}}) \simeq F_2 \times F_2 \tag{9.33}
$$

which is a special instance of **Theorem 4.11** of [\[156\]](#page-157-7) (in turn a special case of [\[157\]](#page-157-8)).

**Remark 26.** As an abstract group  $\pi_1(\mathbb{P}^2 \setminus \mathcal{S}_{(3)})$  is the Artin braid group of the triangle graph [\[158,](#page-157-9) [159\]](#page-157-10)

$$
\begin{array}{c}\n \stackrel{4}{\overbrace{\qquad \qquad }} \quad \stackrel{4}{\overbrace{\qquad \qquad }} \quad \stackrel{4}{\overbrace{\qquad \qquad }} \quad \ \ (9.34)\n \end{array}
$$

For a van Kampen-like presentation of  $\pi_1(\mathbb{P}^2 \setminus \mathcal{S}_{(3)})$ , see instead [\[160\]](#page-157-4).

<span id="page-133-1"></span> $87$  Compare the line arrangement in figure [\(9.22\)](#page-130-0) with the one in the figure 13.8 of [\[100\]](#page-154-0).

<span id="page-133-2"></span><sup>&</sup>lt;sup>88</sup> As before,  $F_2$  is the free group in two generators (isomorphic to  $\Gamma(2)$  as abstract groups).

#### <span id="page-134-0"></span>9.3.1 Comparison with Appell's hypergeometric PDEs

Above we have followed our physical intuition to get "economic" examples of complements of planar curves which may be seen, at least informally, as higher dimensional counterparts of the sphere with three punctures. It turns out that the 3 varieties we got realize the desired analogy in a much deeper sense than expected a priori. Indeed, as stressed by Riemann in 1851, the most important property of the thrice-punctured sphere is that the degree-2 representations of its fundamental group are the monodromies of the order-2 hypergeometric ODE, and these representations are rigid. Appell [\[161\]](#page-157-11) extended Riemann ideas to higher dimensions introducing four hypergeometric functions  $F_i(x, y)$   $(i = 1, \ldots, 4)$  depending on two variables  $x, y$  which satisfy certain systems of linear PDEs with regular singularities (a Picard-Fuchs system) which are, in a sense, the 2-variable generalization of Gauss' hypegeometric ODE. One studies the monodromy representation of this Picard-Fuchs systems and finds that they are *rigid* (under some natural condition)  $|100, 154, 162-164|$  $|100, 154, 162-164|$  $|100, 154, 162-164|$ . The monodromy of the Appell function is precisely a (rigid) representation of the fundamental group  $\pi_1(\mathscr{P} \setminus \mathscr{S}_{(1)})$  of the complement of the arrangement of lines in figure [\(9.22\)](#page-130-0) which then plays the same role in the theory of rigid monodromy of Picard-Fuchs PDEs on  $\mathbb{P}^2$  than the thrice-punctured sphere plays classically in the theory of rigid monodromies of Picard-Fuchs ODEs on  $\mathbb{P}^1!$ 

Important examples of rigid monodromy representations of the group [\(9.31\)](#page-133-3) are explic-itly know [\[154,](#page-157-5) [162–](#page-157-12)[164\]](#page-158-0): they are the monodromies of the four Appell's functions  $F_1(x, y)$ ,  $F_2(x, y)$ ,  $F_3(x, y)$ ,  $F_4(x, y)$  which satisfy pairs of second order PDEs [\[161,](#page-157-11) [165\]](#page-158-1). In facts, the divisors  $\mathcal{S}_{(1)}$ ,  $\mathcal{S}_{(2)}$  and  $\mathcal{S}_{(3)}$  are the loci of the regular singularities of the Picard-Fuchs PDEs satisfied, respectively, by  $F_1$ ,  $F_3$ , and  $F_4$  (the divisor for  $F_2$  is obtained from the one for  $F_1$ by a linear redefinition of the coordinates [\[154\]](#page-157-5)). Their monodromy representations have the form

$$
\varrho: \pi_1(\mathbb{P}^2 \setminus \mathcal{S}_{(1)}) \to GL(n, \mathbb{C}) \quad \text{where} \quad n = \begin{cases} 3 & \text{for } F_1 \\ 4 & \text{for } F_2, F_3, F_4 \end{cases} \tag{9.35}
$$

The theory of the Appell PDEs may be used to illustrate the "stratification" at the level of the Picard-Fuchs PDEs (cf.  $\S$ , [4.3\)](#page-53-0). The restrictions of the Appell PDE system for  $F_1$  to each one of the irreducible components of the special divisor  $\mathscr{S}_{(1)}$  is studied in **Example** 12.2 of [\[100\]](#page-154-0). We quote the result: the restriction of the PDE to each line in the special locus yields back the Gauss hypergeometric ODE.

## 9.4 Appell's monodromies?

Appell's monodromy representation cannot be directly identified with the monodromy of a family of Abelian varieties of dimension 3 since the latter has dimension (degree) 6, while in the Appell case we have 3-dimensional representations  $V_3$  and 4-dimensional ones  $V_4$ . Indeed, the Appell functions are the higher dimensional analogue of the Gauss hypergeometric functions  ${}_2F_1(z)$  whereas already for  $r = 2$  the SW periods were expressed in terms of the generalized hypergeometric functions  ${}_4F_3(z)$  which solve Thomae's order-4 linear ODE. These equations have the same singular set, but different order. For our applications we need the higher dimensional analogue of the Thomae hypergeometric – with a dimension-6 monodromy representation – not Appell's analog of the Gauss equation (which is good enough as long as understanding the special divisor  $\mathscr S$  goes).

However we may still think of a "functorial" construction where we get a dimension 6 monodromy representation  $V_6$  of  $\pi_1(\mathbb{P}^2 \setminus \mathscr{S}_{(1)})$  starting from Apell's monodromies  $V_3$  and  $V_4$ of dimension (respectively) 3 and 4. Applying the standard operations of linear algebra to  $V_3$ ,  $V_4$  one constructs "functorially" various 6-dimensional representations such as  $V_3 \oplus V_3$  or  $V_3 \odot V_3$  or  $V_4 \wedge V_4$ . More generally, we may consider non-trivial factorizations of the form

$$
\mu \colon \pi_1(\mathbb{P}^2 \setminus \mathscr{S}_{(1)}) \to G(\mathbb{Z}) \xrightarrow{\iota} Sp(6, \mathbb{Z})
$$
\n(9.36)

where we assume  $G(\mathbb{Z})$  to be the smallest group with this property. The Mumford-Tate group G of the Abelian family  $\mathscr{A} \to \mathscr{\hat{P}}$  [\[38](#page-150-1)[–40\]](#page-151-0) is the connected component of the Q-algebraic group which is the Zariski Q-closure  $\overline{G(\mathbb{Z})}^{\mathbb{Q}}$  of the discrete subgroup  $G(\mathbb{Z}) \subset Sp(6, \mathbb{Q})$ . The real Lie group  $G(\mathbb{R})$  of real-valued points of G is a Hermitian group<sup>[89](#page-135-0)</sup>.  $\iota$  extends to an inclusion of real Lie groups  $G(\mathbb{R}) \stackrel{\iota}{\to} Sp(6, \mathbb{R})$  given by a representation which is symplectic in the "motivic" sense [\[88,](#page-153-2) [101\]](#page-154-2). The allowed pairs of algebraic groups and symplectic representations (for any rank r) may be read for instance in the table on page 10 of [\[88\]](#page-153-2). If  $G(\mathbb{R}) \subset Sp(6, \mathbb{R})$ is a non-trivial subgroup, it can be only one of the semi-simple real Hermitian maximal subgroups

$$
SL(2,\mathbb{R}), \quad SL(2,\mathbb{R}) \times SL(2,\mathbb{R}), \quad SL(2,\mathbb{R}) \times Sp(4,\mathbb{R}), \quad SU(2,1) \tag{9.37}
$$

or a subgroup thereof. The first two have no real dimension 6 representation in the table [\[88\]](#page-153-2). For the third one the only allowed representation is  $2 \oplus 4$  which leads to a reducible representation. As we learned from the reducible examples in rank-2, this may still produce a non-trivial geometry (for instance a quasi-isotrivial geometry with a generic fiber which is isogenic to the product of a fixed elliptic curve with an Abelian surfaces whose 4 periods satisfy an Appell's PDE) but here we are mainly interested in  $\mu$ -irreducible geometries and leave apart this possibility for future work.

We remain with  $G(\mathbb{R}) = SU(2, 1)$  and a real representation  $V_6$  such that  $V_6 \otimes \mathbb{C} \simeq 3 \oplus \overline{3}$ . The embedding of  $SU(2,1)$  into  $Sp(6,\mathbb{R})$  is as follows: identify  $Sp(6,\mathbb{R})$  with the group of real  $6 \times 6$  matrices S satisfying

$$
S^{t} \tilde{\Omega} S = \tilde{\Omega} \text{ where } \tilde{\Omega} = \begin{pmatrix} 0 & \eta \\ -\eta & 0 \end{pmatrix} \text{ and } \eta = \text{diag}(+1, +1, -1) \tag{9.38}
$$

<span id="page-135-0"></span><sup>&</sup>lt;sup>89</sup> A Hermitian group is a real Lie group G such that the symmetric space  $G/K$  ( $K\subset G$  a maximal compact subgroup) is Kähler. All Hermitian groups are of Hodge type (the converse is false).

and  $SU(2, 1)$  with the group of  $3 \times 3$  complex matrices U such that  $U^{\dagger} \eta U = \eta$ . Then

<span id="page-136-0"></span>
$$
SU(2,1) \hookrightarrow Sp(6,\mathbb{R}), \quad U \mapsto \begin{pmatrix} \text{Re } U & \text{Im } U \\ -\text{Im } U & \text{Re } U \end{pmatrix}.
$$
 (9.39)

We have the constraint that the monodromy group should contain a non-trivial unipotent element  $S_0$  i.e. such that  $S_0 - 1$  is nilpotent of rank 1, in facts

$$
S_0 = \mathbf{1} + v \otimes v^t \, \tilde{\Omega} \qquad v \neq 0. \tag{9.40}
$$

But  $S_0$  is not of the form in the right side of [\(9.39\)](#page-136-0), so  $S_0 \notin SU(2,1)$ . We conclude

**Fact 36.** A non-isotrivial,  $\mu$ -irreducible rank-3 special geometry has the standard Mumford-Tate group  $Sp_6$ .

Thus we cannot construct a rank-3  $\mu$ -irreducible special geometry by a simply-minded use of Appell's transcendents. To get a candidate monodromy for a putative  $\mu$ -irreducible (and  $\mu$ -rigid) special geometry based on our "economic" analogue of the 3-punctured sphere we should look for "new" transcendents which generalize to multiple variables the Thomae ones. In the next subsection we present evidence that nice transcendents of the appropriate kind exist.

## <span id="page-136-1"></span>9.5 An "economic"  $r = 4$  example

We look for would-be for rank-4 analogues of the 3-punctured sphere. For simplicity we take  $\mathscr{P}$  to be  $\mathbb{P}^3$ : the physically relevant case of a weighted projective space  $\mathbb{P}(d_1,\ldots,d_4)$  can be reduced to this one by a finite cover as before.

To construct a natural analogue in  $\mathbb{P}^3$  of the planar configuration of a conic with 3 tangent lines we start from the Cayley cubic  $X_3 \subset \mathbb{P}^3$  which is the unique (modulo  $PGL(4,\mathbb{C})$ projective equivalence) singular normal cubic hypersurface with 4 singularities which are of type  $A_1$  (simple nodes) [\[155,](#page-157-6) [166\]](#page-158-2). Choosing the coordinates so that the 4 nodes are at

$$
(1:0:0:0), (0:1:0:0), (0:0:1:0), (0:0:0:0:1), (9.41)
$$

the equation is

$$
x_1x_2x_3 + x_2x_3x_4 + x_3x_4x_1 + x_4x_1x_2 = 0.
$$
\n(9.42)

Since the Cayley cubic is singular, it has only 9 lines and 11 tritangent planes instead of 27 lines and 45 tritangents as a generic cubic. The lines are of two types [\[166\]](#page-158-2):



The 11 tritangent planes (i.e. planes tangent in three lines, counted with multiplicity) are:

$\#$ equation	it contains	
4 $x_i = 0, \quad i = 1, \ldots, 4$ 3 distinct <b>A</b> lines		(9.44)
	6 $x_i + x_j = 0$ , $1 \leq i < j \leq 4$ 1 <b>A</b> line (multiplicity 2) & 1 <b>B</b> line	
1 $x_1 + x_2 + x_3 + x_4 = 0$ 3 lines of type <b>B</b>		

Proposal. As an example of rank 4 analogue to the 3-puncture sphere we propose the complement in  $\mathbb{P}^3$  of the divisor  $\mathscr{S}_8$  whose 8 irreducible components are

- the Cayley cubic  $X_3$ ;
- the 6 tritangent planes containing double lines through two nodes;
- the tritangent plane containing the  $\beta$  lines of type  $\boldsymbol{B}$ ,

that is

$$
(x_1 + x_2)(x_1 + x_3)(x_1 + x_4)(x_2 + x_3)(x_2 + x_4)(x_3 + x_4) \times
$$
  
 
$$
\times (x_1 + x_2 + x_3 + x_4)(x_1x_2x_3 + x_2x_3x_4 + x_3x_4x_1 + x_4x_1x_2) = 0 \quad (9.45)
$$

Clearly this divisor is rigid. Let us check hereditarity. The restriction to a tritangent containing two lines, say  $x_1 + x_2 = 0$ , followed by the change of coordinates  $(x_1, x_3, x_4)$  $(y_3, 2y_1 - y_3, -2y_2 + y_3)$  gives

$$
-128 y_1 y_2 y_3^2 (y_1 - y_3)(y_2 - y_3)(y_1 - y_2)^2 = 0
$$
\n(9.46)

which, after reduction, is exactly the rank-3 line configuration in figure  $(9.22)$  Clearly adding further planes would spoil this property. The restriction to the last tritangent plane yields, after reduction, a plane with 3 lines in general position, which is consistent provided that the restriction of the rank-4 geometry to this plane is a product of rank-1 geometries (actually just isotrivial after taking into account the finite cover).

On the other hand the birational interpretation (as a singular del Pezzo surface) of the Cayley cubic sees it as the image of the linear system of cubics passing through the six points which are the pairwise intersections of 4 lines in general position in  $\mathbb{P}^2$  [\[166\]](#page-158-2). This yields the rational parametrization of  $X_3 \subset \mathbb{P}^3$ 

$$
(x_1:x_2:x_3:x_4) ==(y_2^2y_3+y_3^2y_2+y_1y_2y_3:y_3^2y_1+y_1^2y_3+y_1y_2y_3:y_1^2y_2+y_2^2y_1+y_1y_2y_3:-y_1y_2y_3)
$$
(9.47)

so that  $X_4$  is an admissible Coulomb branch in the sense of **Definition** ??.

#### 9.5.1 Relation to recent work on Fuchsian systems

The divisor  $\mathscr{S}_8$  has recently appeared in the math literature as the singular locus of a Fuchsian system of rank 8 in  $\mathbb{P}^3$ , see ref. [\[167\]](#page-158-3). In other words that paper gives a monodromy representation

$$
\mu \colon \pi_1(\mathbb{P}^3 \setminus \mathscr{S}_8) \to GL(8, \mathbb{C}) \tag{9.48}
$$

which when restricted to a suitable irreducible component of  $\mathscr{S}_8$  yields a *rank 6* representation

$$
\phi \colon \pi_1(\mathbb{P}^2 \setminus \mathcal{S}_{(1)}) \to GL(6, \mathbb{C}).\tag{9.49}
$$

This seems exactly the gadged we need: a dimension 6 representation of the rank-3 group  $\pi_1(\mathbb{P}^2 \setminus \mathscr{S}_{(1)})$  which arises from the restriction of a dimension 8 representation for the rank-4 group  $\pi_1(\mathbb{P}^3 \setminus \mathscr{S}_8)$  as expected from "stratification" of a rank-4 special geometry. Indeed this is almost right but, just as in rank-2, the math literature mostly focuses on representations with semisimple local monodromies, and so we cannot simply borrow their results and get a solution to the inverse problem of special geometry. This remark applies a *fortiori* in higher rank. However the existence of natural Fuchsian equations and monodromies of roughly the correct type, with the proper "stratification" properties yields us confidence that we are on the right track.

### 9.6 Lauricella singular divisors and all that

From the above examples we infer that a plausible class of special divisors  $\mathscr{S} \subset \mathbb{P}^{r-1}$  with all the required properties (the "higher dimensional analogues of the 3-punctured sphere") are given by the loci of (regular) singularities of flat Gauss-Manin connections which generalizes the hypergeometric equation. Lauricella [\[168\]](#page-158-4) gave a list of fours hypergeometric functions  $F_A(x_i)$ ,  $F_B(x_i)$ ,  $F_C(x_i)$ ,  $F_D(x_i)$  depending on n variables (and suitable parameters) for all  $n \in \mathbb{N}$ . The singular loci of the Fuchsian PDEs satisfied by the Lauricella functions of n variables yield examples of rigid special divisors  $\mathscr S$  with the required properties. The singular loci of the PDEs satisfied by the Lauricella functions are studied e.g. in [\[169,](#page-158-5) [170\]](#page-158-6).

The singular locus of the PDE for  $F_C$  with m variables, when restricted to the affine chart  $\mathbb{C}^m \equiv \mathbb{P}^m \setminus H_{\infty}$  is the zero set of the reducible polynomial [\[169\]](#page-158-5)

$$
\prod_{i=1}^{m} x_i \prod_{\varepsilon_i = \pm 1} (1 + \varepsilon_1 \sqrt{x_1} + \dots + \varepsilon_m \sqrt{x_m}) \tag{9.50}
$$

which for  $m = 2$  yields back the three lines  $x_a = 0$  plus the conic [\(9.24\)](#page-130-1) i.e. the special set  $\mathscr{S}_{(3)}$ .

For the Lauricella function  $F_A$  of m variables one gets the singular locus [\[170\]](#page-158-6) is the zero

set of the polynomial [\[170\]](#page-158-6)

$$
\prod_{i=1}^{m} x_i \prod_{\{i_1,\dots,i_r\} \subset \{1,\dots,m\}} \left(1 - \sum_{p=1}^{r} x_{i_p}\right) \tag{9.51}
$$

so for  $m = 2$  we get  $x_1x_2(1-x_1)(1-x_1-x_2) = 0$  which with the change of variables  $x_2 \leftrightarrow 1 - x_2$  gives  $x_1x_2(1-x_1)(1-x_2)(x_2-x_1) = 0$  i.e. the special set  $\mathscr{S}_{(1)}$  (reinserting the line at infinity). Lauricella's function  $F_B$  gives an equivalent singular locus.

Finally Lauricella function  $F_D$  has a singular locus

$$
\prod_{i=1}^{m} x_k (1 - x_k) \prod_{1 \le i < j \le m} (x_i - x_j) = 0 \tag{9.52}
$$

which for  $m = 2$  gives back  $\mathscr{S}_{(1)}$ .

These Lauricella singular loci (together with the hyperplane at infinity) give the basic examples of admissible special divisors for *smooth* special geometry in rank  $r = m + 1$  when  $\mathscr{P} \simeq \mathbb{P}^{r-1}$ . In addition we have the generalizations of the special divisor in §.[9.5](#page-136-1) based on the Klein cubic for  $m = 3$ , and finite covers thereof. The situation is similar when  $\mathscr P$  is a general weighted projective space, and indeed the difference is a finite group cf. §. [9.2.](#page-126-2)

### 9.7 Conclusion

The examples and the discussion above strongly suggest that the classification (possibly up to finite covers) of all complements  $\mathscr{P} \setminus \mathscr{S}$  with the required properties is feasible for all r, and certainly for  $r = 3$ . When we have a list of putative special divisors, we can state the inverse program in quite explicit terms. The monodromy representation is expected to be rigid, and then in principle one can proceed with the recursion in the rank of the inverse problem which however leads to an enormous "vast program".

## Acknowledgments

I have benefited of discussions with Michele Del Zotto, Mario Martone, Robert Moscrop, and Artan Sheshmani. I thank Chris Peters, Carlos Simpson, Alexandru Dimca, Yoshishige Haraoka, and V.P. Kostov for clarifications on their respective works.

# A Sketch of Lemma [1](#page-17-0)

**Lemma.** The connected component ( $\mathscr{X}^{\text{reg}}_0$  $\mathscr{X}_0^{\rm reg}$  of the smooth locus  $\mathscr{X}_0^{\rm reg} \subset \mathscr{X}$  of the fiber over the origin 0 has the structure

$$
(\mathcal{X}_0^{\text{reg}})^0 = A_0 \times \mathbb{C}^{r-\ell}, \quad \ell \equiv \dim A_0 = \dim \mathcal{R}_{\Delta=1},\tag{A.1}
$$

where  $A_0$  is a polarized Abelian variety, while the (general) smooth fibers have the form  $\mathscr{X}_u = A_0 \times \mathscr{Y}_u$  with  $\mathscr{Y}_u$  a polarized Abelian variety of dimension  $r - \ell$ .

We know from the discussion in §. [2.9](#page-32-0) that  $(\mathscr{X}^{\text{reg}}_0)$  $\binom{reg}{0}$ <sup>0</sup> is a connected, commutative, algebraic group over  $\mathbb C$  of dimension r. Hence by the Chevalley-Barsotti theorem it is a fibration over an Abelian variety  $A_0$  with fiber a product of copies of  $\mathbb C$  and copies of  $\mathbb C^{\times}$ .  $\exp(z \mathcal{E})$  acts by automorphisms of the commutative group ( $\mathscr{X}^{\text{reg}}_0$  $\binom{reg}{0}^0$  for all  $z \in \mathbb{C}$ .

Let L be the (commutative) Lie algebra of the group  $(\mathscr{X}^{\text{reg}}_0)$ <sup>reg</sup><sup>0</sup> isomorphic to  $T_0(\mathscr{X}_0^{\text{reg}})$  $\binom{reg}{0}$ which in turn is isomorphic to

$$
L \simeq (\mathcal{R}_+ / \mathcal{R}_+^2)[1] = L_0 \oplus (\bigoplus_{\Delta_i > 1} L_{\Delta_i - 1})
$$
\n(A.2)

The action of  $\exp(z\mathcal{E})$  on the group induces an action on the Lie algebra which acts on each summand as

<span id="page-140-0"></span>
$$
\exp(z\,\mathcal{E})\Big|_{L_d} = e^{z\,d}.\tag{A.3}
$$

We can reconstruct the group from its Lie algebra modulo a discrete subgroup D

<span id="page-140-1"></span>
$$
(\mathscr{X}_0^{\text{reg}})^0 = \left(\mathbb{C}^{\ell_0} \times \prod_{d>0} \mathbb{C}^{\ell_d}\right) / D, \qquad \ell_d = \dim L_d. \tag{A.4}
$$

The subgroup D should be preserved by the automorphism  $\exp(z \mathcal{E})$  for all z. In view of [\(A.3\)](#page-140-0), a point in D has the form  $p \times \{0\} \times \cdots \times \{0\}$  with  $p \in \mathbb{C}^{\ell_0}$  so that

$$
(\mathcal{X}_0^{\text{reg}})^0 = (\mathbb{C}^{\ell_0}/D) \times \mathbb{C}^{r-\ell_0}.
$$
\n(A.5)

We have to show that the group  $\mathbb{C}^{\ell_0}/D$  is an Abelian variety (i.e. that D is a maximal lattice). For simplicity of notation, we assume that there is just one Coulomb coordinate  $u_1$ of dimension 1, while  $\Delta_i > 1$  for  $i > 1$ . The general case is an analogous.

Let  $u \in \mathscr{C}$  be a generic point. In a neighborhood of u preserved by  $\exp(z\mathcal{E})$  we write  $\Omega = dz^i \wedge du_i$ . The Hamiltonian vector field generated by  $u_i$  is  $\partial/\partial z^i$  which acts on the fiber coordinates by translation  $z^j \to z^j + \alpha \delta^j_i$ . The kernel of the exponential map exp: Lie( $\mathscr{X}_u$ )  $\to \mathscr{X}_u$  is then a lattice in  $\mathbb{C}^r$  which we write as equivalence relations for the coordinates of the fiber  $\mathscr{X}_u$ 

<span id="page-140-2"></span>
$$
z^{i} \sim z^{i} + A^{ij}(u) m_{j} + B^{ij}(u) n_{j}, \qquad m_{j}, n_{j} \in \mathbb{Z},
$$
\n(A.6)

where the matrix  $A(u)$  is invertible while the matrix  $A(u)^{-1}B(u)$  is symmetric with positivedefinite imaginary part. Now

$$
A^{ij}(e^{z\mathcal{E}}u) = e^{z(1-\Delta_i)} A^{ij}(u), \qquad B^{ij}(e^{z\mathcal{E}}u) = e^{z(1-\Delta_i)} B^{ij}(u)
$$
 (A.7)

which shows that as  $z \to -\infty$  (that is, as  $e^{z\mathcal{E}}u \to 0$ ), the fiber coordinates  $w_i$  with  $i > 1$ 

become valued in C, while

$$
z^{1} \sim z^{1} + A^{1j}(u) m_{j} + B^{1j}(u) n_{j}.
$$
\n(A.8)

Comparing with eq.[\(A.4\)](#page-140-1), we get  $D \equiv \{A^{1j}(u)m_j + B^{1,j}n_j\} \subset \mathbb{C}$ .  $D \subset \mathbb{C}$  is discrete subgroup, hence a lattice of rank  $\leq 2$ . If the rank is  $\lt 2$ , we get from  $(A.6)$  that  $\mathscr{X}_u$  was not compact, contrary to the assumption. Returning to the general case, we see that the factor  $A_0 \equiv \mathbb{C}^{\ell_0}/D$  in FACTrr  $(\mathscr{X}_0^{\text{reg}})$  $(0 \text{ or } 0)$ <sup>teg</sup>)<sup>0</sup> is an Abelian variety of dimension equal to the multiplicity of 1 as a Coulomb dimension, while the general fiber has the form  $\mathscr{X}_u = A_0 \times \mathscr{Y}_u$  for some Abelian variety  $\mathscr{Y}_u$ . The fiber is then a product of the fixed Abelian variety  $A_0$  with the u-dependent variety  $\mathscr{Y}_u$ . Since the Lie algebra of  $A_0$  is given by the Hamiltonian fields of the  $u_i$ 's with  $\Delta_i = 1$ , the geometry is a global product of a trivial geometry with base  $\mathbb{C}^{\ell_0}$ and fixed fiber  $A_0$  and a rank  $(r - \ell_0)$  interacting geometry.

# B Proof of Lemma [2](#page-93-0)

Lemma. A monodromy representation associated to the graph (e) with local monodromies in the allowed lists is irreducible only when  $\varkappa = 2$  and the three semisimple monodromies have eigenvalues

$$
(1, 1, -1, -1), \quad (i, i, -i, -i), \quad (\pm \zeta_3, \pm \zeta_3, \pm \zeta_3^{-1}, \pm \zeta_3^{-1}), \tag{B.1}
$$

where  $\zeta_3$  is a primitive 3-rd root of 1.

Proof. We consider the graph (e)



The possible fugacities are

$$
\zeta_1^{-2} \pm \zeta_1 \zeta_2 \zeta_3 \zeta_3^{-2} \zeta_1^{-2} \pm \zeta_1 \zeta_1^{-2} \zeta_1^{-2} \pm \zeta_1 - 1
$$
\n
$$
-1 \pm 1 \ -1 \zeta_1^{-2} \pm \zeta_1 \zeta_2 - 1
$$
\n
$$
\zeta_1^{-2} \pm \zeta_1 \zeta_2 - 1
$$
\n(B.2)

where  $\zeta_a$  stands for a root of unity of degree 2 (i.e. a primitive 3-rd, 4-th, or 6-th root of

unity). In the first and second cases, for either sign, we have the  $R_+$  decomposition

$$
1\frac{1}{0}1 + 1\frac{1}{0}1 + 0\frac{0}{1}0 \tag{B.3}
$$

We consider the third case. If the central fugacity is  $+1$ , the central simple root  $\alpha_{\star}$  and  $(\alpha - \alpha_{\star})$  are in  $R_{+}$ . We take the opposite sign -1. Then we have the decomposition in  $R_{+}$ 

$$
1\frac{1}{1}\mathbf{1} + \mathbf{0}\frac{1}{1}\mathbf{0} + \mathbf{1}\frac{0}{1}\mathbf{0} + \mathbf{0}\frac{0}{1}\mathbf{1} + \mathbf{0}\frac{0}{1}\mathbf{0}
$$
(B.4)

It remains the forth case. Suppose  $\pm \zeta_1 \zeta_2 = 1$ . Both  $\alpha_\star$  and  $(\alpha - \alpha_\star)$  are in  $R_+$ . Suppose  $\pm \zeta_1 \zeta_2 = -1$ . Then we have the  $R_+$  decomposition

$$
2\frac{2}{3}1 + 0\frac{1}{0}1 \tag{B.5}
$$

Rearranging the order of the eigenvalues, we get the same conclusion whenever  $\pm \zeta_1 \zeta_2^{-1} = \pm 1$ . Therefore the situation with graph (e) when the representation may be irreducible is the forth case with the minus sign (possible only if  $\varkappa = 2$ ) and  $\zeta_1 = e^{\pi i/2}$  and  $\zeta_2 = \pm e^{2\pi i/3}$ .  $\Box$ 

# C Deferred proof from §. [8.7.1](#page-102-1)

**Lemma 4.** Let  $\Lambda_n$  be  $\mathbb{Z}^{2n}$  equipped with the standard principal integral symplectic form  $\langle -, - \rangle$ . We are given four elements  $n_i \in \Lambda_2$   $(i = 1, ..., 4)$  such that:

- (1) the  $n_i$  are linear independent over  $\mathbb{Q}$ ;
- (2) each  $n_i = (n_{i,a})$  is primitive i.e.  $gcd_a(n_{i,a}) = 1$  for all i;
- (3) for  $1 \leq i < j \leq 4$  we have  $\langle n_i, n_j \rangle = 2$ .

Define  $n_5 \equiv -n_1 + n_2 - n_3 + n_4$ . We have  $n_5 \equiv 0 \mod 2$ .

*Proof.* We write  $e_i$  for the standard basis elements of  $\Lambda_2$  with

$$
\langle e_1, e_2 \rangle = -\langle e_2, e_1 \rangle = \langle e_3, e_4 \rangle = -\langle e_4, e_3 \rangle = 1
$$
 (C.1)

and all other zero. Since  $n_1$  is primitive, by a  $Sp(4, \mathbb{Z})$  rotation we may assume  $n_1 = e_1$ . Then

$$
n_i = 2e_2 + a_i e_1 + z_i \quad i = 2, 3, 4, \quad a_i \in \mathbb{Z}, \quad z_i \in \mathbb{Z}e_3 + \mathbb{Z}e_4 \equiv \Lambda_1.
$$
 (C.2)

We have  $(i, j = 2, 3, 4)$ 

<span id="page-142-0"></span>
$$
\langle n_i, n_j \rangle = 2(a_i - a_j) + \langle z_i, z_j \rangle = \begin{cases} 2 & i < j \\ -2 & i < j \\ 0 & i = j \end{cases} \tag{C.3}
$$

Let  $\bar{z}_i$  be the reduction mod 2 of  $z_i$ . The  $\bar{z}_i$ 's belong to the principal symplectic  $\mathbb{F}_2$ -space  $\mathbb{F}_2^2$ . Eq. [\(C.3\)](#page-142-0) says that the span of the  $\bar{z}_i$ 's is either trivial or a Lagrangian  $\mathbb{F}_2$ -subspace. Let us show that it is not trivial. Suppose that  $\bar{z}_i = 0$  for  $i = 2, 3, 4$ . By (2)  $a_i$  must be odd for  $i = 2, 3, 4$  and so the LHS of [\(C.3\)](#page-142-0) is 0 mod 4 for all i, j, giving a contradiction. We conclude that the  $\mathbb{F}_2$ -span of  $\bar{z}_2, \bar{z}_3, \bar{z}_4$  is one-dimensional. Suppose (say) that  $\bar{z}_3 \neq 0$ . By a  $Sp(2, \mathbb{Z})$ rotation we may assume  $z_3 = (1 + 2m)e_3$  for some  $m \in \mathbb{Z}$ . Now we have

$$
n_1 = e_1
$$
  
\n
$$
n_2 = 2e_2 + a_2e_1 + b_2e_3 + 2c_2e_4
$$
  
\n
$$
n_3 = 2e_2 + a_3e_1 + (1 + 2m)e_3
$$
  
\n
$$
n_4 = 2e_2 + a_4e_1 + b_4e_3 + 2c_4e_4
$$
\n(C.4)

where  $a_2, b_2$  and, respectively,  $a_4, b_4$  cannot be both even. We have

<span id="page-143-0"></span>
$$
1 = \frac{1}{2} \langle n_2, n_3 \rangle = a_2 - a_3 - (1 + 2m)c_2
$$
  
\n
$$
1 = \frac{1}{2} \langle n_3, n_4 \rangle = a_3 - a_4 + (1 + 2m)c_4
$$
  
\n
$$
1 = \frac{1}{2} \langle n_2, n_4 \rangle = a_2 - a_4 + b_2c_4 - b_4c_2.
$$
\n(C.5)

From now on all equalities are meant to be mod 2. We have

$$
0 \equiv (a_2 - a_4) - (c_2 - c_4)
$$
  
\n
$$
1 \equiv (a_2 - a_4)(1 - b_4) + (b_2 - b_4)c_2 \mod 2
$$
 (C.6)

 $b_2$ ,  $b_4$  cannot be both odd. If they are both even,  $a_2$ ,  $a_4$  should be both odd, and the equation is still impossible. So one of the  $b_2$ ,  $b_4$  is even and the other odd. Say  $b_2 \equiv 0$  and  $b_4 \equiv 1$ . Then  $a_2 \equiv 1$ ,  $c_2 \equiv 1$ , and  $c_4 \equiv a_4$ . Eqs.[\(C.5\)](#page-143-0) give (mod 2)

$$
a_3 \equiv 1, \qquad a_4 \equiv 1. \tag{C.7}
$$

Now (mod 2)

$$
n_1 \equiv e_1 \qquad n_2 \equiv e_1
$$
  
\n
$$
n_3 \equiv e_1 + e_3 \qquad n_4 \equiv e_1 + e_3 \qquad (C.8)
$$

and

$$
n_5 \equiv 0 \bmod 2. \tag{C.9}
$$

 $\Box$ 

# D A *priori* restrictions on the allowed Kodaira types

Three out of four allowed Dynkin graphs have a  $\delta = 1$  branch: (a), (d), and (f) with the fugacity in eq.[\(8.21\)](#page-88-0). We know that  $\delta = 1$  corresponds to a non-enhanced divisor component of type  $I_n$  with  $n > 0$ . We wish to give an a priori restriction on the value of n.
Fact 37. In a non-isotrivial irreducible rank-2 special geometry with non-enhanced divisors of type  $I_{n_1}, \ldots, I_{n_t}$  we have:

- (1)  $p \nmid n_k$  for all odd primes p and  $k = 1, \ldots, t$
- (2)  $gcd(n_k) \in \{1,2\}$
- (3) in graph (d) if  $gcd(n_1, n_2) = 2$  then the dimensions are  $\{2, 4\}$

$$
(4) n_k/n_h \in \mathbb{Q}^2
$$

Argument. (1) Let us suppose  $p \mid n$ . The Crawley-Boevey solution of the Deligne-Simpson problem holds on every algebraically closed field K. We take for K the algebraic closure of  $\mathbb{F}_p$ . The reduction mod p of the local monodromy of type  $I_n$  is the identity matrix and the graph loses the corresponding branch. As a K-representation the reduction mod  $p$  of the original monodromy representation  $V_{\mathbb{K}}$  is reducible, since the graph amputated of the  $\delta = 1$  branch is not a root. Suppose  $V_{\mathbb{K}}$  is a direct sum of irreducible representations. Each summand must be a root in  $(\Sigma_{+})_p$  which now is defined using the fugacities (i.e. the eigenvalues) of the local monodromies mod  $p$  or more precisely by the roots of their minimal polynomials mod p. Minkowski theorem (and its generalizations [\[171\]](#page-158-0)) implies that if  $\ell$  is the minimal positive integer such that the integral-valued matrix M satisfies  $(\mu^{\ell}-1)^{k+1}$  for some  $k \in \mathbb{N}$ , the reduction of M mod an odd prime would satisfy a similar condition with the same minimal  $\ell$  (but possibly a different k). Hence, apart for a few special cases, the fugacities will remain roots of unity (in the appropriate algebrically closed field) of the same order after the reduction mod  $p$ . An exception is, for instance, graph (a) where the two long branches represent monodromies with minimal polynomials  $\Phi_1(z)^2 \Phi_6(z)$  and respectively  $\Phi_2(z)^2 \Phi_3(z)$ , without common roots, but which have the same order  $\ell = 6 = 2 \cdot 3$ : in this case we may have  $3 \mid n$ . However these special cases do not correspond to special geometries since no dimension pair  $\{\Delta_1, \Delta_2\}$  is consistent with these minimal polynomials (as we show in §...). For the cases which do are allowed we cannot find a decomposition of the root with the  $\delta = 1$  branch omitted into elements of  $(\Sigma_{+})_p$  (otherwise the original representation would also be reducible) i.e. the monodromy representation with  $p \mid n$  is impossible.

 $(2)(3)$  For graph (f) this follows from the direct analysis of §.[8.7.1.](#page-102-0) Graphs (a) and (d) have precisely two branches with  $\delta \neq 1$  with local monodromies  $\mu_0$ ,  $\mu_{\infty}$ . Let  $\ell = \gcd(n_k)$ which is a power of 2 by  $(1)$ . One has

$$
\det[z - \mu_{\infty}] - \det[z - \mu_0] = 0 \text{ mod } \ell,
$$
\n(D.1)

For graph (a) this relation can be satisfied only for  $\ell = 1, 2$  (with the exception mentioned above), see also §. [8.14.](#page-120-0) For graph (d) we have  $\det[z - \mu_0] = \Phi_m(z)^2$ ,  $\det[z - \mu_0] = (z^2 - 1)^2$ which are congruent modulo  $\ell > 1$  only when  $m = 4$  (i.e.  $\{\Delta_1, \Delta_4\} = \{2, 4\}$ ) in which case

$$
\Phi_4(z) - (z^2 - 1)^2 = 4z^2. \tag{D.2}
$$

However iff  $\mu_{\infty} = \mu_0 \mod 4$ , the image of  $\mu_{\infty}$  in  $GL(4,\mathbb{Z}/4\mathbb{Z})$  would be semisimple with minimal polynomial  $z^2 + 1 = 0 \mod 4$  which is inconsistent with the fact that 1 is an eigenvalue. We remain with  $gcd(n_1, n_2) = 1, 2$ .

(4) By symmetry of the graphs under interchange of any two  $\delta = 1$  branches and rigidity there exists  $U \in Sp(4,\mathbb{C})$  such that

$$
M_{\pi(i)} = UM_iU^{-1}
$$
\n(D.3)

where  $\pi$  is the permutation of the indices  $\{1, 2, \ldots, s\}$  labelling the branches which corresponds to the interchange of the given two  $\delta = 1$  branches. Since the  $M_i$ 's are integral matrices, U is defined over Q, so (by irreducibility)  $U \in Sp(4,\mathbb{Q})$ . Under conjugacy in  $Sp(4,\mathbb{Q})$  the type n of a  $I_n$  conjugacy class will change as

$$
n \rightsquigarrow n'q^2 \quad \text{with} \quad q \in \mathbb{Q}.\tag{D.4}
$$

It follows that if we have a number of non-enhanced discriminants of type  $I_{n_i}$   $(i = 1, \ldots, s)$ we must have

<span id="page-145-0"></span>
$$
\frac{n_i}{n_j} \in \mathbb{Q}^2 \quad \text{for all} \quad i, j. \tag{D.5}
$$

**Remark 27.** The authors of  $[8]$  proposed eq. $(D.5)$  on the physical basis of the Dirac quantization of electromagnetic charge.

## E Proof of Lemma [3](#page-109-0)

**Lemma.** In a non-isotrivial geometry of class  $E_7$  the short  $\delta = 2$  branch cannot stand for a non-enhanced divisor.

*Proof.* We write  $\mu_2$ ,  $\mu_3$  and  $\mu_4$  for the local monodromies associated to the branches with  $\delta = 2, 3$  and 4, respectively.

If the  $\delta = 2$  branch is non-enhanced,  $\mu_2$  is of  $I_0^*$  type i.e. semisimple with two eigenvalues  $+1$  and two  $-1$ . The fugacity of the peripheral node is  $-1$  while the contribution of this branch to the fugacity of the central node is a factor  $\pm 1$  depending on the choice of order of the two eigenvalues. In particular, we can always fix the overall sign of the central fugacity according to convenience.

Suppose  $\mu_3$  is non-semisimple. Then it should have eigenvalues  $(+1, +1, -1, -1)$  and a non-trivial Jordan block for one of the two eigenvalues. The fugacities along the branch (starting from the peripheral node) are  $(1)$ – $(-1)$ <sup>–\*</sup>. The fugacities are

$$
(-1) \qquad (-1)
$$

for some complex numbers  $\xi$ ,  $\lambda$  with  $\lambda \neq \xi^{\pm 1}$ ; when  $\lambda \neq \pm 1$   $\mu_4$  is semisimple, otherwise has a non-trivial Jordan block. Consider the decomposition

$$
1\ 1\ 2\ 2\ 1\ 1\ +\ 0\ 1\ 2\ 1\ 1\ 0\tag{E.2}
$$

Both elements are positive roots in  $R_+$  (cf. PLANCHE VI in [\[172\]](#page-158-1)), so the monodromy representation is not irreducible and should be descarded. We conclude that  $\mu_3$  should be semisimple.

Now assume  $\mu_4$  is non-semisimple. It should have eigenvalues  $(\pm 1, \pm 1, \xi, \xi^{-1})$ . By a clever ordering of the eigenvalues we get the fugacities

$$
(\eta^2) \longrightarrow (\eta) \longrightarrow (-1) \longrightarrow (1) \longrightarrow (\xi) \longrightarrow (\xi^{-2})
$$
\n(E.3)\n  
\n(5.3)

and we have the decomposition

$$
1\ 2\ 3\ 3\ 2\ 1\ +\ 0\ 0\ 1\ 0\ 0\ 0\tag{E.4}
$$

which are positive  $E_7$  roots in  $R_+$  (cf. PLANCHE VI in [\[172\]](#page-158-1)). We conclude that  $\mu_4$  should also be semisimple. Hence all three monodromies  $\mu_2$ ,  $\mu_3$  and  $\mu_4$  are semisimple and the geometry is isotrivial contrary to the assumption.  $\Box$ 

## F Quasi-isotrivial geometries

We consider the regularity conditions along the axes of the putative quasi-isotrivial geome-tries discussed in §.[8.1.3.](#page-83-0) There are two cases: dimensions  $\{3, 4\}$  and dimensions  $\{4, 6\}$ . In this appendix m is an integer attached to the Kodaira type which is equal 1 for  $I_n$ , 2 for  $I_n^*$ , 3 for  $IV^*$ , 4 for  $III^*$  and 6 for  $II^*$ . For types IV, III and II replace in the following fromulae  $1/m \mapsto (m-1)/m$  with, respectively,  $m = 3, 4$ , and 6.

We write  $\alpha_1, \alpha_2$  for the "exponents" of a local  $\mu$ -monodromy  $\mu$  around a special point corresponding to an axis. This means that that eigenvalues of  $\mu$  are  $\{e^{2\pi i\alpha_1}, e^{2\pi i\alpha_2}, e^{-2\pi i\alpha_1}, e^{-2\pi i\alpha_2}\}.$ 

**Dimensions**  $\{3, 4\}$ . Along the first axis  $(u_2 = 0)$  we must have

$$
\frac{u_2}{u_1} \left(\frac{u_1^4}{u_2^3}\right)^{\alpha_1} = u_1^{1/3}, \qquad \frac{u_2}{u_1} \left(\frac{u_1^4}{u_2^3}\right)^{\alpha_2} = \begin{cases} u_2^{1/m} u_1^x & m = 1, 2, 3, 4, 6\\ u_2^{(m-1)/m} u_1^x & m = 3, 4, 6 \end{cases}
$$
(F.1)

i.e.

$$
\{\alpha_1, \alpha_2\} = \begin{cases} \{1/3, (m-1)/3m\} & m = 1, 2, 3, 4, 6\\ \{\alpha_1, \alpha_2\} = \{1/3, 1/3m\} & m = 3, 4, 6 \end{cases}
$$
 (F.2)

 $\exp(2\pi i \alpha_2)$  should be an algebraic number of degree 2, and we have the list of possible exponents

$$
\{1/3, 1/6\}_{m=2}, \quad \{1/3, 1/4\}_{m=4}.
$$
 (F.3)

On the second axis:

$$
\frac{u_2}{u_1} \left(\frac{u_1^4}{u_2^3}\right)^{\alpha_1} = u_2^{1/4}, \qquad \frac{u_2}{u_1} \left(\frac{u_1^4}{u_2^3}\right)^{\alpha_2} = \begin{cases} u_1^{1/m} u_2^x \\ u_1^{(m-1)/m} u_2^x \end{cases}
$$
(F.4)

i.e.

$$
\{\alpha_1, \alpha_2\} = \begin{cases} \{1/4, (m+1)/4m\} & m = 1, 2, 3, 4, 6\\ \{1/4, (2m-1)/4m\} & m = 3, 4, 6 \end{cases}
$$
 (F.5)

and  $m = 1, 3$  and we have

$$
\{1/4, 1/2\}_{m=1}, \quad \{1/4, 1/3\}_{m=3} \tag{F.6}
$$

Since the monodromies around the two exceptional fibers of  $\mathcal{E}_2$  have the same exponents, a pair in this list is allowed only if it has a non-trivial overlap with the list for the first axis. If both exponents are equal the geometry becomes isotrivial, so we remain with two pairs of couples

$$
\left(\{1/3, 1/6\}_{m=2} \& \{1/4, 1/3\}_{m=3}\right) \quad \text{and} \quad \left(\{1/3, 1/4\}_{m=4} \& \{1/4, 1/2\}_{m=1}\right) \tag{F.7}
$$

which IFF they exist would correspond to geometries with dimensions  $\{3, 4\}$  and respectively

- the axis of dimension 4 is of type  $IV^*$  and the axis of dimension 3 has type  $I_0^*$  (it cannot be of type  $I_n^*$  with  $n > 0$  since the  $\mu$ -monodromy, hence the  $\varrho$ -monodromy is semisimple);
- the axis of dimension 3 has type  $III^*$  and the axis of dimension 4 has type  $I_n$  with  $n > 0$  (since  $n = 0$  will produce a "dummy" monodromy summand).

**Dimensions**  $\{4, 6\}$ . Along the first axis  $(u_2 = 0)$  we must have

$$
\frac{u_2^{1/2}}{u_1^{1/2}} \left(\frac{u_1^3}{u_2^2}\right)^{\alpha_1} = u_1^{1/4}, \qquad \frac{u_2^{1/2}}{u_1^{1/2}} \left(\frac{u_1^3}{u_2^2}\right)^{\alpha_2} = \begin{cases} u_2^{1/m} u_1^x & m = 1, 2, 3, 4, 6\\ u_2^{(m-1)/m} u_1^x & m = 3, 4, 6 \end{cases}
$$
 (F.8)

i.e.

$$
\{\alpha_1, \alpha_2\} = \begin{cases} \{1/4, (m-2)/4m\} & m = 1, 2, 3, 4, 6\\ \{1/4, (2-m)/4m\} & m = 3, 4, 6 \end{cases}
$$
 (F.9)

and the list of possible exponents is

$$
\{1/4, -1/4\}_{m=1}, \ \{1/4, 0\}_{m=2}, \ \{1/4, 1/6\}_{m=6}, \ \{1/4, -1/6\}_{m=6} \tag{F.10}
$$

but only the third one seems to be plausible for integral dimensions. On the second axis:

$$
\frac{u_2^{1/2}}{u_1^{1/2}} \left(\frac{u_1^3}{u_2^2}\right)^{\alpha_1} = u_2^{1/6}, \qquad \frac{u_2^{1/2}}{u_1^{1/2}} \left(\frac{u_1^3}{u_2^2}\right)^{\alpha_2} = \begin{cases} u_1^{1/m} u_2^x & m = 1, 2, 3, 4, 6\\ u_1^{(m-1)/m} u_2^x & m = 3, 4, 6 \end{cases}
$$
(F.11)

i.e.

$$
\{\alpha_1, \alpha_2\} = \begin{cases} \{1/6, (m+2)/6m\} & m = 1, 2, 3, 4, 6\\ \{1/6, (3m-2)/6m\} & m = 3, 4, 6 \end{cases}
$$
 (F.12)

and we have

 $\{1/6, 1/2\}_{m=1}$ ,  $\{1/6, 1/3\}_{m=2}$ ,  $\{1/6, 1/4\}_{m=4}$ , (F.13)

We disregard the last possibility because it has two exponents in common with the monodromy around the other axis. We remain with two possibilities

- dimension  $\{4,6\}$  where the axis of dimension 4 has Kodaira type  $II^*$  and the axis of dimension 6 has type  $I_n$  with  $n > 0$  (since  $n = 0$  will produce a "dummy" monodromy summand);
- dimension  $\{4, 6\}$  where the axis of dimension 4 has Kodaira type  $II^*$  and the axis of dimension 6 has type  $I_0^*$ . Indeed the  $\mu$ -monodromy (hence the  $\varrho$ -monodromy) is semisimple around the second axis.

## References

- [1] N. Seiberg and E. Witten, Electric-magnetic duality, monopole condensation, and confinement in N=2 supersymmetric Yang-Mills theory, Nucl. Phys. B 426, 19 (1994) Erratum: Nucl. Phys. B 430, 485 (1994), arXiv:hep-th/9407087.
- [2] N. Seiberg and E. Witten, Monopoles, duality and chiral symmetry breaking in  $N=2$  supersymmetric QCD, Nucl. Phys. B 431, 484 (1994), arXiv:hep-th/9408099.
- [3] R. Donagi and E. Witten, Supersymmetric Yang-Mills theory and integrable systems, Nucl. Phys. B 460, 299 (1996), arXiv:hep-th/9510101.
- [4] R. Donagi, Seiberg-Witten integrable systems, in Surveys in Differential Geometry, Vol. IV (1998) pp. 83-129.
- [5] P. C. Argyres, M. Crescimanno, A. D. Shapere and J. R. Wittig, Classification of N=2 superconformal field theories with two-dimensional Coulomb branches, hep-th/0504070.
- [6] P. C. Argyres and J. R. Wittig, Classification of N=2 superconformal field theories with two-dimensional Coulomb branches. II, hep-th/0510226.
- [7] P. Argyres, M. Lotito, Y. Lü and M. Martone, Geometric constraints on the space of  $N=2$ SCFTs I: physical constraints on relevant deformations, arXiv:1505.04814 [hep-th].
- <span id="page-149-0"></span>[8] P. C. Argyres, M. Lotito, Y. Lü and M. Martone, Geometric constraints on the space of  $N=2$ SCFTs II: Construction of special Kahler geometries and RG flows, JHEP 1802, 002 (2018), arXiv:1601.00011 [hep-th].
- [9] P. C. Argyres, M. Lotito, Y. Lü and M. Martone, Expanding the landscape of  $N = 2$  rank 1 SCFTs, JHEP 1605, 088 (2016) [arXiv:1602.02764 [hep-th]].
- [10] P. Argyres, M. Lotito, Y. Lü and M. Martone, Geometric constraints on the space of  $N=2$ SCFTs III: enhanced Coulomb branches and central charges, arXiv:1609.04404 [hep-th].
- [11] P. C. Argyres and M. Martone, 4d  $N = 2$  theories with disconnected gauge groups, JHEP 1703, 145 (2017)arXiv:1611.08602.
- [12] P. C. Argyres, Y. Lü and M. Martone, Seiberg-Witten geometries for Coulomb branch chiral rings which are not freely generated, JHEP 1706, 144 (2017) [arXiv:1704.05110 [hep-th]].
- [13] P. C. Argyres, C. Long and M. Martone, The singularity structure of scale-invariant rank-2 Coulomb branches, arXiv:1801.01122 [hep-th].
- [14] P. C. Argyres and M. Martone, Coulomb branches with complex singularities, JHEP 06, 045 (2018), arXiv:1804.03152.
- [15] P. Argyres and M. Martone, Construction and classification of Coulomb branch geometries, arXiv:2003.04954.
- [16] M. Martone, Towards the classification of rank-r $\mathcal{N}=2$  SCFTs. Part I. Twisted partition function and central charge formulae, JHEP 12, 021 (2020), arXiv:2006.16255.
- [17] P.C. Argyres and M. Martone, Towards a classification of rank  $r N = 2$  SCFTs Part II: special Kahler stratification of the Coulomb branch, arXiv:2007.00012.
- [18] P. C. Argyres and M. Martone, The rank 2 classification problem I: scale invariant geometries, arXiv:2209.09248.
- [19] P. C. Argyres and M. Martone, The rank 2 classification problem II: mapping scale-invariant solutions to SCFTs, arXiv:2209.09911.
- [20] P. C. Argyres and M. Martone, The rank-2 classification problem III: curves with additional automorphisms, arXiv:2209.10555.
- [21] M. Martone, The constraining power of Coulomb Branch Geometry: lectures on Seiberg-Witten theory, arXiv:2006.14038.
- [22] M. Caorsi and S. Cecotti, Geometric classification of 4d  $\mathcal{N}=2$  SCFTs, JHEP 07, 138 (2018), arXiv:1801.04542.
- [23] M. Caorsi and S. Cecotti, Special Arithmetic of Flavor, JHEP 08, 057 (2018) [arXiv:1803.00531 [hep-th]].
- [24] M. Caorsi and S. Cecotti, Homological classification of 4d  $\mathcal{N}=2$  QFT. Rank-1 revisited," JHEP 10 (2019), 013, arXiv:1906.03912.
- [25] S. Cecotti, M. Del Zotto, M. Martone and R. Moscrop, The characteristic dimension of four-dimensional  $\mathcal{N} = 2$  SCFTs," Commun. Math. Phys. 400, no.1, 519-540 (2023), arXiv:2108.10884 [hep-th].
- [26] S. Cecotti, Hwang-Oguiso invariants and frozen singularities in special geometry, arXiv:2304.04481.
- [27] B. de Wit and A. Van Proeyen, Potentials and symmetries of general gauged  $N = 2$ supergravity-Yang-Mills models, Nucl. Phys. B245 (1984) 89-117.
- [28] S. Cecotti, S. Ferrara and L. Girardello, Geometry of Type II superstrings and the moduli of superconformal field theories, Int. J. Mod. Phys. A 4 (1989) 2475.
- [29] S. Cecotti,  $N = 2$  Supergravity, Type IIB superstrings and Algebraic Geometry, Com- mun. Math. Phys. 131 517 (1990).
- [30] A. Strominger, Special geometry, Comm. Math. Phy. 133 (1990)163-180.
- [31] D.S. Freed, Special Kähler manifolds, Commun. Math. Phys. 203 (1999) 31-52.
- [32] S. Cecotti, Special geometry and the swampland, JHEP 09, 147 (2020), arXiv:2004.06929.
- [33] S. Cecotti, Supersymmetric field theories: Geometric structures and dualities, Cambridge University Press, 2015.
- [34] P.A. Griffiths, Periods of integrals on algebraic manifolds, III (Some global differential- geometric properties of the period mapping), Publ. IHES 38 (1970) 125-180.
- [35] P. Griffiths, Topics in Transcendental Algebraic Geometry, PUP (1984).
- [36] P. Deligne, Travaux de Griffiths, Séminaire Boubaki Exp. 376, Springer Lecture Notes in Mathematics 180, Springer (1970).
- [37] P. Griffiths, Mumford-Tate groups, notes available at https://publications.ias.edu/sites/default/files/Trieste.pdf.
- [38] M. Green, P. Griffiths, M. Kerr, Mumford-Tate domains, Boll. Unione Mat. Ital. 3 (2010) 281-307. Available on-line at https://www.math.wustl.edu/ matkerr/MTD.pdf.
- [39] M. Green, P. Griffiths and M. Kerr, Mumford-Tate groups and domains: their geometry and arithmetic, Annals of Mathematics Studies, PUP 2012.
- [40] J. Carlson, S. Müller-Stach, C. Peters, Period mappings and period domains, Second Edition, Cambridge studies in advanced mathematics 168, CUP 2017.
- [41] R. Bryant and P. Griffiths, Some observations on the infinitesimal period relations for regular threefolds with trivial canonical bundle, in M. Artin and J. Tate (Eds.) Arithmetic and Geometry. Papers dedicated to I.R. Shafarevich, vol. II Birkäuser (1983) pages 77-102.
- [42] M. Green, P. Griffiths and M. Kerr, Hodge theory, complex geometry, and representation theory, AMS (2013).
- [43] A. Cannas da Silva, Lectures on symplectic geometry, Lecture Notes in Mathematics 1764, Springer, 2008.
- [44] C. Boyer, K. Galicki, Sasakian geometry, Oxford Mathematical Monographs, OUP 2008.
- [45] J.-M. Hwang and K. Oguiso, Characteristic foliation on the discriminant hypersurface of a holomorphic Lagrangian fibration, Amer. J. Math. 131 (2009), no. 4, 981-1007, arXiv:0710.2376.
- [46] J.-M. Hwang and K. Oguiso, Multiple fibers of holomorphic Lagrangian fibrations, Commun. Contemp. Math. 13 (2011), no. 2, 309-329. arXiv:0907.4869
- [47] J.-M. Hwang and K. Oguiso, Local structure of principally polarized stable Lagrangian fibrations, arXiv:1007.2043
- [48] D. Matsushita, On singular fibres of Lagrangian fibrations over holomorphic symplectic manifolds, Math. Ann. 321 (2001), no. 4, 755-773.
- [49] D. Matsushita, Higher direct images of dualizing sheaves of Lagrangian fibrations, Amer. J. Math. 127 (2005), no. 2, 243-259.
- [50] J. Sawon, Singular fibres of very general Lagrangian fibrations, arXiv:1905.03386.
- [51] K. Kodaira, On compact analytic surfaces, in Analytic Functions, Princeton University Press, 1960, pp. 121-135.
- [52] K. Kodaira, On compact analytic surfaces, II, Ann. of Math. 77, 563-626 (1963)
- [53] K. Kodaira, On compact analytic surfaces, III, Ann. of Math. 78, 1-40 (1963).
- [54] W. Barth, K. Hulek, C. Peters, and A. van de Ven, Compact complex surfaces, Second enlarged edition. Springer 2004.
- [55] S. Cecotti, M. Del Zotto, M. Martone and R. Moscrop, to appear (hopefully).
- [56] J. Sawon, On the discriminant locus of a Lagrangian fibration, Math. Ann. 341 (2008) 201- 221.
- [57] A. Arakelov, Families of algebraic curves with fixed degeneracies. Izv. Akad. Nauk. SSSR, Ser. Math, 35 (1971) 1277-1302.
- [58] G. Faltings, Arakelov's theorem for Abelian varieties, Invent. Math. 73 (1983) 337-347.
- [59] M.-H. Saito, Classification of non-rigid families of abelian varieties, Tohoku Math. J. 45 (1993) 159-189.
- [60] C. Peters, Rigidity for variations of Hodge structure and Arakelov-type finiteness theorems, Comp. Math. 75 (1990) 113-126.
- [61] C. Peters, Rigidity, past and present. In *Teichmüler theory and moduli problem*, volume 10 of Ramanujan Math. Soc. Lect. Notes Ser., pages 529-548, Mysore, 2010. Ramanujan Math. Soc.
- [62] G. Pearlstein, C. Peters, Holomorphic bisectional curvature and applications to deformations and rigidity for variations of mixed Hodge structure, arXiv:2009.05857.
- [63] C. Simpson, Higgs bundles and local systems, Publ. Math. IHES 75 (1992) 5-95.
- [64] C. Simpson, Moduli of representations of the fundamental group of a smooth variety. Publ. Math. IHES 79 (1994) 47-129.
- [65] C. Simpson, Moduli of representations of the fundamental group of a smooth projective variety, II. Publ. Math. IHES 80 5-79.
- [66] E. Witten, Topological quantum field theory, Comm. Math. Phys. 117 (1988) 353-386.
- [67] E. Witten, "Quantum Field Theory and the Jones Polynomial," Commun. Math. Phys. 121 351 (1989).
- [68] J. Labastida and M. Mariño, Topological Quantum Field Theory and Four Manifolds, Springer, 2005.
- [69] S. Cecotti and C. Vafa, Topological antitopological fusion, Nucl. Phys. B 367, 359-461 (1991).
- [70] D. Eisenbud, Commutative Algebra with a view toward Algebraic Geometry, Graduate Texts in Mathematics 150, Springer 1995.
- [71] I. Dolgachev, Weighted projective varieties, in Group actions and vector fields, Lecture Notes in Mathematics 956, Springer, pp. 34-71.
- [72] E. Witten, Monopoles and four manifolds, Math. Res. Lett. 1 (1994) 769, arXiv:hepth/9411102.
- [73] E. Witten, From superconductors and four-manifolds to weak interactions, Bull. Am. Math. Soc. 44 (2007) 361-391.
- [74] D. Gaiotto, G. W. Moore, and A. Neitzke, Four-dimensional wall-crossing via threedimensional field theory, arXiv:0807.4723.
- [75] S. Bosch, W. Lütkebohmert, M. Raynaud, Néron Models, Modern Surveys in Mathematics 21, Springer, 1990.
- [76] V.I. Arnold, A.B. Givental, and D.P. Novikov, Symplectic geometry, in Dynamical systems IV, Encyclopaedia of Mathematical Sciences 4, Springer 2001, pp. 1-138.
- [77] C. Birkenhake, H. Lange, Complex Abelian varieties, second edition, A Series of Comprehensive Studies in Mathematics 302, Springer 2004.
- [78] B. Hunt, The geometry of some special arithmetic quotients, Lecture Notes in Mathematics 1637, Springer 1996.
- [79] C.A.M. Peters and J.H.M. Steenbrink, Monodromy of variations of Hodge structure, Acta Appl. Math. 75 (2003) 183-194.
- [80] A. Dimca, Singularities and topology of hypersurfaces, Universitext, Springer 1992.
- [81] M. Oka, On the fundamental group of the complement of certain plane curves, J. Math. Soc. Japan 30 (1978) 579-597.
- [82] J. Harris, On the Severi problem, Invent. Math. 84 (1986) 445-461.
- [83] E.R. van Kampen, On the fundamental group of an algebraic curve, Amer. J. Math. 55 (1933) 255-260.
- [84] D. Cheniot, Une demonstration du theoreme de Zariski sur les sections hyperplanes d'une hypersurface projective et du theoreme du Van Kampen sur Ie group fondamental du compiementaire d'une courbe projective plane, Compositio Math. 27 (1973) 141-158.
- [85] R. Miranda, The Basic Theory of Elliptic Surfaces, ETS Editrice, Pisa, 1989.
- [86] M. Schütt, T. Shioda, Mordell-Weil lattices, A Series of Modern Surveys in Mathematics 70, Springer 2019.
- [87] J. Sawon, A finiteness theorem for Lagrangian fibrations, J. Algebraic Geom. 25 (2016), 431- 459.
- [88] J.S. Milne, Shimura varieties and motives (2018), available on line from the author's web-page: www.jmilne.org/math/
- [89] M. Martone, Testing our understanding of SCFTs: a catalogue of rank-2  $\mathcal{N}=2$  theories in four dimensions,' JHEP 07, 123 (2022), arXiv:2102.02443 [hep-th].
- [90] D.W. Morris, Introduction to Arithmetic Groups, book available on-line at arXiv:math/0106063.
- [91] A. Grothendieck, Éléments de géométrie algébrique : II. Étude globale élémentaire de quelques classes de morphismes, Publications mathématiques de l'I.H.É.S., tome 8 (1961), p. 5-222.
- [92] A.L. Besse, Einstein Manifolds, Springer (2008).
- [93] G.C. Shephard and J.A. Todd, Finite unitary reflection groups, Can. J. Math. 6 (1954), 274-304.
- [94] C. Chevalley, Invariants of finite groups generated by reflections, Amer. J. Math. 77 (1955) 778-782.
- [95] L. Bhardway, Y. Tachikawa, Classification of 4d  $N = 2$  gauge theories, JHEP 12 (2013) 100, arXiv:1309.5160.
- [96] P. Deligne, Equations differentielles a points singuliers reguliers, Lecture Notes in Mathematics 163, Springer, 1970.
- [97] D. van Straten, Calabi–Yau Operators, arXiv:1704.00164.
- [98] G. Almkvist, W. Zudilin, Differential equations, mirror maps and zeta values, in: Mirror symmetry. V, 481515, AMS/IP Stud. Adv. Math., 38, Amer. Math. Soc., Providence, RI, (2006). arXiv:math/0402386.
- [99] A. Beauville, Monodromie des systèmes différentiels linéaires à poles simples sur la sphère de Riemann, Astérisque tome 216 (1993), Séminarie Bourbaki, exp. no. 765 pp.103-119.
- [100] Y. Haraoka, Linear differential equations in the complex domain. From classical theory to forefront, Lecture Notes in Mathematics 2271, Springer 2020.
- [101] P. Deligne, Variétés de Shimura: Interprétation modulaire, et techniques de construction de modéles canoniques, Proc. Symp. Pure Math., 33, 1979, part 2, 247-290.
- [102] N.M. Katz, Rigid local systems, PUP 1996.
- [103] W. Schmid, Variations of Hodge structure: the singularities of the period mapping. Inv. Math. 22 (1973) 211-320.
- [104] E. Viehweg, Quasi-projective moduli for polarized manifolds, A Series of Modern Surveys in Mathematics, Springer, 1995.
- [105] S. Lang, Abelian varieties, Interscience, New York 1959; reprinted by Springer-Verlag, 1983
- [106] V. Gonzáles-Aguileira, R.E. Rodríguez, Families of irreducible principally polarized Abelian varieties isomorphic to a product of elliptic curves, Proc. AMS 128 (1999) 629-636.
- [107] A. Carocca, V. Gonzáles-Aguileira, R.E. Rodríguez, Weyl groups and Abelian varieties, arXiv:math/0503340.
- [108] A. Braverman, M. Finkelberg, H. Nakajima, Towards a mathematical definition of Coulomb branches of 3-dimensional  $\mathcal{N} = 4$  gauge theories, II arXiv:1601.03586.
- [109] S. Cecotti, to appear.
- [110] D. Gaiotto, N=2 dualities, JHEP 08 (2012), 034, arXiv:0904.2715 [hep-th].
- [111] D. Gaiotto, G.W. Moore, A. Neitzke, Wall-crossing, Hitchin systems, and the WKB approximation, arXiv:0907.3987.
- [112] P. Deligne, D. Mumford, The irreducibility of the space of curves of given genus, Publications Mathématiques de l'IHÉS (Paris) 36 75-109 (1969).
- [113] J. Harris, I. Morrison, The moduli of curves, Springer-Verlag, 1998.
- [114] P. C. Argyres and N. Seiberg, S-duality in N=2 supersymmetric gauge theories, arXiv:0711.0054.
- [115] J. Minahan and D. Nemeschansky, An  $N=2$  superconformal fixed point with E6 global symmetry, Nucl. Phys. B482 (1996) 142-152, arXiv:hep-th/9608047.
- [116] J. Minahan and D. Nemeschansky, Superconformal fixed points with En global symmetry, Nucl. Phys. B489 (1997) 24-46, arXiv:hep-th/96010076.
- [117] F. Diamond and J. Shurman, A first course in modular forms, Graduate Texts in Mathematics 228, Springer, 2005.
- [118] A. Fujiki, Finite automorphism groups of complex tori of dimension 2, Publ. RIMS 24 (1988) 1-97.
- [119] A. Borel, Some metric properties of arithmetic quotients of symmetric spaces and an extension theorem, J. Diff. Geom. 6 (1972) 543-560.
- [120] A. Borel and L. Ji, Compactifications of symmetric and locally symmetric spaces, Birkäuser (2006).
- [121] S. Cecotti, Solving a puzzle in the rank  $2 \mathcal{N} = 2$  classification by Argyres and Martone, arXiv:2209.15199.
- [122] U. Persson, Configurations of Kodaira fibers on rational elliptic surfaces, Math. Z. 205 (1990) 1-47.
- [123] R. Miranda, Persson's list of singular fibers for a rational elliptic surface, Math. Z. 205 (1990) 191-211.
- [124] P. Deligne, Un théorème de finitude pour la monodromie, in Discrete groups in geometry and analysis (New Haven, Conn., 1984), 1–19, Progr. Math., 67, Birkähuser 1987.
- [125] Y. André, *G-functions and geometry*, Vieweg (1989).
- [126] Y. André and F. Baldassarri, Geometric theory of G-functions. Arithmetic ge- ometry (Cortona, 1994), 1-22. Cambridge (1997).
- [127] S. Cecotti, A. Neitzke and C. Vafa, Twistorial topological strings and a tt\* geometry for  $\mathcal{N}=2$ theories in 4d, Adv. Theor. Math. Phys. 20, 193-312 (2016), arXiv:1412.4793.
- [128] W. Crawley-Boevey, On matrices in prescribed conjugacy classes with no common invariant subspace and sum zero, Duke Math. J. 118 (2003) 339-352, arXiv:math/0103101.
- [129] W. Crawley-Boevey, Indecomposable parabolic bundles and the existence of matrices in prescribed conjugacy class closures with product equal to the identity, Publ. Math. Inst. Hautes Études Sci. 100 (2004) 171-207, arXiv:math/0307246.
- [130] W. Crawley-Boevey and P. Shaw, Multiplicative preprojective algebras, middle convolution and the Deligne-Simpson problem, Adv. Math. 201 (2006) 180-208, arXiv:math/0404186.
- [131] W. Crawley-Boevey, Quiver algebras, weighted projective lines, and the Deligne-Simpson problem, in: Sanz-Solé et al. (Eds.), International Congress of Mathematicians, vol. 2 (Madrid, 2006), European Mathematical Society, 2007, pp. 117-129, arXiv:math/0604273.
- [132] W. Crawley-Boevey, Connections for weighted projective lines, arXiv:0904.3430.
- [133] W. Crawley-Boevey, Monodromy for systems of vector bundles and multiplicative preprojective algebras, arXiv:1109.2018.
- [134] S. Cecotti, Fuchsian ODEs as Seiberg dualities, arXiv:2212.09370.
- [135] J.-P. Serre, Algebraic groups and class fields, Graduate Text in Mathematics 117, Springer 1988.
- [136] P. Deligne, Théorie de Hodge II, Publ. Math. IHES 40 (1971) 5-57.
- [137] C.T. Simpson, Product of matrices, in Differential geometry, global analysis and topology, Proceedings of the Halifax symposium, Canad. Math. Soc. Conf. Proc. 12 AMS (1991) pp. 157-185.
- [138] V.P. Kostov, The Deligne-Simpson problem a survey, J. of Algebra 281 (2004) 83-108, arXiv:math/0206298.
- [139] V. P. Kostov, On the Deligne-Simpson problem, C. R. Acad. Sci. Paris Sér. I Math. 329 (1999), 657-662.
- [140] V. P. Kostov, On some aspects of the Deligne-Simpson problem, arXiv:math.AG/0005016.
- [141] V. P. Kostov, On the Deligne-Simpson problem, arXiv:math.AG/0011013.
- [142] É. Goursat, Sur le fonctions d'une variable analogues aux fonctions hypergéométriques, Annales scientifiques de I.É.N.S. 3e série 3 (1886) 107-136.
- [143] D. Radchenko, F. Rodriguez Villegas, Goursat rigid local systems of rank four. Res. Math. Sci. 5 38 (2018), arXiv:1803.08379.
- [144] V.P. Kostov, Some examples related to the Deligne-Simpson problem, Second International Conference on Geometry, Integrability and Quantization Varna 2000, I.M. Mladenov and G.L. Naber eds., Coral Press, Sofia 2001, arXiv:math/0011015.
- [145] V.G. Kac, Infinite root systems, representations of graphs and invariant theory, Inv. Math. 56 (1980) 57-92.
- [146] Y. Haraoka, Finite monodromy of Pochhammer equation, Annales Inst. Fourier, 44 (1994) 767-810.
- [147] K. Mimachi, Solutions to some families of Fuchsian differential equations free from accessory parameters in terms of the integrals of Euler type, Adv. Studies in Pure Math. 62 (2012) Arrangements of Hyperplanes, pp. 233-260.
- [148] T. Sasai, On a monodromy group and irreducibility conditions of a fourth order Fuchsian differential system of Okubo type, J. Reine Angew. Math. 299/300 (1978) 38-50.
- [149] Y. Haraoka, Monodromy representations of systems of differential equations free from accessory parameters, SIAM J. Math. Anal. 25 (1994) 1595-1621.
- [150] A.H.M. Levelt, Hypergeometric functions, Thesis, University of Amsterdam, 1961.
- [151] F. Beukers, G. Heckman, Monodromy for the hypergeometric function  $nF_{n-1}$ , Inven. math. 95 (1989) 325-354.
- [152] M. Newman, Integral Matrices, Academic Press, New York, 1972.
- [153] R. Randell, The fundamental group of the complement of a union of complex hyperplanes, Inven. Math. 69 (1982) 103-108.
- [154] Y. Haraoka, T. Kikukawa, Rigidity of monodromies for Appell's hypergeometric functions. Opuscula Math. 35 (2015) 567-594.
- [155] I.V. Dolgachev, Classical Algebraic Geometry: a modern view, Cambridge University Press, 2012.
- [156] A. Dimca, Hyperplane arrangements. An introduction, Universitext, Springer, (2017).
- [157] A. Oka and K. Sakamoto, Product theorem of the fundamental group of a reducible curve, J. Math. Soc. Japan 30 (1978) 599-602.
- [158] E. Artal, J.I. Cogolludo-Agustín, D. Matei, Quasi-projectivity, Artin-Tits groups, and pencil maps, arXiv:1005.5225.
- [159] R. Blasco-García, J.I. Cogolludo-Agustín, Quasi-projectivity of even Artin groups, arXiv:1803.05274.
- [160] E.A. Bartolo, J.I. Cogolludo-Agustín, J. Martín-Morales, Cremona transformations of weighted projective planes, Zariski pairs, and rational cuspidal curves, arXiv:2001.07232.
- [161] P. Appell and J. Kampé de Fériet, Fonctions hypergéométriques et hypersphériques. Polynomes d'Hermite, Gauthier-Villars, Paris, 1926.
- [162] Y. Haraoka, Y. Ueno, Rigidity for Appell's hypergeometric series F4. Funkcial. Ekvac. 51 (2008) 149-164.
- [163] M. Kato, The Riemann problem for Appell's F4. Mem. Fac. Sci. Kyushu Univ. Ser. A Math. 47 (1993) 227-243.
- [164] Y. Haraoka, T. Matsumura, Three-dimensional representations of braid groups associated with some finite complex reflection groups, Int. J. Math. 28 (2017) 1750109, 44 p.
- [165] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi, Higher transcendental functions. Vol. I, McGraw-Hill Book Company, Inc. 1953.
- [166] B. Hunt, The geometry of some special arithmetic quotients, Lecture Notes in Mathematics 1637, Springer, 1996.
- [167] A. Ebisu, Y. Haraoka, M. Kaneko, H. Ochiai, T. Sasaki, M. Yoshida, A study of a Fuchsian system of rank 8 in 3 variables and the ordinary differential equations as its restrictions, arXiv:2005.04465.
- [168] G. Lauricella, Sulle funzioni ipergeometriche in più variabili, Rend. Circ. Math. Palermo 7 (1893) 111-158.
- [169] R. Hattori, N. Takayama, The singular locus of Lauricella's  $F_C$ , J. Math. Soc. Japan 66 (2014) 981-995, arXiv:1110.6675.
- [170] Y. Goto, Appell-Lauricella's hypergeometric functions and intersection theory, Proceedings of Science, PoS(MA2019) 008 (2019).
- <span id="page-158-0"></span>[171] J.-P. Serre, Bounds for the orders of the finite subgroups of  $G(k)$ , arXiv:1011.0346.
- <span id="page-158-1"></span>[172] N. Bourbaki, Éléments de Mathématique. Groupes et algèbres de Lie. Chapitres 4 à 6, Springer, 1981.