

NON-BACKTRACKING EIGENVALUES AND EIGENVECTORS OF RANDOM REGULAR GRAPHS AND HYPERGRAPHS

XIANGYI ZHU AND YIZHE ZHU

ABSTRACT. The non-backtracking operator of a graph is a powerful tool in spectral graph theory and random matrix theory. Most existing results for the non-backtracking operator of a random graph concern only eigenvalues or top eigenvectors. In this paper, we take the first step in analyzing its bulk eigenvector behaviors. We demonstrate that for the non-backtracking operator B of a random d -regular graph, its eigenvectors corresponding to nontrivial eigenvalues are completely delocalized with high probability. Additionally, we show complete delocalization for a reduced $2n \times 2n$ non-backtracking matrix \tilde{B} . By projecting all eigenvalues of \tilde{B} onto the real line, we obtain an empirical measure that converges weakly in probability to the Kesten-McKay law for fixed $d \geq 3$ and to a semicircle law as $d \rightarrow \infty$ with $n \rightarrow \infty$. We extend our analysis to random regular hypergraphs, including the limiting measure of the real part of the spectrum for \tilde{B} , ℓ_∞ -norm bounds for the eigenvectors of \tilde{B} and B , and a deterministic relation between eigenvectors of B and the eigenvectors of the adjacency matrix.

As an application, we analyze the non-backtracking spectrum of the (d_1, d_2) -regular stochastic block model (RSBM) and provide a spectral method based on eigenvectors of \tilde{B} to recover the community structure exactly. We also show that there exists an isolated real eigenvalue with an informative eigenvector inside the circle of radius $\sqrt{d_1 + d_2 - 1}$ in the spectrum of B , analogous to the ‘‘eigenvalue insider’’ phenomenon for the Erdős-Rényi stochastic block model conjectured in [23].

1. INTRODUCTION

1.1. Non-backtracking operators of random graphs. The non-backtracking operator is an important object in the study of spectral graph theory [53, 5, 8, 37, 30, 44, 36, 54]. It has recently been used as a powerful tool for studying random matrices [15, 12, 11] and for designing efficient algorithms in community detection and matrix completion [38, 15, 49, 14, 50, 51]. Many recent results on the spectrum of the non-backtracking operator have been established in various graph models, such as Erdős-Rényi graphs [56], stochastic block models [15, 22], inhomogeneous random graphs [11, 27], random regular graphs [12], and bipartite biregular graphs [18]. Very recently, some of these results have also been extended to hypergraphs [26, 50, 20]. However, so far, these results mainly focus on top eigenvalues and eigenvectors and global spectral distribution, while not much is known about bulk eigenvector behavior.

In this paper, we take a first step towards understanding the bulk eigenvectors of the non-backtracking operator in random graphs. Eigenvector delocalization is an important topic in the study of random matrices, demonstrating that many matrix models exhibit behavior similar to their Gaussian analog, a phenomenon known as universality. A unit eigenvector is considered delocalized if its infinity norm is close to that of a uniformly distributed random vector on the unit sphere up to a polylog factor. Eigenvector delocalization has been shown for many random matrix models, including non-Hermitian ones. However, for sparse and non-Hermitian models, the literature includes only a few results, such as those on eigenvector delocalization in random regular digraphs [40] and Erdős-Rényi digraphs [33].




FIGURE 1. Simulation for eigenvalues of \tilde{B} and B for a random 5-regular graph with 200 vertices.

The non-backtracking operator B for a random d -regular graph is a sparse and non-Hermitian operator, where each row and column of B has exactly $(d - 1)$ many nonzero entries. It can also be viewed as the adjacency matrix of a random $(d - 1)$ -regular digraph with (nd) many vertices. However, not all $(d - 1)$ -regular digraphs can be obtained in this way (see [44, Theorem 2.3]), and the eigenvalue distribution of B is very different from the conjectured oriented Kesten-McKay law for a uniformly chosen random $(d - 1)$ -regular digraph [13]. A simulation of the non-backtracking spectrum of a random d -regular graph is shown in Figure 1.

In Theorem 3.3, we show that with high probability, all eigenvectors corresponding to nontrivial non-backtracking eigenvalues are completely delocalized. The main idea behind the proof is to use the delocalization [35] and spectral gap [35, 12] results for the adjacency matrix of a random d -regular graph to understand eigenvector delocalization for the non-backtracking matrix with the help of the Ihara-Bass formula [8]. The algebraic structure of the non-backtracking operator enables the translation of eigenvalue and eigenvector information from the corresponding adjacency matrix. However, this precise algebraic connection between these two operators is only available for regular graphs. Exploring beyond d -regular graphs is an interesting future direction.

In the course of proving non-backtracking eigenvector delocalization, we show in Theorem 3.1 the convergence in probability to the Kesten-McKay law and semicircle law for fixed d and growing d when projecting the eigenvalues of the non-backtracking operator onto the real line. From the Ihara-Bass formula, another $2n \times 2n$ non-Hermitian block matrix \tilde{B} closely related to B emerges, which we call a reduced non-backtracking operator [5, 56, 22, 48]. We also demonstrate that the eigenvectors of \tilde{B} are completely delocalized with high probability in Theorem 3.2.

1.2. Extension to regular hypergraphs. The spectral theory for hypergraphs has drawn considerable interest due to its applications in number theory, community detection, and network analysis. [28, 39, 47, 52] In [6, 50, 20], spectral algorithms based on the non-backtracking eigenvectors were studied for community detection problems in hypergraph stochastic block models. Several results we obtained for random regular graphs can be generalized to random regular hypergraphs, which is a popular hypergraph model studied in combinatorics, statistical physics, and computer science [21, 26, 31, 19, 46].

In Theorem 3.7, we show that by projecting the eigenvalues of a non-backtracking operator B for random regular hypergraphs, we can obtain different limiting measures depending on the regime of d, k . We provide an exact spectral relation between a reduced non-backtracking operator \tilde{B} and the adjacency matrix A of a regular hypergraph Lemma in 7.1. A precise relation between eigenvectors

of B and A is given in Lemma 8.1, which generalizes the result for regular graphs in [41]. These relations are used to show ℓ_∞ -norm bound on eigenvectors of \tilde{B} and B in Theorem 3.8.

1.3. Application in a regular stochastic block model. Community detection in the stochastic block model is an important topic in statistical physics, machine learning, and network science [1]. Using the non-backtracking operator to detect the community structure is proven to achieve the information-theoretical threshold in many settings [15, 50]. However, the spectrum of the non-backtracking operator for stochastic block models is not fully understood. An intriguing behavior of the eigenvalue for the non-backtracking matrix B in the Erdős-Rényi stochastic block model with constant expected degree was observed in [23] that there exists a real eigenvalue inside the bulk spectrum of B close to the ratio of the top two eigenvalues of B . This was justified in [22] in the dense regime when the expected degree is $\omega(\log n)$. We study an analog of the stochastic block model in the random regular graph setting, which is called the *regular stochastic block model* (RSBM) [17, 7, 45] studied in the literature, which exhibits different behaviors from the Erdős-Rényi SBM. The RSBM is constructed by combining a random d_2 -regular bipartite regular graph and a random d_1 -regular graph; see Definition 3.9 for a precise description.

We confirm the *eigenvalue insider* phenomenon in the RSBM in Theorem 3.10 and Corollary 3.11: When $(d_1 - d_2)^2 > 4(d_1 + d_2 - 1)$, a real eigenvalue of \tilde{B} exists inside the circle of radius $\sqrt{d_1 + d_2 - 1}$, and the corresponding eigenvector reveals the community structure exactly. In contrast to the Kesten-Stigum threshold in the Erdős-Rényi SBM [43], even when $(d_1 - d_2)^2 < 4(d_1 + d_2 - 1)$, other methods were conjectured to achieve exact recovery in the RSBM in the statistical physics literature [7].

Organization of the paper. The rest of the paper is organized as follows. In Section 2, we provide some basic notations and definitions for regular graphs, the non-backtracking operator, and hypergraphs. In Section 3, we state the main results for random regular graphs, random regular hypergraphs, and the regular stochastic block model, respectively. Sections 4 to 9 contain all the proofs. In Section 10, we discuss the open problem of generalizing the results to Erdős-Rényi graphs.

2. PRELIMINARIES

2.1. Regular graphs. Let $G = (V, E)$ be a graph. G is d -regular of size n if each vertex has degree d and $|V| = n$. For given n and d , we say G is a *random d -regular graph* if it is uniformly chosen from all d -regular graphs with n vertices.

The (i, j) -th entry of the *adjacency matrix* A of a graph G is defined as

$$A_{ij} = \begin{cases} 1 & \text{if } \{i, j\} \in E, \\ 0 & \text{otherwise.} \end{cases}$$

The degree matrix D of a graph G is a diagonal matrix where $D_{ii} = \sum_{j \in V} A_{ij}$. Define the oriented edge set \vec{E} for G as

$$\vec{E} = \{(i, j) : \{i, j\} \in E\}. \tag{1}$$

Each edge yields two oriented edges; therefore, $|\vec{E}| = 2|E|$.

Definition 2.1 (Non-backtracking operator). The *non-backtracking operator* B of G is a non-Hermitian operator of size $|\vec{E}| \times |\vec{E}|$. For any $(u, v), (x, y) \in \vec{E}$, B is defined as

$$B_{(u,v),(x,y)} = \begin{cases} 1 & v = x, u \neq y, \\ 0 & \text{otherwise.} \end{cases}$$




FIGURE 2. A $(2, 3)$ -regular hypergraph with 9 vertices

In particular, for a d -regular graph with n vertices, the corresponding B is of size $nd \times nd$. A useful identity we will use in this paper is the following Ihara-Bass formula [8].

Lemma 2.2 (Ihara-Bass formula). *For any graph $G = (V, E)$, and any $z \in \mathbb{C}$, the following identity holds:*

$$\det(B - zI) = (z^2 - 1)^{|E|-n} \det(z^2I - zA + D - I). \quad (2)$$

Define a block matrix

$$\tilde{B} = \begin{bmatrix} 0 & D - I \\ -I & A \end{bmatrix} \in \mathbb{R}^{2n \times 2n}.$$

Then from (2), we have

$$\det(B - zI) = (z^2 - 1)^{|E|-n} \det(\tilde{B} - zI). \quad (3)$$

The identity (3) implies that B and \tilde{B} share the same spectrum, up to the multiplicity of trivial eigenvalues ± 1 . We also call \tilde{B} the *reduced non-backtracking matrix* of G . When G is a d -regular graph with n vertices, (2) can be further simplified to

$$\det(B - zI) = (z^2 - 1)^{nd/2-n} \det(z^2I - zA + (d - 1)I). \quad (4)$$

2.2. Regular hypergraphs. In this section, we include some standard definitions of hypergraphs.

Definition 2.3 (Hypergraph). A *hypergraph* H consists of a set V of vertices and a set E of hyperedges such that each hyperedge is a nonempty set of V . A hypergraph H is k -uniform for an integer $k \geq 2$ if every hyperedge $e \in E$ contains exactly k vertices. The *degree* of i , denoted $\deg(i)$, is the number of all hyperedges incident to i .

A hypergraph is d -regular if all of its vertices have degree d . A hypergraph is (d, k) -regular if it is both d -regular and k -uniform. See Figure 2 for an example.

Definition 2.4 (Adjacency matrix of a hypergraph). For a hypergraph H with n vertices, we associate a $n \times n$ symmetric matrix A called the *adjacency matrix* of H . For $i \neq j$, we define A_{ij} as the number of hyperedges containing both i and j ; we define $A_{ii} = 0$ for all $1 \leq i \leq n$. When the hypergraph is 2-uniform, this is the definition for the adjacency matrix of a graph.

3. MAIN RESULTS

3.1. Random regular graphs. Let μ_1, \dots, μ_{2n} be the eigenvalues of \tilde{B} . Define x_i to be the real part of μ_i . The empirical measure of $\{x_i, 1 \leq i \leq 2n\}$ is denoted as

$$\mu = \frac{1}{2n} \sum_{i=1}^{2n} \delta_{x_i}.$$




FIGURE 3. Simulation for the empirical measure of the eigenvalue real parts of \tilde{B} for a random 5-regular graph with 2000 vertices, excluding the deterministic eigenvalue $\mu_1 = 4$.

The following theorem characterizes the eigenvalue distribution of \tilde{B} after projecting all eigenvalues to the real line.

Theorem 3.1 (Projecting eigenvalues of \tilde{B} to the real line). *The following holds for the reduced non-backtracking matrix \tilde{B} of a uniformly chosen random d -regular graph:*

- (1) *When $d \geq 3$ is a fixed integer, μ converges weakly in probability to a rescaled Kesten-McKay distribution μ_{KM} supported on $[-\sqrt{d-1}, \sqrt{d-1}]$, where*

$$\mu_{\text{KM}}(x) = \frac{2d\sqrt{(d-1)-x^2}}{\pi(d^2-4x^2)} \mathbf{1}_{\{|x| \leq \sqrt{d-1}\}}.$$

- (2) *When $d \rightarrow \infty$ as $n \rightarrow \infty$, the empirical measure of $\left\{\frac{2x_i}{\sqrt{d-1}}, i \in [2n]\right\}$ converges weakly in probability to a semicircle law μ_{SC} , where*

$$\mu_{\text{SC}}(x) = \frac{1}{2\pi} \sqrt{4-x^2} \mathbf{1}_{\{|x| \leq 2\}}. \quad (5)$$

Figure 3 is a simulation for the projected eigenvalues on the real line of a random regular graph. Similar results have been established for Erdős-Rényi graphs $G(n, p)$ with $np = \omega(\log n)$ [56] and for stochastic block models [22] in the same regime.

We now move on to study the eigenvectors of B and \tilde{B} . Note that each unit eigenvector of \tilde{B} is in \mathbb{C}^{2n} . We show they are completely delocalized based on the eigenvector delocalization results in [35] for the adjacency matrix of a random d -regular graph.

Theorem 3.2 (Eigenvector delocalization for \tilde{B}). *Let $d \geq 3$ be fixed, and \tilde{B} be the reduced non-backtracking matrix of a random d -regular graph. Let $u_i, i \in [2n]$ be the ℓ_2 -normalized eigenvector associated with μ_i . Then there exist absolute constants $C_1, C_2 > 0$ such that with probability at least $1 - n^{-C_1}$, for all $i \in [2n]$,*

$$\|u_i\|_\infty \leq \frac{\log^{C_2}(n)}{\sqrt{n}}.$$

By a different argument with the help of the Ihara-Bass formula, we can also show that unit eigenvectors of B as a vector in \mathbb{C}^{nd} are completely delocalized.

Theorem 3.3 (Eigenvector delocalization for B). *Let $d \geq 3$ be fixed, and B be the non-backtracking matrix of a random d -regular graph. Let $\mu_i, i \in [2n]$ be the eigenvalues of \tilde{B} . Then there exist*




FIGURE 4. The infinite norms of eigenvectors for A , \tilde{B} , B of a 4-regular random graph with $n = 200$.

absolute constants $C_1, C_2 > 0$ such that with probability at least $1 - n^{-C_1}$, all the unit eigenvectors w_i of B associated with $\mu_i \neq \pm 1$ satisfy

$$\|w_i\|_\infty \leq \frac{\log^{C_2}(n)}{\sqrt{nd}}.$$

The ℓ_∞ -norms of eigenvectors of B and \tilde{B} are illustrated in Figure 4. Similar results for non-backtracking eigenvectors of random d -regular graph for growing d can be proved in the same way by using the eigenvector delocalization bounds for the adjacency matrix in [24, 55, 10, 9].

Remark 3.4 (Eigenvectors corresponding to eigenvalues ± 1 in Equation (4)). In Equation (4), we see there are trivial eigenvalues ± 1 of B with an extra multiplicity $\frac{nd}{2} - n$ which are not given by eigenvalues of \tilde{B} . The structure of the eigenspaces of ± 1 was discussed in [41, Proof of Proposition 3.1].

3.2. Random regular hypergraphs.

Definition 3.5 (Non-backtracking operator of a hypergraph). For a hypergraph $H = (V, E)$, its *non-backtracking operator* B is a square matrix indexed by oriented hyperedges

$$\vec{E} = \{(i, e) : i \in V, e \in E, i \in e\}$$

with entries given by

$$B_{(i,e),(j,f)} = \begin{cases} 1 & \text{if } j \in e \setminus \{i\}, f \neq e, \\ 0 & \text{otherwise,} \end{cases}$$

for any oriented hyperedges $(i, e), (j, f) \in \vec{E}$.

This is a generalization of the graph non-backtracking operators to hypergraphs. For a k -uniform hypergraph $H = (V, E)$, define the $2n \times 2n$ reduced non-backtracking matrix \tilde{B} as

$$\tilde{B} = \begin{pmatrix} 0 & (D - I) \\ -(k-1)I & A - (k-2)I \end{pmatrix},$$

where A is the adjacency matrix of H , and D is the diagonal degree matrix with

$$D_{ii} = \#\{e \in E : i \in e\}.$$

The following Ihara-Bass formula for hypergraphs was proved in [50]. Lemma 3.6 shows that the spectrum of \tilde{B} is identical to that of B , except for possible trivial eigenvalues at -1 and $-(k-1)$.

Lemma 3.6 (Lemma 1 in [50]). *Let $H = (V, E)$ be a k -uniform hypergraph. The following identity holds for any $z \in \mathbb{C}$:*

$$\begin{aligned} \det(B - zI) &= (z-1)^{(k-1)|E|-n} (z+(k-1))^{|E|-n} \det(z^2 + (k-2)z - zA + (k-1)(D-I)) \\ &= (z-1)^{(k-1)|E|-n} (z+(k-1))^{|E|-n} \det(\tilde{B} - zI). \end{aligned}$$

We obtain the following limiting distributions for the real part of eigenvalues in \tilde{B} for a uniformly chosen random (d, k) -regular hypergraph.

Theorem 3.7 (Projecting eigenvalues of \tilde{B} to the real line). *Let \tilde{B} be the reduced non-backtracking operator of a random (d, k) -regular hypergraph. Let x_1, \dots, x_{2n} be the real part of the eigenvalues of \tilde{B} . Define μ to be the empirical measure of $\left\{ \frac{2x_i - (k-2)}{\sqrt{(d-1)(k-1)}} \right\}_{i=1}^{2n}$. Then the following holds:*

- (1) *If d, k are fixed, as $n \rightarrow \infty$, μ converges weakly in probability to a distribution supported on $[-2, 2]$ whose density function is given by*

$$\mu_{d,k}(x) = \frac{1 + \frac{k-1}{q}}{\left(1 + \frac{1}{q} - \frac{x}{\sqrt{q}}\right) \left(1 + \frac{(k-1)^2}{q} + \frac{(k-1)x}{\sqrt{q}}\right) \pi} \sqrt{1 - \frac{x^2}{4}},$$

where $q = (k-1)(d-1)$.

- (2) *If $d/k \rightarrow \alpha$ as $n \rightarrow \infty$ and $d \leq \frac{n}{32}$, μ converges weakly in probability to a distribution supported on $[-2, 2]$ with density function*

$$\mu_\alpha(x) = \frac{\alpha}{(1 + \alpha + \sqrt{\alpha x}) \pi} \sqrt{1 - \frac{x^2}{4}}.$$

- (3) *If $d \rightarrow \infty$, $d = o(n^\varepsilon)$ for any $\varepsilon > 0$ and $\frac{d}{k} \rightarrow \infty$, μ converges weakly in probability to the semicircle law given in (5).*

We obtain the following ℓ_∞ -norm bounds for eigenvectors of \tilde{B} and B .

Theorem 3.8 (ℓ_∞ -norm bound for eigenvectors of \tilde{B} and B). *Let B and \tilde{B} be the non-backtracking operator and reduced non-backtracking operator of a (d, k) -regular hypergraph, respectively. Let d, k be fixed. Then the following holds:*

- (1) *Let v_i be a unit eigenvector of A associated with the eigenvalue λ_i . Let u_i, u'_i be unit eigenvectors of \tilde{B} associated with two eigenvalues μ_i, μ'_i , which are given by the solutions of*

$$\mu^2 - (\lambda_i - k + 2)\mu + (d-1)(k-1) = 0. \quad (6)$$

Then for all $i \in [n]$,

$$\|u_i\|_\infty, \|u'_i\|_\infty \leq \|v_i\|_\infty.$$

(2) For a random (d, k) -regular graph, let w_i, w'_i be unit eigenvectors of B associated with two eigenvalues μ_i, μ'_i given by (6) with $\mu_i, \mu'_i \notin \{1, -(k-1)\}$. Then, for $d > k \geq 3$, with high probability, for all $i \in [n]$,

$$\|w_i\|_\infty, \|w'_i\|_\infty \leq \frac{\sqrt{k-1} + o(1)}{\sqrt{d-1} - \sqrt{k-1}} \|v_i\|_\infty.$$

Theorem 3.8 shows that eigenvector delocalization of A implies eigenvector delocalization for B and \tilde{B} . However, no eigenvector delocalization results are available in the literature for the adjacency matrix random regular hypergraphs. It's possible that by adapting the result of eigenvector delocalization for bipartite biregular graphs [57, Corollary 2.8] together with the connection between random regular hypergraphs and random bipartite biregular graphs established in [26], one can obtain an eigenvector delocalization bound for the adjacency matrix in the random regular hypergraph case.

3.3. Community detection in the RSBM.

Definition 3.9 (Regular stochastic block model (RSBM)). For an even integer n and two integers d_1 and d_2 , the *regular stochastic block model* with vertex set $[n]$ is obtained as follows. Choose a partition V_1, V_2 of equal size $n/2$ of the vertex set $[n]$, uniformly from among the set of all such partitions. Choose two independent copies of uniform d_1 -regular graphs with vertex set V_1 , respectively V_2 . Finally, connect the vertices from V_1 with those from V_2 by a uniformly random d_2 -bipartite regular graph.

Let $\sigma \in \{-1, 1\}^n$ be a vector for the community assignment defined by

$$\sigma_i = \begin{cases} 1 & i \in V_1 \\ -1 & i \in V_2. \end{cases}$$

The *community detection* problem for the regular stochastic block model is to observe a random graph sampled from the model and construct an estimator for the community assignment vector σ .

Note that a graph sampled from this model is a random $(d_1 + d_2 - 1)$ -regular graph but not uniformly distributed. [16, Proposition 1] shows that the RSBM is asymptotically distinguishable from a uniformly chosen random regular graph with the same degree. However, some of our results for deterministic regular graphs can still be applied to this model. We describe the real eigenvalues and eigenvectors of \tilde{B} as follows:

Theorem 3.10 (Real eigenvalues and eigenvectors of \tilde{B}). *Let \tilde{B} be the reduced non-backtracking operator of an (n, d_1, d_2) -regular stochastic block model. Then, the following holds:*

- (1) $d_1 + d_2 - 1$ is an eigenvalue of \tilde{B} with the corresponding eigenvector $v_1 = [1, \dots, 1]^\top \in \mathbb{R}^{2n}$ and 1 is an eigenvalue of \tilde{B} with the corresponding eigenvector

$$v'_1 = \left[1, \dots, 1, \frac{1}{d_1 + d_2 - 1}, \dots, \frac{1}{d_1 + d_2 - 1} \right]^\top \in \mathbb{R}^{2n}.$$

- (2) When $(d_1 - d_2)^2 > 4(d_1 + d_2 - 1)$, there are two real eigenvalues of \tilde{B} given by

$$\mu_2, \mu'_2 = \frac{d_1 - d_2 \pm \sqrt{(d_1 - d_2)^2 - 4(d_1 + d_2 - 1)}}{2}, \quad (7)$$

with the corresponding eigenvector

$$v_2 = \begin{bmatrix} \sigma \\ \frac{\mu_2}{d_1 + d_2 - 1} \sigma \end{bmatrix}, \quad v'_2 = \begin{bmatrix} \sigma \\ \frac{\mu'_2}{d_1 + d_2 - 1} \sigma \end{bmatrix}.$$




FIGURE 5. On the left is the non-backtracking spectrum of an RSBM, where we choose $d_1 = 12$ and $d_2 = 4$ with size 4000. There is an eigenvalue μ'_2 inside the circle matched with (7) given by $\frac{\mu_1}{\mu_2}$. On the right is the non-backtracking spectrum of a stochastic block model where we choose $a = 15$ and $b = 6$ with size 4000. The insider real eigenvalue is close to $\frac{\mu_1}{\mu_2}$.

- (3) When $(d_1 - d_2)^2 > 4(d_1 + d_2 - 1)$ and d_1 is even, with high probability, $\{d_1 + d_2 - 1, 1, \mu_2, \mu'_2\}$ are 4 real eigenvalues of multiplicity one, and the rest of the eigenvalues of \tilde{B} are within $o(1)$ distance from the circle of radius $\sqrt{d_1 + d_2 - 1}$.

See Figure 5 for the simulation on the spectrum of an RSBM and an Erdős-Rényi SBM. From Theorem 3.10, we obtain the following corollary for exact recovery with a spectral method in the RSBM.

Corollary 3.11 (Exact recovery in the RSBM). Assume $(d_1 - d_2)^2 > 4(d_1 + d_2 - 1)$ and d_1 is even. Let $u_2 = \begin{bmatrix} x_2 \\ y_2 \end{bmatrix}, u'_2 = \begin{bmatrix} x'_2 \\ y'_2 \end{bmatrix}$ be an eigenvector of \tilde{B} corresponding to eigenvalue μ_2, μ' , respectively. With high probability, we have $x_2, y_2, x'_2, y'_2 \in \text{span}\{\sigma\}$.

Corollary 3.11 implies both eigenvectors u_2 and u'_2 of \tilde{B} can exactly recover the community assignment σ with high probability. Such a spectral method based on the eigenvector associated with the insider eigenvalue for the Erdős-Rényi SBM was conjectured in [23] and remains open. Compared with the regular stochastic block model, the key difference for the Erdős-Rényi stochastic block model is the lack of degree concentration in the very sparse regime, which makes it challenging to establish a direct connection between eigenvectors of \tilde{B} and A .

Remark 3.12 (The parity constraint of d_1). In Theorem 3.10, the constraint that d_1 is even is a technical condition due to the proof technique in [16], which involves random lifts studied in [12, 29]. It was conjectured in [16] that such a restriction can be removed. We expect by adapting the proof techniques in [12, 18], Theorem 3.10 can be shown for all $(d_1 - d_2)^2 > 4(d_1 + d_2 - 1)$. We leave it as a future direction.

4. PROOF OF THEOREM 3.1

We first derive an algebraic relation between eigenvalues and eigenvectors of A and B when G is a d -regular graph.

Lemma 4.1 (Spectral relation between A and \tilde{B}). Let G be a d -regular graph with adjacency matrix A and non-backtracking matrix B and $d \geq 2$. Let v_i be an eigenvector of A with respect to

an eigenvalue $\lambda_i, i \in [n]$. Then

$$\mu_i, \mu'_i = \frac{\lambda_i \pm \sqrt{\lambda_i^2 - 4(d-1)}}{2}$$

are two eigenvalues of \tilde{B} , with two corresponding eigenvectors of the form

$$u_i = \begin{bmatrix} v_i \\ \frac{\mu_i}{d-1}v_i \end{bmatrix}, \quad u'_i = \begin{bmatrix} v_i \\ \frac{\mu'_i}{d-1}v_i \end{bmatrix}. \quad (8)$$

Proof. Let $\begin{bmatrix} x \\ y \end{bmatrix}, x, y \in \mathbb{C}^n$ be an eigenvector of \tilde{B} with respect to an eigenvalue μ , we have

$$\mu \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 & (d-1)I \\ -I & A \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}.$$

This equation implies

$$Ax = \left(\mu + \frac{d-1}{\mu} \right) x, \quad y = \frac{\mu}{d-1} x. \quad (9)$$

For each eigenvalue λ_i of A with a corresponding eigenvector $v_i \in \mathbb{R}^n$, there are two corresponding eigenvalue μ_i, μ'_i of \tilde{B} satisfying the quadratic equation

$$x^2 - \lambda_i x + d - 1 = 0. \quad (10)$$

In particular, $\lambda_1 = d$ gives two real eigenvalues $\mu_1 = d - 1$ and $\mu'_1 = 1$.

By counting multiplicity, Equation (10) gives all the $2n$ eigenvalues of \tilde{B} . Thus, by Equations (9) and (10), there are n eigenvectors of \tilde{B} of the form

$$u_i = \begin{bmatrix} v_i \\ \frac{\mu_i}{d-1}v_i \end{bmatrix}, \text{ where } \mu_i = \frac{\lambda_i + \sqrt{\lambda_i^2 - 4(d-1)}}{2},$$

and other n eigenvectors of the form,

$$u'_i = \begin{bmatrix} v_i \\ \frac{\mu'_i}{d-1}v_i \end{bmatrix}, \text{ where } \mu'_i = \frac{\lambda_i - \sqrt{\lambda_i^2 - 4(d-1)}}{2}.$$

□

With Lemma 4.1, we are ready to prove Theorem 3.1 for random d -regular graphs.

Proof of Theorem 3.1. From Equation (10), when $|\lambda_i|^2 \leq 4(d-1)$, we have

$$\mu_i + \mu'_i = \lambda_i, \quad |\mu_i| = |\mu'_i| = \sqrt{d-1},$$

and the real parts of μ_i, μ'_i satisfy

$$x_i = x'_i = \frac{\lambda_i}{2}. \quad (11)$$

The Kesten-McKay law in [42] shows that with high probability, the limiting eigenvalue distribution of the adjacency matrix for a random d -regular graph converges weakly to a probability measure with density

$$f(x) = \frac{d\sqrt{4(d-1) - x^2}}{2\pi(d^2 - x^2)} \mathbf{1}\{|x| \leq 2\sqrt{d-1}\}.$$

From the Kesten-McKay law above, with high probability, there are $n - o(n)$ many eigenvalues λ_i satisfies $|\lambda_i|^2 \leq 4(d-1)$. Therefore, with high probability, there are $2n - o(n)$ eigenvalues of \tilde{B}

on the circle of radius $\sqrt{d-1}$, and n pairs of x_i, x'_i satisfies (11). Therefore, the empirical measure of the real parts of μ_i converges weakly in probability to a rescaled Kesten-McKay law given by

$$\mu_{\text{KM}}(x) = \frac{2d\sqrt{(d-1)-x^2}}{\pi(d^2-4x^2)} \mathbf{1}\{|x| \leq \sqrt{d-1}\}.$$

When $d \rightarrow \infty$ as $n \rightarrow \infty$, it is shown in [55, 24] that the empirical spectral distribution of $\frac{A}{\sqrt{d-1}}$ converges weakly in probability to a semicircle law. With Equation (11), following the same argument as in the case for fixed d , the empirical measure $\frac{1}{2n} \sum_{i=1}^{2n} \delta_{2x_i/\sqrt{d-1}}$ converges weakly in probability to the semicircle law. \square

5. PROOF OF THEOREM 3.2

Our proof is based on the following eigenvector delocalization bound from [35].

Lemma 5.1 (Theorem 1.4 in [35]). *For any fixed $d \geq 3$, there exists absolute constants $C_1, C_2 > 0$ such that with probability at least $1 - n^{-C_1}$, all unit eigenvectors v_i of A for a random d -regular graph satisfies*

$$\|v_i\|_\infty \leq \frac{\log^{C_2}(n)}{\sqrt{n}}.$$

With Lemmas 4.1 and 5.1, we are able to prove Theorem 3.2.

Proof of Theorem 3.2. From (8), any eigenvector u_i of \tilde{B} associated with v_i of A satisfies

$$\frac{\|u_i\|_\infty}{\|u_i\|_2} \leq \frac{\max\left\{1, \frac{|\mu_i|}{d-1}\right\}}{\sqrt{1 + \frac{|\mu_i|^2}{(d-1)^2}}} \|v_i\|_\infty \leq \frac{\|v_i\|_\infty}{\|v_i\|_2}.$$

Then Theorem 3.2 follows from Lemma 5.1. \square

6. PROOF OF THEOREM 3.3

We first introduce a lemma contained in [41] for a deterministic relation between eigenvectors of B and eigenvectors of A and provide a proof for completeness.

Lemma 6.1 (Remark 3.4 in [41]). *Let \vec{E} be oriented edge set for a d -regular graph $G = (V, E)$ defined in (1). Let μ_i, μ'_i denote each eigenvalue pair of \tilde{B} corresponding to eigenvalue λ_i of A through the quadratic equation (10) with $|\lambda_i| \neq d$. Let v_i be a unit eigenvector of A associated λ_i . Define $\tilde{w}_i, \tilde{w}'_i \in \mathbb{C}^{nd}$ such that for any $(x, y) \in \vec{E}$,*

$$\tilde{w}_i(x, y) := \mu_i v_i(y) - v_i(x), \quad \tilde{w}'_i(x, y) := \mu'_i v_i(y) - v_i(x). \quad (12)$$

Then, each $\tilde{w}_i, \tilde{w}'_i$ is an eigenvector of B associated with μ_i, μ'_i , respectively.

Proof. It suffices to check $\tilde{w}_i, i \in [n]$, and the same argument works for \tilde{w}'_i . For any $(x, y) \in \vec{E}$, with (12), we are able to calculate

$$\begin{aligned} (B\tilde{w}_i)(x, y) &= \sum_{\substack{z:(y,z) \in \vec{E} \\ z \neq x}} (\mu_i v_i(z) - v_i(y)) \\ &= \mu_i [(Av_i)(y) - v_i(x)] - (d-1)v_i(y) \\ &= [\mu_i \lambda_i - (d-1)]v_i(y) - \mu_i v_i(x) \\ &= [\mu_i^2 + d - 1 - (d-1)]v_i(y) - \mu_i v_i(x) \\ &= \mu_i \tilde{w}_i(x, y), \end{aligned}$$

where in the fourth line $\mu_i \lambda_i$ is replaced by $\mu_i^2 + d - 1$ due to the quadratic equation (10). Then, we have $B\tilde{w}_i = \mu_i \tilde{w}_i$. When $|\lambda_i| \neq d$, $\mu_i \neq \pm 1$. We can check $\tilde{w}_i \neq 0$, which implies w_i is an eigenvalue of B with the corresponding eigenvalue μ_i . \square

We can derive a deterministic ℓ_∞ -norm bound for eigenvectors of B constructed from Lemma 6.1 as follows.

Lemma 6.2 (Deterministic ℓ_∞ -norm bound for eigenvectors of B). *Let B be the non-backtracking operator of a connected d -regular graph, and let \tilde{w}_i be eigenvectors of B associated with μ_i constructed in Lemma 6.1 where $\mu_i \notin \{1, d-1\}$. Then*

$$\frac{\|\tilde{w}_i\|_\infty}{\|\tilde{w}_i\|_2} \leq \frac{\|v_i\|_\infty(|\mu_i| + 1)}{\sqrt{d^2 - \lambda_i^2}}. \quad (13)$$

The same bound holds for \tilde{w}'_i .

Proof. It suffices to prove the estimate for \tilde{w}_i . From (12),

$$\|\tilde{w}_i\|_\infty \leq \sup_{(x,y) \in \vec{E}} |\mu_i v_i(y) - v_i(x)| \leq \|v_i\|_\infty(|\mu_i| + 1). \quad (14)$$

On the other hand, since μ_i, μ'_i is a conjugate pair from (10),

$$\begin{aligned} \|\tilde{w}_i\|_2^2 &= \sum_{(x,y) \in \vec{E}} (\mu_i v_i(y) - v_i(x))(\mu'_i v_i(y) - v_i(x)) \\ &= |\mu_i|^2 \sum_{(x,y) \in \vec{E}} (v_i(y))^2 + \sum_{(x,y) \in \vec{E}} (v_i(x))^2 - (\mu_i + \mu'_i) \sum_{(x,y) \in \vec{E}} v_i(y)v_i(x) \\ &= d|\mu_i|^2 \|v_i\|_2^2 + d\|v_i\|_2^2 - (\mu_i + \mu'_i) \sum_{(x,y) \in \vec{E}} v_i(y)v_i(x) \\ &= d^2 - \lambda_i \sum_{(x,y) \in \vec{E}} v_i(y)v_i(x). \end{aligned}$$

Moreover,

$$\sum_{(x,y) \in \vec{E}} v_i(y)v_i(x) = \sum_{x \in V} (Av_i)(x)v_i(x) = \lambda_i \sum_{x \in V} v_i^2(x) = \lambda_i.$$

This gives us

$$\|\tilde{w}_i\|_2^2 = d^2 - \lambda_i^2. \quad (15)$$

Since $\mu_i \notin \{1, d-1\}$, we have $\lambda_i \neq d$. Equations (14) and (15) imply (13). \square

Now, we are ready to prove Theorem 3.3.

Proof of Theorem 3.3. We consider eigenvectors of B associated with eigenvalues μ , $\mu_i \in \{1, d-1\}$ and $\mu_i \notin \{1, d-1\}$ separately.

Case 1: $\mu_i \in \{1, d-1\}$. The deterministic eigenvalue $\lambda_1 = d$ of A with eigenvector $v_1 = [1, \dots, 1]^\top \in \mathbb{R}^n$ yields two eigenvalues $\mu_1 = d-1, \mu'_1 = 1$. Let \tilde{w}_1 be the eigenvector given by (12) associated with μ_1 . One can directly check that

$$\frac{\|\tilde{w}_1\|_\infty}{\|\tilde{w}_1\|_2} = \frac{1}{\sqrt{nd}}.$$

Case 2: $\mu_i \notin \{1, d-1\}$. By the spectral gap bound from [12, Theorem 1], we have for some constants $C, C_1 > 0$, with probability at least $1 - n^{-C}$, for any eigenvalue μ of \tilde{B} with $\mu \neq d-1$,

$$|\mu| \leq \sqrt{d-1} + C_1 \left(\frac{\log \log n}{\log n} \right)^2,$$

and for any eigenvalue λ of A with $\lambda \neq d$,

$$|\lambda| \leq 2\sqrt{d-1} + C_1 \left(\frac{\log \log n}{\log n} \right)^2.$$

Therefore with (13), with probability at least $1 - n^{-C}$,

$$\frac{\|\tilde{w}_i\|_\infty}{\|\tilde{w}_i\|_2} \leq \frac{C'(\sqrt{d-1} + 1)}{d-2} \|v_i\|_\infty \leq \frac{2C'}{\sqrt{d}} \|v_i\|_\infty,$$

for an absolute constant $C' > 0$. The delocalization bound of v_i in Lemma 5.1 implies that with probability at least $1 - n^{-C_1}$,

$$\frac{\|\tilde{w}_i\|_\infty}{\|\tilde{w}_i\|_2} \leq \frac{\log^{C_2}(n)}{\sqrt{n}}.$$

With the two cases discussed above, taking ℓ_2 -normalized eigenvector $w_i = \frac{\tilde{w}_i}{\|\tilde{w}_i\|_2}$ for $i \in [n]$ completes the proof. \square

7. PROOF OF THEOREM 3.7

Similar to Lemma 4.1, we first establish a spectral relation between A and \tilde{B} for a regular hypergraph graph.

Lemma 7.1 (Spectral relation between A and \tilde{B}). *Let $H = (V, E)$ be a k -uniform, d -regular hypergraph with adjacency matrix A and reduced non-backtracking matrix \tilde{B} . Let λ_i be an eigenvalue of A . Then each λ_i corresponds to two eigenvalues μ_i, μ'_i of \tilde{B} , which satisfy the equation*

$$\mu^2 - (\lambda_i - k + 2)\mu + (d-1)(k-1) = 0, \quad (16)$$

and with the corresponding eigenvectors

$$u_i = \begin{bmatrix} v_i \\ \frac{\mu_i}{d-1} v_i \end{bmatrix}, \quad u'_i = \begin{bmatrix} v_i \\ \frac{\mu'_i}{d-1} v_i \end{bmatrix}. \quad (17)$$

Proof. Let $\begin{bmatrix} x \\ y \end{bmatrix}$, $x, y \in \mathbb{C}^n$ be an eigenvector of \tilde{B} with respect to an eigenvalue μ , we have

$$\mu \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 & (d-1)I \\ -(k-1)I & A - (k-2)I \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}.$$

This equation implies

$$Ax = \left(\mu + \frac{(k-1)(d-1)}{\mu} + k - 2 \right) x, \quad y = \frac{\mu}{d-1} x.$$

For each eigenvalue λ_i of A with a corresponding eigenvector $v_i \in \mathbb{R}^n$, there are two corresponding eigenvalue μ_i, μ'_i of \tilde{B} satisfying the quadratic equation

$$\mu^2 - (\lambda_i - k + 2)\mu + (d-1)(k-1) = 0.$$

This gives all the $2n$ eigenvalues of \tilde{B} . The rest of the proof follows in the same way as in Lemma 4.1. \square

Proof of Theorem 3.7. From Lemma 7.1, all eigenvalues of \tilde{B} satisfies (16). When

$$|\lambda_i - k + 2|^2 \leq 4(d-1)(k-1),$$

we have

$$\mu_i + \mu'_i = \lambda_i - k + 2, \quad |\mu_i| = |\mu'_i| = \sqrt{(d-1)(k-1)},$$

and the real parts of μ_i, μ'_i satisfy

$$x_i = x'_i = \frac{\lambda_i}{2}. \quad (18)$$

For different regimes of (d, k) , the limiting spectral distribution of the normalized adjacency matrix $\tilde{A} := \frac{A - (k-2)}{\sqrt{(d-1)(k-1)}}$ for random regular hypergraphs were obtained in [26, Corollary 6.11]:

- If d, k are fixed, the empirical spectral distribution of \tilde{A} converges in probability to a probability measure supported on $[-2, 2]$ whose density function is given by

$$f(x) = \frac{1 + \frac{k-1}{q}}{(1 + \frac{1}{q} - \frac{x}{\sqrt{q}})(1 + \frac{(k-1)^2}{q} + \frac{(k-1)x}{\sqrt{q}})} \frac{1}{\pi} \sqrt{1 - \frac{x^2}{4}},$$

where $q = (k-1)(d-1)$.

- For $d, k \rightarrow \infty$ with $\frac{d}{k} \rightarrow \alpha \geq 1$ and $d \leq \frac{n}{32}$, the empirical spectral distribution of \tilde{A} converges in probability to a measure supported on $[-2, 2]$ with a density function given by

$$g(x) = \frac{\alpha}{1 + \alpha + \sqrt{\alpha x}} \frac{1}{\pi} \sqrt{1 - \frac{x^2}{4}}.$$

- If $d \rightarrow \infty, d = o(n^\epsilon)$ for any $\epsilon > 0$ and $\frac{d}{k} \rightarrow \infty$, the the empirical spectral distribution of \tilde{A} converges to the semicircle law in probability.

With the relation (18), we can follow the same argument as in the proof of Theorem 3.1 to complete the proof of Theorem 3.7. \square

8. PROOF OF THEOREM 3.8

Proof of Theorem 3.8 (1). From (17), the corresponding unit eigenvectors u_i, u'_i satisfies

$$\|u_i\|_\infty, \|u'_i\|_\infty = \frac{\max\left\{1, \frac{|\mu_i|}{d-1}\right\}}{\sqrt{1 + \frac{|\mu_i|^2}{(d-1)^2}}} \|v_i\|_\infty \leq \|v_i\|_\infty.$$

This proves the first statement of Theorem 3.8. \square

We now consider the second statement of Theorem 3.8. The following lemma provides a relation between eigenvectors of B and A for regular hypergraphs.

Lemma 8.1 (Eigenvectors of B for regular hypergraphs). *Let \vec{E} be oriented hyperedge set for a (d, k) -regular hypergraph $H = (V, E)$. Let μ_i, μ'_i denote each eigenvalue pair of \tilde{B} corresponding to eigenvalue λ_i of the adjacency matrix A , through the quadratic equation (16). Let v_i be a unit eigenvector of A associated λ_i . Define $\tilde{w}_i, \tilde{w}'_i \in \mathbb{C}^{nd}$ such that for any $(x, e) \in \vec{E}$,*

$$\tilde{w}_i(x, e) := \mu_i \left(\sum_{y \in e, y \neq x} v_i(y) \right) - (k-1)v_i(x), \quad \tilde{w}'_i(x, e) := \mu'_i \left(\sum_{y \in e, y \neq x} v_i(y) \right) - (k-1)v_i(x). \quad (19)$$

Assume $\tilde{w}_i, \tilde{w}'_i \neq 0$. Then $\tilde{w}_i, \tilde{w}'_i$ is an eigenvector of B associated with μ_i, μ'_i , respectively.

Proof. It suffices to consider $\tilde{w}_i, i \in [n]$. With (19) and the definition of B for hypergraphs, we have for any $(x, e) \in \vec{E}$,

$$\begin{aligned}
(B\tilde{w}_i)(x, e) &= \sum_{(y, f) \in \vec{E}: y \in e, y \neq x, f \neq e} \tilde{w}_i(y, f) \\
&= \sum_{(y, f) \in \vec{E}: y \in e, y \neq x, f \neq e} \left(\mu_i \left(\sum_{z \in f, z \neq y} v_i(z) \right) - (k-1)v_i(y) \right) \\
&= \mu_i \left(\sum_{(y, f) \in \vec{E}: y \in e, y \neq x, f \neq e} \sum_{z \in f, z \neq y} v_i(z) \right) - (d-1)(k-1) \sum_{y \in e, y \neq x} v_i(y) \\
&= \mu_i \left(\sum_{y \in e, y \neq x} (Av_i)(y) - \sum_{y \in e, y \neq x} \sum_{z \in e, z \neq y} v_i(z) \right) - (d-1)(k-1) \sum_{y \in e, y \neq x} v_i(y) \\
&= \mu_i \left(\sum_{y \in e, y \neq x} \lambda_i v_i(y) - \left(\sum_{y \in e, y \neq x} (k-2)v_i(y) \right) - (k-1)v_i(x) \right) - (d-1)(k-1) \sum_{y \in e, y \neq x} v_i(y) \\
&= (\mu_i \lambda_i - (k-2)\mu_i - (d-1)(k-1)) \left(\sum_{y \in e, y \neq x} v_i(y) \right) - \mu_i(k-1)v_i(x) \\
&= \mu_i^2 \left(\sum_{y \in e, y \neq x} v_i(y) \right) - \mu_i(k-1)v_i(x) \tag{20} \\
&= \mu_i \left(\mu_i \sum_{y \in e, y \neq x} v_i(y) - (k-1)v_i(x) \right) = \mu_i \tilde{w}_i(x, e),
\end{aligned}$$

where in (20), we use the relation between μ_i, λ_i in (16). This implies \tilde{w}_i is an eigenvector of B associated with μ_i . \square

Lemma 8.1 can be used to prove the following deterministic ℓ_∞ -norm bound for eigenvectors. When $k = 2$, Lemma 8.2 reduces to Lemma 6.2.

Lemma 8.2 (Deterministic ℓ_∞ -norm bound for eigenvectors of B). *Let B be the non-backtracking operator of a (d, k) -regular hypergraph. Let \tilde{w}_i be the eigenvector of B associated with μ_i constructed in Lemma 8.1 with $\mu_i \notin \{1, (d-1)(k-1)\}$ and λ_i be the corresponding eigenvalue of A . Then*

$$\frac{\|\tilde{w}_i\|_\infty}{\|\tilde{w}_i\|_2} \leq \frac{\sqrt{k-1}(|\mu_i| + 1)}{\sqrt{(d + \lambda_i)(d(k-1) - \lambda_i)}} \|v_i\|_\infty. \tag{21}$$

The same bound holds for \tilde{w}'_i .

Proof. It suffices to prove the estimate for \tilde{w}_i . From (19),

$$\|\tilde{w}_i\|_\infty \leq \sup_{(x, e) \in \vec{E}} \left| \mu_i \left(\sum_{y \in e, y \neq x} v_i(y) \right) - (k-1)v_i(x) \right| \leq \|v_i\|_\infty \left(|\mu_i|(k-1) + k-1 \right). \tag{22}$$

On the other hand, since μ_i, μ'_i is a conjugate pair in (16),

$$\begin{aligned}
\|\tilde{w}_i\|_2^2 &= \sum_{(x,e) \in \vec{E}} \left(\mu_i \left(\sum_{y \in e, y \neq x} v_i(y) \right) - (k-1)v_i(x) \right) \left(\mu'_i \left(\sum_{y \in e, y \neq x} v_i(y) \right) - (k-1)v_i(x) \right) \\
&= |\mu_i|^2 \sum_{(x,e) \in \vec{E}} \left(\sum_{y \in e, y \neq x} v_i(y) \right)^2 + (k-1)^2 \sum_{(x,e) \in \vec{E}} v_i(x)^2 \\
&\quad - (\mu_i + \mu'_i)(k-1) \sum_{(x,e) \in \vec{E}} \left(\sum_{y \in e, y \neq x} v_i(y) \right) v_i(x) \\
&= (d-1)(k-1) \sum_{(x,e) \in \vec{E}} \left(\sum_{y \in e, y \neq x} v_i(y) \right)^2 + (k-1)^2 \sum_{(x,e) \in \vec{E}} v_i(x)^2 \\
&\quad - (\lambda_i - k + 2) \sum_{(x,e) \in \vec{E}} \left(\sum_{y \in e, y \neq x} v_i(y) \right) v_i(x). \tag{23}
\end{aligned}$$

For the first term in (23),

$$\begin{aligned}
&\sum_{(x,e) \in \vec{E}} \left(\sum_{y \in e, y \neq x} v_i(y) \right)^2 \\
&= \sum_{(x,e) \in \vec{E}} \left(\sum_{y \in e} v_i(y) \right)^2 - 2 \sum_{(x,e) \in \vec{E}} \left(\sum_{y \in e} v_i(y) \right) v_i(x) + \sum_{(x,e) \in \vec{E}} v_i(x)^2. \tag{24}
\end{aligned}$$

Using the fact that v_i is a unit vector, we have

$$\sum_{(x,e) \in \vec{E}} v_i(x)^2 = d.$$

We then calculate the first and second terms separately in the following:

$$\begin{aligned}
\sum_{(x,e) \in \vec{E}} \left(\sum_{y \in e} v_i(y) \right)^2 &= k \sum_{y \in V} v_i(y) \left(\sum_{x \in V} A_{yx} v_i(x) \right) + kd \sum_{y \in V} v_i(y)^2 \\
&= k \sum_{y \in V} v_i(y) \lambda_i v_i(y) + kd = k\lambda_i + kd, \tag{25}
\end{aligned}$$

and

$$\begin{aligned}
\sum_{(x,e) \in \vec{E}} \left(\sum_{y \in e} v_i(y) \right) v_i(x) &= \sum_{(x,e) \in \vec{E}} \left(\sum_{y \in e, y \neq x} v_i(y) \right) v_i(x) + \sum_{(x,e) \in \vec{E}} v_i(x)^2 \\
&= \sum_{x \in V} \lambda_i v_i(x)^2 + d = \lambda_i + d. \tag{26}
\end{aligned}$$

Together, we can use (25) and (26) to get the value for (24), that is

$$\sum_{(x,e) \in \vec{E}} \left(\sum_{y \in e, y \neq x} v_i(y) \right)^2 = (k-2)\lambda_i + (k-1)d.$$

Also, we have from (26),

$$\sum_{(x,e) \in \tilde{E}} \left(\sum_{y \in e, y \neq x} v_i(y) \right) v_i(x) = \lambda_i.$$

From (23), these identities give us

$$\|\tilde{w}_i\|_2^2 = (k-1)(d + \lambda_i)(d(k-1) - \lambda_i). \quad (27)$$

When $\mu_i \notin \{1, (d-1)(k-1)\}$, we have $\lambda_i \notin \{d(k-1), -d\}$ from (16). Hence, \tilde{w}_i is a nonzero vector. Equations (22) and (27) imply (21). \square

With Lemma 8.2, we are ready to prove the second statement of Theorem 3.8.

Proof of Theorem 3.8 (2). Let A be the adjacency matrix of a (d, k) -random regular hypergraph. The largest eigenvalue of A is $d(k-1)$ with an eigenvector $v_1 = \frac{1}{\sqrt{n}}[1, \dots, 1]^\top$. By Lemma 8.2, We consider when $\mu_i \in \{1, (d-1)(k-1)\}$ and $\mu_i \notin \{1, (d-1)(k-1)\}$ separately.

Case 1: When $\mu_i \in \{1, (d-1)(k-1)\}$. $\lambda_1 = d(k-1)$ as a deterministic eigenvalue for A yields two eigenvalues for \tilde{B} : $\mu_1 = (d-1)(k-1)$ and $\mu'_1 = 1$. Let \tilde{w}_1 be the eigenvector given by (19) associated with μ_1 . One can directly check that

$$\frac{\|\tilde{w}_1\|_\infty}{\|\tilde{w}_1\|_2} = \frac{1}{\sqrt{nd}} = \frac{1}{\sqrt{d}} \|v_i\|_\infty.$$

Case 2: When $\mu_i \notin \{1, (d-1)(k-1)\}$. By [26, Theorem 11], when $d \geq k \geq 3$, we have any eigenvalues $\lambda \neq d(k-1)$ of A satisfies

$$-2\sqrt{(k-1)(d-1)} + k - 2 - \varepsilon_n \leq \lambda \leq 2\sqrt{(k-1)(d-1)} + k - 2 + \varepsilon_n, \quad (28)$$

asymptotically almost surely with $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$. With (28) and (27), we have, with high probability,

$$\|\tilde{w}_i\|_2^2 \geq (k-1)(\sqrt{d-1} - \sqrt{k-1})^2(\sqrt{(k-1)(d-1)} + 1)^2 + o(1). \quad (29)$$

By the spectral gap bounded for any eigenvalues $\mu \neq (d-1)(k-1)$ of B from [26, Theorem 5.5], we have

$$|\mu| \leq \sqrt{(k-1)(d-1)} + o(1) \quad (30)$$

with high probability. With (29) and (30), we have, for $d > k \geq 3$, with high probability,

$$\begin{aligned} \frac{\|\tilde{w}_i\|_\infty}{\|\tilde{w}_i\|_2} &\leq \frac{(\sqrt{(k-1)(d-1)} + 1)\sqrt{k-1} + o(1)}{(\sqrt{d-1} - \sqrt{k-1})(\sqrt{(k-1)(d-1)} + 1)} \|v_i\|_\infty \\ &\leq \frac{\sqrt{k-1} + o(1)}{\sqrt{d-1} - \sqrt{k-1}} \|v_i\|_\infty. \end{aligned} \quad (31)$$

Inequality (31) also holds when $\mu_i \in \{1, (d-1)(k-1)\}$. Thus, from the two cases above, Theorem 3.8 (2) holds. \square

9. PROOF OF THEOREM 3.10

We need the following spectral gap estimate for the adjacency matrix of an RSBM from [17].

Lemma 9.1. *Let A be the adjacency matrix of an (n, d_1, d_2) -regular stochastic block model. Then with high probability, $(d_1 + d_2)$, $(d_1 - d_2)$ are two eigenvalues of A with multiplicity one. Moreover, for any $\varepsilon > 0$, all eigenvalues $\lambda \notin \{d_1 + d_2, d_1 - d_2\}$ satisfy*

$$\lim_{n \rightarrow \infty} \mathbb{P} \left(|\lambda| \leq 2\sqrt{d_1 + d_2 - 1} + \varepsilon \right) = 0.$$

Proof. This is shown in [16, Section 4] by using the result of the spectral gap of random lifts from [12, Corollary 24], and contiguity of the permutation model and the configuration model of regular graphs from [32, Theorem 1.3]. \square

With Lemma 9.1, we prove Theorem 3.10 by using the spectral relation between A and \tilde{B} established in Lemma 4.1.

Proof of Theorem 3.10. Since an (n, d_1, d_2) -regular stochastic block model is a $(d_1 + d_2)$ -regular graph, the first statement follows from Lemma 4.1 directly.

For the second statement, let A be the adjacency matrix of an (n, d_1, d_2) regular stochastic block model. Since $d_1 - d_2$ is an eigenvalue for A with associated eigenvector σ , from Lemma 4.1, there are two eigenvalues of \tilde{B} satisfying

$$\mu^2 - (d_1 - d_2)\mu + (d_1 + d_2 - 1) = 0.$$

When $(d_1 - d_2)^2 > 4(d_1 + d_2 - 1)$, the two eigenvalues are real, and the second statement holds.

We now show the third statement. From Lemma 9.1, since $d_1 + d_2, d_1 - d_2$ are two eigenvalues of A with multiplicity one with high probability, Lemma 4.1 shows that $d_1 + d_2 - 1, 1, \mu_2, \mu'_2$ are four real eigenvalues of \tilde{B} with multiplicity one with high probability when $(d_1 - d_2)^2 > 4(d_1 + d_2 - 1)$. Let λ be an eigenvalue of A with $\lambda \notin \{d_1 + d_2, d_1 - d_2\}$. With Lemma 4.1, $|\lambda| \leq 2\sqrt{d_1 + d_2 - 1} + \varepsilon$ with high probability, and the corresponding two eigenvalues μ, μ' of \tilde{B} satisfies

$$x^2 - \lambda x + (d_1 + d_2 - 1) = 0.$$

We consider two cases:

- (1) If $|\lambda| \leq 2\sqrt{d_1 + d_2 - 1}$, then the associated two eigenvalues have $|\mu| = |\mu'| = \sqrt{d_1 + d_2 - 1}$.
- (2) If $|\lambda| \in [2\sqrt{d_1 + d_2 - 1}, 2\sqrt{d_1 + d_2 - 1} + \varepsilon]$, μ, μ' are real and

$$\mu, \mu' = \frac{\lambda \pm \sqrt{\lambda^2 - 4(d_1 + d_2 - 1)}}{2}.$$

We have for any $\varepsilon \in (0, 1)$, with high probability,

$$|\mu - \sqrt{d_1 + d_2 - 1}|, |\mu' - \sqrt{d_1 + d_2 - 1}| \leq C\sqrt{\varepsilon}$$

for a constant C depending only on $(d_1 + d_2 - 1)$.

Therefore, all eigenvalues $\lambda \notin \{d_1 + d_2 - 1, 1, \mu_2, \mu'_2\}$ are within $o(1)$ distance from the circle of radius $\sqrt{d_1 + d_2 - 1}$. This completes the proof of Theorem 3.10. \square

10. ERDŐS-RÉNYI GRAPHS

Studying bulk non-backtracking eigenvectors for Erdős-Rényi graphs $G(n, p)$ is an interesting open problem. Previously, using the fact that when $np = \omega(\log n)$ the degrees in $G(n, p)$ are concentrated around np , [56, 22] studied the eigenvalues of \tilde{B} by approximating the degree matrix D with $(np)I_n$, and analyzed a perturbed quadratic equation from the Ihara-Bass formula. However, it is not clear to us how to generalize this partial de-randomization argument introduced in [56] to study eigenvector delocalization. We conjecture that non-backtracking eigenvector delocalization holds when $np = \omega(\log n)$ and does not hold for $np = o(\log n)$, similar to the results established for the adjacency matrix [55, 25, 34, 3, 2, 4]. See Figures 6 for simulation results.




FIGURE 6. Simulation for Erdős-Rényi graphs. The upper graph is a simulation for $np = \log^2 n$ with $n = 300$; the lower graph is a simulation for $np = 3$ with $n = 2000$.

Acknowledgments. Y.Z. thanks Paolo Barucca, Ludovic Stephan, and Ke Wang for helpful discussions. X.Z. was supported by an AMS Undergraduate Travel Grant to present this work at Joint Mathematics Meetings 2024. Y.Z. was partially supported by the NSF-Simons Research Collaborations on the Mathematical and Scientific Foundations of Deep Learning and an AMS-Simons Travel Grant. Part of this work was completed during Y.Z.’s visit to Université Paris Cité, for which Y.Z. thanks Simon Coste for his warm hospitality.

REFERENCES

- [1] Emmanuel Abbe. Community detection and stochastic block models: Recent developments. *Journal of Machine Learning Research*, 18(177):1–86, 2018.
- [2] Johannes Alt, Raphael Ducatez, and Antti Knowles. Delocalization transition for critical Erdős-Rényi graphs. *Communications in Mathematical Physics*, 388(1):507–579, 2021.
- [3] Johannes Alt, Raphaël Ducatez, and Antti Knowles. The completely delocalized region of the Erdős-Rényi graph. *Electronic Communications in Probability*, 27:1–9, 2022.
- [4] Johannes Alt, Raphael Ducatez, and Antti Knowles. Poisson statistics and localization at the spectral edge of sparse erdős-rényi graphs. *The Annals of Probability*, 51(1):277–358, 2023.
- [5] Omer Angel, Joel Friedman, and Shlomo Hoory. The non-backtracking spectrum of the universal cover of a graph. *Transactions of the American Mathematical Society*, 367(6):4287–4318, 2015.
- [6] Maria Chiara Angelini, Francesco Caltagirone, Florent Krzakala, and Lenka Zdeborová. Spectral detection on sparse hypergraphs. In *2015 53rd Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, pages 66–73. IEEE, 2015.
- [7] Paolo Barucca. Spectral partitioning in equitable graphs. *Physical Review E*, 95(6):062310, 2017.
- [8] Hyman Bass. The Ihara-Selberg Zeta function of a tree lattice. *International Journal of Mathematics*, 03(06):717–797, December 1992.
- [9] Roland Bauerschmidt, Jiaoyang Huang, and Horng-Tzer Yau. Local Kesten-McKay law for random regular graphs. *Communications in Mathematical Physics*, 369:523–636, 2019.
- [10] Roland Bauerschmidt, Antti Knowles, and Horng-Tzer Yau. Local semicircle law for random regular graphs. *Communications on Pure and Applied Mathematics*, 70(10):1898–1960, 2017.
- [11] Florent Benaych-Georges, Charles Bordenave, and Antti Knowles. Spectral radii of sparse random matrices. *Annales de l’Institut Henri Poincaré, Probabilités et Statistiques*, 56(3):2141–2161, 2020.

- [12] Charles Bordenave. A new proof of Friedman’s second eigenvalue theorem and its extension to random lifts. *Annales Scientifiques de l’École Normale Supérieure*, 4(6):1393–1439, 2020.
- [13] Charles Bordenave and Djailil Chafai. Around the circular law. *Probability Surveys*, 9:1–89, 2012.
- [14] Charles Bordenave, Simon Coste, and Raj Rao Nadakuditi. Detection thresholds in very sparse matrix completion. *Foundations of Computational Mathematics*, 23(5):1619–1743, 2023.
- [15] Charles Bordenave, Marc Lelarge, and Laurent Massoulié. Nonbacktracking spectrum of random graphs: Community detection and nonregular ramanujan graphs. *Annals of probability*, 46(1):1–71, 2018.
- [16] Gerandy Brito, Ioana Dumitriu, Shirshendu Ganguly, Christopher Hoffman, and Linh V Tran. Recovery and rigidity in a regular stochastic block model. *arXiv preprint arXiv:1507.00930*, 2015.
- [17] Gerandy Brito, Ioana Dumitriu, Shirshendu Ganguly, Christopher Hoffman, and Linh V Tran. Recovery and rigidity in a regular stochastic block model. In *Proceedings of the twenty-seventh annual ACM-SIAM symposium on Discrete algorithms*, pages 1589–1601. Society for Industrial and Applied Mathematics, 2016.
- [18] Gerandy Brito, Ioana Dumitriu, and Kameron Decker Harris. Spectral gap in random bipartite biregular graphs and applications. *Combinatorics, Probability and Computing*, 31(2):229–267, 2022.
- [19] Evan Chang, Neel Kolhe, and Youngtak Sohn. Upper bounds on the 2-colorability threshold of random d -regular k -uniform hypergraphs for $k \geq 3$. *arXiv preprint arXiv:2308.02075*, 2023.
- [20] Philip Chodrow, Nicole Eikmeier, and Jamie Haddock. Nonbacktracking spectral clustering of nonuniform hypergraphs. *SIAM Journal on Mathematics of Data Science*, 5(2):251–279, 2023.
- [21] Colin Cooper, Alan Frieze, Michael Molloy, and Bruce Reed. Perfect matchings in random r -regular, s -uniform hypergraphs. *Combinatorics, Probability and Computing*, 5(1):1–14, 1996.
- [22] Simon Coste and Yizhe Zhu. Eigenvalues of the non-backtracking operator detached from the bulk. *Random Matrices: Theory and Applications*, 10(03):2150028, 2021.
- [23] Lorenzo Dall’Amico, Romain Couillet, and Nicolas Tremblay. Revisiting the Bethe-Hessian: Improved community detection in sparse heterogeneous graphs. *Advances in neural information processing systems*, 32, 2019.
- [24] Ioana Dumitriu and Soumik Pal. Sparse regular random graphs: Spectral density and eigenvectors. *Annals of Probability*, 40(5):2197–2235, 2012.
- [25] Ioana Dumitriu and Yizhe Zhu. Sparse general Wigner-type matrices: Local law and eigenvector delocalization. *Journal of Mathematical Physics*, 60(2), 2019.
- [26] Ioana Dumitriu and Yizhe Zhu. Spectra of random regular hypergraphs. *Electronic Journal of Combinatorics*, 28(3):P3.36, 2021.
- [27] Ioana Dumitriu and Yizhe Zhu. Extreme singular values of inhomogeneous sparse random rectangular matrices. *arXiv preprint arXiv:2209.12271*, 2022.
- [28] Keqin Feng and Wen-Ching Winnie Li. Spectra of hypergraphs and applications. *Journal of number theory*, 60(1):1–22, 1996.
- [29] Joel Friedman and David-Emmanuel Kohler. The relativized second eigenvalue conjecture of alon. *arXiv preprint arXiv:1403.3462*, 2014.
- [30] Cory Glover and Mark Kempton. Some spectral properties of the non-backtracking matrix of a graph. *Linear Algebra and its Applications*, 618:37–57, 2021.
- [31] Catherine Greenhill, Mikhail Isaev, and Gary Liang. Spanning trees in random regular uniform hypergraphs. *Combinatorics, Probability and Computing*, 31(1):29–53, 2022.
- [32] Catherine Greenhill, Svante Janson, Jeong Han Kim, and Nicholas C Wormald. Permutation pseudographs and contiguity. *Combinatorics, Probability and Computing*, 11(3):273–298, 2002.
- [33] Yukun He. Edge universality of sparse Erdős-Rényi digraphs. *arXiv preprint arXiv:2304.04723*, 2023.
- [34] Yukun He, Antti Knowles, and Matteo Marozzi. Local law and complete eigenvector delocalization for supercritical Erdős-Rényi graphs. *Annals of Probability*, 47(5):3278–3302, 2019.
- [35] Jiaoyang Huang and Horng-Tzer Yau. Spectrum of random d -regular graphs up to the edge. *arXiv preprint arXiv:2102.00963*, 2021.
- [36] Jürgen Jost, Raffaella Mulas, and Leo Torres. Spectral theory of the non-backtracking laplacian for graphs. *Discrete Mathematics*, 346(10):113536, 2023.
- [37] Mark Kempton. Non-backtracking random walks and a weighted Ihara’s theorem. *Open Journal of Discrete Mathematics*, 6(4):207–226, 2016.
- [38] Florent Krzakala, Cristopher Moore, Elchanan Mossel, Joe Neeman, Allan Sly, Lenka Zdeborová, and Pan Zhang. Spectral redemption in clustering sparse networks. *Proceedings of the National Academy of Sciences*, 110(52):20935–20940, 2013.
- [39] Wen-Ching Winnie Li. Ramanujan hypergraphs. *Geometric & Functional Analysis GFA*, 14(2):380–399, 2004.
- [40] Alexander Litvak, Anna Lytova, Konstantin Tikhomirov, Nicole Tomczak-Jaegermann, and Pierre Youssef. Structure of eigenvectors of random regular digraphs. *Transactions of the American Mathematical Society*, 371(11):8097–8172, 2019.

- [41] Eyal Lubetzky and Yuval Peres. Cutoff on all Ramanujan graphs. *Geometric and Functional Analysis*, 26(4):1190–1216, 2016.
- [42] Brendan D. McKay. The expected eigenvalue distribution of a large regular graph. *Linear Algebra and its Applications*, 40:203–216, 1981.
- [43] Elchanan Mossel, Joe Neeman, and Allan Sly. Reconstruction and estimation in the planted partition model. *Probability Theory and Related Fields*, 162(3-4):431–461, 2015.
- [44] Raffaella Mulas, Dong Zhang, and Giulio Zucal. There is no going back: Properties of the non-backtracking laplacian. *Linear Algebra and its Applications*, 680:341–370, 2024.
- [45] MEJ Newman and Travis Martin. Equitable random graphs. *Physical Review E*, 90(5):052824, 2014.
- [46] Subhabrata Sen. Optimization on sparse random hypergraphs and spin glasses. *Random Structures & Algorithms*, 53(3):504–536, 2018.
- [47] Patrick Solé. Spectra of regular graphs and hypergraphs and orthogonal polynomials. *European Journal of Combinatorics*, 17(5):461–477, 1996.
- [48] Ludovic Stephan and Laurent Massoulié. Robustness of spectral methods for community detection. In *Conference on Learning Theory*, pages 2831–2860. PMLR, 2019.
- [49] Ludovic Stephan and Laurent Massoulié. Non-backtracking spectra of weighted inhomogeneous random graphs. *Mathematical Statistics and Learning*, 5(3):201–271, 2022.
- [50] Ludovic Stephan and Yizhe Zhu. Sparse random hypergraphs: Non-backtracking spectra and community detection. In *2022 IEEE 63rd Annual Symposium on Foundations of Computer Science (FOCS)*, pages 567–575. IEEE, 2022.
- [51] Ludovic Stephan and Yizhe Zhu. A non-backtracking method for long matrix and tensor completion. *arXiv preprint arXiv:2304.02077*, 2023.
- [52] Christopher K Storm. The Zeta function of a hypergraph. *The Electronic Journal of Combinatorics*, pages R84–R84, 2006.
- [53] Audrey Terras. *Zeta functions of graphs: A stroll through the garden*, volume 128. Cambridge University Press, 2010.
- [54] Leo Torres, Kevin S Chan, Hanghang Tong, and Tina Eliassi-Rad. Nonbacktracking eigenvalues under node removal: X-centrality and targeted immunization. *SIAM Journal on Mathematics of Data Science*, 3(2):656–675, 2021.
- [55] Linh V Tran, Van H Vu, and Ke Wang. Sparse random graphs: Eigenvalues and eigenvectors. *Random Structures & Algorithms*, 42(1):110–134, 2013.
- [56] Ke Wang and Philip Matchett Wood. Limiting empirical spectral distribution for the non-backtracking matrix of an erdős-rényi random graph. *Combinatorics, Probability and Computing*, 32(6):956–973, 2023.
- [57] Kevin Yang. Bulk eigenvalue correlation statistics of random biregular bipartite graphs. *arXiv preprint arXiv:1705.00083*, 2017.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA IRVINE, IRVINE, CA 92697
 Email address: `xiangz27@uci.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA IRVINE, IRVINE, CA 92697
 Email address: `yizhe.zhu@uci.edu`