NORM CONVERGENCE OF CONFINED FERMIONIC SYSTEMS AT ZERO TEMPERATURE

ESTEBAN CÁRDENAS

ABSTRACT. The semi-classical limit of ground states of large systems of fermions was studied by Fournais, Lewin and Solovej in (Calc. Var. Partial Differ. Equ., 2018). In particular, the authors prove weak convergence towards classical states associated to the minimizers of the Thomas-Fermi functional. In this paper, we revisit this limit and show that under additional assumptions—and, using simple arguments—it is possible to prove that strong convergence holds in relevant normed spaces.

1. INTRODUCTION

In this article, we study a system of N identical fermions that move in Euclidean space \mathbb{R}^d , where $d \ge 1$ is the spatial dimension. We assume that the particles interact through a two-body potential V(x - y), and move under the action of an external trap U(x). Neglecting spin variables, the Hamiltonian of the system takes the form (in appropriate units)

$$H_N \equiv \sum_{i=1}^N \hbar^2(-\Delta_{x_i}) + \sum_{i=1}^N U(x_i) + \lambda \sum_{i< j} V(x_i - x_j) .$$
(1.1)

Here, $\hbar > 0$ plays the role of Planck's constant, and λ is the interaction strength. The Hilbert space of the system corresponds to the subspace $\mathfrak{h}_N \equiv L^2_a(\mathbb{R}^{dN})$ of functions in L^2 , that are antisymmetric with respect to the permutation of their variables. Namely

$$L^2_a(\mathbb{R}^{dN}) = \{\Psi \in L^2(\mathbb{R}^{dN}) : \Psi(x_{\sigma(1)}, \dots, x_{\sigma(N)}) = \operatorname{sgn} \sigma \Psi(x_1, \dots, x_N) \ \forall \sigma \in S_N\} , \quad (1.2)$$

where S_N stands for the permutation group of N elements.

In this work, we are interested in the case in which the external potential U(x) acts as a trap, and focus on the associated ground state problem. Indeed, our focus here will be on the ground state energy, for large number of particles $N \ge 1$, and on which the scales of the system are semi-classical, and the interaction is of mean-field type. In other words, we consider

$$E(N) \equiv \inf \sigma(H_N)$$
 for $\hbar \equiv \frac{1}{N^{1/d}}$ and $\lambda \equiv \frac{1}{N}$. (1.3)

In this situation, the Fermi momentum is of $\mathcal{O}(1)$ and, consequently, the energy per particle is $\mathcal{O}(1)$ as $N \to \infty$. Thus, it makes mathematical sense to analyze the asymptotics of the ratio E(N)/N, describing the average energy per particle in the system.

1.1. Thomas-Fermi theory. Heuristically, in the large N limit one is able to introduce a mean-field description by means of the Thomas-Fermi energy [8, 23]. In this theory, one replaces the N-particle wave function $\Psi_N \in \mathfrak{h}_N$ in the energy functional, with its average over the positions

$$\rho_{\Psi_N}(x) \equiv \int_{\mathbb{R}^{d(N-1)}} |\Psi_N(x, x_1, \cdots, x_{N-1})|^2 \mathrm{d}x_1 \cdots \mathrm{d}x_{N-1} , \qquad x \in \mathbb{R}^d .$$
(1.4)

In this approximation, the leading order term of E(N)/N is then expected to be described in terms of the *Thomas-Fermi functional*

$$\mathcal{E}(\rho) \equiv \frac{d}{d+2} C_{TF} \int_{\mathbb{R}^d} \rho(x)^{1+\frac{2}{d}} \mathrm{d}x + \int_{\mathbb{R}^d} U(x)\rho(x) \mathrm{d}x + \frac{1}{2} \int_{\mathbb{R}^{2d}} \rho(x)V(x-y)\rho(y) \mathrm{d}x \mathrm{d}y \quad (1.5)$$

where $\rho(x)$ denotes the number density of the particles at the point $x \in \mathbb{R}^d$, and $C_{TF} = 4\pi^2 (d/|S_{d-1}|)^{2/d}$ stands for the Thomas-Fermi constant. The first term in (1.5) corresponds to the kinetic energy of the system after taking into account Pauli's Exclusion Principle, and is sometimes referred to as the quantum pressure; the second and third terms on the other hand correspond to the one- and two-body potential energies, respectively. The variational problem now reads

$$e_{TF} \equiv \inf \left\{ \mathcal{E}(\rho) : 0 \leqslant \rho \in L^1 \cap L^{1+2/d}(\mathbb{R}^d), \|\rho\|_{L^1} = 1 \right\} .$$

$$(1.6)$$

The analysis of the relationship between E(N)/N and e_{TF} has been the focus of extensive research. In particular, it has been proven under general assumptions on the interaction potentials that

$$\lim_{N \to \infty} \frac{E(N)}{N} = e_{TF} . \tag{1.7}$$

The first rigorous derivation of (1.7) goes back to the works of Lieb and Simon [17, 18] for Coulomb systems, and has been recently revisited by Fournais, Lewin and Solovej [9, Theorem 1.1] for more general models.

On the other hand, one may also study the problem of convergence of the spatial distributions of the particles, as $N \to \infty$. In other words, here one is interested in proving the following convergence statement for the position densities

$$\lim_{N \to \infty} \rho_{\Psi_N} = \rho_{TF} , \qquad (1.8)$$

where Ψ_N is extracted from the many-body Hamiltonian, and ρ_{TF} is the minimizer of e_{TF} . The first result in this direction was again proven by Lieb and Simon [18, Theorem III.3] for Coulomb systems, where convergence holds in the weak- L^1 sense. Since then, convergence has been further improved and understood. For instance, more recently, the authors in [9] have proven convergence in the weak- $L^{1+d/2}$ sense, for a significantly larger class of potentials. In a similar spirit, Gottschling and Nam [12] have proven convergence of the energy functionals in the sense of Gamma-convergence. See also the work of Nguyen [19], where convergence is understood in the context of Weyl's semiclassical law, starting from the Hartree-Fock functional.

1.2. Convergence of states. While the analysis of the limit of the spatial distribution ρ_{Ψ_N} has received considerable attention in the literature in the last few decades, the question of convergence of *states* has only recently started to be the focus of mathematical research. Note that the interest in the former arises in the context of Density Functional Theory and its applications to quantum chemistry, where one is interested in determining the *static* electronic structure of many-body ground states.

Let us further explain and, at the same time, fix the notation that we use in the rest of this paper. Namely, let us consider a sequence of normalized states $\Psi_N \in \mathfrak{h}_N$ that satisfies

$$\langle \Psi_N, H_N, \Psi_N \rangle_{\mathfrak{h}_N} = E(N) (1 + o(1)) \qquad N \to \infty .$$
 (1.9)

We shall refer to Ψ_N as an approximate ground state. We define its one-particle reduced density matrix as the trace-class operator on $L^2(\mathbb{R}^d)$ with kernel

$$\gamma_{\Psi_N}(x, x') \equiv N \int_{\mathbb{R}^{d(N-1)}} \Psi_N(x, x_1, \dots, x_{N-1}) \overline{\Psi_N(x', x_1, \dots, x_{N-1})} \, \mathrm{d}x_1 \cdots \mathrm{d}x_{N-1} \quad , \quad (1.10)$$

for $(x, x') \in \mathbb{R}^{2d}$. In particular, $\operatorname{Tr}\gamma_{\Psi_N} = N$ and $0 \leq \gamma_{\Psi_N} \leq 1$ due to Fermi statistics. Because all particles are identical, the operator γ_{Ψ_N} contains all the "one-particle" information of the system, i.e. expectation values of one-particle observables can be written in terms of traces over γ_{Ψ_N} . Furthermore, for weakly interacting systems, higher-order reduced density matrices (containing information about particle correlations) can be expected to be written in terms of γ_{Ψ_N} plus an error that vanishes in the limit $N \to \infty$.¹ Thus, it is natural to study the limit of γ_{Ψ_N} for large N in order to understand the physical behaviour of the system.

Since our scaling regime is of semi-classical type $\hbar = N^{-1/d}$, we shall analyze the asymptotics of γ_{Ψ_N} by looking at its *Wigner function*

$$f_{\Psi_N}(x,p) \equiv \int_{\mathbb{R}^d} \gamma_{\Psi_N}\left(x + \frac{y}{2}, x - \frac{y}{2}\right) e^{-i\frac{y\cdot p}{\hbar}} \mathrm{d}y , \qquad (1.11)$$

where $(x, p) \in \mathbb{R}^{2d}$ should be regarded as varying over macroscopic scales. With the present definition, it follows from $\hbar^d N = 1$ that the following normalization holds

$$\frac{1}{(2\pi)^d} \int_{\mathbb{R}^{2d}} f_{\Psi_N}(x, p) \mathrm{d}x \mathrm{d}p = 1 .$$
 (1.12)

While the previous equation suggests that one could regard $(2\pi)^{-d} f_{\Psi_N}(x,p)$ as a classical state (i.e. a probability measure on \mathbb{R}^{2d}), it is well-known that in general it is not positive, and may take on negative values. On the other hand, its associated *Husimi measure*, is a well-defined measure on phase space. Here, we adopt the following definition

$$m_{\Psi_N} \equiv f_{\Psi_N} * \mathscr{G}_\hbar \tag{1.13}$$

¹This approximation can be justified for instance in Hartree-Fock theory, where Ψ_N is an N-particle quasi-free state, i.e. a Slater determinant. Higher-order particle distributions are written in terms of γ_{Ψ_N} thanks to Wick's theorem.

where $\mathscr{G}_{\hbar}(z) \equiv \hbar^{-2d} \mathscr{G}_{1}(z/\hbar)$ is a mollifier with Gaussian profile $\mathscr{G}_{1} \in L^{1}(\mathbb{R}^{2d})$, at scale $\hbar > 0$. In particular, the following holds for the Husimi measure

$$0 \leqslant m_{\Psi_N}(x,p) \leqslant 1 \qquad \text{and} \qquad \frac{1}{(2\pi)^d} \int_{\mathbb{R}^{2d}} m_{\Psi_N}(x,p) \mathrm{d}x \mathrm{d}p = 1 , \qquad (1.14)$$

see for reference [9, Section 2]. The first inequality is nothing but the Pauli Exclusion Principle, whereas the second identity is a normalization condition. Let us remark here that, at least formally, these two functions have the same limit: $\lim_{N\to\infty} m_{\Psi_N} = \lim_{N\to\infty} f_{\Psi_N}$. Hence, it is convenient to study them simultaneously.

The question that one now may ask is what the asymptotics of f_{Ψ_N} is, for large N. In order to motivate this limit, we observe that one may re-write the energy functional E(N)in terms of reduced density matrices, and then in terms of the Wigner function. In the limit, one formally obtains the Vlasov energy

$$\mathcal{V}(f) \equiv \frac{1}{(2\pi)^d} \int_{\mathbb{R}^{2d}} p^2 f(x, p) \mathrm{d}x \mathrm{d}p + \int_{\mathbb{R}^d} \rho_f(x) U(x) \mathrm{d}x + \frac{1}{2} \int_{\mathbb{R}^{2d}} \rho_f(x) V(x - y) \rho_f(y) \mathrm{d}x \mathrm{d}y ,$$
(1.15)

where $\rho_f(x) \equiv (2\pi)^{-d} \int_{\mathbb{R}^d} f(x, p) dp$ is the associated position density of f. The minimization is carried out over all functions in L^1 , complying with the Pauli Exclusion Principle $0 \leq f \leq 1$. The connection between the Vlasov and Thomas-Fermi energies can be understood as follows. Let $\rho \in L^1(\mathbb{R}^d)$ be a position density and define the classical state

$$f_{\rho}(x,p) \equiv \mathbb{1}(|p|^2 \leqslant C_{TF}\rho(x)^{2/d}), \qquad (x,p) \in \mathbb{R}^{2d}.$$
 (1.16)

Then, there holds $\mathcal{V}(f_{\rho}) = \mathcal{E}(\rho)$. Note that (1.16) is nothing but a local version of a Fermi-Dirac distribution at zero temperature: at position $x \in \mathbb{R}^d$ all the electrons momenta accomodate in order to fill a ball of radius $p_F(x) \sim \rho(x)^{1/d}$. Letting ρ_{TF} be the minimizer of e_{TF} , we can then expect the relevant classical state to be $f_{TF} \equiv f_{\rho_{TF}}$.

With the above notations, we are now ready to state the main question of interest in this work. Namely:

Assuming that Ψ_N is an approximate ground state of H_N , in what sense can we prove that its Wigner function f_{Ψ_N} converges to the state f_{TF} ?

1.3. Our contribution. Recently, Fournais, Lewin and Solovej [9, Theorem 1.2] have studied this problem in great generality. In our context, their most notable result is the fact that convergence holds in a weak sense, even without requiring the uniqueness of the minimizers. More precisely, it is shown that there exits a probability measure \mathscr{P} over the set \mathcal{M} of all minimizers of the Thomas-Fermi functional, such that the following limit holds (up to the extraction of a subsequence)

$$\int_{\mathbb{R}^{2d}} f_{\Psi_N}(x,p)\varphi(x,p)\mathrm{d}x\mathrm{d}p \longrightarrow \int_{\mathcal{M}} \left(\int_{\mathbb{R}^{2d}} f_{TF}(x,p)\varphi(x,p)\mathrm{d}x\mathrm{d}p\right)\mathrm{d}\mathscr{P}(\rho)$$
(1.17)

for all test functions φ with bounded first derivatives. Their main ingredient is a fermionic version of the de Finetti–Hewitt–Savage theorem for classical measures–see their Theorem 2.7. Let us note that their results also include convergence for Husimi measures (in a stronger mode of convergence), as well as for higher-order density matrices, and additional analysis in the unconfined case (i.e where some particles may be lost in the limit).

The work of [9] motivated later studies on the semi-classical limit of similar fermionic systems. For instance, the case of *positive* temperature has been considered by Lewin, Madsen and Triay in [14]. Most notably, strong convergence in L^1 is proven for Husimi measures, and analogous weak convergence results are proven for Wigner functions and higher-order density matrices. Here, the additional assumption of minimizers being unique is made. Three-dimensional systems in the presence of a homogeneous magnetic fields were studied by Fournais and Madsen [22] where various scalings for the strength of the magnetic field are considered. Girardot and Rougerie [11] have also analyzed the semi-classical limit of fermionic anyons. Here, analogous results on the weak convergence of states is proven for k-particle Husimi measures. Finally, let us mention that the study of convergence of states of large quantum-mechanical systems in the style presented above was first analyzed for bosons. See e.g. the work of Lewin, Nam and Rougerie [15].

The main contribution of this article is to revisit the convergence (1.17) assuming zero magnetic fields and $\hat{V} \ge 0$ (in particular, minimizers of e_{TF} are unique). We summarize our main results as follows.

- (1) In Theorem 1 we prove that the Husimi measure converges strongly in L^p for all $p \in [1, \infty)$, and the Wigner function converges strongly with respect to certain Fourierbased norms, which include the negative Sobolev space H^s for every s < 0. The latter Fourier-based norms have been previously considered in the derivation of the Vlasov equation from many-body systems (see e.g. [4, Theorem 2.5]), as well as in the analysis of the space homogeneous Boltzmann equation [24].
- (2) While we are not able to prove in full generality that f_{Ψ_N} converges strongly to f_{TF} in L^p spaces, in Theorem 2 we provide additional conditions under which convergence holds. Most notably, we show that L^2 convergence holds if the sequence Ψ_N consists of Slater determinants (i.e. Hartree-Fock states). Additionally, we prove that whenever the system has uniform moments in phase-space, then strong convergence in L^p holds for additional values of p < 2. This condition on the moments is verified for the harmonic trap $U(x) = x^2$.
- (3) In Theorem 3 we investigate the convergence of higher-order distribution functions, first established by the authors in [9] in a weak sense. Here, we show that L^p convergence can be extended to the Husimi functions to all $p \in [1, \infty)$. For Slater determinants, we show that the Wigner functions converge strongly in H^s for all s < 0, but not in L^2 .

The reader may wonder if upgrading the mode of convergence has value beyond pure mathematical interest. It turns out this question has its motivation in the *quantitative* derivation of effective equations for large systems of interacting fermions. Indeed, here one considers an externally prepared system of initial data, that converges with respect to some norm, as $N \to \infty$. Subsequently, one wishes to prove (and, quantify) that convergence is propagated at later times along the solutions of the equations under consideration; see e.g [3, 4, 5, 10] and the references therein. In this context, our main result provides

examples of initial data for the derivation of effective equations, which converges with respect to appropriate norms. Most importantly, these examples stem from states at zero temperature and include the orthogonal projections (i.e. the Slater determinants). In particular, one must be careful with the choice of norm in this situation, since regularity of the limiting function is not abundant-the functions f_{ρ} introduced in (1.16) are only of bounded variation.

2. Main results

In this section, we state the main results of this article. Namely, Theorem 1 and 2. First, we fix the assumptions on the potentials that we work with, as well as fixing the notation to be used throughout this paper.

2.1. Assumptions, definitions and notations. We will work with the following set of assumptions. These are taken from [9] up to the following modifications. First, we take $A \equiv 0$. Second, we assume $\hat{V} \ge 0$.

Assumption 1. $U : \mathbb{R}^d \to \mathbb{R}$ and $V : \mathbb{R}^d \to \mathbb{R}$ satisfy the following conditions.

- (1) $U_{-} \in L^{1+d/2}(\mathbb{R}^d) + L^{\infty}_{\varepsilon}(\mathbb{R}^d)$ and $U_{+} \in L^{1}_{\text{loc}}(\mathbb{R}^d)$, $\lim_{|x|\to\infty} U_{+}(x) = +\infty$. (2) V is even, $V \in L^{1+d/2}(\mathbb{R}^d) + L^{\infty}_{\varepsilon}(\mathbb{R}^d)$, and $\hat{V} \ge 0$.

Remark 2.1. Given $p, q \in [1, \infty]$, we write $U \in L^p + L^q_{\varepsilon}$ if the following is satisfied: for all $\varepsilon > 0$ there exists $U_1 \in L^p$ and $U_2 \in L^q$ such that $U = U_1 + U_2$ and $||U_2||_{L^q} \leq \varepsilon$.

Remark 2.2. Under these conditions, there always exists a minimizer of the Thomas-Fermi functional. In addition, thanks to $V \ge 0$, the functional $\mathcal{E}(\rho)$ is strictly convex in ρ . Hence, the minimizer is unique and is denoted by ρ_{TF} .

Remark 2.3. The additional conditions A = 0 and $\hat{V} \ge 0$ can be relaxed. Namely, as long as the Thomas-Fermi functional has a unique minimizer, then Theorem 1 and 2 still hold. These have only been included here for notational convenience and simplicity of the exposition. The only result that makes explicit use of these additional conditions is Lemma 4.1 (see Remark 2.9 below), although we believe the result of the lemma is still true for potentials A and \hat{V} verifying the more general assumptions considered in [9].

States. Let $(\Psi_N)_{N \ge 1}$ be a sequence of approximate ground states satisfying (1.9). We let γ_{Ψ_N} be its one-particle reduced density matrix, f_{Ψ_N} its Wigner function and m_{Ψ_N} its Husimi measure. In order to simplify the notation, here and in the sequel we write for all $N \ge 2$

 $f_N \equiv f_{\Psi_N}$, $m_N \equiv m_{\Psi_N}$, $\gamma_N \equiv \gamma_{\Psi_N}$, and $f \equiv f_{\rho_{TF}}$, (2.1)where ρ_{TF} is the unique minimizer of e_{TF} and f is defined through (1.16).

Weak convergence. Let us denote $\langle \varphi, g \rangle = \int_{\mathbb{R}^{2d}} \varphi(z) g(z) dz$. Then, with the above notation, it follows from [9, Theorem 1.2] and uniqueness of the Thomas-Fermi minimizer that

$$\lim_{N \to \infty} \langle \varphi, f_N \rangle = \langle \varphi, f \rangle \tag{2.2}$$

for all test functions φ with φ , $\nabla \varphi \in L^{\infty}(\mathbb{R}^{2d})$. On the other hand, for the Husimi measures

$$\lim_{N \to \infty} \langle \varphi, m_N \rangle = \langle \varphi, f \rangle \tag{2.3}$$

for all test functions $\varphi \in L^1(\mathbb{R}^{2d}) + L^{\infty}(\mathbb{R}^{2d})$. Let us note that the original result in [9] includes extraction of a subsequence via a compactness argument. Since the limit here is unique and independent of the subsequence, convergence holds for the whole sequence.

Wigner transform. We denote by $W^{\hbar}: L^2_{x,x'}(\mathbb{R}^{2d}) \to L^2_{x,p}(\mathbb{R}^{2d})$ the linear map

$$W^{\hbar}[\gamma](x,p) \equiv \int_{\mathbb{R}^d} \gamma \left(x - \frac{y}{2}, x + \frac{y}{2} \right) e^{-\frac{ip \cdot y}{\hbar}} \mathrm{d}y \ , \qquad (x,p) \in \mathbb{R}^{2d}$$
(2.4)

which we refer to as the Wigner transform. We abuse notation and identify Hilbert-Schmidt operators with their L^2 kernels. In particular, $f_N = W^{\hbar}[\gamma_N]$. The map W^{\hbar} is an L^2 -isomorphism, and the normalization is chosen so that $\|W^{\hbar}[\gamma]\|_{L^2} = (2\pi\hbar)^{d/2} \|\gamma\|_{L^2}$. See e.g [7, Proposition 13].

Fourier-based norms. Let $n \in \mathbb{N}$. Throughout this article, we let $\langle \zeta \rangle \equiv (1 + \zeta^2)^{\frac{1}{2}}$ be the Japanese bracket, and $\hat{g}(\zeta) = (2\pi)^{-n/2} \int_{\mathbb{R}^n} e^{-i\zeta \cdot z} g(z) dz$ the Fourier transform of g.

Definition 1. Let $s \in \mathbb{R}$ and $q \in [1, \infty]$. We denote by $H_{s,q}(\mathbb{R}^n)$ the space of tempered distributions $g \in \mathcal{S}'(\mathbb{R}^n)$ whose Fourier transform is regular $\hat{g} \in L^1_{loc}(\mathbb{R}^n)$, and for which the norm

$$\|g\|_{s,q} \equiv \|\langle \zeta \rangle^s \, \hat{g}\|_{L^q(\mathbb{R}^n)} \tag{2.5}$$

is finite.

We will use these spaces only in the case of negative order s < 0.

Remark 2.4. A few comments are in order regarding the Fourier-based spaces $H_{s,q}$.

- (1) For all s < 0 there holds $|g|_{s,2} \leq ||g||_{L^2}$ and $|g|_{s,\infty} \leq ||g||_{L^1}$.
- (2) For q = 2, $H_{s,2} = H^s$ is the standard Sobolev space of order $s \in \mathbb{R}$.
- (3) For $q = \infty$, the spaces $H_{s,\infty}$ are quite useful in the study of fermionic systems. Namely, for the Fourier transform of f_N the following formula holds, sometimes known as *Groenewold's formula*:

$$\hat{f}_N(\zeta) = \frac{1}{N} \operatorname{Tr} \left[\mathcal{O}_{\xi,\eta} \gamma_N \right], \qquad \zeta = (\xi,\eta) \in \mathbb{R}^{2d}.$$
 (2.6)

Here, $\mathcal{O}_{\xi,\eta} \equiv \exp(i\xi \cdot \hat{x} + i\eta \cdot \hat{p})$ is a semi-classical observable, with \hat{x} and $\hat{p} \equiv -i\hbar\nabla_x$ the standard position and momentum observables on $L^2(\mathbb{R}^d)$. In particular, while the L^1 norm of f_N cannot be controlled by trace norms of γ_N , it follows easily from (2.6) that

$$|\hat{f}_N(\zeta)| \leq \frac{1}{N} \|\mathcal{O}_{\xi,\eta}\|_{B(L^2)} \operatorname{Tr}\gamma_N \leq 1 , \qquad \zeta = (\xi,\eta) \in \mathbb{R}^{2d} .$$

$$(2.7)$$

Consequently, $|f|_{s,\infty} \leq 1$ for all $s \leq 0$. These uniform bounds will replace the possible lack of L^1 boundedness in our analysis.

Slater determinants. We say that $\Phi \in \mathfrak{h}_N$ is a Slater determinant if there exists an orthonormal set $(\varphi_i)_{i=1}^N \subset L^2(\mathbb{R}^d)$ such that

$$\Phi(x_1,\ldots,x_N) = \frac{1}{\sqrt{N!}} \det_{1 \le i,j \le N} \left[\varphi_i(x_j) \right], \qquad (x_1,\ldots,x_N) \in \mathbb{R}^{dN}, \tag{2.8}$$

and we write $\Phi = \varphi_1 \wedge \cdots \wedge \varphi_N$. In particular, Φ is a Slater determinant if and only if its one-particle reduced density matrix is an orthogonal projection, i.e. if $\gamma_N^2 = \gamma_N$.

2.2. Main results. We are now ready to state our main result, which is the content of the next theorem. Here, we make no additional assumptions on the sequence Ψ_N under consideration. The proof, given in Section 3, uses only elementary L^p inequalities and contains the main idea of the paper. Namely, that weak convergence can be improved to strong convergence, using the fact that the limiting function solves the equation $f^2 = f$.

Theorem 1 (Norm convergence). Let f_N , m_N and f be as in (2.1), and let Assumption 1 hold. Then, the following statements are true.

- (1) The sequence m_N converges to f strongly in $L^p(\mathbb{R}^{2d})$ for all $p \in [1, \infty)$.
- (2) The sequence f_N converges to f strongly in $H_{s,q}(\mathbb{R}^{2d})$ for all $q \in [2,\infty]$ and s < 0.

Remark 2.5 (k-particle functions). Originally, the authors in [9] proved that (2.2) and (2.3) hold for higher-order Wigner and Husimi functions. Hence, the reader may wonder if the content of Theorem 1 extend to these functions. The answer to this question is rather subtle and is explained further in Section 5.

Remark 2.6 (Positive temperature). Note that the equation $f = f^2$ holds only at zero temperature, because f is a characteristic function of the phase-space. In particular, the proof of Theorem 1 does not apply directly to the positive temperature setting [14].

Remark 2.7 (Position densities). Let ρ_{TF} be the unique minimizer of the Thomas-Fermi functional e_{TF} . Then, Theorem 1 implies that

$$\int_{\mathbb{R}^d} m_N(\cdot, p) \mathrm{d}p \to \rho_{TF} \qquad \text{strongly in } L^1(\mathbb{R}^d) \ . \tag{2.9}$$

On the other hand, for the position density $\rho_N(x) = \int_{\mathbb{R}^d} f_N(x, p) dp$ defined initially in terms of Ψ_N as (1.4), we have

$$\rho_N \to \rho_{TF} \quad \text{strongly in } H_{s,\infty}(\mathbb{R}^d)$$
(2.10)

for all s < 0. This follows from $\hat{\rho}_N(\xi) = \hat{f}_N(\xi, 0)$ and $\langle \xi \rangle^{-s} \leq \langle \zeta \rangle^{-s}$ for $\zeta = (\xi, \eta) \in \mathbb{R}^{2d}$.

A natural question is if the Wigner function f_N converges to f in any L^p space. We are not able to prove this statement in the most general case, but we are able to show convergence under additional assumptions on the sequence Ψ_N . The first result in this direction establishes L^p convergence whenever the sequence f_N is known to verify a uniform *smoothness* assumption. We record this in the following corollary. **Corollary 2.1.** Under the same conditions of Theorem 1, assume additionally that there exist r > 0, $p \in [1, \infty)$ and $q \in [1, \infty]$ such that

$$\lim_{N \to \infty} \left\| \frac{\|f_N - f_N(\bullet + \hbar z)\|_{L^p}}{|z|^r} \right\|_{L^q(dz)} = 0 , \qquad (2.11)$$

then f_N converges to f strongly in $L^p(\mathbb{R}^{2d})$.

Remark 2.8. The condition given in (2.11) can be interpreted as smoothness of the sequence that is *uniform* with respect to the number of particles N. For illustration, we consider two situations have been considered in the literature.

(1) Assume that $\sup_{N \ge 1} \|\nabla f_N\|_{L^1} < \infty$. The Taylor formula $f(z+\hbar z')-f(z) = \hbar z' \int_0^1 \nabla f(z+t\hbar z') dt$ and a change of variables $z \mapsto z - t\hbar z'$ implies for r = 1, p = 1 and $q = \infty$

$$\sup_{z' \in \mathbb{R}^{2d}} \frac{1}{|z'|} \int_{\mathbb{R}^{2d}} |f_N(z + \hbar z') - f_N(z)| \mathrm{d}z \leqslant \hbar \|\nabla f_N\|_{L^1},$$
(2.12)

which yields L^1 convergence. The condition $\sup_N \|\nabla f_N\|_{L^1} < \infty$ has been previously considered in the derivation of mean-field dynamics for fermionic systems, see e.g [4, Theorem 2.5] and the remark after the theorem.

(2) Given $z_0 = (x_0, p_0) \in \mathbb{R}^{2d}$, it is possible to verify the following relation between translations in phase-space distributions, and on quantum density matrices

$$\|f_N - f_N(\bullet + \hbar z_0)\|_{L^2} = (2\pi\hbar)^{d/2} \|[\mathcal{O}_{p_0, -x_0}, \gamma_N]\|_{HS} .$$
(2.13)

In particular, if γ_N satisfies the commutator estimates

$$\|[\mathcal{O}_{p,x},\gamma_N]\|_{HS}^2 \leqslant CN\hbar|z| , \qquad \forall z = (x,p) \in \mathbb{R}^{2d}$$
(2.14)

then f_N satisfies (2.11) for p = 2, $q = \infty$ and r = 1/2, and yields L^2 convergence. Estimates of the form (2.14) (as well as their stronger *trace-class* variants) arise in practice for initial data in the derivation of effective dynamics for fermion systems [3, 4, 5, 10], and have been shown to be satisfied in a few special cases [2, 21].

In practice, verifying the smoothness condition (2.11) is a challenging task. Our next result brings an alternative. Namely, if one assumes that Ψ_N is a Slater determinant, then L^2 convergence is immediate. Further, we show that uniform control on the *moments* of the system yields L^p convergence for smaller values of p.

Theorem 2 (Slater determinants). Under the same conditions of Theorem 1, assume additionally that Ψ_N is a Slater determinant for all $N \ge 1$. Then, the following holds.

- (1) The sequence f_N converges to f strongly in $L^2(\mathbb{R}^{2d})$.
- (2) (Moments) If there exists m > 0 such that

$$\sup_{N \ge 1} \| (|x| + |p|)^m f_N \|_{L^2} < \infty,$$
(2.15)

then f_N converges to f strongly in $L^p(\mathbb{R}^{2d})$ for all $p \in [1,2] \cap (\frac{2}{1+2m/d},2]$.

Remark 2.9 (The harmonic trap). For the external potential $U(x) = x^2$ we verify in Section 4 that the bounds on the moments (2.15) hold for m = 1, under Assumption 1. In particular, this yields

$$f_N \to f$$
 strongly in $L^p(\mathbb{R}^{2d})$ for $\begin{cases} p \in [1,2] & \text{if } d = 1 \ , \\ p \in (1,2] & \text{if } d = 2 \ , \\ p \in (6/5,2) & \text{if } d = 3 \ , \end{cases}$ (2.16)

for Slater determinants. We would like to note that for d = 3 our assumptions include systems that interact through repulsive Coulomb potentials $V(x) = \lambda |x|^{-1}$ with $\lambda > 0$.

Remark 2.10. The L^2 convergence result does not apply to the higher-order Wigner functions $f_N^{(k)}$ (see (5.1) for a definition). In particular, we prove in Section 5 that for Slater determinants

$$\liminf_{N \to \infty} \|f_N^{(k)} - f^{\otimes k}\|_{L^2} \ge (2\pi)^{\frac{dk}{2}} \left(\sqrt{k!} - 1\right) \qquad \forall k \ge 2$$

$$(2.17)$$

even though convergence $f_N^{(k)} \to f^{\otimes k}$ holds in $H^s(\mathbb{R}^{2dk})$ for all s < 0.

The last of our main results is the following interesting corollary. Here, we derive an equivalent formulation of convergence in L^p norm for Slater determinants.

Corollary 2.2. Assume that Ψ_N is a Slater determinant for all $N \ge 1$, and let $p \in [1, 2]$. Then, f_N converges to f in $L^p(\mathbb{R}^{2d})$ if and only if

$$\lim_{N \to \infty} \|f_N\|_{L^p} = (2\pi)^{d/p} .$$
(2.18)

3. Proof of the main results

In this section we prove our main results, Theorem 1 and Theorem 2.

3.1. **Proof of Theorem 1.** The heart of the proof of Theorem 1 has as a starting point the modes of convergences (2.2) and (2.3), proven in [9]. We establish strong convergence by making heavy use of the identity $f^2 = f$, which follows from the fact that f is a characteristic function. In particular, it holds that f(1-f) = 0, i.e. we regard f and 1-f as orthogonal projections.

It turns out that the proof can be formulated using rather general arguments, and is independent of the fact that f_N is extracted from an N-body quantum system. For transparency, we prove the following abstract lemma which may be of interest in its own right. To the authors best knowledge, this result (as well as its application to fermion systems) is new. Essentially, it states that an appropriate notion of weak convergence towards a characteristic function can always be upgraded to strong convergence in L^p .

Lemma 3.1. Let (X, \mathscr{F}, μ) be a measure space. Consider a sequence $(F_n)_{n=1}^{\infty}$ of nonnegative functions F_n in $L^1(X) \cap L^{\infty}(X)$ that satisfies

$$0 \leqslant F_n \leqslant 1 \ \mu\text{-a.e}$$
 and $\limsup_{N \to \infty} \int_X F_n d\mu = 1$. (3.1)

Let $F \in L^1(X) \cap L^{\infty}(X)$ be a non-negative function satisfying

$$0 \leqslant F \leqslant 1 \ \mu\text{-a.e}$$
 and $\int_X F d\mu = 1$, (3.2)

and assume that for all $\phi \in L^1(X) + L^{\infty}(X)$

$$\lim_{n \to \infty} \langle \phi, F_n \rangle = \langle \phi, F \rangle .$$
(3.3)

Then, if $F = F^2 \mu$ -a.e., there holds

$$\lim_{n \to \infty} \|F_n - F\|_{L^p} = 0 \qquad \forall p \in [1, \infty) .$$

$$(3.4)$$

Proof of Lemma 3.1. First, we prove the case p = 2. Then, we prove the case p = 1. Every other case $p \in (1, \infty)$ then follows from the p = 1 case, and uniform L^{∞} bounds.

Let us prove the p = 2 case. Since the L^2 norm is generated by an inner product, we find that

$$||F_n - F||_{L^2}^2 = ||F||_{L^2}^2 + ||F_n||_{L^2}^2 - 2\langle F, F_n \rangle .$$
(3.5)

For the first term in the right hand side of (3.5), we may use $F = F^2$ and calculate $||F||_{L^2} = 1$. For the second term, we use the upper bound $||F_n||_{L^2} \leq ||F_n||_{L^\infty}^{1/2} ||F_n||_{L^1}^{1/2}$. Hence, $\limsup_{n\to\infty} ||F_n||_{L^2} \leq 1$. For the third term, we note that since the limiting function F is in L^1 we may use the test function $\phi = F$ and conclude that

$$\lim_{N \to \infty} \langle F, F_n \rangle = \langle F, F \rangle = 1 .$$
(3.6)

Thus, we put everything together to find that

$$\limsup_{N \to \infty} \|F_n - F\|_{L^2}^2 \leqslant 2 - 2 \lim_{N \to \infty} \langle F_n, F \rangle = 0 .$$
(3.7)

This finishes the proof of the p = 2 case.

Let us prove the p = 1 case. In view of the identity F(1-F) = 0 and the non-negativity of F_n and 1 - F we may consider the following decomposition

$$\|F_n - F\|_{L^1} = \|F(F - F_n)\|_{L^1} + \|(1 - F)F_n\|_{L^1} \leq \|F\|_{L^2}\|F - F_n\|_{L^2} + \langle 1 - F, F_n \rangle .$$
(3.8)

The first term in the right hand side of (3.8) is controlled by the difference in L^2 norm, analyzed above. For the second term, we use again weak convergence with the test function $\phi = 1 - F \in L^{\infty}$. It suffices now to take the $n \to \infty$ limit, which finishes the p = 1 case.

Let us prove the $p \in (1, \infty)$ case. For all such p, we use the fact that $||F_n - F||_{L^{\infty}} \leq 2$ to find that

$$\|F_n - F\|_{L^p}^p = \int |F_n - F|^p d\mu = \int |F_n - F| |F_n - F|^{p-1} d\mu \leqslant 2^{p-1} \|F - F_n\|_{L^1} .$$
(3.9)

Because of the p = 1 case, the right hand side now vanishes in the limit $n \to \infty$. This finishes the proof of the lemma.

Let us now turn to the proof of our main result, Theorem 1. The proof of its first part is a direct application of Lemma 3.1. The proof of the second part analyzes the difference between f_N and m_N .

Proof of Theorem 1. Let m_N , f_N and f be as in the statement of the theorem.

(1) It suffices to use Lemma 3.1 with $X = \mathbb{R}^{2d}$, \mathscr{F} the Borel sets and μ the 2*d*dimensional Lebesgue measure. We consider the functions $F_N(z) = m_N((2\pi)^{1/2}z)$ and $F(z) = f((2\pi)^{1/2}z)$. In particular, (3.1) and (3.2) are readily verified after a change of variables thanks to (1.14) and (1.16). The weak convergence follows from (2.2) and (2.3), proven in [9].

(2) Let $|g|_{s,q}$ be the norm introduced in (2.5). First, we prove the $q = \infty$ case. Second, we prove the q = 2 case. Every other $q \in (2, \infty)$ then follows by interpolation.

The $q = \infty$ case. Let s < 0. We use the triangle inequality together with the elementary bound $|\cdot|_{s,\infty} \leq ||\cdot||_{L^1}$ to find that

$$|f_N - f|_{s,\infty} \leqslant |f_N - m_N|_{s,\infty} + ||m_N - f||_{L^1}$$
(3.10)

The second term in (3.10) converges to zero, in view of part (1) of the Theorem. Thus, it suffices to analyze the first term in (3.10). In particular, we consider only the case $|s| \in (0, 1]$, since otherwise we can use $|f_N - m_N|_{s,\infty} \leq |f_N - m_N|_{1,\infty}$ in (3.10). To this end, we look at its Fourier transform at $\zeta \in \mathbb{R}^{2d}$. Namely, let $|g|_{C^{0,\alpha}} = \sup_{\zeta_1,\zeta_2 \in \mathbb{R}^{2d}} |g(\zeta_1) - g(\zeta_2)| |\zeta_1 - \zeta_2|^{-\alpha}$ the Hölder norm of a function $g : \mathbb{R}^{2d} \to \mathbb{C}$. Then, we find

$$|(\hat{f}_N - \hat{m}_N)(\zeta)| = |\hat{f}_N(\zeta)| |1 - \hat{\mathscr{G}}_1(\hbar\zeta)| \leqslant |\hat{f}_N(\zeta)| |\hat{\mathscr{G}}_1|_{C^{0,|s|}} \hbar^{|s|} |\zeta|^{|s|} \leqslant |\hat{\mathscr{G}}_1|_{C^{0,|s|}} \hbar^{|s|} |\zeta|^{|s|}$$
(3.11)

where we used $\mathscr{G}_1(0) = 1$, together with the uniform bound $\|f_N\|_{L^{\infty}} \leq 1$ (see (2.7)). The last inequality now implies that

$$|f_N - m_N|_{s,\infty} = \sup_{\zeta \in \mathbb{R}^{2d}} \langle \zeta \rangle^s \left| (\hat{f}_N - \hat{m}_N)(\zeta) \right| \leqslant |\hat{\mathscr{G}}_1|_{C^{0,s}} \hbar^{|s|}.$$
(3.12)

This finishes the proof thanks to the fact that $|\hat{\mathscr{G}}_1|_{C^{0,|s|}} < \infty$ for $|s| \in (0, 1]$.

The q = 2 case. The proof is identical to the proof of (2), with the following changes. First, one replaces $||m_N - f||_{L^1}$ with $||m_N - f||_{L^2}$ and uses part (1) for p = 2. Second, one replaces the uniform bound $||\hat{f}_N||_{L^{\infty}} \leq 1$ with the alternative uniform L^2 -bound $||\hat{f}_N||_{L^2} \leq (2\pi)^{\frac{d}{2}}$. The latter bounds follows from the fact that (up to scaling) the Wigner transformation is a unitary map between $L^2_{x,x'}$ and $L^2_{x,p}$. More precisely, in our setting we obtain

$$\|f_N\|_{L^2} = (2\pi\hbar)^{\frac{d}{2}} \|\gamma_N\|_{L^2} = (2\pi\hbar)^{\frac{d}{2}} \|\gamma_N\|_{HS} \leqslant (2\pi\hbar)^{\frac{d}{2}} \|\gamma_N\|_{\mathrm{Tr}}^{\frac{1}{2}} = (2\pi)^{\frac{d}{2}}.$$
 (3.13)

In the last two inequalities we used the fact that $\|\gamma\|_{HS} = (\text{Tr}\gamma^*\gamma)^{\frac{1}{2}} \leq \|\gamma\|_{B(L^2)}^{\frac{1}{2}} \|\gamma\|_{\text{Tr}}^{\frac{1}{2}}$ for arbitrary trace-class operators on L^2 , the fact that for fermions $0 \leq \gamma \leq 1$, and the scaling $\text{Tr}\gamma = N = \hbar^{-d}$.

Proof of Corollary 2.1. Let p, q and r be as in the statement of the corollary. Then, we use the triangle inequality to find that

$$||f_N - f||_{L^p} \leq ||f_N - m_N||_{L^p} + ||m_N - f||_{L^p} .$$
(3.14)

Thanks to Theorem 1, $\lim_{N\to\infty} ||m_N - f||_{L^p} = 0$. Thus, we only look at the first term. We write for $z = (x, p) \in \mathbb{R}^{2d}$

$$f_N(z) - m_N(z) = \int_{\mathbb{R}^{2d}} \left(f_N(z) - f_N(z - \hbar z') \right) \mathscr{G}_1(z') dz' .$$
 (3.15)

Hence, Minkowski's and Hölder's inequality implies

$$\|f_{N} - m_{N}\|_{L^{p}} \leq \int_{\mathbb{R}^{2d}} \|f_{N} - f_{N}(\bullet + \hbar z')\|_{L^{p}} \mathscr{G}_{1}(z') dz'$$

$$= \int_{\mathbb{R}^{2d}} \frac{\|f_{N} - f_{N}(\bullet + \hbar z')\|_{L^{p}}}{|z'|^{r}} |z'|^{r} \mathscr{G}_{1}(z') dz'$$

$$\leq \left\|\frac{\|f_{N} - f_{N}(\bullet + \hbar z')\|_{L^{p}}}{|z'|^{r}}\right\|_{L^{q}(dz')} \||z'|^{r} \mathscr{G}_{1}\|_{L^{q'}(dz')}$$
(3.16)

where 1/q' = 1 - 1/q is the dual exponent of q. Note that $|||z'|^r \mathscr{G}_1||_{L^{q'}(dz')} < \infty$ because \mathscr{G}_1 is a Schwartz-class function. In view of the assumption of the theorem, it suffices to take the limit $N \to \infty$.

3.2. **Proof of Theorem 2.** Recall that in order to prove that f_N converges to f in a Banach space E, it suffices to show that from all subsequences $(f_{N_k})_{k\geq 1}$ we can extract a further subsequence converging to f in E. We shall be using this fact in the proof of Theorem 2 part (1) with $E = L^2$ and Corollary 2.2 with $E = L^1$, but make no explicit reference to it in order to simplify the exposition.

Proof of Theorem 2. Let γ_N , f_N and f be as in the statement of the theorem.

(1) When γ_N is a rank-*N* orthogonal projection, its Hilbert-Schmidt norm can be calculated to be $\|\gamma_N\|_{HS} = (\text{Tr}\gamma_N^*\gamma_N)^{\frac{1}{2}} = (\text{Tr}\gamma_N)^{\frac{1}{2}} = N^{\frac{1}{2}}$. Thus, the same reasoning that led to (3.13) now yields

$$\|f_N\|_{L^2} = (2\pi)^{\frac{d}{2}} . aga{3.17}$$

Thus, we may assume that f_N converges weakly in L^2 . Of course, in view of (2.2), this limit is given by f. Further, $f = f^2$ implies that $||f||_{L^2} = (2\pi)^{d/2}$. This shows that $\lim_{N\to\infty} ||f_N||_{L^2} = ||f||_{L^2}$. Since L^2 is a Uniformly Convex Space, we may conclude that convergence is strong in L^2 , and this finishes the proof of the first part of the theorem.

(2) Let m > 0 and $p \in [1, 2] \cap (\frac{2}{1+2m/d}, 2]$ be as in the statement of the theorem, and let us denote by

$$K_m \equiv \sup_{N \ge 1} \| \langle z \rangle^m f_N \|_{L^2} < \infty .$$
(3.18)

Let R > 0 and denote $B_R \equiv \{z \in \mathbb{R}^{2d} : |z| \leq R\}$ and $B_R^c \equiv \mathbb{R}^{2d}/B_R$. We let $\chi_R \equiv \mathbb{1}_{B_R}$ and $\chi_R^c \equiv 1 - \chi_R$.

The proof starts with the decomposition

$$\|f - f_N\|_{L^p} \le \|\chi_R(f - f_N)\|_{L^p} + \|\chi_R^c(f - f_N)\|_{L^p} .$$
(3.19)

We estimate each term separately. The first term can be estimated in terms of the L^2 norm, using Hölder's inequality

$$\|\chi_R(f - f_N)\|_{L^p} \leqslant |B_R|^{\frac{2-p}{2p}} \|f - f_N\|_{L^2} .$$
(3.20)

Note that thanks to the first part of the theorem, the right hand side vanishes in the limit $N \to \infty$. For the second term of (3.19), we use the triangle inequality

$$\|\chi_R^c(f - f_N)\|_{L^p} \le \|\chi_R^c f\|_{L^p} + \|\chi_R^c f_N\|_{L^p} .$$
(3.21)

Note that $f \in L^p$ and so the first term converges to zero as $R \to \infty$. In order to control the second one, we use the moments of f_N . Namely, Hölder's inequality gives

$$|\chi_R^c f_N||_{L^p} = ||\chi_R^c \langle z \rangle^{-m} \langle z \rangle^m f_N||_{L^p} \leqslant ||\chi_R^c \langle z \rangle^{-m} ||_{L^{r'}} ||\langle z \rangle^m f_N||_{L^r}$$
(3.22)

where 1/p = 1/r + 1/r'. We choose r = 2 and r' = 2p/(2-p). In particular, we observe that $\langle z \rangle^{-m} \in L^{r'}$. Indeed, this is equivalent to mr' > d, which holds for $p > 2/(1 + \frac{2m}{d})$.

To finish the proof, we take the $\limsup_{N\to\infty}$ and use the inequalities (3.20), (3.21) and (3.22) to find that

$$\limsup_{N \to \infty} \|f - f_N\|_{L^p} \leqslant \|\chi_R^c f\|_{L^p} + K_m \|\chi_R^c \langle z \rangle^{-m}\|_{L^{r'}}.$$
(3.23)

It suffices to take the limit $R \to \infty$. This finishes the proof of the theorem.

Proof of Corollary 2.2. For simplicity we only present the proof for p = 1, the other cases being analogous with only minor modifications. Since one implication is obvious, we only prove the second one.

We start by using the fact $f = f^2$ through the formula f(1 - f) = 0, and estimate the L^1 norm difference as follows

$$\|f - f_N\|_{L^1} = \int f|f - f_N| + \int (1 - f)|f - f_N| = \int f|f - f_N| + \int (1 - f)|f_N| . \quad (3.24)$$

The first term in (3.24) can be estimated in view of the Cauchy-Schwarz inequality, and the p = 2 case established in Theorem 2. Namely

$$\int f|f - f_N| \leq ||f||_{L^2} ||f - f_N||_{L^2} = (2\pi)^{\frac{d}{2}} ||f - f_N||_{L^2} .$$
(3.25)

The second term in (3.24) is more involved. First, because of L^2 convergence, we may assume that convergence holds pointwise almost everywhere, up to extraction of a subsequence (which we do not display explicitly). Hence, Fatou's Lemma implies that

$$(2\pi)^d = \int f|f| \leqslant \liminf_{N \to \infty} \int f|f_N| . \tag{3.26}$$

14

On the other hand, $||f_N||_{L^2} = ||f||_{L^2} = (2\pi)^{\frac{d}{2}}$ combined with the Cauchy-Schwarz inequality implies

$$\limsup_{N \to \infty} \int f|f_N| \leq \|f\|_{L^2} \limsup_{N \to \infty} \|f_N\|_{L^2} = (2\pi)^d .$$
 (3.27)

Hence, (3.26) and (3.27) imply that $\lim_{N\to\infty} \int f|f_N| = (2\pi)^d$. Thus, going back to (3.24) we see that

$$\limsup_{N \to \infty} \|f - f_N\|_{L^1} \leqslant \limsup_{N \to \infty} \|f_N\|_{L^1} - (2\pi)^d .$$
(3.28)

Thus, thanks to the assumption in the statement of the Corollary, we find $\lim_{N\to\infty} ||f_N||_{L^1} = (2\pi)^d$. Finally, we note that the limit f is independent of the chosen subsequence. Hence, L^1 -convergence holds for the entire sequence (see also the comment at the beginning of Subsection 3.2).

4. Application: the harmonic trap

Throughout this section, we consider the external potential $U(x) = x^2$. Using the same notation as in the previous section, we denote by f_N the sequence of Wigner functions of the approximate ground state Ψ_N , taken to be a Slater determinant in the context of Theorem 2. We denote by $f = f_{TF}$ the classical state determined by ρ_{TF} , the minimizer of e_{TF} .

In the following lemma, we prove that the condition on the moments (2.15) is verified for m = 1. Note this is the only point in this article where we make direct use of the conditions A = 0 and $\hat{V} \ge 0$. We believe further analysis would show these can be relaxed to those considered originally in [9].

Lemma 4.1. Assume that Ψ_N is a Slater determinant for all $N \ge 1$. Then, for $U(x) = x^2$ there holds

$$\sup_{N \ge 1} \|(|x| + |p|) f_N\|_{L^2} < \infty .$$
(4.1)

Remark 4.1. As it will be clear from the proof, one only needs a lower bound $U(x) \ge Cx^2$ for the external trap, and the same result is true. For simplicity of the exposition we choose the harmonic trap $U(x) = x^2$.

Remark 4.2. The estimate (4.10) contained in the proof of Lemma 4.1 below is borrowed from the proof of [13, Theorem 3]. Here, the author considers the convergence of the Wigner transform of the following sequence of one-particle density matrices

$$\rho_{\hbar} = \mathbb{1}_{(-\infty,0]} \left(\hat{p}^2 + W(\hat{x}) \right)$$
(4.2)

where $W : \mathbb{R}^d \to \mathbb{R}$ is a nice potential chosen so that the spectrum of $p^2 + W(x)$ is discrete in $(-\infty, 0]$. These states were first studied in [21] for $d \ge 2$ where it is shown that they satisfy optimal semi-classical commutator estimates.

Corollary 4.1. Under the same assumptions as in Theorem 2 and Lemma 4.1, there holds

$$f_N \to f$$
 strongly in $L^p(\mathbb{R}^{2d})$ for $p \in [1,2] \cap \left(\frac{2}{1+2/d},2\right]$. (4.3)

Remark 4.3 (One-dimensional harmonic oscillator). Let d = 1 and consider the harmonic oscillator $h = -\hbar^2 d^2/dx^2 + x^2$ on $L^2(\mathbb{R})$, with spectrum $\sigma(h) = \bigcup_{n=0}^{\infty} \{E_n\}$ where $E_n = \hbar(n+1/2)$. Each eigenvalue E_n is non-degenerate and corresponds to the eigenfunction ψ_n being the *n*-th Hermite function. Since $(\psi_n)_n$ is an orthonormal basis for $L^2(\mathbb{R})$, it is possible to show that the ground state of $H_N = \sum_{i=1}^N -\hbar^2 d/dx_i^2 + x_i^2$ on $\bigwedge_{i=1}^N L^2(\mathbb{R})$ corresponds to the Slater determinant

$$\Psi_{N,0} = \psi_0 \wedge \dots \wedge \psi_{N-1} , \qquad (4.4)$$

with ground state energy $\sum_{n=0}^{N-1} E_n$. Letting $f_{N,0}$ be the Wigner transform of the ground state $\Psi_{N,0}$, Corollary 4.1 implies

$$f_{N,0} \to f_{\rho_0}$$
 strongly in $L^1(\mathbb{R}^2)$. (4.5)

where we denote by $\rho_0 \in L^1(\mathbb{R})$ the minimizer of the Thomas-Fermi functional.

Proof of Lemma 4.1. Let us denote by \hat{x} and $\hat{p} = -i\hbar\nabla_x$ the standard position and momentum observables in $L^2(\mathbb{R}^d)$. We start the proof with the identity

$$xf_N = (1/2) W^{\hbar} [\hat{x}\gamma_N + \gamma_N \hat{x}], \qquad (4.6)$$

which follows from the decomposition x = 1/2(x + y/2) + 1/2(x - y/2), the definition (2.4) and the relations $(\hat{x}\gamma)(x,y) = x\gamma(x,y)$ and $(\gamma\hat{x})(x,y) = y\gamma(x,y)$. Thus, $||W^{\hbar}[\gamma]||_{L^2} = (2\pi\hbar)^{d/2} ||\gamma||_{L^2}$ implies

$$\|xf_N\|_{L^2} \leqslant (1/2)(2\pi\hbar)^{d/2} \Big(\|\hat{x}\gamma_N\|_{L^2} + \|\gamma_N\hat{x}\|_{L^2}\Big) .$$
(4.7)

We now calculate the right hand side as follows

$$\|\hat{x}\gamma_N\|_{L^2}^2 = \|\hat{x}\gamma_N\|_{HS}^2 = \operatorname{Tr}(\hat{x}\gamma_N)^*(\hat{x}\gamma_N) = \operatorname{Tr}\left(\gamma_N\hat{x}^2\gamma_N\right) = \operatorname{Tr}\left(\hat{x}^2\gamma_N\right), \quad (4.8)$$

wher we used cyclicity of the trace, and $\gamma_N = \gamma_N^* = \gamma_N^2$. The same identity holds for the second term of (4.7). Thus,

$$\|xf_N\|_{L^2}^2 \leqslant (2\pi\hbar)^d \operatorname{Tr}(\hat{x}^2 \gamma_N) .$$
(4.9)

In a similar fashion, the same argument can be repeated for the momentum variable and can be combined with the last estimate. Thus, the triangle inequality and the scaling $\hbar^d = N^{-1}$ gives

$$\|(|x|+|p|)f_N\|_{L^2}^2 \leqslant \frac{C}{N} \operatorname{Tr}\left((\hat{x}^2+\hat{p}^2)\gamma_N\right), \qquad (4.10)$$

for a constant C > 0. It suffices to show that the right hand side of (4.10) is bounded uniformly in N. To this end, we use two facts: that γ_N corresponds to an approximate ground state; and that the two-body interaction is dominated by the kinetic energy. We do this in the following two steps. Step 1. Let Ψ_N be the approximate ground state under consideration, which we assume is a Slater determinant. Then, a standard calculation shows that

$$\langle \Psi_N, H_N \Psi_N \rangle_{\mathfrak{h}_N}$$

$$(4.11)$$

$$= \operatorname{Tr}(\hat{p}^{2} + \hat{x}^{2})\gamma_{N} + N \int_{\mathbb{R}^{2d}} \rho_{N}(x)V(x-y)\rho_{N}(y)\mathrm{d}x\mathrm{d}y - \frac{1}{N}\int_{\mathbb{R}^{2d}} V(x-y)|\gamma(x,y)|^{2}\mathrm{d}x\mathrm{d}y \ .$$

where we denote $\rho_N(x) \equiv \frac{1}{N}\gamma_N(x,x)$. Recall that in Assumption 1 we consider $\hat{V} \ge 0$. Hence, the second term on the right hand side above is non-negative, in view of the following representation

$$\int_{\mathbb{R}^{2d}} \rho(x) V(x-y) \rho(y) \mathrm{d}x \mathrm{d}y = \int_{\mathbb{R}^d} \hat{V}(k) |\hat{\rho}(k)|^2 \mathrm{d}k \ge 0.$$
(4.12)

Therefore, thanks to (4.11), (4.12) and (1.7) we conclude that for $N \ge N_0$ large enough

$$\operatorname{Tr}(\hat{p}^{2} + \hat{x}^{2})\gamma_{N} \leqslant CNe_{TF} + \frac{1}{N} \int_{\mathbb{R}^{2d}} V(x-y)|\gamma_{N}(x,y)|^{2} \mathrm{d}x \mathrm{d}y .$$

$$(4.13)$$

Step 2. Here, we borrow the following estimate from the proof of [9, Proposition 3.1]

$$\left|\frac{1}{N}\int_{\mathbb{R}^d\times\mathbb{R}^d}V(x-y)|\gamma(x,y)|^2\mathrm{d}x\mathrm{d}y\right|\leqslant\varepsilon_N\mathrm{Tr}(\hat{p}^2+\mathbb{1})\gamma_N\tag{4.14}$$

for a sequence of positive numbers $\varepsilon_N \to 0$ as $N \to \infty$. Consequently, for $N \ge N_0$ large enough we may assume $\varepsilon_N \le 1/2$. In particular, it follows from (4.13), (4.14) and $\text{Tr}\gamma_N = N$ that for a constant C > 0

$$\operatorname{Tr}(\hat{p}^2 + \hat{x}^2)\gamma_N \leqslant CN(e_{TF} + 1) . \tag{4.15}$$

The proof is finished once we combine the last inequality with (4.10).

5. Results on k-particle distribution functions

In this section we address the problem of convergence of higher-order distribution functions, as originally considered by the authors in [9].

5.1. **Definitions.** Let $2 \leq k < N$ and denote by $\Psi_N \in \mathfrak{h}_N$ the approximate ground state. We will work with the following definitions.

(1) Wigner functions. Following [9, Eq. (1.18)] we define the k-particle Wigner function as follows

$$f_N^{(k)}(X_k, P_k) \equiv \int_{\mathbb{R}^{2dk} \times \mathbb{R}^{d(N-k)}} \Psi_N\left(X_k + \frac{\hbar}{2}Y_k, X_{N-k}\right) \overline{\Psi_N}\left(X_k + \frac{\hbar}{2}Y_k, X_{N-k}\right) e^{-iY_k \cdot P_k} \mathrm{d}X_{N-k} \mathrm{d}Y_k$$
(5.1)

where $(X_k, P_k) \in \mathbb{R}^{2dk}$, and we write $X_{N-k} = (x_{k+1}, \ldots, x_N) \in \mathbb{R}^{d(N-k)}$ for the integration variable. In particular, the normalization is chosen so that

$$\frac{1}{(2\pi)^{dk}} \int_{\mathbb{R}^{2dk}} f_N^{(k)}(X_k, P_k) \mathrm{d}X_k \mathrm{d}P_k = 1 \quad .$$
(5.2)

(2) Husimi measures. In this article, we define the k-particle Husimi measure as the convolution

$$m_N^{(k)} \equiv \frac{N \cdots (N-k+1)}{N^k} f_N^{(k)} * \mathscr{G}_{\hbar}^{\otimes k}$$
(5.3)

where $\mathscr{G}_{\hbar}^{\otimes k} = \mathscr{G}_{\hbar} \otimes \cdots \otimes \mathscr{G}_{\hbar}$ is the k-fold tensor product of the one-particle Gaussian mollifier $\mathscr{G}_{\hbar}(z) = \hbar^{-d} \mathscr{G}_{1}(\hbar^{-1/2}z)$. As explained by the authors in [9, Section 2.4], the convolution (5.3) agrees with their original definition. In particular, we have [9, Lemma 2.2]

$$0 \leqslant m_N^{(k)} \leqslant 1$$
 and $\frac{1}{(2\pi)^{dk}} \|m_N^{(k)}\|_{L^1} = \frac{N \cdots (N-k+1)}{N^k}$. (5.4)

(3) Reduced densities. We define $\gamma_N^{(k)}$, the k-particle reduced density matrix, as the trace-class operator on $L^2(\mathbb{R}^{dk})$ with kernel

$$\gamma_N^{(k)}(X_k, X_k') \equiv \frac{N!}{(N-k)!} \int_{\mathbb{R}^{d(N-k)}} \Psi_N(X_k, X_{N-k}) \overline{\Psi_N(X_k', X_{N-k})} \, \mathrm{d}X_{N-k}$$
(5.5)

where $X_k = (x_1, ..., x_k), X'_k = (x'_1, ..., x'_k) \in \mathbb{R}^{dk}$ and we denote $X_{N-k} = (x_{k+1}, ..., x_N)$. In particular, the normalization is chosen so that

$$\operatorname{Tr}\gamma_N^{(k)} = \frac{N!}{(N-k)!}$$
 (5.6)

Note the definition is not the same as [9], but we take the one from [16] since we shall borrow a calculation in the proof of Lemma 5.1 below.

5.2. Statements. In [9] the authors prove that the following convergence statement holds

$$\left\langle \varphi, m_N^{(k)} \right\rangle \to \left\langle \varphi, f^{\otimes k} \right\rangle$$
 (5.7)

for all $\varphi \in L^1(\mathbb{R}^{2dk}) + L^{\infty}(\mathbb{R}^{2dk})$, where $f = f_{\rho_{TF}}$ is defined in terms of the unique minimizer of the Thomas-Fermi functional (1.5). In particular, one may then ask if the results contained in Theorem 1 and 2 extend to higher-order distribution functions.

We address this question in the next Theorem. While some convergence results are easily translated into higher-order distribution functions, it turns out that some are not. Most notably, we prove that for Slater determinants the strong L^2 convergence $f_N^{(k)} \to f^{\otimes k}$ holds only for k = 1, although convergence holds in H^s for all s < 0.

Theorem 3 (k-particle functions). Let $k \ge 2$. Then, the following statements are true.

- (1) The sequence $m_N^{(k)}$ converges to $f^{\otimes k}$ strongly in $L^p(\mathbb{R}^{2dk})$ for all $p \in [1, \infty)$. (2) The sequence $f_N^{(k)}$ converges to $f^{\otimes k}$ strongly in $H_{s,\infty}(\mathbb{R}^{2dk})$ for all s < 0.
- Additionally, if Ψ_N is a Slater determinant for all $N \ge 1$, the following is true.
 - (3) The sequence $f_N^{(k)}$ converges to $f^{\otimes k}$ strongly in $H_{s,q}(\mathbb{R}^{2dk})$ for all s < 0 and $q \in [2, \infty]$.
 - (4) The sequence $f_N^{(k)}$ does not converge to $f^{\otimes k}$ strongly in $L^2(\mathbb{R}^{2dk})$. In fact, there holds

$$\liminf_{N \to \infty} \|f_N^{(k)} - f^{\otimes k}\|_{L^2} \ge (2\pi)^{dk/2} (\sqrt{k!} - 1).$$
(5.8)

Remark 5.1 (Lack of uniform L^2 bounds). Note that in item (2) we are not able to obtain convergence in $H^s(\mathbb{R}^{2dk})$, as opposed to the k = 1 case in Theorem 1. This is a consequence of our lack of control of the L^2 norms of $f_N^{(k)}$ for $k \ge 2$. Recall that for density matrices $\gamma_N^{(k)}$ for $k \ge 2$ one does not have operator bounds which are uniform in N. For illustration, for k = 2 it is in general known that

$$0 \leqslant \gamma_N^{(2)} \leqslant (N-1)\mathbb{1} , \qquad (5.9)$$

see e.g [1, Lemma 1] or [20, Theorem 8.12]. In contrast, for the case k = 1, there holds $0 \leq \gamma_N^{(1)} \leq 1$, which implies $\|f_N^{(1)}\|_{L^2} \leq (2\pi)^{d/2}$. On the other hand, the operator bound (5.9) only gives the estimate $\|\gamma_N^{(2)}\|_{HS} \leq \|\gamma_N^{(2)}\|_2^{\frac{1}{2}} \|\gamma_N^{(2)}\|_{\mathrm{Tr}}^{\frac{1}{2}} \leq N^{3/2}$, which is not sufficient to guarantee a uniform L^2 bound for $f_N^{(2)}$. Let us further make two comments.

- (1) In item (3), the uniform L^2 bounds are recovered for the Slater determinants, for which an explicit formula can be given for all $k \ge 2$; see Lemma 5.1. This allows for convergence in the larger class of spaces.
- (2) Upon completion of this work, the author became aware of the recent result of Christiansen [6, Theorem 1]², where it is proven that $\|\gamma_N^{(2)}\|_{HS} \leq \sqrt{5}N$. The proof bypasses the operator norm (5.9) and holds for any normalized state Ψ_N . Consequently, $\sup_{N\geq 1} \|f_N^{(2)}\|_{L^2} < \infty$ using the unitarity of the Wigner transform (see also (5.16)). Hence, we can relax the assumption of Ψ_N being a Slater determinant in item (3) in Theorem 3 for k = 2. In particular, $f_N^{(2)} \to f \otimes f$ in $H^s(\mathbb{R}^{4d})$ for all s < 0.

The proof of items (3) and (4) of Theorem 3 are based on the following calculation. While the result may have been known to experts, we record it here for completeness of the exposition.

Lemma 5.1. Assume that Ψ_N is a Slater determinant for all $N \ge 2$. Then, for all $k \ge 2$

$$\lim_{N \to \infty} \|f_N^{(k)}\|_{L^2} = \left((2\pi)^{dk} k! \right)^{\frac{1}{2}} .$$
 (5.10)

Remark 5.2. The L^2 norm of $f_N^{(k)}$ can be computed explicitly for finite N, see (5.20).

5.3. **Proof of Theorem 3.** First, we shall give a proof of Theorem 3 which is based on Lemma 5.1. Then, we prove Lemma 5.1.

Before we turn to the proof of Theorem 3, let us introduce some notation. Namely, it will be extremely convenient to recast the Wigner function $f_N^{(k)}$ as the Wigner transform of the reduced density matrix $\gamma_N^{(k)}$. To this end, we define the following linear map, which we refer to as the Wigner transform of order k:

$$W_k^{\hbar}[\gamma_k](X_k, P_k) \equiv \int_{\mathbb{R}^{dk}} \gamma_k \left(X_k + \frac{1}{2} Y_k, X_k - \frac{1}{2} Y_k \right) \exp\left(-\frac{i}{\hbar} Y_k \cdot P_k \right) \, \mathrm{d}Y_k \tag{5.11}$$

where $\gamma_k \in L^2(\mathbb{R}^{2dk})$ and $(X_k, P_k) \in \mathbb{R}^{2dk}$. In particular, the following is true.

²The author would like to thank the anonymous referee that pointed this out

(i) The map W_k^{\hbar} is an L²-isomorphism, and with the present normalization

$$\|W_k^{\hbar}[\gamma_k]\|_{L^2} = (2\pi\hbar)^{\frac{dk}{2}} \|\gamma_k\|_{L^2} , \qquad \gamma_k \in L^2(\mathbb{R}^{2dk}) .$$
(5.12)

See e.g [7, Proposition 13].

(ii) Starting from (5.1), we use the definition (5.5) of $\gamma_N^{(k)}$, change variables $Y_k \mapsto \hbar^{-1}Y_k$ and use the scaling $\hbar^{-dk} = N^k$ to find that

$$f_N^{(k)} = N^k \frac{(N-k)!}{N!} W_k^{\hbar} [\gamma_N^{(k)}] .$$

$$\frac{(N-k)!}{N!} = 1$$
(5.13)

Note that $\lim_{N\to\infty} N^k \frac{(N-k)!}{N!} = 1.$

Proof of Theorem 3. In what follows we fix $k \ge 2$ and assume N > k.

(1) The proof relies on Lemma 3.1. Here, we take $F_n(Z_k) = m_N^{(k)}((2\pi)^{\frac{1}{2}}Z_k)$ and similarly $F(Z_k) = (f^{\otimes k})((2\pi)^{\frac{1}{2}}Z_k)$ for $Z_k \in \mathbb{R}^{2dk}$. The bounds (3.1) and (3.2) are straightforward to verify and follow from (5.4) and $\int f = (2\pi)^{d/2}$ and $f = f^2$, respectively. The weak convergence (5.7) has been proven in [9].

(2) Here, we repeat the argument used in the proof of Theorem 1. The proof is identical, and the only modification comes from justifying the uniform bound $\|\hat{f}_N\|_{L^{\infty}} \leq 1$. To justify this in the case $k \geq 2$, we note that a calculation using (5.13) and the scaling $\hbar^{dk} = N^{-k}$ yields the analogous Groenewold's formula (2.6) for k-particle Wigner functions

$$\widehat{f_N^{(k)}}(\zeta_k) = \frac{(N-k)!}{N!} \operatorname{Tr} \left[\mathcal{O}_{\zeta_k}^{(k)} \gamma_N^{(k)} \right], \qquad \zeta_k = (\xi_k, \eta_k) \in \mathbb{R}^{dk} \times \mathbb{R}^{dk}$$
(5.14)

where the trace is over $L^2(\mathbb{R}^{dk})$, and $\mathcal{O}_{\zeta_k}^{(k)} = \exp(i(\xi_k \hat{X}_k + \eta_k \hat{P}_k))$ is a semi-classical observable in $L^2(\mathbb{R}^{2dk})$. Hence, thanks to (5.6) we find $\|\widehat{f_N^{(k)}}\|_{L^{\infty}} \leq 1$.

(3) We repeat the argument used in the proof of Theorem 1. Namely, we prove the q = 2 case. The only modification comes from the uniform L^2 bound $||f_N||_{L^2} \leq (2\pi)^{d/2}$. For $k \geq 2$, this uniform bound is now provided by Lemma 5.1, i.e. there holds $\sup_{N\geq 1} ||f_N^{(k)}||_{L^2} < \infty$. We can then interpolate between q = 2 and $q = \infty$.

(4) It suffices to use the triangle inequality, Lemma 5.1 and $||f^{\otimes k}||_{L^2} = ||f||_{L^2}^k = (2\pi)^{dk/2}$ to find that

$$\liminf_{N \to \infty} \|f_N^{(k)} - f^{\otimes k}\|_{L^2} \ge \liminf_{N \to \infty} \|f_N^{(k)}\|_{L^2} - \|f^{\otimes k}\|_{L^2} = (2\pi)^{dk/2} \left(\sqrt{k!} - 1\right) \,. \tag{5.15}$$

This finishes the proof of the theorem.

Proof of Lemma 5.1. Fix $k \ge 2$ and let N > k. We split the proof into two parts. In the first part, we calculate a relation between $\|f_N^{(k)}\|_{L^2}$ and $\|\gamma_N^{(k)}\|_{HS}$, which is independent of Ψ_N being a Slater determinant. In the second part, we use the fact that Ψ_N is a Slater determinant to compute the Hilbert-Schmidt norm of $\gamma_N^{(k)}$.

20

For the first part, we use (5.13) and then (5.12) to find that

$$\|f_{N}^{(k)}\|_{L^{2}} = N^{k} \frac{(N-k)!}{N!} \|W_{k}^{\hbar}[\gamma_{N}^{(k)}]\|_{L^{2}}$$

$$= N^{k} \frac{(N-k)!}{N!} (2\pi\hbar)^{\frac{dk}{2}} \|\gamma_{N}^{(k)}\|_{L^{2}}$$

$$= N^{k} \frac{(N-k)!}{N!} (2\pi\hbar)^{\frac{dk}{2}} \|\gamma_{N}^{(k)}\|_{HS}$$
(5.16)

where in the last line we used $\|\gamma_k\|_{HS} = \|\gamma_k\|_{L^2}$.

For the second part, we observe that when Ψ_N is a Slater determinant, the k-particle reduced density matrix $\gamma_N^{(k)}$ can be calculated explicitly. Namely, assume that

$$\Psi_N(x_1,\ldots,x_N) = \frac{1}{\sqrt{N!}} \det_{1 \le i,j \le N} \left[\varphi_i(x_j) \right], \qquad (x_1,\ldots,x_N) \in \mathbb{R}^{dN} . \tag{5.17}$$

The orbitals $(\varphi_i)_{i=1}^N$ can depend on N but we do not display such dependence in the notation, for it has no effect in the upcoming calcuation. Next, we note that the kernel of $\gamma_N^{(k)}$ is given by (see for reference [16, Section 3.1.5])

$$\gamma_N^{(k)}(x_1,\ldots,x_k,x_1',\ldots,x_k') = k! \sum_{1 \le \ell_1 < \cdots < \ell_k \le N} \frac{1}{\sqrt{k}} \det_{1 \le i,j \le N} \left[\varphi_{\ell_i}(x_j) \right] \frac{1}{\sqrt{k}} \det_{1 \le i,j \le N} \left[\varphi_{\ell_i}(x_j') \right].$$

Hence, we write in operator form

$$\gamma_N^{(k)} = k! \sum_{1 \leqslant \ell_1 < \dots < \ell_k \leqslant N} |\varphi_{\ell_1} \wedge \dots \wedge \varphi_{\ell_k}\rangle \langle \varphi_{\ell_1} \wedge \dots \wedge \varphi_{\ell_k}| \quad .$$
(5.18)

In particular, note that $\operatorname{Tr}\gamma_N^{(k)} = k! \binom{N}{k} = \frac{N!}{(N-k)!}$. Furthermore, it follows from (5.18) that $\gamma_N^{(k)}\gamma_N^{(k)} = k!\gamma_N^{(k)}$. Thus, we can compute the Hilbert-Schmidt norm as follows

$$\|\gamma_N^{(k)}\|_{HS}^2 = \operatorname{Tr}\left[(\gamma_N^{(k)})^* \gamma_N^{(k)}\right] = k! \operatorname{Tr}\gamma_N^{(k)} = k! \frac{N!}{(N-k)!} , \qquad (5.19)$$

where we have used self-adjointness of $\gamma_N^{(k)}$.

Let us now put everything together. Namely (5.19) and (5.16) imply

$$\|f_N^{(k)}\|_{L^2} = N^k \frac{(N-k)!}{N!} (2\pi\hbar)^{\frac{dk}{2}} \left(k! \frac{N!}{(N-k)!}\right)^{\frac{1}{2}} \\ = \left((2\pi)^{dk} k!\right)^{\frac{1}{2}} \left[N^k \hbar^{\frac{dk}{2}} \left(\frac{(N-k)!}{N!}\right)^{\frac{1}{2}}\right].$$
(5.20)

It suffices now to use the scaling $\hbar^d = N^{-1}$, take the limit $N \to \infty$ and observe that the factor in squared brackets $[\cdots]$ in (5.20) converges to 1.

Acknowledgements. I am grateful to J.K Miller and N. Pavlović for discussions that ultimately led to the study of the problem in this article. I am also very thankful to D. Hundertmark for his comments that helped improve an earlier version of this manuscript.

I would also like to acknowledge important remarks from two anonymous referees that helped significantly improve the conclusion of Theorem 1, covering now the additional range $p \in (2, \infty)$. The author gratefully acknowledges support from the Provost's Graduate Excellence Fellowship at The University of Texas at Austin and from the NSF grant DMS-2009549, and the NSF grant DMS-2009800 through T. Chen.

Data availability. This manuscript has no associated data.

Conflict of interest. On behalf of all authors, the corresponding author states that there is no conflict of interest.

References

- V. Bach. Hartree-Fock Theory, Lieb's Variational Principle, and their Generalizations. arXiv:2209.10189.
- [2] N. Benedikter. Effective dynamics of interacting fermions from semiclassical theory to the random phase approximation. J. Math. Phys. 1 August 2022; 63 (8): 081101.
- [3] N. Benedikter, M. Porta, B. Schlein. Mean-Field Evolution of Fermionic Systems. Commun. Math. Phys. 331, 1087–1131 (2014).
- [4] N. Benedikter, M. Porta, C. Saffirio, B. Schlein. From the Hartree Dynamics to the Vlasov Equation. Arch. Rational Mech. Anal. 221 (2016) 273–334.
- [5] N. Benedikter, V. Jakšić, M. Porta, C. Saffirio, B. Schlein. Mean-Field Evolution of Fermionic Mixed States. Comm. Pure Appl. Math., 69: 2250-2303 (2016).
- [6] M. R. Christiansen. Hilbert-schmidt estimates for fermionic 2-body operators. Commun. Math. Phys., 405 (2024), pp. 1–9.
- [7] M. Combescure, D. Robert. Coherent States and Applications in Mathematical Physics. Theoretical and Mathematical Physics. Springer, Dordrecht (2012).
- [8] E. Fermi. Un metodo statistico per la determinazione di alcune priorieta dell'atome. Rend. Accad. Naz. Lincei 6, 602–607 (1927)
- [9] S. Fournais, M. Lewin, J. P. Solovej, The semi-classical limit of large fermionic systems. Calc. Var. (2018) 57:105.
- [10] L. Fresta, M. Porta, B. Schlein. Effective Dynamics of Extended Fermi Gases in the High-Density Regime. Commun. Math. Phys. 401, 1701–1751 (2023).
- [11] T. Girardot, N. Rougerie. Semiclassical Limit for Almost Fermionic Anyons. Commun. Math. Phys. 387, 427–480 (2021).
- [12] N. Gottschling, P. T. Nam. Convergence of Levy-Lieb to Thomas-Fermi density functional. Calc. Var. (2018) 57:146.
- [13] L. Lafleche. Optimal Semiclassical Regularity of Projection Operators and Strong Weyl Law. arXiv:2302.04816.
- [14] M. Lewin, P. S. Madsen, A. Triay. Semi-classical limit of large fermionic systems at positive temperature. J. Math. Phys. 1 September 2019; 60 (9): 091901.
- [15] M. Lewin, P. T. Nam, N. Rougerie. Derivation of Hartree's theory for generic mean-field Bose systems. Advances in Mathematics, Volume 254, 2014, Pages 570-621.
- [16] E. Lieb, R. Seiringer. The Stability of Matter in Quantum Mechanics. Cambridge: Cambridge University Press. (2009).
- [17] E. H. Lieb, B. Simon. Thomas-Fermi theory revisited. Physical Review Letters, 31 (1973), p. 681.
- [18] E. H. Lieb, B. Simon. The Thomas-Fermi theory of atoms, molecules and solids. Advances in mathematics, 23 (1977), pp. 22–116
- [19] N. N. Nguyen. Weyl laws for interacting particles. arXiv:2305.06237.

22

- [20] J.P. Solovej. Many Body Quantum Mechanics. March 2014.
- [21] S. Fournais, S. Mikkelsen. An optimal semiclassical bound on commutators of spectral projections with position and momentum operators. Lett Math Phys 110, 3343–3373 (2020).
- [22] S. Fournais, P.S. Madsen. Semi-classical Limit of Confined Fermionic Systems in Homogeneous Magnetic Fields. Ann. Henri Poincaré 21, 1401–1449 (2020).
- [23] L.H. Thomas. The calculation of atomic fields. Proc. Camb. Philos. Soc. 23, 542–548 (1927)
- [24] G. Toscani, C. Villani. Probability Metrics and Uniqueness of the Solution to the Boltzmann Equation for a Maxwell Gas. Journal of Statistical Physics 94, 619–637 (1999).

(Esteban Cárdenas) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEXAS AT AUSTIN, 2515 SPEEDWAY, AUSTIN TX, 78712, USA

Email address: eacardenas@utexas.edu