A sufficient condition for superstatistics in steady state ensembles

Constanza Farías¹ and Sergio Davis^{2,1}

E-mail: m.fariasparra@uandresbello.edu

¹Departamento de Física, Facultad de Ciencias Exactas, Universidad Andres Bello. Sazié 2212, piso 7, Santiago, 8370136, Chile.

²Research Center in the Intersection of Plasma Physics, Matter and Complexity (P²mc), Comisión Chilena de Energía Nuclear, Casilla 188-D, Santiago, Chile

Abstract. In recent years, the theory of superstatistics, which aims to describe nonequilibrium steady state systems, has gained attention due to its different real world applications, highlighting its versatility and concise mathematical formulation in terms of a probability density for the inverse temperature $\beta = 1/k_BT$. When exploring the domain of application of the superstatistical theory, recent works have shown some necessary conditions for a superstatistical description of a given steady state, in terms of the fundamental and microcanonical inverse temperature. In this work, a new theorem that establishes a sufficient condition for the existence of a superstatistical description of a particular steady state is presented, using the language of momentgenerating functions and connecting them with properties of the derivatives of the fundamental inverse temperature.

1. Introduction

The study of complex systems has gained relevant interest in the latest years because of its importance in Nature, such as in weather prediction, human behavior, among others. There exist several ways to describe the complexity of systems, for example, stochastic processes or graph theory for network science are often used. In statistical mechanics some generalizations have been proposed, such as non-extensive statistical physics or the more recent theory of *superstatistics* [1,2], that generalizes Boltzmann-Gibbs statistical mechanics and aims to explain the statistics of different types of systems characterized to be out-of-equilibrium. From its beginning, this theory has been applied in different areas of knowledge, such as, the study of turbulent fluids [3,4], plasma physics [5,6], astrophysics [7–9], environmental science [10–13], spins models [14,15], and most recently in describing the statistical properties of power grids [16,17].

In a particular way, the theory of superstatistics has been used based on statistical inference to determine probability distributions of complex systems, being its goal to give a complete description of systems that are characterized by a fluctuating quantity, the inverse temperature β , when these systems are in contact with an environment. It

is known from the theory of probability and statistics, that a probability distribution could model most aspects of a system of interest, as the probability density function provides relevant information about all degrees of freedom in a system. In addition, the analysis of the moments of a distribution can give essential information, by describing in a complete manner the probability distribution. Based on the essence of the last argument, the superstatistical framework uses these mathematical tools to describe and find the properties of the system of interest just by analyzing the inverse temperature, which in this case is the relevant fluctuating quantity in such a system. Until now, recent works have developed the idea of exploring the limitations of the superstatistical framework [14,18], introducing novel ways of classifying out-of-equilibrium steady state systems given by its inverse temperature covariance \mathcal{U} . In particular, the fact that $\mathcal{U} \leq 0$ in superstatistics constitutes a necessary condition for its validity.

In this work we introduce a sufficient condition for the applicability of the superstatistical framework, which is based on the moment generating function [19-24] and its role regarding a probability distribution. In order to prove the soundness of this condition, we present two different applications. The paper is organized as follows. In Section 2, we briefly describe the theory of superstatistics and give the definition of the fundamental and microcanonical inverse temperatures, showing in addition the actual boundaries proposed for superstatistics. In Section 3, we introduce the sufficient conditions for superstatistics as a new theorem, and in 4 we present the proof of said theorem. Section 6 shows two different examples in order to illustrate the sufficient condition, and finally in Section 7, we provide some concluding remarks.

2. Superstatistics and its boundaries

2.1. Superstatistics

The theory of superstatistics is considered a concise and powerful formalism to structure in a more general manner the framework of statistical physics, and is sometimes called a *statistics of statistics*. This theory could be understood as the superposition of different canonical ensembles, each of them based on the traditional statistical mechanics, in which the temperature T is a fixed intensive quantity. In superstatistics theory, the inverse temperature $\beta := 1/T$, becomes a fluctuating quantity, therefore it is taken as a random variable whose probability distribution we will denote as $P(\beta|S)$. The distribution of microstates is defined as usual in the canonical ensemble,

$$P(\boldsymbol{x}|\beta) = \frac{\exp(-\beta \mathcal{H}(\boldsymbol{x}))}{Z(\beta)},\tag{1}$$

where $Z(\beta) = \int d\boldsymbol{x} \exp(-\beta \mathcal{H}(\boldsymbol{x}))$ is known as the *partition function* and $\mathcal{H}(\boldsymbol{x})$ is the Hamiltonian of the system.

The probability distribution (1) becomes modified as the system experiences fluctuations of the inverse temperature β in such a way that it is replaced by a joint

A sufficient condition for superstatistics in steady state ensembles

probability distribution, upon which we can use *Bayes' theorem* [25],

$$P(\boldsymbol{x}, \beta|S) = P(\boldsymbol{x}|S)P(\beta|S)$$

= $\left[\frac{\exp(-\beta \mathcal{H}(\boldsymbol{x}))}{Z(\beta)}\right]P(\beta|S),$ (2)

where S represents the parameters that describe the system and $P(\beta|S)$ is known as the *temperature distribution*, which contains all the information of the temperature fluctuations relevant to the system.

In the case of non-equilibrium steady state systems, the probability density of microstates is of the form

$$P(\boldsymbol{x}|S) = \rho(\mathcal{H}(\boldsymbol{x})), \qquad (3)$$

where the function $\rho(E)$ is known as the generalized Boltzmann factor or ensemble function, which is considered as the Laplace transform of $f(\beta)$,

$$\rho(E;S) = \int_0^\infty d\beta f(\beta) \exp\left(-\beta E\right),\tag{4}$$

where

$$f(\beta) := \frac{P(\beta|S)}{Z(\beta)}.$$
(5)

In a similar way as *Tsallis statistics* [26], in the limit of $q \rightarrow 1$ superstatistics recovers the classical canonical ensemble in the case where

$$P(\beta|S) \longrightarrow \delta(\beta - \beta_0). \tag{6}$$

2.2. Temperature definition in superstatistics

For every non-equilibrium steady state described by Eq. 4 we can obtain the temperature by two different paths. On the one hand we have the *fundamental inverse* temperature [27, 28], defined by

$$\beta_F(E) := -\frac{\partial}{\partial E} \ln \rho(E), \tag{7}$$

which comes from the definition of the Shannon-Jaynes entropy. On the other hand, we have the *microcanonical inverse temperature* [29-36]

$$\beta_{\Omega}(E) := \frac{\partial}{\partial E} \ln \Omega(E), \tag{8}$$

which is related to the logarithm of the density of states, i.e. the Boltzmann entropy of the system, where

$$\Omega(E) := \int d\boldsymbol{x} \delta(E - \mathcal{H}(\boldsymbol{x})).$$
(9)

2.3. When is a system superstatistical?

In recent works [37,38], the fundamental temperature defined earlier and its connection with superstatistics theory has been explored. In addition, its properties can be determined using different fluctuation theorems, such as the conjugate variables theorem (CVT), providing a different path to find some features of the inverse temperature and other relevant information for the description of the system. Particularly, from Eq. 7 we see that β_F agrees with the first moment of the $P(\beta|E, S)$ distribution, and in general the moments of β given E can be written as

$$\left\langle \beta^n \right\rangle_{E,S} = \frac{(-1)^n}{\rho} \frac{\partial^n \rho}{\partial E^n},\tag{10}$$

from which $\langle \beta \rangle_{E,S} = \beta_F(E)$ can be easily seen by using n = 1. On the other hand, if we take n = 2 we obtain

$$\left\langle \beta^2 \right\rangle_{E,S} = \frac{1}{\rho} \frac{\partial^2 \rho}{\partial E^2},$$
(11)

and after using both of them, we can build the conditional variance for β as

$$\langle (\delta\beta)^2 \rangle_{E,S} = \langle \beta^2 \rangle_{E,S} - \langle \beta \rangle_{E,S}^2 = \frac{1}{\rho} \frac{\partial^2 \rho}{\partial E^2} - \left(\frac{\partial}{\partial E} \ln \rho \right)^2 = \frac{\partial^2}{\partial E^2} \ln \rho = -\frac{\partial \beta_F(E)}{\partial E},$$
(12)

then it follows that, for every superstatistical model it must hold that

$$\frac{\partial \beta_F(E)}{\partial E} = -\left\langle (\delta\beta)^2 \right\rangle_{E,S} \le 0.$$
(13)

This condition still applies if we consider a system in contact with its environment in which it has $C_V > 0$: such a system will not follow superstatistics [14,38], as we have

$$\frac{\partial \beta_F^{(\boldsymbol{x})}}{\partial E} = \frac{\left(\beta_{\Omega}^{(\boldsymbol{y})}\right)^2}{C_V^{(\boldsymbol{y})}} \ge 0, \tag{14}$$

where we have expressed C_V in terms of the derivative of β_{Ω} using

$$C_V(\varepsilon) = \left(\frac{\partial T(\varepsilon)}{\partial \varepsilon}\right)^{-1} = -\frac{\beta_{\Omega}(\varepsilon)^2}{\beta'_{\Omega}(\varepsilon)}.$$
(15)

Another necessary condition for superstatistics is related to the covariance of the inverse temperature [18], defined by

$$\mathcal{U} := \langle \delta \beta_F \delta \beta_\Omega \rangle$$

and which allows us to classify supercanonical or subcanonical steady states. This quantity is connected to the variance of both inverse temperatures, microcanonical (8) and fundamental (7). In the case of a subcanonical state, $\mathcal{U} < 0$ and therefore \mathcal{U} cannot be equal to a variance, however it is always equal to the covariance between β_F and β_{Ω} , which can be negative. For this reason, superstatistics is only possible when $\mathcal{U} \geq 0$.

3. Existence of Superstatistics

In the last section we have reviewed, according to the literature, two of the necessary conditions for the existence of superstatistics in non-equilibrium systems. In this work we go further and present the proof of a sufficient condition. The motivation of this theorem comes from the relevance of the analysis of different *moments* and *cumulants* [39] of a distribution in several works related to superstatistics [40–43].

In order to introduce the required tools, let us consider the definition of β_F according to Eq. 7, which in its integral form is

$$\beta_F(E) = \frac{1}{\rho(E|S)} \int_0^\infty \beta \exp\left(-\beta E\right) f(\beta) d\beta = \langle\beta\rangle_{E,S}.$$
 (16)

Then, a relation between the moments of the distribution $P(\beta|S)$ with β_F could be obtained by the *Fluctuation-Dissipation theorem*, given by,

$$\partial_E \langle \omega \rangle_{E,S} = \langle \partial_E \omega \rangle_{E,S} + \langle \omega \partial_E \ln P(\beta | E, S) \rangle_{E,S}.$$
(17)

It is straightforward to prove, using $\omega = \beta^k$, that for the k-th power of β we have

$$\partial_E \left\langle \beta^k \right\rangle_{E,S} = \beta_F(E) \left\langle \beta^k \right\rangle_{E,S} - \left\langle \beta^{k+1} \right\rangle_{E,S}.$$
(18)

As we see shortly, the moments of the inverse temperature β given E are functions of β_F and its derivatives only. In order to illustrate how is it so, we will use the fluctuation-dissipation theorem to obtain a few of those moments,

$$\langle \beta \rangle_{E,S} = \beta_F,$$

$$\langle \beta^2 \rangle_{E,S} = \beta_F^2 - \beta'_F,$$

$$\langle \beta^3 \rangle_{E,S} = \beta''_F - 3\beta_F \beta'_F + \beta_F^3,$$

$$\langle \beta^4 \rangle_{E,S} = -\beta'''_F + 4\beta_F \beta''_F - 6\beta_F^2 \beta'_F + 3(\beta'_F)^2 + \beta_F^4,$$

$$(19)$$

and so on. Our main result is the following theorem, that we will prove in the next section.

Theorem 1. Provided that

- (i) β_F and all of its derivatives exist and are all finite, and
- (ii) The odd derivatives of β_F are negative or zero and even derivatives of β_F are positive or zero,

then a superstatistical representation exists for $\rho(E)$.

4. Proof of the theorem

The moments of the distributions [44, 45] allow us to describe in a complete form the probability density function and sometimes provide a better understanding of phenomena as demonstrated several works [46, 47]. These objects could be computed given the moment generation function M(t, E) [48], defined for a continuus, non-negative variable x as

$$M_x(t) = \langle \exp(tx) \rangle_I = \int_0^\infty \exp(tx) P(x|I) dx$$

The mathematical background of the superstatistical framework as we introduce in the previous section, is based on statistics, in particular this framework requires information of the system given by the density of states to model the system through the probability distribution, or in statistical physics language, the ensemble. In the context of superstatistics, we take Eq. (4) and generalize it for the *n*-th derivative of $\rho(E; S)$, which is given by

$$\frac{\partial^n}{\partial E^n}\rho(E;S) = \int_0^\infty d\beta \frac{P(\beta|S)}{Z(\beta)} (-1)^n \beta^n \exp\left(-\beta E\right).$$
(20)

Using Bayes' theorem, we can find the conditional temperature distribution with the energy of the system

$$P(\beta|E,S) = \frac{\exp\left(-\beta E\right)f(\beta)}{\rho(E|S)}.$$
(21)

Therefore, we can write

$$\langle \beta^n \rangle_{E,S} = \int_0^\infty d\beta \left[\frac{P(\beta|S)}{\rho(E;S)} \frac{\exp(-\beta E)}{Z(\beta)} \right] \beta^n = \frac{(-1)^n}{\rho(E;S)} \frac{\partial^n \rho(E;S)}{\partial E^n}.$$
 (22)

In order to proceed to the proof, we will first prove the following lemma.

Lemma 1. The n-th derivative of the moment generating function for $P(\beta|E, S)$ in the neighbourhood of t = 0, that is, the n-th moment of β given E, is a function of β_F and its derivatives only, of the type

$$\langle \beta^n \rangle_{E,S} = \partial_t^n M_\beta(t) \big|_{t=0} = G_n \big(\beta_F(E), \beta'_F, \dots, \beta_F^{n-1} \big), \tag{23}$$

where G_n is a well-defined target function to be determined.

On the other hand, the moment generating function for $P(\beta|E, S)$ could be written as

$$M_{\beta}(t; E) := \langle \exp(t\beta) \rangle_{E,S} = \sum_{k=0}^{\infty} \frac{t^k}{k!} \left\langle \beta^k \right\rangle_{E,S}, \qquad (24)$$

that is,

$$M_{\beta}(t;E) = \int_0^\infty \exp\left(\beta(t-E)\right) \frac{f(\beta)}{\rho(E|S)} d\beta,$$
(25)

A sufficient condition for superstatistics in steady state ensembles

which can be compactly written as

$$M_{\beta}(t;E) = \frac{\rho(E-t|S)}{\rho(E|S)},\tag{26}$$

depending only on the ensemble function. Moreover, we can write

$$M_{\beta}(t) = \exp\left(\ln\frac{\rho(E-t|S)}{\rho(E|S)}\right)$$
(27)

in such a way that we can compute its n-th derivative as

$$\langle \beta^n \rangle_{E,S} = \partial_t^n M_\beta(t)|_{t=0} = \partial_t^n \left[\exp\left(\ln \frac{\rho(E-t|S)}{\rho(E|S)} \right) \right]_{t=0},$$
(28)

Using the generalized chain rule for the n-th derivative and the definition of the *Bell* polynomials [49], we obtain

$$\partial_t^n \left[\exp\left(\ln \frac{\rho(E-t|S)}{\rho(E|S)} \right) \right]_{t=0} = B_n \left(\ln^{(1)}(\rho_{Et}), \ln^{(2)}(\rho_{Et}), ..., \ln^{(n)}(\rho_{Et}) \right) \Big|_{t=0}, \quad (29)$$

where we have introduced the shortcut notation

$$\rho_{Et} := \frac{\rho(E - t|S)}{\rho(E|S)},\tag{30}$$

$$\ln^{(n)}(\rho_{Et}) := \frac{\partial^n}{\partial t_n} \ln \rho_{Et}.$$
(31)

By definition, the *n*-th complete exponential Bell polynomials are given by,

$$B_n(x_1, \dots, x_n) = \sum_{k=1}^n \sum \frac{n!}{j_1! j_2! \dots j_{n-k+1}!} \left(\frac{x_1}{1!}\right)^{j_1} \left(\frac{x_2}{2!}\right)^{j_2} \dots \left(\frac{x_{n-k+1}}{(n-k+1)!}\right)^{j_{n-k+1}}.$$
 (32)

In our case, replacing $x_i \longrightarrow \ln^{(i)}(\rho_{Et})$, we obtain

$$B_{n,k} = \sum \frac{n!}{j_1! j_2! \dots j_{n-k+1}!} \left(\frac{\ln^{(1)}(\rho_{Et})}{1!} \right)^{j_1} \left(\frac{\ln^{(2)}(\rho_{Et})}{2!} \right)^{j_2} \dots \left(\frac{\ln^{(n-k)}(\rho_{Et})}{(n-k+1)!} \right)^{j_{n-k+1}}.$$
(33)

In addition, recalling the definition of $\beta_F := -\partial_E \ln \rho(E)$, then the general form of $B_{n,k}$:

$$B_{n,k} = \sum \frac{n!}{j_1! j_2! \dots j_{n-k+1}!} \left(\frac{(-1)\beta_F^{(0)}}{(1)!} \right)^{j_1} \left(\frac{(1)\beta_F^{(1)}}{(2)!} \right)^{j_2} \left(\frac{(-1)\beta_F^{(2)}}{(3)!} \right)^{j_3} \dots \left(\frac{(-1)^n \beta_F^{(n-1)}}{(n-k+1)!} \right)^{j_{n-k+1}}.$$
(34)

Given this definition, is clear that,

$$\langle \beta^n \rangle_{E,S} = \partial_t^n M_\beta(t) \big|_{t=0} = \sum_{k=1}^n B_{n,k}$$
(35)

is a function only of β_F and its n-1 derivatives.

Here we can see that the moments only depend on β_F and its derivatives, and moreover, that odd derivatives of β_F being negative while β_F and its even derivatives being positive ensure that all moments are non-negative.

Eq. 35 proves our lemma and the expression of $B_{n,k}$ in Eq. 33 proves that all the odd derivatives of β_F being negative implies the existence of superstatistics. Note that we fixed our function $G_n = \sum_{k=1}^n B_{n,k}$.

5. Completely monotone $\beta_F(E)$ and $\rho(E|S)$

Based on the relation of the ensemble function $\rho(E|S)$ and the fundamental inverse temperature $\beta_F(E)$ lately presented, in this section we present the completely monotone character of both functions. We will prove the following theorem.

Theorem 2. Suppose a function

$$\beta_F(E) := -\frac{\partial}{\partial E} \ln \rho(E|S) \tag{36}$$

is completely monotone, that is, $(-1)^n \beta_F^{(n)}(E) \ge 0$ for all $n = 0, 1, 2, 3, \ldots$ Then, $\rho(E|S)$ must also be a completely monotone function.

Proof. If $\beta_F(E)$ is completely monotone, then

$$(-1)^{(n)} \frac{\partial^{n}}{\partial E^{n}} \beta_{F}(E) \ge 0, \quad \forall E \ge 0$$

$$(-1)^{(n)} \frac{\partial^{n}}{\partial E^{n}} \left[-\frac{\partial}{\partial E} \ln \left(\rho(E) \right) \right] = (-1)^{(n)} \frac{\partial^{n+1}}{\partial E^{n+1}} \left[-\ln \left(\rho(E) \right) \right] \ge 0$$

$$\implies (-1)^{(n-1)} \frac{\partial^{n}}{\partial E^{n}} \left[-\ln \rho(E) \right] \ge 0 \quad (37)$$

In Eq. 37, we note that $E \mapsto -\ln \rho(E)$ is a Bernstein function. Given by the points (ii) and (iii) of Theorem 3.7, p.27 in [50], we can define a function g such that

$$g(E) = \exp(\ln\rho(E)) = \exp(-(-\ln\rho(E)))$$
 (38)

which is a completely monotone function, because $\exp(-\bullet)$ is a completely monotone function and $-\ln \rho(\bullet)$ is a Bernstein function, then the composite function

$$\exp(-\bullet) \circ - \ln \rho(\bullet)$$

is a completely monotone function.

In conclusion, we have shown that, if $\beta_F(E)$ is completely monotone, then $\rho(E|S)$ must be completely monotone, and $\rho(E|S)$ will have a superstatistical representation, because every completely monotone function is a Laplace transform of a non-negative function.

8

6. Some illustrations

Having proved both theorems, we apply this result to two different models, were one of them is characterized to be a superstatistical system, namely the q-canonical ensemble and other which is not, the Gaussian ensemble.

In the case of the Gaussian ensemble, it is characterized by the ensemble function $\rho(E; A, E_0)$ which is defined as

$$\rho(E; A, E_0) = \frac{1}{\eta_A(E_0)} \exp\left(-A(E - E_0)\right),\tag{39}$$

where A > 0. Then, we proceed to determine the moment generating function

$$M_{\beta}(t) = \rho(E - t|S) / \rho(E|S),$$

which in this case, is given by

$$M_{\beta}(t) = \frac{\exp\left(-A(E-t-E_0)^2\right)}{\exp\left(-A(E-E_0)^2\right)} = \exp\left(-A[t^2 - 2t(E-E_0)]\right).$$
(40)

Then for a general case, considering A > 0, we have that

$$\beta_F = 2A(E - E_0),$$

$$\beta'_F = 2A,$$

$$\beta''_F = 0,$$

$$\dots$$

$$\beta^n_F = 0.$$

(41)

Noting that $\beta'_F = A > 0$ and the values of the higher derivatives are zero, we see that, this model does not follow superstatistics. On the other hand, the *q*-canonical ensemble, is described by the ensemble function $\rho(E; \beta_0, q)$, where *q* is a generalized parameter defined in non-extensive Tsallis statistics, given by

$$\rho(E;\beta_0,q) = \rho(0) \left[1 - (1-q)\beta_0 E\right]^{\frac{1}{1-q}}.$$
(42)

For this case, $M_{\beta}(t) = \rho(E - t|S)/\rho(E|S)$, then we obtain

$$M_{\beta}(t) = \left[\frac{1 - (1 - q)\beta_0(E - t)}{1 - (1 - q)\beta_0 E}\right]^{\frac{1}{1 - q}} = \left[1 + \frac{1 + (1 - q)\beta_0 t}{1 - (1 - q)\beta_0 E}\right]^{\frac{1}{1 - q}}$$

After this, the first derivatives of the moment generating function at t = 0, corresponding to β_F , is

$$\partial_E M_\beta(t)|_{t=0} = \frac{\beta_0}{1 - (1 - q)\beta_0 E} = \beta_F(E).$$

In the following, we show that the first derivatives of β_F for the q-canonical ensemble. In the case of $q \ge 1$ they are given by

$$\beta_{F}^{(1)} = \beta_{F}^{2}(1-q) \leq 0,
\beta_{F}^{(2)} = 2\beta_{F}^{3}(1-q)^{2} \geq 0,
\beta_{F}^{(3)} = 6\beta_{F}^{4}(1-q)^{3} \leq 0, \qquad (43)
\beta_{F}^{(4)} = 24\beta_{F}^{5}(1-q)^{4} \geq 0,
\beta_{F}^{(5)} = 120\beta_{F}^{6}(1-q)^{5} \leq 0,$$

considering that $q \ge 1$, and we can verify that the general form of the *n*-th derivative for this case is given by

$$\beta_F^{(n)} = n! \beta_F^{n+1} (1-q)^n = n! \beta_F^{n+1} (q-1)^n (-1)^n.$$

Finally, it is straightforward to note that this model follow superstatistics only for $q \ge 1$, because $\beta_F > 0$ implies $\operatorname{sgn}(\beta_F^{(n)}) = (-1)^n$ for q > 1.

7. Concluding Remarks

In this work, we have reviewed some of the necessary conditions for the existence of a superstatistical description. More importantly, we have proved a theorem establishing a sufficient condition, namely that all the odd derivatives of β_F must be negative or zero, while β_F and its even derivatives must be positive or zero. This new result of the superstatistical theory was developed and proved using the fluctuation-dissipation theorem for the conditional distribution $P(\beta|E, S)$. We present two different illustrations of this sufficient condition: on the one hand refuting the existence of superstatistics for the Gaussian ensemble, while on the other ensuring a superstatistical representation for the q-canonical ensemble if $q \ge 1$. Another path showing the relation between the inverse temperature moments and β_F and its derivatives was presented in terms of a differential equation for the moment-generating function $M_{\beta}(t; E)$ for a given function β_F .

These results widen the domain of application of the theory of superstatistics, in such a way that they allow the exploration of complex systems and their classification by means of the presented theorem.

Acknowledgements

The authors gratefully acknowledge funding from ANID FONDECYT Regular 1220651. CF also acknowledges Beca ANID Doctorado Nacional/(2021) - 21210658.

Data Availability Statement

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

Appendix A. Differential equation for the moment generating function $M_{\beta}(t; E)$

There exists another way to relate the moments of a distribution with β_F and its derivatives. Using the fluctuation-displation theorem of Eq.18, we can rewrite it as

$$\left\langle \beta^{k+1} \right\rangle_{E,S} = \left(\beta_F - \partial_E \right) \left\langle \beta^k \right\rangle_{E,S},$$
 (A.1)

where the term $(\beta_F - \partial_E)$ is an operator. Then, multiplying Eq.A.1 by $t^k/(k!)$ and performing the summation over k, we obtain

$$\sum_{k=0}^{\infty} \frac{t^k}{k!} \left\langle \beta^{k+1} \right\rangle_{E,S} = (\beta_F - \partial_E) M_\beta(t; E), \tag{A.2}$$

and changing index such as k + 1 = m and using

$$\partial_t \left(\frac{t^m}{m!}\right) = \frac{t^{(m-1)}}{(m-1)!},$$

we obtain

$$\partial_t \left(\sum_{m=0}^{\infty} \frac{t^m}{m!} \left< \beta^m \right>_{E,S} - 1 \right) = (\beta_F - \partial_E) M_\beta(t; E).$$

Finally, we arrive at a differential equation for M_{β} , namely

$$\partial_t M_\beta(t; E) = \partial_t M_\beta(t; E) = (\beta_F - \partial_E) M_\beta(t; E).$$
(A.3)

This result confirms that β_F determines completely the function $M_{\beta}(t; E)$. We can easily verify by direct evaluation that

$$M_{\beta}(t; E) = \frac{\rho(E - t|S)}{\rho(E|S)}$$

is a solution of Eq. A.3, the left-hand side,

$$\partial_t M_\beta(t; E) = \partial_t \left(\frac{\rho(E - t|S)}{\rho(E|S)} \right) = -\frac{\rho'(E - t|S)}{\rho(E|S)},$$

while for the right-hand side, we have

$$(\beta_F - \partial_E) \left(\frac{\rho(E - t|S)}{\rho(E|S)} \right) = \beta_F \left(\frac{\rho(E - t|S)}{\rho(E|S)} \right) - \left[\frac{\rho'(E - t|S)\rho(E|S) - \rho(E - t|S)\rho'(E|S)}{\rho(E|S)^2} \right]$$
$$= -\frac{\rho'(E - t|S)}{\rho(E|S)} \quad \blacksquare$$
(A.4)

References

- C. Beck and E.G.D. Cohen. Superstatistics. Physica A: Statistical Mechanics and its Applications, 322:267–275, 2003.
- [2] C. Beck. Superstatistics: theory and applications. Continuum Mech. Thermodyn., 16:293–304, 2004.
- [3] C. Beck. Statistics of three-dimensional lagrangian turbulence. *Physical review letters*, 98(6):064502, 2007.
- [4] G. Li, W. He, and B. et al. Yang. Application of superstatistical analysis on fluctuant surface shear in particle-laden turbulence boundary layer. *Eur. Phys. J. E*, 45:5, 2022.
- [5] S. Davis, G. Avaria, B. Bora, J. Jain, J. Moreno, C. Pavez, and L. Soto. Single-particle velocity distributions of collisionless, steady-state plasmas must follow superstatistics. *Phys. Rev. E*, 100:023205, 2019.
- [6] S. Dilmi, F. Khalfaoui, and A. Boumali. The effects of superstatistics properties on hot plasma. Eng. Technol. Appl. Sci. Res., 12(5):9342–9346, 2022.
- [7] P. Chavanis. Coarse-grained distributions and superstatistics. *Physica A: Statistical Mechanics and its Applications*, 359:177–212, 2006.
- [8] A. Martínez-Merino and M. Sabido. On superstatistics and black hole quasinormal modes. *Physics Letters B*, 829:137085, 2022.
- [9] E. Gravanis, E. Akylas, and G. Livadiotis. Physical meaning of temperature in superstatistics. Europhysics Letters, 130(3):30005, jun 2020.
- [10] J. Weber, M. Reyers, and C. et al. Beck. Wind power persistence characterized by superstatistics. Sci Rep, 9:19971, 2019.
- [11] B. Schäfer, C. M. Heppell, H. I Rhys, and C. Beck. Fluctuations of water quality time series in rivers follow superstatistics. *iScience*, 24, 2021.
- [12] C. De Michele and F. Avanzi. Superstatistical distribution of daily precipitation extremes: a worldwide assessment. sci. rep. 8 (1), 1–11, 2018.
- [13] G. Cigdem Yalcin and C. Beck. Environmental superstatistics. Physica A: Statistical Mechanics and its Applications, 392(21):5431–5452, 2013.
- [14] C. Farías and S. Davis. Temperature distribution in finite systems: application to the onedimensional ising chain. The European Physical Journal B, 96:39, 2023.
- [15] J. Cheraghalizadeh, M. Seifi, Z. Ebadi, H. Mohammadzadeh, and MN. Najafi. Superstatistical two-temperature ising model. *Physical Review E*, 103(3):032104, 2021.
- [16] L. Rydin Gorjão, B. Schäfer, D. Witthaut, and C. Beck. Spatio-temporal complexity of power-grid frequency fluctuations. New Journal of Physics, 23(7):073016, jul 2021.
- [17] B. Schäfer, C. Beck, and K. et al. Aihara. Non-gaussian power grid frequency fluctuations characterized by lévy-stable laws and superstatistics. *Nat Energy*, 3:119–126, 2018.
- [18] S. Davis. A classification of nonequilibrium steady states based on temperature correlations. *Physica A: Statistical Mechanics and its Applications*, 608:128249, 2022.
- [19] N. Cressie, A. S. Davis, J. L. Folks, and G. E. Policello. The moment-generating function and negative integer moments. *The American Statistician*, 35(3):148–150, 1981.
- [20] N. Cressie and M. Borkent. The moment generating function has its moments. Journal of Statistical Planning and Inference, 13:337–344, 1986.
- [21] A. Mukherjea, M. Rao, and S. Suen. A note on moment generating functions. Statistics & Probability Letters, 76(11):1185–1189, 2006.
- [22] N.G. Ushakov and V.G. Ushakov. On convergence of moment generating functions. Statistics & Probability Letters, 81(4):502–505, 2011.
- [23] C. Stone. On moment generating functions and renewal theory. The Annals of Mathematical Statistics, 36(4):1298–1301, 1965.
- [24] W. Kozakiewicz. On the convergence of sequences of moment generating functions. The Annals of Mathematical Statistics, 18(1):61–69, 1947.

- [25] E. T. Jaynes. "Probability theory: The Logic Of Science". Cambridge University Press, 2003.
- [26] C. Tsallis. Possible generalization of Boltzmann-Gibbs statistics. Journal of Statistical Physics, 52:479–487, 1988.
- [27] S. Davis and G. Gutiérrez. Temperature is not an observable in superstatistics. Physica A: Statistical Mechanics and its Applications, 505:864–870, 2018.
- [28] S. Davis and G. Gutiérrez. Emergence of Tsallis statistics as a consequence of invariance. *Physica A*, 533:122031, 2019.
- [29] U. A. Salian. Microcanonical temperature and "heat capacity" computation of Lennard-Jones clusters under isoergic molecular dynamics simulation. J. Chem. Phys., 108:6342–6346, 1998.
- [30] M. Pleimling and H. Behringer. Microcanonical analysis of small systems. *Phase Transitions*, 78:787–797, 2005.
- [31] H. H. Rugh. Dynamical approach to temperature. Phys. Rev. Lett., 78(5):772 774, 1997.
- [32] G. Rickayzen and J. G. Powles. Temperature in the classical microcanonical ensemble. J. Chem. Phys., 114(9):4333 – 4334, 2001.
- [33] G. P. Morriss and L. Rondoni. Definition of temperature in equilibrium and nonequilibrium systems. *Phys. Rev. E*, 59:R5–R8, Jan 1999.
- [34] G. Palma, G. Gutiérrez, and S. Davis. Ensemble-free configurational temperature for spin systems. *Phys. Rev. E*, 94:062113, 2016.
- [35] P. Buonsante, R. Franzosi, and A. Smerzi. Phase transitions at high energy vindicate negative microcanonical temperature. *Phys. Rev. E*, 95:052135, 2017.
- [36] G. Gutiérrez, S. Davis, and G. Palma. Configurational temperature in constrained systems: the case of spin dynamics. *Journal of Physics A: Mathematical and Theoretical*, 51(45):455003, 2018.
- [37] S. Davis. On the possible distributions of temperature in nonequilibrium steady states. Journal of Physics A: Mathematical and Theoretical, 53:045004, 2020.
- [38] S. Davis. Fluctuating temperatures outside superstatistics: Thermodynamics of small systems. Physica A: Statistical Mechanics and its Applications, 589:126665, 2022.
- [39] R. Kubo. Generalized cumulant expansion method. Journal of the Physical Society of Japan, 17(7):1100–1120, 1962.
- [40] K. Kiyono and H. Konno. Log-amplitude statistics for beck-cohen superstatistics. Phys. Rev. E, 87:052104, May 2013.
- [41] A. Saboor, S. B. Provost, and M. Ahmad. The moment generating function of a bivariate gammatype distribution. Applied Mathematics and Computation, 218(24):11911–11921, 2012.
- [42] X. I. A. Yang, I Marusic, and C Meneveau. Moment generating functions and scaling laws in the inertial layer of turbulent wall-bounded flows. *Journal of Fluid Mechanics*, 791:R2, 2016.
- [43] E. R. Villa and L. A. Escobar. Using moment generating functions to derive mixture distributions. The American Statistician, 60(1):75–80, 2006.
- [44] G. Grimmet and D. Welsh. Probability: An Introduction. Oxford University Press, 2014.
- [45] L. Wasserman. All of Statistics: A Concise Course in Statistical Inference. Springer New York, NY, 2004.
- [46] D. Hainaut. Moment generating function of non-Markov self-excited claims processes. Insurance: Mathematics and Economics, 101:406–424, 2021.
- [47] S. Ghosh. Normality testing for a long-memory sequence using the empirical moment generating function. Journal of Statistical Planning and Inference, 143(5):944–954, 2013.
- [48] J. H. Curtiss. A note on the theory of moment generating functions. The Annals of Mathematical Statistics, 13(4):430 – 433, 1942.
- [49] E. T. Bell. Partition polynomials. Annals of Mathematics, 29(1/4):38–46, 1927.
- [50] R. L. Schilling, R. Song, and Z. Vondracek. Bernstein Functions. De Gruyter, Berlin, New York, 2010.